

Density-Induced Hadron-Quark Crossover via the Formation of Cooper Triples

Hiroyuki Tajima,¹ Shoichiro Tsutsui,^{2,3} Takahiro M. Doi,^{4,3} and Kei Iida⁵

¹*Department of Physics, Graduate School of Science,
The University of Tokyo, Tokyo 113-0033, Japan*

²*QunaSys Inc., Aqua Hakusan Building 9F, 1-13-7 Hakusan, Bunkyo, Tokyo 113-0001, Japan*

³*RIKEN iTHEMS, Wako, Saitama 351-0198, Japan*

⁴*Research Center for Nuclear Physics (RCNP), Osaka University, 567-0047, Japan*

⁵*Department of Mathematics and Physics, Kochi University, Kochi 780-8520, Japan*

(Dated: November 28, 2022)

We discuss the hadron-quark crossover accompanied by the formation of Cooper triples (three-body counterpart of Cooper pairs), by analogy with the Bose-Einstein condensate to Bardeen-Cooper-Schrieffer crossover in two-component fermionic systems. Such a crossover is different from a phase transition, which often involves symmetry breaking. We calculate the in-medium three-body energy from the three-body T -matrix with a phenomenological three-body force characterizing a bound hadronic state in vacuum. With increasing density, the hadronic bound-state pole smoothly undergoes a crossover toward the Cooper triple phase where the in-medium three-body clusters coexist with the quark Fermi sea.

I. INTRODUCTION

After the discovery of massive neutron stars and dramatic development of astrophysical observatories, the properties of dense matter have attracted huge attention [1]. In particular, the equation of state for extremely dense matter has been determined from the astrophysical observations such as X -ray and gravitational-wave measurements [2, 3] and is regarded as an important testing platform of many-body nuclear theories [4].

Along this direction, it may well be an exciting and challenging problem to clarify how nuclear matter changes into possible quark matter in the central core region of massive compact stars. Such a high-density regime is out of the range accessible by the lattice quantum chromodynamics (QCD) simulation due to the severe sign problem [5], and there are various predictions such as first-order phase transition and appearance of quarkyonic matter [6]. A fascinating idea to resolve this problem is a crossover scenario called "hadron-quark crossover" [7], where hadrons consisting of three quarks, i.e., baryons, gradually change into deconfined quarks without phase transitions along the evolution of the chemical potential (or the baryon density). Interestingly, it is reported that such a crossover can be probed via future observations of the gravitational waves emitted from binary neutron-star mergers [8]. In particular, the nuclear equation of state constrained from the recent simultaneous measurements of neutron star masses and radii implies a non-monotonic behavior of the speed of sound in the high-density regime [9].

Recently, the existence of the peak in the density dependence of the speed of sound has been discussed in the context of hadron-quark crossover [10] as well as quarkyonic matter [11]. However, the elucidation of the microscopic crossover mechanism leading to the peak is still challenging because of the sign problem in the finite-density lattice QCD [5]. In this regard, the peak of the speed of sound in two-color dense QCD, for which sign problems are avoidable even at finite density in the lattice Monte Carlo simulations, has been discussed by analogy with the Bose-Einstein condensate (BEC) to Bardeen-Cooper-Schrieffer (BCS) crossover [12]. Moreover, the latest results of the lattice simulations also indicate the existence of the peak of the speed of sound in the crossover regime [13, 14]. A similar strong-coupling crossover phenomenon has been extensively studied in cold atomic systems [15–19], where the molecular BEC continuously changes to the BCS-type Cooper-pair condensate in two-component Fermi gases. The possible occurrence of the peak of the speed of sound has also been discussed in the density-induced BEC-BCS crossover with the finite-range interaction [20]. We note that the carrier-density-induced BEC-BCS crossover has been realized in strongly-correlated electron systems recently [21–24].

Going back to the three-color QCD at finite density, we have to consider in-medium three-body correlations unavoidably and seriously because of the formation of hadrons consisting of three quarks [25], just like the case of the BEC-BCS crossover in which one has to consider in-medium two-body correlations. Such a three-quark bound state accompanies famous confinement effects associated with the $SU(3)$ gauge invariance [26, 27]. While the three-body correlations in finite-density fermionic systems are still elusive, several ideas appear in the context of cold atomic physics as a quantum simulation of matter in extreme conditions. Indeed, although there are several difficulties such as atomic losses, three-component ultracold Fermi gases (mimicking three-color quarks) can be prepared experimentally by using the different three hyperfine states [28]. Also, a similarity between the three-body crossover in three-component Fermi gases and the hadron-quark continuity in the three-color QCD has been pointed out theoretically [29, 30]. To understand a non-trivial state of matter in the Fermi-degenerate state with three-body correlations,




FIG. 1. Schematics of the density-induced two- and three-body crossovers. While the gases of tightly bound states (i.e., dimers and trimers) exhibit smaller cluster size ℓ compared to the mean interparticle distance $d \sim k_F^{-1}$ in the low-density regime, these bound states start to be overlapped with each other ($d \lesssim \ell$) with increasing density (i.e., decreasing d) and gradually change into Cooper pairs and triples. In ultracold Fermi gases near the Feshbach resonance, such an in-medium two-body crossover is triggered by the change of the attractive interaction, where ℓ increases when the attractive interaction decreases.

one may borrow the knowledge of the BCS-BEC-crossover physics well established in cold-atom communities. In this regard, just as Cooper pairs are formed in the BCS side of the density-induced BEC-BCS crossover [31], one may imagine its three-body counterpart, that is, a Cooper triple [32–37], in dense Fermi-degenerate regime but with non-negligible three-body correlations.

Figure 1 shows a comparison between the two-body crossover (i.e., BEC-BCS crossover) and the three-body crossover accompanied by Cooper triples when the number density increases or the attractive interaction decreases. First, let us review the famous BEC-BCS-crossover case; if the density increases, the mean interparticle distance given by $d \sim k_F^{-1}$ decreases where k_F is the Fermi momentum. In the dilute regime, d is much longer than the cluster size ℓ which is determined by the interaction (e.g., scattering length) [19]. Such a state is well described by the gas of two-body molecules and undergoes the BEC at low temperature. At sufficiently high density, d becomes smaller than the cluster size ℓ , indicating the tremendous overlap among clusters, that is, the formation of Cooper pairs. These Cooper pairs are repeatedly formed and dissociated by coexisting with the underlying Fermi sphere, although we do not draw such a dynamical picture explicitly in Fig. 1. While Cooper pairs form the BCS condensate, which spontaneously breaks U(1) gauge symmetry, in conventional superconductors, we note that the spontaneous symmetry breaking is not necessary to form Cooper pairs themselves as the so-called preformed Cooper pairs are found in strongly-correlated or disordered superconductors above the critical temperature [38, 39].

Then, let us move to the three-body case as shown in the lower side of Fig. 1. One would see tightly bound three-body states with the cluster size ℓ if the strong two- or three-body attractive interactions exist in three-color fermionic systems. When the number density increases, these trimer states start to overlap at certain densities and eventually may form an unconventional state that can be analogous to the Cooper-pairing state. In the case of QCD, we recall again that such a three-body state should be kept due to the SU(3) gauge invariance. These overlapped three-body states in medium are similar to Cooper triples, which are the three-body counterpart of Cooper pairs predicted in the context of cold atomic physics [32–35, 37]. The Cooper triple state does not necessarily form a condensate (however note Ref. [40]) but resulting three-body correlations can modify the equation of state substantially [35, 41].

In this paper, motivated by the recent astronomical observations of compact stars as well as the theoretical progress in the in-medium three-body correlations in cold atomic systems, we discuss the possible scenario of the density-induced hadron-quark crossover accompanied by the formation of Cooper triples by using a simplified toy model with three-quark attractive interaction responsible for the baryon formation. We show that the in-medium three-body energy obtained from the three-body T -matrix exhibits a unique crossover from baryonic tightly-bound states to Cooper triples with increasing quark chemical potential.

This paper is organized as follows. In Sec. II, we introduce the theoretical model and explain how to address the crossover via the three-body T -matrix approach [35]. In Sec. III, we show the numerical result for the in-medium three-body energy and discuss the density-induced hadron-quark crossover. We summarize this paper in Sec. IV. Throughout this paper, we take $\hbar = k_B = 1$ and the system volume is set to unity.

II. MODEL AND METHODS

In a way similar to the study of quark-gluon plasmas in Ref. [42], we start from the Hamiltonian based on the Thompson scheme [43] as given by

$$H = \sum_{\mathbf{p}} \sum_j \varepsilon_{\mathbf{p}} \psi_{\mathbf{p},j}^\dagger \psi_{\mathbf{p},j} + \sum_{\mathbf{k},\mathbf{q},\mathbf{k}',\mathbf{q}',\mathbf{P}} V_{\mathbf{k},\mathbf{q},\mathbf{k}',\mathbf{q}',\mathbf{P}} \psi_{\mathbf{k},r}^\dagger \psi_{\mathbf{q},g}^\dagger \psi_{\mathbf{P}-\mathbf{k}-\mathbf{q},b}^\dagger \psi_{\mathbf{P}-\mathbf{k}'-\mathbf{q}',b} \psi_{\mathbf{q}',g} \psi_{\mathbf{k}',r}, \quad (1)$$

where $\psi_{\mathbf{p},j}^{(\dagger)}$ is the fermionic annihilation (creation) operator with three colors ($j = r, g, b$), and $\varepsilon_{\mathbf{p}}$ is the relativistic dispersion $\varepsilon_{\mathbf{p}} = \sqrt{p^2 + m_q^2}$ with the nonzero constituent quark mass $m_q \simeq 0.34$ GeV. For simplicity, we neglect the spin and flavor degrees of freedom, which are not important for our purpose and will be considered in the future work. The second term of Eq. (1) is the three-body attractive interaction responsible for the color-singlet baryon formation. While the coupling constant should be momentum-dependent due to the color confinement [44–46] as well as relativistic effects (the projection operator for the positive energy states and the coupling with negative-energy states) [42, 43], we employ the constant coupling $V_{\mathbf{k},\mathbf{q},\mathbf{k}',\mathbf{q}',\mathbf{P}} \simeq V_3$ for the purpose of qualitatively understanding the hadron-quark crossover accompanied by the Cooper triple formation in the relatively high-density regime. Instead, we will adjust the value of V_3 later in such a way as to roughly reproduce the baryon mass in vacuum M_B as $\simeq 0.91$ GeV. Incidentally, we note that non-relativistic or relativistic quark models with three-body forces have been introduced in Refs. [47–49]. We obtain the in-medium three-body T -matrix at the zero center-of-mass momentum [35],

$$T_3(\Omega) = \frac{V_3}{1 - V_3 \Xi(\Omega)}, \quad (2)$$

where $\Omega_+ = \Omega + i\delta$ is the three-body energy with a small imaginary part $i\delta$. The in-medium three-body propagator $\Xi(\Omega)$ reads [50]

$$\Xi(\Omega) = \sum_{\mathbf{k},\mathbf{q}} \frac{(1 - f_{\mathbf{k}})(1 - f_{\mathbf{q}})(1 - f_{\mathbf{k}+\mathbf{q}}) + f_{\mathbf{k}} f_{\mathbf{q}} f_{\mathbf{k}+\mathbf{q}}}{\Omega_+ + 3\mu - \varepsilon_{\mathbf{k}} - \varepsilon_{\mathbf{q}} - \varepsilon_{\mathbf{k}+\mathbf{q}}}, \quad (3)$$

where

$$f_{\mathbf{k}} = \frac{1}{\exp\left(\frac{\varepsilon_{\mathbf{k}} - \mu}{T}\right) + 1} \quad (4)$$

is the Fermi distribution function. We note that while the antiquarks are ignored in our toy model, such antiparticle contributions involve the antiparticle distribution function $\tilde{f}_{\mathbf{k}} = \frac{1}{\exp\left(\frac{\varepsilon_{\mathbf{k}} + \mu}{T}\right) + 1}$, which can be small at large chemical potential and low temperature.

We set $V_3 = 2.16 \times 10^2$ (GeV)⁻⁵ such that $T_3(\Omega)$ has a pole at $\Omega \simeq M_B$ in vacuum. The cutoff and the small imaginary energy for the analytic continuation are adapted as $\Lambda = 2$ GeV and $\delta = 2 \times 10^{-2}$ GeV. We note that the continuum bottom of three quarks is given by

$$E_{\text{th}} = 3m_q - 3\mu. \quad (5)$$

In this regard, the in-medium three-body binding energy E_B^M (i.e., energy gain due to the formation of the in-medium three-body cluster) can be defined as

$$\frac{1}{V_3} - \Xi(\Omega_{\text{pole}} = E_{\text{th}} - E_B^M) = 0, \quad (6)$$

which is the pole of the in-medium three-body T -matrix. In the vacuum case, such an energy gain is given by $E_B = 3m_q - M_B \simeq 0.11$ GeV.




FIG. 2. The in-medium three-body energy in the hadron-quark crossover regime at different temperatures. For reference, we plot $\Omega - E_{\text{th}} = 3\mu - 3m_q$, that is, $\Omega = 0$ (at the Fermi level). The lower panel presents the schematic comparison between the single-particle energy $\varepsilon_k = \sqrt{k^2 + m_q^2}$ and the three-body energy per particle $(\Omega_{\text{pole}} - E_{\text{th}})/3$ (dashed line) along the hadron-quark crossover.

III. RESULTS AND DISCUSSIONS

Figure 2 shows the in-medium three-body energy pole Ω_{pole} measured from the bottom of the continuum $E_{\text{th}} = 3m_q - 3\mu$ as a function of μ at different temperatures T . One can see that the the three-body energy located in $\Omega_{\text{pole}} - E_{\text{th}} \simeq M_B - 3m_q \equiv -E_B$ (i.e., $E_B^M \simeq E_B$) at $\mu = 0$ moves toward the continuum bottom $\Omega_{\text{pole}} - E_{\text{th}} = 0$ when μ increases. Moreover, this three-body pole remains at larger μ and locates below the Fermi level given by $\Omega - E_{\text{th}} = 3\mu - 3m_q$, indicating the coexistence of the quark Fermi sea and in-medium three-body clusters. This result manifests a picture of the smooth crossover from the tightly bound baryonic trimer state to the Cooper triple state, being analogous to the density-induced BEC-BCS crossover [18, 31, 51]. While the existence of the positive pole in the continuum is in contrast to the non-relativistic one-dimensional case [35], the similar behavior of the three-body energy can be found in the variational approach to the non-relativistic Fermi gases in three dimensions [34]. At higher temperatures, Ω_{pole} is shifted toward an upper energy level around the crossover because of thermal agitation effect on the binding energy [35, 52]. However, one can see that the three-body pole is still located below the Fermi level ($\Omega - E_{\text{th}} < -E_{\text{th}}$, i.e., $\Omega < 0$) at low temperatures, implying the presence of the stable three-body bound state in medium.

To see the physical picture of this crossover from a different point of view, we show the schematic single-particle energy level diagram in the lower panel of Fig. 2. At $\mu = 0$, the three-body bound state is located in the energy level $-E_B/3$ measured from the continuum bottom $\varepsilon_{k=0} = m_q$. At finite μ , this three-body bound state is shifted toward an upper energy level because of the reduced binding energy associated with the Pauli-blocking effect. However, such a Fermi-surface effect does not suppress the three-body pole but leads eventually to the in-medium bound state in continuum at high density as in the case of the famous Cooper pairing in superconductors. This state can be regarded as the Cooper triple phase, but we note that there are no clear boundaries between the hadronic bound-trimer phase and Cooper triple phase, a feature similar to the BEC-BCS crossover [34, 35].

While we have fixed $m_q = 0.34$ GeV in Fig. 2, it may have a density dependence through the quark-antiquark pairing [53]. In Fig. 3, we show a comparison between the results with massive ($m_q = 0.34$ GeV) and massless ($m_q = 0$) quarks; in the latter case, the chiral symmetry is assumed to be restored due to the Pauli blocking effect




FIG. 3. Comparison of the in-medium three-body energy with $m_q = 0.34$ GeV and $m_q = 0$.

on the quark-antiquark pairing. While the result with $m_q = 0$ (dashed curve) is found to be larger than that with $m_q = 0.34$ GeV (solid curve), this difference can be approximately given by $3m_q$ with $m_q = 0.34$ GeV. On the other hand, even the massless case is well below the corresponding Fermi level (3μ in the absence of m_q) at high density, where the Cooper triple phase is expected to occur. For more realistic case, m_q changes from the in-vacuum value down to the current quark mass across the evolution of μ , leading to the in-medium three-body pole in between.

In the high-density regime, the three-body pole measured from the Fermi level (i.e., $3\mu - 3m_q - (\Omega_{\text{pole}} - E_{\text{th}}) \equiv -\Omega_{\text{pole}}$) can be regarded as the energy gain of the formation of a Cooper triple. The ratio of this energy gain to the Fermi energy may well become smaller as the density increases. However, the non-local properties of the three-body force, if remaining, would be important in such a high-density regime. Indeed, if the three-body force is replaced by the linear potential $V_{qqq}(R) \simeq \sigma R$ [44–46] where σ is the string tension and R is the characteristic length scale for the three-body state, the energy gain might be corrected due to a significant spatial overlap of Cooper triples. For example, considering the density-dependent interaction given by $V_3 = V_{qqq}(R \simeq d)$ where $d \simeq k_F^{-1}$ is the mean interquark distance, one finds $V_3 \simeq \sigma/\mu$ at $m_q \rightarrow 0$. Combined effects of the density dependence of the dynamical quark mass and of the linear-shaped three-quark interaction would cause the three-body energy pole to increase more rapidly. It will be left for an interesting future work. We remark in passing that at asymptotically high density, the confining three-body force would be suppressed, while two-body interactions due to one-gluon exchange would take over. Even in the density regime of interest here, instanton-induced two-body interactions would take effect. Such two-body interactions would lead to quark Cooper pairs and hence color superconductivity [54], which are ignored in the present analysis.

IV. CONCLUSIONS

In this paper, we have discussed the density-induced hadron-quark crossover via the formation of Cooper triples by analogy with BEC-BCS crossover realized in cold-atomic and in condensed-matter systems. Applying the in-medium three-body T -matrix approach to the toy model where relativistic quarks interact with each other via the phenomenological three-body forces leading to the baryonic three-body bound state in vacuum, we show that the in-medium three-body pole exhibits a smooth crossover from the state of baryonic three-body trimers to the state of Cooper triples with increasing chemical potential, which can be regarded as the three-body counterpart of the density-induced BEC-BCS crossover.

For future perspectives, it is interesting to examine the equation of state and the speed of sound in a more realistic model including e.g., density dependence of the dynamical quark mass, effects of gluons, and competition with diquark formation and color superconductivity [54]. In the presence of additional degrees of freedom such as spin and flavor, dibaryonic superconducting quark matter might also appear at low temperature [55, 56]. For ab initio calculations to elucidate the hadron-quark crossover, the complex Langevin method would be one of the promising methods to address finite-density QCD by avoiding the sign problem as demonstrated in Ref. [57, 58]. The complex Langevin method can also be used for clarifying multi-body correlations in ultracold Fermi gases [59].

ACKNOWLEDGMENTS

The authors thank E. Nakano for useful discussions. This research was funded by Grants-in-Aid for Scientific Research provided by JSPS through No. 18H05406, No. 20K14480, No. 22H01158, and No. 22K13981.

-
- [1] G. Baym, T. Hatsuda, T. Kojo, P. D. Powell, Y. Song, and T. Takatsuka, *Reports on Progress in Physics* **81**, 056902 (2018).
- [2] A. L. Watts, N. Andersson, D. Chakrabarty, M. Feroci, K. Hebeler, G. Israel, F. K. Lamb, M. C. Miller, S. Morsink, F. Özel, A. Patrino, J. Poutanen, D. Psaltis, A. Schwenk, A. W. Steiner, L. Stella, L. Tolos, and M. van der Klis, *Rev. Mod. Phys.* **88**, 021001 (2016).
- [3] L. Baiotti, *Progress in Particle and Nuclear Physics* **109**, 103714 (2019).
- [4] J. M. Lattimer, *Annual Review of Nuclear and Particle Science* **62**, 485 (2012).
- [5] K. Nagata, *Progress in Particle and Nuclear Physics*, 103991 (2022).
- [6] K. Fukushima and T. Hatsuda, *Reports on Progress in Physics* **74**, 014001 (2010).
- [7] K. Masuda, T. Hatsuda, and T. Takatsuka, *The Astrophysical Journal* **764**, 12 (2013).
- [8] Y.-J. Huang, L. Baiotti, T. Kojo, K. Takami, H. Sotani, H. Togashi, T. Hatsuda, S. Nagataki, and Y.-Z. Fan, *Phys. Rev. Lett.* **129**, 181101 (2022).
- [9] T. Kojo, G. Baym, and T. Hatsuda, *The Astrophysical Journal* **934**, 46 (2022).
- [10] T. Kojo, *AAPPS Bulletin* **31**, 1 (2021).
- [11] L. McLerran and S. Reddy, *Phys. Rev. Lett.* **122**, 122701 (2019).
- [12] T. Kojo and D. Suenaga, *Phys. Rev. D* **105**, 076001 (2022).
- [13] K. Iida, E. Itou, and T.-G. Lee, *Journal of High Energy Physics* **2020**, 1 (2020).
- [14] K. Iida and E. Itou, *Progress of Theoretical and Experimental Physics* 10.1093/ptep/ptac137 (2022).
- [15] Q. Chen, J. Stajic, S. Tan, and K. Levin, *Physics Reports* **412**, 1 (2005).
- [16] W. Zwerger, *The BCS-BEC crossover and the unitary Fermi gas*, Vol. 836 (Springer Science & Business Media, 2011).
- [17] M. Randeria and E. Taylor, *Annu. Rev. Condens. Matter Phys.* **5**, 209 (2014).
- [18] G. C. Strinati, P. Pieri, G. Röpke, P. Schuck, and M. Urban, *Physics Reports* **738**, 1 (2018), the BCS–BEC crossover: From ultra-cold Fermi gases to nuclear systems.
- [19] Y. Ohashi, H. Tajima, and P. van Wyk, *Progress in Particle and Nuclear Physics* **111**, 103739 (2020).
- [20] H. Tajima and H. Liang, *Phys. Rev. A* **106**, 043308 (2022).
- [21] S. Kasahara, T. Watashige, T. Hanaguri, Y. Kohsaka, T. Yamashita, Y. Shimoyama, Y. Mizukami, R. Endo, H. Ikeda, K. Aoyama, T. Terashima, S. Uji, T. Wolf, H. von Löhneysen, T. Shibauchi, and Y. Matsuda, *Proceedings of the National Academy of Sciences* **111**, 16309 (2014).
- [22] T. Hashimoto, Y. Ota, A. Tsuzuki, T. Nagashima, A. Fukushima, S. Kasahara, Y. Matsuda, K. Matsuura, Y. Mizukami, T. Shibauchi, *et al.*, *Sci. Adv.* **6**, eabb9052 (2020).
- [23] Y. Nakagawa, Y. Kasahara, T. Nomoto, R. Arita, T. Nojima, and Y. Iwasa, *Science* **372**, 190 (2021).
- [24] Y. Suzuki, K. Wakamatsu, J. Ibuka, H. Oike, T. Fujii, K. Miyagawa, H. Taniguchi, and K. Kanoda, *Phys. Rev. X* **12**, 011016 (2022).
- [25] S. Pittel, J. Engel, J. Dukelsky, and P. Ring, *Physics Letters B* **247**, 185 (1990).
- [26] K. Yagi, T. Hatsuda, and Y. Miake, *Quark-gluon plasma: From big bang to little bang*, Vol. 23 (Cambridge University Press, 2005).
- [27] W. Greiner, S. Schramm, and E. Stein, *Quantum chromodynamics* (Springer Science & Business Media, 2007).
- [28] K. M. O’Hara, *New Journal of Physics* **13**, 065011 (2011).
- [29] Y. Nishida, *Phys. Rev. Lett.* **109**, 240401 (2012).
- [30] Y. Nishida, *Phys. Rev. Lett.* **114**, 115302 (2015).
- [31] N. Andrenacci, A. Perali, P. Pieri, and G. C. Strinati, *Phys. Rev. B* **60**, 12410 (1999).
- [32] P. Niemann and H.-W. Hammer, *Phys. Rev. A* **86**, 013628 (2012).
- [33] T. Kirk and M. M. Parish, *Phys. Rev. A* **96**, 053614 (2017).
- [34] H. Tajima, S. Tsutsui, T. M. Doi, and K. Iida, *Phys. Rev. A* **104**, 053328 (2021).
- [35] H. Tajima, S. Tsutsui, T. M. Doi, and K. Iida, *Phys. Rev. Research* **4**, L012021 (2022).
- [36] Y. Guo and H. Tajima, *Phys. Rev. A* **106**, 043310 (2022).
- [37] Y. Guo and H. Tajima, *arXiv preprint arXiv:2210.07042* (2022).
- [38] B. L. Kang, M. Z. Shi, S. J. Li, H. H. Wang, Q. Zhang, D. Zhao, J. Li, D. W. Song, L. X. Zheng, L. P. Nie, T. Wu, and X. H. Chen, *Phys. Rev. Lett.* **125**, 097003 (2020).
- [39] K. M. Bastiaans, D. Chatzopoulos, J.-F. Ge, D. Cho, W. O. Tromp, J. M. van Ruitenbeek, M. H. Fischer, P. J. de Visser, D. J. Thoen, E. F. Driessen, *et al.*, *Science* **374**, 608 (2021).
- [40] S. Akagami, H. Tajima, and K. Iida, *Phys. Rev. A* **104**, L041302 (2021).
- [41] J. R. McKenney, A. Jose, and J. E. Drut, *Phys. Rev. A* **102**, 023313 (2020).
- [42] S. Y. F. Liu and R. Rapp, *Phys. Rev. C* **97**, 034918 (2018).
- [43] R. H. Thompson, *Phys. Rev. D* **1**, 110 (1970).

- [44] S. Capstick and N. Isgur, Phys. Rev. D **34**, 2809 (1986).
- [45] T. T. Takahashi, H. Matsufuru, Y. Nemoto, and H. Suganuma, Phys. Rev. Lett. **86**, 18 (2001).
- [46] T. T. Takahashi, H. Suganuma, Y. Nemoto, and H. Matsufuru, Phys. Rev. D **65**, 114509 (2002).
- [47] J. Richard and P. Taxil, Annals of Physics **150**, 267 (1983).
- [48] J. Carlson, J. Kogut, and V. R. Pandharipande, Phys. Rev. D **27**, 233 (1983).
- [49] W. Blask, U. Bohn, M. Huber, B. C. Metsch, and H. Petry, Zeitschrift für Physik A Atomic Nuclei **337**, 327 (1990).
- [50] T. Sogo, G. Röpke, and P. Schuck, Phys. Rev. C **81**, 064310 (2010).
- [51] M. Jin, M. Urban, and P. Schuck, Phys. Rev. C **82**, 024911 (2010).
- [52] H. Tajima and P. Naidon, New Journal of Physics **21**, 073051 (2019).
- [53] B. Holdom, J. Ren, and C. Zhang, Phys. Rev. Lett. **120**, 222001 (2018).
- [54] M. G. Alford, A. Schmitt, K. Rajagopal, and T. Schäfer, Rev. Mod. Phys. **80**, 1455 (2008).
- [55] B. C. Barrois, Nuclear Physics B **129**, 390 (1977).
- [56] M. Shahrabaf, D. Blaschke, S. Typel, G. R. Farrar, and D. E. Alvarez-Castillo, Phys. Rev. D **105**, 103005 (2022).
- [57] K. Nagata, J. Nishimura, and S. Shimasaki, Phys. Rev. D **98**, 114513 (2018).
- [58] Y. Ito, H. Matsufuru, Y. Namekawa, J. Nishimura, S. Shimasaki, A. Tsuchiya, and S. Tsutsui, Journal of High Energy Physics **2020**, 1 (2020).
- [59] C. Berger, L. Rammelmüller, A. Loheac, F. Ehmman, J. Braun, and J. Drut, Physics Reports **892**, 1 (2021).