

A CONNECTION BETWEEN THE MONOGENICITY OF CERTAIN POWER-COMPOSITIONAL TRINOMIALS AND k -WALL-SUN-SUN PRIMES

LENNY JONES

ABSTRACT. We say that a monic polynomial $f(x) \in \mathbb{Z}[x]$ of degree N is *monogenic* if $f(x)$ is irreducible over \mathbb{Q} and

$$\{1, \theta, \theta^2, \dots, \theta^{N-1}\}$$

is a basis for the ring of integers of $\mathbb{Q}(\theta)$, where $f(\theta) = 0$.

Let k be a positive integer, and let $U_n := U_n(k, -1)$ be the Lucas sequence $\{U_n\}_{n \geq 0}$ of the first kind defined by

$$U_0 = 0, \quad U_1 = 1 \quad \text{and} \quad U_n = kU_{n-1} + U_{n-2} \quad \text{for } n \geq 2.$$

A k -Wall-Sun-Sun prime is a prime p such that

$$U_{\pi_k(p)} \equiv 0 \pmod{p^2},$$

where $\pi_k(p)$ is the length of the period of $\{U_n\}_{n \geq 0}$ modulo p .

Let $\mathcal{D} = k^2 + 4$ if $k \equiv 1 \pmod{2}$, and $\mathcal{D} = (k/2)^2 + 1$ if $k \equiv 0 \pmod{2}$. Suppose that $k \not\equiv 0 \pmod{4}$ and \mathcal{D} is squarefree, and let h denote the class number of $\mathbb{Q}(\sqrt{\mathcal{D}})$. Let $s \geq 1$ be an integer such that, for every odd prime divisor p of s , \mathcal{D} is not a square modulo p and $\gcd(p, h\mathcal{D}) = 1$. In this article, we prove that $x^{2s^n} - kx^{s^n} - 1$ is monogenic for all integers $n \geq 1$ if and only if no prime divisor of s is a k -Wall-Sun-Sun prime.

1. INTRODUCTION

For a positive integer k , we let $U_n := U_n(k, -1)$ denote the n th term of the Lucas sequence $\{U_n\}_{n \geq 0}$ of the first kind defined by

$$(1.1) \quad U_0 = 0, \quad U_1 = 1 \quad \text{and} \quad U_n = kU_{n-1} + U_{n-2} \quad \text{for } n \geq 2.$$

The sequence $\{U_n\}_{n \geq 0}$ is periodic modulo any prime p , and we let $\pi_k(p)$ denote the length of the period of $\{U_n\}_{n \geq 0}$ modulo p .

A k -Wall-Sun-Sun prime [19] is a prime p such that

$$(1.2) \quad U_{\pi_k(p)} \equiv 0 \pmod{p^2}.$$

When $k = 1$, the sequence $\{U_n\}_{n \geq 0}$ is the well-known Fibonacci sequence, and the k -Wall-Sun-Sun primes in this case are also known as *Fibonacci-Wieferich* primes [18], or simply *Wall-Sun-Sun* primes [4, 19]. However, at the time this article was written, no such primes were known to exist. The existence of Wall-Sun-Sun primes was first investigated by D. D. Wall [16] in 1960, and subsequently studied by the Sun brothers [14], who showed a connection with Fermat's Last Theorem.

Date: November 29, 2022.

2020 Mathematics Subject Classification. Primary 11R04, 11B39, Secondary 11R09, 12F05.

Key words and phrases. k -Wall-Sun-Sun prime, monogenic, power-compositional.

Throughout this article, we let $\Delta(f)$ and $\Delta(K)$ denote, respectively, the discriminants over \mathbb{Q} of $f(x) \in \mathbb{Z}[x]$ and a number field K . We define $f(x) \in \mathbb{Z}[x]$ to be *monogenic* if $f(x)$ is monic, irreducible over \mathbb{Q} and $\{1, \theta, \theta^2, \dots, \theta^{\deg(f)-1}\}$ is a basis for the ring of integers \mathbb{Z}_K of $K = \mathbb{Q}(\theta)$, where $f(\theta) = 0$. If $f(x)$ is irreducible over \mathbb{Q} with $f(\theta) = 0$, then [3]

$$(1.3) \quad \Delta(f) = [\mathbb{Z}_K : \mathbb{Z}[\theta]]^2 \Delta(K).$$

Observe then, from (1.3), that $f(x)$ is monogenic if and only if $\Delta(f) = \Delta(K)$. Thus, if $\Delta(f)$ is squarefree, then $f(x)$ is monogenic from (1.3). However, the converse does not hold in general, and when $\Delta(f)$ is not squarefree, it can be quite difficult to determine whether $f(x)$ is monogenic.

Throughout this article, we also let $\{U_n\}_{n \geq 0}$ be the sequence as defined in (1.1), and let \mathcal{D} be defined as follows:

Definition 1.1. Let $k \geq 1$ be an integer, with $k \neq 4$. Define

$$\mathcal{D} := \begin{cases} k^2 + 4 & \text{if } k \equiv 1 \pmod{2} \\ (k/2)^2 + 1 & \text{if } k \equiv 0 \pmod{2}. \end{cases}$$

We let δ denote the Legendre symbol $\left(\frac{\mathcal{D}}{p}\right)$, where p is a prime determined by the context.

In this article, we establish a connection between the monogenicity¹ of certain power-compositional trinomials and k -Wall-Sun-Sun primes. More precisely, we prove

Theorem 1.2. Let $f(x) = x^2 - kx - 1 \in \mathbb{Z}[x]$, such that $k \geq 1$, $k \not\equiv 0 \pmod{4}$ and \mathcal{D} is squarefree. Let $s \geq 1$ be an integer such that $\delta = -1$ and $\gcd(p, h\mathcal{D}) = 1$ for every prime divisor $p \geq 3$ of s , where h is the class number of $\mathbb{Q}(\sqrt{\mathcal{D}})$. Then $f(x^{s^n})$ is monogenic for all integers $n \geq 1$ if and only if no prime divisor of s is a k -Wall-Sun-Sun prime.

Theorem 1.2 is motivated in part by the recent result [9, Theorem 4.5] that if $f(x) = x^2 + ax + a$ with $a \in \{2, 3\}$, then $f(x^{s^n})$ is monogenic for all integers $n \geq 0$ if and only if $s \geq 2$ has no prime divisors p with the property that $a^{p-1} - 1 \equiv 0 \pmod{p^2}$ for some integer $a > 1$. If $a^{p-1} - 1 \equiv 0 \pmod{p^2}$ for some integer $a > 1$, then p is called a *base- a Wieferich prime* [18].

A second motivation for this article arises from recent results of Bouazzaoui [1, 2]. Let $p \geq 3$ be a rational prime. A number field K is said to be *p -rational* if the Galois group of the maximal pro- p -extension of K which is unramified outside p is a free pro- p -group of rank $r_2 + 1$, where r_2 is the number of pairs of complex embeddings of K . Let $d > 0$ be a fundamental discriminant [17], and let h_d be the class number of the real quadratic field $K = \mathbb{Q}(\sqrt{d})$. For any unit $u \in K$ with $u \notin \{\pm 1\}$, Bouazzaoui defines a rational prime $p \geq 3$ to be *Wieferich of basis u* if

$$u^{p^r-1} - 1 \equiv 0 \pmod{p^2},$$

where r is the residual degree of p in K . Let ε be the fundamental unit of K , and let $\left(\frac{d}{p}\right)$ be the Legendre symbol. Bouazzaoui proves

¹Although the terms *monogenity* and *monogeneity* are more common in the literature, we have decided to use the more grammatically-correct term *monogenicity*.

Theorem 1.3. Suppose that $p \geq 3$ is a prime such that $p \nmid (\varepsilon - \bar{\varepsilon})^2 h_d$. Then

$$(1.4) \quad \begin{aligned} K \text{ is not } p\text{-rational} &\iff p \text{ is Wieferich of basis } \varepsilon \\ &\iff \pi(p) = \pi(p^2), \\ &\iff F_{p - (\frac{d}{p})} \equiv 0 \pmod{p^2}, \end{aligned}$$

where $\{F_n\}_{n \geq 0}$ is the Lucas sequence of the first kind defined by

$$F_0 = 0, \quad F_1 = 1 \quad \text{and} \quad F_n = (\varepsilon + \bar{\varepsilon})U_{n-1} - \mathcal{N}_{K/\mathbb{Q}}(\varepsilon)U_{n-2} \quad \text{for } n \geq 2,$$

with $\pi(p)$ and $\pi(p^2)$ the respective period lengths of $\{F_n\}_{n \geq 0}$ modulo p and p^2 .

Remark 1.4. Theorem 1.3 generalizes a theorem of Greenberg [6].

2. PRELIMINARIES

The formula for the discriminant of an arbitrary monic trinomial, due to Swan [15], is given in the following theorem.

Theorem 2.1. Let $f(x) = x^N + Ax^M + B \in \mathbb{Z}[x]$, where $0 < M < N$. Let $r = \gcd(N, M)$, $N_1 = N/r$ and $M_1 = M/r$. Then

$$\Delta(f) = (-1)^{N(N-1)/2} B^{M-1} D^r,$$

where

$$(2.1) \quad D := N^{N_1} B^{N_1 - M_1} - (-1)^{N_1} M^{M_1} (N - M)^{N_1 - M_1} A^{N_1}.$$

The next two theorems are due to Capelli [13].

Theorem 2.2. Let $f(x)$ and $h(x)$ be polynomials in $\mathbb{Q}[x]$ with $f(x)$ irreducible. Suppose that $f(\alpha) = 0$. Then $f(h(x))$ is reducible over \mathbb{Q} if and only if $h(x) - \alpha$ is reducible over $\mathbb{Q}(\alpha)$.

Theorem 2.3. Let $c \in \mathbb{Z}$ with $c \geq 2$, and let $\alpha \in \mathbb{C}$ be algebraic. Then $x^c - \alpha$ is reducible over $\mathbb{Q}(\alpha)$ if and only if either there is a prime p dividing c such that $\alpha = \beta^p$ for some $\beta \in \mathbb{Q}(\alpha)$ or $4 \mid c$ and $\alpha = -4\beta^4$ for some $\beta \in \mathbb{Q}(\alpha)$.

The following theorem is a compilation of results from various sources.

Theorem 2.4. Let p be a prime.

(1) $\pi_k(p) = 2$ if and only if $k \equiv 0 \pmod{p}$.

(2) If $p = 2$, then

$$\pi_k(p) = \begin{cases} 2 & \text{if } k \equiv 0 \pmod{2} \\ 3 & \text{if } k \equiv 1 \pmod{2} \end{cases} \quad \text{and} \quad \pi_k(p^2) = \begin{cases} 2 & \text{if } k \equiv 0 \pmod{4} \\ 6 & \text{if } k \equiv 1 \pmod{4} \\ 4 & \text{if } k \equiv 2 \pmod{4} \\ 6 & \text{if } k \equiv 3 \pmod{2}. \end{cases}$$

(3) If $p \geq 3$, then $\pi_k(p) \equiv 0 \pmod{2}$.

(4) If $p \geq 3$, then $\pi_k(p^2) \in \{\pi_k(p), p\pi_k(p)\}$.

(5) If $p \geq 3$ and $\delta = 1$, then $p - 1 \equiv 0 \pmod{\pi_k(p)}$.

(6) If $\delta = -1$, then $2(p + 1) \equiv 0 \pmod{\pi_k(p)}$.

Proof. Items (1) and (2) of Theorem 2.4 follow easily by direct calculation, item (3) is a special case of work found in [5], item (4) is a special case of a result in [11], while items (5) and (6) follow from two theorems in [7]. \square

The next proposition appears as Proposition 1 in [20].

Proposition 2.5. *Let $k \geq 1$ be an integer, such that $k \neq 4$ and \mathcal{D} is squarefree. Then $\varepsilon := (k + \sqrt{k^2 + 4})/2$ is the fundamental unit of $\mathbb{Q}(\sqrt{\mathcal{D}})$ with $\mathcal{N}(\varepsilon) = -1$, where $\mathcal{N} := \mathcal{N}_{\mathbb{Q}(\varepsilon)/\mathbb{Q}}$ denotes the algebraic norm.*

Whenever the hypotheses of Proposition 2.5 hold in the sequel, we assume that ε denotes the fundamental unit $(k + \sqrt{k^2 + 4})/2$ of $\mathbb{Q}(\sqrt{\mathcal{D}})$.

The following theorem, known as *Dedekind's Index Criterion*, or simply *Dedekind's Criterion* if the context is clear, is a standard tool used in determining the monogenicity of a polynomial.

Theorem 2.6 (Dedekind [3]). *Let $K = \mathbb{Q}(\theta)$ be a number field, $T(x) \in \mathbb{Z}[x]$ the monic minimal polynomial of θ , and \mathbb{Z}_K the ring of integers of K . Let p be a prime number and let $\overline{}$ denote reduction of $*$ modulo p (in \mathbb{Z} , $\mathbb{Z}[x]$ or $\mathbb{Z}[\theta]$). Let*

$$\overline{T}(x) = \prod_{i=1}^k \overline{\tau}_i(x)^{e_i}$$

be the factorization of $T(x)$ modulo p in $\mathbb{F}_p[x]$, and set

$$g(x) = \prod_{i=1}^k \tau_i(x),$$

where the $\tau_i(x) \in \mathbb{Z}[x]$ are arbitrary monic lifts of the $\overline{\tau}_i(x)$. Let $h(x) \in \mathbb{Z}[x]$ be a monic lift of $\overline{T}(x)/\overline{g}(x)$ and set

$$F(x) = \frac{g(x)h(x) - T(x)}{p} \in \mathbb{Z}[x].$$

Then

$$[\mathbb{Z}_K : \mathbb{Z}[\theta]] \not\equiv 0 \pmod{p} \iff \gcd(\overline{F}, \overline{g}, \overline{h}) = 1 \text{ in } \mathbb{F}_p[x].$$

The next result is essentially an algorithmic adaptation of Theorem 2.6 specifically for trinomials.

Theorem 2.7. [8] *Let $N \geq 2$ be an integer. Let $K = \mathbb{Q}(\theta)$ be an algebraic number field with $\theta \in \mathbb{Z}_K$, the ring of integers of K , having minimal polynomial $f(x) = x^N + Ax^M + B$ over \mathbb{Q} , with $\gcd(M, N) = r$, $N_1 = N/r$ and $M_1 = M/r$. Let D be as defined in (2.1). A prime factor p of $\Delta(f)$ does not divide $[\mathbb{Z}_K : \mathbb{Z}[\theta]]$ if and only if p satisfies one of the following items:*

- (1) *when $p \mid A$ and $p \mid B$, then $p^2 \nmid B$;*
- (2) *when $p \mid A$ and $p \nmid B$, then*

$$\text{either } p \mid A_2 \text{ and } p \nmid B_1 \quad \text{or} \quad p \nmid A_2 \left((-B)^{M_1} A_2^{N_1} - (-B_1)^{N_1} \right),$$

$$\text{where } A_2 = A/p \text{ and } B_1 = \frac{B + (-B)^{p^e}}{p} \text{ with } p^e \parallel N;$$

- (3) *when $p \nmid A$ and $p \mid B$, then*

$$\text{either } p \mid A_1 \text{ and } p \nmid B_2 \quad \text{or} \quad p \nmid A_1 B_2^{M-1} \left((-A)^{M_1} A_1^{N_1-M_1} - (-B_2)^{N_1-M_1} \right),$$

$$\text{where } A_1 = \frac{A + (-A)^{p^j}}{p} \text{ with } p^j \parallel (N - M), \text{ and } B_2 = B/p;$$

- (4) when $p \nmid AB$ and $p \mid M$ with $N = up^m$, $M = vp^m$, $p \nmid \gcd(u, v)$, then the polynomials

$$x^{N/p^m} + Ax^{M/p^m} + B \quad \text{and} \quad \frac{Ax^M + B + (-Ax^{M/p^m} - B)^{p^m}}{p}$$

are coprime modulo p ;

- (5) when $p \nmid ABM$, then $p^2 \nmid D/r^{N_1}$.

Remark 2.8. We will find both Theorem 2.6 and Theorem 2.7 useful in our investigations.

The next theorem follows from Corollary (2.10) in [10].

Theorem 2.9. Let K and L be number fields with $K \subset L$. Then

$$\Delta(K)^{[L:K]} \mid \Delta(L).$$

3. THE PROOF OF THEOREM 1.2

We first prove some lemmas.

Lemma 3.1. Let k and s be positive integers with $k \neq 4$. Let $f(x) = x^2 - kx - 1$. If \mathcal{D} is squarefree, then $f(x^{s^n})$ is irreducible over \mathbb{Q} for all integers $n \geq 1$.

Proof. Since $\mathcal{D} > 1$ is squarefree, it follows that $f(x)$ is irreducible over \mathbb{Q} , and the trivial case of $s = 1$ is true. Suppose then that $s \geq 2$. Note that $f(\varepsilon) = 0$. Let $h(x) = x^{s^n}$ and assume, by way of contradiction, that $f(h(x))$ is reducible. Then, by Theorems 2.2 and 2.3 (with $\alpha = \varepsilon$), we have, for some $\beta \in \mathbb{Q}(\varepsilon)$, that either $\varepsilon = \beta^p$ for some prime p dividing s , or $\varepsilon = -4\beta^4$ if $s^n \equiv 0 \pmod{4}$. Thus, it is immediate that $\varepsilon = -4\beta^4$ is impossible since $\mathcal{N}(\varepsilon) = -1$ and $\mathcal{N}(-4\beta) \equiv 0 \pmod{16}$. Hence, $\varepsilon = \beta^p$ for some prime divisor p of s . Then, we see by taking norms that

$$-1 = \mathcal{N}(\varepsilon) = \mathcal{N}(\beta)^p,$$

which implies that $p \equiv 1 \pmod{2}$ and $\mathcal{N}(\beta) = -1$, since $\mathcal{N}(\beta) \in \mathbb{Z}$. Thus, β is a unit, and therefore $\beta = \pm \varepsilon^j$ for some $j \in \mathbb{Z}$, since ε is the fundamental unit of $\mathbb{Q}(\sqrt{\mathcal{D}})$ by Proposition 2.5. Consequently,

$$\varepsilon = \beta^p = (\pm 1)^p \varepsilon^{jp},$$

which implies that $(\pm 1)^p \varepsilon^{jp-1} = 1$, contradicting the fact that ε has infinite order in the unit group of the ring of algebraic integers of the real quadratic field $\mathbb{Q}(\sqrt{\mathcal{D}})$. \square

Lemma 3.2. Let $k \geq 1$ be an integer, and let $f(x) = x^2 - kx - 1$. Then $f(x)$ is monogenic if and only if $k \not\equiv 0 \pmod{4}$ and \mathcal{D} is squarefree.

Proof. By Lemma 3.1, $f(x)$ is irreducible over \mathbb{Q} . Let $f(\theta) = 0$, and let p be a prime divisor of $\Delta(f) = k^2 + 4$. To examine the monogenicity of $f(x)$, we use Theorem 2.7. Suppose first that $p \mid k$. Then $p = 2$, and item (2) of Theorem 2.7 applies. Since $A_2 = k/2$ and $B_1 = 0$, it is easy to see that

$$[\mathbb{Z}_K : \mathbb{Z}[\theta]] \not\equiv 0 \pmod{2} \iff k \not\equiv 0 \pmod{4}.$$

Suppose next that $p \nmid k$, so that $p \neq 2$. Then, by item (5) of Theorem 2.7, we deduce that

$$[\mathbb{Z}_K : \mathbb{Z}[\theta]] \not\equiv 0 \pmod{p} \iff k^2 + 4 \not\equiv 0 \pmod{p^2},$$

which completes the proof. \square

Lemma 3.3. *Let $k \in \mathbb{Z}$ with $k \geq 1$, and let p be a prime.*

- (1) *If $p = 2$, then p is a k -Wall-Sun-Sun prime if and only if $k \equiv 0 \pmod{4}$.*
- (2) *If $p \geq 3$ and $k \equiv 0 \pmod{p^2}$, then p is a k -Wall-Sun-Sun prime.*

Proof. Item (1) is easily verified by direct calculation using Theorem 2.4. For item (2), note that $U_2 = k$ and $U_3 = k^2 + 1$. Thus, since $k \equiv 0 \pmod{p^2}$, we have that $\pi_k(p) = 2$ and $U_2 \equiv 0 \pmod{p^2}$. \square

For the next lemma, we let $\text{ord}_m(*)$ denote the order of $*$ modulo the integer $m \geq 2$.

Lemma 3.4. *Let $k \geq 1$ be an integer, such that $k \neq 4$ and \mathcal{D} is squarefree. Let $p \geq 3$ be a prime such that $\delta = -1$. Then*

- (1) $\text{ord}_m(\varepsilon) = \text{ord}_m(\bar{\varepsilon}) = \pi_k(m)$ for $m \in \{p, p^2\}$,
- (2) $\varepsilon^{p+1} + 1 \equiv 0 \pmod{p}$.

Proof. It follows from [12] that the order modulo an odd integer $m \geq 3$ of the companion matrix \mathcal{C} for the characteristic polynomial of $\{U_n\}_{n \geq 0}$ is $\pi_k(m)$. The characteristic polynomial of $\{U_n\}_{n \geq 0}$ is $f(x) = x^2 - kx - 1$, so that

$$\mathcal{C} = \begin{bmatrix} 0 & 1 \\ 1 & k \end{bmatrix}.$$

Since the eigenvalues of \mathcal{C} are ε and $\bar{\varepsilon}$, we conclude that

$$\text{ord}_m \left(\begin{bmatrix} \varepsilon & 0 \\ 0 & \bar{\varepsilon} \end{bmatrix} \right) = \text{ord}_m(\mathcal{C}) = \pi_k(m), \quad \text{for } m \in \{p, p^2\}.$$

Let $z \geq 1$ be an integer, and suppose that $\varepsilon^z = a + b\sqrt{\mathcal{D}} \in \mathbb{Q}(\sqrt{\mathcal{D}})$. Then $\mathcal{N}(\varepsilon^z) = a^2 - \mathcal{D}b^2$. But $\mathcal{N}(\varepsilon^z) = \mathcal{N}(\varepsilon)^z = (-1)^z$, so that $a^2 - \mathcal{D}b^2 = (-1)^z$. Thus,

$$\bar{\varepsilon}^z = (-1/\varepsilon)^z = (-1)^z/(a + b\sqrt{\mathcal{D}}) = (-1)^z(a - b\sqrt{\mathcal{D}})/(a^2 - \mathcal{D}b^2) = a - b\sqrt{\mathcal{D}}.$$

Hence, since $\delta = -1$, it follows that

$$\varepsilon^z \equiv 1 \pmod{m} \quad \text{if and only if} \quad \bar{\varepsilon}^z \equiv 1 \pmod{m}$$

for $m \in \{p, p^2\}$, which establishes item (1).

For item (2), since $\delta = -1$, we have by Euler's criterion that

$$\left(\sqrt{k^2 + 4} \right)^{p+1} = (k^2 + 4)^{(p-1)/2} (k^2 + 4) \equiv \delta(k^2 + 4) \equiv -(k^2 + 4) \pmod{p},$$

which implies

$$\left(\sqrt{k^2 + 4} \right)^p \equiv -\sqrt{k^2 + 4} \pmod{p}.$$

Hence,

$$\begin{aligned}
\varepsilon^{p+1} &= \left(\frac{k + \sqrt{k^2 + 4}}{2} \right) \left(\frac{k + \sqrt{k^2 + 4}}{2} \right)^p \\
&= \left(\frac{k + \sqrt{k^2 + 4}}{2} \right) \sum_{j=0}^p \binom{p}{j} \left(\frac{k}{2} \right)^j \left(\frac{\sqrt{k^2 + 4}}{2} \right)^{p-j} \\
&\equiv \left(\frac{k + \sqrt{k^2 + 4}}{2} \right) \left(\left(\frac{k}{2} \right)^p + \left(\frac{\sqrt{k^2 + 4}}{2} \right)^p \right) \pmod{p} \\
&\equiv \left(\frac{k + \sqrt{k^2 + 4}}{2} \right) \left(\frac{k - \sqrt{k^2 + 4}}{2} \right) \pmod{p} \\
&\equiv -1 \pmod{p},
\end{aligned}$$

which completes the proof of the lemma. \square

Lemma 3.5. *Let $p \geq 3$ be a prime. Then p is a k -Wall-Sun-Sun prime if and only if $U_{p-\delta} \equiv 0 \pmod{p^2}$.*

Proof. We provide details only for $\delta = -1$ since the proof is similar when $\delta = 1$.

Suppose first that p is a k -Wall-Sun-Sun prime. Then

$$(3.1) \quad U_{\pi_k(p)} \equiv 0 \pmod{p^2}.$$

Using the Binet-formula representation and item (3) of Theorem 2.4, we have that

$$(3.2) \quad U_{\pi_k(p)} = \frac{\varepsilon^{\pi_k(p)} - (-1/\varepsilon)^{\pi_k(p)}}{\varepsilon + 1/\varepsilon} = \frac{(\varepsilon^{\pi_k(p)} - 1)(\varepsilon^{\pi_k(p)} + 1)}{\varepsilon^{\pi_k(p)}(\varepsilon + 1/\varepsilon)}.$$

By Lemma 3.4, $\varepsilon^{\pi_k(p)} \equiv 1 \pmod{p}$, so that $\varepsilon^{\pi_k(p)} + 1 \equiv 2 \pmod{p}$. Hence,

$$\varepsilon^{\pi_k(p)} - 1 \equiv 0 \pmod{p^2}$$

by (3.1). Thus, if $\delta = -1$, then

$$(3.3) \quad U_{p-\delta} = \frac{\varepsilon^{2(p+1)-1} - 1}{\varepsilon^{p+1}(\varepsilon + 1/\varepsilon)} \equiv 0 \pmod{p^2},$$

by item (6) of Theorem 2.4, which completes the proof in this direction.

Conversely, with $\delta = -1$, suppose that (3.3) holds. From item (6) of Theorem 2.4, we can write $2(p+1) = z\pi_k(p)$. Then

$$\varepsilon^{2(p+1)} - 1 = (\varepsilon^{\pi_k(p)} - 1)S \equiv 0 \pmod{p^2},$$

where

$$S = (\varepsilon^{\pi_k(p)})^{z-1} + (\varepsilon^{\pi_k(p)})^{z-2} + \dots + \varepsilon^{\pi_k(p)} + 1 \equiv z \not\equiv 0 \pmod{p},$$

since $\varepsilon^{\pi_k(p)} \equiv 1 \pmod{p}$ by Lemma 3.4. Thus, $\varepsilon^{\pi_k(p)} - 1 \equiv 0 \pmod{p^2}$, which implies that $U_{\pi_k(p)} \equiv 0 \pmod{p^2}$ by (3.2), completing the proof of the lemma. \square

Note that \mathcal{D} and $4\mathcal{D}$ are fundamental discriminants when $k \equiv 1 \pmod{2}$ and $k \equiv 0 \pmod{2}$, respectively. The next lemma then follows from Theorem 1.3 and Lemma 3.5.

Lemma 3.6. *Let $k \in \mathbb{Z}$ with $k \geq 1$, and let p be a prime such that $p \geq 3$ and $\gcd(\mathcal{D}h, p) = 1$, where h is the class number of $\mathbb{Q}(\sqrt{\mathcal{D}})$. Then*

$$\begin{aligned} p \text{ is a } k\text{-Wall-Sun-Sun prime} &\iff \varepsilon^{p^r-1} - 1 \equiv 0 \pmod{p^2}, \\ &\iff \pi_k(p) = \pi_k(p^2), \\ &\iff U_{p-\delta} \equiv 0 \pmod{p^2}, \end{aligned}$$

where r is the residual degree of p in $\mathbb{Q}(\sqrt{\mathcal{D}})$.

Lemma 3.7. *Let $k \in \mathbb{Z}$, such that $k \geq 1$, $k \not\equiv 0 \pmod{4}$ and \mathcal{D} is squarefree. Let p be a prime such that $p \geq 3$, $p \nmid k$ and $\gcd(p, \mathcal{D}h) = 1$, where h is the class number of $\mathbb{Q}(\sqrt{\mathcal{D}})$. If $\delta = -1$, then the following conditions are equivalent:*

- (1) p is a k -Wall-Sun-Sun prime,
- (2) $\varepsilon^{2p^m} - k\varepsilon^{p^m} - 1 \equiv 0 \pmod{p^2}$ for all integers $m \geq 1$,
- (3) $\varepsilon^{2p^m} - k\varepsilon^{p^m} - 1 \equiv 0 \pmod{p^2}$ for some integer $m \geq 1$.

Proof. First, observe that item (2) clearly implies item (3).

We show next that item (1) implies item (2). Since $\delta = -1$, we see from Theorem 2.4 that $2(p+1) \equiv 0 \pmod{\pi_k(p)}$. Thus, since p is a k -Wall-Sun-Sun prime, it follows from Lemma 3.6 that $\pi_k(p) = \pi_k(p^2)$. Hence, $2(p+1) \equiv 0 \pmod{\pi_k(p^2)}$ and

$$(3.4) \quad \varepsilon^{2(p+1)} - 1 \equiv (\varepsilon^{p+1} - 1)(\varepsilon^{p+1} + 1) \equiv 0 \pmod{p^2}$$

by part (1) of Lemma 3.4. Since

$$(3.5) \quad \gcd(\varepsilon^{p+1} - 1, \varepsilon^{p+1} + 1) \not\equiv 0 \pmod{p},$$

we conclude from (3.4) and part (2) of Lemma 3.4 that

$$(3.6) \quad \varepsilon^p \equiv -\varepsilon^{-1} \pmod{p^2}.$$

Let $m \geq 1$ be an integer. Since

$$p^m \equiv \begin{cases} 1 & \pmod{2(p+1)} & \text{if } m \equiv 0 \pmod{2}, \\ p & \pmod{2(p+1)} & \text{if } m \equiv 1 \pmod{2}, \end{cases}$$

we have from item (1) of Lemma 3.4 that

$$\varepsilon^{2p^m} - k\varepsilon^{p^m} - 1 \equiv \begin{cases} \varepsilon^2 - k\varepsilon - 1 & \pmod{p^2} & \text{if } m \equiv 0 \pmod{2}, \\ \varepsilon^{2p} - k\varepsilon^p - 1 & \pmod{p^2} & \text{if } m \equiv 1 \pmod{2}. \end{cases}$$

Using (3.6), we deduce that

$$\varepsilon^{2p} - k\varepsilon^p - 1 \equiv \frac{-(\varepsilon^2 - k\varepsilon - 1)}{\varepsilon^2} \pmod{p^2},$$

which completes the proof in this direction since $\varepsilon^2 - k\varepsilon - 1 = 0$.

Finally, we show that item (3) implies item (1). By items (6) and (4) of Theorem 2.4, we deduce that $2p(p+1) \equiv 0 \pmod{\pi_k(p^2)}$. Let $m \geq 1$ be any integer. Then, since

$$p^m \equiv \begin{cases} p & \pmod{2p(p+1)} & \text{if } m \equiv 1 \pmod{2}, \\ p^2 & \pmod{2p(p+1)} & \text{if } m \equiv 0 \pmod{2}, \end{cases}$$

it follows from item (1) of Lemma 3.4 that

$$\varepsilon^{2p^m} - k\varepsilon^{p^m} - 1 \equiv \begin{cases} \varepsilon^{2p} - k\varepsilon^p - 1 & \pmod{p^2} & \text{if } m \equiv 1 \pmod{2}, \\ \varepsilon^{2p^2} - k\varepsilon^{p^2} - 1 & \pmod{p^2} & \text{if } m \equiv 0 \pmod{2}. \end{cases}$$

By assumption, we have that

$$(3.7) \quad \text{either } \varepsilon^{2p} - k\varepsilon^p - 1 \equiv 0 \pmod{p^2} \quad \text{or} \quad \varepsilon^{2p^2} - k\varepsilon^{p^2} - 1 \equiv 0 \pmod{p^2}.$$

The zeros of $f(x) = x^2 - kx - 1$ in the ring $(\mathbb{Z}/p\mathbb{Z})[\mathcal{D}]$ are ε and $\bar{\varepsilon} = -1/\varepsilon$. By Hensel, these zeros lift to the zeros ε and $\bar{\varepsilon}$ of $f(x)$ in $(\mathbb{Z}/p^2\mathbb{Z})[\mathcal{D}]$. Hence, if the first possibility of (3.7) is true, then

$$(3.8) \quad \varepsilon^p \equiv \varepsilon \pmod{p^2} \quad \text{or} \quad \varepsilon^p \equiv -1/\varepsilon \pmod{p^2}.$$

Suppose the first possibility of (3.8) is true. Then, by item (2) of Lemma 3.4, we have that

$$-1 \equiv \varepsilon^{p+1} \equiv \varepsilon^p \varepsilon \equiv \varepsilon^2 \equiv \frac{k^2 + 2 + k\sqrt{k^2 + 4}}{2} \pmod{p}.$$

Thus,

$$\frac{k^2 + 4 + k\sqrt{k^2 + 4}}{2} \equiv 0 \pmod{p},$$

which implies that $k^2 + 4 \equiv 0 \pmod{p}$, contradicting the fact that $\gcd(p, \mathcal{D}) = 1$. Thus, the second possibility of (3.8) holds, which implies that $\varepsilon^{p+1} \equiv -1 \pmod{p^2}$. Since $\delta = -1$, the residual degree of p is $r = 2$. Hence,

$$\varepsilon^{p^r-1} - 1 \equiv \varepsilon^{p^2-1} - 1 \equiv (\varepsilon^{p+1})^{p-1} - 1 \equiv (-1)^{p-1} - 1 \equiv 0 \pmod{p^2},$$

which implies that p is a k -Wall-Sun-Sun prime by Lemma 3.6.

Suppose now that the second possibility,

$$(3.9) \quad \varepsilon^{2p^2} - k\varepsilon^{p^2} - 1 \equiv 0 \pmod{p^2},$$

of (3.7) is true. Then

$$(3.10) \quad \varepsilon^{p^2} \equiv \varepsilon \pmod{p^2} \quad \text{or} \quad \varepsilon^{p^2} \equiv -1/\varepsilon \pmod{p^2}.$$

If the second possibility in (3.10) holds, then $\varepsilon^{2(p^2+1)} \equiv 1 \pmod{p}$. Since $\text{ord}_p(\varepsilon) = \pi_k(p)$ by item (1) of Lemma 3.4, it follows that $2p^2 + 2 \equiv 0 \pmod{\pi_k(p)}$. By item (6) of Theorem 2.4, we have that $2p + 2 \equiv 0 \pmod{\pi_k(p)}$. Hence,

$$\pi_k(p) \text{ divides } (2p^2 + 2) - (p - 1)(2p + 2) = 4,$$

so that $\pi_k(p) \in \{2, 4\}$. Recall that $p \nmid k$ by hypothesis. Thus, $\pi_k(p) \neq 2$ by item (1) of Theorem 2.4. If $\pi_k(p) = 4$, then

$$U_4 = k(k^2 + 2) \equiv 0 \pmod{p} \quad \text{and} \quad U_5 = k^2(k^2 + 3) + 1 \equiv 1 \pmod{p}.$$

Hence, it follows that

$$p \text{ divides } (k^2 + 3) - (k^2 + 2) = 1,$$

which is impossible. Therefore, the first possibility in (3.10) holds, which implies that $\varepsilon^{p^2-1} - 1 \equiv 0 \pmod{p^2}$, and the proof is complete by Lemma 3.6. \square

Corollary 3.8. *Let $k \in \mathbb{Z}$, such that $k \geq 1$, $k \not\equiv 0 \pmod{4}$ and \mathcal{D} is squarefree. Let p be a prime such that $p \geq 3$, $p \nmid k$ and $\gcd(p, \mathcal{D}h) = 1$, where h is the class number of $\mathbb{Q}(\sqrt{\mathcal{D}})$. If $\delta = -1$, then the following conditions are equivalent:*

- (1) p is a k -Wall-Sun-Sun prime,
- (2) $\bar{\varepsilon}^{2p^m} - k\bar{\varepsilon}^{p^m} - 1 \equiv 0 \pmod{p^2}$ for all integers $m \geq 1$,
- (3) $\bar{\varepsilon}^{2p^m} - k\bar{\varepsilon}^{p^m} - 1 \equiv 0 \pmod{p^2}$ for some integer $m \geq 1$.

Proof. Since $\bar{\varepsilon} = -1/\varepsilon$, we have that

$$\bar{\varepsilon}^{2p^m} - k\bar{\varepsilon}^{p^m} - 1 = -\frac{\varepsilon^{2p^m} - k\varepsilon^{p^m} - 1}{\varepsilon^{2p^m}}.$$

Thus, the corollary follows from Lemma 3.7. \square

Proof of Theorem 1.2. For brevity of notation, define

$$\mathcal{F}_n(x) := f(x^{s^n}) = x^{2s^n} - kx^{s^n} - 1.$$

Since \mathcal{D} is squarefree and $k \neq 4$, we have that $\mathcal{F}_n(x)$ is irreducible for all $n \geq 1$ by Lemma 3.1. The case $s = 1$ follows from Lemma 3.2. So assume that $s \geq 2$.

(\Rightarrow) Suppose that s has a prime divisor p that is a k -Wall-Sun-Sun prime. We claim that $\mathcal{F}_1(x)$ is not monogenic. Let $\mathcal{F}_1(\theta) = 0$, $K = \mathbb{Q}(\theta)$ and \mathbb{Z}_K be the ring of integers of K . Since $k \not\equiv 0 \pmod{4}$, it follows from item (1) of Lemma 3.3 that $p \geq 3$. If $p \mid k$, then $\pi_k(p) = 2$, so that $U_2 = k \equiv 0 \pmod{p^2}$, since p is a k -Wall-Sun-Sun prime. Thus, applying item (2) of Theorem 2.7 to $\mathcal{F}_1(x)$, we see that

$$B_1 \equiv 0 \pmod{p} \quad \text{and} \quad A_2 = k/p \equiv 0 \pmod{p},$$

from which we conclude that $[\mathbb{Z}_K : \mathbb{Z}[\theta]] \equiv 0 \pmod{p}$ and $\mathcal{F}_1(x)$ is not monogenic.

Suppose next that $p \nmid k$ and $p^m \parallel s$, with $m \geq 1$. We apply Theorem 2.6 to $T(x) := \mathcal{F}_1(x)$ using the prime p . Let $\tau(x) = x^{2s/p^m} - kx^{s/p^m} - 1$, and suppose that

$$\bar{\tau}(x) = x^{2s/p^m} - \bar{k}x^{s/p^m} - 1 = \prod_{i=1}^k \bar{\tau}_i(x)^{e_i},$$

where the $\bar{\tau}_i(x)$ are irreducible. Then $\bar{T}(x) = \prod_{i=1}^k \bar{\tau}_i(x)^{p^m e_i}$. Thus, we can let

$$g(x) = \prod_{i=1}^k \tau_i(x) \quad \text{and} \quad h(x) = \prod_{i=1}^k \tau_i(x)^{p^m e_i - 1},$$

where the $\tau_i(x)$ are monic lifts of the $\bar{\tau}_i(x)$. Note also that

$$g(x)h(x) = \prod_{i=1}^k \tau_i(x)^{e_i} = \bar{\tau}(x) + pr(x),$$

for some $r(x) \in \mathbb{Z}[x]$. Then, in Theorem 2.6, we have that

$$\begin{aligned} F(x) &= \frac{g(x)h(x) - T(x)}{p} \\ &= \frac{(\bar{\tau}(x) + pr(x))^{p^m} - T(x)}{p} \\ &= \sum_{j=1}^{p^m-1} \frac{\binom{p^m}{j}}{p} \bar{\tau}(x)^j (pr(x))^{p^m-j} + p^{p^m-1} r(x)^{p^m} + \frac{\bar{\tau}(x)^{p^m} - T(x)}{p}, \end{aligned}$$

which implies that

$$\bar{F}(x) = \overline{\left(\frac{\bar{\tau}(x)^{p^m} - T(x)}{p} \right)}.$$

Suppose that $\bar{\tau}(\alpha) = 0$. Then

$$\bar{\tau}(\alpha)^{p^m} = (\beta^2 - \bar{k}\beta - 1)^{p^m} = 0,$$

where $\beta = \alpha^{s/p^m}$, so that $\alpha^s = \beta^{p^m}$. Since $f(\varepsilon) = f(\bar{\varepsilon}) = 0$, we assume, without loss of generality, that $\alpha^s = \varepsilon^{p^m}$. Thus,

$$\overline{F}(\alpha) = -\overline{\left(\frac{T(\alpha)}{p}\right)} = -\overline{\left(\frac{\alpha^{2s} - k\alpha^s - 1}{p}\right)} = -\overline{\left(\frac{\varepsilon^{2p^m} - k\varepsilon^{p^m} - 1}{p}\right)} = 0,$$

since $\varepsilon^{2p^m} - k\varepsilon^{p^m} - 1 \equiv 0 \pmod{p^2}$ by Lemma 3.7. Therefore, by Theorem 2.6, we conclude that $[\mathbb{Z}_K : \mathbb{Z}[\theta]] \equiv 0 \pmod{p}$ and $\mathcal{F}_1(x)$ is not monogenic, which completes the proof in this direction.

(\Leftarrow) Assume now that no prime divisor of s is a k -Wall-Sun-Sun prime. By Lemma 3.2, we see that $\mathcal{F}_0(x) = f(x)$ is monogenic. Note that $\mathcal{F}_0(\varepsilon) = 0$. For $n \geq 0$, define

$$\varepsilon_n := \varepsilon^{1/s^n} \quad \text{and} \quad K_n := \mathbb{Q}(\varepsilon_n).$$

Then $\varepsilon_0 = \varepsilon$ and, since $\mathcal{F}_0(x)$ is monogenic, we have that $\Delta(\mathcal{F}_0) = \Delta(K_0)$. Additionally, by Lemma 3.1,

$$\mathcal{F}_n(\varepsilon_n) = 0 \quad \text{and} \quad [K_{n+1} : K_n] = s$$

for all $n \geq 0$. We assume that $\mathcal{F}_n(x)$ is monogenic, so that $\Delta(\mathcal{F}_n) = \Delta(K_n)$, and we proceed by induction on n to show that $\mathcal{F}_{n+1}(x)$ is monogenic. Let $\mathbb{Z}_{K_{n+1}}$ denote the ring of integers of K_{n+1} . Consequently, by Theorem 2.9, it follows that

$$\Delta(\mathcal{F}_n)^s \text{ divides } \Delta(K_{n+1}) = \frac{\Delta(\mathcal{F}_{n+1})}{[\mathbb{Z}_{K_{n+1}} : \mathbb{Z}[\varepsilon_{n+1}]]^2},$$

which implies that

$$[\mathbb{Z}_{K_{n+1}} : \mathbb{Z}[\varepsilon_{n+1}]]^2 \text{ divides } \frac{\Delta(\mathcal{F}_{n+1})}{\Delta(\mathcal{F}_n)^s}.$$

We see from Theorem 2.1 that

$$|\Delta(\mathcal{F}_n)^s| = s^{2ns^{n+1}}(k^2 + 4)^{s^{n+1}} \quad \text{and} \quad |\Delta(\mathcal{F}_{n+1})| = s^{2(n+1)s^{n+1}}(k^2 + 4)^{s^{n+1}}.$$

Hence,

$$\left| \frac{\Delta(\mathcal{F}_{n+1})}{\Delta(\mathcal{F}_n)^s} \right| = s^{2s^{n+1}}.$$

Thus, it is enough to show that $\gcd(s, [\mathbb{Z}_{K_{n+1}} : \mathbb{Z}[\varepsilon_{n+1}]]) = 1$. Suppose then that p is a prime divisor of s .

If $p \mid k$, then we can apply item (2) of Theorem 2.7 to $\mathcal{F}_{n+1}(x)$. Observe that $A_2 = k/p$ and $B_1 = 0$, so that the first condition of item (2) does not hold. If

$$A_2 \left((-B)^{M_1} A_2^{N_1} - (-B_1)^{N_1} \right) = k^3/p^3 \equiv 0 \pmod{p},$$

then $k \equiv 0 \pmod{p^2}$, and p is a k -Wall-Sun-Sun prime by Lemma 3.3, contradicting the fact that s has no such prime divisors. Hence, the second condition of item (2) holds and therefore, $[\mathbb{Z}_{K_{n+1}} : \mathbb{Z}[\varepsilon_{n+1}]] \not\equiv 0 \pmod{p}$ in this case.

Suppose next that $p \nmid k$ and $p^m \parallel s$, with $m \geq 1$. We apply Theorem 2.6 to $T(x) := \mathcal{F}_{n+1}(x)$ using the prime p . We assume that $\overline{\tau}(\alpha) = 0$, with

$$\tau(x) = x^{2s^n/p^{mn}} - kx^{s^n/p^{mn}} - 1,$$

and use the same argument as in the other direction of the proof where we showed that $\mathcal{F}_1(x)$ is not monogenic when $p \nmid k$. Omitting the details, we arrive at

$$\overline{F}(\alpha) = -\overline{\left(\frac{T(\alpha)}{p}\right)} = -\overline{\left(\frac{\alpha^{2s^n} - k\alpha^{s^n} - 1}{p}\right)} = -\overline{\left(\frac{\varepsilon^{2p^{mn}} - k\varepsilon^{p^{mn}} - 1}{p}\right)}.$$

Therefore, $\overline{F}(\alpha) = 0$ if and only if $\varepsilon^{2p^{mn}} - k\varepsilon^{p^{mn}} - 1 \equiv 0 \pmod{p^2}$, which is true if and only if p is a k -Wall-Sun-Sun prime by Lemma 3.7. Since s has no prime divisors that are k -Wall-Sun-Sun primes, it follows that $\gcd(\overline{F}, \overline{g}, \overline{h}) = 1$. Hence, by Theorem 2.6, we conclude that $[\mathbb{Z}_{K_{n+1}} : \mathbb{Z}[\varepsilon_{n+1}]] \not\equiv 0 \pmod{p}$ and $\mathcal{F}_{n+1}(x)$ is monogenic, which completes the proof of the theorem. \square

DATA AVAILABILITY STATEMENT

The author confirms that all relevant data are included in the article.

REFERENCES

- [1] Z. Bouazzaoui, *Fibonacci numbers and real quadratic p -rational fields*, Period. Math. Hungar. **81** (2020), no. 1, 123–133.
- [2] Z. Bouazzaoui, *On periods of Fibonacci sequences and real quadratic p -rational fields*, Fibonacci Quart. **58** (2020), no. 5, 103–110.
- [3] H. Cohen, *A Course in Computational Algebraic Number Theory*, Springer-Verlag, 2000.
- [4] R. Crandall, K. Dilcher and C. Pomerance, *A search for Wieferich and Wilson primes*, Math. Comp. **66** (1997), no. 217, 433–449.
- [5] S. Falcón and Á. Plaza, *On k -Fibonacci sequences modulo m* , Chaos, Solitons and Fractals **41** (2009), 497–504.
- [6] R. Greenberg, *Galois representations with open image*, Ann. Math. Qué **40** (2016), no. 1, 83–119.
- [7] S. Gupta, P. Rockstroh and F. E. Su, *Splitting fields and periods of Fibonacci sequences modulo primes*, Math. Mag. **85** (2012), no. 2, 130–135.
- [8] A. Jakhar, S. Khanduja and N. Sangwan, *Characterization of primes dividing the index of a trinomial*, Int. J. Number Theory **13** (2017), no. 10, 2505–2514.
- [9] L. Jones, *The monogenity of power-compositional Eisenstein polynomials*, Ann. Math. Inform. (2022) <https://ami.uni-eszterhazy.hu/index.php?vol=Latest>.
- [10] J. Neukirch, *Algebraic Number Theory*, Springer-Verlag, Berlin, 1999.
- [11] M. Renault, *The period, rank, and order of the (a, b) -Fibonacci sequence mod m* , Math. Mag. **86** (2013), no. 5, 372–380.
- [12] D. W. Robinson, *A note on linear recurrent sequences modulo m* , Amer. Math. Monthly **73** (1966), 619–621.
- [13] A. Schinzel, *Polynomials with Special Regard to Reducibility*, Encyclopedia of Mathematics and its Applications, **77**, Cambridge University Press, Cambridge, 2000.
- [14] Zhi Hong Sun and Zhi Wei Sun, *Fibonacci numbers and Fermat’s last theorem*, Acta Arith. **60** (1992), no. 4, 371–388.
- [15] R. Swan, *Factorization of polynomials over finite fields*, Pacific J. Math. **12** (1962), 1099–1106.
- [16] D. D. Wall, *Fibonacci series modulo m* , Amer. Math. Monthly **67** (1960), 525–532.
- [17] *Fundamental Discriminant* https://en.wikipedia.org/wiki/Fundamental_discriminant
- [18] *Wieferich Prime* https://en.wikipedia.org/wiki/Wieferich_prime
- [19] *Wall-Sun-Sun Prime* https://en.wikipedia.org/wiki/Wall%E2%80%93Sun%E2%80%93Sun_prime
- [20] H. Yokoi, *On real quadratic fields containing units with norm -1* , Nagoya Math. J. **33** (1968), 139–152.

PROFESSOR EMERITUS, DEPARTMENT OF MATHEMATICS, SHIPPENSBURG UNIVERSITY, SHIPPENSBURG, PENNSYLVANIA 17257, USA

Email address, Lenny Jones: doctorlennyjones@gmail.com