

Topological classes of rotating black holes

Di Wu*

*School of Physics and Astronomy, China West Normal University,
Nanchong, Sichuan 637002, People's Republic of China*

(Dated: December 1, 2022)

In this paper, we investigate the topological numbers for singly rotating Kerr black holes in arbitrary dimensions and four-dimensional Kerr-Newman black hole. We show that for uncharged black holes, the rotation parameter has a significant effect on the topological number, and for rotating black holes, the dimension of spacetime has a remarkable effect on the topological number too. In addition, we find that the topological numbers of the four-dimensional Kerr and Kerr-Newman black holes are the same, which seems to indicate that the electric charge parameter has no effect on the topological number of rotating black holes. Our current research provides more evidence that the conjecture put forward in [Phys. Rev. Lett. **129**, 191101 (2022)], according to which all black hole solutions should be separated into three different topological classes, is accurate, at least in the pure Einstein-Maxwell gravity theory.

I. INTRODUCTION

As one of the most remarkable and fascinating objects in nature, black hole has always been a subject of extensive theoretical and observational studies. On the observational side, recent years have witnessed remarkable successes in “seeing” the shadow of a black hole via the event horizon telescope (EHT) [1, 2] and “hearing” gravitational waves from black hole coalescence [3, 4]. On the theoretical side, the study of the topology of black holes has very recently shed new light on the nature of gravity by means of light rings [5–8], timelike circular orbit [9] and thermodynamic properties [10–19].

In particular, in Ref. [10], the black holes are treated as topological thermodynamic defects by using the generalized off-shell free energy, and thus black hole solutions are divided into three different topological classes according to their different topological numbers, which shed new light on the fundamental nature of quantum gravity. Because of its simplicity and easy maneuverability of the procedure, the topological method proposed in Ref. [10] soon attracted a great deal of attention and was then successfully applied to investigate the topological numbers of some known black solutions [15–17, 19], i.e., the Schwarzschild-AdS black hole solutions [15], the static black hole solutions in Lovelock gravity [16], the static Gauss-Bonnet-AdS black hole solutions [17], and the static black hole solution in nonlinear electrodynamics [19]. However, all of the above-mentioned progresses [10, 15–17, 19] are only restricted to the static cases, leaving the topological numbers of the rotating black holes unexplored. Up to date, the astronomical observation of black holes are basically rotating black holes, so it is very important and necessary to study the topological number of rotating black holes, which also provides the motivation of this paper.

In this paper, we investigate the topological numbers for singly rotating Kerr black holes in arbitrary dimensions and four-dimensional Kerr-Newman black hole. We show that for uncharged black holes, the rotation parameter has a significant effect on the topological number, and for rotating black

holes, the dimension of spacetime has a remarkable effect on the topological number too. The remaining part of this paper is organized as follows. In Sec. II, we first give a brief review of the topological approach, and then investigate the topological number of the four-dimensional Kerr black hole. In Sec. III, we extend the discuss in Sec. II to the cases of the d -dimensional singly rotating Kerr black holes. In Sec. IV, we turn to discuss the topological number of the four-dimensional Kerr-Newman black hole. Finally, we present our conclusions in Sec. V.

II. FOUR-DIMENSIONAL KERR BLACK HOLE

We first investigate the effect of the rotation parameter on the topological number of black holes. We consider the four-dimensional Kerr black hole case [20]. The metric of Kerr black hole has the form

$$ds^2 = -\frac{\Delta_r}{\Sigma} (dt - a \sin^2 \theta d\phi)^2 + \frac{\Sigma}{\Delta_r} dr^2 + \Sigma d\theta^2 + \frac{\sin^2 \theta}{\Sigma} [adt - (r^2 + a^2)d\phi]^2 \quad (1)$$

in the Boyer-Lindquist coordinates, where

$$\Delta_r = r^2 + a^2 - 2mr, \quad \Sigma = r^2 + a^2 \cos^2 \theta,$$

m and a are the mass and the rotation parameters of the black hole. The thermodynamic quantities associated with the above solution (1) can be computed via the standard method and have the following exquisite expressions:

$$M = m, \quad J = ma, \quad \Omega = \frac{a}{r_h^2 + a^2}, \quad (2)$$

$$S = \pi(r_h^2 + a^2), \quad T = \frac{r_h^2 - a^2}{4\pi r_h(r_h^2 + a^2)},$$

where $r_h = m \pm \sqrt{m^2 - a^2}$ are the location of the event and Cauchy horizons.

With the expressions of the thermodynamical quantities in hand, it is now the position to investigate the topological number of the four-dimensional Kerr black hole. As is shown in

* wdcwnu@163.com

Ref. [10], one can first introduce the generalized off-shell free energy

$$\mathcal{F} = M - \frac{S}{\tau} \quad (3)$$

for a black hole system with mass M and entropy S , where τ is an extra variable and can be considered as the inverse temperature of the cavity surrounding the black hole. Only when $\tau = 1/T$, the generalized free energy is on-shell.

In Ref. [10], a vector ϕ is established as

$$\phi = \left(\frac{\partial \mathcal{F}}{\partial r_h}, -\cot \Theta \csc \Theta \right), \quad (4)$$

where the parameter $0 \leq \Theta \leq \pi$ for convenience and intuition. The component ϕ^Θ is divergent at $\Theta = 0, \pi$, and thus the direction of the vector points outward. Furthermore, as discussed in Ref. [17], the extremal points of the generalized free energy landscape exactly correspond to the on-shell black holes, so the zero point of the component ϕ^{r_h} exactly meets the black hole solution. The component $\phi^\Theta = 0$ will yield $\Theta = \pi/2$.

According to Duan's ϕ -mapping topological current theory, a topological current can be stated as [21–24]

$$j^\mu = \frac{1}{2\pi} \varepsilon^{\mu\nu\rho} \varepsilon_{ab} \partial_\nu n^a \partial_\rho n^b, \quad \mu, \nu, \rho = 0, 1, 2, \quad (5)$$

where $\partial_\nu = (\partial/\partial x^\nu)$ and $x^\nu = (\tau, r_h, \Theta)$. The unit vector n is obtained as $n = (n^r, n^\Theta)$, where $n^r = \frac{\phi^{r_h}}{\|\phi\|}$ and $n^\Theta = \frac{\phi^\Theta}{\|\phi\|}$. It is simple to prove that the above topological current (5) is conserved, thus one can easily derive $\partial_\mu j^\mu = 0$. It is further shown that the topological current j^μ is a δ -function of the field configuration [7, 23, 24]

$$j^\mu = \delta^2(\phi) J^\mu \left(\frac{\phi}{x} \right), \quad (6)$$

where the three dimensional Jacobian $J^\mu \left(\frac{\phi}{x} \right)$ is defined as $\varepsilon^{ab} J^\mu \left(\frac{\phi}{x} \right) = \varepsilon^{\mu\nu\rho} \partial_\nu \phi^a \partial_\rho \phi^b$. It is easy to see that j^μ equals zero except when $\phi^a(x_i) = 0$, and one can derived the expressions of the topological number W as

$$W = \int_\Sigma j^0 d^2x = \sum_{i=1}^N \beta_i \eta_i = \sum_{i=1}^N w_i, \quad (7)$$

where β_i is the positive Hopf index counting the number of the loops that ϕ^a makes in the vector ϕ space when x^μ goes around the zero point z_i , $\eta_i = \text{sign}(J^0(\phi/x)_{z_i}) = \pm 1$ is the Brouwer degree, and last but not least, w_i is the winding number for the i -th zero point of ϕ contained in Σ . It is worth to note that if two loops Σ_1 and Σ_2 surround the same zero points of ϕ , they have the same winding number. Alternatively, if there is no zero point in the enclosed region, one can have $W = 0$.

From the results already given before in Eq. (2), one can easily obtain the generalized free energy

$$\mathcal{F} = \frac{r_h^2 + a^2}{2r_h} - \frac{\pi(r_h^2 + a^2)}{\tau} \quad (8)$$




FIG. 1: Zero points of the vector ϕ^{r_h} shown on the $r_h - \tau$ plane. To facilitate comparison with previous results in Ref. [10], the blue and red solid lines are for the Schwarzschild black hole (Sch BH) and Kerr black hole (Kerr BH) with $a/r_0 = 1$. The black dot with $\tau_c = 6\sqrt{3}\pi a$ represents the generation point for the Kerr black hole. there is only one Schwarzschild black hole when $\tau = \tau_1$, and there is one Schwarzschild black hole and two Kerr black holes when $\tau = \tau_2$.

of the four-dimensional Kerr black hole. Then the components of the vector ϕ can be computed as

$$\phi^{r_h} = \frac{1}{2} - \frac{a^2}{2r_h^2} - \frac{2\pi r_h}{\tau}, \quad (9)$$

$$\phi^\Theta = -\cot \Theta \csc \Theta. \quad (10)$$

By solving the equation $\phi^{r_h} = 0$, one can obtain a curve on the $r_h - \tau$ plane. For four-dimensional Kerr black hole, the result is

$$\tau = \frac{4\pi r_h^3}{r_h^2 - a^2}. \quad (11)$$

In Fig. 1, we show the zero points of the component ϕ^{r_h} for four-dimensional Kerr black hole with $a = r_0$, where r_0 is an arbitrary length scale set by the size of a cavity enclosing the black hole. For large τ , such as $\tau = \tau_2$, there are one and two intersection points for the Schwarzschild and Kerr black holes, respectively. The intersection points exactly satisfy the condition $\tau = 1/T$, and therefore represent the on-shell black hole solutions with the characteristic temperature $T = 1/\tau$. Similar to the RN black hole [10], but in contrast to the Schwarzschild black hole, when $\tau < \tau_c$, the two intersection points for the Kerr black hole coincide and then vanish. Especially, at the point $\tau_c = 6\sqrt{3}\pi a$, one can easily obtain $(d^2\tau/dr_h^2) = 6\sqrt{3}\pi/a > 0$ for four-dimensional Kerr black hole. It indicates that τ_c is a generation point, which is also straightforward from Fig. 1. Furthermore, the generation point τ_c divides the Kerr black hole into upper and lower branches, and one can find that the upper branch has the winding number $w = -1$, the lower branch has the winding number $w = 1$, thus the topological number $W = 0$ for four-dimensional Kerr black hole. Alternatively, one can also obtain the topological number for four-dimensional Kerr black




FIG. 2: The red arrows represent the unit vector field n on a portion of the $r_h - \Theta$ plane. The unit vector field for the four-dimensional Kerr black hole with $\tau/r_0 = 34.48$ and $a/r_0 = 1$. The zero points (ZPs) marked with black dots are at $(r_h/r_0, \Theta) = (1.46, (\pi/2))$, and $2.15, (\pi/2)$, for ZP_1 , and ZP_2 , respectively. Thus the two winding numbers $w_1 = 1$ and $w_2 = -1$, and the topological number $W = w_1 + w_2 = 0$.

hole by plot the unit vector field n in Fig. 2 for arbitrarily chosen typical values $\tau/r_0 = 34.48$ (τ must be larger than τ_c) and $a/r_0 = 1$. In Fig. 2, we find two zero points, ZP_1 and ZP_2 , at $r_h = 1.46r_0$ and $2.15r_0$, respectively. The winding numbers of these two zero points are $w_1 = 1$, $w_2 = -1$, thus one can get the topological number $W = w_1 + w_2 = 0$ for four-dimensional Kerr black hole. According to the classification proposal of black hole solutions, which is based on their different topological numbers given in Ref. [10], Kerr black hole and RN black hole are the same kind of black hole solutions since the topological numbers of both black holes are equal to zero. In addition, since the topological number of Schwarzschild black hole is -1, while that of Kerr black hole is 0, it can be easily found that the existence of rotation parameter has an important effect on the topological number for the uncharged black hole.

III. HIGHER DIMENSIONAL SINGLY-ROTATING KERR BLACK HOLES

In this section, we will extend the above discussion to the cases of a rotating black hole with higher dimensions by considering the d -dimensional singly rotating Kerr black holes. For the singly rotating Kerr black holes in arbitrary dimensions, the metric has the form [25, 26]

$$ds^2 = -\frac{\Delta_r}{\Sigma} (dt - a \sin^2 \theta d\phi)^2 + \frac{\Sigma}{\Delta_r} dr^2 + \Sigma d\theta^2 + \frac{\sin^2 \theta}{\Sigma} [adt - (r^2 + a^2)d\phi]^2 + r^2 \cos^2 \theta d\Omega_{d-4}^2, \quad (12)$$

where $d\Omega_d$ represents the metric element on a d -dimensional sphere, and

$$\Delta_r = r^2 + a^2 - 2mr^{5-d}, \quad \Sigma = r^2 + a^2 \cos^2 \theta.$$

The thermodynamic quantities are [26]

$$\begin{aligned} M &= \frac{d-2}{8\pi} \omega_{d-2} m, & J &= \frac{\omega_{d-2}}{4\pi} ma, \\ \Omega &= \frac{a}{r_h^2 + a^2}, & S &= \frac{\omega_{d-2}}{4} (r_h^2 + a^2) r_h^{d-4}, \\ T &= \frac{r_h}{2\pi} \left(\frac{1}{r_h^2 + a^2} + \frac{d-3}{2r_h^2} \right) - \frac{1}{2\pi r_h}, \end{aligned} \quad (13)$$

where $\omega_{d-2} = 2\pi^{(d-1)/2}/\Gamma[(d-1)/2]$, and the horizon radius r_h of the black hole is determined by the equation $\Delta_r = 0$.

From Eq. (13), one can get the generalized free energy

$$\begin{aligned} \mathcal{F} &= M - \frac{S}{\tau} \\ &= \frac{\omega_{d-2}(d-2)(r_h^2) + a^2}{16\pi r_h^{5-d}} - \frac{\omega_{d-2}(r_h^2 + a^2)r_h^{d-4}}{4\tau}. \end{aligned} \quad (14)$$

The components of the vector ϕ can be calculated as

$$\begin{aligned} \phi^{r_h} &= \frac{\omega_{d-2} r_h^{d-6}}{16\pi\tau} \left\{ (d-2)[(d-3)\tau - 4\pi r_h] r_h^2 \right. \\ &\quad \left. + a^2 [\tau(d-2)(d-5) - 4(d-4)\pi r_h] \right\}, \end{aligned} \quad (15)$$

$$\phi^\Theta = -\cot \Theta \csc \Theta. \quad (16)$$

By solving the equation $\phi^{r_h} = 0$, one can obtain

$$\tau = \frac{4\pi[(d-2)r_h^3 + (d-4)a^2 r_h]}{(d-2)[(d-3)r_h^2 + (d-5)a^2]} \quad (17)$$

as the zero point of the vector field ϕ .

As some examples, we show the zero points of the component ϕ^{r_h} for five- to nine-dimensional singly rotating Kerr black holes with $a = r_0$ in Fig. 3, and the unit vector field n for five- to seven-dimensional singly rotating Kerr black holes in Fig. 4 with $a = r_0$, $\tau = 20r_0$ ($d = 5$), $3\pi r_0/2$ ($d = 6$), and $16\pi r_0/15$ ($d = 7$), respectively. It is worth to noted that the unit vector field n for eight- and nine-dimensional singly rotating Kerr black holes is similar to that for six- and seven-dimensional singly rotating Kerr black holes. The only difference is that the typical values $\tau = 5\pi/6$ ($d = 8$) and $\tau = 24\pi/35$ ($d = 9$). Here we are not going to plot them again.

From Figs. 3-4, one can obtain the topological numbers of the higher dimensional singly-rotating Kerr black holes, where the topological number of the five-dimensional singly-rotating Kerr black hole is $W = 1 - 1 = 0$, which is same as that of the four-dimensional Kerr black hole. In particular, comparing Figs. 1-4, the topological numbers of the $d \geq 6$ singly-rotating Kerr black holes are both equal to -1, which are different from the four- and five-dimensional Kerr black holes. To summarize, according to the topological approach proposed in Ref. [10], we find that the $d \geq 6$ singly-rotating




FIG. 3: Zero points of the vector ϕ^{r_h} shown on the $r_h - \tau$ plane for five- to nine-dimensional singly rotating Kerr black holes. The red, blue, green, brown and purple solid lines are for five- to nine-dimensional singly rotating Kerr black holes with $a/r_0 = 1$, respectively. The black dot with $\tau_c = 4\sqrt{3}\pi a/3$ represents the generation point for the five-dimensional singly rotating Kerr black hole.

Kerr black holes and the $d = 4, 5$ singly-rotating Kerr black holes are belong to two different topological classes. Therefore, the dimension of spacetime has an important effect on the topological number of the rotating black holes, which can also be found in Ref. [16] for higher-dimensional static uncharged black holes in Lovelock gravity theory.

IV. FOUR-DIMENSIONAL KERR-NEWMAN BLACK HOLE

On the one hand, we want to explore the effect of the electric charge parameters on the topological number of rotating black holes, on the other hand, for the general consideration of the solution of four-dimensional black holes in the pure Einstein-Maxwell gravity theory, in this section, we turn to investigate the topological number of four-dimensional Kerr-Newman black hole [27, 28], whose metric and Abelian gauge potential are

$$ds^2 = -\frac{\Delta_r}{\Sigma} (dt - a \sin^2 \theta d\phi)^2 + \frac{\Sigma}{\Delta_r} dr^2 + \Sigma d\theta^2 + \frac{\sin^2 \theta}{\Sigma} [adt - (r^2 + a^2)d\phi]^2, \quad (18)$$


$$A = \frac{qr}{\Sigma} (dt - a \sin^2 \theta d\phi), \quad (19)$$

where


$$\Delta_r = r^2 + a^2 - 2mr + q^2, \quad \Sigma = r^2 + a^2 \cos^2 \theta.$$

Here, m is the mass parameter, a and q are the rotation and electric charge parameters, respectively.


The thermodynamic quantities that can be evaluated via the



(a) The unit vector field for the five-dimensional singly rotating Kerr black hole with $\tau/r_0 = 20$ and $a/r_0 = 1$



(b) The unit vector field for the six-dimensional singly rotating Kerr black hole with $\tau/r_0 = 3\pi/2$ and $a/r_0 = 1$



(c) The unit vector field for the seven-dimensional singly rotating Kerr black hole with $\tau/r_0 = 16\pi/15$ and $a/r_0 = 1$

FIG. 4: The red arrows represent the unit vector field n on a portion of the $r_h - \Theta$ plane. The zero points (ZPs) marked with black dots are at $(r_h/r_0, \Theta) = (0.28, (\pi/2))$, $(3.14, (\pi/2))$, $(1, (\pi/2))$, and $(1, (\pi/2))$, for ZP_1 , ZP_2 , ZP_3 , and ZP_4 , respectively. The blue contours C_i are closed loops surrounding the zero points.




FIG. 5: Zero points of the vector ϕ^{r_h} shown on the $r_h - \tau$ plane for the Kerr-Newman black hole, where $a/r_0 = 1$ and $Q/r_0 = 1$. The black dot with $\tau_c = 6\pi\sqrt{3(a^2 + Q^2)}$ denotes the generation point for the Kerr-Newman black hole. At $\tau = \tau_c$, there is only one Kerr-Newman black hole, and at $\tau = \tau_1$, there are two Kerr-Newman black holes.

standard method as follows

$$\begin{aligned} M &= m, & J &= ma, & \Omega &= \frac{a}{r_h^2 + a^2}, \\ Q &= q, & \Phi &= \frac{qr}{r_h^2 + a^2}, \\ S &= \pi(r_h^2 + a^2), & T &= \frac{r_h^2 - a^2 - q^2}{4\pi r_h(r_h^2 + a^2)}, \end{aligned} \quad (20)$$

where $r_h = m \pm \sqrt{m^2 - a^2 - q^2}$ are the location of the outer and inner horizons.

Using the results of Eq. (20), one can directly obtain the generalized free energy of the Kerr-Newman black hole as

$$\mathcal{F} = M - \frac{S}{\tau} = \frac{r_h^2 + a^2 + Q^2}{2r_h} - \frac{\pi(r_h^2 + a^2)}{\tau}, \quad (21)$$

and the components of the vector ϕ can be computed as

$$\phi^{r_h} = 1 - \frac{r_h^2 + a^2 + Q^2}{2r_h^2} - \frac{2\pi r_h}{\tau}, \quad (22)$$

$$\phi^\Theta = -\cot\Theta \csc\Theta. \quad (23)$$

One can also obtain

$$\tau = \frac{4\pi r_h^3}{r_h^2 - a^2 - Q^2} \quad (24)$$

as the zero point of the vector field ϕ .

Similar to the procedure done in the last two sections, we show the the zero points of the component ϕ^{r_h} with $a = r_0$ and $Q = r_0$ in Fig. 5, and the unit vector field n with $\tau = 50r_0$, $a = r_0$, and $Q = r_0$ in Fig. 6. These two figures allow us to determine that the topological number of the four-dimensional Kerr-Newman black hole is $W = 0$, which is the same as the findings for four-dimensional Kerr black hole in Sec. II and the five-dimensional singly rotating Kerr black hole in




FIG. 6: The red arrows denote the unit vector field n on a portion of the $r_h - \Theta$ plane. The unit vector field for the Kerr-Newman black hole with $\tau/r_0 = 50$, $a/r_0 = 1$, and $Q/r_0 = 1$. The zero points (ZPs) marked with black dots are at $(r_h/r_0, \Theta) = (1.95, (\pi/2))$, and $(3, (\pi/2))$, for ZP_1 , and ZP_2 , respectively.

Sec. III. In addition, this also seems to imply that the electric charge parameter has no effect on the topological number of rotating black holes. This conclusion needs to be tested by investigating more topological numbers of the rotating charged black holes.

V. CONCLUSIONS

TABLE I: The topological number W , numbers of generation and annihilation points for black holes.

BH solution	W	Generation point	Annihilation point
Schwarzschild BH [10]	-1	0	0
$d \geq 6$ singly-rotating Kerr BH	-1	0	0
$d = 5$ singly-rotating Kerr BH	0	1	0
Kerr BH	0	1	0
Kerr-Newman BH	0	1	0
RN BH [10]	0	1	0
RN-AdS BH [10]	1	1 or 0	1 or 0

In this paper, we have investigate the topological numbers of the singly rotating Kerr black holes in arbitrary dimensions and the four-dimensional Kerr-Newman black hole. Considering our results and those in Ref. [10], Table I can be summarized. The $d \geq 6$ singly-rotating Kerr black holes and the Schwarzschild black hole belong to the first kind of topological classes since their topological number $W = -1$; the RN black hole, the Kerr-Newman black hole, and the $d = 4, 5$ singly-rotating Kerr black holes belong to the second kind of topological classes due to their topological number $W = 0$; the RN-AdS black hole belong to the third kind of topological classes since its topological number equals to 1. In this work, we come to two particularly exciting conclusions: that the ex-

istence of the rotation parameter has an important effect on the topological number of the uncharged black hole, as well as that the dimension of spacetime has an important effect on the topological numbers of the rotating black holes. In addition, the fact that the topological numbers of the four-dimensional Kerr and Kerr-Newman black holes are the same seems to suggest that the electric charge parameter has no effect on the topological number of rotating black holes. Our current work also support the conjecture proposed in Ref. [10] that all black hole solution should be classified into three different topological classes, at least in the pure Einstein-Maxwell gravity theory.

There are two promising further topics to be pursued in the future. One intriguing topic is to extend the present work to the more general cases with nonzero cosmological constant, and the black hole solutions in the supergravity and modified

gravity theories. And the other is to investigate the topological number of the black hole solutions with the planar [29], toroidal [30], hyperbolic [31], and spindle [32–36] horizons to explore whether there is a relation between topological number and horizon topology of the black holes. We hope to report the related progress along these two directions soon.

ACKNOWLEDGMENTS

We thank Prof. Shao-Wen Wei for very helpful discussions. This work is supported by the National Natural Science Foundation of China (NSFC) under Grant No. 12205243, No. 11675130, and by the Doctoral Research Initiation Project of China West Normal University under Grant No. 21E028.

-
- [1] The Event Horizon Telescope Collaboration, First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole, *Astrophys. J. Lett.* **875**, L1 (2019).
 - [2] The Event Horizon Telescope Collaboration, First Sagittarius A* Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole in the Center of the Milky Way, *Astrophys. J. Lett.* **930**, L12 (2022).
 - [3] B.P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), Observation of Gravitational Waves from a Binary Black Hole Merger, *Phys. Rev. Lett.* **116**, 061102 (2016).
 - [4] B.P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), GW151226: Observation of Gravitational Waves from a 22-Solar-Mass Binary Black Hole Coalescence, *Phys. Rev. Lett.* **116**, 241103 (2016).
 - [5] P.V.P. Cunha, E. Berti, and C.A.R. Herdeiro, Light Ring Stability in Ultra-Compact Objects, *Phys. Rev. Lett.* **119**, 251102 (2017).
 - [6] P.V.P. Cunha and C.A.R. Herdeiro, Stationary Black Holes and Light Rings, *Phys. Rev. Lett.* **124**, 181101 (2017).
 - [7] S.-W. Wei, Topological charge and black hole photon spheres *Phys. Rev. D* **102**, 064039 (2020).
 - [8] M. Guo and S. Gao, Universal properties of light rings for stationary axisymmetric spacetimes, *Phys. Rev. D* **103**, 104031 (2021).
 - [9] S.-W. Wei and Y.-X. Liu, Topology of equatorial timelike circular orbits around stationary black holes, [arXiv: 2207.08397](https://arxiv.org/abs/2207.08397).
 - [10] S.-W. Wei, Y.-X. Liu and R.B. Mann, Black Hole Solutions as Topological Thermodynamic Defects, *Phys. Rev. Lett.* **129**, 191101 (2022).
 - [11] S.-W. Wei and Y.-X. Liu, Topology of black hole thermodynamics, *Phys. Rev. D* **105**, 104003 (2022).
 - [12] P.K. Yerra and C. Bhamidipati, Topology of black hole thermodynamics in Gauss-Bonnet gravity, *Phys. Rev. D* **105**, 104053 (2022).
 - [13] M.B. Ahmed, D. Kubiznak, and R. B. Mann, Vortex/anti-vortex pair creation in black hole thermodynamics, [arXiv: 2207.02147](https://arxiv.org/abs/2207.02147).
 - [14] P.K. Yerra and C. Bhamidipati, Topology of Born-Infeld AdS black holes in 4D novel Einstein-Gauss-Bonnet gravity, [arXiv: 2207.10612](https://arxiv.org/abs/2207.10612).
 - [15] P.K. Yerra, C. Bhamidipati and S. Mukherji, Topology of critical points and Hawking-Page transition, *Phys. Rev. D* **106**, 064059 (2022).
 - [16] N.-C. Bai, L. Li, and J. Tao, Topology of black hole thermodynamics in Lovelock gravity, [arXiv: 2208.10177](https://arxiv.org/abs/2208.10177).
 - [17] C.H. Liu and J. Wang, The topological natures of the Gauss-Bonnet black hole in AdS space, [arXiv: 2211.05524](https://arxiv.org/abs/2211.05524).
 - [18] Z.-Y. Fan, Topological interpretation for phase transitions of black holes, [arXiv: 2211.12957](https://arxiv.org/abs/2211.12957).
 - [19] C.X. Fang, J. Jiang, and M. Zhang, Revisiting thermodynamic topologies of black holes, [arXiv: 2211.15534](https://arxiv.org/abs/2211.15534).
 - [20] R.P. Kerr, Gravitational field of a spinning mass as an example of algebraically special metrics, *Phys. Rev. Lett.* **11**, 237 (1963).
 - [21] Y.-S. Duan and M.-L. Ge, $SU(2)$ gauge theory and electro-dynamics of N moving magnetic monopoles, *Sci. Sin.* **9**, 1072 (1979).
 - [22] Y.-S. Duan, The structure of the topological current, Report No. SLAC-PUB-3301, 1984.
 - [23] Y.-S. Duan, S. Li, and G.-H. Yang, The bifurcation theory of the Gauss-Bonnet-Chern topological current and Morse function, *Nucl. Phys. B* **514**, 705 (1998).
 - [24] L.-B. Fu, Y.-S. Duan, and H. Zhang, Evolution of the Chern-Simons vortices, *Phys. Rev. D* **61**, 045004 (2000).
 - [25] R.C. Myers and M.J. Perry, Black holes in higher dimensional space-times, *Annals Phys.* **172**, 304 (1986).
 - [26] S.-W. Wei, P. Cheng, and Y.-X. Liu, Analytical and exact critical phenomena of d -dimensional singly spinning Kerr-AdS black holes, *Phys. Rev. D* **93**, 084015 (2016).
 - [27] E.T. Newman and A.I. Janis, Note on the Kerr spinning particle metric, *J. Math. Phys.* **6**, 915 (1965).
 - [28] E.T. Newman, E. Couch, K. Chinnapared, A. Exton, A. Prakash, R. Torrence, Metric of a rotating, charged mass, *J. Math. Phys.* **6**, 918 (1965).
 - [29] R.G. Cai and Y.Z. Zhang, Black plane solutions in four-dimensional space-times, *Phys. Rev. D* **54**, 4891 (1996).
 - [30] D.R. Brill and J. Louko, Thermodynamics of (3+1)-dimensional black holes with toroidal or higher genus horizons, *Phys. Rev. D* **56**, 3600 (1997).
 - [31] Y. Chen, Y.K. Lim, and E. Teo, Deformed hyperbolic black holes, *Phys. Rev. D* **92**, 044058 (2015).
 - [32] D. Klemm, Four-dimensional black holes with unusual horizons, *Phys. Rev. D* **89**, 084007 (2014).
 - [33] R.A. Hennigar, R.B. Mann, and D. Kubizňák, Entropy Inequality Violations from Ultraspinning Black Holes, *Phys. Rev.*

- [Lett. **115**, 031101 \(2015\).](#)
- [34] D. Wu, P. Wu, H. Yu, and S.-Q. Wu, Are ultraspinning Kerr-Sen-AdS₄ black holes always superentropic?, [Phys. Rev. D **102**, 044007 \(2020\).](#)
- [35] D. Wu, S.-Q. Wu, P. Wu, and H. Yu, Aspects of the dyonic Kerr-Sen-AdS₄ black hole and its ultraspinning version, [Phys. Rev. D **103**, 044014 \(2021\).](#)
- [36] D. Wu and S.-Q. Wu, Ultraspinning Chow's black holes in six-dimensional gauged supergravity and their thermodynamical properties, [J. High Energy Phys. **11** \(2021\) 031.](#)