Provably Efficient Model-free RL in Leader-Follower MDP with Linear Function Approximation

Arnob Ghosh

Electrical and Computer Engg., The Ohio State University

GHOSH.244@OSU.EDU

Abstract

We consider a multi-agent episodic MDP setup where an agent (leader) takes action at each step of the episode followed by another agent (follower). The state evolution and rewards depend on the joint action pair of the leader and the follower. Such type of interactions can find applications in many domains such as smart grids, mechanism design, security, and policymaking. We are interested in how to learn policies for both the players with provable performance guarantee under a bandit feedback setting. We focus on a setup where both the leader and followers are non-myopic, i.e., they both seek to maximize their rewards over the entire episode and consider a linear MDP which can model continuous state-space which is very common in many RL applications. We propose a *model-free* RL algorithm and show that $\tilde{O}(\sqrt{d^3H^3T})$ regret bounds can be achieved for both the leader and the follower, where d is the dimension of the feature mapping, H is the length of the episode, and T is the total number of steps under the bandit feedback information setup. Thus, our result holds even when the number of states becomes infinite. The algorithm relies on novel adaptation of the LSVI-UCB algorithm. Specifically, we replace the standard greedy policy (as the best response) with the soft-max policy for both the leader and the follower. This turns out to be key in establishing uniform concentration bound for the value functions. To the best of our knowledge. this is the first sub-linear regret bound guarantee for the Markov games with non-myopic followers with function approximation.

1. Introduction

Multi-agent Reinforcement Learning (MARL) has become important tool for decision making in a Markov game involving multiple agents. In many real-world problems, agents often have asymmetric roles. For example, one agent (leader) can act first and observing the action of the leader, the other agent (follower) reacts at each step of the MDP. This type of interaction requires two levels of thinking: the leader must reason what the follower would do in order to find its optimal decision. For example, an electric utility company (the *leader*) seeks to maximize the social welfare by selecting prices at different times over a day while the users (the *followers*) seek to optimize their own consumption based on the prices set by the utility company. Here, the sequential decision making process can be modeled as MDP where at each period the utility company sets its price first, then, the users decide their consumption. The reward and the underlying transition probability depend on both the leader's and the follower's action. Such leader-follower interactions appear in other applications as well such as in AI Economist Zheng et al. (2020), Mechanism Design Conitzer and Sandholm (2004), optimal auction Cole and Roughgarden (2014), and security games Tambe (2011).

Such kind of leader-follower interaction is different from the simultaneous play in the Markov setting as considered in Jin et al. (2021b); Tian et al. (2021). In general, the Stackelberg equilibrium

is the relevant concept for this type of leader-follower interaction Conitzer and Sandholm (2006) compared to the Nash equilibrium considered in the above paper. Letchford et al. (2009); Peng et al. (2019) considered a learning framework in order to learn the Stackelberg equilibrium with a best-response oracle for the follower. However, these works can not be generalized to the *bandit feedback* setting where the leader and the follower are unaware of the transition probabilities and the rewards, and can only observe those rewards corresponding to the state-actions pairs encountered. Efficient learning in this leader-follower MDP setting under the bandit feedback (which is more natural) is fundamentally more challenging compared to the single agent setting due to the more challenging exploration-exploitation trade-off.

Few recent works have focused on such kind of leader-follower interaction in MDP with the bandit feedback setting. Bai et al. (2021) considered a model where the leader selects the underlying MDP on which the follower acts on, i.e., the leader only acts once at the start of the episode. However, we consider the setup where both the leader and the follower interact at every step of the episode. Kao et al. (2022) considers a setup where both the leader and the follower interact at every step similar to ours. However, Kao et al. (2022) considered the setup where both the leader and the follower receive the same reward whereas in our setting the rewards can be different. Further, both the above papers consider the finite state-space (a.k.a. tabular setup) where the sample complexity scales with the state space. Thus, the above approach would not be useful for large-scale RL applications where the number of states could even be infinite. To address this curse of dimensionality, modern RL has adopted *function approximation* techniques to approximate the (action-) value function or a policy, which greatly expands the potential reach of RL, especially via deep neural networks. For large state-space the model-based approaches as considered in the above papers have limited application Wei et al. (2020), thus, we focus on developing model-free algorithm. Only Zhong et al. (2021) considers a leader-follower Markov setup with function approximation. However, they consider *myopic* followers which seek to maximize instantaneous reward and also consider the followers' reward are known. Hence, it greatly alleviates the exploration challenge as it is only limited to the leader's side. Rather, we consider the setup where the followers are also non-myopic with bandit feedback. Thus, we seek to answer the following question

Can we achieve provably optimal regret for model-free exploration for leader-follower (non-myopic) interaction in MDP with function approximation under bandit feedback?

Our Contribution: To answer the above question, we consider the Markov game with linear function approximation (bandit feedback) where at each step of the sequential decision process the leader takes an action and observing the action the follower reacts. The transition probability and the reward functions can be represented as a *linear function* of some known feature mapping adapted from the single agent set up in Jin et al. (2020). Our main contributions are:

- We show that with a proper parameter choice, our proposed model-free algorithm achieves $\tilde{O}(\sqrt{d^3H^3T})$ regret for both leader and follower, where d is the dimension of the feature mapping, H is the length of the episode, and T is the total number of steps. Note that for the single agent setup, the regret is of the same order achieved in Jin et al. (2020). Hence, our result matches the regret bound for the single agent setup.
- Our bounds are attained without explicitly estimating the unknown transition model or *requiring any simulator or best-response oracle*, and they depend on the state space only through the dimension of the feature mapping. To the best of knowledge, these sub-linear regret bounds are

the first results for leader-follower non-myopic MDP game with function approximations under bandit feedback.

- Since linear MDP contains tabular setup, as a by-product, we provide the first result on the regret bound for the leader-follower (non myopic) MDP game under bandit feedback using model-free RL algorithm.
- We adapt the classic model-free LSVI-UCB algorithm proposed in Jin et al. (2020) in a novel manner. Due to the nature of the leader-follower interaction, a key challenge arises while establishing the value-aware uniform concentration, which lies at the heart of the performance analysis of model-free exploration. In particular, for a single agent set-up the greedy selection with respect to the standard *Q*-function achieves a small covering number. However, in the game setting, for a given policy, the best response strategy of a player fails to achieve such non-trivial covering number for the value-function class of the players (i.e., *V*-function). To address this fundamental issue, we instead adopt a soft-max policy for the players by utilizing its nice property of approximation-smoothness trade-off via its parameter, i.e., temperature coefficient.

Related Literature: Provably efficient RL algorithms for zero-sum Markov games have been proposed Wei et al. (2017); Bai et al. (2020); Liu et al. (2021); Xie et al. (2020); Chen et al. (2021); Sayin et al. (2021). Provably efficient algorithms to obtain coarse correlated equilibrium have also been proposed for the general sum-game as well Bai and Jin (2020); Jin et al. (2021b); Mao and Başar (2022). In contrast to the above papers, we consider a leader-follower setup. Hence, our work is not directly comparable.

For learning Stackelberg equilibrium, most of the works focus on the normal form game which is equivalent to step size H = 1 in our setting Balcan et al. (2015); Blum et al. (2014); Peng et al. (2019); Letchford et al. (2009). Further, in the above papers, it is assumed that the followers' responses are known, in contrast, in our setting the follower is also learning its optimal policy. Zhong et al. (2021) considered a leader-follower setup with myopic follower and known reward for linear MDP which alleviates the challenges of exploration for the follower. Zhong et al. (2021) also proposed a model-based approach for tabular setup with myopic follower and unknown reward. In contrast, we consider linear MDP and non-myopic follower with bandit feedback, and proposed model-free RL algorithm. Recently, Kao et al. (2022) proposed a decentralized cooperative RL algorithm with hierarchical information structure for a tabular set-up. In contrast, we consider linear function approximation setup. In our model, the rewards of the leader and the follower can be different unlike in Kao et al. (2022). Bai et al. (2021) provides sample complexity guarantee for Stackelberg equilibrium for bandit-RL game where leader only takes action at the start of the episode which is quite different from our setup. Zheng et al. (2022) modeled the actor-critic framework as a Stackelberg game which is quite different from our framework.

2. Leader-Follower MDP Game

The Model: We consider an episodic MDP with the tuple $(S, A, B, P, H, \mathbf{R})$ where S is the state space, A is the action space for leader, and B is the action space for followers. Each MDP starts from the state x_1 . At every step h, observing the state x_h , the leader first takes an action $a_h \in A$, then the follower takes an action $b_h \in B$ observing the action of the leader and the state x_h . The state transitions to $x_{h+1} \in S$ depending on x_h, a_h , and b_h . The process continues for H steps. The

transition probability kernel is defined as the following, $\mathcal{P} = \{\mathbb{P}_h\}_{h=1}^H \mathbb{P}_h : \mathcal{S} \times \mathcal{A} \times \mathcal{B} \to \mathcal{S}$. The reward vector for leader and follower are defined as $\{r_{l,h}\}_{h=1}^H$ and $\{r_{f,h}\}_{h=1}^H$ respectively. The reward for agent $m = l, f r_{m,h}(x_h, a_h, b_h)$ denotes the reward received by agent m when the leader selects the action a_h and the follower selects the action b_h at step h.

Several important points on the model should be noted here. This setup is also known as hierarchical MDP Kao et al. (2022). Our setup can model various real-world applications. For example, consider a dynamic electricity market where a social planner sets a price at every hour of a day. Observing the price at current period, the users decide how much to consume at that period. Here, the state can represent the demand of the user, and the supply. The user's objective is to maximize its total utility over the day whereas the utility company's objective is to maximize the social welfare. Note that such kind of leader-follower decision hierarchy can also occur in MARL when one agent has some advantage (e.g., can compute decision faster compared to the other agent) over the other agent.

Throughout this paper, we consider deterministic (unknown) reward. Without loss of generality, we also assume $|r_{m,h}| \le 1$ for all m and h. Our model can be easily extended to the setup where the rewards are random yet bounded. Our model can also be extended to the setup where the initial state of each episodic MDP is drawn from a distribution.

Note that we consider the state x as a joint state of both the leader and the follower. Our work can be extended to the setup where the leader and the follower's states are decoupled with the underlying assumption that the leader can observe the follower's state.

Policy: The agents interact repeatedly over K episodes. The policy of the leader at step $h \in [H]$ at episode $k \in [K]$ is $\pi_{l,h}^k(a|x_h^k)$ that denotes the probability with which action $a \in \mathcal{A}$ is taken at step h at episode k when the state is x_h^k . The policy for the follower at step $h \in [H]$ at episode [K] is $\pi_{f,h}^k(b|x_h, a_h^k)$ that denotes the probability with action $b \in \mathcal{B}$ is chosen by the follower at state x_h at the h-th step of episode k and when the leader's action is a_h^k . Note the difference with the simultaneous play, here, the follower's policy is a function of the leader's action at step h whereas in the simultaneous game, it is independent of the other players' actions. Let $\pi_l = {\pi_{l,h}}_{h=1}^{H}$, and $\pi_f = {\pi_{f,h}}_{h=1}^{H}$ be the collection of the policies of leader and follower respectively across the episode.

Q-function and Value function The joint state-action value function for any player m, for m = l, f at step h is

$$Q_{m,h}^{\pi_l,\pi_f}(x, a, b) = \mathbb{E}\left[\sum_{i=h}^{H} r_{m,i}(x_i, a_i, b_i) | x_h = x, a_h = a, b_h = b\right]$$

Here, the expectation is taken over the transition probability kernel, and the policies of both the leader and the follower. We can also define a *marginal Q*-function for the leader as

$$q_{l,h}^{\pi_l,\pi_f}(x,a) = \sum_b \pi_{f,h}(b|x,a) Q_{l,h}^{\pi_l,\pi_f}(x,a,b).$$
(1)

The above denotes the expected cumulative reward starting from step h after playing action a, and then following the policy π_l (from step h + 1) while the follower following the policy π_f from step h. We later show that marginal q plays an important role in decision making.

For compactness of operator, we introduce the following notations

$$\mathbb{D}^{\pi_l,\pi_f}[Q](x) = \mathbb{E}_{a \sim \pi_l(\cdot|x), b \sim \pi_f(\cdot|x,a)} Q(x,a,b),$$
$$\mathbb{D}^{\pi_f}[Q](x,a) = \mathbb{E}_{b \sim \pi_f(\cdot|x,a)} Q(x,a,b),$$
$$\mathbb{P}_h V(x,a,b) = \mathbb{E}_{x' \sim \mathbb{P}_h(\cdot|x,a,b)} V(x').$$

The value function or expected cumulative reward starting from step h for the leader is defined as

$$V_{l,h}^{\pi_{l},\pi_{f}}(x) = \mathbb{D}^{\pi_{l,h},\pi_{f,h}} Q_{l,h}^{\pi_{l},\pi_{f}}(x)$$
(2)

here the expectation is first taken over the follower's policy for a given leader's action then expectation is taken over the leader's action.

We now consider the follower's value functions. The leader's action dependent value function for the follower at step h when the leader takes action a at step h is given by

$$\bar{V}_{f,h}^{\pi_l,\pi_f}(x,a) = \mathbb{D}^{\pi_{f,h}} Q_{f,h}^{\pi_l,\pi_f}(x,a,b)$$
(3)

The above denotes the expected cumulative reward the follower would get from step h after observing the leader's action at h, and then following its own policy at step h.

We also define the following value function for the follower by taking the expectation over a on \bar{V}

$$V_{f,h}^{\pi_l,\pi_f}(x) = \mathbb{D}^{\pi_{l,h},\pi_{f,h}} Q_{f,h}^{\pi_l,\pi_f}(x,a,b)$$
(4)

Bellman's Equation: We now describe the Bellman's equations for a given state and the joint action pair

$$Q_{f,h}^{\pi_f}(x,a,b) = r_{f,h}(x,a,b) + \mathbb{P}_h V_{f,h+1}^{\pi_l,\pi_f}(x,a,b)$$
(5)

Similarly, we have the following relationship

$$Q_{l,h}^{\pi_f}(x,a,b) = r_{l,h}(x,a,b) + \mathbb{P}_h V_{l,h+1}^{\pi_l,\pi_f}(x,a,b)$$
(6)

Information Structure: We assume the following information structure-

Assumption 1 The transition probability kernel, and rewards are unknown to both the leader and the follower. The leader and follower observe the rewards of each other only for the encountered state-action pairs (*i.e.*, bandit feedback).

Note that the agent can not access the rewards for the states and action which have not encountered before. This is also known as the *bandit-feedback* setting Bai et al. (2021). Thus, both the agents need to employ an exploratory policy. Also, note that we consider an information structure where the leader and follower can observe each other's reward. Extending our analysis to the setup where the leader does not observe the action and/or reward of the follower constitutes a future research direction.

Objective: After observing the action of the leader, the follower seeks to optimize its own leader's action dependent value function: $\max_{\pi_f} \bar{V}_{f,1}^{\pi_l,\pi_f}(x_1,a)$

Given the policy of the follower, the leader seeks to optimize its own value function

$$\max_{\pi_l} V_{l,1}^{\pi_l,\pi_f}(x_1), \quad \pi_l^* = \arg\max_{\pi^l} V_{l,1}^{\pi_l,\pi_f}(x_1)$$

Now, we describe the relationship with Q-function and the value functions for the optimal policies of both the leader and the follower which will also specify how to select optimal policy at every step. For the follower, we have Bellman's optimality equation–

$$Q_{f,h}^{\pi_l,\pi_f^*}(x,a,b) = r_{f,h}(x,a,b) + \mathbb{P}_h V_{f,h+1}^{\pi_l,\pi_f^*}(x,a,b)$$
(7)

Hence, given the action a of the leader at step h, the follower's policy is $\pi_{f,h}^*(b|x,a) = 1$ where $b = \max_b Q_{f,h}^{\pi_l,\pi_f}(x,a,b)$. Thus, the optimal policy is greedy with respect to the joint state-action Q function. The follower's policy at step h depends on the leader's action at step h, the leader's policy and the follower's own policy starting from h + 1.

The optimal policy $\pi_{l,h}^*$ for the leader if the follower selects the policy π_f is given by

$$V_{l,h}^{\pi_l^*,\pi_f}(x_h) = \max_a q_{l,h}^{\pi_l^*,\pi_f}(x_h,a)$$
(8)

Hence, the optimal policy for the leader is greedy with respect to its marginal Q-function $q_{l,h}^{\pi_l^*,\pi_f}(x_h,\cdot)$. The optimal policy at step h depends on the follower's policy starting at step h, and the leader's policy starting from step h + 1.

If the Q-functions are known, the optimal policy for the leader and follower are obtained using backward induction. At step H, the follower's best response for every leader's action is computed ((as in (7). Then, leader's best response is computed based on the marginal q function (8). Once the policy is computed for the step H, the policy for step H - 1 is computed in the similar manner and so on.

Learning Metric: Since the leader and the follower are unaware of the rewards and the transition probability, selecting optimal policy from the start is difficult. Rather, they seek to learn policies with good performance guarantee. Thus, instead of finding the Stackelberg equilibrium, we consider the following learning metric

Definition 1 Regret for the leader is defined as

$$\operatorname{Regret}_{\ell}(K) = \sum_{k=1}^{K} (V_{l,1}^{\pi_{l}^{k,*},\pi_{f}^{k}}(x_{1}) - V_{1}^{\pi_{l}^{k},\pi_{f}^{k}}(x_{1}))$$
(9)

where $\pi_l^{k,*}$ is the optimal policy for the leader when the follower plays the policy π_f^k .

Note that the regret for the leader defines the optimality gap between the policies employed by the leader and the best response policy (in hindsight) of the leader given the follower's policy at an episode. Note that the follower's policy is unknown, rather, the leader needs to reason about the follower's policy at episode k. Also note that the leader's regret measures the gap between the policy π_l^k and the best policy $\pi_l^{k,*}$ of the leader in response to the follower's policy at episode k, rather than the optimal policy of the follower. This is because the follower is also learning its optimal policy, hence, the policy π_f^k would not be optimal. Thus, the regret considers how good the leader is doing compared to the policy employed by the follower across the episodes. Obviously, at different episodes, the optimal policy of the leader may be different.

Definition 2 Regret for the follower is defined as

$$\operatorname{Regret}_{f} = \sum_{k=1}^{K} (\bar{V}_{f,1}^{\pi_{l},\pi_{f}^{*}}(x_{1},a_{1}^{k}) - \bar{V}_{f,1}^{\pi_{l},\pi_{f}^{k}}(x_{1},a_{1}^{k}))$$
(10)

where π_f^* denotes the optimal policy for the follower and a_1^k is the action of the leader at the first step of episode k.

The regret for the follower captures how good the follower's policy is compared to the optimal policy for a given initial action and the policy of the leader. Even though the follower knows the initial action of the leader, it is unaware of the leader's policy. Rather, the follower also needs to reason about the leader's policy starting from step 2. Both the leader and the follower seek to minimize their respective regrets. Initial action specific regret is unique to the leader-follower setup.

Note that Zhong et al. (2021) considers the setup where the leader takes action at every step of the MDP whereas the followers are myopic. They also consider that the rewards are known, thus, one can compute the best response of the follower at every step. Thus, the regret for the follower does not arise there. Kao et al. (2022) considered the reward is the same, thus, they consider the regret of the joint policy rather than the individual agent's regret.

In general, achieving sub-linear regret in multi-agent RL setup is more challenging compared to the single agent setup since the underlying environment of an agent may change depending on the other agent's policy Tian et al. (2021); Bai et al. (2021). Nevertheless, we obtain sub-linear regret for both the leader and follower.

Leader-follower Linear MDP: We consider a linear MDP set-up in order to handle large state-space.

Assumption 2 We consider a linear MDP with the known (to both the players) feature map ϕ : $S \times A \times B \rightarrow \mathbb{R}^d$, if for any h, there exists d unknown signed measures $\mu_h = {\mu_h^1, \ldots, \mu_h^d}$ over S such that for any $(x, a, x') \in S \times A_1 \times B$,

$$\mathbb{P}_h(x'|x,a,b) = \langle \phi(x,a,b), \mu_h(x') \rangle$$

and there exists vectors $\theta_{l,h}$, $\theta_{f,h} \in \mathbb{R}^d$ such that for any $(x, a, b) \in \mathcal{S} \times \mathcal{A}_1 \times \mathcal{A}_2$,

$$r_{l,h}(x,a,b) = \langle \phi(x,a,b), \theta_{l,h} \rangle \quad r_{f,h}(x,a) = \langle \phi(x,a,b), \theta_{f,h} \rangle$$

Note that such a linear MDP setup is considered for single agent scenario Jin et al. (2020); Yang and Wang (2019). Examples of linear MDP includes tabular setup.

For the leader-follower linear MDP setup, we have-

Lemma 1
$$Q_{m,h}^{\pi_l,\pi_f}(x,a,b) = \langle \phi(x,a,b), w_{m,h}^{\pi_l,\pi_f} \rangle \ \forall m.$$

Thus, the Q-functions of both the leader and follower are linear in the feature space. We can thus search over w in order to find the optimal Q-function.

Algorithm 1	l Leader's	Model Free	RL Algorithm
-------------	------------	------------	--------------

 $w_{l,h} = 0, w_{f,h} = 0, \alpha_f = \frac{\log(|\mathcal{B}|)\sqrt{K}}{H}, \alpha_l = \frac{\log(|\mathcal{A}|)\sqrt{K}}{H}, \beta =$ 1: Initialization: $C_1 dH \sqrt{\log(4(\log(|\mathcal{B}||\mathcal{A}|) + 2\log(|\mathcal{B}|)\log(|\mathcal{A}|))} dT_{\mathcal{A}}$ 2: for episodes $k = 1, \ldots, K$ do Receive the initial state x_1^k . 3: for step h = H, H - 1, ..., 1 do 4:
$$\begin{split} & \Lambda_{h}^{k} \leftarrow \sum_{\tau=1}^{k-1} \phi(x_{h}^{\tau}, a_{h}^{\tau}, b_{h}^{\tau}) \phi(x_{h}^{\tau}, a_{h}^{\tau}, b_{h}^{\tau})^{T} + \lambda \mathbf{I} \\ & w_{m,h}^{k} \leftarrow (\Lambda_{h}^{k})^{-1} [\sum_{\tau=1}^{k-1} \phi(x_{h}^{\tau}, a_{h}^{\tau}, b_{h}^{\tau}) [r_{m,h}(x_{h}^{\tau}, a^{\tau}, b_{h}^{\tau}) + V_{m,h+1}^{k}(x_{h+1}^{\tau})]] \\ & Q_{m,h}^{k}(\cdot, \cdot, \cdot) \leftarrow \min\{\langle w_{m,h}^{k}, \phi(\cdot, \cdot, \cdot) \rangle + \beta(\phi(\cdot, \cdot, \cdot)^{T} (\Lambda_{h}^{k})^{-1} \phi(\cdot, \cdot, \cdot))^{1/2}, H\} \end{split}$$
5: 6: 7: for $a \in \mathcal{A}$ do 8: $\pi_{f,h}^{k}(b|\cdot,a) = \frac{\exp(\alpha_{f}(Q_{f,h}^{k}(\cdot,a,b)))}{\sum_{b_{m}}\exp(\alpha_{f}(Q_{f,h}^{k}(\cdot,a,b_{m})))}$ 9: $q_{l,h}^{k}(\cdot,\cdot) \leftarrow \langle \pi_{f,h}^{k}(\cdot|\cdot,\cdot), Q_{l,h}^{k}(\cdot,\cdot,\cdot) \rangle$ 10: $\bar{V}_{f,h}^{k}(\cdot,\cdot) \leftarrow \langle \pi_{f,h}^{k}(\cdot|\cdot,\cdot), Q_{f,h}^{k}(\cdot,\cdot,\cdot) \rangle$ 11: $\pi_{l,h}^{k}(a|\cdot) = \frac{\exp(\alpha_{l}(q_{l,h}^{k}(\cdot,a)))}{\sum_{a_{m}}\exp(\alpha_{l}(q_{l,h}^{k}(\cdot,a_{m})))}$ 12:
$$\begin{split} V_{l,h}^k(\cdot) &= \langle \pi_{l,h}^k, q_{l,h}^k(\cdot, \cdot) \rangle \\ V_{f,h}^k(\cdot) &= \langle \pi_{l,h}^k, \bar{V}_{f,h}^k(\cdot, \cdot) \rangle \\ \text{for } h = 1, \dots, H \text{ do} \end{split}$$
13: 14: 15: for $a \in \mathcal{A}$ do 16: Compute $Q_{f,h}^k(x_h^k, a, b)$, $Q_{l,h}^k(x_h^k, a, b)$ for all b. 17: Compute policy $\pi_{f,h}^k(b|x_h^k, a)$ according to the Soft-max for $Q_{f,h}^k(x_h^k, a, \cdot)$ with param-18: $q_{l,h}(x_1^k, a) = \sum_b \pi_{f,h}^k(b|x_h^k, a) Q_{l,h}^k(x_h^k, a, b)$ 19: The leader takes action a_h^k according to Soft-max policy with respect to $q_{l,h}^k(x_h^k, \cdot)$ with 20: parameter α_l . The follower takes an action b_h^k (Algorithm 2) and observe $x_{h+1}^k \sim \mathbb{P}_h(x_h^k, a_h^k, b_h^k)$ 21:

Algorithm 2 Follower's Model Free RL Algorithm

1: Execute steps 1-14 of Algorithm 1.

- 2: **for** step h = 1, ..., H **do**
- 3:
- 4:
- Observe the action a_h^k of the leader. Compute $Q_{f,h}^k(x_h^k, a_h^k, b), \pi_{f,h}^k(b|x_h^k, a^k)$ for all b based on $w_{f,h}^k$. The follower takes action $b_h^k \sim \pi_{f,h}^k(\cdot|x_h^k, a_h^k)$ and observe x_{h+1}^k . 5:

3. Proposed Algorithm

We now describe our proposed algorithms for the leader (Algorithm 1) and the follower (Algorithm 2). The algorithm is based on the LSVI-UCB Jin et al. (2020) with some subtle differences which we will point out along our description.

We first describe the leader's algorithm. Note that in order to obtain its policy, the leader also needs to reason the policy the follower would play. Hence, the leader's algorithm also consists of how the follower selects its policy. The first part (steps 5-6) consists of updating the parameters $\Lambda_h^k, w_{l,h}^k, w_{f,h}^k$ which are used to update the joint state-action value functions $Q_{m,h}^k$ and value functions $V_{m,h}^k$ for m = l, f. Note that the Steps 7-14 are not evaluated for each state, rather, they are evaluated only for the encountered states till episode k - 1. Hence, we do not need to iterate over potentially infinite number of states. $V_{m,H+1}^k = 0$ for all k.

We now discuss the rationale behind updating $w_{m,h}^k$. We seek to obtain w such that it approximates the Q-function since the Q-function is inner product of w and ϕ (Lemma 1). Thus, we parameterize $Q_{m,h}^{\pi_l,\pi_f}(\cdot,\cdot,\cdot)$ by a linear form $\langle w_{m,h}^k,\phi(\cdot,\cdot,\cdot)\rangle$. The intuition is to obtain $w_{m,h}^k$ from the Bellman's equation using the regularized least-square regression. However, there are challenges. We do not know \mathbb{P}_h in Bellman's equations ((5), and (6)) rather $\mathbb{P}_h V_{m,h+1}^{\pi_l,\pi_f}$ should be replaced by the empirical samples. We obtain $w_{m,h}^k$ for m = l, f by solving the following regularized least-square problem

$$w_{m,h}^{k} \leftarrow \arg\min_{w \in \mathbb{R}^{d}} \sum_{\tau=1}^{k-1} [r_{m,h}(x_{h}^{\tau}, a_{h}^{\tau}, b_{h}^{\tau}) + V_{m,h+1}^{k}(x_{h+1}^{\tau}) - w^{T}\phi(x_{h}^{\tau}, a_{h}^{\tau})]^{2} + \lambda ||w||_{2}^{2}$$
(11)

where $V_{m,h+1}^k$ is the estimate of the value function $V_{m,h+1}$. After we obtain $w_{m,h}^k$, we add an additional bonus term $\beta(\phi(\cdot,\cdot,\cdot)^T(\Lambda_h^k)^{-1}\phi(\cdot,\cdot,\cdot))^{1/2}$ similar to Jin et al. (2020) to obtain $Q_{m,h}^k$. β is constant which we will characterize in the next section. Λ_h^k is the Gram matrix for the regularized least square problem. Such an additional term is used for upper confidence bound in LSVI-UCB Jin et al. (2020) as well. The same additional term is used for both $Q_{l,h}^k$ and $Q_{f,h}^k$. This bonus term would ensure the exploration for both the leader and the follower.

Now, we describe how we estimate the value function $V_{m,h+1}^k$ function which we use in (11). In order to update the value function, we need to compute the policy for the follower and the leader (cf.(2)). Unlike LSVI-UCB, we use the soft-max policy for both the leader and follower. Soft-max policy SOFT-MAX_{α}(**X**) = {SOFT-MAX_{α}^{*i*}(**X**)} $_{i=1}^{|\mathcal{L}|}$ for any vector **X** $\in \mathbb{R}^{|\mathcal{L}|}$ is a vector with the same dimension as in **X** with parameter α where the *i*-th component

SOFT-MAX^{*i*}_{$$\alpha$$}(**X**) = $\frac{\exp(\alpha X_i)}{\sum_{n=1}^{|\mathcal{L}|} \exp(\alpha X_n)}$ (12)

In order to estimate value function, first, one needs to compute follower's policy at a given leader's action, and subsequently, the leader's policy needs to computed (cf.(2) & (4)). At step h, for every leader's action a, $\pi_{f,h,k}(b|x_h^{\tau}, a)$ is computed based on the soft-max policy on the estimated Q-function for the follower $Q_{f,h}^k(x_h^{\tau}, a, b)$ at step 9. Based on the follower's policy, the leader updates its marginal Q function q_l at step 10 (cf.(1)) and the follower's leader's action dependent value function $\bar{V}_{f,h}^k$ (cf.(3)) at step 11. Now, the leader computes its policy based on $q_{l,h}^k(\cdot, \cdot)$. Finally, We update the leader's and follower's value function based on the leader's policy at steps 13 and 14.

Note that when $\alpha_l = \alpha_f = \infty$, the policy of the follower and leader become equal to the greedy policy. The greedy-policy is optimal for leader with respect to its marginal q-function (Eq.(8)) and for the follower with respect to its joint Q-function (cf.(7)). However, we can not obtain optimal regret for the leader and follower with the greedy policy unlike the single agent scenario.

The last part consists of execution of the policy. In order to find its optimal policy, the leader computes $Q_{f,h}^k$ for each action of the leader *a* (Line 17), based on the already computed $w_{f,h}^k$. Once

 $Q_{f,h}^k$ is computed, the follower's policy is also computed based on the soft-max function (Step 18). Once the follower's policy is computed, the marginal Q-function for the leader $q_{l,h}^k(x_h^k, \cdot)$ is computed. The leader then takes an action a_h^k based on the soft-max policy with respect to $q_{l,1}^k$. The follower takes an action b_h^k by observing the action a_h^k which we will describe next.

As mentioned before, the steps of the follower (Algorithm 2) are already contained in the leader's algorithm. The follower also obtains its w by solving (11). Hence, the leader and the follower have the same updates on w. The only difference is the execution as the follower executes its action based on the action taken by the leader at every step. At step 4 of Algorithm 2, the follower computes Q-function based on the current state x_h^k and action a_h^k of the leader. The follower then chooses its action based on the soft-max policy on the Q-function.

The space and time complexities of Algorithms 1 and 2 are of the same order as the LSVI-UCB. To be precise, the space complexity is $\mathcal{O}(d^2H + d|\mathcal{A}||\mathcal{B}|T)$. When we compute $(\Lambda_h^k)^{-1}$ using Sherman-Morrison formula, the computation of $V_{m,h+1}^k$ is dominated by computing $Q_{m,h+1}^k$ and the policy π_l^k , and π_f^k . Hence, it takes $\mathcal{O}(d^2|\mathcal{A}||\mathcal{B}|T)$ time.

4. Analysis

4.1. Main Results

Theorem 3 Fix p > 0. If we set $\beta = C_1 dH \sqrt{\iota}$ in Algorithm 1 and Algorithm 2 where $\iota = \log((\log(|\mathcal{A}||\mathcal{B}|) + 2\log(|\mathcal{A}|)\log(|\mathcal{B}|))4dT/p)$ for some absolute constant C_1 , then with probability (1-p),

$$\begin{aligned} \operatorname{Regret}_{l}(K) &\leq C\sqrt{d^{3}H^{3}T\iota^{2}}\\ \operatorname{Regret}_{f}(K) &\leq C'\sqrt{d^{3}H^{3}T\iota^{2}} \end{aligned}$$

where T = KH for some absolute constants C, and C'.

The result shows that regret for both the leader and the follower scale with $\tilde{\mathcal{O}}(\sqrt{d^3H^3T})$. Note that for the single-agent scenario Jin et al. (2020), the order of the regret is the same. However, compared to Jin et al. (2020), there is an additional multiplicative $\log(|\mathcal{A}|)$ and $\log(|\mathcal{B}|)$ factor in the value of ι which arises because we use soft-max policy for the leader and the follower instead of the greedy policy. The regret bounds do not depend on the dimension of the state space, rather, it depends on the dimension of the feature space d. If there is no follower, we set $|\mathcal{B}|= 1$, and can achieve single agent's regret for the soft-max policy as well. To the best of our knowledge this is the first result which shows $\tilde{\mathcal{O}}(\sqrt{T})$ regret for both the leader and the follower in the model-free with function approximation.

We assume that the leader observes the follower's action (this is known as informed game Tian et al. (2021)). Under uninformed setting (where a player may not observe the action of other player), it is statistically hard to obtain sub-linear regret even in zero-sum game Tian et al. (2021). Thus, Tian et al. (2021) shows a sub-linear regret under a weaker notion of regret. It remains to be seen under such information structure, whether such result holds in the leader-follower setup.

4.2. Outline of the Proof

In this section, we provide an outline of our proof. First, we analyze the regret-bound for the leaders.

Regret for Leader: We decompose the regret term for the leader as the following

$$\sum_{k} \underbrace{(V^{\pi_{l}^{k,*},\pi_{f}^{k}}(x_{1}) - V_{l,1}^{k}(x_{1}))}_{\mathcal{T}_{1}} + \underbrace{(V_{l,1}^{k}(x_{1}) - V^{\pi_{l}^{k},\pi_{f}^{k}}(x_{1}))}_{\mathcal{T}_{2}}$$

 \mathcal{T}_1 denotes the optimism term, and \mathcal{T}_2 denotes the model-prediction error.

Uniform Concentration: In order to bound \mathcal{T}_1 and \mathcal{T}_2 , we need to bound the difference between the estimated value function $V_{l,h}^k$ and the value function $V_{l,h}^{\pi_l,\pi_f}$ for any joint policy π_l and π_f . As in the single agent case, the key step is to control the fluctuations in least-squares value iteration. In particular, we need to show that for all $(k, h) \in [K] \times [H]$ with high probability

$$\left\|\sum_{\tau=1}^{k-1} \phi(x_h^{\tau}, a_h^{\tau}, b_h^{\tau}) \left[V_{l,h+1}^k(x_{h+1}^{\tau}) - \mathbb{P}_h V_{l,h+1}^k(x_h^{\tau}, a_h^{\tau}, b_h^{\tau}) \right] \right\|_{(\Lambda_h^k)^{-1}}$$

is upper bounded by $\mathcal{O}(d\sqrt{\log K})$. To this end, value-aware uniform concentration is required to handle the dependence between $V_{l,h+1}^k$ and samples $\{x_{h+1}^{\tau}\}_{\tau=1}^{k-1}$, which renders the standard selfnormalized inequality infeasible in the model-free setting. The general idea here is to fix a function class $\mathcal{V}_{l,h}$ in advance and then show that each possible value function in our algorithm $V_{l,h}^k$ is within this class and has log ϵ -covering number that only scales with $\mathcal{O}(\log(K))$. In the following, we fix an $h \in [H]$ and drop the subscript h for notation simplicity.

Why Soft-max?: We first note that this uniform concentration bound is the main motivation for us to choose a soft-max policy for the leader and the follower as we will see that the standard greedy policy would fail in this case. That is, in order to guarantee that for each possible V_l^k , there is an ϵ -close function in \mathcal{V}_l , it would basically lead to a very large covering number. Soft-max resolves the issue.

We first define the following class for Q_m -function for m = l, f. $Q_m = \{Q_m | Q_m(\cdot, \cdot, \cdot) =$ $\min\{\langle w_m, \phi(\cdot, \cdot, \cdot) \rangle + \beta \sqrt{\phi(\cdot, \cdot, \cdot)^T (\Lambda_h^k)^{-1} \phi(\cdot, \cdot, \cdot)}, H\}.$ Then, we define the class of marginal q function of the leader \hat{Q}_l for a given leader's action a. $\hat{Q}_l(a) = \{q_l | q_l(\cdot, a) = \sum_b \pi(b|\cdot, a) Q_l(\cdot, a, b); Q_l \in \mathbb{C}\}$ $\mathcal{Q}_l, \pi \in \Pi$ }, where Π is given by the following class $\Pi = \{\pi | \pi(b|\cdot, a) = \text{SOFT-MAX}_{\alpha_f}^b((Q_f(\cdot, a, \cdot)); \forall b \in \mathbb{C}\})$ $\mathcal{B}, Q_f \in \mathcal{Q}_f$, where SOFT-MAX is defined in (12). The necessity of the above function class is that the value-function class depends on q_l , i.e., the class of the value function for leader is $\mathcal{V}_l = \{V_l | V_l(\cdot) = \sum_a \pi_l(a | \cdot) q_l(\cdot, a), q_l \in \hat{Q}_l(a), \forall a \in \mathcal{A}\}$ where π_l is again Soft-max policy with parameter α_l . In order to compute ϵ -covering for the class \mathcal{V}_l we need to compute ϵ -covering for the class \hat{Q}_l . At this moment, we can explain why the introduction of soft-max in our algorithm is critical. Suppose we follow the standard greedy selection for the follower, which corresponds to $\alpha_f = \infty$ in above. The key issue in this approach is that one needs a large ϵ -covering for \hat{Q}_l so that each possible q_l^k can be well-approximated (i.e., ϵ -close) by function in \hat{Q}_l . This is in sharp contrast to the single-agent case where \mathcal{V}_l has a covering number polynomial in K. To see this difference, in the single agent case, we do not have the policy of the follower. Since the value function for the single agent only depends on the agent's own policy, by the fact that max is a contraction map, an ϵ -covering of Q_l^k implies an ϵ -covering of V_l^k and meanwhile the ϵ - covering number of \mathcal{Q}_m is polynomial in K (Lemma 16) by standard arguments. This no longer holds in the leader-follower setup as the value function now inherently depends on the follower's policy. In particular, note that if the policy is greedy for the follower then an ϵ -covering of Q_l^k fails to be an ϵ -covering of q_l^k in general since the greedy policy is not smooth in that a slight change of the follower's Q-function could

lead to a substantial change of the follower's policy eventually the leader's marginal q-function q_l^k . This leads to a large distance for *leader's* value functions due to the different action choices for the follower, even though the Q-function for the follower are close (See Example 1 for an example). Hence, one can not approximate the leader's value function within ϵ -bound using greedy policy based on the follower's Q-function. This fact motivates us to turn to SOFT-MAX_{α}, which is Lipschitz continuous with a Lipschitz constant at most 2α . Thus, our main idea is as follows. Given Q_f^k , we can first find fixed $\tilde{Q}_f \in Q_f$, such that $\left\|Q_f^k - \tilde{Q}_f\right\|_{\infty} \leq \epsilon_1$, then, thanks to the smoothness of soft-max function, we have $\|\pi_k - \tilde{\pi}\|_1 \leq 2\alpha\epsilon_1$ (Lemma 17) where $\tilde{\pi}$ is the soft-max policy based on \tilde{Q}_f . Hence, using the above we obtain $\sum_b \pi(b|x, a)Q_l^k(x, a, b) - \sum_b \tilde{\pi}(b|x, a)\tilde{Q}_l(x, a, b) \leq \epsilon'$ where $\left\|Q_l^k - \tilde{Q}_l\right\|_{\infty} \leq \epsilon_2$ by carefully choosing ϵ_1 and ϵ_2 as the family of Q functions have small ϵ -covering number. Thus, one can show that $\hat{Q}_l(a)$ has a log ϵ -covering number which scales as $\mathcal{O}(\log(K))$ for every a as well.

Once we show that $\hat{\mathcal{Q}}_l(a)$ has $\log_{\epsilon} \epsilon$ covering number which scales with $\log(K)$, we can show that the $\log_{\epsilon} \epsilon$ covering number for \mathcal{V}_l also scales at most with $\log(K)$. In particular, we find \tilde{q}_l for a given q_l such that $||\tilde{q}_l - q_l||_{\infty} \leq \epsilon'$. Now using the Lipschitz continuity of the soft-max policy for the leader, we then show that $||\pi_l - \tilde{\pi}_l||_1 \leq 2\alpha_l \epsilon'$. Hence, we can find \tilde{V}_l such that $||V_l - \tilde{V}_l||_{\infty} \leq \epsilon$ by carefully choosing ϵ' which is enough to show that $\log \epsilon$ -covering number for the value function class scales at most with $\mathcal{O}(\log(K))$. To summarize, the soft-max policies enable to obtain ϵ covering number for the value function class from the ϵ -covering number for the joint state-action value classes Q. Using the fact that Q_m has $\log \epsilon$ -covering number of the order $\log(K)$ we can also achieve the log ϵ -covering number for the value function class as $\log(K)$.

In fact, a larger value of α_f and α_l means that we need a smaller ϵ , hence a larger covering number. Then, one may wonder if we can choose an arbitrarily small value for α_f or α_l . However, the leader's regret will be large if we choose too small α_l (similarly, if we choose too small α_f the regret for the follower would be large). In particular, we obtain

Lemma 2 With probability 1 - p/2,

$$\mathcal{T}_1 \le \frac{KH\log(|\mathcal{A}|)}{\alpha_l} \tag{13}$$

For $\alpha_l = \frac{\log(|\mathcal{A}|\sqrt{K})}{H}$, we have $\mathcal{T}_1 \leq \sqrt{K}H^2$

Thus, we can not set α_l too small. Note that unlike the single agent scenario, \mathcal{T}_1 is not upper bounded by 0 as the leader does not select policy based on the greedy action.

Combining the Azuma-Hoeffding inequality with the above, we obtain

Lemma 3 With probability 1 - p/2, $\mathcal{T}_2 \leq \mathcal{O}(\sqrt{d^3TH^3\iota^2})$.

Regret for Follower: Now, we analyze the regret bound for the followers. We decompose the regret term similar to the leader in the following manner

$$\sum_{k} \underbrace{\bar{V}_{f,1}^{\pi_{k}^{k},\pi_{f}^{*}}(x_{1},a_{1}^{k}) - \bar{V}_{f,1}^{k}(x_{1},a_{1}^{k})}_{\mathcal{T}_{3}} + \underbrace{\bar{V}_{f,1}^{k}(x_{1},a_{1}^{k}) - \bar{V}_{f,1}^{\pi_{k}^{k},\pi_{f}^{*}}(x_{1},a_{1}^{k})}_{\mathcal{T}_{4}}$$

The analysis again hinges on showing that the value function class \mathcal{V}_f has log ϵ -covering number that is upper bounded by $\mathcal{O}(\log(K))$. Since the value function V_f^k depends on the joint policy of the leader and the follower, one needs soft-max policy of the leader and the follower to show that ϵ -covering number of the Q-function class is enough to bound ϵ -covering number for \mathcal{V}_f by $\mathcal{O}(\log(K))$. Similar to \mathcal{T}_1 and \mathcal{T}_2 we obtain

Lemma 4 With probability 1 - p/2

$$\mathcal{T}_3 \le KH \frac{\log(|\mathcal{B}|)}{\alpha_f}, \quad \mathcal{T}_4 \le \sqrt{d^3 H^3 T \iota^2} \tag{14}$$

By choosing $\alpha_f = \frac{\log(|\mathcal{B}|)\sqrt{K}}{H}$, we obtain $\mathcal{T}_3 \leq \sqrt{KH^4}$.

As one can see one can not set too small value of α_f which would increase the regret. On the other hand, too large α_f would not enable us to obtain optimal regret bound as well.

Remark 4 We have assumed that the feature space ϕ is known. Note that feature space learning is an active area of research Zhang et al. (2022); Agarwal et al. (2020); Modi et al. (2021) for linear approximation setup. The most promising technique is to estimate the Q-function by jointly optimizing over w and ϕ . Neural networks can be used to obtain such w and ϕ . Using similar technique we can learn ϕ , however, such a characterization is left for the future.

Remark 5 *Recently, regret analysis for zero-sum game setup has been extended to the non-linear MDP setup Jin et al. (2021a); Huang et al. (2021). Extending our analysis towards non-linear MDP by combining their approach constitutes an interesting future research direction.*

5. Equilibrium Learning

The Algorithms 1 and 2 also enables us to obtain equilibrium policies (ϵ -close).

Finding Coarse Correlated Stackelberg Equilibrium: We first define CCSE.

Definition 6 Joint policy (π_l, π_f) is coarse-correlated Stackelberg equilibrium (CCSE) if for all x

$$V_{l,1}^{\pi_l,\pi_f}(x) \ge V_{l,1}^{\pi'_l,\pi_f}(x), \quad V_{f,1}^{\pi_l,\pi_f}(x) \ge V_{f,1}^{\pi_l,\pi'_f}(x)$$

for any leader's policy π'_l and follower's policy π'_f . The joint policy is ϵ -CCSE if $V_{l,1}^{\pi_l,\pi_f}(x) \geq V_{l,1}^{\pi'_l,\pi_f}(x) - \epsilon$.

Now, we show that our algorithms indeed return a joint policy which is ϵ -CCSE.

Corollary 1 Consider the joint policy: the leader and follower jointly choose a $k \in [K]$ with prob. 1/K, and then the leader and follower select the policy π_l^k and π_f^k respectively (returned by Algorithms 1 and 2). Such a joint policy is $\tilde{O}(\sqrt{d^3H^4/K})$ -CCSE with probability 1 - p.

The above result entails that in order to achieve ϵ -CCSE one needs $\tilde{O}(1/\epsilon^2)$ episodes. This is the first such result for non-myopic leader and follower Markov game setup with function approximation. The agent only needs to coordinate on the random number to choose the episode index k.

Finding Stackelberg Equilibrium: For a zero-sum game, CCSE coincides with the Stackelberg equilibrium. Hence, by Corollary 1, we also obtain ϵ - Stackelberg equilibrium for zero-sum game.

For more general setting, we can combine the reward-free exploration proposed in Wang et al. (2020); Liu et al. (2021) and the soft-max policy to obtain SE with self-play. In particular, we divide the total episodes in two-phases, exploration phase and exploitation phase. In exploration phase, both the leader and the follower only explore to reduce the confidence bound. Instead of true reward, the reward will be the bonus term, $||\phi(x, a, b)||_{(\Lambda_h^k)^{-1}}$ to both the leader and follower which will incentivize the players to explore similar to Wang et al. (2020) (however, we have to use soft-max policy instead of greedy policy). In exploitation phase, the leader and follower obtain policy similar to Algorithm 1 and 2. The complete characterization is left for the future.

6. Conclusion and Future Work

We propose a model-free RL-based algorithm for both the leader and the follower. We have achieved $\tilde{O}(\sqrt{d^3H^3T})$ regret for both the leader and the follower. We have extended the LSVI-UCB algorithm towards the leader-follower setup. We have underlined the technical challenges in doing so and explained how the greedy policy for the players fail to achieve an uniform concentration bound for individual value function. We show that a soft-max policy for the players can achieve the regret bound.

Whether we can tighten this dependence on d remains an important future research direction. Whether we can the tighten the dependence on H also constitutes a future research direction. Finally, we consider one leader and one follower setup. Extending our setup to multiple leaders and followers constitutes an important future research direction.

References

- Yasin Abbasi-Yadkori, Dávid Pál, and Csaba Szepesvári. Improved algorithms for linear stochastic bandits. Advances in neural information processing systems, 24:2312–2320, 2011.
- Alekh Agarwal, Sham Kakade, Akshay Krishnamurthy, and Wen Sun. Flambe: Structural complexity and representation learning of low rank mdps. *Advances in neural information processing systems*, 33:20095–20107, 2020.
- Yu Bai and Chi Jin. Provable self-play algorithms for competitive reinforcement learning. In *International conference on machine learning*, pages 551–560. PMLR, 2020.
- Yu Bai, Chi Jin, and Tiancheng Yu. Near-optimal reinforcement learning with self-play. *Advances in neural information processing systems*, 33:2159–2170, 2020.
- Yu Bai, Chi Jin, Huan Wang, and Caiming Xiong. Sample-efficient learning of stackelberg equilibria in general-sum games. Advances in Neural Information Processing Systems, 34:25799–25811, 2021.
- Maria-Florina Balcan, Avrim Blum, Nika Haghtalab, and Ariel D Procaccia. Commitment without regrets: Online learning in stackelberg security games. In *Proceedings of the sixteenth ACM conference on economics and computation*, pages 61–78, 2015.
- Avrim Blum, Nika Haghtalab, and Ariel D Procaccia. Learning optimal commitment to overcome insecurity. *Advances in Neural Information Processing Systems*, 27, 2014.
- Zixiang Chen, Dongruo Zhou, and Quanquan Gu. Almost optimal algorithms for two-player markov games with linear function approximation. *arXiv preprint arXiv:2102.07404*, 2021.
- Richard Cole and Tim Roughgarden. The sample complexity of revenue maximization. In *Proceed*ings of the forty-sixth annual ACM symposium on Theory of computing, pages 243–252, 2014.
- Vincent Conitzer and Tuomas Sandholm. Self-interested automated mechanism design and implications for optimal combinatorial auctions. In *Proceedings of the 5th ACM Conference on Electronic Commerce*, pages 132–141, 2004.
- Vincent Conitzer and Tuomas Sandholm. Computing the optimal strategy to commit to. In *Proceedings of the 7th ACM conference on Electronic commerce*, pages 82–90, 2006.
- Alessandro Epasto, Mohammad Mahdian, Vahab Mirrokni, and Manolis Zampetakis. Optimal approximation–smoothness tradeoffs for soft-max functions. *arXiv preprint arXiv:2010.11450*, 2020.
- Baihe Huang, Jason D Lee, Zhaoran Wang, and Zhuoran Yang. Towards general function approximation in zero-sum markov games. arXiv preprint arXiv:2107.14702, 2021.
- Chi Jin, Zhuoran Yang, Zhaoran Wang, and Michael I Jordan. Provably efficient reinforcement learning with linear function approximation. In *Conference on Learning Theory*, pages 2137– 2143. PMLR, 2020.

- Chi Jin, Qinghua Liu, and Sobhan Miryoosefi. Bellman eluder dimension: New rich classes of rl problems, and sample-efficient algorithms. *Advances in neural information processing systems*, 34:13406–13418, 2021a.
- Chi Jin, Qinghua Liu, Yuanhao Wang, and Tiancheng Yu. V-learning–a simple, efficient, decentralized algorithm for multiagent rl. *arXiv preprint arXiv:2110.14555*, 2021b.
- Hsu Kao, Chen-Yu Wei, and Vijay Subramanian. Decentralized cooperative reinforcement learning with hierarchical information structure. In *International Conference on Algorithmic Learning Theory*, pages 573–605. PMLR, 2022.
- Joshua Letchford, Vincent Conitzer, and Kamesh Munagala. Learning and approximating the optimal strategy to commit to. In *International symposium on algorithmic game theory*, pages 250–262. Springer, 2009.
- Qinghua Liu, Tiancheng Yu, Yu Bai, and Chi Jin. A sharp analysis of model-based reinforcement learning with self-play. In *International Conference on Machine Learning*, pages 7001–7010. PMLR, 2021.
- Weichao Mao and Tamer Başar. Provably efficient reinforcement learning in decentralized generalsum markov games. *Dynamic Games and Applications*, pages 1–22, 2022.
- Aditya Modi, Jinglin Chen, Akshay Krishnamurthy, Nan Jiang, and Alekh Agarwal. Model-free representation learning and exploration in low-rank mdps. *arXiv preprint arXiv:2102.07035*, 2021.
- Ling Pan, Qingpeng Cai, Qi Meng, Wei Chen, Longbo Huang, and Tie-Yan Liu. Reinforcement learning with dynamic boltzmann softmax updates. *arXiv preprint arXiv:1903.05926*, 2019.
- Binghui Peng, Weiran Shen, Pingzhong Tang, and Song Zuo. Learning optimal strategies to commit to. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 33, pages 2149–2156, 2019.
- Muhammed Sayin, Kaiqing Zhang, David Leslie, Tamer Basar, and Asuman Ozdaglar. Decentralized q-learning in zero-sum markov games. *Advances in Neural Information Processing Systems*, 34:18320–18334, 2021.
- Milind Tambe. Security and game theory: algorithms, deployed systems, lessons learned. Cambridge university press, 2011.
- Yi Tian, Yuanhao Wang, Tiancheng Yu, and Suvrit Sra. Online learning in unknown markov games. In *International conference on machine learning*, pages 10279–10288. PMLR, 2021.
- Roman Vershynin. Introduction to the non-asymptotic analysis of random matrices. *arXiv preprint arXiv:1011.3027*, 2010.
- Ruosong Wang, Simon S Du, Lin Yang, and Russ R Salakhutdinov. On reward-free reinforcement learning with linear function approximation. *Advances in neural information processing systems*, 33:17816–17826, 2020.

- Chen-Yu Wei, Yi-Te Hong, and Chi-Jen Lu. Online reinforcement learning in stochastic games. Advances in Neural Information Processing Systems, 30, 2017.
- Chen-Yu Wei, Mehdi Jafarnia Jahromi, Haipeng Luo, Hiteshi Sharma, and Rahul Jain. Modelfree reinforcement learning in infinite-horizon average-reward markov decision processes. In *International conference on machine learning*, pages 10170–10180. PMLR, 2020.
- Qiaomin Xie, Yudong Chen, Zhaoran Wang, and Zhuoran Yang. Learning zero-sum simultaneousmove markov games using function approximation and correlated equilibrium. *arXiv preprint arXiv:2002.07066*, 2020.
- Lin Yang and Mengdi Wang. Sample-optimal parametric q-learning using linearly additive features. In *International Conference on Machine Learning*, pages 6995–7004. PMLR, 2019.
- Xuezhou Zhang, Yuda Song, Masatoshi Uehara, Mengdi Wang, Alekh Agarwal, and Wen Sun. Efficient reinforcement learning in block mdps: A model-free representation learning approach. In *International Conference on Machine Learning*, pages 26517–26547. PMLR, 2022.
- Liyuan Zheng, Tanner Fiez, Zane Alumbaugh, Benjamin Chasnov, and Lillian J Ratliff. Stackelberg actor-critic: Game-theoretic reinforcement learning algorithms. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 36, pages 9217–9224, 2022.
- Stephan Zheng, Alexander Trott, Sunil Srinivasa, Nikhil Naik, Melvin Gruesbeck, David C Parkes, and Richard Socher. The ai economist: Improving equality and productivity with ai-driven tax policies. arXiv preprint arXiv:2004.13332, 2020.
- Han Zhong, Zhuoran Yang, Zhaoran Wang, and Michael I Jordan. Can reinforcement learning find stackelberg-nash equilibria in general-sum markov games with myopic followers? *arXiv preprint arXiv:2112.13521*, 2021.

Notations: Throughout the rest of this paper, we denote $V_{l,h}^k$, $V_{f,h}^k$, $Q_{l,h}^k$, $Q_{f,h}^k$, $w_{l,h}^k$, $w_{f,h}^k$, Λ_h^k as the Q-value and the parameter values estimated at the episode k. We also denote the marginal q-function for leader as $q_{l,h}^k$ and the leader-action dependent estimated value function as $\overline{V}_{f,h}^k$ at episode k. To simplify the presentation, we denote $\phi_h^k = \phi(x_h^k, a_h^k, b_h^k)$. To simplify notation, we also sometimes denote $\pi = \{\pi_l, \pi_f\}$.

Without loss of generality, we assume $||\phi(x, a)||_2 \leq 1$ for all $(x, a) \in \mathcal{S} \times \mathcal{A}$, $||\mu_h(\mathcal{S})||_2 \leq \sqrt{d}$, $||\theta_{m,h}||_2 \leq \sqrt{d}$ for m = l, f and all $h \in [H]$.

Appendix A. Preliminary Results

Lemma 5 Under Assumption 2, for any fixed policy π_f , π_l , let $w_{m,h}^{\pi}$ be the corresponding weights such that $Q_{m,h}^{\pi}(x, a, b) = \langle \phi(x, a, b), w_{m,h}^{\pi} \rangle$, for $m \in \{l, f\}$, then we have for all $h \in [H]$,

$$||w_{m,h}^{\pi}|| \le 2H\sqrt{d} \tag{15}$$

Proof From the linearity of the action-value function, we have

$$Q_{m,h}^{\pi}(x,a,b) = r_{m,h}(x,a,b) + \mathbb{P}_h V_{m,h}^{\pi}(x,a,b)$$

= $\langle \phi(x,a,b), \theta_{m,h} \rangle + \int_{\mathcal{S}} V_{m,h+1}^{\pi}(x',a) \langle \phi(x,a,b), d\mu_h(x') \rangle$
= $\langle \phi(x,a,b), w_{m,h}^{\pi} \rangle$ (16)

where $w_{m,h}^{\pi} = \theta_{m,h} + \int_{S} V_{m,h+1}^{\pi}(x',a) d\mu_h(x').$

Now, $||\theta_{m,h}|| \leq \sqrt{d}$, and $||\int_{\mathcal{S}} V_{j,h+1}^{\pi}(x')d\mu_h(x')|| \leq H\sqrt{d}$. Thus, the result follows from (16).

Lemma 6 For any (k, h), the weight $w_{j,h}^k$ satisfies

$$||w_{m,h}^k|| \le 2H\sqrt{dk/\lambda} \tag{17}$$

a

Proof For any vector $v \in \mathcal{R}^d$ we have from the definition of $w_{m,h}^k$ and $V_{m,h+1}^k$ as

$$|v^{T}w_{m,h}^{k}| = |v^{T}(\Lambda_{h}^{k})^{-1} \sum_{\tau=1}^{k-1} \phi_{h}^{\tau}(x_{h}^{\tau}, a^{\tau}, b_{h}^{\tau}) (m_{h}(x_{h}^{\tau}, a^{\tau}, b_{h}^{\tau}) + \sum_{a} \pi_{l,h+1}^{k}(a|x_{h+1}^{\tau}) \sum_{b} \pi_{f,h+1}^{k}(b|x_{h+1}^{\tau}, a) Q_{m,h+1}^{k}(x_{h+1}^{\tau}, a) Q_{m,h$$

here $\pi_{l,h+1}^k$, and $\pi_{f,h+1}^k(\cdot|x,a)$ are the Soft-max policies with respect to the marginal q-function and the joint state-action value function.

Note that $Q_{m,h+1}^k(x,a,b) \leq H$ for any (x,a,b). Hence, from (18) we have

$$v^{T}w_{j,h}^{k}| \leq \sum_{\tau=1}^{k-1} |v^{T}(\Lambda_{h}^{k})^{-1}\phi_{h}^{\tau}|.2H$$

$$\leq \sqrt{\sum_{\tau=1}^{k-1} v^{T}(\Lambda_{h}^{k})^{-1}v} \sqrt{\sum_{\tau=1}^{k-1} \phi_{h}^{\tau}(\Lambda_{h}^{k})^{-1}\phi_{h}^{\tau}}.2H$$

$$\leq 2H||v||\frac{\sqrt{dk}}{\sqrt{\lambda}}$$
(19)

Note that $||w_{m,h}^k|| = \max_{v:||v||=1} |v^T w_{m,h}^k|$. Hence, the result follows.

Appendix B. Regret for Leader

Let us recall the decomposition of the regret for the leader.

$$\operatorname{Regret}(K) = \sum_{k} \underbrace{(V^{\pi_{l}^{k,*},\pi_{f}^{k}}(x_{1}) - V_{l,1}^{k}(x_{1}))}_{\mathcal{T}_{1}} + \underbrace{(V_{l,1}^{k}(x_{1}) - V^{\pi_{l}^{k},\pi_{f}^{k}}(x_{1}))}_{\mathcal{T}_{2}}$$
(20)

We now bound \mathcal{T}_1 and \mathcal{T}_2 .

B.1. Proof Outline

Here, we provide a outline of the Proof. The common step in obtaining the bound for \mathcal{T}_1 and \mathcal{T}_2 is to bound the gap between $Q_{l,h}^{\pi_l,\pi_f}$ and $Q_{l,h}^k$ for every h and k. In particular, we show that the gap between $\langle \phi(x,a,b), w_{l,h}^k \rangle$ and $\langle \phi(x,a,b), w_{l,h}^{\pi_l,\pi_f}(x,a,b) \rangle$ is upper bounded by expected gap between $V_{l,h+1}^k$ and $V_{l,h+1}^{\pi_l,\pi_f}$ plus a term (which is exactly equal to the bonus term) with high probability (Lemma 8). In order to prove Lemma 8, we state and prove Lemma 7 which lies in the heart of the proof. Further, we show that if one use soft-max, the sub-optimality gap can be bounded (Lemma 9). Now, recall that we add the bonus term to $\langle \phi(x,a,b), w_{l,h}^k \rangle$ to obtain $Q_{l,h}^k$. Thus, combining all of the above we bound \mathcal{T}_1 using backward induction. Using Lemma 8, we show that the difference between $V_{l,h}^k - V_{l,h}^{\pi_l,\pi_f}$ can be upper bounded by sum of Martingale differences and sum of the bonus terms (Lemma 10). Thus, \mathcal{T}_2 can be bounded using Azuma-Hoeffding inequality and elliptical potential lemma.

B.2. Base Results

First, we state and prove Lemma 7. In order to bound T_1 and T_2 , we state and prove some base results.

Lemma 7 There exists a constant C_2 such that for any fixed $p \in (0, 1)$, if we let \mathcal{E} be the event that

$$\left\|\sum_{\tau=1}^{k-1} \phi_h^{\tau} [V_{l,h+1}^k(x_{h+1}^{\tau}) - \mathbb{P}_h V_{l,h+1}^k(x_h^{\tau}, a_h^{\tau}, b_h^{\tau})]\right\|_{(\Lambda_h^k)^{-1}} \le C_2 dH \sqrt{\chi}$$
(21)

for all $j \in \{r, g\}$, $\chi = \log[4(C_1 + 1)(\log(|\mathcal{A}||\mathcal{B}|) + 2\log(|\mathcal{A}|)\log(|\mathcal{B}|))dT/p]$, for some constant C_2 , then $\Pr(\mathcal{E}) = 1 - p/2$.

The proof of Lemma 7 is technical and relegated later. An extra $\log(|\mathcal{A}|)$ term appears compared to the single agent case Jin et al. (2020) because the temp. coefficients of the soft-max policies (α_l, α_f) appears in the covering number.

We now recursively bound the difference between the value function maintained in Algorithm 1 (without the bonus term) and the value function $V_{l,h+1}^{\pi_l,\pi_f}$ for any follower's policy. We bound this using the expected difference at the next step plus an error term. This error term can be upper bounded by the bonus term with a high-probability.

Lemma 8 There exists an absolute constant $\beta = C_1 dH \sqrt{\iota}$, $\iota = \log((\log(|\mathcal{A}||\mathcal{B}|) + 2\log(|\mathcal{A}|)\log(|\mathcal{B}|))4dT/p)$, and for any fixed policy π_f , on the event \mathcal{E} defined in Lemma 7, we have

$$\langle \phi(x,a,b), w_{l,h}^k \rangle - Q_{l,h}^{\pi_l,\pi_f}(x,a,b) = \mathbb{P}_h(V_{l,h+1}^k - V_{l,h+1}^{\pi_l,\pi_f})(x,a,b) + \Delta_h^k(x,a,b)$$
(22)

for some $\Delta_h^k(x, a)$ that satisfies $|\Delta_h^k(x, a, b)| \leq \beta \sqrt{\phi(x, a, b)^T (\Lambda_h^k)^{-1} \phi(x, a, b)}$.

Proof Note that $Q_{l,h}^{\pi_l,\pi_f}(x,a,b) = \langle \phi(x,a,b), w_{l,h}^{\pi_l,\pi_f} \rangle = r_{l,h}(x,a,b) + \mathbb{P}_h V_{l,h+1}^{\pi_l,\pi_f}(x,a,b)$. Hence, we have

$$w_{l,h}^{k} - w_{l,h}^{\pi_{l},\pi_{f}} = (\Lambda_{h}^{k})^{-1} \sum_{\tau=1}^{k-1} \phi_{h}^{\tau} [r_{l,h}^{\tau} + V_{l,h+1}^{k}(x_{h+1}^{\tau})] - w_{l,h}^{\pi_{l},\pi_{f}}$$

$$= -\lambda (\Lambda_{h}^{k})^{-1} (w_{l,h}^{\pi_{l},\pi_{f}}) + (\Lambda_{h}^{k})^{-1} \sum_{\tau=1}^{k-1} \phi_{h}^{\tau} [V_{l,h+1}^{k}(x_{h+1}^{\tau}) - \mathbb{P}_{h} V_{l,h+1}^{k}(x_{h}^{\tau}, a_{h}^{\tau}, b_{h}^{\tau})]$$

$$+ (\Lambda_{h}^{k})^{-1} \sum_{\tau=1}^{k-1} \phi_{h}^{\tau} [\mathbb{P}_{h} V_{l,h+1}^{k}(x_{h}^{\tau}, a_{h}^{\tau}, b_{h}^{\tau}) - \mathbb{P}_{h} V_{l,h+1}^{\pi}(x_{h}^{\tau}, a_{h}^{\tau}, b_{h}^{\tau})]$$
(23)

Now, we bound each term in the right hand side of expression in (23). We call those terms as q_1 , q_2 , and q_3 respectively.

First, note that

$$\begin{aligned} |\langle \phi(x,a,b), \mathbf{q}_1 \rangle| &= |\lambda \langle \phi(x,a,b), (\Lambda_h^k)^{-1} (w_{l,h}^{\pi_l,\pi_f}) \rangle| \\ &\leq \sqrt{\lambda} ||w_{l,h}^{\pi_l,\pi_f}|| \sqrt{\phi(x,a,b)^T (\Lambda_h^k)^{-1} \phi(x,a,b)} \end{aligned}$$
(24)

Second, from Lemma 7, for the event in \mathcal{E} , we have

$$|\langle \phi(x,a,b), \mathbf{q}_2 \rangle| \le C dH \sqrt{\chi} \sqrt{\phi(x,a,b)^T (\Lambda_h^k)^{-1} \phi(x,a,b)}$$
(25)

where $\chi = \log[4(C_1 + 1)(\log(|\mathcal{A}||\mathcal{B}|) + 2\log(|\mathcal{A}|)\log(|\mathcal{B}|))dT/p].$ Third,

$$\begin{aligned} \langle \phi(x,a,b), \mathbf{q}_{3} \rangle &= \langle \phi(x,a,b), (\Lambda_{h}^{k})^{-1} \sum_{\tau=1}^{k-1} \phi_{h}^{\tau} [\mathbb{P}_{h} (V_{l,h+1}^{k} - V_{l,h+1}^{\pi_{l},\pi_{f}}) (x_{h}^{\tau}, a_{h}^{\tau}, b_{h}^{\tau})] \rangle \\ &= \langle \phi(x,a,b), (\Lambda_{h}^{k})^{-1} \sum_{\tau=1}^{k-1} \phi_{h}^{\tau} (\phi_{h}^{\tau})^{T} \int (V_{l,h+1}^{k} - V_{l,h+1}^{\pi_{l},\pi_{f}}) (x') d\mu_{h}(x') \rangle \\ &= \langle \phi(x,a,b), \int (V_{l,h+1}^{k} - V_{l,h+1}^{\pi_{l},\pi_{f}}) (x') d\mu_{h}(x') \rangle - \langle \phi(x,a,b), \lambda (\Lambda_{h}^{k})^{-1} \int (V_{l,h+1}^{k} - V_{l,h+1}^{\pi_{l},\pi_{f}}) (x') d\mu_{h}(x') \rangle \end{aligned}$$

$$(26)$$

The second term in (26) can be bounded as the following

$$|\langle \phi(x,a,b), \lambda(\Lambda_{h}^{k})^{-1} \int (V_{m,h+1}^{k} - V_{m,h+1}^{\pi_{l},\pi_{f}})(x') d\mu_{h}(x') \rangle| \leq 2H\sqrt{d\lambda} \sqrt{\phi(x,a,b)^{T}(\Lambda_{h}^{k})^{-1}\phi(x,a,b)}$$
(27)

since $||\int (V_{r,h+1}^k - V_{r,h+1}^\pi)(x')d\mu_h(x')||_2 \le 2H\sqrt{d}$ as $||\mu_h(\mathcal{S})|| \le \sqrt{d}$. The first term in (26) is equal to

$$\mathbb{P}_{h}(V_{m,h+1}^{k} - V_{m,h+1}^{\pi_{l},\pi_{f}})(x,a,b)$$
(28)

Note that $\langle \phi(x, a, b), w_{l,h}^k \rangle - Q_{l,h}^{\pi_l, \pi_f}(x, a, b) = \langle \phi(x, a, b), w_{l,h}^k - w_{l,h}^{\pi_l, \pi_f} \rangle = \langle \phi(x, a, b), \mathbf{q_1} + \mathbf{q_2} + \mathbf{q_3} \rangle$. Since $\lambda = 1$, we have from (24), (25),(27), and (28)

$$|\langle \phi(x,a,b), w_{l,h}^k \rangle - Q_{l,h}^{\pi_l,\pi_f}(x,a,b) - \mathbb{P}_h(V_{l,h+1}^k - V_{l,h+1}^{\pi_l,\pi_f})(x,a,b)| \le C_3 dH \sqrt{\chi} \sqrt{\phi(x,a,b)^T (\Lambda_h^k)^{-1} \phi(x,a,b)}$$
(29)

for some constant C_3 which is independent of C_1 . Finally, note that

$$C_{3}\sqrt{\chi} = \sqrt{\log[4(C_{1}+1)(\log(|\mathcal{A}||\mathcal{B}|)+2\log(|\mathcal{A}|)\log(|\mathcal{B}|))]}dT/p]$$

= $C_{3}\sqrt{\iota + \log(C_{1}+1)}$
 $\leq C_{1}\sqrt{\iota}$ (30)

B.3. Proof of Lemma 2

Using the above, given a follower's policy, we bound the gap between the optimal value function and the estimated value function $V_{l,h}^k$ maintained by our algorithm (Lemma 2).

. . .

Before that we state and prove another result.

Lemma 9 Then,
$$\tilde{V}_{l,h}^k(x) - V_{l,h}^k(x) \le \frac{\log |\mathcal{A}|}{\alpha_l}$$

where

Definition 7
$$\tilde{V}_{l,h}^k(\cdot) = \max_a \left[\sum_b \pi_{f,h}^k(b|\cdot,a) Q_{l,h}^k(\cdot,a,b)\right] = \max_a q_{l,h}^k(\cdot,a).$$

 $\bar{V}_{l,h}^k(\cdot)$ is the value function corresponds to the greedy-policy with respect to the marginal Q function.

Proof Note that

$$V_{l,h}^{k}(x) = \sum_{a} \pi_{l,h}(a|x) \sum_{b} \pi_{f,h}^{k}(b|x,a) Q_{l,h}^{k}(x,a,b) = \sum_{a} \pi_{l,h}(a|x) q_{l,h}^{k}(x,a)$$
(31)

where

$$\pi_{h,k}(a|x) = \frac{\exp(\alpha_l[q_{l,h}^k(x,a)])}{\sum_a \exp(\alpha_l[q_{l,h}^k(x,a)])}$$
(32)

Denote $a_x = \arg \max_a q_{l,h}^k(x,a)$

Now, recall from Definition 7 that $\tilde{V}_h^k(x) = q_{l,h}^k(a_x)$. Then,

$$\tilde{V}_{l,h}^{k}(x) - V_{l,h}^{k}(x) = q_{l,h}^{k}(a_{x}) - \sum_{a} \pi_{l,h}^{k}(a|x)q_{l,h}^{k}(x,a)$$

$$\leq \left(\frac{\log(\sum_{a} \exp(\alpha_{l}(q_{l,h}^{k}(x,a))))}{\alpha_{l}}\right) - \sum_{a} \pi_{l,h}^{k}(a|x)q_{l,h}^{k}(x,a)$$

$$\leq \frac{\log(|\mathcal{A}|)}{\alpha_{l}}$$
(33)

where the last inequality follows from Proposition 1 in Pan et al. (2019).

We are now ready to prove Lemma 2.

Proof We prove the lemma by Induction.

First, we prove for the step H. Note that $Q_{l,H+1}^k = 0 = Q_{l,H+1}^{\pi}$. Under the event in \mathcal{E} as described in Lemma 7 and 8, we have

$$|\langle \phi(x,a,b), w_{l,H}^k \rangle - Q_{l,H}^{\pi}(x,a,b)| \le \beta \sqrt{\phi(x,a,b)^T (\Lambda_H^k)^{-1} \phi(x,a,b)}$$

Hence, for any (x, a, b),

$$Q_{l,H}^{\pi}(x,a,b) \leq \min\{\langle \phi(x,a,b), w_{l,H}^k \rangle + \beta \sqrt{\phi(x,a,b)^T (\Lambda_H^k)^{-1} \phi(x,a,b), H}\}$$
$$= Q_{l,H}^k(x,a,b)$$
(34)

Since π_f^k derived by the leader and the played by the follower is the same, thus, for any (x, a)

$$q_{l,H}^{\pi}(x,a) = \sum_{b} \pi_{f}^{k}(b|x,a)Q_{l,H}^{\pi}(x,a,b)$$

$$\leq \sum_{b} \pi_{f}^{k}(b|x,a)Q_{l,H}^{k}(x,a,b) = q_{l,H}^{k}(x,a)$$
(35)

Hence, from the definition of \tilde{V}_h^k ,

$$\tilde{V}_{H}^{k}(x) = \max_{a} q_{l,H}^{k}(x,a,b) \ge \sum_{a} \pi_{l,h}^{k}(a|x)q_{l,h}^{k}(x,a)
= V_{l,H}^{\pi_{l},\pi_{f}}(x)$$
(36)

for any policy $\{\pi_l, \pi_f\}$. Thus, it also holds for $\{\pi_l^{k,*}, \pi_f\}$, the optimal policy. Hence, from Lemma 9, we have

$$V_{l,H}^{\pi_l^{k,*},\pi_f}(x) - V_{l,H}^k(x) \le \frac{\log(|\mathcal{A}|)}{\alpha_l}$$

Now, suppose that it is true till the step h + 1 and consider the step h.

Since, it is true till step h + 1, thus, for any policy π ,

$$\mathbb{P}_{h}(V_{l,h+1}^{\pi_{l}^{k,*},\pi_{f}} - V_{l,h+1}^{k})(x,a,b) \leq \frac{(H-h)\log(|\mathcal{A}|)}{\alpha_{l}}$$
(37)

From Lemma 8 and the above result, we have for any (x, a, b)

$$Q_{l,h}^{\pi_l^{k,*},\pi_f}(x,a,b) \le Q_{l,h}^k(x,a,b) + \frac{(H-h)\log(|\mathcal{A}|)}{\alpha_l}$$
(38)

Hence, for any (x,a) and the fact that $\sum_b \pi_{f,h}(b|x,a)=1,$ we have

$$q_{l,h}^{\pi_l^{k,*},\pi_f}(x,a) \le q_{l,h}^k(x,a) + \frac{(H-h)\log(|\mathcal{A}|)}{\alpha_l}$$
(39)

Thus, from the definition of $\tilde{V}_{l,h}^k$

$$V_{l,h}^{\pi_{l}^{k,*},\pi_{f}}(x) \leq \tilde{V}_{l,h}^{k}(x) + \frac{(H-h)\log(|\mathcal{A}|)}{\alpha_{l}}$$
(40)

Now, again from Lemma 9, we have $\tilde{V}_{l,h}^k(x) - V_{l,h}^k(x) \le \frac{\log(|\mathcal{A}|)}{\alpha}$. Thus,

$$V_{l,h}^{\pi_l^{k,*},\pi_f}(x) - V_{l,h}^k(x) \le \frac{(H-h+1)\log(|\mathcal{A}|)}{\alpha_l}$$
(41)

Thus, we have

$$V_{l,h}^{\pi_l^*,\pi_f}(x) - V_{l,h}^k(x) \le \frac{(H-h+1)\log(|\mathcal{A}|)}{\alpha_l}$$

Hence, the result follows by summing over K and considering h = 1.

In order to prove the Lemma 3, we state and prove another result. First, we introduce a notation. Let

$$D_{l,h,1}^{k} = \mathbb{D}^{\pi_{l}^{k},\pi_{f}^{k}}(Q_{l,h}^{k} - Q_{l,h}^{\pi_{l}^{k},\pi_{f}^{k}}) - (Q_{l,h}^{k}(x_{h}^{k},a_{h}^{k},b_{h}^{k}) - Q_{l,h}^{\pi_{k}^{k}}(x_{h}^{k},a_{h}^{k},b_{h}^{k}))$$
$$D_{l,h,2}^{k} = \mathbb{P}_{h}(V_{l,h+1}^{k} - V_{j,h+1}^{\pi_{k}})(x_{h}^{k},a_{h}^{k},b_{h}^{k}) - [V_{l,h+1}^{k} - V_{l,h+1}^{\pi_{k}}](x_{h+1}^{k})$$
(42)

We will show that the above two terns are Martingale under proper filtration which we use to prove Lemma 3. First, we show that the difference between $V_{l,h}^k(x_1) - V_{l,h}^{\pi_l^k,\pi_f^k}$ as the sum of the above two martingales plus the bonus term. In the following, we denote $\pi^k = {\pi_l^k, \pi_f^k}$.

Lemma 10 On the event defined in \mathcal{E} in Lemma 7, we have

$$V_{l,1}^k(x_1) - V_{l,1}^{\pi_l^k, \pi_f^k}(x_1) \le \sum_{h=1}^H (D_{l,h,1}^k + D_{l,h,2}^k) + \sum_{h=1}^H 2\beta \sqrt{\phi(x_h^k, a_h^k, b_h^k)^T (\Lambda_h^k)^{-1} \phi(x_h^k, a_h^k, b_h^k)}$$

Proof By Lemma 8, for any x, h, a, b, k

$$\langle w_{l,h}^{k}, \phi(x,a,b) \rangle + \beta \sqrt{\phi(x,a,b)^{T} (\Lambda_{h}^{k})^{-1} \phi(x,a,b)} - Q_{j,h}^{\pi^{k}}(x,a,b)$$

$$\leq \mathbb{P}_{h} (V_{l,h+1}^{k} - V_{l,h+1}^{\pi^{k}})(x,a,b) + 2\beta \sqrt{\phi(x,a,b)^{T} (\Lambda_{h}^{k})^{-1} \phi(x,a,b)}$$
 (43)

Thus,

$$Q_{l,h}^{k}(x,a,b) - Q_{l,h}^{\pi^{k}}(x,a,b) \leq \mathbb{P}_{h}(V_{l,h+1}^{k} - V_{l,h+1}^{\pi^{k}})(x,a,b) + 2\beta\sqrt{\phi(x,a,b)^{T}(\Lambda_{h}^{k})^{-1}\phi(x,a,b)} \\ \mathbb{P}_{h}(V_{l,h+1}^{k} - V_{l,h+1}^{\pi^{k}})(x,a,b) + 2\beta\sqrt{\phi(x,a,b)^{T}(\Lambda_{h}^{k})^{-1}\phi(x,a,b)} - (Q_{l,h}^{k}(x,a,b) - Q_{l,h}^{\pi^{k}}(x,a,b)) \geq 0$$

$$(44)$$

Since $V_{l,h}^k(x) = \mathbb{D}^{\pi_{l,h}^k, \pi_{f,h}^k} Q_{l,h}^k(x)$ and $V_{l,h}^{\pi^k}(x) = \mathbb{D}^{\pi_{l,h}^k, \pi_{f,h}^k} Q_{l,h}^{\pi^k}(x)$. Thus, from (44), $V_{l,h}^k(x_h^k) - V_{l,h}^{\pi_k}(x_h^k) = \sum_{a} \sum_{b} \pi_{l,h}^k(a|x_h^k) \pi_{f,h}^k(b|x_h^k,) [Q_{l,h}^k(x_h^k,a,b) - Q_{l,h}^{\pi^k}(x_h^k,a,b)]$ $\leq \sum_{a} \sum_{b} \pi^{k}_{l,h}(a|x^{k}_{h}) \pi^{k}_{f,h}(b|x^{k}_{h},a) [Q^{k}_{j,h}(x^{k}_{h},a,b) - Q^{\pi^{k}}_{j,h}(x^{k}_{h},a,b)]$ $+2\beta\sqrt{\phi(x_{h}^{k},a_{h}^{k},b_{h}^{k})^{T}(\Lambda_{h}^{k})^{-1}\phi(x_{h}^{k},a_{h}^{k},b_{h}^{k})} + \mathbb{P}_{h}(V_{l,h+1}^{k} - V_{l,h+1}^{\pi^{k}})(x_{h}^{k},a_{h}^{k},b_{h}^{k}) - (Q_{l,h}^{k}(x_{h}^{k},a_{h}^{k},b_{h}^{k}) - Q_{l,h}^{\pi^{k}}(x_{h}^{k},a_{h}^{k})) + (Q_{l,h+1}^{k}(x_{h}^{k},a_{h}^{k},b_{h}^{k}) - (Q_{l,h}^{k}(x_{h}^{k},a_{h}^{k},b_{h}^{k})) - (Q_{l,h}^{k}(x_{h}^{k},a_{h}^{k},b_{h}^{k})) + (Q_{l,h+1}^{k}(x_{h}^{k},a_{h}^{k},b_{h}^{k})) + (Q_{l,h+1}^{k}(x_{h}^{k},a_{h$

(45)

Thus, from (45), we have

$$V_{l,h}^{k}(x_{h}^{k}) - V_{l,h}^{\pi^{k}}(x_{h}^{k}) \leq D_{l,h,1}^{k} + D_{l,h,2}^{k} + [V_{l,h+1}^{k} - V_{l,h+1}^{\pi^{k}}](x_{h+1}^{k}) + 2\beta\sqrt{\phi(x_{h}^{k}, a_{h}^{k}, b_{h}^{k})^{T}(\Lambda_{h}^{k})^{-1}\phi(x_{h}^{k}, a_{h}^{k}, b_{h}^{k})}$$

Hence, by iterating recursively, we have

$$V_{l,1}^k(x_1) - V_{l,1}^{\pi^k}(x_1) \le \sum_{h=1}^H (D_{l,h,1}^k + D_{l,h,2}^k) + \sum_{h=1}^H 2\beta \sqrt{\phi(x_h^k, a_h^k, b_h^k)^T (\Lambda_h^k)^{-1} \phi(x_h^k, a_h^k, b_h^k)^{-1} \phi(x_h^k, a_h^k)^{-1} \phi(x_h^k)^{-1} \phi(x_h^k)^{-1}$$

The result follows.

Proof Note from Lemma 10, we have

$$\sum_{k=1}^{K} V_{l,1}^{k}(x_{1}) - V_{l,1}^{\pi_{k}}(x_{1}) \leq \sum_{k=1}^{K} \sum_{h=1}^{H} (D_{l,h,1}^{k} + D_{l,h,2}^{k}) + \sum_{k=1}^{K} \sum_{h=1}^{H} 2\beta \sqrt{\phi(x_{h}^{k}, a_{h}^{k}, b_{h}^{k})^{T} (\Lambda_{h}^{k})^{-1} \phi(x_{h}^{k}, a_{h}^{k}, b_{h}^{k})}$$

$$(46)$$

We, now, bound the individual terms. First, we show that the first term corresponds to a Martingale difference.

For any $(k,h) \in [K] \times [H]$, we define $\mathcal{F}_{h,1}^k$ as σ -algebra generated by the joint state-action sequences, $\{(x_i^{\tau}, a_i^{\tau}, b_i^{\tau})\}_{(\tau,i)\in[k-1]\times[H]} \cup \{(x_i^k, a_i^k, b_i^k)\}_{i\in[h]}$. Similarly, we define the $\mathcal{F}_{h,2}^k$ as the σ -algebra generated by $\{(x_i^{\tau}, a_i^{\tau}, b_i^{\tau})\}_{(\tau,i)\in[k-1]\times[H]} \cup \{(x_i^{\tau}, a_i^{\tau}, b_i^{\tau})\}_{(\tau,i)\in[k-1]\times[H]}$

 $\{(x_i^k, a_i^k, b_i^k)\}_{i \in [h]} \cup \{x_{h+1}^k\}. x_{H+1}^k$ is a null state for any $k \in [K]$.

A filtration is a sequence of σ -algebras $\{\mathcal{F}_{h,m}^k\}_{(k,h,m)\in[K]\times[H]\times[2]}$ in terms of time index

$$t(k, h, m) = 2(k - 1)H + 2(h - 1) + m$$

which holds that $\mathcal{F}_{h,m}^k \subset \mathcal{F}_{h',m'}^{k'}$ for any $t \leq t'$.

Note from the definitions in (42) that $D_{l,h,1}^k \in \mathcal{F}_{h,1}^k$ and $D_{l,h,2}^k \in \mathcal{F}_{h,2}^k$. Thus, for any $(k,h) \in [K] \times [H]$,

$$\mathbb{E}[D_{l,h,1}^{k}|\mathcal{F}_{h-1,2}^{k}] = 0, \quad \mathbb{E}[D_{l,h,2}^{k}|\mathcal{F}_{h,1}^{k}] = 0$$

Notice that t(k, 0, 2) = t(k - 1, H, 2) = 2(H - 1)k. Clearly, $\mathcal{F}_{0,2}^k = \mathcal{F}_{H,2}^{k-1}$ for any $k \ge 2$. Let $\mathcal{F}_{0,2}^1$ be empty. We define a Martingale sequence

$$\begin{split} M_{l,h,m}^{k} &= \sum_{\tau=1}^{k-1} \sum_{i=1}^{H} (D_{l,i,1}^{\tau} + D_{l,i,2}^{\tau}) + \sum_{i=1}^{h-1} (D_{l,i,1}^{k} + D_{l,i,2}^{k}) + \sum_{l'=1}^{m} D_{l,h,l'}^{k} \\ &= \sum_{(\tau,i,l') \in [K] \times [H] \times [2], t(\tau,i,l') \le t(k,h,m)} D_{l,i,l'}^{\tau} \end{split}$$

where t(k, h, m) = 2(k-1)H + 2(h-1) + m is the time index. Clearly, this martingale is adopted to the filtration $\{\mathcal{F}_{h,m}^k\}_{(k,h,m)\in[K]\times[H]\times[2]}$, and particularly

$$\sum_{k=1}^{K} \sum_{h=1}^{H} (D_{l,h,1}^{k} + D_{l,h,2}^{k}) = M_{l,H,2}^{K}$$
(47)

Thus, $M_{l,H,2}^K$ is a Martingale difference satisfying $|M_{l,H,2}^K| \le 4H$ since $|D_{l,h,1}^k|, |D_{l,h,2}^k| \le 2H$ From the Azuma-Hoeffding inequality, we have

$$\Pr(M_{l,H,2}^K > s) \le 2\exp(-\frac{s^2}{16TH^2})$$

With probability 1 - p/4 at least

$$\sum_{k} \sum_{h} M_{l,H,2}^{K} \le \sqrt{16TH^2 \log(4/p)}$$
(48)

Now, we bound the second term in (46). Note that the minimum eigen value of Λ_h^k is at least $\lambda = 1$ for all $(k, h) \in [K] \times [H]$. By Lemma 21,

$$\sum_{k=1}^{K} (\phi_h^k)^T (\Lambda_h^k)^{-1} \phi_h^k \le 2 \log \left[\frac{\det(\Lambda_h^{k+1})}{\det(\Lambda_h^1)} \right]$$

Moreover, note that $||\Lambda_h^{k+1}|| = ||\sum_{\tau=1}^k \phi_h^k (\phi_h^k)^T + \lambda \mathbf{I}|| \le \lambda + k$, hence,

$$\sum_{k=1}^{K} (\phi_h^k)^T (\Lambda_h^k)^{-1} \phi_h^k \le 2d \log \left[\frac{\lambda+k}{\lambda}\right] \le 2d\iota$$

Now, by Cauchy-Schwartz inequality, we have

$$\sum_{k=1}^{K} \sum_{h=1}^{H} \sqrt{(\phi_{h}^{k})^{T} (\Lambda_{h}^{k})^{-1} \phi_{h}^{k}} \leq \sum_{h=1}^{H} \sqrt{K} [\sum_{k=1}^{K} (\phi_{h}^{k})^{T} (\Lambda_{h}^{k})^{-1} \phi_{h}^{k}]^{1/2} \leq H \sqrt{2dK\iota}$$
(49)

Note that $\beta = C_1 dH \sqrt{\iota}$.

Thus, combining (47), (48), and (49), we have with probability 1 - p/2,

$$\sum_{k=1}^{K} V_{l,1}^{k}(x_{1}^{k}) - V_{l,1}^{\pi_{k}}(x_{1}^{k}) \leq \left[\sqrt{16TH^{2}\log(4/p)} + C_{4}\sqrt{d^{3}H^{3}T\iota^{2}}\right]$$

Hence, the result follows.

Appendix C. Regret for Follower

We now prove the regret bound for the followers. Recall the decomposition of the regret term for follower.

$$\sum_{k} \underbrace{\bar{V}_{f,1}^{\pi_{l}^{k},\pi_{f}^{*}}(x_{1},a_{1}^{k}) - \bar{V}_{f,1}^{k}(x_{1},a_{1}^{k})}_{\mathcal{T}_{3}} + \underbrace{\bar{V}_{f,1}^{k}(x_{1},a_{1}^{k}) - \bar{V}_{f,1}^{\pi_{l}^{k},\pi_{f}^{k}}(x_{1},a_{1}^{k})}_{\mathcal{T}_{4}} + \underbrace{\bar{V}_{f,1}^{k}(x_{1},a_{1}^{k}) - \bar{V}_{f,1}^{k}(x_{1},a_{1}^{k})}_{\mathcal{T}_{4}} + \underbrace{\bar{V}_{f,1}^{k}(x_{1},a_{1}^{k}) - \underbrace{\bar{V}_{f,1}^{k}(x_{1},a_{1}^{k})}_{\mathcal{T}_{4}} + \underbrace{\bar{V}_{f,1}^{k}(x_{1},a_{1}^{k}) - \underbrace{\bar{V}_{f,1}^{k}(x_{1},a_{1}^{k})}_{\mathcal{T}_{4}} + \underbrace{\bar{V}_{f,1}^{k}(x_{1},a_{1}^{k}) - \underbrace{\bar{V}_{f,1}^{k}(x_{1},a_{1}^{k})}_{\mathcal{T}_{4}} + \underbrace{\bar{V}_{f,1}^{k}(x_{1},a$$

Similar to the way we proved the regret bound for the leader, we are going to prove the regret bound for the follower. There are some subtle differences as we need to consider the fact the follower takes action after observing the leader's action. In order to bound T_3 and T_4 , we state and prove some base results.

First, similar to Lemma 8, we state and prove Lemma 12 which characterizes the difference between $\langle w_{f,h}^k, \phi(x,a,b) \rangle$ and $Q_{f,h}^{\pi_l,\pi_f}$. In particular, we show that the gap between $\langle \phi(x,a,b), w_{l,h}^k \rangle$ and $\langle \phi(x,a,b), w_{l,h}^{\pi_l,\pi_f}(x,a,b) \rangle$ is upper bounded by expected gap between $V_{f,h+1}^k$ and $V_{f,h+1}^{\pi_l,\pi_f}$ plus the bonus term (Lemma 12). In order to prove Lemma 12, we state and prove Lemma 11 which lies in the heart of the proof of regret. Further, we show that if one use soft-max, the sub-optimality gap can be bounded (Lemma 13). Now, recall that we add the bonus term to $\langle \phi(x,a,b), w_{f,h}^k \rangle$ to obtain $Q_{f,h}^k$. Thus, combining all of the above we bound \mathcal{T}_3 using backward induction since the follower also computes the leader's policy. Using Lemma 12, we show that the difference between $\bar{V}_{f,h}^k - \bar{V}_{f,h}^{\pi_l,\pi_f}$ can be upper bounded by sum of Martingale differences and sum of the bonus terms (Lemma 14). Thus, \mathcal{T}_4 can be bounded using Azuma-Hoeffding inequality and elliptical potential lemma.

C.1. Fomral Proof

Similar to Lemma 7, we can show the following

Lemma 11 There exists a constant C_2 such that for any fixed $p \in (0,1)$, if we let \mathcal{E} be the event that

$$\left\|\sum_{\tau=1}^{k-1} \phi_h^{\tau} [V_{f,h+1}^k(x_{h+1}^{\tau}) - \mathbb{P}_h V_{f,h+1}^k(x_h^{\tau}, a_h^{\tau}, b_h^{\tau})]\right\|_{(\Lambda_h^k)^{-1}} \le C_2 dH \sqrt{\chi}$$
(50)

 $\chi = \log[4(C_1 + 1)(\log(|\mathcal{A}||\mathcal{B}|) + 2\log(|\mathcal{A}|)\log(|\mathcal{B}|))dT/p], \text{ for some constant } C_2, \text{ then } \Pr(\mathcal{E}) = 1 - p/2.$

Similar to Lemma 8, we can also show the following for the follower as well.

Lemma 12 There exists an absolute constant $\beta = C_1 dH \sqrt{\iota}$, $\iota = \log((\log(|\mathcal{A}||\mathcal{B}|) + 2\log(|\mathcal{A}|)\log(|\mathcal{B}|))4dT/p)$, and for any fixed policy π_f , on the event \mathcal{E} defined in Lemma 11, we have

$$\langle \phi(x,a,b), w_{l,h}^k \rangle - Q_{f,h}^{\pi_l,\pi_f}(x,a,b) = \mathbb{P}_h(V_{f,h+1}^k - V_{f,h+1}^{\pi_l,\pi_f})(x,a,b) + \Delta_h^k(x,a,b)$$
(51)

for some $\Delta_h^k(x, a)$ that satisfies $|\Delta_h^k(x, a, b)| \leq \beta \sqrt{\phi(x, a, b)^T (\Lambda_h^k)^{-1} \phi(x, a, b)}$.

Similar to Lemma 9, we also obtain the following

Lemma 13 Then, $\tilde{V}_{f,h}^k(x,a) - \bar{V}_{f,h}^k(x,a) \le \frac{\log |\mathcal{B}|}{\alpha}$

where

Definition 8
$$\tilde{V}_{f,h}^k(\cdot, a) = \sum_b \pi_{f,h}^k(b|\cdot, a) Q_{f,h}^k(\cdot, a, b)].$$

 $\tilde{V}_{f,h}^k(\cdot)$ is the value function corresponds to the greedy-policy with respect to the Q function given the leader's action. There is a slight difference with Lemma 9. Here, we are bounding the gap for a given leader's action.

Proof Note that

$$\bar{V}_{f,h}^{k}(x,a) = \sum_{b} \pi_{f,h}^{k}(b|x,a) Q_{f,h}^{k}(x,a,b)$$

where

$$\pi_{f,h}^{k}(b|x,a) = \frac{\exp(\alpha_{f}[Q_{f,h}^{k}(x,a,b)])}{\sum_{b}\exp(\alpha_{f}[Q_{f,h}^{k}(x,a,b)])}$$
(52)

Denote $b_x = \arg \max_b Q_{f,h}^k(x, a, b)$

Now, recall from Definition 8 that $\tilde{V}_{f,h}^k(x,a) = Q_{f,h}^k(x,a,b_x)$. Then,

$$\tilde{V}_{f,h}^{k}(x,a) - \bar{V}_{f,h}^{k}(x,a) = Q_{f,h}^{k}(x,a,b_{x}) - \sum_{b} \pi_{f,h}^{k}(b|x,a)Q_{f,h}^{k}(x,a,b)$$

$$= \left(\frac{\log(\sum_{b} \exp(\alpha_{f}(Q_{f,h}^{k}(x,a,b))))}{\alpha_{f}}\right) - \sum_{b} \pi_{f,h}^{k}(b|x,a)Q_{f,h}^{k}(x,a,b)$$

$$\leq \frac{\log(|\mathcal{B}|)}{\alpha_{f}} \tag{53}$$

where the last inequality follows from Proposition 1 in Pan et al. (2019).

We are now ready to prove the first part of Lemma 4, i.e., we obtain the bound for \mathcal{T}_3 . **Proof** We prove the lemma by Induction.

First, we prove for the step H.

Note that $Q_{f,H+1}^k = 0 = Q_{f,H+1}^{\pi}$. Under the event in \mathcal{E} as described in Lemma 11 and 12, we have

$$|\langle \phi(x,a,b), w_{f,H}^k \rangle - Q_{f,H}^{\pi}(x,a,b)| \le \beta \sqrt{\phi(x,a,b)^T (\Lambda_H^k)^{-1} \phi(x,a,b)}$$

Hence, for any (x, a, b),

$$\begin{aligned} Q_{f,H}^{\pi}(x,a,b) &\leq \min\{\langle \phi(x,a,b), w_{f,H}^k \rangle + \beta \sqrt{\phi(x,a,b)^T (\Lambda_H^k)^{-1} \phi(x,a,b), H} \} \\ &= Q_{f,H}^k(x,a,b) \end{aligned}$$

Hence, from the definition of \tilde{V}_h^k , for any (x, a)

$$\begin{split} \tilde{V}_{H}^{k}(x,a) &= \max_{b} Q_{f,H}^{k}(x,a,b) \geq \sum_{b} \pi_{f,h}^{k}(b|x,a) Q_{f,h}^{\pi}(x,a,b) \\ &= \bar{V}_{f,H}^{\pi}(x,a) \end{split}$$

for any policy π . Thus, it also holds for $(\pi_l, \pi_f^{k,*})$, the optimal policy for the follower for a given leader's policy π_l . Hence, from Lemma 13, we have for any a

$$\bar{V}_{f,H}^{\pi_l,\pi_f^*}(x,a) - \bar{V}_{f,H}^k(x,a) \le \frac{\log(|\mathcal{B}|)}{\alpha_f}$$

Now, suppose that it is true till the step h + 1 and consider the step h. Thus,

$$\bar{V}_{f,h+1}^{\pi_l^k,\pi_f^{k,*}}(x,a) - \bar{V}_{f,h+1}^k(x,a) \le \frac{(H-h)\log(|\mathcal{B}|)}{\alpha_f}$$
(54)

Hence, for a given policy π_l , we have

$$\sum_{a} \pi_{l,h+1}(a|x) (V_{f,h+1}^{\pi_l,\pi_f^*}(x,a) - V_{f,h+1}^k(x,a)) \le \frac{(H-h)\log(|\mathcal{B}|)}{\alpha_f}$$
(55)

since $\sum_{a} \pi_{l,h+1}(a|x) = 1$. However, in (55, the left-hand side is exactly equal to $V_{f,h+1}^{\pi_l,\pi_f^*}(x) - V_{f,h+1}^{\pi_l,\pi_f^*}(x)$ $V_{f,h+1}^k(x)$. Hence, we have

$$V_{f,h+1}^{\pi_l,\pi_f^*}(x) - V_{f,h+1}^k(x) \le \frac{(H-h)\log(|\mathcal{B}|)}{\alpha_f}$$
(56)

From Lemma 12 and the above result, we have for any (x, a, b) for any policy $\pi = \{\pi_l, \pi_f\}$

$$Q_{f,h}^{\pi}(x,a,b) \le Q_{f,h}^{k}(x,a,b) + \frac{(H-h)\log(|\mathcal{B}|)}{\alpha}$$
(57)

From the definition of $\tilde{V}_{f,h}^k$, we have

$$\sum_{b} \pi_{f}(b|x,a) Q_{f,h}^{\pi}(x,a,b) \le \max_{b} Q_{f,h}^{k}(x,a,b) + \sum_{b} \pi_{f}(b|x,a) \frac{(H-h)\log(|\mathcal{B}|)}{\alpha}$$
(58)

Since $\sum_b \pi_f(b|x,a) = 1$, and from the Definition of $\tilde{V}^k_{f,h}$, we obtain

$$\bar{V}_{f,h}^{\pi}(x,a) \leq \tilde{V}_{f,h}^{k}(x,a) + \frac{(H-h)\log(|\mathcal{B}|)}{\alpha_f}$$
(59)

Now, from Lemma 13, we have $\tilde{V}_{f,h}^k(x,a) - \bar{V}_{f,h}^k(x,a) \le \frac{\log(|\mathcal{B}|)}{\alpha_f}$. Thus, for any (x,a)

$$\bar{V}_{f,h}^{\pi}(x,a) - \bar{V}_{f,h}^{k}(x,a) \le \frac{(H-h+1)\log(|\mathcal{B}|)}{\alpha_{f}}$$

Now, since it is true for any policy π , it will be true for $\{\pi_l, \pi_f^*\}$. From the definition of $\bar{V}_{f,h}^{\pi}$, we have

$$\bar{V}_{f,h}^{\pi_l,\pi_f^*}(x,a_h) - \bar{V}_{f,h}^k(x,a_h) \le \frac{(H-h+1)\log(|\mathcal{B}|)}{\alpha_f}$$
(60)

Hence, the result follows by summing over K and considering h = 1.

In order to prove the second part of Lemma 4 (i.e., in order to bound T_4), we state and prove a result.

First, we introduce some notations. Let

$$D_{f,h,1}^{k} = \langle (Q_{f,h}^{k}(x_{h}^{k}, a_{h}^{k}, \cdot) - Q_{f,h}^{\pi^{k}}(x_{h}^{k}, a_{h}^{k}, \cdot)), \pi_{f,h}^{k}(\cdot | x_{h}^{k}, a_{h}^{k}) \rangle - (Q_{f,h}^{k}(x_{h}^{k}, a_{h}^{k}, b_{h}^{k}) - Q_{f,h}^{\pi^{k}}(x_{h}^{k}, a_{h}^{k}, b_{h}^{k})) \\ D_{f,h,2}^{k} = \mathbb{P}_{h}(V_{f,h+1}^{k} - V_{f,h+1}^{\pi^{k}})(x_{h}^{k}, a_{h}^{k}, b_{h}^{k}) - [V_{f,h+1}^{k} - V_{f,h+1}^{\pi^{k}}](x_{h+1}^{k})$$
(61)

Lemma 14 On the event defined in \mathcal{E} in Lemma 11, we have

$$\bar{V}_{f,1}^k(x_1, a_1^k) - \bar{V}_{f,1}^{\pi^k}(x_1, a_1^k) \le \sum_{h=1}^H (D_{f,h,1}^k + D_{f,h,2}^k) + \sum_{h=1}^H 2\beta \sqrt{\phi(x_h^k, a_h^k, b_h^k)^T (\Lambda_h^k)^{-1} \phi(x_h^k, a_h^k, b_h^k)}$$

Proof By Lemma 12, for any x, h, a, b, k

$$\langle w_{f,h}^{k}, \phi(x,a,b) \rangle + \beta \sqrt{\phi(x,a,b)^{T}(\Lambda_{h}^{k})^{-1}\phi(x,a,b)} - Q_{f,h}^{\pi^{k}}(x,a,b)$$

$$\leq \mathbb{P}_{h}(V_{f,h+1}^{k} - V_{f,h+1}^{\pi^{k}})(x,a,b) + 2\beta \sqrt{\phi(x,a,b)^{T}(\Lambda_{h}^{k})^{-1}\phi(x,a,b)}$$
 (62)

Thus,

$$Q_{f,h}^{k}(x,a,b) - Q_{f,h}^{\pi^{k}}(x,a,b) \leq \mathbb{P}_{h}(V_{f,h+1}^{k} - V_{f,h+1}^{\pi^{k}})(x,a,b) + 2\beta \sqrt{\phi(x,a,b)^{T}(\Lambda_{h}^{k})^{-1}\phi(x,a,b)} \\ \mathbb{P}_{h}(V_{f,h+1}^{k} - V_{f,h+1}^{\pi^{k}})(x,a,b) + 2\beta \sqrt{\phi(x,a,b)^{T}(\Lambda_{h}^{k})^{-1}\phi(x,a,b)} - (Q_{f,h}^{k}(x,a,b) - Q_{f,h}^{\pi^{k}}(x,a,b)) \geq 0$$

$$(63)$$

$$\begin{aligned} \text{Since } \bar{V}_{f,h}^{k}(x,a) &= \sum_{b} \pi_{f,h}^{k}(b|x,a)Q_{f,h}^{k}(x,a,b) \text{ and } \bar{V}_{f,h}^{\pi^{k}}(x,a) &= \sum_{b} \pi_{f,h}^{k}(b|x,a)Q_{f,h}^{\pi^{k}}(x,a,b). \\ \text{Thus, from (63),} \end{aligned}$$

$$\begin{aligned} \bar{V}_{f,h}^{k}(x_{h}^{k},a_{h}^{k}) &- \bar{V}_{f,h}^{\pi^{k}}(x_{h}^{k},a_{h}^{k}) &= \sum_{b} \pi_{f,h}^{k}(b|x_{h}^{k},a_{h}^{k})[Q_{f,h}^{k}(x_{h}^{k},a_{h}^{k},b) - Q_{f,h}^{\pi^{k}}(x_{h}^{k},a_{h}^{k},b)] \end{aligned}$$

$$\leq \sum_{b} \pi_{f,h}^{k}(b|x_{h}^{k},a_{h}^{k})[Q_{f,h}^{k}(x_{h}^{k},a_{h}^{k},b) - Q_{f,h}^{\pi^{k}}(x_{h}^{k},a_{h}^{k},b)] \end{aligned}$$

$$+ 2\beta \sqrt{\phi(x_{h}^{k},a_{h}^{k},b_{h}^{k})^{T}(\Lambda_{h}^{k})^{-1}\phi(x_{h}^{k},a_{h}^{k},b_{h}^{k})} + \mathbb{P}_{h}(V_{f,h+1}^{k} - V_{f,h+1}^{\pi^{k}})(x_{h}^{k},a_{h}^{k},b_{h}^{k}) - (Q_{f,h}^{k}(x_{h}^{k},a_{h}^{k},b_{h}^{k}) - Q_{f,h}^{\pi^{k}}(x_{h}^{k},a_{h}^{k},b_{h}^{k})) \end{aligned}$$

$$(64)$$

Thus, from (64), we have

$$\bar{V}_{f,h}^{k}(x_{h}^{k},a_{h}^{k}) - \bar{V}_{f,h}^{\pi^{k}}(x_{h}^{k},a_{h}^{k}) \leq D_{f,h,1}^{k} + D_{f,h,2}^{k} + [V_{f,h+1}^{k} - V_{f,h+1}^{\pi^{k}}](x_{h+1}^{k}) + 2\beta\sqrt{\phi(x_{h}^{k},a_{h}^{k},b_{h}^{k})^{T}(\Lambda_{h}^{k})^{-1}\phi(x_{h}^{k},a_{h}^{k},b_{h}^{k})}$$

$$(65)$$

Hence, by iterating recursively, we have

$$\bar{V}_{f,1}^k(x_1, a_1^k) - \bar{V}_{f,1}^{\pi^k}(x_1, a_1^k) \le \sum_{h=1}^H (D_{f,h,1}^k + D_{f,h,2}^k) + \sum_{h=1}^H 2\beta \sqrt{\phi(x_h^k, a_h^k, b_h^k)^T (\Lambda_h^k)^{-1} \phi(x_h^k, a_h^k, b_h^k)}$$
(66)

The result follows.

Now,, we are ready to obtain the upper bound for \mathcal{T}_4 . **Proof** Note from Lemma 14, we have

$$\sum_{k=1}^{K} \bar{V}_{f,1}^{k}(x_{1}, a_{1}^{k}) - \bar{V}_{f,1}^{\pi^{k}}(x_{1}, a_{1}^{k}) \leq \sum_{k=1}^{K} \sum_{h=1}^{H} (D_{f,h,1}^{k} + D_{f,h,2}^{k}) + \sum_{k=1}^{K} \sum_{h=1}^{H} 2\beta \sqrt{\phi(x_{h}^{k}, a_{h}^{k}, b_{h}^{k})^{T} (\Lambda_{h}^{k})^{-1} \phi(x_{h}^{k}, a_{h}^{k}, b_{h}^{k})}$$

$$(67)$$

We, now, bound the individual terms. First, we show that the first term corresponds to a Martingale difference.

For any $(k,h) \in [K] \times [H]$, we define $\mathcal{F}_{h,1}^k$ as σ -algebra generated by the state-action sequences,

rewards for leader and follower $\{(x_i^{\tau}, a_i^{\tau}, b_i^{\tau})\}_{(\tau,i)\in[k-1]\times[H]} \cup \{(x_i^k, a_i^k, b_i^k)\}_{i\in[h]}$. Similarly, we define the $\mathcal{F}_{h,2}^k$ as the σ -algebra generated by $\{(x_i^{\tau}, a_i^{\tau}, b_i^{\tau})\}_{(\tau,i)\in[k-1]\times[H]} \cup \{(x_i^{\tau}, a_i^{\tau}, b_i^{\tau})\}_{(\tau,i)\in[k-1]\times[H]} \cup \{(x_i$ $\{(x_i^k, a_i^k, b_i^k)\}_{i \in [h]} \cup \{x_{h+1}^k, a_{h+1}^k\}. x_{H+1}^k \text{ is a null state for any } k \in [K].$ A filtration is a sequence of σ -algebras $\{\mathcal{F}_{h,m}^k\}_{(k,h,m)\in [K]\times [H]\times [2]}$ in terms of time index

$$t(k, h, m) = 2(k-1)H + 2(h-1) + m$$

which holds that $\mathcal{F}_{h,m}^k \subset \mathcal{F}_{h',m'}^{k'}$ for any $t \leq t'$. Also, $\mathcal{F}_{h,m}^k \subset \mathcal{F}_{h',m'}^{k'}$ for any $(h,m,k) \leq t'$. (h', m', k').

Note from the definitions in (61) that $D_{f,h,1}^k \in \mathcal{F}_{h,1}^k$ and $D_{f,h,2}^k \in \mathcal{F}_{h,2}^k$. Thus, for any $(k,h) \in \mathcal{F}_{h,2}^k$. $[K] \times [H],$

$$\mathbb{E}[D_{f,h,1}^{k}|\mathcal{F}_{h-1,2}^{k}] = 0, \quad \mathbb{E}[D_{f,h,2}^{k}|\mathcal{F}_{h,1}^{k}] = 0$$

Notice that t(k, 0, 2) = t(k - 1, H, 2) = 2(H - 1)k. Clearly, $\mathcal{F}_{0,2}^k = \mathcal{F}_{H,2}^{k-1}$ for any $k \ge 2$. Let $\mathcal{F}_{0,2}^1$ be empty. We define a Martingale sequence

$$M_{f,h,m}^{k} = \sum_{\tau=1}^{k-1} \sum_{i=1}^{H} (D_{f,i,1}^{\tau} + D_{f,i,2}^{\tau}) + \sum_{i=1}^{h-1} (D_{f,i,1}^{k} + D_{f,i,2}^{k}) + \sum_{l'=1}^{m} D_{f,h,l'}^{k}$$
$$= \sum_{(\tau,i,l')\in[K]\times[H]\times[2]} D_{f,i,l'}^{\tau}$$
(68)

Clearly, this martingale is adopted to the filtration $\{\mathcal{F}_{h,m}^k\}_{(k,h,m)\in[K]\times[H]\times[2]}$, and particularly

$$\sum_{k=1}^{K} \sum_{h=1}^{H} (D_{f,h,1}^{k} + D_{f,h,2}^{k}) = M_{f,H,2}^{K}$$
(69)

Thus, $M_{f,H,2}^{K}$ is a Martingale difference satisfying $|M_{f,H,2}^{K}| \le 4H$ since $|D_{f,h,1}^{k}|, |D_{f,h,2}^{k}| \le 2H$ From the Azuma-Hoeffding inequality, we have

$$\Pr(M_{f,H,2}^K > s) \le 2\exp(-\frac{s^2}{16TH^2})$$
(70)

With probability 1 - p/4 at least,

$$\sum_{k} \sum_{h} M_{f,H,2}^{K} \le \sqrt{16TH^2 \log(4/p)}$$
(71)

Now, we bound the second term. Note that the minimum eigen value of Λ_h^k is at least $\lambda = 1$ for all $(k,h) \in [K] \times [H]$. By Lemma 21,

$$\sum_{k=1}^{K} (\phi_h^k)^T (\Lambda_h^k)^{-1} \phi_h^k \le 2 \log \left[\frac{\det(\Lambda_h^{k+1})}{\det(\Lambda_h^1)} \right]$$
(72)

Moreover, note that $||\Lambda_h^{k+1}|| = ||\sum_{\tau=1}^k \phi_h^k (\phi_h^k)^T + \lambda \mathbf{I}|| \le \lambda + k$, hence,

$$\sum_{k=1}^{K} (\phi_h^k)^T (\Lambda_h^k)^{-1} \phi_h^k \le 2d \log\left[\frac{\lambda+k}{\lambda}\right] \le 2d\iota$$
(73)

Now, by Cauchy-Schwartz inequality, we have

$$\sum_{k=1}^{K} \sum_{h=1}^{H} \sqrt{(\phi_{h}^{k})^{T} (\Lambda_{h}^{k})^{-1} \phi_{h}^{k}} \leq \sum_{h=1}^{H} \sqrt{K} [\sum_{k=1}^{K} (\phi_{h}^{k})^{T} (\Lambda_{h}^{k})^{-1} \phi_{h}^{k}]^{1/2} \leq H \sqrt{2dK\iota}$$
(74)

Note that $\beta = C_1 dH \sqrt{\iota}$.

Thus, we have with probability 1 - p/2,

$$\sum_{k=1}^{K} V_{f,1}^{k}(x_{1}^{k}, a_{1}^{k}) - V_{f,1}^{\pi_{k}}(x_{1}^{k}, a_{1}^{k}) \le \left[\sqrt{16TH^{2}\log(4/p)} + C_{4}\sqrt{d^{3}H^{3}T\iota^{2}}\right]$$
(75)

Hence, the result follows.

Appendix D. Proof of Lemma 7

In this section, we prove Lemma 7. First, we provide some supporting result.

D.1. Supporting Result

In order to prove Lemma 7, we leverage on the following result which has been proved in Abbasi-Yadkori et al. (2011).

Theorem 9 [Concentration of Self-Normalized Process Abbasi-Yadkori et al. (2011)] Let $\{\epsilon_t\}_{t=1}^{\infty}$ be a real-valued stochastic process with corresponding filtration $\{\mathcal{F}_t\}_{t=0}^{\infty}$. Let $\epsilon_t | \mathcal{F}_{t-1}$ be a zero mean and σ sub-Gaussian, i.e., $\mathbb{E}[\epsilon_t | \mathcal{F}_{t-1}] = 0$, and

$$\forall \zeta \in \Re, \quad \mathbb{E}[e^{\zeta \epsilon_t} | \mathcal{F}_{t-1}] \le e^{\zeta^2 \sigma^2/2}. \tag{76}$$

Let $\{\phi_t\}_{t=1}^{\infty}$ be a \Re^d -valued Stochastic process where $\phi_t \in \mathcal{F}_{t-1}$. Assume $\Lambda_0 \in \Re^{d \times d}$ is a positivedefine matrix, let, $\Lambda_t = \Lambda_0 + \sum_{j=0}^t \phi_j \phi_j^T \phi_j$. Then for any $\delta > 0$ with probability at least $1 - \delta$, we have

$$\left|\sum_{s=1}^{t} \phi_s \epsilon_s\right|_{\Lambda_t^{-1}}^2 \le 2\sigma^2 \log\left[\frac{\det(\Lambda_t)^{1/2} \det(\Lambda_0)^{-1/2}}{\delta}\right]$$
(77)

Further, we also use the following result from Jin et al. (2020) (Lemma D.4)

Proposition 10 Let $\{x_{\tau}\}_{\tau=1}^{\infty}$ be a stochastic process on state space S with corresponding Filtration $\{\mathcal{F}_{\tau}\}_{\tau=0}^{\infty}$. Let $\{\phi_{\tau}\}_{\tau=1}^{\infty}$ be a \mathbb{R}^{d} -valued stochastic process where $\phi_{\tau} \in \mathcal{F}_{\tau-1}$, and $||\phi_{\tau}|| \leq 1$. Let $\Lambda_{k} = \lambda \mathbf{I} + \sum_{\tau=1}^{k-1} \phi_{\tau} \phi_{\tau}^{T}$. Then for any $\delta > 0$ with probability $1 - \delta$, for all $k \geq 0$, and any $V \in \mathcal{V}$ so that $\sup_{x} |V(x)| \leq H$, we have

$$\left\|\sum_{\tau=1}^{k-1} \phi_{\tau} [V(x_{\tau}) - \mathbb{E}[V(x_{\tau}|\mathcal{F}_{\tau-1})]]\right\|_{(\Lambda_{h}^{k})^{-1}} \leq 4H^{2} \left[\frac{d}{2}\log(\frac{k+\lambda}{\lambda}) + \log\frac{N_{\epsilon}^{V}}{\delta}\right] + 8k^{2}\epsilon^{2}/\lambda^{2} \quad (78)$$

where N_{ϵ}^{V} is the ϵ -covering number for the value function class \mathcal{V} .

D.2. Proof Outline

Recognizing that (78) is the same as in the statement of Lemma 7, we only need to compute the ϵ -covering number N_{ϵ}^{V} for the value function class of leader. In the following, we compute the ϵ -covering number for value function class (Lemma 15).

In order to compute that we first compute the ϵ -covering number of the individual joint stateaction pair value function (Q-functions) (Lemma 16). Subsequently, we show that if the two Qfunctions are close, the policies are also close (Lemma 17). Using the above, we compute the ϵ covering number for the class of marginal Q-function (q_l) (Lemma 18). Finally, again applying the fact that the leader's policies are close as the leader is employing soft-max policy, and ϵ -covering number for the marginal Q function we compute the ϵ -covering number for value function class. Finally, combining Proposition 10 and Lemma 15, we prove Lemma 7.

D.3. Formal Proof

In the following we drop the subscript h from $Q_{l,h}^k, Q_{f,h}^k, V_{l,h}^k$, and $q_{l,h}^k$ for notational simplicity.

We first define the value function. In order to do that, we need to define the Q function class for both the leader and the follower

Definition 11 Let $\mathcal{Q}_m = \{Q|Q(\cdot, \cdot, \cdot) = \min\{w_m^T \phi(\cdot, \cdot, \cdot) + \beta \sqrt{\phi^T(\cdot, \cdot, \cdot)^T \Lambda^{-1} \phi(\cdot, \cdot, \cdot)}, H\}\}$ for m = l, f, where $\{w \in \mathbb{R}^d |||w|| \leq 2H\sqrt{dk/\lambda}\}$,

Note from Lemma 6 such class of Q function indeed covers the estimated Q-functions we obtain in our algorithm.

We now define the marginal (or,induced) Q-function class for the leader

Definition 12 Consider the class of function $\hat{Q}_l = \{q_l | q_l(\cdot, a) = \langle \pi_f(\cdot | \cdot, a) Q_l(\cdot, a, \cdot), \pi_f \in \Pi_{\mathcal{B}} \rangle \}$ where

$$\Pi_{\mathcal{B}} = \{ \pi | \forall b \in \mathcal{B}, \pi(b|\cdot, \cdot) = \text{Soft-Max}_{\alpha_f}^b((Q_f(\cdot, \cdot, \cdot)), Q_f \in \mathcal{Q}_f \},\$$

We can now introduce the class of value function for V_l .

Definition 13 Let $\mathcal{V}_l = \{V_l | V_l(\cdot) = \mathbb{D}^{\pi_l, \pi_f} Q_m(\cdot, \cdot, \cdot), \pi_l \in \Pi_{\mathcal{A}}, \pi_f \in \Pi_{\mathcal{B}}\}$ where

$$\Pi_{\mathcal{A}} = \{ \pi | \forall a \in \mathcal{A}, \pi(a|\cdot) = \text{SOFT-MAX}^a_{\alpha_l}((q_l(\cdot, \cdot)), q_l \in \mathcal{Q}_l \} \}$$

The class of value function \mathcal{V}_l is parameterized by w_l , w_f , and Λ . It is needless to say we can define the value function for the follower in the similar manner.

Lemma 15 There exists a $\tilde{V}_l \in \mathcal{V}_l$ parameterized by $(\tilde{w}_l, \tilde{w}_f, \tilde{\beta}, \Lambda)$ such that DIST $(V_l, \tilde{V}_l) \leq \epsilon$ where

$$DIST(V_l, \tilde{V}_l) = \sup_{x} |V_l(x) - \tilde{V}_l(x)|.$$
(79)

Let $N_{\epsilon}^{V_l}$ be the ϵ -covering number for the set \mathcal{V}_l , then,

$$\log N_{\epsilon}^{V_l} \le d \log \left(1 + 8H \frac{\sqrt{dk}}{\sqrt{\lambda}\epsilon'} \right) + d^2 \log \left[1 + 8d^{1/2}\beta^2 / (\lambda(\epsilon')^2) \right]$$
(80)

where $\epsilon' = \frac{\epsilon}{1 + 2(\alpha_l + \alpha_f)H + 4\alpha_l\alpha_f H^2}$.

In order to prove the above result, we need to state and prove few results first.

We, first, obtain the N_{ϵ}^Q covering number for the joint state-action value action Q_j . Towards this end, we first, introduce some notations.

Definition 14 Let C_w^{ϵ} be an $\epsilon/2$ - cover of the set $\{w \in \mathbb{R}^d | ||w|| \le 2H\sqrt{dk/\lambda}\}$ with respect to the 2-norm. Let $C_{\mathbf{A}}^{\epsilon}$ be an $\epsilon^2/4$ -cover of the set $\{\mathbf{A} \in \mathbb{R}^{d \times d} | ||\mathbf{A}||_F \le d^{1/2}\beta^2\lambda^{-1}\}$ with respect to the Frobenius norm.

Lemma 16

$$|\mathcal{C}_w^{\epsilon}| \le (1 + 8H\sqrt{dk/\lambda}/\epsilon)^d, \quad |\mathcal{C}_{\mathbf{A}}^{\epsilon}| \le [1 + 8d^{1/2}\beta^2/(\lambda\epsilon^2)]^{d^2}$$
(81)

The ϵ -covering number for the set Q_m , for $m = l, f, N_{\epsilon}^{Q_m}$ of the set Q_m for m = l, f satisfies the following

$$\log N_{\epsilon}^{Q_m} \le d \log \left(1 + \frac{8H\sqrt{dk}}{\sqrt{\lambda}\epsilon} \right) + d^2 \log[1 + 8d^{1/2}\beta^2/(\lambda\epsilon)^2]$$
(82)

The distance metric is the ∞ -norm, i.e., $\operatorname{dist}(Q_1, Q_2) = \sup_{x,a} |Q_1(x, a) - Q_2(x, a)|$.

Proof For notational simplicity, we represent $\mathbf{A} = \beta^2 \Lambda^{-1}$, and reparamterized the class \mathcal{Q}_m by (w_m, \mathbf{A}) . Now,

$$dist(Q_{1},Q_{2}) = \sup_{x,a} |[w_{1}^{T}\phi(x,a,b) + \sqrt{\phi^{T}(x,a,b)\mathbf{A}_{1}\phi(x,a,b)}] - [w_{2}^{T}\phi(x,a,b) + \sqrt{\phi^{T}(x,a)\mathbf{A}_{2}\phi(x,a,b)}]$$

$$\leq \sup_{\phi:||\phi|| \leq 1} |[w_{1}^{T}\phi + \sqrt{\phi^{T}\mathbf{A}_{1}\phi}] - [w_{2}^{T}\phi + \sqrt{\phi^{T}\mathbf{A}_{2}\phi}]|$$

$$\leq \sup_{\phi:||\phi|| \leq 1} |(w_{1} - w_{2})^{T}\phi| + \sup_{\phi:||\phi|| \leq 1} \sqrt{|\phi^{T}(\mathbf{A}_{1} - \mathbf{A}_{2})\phi|}$$

$$= ||w_{1} - w_{2}|| + \sqrt{||\mathbf{A}_{1} - \mathbf{A}_{2}||} \leq ||w_{1} - w_{2}|| + \sqrt{||\mathbf{A}_{1} - \mathbf{A}_{2}||_{F}}$$
(83)

where the second-last inequality follows from the fact that $|\sqrt{x} - \sqrt{y}| \le \sqrt{|x - y|}$. For matrices $||\cdot||_F$ denote matrix operator norm and the Frobenius norm respectively.

Recall that C_w is an $\epsilon/2$ - cover of the set $\{w \in \mathbb{R}^d | ||w|| \le 2H\sqrt{dk/\lambda}\}$ with respect to the 2-norm. Also recall that $C_{\mathbf{A}}$ be an $\epsilon^2/4$ -cover of the set $\{\mathbf{A} \in \mathbb{R}^{d \times d} | ||\mathbf{A}||_F \le d^{1/2}\beta^2\lambda^{-1}\}$. Thus, from Lemma 22,

$$|\mathcal{C}_w^{\epsilon}| \le (1 + 8H\sqrt{dk/\lambda}/\epsilon)^d, \quad |\mathcal{C}_{\mathbf{A}}^{\epsilon}| \le [1 + 8d^{1/2}\beta^2/(\lambda\epsilon^2)]^{d^2}$$

For any $Q_m \in \mathcal{Q}_m$, there exists a \tilde{Q}_m parameterized by (w_2, \mathbf{A}_2) where $w_2 \in \mathcal{C}_w^{\epsilon}$ and $\mathbf{A}_2 \in \mathcal{C}_{\mathbf{A}}^{\epsilon}$ such that $\operatorname{dist}(Q_m, \tilde{Q}_m) \leq \epsilon$. Hence, $N_{\epsilon}^{Q_m} \leq |\mathcal{C}_w^{\epsilon}| |\mathcal{C}_{\mathbf{A}}^{\epsilon}|$, which gives the result since $\log(\cdot)$ is an increasing function.

We now show that when Q-functions are close soft-max policy based on the Q-functions are close. In fact the soft-max policy is at most 2α - Lipschitz.

Lemma 17 Suppose that π is the soft-max policy (temp. coefficient $1/\alpha$) corresponding to the joint state-action value function Q- i.e., $\forall b \in \mathcal{B}$

$$\pi(b|\cdot, a) = \text{SOFT-MAX}^b_{\alpha_f}(Q_f^k(\cdot, a, \cdot)).$$

 $\tilde{\pi}$ is the soft-max policy vector with the same temp. coefficient $1/\alpha$ corresponding to the composite Q-function \tilde{Q}_m , i.e, $\forall a \in \mathcal{A}$,

$$\tilde{\pi}(b|\cdot,a) = \text{Soft-Max}^b_{\alpha_f}(\tilde{Q}_f(\cdot,a,\cdot)).$$

then, for any state x,

$$||\pi(\cdot|x,a) - \tilde{\pi}(\cdot|x,a)||_1 \le 2\alpha_f \epsilon' \tag{84}$$

where $\pi(\cdot|x,a) = \{\pi(b|x,a)\}_{b\in\mathcal{B}}$ and $\tilde{\pi}(\cdot|x,a) = \{\tilde{\pi}(b|x,a)\}_{b\in\mathcal{B}}$ when $\operatorname{dist}(Q_f^k, \tilde{Q}_f) \leq \epsilon'$.

Proof Let $\text{Exp}^{\alpha}(P)$ be a soft-max corresponding to the vector P, i.e., the *i*-th component of $\text{Exp}^{\alpha}(P)$ is

$$\frac{\exp(\alpha P_i)}{\sum_i \exp(\alpha P_i)}.$$

Note from Theorem 4.4 in Epasto et al. (2020) then, we have

$$||\operatorname{Exp}^{\alpha}(P_1) - \operatorname{Exp}^{\alpha}(P_2)||_1 \le 2\alpha ||P_1 - P_2||_{\infty}$$
(85)

for two vectors P_1 and P_2 .

Now note that in our case for a given state x, π is equivalent to $\text{Exp}^{\alpha_f}(Q_f^k(x, a, \cdot))$, and $\tilde{\pi}$ is equivalent to $\text{Exp}^{\alpha_f}(\tilde{Q}_f(x, a, \cdot))$. Then from (85) and the fact that $\text{dist}(Q_f^k, \tilde{Q}_f) \leq \epsilon'$ we have

$$||\pi(\cdot|x,a) - \tilde{\pi}(\cdot|x,a)||_1 \le 2\alpha_f \epsilon' \tag{86}$$

Hence, the result follows.

Now, using the above result, we now compute the ϵ -covering number for the marginal q-function (Definition 12).

Lemma 18 If $||Q_l - \tilde{Q}_l|| \le \epsilon'$, and $||Q_f - \tilde{Q}_f|| \le \epsilon'$, then $||q_l - \tilde{q}_l|| \le \epsilon' + 2\alpha_f \epsilon' H$ where $q_l, \tilde{q}_l \in \hat{Q}_l$, $Q_m, \tilde{Q}_m \in Q_m$, for m = l, f.

Proof Consider Q_f and \tilde{Q}_f such that $||Q_f - \tilde{Q}_f||_{\infty} \le \epsilon'$. Hence, from Lemma 17, $||\pi - \tilde{\pi}||_1 \le 2\alpha_f \epsilon^{\epsilon}$. Consider Q_l and \tilde{Q}_l such that $||Q_l - \tilde{Q}_l||_{\infty} \le \epsilon'$. Now,

$$\begin{aligned} |q_{l}(x,a) - \tilde{q}_{l}(x,a)| &= |\sum_{b} \pi(b|x,a)Q_{l}(x,a,b) - \sum_{b} \tilde{\pi}(b|x,a)\tilde{Q}_{l}(x,a,b)| \\ &\leq |\sum_{b} \pi(b|x,a)(Q_{l}(x,a,b) - \tilde{Q}_{l}(x,a,b))| + |\sum_{b} \pi(b|x,a)\tilde{Q}_{l}(x,a,b) - \sum_{b} \tilde{\pi}(b|x,a)\tilde{Q}_{l}(x,a,b)| \\ &\leq \epsilon' + ||\pi(\cdot|x,a) - \tilde{\pi}(\cdot|x,a)||_{1} ||\tilde{Q}_{l}||_{\infty} \\ &\leq \epsilon' + 2\alpha_{f}\epsilon' H \end{aligned}$$
(87)

Hence, the result follows.

Now, we show that if Q-functions are close, the leaders' policies are also close based on those Q-functions.

Lemma 19 Suppose that π_l is the soft-max policy (temp. coefficient $1/\alpha_l$) corresponding to the marginal state-action value function q- i.e., $\forall a \in A$

$$\pi_l(a|\cdot) = \text{SOFT-MAX}^a_{\alpha_l}(q_l(\cdot, \cdot)).$$

 $\tilde{\pi}_l$ is the soft-max policy vector with the same temp. coefficient $1/\alpha_l$ corresponding to the marginal *q*-function \tilde{q}_l , i.e, $\forall a \in \mathcal{A}$,

$$\tilde{\pi}_l(a|\cdot) = \text{SOFT-MAX}^a_{\alpha_l}(\tilde{q}_l(\cdot, \cdot)).$$

then, for any state x,

$$||\pi_l(\cdot|x) - \tilde{\pi}_l(\cdot|x)||_1 \le 2\alpha_l \epsilon'(\epsilon' + 2\alpha_f \epsilon' H)$$
(88)

when $\operatorname{dist}(Q_f^k, \tilde{Q}_f) \leq \epsilon'$, $\operatorname{dist}(Q_l^k, \tilde{Q}_l^k) \leq \epsilon'$.

Proof The proof follows the same steps as in Lemma 17. Now, $||q_l - \tilde{q}_l|| \le \epsilon' + 2\alpha_f \epsilon' H$ (from Lemma 18). Hence, the result follows.

Lemma 20 There exists $\tilde{V}_m \in \mathcal{V}_m$ such that

$$DIST(V_m^k, \widetilde{V}_m) \le \epsilon' + 2\alpha_f \epsilon' H + 2\alpha_l \epsilon' H(\epsilon' + 2\alpha_f \epsilon' H)$$
(89)

where $\operatorname{dist}(\tilde{Q}_m, Q_m) \leq \epsilon', \, \tilde{Q}_m \in \mathcal{Q}_m \text{ for all } j;$

$$\widetilde{V}_m(\cdot) = \sum_a [\widetilde{\pi}(a|\cdot)\widetilde{q}_l(\cdot)]$$

$$\tilde{\pi}_l(a|\cdot) =$$
SOFT-MAX $^a_{\alpha_l}(\tilde{q}_l(\cdot, a)), \quad \forall a \in \mathcal{A}$

Proof For any *x*,

$$\begin{aligned} V_{l}^{k}(x) &- \widetilde{V}_{l}(x) \\ &= \left| \sum_{a} \pi(a|x) q_{l}^{k}(x,a) - \sum_{a} \tilde{\pi}(a|x) \tilde{q}_{l}(x,a) \right| \\ &= \left| \sum_{a} \pi(a|x) q_{l}^{k}(x,a) - \sum_{a} \pi(a|x) \tilde{q}_{l}(x,a) + \sum_{a} \pi(a|x) \tilde{q}_{l}(x,a) - \sum_{a} \tilde{\pi}(a|x) \tilde{q}_{l}(x,a) \right| \\ &\leq \left| \sum_{a} \pi(a|x) q_{l}^{k}(x,a) - \sum_{a} \pi(a|x) \tilde{q}_{l}(x,a) \right| + \left| \sum_{a} \pi(a|x) \tilde{q}_{l}(x,a) - \sum_{a} \tilde{\pi}(a|x) \tilde{q}_{l}(x,a) \right| \\ &\leq \epsilon' + 2\alpha_{f} \epsilon' H + \left| |\pi(\cdot|x) - \tilde{\pi}(\cdot|x)| |_{1} \right| |\tilde{q}_{l}(x)| |_{\infty} \\ &\leq \epsilon' + 2\alpha_{f} \epsilon' H + 2\alpha_{l} \epsilon' H(\epsilon' + 2\alpha_{f} \epsilon' H) \end{aligned}$$
(90)

where we use the fact that $\operatorname{dist}(q_l^k, \tilde{q}_l) \leq \epsilon' + 2\alpha_f \epsilon' H$, and $\sum_a \pi(a|x) = 1$ for the first term and the Holder's inequality in the second term for the second last inequality. For the last inequality, we use Lemma 17, and the fact that $\tilde{q}_l(x, a) \leq H$ for any (x, a). Hence, we have the result.

Assuming that $\epsilon' \leq 1$, we have $\text{DIST}(V_m^k, \tilde{V}_m) \leq \epsilon' + 2H(\alpha_f + \alpha_l)\epsilon' + 4\alpha_l\alpha_f H^2\epsilon'$. Now, we are ready to prove Lemma 15. **Proof** Fix an ϵ . Let $\epsilon' = \frac{\epsilon}{1 + 2(\alpha_l + \alpha_f)H + 4\alpha_l\alpha_f H^2}$, then from Lemma 20, we have $\text{DIST}(V_l^k, \tilde{V}_l) \leq \epsilon$. Thus, we only need to find parameters in the ϵ' -covering of the Q-functions as described in Lemma 16 in order to obtain ϵ -close value function.

Recall the Definition 14. Then, there exists $\tilde{w}_l, \tilde{w}_f \in C_w^{\epsilon'}$ such that $||\tilde{w}_l - w_l|| \leq \frac{\epsilon'}{2}, ||\tilde{w}_f - w_f|| \leq \frac{\epsilon'}{2}$. Further, there exists $\mathbf{A}_2 \in C_{\mathbf{A}}^{\epsilon'}$ such that $||\mathbf{A} - \tilde{\mathbf{A}}||_F \leq \frac{\epsilon'^2}{4}$, $\mathbf{A} = \beta^2 (\Lambda^k)^{-1}$, $\tilde{\mathbf{A}} = \beta^2 (\tilde{\Lambda})^{-1}$, for some $\tilde{\Lambda}$. Then we obtain \tilde{Q}_m parameterized by $(\tilde{w}_m, \beta, \tilde{\Lambda})$ for m = l, f, such that $\operatorname{dist}(Q_m, \tilde{Q}_m) \leq \epsilon'$ (by Lemma 16).

Now, from Lemma 20, we have $\text{DIST}(V_l^k, \tilde{V}_l) \leq \epsilon$. Hence, there exists \tilde{V}_l parameterized by $\tilde{w}_l, \tilde{w}_f, \tilde{\mathbf{A}}$, such that $\text{Dist}(\tilde{V}_l, V_l^k) \leq \epsilon$. Hence, $N_{\epsilon}^{V_l} \leq |\mathcal{C}_{\mathbf{A}}^{\epsilon'}||\mathcal{C}_{\mathbf{A}}^{\epsilon'}|$. Thus, from Lemma 16, the ϵ -covering number $N_{\epsilon}^{V_l}$ for the set \mathcal{V}_l satisfies the following

$$\log N_{\epsilon}^{V_l} \le d \log \left(1 + 8H \frac{\sqrt{dk}}{\sqrt{\lambda}\epsilon'} \right) + d^2 \log \left[1 + 8d^{1/2}\beta^2 / (\lambda(\epsilon')^2) \right)].$$

Hence, the result follows.

From Lemma 15, note that we need ϵ' covering for the Q-functions where $\epsilon' = \frac{\epsilon}{1 + 2(\alpha_l + \alpha_f)H + 4\alpha_l\alpha_f H^2}$ if we need to bound DIST (V_j, \tilde{V}_j) by ϵ .

Now, we are ready to prove Lemma 7.

Proof By Lemma 15, we know that there exists \tilde{V}_i in the ϵ -covering for \mathcal{V}_i such that for every x,

$$V_l(x) = \tilde{V}_l(x) + \Delta V(x) \tag{91}$$

where $\sup_x \Delta V(x) \le \epsilon$.

Hence,

$$\left\|\sum_{\tau=1}^{k} \phi^{\tau}(V_{l}(x_{\tau}) - \mathbb{E}[V_{l}(x_{\tau})|\mathcal{F}_{\tau-1}])\right\|_{(\Lambda^{k})^{-1}}^{2} \leq 2\left\|\sum_{\tau=1}^{k} \phi^{\tau}(\tilde{V}_{l}(x_{\tau}) - \mathbb{E}[\tilde{V}_{l}(x_{\tau})|\mathcal{F}_{\tau-1}])\right\|_{(\Lambda^{k})^{-1}}^{2} + 2\left\|\sum_{\tau=1}^{k} \phi^{\tau}(\Delta V(x_{\tau}) - \mathbb{E}[\Delta V(x_{\tau})|\mathcal{F}_{\tau-1}])\right\|_{(\Lambda^{k})^{-1}}^{2}$$
(92)

The last expression is bounded by $\frac{8k^2\epsilon^2}{\lambda}$.

Now, we bound the first term. Note from Lemma 15 that in order to obtain \tilde{V}_l which satisfies (91), we need to obtain we need $N_{\epsilon}^{V_l}$ number of elements to obtain such $(\tilde{w}_l, \tilde{w}_f, \beta, \tilde{\Lambda})$. Such \tilde{V}_l is independent of samples. Hence, we can use the self-normalization lemma (Theorem 9). From

Proposition 10 we obtain

$$\left\|\sum_{\tau=1}^{k} \phi^{\tau}(\tilde{V}_{l}(x_{\tau}) - \mathbb{E}[\tilde{V}_{l}(x_{\tau})|\mathcal{F}_{\tau-1}])\right\|_{(\Lambda^{k})^{-1}}^{2} \leq 2H^{2} \left[d\log\left(\frac{k+\lambda}{\lambda}\right) + \log\left(\frac{N_{\epsilon}^{V_{l}}}{\delta}\right)\right]$$
(93)

where $N_{\epsilon_l}^V$ is upper bounded in (80). β is equal to $C_1 dH \sqrt{\iota}$ for some constant C_1 , and $\iota = \log(\log(|\mathcal{A}||\mathcal{B}|) + \log(|\mathcal{B}|) \log(|\mathcal{A}|) 4dT/p)$. We obtain from (93)

$$\left\|\sum_{\tau=1}^{k} \phi^{\tau}(\tilde{V}_{l}(x_{\tau}) - \mathbb{E}[\tilde{V}_{l}(x_{\tau})|\mathcal{F}_{\tau-1}])\right\|_{(\Lambda^{k})^{-1}}^{2} \leq 4H^{2}\left[\frac{d}{2}\log\left(\frac{k+\lambda}{\lambda}\right) + d\log\left(1 + \frac{8H\sqrt{dk}}{\epsilon'\sqrt{\lambda}}\right) + d^{2}\log\left(1 + \frac{8d^{1/2}\beta^{2}}{\epsilon'^{2}\lambda}\right) + \log\left(\frac{4}{p}\right)\right]$$
(94)

where $\epsilon' = \frac{\epsilon}{1 + 2(\alpha_l + \alpha_f)H + 4\alpha_l\alpha_f H^2}$. Set $\epsilon = \frac{1}{k}$, $\lambda = 1$. Thus, $\epsilon' = \frac{1}{(1 + 2(\alpha_l + \alpha_f)H + 4\alpha_l\alpha_f H^2)k}$. Plugging in the above, and putting $\alpha_l = \frac{\log(|\mathcal{A}|)K}{2H}$, and $\alpha_f = \frac{\log(|\mathcal{B}|)K}{2H}$, we obtain from (94)

$$||\sum_{\tau=1}^{k} \phi^{\tau} (\tilde{V}_{l}(x_{\tau}) - \mathbb{E}[\tilde{V}_{l}(x_{\tau})|\mathcal{F}_{\tau-1}])||_{\Lambda_{k}^{-1}}^{2} \leq C_{2}H^{2}d^{2}\log\left(\frac{4(C_{1}+1)(\log(|\mathcal{A}||\mathcal{B}|)+2\log(|\mathcal{A}|)\log(|\mathcal{B}|))dT}{p}\right)$$
(95)

for some constant C_2 . Hence, the result follows.

Appendix E. Supporting Results

The following result is shown in Abbasi-Yadkori et al. (2011) and in Lemma D.2 in Jin et al. (2020).

Lemma 21 Let $\{\phi_t\}_{t\geq 0}$ be a sequence in \Re^d satisfying $\sup_{t\geq 0} ||\phi_t|| \leq 1$. For any $t\geq 0$, we define $\Lambda_t = \Lambda_0 + \sum_{j=0}^t \phi_j \phi_j^T \phi_j$. Then if the smallest eigen value of Λ_0 be at least 1, we have

$$\log\left[\frac{\det(\Lambda_h^{k+1})}{\det(\Lambda_h^1)}\right] \le \sum_{k=1}^K (\phi_h^k)^T (\Lambda_h^k)^{-1} \phi_h^k \le 2\log\left[\frac{\det(\Lambda_h^{k+1})}{\det(\Lambda_h^1)}\right]$$
(96)

The next result characterizes the covering number of an Euclidean ball (Lemma 5.2 in Vershynin (2010)).

Lemma 22 [Covering Number of Euclidean Ball] For any $\epsilon > 0$, the ϵ -covering number of the Euclidean ball in \mathbb{R}^d with radius R is upper bounded by $(1 + 2R/\epsilon)^d$.

Appendix F. Why does the Greedy Policy fail?

In this section, using an example we show that the greedy-policy is not Lipschitz. Further, we illustrate that (for leader) we can not use the greedy-policy for the follower based on the joint Q-function to obtain the ϵ -close covering for the leader's value function. In the similar realm, we can construct an example where the greedy-policy based on the leader's marginal q-function would not provide ϵ -covering number for the follower's value function.

Consider the following toy-example:

Example 1 We basically show that we can not obtain ϵ -covering number for marginal Q-function (q_l) for the leader if the follower's action is based on the greedy policy. Suppose that the cardinality of the action space $|\mathcal{B}|$, $|\mathcal{A}|$ are 2. Further, suppose that Q-table is obtained as in Table 1. Now,

 Table 1: The Q-table for leader and follower, first number are for leader, second numbers are for follower

ACTIONS	b_1	b_2
$a_1 \\ a_2$	$\begin{array}{c} M-\epsilon, 1-\epsilon/2\\ 0,1 \end{array}$	$\begin{array}{c} 0,1+\epsilon/2\\ 0,1-\epsilon/2 \end{array}$

consider ϵ -close values (\tilde{Q}) are obtained as in Table 2. Now, we will be able to show how the

Table 2: The ϵ close values \tilde{Q} -table for leader and follower. The first numbers are for leader, second numbers are for follower

ACTIONS	b_1	b_2
$egin{array}{c} a_1 \ a_2 \end{array}$	$\begin{array}{c} M, 1+\epsilon/2 \\ 0, 1+\epsilon/2 \end{array}$,

marginal q_l values change if the policy is greedy. For example for a_1 , if the follower's policy would have been greedy, $q_l(x, a_1) = M$ for Table 1, and $q_l(x, a_1) = 0$ for values in Table 2. Thus, slight changes in the values of Q can lead to a drastic changes in the values of the marginal q values. Also note that M is an arbitrary number, thus, such change can be arbitrary.

If the policy is soft-max, we can easily show that q_l values are also close.

Appendix G. Why does the approach proposed in Zhong et al. (2021) fail?

In Zhong et al. (2021), at every step they find a stackelberg equilibrium using an oracle for an ϵ approximated Q function for the leader and known rewards of the follower (they consider myopic
follower with known rewards). In particular, Zhong et al. (2021) proposed an approach where they
truncate $w_{l,h}^k$, Λ_h^k to ϵ -close value of w_l , and Λ respectively. Subsequently, they obtained equilibrium
policies using these ϵ -close values. Then the proposed algorithm uses the above equilibrium strategy

attained using ϵ -close values as their policies. Since these ϵ -close values are predetermined using the ϵ -covering set of w and Λ (as we have described the ϵ -covering set in Lemma 15), hence, one can apply uniform concentration lemma for the leader Zhong et al. (2021) with error of at most ϵ .

However, we can not apply the similar trick since here the followers are non-myopic, and the rewards are unknown. In particular, in Zhong et al. (2021), the followers' Q-functions are not required to be estimated as the rewards are known. Thus, the followers' policies are fixed given the action of the leader. However, in our case, as shown in Example 1, slight change in the Q-function can lead to a change in the action for the follower. Thus, an oracle which computes Stackelberg equilibrium, it would also gives a different action which in turn would make the marginal Q-function for the leader arbitrary large as shown in Example 1. Thus, one can not show that the log ϵ -covering number for the leader's value functions scales with $\log(K)$.

Appendix H. Proof of Corollary 1

First, consider the policy– $(\tilde{\pi}_l, \tilde{\pi}_f)$ where both the leader and the follower pick any $k \in [K]$ randomly and then follow the policy π_l^k, π_f^k given by the Algorithms 1 and 2 at episode k.

Note from Theorem 3 that

$$\frac{1}{K} \left(\sum_{k} \sum_{a_{1}^{k}} \pi_{l}^{k}(a_{1}^{k}|x_{1})(\bar{V}_{f,1}^{\pi_{l}^{k},\pi_{f}^{k,*}}(x_{1},a_{1}^{k}) - \bar{V}_{f,1}^{\pi_{l}^{k},\pi_{f}^{k}}(x_{1},a_{1}^{k})) \right) \leq \tilde{\mathcal{O}}(\sqrt{d^{3}H^{4}})/\sqrt{K}$$

$$V_{f,1}^{\tilde{\pi}_{l},\pi_{f}^{*}}(x_{1}) - V_{f,1}^{\tilde{\pi}_{l},\tilde{\pi}_{f}}(x_{1}) \leq \tilde{\mathcal{O}}(\sqrt{d^{3}H^{4}})/\sqrt{K}$$
(97)

By selecting $K = \tilde{\mathcal{O}}(d^3H^4/\epsilon^2)$ we have the result.

Similarly, for the leader we have

$$\frac{1}{K} \left(\sum_{k} \bar{V}_{l,1}^{\pi_{l}^{k,*},\pi_{f}^{k}}(x_{1}) - \bar{V}_{l,1}^{\pi_{l}^{k},\pi_{f}^{k}}(x_{1}) \right) \leq \tilde{\mathcal{O}}(\sqrt{d^{3}H^{4}}) / \sqrt{K}$$

$$V_{l,1}^{\pi_{l}^{*},\tilde{\pi}_{f}}(x_{1}) - V_{l,1}^{\tilde{\pi}_{l},\tilde{\pi}_{f}}(x_{1}) \leq \tilde{\mathcal{O}}(\sqrt{d^{3}H^{4}}) / \sqrt{K}$$
(98)

Again by plugging in $K = \tilde{\mathcal{O}}(d^3H^4/\epsilon^2)$ we have the result.