

On the multiplicative independence between n and $\lfloor \alpha n \rfloor$

DAVID CRNČEVIĆ and FELIPE HERNÁNDEZ and KEVIN RIZK
and KHUNPOB SEREESUCHART and RAN TAO

December 8, 2023

Abstract

In this article we investigate different forms of multiplicative independence between the sequences n and $\lfloor n\alpha \rfloor$ for irrational α . Our main theorem shows that for a large class of arithmetic functions $a, b: \mathbb{N} \rightarrow \mathbb{C}$ the sequences $(a(n))_{n \in \mathbb{N}}$ and $(b(\lfloor \alpha n \rfloor))_{n \in \mathbb{N}}$ are asymptotically uncorrelated. This new theorem is then applied to prove a 2-dimensional version of the Erdős-Kac theorem, asserting that the sequences $(\omega(n))_{n \in \mathbb{N}}$ and $(\omega(\lfloor \alpha n \rfloor))_{n \in \mathbb{N}}$ behave as independent normally distributed random variables with mean $\log \log n$ and standard deviation $\sqrt{\log \log n}$. Our main result also implies a variation on Chowla's Conjecture asserting that the logarithmic average of $(\lambda(n)\lambda(\lfloor \alpha n \rfloor))_{n \in \mathbb{N}}$ tends to 0.

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 1 |
| 2 | Preliminaries | 4 |
| 2.1 | Notation | 4 |
| 2.2 | Averaging | 5 |
| 2.3 | Almost Periodicity | 12 |
| 2.4 | BMAI sequences | 14 |
| 3 | Main Result | 18 |
| 3.1 | Basic Case | 18 |
| 3.2 | Main Theorem | 20 |
| 4 | Consequences of the main theorem | 25 |
| 4.1 | Erdős-Kac generalization | 25 |
| 4.2 | Logarithmic averages of the Liouville function | 31 |
| 5 | Open Questions | 32 |

1. Introduction

For a set $A \subseteq \mathbb{N}$ we say A has *density* $d(A)$ if $d(A) := \lim_{n \rightarrow \infty} \frac{|A \cap \{1, \dots, N\}|}{N}$ exists. Even though d is not a probability measure, since it is not countably additive, it can be used to measure

the size of infinite sets of integers. For instance, if we define A_p to be the set of positive integers divisible by $p \in \mathbb{P}$, then A_p has density $d(A_p) = 1/p$.

It is believed that the sequences $(n)_{n \in \mathbb{N}}$ and $(\lfloor \alpha n \rfloor)_{n \in \mathbb{N}}$, for $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, have independent multiplicative structure. The following classical result of G. L. Watson [Wat53] from 1953 supports this claim, showing that the proportion of natural numbers $n \in \mathbb{N}$ such that n and $\lfloor \alpha n \rfloor$ are coprime is $6/\pi^2$, which is the same probability of two arbitrary integers being relatively prime.

Theorem 1.1 (see [Wat53]). *For every $\alpha \in \mathbb{R} \setminus \mathbb{Q}$:*

$$d(\{n \in \mathbb{N} \mid (n, \lfloor \alpha n \rfloor) = 1\}) = \frac{6}{\pi^2},$$

where (a, b) denotes the greatest common divisor between $a, b \in \mathbb{N}$.

In 1917, Hardy and Ramanujan [HR17] showed the following theorem which describes the asymptotic behaviour of the function $\omega(n)$ which counts distinct prime factors of n .

Theorem 1.2 (see [HR17]). *Let $\varepsilon > 0$, then for almost all $n \in \mathbb{N}$*

$$|\omega(n) - \log \log n| < (\log \log n)^{\frac{1}{2} + \varepsilon}.$$

In 1940 P. Erdős and M. Kac proved their well-known result about the Gaussian law associated to the function ω , stated here.

Theorem 1.3 (see [EK40]). *For all $a, b \in \mathbb{R}$ where $a < b$,*

$$d\left(\left\{n \in \mathbb{N} : a \leq \frac{\omega(n) - \log \log n}{(\log \log n)^{1/2}} \leq b\right\}\right) = \frac{1}{\sqrt{2\pi}} \int_a^b e^{-t^2/2} dt.$$

Later in 2007, William D. Banks and Igor E. Shparlinski showed that one can replace n in the Erdős-Kac theorem by any Beatty sequence (see Definition 2.3).

Theorem 1.4 (see [BS07, Theorem 4]). *Let $\alpha \in \mathbb{R}$ be a positive irrational real number, then*

$$d\left(\left\{n \in \mathbb{N} : a \leq \frac{\omega(\lfloor \alpha n \rfloor) - \log \log n}{(\log \log n)^{1/2}} \leq b\right\}\right) = \frac{1}{\sqrt{2\pi}} \int_a^b e^{-t^2/2} dt.$$

Our first result unifies the theorem of Banks and Shparlinski with the original result of Erdős-Kac, following the heuristic that the sequences n and $\lfloor \alpha n \rfloor$, for α irrational, behave independently.

Theorem A. *Let $\alpha \in \mathbb{R}$ be a positive irrational real number and a, b, c, d real numbers. Then*

$$\begin{aligned} d\left(\left\{n \in \mathbb{N} : a \leq \frac{\omega(\lfloor \alpha n \rfloor) - \log \log n}{(\log \log n)^{1/2}} \leq b\right\} \cap \left\{n \in \mathbb{N} : c \leq \frac{\omega(n) - \log \log n}{(\log \log n)^{1/2}} \leq d\right\}\right) \\ = \left(\frac{1}{\sqrt{2\pi}} \int_a^b e^{-t^2/2} dt\right) \left(\frac{1}{\sqrt{2\pi}} \int_c^d e^{-t^2/2} dt\right). \end{aligned}$$

For our second result, we took inspiration from a long-standing conjecture in number theory attributed to Chowla.

Conjecture 1.5 (Chowla's conjecture [see [Cho65, Problem 57], [Tao16]]). *Let $k \geq 1$ be a natural number.*

1. (Chowla's conjecture) *If $h_1, \dots, h_k \in \mathbb{Z}$ are distinct integers, then*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \lambda(n + h_1) \cdots \lambda(n + h_k) = 0.$$

2. (Logarithmically averaged Chowla's conjecture) *If $h_1, \dots, h_k \in \mathbb{Z}$ are distinct integers, then*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N}^{\log} \lambda(n + h_1) \cdots \lambda(n + h_k) = 0,$$

where λ is the Liouville function (see Definition 2.2) and $\mathbb{E}_{n \leq N}$, $\mathbb{E}_{n \leq N}^{\log}$ are standard and logarithmic averages (see Definition 2.5).

Remark 1.6. The logarithmically averaged Chowla's conjecture was first stated by Tao in [Tao16]. Furthermore, Chowla's conjecture (with Cesàro averages) implies the logarithmically averaged version.

Chowla's conjecture is known to be true only for $k = 1$, which case is elementally equivalent to the prime number theorem. In addition, due to recent developments, we know that the logarithmically averaged Chowla's conjecture holds for $k = 2$ [Tao16] and for odd values of k [TT18]. Inspired by this recent breakthrough, we established the following result.

Theorem B. *If α is an irrational positive real number, then*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N}^{\log} \lambda(n) \lambda(\lfloor \alpha n \rfloor) = 0,$$

where $[N] := \{1, \dots, N\}$.

Both Theorem A and Theorem B are consequences of a more general theorem about a class of bounded sequences $a : \mathbb{N} \rightarrow \mathbb{C}$ which we call **BMAI** (*Bounded Multiplicative Almost Invariant*). Among other things, this class of sequences includes all multiplicative functions which don't take the value 0; for more details and further examples, see Definition 2.23.

Theorem C. *Let $a : \mathbb{N} \rightarrow \mathbb{C}$ and $b : \mathbb{N} \rightarrow \mathbb{S}^1$ be BMAI sequences such that:*

1. $\lim_{H \rightarrow \infty} \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |\mathbb{E}_{h \leq H} a(n + h)| = 0$; and
2. for all $\beta \in \mathbb{R} \setminus \mathbb{Q}$ and $h \in \mathbb{N}$ we have

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} a(n) \overline{a(n + h)} e(n\beta) = 0 \quad (\text{resp. } \mathbb{E}_{n \leq N}^{\log} \bullet).$$

Then, for any irrational $\alpha \in \mathbb{R}$,


$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} b(n) a(\lfloor \alpha n \rfloor) = 0. \quad (\text{resp. replacing } \mathbb{E}_{n \leq N} \text{ with } \mathbb{E}_{n \leq N}^{\log} \bullet) \quad (1.1)$$

To derive Theorem A from Theorem C, we take $a(n) = G(\frac{\omega(n) - \log \log n}{(\log \log n)^{1/2}})$ and $b(n) = F(\frac{\omega(n) - \log \log n}{(\log \log n)^{1/2}})$ for certain functions $F, G : \mathbb{N} \rightarrow \mathbb{C}$, and employ a generalization of the

Erdős-Kac theorem on short intervals to verify that the hypothesis of Theorem C are satisfied for this choice of a and b . The details of this derivation are given in Section 3 and Section 4.

Likewise, for the proof of Theorem B we take $a(n) = b(n) = \lambda(n)$ in Theorem C and use Theorem 4.5 and Lemma 4.6. For the details of the proof of Theorem B, we refer the reader to Section 3 and Section 4 as well.

Finally, since there are many intermediate steps involved in the proof of Theorem C, we include here a diagram illustrating the path to our main results.



Acknowledgements

We would like to thank the Bernoulli Center for funding the Young Researchers in Mathematics Summer Program, which allowed us to develop such results. We would also like to thank Florian Richter and Joel Moreira for their guidance throughout the program and for making this project possible. Additionally, we owe a special thank you to Noah Kravitz and Katharine Woo for providing detailed comments and suggestions.

2. Preliminaries

2.1. Notation

First, we introduce some classical functions in number theory.

Definition 2.1 (Prime omega functions). We define the prime omega functions $\Omega, \omega : \mathbb{N} \rightarrow \mathbb{N}$ as follows. For $n \in \mathbb{N}$, we write $n = p_1^{a_1} \cdots p_k^{a_k}$ where p_1, \dots, p_k are distinct primes numbers

and $a_j \in \mathbb{N}$ for $j = 1, \dots, k$. We define

$$\omega(n) := k.$$

In other words, $\omega(n)$ is the number of distinct prime divisors of n . We also define

$$\Omega(n) := \sum_{j=1}^k a_j,$$

which is the numbers of prime divisors with multiplicity.

Definition 2.2 (Liouville function). The classical Liouville function is defined as

$$\lambda(n) := (-1)^{\Omega(n)}.$$

Throughout the paper we will use $e(x) := e^{2\pi i x}$, $x \in \mathbb{R}$, for the complex exponential function. For a real number α , we will denote $\lfloor \alpha \rfloor$ the greatest integer less than or equal to α , and $\{\alpha\} = \alpha - \lfloor \alpha \rfloor$ the fractional part. We now define the notion of a Beatty sequence.

Definition 2.3 (Beatty Sequence). For $\alpha \in \mathbb{R}_+$ the corresponding Beatty sequence is the sequence of integers defined by

$$\mathcal{B}_\alpha = (\lfloor \alpha n \rfloor)_{n \in \mathbb{N}}.$$

Finally, we introduce some classical definitions to measure how fast a sequence goes to 0 and how bounded the growth rate of a sequence is.

Definition 2.4. For a function $g : \mathbb{N} \rightarrow \mathbb{R}$, which is strictly positive for all large enough values of n , we use the classic notation $f(N) = o_{N \rightarrow \infty}(g(N))$ to denote a function $f : \mathbb{N} \rightarrow \mathbb{R}$ such that

$$\lim_{N \rightarrow \infty} \frac{f(N)}{g(N)} = 0.$$

Additionally, we denote $f(N) = O(g(N))$, if there exists a positive real number C and an integer $M \in \mathbb{N}$ such that

$$|f(N)| \leq Cg(N), \forall N \geq M.$$

2.2. Averaging

In our proofs, we will sometimes use expectation notation to denote averages. In particular, we define here the Cesàro and the logarithmic averages.

Definition 2.5. For a function $f : \mathbb{N} \rightarrow \mathbb{C}$ and $N, S \in \mathbb{N}$, we define its

- *average* as

$$\mathbb{E}_{n \leq N}(f) := \frac{1}{N} \sum_{n=1}^N f(n),$$

- *logarithmic average* as

$$\mathbb{E}_{n \leq N}^{\log}(f) := \frac{1}{\log N} \sum_{n=1}^N \frac{f(n)}{n},$$

- *logarithmic expectation over the primes* as

$$\mathbb{E}_{p \leq S}^{\log}(g) := \frac{1}{\log \log S} \sum_{p \leq S} \frac{1}{p} g(p),$$

where p only takes prime values less than or equal to S .

Remark 2.6. When the limit $\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N}(f)$ exists, it is called the *Cesàro average* of the sequence $f : \mathbb{N} \rightarrow \mathbb{C}$. In this case, the limit $\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N}^{\log}(f)$ exists as well.

We will start by stating and proving results which relates the Cesàro average and the logarithmic average.

Lemma 2.7. *We have for $f : \mathbb{N} \rightarrow \mathbb{C}$ bounded,*

$$\mathbb{E}_{n \leq N}^{\log} f(n) = \mathbb{E}_{M \leq N}^{\log} \mathbb{E}_{n \leq M} f(n) + o_{N \rightarrow \infty}(1).$$

Proof. By expanding the sum $\mathbb{E}_{M \leq N}^{\log} \mathbb{E}_{n \leq M} f(n)$, we get

$$\begin{aligned} \mathbb{E}_{M \leq N}^{\log} \mathbb{E}_{n \leq M} f(n) &= \frac{1}{\log N} \sum_{M=1}^N \frac{1}{M^2} \sum_{n=1}^M f(n) \\ &= \frac{1}{\log N} \sum_{n=1}^N f(n) \sum_{M=n}^N \frac{1}{M^2}. \end{aligned}$$

Using

$$\sum_{M=n}^N \frac{1}{M^2} = \frac{1}{n} + \frac{1}{N} + O(1/n^2),$$

and that f is bounded, we get:

$$\begin{aligned} \mathbb{E}_{M \leq N}^{\log} \mathbb{E}_{n \leq M} f(n) &= \frac{1}{\log N} \sum_{n=1}^N \left(\frac{f(n)}{n} + \frac{f(n)}{N} + O(1/n^2) \right) \\ &= \mathbb{E}_{n \leq N}^{\log} f(n) + \frac{1}{N \log N} \sum_{n=1}^N O(1) + \frac{1}{\log N} \sum_{n=1}^N O(1/n^2) \\ &= \mathbb{E}_{n \leq N}^{\log} f(n) + o_{N \rightarrow \infty}(1). \end{aligned}$$

□

Lemma 2.8. *Let $f : \mathbb{N} \rightarrow \mathbb{C}$ a bounded function. Then*

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N}^{\log} |f(n)| \leq \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |f(n)|.$$

Proof. For each $N \in \mathbb{N}$, define

$$M_N = \operatorname{argmax}\{\mathbb{E}_{n \leq M} |f(n)| \mid \log \log N \leq M \leq N\}.$$

By Lemma 2.7 we have that:

$$\begin{aligned}
\mathbb{E}_{n \leq N}^{\log} |f(n)| &= \mathbb{E}_{M \leq N}^{\log} \mathbb{E}_{n \leq M} |f(n)| + o_{N \rightarrow \infty}(1) \\
&= \frac{1}{\log N} \sum_{M < \log \log N} \frac{1}{M} \mathbb{E}_{n \leq M} |f(n)| + \frac{1}{\log N} \sum_{M = \log \log N}^N \frac{1}{M} \mathbb{E}_{n \leq M} |f(n)| + o_{N \rightarrow \infty}(1) \\
&\leq \frac{\|f\|_{\ell^\infty} \log \log N}{\log N} + \left(\frac{1}{\log N} \sum_{M = \log \log N}^N \frac{1}{M} \right) \left(\mathbb{E}_{n \leq M_N} |f(n)| \right) + o_{N \rightarrow \infty}(1),
\end{aligned}$$

applying \limsup_N yields

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N}^{\log} |f(n)| \leq \limsup_N \mathbb{E}_{n \leq M_N} |f(n)| \leq \limsup_N \mathbb{E}_{n \leq N} |f(n)|,$$

as desired. \square

The following corollary will be useful later on.

Corollary 2.9. *Let $S \subseteq \mathbb{N}$ such that $|S| = \infty$ and let $f : \mathbb{N} \times S \rightarrow \mathbb{C}$ such that for all $t \in S$, $f(\cdot, t)$ is a bounded function. We have that*

$$\limsup_{T \rightarrow \infty, T \in S} \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N}^{\log} |f(n, T)| \leq \limsup_{T \rightarrow \infty, T \in S} \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |f(n, T)|.$$

The rest of this subsection will contain some useful known results.

Lemma 2.10 (see [BM16, Theorem 2.4]). *Let $(u_n)_{n \in \mathbb{N}}$ be a bounded sequence in \mathbb{C} . If for all $h \in \mathbb{N}$*

$$\frac{1}{N} \sum_{n=1}^N u_{n+h} \overline{u_n} \rightarrow 0, \tag{2.1}$$

then

$$\frac{1}{N} \sum_{n=1}^N u_n \rightarrow 0. \tag{2.2}$$

The following lemmas, scattered throughout the literature, will allow us to transition between Cesàro averages and logarithmic expectations over the primes.

Lemma 2.11. *For $S, N \in \mathbb{N}$ with $N > S$, let $L(S) = \sum_{p \leq S} \frac{1}{p}$, we have*

$$\mathbb{E}_{n \leq N} \left| \sum_{p \leq S, p|n} 1 - L(S) \right|^2 \leq E(S, N), \tag{2.3}$$

with $E(S, N) = O(\log \log S) + \frac{1}{N} O(S^2)$. We also have

$$\mathbb{E}_{n \leq N}^{\log} \left| \sum_{p \leq S, p|n} 1 - L(S) \right|^2 \leq E'(S, N), \tag{2.4}$$

with $E'(S, N) = O(\log \log S) + \frac{1}{\log(N)} O(S^2)$.

Proof. Eq. (2.3) follows from [Eil79, Lemma 4.1] applied to the additive arithmetic function $f(n) := \sum_{p \leq S, p|n} 1$.

Now let's prove Eq. (2.4).

First, we estimate $\sum_{n \leq N} \sum_{p \leq S, p|n} \frac{1}{n}$:

$$\begin{aligned} \sum_{p \leq S} \sum_{n \leq N, p|n} \frac{1}{n} &= \sum_{p \leq S} \sum_{n \leq N/p} \frac{1}{np} = \sum_{p \leq S} \frac{1}{p} \sum_{n \leq N/p} \frac{1}{n} \\ &= \sum_{p \leq S} \frac{1}{p} (\log(N) - \log(p) + O(1)) = L(S) \log(N) + O(S). \end{aligned}$$

Second, we estimate $\sum_{n \leq N} \sum_{p, q \leq S, p, q|n} \frac{1}{n}$:

$$\begin{aligned} \sum_{n \leq N} \sum_{p, q \leq S, p, q|n} \frac{1}{n} &= \sum_{p \leq S} \sum_{q \leq S} \sum_{n \leq N, p, q|n} \frac{1}{n} \\ &= \sum_{p \leq S} \sum_{n \leq N, p|n} \frac{1}{n} + \sum_{p \leq S} \sum_{q \leq S, q \neq p} \left(\sum_{n \leq N/pq} \frac{1}{pqn} \right) \\ &\leq L(S) \log(N) + O(S) + L(S)^2 \log(N) + O(S^2). \end{aligned}$$

Finally, we estimate $\frac{1}{\log N} \sum_{n \leq N} \frac{1}{n} \left| \sum_{p \leq S, p|n} 1 - L(S) \right|^2$:

$$\begin{aligned} \frac{1}{\log N} \sum_{n \leq N} \frac{\left| \sum_{p \leq S, p|n} 1 - L(S) \right|^2}{n} \\ = \frac{1}{\log N} \sum_{n \leq N} \sum_{p, q \leq S, p, q|n} \frac{1}{n} - \left(2L(S) \frac{1}{\log N} \sum_{n \leq N} \sum_{p \leq S, p|n} \frac{1}{n} \right) + L(S)^2. \end{aligned}$$

Now using our estimations, we get

$$\begin{aligned} \frac{1}{\log N} \sum_{n \leq N} \frac{\left| \sum_{p \leq S, p|n} 1 - L(S) \right|^2}{n} &\leq L(S) + \frac{1}{\log N} O(S) + \frac{1}{\log N} O(S^2) \\ &\quad + L(S)^2 - 2L(S)^2 + \frac{L(S)}{\log N} O(S) + L(S)^2 \\ &= L(S) + \frac{1}{\log N} O(S^2), \end{aligned}$$

and as $L(S) = O(\log \log S)$, we get the desired result. □

Remark 2.12. We note that the above lemma still holds when dropping the condition $N > S$ by observing that when $p > N$, $\sum_{n \leq \frac{N}{p}} \frac{1}{n} = 0$.

Lemma 2.13. For any bounded function $f : \mathbb{N} \rightarrow \mathbb{C}$, we have that

$$\frac{1}{N} \sum_{n=1}^N f(n) = \mathbb{E}_{p \leq S}^{\log} \frac{1}{N/p} \sum_{n=1}^{N/p} f(pn) + \varepsilon(S, N), \quad (2.5)$$

with $\limsup_{S \rightarrow \infty} \limsup_{N \rightarrow \infty} \varepsilon(S, N) = 0$. We also have that

$$\mathbb{E}_{n \leq N}^{\log} f(n) = \mathbb{E}_{p \leq S}^{\log} \mathbb{E}_{n \leq N/p}^{\log} f(pn) + \varepsilon'(S, N), \quad (2.6)$$

with $\limsup_{S \rightarrow \infty} \limsup_{N \rightarrow \infty} \varepsilon'(S, N) = 0$.

Proof. Let $L(S) = \sum_{p \leq S} \frac{1}{p}$, Lemma 2.11 gives

$$\frac{1}{N} \sum_{n \leq N} \left| \sum_{p \leq S, p|n} 1 - L(S) \right|^2 \leq E(S, N).$$

By Cauchy-Schwarz, and as f is bounded we get

$$\left| \frac{1}{N} \sum_{n \leq N} f(n) \left(\sum_{p \leq S, p|n} 1 - L(S) \right) \right| \leq CE(S, N)^{\frac{1}{2}},$$

for some constant C independent of N and S . Now dividing by $L(S)$, we get

$$\left| \frac{1}{N} \sum_{n \leq N} f(n) - \frac{1}{L(S)} \frac{1}{N} \sum_{n \leq N} \sum_{p \leq S, p|n} f(n) \right| \leq \frac{1}{L(S)} CE(S, N)^{\frac{1}{2}},$$

where the second term on the left can be written as

$$\begin{aligned} \frac{1}{L(S)N} \sum_{n \leq N} \sum_{p \leq S, p|n} f(n) &= \frac{1}{L(S)} \sum_{p \leq S} \frac{1}{N} \sum_{n \leq N, p|n} f(n) = \frac{1}{L(S)} \sum_{p \leq S} \frac{1}{p} \frac{1}{N/p} \sum_{n \leq N/p} f(pn) \\ &= \mathbb{E}_{p \leq S}^{\log} \frac{1}{N/p} \sum_{n=1}^{N/p} f(pn). \end{aligned}$$

The only thing left to show is that the error term goes to zero as desired. Indeed,

$$\begin{aligned} \limsup_{S \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{1}{L(S)} CE(S, N)^{\frac{1}{2}} &= \limsup_{S \rightarrow \infty} \frac{1}{L(S)} \left(\limsup_{N \rightarrow \infty} O(\log \log S) + \frac{1}{N} O(S^2) \right)^{\frac{1}{2}} \\ &= \limsup_{S \rightarrow \infty} O\left((\log \log S)^{-\frac{1}{2}}\right) = 0, \end{aligned}$$

where we used that $L(S) = O(\log \log S)$ to conclude.

The proof for (2.6) is similar. □

Lemma 2.14. *For any bounded $f : \mathbb{N} \rightarrow \mathbb{C}$ and for any $k, N \in \mathbb{N}$, we have*

$$\mathbb{E}_{n \leq N}^{\log} f(n) = \mathbb{E}_{n \leq N/k}^{\log} f(n) + E(N, k)$$

such that $\limsup_{N \rightarrow \infty} E(N, k) = 0$.

Proof. By using the definition of logarithmic average, we have

$$E(N, k) = \left(\frac{1}{\log N} - \frac{1}{\log(N/k)} \right) \sum_{n=1}^{N/k} \frac{f(n)}{n} + \frac{1}{\log N} \sum_{n=N/k}^N \frac{f(n)}{n}.$$

We will start by showing that the first part of the sum tends to zero as N goes to infinity. Indeed, we have

$$\left(\frac{1}{\log N} - \frac{1}{\log(N/k)} \right) \sum_{n=1}^{N/k} \frac{f(n)}{n} = \frac{1}{\log N} \sum_{n=1}^{N/k} \frac{f(n)}{n} \left(\frac{\log k}{\log N - \log k} \right), \quad (2.7)$$

but as f is bounded, we have that $\frac{1}{\log N} \sum_{n=1}^{N/k} \frac{f(n)}{n}$ is bounded, and as $\left(\frac{\log k}{\log N - \log k} \right)$ tends to 0 as N tends to infinity, we get that (2.7) goes to zero as $N \rightarrow \infty$.

As for the second part of the sum, as f is bounded, we have that $|\limsup_{N \rightarrow \infty} \sum_{n=N/k}^N \frac{f(n)}{n}| \leq \max_{n \in \mathbb{N}} |f(n)| \log k$ and so we indeed get that

$$\limsup_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=N/k}^N \frac{f(n)}{n} = 0.$$

□

Using Lemma 2.13, we can derive the following orthogonality criterion, which has appeared in [Kát86], [DD82] and [BSZ13].

Lemma 2.15. *There exists C such that for every bounded function $f : \mathbb{N} \rightarrow \mathbb{C}$ and $N, S \in \mathbb{N}$*

$$|\mathbb{E}_{n \leq N} f(n)| \leq C \sum_{l=1}^{\log_2 S} \frac{1}{l \log \log S} \left(\max_{p, q \in [2^l, 2^{l+1})} \left| \mathbb{E}_{n \leq \frac{N}{\max\{p, q\}}} f(pn) \overline{f(qn)} \right| \right)^{1/2} + \varepsilon(S, N), \quad (2.8)$$

where $\limsup_{S \rightarrow \infty} \limsup_{N \rightarrow \infty} \varepsilon(S, N) = 0$.

We also have the following logarithmic version of the inequality:

$$\left| \mathbb{E}_{n \leq N}^{\log} f(n) \right| \leq \left(\mathbb{E}_{p \leq S}^{\log} \mathbb{E}_{q \leq S}^{\log} \left| \mathbb{E}_{n \leq N}^{\log} f(pn) \overline{f(qn)} \right| \right)^{1/2} + \varepsilon'(S, N) \quad (2.9)$$

where $\limsup_{S \rightarrow \infty} \limsup_{N \rightarrow \infty} \varepsilon'(S, N) = 0$.

Proof. Let $A_l := [2^l, 2^{l+1})$.

$$\begin{aligned}
|\mathbb{E}_{n \leq N} f(n)| &= \left| \mathbb{E}_{p \leq S}^{\log} \mathbb{E}_{n \leq N/p} f(pn) \right| + \varepsilon(S, N) \text{ by Lemma 2.13} \\
&\leq \sum_{l=1}^{\log_2 s} \left| \mathbb{E}_{p \leq S}^{\log} \mathbb{1}_{A_l}(p) \left(\frac{2^l}{p} \mathbb{E}_{n \leq N/2^l} \mathbb{1}_{pn \leq N} f(pn) \right) \right| + \varepsilon(S, N) \\
&= \sum_{l=1}^{\log_2 s} \left| \mathbb{E}_{n \leq N/2^l} \mathbb{E}_{p \leq S}^{\log} \mathbb{1}_{A_l}(p) \left(\frac{2^l}{p} \mathbb{1}_{pn \leq N} f(pn) \right) \right| + \varepsilon(S, N) \\
&\leq \sum_{l=1}^{\log_2 s} \left(\mathbb{E}_{n \leq N/2^l} \left| \mathbb{E}_{p \leq S}^{\log} \mathbb{1}_{A_l}(p) \left(\frac{2^l}{p} \mathbb{1}_{pn \leq N} f(pn) \right) \right|^2 \right)^{1/2} + \varepsilon(S, N) \\
&= \sum_{l=1}^{\log_2 s} \left(\mathbb{E}_{n \leq N/2^l} \mathbb{E}_{p, q \leq S}^{\log} \mathbb{1}_{A_l}(p) \mathbb{1}_{A_l}(q) \left(\frac{4^l}{pq} \mathbb{1}_{pn \leq N} \mathbb{1}_{qn \leq N} f(pn) \overline{f(qn)} \right) \right)^{1/2} + \varepsilon(S, N) \\
&\leq 2 \sum_{l=1}^{\log_2 s} \left(\mathbb{E}_{p, q \leq S}^{\log} \mathbb{1}_{A_l}(p) \mathbb{1}_{A_l}(q) \left(\max_{p, q \in A_l} |\mathbb{E}_{n \leq N/2^l} \mathbb{1}_{\leq N}(pn) \mathbb{1}_{\leq N}(qn) f(pn) \overline{f(qn)}| \right) \right)^{1/2} \\
&\quad + \varepsilon(S, N).
\end{aligned}$$

We can expand

$$\mathbb{E}_{p, q \leq S}^{\log} \mathbb{1}_{A_l}(p) \mathbb{1}_{A_l}(q) = \frac{1}{(\log \log S)^2} \sum_{p \in A_l} \frac{1}{p} \sum_{q \in A_l} \frac{1}{q}.$$

Using the prime number theorem, we also have

$$\sum_{p \in A_l} \frac{1}{p} = \sum_{q \in A_l} \frac{1}{q} \leq \sum_{q \in A_l} \frac{1}{2^l} \leq \theta \frac{1}{l},$$

giving (3.2).

Now let's prove (2.9):

$$\begin{aligned}
\left| \mathbb{E}_{n \leq N}^{\log} f(n) \right| &= \left| \mathbb{E}_{p \leq S}^{\log} \mathbb{E}_{n \leq N/p}^{\log} f(pn) \right| + \varepsilon(S, N) \text{ by Lemma 2.13} \\
&\leq \left| \mathbb{E}_{n \leq N}^{\log} \mathbb{E}_{p \leq S}^{\log} f(pn) \right| + \left| \mathbb{E}_{p \leq S}^{\log} E(N, p) \right| + \varepsilon(S, N) \text{ by Lemma 2.14} \\
&\leq \left(\mathbb{E}_{n \leq N}^{\log} \left| \mathbb{E}_{p \leq S}^{\log} f(pn) \right|^2 \right)^{1/2} + \left| \mathbb{E}_{p \leq S}^{\log} E(N, p) \right| + \varepsilon(S, N) \\
&= \left(\mathbb{E}_{n \leq N}^{\log} \mathbb{E}_{p \leq S}^{\log} \mathbb{E}_{q \leq S}^{\log} f(pn) \overline{f(qn)} \right)^{1/2} + \varepsilon'(S, N) \\
&\leq \left(\mathbb{E}_{p \leq S}^{\log} \mathbb{E}_{q \leq S}^{\log} \left| \mathbb{E}_{n \leq N}^{\log} f(pn) \overline{f(qn)} \right| \right)^{1/2} + \varepsilon'(S, N).
\end{aligned}$$

□

Next is a powerful theorem from [Tan60], formalizing the independent behavior of polynomials with regard to the function Ω .

Theorem 2.16 (see [Tan60, Theorem 1]). Let $\{f_i : i \leq k\}$ be a finite family of pairwise relatively prime non-constant polynomials, where f_i is a product of $r_i \geq 1$ irreducible polynomials.

Let

$$u_i(n) := \frac{\Omega(f_i(n)) - r_i \log \log n}{\sqrt{r_i \log \log n}}, \quad (2.10)$$

then for $E \subseteq \mathbb{R}^k$ Jordan measurable,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \#\{n \leq N : (u_i(n))_i \in E\} = \frac{1}{2\pi^{\frac{k}{2}}} \int_E \exp\left(\frac{-1}{2} \sum_{i=1}^k x_i^2\right) dx_1 \dots dx_k. \quad (2.11)$$

Remark 2.17. Theorem 2.16 is also true when Ω is replaced by ω , proven in the same paper.

2.3. Almost Periodicity

We now recall the notion of almost periodic function, which plays an important role in the sequel. Given $g : \mathbb{N} \rightarrow \mathbb{C}$, we define its 1-norm as

$$\|g\|_1 := \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |g(n)|.$$

Definition 2.18. Let $f : \mathbb{N} \rightarrow \mathbb{C}$ be an arithmetic function. We say that f is *almost periodic* (sometimes also referred to as *Besicovitch almost periodic* in the literature) if for each $\varepsilon > 0$ there is some linear combination h over \mathbb{C} of exponential functions $e(\alpha n)$, $\alpha \in \mathbb{R}$, such that $\|f - h\|_1 < \varepsilon$.

Lemma 2.19. Let $f : \mathbb{N} \rightarrow \mathbb{C}$ and $g : \mathbb{R} \rightarrow \mathbb{C}$ be two functions such that $f = g$ on \mathbb{N} and g is a Riemann integrable bounded periodic function with period $\alpha \in \mathbb{R} \setminus \mathbb{Q}$. Then, for all $\varepsilon > 0$, $\exists M \in \mathbb{N}$ and $c_k \in \mathbb{C}$

$$P(x) = \sum_{k=0}^M c_k e(kx/\alpha),$$

such that $\|f - P\|_1 \leq \varepsilon$. In particular, f is almost periodic.

Remark 2.20. If $\alpha \in \mathbb{Q}$, the result still holds. In fact, in that case f is periodic.

Proof. Consider the function $h : \mathbb{R} \rightarrow \mathbb{C}$ defined as $h(x) = g(x\alpha)$. Note that h is a 1-periodic function and piece-wise continuous. As h is bounded, it can be seen as a function of $L^2([0, 1])$. Thus, there exists $c_k := \hat{h}(k) = \frac{1}{2\pi} \int_{-\pi}^{\pi} h(x) e^{ikx} dx$,

where $\hat{h}(k)$ is the k th Fourier coefficient of h , such that $\|h - \sum_{k=0}^M c_k e(kx)\|_{L^1[0,1]} \rightarrow_M 0$.

Take M big enough such that $\|h - \sum_{k=0}^M c_k e(kx)\|_1 \leq \varepsilon$, and define $P(x) = \sum_{k=0}^M c_k e(kx)$. Since $f(n) = h(n/\alpha)$, we have that

$$f(n) - \sum_{k=0}^M c_k e(kn/\alpha) = h(n/\alpha) - P(n/\alpha),$$

thus

$$\frac{1}{N} \sum_{n=1}^N |f(n) - \sum_{k=0}^M c_k e(kn/\alpha)| = \frac{1}{N} \sum_{n=1}^N |h - P|(n\alpha^{-1}). \quad (2.12)$$

As $(n\alpha^{-1})_n$ is equidistributed in \mathbb{T} (as α^{-1} is irrational), and $|h - P|$ is a piece-wise continuous function on the torus, by the Weyl equidistribution criteria we have that

$$\frac{1}{N} \sum_{n=1}^N |h - P|(n\alpha^{-1}) \xrightarrow{N \rightarrow \infty} \int_{\mathbb{T}} |h - P| d\mu = \|h - P\|_1 \leq \varepsilon.$$

Therefore, applying limsup in (2.12), we conclude that $\|f - P\|_1 \leq \varepsilon$. □

Lemma 2.21. *If $\alpha \in \mathbb{R}$, then $\mathbb{1}_{[\alpha\mathbb{Z}]} : \mathbb{N} \rightarrow \mathbb{C}$ is an almost periodic function, which can be approximated with linear combination of exponential functions $e(\alpha n)$ with $\alpha \in \mathbb{R} \setminus \mathbb{Q}$.*

Proof. First, we note that for $m \in \mathbb{Z}$,

$$\begin{aligned} m \in [\alpha\mathbb{Z}] &\iff \exists n \in \mathbb{Z} \text{ such that } m \leq \alpha n < m + 1 \\ &\iff \exists n \in \mathbb{Z} \text{ such that } \alpha n - 1 < m \leq \alpha n. \end{aligned} \quad (2.13)$$

Consider the set

$$R_\alpha := \bigcup_{n \in \mathbb{Z}} (\alpha n - 1, \alpha n].$$

We note that $\mathbb{1}_{[\alpha\mathbb{Z}]} = \mathbb{1}_{R_\alpha}$ on \mathbb{N} . Treating $\mathbb{1}_{R_\alpha}$ as a function over \mathbb{R} , we have that it is a Riemann integrable bounded periodic function with period $\alpha \in \mathbb{R}$. Therefore, Lemma 2.19 yields the conclusion. □

Lemma 2.22. *For $p \in \mathbb{P}$, fix $0 \leq i < p$, $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and let*

$$D_{p,i} := \{n \in [\alpha\mathbb{N}] : n = \lfloor m\alpha \rfloor \text{ and } \lfloor p\alpha m \rfloor = p\lfloor \alpha m \rfloor + i\}.$$

There is a Riemann integrable α -periodic function f over the reals such that $f = \mathbb{1}_{D_{p,i}}$ when restricted to \mathbb{N} .

Proof. First, for any $m \in \mathbb{N}$,

$$\begin{aligned} \lfloor p\alpha m \rfloor = p\lfloor \alpha m \rfloor + i &\iff \{\alpha m\} \in \left[\frac{i}{p}, \frac{i+1}{p}\right) \\ &\iff \alpha m - \frac{i+1}{p} < \lfloor \alpha m \rfloor \leq \alpha m - \frac{i}{p}. \end{aligned} \quad (2.14)$$

Therefore, we define the set

$$D'_{p,i} := \bigcup_{m \in \mathbb{Z}} \left(\alpha m - \frac{i+1}{p}, \alpha m - \frac{i}{p} \right],$$

and note that $\mathbb{1}_{D_{p,i}} = \mathbb{1}_{D'_{p,i}}$ on \mathbb{N} . Furthermore, $f := \mathbb{1}_{D'_{p,i}}$ is a Riemann integrable function, which is also α -periodic. □

2.4. BMAI sequences

The goal of this subsection is to introduce the notion of BMAI sequences and list some properties. We let \mathbb{P} denote the set of prime numbers.

Definition 2.23. A bounded sequence $a : \mathbb{N} \rightarrow \mathbb{C}$ is called *BMAI* if there is a bounded sequence $(\theta_p)_{p \in \mathbb{P}} \subseteq \mathbb{C} \setminus \{0\}$ such that

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |a(pn) - \theta_p a(n)| = 0. \quad (2.15)$$

The idea behind this definition is that the limiting behaviour of a BMAI sequence approximates that of a multiplicative function.

Remark 2.24. If $a : \mathbb{N} \rightarrow \mathbb{C}$ is a BMAI sequence we also have by Corollary 2.9 using $S = \mathbb{P}$ and for $T \in S$, $f(n, T) = a(Tn) - \theta_T a(n)$, that

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N}^{\log} |a(pn) - \theta_p a(n)| = 0.$$

Example 2.25. The following are BMAI functions.

- The Liouville function $\lambda : \mathbb{N} \rightarrow \mathbb{C}$ with $\theta_p = -1$, $\forall p \in \mathbb{P}$.
- The function $\psi_F(n) := F\left(\frac{\Omega(n) - \log \log n}{\sqrt{\log \log n}}\right)$, for a bounded continuous $F : \mathbb{R} \rightarrow \mathbb{C}$ with $\theta_p = 1$, $\forall p \in \mathbb{P}$.
- Any bounded multiplicative arithmetic function f , with $\theta_p = f(p)$ nonzero, $\forall p \in \mathbb{P}$.

We prove some basic properties.

Proposition 2.26. Let $a, b : \mathbb{N} \rightarrow \mathbb{C}$ BMAI sequences. Then for primes $p, q \in \mathbb{P}$ we have

$$\limsup_N \mathbb{E}_{n \leq N} (|a(pn)b(qn) - \theta_p^a \theta_q^b a(n)b(n)|) = 0,$$

and also

$$\limsup_N \mathbb{E}_{n \leq N}^{\log} (|a(pn)b(qn) - \theta_p^a \theta_q^b a(n)b(n)|) = 0.$$

Proof. We have the following inequality

$$\begin{aligned} |a(pn)b(qn) - \theta_p^a \theta_q^b a(n)b(n)| &\leq |a(pn)b(qn) - \theta_p^a a(n)b(qn)| + |\theta_p^a a(n)b(qn) - \theta_p^a \theta_q^b a(n)b(n)| \\ &\leq \|b\|_{\ell^\infty} |a(pn) - \theta_p^a a(n)| + \|a\|_{\ell^\infty} \|(\theta_p^a)_p\|_{\ell^\infty} |b(pn) - \theta_q^b b(n)| \end{aligned}$$

which gives

$$\limsup_N |a(pn)b(pn) - \theta_p^a \theta_p^b a(n)b(n)| = 0.$$

□

Corollary 2.27. BMAI sequences are closed under coordinate-wise multiplication, i.e. if $(a(n))_n$ and $(b(n))_n$ are BMAI then so is $(a(n)b(n))_n$.

Proof. Taking $p = q$, Proposition 2.26 gives

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |(a \cdot b)(pn) - \theta_p^a \theta_p^b(a \cdot b)(n)| = 0.$$

□

Corollary 2.28. For a BMAI sequence $a : \mathbb{N} \rightarrow \mathbb{C}$,

$$\limsup_N \mathbb{E}_{n \leq N} |a(pn) \overline{a(qn)} - \theta_p \overline{\theta_q} |a(n)|^2| = 0,$$

and

$$\limsup_N \mathbb{E}_{n \leq N}^{\log} |a(pn) \overline{a(qn)} - \theta_p \overline{\theta_q} |a(n)|^2| = 0,$$

for all $p, q \in \mathbb{P}$.

Proof. Take $b = \bar{a}$ in Proposition 2.26. □

Proposition 2.29. Let $a : \mathbb{N} \rightarrow \mathbb{C}$ be a BMAI sequence and $f : \mathbb{N} \rightarrow \mathbb{C}$ a bounded function. Then for any $d_1, d_2 \in \mathbb{N}$

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |a(p(d_1n + d_2))f(n) - \theta_p a(d_1n + d_2)f(n)| = 0,$$

and

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N}^{\log} |a(p(d_1n + d_2))f(n) - \theta_p a(d_1n + d_2)f(n)| = 0.$$

Proof. Without loss of generality, we can take $f \equiv 1$ since

$$\mathbb{E}_{n \leq N} |a(p(d_1n + d_2))f(n) - \theta_p a(d_1n + d_2)f(n)| \leq \|f\|_{\ell^\infty} \mathbb{E}_{n \leq N} |a(p(d_1n + d_2)) - \theta_p a(d_1n + d_2)|.$$

Note that

$$\begin{aligned} \mathbb{E}_{n \leq N} |a(p(d_1n + d_2)) - \theta_p a(d_1n + d_2)| &= \frac{1}{N} \sum_{m=1}^{d_1N+d_2} |a(pm) - \theta_p a(m)| \mathbb{1}_{d_1\mathbb{N}+d_2}(m) \\ &= \left(d_1 + \frac{d_2}{N}\right) \mathbb{E}_{m \leq d_1N+d_2} |a(pm) - \theta_p a(m)| \mathbb{1}_{(d_1\mathbb{N}+d_2)}(m) \\ &\leq \left(d_1 + \frac{d_2}{N}\right) \mathbb{E}_{m \leq d_1N+d_2} |a(pm) - \theta_p a(m)|. \end{aligned}$$

Taking \limsup , we derive

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |a(p(d_1n + d_2))f(n) - \theta_p a(d_1n + d_2)f(n)| = 0.$$

□

Proposition 2.30. *Let $a : \mathbb{N} \rightarrow \mathbb{C}$ be a BMAI sequence, then there exists a (actual) multiplicative function $\theta : \mathbb{N} \rightarrow \mathbb{C}$ such that for every $m \in \mathbb{N}$*

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |a(mn) - \theta(m)a(n)| = 0.$$

Additionally, $\|a\|_1 \neq 0$ if and only if θ is unique.

Proof. Let $(\theta_p)_{p \in \mathbb{P}}$ be a sequence given by a being BMAI. We define $\theta : \mathbb{N} \rightarrow \mathbb{C}$, as the multiplicative function defined by $\theta(p) = \theta_p, \forall p \in \mathbb{P}$.

We prove the statement using induction over $\Omega(m)$. The definition of a BMAI sequence yields the case $\Omega(m) = 1$. We assume that the statement holds for all $m \in \mathbb{N}$ with $\Omega(m) \leq k$. Let $m \in \mathbb{N}$ with $\Omega(m) = k$ and $p \in \mathbb{P}$. We use Proposition 2.29 with $d_1 = m, d_2 = 0$ and $f \equiv 1$ to obtain

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |a(pmn) - \theta_p a(mn)| = 0. \quad (2.16)$$

Using (2.16) and the inductive hypothesis:

$$\begin{aligned} & \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |a(pmn) - \theta(pn)a(n)| \\ & \leq \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |a(pmn) - \theta_p a(mn)| + |\theta_p| \mathbb{E}_{n \leq N} |a(mn) - \theta(m)a(n)| \\ & = 0, \end{aligned}$$

and as every $n \in \mathbb{N}$ with $\Omega(n) = k + 1$ can be written as $n = pm$ for $p \in \mathbb{P}$ with $p \mid n$ and $m = n/p$, thus concluding existence.

For uniqueness, take $p \in \mathbb{P}$ and let θ'_p such that

$$\limsup_{N \rightarrow \infty} |a(pn) - \theta'_p a(n)| = 0.$$

Note that

$$\limsup_{N \rightarrow \infty} \mathbb{E} |\theta_p a(n) - \theta'_p a(n)| \leq \limsup_{N \rightarrow \infty} |a(pn) - \theta'_p a(n)| + |a(pn) - \theta_p a(n)| = 0.$$

Therefore

$$|\theta_p - \theta'_p| \cdot \|a\|_1 = 0, \forall p \in \mathbb{P},$$

so if $\|a\|_1 \neq 0$ then $\theta_p = \theta'_p, \forall p \in \mathbb{P}$. For the other direction, if $\|a\|_1 = 0$, then $\theta_p \in \mathbb{C}$ can be any value. In fact, notice that

$$\begin{aligned} \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |a(pn) - \theta_p a(n)| & \leq \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |a(pn)| + |\theta_p| \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |a(n)| \\ & \leq p \|a\|_1 + |\theta_p| \cdot \|a\|_1 = 0, \end{aligned}$$

which doesn't depend on θ_p . □

Lemma 2.31. Let $a : \mathbb{N} \rightarrow \mathbb{C}$ be a BMAI sequence such that for all natural numbers $h \in \mathbb{N}$ and irrational β ,

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} a(n) \overline{a(n+h)} e(\beta n) = 0 \quad (\text{resp. } \mathbb{E}_{n \leq N}^{\log} \bullet).$$

Then, $\forall p, q \in \mathbb{P}$, $\beta \in \mathbb{R} \setminus \mathbb{Q}$ and $h_1, h_2 \in \mathbb{N}$ with $qh_1 - ph_2 \neq 0$,

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} a(pn + h_1) \overline{a(qn + h_2)} e(\beta n) = 0.$$

Proof. We write out the proof for Cesàro averages only, as the logarithmic case is analogous. Without loss of generality, assume $ph_2 - qh_1 > 0$. Then

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} a(n + qh_1) \overline{a(n + ph_2)} e(\beta n) e(\beta qh_1) = \lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} a(n) \overline{a(n + ph_2 - qh_1)} e(\beta n) = 0.$$

Thus

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} a(n + qh_1) \overline{a(n + ph_2)} e(\beta n) = 0, \quad (2.17)$$

for every $\beta \in \mathbb{R} \setminus \mathbb{Q}$ and $p, q \in \mathbb{P}$ and $h_1, h_2 \in \mathbb{N}$ with $qh_1 - ph_2 \neq 0$. On the other hand, using Proposition 2.29 twice, we get

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \left| \overline{\theta}_p \theta_q a(pn + h_1) \overline{a(qn + h_2)} e(\beta n) - a(pqn + qh_1) \overline{a(pqn + ph_2)} e(\beta n) \right| = 0.$$

Hence

$$\begin{aligned} & \limsup_{N \rightarrow \infty} \left| \mathbb{E}_{n \leq N} \overline{\theta}_p \theta_q a(pn + h_1) \overline{a(qn + h_2)} e(\beta n) \right| \\ &= \limsup_{N \rightarrow \infty} \left| \mathbb{E}_{n \leq N} a(pqn + qh_1) \overline{a(pqn + ph_2)} e(\beta n) \right| \\ &= \limsup_{N \rightarrow \infty} \left| \frac{1}{N} \sum_{n=1}^{Npq} a(n + qh_1) \overline{a(n + ph_2)} e\left(\frac{\beta}{pq}n\right) \mathbb{1}_{pq\mathbb{N}}(n) \right|. \end{aligned}$$

Since $\mathbb{1}_{pq\mathbb{N}}(n) = \frac{1}{pq} \sum_{k=1}^{pq} e(nk/pq)$, we have

$$\begin{aligned} & \limsup_{N \rightarrow \infty} \left| \mathbb{E}_{n \leq N} a(pn + h_1) \overline{a(qn + h_2)} e(\beta n) \right| \\ &= \left| \frac{1}{\overline{\theta}_p \theta_q} \right| \cdot \left(\sum_{k=1}^{pq} \lim_{N \rightarrow \infty} \frac{1}{N} \left| \sum_{n=1}^{Npq} a(n + qh_1) \overline{a(n + ph_2)} e\left(\frac{\beta + k}{pq}n\right) \right| \right) \\ &= 0, \end{aligned}$$

in which we used equation (2.17) in the last equality to conclude. \square

3. Main Result

3.1. Basic Case

The following result is well known, as discussed in [JLW20], for multiplicative sequences. We generalize it to all BMAI sequences.

Lemma 3.1. *Let $a : \mathbb{N} \rightarrow \mathbb{C}$ be a BMAI sequence, then for any irrational $\beta \in (0, 1)$ we have*

$$\lim_{M \rightarrow \infty} \mathbb{E}_{m \leq M} a(m) e(m\beta) = 0.$$

Proof. By Lemma 2.15 we have a universal constant $C > 0$ such that for $N, S \in \mathbb{N}$

$$|\mathbb{E}_{n \leq N} a(n) e(n\beta)| \leq C \sum_{l=1}^{\frac{\log S}{\log 2}} \frac{1}{l \log \log S} \left(\max_{p, q \in [2^l, 2^{l+1}]} \left| \mathbb{E}_{n \leq \frac{N}{\max\{p, q\}}} a(pn) e(pn\beta) \overline{a(qn) e(qn\beta)} \right| \right)^{1/2} + \varepsilon(S, N), \quad (3.1)$$

with $\limsup_{S \rightarrow \infty} \limsup_{N \rightarrow \infty} \varepsilon(S, N) = 0$. Notice that

$$\begin{aligned} & \lim_{M \rightarrow \infty} \left| \mathbb{E}_{m \leq M} a(pm) \overline{a(qm)} e(\beta(p-q)m) \right| \\ & \leq \lim_{M \rightarrow \infty} \mathbb{E}_{m \leq M} |(a(pm) \overline{a(qm)} - \theta_p \overline{\theta_q}) e(\beta(p-q)m)| + \lim_{M \rightarrow \infty} |\theta_p \overline{\theta_q}| \cdot |\mathbb{E}_{m \leq M} e(\beta(p-q)m)|. \end{aligned}$$

By lemma 2.28, we have that the first term of the sum is $o_{p \rightarrow \infty}(1) + o_{q \rightarrow \infty}(1)$. And as β is irrational, we have that $\beta(p-q)$ is not an integer and so $\lim_{M \rightarrow \infty} |\mathbb{E}_{m \leq M} e(\beta(p-q)m)| = 0$. Therefore, we have

$$\limsup_N |\mathbb{E}_{n \leq N} a(n) e(n\beta)| \leq C \sum_{l=1}^{\frac{\log S}{\log 2}} \frac{1}{l \log \log S} \left(\max_{p, q \in [2^l, 2^{l+1}]} o_{p \rightarrow \infty}(1) + o_{q \rightarrow \infty}(1) \right)^{1/2} + \limsup_N \varepsilon(S, N). \quad (3.2)$$

Take L big enough such that $(\max_{p, q \in [2^l, 2^{l+1}]} o_{p \rightarrow \infty}(1) + o_{q \rightarrow \infty}(1))^{1/2} \leq \varepsilon$, $\forall l \geq L$. Then, taking \limsup_S in Eq. (3.2)

$$\begin{aligned} \limsup_N |\mathbb{E}_{n \leq N} a(n) e(n\beta)| & \leq C \limsup_S \sum_{l=L}^{\frac{\log S}{\log 2}} \frac{1}{l \log \log S} \left(\max_{p, q \in [2^l, 2^{l+1}]} o_{p \rightarrow \infty}(1) + o_{q \rightarrow \infty}(1) \right)^{1/2} \\ & \leq C\varepsilon \limsup_S \frac{1}{\log \log S} \sum_{l=L}^{\frac{\log S}{\log 2}} \frac{1}{l} = C\varepsilon. \end{aligned}$$

Taking $\varepsilon \rightarrow 0$, we conclude that

$$\lim_N \mathbb{E}_{n \leq N} a(n) e(n\beta) = 0.$$

□

Proposition 3.2. Suppose $a : \mathbb{N} \rightarrow \mathbb{C}$ is a BMAI sequence such that $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N a(n) = c$ with $c \in \mathbb{C}$. Then for any irrational $\alpha > 1$, we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N a(\lfloor \alpha n \rfloor) = c.$$

Proof. First notice that

$$\frac{1}{N} \sum_{n=1}^N a(\lfloor \alpha n \rfloor) = \alpha \frac{1}{\alpha N} \sum_{m=1}^{\alpha N} a(m) \mathbb{1}_{[\alpha \mathbb{N}]}(m).$$

And so to prove that $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N a(\lfloor \alpha n \rfloor) = c$, it suffices to show that

$$\lim_{M \rightarrow \infty} \frac{1}{M} \sum_{m=1}^M a(m) \mathbb{1}_{[\alpha \mathbb{N}]}(m) = \alpha^{-1} c. \quad (3.3)$$

By Lemma 2.21, $\mathbb{1}_{[\alpha \mathbb{N}]}(m)$ is an almost periodic function such that for $\varepsilon > 0$, there is $(c_i)_{i=0, \dots, k} \in \mathbb{C}^k$ and irrationals $(\beta_i)_{i=1, \dots, k} \in (0, 1)^k$ such that

$$\left\| \mathbb{1}_{[\alpha \mathbb{N}]}(m) - \left(c_0 + \sum_{i=1}^k c_i e(m\beta_i) \right) \right\|_1 < \varepsilon.$$

In addition, as $d([\alpha N]) = \alpha^{-1}$ and for each $\beta \in (0, 1)$, $\mathbb{E}_{n \leq N} e(m\beta_i) \rightarrow 0$, we have that $c_0 = \alpha^{-1}$.

Hence, we have

$$\begin{aligned} & \limsup_M |\mathbb{E}_{m \leq M} a(m) \mathbb{1}_{[\alpha \mathbb{N}]}(m) - \alpha^{-1} c| \\ & \leq \|a\|_\infty \left\| \mathbb{1}_{[\alpha \mathbb{N}]}(m) - \left(c_0 + \sum_{i=1}^k c_i e(m\beta_i) \right) \right\|_1 \\ & \quad + \limsup_M \left| \mathbb{E}_{n \leq M} a(m) \left(c_0 + \sum_{i=1}^k c_i e(m\beta_i) \right) - \alpha^{-1} c \right| \\ & \leq \|a\|_\infty \varepsilon + \underbrace{\limsup_M |\mathbb{E}_{n \leq M} a(m) \alpha^{-1} - c \alpha^{-1}|}_{=0} + \sum_{i=1}^k |c_i| \limsup_M |\mathbb{E}_{n \leq M} a(m) e(m\beta_i)|. \end{aligned}$$

By the assumptions on a , $\lim_{M \rightarrow \infty} \frac{1}{M} \sum_{m=1}^M a(m) \alpha^{-1} = \alpha^{-1} c$. By Lemma 3.1, the last term of the sum tends to 0, giving

$$\limsup_M |\mathbb{E}_{n \leq M} a(m) \mathbb{1}_{[\alpha \mathbb{N}]}(m) - \alpha^{-1} c| \leq \|a\|_\infty \varepsilon.$$

We conclude by taking $\varepsilon \searrow 0$. □

Remark 3.3. Note that the condition of the above proposition can be weakened. Indeed, we can remove the conditions of a being BMAI and only ask for a bounded $a : \mathbb{N} \rightarrow \mathbb{C}$ which is orthogonal to $e(n\beta)$ for β irrational, i.e. $\lim_{M \rightarrow \infty} \frac{1}{M} \sum_{m=1}^M a(m)e(m\beta) = 0$.

Remark 3.4. By von Mangoldt [vM97, p. 852] and Landau [Lan09, pp. 571–572, 620–621], the prime number theorem is equivalent to

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \lambda(n) = 0.$$

Corollary 3.5. For any irrational $\alpha > 1$, we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \lambda(\lfloor \alpha n \rfloor) = 0.$$

Proof. Using that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \lambda(n) = 0,$$

and the fact that λ is BMAI, by Proposition 3.2, we obtain the desired result. \square

3.2. Main Theorem

We now state the main technical result, whose proof will come later. The main theorem follows from this result using Lemma 2.31.

Theorem 3.6. Let $a : \mathbb{N} \rightarrow \mathbb{C}$ be a bounded sequence and $b : \mathbb{N} \rightarrow \mathbb{S}^1$ be a BMAI sequence such that:

1. $\lim_{H \rightarrow \infty} \lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |\mathbb{E}_{h \leq H} a(n+h)| = 0$,
2. for all $\beta \in \mathbb{R} \setminus \mathbb{Q}$, $p \neq q$ primes and $(i, j) \in \{0, \dots, p-1\} \times \{0, \dots, q-1\} \setminus (0, 0)$,

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} a(pn+i) \overline{a(qn+j)} e(n\beta) = 0 \quad (\text{resp. } \mathbb{E}_{n \leq N}^{\log} \bullet).$$

Then for any irrational $\alpha \in \mathbb{R}_+$,

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} b(n) a(\lfloor \alpha n \rfloor) = 0 \quad (\text{resp. } \mathbb{E}_{n \leq N}^{\log} \bullet).$$

We now use Theorem 3.6 to prove Theorem C.

Proof of Theorem C using Theorem 3.6. We just need to prove condition 2 from Theorem 3.6. Let $\beta \in \mathbb{R} \setminus \mathbb{Q}$, $p \neq q$ primes and $(i, j) \in \{0, \dots, p-1\} \times \{0, \dots, q-1\} \setminus (0, 0)$. If $iq = pj$, then $p|i$, which implies that $i = 0$ and, therefore, $j = 0$. Nevertheless, this contradicts that $(i, j) \neq (0, 0)$ and, in consequence, $iq - pj \neq 0$. Finally, using Lemma 2.31 we immediately obtain condition 2. \square

The following example illustrates the necessity of the first condition in both theorems.

Example 3.7. Consider $a \equiv b \equiv 1$. Clearly both functions are BMAI with values in \mathbb{S}^1 , and satisfy condition 2 of Theorem C by

$$\left| \frac{1}{N} \sum_{n=1}^N e(n\beta) \right| \leq \frac{1}{N} \cdot \frac{2}{|1 - e(\beta)|} \rightarrow 0.$$

However, a clearly doesn't satisfy condition 1 as the associated limit is 1.

In contrast, the BMAI condition can be relaxed. Corollary 3.8 below is an example of a function which is not BMAI but satisfies the conditions of Theorem C.

Corollary 3.8. *For an irrational α and a real number c ,*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \lambda(n) e(c \lfloor \alpha n \rfloor) = 0. \quad (3.4)$$

Furthermore, for any almost periodic function $f : \mathbb{N} \rightarrow \mathbb{C}$,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \lambda(n) f(\lfloor \alpha n \rfloor) = 0. \quad (3.5)$$

Remark 3.9. Similar results have been obtained in other contexts such as [GT12].

Proof. If $c \in \mathbb{Z}$, then (3.4) reduces to Conjecture 1.5 (Chowla's conjecture) for the case $k = 1$, which is well known to be equivalent to the prime number theorem.

Suppose $c \in \mathbb{R} \setminus \mathbb{Z}$, then

$$\frac{1}{N} \sum_{n=1}^N \left| \frac{1}{H} \sum_{h=1}^H e(cn + ch) \right| = \frac{1}{N} \sum_{n=1}^N \left| \frac{1}{H} \sum_{h=1}^H e(ch) \right| = \left| \frac{1}{H} \sum_{h=1}^H e(ch) \right|$$

as $c \notin \mathbb{Z}$, $\lim_{H \rightarrow \infty} \left| \frac{1}{H} \sum_{h=1}^H e(ch) \right| = 0$ and $e(cn)$ satisfies condition 1 and 2 of Theorem C. Since λ is a BMAI sequence, the result follows from Theorem C.

As for (3.5), note that one can approximate almost periodic functions by a finite sum of $e(cn)$ functions. \square

The following lemma will be useful in proving Theorem 3.6.

Lemma 3.10. *For any bounded $a : \mathbb{N} \rightarrow \mathbb{C}$ that satisfies condition 1 of Theorem C,*

$$\lim_{H \rightarrow \infty} \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |\mathbb{E}_{h \leq H} a(Hn + h)| = 0 \quad (\text{resp. } \mathbb{E}_{n \leq N}^{\log} |\mathbb{E}_{h \leq H} \bullet|).$$

Proof. Consider the function

$$\gamma_H(n) = \left| \frac{1}{H} \sum_{h=1}^H a(n + h) \right|,$$

and let $0 < \varepsilon < 1$. By assumption, we can find an $\tilde{H} \in \mathbb{N}$ such that for every $H \geq \tilde{H}$,

$$\limsup_N \mathbb{E}_{n \leq N} \gamma_H(n) \leq \varepsilon^3.$$

In the following, we will omit the subscript H in γ_H and just write γ . All statements will hold for any $H \geq \tilde{H}$. Let $E = \{n \in \mathbb{N} \mid \gamma(n) \leq \varepsilon\}$, and note that

$$\mathbb{E}_{n \leq N} \gamma(n) \geq \mathbb{E}_{n \leq N} \gamma(n) \mathbb{1}_{E^c}(n) \geq \mathbb{E}_{n \leq N} \varepsilon \mathbb{1}_{E^c}(n). \quad (3.6)$$

Taking \limsup in Eq. (3.6),

$$\varepsilon^3 \geq \limsup_N \mathbb{E}_{n \leq N} \gamma(n) \geq \varepsilon d(E^c) \implies \varepsilon^2 \geq d(E^c).$$

Let

$$D = \{n \in \mathbb{N} \mid \frac{1}{H} |[Hn, H(n+1)) \cap E| \geq (1 - \varepsilon)\} = \{n \in \mathbb{N} \mid \frac{1}{H} |[Hn, H(n+1)) \cap E^c| \leq \varepsilon\}$$

be the set of numbers n for which the H points after starting points Hn are covered by E up to ε . Note that

$$\begin{aligned} \mathbb{E}_{n \leq (N+1)H-1} \mathbb{1}_{E^c} &= \frac{NH}{(N+1)H-1} \cdot \frac{1}{NH} \sum_{n \leq N} |[Hn, H(n+1)) \cap E^c| \\ &\geq \frac{NH}{(N+1)H-1} \cdot \varepsilon \frac{1}{N} \sum_{n \leq N} \mathbb{1}_{D^c}(n). \end{aligned}$$

Taking \limsup , we conclude that

$$\varepsilon^2 \geq \varepsilon d(D^c) \implies \varepsilon \geq d(D^c).$$

In other words, since E has high density, “good” starting points are also “dense”. If $n \in D$ then Hn is at most at εH distance of an element of $m \in E$. Using that $\gamma(m) \leq \varepsilon \implies \gamma(m-k) \leq \varepsilon + \frac{2k\|a\|_\infty}{H}$ for $0 \leq k \leq \varepsilon H$ we conclude that $\gamma(Hn) \leq \varepsilon + 2\varepsilon\|a\|_\infty$ for every $n \in D$. Finally, we note that

$$\begin{aligned} \mathbb{E}_{n \leq N} \gamma(nH) &= \mathbb{E}_{n \leq N} \gamma(nH) \mathbb{1}_D(n) + \mathbb{E}_{n \leq N} \gamma(nH) \mathbb{1}_{D^c}(n) \leq \varepsilon + 2\varepsilon\|a\|_\infty + \|a\|_\infty \mathbb{E}_{n \leq N} \mathbb{1}_{D^c}(n) \\ &\implies \limsup_N \mathbb{E}_{n \leq N} \gamma(nH) \leq \varepsilon + 2\varepsilon\|a\|_\infty + \|a\|_\infty d(D^c) \leq \varepsilon(1 + 3\|a\|_\infty). \end{aligned}$$

As this is valid for every $H \geq \tilde{H}$, we conclude that

$$\lim_{H \rightarrow \infty} \limsup_N \mathbb{E}_{n \leq N} \gamma(nH) = 0. \quad (3.7)$$

For the logarithmic version, we use Eq. (3.7) combined with Corollary 2.9, taking $S = \mathbb{N}$ and $f(n, T) = \gamma_T(n)$.

□

Proof of Theorem 3.6. For simplicity, we restrict our attention to Cesaró averages, as the proof for logarithmic averages is analogous and all the necessary lemmas used in the proof hold for both types of averages. We will apply Lemma 2.15 to $f(n) = b(n)a(\lfloor \alpha n \rfloor)$ to get

$$\begin{aligned} & |\mathbb{E}_{n \leq N} b(n)a(\lfloor \alpha n \rfloor)| \\ & \leq C \frac{1}{\log \log S} \sum_{l=1}^{\frac{\log S}{\log 2}} \frac{1}{l} \left(\max_{p,q \in [2^l, 2^{l+1})} |\mathbb{E}_{n \leq \frac{N}{\max\{p,q\}}} b(pn) \overline{b(qn)} a(\lfloor p\alpha n \rfloor) \overline{a(\lfloor q\alpha n \rfloor)}| \right)^{\frac{1}{2}} + \varepsilon(S, N), \end{aligned}$$

with $\limsup_{S \rightarrow \infty} \limsup_{N \rightarrow \infty} \varepsilon(S, N) = 0$. Similarly as in the proof of Proposition 3.2, we take limits and use Corollary 2.28 to get

$$\begin{aligned} & \limsup_{N \rightarrow \infty} |\mathbb{E}_{n \leq N} b(n)a(\lfloor \alpha n \rfloor)| \leq \\ & \limsup_{S \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{C \|b\|_{\ell^\infty}}{\log \log S} \sum_{l=1}^{\frac{\log S}{\log 2}} \frac{1}{l} \left(\max_{p,q \in [2^l, 2^{l+1})} |\theta_p \overline{\theta_q}| |\mathbb{E}_{n \leq \frac{N}{\max\{p,q\}}} |b(n)|^2 a(\lfloor p\alpha n \rfloor) \overline{a(\lfloor q\alpha n \rfloor)}| \right)^{\frac{1}{2}}. \end{aligned} \quad (3.8)$$

Moreover, as $|\theta_p \overline{\theta_q}| \leq C_\theta$ for all $p \neq q$, we get that

$$\begin{aligned} & \limsup_{N \rightarrow \infty} |\mathbb{E}_{n \leq N} b(n)a(\lfloor \alpha n \rfloor)| \leq \\ & \limsup_{S \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{C \|b\|_{\ell^\infty} C_\theta}{\log \log S} \sum_{l=1}^{\frac{\log S}{\log 2}} \frac{1}{l} \left(\max_{p,q \in [2^l, 2^{l+1})} |\mathbb{E}_{n \leq \frac{N}{\max\{p,q\}}} a(\lfloor p\alpha n \rfloor) \overline{a(\lfloor q\alpha n \rfloor)}| \right)^{\frac{1}{2}}. \end{aligned} \quad (3.9)$$

It is thus sufficient to show

$$\limsup_{S \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{1}{\log \log S} \sum_{l=1}^{\frac{\log S}{\log 2}} \frac{1}{l} \left(\max_{p,q \in [2^l, 2^{l+1})} |\mathbb{E}_{n \leq \frac{N}{\max\{p,q\}}} a(\lfloor p\alpha n \rfloor) \overline{a(\lfloor q\alpha n \rfloor)}| \right)^{\frac{1}{2}} = 0. \quad (3.10)$$

To begin, note that

$$\frac{1}{\log \log S} \sum_{l=1}^{\log \frac{1}{\alpha}} \frac{1}{l} \left(\max_{p,q \in [2^l, 2^{l+1})} |\mathbb{E}_{n \leq \frac{N}{\max\{p,q\}}} a(\lfloor p\alpha n \rfloor) \overline{a(\lfloor q\alpha n \rfloor)}| \right)^{\frac{1}{2}} \quad (3.11)$$

$$\leq \frac{1}{\log \log S} \sum_{l=1}^{\log \frac{1}{\alpha}} \frac{1}{l} \left(\max_{p,q \in [2^l, 2^{l+1})} |\mathbb{E}_{n \leq \frac{N}{\max\{p,q\}}} \|a\|_\infty| \right)^{\frac{1}{2}} \quad (3.12)$$

$$\leq \frac{\|a\|_\infty^2}{\log \log S} \sum_{l=1}^{\log \frac{1}{\alpha}} \frac{1}{l} \quad (3.13)$$

$$\leq O_\alpha \left(\frac{1}{\log \log S} \right), \quad (3.14)$$

which tends to 0 as S goes to infinity. Thus, it remains to investigate the case where $p, q > \frac{1}{\alpha}$. Let $i(p, n) := \lfloor p\alpha n \rfloor - p\lfloor \alpha n \rfloor$, thus

$$a(\lfloor p\alpha n \rfloor) \overline{a(\lfloor q\alpha n \rfloor)} = a(p\lfloor \alpha n \rfloor + i(p, n)) \overline{a(q\lfloor \alpha n \rfloor + i(q, n))}.$$

Notice that $i(p, n)$ only takes values in $\{0, \dots, p-1\}$. Let

$$D_{p,i} := \{n \in \lfloor \alpha \mathbb{N} \rfloor : n = \lfloor m\alpha \rfloor \text{ and } \lfloor p\alpha m \rfloor = p\lfloor \alpha m \rfloor + i\}.$$

$D_{p,0}, \dots, D_{p,p-1}$ form a partition of $\lfloor \alpha \mathbb{N} \rfloor$. We have

$$\mathbb{E}_{n \leq \frac{N}{\max\{p,q\}}} a(\lfloor p\alpha n \rfloor) \overline{a(\lfloor q\alpha n \rfloor)} = \alpha \mathbb{E}_{n \leq \frac{\alpha N}{\max\{p,q\}}} \sum_{i(p)} \sum_{j(q)} a(pn + i) \overline{a(qn + j)} \mathbb{1}_{D_{p,i}}(n) \mathbb{1}_{D_{q,i}}(n) \quad (3.15)$$

where $i(p)$ denotes $0 \leq i < p$ and similarly for $j(q)$. We consider two cases separately, that of $(i, j) = (0, 0)$ and $(i, j) \neq (0, 0)$.

Firstly, for the case $(i, j) = (0, 0)$, since α is irrational, the density of the set $D_{p,0} \cap D_{q,0}$ is $\frac{1}{pq}$. Hence

$$\limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq \frac{N}{\max\{p,q\}}} |a(pn + i) \overline{a(qn + j)} \mathbb{1}_{D_{p,0} \cap D_{q,0}}(n)| \leq \frac{\|a\|_\infty^2}{pq}.$$

Therefore, the contribution of $(i, j) = (0, 0)$ in Eq. (3.10) is

$$\leq \frac{\alpha \|a\|_\infty}{\log \log S} \sum_{l=\log \frac{1}{\alpha}}^{\frac{\log S}{\log 2}} \frac{1}{l} \max_{p,q \in [2^l, 2^{l+1})} \frac{1}{\sqrt{pq}} \leq \frac{\alpha \|a\|_\infty}{\log \log S} \sum_{l=1}^{\frac{\log S}{\log 2}} \frac{1}{l 2^l} \rightarrow_{S \rightarrow \infty} 0.$$

We now consider the case $(i, j) \neq (0, 0)$. We want to estimate

$$\limsup_N \left| \sum_{\substack{i(p), j(q) \\ (i,j) \neq (0,0)}} \alpha \mathbb{E}_{m \leq \frac{N\alpha}{2^l}} a(pm + i) \overline{a(qm + j)} \mathbb{1}_{D_{p,i} \cap D_{q,i}}(m) \right|. \quad (3.16)$$

For each $(i, j) \neq (0, 0)$, $\mathbb{1}_{D_{p,i} \cap D_{q,i}}(m)$ is an almost periodic function that can be extended to a piece-wise continuous function on all of \mathbb{R} with period α , using Lemma 2.22. In particular, for $\varepsilon > 0$, by Lemma 2.19 there exist $c_1, \dots, c_K \neq 0$ and $\beta_1, \dots, \beta_K \in \mathbb{R} \setminus \mathbb{Q}$ such that

$$\|\mathbb{1}_{D_{p,i} \cap D_{q,i}}(m) - \left(\frac{1}{pq} + \sum_{k=1}^K c_k e(m\beta_k) \right)\|_1 < \frac{\varepsilon}{\|a\|_\infty^2}.$$

Using the triangle inequality,

$$\text{Eq. (3.16)} \leq \limsup_N \alpha \mathbb{E}_{m \leq \frac{N\alpha}{2^l}} \left| \sum'_{i(p), j(q)} \frac{1}{pq} a(pm + i) \overline{a(qm + j)} \right|$$

$$+ \limsup_N \alpha \sum_{\substack{i(p), j(q) \\ (i,j) \neq (0,0)}} \sum_{k=1}^K c_k |\mathbb{E}_{m \leq \frac{N\alpha}{2^l}} a(pm+i) \overline{a(qm+j)} e(m\beta_k)| + \varepsilon.$$

The second \limsup_N is 0 from our assumptions on a . We are thus left with the first term of the sum. Rewriting,

$$|\mathbb{E}_{m \leq \frac{N\alpha}{2^l}} \sum_{\substack{i(p), j(q) \\ (i,j) \neq (0,0)}} \frac{1}{pq} a(pm+i) \overline{a(qm+j)}| \leq |\mathbb{E}_{m \leq \frac{N\alpha}{2^l}} \frac{1}{p} \sum_{i=0}^{p-1} a(pm+i)| \cdot |\frac{1}{q} \sum_{j=0}^{q-1} \overline{a(qm+j)}|. \quad (3.17)$$

Using Lemma 3.10 we have that

$$\lim_{H \rightarrow \infty} \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} |\mathbb{E}_{h \leq H} a(Hn+h)| = 0,$$

and, taking l large enough (and, therefore, $H = p$ large enough) we derive that

$$\begin{aligned} \limsup_N \mathbb{E}_{m \leq \frac{N\alpha}{2^l}} \left| \frac{1}{p} \sum_{i=0}^{p-1} a(pm+i) \right| &\leq \frac{\varepsilon}{\|a\|_\infty} \\ \implies \limsup_N \mathbb{E}_{m \leq \frac{N\alpha}{2^l}} \left| \frac{1}{p} \sum_{i=0}^{p-1} a(pm+i) \right| \cdot \left| \frac{1}{q} \sum_{j=0}^{q-1} \overline{a(qm+j)} \right| &\leq \varepsilon. \end{aligned}$$

Thus for l large enough

$$\limsup_N \left| \sum_{\substack{i(p), j(q) \\ (i,j) \neq (0,0)}} \alpha \mathbb{E}_{m \leq \frac{N\alpha}{2^l}} a(pm+i) \overline{a(qm+j)} \mathbb{1}_{D_{p,i} \cap D_{q,i}}(m) \right| \leq 3\varepsilon.$$

Finally, together with Eq. (3.11), we have that

$$\limsup_{S \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{1}{\log \log S} \sum_{l=1}^{\frac{\log S}{\log 2}} \frac{1}{l} \left(\max_{p,q \in [2^l, 2^{l+1})} |\mathbb{E}_{n \leq \frac{N}{2^l}} a(\lfloor p\alpha n \rfloor) \overline{a(\lfloor q\alpha n \rfloor)}| \right)^{\frac{1}{2}} \leq 3\varepsilon,$$

whence taking $\varepsilon \rightarrow 0$ gives the desired result. \square

4. Consequences of the main theorem

4.1. Erdős-Kac generalization

For the following sections, we denote for every $n \in \mathbb{N}$,

$$B_n := \frac{\omega(n) - \log \log n}{(\log \log n)^{1/2}}, \quad A_n := \frac{\omega(\lfloor \alpha n \rfloor) - \log \log n}{(\log \log n)^{1/2}},$$

and for a function $F : \mathbb{R} \rightarrow \mathbb{C}$,

1. $\Psi_F(n) := F(B_n)$,
2. $I(F) := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} F(x) e^{-\frac{x^2}{2}} dx$.

First, we prove that $a = \Psi_G$ satisfies the conditions of theorem Theorem 3.6 whenever $I(G) = 0$.

Lemma 4.1. *Let $G : \mathbb{R} \rightarrow \mathbb{C}$ be a bounded continuous function such that $I(G) = 0$. Then Ψ_G satisfies the conditions of Theorem C.*

Proof. First, we show that Ψ_G satisfies the first condition. Proving the following will be sufficient:

$$\lim_{H \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \left| \frac{1}{H} \sum_{h=1}^H \Psi_G(n+h) \right|^2 = 0.$$

Indeed, by expanding the squares and switching the order of summation, we get

$$\lim_{H \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \left| \frac{1}{H} \sum_{h=1}^H \Psi_G(n+h) \right|^2 = \lim_{H \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{1}{H^2} \sum_{h=1}^H \sum_{h'=1}^H \frac{1}{N} \sum_{n=1}^N \Psi_G(n+h) \overline{\Psi_G(n+h')}.$$

Splitting the sum into the diagonal terms and the non diagonal terms,

$$\lim_{H \rightarrow \infty} \frac{1}{H^2} \sum_{h \neq h'} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \Psi_G(n+h) \overline{\Psi_G(n+h')} + \lim_{H \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{1}{H^2} \sum_{h=1}^H \frac{1}{N} \sum_{n=1}^N |\Psi_G(n+h)|^2.$$

The first sum tends to 0. Indeed let $\varepsilon > 0$ and $(E_i)_{i=1}^I$ be Jordan measurable sets, where $E_i \subseteq \mathbb{R}$ for each $i \in I$, and $(c_i)_{i=1}^I \subseteq \mathbb{C}^I$ such that

$$\left\| \sum_{i=1}^I c_i \mathbb{1}_{E_i} - G \right\|_{L^1(\mathbb{R})} \leq \sqrt{2\pi\varepsilon}, \quad \left\| \sum_{i=1}^I c_i \mathbb{1}_{E_i} \right\|_{L^\infty(\mathbb{R})} \leq \|G\|_{L^\infty(\mathbb{R})}.$$

We may assume $\|\mathbb{1}_{[-M,M]}(\sum_{i=1}^I c_i \mathbb{1}_{E_i} - G)\|_{L^\infty(\mathbb{R})} \leq \varepsilon$, where $M > 0$ is such that

$$\left| \int_{[-M,M]^c} e^{-x^2/2} dx \right| \leq \frac{1}{2\|G\|_{L^\infty(\mathbb{R})}} \varepsilon.$$

Notice that

$$\begin{aligned} & \limsup_{n \leq N} \mathbb{E}_{n \leq N} |\Psi_G(n+h) - \sum_{i=1}^I c_i \mathbb{1}_{E_i}(B_{n+h})| \\ & \leq \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \left| \left(\mathbb{1}_{[-M,M]} \left(G - \sum_{i=1}^I c_i \mathbb{1}_{E_i} \right) \right) (B_{n+h}) \right| \\ & + \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \left| \left(\mathbb{1}_{[-M,M]^c} \left(G - \sum_{i=1}^I c_i \mathbb{1}_{E_i} \right) \right) (B_{n+h}) \right| \end{aligned}$$

$$\begin{aligned}
&\leq \|\mathbb{1}_{[-M,M]}(\sum_{i=1}^I c_i \mathbb{1}_{E_i} - G)\|_{L^\infty(\mathbb{R})} + 2\|G\|_{L^\infty(\mathbb{R})} \limsup_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \mathbb{1}_{[-M,M]^c}(B_{n+h}) \\
&\leq \varepsilon + 2\|G\|_{L^\infty(\mathbb{R})} \left| \int_{[-M,M]^c} e^{-x^2/2} dx \right| \leq 2\varepsilon,
\end{aligned}$$

where we used Theorem 2.16. Hence

$$\limsup_{N \rightarrow \infty} |\mathbb{E}_{n \leq N} (\Psi_G(n+h) \overline{\Psi_G(n+h')}) - \sum_{i,j \in [I]^2} c_i \overline{c_j} \mathbb{1}_{E_i}(B_{n+h}) \mathbb{1}_{E_j}(B_{n+h'})| \leq 4\varepsilon \|G\|_\infty$$

Using Theorem 2.16 again, we have

$$\begin{aligned}
\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \sum_{i,j \in [I]^2} c_i \overline{c_j} \mathbb{1}_{E_i}(B_{n+h}) \mathbb{1}_{E_j}(B_{n+h'}) &= \frac{1}{2\pi} \sum_{i,j \in [I]^2} c_i \overline{c_j} \int_{E_i \times E_j} e^{-\frac{1}{2}(x_1^2 + x_2^2)} dx_1 dx_2 \\
&= \frac{1}{2\pi} \sum_{i,j \in [I]^2} c_i \overline{c_j} \left(\int_{E_i} e^{-x^2/2} dx \right) \left(\int_{E_j} e^{-x^2/2} dx \right) \\
&= \frac{1}{2\pi} \left| \int_{-\infty}^{\infty} \sum_{i=1}^I c_i \mathbb{1}_{E_i} e^{-x^2/2} dx \right|^2.
\end{aligned}$$

Using this, we derive

$$\begin{aligned}
\limsup_{n \leq N} |\mathbb{E}_{n \leq N} \Psi_G(n+h) \overline{\Psi_G(n+h')}| &\leq 4\varepsilon \|G\|_\infty + \lim_{n \rightarrow N} |\mathbb{E}_{n \leq N} \sum_{i,j \in [I]^2} c_i \overline{c_j} \mathbb{1}_{E_i}(B_{n+h}) \mathbb{1}_{E_j}(B_{n+h'})| \\
&= 4\varepsilon \|G\|_\infty + \frac{1}{2\pi} \left| \int_{-\infty}^{\infty} \sum_{i=1}^I c_i \mathbb{1}_{E_i}(x) e^{-x^2/2} dx \right|^2 \\
&\leq 4\varepsilon \|G\|_\infty + \frac{1}{2\pi} \left| \int_{-\infty}^{\infty} \left(\sum_{i=1}^I c_i \mathbb{1}_{E_i}(x) - G(x) \right) e^{-x^2/2} dx \right|^2 \\
&\quad (\text{since } I(G) = 0) \\
&\leq 4\varepsilon \|G\|_\infty + \frac{1}{2\pi} \left\| \sum_{i=1}^I c_i \mathbb{1}_{E_i} - G \right\|_1^2 \\
&\leq 4\varepsilon \|G\|_\infty + \varepsilon.
\end{aligned}$$

Thus as $\varepsilon > 0$ was arbitrary, we get that $\lim_{N \rightarrow \infty} |\mathbb{E}_{n \leq N} \Psi_G(n+h) \overline{\Psi_G(n+h')}| = 0$.

For the second sum, notice that $\frac{1}{N} \sum_{n=1}^N |\Psi_G(n+h)|^2 \leq C$ for some absolute constant as G is bounded, and thus

$$\lim_{H \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{1}{H^2} \sum_{h=1}^H \frac{1}{N} \sum_{n=1}^N |\Psi_G(n+h)|^2 \leq \lim_{H \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{1}{H^2} \sum_{h=1}^H C = \lim_{H \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{C}{H} = 0.$$

Hence the first condition is satisfied.

To verify the second condition, in light of Lemma 2.10, it is enough to show that for all $h \in \mathbb{N}$

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \Psi_G(p(n+h) + i) \overline{\Psi_G(q(n+h) + j) e((n+h)\beta)} \overline{\Psi_G(pn + i) \Psi_G(qn + j) e(n\beta)} = 0,$$

which is equivalent to showing

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \Psi_G(pn + ph + i) \overline{\Psi_G(qn + qh + j)} \overline{\Psi_G(pn + i) \Psi_G(qn + j)} = 0.$$

As $p \neq q$, $(i, j) \in \{0, \dots, p-1\} \times \{0, \dots, q-1\} \setminus (0, 0)$ and $h > 0$, we have that $pn + ph + i$, $qn + qh + j$, $pn + i$ and $qn + j$ are polynomials over \mathbb{Z} that are pairwise relatively prime. We can thus apply Theorem 2.16 to get the desired result. \square

Lemma 4.2. *Let $\alpha \in \mathbb{R}$ be a positive irrational real number. Then for every continuous $F : \mathbb{R} \rightarrow \mathbb{S}^1$ and bounded $G : \mathbb{R} \rightarrow \mathbb{C}$ with $I(G) = 0$,*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \Psi_F(n) \Psi_G(\lfloor \alpha n \rfloor) = 0. \quad (4.1)$$

Proof. Let $a(n) = \Psi_G(n)$ and $b(n) = \Psi_F(n)$. By Lemma 4.1, a satisfies the conditions of Theorem C and b is bounded. Hence

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \Psi_F(n) \Psi_G(\lfloor \alpha n \rfloor) = \lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} b(n) a(\lfloor \alpha n \rfloor) = 0.$$

\square

Lemma 4.3. *Suppose that for every continuous function with compact support $F : \mathbb{R} \rightarrow \mathbb{S}^1$, and bounded function $G : \mathbb{R} \rightarrow \mathbb{C}$ with $I(G) = 0$,*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} F(A_n) G(B_n) = 0.$$

Then, for all $a, b, c, d \in \mathbb{R}$,

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \mathbb{1}_{[a,b]}(A_n) \mathbb{1}_{[c,d]}(B_n) = I(\mathbb{1}_{[a,b]}) I(\mathbb{1}_{[c,d]}).$$

Proof. First, notice that the family of functions

$$\mathcal{A} = \text{span} \left(F \in \mathcal{C}(\mathbb{R}, \mathbb{S}^1) \mid F : \mathbb{R} \rightarrow \mathbb{S}^1 \text{ has compact support} \right),$$

is such that

- \mathcal{A} separates points, in other words, for any $x \neq y \in X$, there exists $f \in \mathcal{A}$ such that $f(x) \neq f(y)$,
- for every $x \in X$, there exists $f \in \mathcal{A}$ such that $f(x) = 0$,
- \mathcal{A} is self-adjoint i.e. if $f \in \mathcal{A}$ then $\overline{f} \in \mathcal{A}$.

Then by the Stone-Weierstrass theorem, \mathcal{A} is dense in $C_\infty(\mathbb{R}, \mathbb{C})$. If $H \in C_\infty(\mathbb{R}, \mathbb{C})$ and $\varepsilon > 0$, let $F_1, \dots, F_r \in \mathcal{A}$ and $\{c_i\}_{i=1}^r \subseteq \mathbb{C}$ such that

$$\|H - \sum_{i=1}^r c_i F_i\|_\infty \leq \varepsilon.$$

Then, denoting $F_\varepsilon = \sum_{i=1}^r c_i F_i$ and $M = \max_{i=1, \dots, r} |c_i|$,

$$\begin{aligned} |\mathbb{E}_{n \leq N} H(A_n) G(B_n)| &\leq |\mathbb{E}_{n \leq N} \sum_{i=1}^r c_i F_i(A_n) G(B_n)| + \mathbb{E}_{n \leq N} |F_\varepsilon(A_n) G(B_n) - H(A_n) G(B_n)| \\ &\leq M \sum_{i=1}^r |\mathbb{E}_{n \leq N} F_i(A_n) G(B_n)| + \|G\|_\infty \|F_\varepsilon - H\|_\infty \\ &\leq M \sum_{i=1}^r |\mathbb{E}_{n \leq N} F_i(A_n) G(B_n)| + \|G\|_\infty \varepsilon. \end{aligned}$$

Thus, using the hypothesis,

$$\limsup_{N \rightarrow \infty} |\mathbb{E}_{n \leq N} H(A_n) G(B_n)| \leq \|G\|_\infty \varepsilon,$$

and as this is valid for all $\varepsilon > 0$, we conclude that

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} H(A_n) G(B_n) = 0, \quad \forall H \in C_\infty(\mathbb{R}, \mathbb{C}).$$

Now, given $a, b, c, d \in \mathbb{R}$ such that $a < b, c < d$ (otherwise it is trivial) and $\varepsilon > 0$ such that $\varepsilon < (b - a)/2$, we choose $H \in C_\infty(\mathbb{R})$ such that

$$H(x) = \begin{cases} 1 & x \in [a + \varepsilon, b - \varepsilon], \\ 0 & x \in (-\infty, a - \varepsilon] \cup [b + \varepsilon, \infty), \end{cases}$$

and such that H is monotone in $(a - \varepsilon, a + \varepsilon)$ and in $(b - \varepsilon, b + \varepsilon)$. Therefore, we have that

$$\begin{aligned} |\mathbb{E}_{n \leq N} (\mathbb{1}_{[a, b]}(A_n) G(B_n))| &\leq |\mathbb{E}_{n \leq N} (\mathbb{1}_{[a, b]}(A_n) G(B_n)) - H(A_n) G(B_n)| + |\mathbb{E}_{n \leq N} (H(A_n) G(B_n))| \\ &\leq \|G\|_\infty \mathbb{E}_{n \leq N} |\mathbb{1}_{[a, b]}(A_n) - H(A_n)| + |\mathbb{E}_{n \leq N} (H(A_n) G(B_n))| \\ &\leq \|G\|_\infty \mathbb{E}_{n \leq N} \mathbb{1}_{(a - \varepsilon, a + \varepsilon) \cup (b - \varepsilon, b + \varepsilon)}(A_n) + |\mathbb{E}_{n \leq N} (H(A_n) G(B_n))|. \end{aligned}$$

Taking \limsup and using Theorem 1.4 yields

$$\limsup_{N \rightarrow \infty} |\mathbb{E}_{n \leq N} (\mathbb{1}_{[a, b]}(A_n) G(B_n))| \leq \|G\|_\infty I(\mathbb{1}_{(a - \varepsilon, a + \varepsilon) \cup (b - \varepsilon, b + \varepsilon)}) \leq C \|G\|_\infty \varepsilon,$$

with $C = 4/\sqrt{2\pi}$. Therefore, as this is valid for every $\varepsilon > 0$, we conclude that

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} (\mathbb{1}_{[a, b]}(A_n) G(B_n)) = 0.$$

Taking $G(x) = \mathbb{1}_{[c, d]}(x) - I(\mathbb{1}_{[c, d]})$, we have that

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \mathbb{1}_{[a, b]}(A_n) (\mathbb{1}_{[c, d]}(B_n) - I(\mathbb{1}_{[c, d]})) = 0,$$

and thanks to Erdős-Kac (Theorem 1.3) and Theorem 1.4, this is equivalent to

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \mathbb{1}_{[a,b]}(A_n) \mathbb{1}_{[c,d]}(B_n) = I(\mathbb{1}_{[a,b]})I(\mathbb{1}_{[c,d]}),$$

concluding the proof. \square

Lemma 4.4. *There exists a set $E \subseteq \mathbb{N}$ of density 1 such that*

$$|A_n - B_{[\alpha n]}| \xrightarrow[n \in E]{n \rightarrow \infty} 0.$$

Proof. Notice that $\log \log [\alpha n] = \log \log n + b_n$, where $(b_n)_n$ is a bounded sequence (in which we used that $\log : (1, \infty) \rightarrow \mathbb{R}$ is 1-Lipschitz given that its derivative is bounded by 1), and let $C_n := \frac{w([\alpha n]) - \log \log n}{(\log \log [\alpha n])^{1/2}}$. Note that

$$|B_{[\alpha n]} - C_n| \leq \frac{|b_n|}{\log \log [\alpha n]} \xrightarrow[n \rightarrow \infty]{} 0.$$

Therefore, it is enough to see that $|A_n - C_n| \rightarrow_n 0$ with n taking values in a set of density 1. Using similar arguments, we have that $(\log \log [\alpha n])^{1/2} = (\log \log n)^{1/2} + b'_n$, where $(b'_n)_n$ is a bounded sequence. Let $M = \max(\|(b_n)_n\|_{\ell^\infty}, \|(b'_n)_n\|_{\ell^\infty})$. Hence

$$\begin{aligned} |A_n - C_n| &= |b'_n \frac{w([\alpha n]) - \log \log n}{\log \log n + b_n(\log \log n)^{1/2}}| \\ &\leq M \left| \frac{1}{1 + b_n/(\log \log n)} \right| \cdot \left| \frac{w([\alpha n]) - \log \log n}{\log \log n} \right|. \end{aligned}$$

The last term tends to 0 with n taking values in a set E of density 1 given by the Hardy–Ramanujan theorem (Theorem 1.2). Meanwhile, the first term tends to $M \cdot 1$ as $n \rightarrow \infty$, therefore we conclude that

$$|A_n - C_n| \xrightarrow[n \in E]{n \rightarrow \infty} 0.$$

By the triangle inequality, we get the desired result

$$|A_n - B_{[\alpha n]}| \xrightarrow[n \in E]{n \rightarrow \infty} 0.$$

\square

We now restate and prove Theorem A.

Theorem A. *Let $\alpha \in \mathbb{R}$ be a positive irrational real number and $a, b, c, d \in \mathbb{R}$. Then*

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \#\{n \leq N : a \leq \frac{\omega([\alpha n]) - \log \log n}{(\log \log n)^{1/2}} \leq b, c \leq \frac{\omega(n) - \log \log n}{(\log \log n)^{1/2}} \leq d\} \\ = \left(\frac{1}{\sqrt{2\pi}} \int_a^b e^{-t^2/2} dt \right) \left(\frac{1}{\sqrt{2\pi}} \int_c^d e^{-t^2/2} dt \right). \end{aligned}$$

Proof. We want to show that

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \mathbb{1}_{[a,b]}(A_n) \mathbb{1}_{[c,d]}(B_n) = I(\mathbb{1}_{[a,b]})I(\mathbb{1}_{[c,d]}).$$

Using Lemma 4.3, it is enough to prove that for every continuous and compactly supported $F : \mathbb{R} \rightarrow \mathbb{S}^1$ and bounded $G : \mathbb{R} \rightarrow \mathbb{C}$ with $I(G) = 0$,

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} F(A_n)G(B_n) = 0. \quad (4.2)$$

In order to prove Eq. (4.2), we show that

$$\lim_{N \rightarrow \infty} |\mathbb{E}_{n \leq N} F(A_n)G(B_n) - F(B_{\lfloor \alpha n \rfloor})G(B_n)| = 0.$$

Indeed, since G is bounded,

$$|\mathbb{E}_{n \leq N} F(A_n)G(B_n) - F(B_{\lfloor \alpha n \rfloor})G(B_n)| \leq \|G\|_{\infty} \mathbb{E}_{n \leq N} |F(A_n) - F(B_{\lfloor \alpha n \rfloor})|, \quad (4.3)$$

and as F is uniformly continuous and has compact support, it is sufficient to show

$$|A_n - B_{\lfloor \alpha n \rfloor}| \xrightarrow{n \rightarrow \infty} 0,$$

with n taking values in a set of density 1, which is given by Lemma 4.4. In this way, it's enough to prove that for every continuous with compact support $F : \mathbb{R} \rightarrow \mathbb{S}^1$ and bounded $G : \mathbb{R} \rightarrow \mathbb{C}$ with $I(G) = 0$ we have that

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} \Psi_F(n) \Psi_G(\lfloor \alpha n \rfloor) = 0, \quad (4.4)$$

which follows from Lemma 4.2, concluding the proof. \square

4.2. Logarithmic averages of the Liouville function

We now cite a relatively recent theorem of Matomaki and Radziwiłł.

Theorem 4.5 (see [MR16]).

$$\lim_{H \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \left| \frac{1}{H} \sum_{h=1}^H \lambda(n+h) \right| = 0.$$

Note that this is the first condition of our main theorem, Theorem C, satisfied by the Liouville function. The following lemma from [FH18] shows that the Liouville function and the Möbius function also satisfy the second condition of Theorem C.

Lemma 4.6 (see [FH18, Corollary 1.5]). *Let $\beta \in \mathbb{R}$ be irrational, then*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N}^{\log} e(n\beta) \prod_{j=1}^j \lambda(n+h_j) = 0$$

for all $l \in \mathbb{N}$ and $h_1, \dots, h_l \in \mathbb{Z}$. Moreover, a similar statement holds with the Möbius function μ in place of λ .

We now restate and prove Theorem B.

Theorem B. *Let $\alpha \in \mathbb{R}$ an irrational positive real number, then*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N}^{\log} \lambda(n) \lambda(\lfloor \alpha n \rfloor) = 0.$$

Proof. Immediate by taking $b = \lambda$ and $a = \lambda$ in Theorem C and using Lemma 4.6. \square

5. Open Questions

While we proved the relative independence of n and $\lfloor \alpha n \rfloor$ under specific conditions, we have yet to show independence of different Beatty sequences. Namely, one may consider $\lfloor \alpha n \rfloor$ and $\lfloor \beta n \rfloor$ instead, which leads to the following conjecture.

Conjecture 5.1. *Let $a : \mathbb{N} \rightarrow \mathbb{C}$ and $b : \mathbb{N} \rightarrow \mathbb{S}^1$ be a BMAI sequences such that we also have that*

1. $\lim_{H \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \left| \frac{1}{H} \sum_{h=1}^H a(n+h) \right| = 0$
2. *if for all $\beta \in \mathbb{R} \setminus \mathbb{Q}$, $h \in \mathbb{N}$ we have that :*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} a(n) \overline{a(n+h)} e(n\beta) = 0 \quad (\text{resp } \mathbb{E}_{n \leq N}^{\log})$$

Then for any irrational $\alpha, \beta \in \mathbb{R}$ such that α and β are rationally independent,

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N} b(\lfloor \beta n \rfloor) \overline{a(\lfloor \alpha n \rfloor)} = 0. \quad (\text{resp } \mathbb{E}_{n \leq N}^{\log})$$

Once we have pairwise independence of Beatty sequences, we can then consider a higher-order version. For example, the following conjecture offers a higher-order generalization of Theorem B.

Conjecture 5.2. *For $\{\alpha_i\}_{i=2}^\ell \in \mathbb{R} \setminus \mathbb{Q}$ rationally independent, we have*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N}^{\log} \lambda(n) \lambda(\lfloor \alpha_2 n \rfloor) \dots \lambda(\lfloor \alpha_\ell n \rfloor) = 0.$$

A more recent work of Joni Teräväinen and Aled Walker ([[TW23](#)]) proves a generalization of Theorem B, namely for $\alpha_1, \alpha_2 > 0$ and $\beta_1, \beta_2 \in \mathbb{R}$ such that α_1/α_2 is irrational,

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \leq N}^{\log} \lambda(\lfloor \alpha_1 n + \beta_1 \rfloor) \lambda(\lfloor \alpha_2 n + \beta_2 \rfloor) = 0.$$

References

- [BS07] W. D. BANKS and I. E. SHPARLINSKI, Prime divisors in beatty sequences, *Journal of Number Theory* **123** (2007), 413–425. <https://doi.org/10.1016/j.jnt.2006.07.011>.
- [BM16] V. BERGELSON and J. MOREIRA, Van der Corput’s difference theorem: some modern developments, *Indag. Math. (N.S.)* **27** no. 2 (2016), 437–479. <https://doi.org/10.1016/j.indag.2015.10.014>.
- [BSZ13] J. BOURGAIN, P. SARNAK, and T. ZIEGLER, Disjointness of moebius from horocycle flows, in *From Fourier analysis and number theory to radon transforms and geometry*, *Dev. Math.* **28**, Springer, New York, 2013, pp. 67–83. https://doi.org/10.1007/978-1-4614-4075-8_5.

- [Cho65] S. CHOWLA, *The Riemann Hypothesis and Hilbert's Tenth Problem*, *Mathematics and Its Applications* **4**, Gordon and Breach Science Publishers, 1965. Available at <https://mathscinet.ams.org/mathscinet-getitem?mr=177943>.
- [DD82] H. DABOUSSI and H. DELANGE, On multiplicative arithmetical functions whose modulus does not exceed one, *J. London Math. Soc. (2)* **26** no. 2 (1982), 245–264. <https://doi.org/10.1112/jlms/s2-26.2.245>.
- [Ell79] P. D. T. A. ELLIOTT, *Probabilistic number theory. I, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Science]* **239**, Springer-Verlag, New York-Berlin, 1979, Mean-value theorems. <https://doi.org/10.1007/978-1-4612-9989-9>.
- [EK40] P. ERDÖS and M. KAC, The Gaussian law of errors in the theory of additive number theoretic functions, *Amer. J. Math.* **62** (1940), 738–742. <https://doi.org/10.2307/2371483>.
- [FH18] N. FRANTZIKINAKIS and B. HOST, The logarithmic Sarnak conjecture for ergodic weights, *Ann. of Math. (2)* **187** no. 3 (2018), 869–931. <https://doi.org/10.4007/annals.2018.187.3.6>.
- [GT12] B. GREEN and T. TAO, The Möbius function is strongly orthogonal to nilsequences, *Ann. of Math. (2)* **175** no. 2 (2012), 541–566. <https://doi.org/10.4007/annals.2012.175.2.3>.
- [HR17] G. H. HARDY and S. RAMANUJAN, The normal number of prime factors of a number n , *Quar. J. Pure. Appl. Math* **48** (1917), 76–97.
- [JLW20] Y. JIANG, G. LÜ, and Z. WANG, Exponential sums with multiplicative coefficients without the ramanujan conjecture, *Mathematische Annalen* **379** no. 1-2 (2020), 589–632. <https://doi.org/10.1007/s00208-020-02108-z>.
- [Kát86] I. KÁTAI, A remark on a theorem of H. Daboussi, *Acta Math. Hungar.* **47** no. 1-2 (1986), 223–225. <https://doi.org/10.1007/BF01949145>.
- [Lan09] E. LANDAU, *Handbuch der Lehre von der Verteilung der Primzahlen - Zweiter Band*, **2**, Leipzig Und Berlin: B. G. Teubner, 1909. Available at <http://name.umd.umich.edu/ABV2766.0001.001>.
- [vM97] H. VON MANGOLDT, Beweis der Gleichung $\sum_{k=1}^{\infty} \frac{\mu(k)}{k} = 0$, *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin, Zweiter Halbband – Juli bis Dezember* (1897), 835–852. Available at <https://biodiversitylibrary.org/page/29982413>.
- [MR16] K. MATOMÄKI and M. RADZIWIŁŁ, Multiplicative functions in short intervals, *Ann. of Math. (2)* **183** no. 3 (2016), 1015–1056. <https://doi.org/10.4007/annals.2016.183.3.6>.
- [Tan60] M. TANAKA, On the number of prime factors of integers. IV, *Jpn. J. Math.* **30** (1960), 55–83. https://doi.org/10.4099/jjm1924.30.0_55.
- [Tao16] T. TAO, The logarithmically averaged Chowla and Elliott conjectures for two-point correlations, *Forum Math. Pi* **4** (2016), e8, 36. <https://doi.org/10.1017/fmp.2016.6>.
- [TT18] T. TAO and J. TERÄVÄINEN, Odd order cases of the logarithmically averaged chowla conjecture, *J. Théor. Nombres Bordeaux* **30** no. 3 (2018), 997–1015. <https://doi.org/10.5802/jtnb.1062>.
- [TW23] J. TERÄVÄINEN and A. WALKER, On a bohr set analogue of chowla’s conjecture, 2023.

[Wat53] G. L. WATSON, On integers n relatively prime to $[an]$, *Canadian J. Math.* **5** (1953), 451–455.

(DAVID CRNČEVIĆ) ÉCOLE POLYTECHNIQUE, INSTITUT POLYTECHNIQUE DE PARIS, RTE DE SACLAY, PALAISEAU 91128, FRANCE

E-mail address: david.crncevic@polytechnique.edu

(FELIPE HERNÁNDEZ) DEPARTMENT OF MATHEMATICS, ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE, RTE CANTONALE, 1015 LAUSANNE, SWITZERLAND

E-mail address: felipe.hernandezcastro@epfl.ch

(KEVIN RIZK) DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, 450 JANE STANFORD WAY, STANFORD, CA 94305

E-mail address: kevin.b.rizk@gmail.com

(KHUNPOB SEREESUCHART) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, 520 PORTOLA PLAZA, LOS ANGELES, CA 90095

E-mail address: ksereesu2417@math.ucla.edu

(RAN TAO) DEPARTMENT OF MATHEMATICAL SCIENCES, CARNEGIE MELLON UNIVERSITY, 5000 FORBES AVE, PITTSBURGH, PA 15213

E-mail address: rant2@andrew.cmu.edu