

Dirac Kondo effect under magnetic catalysis

Koichi Hattori^{1,2,*}, Daiki Suenaga^{3,2,†}, Kei Suzuki^{4,‡} and Shigehiro Yasui^{5,6,§}

¹Zhejiang Institute of Modern Physics, Department of Physics, Zhejiang University, Hangzhou, 310027, China

²Research Center for Nuclear Physics (RCNP), Osaka University, Osaka 567-0047, Japan

³Strangeness Nuclear Physics Laboratory, RIKEN Nishina Center, Wako 351-0198, Japan

⁴Advanced Science Research Center, Japan Atomic Energy Agency (JAEA), Tokai 319-1195, Japan

⁵International Institute for Sustainability with Knotted Chiral Meta Matter (WPI-SKCM²),

Hiroshima University, Higashi-Hiroshima, Hiroshima 739-8526, Japan

⁶Research and Education Center for Natural Sciences, Keio University,

Hiyoshi 4-1-1, Yokohama, Kanagawa 223-8521, Japan

(Dated: December 5, 2023)

We develop a mean-field theory of a novel Kondo effect emerging in systems without a Fermi surface, which instead emerges under strong magnetic fields. We determine the magnitude of the Kondo condensate which is a particle pairing composed of conducting Dirac fermions and localized impurities. We focus on the competition between the Kondo effect and the energy gap formation that stems from the pairing among the Dirac fermions leading to the dynamical chiral symmetry breaking. We find that this competition induces a quantum critical point. We also investigate finite-temperature effects. This system at vanishing fermion density can be studied with Monte Carlo lattice simulations which do not suffer from the sign problem.

I. INTRODUCTION

Quantum systems under strong magnetic fields have attracted much attention over the last century in various physical systems (see, e.g., Refs. [1–6] for reviews). In the last decade, even more, common interests have been developed with the aid of advanced engineering of ultra-relativistic heavy-ion collisions [7–11] and of Dirac/Weyl semimetals [12–15] as well as the tremendous progress in the numerical lattice simulations [16, 17] (see also references therein). These situations motivate us to study phases of matter under strong magnetic fields. As we will discuss in detail, it has been pointed out that strong magnetic fields *catalyze* particle pairing phenomena, inducing nonperturbative modification of the ground state of the system when the Landau quantization becomes sizable.

In this paper, we investigate an interacting system composed of relativistic light fermions and nonrelativistic heavy impurities under a strong magnetic field at vanishing fermion density. Such relativistic fermions, governed by the Dirac equation, appear not only as elementary particles in high-energy physics but also in Dirac semimetals in condensed matter physics. Light and heavy fermions correspond to conducting and localized states, respectively. The conventional Kondo effect is absent at vanishing fermion density, whereas our system exhibits the Kondo effect and the chiral symmetry breaking induced by strong external magnetic fields. We call the Kondo effect appearing in Dirac fermion systems the *Dirac Kondo effect*. These phenomena are induced

by two pairing patterns: the Kondo condensate, pairing between the light fermion and the heavy impurity, and the chiral condensate, pairing between the light fermion and its antiparticle. In fact, each condensate is induced even with infinitesimal attractive interactions between the pair. That is, the growth of each condensate is an inevitable fate of the system when a magnetic field increases and temperature decreases. The common concept behind this statement is the *magnetic catalysis* as first pointed out in the formation of the chiral condensate [18–21] and refined in the formation of the Kondo condensate [22].¹

This system provides an interesting question arising from the competition between the chiral and Kondo condensates, which can drastically change the fate of the system. We will find that one condensate interrupts the growth of the other destructively, and there occurs a transition from the vacuum completely dominated by the chiral condensate to that accompanied by the Kondo condensate. At zero temperature, these two phases are divided by a quantum critical point at a certain critical magnetic-field strength where the Kondo condensate starts to grow. The chiral condensate no longer grows above the critical point and saturates at a constant value.

Our findings have an impact on the heart of many-body quantum physics. It is a general and central issue to determine the ground state of a system with condensates. The ground states can exhibit nonperturbative quantum phenomena such as superconductivity [23–25]

* koichi.hattori@zju.edu.cn

† daiki.suenaga@riken.jp

‡ k.suzuki.2010@th.phys.titech.ac.jp

§ yasuis@keio.jp

¹ While the magnetic catalysis is conventionally meant for the chiral condensate, the key concept, the *dimensional reduction* in the phase space stemming from the Landau degeneracy, is found to be more general and can lead to the realization of various pairing patterns in analogy with the Fermi surface effect [23, 24].

and the Kondo effect [26–33]. Relativistic counterparts are the spontaneous chiral symmetry breaking [34–37], the color superconductivity [38–42], and the newly proposed QCD Kondo effect [43, 44] (see also Refs. [45–63] for the developments in the QCD Kondo effect). Relativistic systems can be also realized with graphene and Dirac/Weyl semimetals that provide useful platforms in the study of the magnetic catalysis [64–81] and the Kondo effect [45, 82–97] (see Refs. [14, 98] for reviews).²

Despite the many foregoing studies, it should be stressed that the competition of the Kondo condensate with other condensates is yet elusive; The interplay appears both in destructive and constructive manners. There is a destructive competition between the Kondo effect and superconductivity [45, 99–110] (see also, e.g., Refs. [111–115] for experiments). Two of the present authors have also discussed the competition between the chiral symmetry breaking and the QCD Kondo effect at finite baryon density [50] (see also Refs. [57, 62]). On the other hand, constructive interplay can induce “heavy fermion” superconductivity [30, 115–118]. It is an interesting question how the Kondo condensate could affect the critical point from the interplay between the chiral condensate and the color superconductivity [42, 119, 120].

This paper is organized as follows. In Sec. II, we introduce a model Lagrangian and construct an effective potential within the mean-field approximation. Based on this, we discuss the phase structure with respect to a magnetic-field strength at zero temperature in Sec. III and finite-temperature effects at Sec. IV. In Sec. V, we conclude the paper and discuss possible applications and perspectives. In appendices, we provide more detailed accounts of the model and analytic computations.

II. FORMULATION

Nonperturbative phenomena, such as the formation of condensates, can be studied by the mean-field method (See Refs. [121, 122] for early works on the Kondo effect). In Refs. [47, 49], the QCD Kondo effect was investigated with the mean-field approach applied to a four-point interaction model analogous to the Nambu–Jona-Lasinio (NJL) model (for applications, see Refs. [50, 51, 53, 55–58, 61–63]).

Referring to those studies, we extract the essence of the chiral symmetry breaking and the Kondo effect in the Dirac fermion systems in strong magnetic fields. For this purpose, we use the following Lagrangian in the (3 + 1)-

dimensional spacetime (See Appendix A)

$$\begin{aligned} \mathcal{L} = & \bar{\psi}(i\cancel{\partial}_{\parallel} - m_l)\psi + \frac{G_{ll}}{2N} [(\bar{\psi}\psi)^2 + (\bar{\psi}i\gamma_5\psi)^2] \\ & + \sum_{c=\pm} \left[c\bar{\Psi}_v^c i\partial_0 \Psi_v^c \right. \\ & \left. + \frac{G_{hl}}{N} \{(\bar{\psi}\Psi_v^c)(\bar{\Psi}_v^c\psi) + (\bar{\psi}i\gamma_5\Psi_v^c)(\bar{\Psi}_v^c i\gamma_5\psi)\} \right], \end{aligned} \quad (1)$$

where $\gamma_5 \equiv i\gamma^0\gamma^1\gamma^2\gamma^3$. This Lagrangian contains four eigenmodes of ψ for the spin-polarized light fermion and its antifermion in the lowest Landau level (LLL), where m_l is the light-fermion mass. It has the index for the inner degrees of freedom such as the color for quarks ($N = 3$) and the pseudospin in condensed matter, e.g., graphene ($N = 2$). We assume that a magnetic field is applied in the third (z) direction. Then, ψ has the kinetic term $\cancel{\partial}_{\parallel} \equiv \gamma^0\partial_0 + \gamma^3\partial_3$ and is an eigenstate of the spin along the magnetic field due to the Zeeman effect, i.e., $i\gamma^1\gamma^2\psi = \text{sgn}(q_l B)\psi$ with an electric charge q_l . Note that planar systems such as graphene can be discussed with straightforward modifications of the present formulation.

The Lagrangian (1) also contains Ψ_v^+ and Ψ_v^- for the heavy-fermion and its antifermion fields introduced as impurities. They stem from the original Dirac field Ψ as $\Psi_v^{\pm} \equiv e^{-i\frac{\cancel{\partial}_{\perp}}{2m_h}} e^{\pm im_h v^{\mu} x_{\mu}} \mathcal{Q}_{\pm} \Psi$ with $\cancel{\partial}_{\perp} \equiv (\gamma^{\mu} - \not{v}v^{\mu})\partial_{\mu}$ after the factorization of the plane-wave components at the position x^{μ} and the projection by $\mathcal{Q}_{\pm} = \frac{1}{2}(1 \pm v^{\mu}\gamma_{\mu})$ as explained in Appendix A3. Here, the four-momentum is given as $m_h v^{\mu}$ with m_h and v^{μ} being the heavy-fermion mass and the four velocity, respectively. In the rest frame of heavy fermions, $v^{\mu} = (1, 0, 0, 0)$, $\mathcal{Q}_{\pm} = \frac{1}{2}(1 \pm \gamma_0)$ is the projection operator to the particle and antiparticle components.³ $G_{hl} > 0$ and $G_{ll} > 0$ are the four-point coupling constants for the interactions between the heavy and light fermions and between the light fermions, respectively.

We analyze the Lagrangian (1) with the mean-field approximation. Analogously to Ref. [58] in the (3+1)-dimensional spacetime, the chiral and Kondo condensates are assumed to be

$$\langle \bar{\psi}\psi \rangle_{\text{LLL}} \equiv -\frac{N}{G_{ll}} M, \quad \langle \bar{\psi}\Psi_v^{\pm} \rangle_{\text{LLL}} \equiv \frac{N}{G_{hl}} \Delta, \quad (2)$$

with the gaps M and Δ . By explicitly diagonalizing the mean-field Lagrangian, we obtain the energy-momentum

² References [94, 96] studied the Zeeman effect on the Kondo effect at *finite* density, while we discuss the magnetic catalysis for the Kondo effect at *vanishing* density.

³ This treatment is similar to but slightly different from the conventional framework of the heavy-quark effective theory (HQET) [123–126] in high-energy physics, which is an effective field theory for heavy quarks in QCD. In the conventional HQET, one retains either Ψ_v^+ or Ψ_v^- . On the other hand, in this work we retain both of Ψ_v^+ and Ψ_v^- to include the heavy fermion and antifermion in a charge-conjugation invariant manner [127, 128].

dispersion relations of the four eigenmodes

$$E_{\pm}(p_z) \equiv \pm \frac{1}{2} \left(\sqrt{E_{p_z}^2 + |2\Delta|^2} \pm E_{p_z} \right), \quad (3a)$$

$$\tilde{E}_{\pm}(p_z) \equiv \pm \frac{1}{2} \left(\sqrt{E_{p_z}^2 + |2\Delta|^2} \mp E_{p_z} \right), \quad (3b)$$

where $E_{p_z} \equiv \sqrt{p_z^2 + (m_l + M)^2}$. Straightforwardly, we obtain the thermodynamic potential $\Omega(M, \Delta) \equiv -T \ln Z$ with temperature $T = 1/\beta$ and the partition function Z at the one-loop level [129]:

$$\begin{aligned} \Omega(M, \Delta) &= N \left[\frac{M^2}{2G_{ll}} + \frac{|2\Delta|^2}{2G_{hl}} + \Omega_{\text{vac}} + \Omega_T \right], \\ \Omega_{\text{vac}}(M, \Delta) &= -\rho_B \sum_{\mathcal{E}_i} \int_{-\Lambda}^{\Lambda} \frac{dp_z}{2\pi} \frac{1}{2} |\mathcal{E}_i|, \\ \Omega_T(M, \Delta) &= -\rho_B \sum_{\mathcal{E}_i} \int_{-\infty}^{\infty} \frac{dp_z}{2\pi} \frac{1}{\beta} \ln(1 + e^{-\beta|\mathcal{E}_i|}), \end{aligned} \quad (4)$$

where $\Omega_{\text{vac}/T}$ are the contributions from the vacuum bubbles and the thermal excitation, respectively, and $\rho_B = |q_l B|/(2\pi)$ is the Landau degeneracy factor. The sum runs over the four eigenmodes $\mathcal{E}_i = \{E_+, E_-, \tilde{E}_+, \tilde{E}_-\}$. We are left with evaluating the one-dimensional momentum integral whose interval is limited within the ultraviolet cutoff Λ . All the dimensionful quantities can be scaled by Λ because of the dimensional reduction.

The thermodynamic potential (4) is inevitably unstable at the origin $M = 0 = \Delta$ due to the dimensional reduction at zero temperature, which implies that the system always favors nonzero condensates. This can be confirmed with an analytic expression of Ω_{vac} as follows. We take the massless limit ($m_l = 0$) for a demonstration. One can simply perform the p_z integral and expand the result as

$$\Omega_{\text{vac}}(\Phi) = -\frac{\rho_B}{2\pi} \left(\Lambda^2 + \Phi^2 \left(\frac{1}{2} + \ln \frac{2\Lambda}{\Phi} \right) + \mathcal{O}(\Phi^4) \right), \quad (5)$$

where $\Phi^2 = M^2 + |2\Delta|^2$. Stability at the origin is determined by the sign of the quadratic terms in Φ . The quadratic term enhanced by a logarithmic factor $\ln \Lambda/\Phi$, diverging as $\Phi \rightarrow 0$, makes the potential convex upward at the origin for *any coupling strengths* [18–21, 130]. At finite temperature, the origin can be a stable point since the thermal contribution Ω_T yields a term that can cancel the above-mentioned logarithmic term.

III. ZERO-TEMPERATURE RESULTS

We first investigate the ground state at zero temperature. We consider several cases classified by the relative magnitude between G_{hl} and G_{ll} in Eq. (1). When $G_{hl} = G_{ll}$, the thermodynamic potential (4) is invariant under a rotation in the M - Δ plane in the chiral limit, i.e., the vanishing light-fermion mass limit ($m_l = 0$), and

has a degenerate minimum on a circle at $\Phi^2 = \text{constant}$. A finite m_l breaks the rotational invariance. As a result, the potential is tilted toward the M axis, and has a minimum at a nonzero M and vanishing Δ . This is an analog of the chiral condensate in the effective models of QCD with the nonzero current quark mass (see, e.g., Ref. [37]). When $G_{hl} \neq G_{ll}$, there is no longer the degeneracy on the circle. When $G_{hl} < G_{ll}$, the tendency that M is favored over Δ is enhanced due to the suppression of the potential energy along the M axis for any m_l . However, when $G_{hl} > G_{ll}$, competition appears between the suppression of the potential energy and the tilting effect. This is a nontrivial case as we investigate in detail below.

First, we consider a special case $G_{hl} > G_{ll} = 0$ and elucidate the effects of the light-fermion mass m_l on the Kondo effect. In the chiral limit, we find that there is no phase transition characterized by a singular behavior of Δ , as shown by the dotted line in Fig. 1. On the other hand, when a nonzero m_l is switched on, the monotonic increase changes to a second-order phase transition, as shown by the blue dashed line, at a critical magnetic-field strength B_{cK} . Thus, the light-fermion mass plays an important role in the Kondo effect.

Next, we investigate the competition between the chiral and Kondo condensates with nonzero G_{ll} and G_{hl} . One can find analytic solutions of the stationary conditions $\partial\Omega/\partial M = 0 = \partial\Omega/\partial\Delta$ as follows (see Appendix B for details). One immediately notices a trivial solution $\Delta = 0$ accompanied by a nontrivial solution $M \neq 0$. This set is denoted as $(M_1, \Delta_1 = 0)$. One can find another set of solutions (M_2, Δ_2) as

$$M_2 = \frac{m_l G_{ll}}{G_{hl} - G_{ll}}, \quad (6a)$$

$$\Delta_2 = \frac{1}{2} \sqrt{\frac{\Lambda^2}{\sinh^2\left(\frac{2\pi^2}{\rho_B G_{hl}}\right)} - \frac{m_l^2 G_{hl}^2}{(G_{hl} - G_{ll})^2}}. \quad (6b)$$

Both the two sets, $(M_1, \Delta_1 = 0)$ and (M_2, Δ_2) , satisfy the stationary conditions for any values of the parameters as long as $G_{hl} > G_{ll}$.

We numerically investigated which, or any other set of solutions, serves as the global minimum of the thermodynamic potential $\Omega(M, \Delta)$. The numerical results are shown with the solid lines in Fig. 1. We find that, below a certain critical magnetic-field strength B_c , the first solutions, $(M_1, \Delta_1 = 0)$, are realized: The Kondo effect is prohibited by the existence of the chiral condensate. By solving $\Delta_2 = 0$ in Eq. (6b), we obtain the critical strength of magnetic field:

$$q_l B_c = \frac{4\pi^3}{G_{hl} \operatorname{arcsinh}\left(\Lambda \frac{|G_{hl} - G_{ll}|}{m_l G_{hl}}\right)} \sim \frac{4\pi^3 m_l}{\Lambda |G_{hl} - G_{ll}|}, \quad (7)$$

where the rightmost side holds with a small value of $|G_{hl} - G_{ll}|$. The true vacuum switches over from $(M_1, \Delta_1 = 0)$ to (M_2, Δ_2) at B_c , which is thus identified as a quantum critical point. The analytic solution (6) shows a nontrivial behavior of the condensates above B_c :




FIG. 1. Magnetic-field dependences of Kondo condensate Δ (blue) and chiral condensate M (red). The solid lines show the full results with fixed parameters, $m_l/\Lambda = 0.01$, $G_{ll}\Lambda^2 = 1.0$, and $G_{hl}\Lambda^2 = 3.0$. The dotted and dashed lines show Δ with $m_l = G_{ll} = 0$ and with $G_{ll} = 0$, respectively.




FIG. 2. Temperature dependences of Kondo condensate Δ and chiral condensate M at a fixed magnetic-field strength $|q_l B|/\Lambda^2 = 1.5$. The legends are the same as in Fig. 1. The red dashed line shows M at $G_{hl} = 0$.

The Kondo condensate grows up as the magnetic field increases, while the chiral condensate is forced to be a constant value (M_2). This is an anomalous saturation of the chiral condensate due to the formation of the Kondo condensate. As in Fig. 1, a “plateau” of the chiral condensate above B_c serves as a clear signal of the appearance of the Kondo condensate at zero temperature.

IV. FINITE-TEMPERATURE RESULTS

Finally, we investigate the phase diagram at finite temperature. In Fig. 2, we show the numerical results for the temperature dependence of the condensates, where the magnetic-field strength and the heavy-light coupling are fixed at $|q_l B|/\Lambda^2 = 1.5$ and $G_{hl}\Lambda^2 = 3.0$, respectively. The Kondo condensate at $G_{ll} = 0$ and $m_l = 0$ is shown

by the blue dotted line, where we find a phase transition at the critical temperature T_{cK0} . When we switch on a light-fermion mass $m_l/\Lambda = 0.01$, as shown by the blue dashed line, we observe that the magnitude of condensate is slightly suppressed, and the critical temperature decreases to T_{cK} . This suppression effect by m_l is consistent with our result at zero temperature.

Before discussing the competition, we check the spontaneous chiral symmetry breaking with nonzero G_{ll} at $G_{hl} = 0$. The chiral condensate M is shown by the red dashed line in Fig. 2. The chiral condensate hardly changes within the plot range of Fig. 2 but decreases at higher temperature. The chiral condensate at nonzero m_l shows a crossover transition, and its pseudocritical temperature is located in the higher-temperature region, not shown in the figure.

The numerical results including both nonzero G_{ll} and G_{hl} are shown by solid lines in Fig. 2. We find that the chiral and Kondo condensates coexist at low temperature below T_c . Here, the critical temperature T_c for the Kondo condensate decreases from T_{cK} due to the competition. As we increase temperature across T_c , the Kondo condensate melts away and the chiral condensate starts to *increase* abruptly at T_c since the chiral condensate is released from the competition with the Kondo condensate. Such a steep increase in the chiral condensate can be indirect evidence of the Kondo condensate.

In Fig. 3, we show the condensate values on the T - B plane. The plateau, which we have observed in the chiral condensate M at zero temperature, also extends to nonzero temperature. In addition, one can read off the orders of phase transitions: While both condensates indicate second-order phase transitions at zero temperature, switching on a nonzero temperature leads to the first-order phase transitions. In Fig. 4, we show a schematic phase diagram, where the first-order phase transition line at finite temperature is connected to the quantum critical point at zero temperature.

V. CONCLUSIONS

We have investigated the novel phase diagram of Dirac fermions, which is characterized by the chiral and Kondo condensates. We analytically found the critical magnetic field B_c where the new quantum critical point emerges from the competition between the two condensates. This competition also gives rise to a *saturation behavior* of the chiral condensate above B_c at low temperature. At higher temperature, the Kondo condensate melts away. This is the end of the competition, which is signaled by a steep *increase* of the chiral condensate.

For applications, our study can be applied to condensed matter systems as well as quark systems in high-energy physics. Dirac fermions are known to be realized in a variety of condensed matter systems including graphene. The magnetic catalysis in spatial two dimensions has been discussed in Refs. [18, 19, 67–69] and was




FIG. 3. Magnetic-field and temperature dependences of chiral condensate M (left) and Kondo condensate Δ (right).




FIG. 4. Schematic phase diagram extracted from Fig. 3.

further investigated with lattice simulations for graphene in Refs. [78–80]. In addition, a flat band emerges in twisted bilayer graphene with a magic angle due to mixing between the Dirac cones from different layers [131–135] and holds the $SU(2)$ valley degeneracy. Therefore, stacking this bilayer graphene and another monolayer graphene with a Dirac band will create a platform for the magnetic-field induced Kondo effect. Modeling the inter-layer interactions is a hot topic (see, e.g., Ref. [136]), which we leave as a future work.

One of the relevant questions is the excitation spectrum when the ground state varies with an increasing magnetic field. It is interesting to investigate an extension of the so-called Yu-Shiba-Rusinov state in superconductor [137–140] (see also Refs. [104–110] and Ref. [45] for an analog in QCD), the Nambu-Goldstone bosons, and non-Fermi liquid behavior near quantum critical points [141–143] pursued with the experimental progress (see,

e.g., Refs. [144–147]). It is important to investigate the critical exponents with possible excitations. One can also study an extension to the multi-channel Kondo effect [148], where a non-Fermi liquid system can appear depending on the number of channels (see, e.g., Refs. [45, 46, 60, 148, 149]).

We also draw attention to the heavy-quark dynamics in ultrarelativistic heavy-ion collisions [10, 150]. It is an interesting question how the heavy-quark diffusion can be modified by the strong correlation induced by the Kondo effect.

Numerical simulations on the lattice have provided useful approaches to nonperturbative many-body phenomena. The intertwined dynamics of the chiral symmetry breaking and the Kondo effect may be simulated in strong magnetic fields. Lattice QCD simulations elucidated the magnetic catalysis in QCD with high precision (see, e.g., Refs. [17, 151–156]), and they can be also applied to the counterpart in solid-state physics. The magnetically induced Kondo effect deserves further study beyond the mean-field model calculation and should be confirmed by direct measurements of the Kondo condensate constructed from the correlation function of the heavy and light fermions or by indirect ones through special behaviors of the chiral condensate pointed out in this paper. Recall also that the systems in a magnetic field and at finite density share analogous low-energy excitations stemming from the degeneracy in the Landau levels and on the Fermi surface. Therefore, the Monte Carlo simulation in magnetic fields serves as a “dual” set-up for that at finite density which is highly restricted due to the sign problem.

ACKNOWLEDGMENTS

The authors thank Yasufumi Araki and Kazunori Itakura for helpful discussions. This work was supported by Japan Society for the Promotion of Science (JSPS) KAKENHI under grant Nos. JP17K14277, JP20K14476, 20K03948 and 22H02316, and by the RIKEN special postdoctoral researcher program.

Appendix A: Model Lagrangian

Here, we provide a detailed explanation of the model Lagrangian (1) used in the main text.

1. Light-fermion kinetic term

The kinetic and mass terms for the light fermions is given by the Dirac Lagrangian in the 3 + 1 dimensional spacetime

$$\mathcal{L}_l \equiv \bar{\psi}(i\mathcal{D} - m_l)\psi, \quad (\text{A1})$$

where $\mathcal{D} \equiv \gamma^\mu D_\mu$ with the Dirac (gamma) matrix γ^μ defined by the Clifford algebra [15, 157]. We assume the Einstein notation for the repeated Lorentz indices ($\mu = 0, 1, 2, 3$). The four-component Dirac spinor ψ is composed of the two spin states of particles and antiparticles (holes). The covariant derivative, $D^\mu = \partial^\mu + q_l A_{\text{ext}}^\mu$, contains the gauge field for an external magnetic field, and satisfies the commutation relation $[iD^1, iD^2] = iq_l B$ with all the other vanishing combinations. Here, without loss of generality, we assume a constant magnetic field in the third direction with the strength B .

We briefly explain how the Lagrangian (A1) reduces to that given in Eq. (1) in the lowest Landau level (LLL). By the use of the above commutation relation, one can introduce the annihilation and creation operators

$$a, a^\dagger = \frac{i}{\sqrt{2|q_l B|}} [D^1 \pm \text{sgn}(q_l B) iD^2], \quad (\text{A2})$$

where the upper and lower signs are for a and a^\dagger , respectively. These operators satisfy $[a, a^\dagger] = 1$. One can lower and raise the Landau levels with these operators exactly in the same manner as for the harmonic oscillator.

The Dirac operator is identically cast into the form

$$i\mathcal{D} - m_l = i\mathcal{D}_\parallel - m_l - \sqrt{2|q_l B|} \gamma^1 (a\mathcal{P}_+ + a^\dagger\mathcal{P}_-), \quad (\text{A3})$$

with the spin-projection operator

$$\mathcal{P}_\pm = \frac{1}{2} (1 \pm i \text{sgn}(q_l B) \gamma^1 \gamma^2). \quad (\text{A4})$$

In Eq. (A3), the subscript of \mathcal{D}_\parallel means the parallel direction along the magnetic field, i.e., $\mathcal{D}_\parallel \equiv \gamma^0 \partial_0 + \gamma^3 \partial_3$. The LLL is the ground state that is annihilated by the

operator a and is a spin eigenstate with the lower Zeeman energy projected by \mathcal{P}_+ . Therefore, the last term in Eq. (A3) vanishes for the LLL, and we are left with the Lagrangian (1).

It is instructive to see the Landau levels explicitly. The Klein-Gordon operator reads

$$(i\mathcal{D} + m_l)(i\mathcal{D} - m_l)\psi_\pm \quad (\text{A5}) \\ = - [\partial_t^2 - \partial_z^2 + (2a^\dagger a + 1 \mp 1)|q_l B| + m_l^2] \psi_\pm,$$

where we defined the spin eigenstates $\psi_\pm = \mathcal{P}_\pm \psi$. We readily find the energy levels $E^2 = p_z^2 + (2n + 1 \pm 1)|q_l B|$ with a non-negative integer $n \geq 0$ when the Klein-Gordon equation is satisfied by ψ_\pm . Note that the dispersion relation for the LLL, and thus the kinetic term in the Lagrangian (1), do not depend on the magnetic-field strength B .

2. Heavy-fermion kinetic term

We also briefly summarize the derivation of the low-energy heavy-fermion kinetic term in terms of the large-mass expansion [126–128]. We decompose the heavy-fermion momentum as

$$p^\mu = m_h v^\mu + k^\mu, \quad (\text{A6})$$

where the four velocity, $v^\mu = (v^0, \mathbf{v})$, is normalized as $v^\mu v_\mu = 1$, and k^μ is the residual momentum with a scale much smaller than the heavy-fermion mass m_h . We introduce projection operators

$$\mathcal{Q}_\pm = \frac{1}{2} (1 \pm \not{v}), \quad (\text{A7})$$

with $\not{v} = \gamma^\mu v_\mu$. In the rest frame ($\mathbf{v} = 0$), these operators project out the particle and antiparticle states. The low-energy excitation will be an eigenstate of \mathcal{Q}_\pm since the coupling between the particle and antiparticle states is highly suppressed by the large mass.

One can organize a low-energy effective theory in terms of the $1/m_h$ expansion in such a way that there is no mixing between the eigenstates of \mathcal{Q}_\pm [127, 128]. To get the leading-order term in the expansion, one can factorize the heavy-fermion field Ψ as

$$\Psi = e^{-im_h v^\mu x_\mu} e^{i\frac{\not{v}_\perp}{2m_h}} \Psi_v^+ + e^{+im_h v^\mu x_\mu} e^{i\frac{\not{v}_\perp}{2m_h}} \Psi_v^- \\ = e^{-im_h v^\mu x_\mu} (1 + \frac{i\not{v}_\perp}{2m_h}) \Psi_v^+ \\ + e^{+im_h v^\mu x_\mu} (1 + \frac{i\not{v}_\perp}{2m_h}) \Psi_v^- + \mathcal{O}(m_h^{-2}). \quad (\text{A8})$$

In other words, one can define eigenstates of \mathcal{Q}_\pm as $\Psi_v^\pm \equiv e^{-i\frac{\not{v}_\perp}{2m_h}} e^{\pm im_h v^\mu x_\mu} \mathcal{Q}_\pm \Psi$ that satisfy $\mathcal{Q}_\pm \Psi_v^\pm = \Psi_v^\pm$. The exponential factor works for the leading-order result with $\not{v}_\perp \equiv (\gamma^\mu - \not{v} v^\mu) \partial_\mu$, and needs to be improved on

an order-by-order basis (see Refs. [127, 128]). Multiplying the above expansion (A8) by the Dirac operator, we obtain an expansion of the heavy-fermion kinetic term

$$\begin{aligned}\mathcal{L}_h &\equiv \bar{\Psi}(i\not{\partial} - m_h)\Psi \\ &= \sum_{c=\pm} c\bar{\Psi}_v^c (iv^\mu\partial_\mu)\Psi_v^c + \mathcal{O}(m_h^{-1}),\end{aligned}\quad (\text{A9})$$

where we used identities

$$\begin{aligned}\mathcal{Q}_\pm\mathcal{Q}_\pm &= \mathcal{Q}_\pm, & \mathcal{Q}_\pm\mathcal{Q}_\mp &= 0, & \mathcal{Q}_\pm\gamma^\mu\mathcal{Q}_\pm &= \pm v^\mu\mathcal{Q}_\pm, \\ \mathcal{Q}_\pm\gamma^\mu\mathcal{Q}_\mp &= (\gamma^\mu - \not{v}v^\mu)\mathcal{Q}_\mp.\end{aligned}\quad (\text{A10})$$

The rightmost side in Eq. (A9) is used in the Lagrangian (1) as the low-energy heavy-fermion kinetic term.

3. Interaction Lagrangian

We examine a sequence of the terms generated from an interaction Lagrangian that has the Lorentz and $SU(N)$ symmetries [43, 47, 49, 158]:

$$\mathcal{L}_{\text{int}} = -G(\bar{\psi}\gamma^\mu T^a\psi)(\bar{\Psi}\gamma_\mu T^a\Psi),\quad (\text{A11})$$

where γ^μ ($\mu = 0, 1, 2, 3$) are the Dirac matrices, and T^a ($a = 1, 2, \dots, N^2 - 1$) are the generator matrices of the $SU(N)$ group. The light and heavy fermions, ψ and Ψ , belong to the fundamental representation of the $SU(N)$ symmetry. The sums over μ and a are implicitly taken. Here, $G > 0$ is the coupling constant. This is a simple analog of the NJL interaction [34–37], where the inter-fermion interactions are understood as the mimic of the gluon exchange interactions between quarks in QCD.

The matrix structures in Eq. (A11) can be decomposed by utilizing the Fierz transformations, i.e.,

$$\begin{aligned}(\gamma^\mu)_{\alpha\beta}(\gamma_\mu)_{\gamma\delta} &= \delta_{\alpha\delta}\delta_{\gamma\beta} + (i\gamma_5)_{\alpha\delta}(i\gamma_5)_{\gamma\beta} \\ &\quad - \frac{1}{2}(\gamma^\mu)_{\alpha\delta}(\gamma_\mu)_{\gamma\beta} - \frac{1}{2}(\gamma^\mu\gamma_5)_{\alpha\delta}(\gamma_\mu\gamma_5)_{\gamma\beta},\end{aligned}\quad (\text{A12})$$

for the Dirac matrices and

$$(T^a)_{ij}(T^a)_{kl} = \frac{N^2 - 1}{2N^2}\delta_{il}\delta_{kj} - \frac{1}{N}(T^a)_{il}(T^a)_{kj},\quad (\text{A13})$$

for the $SU(N)$ generators. As we briefly discuss below (cf. Refs. [47–49]), we identify the dominant terms, that are maintained in Eq. (1) in the main text, as the singlet channel $\delta_{il}\delta_{kj}$ in the $SU(N)$ generators. Then, we have a relation between the coupling strengths, G_{hl} in Eq. (1) and G in Eq. (A11), as $G_{hl}/N = G(N^2 - 1)/(2N^2)$. As for the Dirac matrices, we focus on the scalar $\delta_{\alpha\delta}\delta_{\gamma\beta}$ and pseudoscalar $(i\gamma_5)_{\alpha\delta}(i\gamma_5)_{\gamma\beta}$ terms, and potential realization of the vector and axial-vector condensates in magnetic fields are not considered in Eq. (1).

We note that the above singlet channel is the most dominant one in the large N limit [47–49], where the

mean-field approximation can be justified.⁴ The other channel, that is, the adjoint representation from the second term on the right-hand side of Eq. (A13), is suppressed in comparison to the singlet channel in the large N limit.

In addition to the ‘‘particle-antiparticle’’ channel ($\psi\Gamma'\bar{\Psi}$ and $\bar{\Psi}\bar{\Gamma}'\psi$ types) in Eq. (1), one can get the ‘‘particle-particle’’ channels ($\psi^t\Gamma\Psi$ and $\Psi^t\Gamma^t\psi$ types) from Eq. (A11) as interactions between the light and heavy fermions (see, e.g., Ref. [159] for similar discussions in case of light-light fermion interactions). Here Γ , Γ^t (the transpose of Γ), Γ' , and $\bar{\Gamma}' = \gamma_0\Gamma'^t\gamma_0$ (the complex conjugate of Γ') are appropriate combinations of the Dirac matrices resulting from the Fierz transformations. These terms are also suppressed in the large N limit.

Appendix B: Solutions for gap equations

From Eq. (4) in the main text, the stationary conditions $\partial\Omega/\partial M = 0 = \partial\Omega/\partial\Delta$ lead to the gap equations

$$\frac{M}{G_{ll}} + \frac{M + m_l}{2\pi}\rho_B \ln \Xi = 0, \quad \frac{\Delta}{G_{hl}} + \frac{\Delta}{2\pi}\rho_B \ln \Xi = 0, \quad (\text{B1})$$

where

$$\Xi = \frac{(M + m_l)^2 + 4|\Delta|^2}{(\Lambda + \sqrt{(M + m_l)^2 + 4|\Delta|^2 + \Lambda^2})^2}.\quad (\text{B2})$$

By solving these gap equations, we can determine M and Δ .

1. Without competition

We briefly investigate special cases where $G_{hl} = 0$ or $G_{ll} = 0$. According to the discussion around Eq. (5), we find nonzero solutions for the gap equations at any coupling strengths. When $G_{hl} = 0$ in the Lagrangian (1), the gap equation becomes the first form in Eq. (B1) with $\Delta = 0$. In the chiral limit $m_l = 0$, we find an analytic solution ($M > 0$):

$$M = \frac{\Lambda}{\sinh\left(\frac{\pi}{\rho_B G_{ll}}\right)} \rightarrow 2\Lambda e^{-\frac{\pi}{\rho_B G_{ll}}},\quad (\text{B3})$$

which approaches the rightmost side for a small value of G_{ll} . The chiral symmetry is always broken under a strong magnetic field and a nonzero coupling constant by the mechanism of *magnetic catalysis* [18–21].

On the other hand, when $G_{ll} = 0$ in the Lagrangian (1), the gap equation becomes the second form of

⁴ See, e.g., Refs. [121, 122] for early works on the large N limit applied to the Kondo effect and Ref. [33] for generalities.

Eq. (B1) with $M = 0$. In the chiral limit, we find an analytic solution ($\Delta > 0$):

$$\Delta = \frac{\Lambda}{2 \sinh\left(\frac{\pi}{\rho_B G_{hl}}\right)} \rightarrow \Lambda e^{-\frac{\pi}{\rho_B G_{hl}}}, \quad (\text{B4})$$

which approaches the rightmost side for a small value of G_{hl} . This solution clearly indicates the occurrence of the Kondo effect by a strong magnetic field. The novel magnetically induced Kondo effect was previously suggested also by a renormalization-group (RG) analysis [22] and further analyzed by a conformal-field theory approach [52]. Our mean-field solution (B4) is consistent with the emergent scale in the previous RG analysis [22] and determines the magnitude of the Kondo condensate for the first time. The magnetic-field dependence of Δ at

$G_{ll}\Lambda^2 = 1.0$ is shown with the blue dotted line in Fig. 1.

2. With competition

We also obtain analytic solutions in the presence of the competition between the Kondo effect and the magnetic catalysis with nonzero G_{hl} and G_{ll} . We immediately notice that the gap equations (B1) have a trivial solution $\Delta = 0$. In this case, M takes a nontrivial solution which is an extension of Eq. (B3) with a nonzero m_l . We call these solutions $(M_1, \Delta_1 = 0)$. By eliminating $\ln \Xi$ in Eq. (B1), one can find another set of solutions that we call (M_2, Δ_2) and show in Eq. (6).

-
- [1] D. Grasso and H. R. Rubinstein, Magnetic fields in the early universe, *Phys. Rept.* **348**, 163 (2001), [arXiv:astro-ph/0009061 \[astro-ph\]](#).
- [2] W. Dittrich and H. Gies, Probing the quantum vacuum. Perturbative effective action approach in quantum electrodynamics and its application, *Springer Tracts Mod. Phys.* **166**, 1 (2000).
- [3] D. Yoshioka, *The quantum Hall effect*, Vol. 133 (Springer Berlin, Heidelberg, 2002).
- [4] G. V. Dunne, Heisenberg-Euler effective Lagrangians: Basics and extensions, in *From fields to strings: Circumnavigating theoretical physics. Ian Kogan memorial collection (3 volume set)*, edited by M. Shifman, A. Vainshtein, and J. Wheeler (2004) pp. 445–522, [arXiv:hep-th/0406216 \[hep-th\]](#).
- [5] A. K. Harding and D. Lai, Physics of Strongly Magnetized Neutron Stars, *Rept. Prog. Phys.* **69**, 2631 (2006), [arXiv:astro-ph/0606674 \[astro-ph\]](#).
- [6] G. A. Mourou, T. Tajima, and S. V. Bulanov, Optics in the relativistic regime, *Rev. Mod. Phys.* **78**, 309 (2006).
- [7] D. Kharzeev, K. Landsteiner, A. Schmitt, and H.-U. Yee, Strongly Interacting Matter in Magnetic Fields, *Lect. Notes Phys.* **871**, pp.1 (2013).
- [8] D. E. Kharzeev, The Chiral Magnetic Effect and Anomaly-Induced Transport, *Prog. Part. Nucl. Phys.* **75**, 133 (2014), [arXiv:1312.3348 \[hep-ph\]](#).
- [9] D. E. Kharzeev, J. Liao, S. A. Voloshin, and G. Wang, Chiral magnetic and vortical effects in high-energy nuclear collisions—A status report, *Prog. Part. Nucl. Phys.* **88**, 1 (2016), [arXiv:1511.04050 \[hep-ph\]](#).
- [10] K. Hattori and X.-G. Huang, Novel quantum phenomena induced by strong magnetic fields in heavy-ion collisions, *Nucl. Sci. Tech.* **28**, 26 (2017), [arXiv:1609.00747 \[nucl-th\]](#).
- [11] V. Koch, S. Schlichting, V. Skokov, P. Sorensen, J. Thomas, S. Voloshin, G. Wang, and H.-U. Yee, Status of the chiral magnetic effect and collisions of isobars, *Chin. Phys. C* **41**, 072001 (2017), [arXiv:1608.00982 \[nucl-th\]](#).
- [12] A. A. Burkov, Chiral anomaly and transport in Weyl metals, *J. Phys. Condens. Matter* **27**, 113201 (2015), [arXiv:1502.07609 \[cond-mat.mes-hall\]](#).
- [13] D. E. Kharzeev, Topology, magnetic field, and strongly interacting matter, *Ann. Rev. Nucl. Part. Sci.* **65**, 193 (2015), [arXiv:1501.01336 \[hep-ph\]](#).
- [14] V. A. Miransky and I. A. Shovkovy, Quantum field theory in a magnetic field: From quantum chromodynamics to graphene and Dirac semimetals, *Phys. Rept.* **576**, 1 (2015), [arXiv:1503.00732 \[hep-ph\]](#).
- [15] N. P. Armitage, E. J. Mele, and A. Vishwanath, Weyl and Dirac Semimetals in Three Dimensional Solids, *Rev. Mod. Phys.* **90**, 015001 (2018), [arXiv:1705.01111 \[cond-mat.str-el\]](#).
- [16] M. D’Elia, Lattice QCD Simulations in External Background Fields, *Lect. Notes Phys.* **871**, 181 (2013), [arXiv:1209.0374 \[hep-lat\]](#).
- [17] G. Endrődi, Critical point in the QCD phase diagram for extremely strong background magnetic fields, *JHEP* **07**, 173, [arXiv:1504.08280 \[hep-lat\]](#).
- [18] V. P. Gusynin, V. A. Miransky, and I. A. Shovkovy, Catalysis of Dynamical Flavor Symmetry Breaking by a Magnetic Field in 2 + 1 Dimensions, *Phys. Rev. Lett.* **73**, 3499 (1994), [Erratum: *Phys. Rev. Lett.* **76**, 1005 (1996)], [arXiv:hep-ph/9405262 \[hep-ph\]](#).
- [19] V. P. Gusynin, V. A. Miransky, and I. A. Shovkovy, Dynamical flavor symmetry breaking by a magnetic field in 2+1 dimensions, *Phys. Rev. D* **52**, 4718 (1995), [arXiv:hep-th/9407168 \[hep-th\]](#).
- [20] V. P. Gusynin, V. A. Miransky, and I. A. Shovkovy, Dimensional reduction and dynamical chiral symmetry breaking by a magnetic field in 3 + 1 dimensions, *Phys. Lett. B* **349**, 477 (1995), [arXiv:hep-ph/9412257 \[hep-ph\]](#).
- [21] V. P. Gusynin, V. A. Miransky, and I. A. Shovkovy, Dimensional reduction and catalysis of dynamical symmetry breaking by a magnetic field, *Nucl. Phys. B* **462**, 249 (1996), [arXiv:hep-ph/9509320 \[hep-ph\]](#).
- [22] S. Ozaki, K. Itakura, and Y. Kuramoto, Magnetically induced QCD Kondo effect, *Phys. Rev. D* **94**, 074013 (2016), [arXiv:1509.06966 \[hep-ph\]](#).
- [23] J. Polchinski, Effective field theory and the Fermi surface, in *Theoretical Advanced Study Institute (TASI 92): From Black Holes and Strings to Particles Boulder, Colorado, June 3-28, 1992* (1992) pp. 0235–276, [arXiv:hep-](#)

- th/9210046 [hep-th].
- [24] R. Shankar, Renormalization group approach to interacting fermions, *Rev. Mod. Phys.* **66**, 129 (1994), [arXiv:cond-mat/9307009](#).
- [25] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Theory of superconductivity, *Phys. Rev.* **108**, 1175 (1957).
- [26] J. Kondo, Resistance Minimum in Dilute Magnetic Alloys, *Prog. Theor. Phys.* **32**, 37 (1964).
- [27] A. A. Abrikosov, Electron scattering on magnetic impurities in metals and anomalous resistivity effects, *Physics Physique Fizika* **2**, 5 (1965).
- [28] P. W. Anderson, A poor man's derivation of scaling laws for the Kondo problem, *J. Phys. C: Solid State Phys.* **3**, 2436 (1970).
- [29] K. G. Wilson, The renormalization group: Critical phenomena and the Kondo problem, *Rev. Mod. Phys.* **47**, 773 (1975).
- [30] A. C. Hewson, *The Kondo Problem to Heavy Fermions* (Cambridge University Press, Cambridge, England, 1993).
- [31] K. Yosida, *Theory of Magnetism* (Springer-Verlag, Berlin, 1996).
- [32] K. Yamada, *Electron Correlation in Metals* (Cambridge University Press, Cambridge, England, 2004).
- [33] P. Coleman, *Introduction to Many-Body Physics* (Cambridge University Press, Cambridge, England, 2015).
- [34] Y. Nambu and G. Jona-Lasinio, Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. I, *Phys. Rev.* **122**, 345 (1961).
- [35] Y. Nambu and G. Jona-Lasinio, Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. II, *Phys. Rev.* **124**, 246 (1961).
- [36] S. P. Klevansky, The Nambu-Jona-Lasinio model of quantum chromodynamics, *Rev. Mod. Phys.* **64**, 649 (1992).
- [37] T. Hatsuda and T. Kunihiro, QCD phenomenology based on a chiral effective Lagrangian, *Phys. Rept.* **247**, 221 (1994), [arXiv:hep-ph/9401310](#) [hep-ph].
- [38] M. Iwasaki and T. Iwado, Superconductivity in quark matter, *Phys. Lett. B* **350**, 163 (1995).
- [39] M. G. Alford, K. Rajagopal, and F. Wilczek, QCD at finite baryon density: Nucleon droplets and color superconductivity, *Phys. Lett. B* **422**, 247 (1998), [arXiv:hep-ph/9711395](#).
- [40] R. Rapp, T. Schäfer, E. V. Shuryak, and M. Velkovsky, Diquark Bose Condensates in High Density Matter and Instantons, *Phys. Rev. Lett.* **81**, 53 (1998), [arXiv:hep-ph/9711396](#) [hep-ph].
- [41] M. G. Alford, A. Schmitt, K. Rajagopal, and T. Schäfer, Color superconductivity in dense quark matter, *Rev. Mod. Phys.* **80**, 1455 (2008), [arXiv:0709.4635](#) [hep-ph].
- [42] K. Fukushima and T. Hatsuda, The phase diagram of dense QCD, *Rept. Prog. Phys.* **74**, 014001 (2011), [arXiv:1005.4814](#) [hep-ph].
- [43] S. Yasui and K. Sudoh, Heavy-quark dynamics for charm and bottom flavor on the Fermi surface at zero temperature, *Phys. Rev. C* **88**, 015201 (2013), [arXiv:1301.6830](#) [hep-ph].
- [44] K. Hattori, K. Itakura, S. Ozaki, and S. Yasui, QCD Kondo effect: Quark matter with heavy-flavor impurities, *Phys. Rev. D* **92**, 065003 (2015), [arXiv:1504.07619](#) [hep-ph].
- [45] T. Kanazawa and S. Uchino, Overscreened Kondo effect, (color) superconductivity, and Shiba states in Dirac metals and quark matter, *Phys. Rev. D* **94**, 114005 (2016), [arXiv:1609.00033](#) [cond-mat.str-el].
- [46] T. Kimura and S. Ozaki, Fermi/non-Fermi mixing in $SU(N)$ Kondo effect, *J. Phys. Soc. Jpn.* **86**, 084703 (2017), [arXiv:1611.07284](#) [cond-mat.str-el].
- [47] S. Yasui, K. Suzuki, and K. Itakura, Kondo phase diagram of quark matter, *Nucl. Phys. A* **983**, 90 (2019), [arXiv:1604.07208](#) [hep-ph].
- [48] S. Yasui, Kondo cloud of single heavy quark in cold and dense matter, *Phys. Lett. B* **773**, 428 (2017), [arXiv:1608.06450](#) [hep-ph].
- [49] S. Yasui, K. Suzuki, and K. Itakura, Topology and stability of the Kondo phase in quark matter, *Phys. Rev. D* **96**, 014016 (2017), [arXiv:1703.04124](#) [hep-ph].
- [50] K. Suzuki, S. Yasui, and K. Itakura, Interplay between chiral symmetry breaking and the QCD Kondo effect, *Phys. Rev. D* **96**, 114007 (2017), [arXiv:1708.06930](#) [hep-ph].
- [51] S. Yasui and S. Ozaki, Transport coefficients from the QCD Kondo effect, *Phys. Rev. D* **96**, 114027 (2017), [arXiv:1710.03434](#) [hep-ph].
- [52] T. Kimura and S. Ozaki, Conformal field theory analysis of the QCD Kondo effect, *Phys. Rev. D* **99**, 014040 (2019), [arXiv:1806.06486](#) [hep-ph].
- [53] R. Fariello, J. C. Macías, and F. S. Navarra, The QCD Kondo phase in quark stars, (2019), [arXiv:1901.01623](#) [nucl-th].
- [54] K. Hattori, X.-G. Huang, and R. D. Pisarski, Emergent QCD Kondo effect in two-flavor color superconducting phase, *Phys. Rev. D* **99**, 094044 (2019), [arXiv:1903.10953](#) [hep-ph].
- [55] D. Suenaga, K. Suzuki, and S. Yasui, QCD Kondo excitons, *Phys. Rev. Research* **2**, 023066 (2020), [arXiv:1909.07573](#) [nucl-th].
- [56] D. Suenaga, K. Suzuki, Y. Araki, and S. Yasui, Kondo effect driven by chirality imbalance, *Phys. Rev. Research* **2**, 023312 (2020), [arXiv:1912.12669](#) [hep-ph].
- [57] T. Kanazawa, Random matrix model for the QCD Kondo effect, (2020), [arXiv:2006.00200](#) [hep-th].
- [58] Y. Araki, D. Suenaga, K. Suzuki, and S. Yasui, Two relativistic Kondo effects: Classification with particle and antiparticle impurities, *Phys. Rev. Research* **3**, 013233 (2021), [arXiv:2008.08434](#) [hep-ph].
- [59] Y. Araki, D. Suenaga, K. Suzuki, and S. Yasui, Spin-orbital magnetic response of relativistic fermions with band hybridization, *Phys. Rev. Research* **3**, 023098 (2021), [arXiv:2011.00882](#) [cond-mat.mes-hall].
- [60] T. Kimura, ABCD of Kondo effect, *J. Phys. Soc. Jpn.* **90**, 024708 (2021), [arXiv:2011.08301](#) [cond-mat.str-el].
- [61] D. Suenaga, Y. Araki, K. Suzuki, and S. Yasui, Chiral separation effect catalyzed by heavy impurities, *Phys. Rev. D* **103**, 054041 (2021), [arXiv:2012.15173](#) [hep-ph].
- [62] T. Ishikawa, K. Nakayama, and K. Suzuki, Kondo effect with Wilson fermions, *Phys. Rev. D* **104**, 094515 (2021), [arXiv:2107.07270](#) [hep-lat].
- [63] D. Suenaga, Y. Araki, K. Suzuki, and S. Yasui, Heavy-quark spin polarization induced by the Kondo effect in a magnetic field, *Phys. Rev. D* **105**, 074028 (2022), [arXiv:2109.14799](#) [hep-ph].
- [64] G. W. Semenoff, I. A. Shovkovy, and L. C. R. Wijewardhana, Universality and the magnetic catalysis of chiral symmetry breaking, *Phys. Rev. D* **60**, 105024 (1999), [arXiv:hep-th/9905116](#).

- [65] G. W. Semenoff, I. A. Shovkovy, and L. C. R. Wijewardhana, Phase transition induced by a magnetic field, *Mod. Phys. Lett. A* **13**, 1143 (1998), [arXiv:hep-ph/9803371](#).
- [66] D. V. Khveshchenko, Magnetic-Field-Induced Insulating Behavior in Highly Oriented Pyrolytic Graphite, *Phys. Rev. Lett.* **87**, 206401 (2001), [arXiv:cond-mat/0106261](#).
- [67] E. V. Gorbar, V. P. Gusynin, V. A. Miransky, and I. A. Shovkovy, Magnetic field driven metal insulator phase transition in planar systems, *Phys. Rev. B* **66**, 045108 (2002), [arXiv:cond-mat/0202422](#).
- [68] V. P. Gusynin, V. A. Miransky, S. G. Sharapov, and I. A. Shovkovy, Excitonic gap, phase transition, and quantum Hall effect in graphene, *Phys. Rev. B* **74**, 195429 (2006), [arXiv:cond-mat/0605348 \[cond-mat.str-el\]](#).
- [69] I. F. Herbut, Theory of integer quantum Hall effect in graphene, *Phys. Rev. B* **75**, 165411 (2007), [arXiv:cond-mat/0610349 \[cond-mat.mes-hall\]](#).
- [70] I. F. Herbut and B. Roy, Quantum critical scaling in magnetic field near the Dirac point in graphene, *Phys. Rev. B* **77**, 245438 (2008), [arXiv:0802.2546 \[cond-mat.mes-hall\]](#).
- [71] V. P. Gusynin, V. A. Miransky, S. G. Sharapov, I. A. Shovkovy, and C. M. Wyenberg, Edge states on graphene ribbons in magnetic field: Interplay between Dirac and ferromagnetic-like gaps, *Phys. Rev. B* **79**, 115431 (2009), [arXiv:0801.0708 \[cond-mat.mes-hall\]](#).
- [72] B. Roy and I. F. Herbut, Inhomogeneous magnetic catalysis on graphene's honeycomb lattice, *Phys. Rev. B* **83**, 195422 (2011), [arXiv:1102.3481 \[cond-mat.mes-hall\]](#).
- [73] B. Roy, M. P. Kennett, and S. D. Sarma, Chiral symmetry breaking and the quantum Hall effect in monolayer graphene, *Phys. Rev. B* **90**, 201409 (2014), [arXiv:1406.5184 \[cond-mat.mes-hall\]](#).
- [74] B. Roy and J. D. Sau, Magnetic catalysis and axionic charge-density-wave in Weyl semimetals, *Phys. Rev. B* **92**, 125141 (2015), [arXiv:1406.4501 \[cond-mat.mes-hall\]](#).
- [75] X.-T. Zhang and R. Shindou, Transport properties of density wave phases in three-dimensional metals and semimetals under high magnetic field, *Phys. Rev. B* **95**, 205108 (2017).
- [76] Z. Song, Z. Fang, and X. Dai, Instability of Dirac semimetal phase under a strong magnetic field, *Phys. Rev. B* **96**, 235104 (2017).
- [77] C. DeTar, C. Winterowd, and S. Zafeiropoulos, Magnetic Catalysis in Graphene Effective Field Theory, *Phys. Rev. Lett.* **117**, 266802 (2016), [arXiv:1607.03137 \[hep-lat\]](#).
- [78] D. L. Boyda, V. V. Braguta, S. N. Valgushev, M. I. Polikarpov, and M. V. Ulybyshev, Numerical simulation of graphene in an external magnetic field, *Phys. Rev. B* **89**, 245404 (2014), [arXiv:1308.2814 \[hep-lat\]](#).
- [79] C. DeTar, C. Winterowd, and S. Zafeiropoulos, Lattice field theory study of magnetic catalysis in graphene, *Phys. Rev. B* **95**, 165442 (2017), [arXiv:1608.00666 \[hep-lat\]](#).
- [80] Y. Otsuka, S. Yunoki, and S. Sorella, Universal Quantum Criticality in the Metal-Insulator Transition of Two-Dimensional Interacting Dirac Electrons, *Phys. Rev. X* **6**, 011029 (2016), [arXiv:1510.08593 \[cond-mat.str-el\]](#).
- [81] Y. Tada, Quantum criticality of magnetic catalysis in two-dimensional correlated Dirac fermions, *Phys. Rev. Research* **2**, 033363 (2020), [arXiv:2005.01990 \[cond-mat.str-el\]](#).
- [82] A. Principi, G. Vignale, and E. Rossi, Kondo effect and non-Fermi-liquid behavior in Dirac and Weyl semimetals, *Phys. Rev. B* **92**, 041107(R) (2015), [arXiv:1410.8532 \[cond-mat.mes-hall\]](#).
- [83] T. Yanagisawa, Dirac fermions and Kondo effect, *J. Phys. Conf. Ser.* **603**, 012014 (2015), [arXiv:1502.07898 \[cond-mat.str-el\]](#).
- [84] T. Yanagisawa, Kondo Effect in Dirac Systems, *J. Phys. Soc. Jpn.* **84**, 074705 (2015), [arXiv:1505.05295 \[cond-mat.str-el\]](#).
- [85] A. K. Mitchell and L. Fritz, Kondo effect in three-dimensional Dirac and Weyl systems, *Phys. Rev. B* **92**, 121109(R) (2015), [arXiv:1506.05491 \[cond-mat.str-el\]](#).
- [86] J.-H. Sun, D.-H. Xu, F.-C. Zhang, and Y. Zhou, A magnetic impurity in a Weyl semimetal, *Phys. Rev. B* **92**, 195124 (2015), [arXiv:1509.05180 \[cond-mat.str-el\]](#).
- [87] X.-Y. Feng, H. Zhong, J. Dai, and Q. Si, Dirac-Kondo semimetals and topological Kondo insulators in the dilute carrier limit, [arXiv:1605.02380 \[cond-mat.str-el\]](#).
- [88] H.-H. Lai, S. E. Grefe, S. Paschen, and Q. Si, Weyl-Kondo semimetal in heavy-fermion systems, *Proc. Natl. Acad. Sci. U.S.A.* **115**, 93 (2018), [arXiv:1612.03899 \[cond-mat.str-el\]](#).
- [89] S. Ok, M. Legner, T. Neupert, and A. M. Cook, Magnetic Weyl and Dirac Kondo semimetal phases in heterostructures, [arXiv:1703.03804 \[cond-mat.str-el\]](#).
- [90] D. Ma, H. Chen, H. Liu, and X. C. Xie, Kondo effect with Weyl semimetal Fermi arcs, *Phys. Rev. B* **97**, 045148 (2018), [arXiv:1709.08008 \[cond-mat.str-el\]](#).
- [91] L. Li, J.-H. Sun, Z.-H. Wang, D.-H. Xu, H.-G. Luo, and W.-Q. Chen, Magnetic states and Kondo screening in Weyl semimetals with chiral anomaly, *Phys. Rev. B* **98**, 075110 (2018).
- [92] S. Dzsaber, X. Yan, M. Taupin, G. Eguchi, A. Prokofiev, T. Shiroka, P. Blaha, O. Rubel, S. E. Grefe, H.-H. Lai, Q. Si, and S. Paschen, Giant spontaneous Hall effect in a nonmagnetic Weyl-Kondo semimetal, *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2013386118 (2021), [arXiv:1811.02819 \[cond-mat.str-el\]](#).
- [93] H.-F. Lü, Y.-H. Deng, S.-S. Ke, Y. Guo, and H.-W. Zhang, Quantum impurity in topological multi-Weyl semimetals, *Phys. Rev. B* **99**, 115109 (2019).
- [94] K.-S. Kim and J.-H. Han, Interplay between chiral magnetic and Kondo effects in Weyl metal phase, *Curr. Appl. Phys.* **19**, 236 (2019).
- [95] S. E. Grefe, H.-H. Lai, S. Paschen, and Q. Si, Weyl-Kondo semimetals in nonsymmorphic systems, *Phys. Rev. B* **101**, 075138 (2020), [arXiv:1911.01400 \[cond-mat.str-el\]](#).
- [96] S. E. Grefe, H.-H. Lai, S. Paschen, and Q. Si, Extreme topological tunability of Weyl-Kondo semimetal to Zeeman coupling, [arXiv:2012.15841 \[cond-mat.str-el\]](#).
- [97] G. T. D. Pedrosa, J. F. Silva, and E. Vernek, Kondo screening regimes in multi-Dirac and Weyl systems, *Phys. Rev. B* **103**, 045137 (2021), [arXiv:2101.12531 \[cond-mat.str-el\]](#).
- [98] L. Fritz and M. Vojta, The physics of Kondo impurities in graphene, *Rep. Prog. Phys.* **76**, 032501 (2013), [arXiv:1208.3113 \[cond-mat.str-el\]](#).

- [99] T. Soda, T. Matsuura, and Y. Nagaoka, *s-d* exchange interaction in a superconductor, *Prog. Theor. Phys.* **38**, 551 (1967).
- [100] M. Fowler and K. Maki, Conditions for bound states in a superconductor with a magnetic impurity, *Phys. Rev.* **164**, 484 (1967); Conditions for Bound States in a Superconductor with a Magnetic Impurity. II, *Phys. Rev. B* **1**, 181 (1970).
- [101] J. Zittartz and E. Müller-Hartmann, Theory of magnetic impurities in superconductors. I, *Z. Physik* **232**, 11; E. Müller-Hartmann and J. Zittartz, Theory of magnetic impurities in superconductors. II, *Z. Physik* **234**, 58 (1970); Kondo Effect in Superconductors, *Phys. Rev. Lett.* **26**, 428 (1971).
- [102] A. Sakurai, Comments on superconductors with magnetic impurities, *Prog. Theor. Phys.* **44**, 1472 (1970).
- [103] T. Matsuura and Y. Nagaoka, Depression of superconducting critical temperature by impurities with Kondo effect, *Solid State Commun.* **18**, 1583 (1976); T. Matsuura, The effects of impurities on superconductors with Kondo effect, *Prog. Theor. Phys.* **57**, 1823 (1977); T. Matsuura, S. Ichinose, and Y. Nagaoka, Theory of Kondo effect in superconductors. I: transition temperature and upper critical field, *Prog. Theor. Phys.* **57**, 713 (1977).
- [104] M. Jarrell, D. S. Sivia, and B. Patton, Gap states in dilute-magnetic-alloy superconductors: A quantum Monte Carlo study, *Phys. Rev. B* **42**, 4804 (1990).
- [105] K. Satori, H. Shiba, O. Sakai, and Y. Shimizu, Numerical renormalization group study of magnetic impurities in superconductors, *J. Phys. Soc. Jpn.* **61**, 3239 (1992); H. Shiba, K. Satori, O. Sakai, and Y. Shimizu, Numerical renormalization group study of the Kondo effect in superconductors, *Physica B Condens. Matter* **186**, 239 (1993); O. Sakai, Y. Shimizu, H. Shiba, and K. Satori, Numerical renormalization group study of magnetic impurities in superconductors. ii. dynamical excitation spectra and spatial variation of the order parameter, *J. Phys. Soc. Jpn.* **62**, 3181 (1993).
- [106] M. I. Salkola, A. V. Balatsky, and J. R. Schrieffer, Spectral properties of quasiparticle excitations induced by magnetic moments in superconductors, *Phys. Rev. B* **55**, 12648 (1997), [arXiv:cond-mat/9610198](https://arxiv.org/abs/cond-mat/9610198) [cond-mat.supr-con].
- [107] J. Bauer, J. I. Pascual, and K. J. Franke, Microscopic resolution of the interplay of Kondo screening and superconducting pairing: Mn-phthalocyanine molecules adsorbed on superconducting Pb(111), *Phys. Rev. B* **87**, 075125 (2013), [arXiv:1208.3211](https://arxiv.org/abs/1208.3211) [cond-mat.str-el].
- [108] G. Kiršanskas, M. Goldstein, K. Flensberg, L. I. Glazman, and J. Paaske, Yu-Shiba-Rusinov states in phase-biased superconductor–quantum dot–superconductor junctions, *Phys. Rev. B* **92**, 235422 (2015), [arXiv:1509.02770](https://arxiv.org/abs/1509.02770) [cond-mat.mes-hall].
- [109] E. J. H. Lee, X. Jiang, R. Žitko, R. Aguado, C. M. Lieber, and S. De Franceschi, Scaling of subgap excitations in a superconductor–semiconductor nanowire quantum dot, *Phys. Rev. B* **95**, 180502 (2017), [arXiv:1609.07582](https://arxiv.org/abs/1609.07582) [cond-mat.mes-hall].
- [110] C. P. Moca, I. Weymann, M. A. Werner, and G. Zaránd, Kondo cloud in a superconductor, *Phys. Rev. Lett.* **127**, 186804 (2021), [arXiv:2103.05286](https://arxiv.org/abs/2103.05286) [cond-mat.str-el].
- [111] M. B. Maple, W. A. Fertig, A. C. Mota, L. E. DeLong, D. Wohlleben, and R. Fitzgerald, The re-entrant superconducting–normal phase boundary of the Kondo system (La,Ce) Al₂, *Solid State Commun.* **11**, 829 (1972).
- [112] A. Yazdani, B. A. Jones, C. P. Lutz, M. F. Crommie, and D. M. Eigler, Probing the local effects of magnetic impurities on superconductivity, *Science* **275**, 1767 (1997).
- [113] K. J. Franke, G. Schulze, and J. I. Pascual, Competition of superconducting phenomena and Kondo screening at the nanoscale, *Science* **332**, 940 (2011).
- [114] N. Hatter, B. W. Heinrich, M. Ruby, J. I. Pascual, and K. J. Franke, Magnetic anisotropy in Shiba bound states across a quantum phase transition, *Nat Commun.* **6**, 1 (2015).
- [115] A. V. Balatsky, I. Vekhter, and J.-X. Zhu, Impurity-induced states in conventional and unconventional superconductors, *Rev. Mod. Phys.* **78**, 373 (2006), [arXiv:cond-mat/0411318](https://arxiv.org/abs/cond-mat/0411318) [cond-mat.supr-con].
- [116] P. Coleman, Heavy fermions: Electrons at the edge of magnetism, *Handbook of Magnetism and Advanced Magnetic Materials* (2006), [arXiv:cond-mat/0612006](https://arxiv.org/abs/cond-mat/0612006) [cond-mat.str-el].
- [117] C. Pfleiderer, Superconducting phases of *f*-electron compounds, *Rev. Mod. Phys.* **81**, 1551 (2009), [arXiv:0905.2625](https://arxiv.org/abs/0905.2625) [cond-mat.supr-con].
- [118] G. R. Stewart, Unconventional superconductivity, *Adv. in Phys.* **66**, 75 (2017), [arXiv:1705.05593](https://arxiv.org/abs/1705.05593) [cond-mat.supr-con].
- [119] T. Hatsuda, M. Tachibana, N. Yamamoto, and G. Baym, New Critical Point Induced By the Axial Anomaly in Dense QCD, *Phys. Rev. Lett.* **97**, 122001 (2006), [arXiv:hep-ph/0605018](https://arxiv.org/abs/hep-ph/0605018).
- [120] N. Yamamoto, M. Tachibana, T. Hatsuda, and G. Baym, Phase structure, collective modes, and the axial anomaly in dense QCD, *Phys. Rev. D* **76**, 074001 (2007), [arXiv:0704.2654](https://arxiv.org/abs/0704.2654) [hep-ph].
- [121] N. Read and D. M. Newns, On the solution of the Coqblin-Schrieffer Hamiltonian by the large-N expansion technique, *J. Phys. C* **16**, 3273 (1983).
- [122] P. Coleman, $\frac{1}{N}$ expansion for the Kondo lattice, *Phys. Rev. B* **28**, 5255 (1983), [Erratum: *Phys. Rev. B* **29**, 2829(E) (1984)].
- [123] E. Eichten and B. R. Hill, An effective field theory for the calculation of matrix elements involving heavy quarks, *Phys. Lett. B* **234**, 511 (1990).
- [124] H. Georgi, An effective field theory for heavy quarks at low energies, *Phys. Lett. B* **240**, 447 (1990).
- [125] M. Neubert, Heavy-quark symmetry, *Phys. Rep.* **245**, 259 (1994), [arXiv:hep-ph/9306320](https://arxiv.org/abs/hep-ph/9306320) [hep-ph].
- [126] A. V. Manohar and M. B. Wise, *Heavy Quark Physics* (Cambridge University Press, Cambridge, England, 2000).
- [127] J. G. Körner and G. Thompson, The heavy mass limit in field theory and the heavy quark effective theory, *Phys. Lett. B* **264**, 185 (1991).
- [128] S. Balk, J. G. Körner, and D. Pirjol, Quark effective theory at large orders in $1/m$, *Nucl. Phys. B* **428**, 499 (1994), [arXiv:hep-ph/9307230](https://arxiv.org/abs/hep-ph/9307230) [hep-ph].
- [129] J. I. Kapusta and C. Gale, *Finite-temperature field theory: Principles and applications*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 2011).
- [130] K. Fukushima and J. M. Pawłowski, Magnetic catalysis in hot and dense quark matter and quantum fluctua-

- tions, *Phys. Rev. D* **86**, 076013 (2012), [arXiv:1203.4330 \[hep-ph\]](#).
- [131] R. Bistritzer and A. H. MacDonald, Moiré bands in twisted double-layer graphene, *Proc. Natl. Acad. Sci. U.S.A.* **108**, 12233 (2011), [arXiv:1009.4203 \[cond-mat.mes-hall\]](#).
- [132] Z.-D. Song and B. A. Bernevig, Magic-Angle Twisted Bilayer Graphene as a Topological Heavy Fermion Problem, *Phys. Rev. Lett.* **129**, 047601 (2022), [arXiv:2111.05865 \[cond-mat.str-el\]](#).
- [133] H. Shi and X. Dai, Heavy-fermion representation for twisted bilayer graphene systems, *Phys. Rev. B* **106**, 245129 (2022), [arXiv:2209.09515 \[cond-mat.str-el\]](#).
- [134] Y.-Z. Chou and S. D. Sarma, Kondo Lattice Model in Magic-Angle Twisted Bilayer Graphene, *Phys. Rev. Lett.* **131**, 026501 (2023), [arXiv:2211.15682 \[cond-mat.str-el\]](#).
- [135] H. Hu, B. A. Bernevig, and A. M. Tsvelik, Kondo Lattice Model of Magic-Angle Twisted-Bilayer Graphene: Hund's Rule, Local-Moment Fluctuations, and Low-Energy Effective Theory, *Phys. Rev. Lett.* **131**, 026502 (2023), [arXiv:2301.04669 \[cond-mat.str-el\]](#).
- [136] S. Chen, M. He, Y.-H. Zhang, V. Hsieh, Z. Fei, K. Watanabe, T. Taniguchi, D. H. Cobden, X. Xu, C. R. Dean, *et al.*, Electrically tunable correlated and topological states in twisted monolayer–bilayer graphene, *Nature Physics* **17**, 374 (2021).
- [137] L. Yu, Bound state in superconductors with paramagnetic impurities, *Acta Phys. Sin.* **21**, 75 (1965).
- [138] H. Shiba, Classical spins in superconductors, *Prog. Theor. Phys.* **40**, 435 (1968).
- [139] A. I. Rusinov, Superconductivity near a paramagnetic impurity, *JETP Lett.* **9**, 146 (1969).
- [140] A. I. Rusinov, Theory of gapless superconductivity in alloys containing paramagnetic impurities, *Sov. Phys. JETP* **29**, 1101 (1969), [*Zh. Eksp. Teor. Fiz.* **56**, 2047 (1969)].
- [141] J. A. Hertz, Quantum critical phenomena, *Phys. Rev. B* **14**, 1165 (1976).
- [142] S. Doniach, The Kondo lattice and weak antiferromagnetism, *Physica B+ C* **91**, 231 (1977).
- [143] A. J. Millis, Effect of a nonzero temperature on quantum critical points in itinerant fermion systems, *Phys. Rev. B* **48**, 7183 (1993).
- [144] M. Vojta, Quantum phase transitions, *Rep. Prog. Phys.* **66**, 2069 (2003), [arXiv:cond-mat/0309604](#).
- [145] H. v. Löhneysen, A. Rosch, M. Vojta, and P. Wölfle, Fermi-liquid instabilities at magnetic quantum phase transitions, *Rev. Mod. Phys.* **79**, 1015 (2007), [arXiv:cond-mat/0606317 \[cond-mat.str-el\]](#).
- [146] P. Gegenwart, Q. Si, and F. Steglich, Quantum criticality in heavy-fermion metals, *Nat. Phys.* **4**, 186 (2008), [arXiv:0712.2045 \[cond-mat.str-el\]](#).
- [147] Q. Si and F. Steglich, Heavy fermions and quantum phase transitions, *Science* **329**, 1161 (2010), [arXiv:1102.4896 \[cond-mat.str-el\]](#).
- [148] P. Nozières and A. Blandin, Kondo effect in real metals, *J. Phys. France* **41**, 193 (1980).
- [149] I. Affleck, Conformal field theory approach to the Kondo effect, *Bosonization and conformal field theories in high-energy and condensed matter physics. Proceedings, 35th Cracow School of Theoretical Physics, Zakopane, Poland, June 4-14, 1995*, *Acta Phys. Polon.* **B26**, 1869 (1995), [arXiv:cond-mat/9512099 \[cond-mat\]](#).
- [150] K. Fukushima, K. Hattori, H.-U. Yee, and Y. Yin, Heavy Quark Diffusion in Strong Magnetic Fields at Weak Coupling and Implications for Elliptic Flow, *Phys. Rev. D* **93**, 074028 (2016), [arXiv:1512.03689 \[hep-ph\]](#).
- [151] G. S. Bali, F. Bruckmann, G. Endrődi, Z. Fodor, S. D. Katz, S. Krieg, A. Schafer, and K. K. Szabó, The QCD phase diagram for external magnetic fields, *JHEP* **02**, 044, [arXiv:1111.4956 \[hep-lat\]](#).
- [152] G. S. Bali, F. Bruckmann, G. Endrődi, Z. Fodor, S. D. Katz, and A. Schafer, QCD quark condensate in external magnetic fields, *Phys. Rev. D* **86**, 071502 (2012), [arXiv:1206.4205 \[hep-lat\]](#).
- [153] M. D'Elia, F. Manigrasso, F. Negro, and F. Sanfilippo, QCD phase diagram in a magnetic background for different values of the pion mass, *Phys. Rev. D* **98**, 054509 (2018), [arXiv:1808.07008 \[hep-lat\]](#).
- [154] G. Endrődi, M. Giordano, S. D. Katz, T. G. Kovács, and F. Pittler, Magnetic catalysis and inverse catalysis for heavy pions, *JHEP* **07**, 007, [arXiv:1904.10296 \[hep-lat\]](#).
- [155] H. T. Ding, S. T. Li, A. Tomiya, X. D. Wang, and Y. Zhang, Chiral properties of (2+1)-flavor QCD in strong magnetic fields at zero temperature, *Phys. Rev. D* **104**, 014505 (2021), [arXiv:2008.00493 \[hep-lat\]](#).
- [156] M. D'Elia, L. Maio, F. Sanfilippo, and A. Stanzione, Phase diagram of QCD in a magnetic background, *Phys. Rev. D* **105**, 034511 (2022), [arXiv:2111.11237 \[hep-lat\]](#).
- [157] E. Fradkin, *Field theories of condensed matter physics* (Cambridge University Press, 2013).
- [158] D. Ebert, T. Feldmann, R. Friedrich, and H. Reinhardt, Effective meson lagrangian with chiral and heavy quark symmetries from quark flavor dynamics, *Nucl. Phys. B* **434**, 619 (1995), [arXiv:hep-ph/9406220](#).
- [159] M. Buballa, NJL model analysis of quark matter at large density, *Phys. Rept.* **407**, 205 (2005), [arXiv:hep-ph/0402234](#).