

QED Fermions in a noisy magnetic field background

Jorge David Castaño-Yepes,^{1,*} Marcelo Loewe,^{1,2,3,4,†} Enrique Muñoz,^{1,5,‡} Juan Cristóbal Rojas,^{6,§} and Renato Zamora^{7,8,¶}

¹*Facultad de Física, Pontificia Universidad Católica de Chile, Vicuña Mackenna 4860, Santiago, Chile*

²*Centre for Theoretical and Mathematical Physics, and Department of Physics,
University of Cape Town, Rondebosch 7700, South Africa*

³*Centro Científico Tecnológico de Valparaíso CCTVAL,
Universidad Técnica Federico Santa María, Casilla 110-V, Valparaíso, Chile*

⁴*Facultad de Ingeniería, Arquitectura y Diseño, Universidad San Sebastián, Santiago, Chile*

⁵*Center for Nanotechnology and Advanced Materials CIEN-UC, Avenida Vicuña Mackenna 4860, Santiago, Chile*

⁶*Departamento de Física, Universidad Católica del Norte, Angamos 610, Antofagasta, Chile*

⁷*Centro de Investigación y Desarrollo en Ciencias Aeroespaciales (CIDCA),
Fuerza Aérea de Chile, Casilla 8020744, Santiago, Chile.*

⁸*Instituto de Ciencias Básicas, Universidad Diego Portales, Casilla 298-V, Santiago, Chile.*

(Dated: December 1, 2022)

We consider the effects of a noisy magnetic field background over the fermion propagator in QED, as an approximation to the spatial inhomogeneities that would naturally arise in certain physical scenarios, such as heavy-ion collisions or the quark-gluon plasma in the early stages of the evolution of the Universe. We considered a classical, finite and uniform average magnetic field background $\langle \mathbf{B}(\mathbf{x}) \rangle = \mathbf{B}$, subject to white-noise spatial fluctuations with auto-correlation of magnitude Δ_B . By means of the Schwinger representation of the propagator in the average magnetic field as a reference system, we used the replica formalism to study the effects of the magnetic noise in the form of renormalized quasi-particle parameters, leading to an effective charge and an effective refraction index, that depend not only on the energy scale, as usual, but also on the magnitude of the noise Δ_B and the average field \mathbf{B} .

I. INTRODUCTION

High-energy physics under the presence of strong magnetic fields is an important subject of research in many scenarios, such as heavy-ion collisions [1–5], the quark-gluon plasma [6–9] and the early-universe evolution [3, 10]. In such systems, rather strong magnetic fields can emerge in comparatively small regions of space and, moreover, strong spatial anisotropies and fluctuations can develop in the magnitude of such fields [1, 3].

Remarkably, magnetic fields can influence the physical properties of both charged as well as neutral particles, the later due to the quantum mechanical fluctuations of the vacuum, that lead to the creation of virtual charged fermion-antifermion pairs. In the context of high-energy physics, the effect of a constant and “classical” magnetic field background has been studied since the seminal work of Schwinger [11], followed with extensive discussions in the literature in the context of semi-classical effective Lagrangians [12, 13]. More recently, the effect of magnetic fields on nucleon parameters have been discussed in the context of QCD [14]. In addition, several studies have been reported concerning the effects of a classical, static and uniform background magnetic field on the charged vacuum fluctuations leading to the gluon polarization tensor [7, 8, 15, 16], in particular the role of the field in the breaking of the Lorentz invariance, thus predicting the

emergence of the vacuum birefringence phenomena [15, 16]. On the other hand, vacuum fluctuations also affect the propagation of fermions in such magnetized background [17], as expressed by the self-energy, that leads to the definition of a magnetic mass and, according to recent studies in QED [18], to an spectral width involving the contribution of all the magnetic Landau levels. Moreover, non-perturbative theoretical approaches [19] reveal the magnetic catalysis effect, where the presence of a strong uniform magnetic field leads to the emergence of effective masses for the fermion species, regardless of their bare mass.

Interestingly, in most of these studies, the background magnetic field is always idealized as static and uniform, and hence the presence of spatial anisotropies or fluctuations in its magnitude are disregarded in the state of the art of such calculations. A non-uniform but deterministic background magnetic field has been studied in QED by means of a path-integral formulation [20]. On the other hand, statistical fluctuations near a zero average magnetic field have been studied in the context of QED₂₊₁ [21, 22], as it arises as an effective continuum theory for certain low-dimensional Dirac materials such as Graphene [19]. In the later, however, the average background field is assumed to be zero, and hence the reference system is characterized by a free fermion propagator rather than by a Schwinger propagator as we consider in the present study. Since spatial fluctuations with respect to a *finite* background magnetic field may indeed exist in the different aforementioned physical scenarios [3], in the present work we shall study their effect over the renormalization of the fermion propagator itself in QED. As we shall discuss in this article, a perturbative treatment of such fluctuations in the framework of the replica method [23, 24] allows us to show that their effect can be captured in terms of a renormaliza-

* jcastano@uc.cl

† mloewe@fis.puc.cl

‡ munozt@fis.puc.cl

§ jurojas@ucn.cl

¶ rzamorajofre@gmail.com

tion of the charge $e \rightarrow z_3 e$ and an effective refraction index $v'/c = z^{-1}$. Moreover, we show that z_3 and z depend not only on the energy scale, as usual, but also on the magnitude of the average magnetic field $|\mathbf{B}|$, as well as on the strength of its spatial fluctuations, that we define as Δ_B .

II. THE MODEL

We shall consider a physical scenario where a classical and static magnetic field background, possessing random spatial fluctuations, modifies the quantum dynamics of a system of fermions. For this purpose, we shall assume the standard QED theory involving fermionic fields $\psi(x)$, as well as gauge fields $A^\mu(x)$. In the later, we shall distinguish three physically different contributions

$$A^\mu(x) \rightarrow A^\mu(x) + A_{\text{BG}}^\mu(x) + \delta A_{\text{BG}}^\mu(\mathbf{x}). \quad (1)$$

Here, $A^\mu(x)$ represents the dynamical photonic quantum field, while BG stands for “background”, representing the presence of a classical external field imposed by the experimental conditions. Moreover, for this BG contribution, we consider the effect of static (quenched) white noise spatial fluctuations $\delta A_{\text{BG}}^\mu(\mathbf{x})$ with respect to the mean value $A_{\text{BG}}^\mu(x)$, satisfying the statistical properties

$$\begin{aligned} \langle \delta A_{\text{BG}}^j(\mathbf{x}) \delta A_{\text{BG}}^k(\mathbf{x}') \rangle &= \Delta_B \delta_{j,k} \delta^3(\mathbf{x} - \mathbf{x}'), \\ \langle \delta A_{\text{BG}}^\mu(\mathbf{x}) \rangle &= 0. \end{aligned} \quad (2)$$

These statistical properties are represented by a Gaussian functional distribution of the form

$$dP[\delta A_{\text{BG}}^\mu] = \mathcal{N} e^{-\int d^3x \frac{[\delta A_{\text{BG}}^\mu(\mathbf{x})]^2}{2\Delta_B}} \mathcal{D}[\delta A_{\text{BG}}^\mu(\mathbf{x})]. \quad (3)$$

Therefore, we write the Lagrangian for this model as a superposition of two terms

$$\mathcal{L} = \mathcal{L}_{\text{FBG}} + \mathcal{L}_{\text{NBG}}, \quad (4)$$

where the first represents the system of Fermions (and photons) immersed in the deterministic background field (FBG)

$$\mathcal{L}_{\text{FBG}} = \bar{\psi} (i\not{\partial} - e\not{A}_{\text{BG}} - e\not{A} - m) \psi - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}, \quad (5)$$

while the second term represents the interaction between the Fermions and the classical noise (NBG)

$$\mathcal{L}_{\text{NBG}} = \bar{\psi} (-e\delta\not{A}_{\text{BG}}) \psi. \quad (6)$$

The generating functional (in the absence of sources) for a given realization of the noisy fields is given by

$$Z[A] = \int \mathcal{D}[\bar{\psi}, \psi] e^{i \int d^4x [\mathcal{L}_{\text{FBG}} + \mathcal{L}_{\text{NBG}}]}. \quad (7)$$

To study the physics of this system, we need to calculate the statistical average over the magnetic background noise δA_{BG}^μ of the $\ln Z$. For this purpose, we apply the replica method, which is based on the following identity [23]

$$\overline{\ln Z[A]} = \lim_{n \rightarrow 0} \frac{\overline{Z^n[A]} - 1}{n}. \quad (8)$$

Here, we defined the statistical average according to the Gaussian functional measure of Eq. (3), and Z^n is obtained by incorporating an additional “replica” component for each of the Fermion fields, i.e. $\psi(x) \rightarrow \psi^a(x)$, for $1 \leq a \leq n$. The “replicated” Lagrangian has the same form as Eqs. (5) and (6), but with an additional sum over the replica components of the Fermion fields. Therefore, the averaging procedure leads to

$$\begin{aligned} \overline{Z^n[A]} &= \int \prod_{a=1}^n \mathcal{D}[\bar{\psi}^a, \psi^a] \int \mathcal{D}[\delta A_{\text{BG}}^\mu] e^{-\int d^3x \frac{[\delta A_{\text{BG}}^\mu(\mathbf{x})]^2}{2\Delta_B}} \\ &\quad \times e^{i \int d^4x \sum_{a=1}^n (\mathcal{L}_{\text{FBG}}[\bar{\psi}^a, \psi^a] + \mathcal{L}_{\text{NBG}}[\bar{\psi}^a, \psi^a])} \\ &= \int \prod_{a=1}^n \mathcal{D}[\bar{\psi}^a, \psi^a] e^{i \bar{S}[\bar{\psi}^a, \psi^a; A]}, \end{aligned} \quad (9)$$

where in the last step we explicitly performed the Gaussian integral over the background noise, leading to the definition of the effective averaged action for the replica system

$$\begin{aligned} \bar{S}[\bar{\psi}^a, \psi^a; A] &= \int d^4x \left(\sum_a \bar{\psi}^a (i\not{\partial} - e\not{A}_{\text{BG}} - e\not{A} - m) \psi^a - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \right) \\ &\quad + i \frac{e^2 \Delta_B}{2} \int d^4x \int d^4y \sum_{a,b} \sum_{j=1}^3 \bar{\psi}^a(x) \gamma^j \psi^a(x) \bar{\psi}^b(y) \gamma_j \psi^b(y) \delta^3(\mathbf{x} - \mathbf{y}). \end{aligned} \quad (10)$$

Clearly, we end up with an effective interacting theory, with an instantaneous local interaction proportional to the fluctuation amplitude Δ_B that characterizes the magnetic noise, as defined in Eq. (2). The “free” part of the action corresponds to Fermions in the average background classical field $A_{\text{BG}}^\mu(x)$. We choose this background to represent a uniform, static magnetic field along the z -direction $\mathbf{B} = \hat{e}_3 B$, using the

gauge [12]

$$A_{\text{BG}}^\mu(x) = \frac{1}{2}(0, -Bx^2, Bx^1, 0). \quad (11)$$

Therefore, this allows us to use directly the Schwinger proper-time representation of the free-Fermion propagator dressed by

the background field, as follows [11, 12]

$$[S_F(k)]_{a,b} = -i\delta_{a,b} \int_0^\infty \frac{d\tau}{\cos(eB\tau)} e^{i\tau(k_\parallel^2 - \mathbf{k}_\perp^2 \frac{\tan(eB\tau)}{eB\tau} - m^2 + i\epsilon)} \times \left\{ [\cos(eB\tau) + i\gamma^1\gamma^2 \sin(eB\tau)] (m + \not{k}_\parallel) + \frac{\not{k}_\perp}{\cos(eB\tau)} \right\}, \quad (12)$$

which is clearly diagonal in replica space. Here, as usual, we separated the parallel from the perpendicular directions with respect to the background external magnetic field by splitting the metric tensor as $g^{\mu\nu} = g_\parallel^{\mu\nu} + g_\perp^{\mu\nu}$, with

$$\begin{aligned} g_\parallel^{\mu\nu} &= \text{diag}(1, 0, 0, -1), \\ g_\perp^{\mu\nu} &= \text{diag}(0, -1, -1, 0), \end{aligned} \quad (13)$$

thus implying that for any 4-vector, such as the momentum k^μ , we write

$$\not{k} = \not{k}_\perp + \not{k}_\parallel, \quad (14)$$

and

$$k^2 = k_\parallel^2 - \mathbf{k}_\perp^2. \quad (15)$$

In particular, $k_\parallel^2 = k_0^2 - k_3^2$, while $\mathbf{k}_\perp = (k^1, k^2)$ is the Euclidean 2-vector lying in the plane perpendicular to the field, such that its square-norm is $\mathbf{k}_\perp^2 = k_1^2 + k_2^2$. The Schwinger propagator can be expressed as

$$\begin{aligned} [S_F(k)]_{a,b} &= -i\delta_{a,b} [(m + \not{k}) \mathcal{A}_1 \\ &+ (ieB)i\gamma^1\gamma^2 (m + \not{k}_\parallel) \frac{\partial \mathcal{A}_1}{\partial \mathbf{k}_\perp^2} + (ieB)^2 \not{k}_\perp \frac{\partial^2 \mathcal{A}_1}{\partial (\mathbf{k}_\perp^2)^2}] \\ &= -i\delta_{a,b} \left[(m + \not{k}_\parallel) \mathcal{A}_1 + i\gamma^1\gamma^2 (m + \not{k}_\parallel) \mathcal{A}_2 + \mathcal{A}_3 \not{k}_\perp \right] \end{aligned} \quad (16)$$

Here, we defined the function

$$\mathcal{A}_1(k, B) = \int_0^\infty d\tau e^{i\tau(k_\parallel^2 - m^2 + i\epsilon) - i\frac{\mathbf{k}_\perp^2}{eB} \tan(eB\tau)}, \quad (17)$$

that clearly reproduces the inverse scalar propagator (with Feynman prescription) in the zero-field limit

$$\lim_{B \rightarrow 0} \mathcal{A}_1(k, B) = \frac{i}{k^2 - m^2 + i\epsilon} \equiv \frac{i}{\mathcal{D}_0(k)}, \quad (18)$$

with

$$\mathcal{D}_0(k) = k^2 - m^2 + i\epsilon, \quad (19)$$

and its derivatives

$$\begin{aligned} \mathcal{A}_2(k, B) &\equiv \int_0^\infty d\tau \tan(eB\tau) e^{i\tau(k_\parallel^2 - t_B(\tau)\mathbf{k}_\perp^2 - m^2 + i\epsilon)} \\ &= ieB \frac{\partial \mathcal{A}_1}{\partial (\mathbf{k}_\perp^2)}, \end{aligned} \quad (20a)$$

$$\begin{aligned} \mathcal{A}_3(k, B) &\equiv \int_0^\infty \frac{d\tau}{\cos^2(eB\tau)} e^{i\tau(k_\parallel^2 - t_B(\tau)\mathbf{k}_\perp^2 - m^2 + i\epsilon)} \\ &= \mathcal{A}_1 + (ieB)^2 \frac{\partial^2 \mathcal{A}_1}{\partial (\mathbf{k}_\perp^2)^2}. \end{aligned} \quad (20b)$$

Moreover, with these definitions it is straightforward to verify that the inverse of the Schwinger propagator Eq. (16) is given by:

$$\hat{S}_F^{-1}(k) = \frac{i}{\mathcal{D}(k)} \left[(m - \not{k}_\parallel) \mathcal{A}_1 - i\gamma^1\gamma^2 (m - \not{k}_\parallel) \mathcal{A}_2 - \mathcal{A}_3 \not{k}_\perp \right], \quad (21)$$

where

$$\mathcal{D}(k) = \mathcal{A}_3^2 \mathbf{k}_\perp^2 - (\mathcal{A}_1^2 - \mathcal{A}_2^2) (k_\parallel^2 - m^2). \quad (22)$$

Then, all the relevant expressions will be given in terms of \mathcal{A}_1 .

III. PERTURBATION THEORY: SELF-ENERGY AND VERTEX CORRECTIONS

Our goal is to develop a perturbation theory in powers of Δ , where as described in Section II and particularly in Eq. (10), the effective fermion-fermion interaction arises as a result of averaging over the background magnetic noise. Starting from a free Fermion propagator, as defined by Eq. (12), we include the magnetic noise-induced interaction effects by “dressing” the propagator with a self-energy, as shown diagrammatically in the Dyson equation depicted in Fig. 1. We remark that for this theory, the skeleton diagram for the self-energy is represented in Fig. 2.




FIG. 1. Dyson equation for the “dressed” propagator (double-line), in terms of the free propagator (single-line) and the self-energy Σ .




FIG. 2. Skeleton diagram representing the self-energy for the effective interacting theory. The dashed line is the disorder-induced interaction Δ_B , while the box $\hat{\Gamma}$ represents the 4-point vertex function.

IV. SELF-ENERGY AT ORDER Δ

It is possible to express the background-noise contribution to the self-energy (where for notational simplicity, we define




FIG. 3. Self-energy diagram at first order in $\Delta = e^2 \Delta_B$.

the parameter $\Delta \equiv e^2 \Delta_B$, as depicted in the Feynman diagram in Fig. (3), by the integral expression

$$\begin{aligned} \hat{\Sigma}_\Delta(q) &= (i\Delta) \int \frac{d^3 p}{(2\pi)^3} \gamma^j \hat{S}_F(p+q; p_0=0) \gamma_j \\ &= \frac{i(i\Delta)}{(2\pi)^3} \int d^3 p \left\{ 3(\gamma^0 q_0 - m) \mathcal{A}_1(q_0, p_3; \mathbf{p}_\perp) \right. \\ &\quad \left. + i\gamma^1 \gamma^2 (m - q_0 \gamma^0) (ieB) \frac{\partial}{\partial \mathbf{p}_\perp^2} \mathcal{A}_1(q_0, p_3; \mathbf{p}_\perp) \right\}. \end{aligned} \quad (23)$$

The derivative term in the last expression can be integrated in cylindrical coordinates $d^3 p = \pi dp_3 d(\mathbf{p}_\perp^2)$, as follows

$$\begin{aligned} &\int d^3 p \frac{\partial}{\partial \mathbf{p}_\perp^2} \mathcal{A}_1(q_0, p_3; \mathbf{p}_\perp) \\ &= \pi \int_{-\infty}^{+\infty} dp_3 \int_0^\infty d(\mathbf{p}_\perp^2) \frac{\partial}{\partial \mathbf{p}_\perp^2} \mathcal{A}_1(q_0, p_3; \mathbf{p}_\perp) \\ &= -\pi \int_{-\infty}^{+\infty} dp_3 \mathcal{A}_1(q_0, p_3; \mathbf{p}_\perp = 0), \end{aligned} \quad (24)$$

where the identity $\lim_{\mathbf{p}_\perp^2 \rightarrow \infty} \mathcal{A}_1(q_0, p_3; \mathbf{p}_\perp) = 0$ was applied. Substituting this result into Eq. (24), we finally obtain the exact expression (valid at all orders in the background average magnetic field B)

$$\begin{aligned} \hat{\Sigma}_\Delta(q) &= \frac{i(i\Delta)}{(2\pi)^3} \left[3(\gamma^0 q_0 - m) \tilde{\mathcal{A}}_1(q_0) \right. \\ &\quad \left. - i\gamma^1 \gamma^2 (i\pi eB) (m - q_0 \gamma^0) \tilde{\mathcal{A}}_2(q_0) \right], \end{aligned} \quad (25)$$

where we have defined:

$$\begin{aligned} \tilde{\mathcal{A}}_1(q_0) &\equiv \int d^3 p \mathcal{A}_1(q_0, p_3; \mathbf{p}_\perp), \\ \tilde{\mathcal{A}}_2(q_0) &\equiv \int_{-\infty}^{+\infty} dp_3 \mathcal{A}_1(q_0, p_3; \mathbf{p}_\perp = 0). \end{aligned} \quad (26)$$

Inserting the first-order in Δ expression for the self-energy Eq. (25) into the Dyson equation, as depicted diagrammatically in Fig. 1, we obtain the dressed inverse propagator at first-order in Δ

$$\hat{S}_\Delta^{-1}(k) = \hat{S}_F^{-1}(k) - \hat{\Sigma}_\Delta, \quad (27)$$

so that by using Eqs. (21) and (25) we explicitly obtain:

$$\begin{aligned} \hat{S}_\Delta^{-1}(q) &= i \left[\frac{m \mathcal{A}_1(q)}{\mathcal{D}(q)} + \frac{3m(i\Delta)}{(2\pi)^3} \tilde{\mathcal{A}}_1(q_0) \right] \\ &\quad - i \left[\frac{\mathcal{A}_1(q)}{\mathcal{D}(q)} + \frac{3(i\Delta)}{(2\pi)^3} \tilde{\mathcal{A}}_1(q_0) \right] (q_0 \gamma^0) \\ &\quad - i \left[\frac{m \mathcal{A}_2(q)}{\mathcal{D}(q)} - i \frac{m\pi(i\Delta)(eB)}{(2\pi)^3} \tilde{\mathcal{A}}_2(q_0) \right] (i\gamma^1 \gamma^2) \\ &\quad + i \left[\frac{\mathcal{A}_2(q)}{\mathcal{D}(q)} - \frac{i\pi(i\Delta)(eB)}{(2\pi)^3} \tilde{\mathcal{A}}_2(q_0) \right] (i\gamma^1 \gamma^2 q_0 \gamma^0) \\ &\quad - i \frac{\mathcal{A}_1(q)}{\mathcal{D}(q)} (q_3 \gamma^3) + i \frac{\mathcal{A}_2(q)}{\mathcal{D}(q)} (i\gamma^1 \gamma^2 q_3 \gamma^3) - i \frac{\mathcal{A}_3(q)}{\mathcal{D}(q)} \not{q}_\perp. \end{aligned} \quad (28)$$

V. RENORMALIZATION OF THE PROPAGATOR

Let us define by m' , z and z_3 as the renormalization factors for the mass, the wave function and the charge, respectively. While z will emerge as a global factor in the dressed propagator, the factor z_3 will only be associated to the tensor structures involving the spin-magnetic field interaction $e\sigma_{\mu\nu} F_{BG}^{\mu\nu} = i\gamma_1 \gamma_2 eB$. Therefore, we can compare Eq. (21) with Eq. (28), in order to identify the corresponding scalar factors for each tensor structure in both expressions, thus leading to the definition of the renormalized coefficients as follows

- For $\mathbb{1}$:

$$\frac{m \mathcal{A}_1(q)}{\mathcal{D}(q)} + \frac{3m(i\Delta)}{(2\pi)^3} \tilde{\mathcal{A}}_1(q_0) \equiv z \frac{m' \mathcal{A}_1(q)}{\mathcal{D}(q)} \quad (29a)$$

- For $\gamma^1 \gamma^2 \gamma^0$:

$$\frac{\mathcal{A}_2(q)}{\mathcal{D}(q)} - \frac{i\pi(i\Delta)(eB)}{(2\pi)^3} \tilde{\mathcal{A}}_2(q_0) \equiv z \cdot z_3 \frac{\mathcal{A}_2(q)}{\mathcal{D}(q)} \quad (29b)$$

- For $\gamma^1 \gamma^2$:

$$\frac{m \mathcal{A}_2(q)}{\mathcal{D}(q)} - \frac{im\pi(i\Delta)(eB)}{(2\pi)^3} \tilde{\mathcal{A}}_2(q_0) \equiv z \cdot z_3 \frac{m' \mathcal{A}_2(q)}{\mathcal{D}(q)} \quad (29c)$$

Then, solving the system of equations we obtain

$$z = 1 + \frac{3i\Delta}{(2\pi)^3} \frac{\tilde{\mathcal{A}}_1(q_0)}{\mathcal{A}_1(q)} \mathcal{D}(q), \quad (30a)$$

$$z_3 = \frac{1 - \frac{i\pi(i\Delta)(eB)}{(2\pi)^3} \frac{\tilde{\mathcal{A}}_2(q_0)}{\mathcal{A}_2(q)} \mathcal{D}(q)}{1 + \frac{3i\Delta}{(2\pi)^3} \frac{\tilde{\mathcal{A}}_1(q_0)}{\mathcal{A}_1(q)} \mathcal{D}(q)}, \quad (30b)$$

$$m' = m, \quad (30c)$$

and

$$\frac{v'}{c} = z^{-1} = \left(1 + \frac{3i\Delta}{(2\pi)^3} \frac{\tilde{\mathcal{A}}_1(q_0)}{\mathcal{A}_1(q)} \mathcal{D}(q) \right)^{-1} \quad (30d)$$

With these definitions into the “magnetic noise-dressed” inverse propagator Eq. (28), after organizing the different tensor structures, we obtain the expression

$$\begin{aligned} S_{\Delta}^{-1}(q) &= \frac{iz}{\mathcal{D}(q)} \left[(m - q_0\gamma^0 - z^{-1}q_3\gamma^3) \mathcal{A}_1(q) \right. \\ &\quad \left. - z_3 (i\gamma^1\gamma^2) (m - q_0\gamma^0 - z^{-1}q_3\gamma^3) \mathcal{A}_2(q) \right. \\ &\quad \left. - i\mathcal{A}_3(q)z^{-1}\not{q}_{\perp} \right] \\ &= \frac{iz}{\mathcal{D}(q)} \left[(m - \tilde{q}_{\parallel}) \mathcal{A}_1(q) - z_3 (i\gamma^1\gamma^2) (m - \tilde{q}_{\parallel}) \mathcal{A}_2(q) \right. \\ &\quad \left. - i\mathcal{A}_3(q)\tilde{q}_{\perp} \right], \end{aligned} \quad (31)$$

where in the last line we defined the four-vector $\tilde{q}^{\mu} = (q^0, z^{-1}\mathbf{q})$ that incorporates the definition of the effective refraction index $v'/c = z^{-1}$ due to the random magnetic fluctuations. By comparing Eq. (31) with Eq. (21), it is clear that they possess the same tensor structure. Therefore, by means of the elementary properties of the Dirac matrices, this expression can be readily inverted to obtain the “magnetic noise-dressed” fermion propagator

$$\begin{aligned} S_{\Delta}(q) &= -iz^{-1} \frac{\mathcal{D}(q)}{\tilde{\mathcal{D}}(q)} \left[(m + \tilde{q}_{\parallel}) \mathcal{A}_1(q) \right. \\ &\quad \left. + iz_3\gamma^1\gamma^2 (m + \tilde{q}_{\parallel}) \mathcal{A}_2(q) + \mathcal{A}_3(q)\tilde{q}_{\perp} \right], \end{aligned} \quad (32)$$

where $\mathcal{D}(q)$ was defined in Eq. (22), and

$$\tilde{\mathcal{D}}(q) = \mathcal{A}_3^2 z^{-2} \mathbf{q}_{\perp}^2 - (\mathcal{A}_1^2 - \mathcal{A}_2^2) (z^{-2} q_{\parallel}^2 - m^2) \quad (33)$$

Let us now discuss the explicit magnetic field and magnetic noise dependence of the renormalized parameters defined in Eqs. (30a) and (30b), i.e z and z_3 . For this purpose, we shall distinguish three different regimes, corresponding to the very weak, the intermediate and the ultra-intense magnetic field, respectively.

A. Very weak field $eB/m^2 \ll 1$

As shown in detail in the Appendix A, for very weak fields $eB/m^2 \ll 1$ the function $\mathcal{A}_1(k, B)$ can be expanded in terms of the power series

$$\mathcal{A}_1(k, B) = \frac{i}{\mathcal{D}_{\parallel}} \left(1 + \sum_{j=1}^{\infty} \left(\frac{ieB}{\mathcal{D}_{\parallel}} \right)^j \mathcal{E}_j(x) \right), \quad (34)$$

where for notational simplicity, we defined the “parallel” inverse scalar propagator

$$\mathcal{D}_{\parallel} = k_{\parallel}^2 - m^2 + i\epsilon, \quad (35)$$

and the dimensionless variable $x = \mathbf{k}_{\perp}^2/eB$. We also defined the polynomials $\mathcal{E}_j(x)$, as those generated by the function $e^{-ix \tan v}$, i.e.

$$\mathcal{E}_j(x) = \lim_{v \rightarrow 0} \frac{\partial^j}{\partial v^j} (e^{-ix \tan v}). \quad (36)$$

For instance, the explicit analytical expressions for the first three polynomials ($j = 1, 2, 3$) are as follows

$$\begin{aligned} \mathcal{E}_1(x) &= -ix, \\ \mathcal{E}_2(x) &= -x^2, \\ \mathcal{E}_3(x) &= -2ix + ix^3. \end{aligned}$$

At the lowest order, after subtracting the divergent vacuum contribution from Eq. (34), we have after Eq. (A15) (see Appendix A for details),

$$\mathcal{A}_1(k, B) - \mathcal{A}_1(k, 0) = \frac{-2i(eB)^2 \mathbf{k}_{\perp}^2}{[k^2 - m^2 + i\epsilon]^4} + O((eB)^4). \quad (37)$$

Therefore, using this weak field expansion of the propagator, we calculate the integral (details in Appendix C)

$$\begin{aligned} \tilde{\mathcal{A}}_1(q_0) &= -2i(eB)^2 \int d^3p \frac{\mathbf{p}_{\perp}^2}{(q_0^2 - p_3^2 - \mathbf{p}_{\perp}^2 - m^2 + i\epsilon)^4} \\ &= -\frac{\pi^2}{6} \frac{(eB)^2}{(q_0^2 - m^2)^{3/2}}. \end{aligned} \quad (38)$$

In addition, at order $\mathcal{O}((eB)^2)$ we also need to evaluate the integral (details in Appendix C)

$$\begin{aligned} \tilde{\mathcal{A}}_2(q_0) &= i \int_{-\infty}^{+\infty} \frac{dp_3}{q_0^2 - (p^3)^2 - m^2 + i\epsilon} \\ &= \frac{\pi}{\sqrt{q_0^2 - m^2}}. \end{aligned} \quad (39)$$

Therefore, from Eqs. (30a), (37), (38), and (39), we can directly evaluate the renormalization parameters to obtain

$$\begin{aligned} z &= 1 + \frac{3i\Delta}{(2\pi)^3} \frac{\tilde{\mathcal{A}}_1(q_0)}{\mathcal{A}_1(q)} \mathcal{D}(q) \\ &= 1 + \frac{\Delta(eB)^4}{8\pi} \frac{\mathbf{q}_{\perp}^2}{(q^2 - m^2 + i\epsilon)^3 (q_0^2 - m^2 + i\epsilon)^{3/2}} \\ &= 1 + O((eB)^4), \end{aligned} \quad (40)$$

and similarly from Eq. (30b)

$$z_3 = 1 + O((eB)^4). \quad (41)$$

B. Intermediate field

For intermediate magnetic field intensities, we can calculate the integral \mathcal{A}_1 by means of an expansion in terms of Landau levels. For this purpose, let us consider the generating function of the Laguerre polynomials[25]

$$e^{-\frac{x}{2} \frac{1-t}{1+t}} = (1+t)e^{-x/2} \sum_{n=0}^{\infty} (-t)^n L_n^0(x), \quad (42)$$

since

$$\begin{aligned} e^{-ix \tan v} &= \exp \left[-x (1 - e^{-2iv}) / (1 + e^{-2iv}) \right] \\ &= (1 + e^{-2iv}) e^{-x} \sum_{n=0}^{\infty} (-1)^n e^{-2in v} L_n^0(2x) \end{aligned} \quad (43)$$

Therefore, we have (for $x = \mathbf{k}_\perp^2 / eB$)

$$\begin{aligned} \mathcal{A}_1(k) &= e^{-x} \left(\sum_{n=0}^{\infty} (-1)^n L_n^0(2x) \int_0^\infty d\tau e^{i(\mathcal{D}_\parallel - 2(n+1)eB)\tau} \right. \\ &\quad \left. + \sum_{n=0}^{\infty} (-1)^n L_n^0(2x) \int_0^\infty d\tau e^{i(\mathcal{D}_\parallel - 2neB)\tau} \right) \end{aligned} \quad (44)$$

Evaluating the exponential integrals, we obtain

$$\begin{aligned} \mathcal{A}_1(k) &= ie^{-x} \left(\sum_{n=0}^{\infty} (-1)^n \frac{L_n^0(2x)}{\mathcal{D}_\parallel - 2(n+1)eB} \right. \\ &\quad \left. + \sum_{n=0}^{\infty} (-1)^n \frac{L_n^0(2x)}{\mathcal{D}_\parallel - 2neB} \right) \\ &= i \frac{e^{-x}}{\mathcal{D}_\parallel} \left[1 + \sum_{n=1}^{\infty} \frac{(-1)^n [L_n^0(2x) - L_{n-1}^0(2x)]}{1 - 2n \frac{eB}{\mathcal{D}_\parallel}} \right] \end{aligned} \quad (45)$$

Inserting this expression into the definition of $\tilde{\mathcal{A}}_1$ of Eq. (26), we obtain

$$\begin{aligned} \tilde{\mathcal{A}}_1(q_0) &= \int d^3p \mathcal{A}_1(q_0, p_3; \mathbf{p}_\perp) \\ &= i \int d^3p \left[\frac{e^{-\mathbf{p}_\perp^2 / eB}}{q_0^2 - p_3^2 - m^2 + i\epsilon} \right. \\ &\quad \left. + \sum_{n=1}^{\infty} (-1)^n e^{-\mathbf{p}_\perp^2 / eB} \frac{L_n \left(\frac{2\mathbf{p}_\perp^2}{eB} \right) - L_{n-1} \left(\frac{2\mathbf{p}_\perp^2}{eB} \right)}{q_0^2 - p_3^2 - m^2 - 2neB + i\epsilon} \right] \\ &= \mathcal{I}_1 + \sum_{n=1}^{\infty} (-1)^n \mathcal{I}_{2,n} \end{aligned} \quad (46)$$

Here, we defined

$$\begin{aligned} \mathcal{I}_1 &= i \int d^3p \frac{e^{-\mathbf{p}_\perp^2 / eB}}{q_0^2 - p_3^2 - m^2 + i\epsilon} \\ &= \frac{\pi^2 eB}{\sqrt{q_0^2 - m^2 + i\epsilon}}, \end{aligned} \quad (47)$$

and

$$\mathcal{I}_{2,n} = i \int d^3p e^{-\mathbf{p}_\perp^2 / eB} \frac{L_n \left(\frac{2\mathbf{p}_\perp^2}{eB} \right) - L_{n-1} \left(\frac{2\mathbf{p}_\perp^2}{eB} \right)}{q_0^2 - p_3^2 - m^2 - 2neB + i\epsilon} \quad (48)$$

We calculate the momentum integrals in cylindrical coordinates, making use of the azimuthal symmetry, such that $d^3p = dp_3 \pi d(\mathbf{p}_\perp^2)$. Moreover, in the integral over \mathbf{p}_\perp , we




FIG. 4. Spectral density for the Landau level spectrum $\rho(E)$, as a function of the dimensionless energy scale E/m , for different values of the average background magnetic field eB/m^2 . The inset is shown in order to appreciate in detail the staircase pattern produced by the discrete Landau levels.

define the auxiliary variable $x = \frac{2\mathbf{p}_\perp^2}{eB}$, such that

$$\begin{aligned} \mathcal{I}_{2,n} &= \frac{\pi eB}{2} \int_{-\infty}^{\infty} \frac{dp_3}{q_0^2 - m^2 - p_3^2 - 2nqB + i\epsilon} \\ &\quad \times \int_0^\infty dx e^{-x/2} [L_n(x) - L_{n-1}(x)] \\ &= 2\pi eB (-1)^n \int_{-\infty}^{\infty} \frac{dp_3}{q_0^2 - m^2 - p_3^2 - 2nqB + i\epsilon} \end{aligned} \quad (49)$$

where we used the identity [26]

$$\int_0^\infty dx e^{-bx} L_n(x) = (b-1)^n b^{-n-1} \quad \text{Re } b > 0. \quad (50)$$

Inserting Eq. (47) and Eq. (49) into Eq. (46), we obtain (after shifting the index $n \rightarrow n+1$)

$$\begin{aligned} \tilde{\mathcal{A}}_1(q_0) &= \frac{\pi^2(eB)}{\sqrt{q_0^2 - m^2 + i\epsilon}} \\ &\quad + 2\pi eB \int_{-\infty}^{+\infty} dp_3 \sum_{n=0}^{\infty} \frac{1}{q_0^2 - m^2 - p_3^2 - 2(n+1)eB + i\epsilon} \end{aligned} \quad (51)$$

Let us introduce the density of states for Landau levels

$$\rho(E) = \int_{-\infty}^{\infty} \frac{dp_3}{2\pi} \sum_{n=0}^{\infty} \delta(E - E_n(p_3)), \quad (52)$$

with the dispersion relation for the spectrum

$$E_n(p_3) = \sqrt{p_3^2 + m^2 + 2(n+1)eB}. \quad (53)$$

As shown in detail in Appendix B

$$\begin{aligned} \rho(E) &= \Theta(E - \sqrt{m^2 + 2eB}) \frac{E}{\pi\sqrt{eB}} \\ &\times \left[\zeta\left(\frac{1}{2}, \frac{E^2 - m^2 - 2eB}{2eB}\right) - N_{max}(E) \right. \\ &\quad \left. - \zeta\left(\frac{1}{2}, \frac{E^2 - m^2}{2eB}\right) \right], \end{aligned} \quad (54)$$

where we defined

$$N_{max}(E) = \lfloor \frac{E^2 - m^2}{2eB} - 1 \rfloor, \quad (55)$$

with $\lfloor x \rfloor$ the integer part of x , and $\zeta(n, z)$ is the Riemann zeta function. The spectral density Eq. (54) is represented in Fig. 4, as a function of the dimensionless energy scale E/m . For large magnetic fields $eB/m^2 \gg 1$, the spectral density displays a clear staircase pattern, where each step represents the contribution arising from a single Landau level $n = 0, 1, \dots$. On the other hand, for weak magnetic fields $eB/m^2 \ll 1$, the spectral density exhibits a denser, quasi-continuum behavior.

With these definitions, we obtain from Eq. (52) the exact

expression

$$\tilde{\mathcal{A}}_1(q_0) = \frac{\pi^2 eB}{\sqrt{q_0^2 - m^2 + i\epsilon}} + 4\pi^2 eB \int_{-\infty}^{+\infty} dE \frac{\rho(E)}{q_0^2 - E^2 + i\epsilon}, \quad (56)$$

On the other hand, from Eq. (26) we have

$$\begin{aligned} \tilde{\mathcal{A}}_2(q_0) &= \int_{-\infty}^{\infty} dp_3 p \mathcal{A}_1(q_0, p_3; \mathbf{p}_\perp = 0) \\ &= i \int_{-\infty}^{\infty} dp_3 \frac{1}{q_0^2 - p_3^2 - m^2 + i\epsilon} \\ &\quad + i \int_{-\infty}^{\infty} dp_3 \sum_{n=1}^{\infty} (-1)^n \frac{L_n(0) - L_{n-1}(0)}{q_0^2 - p_3^2 - m^2 - 2neB + i\epsilon}, \end{aligned} \quad (57)$$

where the second term vanishes, given that $L_n(0) = 1 \forall n$. Hence, we end up with the simple expression

$$\tilde{\mathcal{A}}_2(q_0) = \frac{\pi}{\sqrt{q_0^2 - m^2 + i\epsilon}}. \quad (58)$$

In order to evaluate the formulas for z in Eq. (30a) and z_3 in Eq. (30b), we also need to evaluate the following coefficients (for $x = \mathbf{k}_\perp^2/(eB)$)

$$\begin{aligned} \mathcal{A}_2 &= ieB \frac{\partial \mathcal{A}_1}{\partial \mathbf{k}_\perp^2} = i \frac{\partial \mathcal{A}_1}{\partial x} \\ &= \frac{e^{-x}}{\mathcal{D}_\parallel} \left[1 + \sum_{n=1}^{\infty} \frac{(-1)^n (L_n(2x) - 2L'_n(2x) - L_{n-1}(2x) + 2L'_{n-1}(2x))}{1 - 2\frac{neB}{\mathcal{D}_\parallel}} \right] \\ \mathcal{A}_3 &= \mathcal{A}_1 + (ieB)^2 \frac{\partial^2 \mathcal{A}_1}{\partial (\mathbf{k}_\perp^2)^2} = \mathcal{A}_1 - \frac{\partial^2 \mathcal{A}_1}{\partial x^2} \\ &= i \frac{e^{-x}}{\mathcal{D}_\parallel} \sum_{n=1}^{\infty} \frac{(-1)^n (L_n(2x) - L_{n-1}(2x) - 2L'_n(2x) + 4L''_n(2x) + 2L'_{n-1}(2x) - 4L''_{n-1}(2x))}{1 - 2\frac{neB}{\mathcal{D}_\parallel}} \end{aligned} \quad (59)$$

Finally, in order to further simplify these expressions, it is convenient to use the identities

$$\begin{aligned} L'_n(2x) &= \begin{cases} -L_{n-1}^{(1)}(2x), & n \geq 1 \\ 0, & \text{otherwise} \end{cases} \\ &= -\theta_{n-1} \cdot L_{n-1}^{(1)}(2x), \end{aligned} \quad (60)$$

$$L''_n(2x) = \theta_{n-2} L_{n-2}^{(2)}(2x), \quad (61)$$

where θ_{n-k} is the Heaviside step function

$$\theta_{n-k} = \begin{cases} 1, & n \geq k \\ 0, & \text{otherwise} \end{cases} \quad (62)$$

With these identities, we obtain the final expressions

$$\begin{aligned} \mathcal{A}_2 &= \frac{e^{-x}}{\mathcal{D}_\parallel} \left[1 + \sum_{n=1}^{\infty} \frac{(-1)^n (L_n(2x) + 2L_{n-1}^{(1)}(2x) - L_{n-1}(2x) - 2\theta_{n-2} \cdot L_{n-2}^{(1)}(2x))}{1 - 2\frac{neB}{\mathcal{D}_\parallel}} \right] \\ \mathcal{A}_3 &= i \frac{e^{-x}}{\mathcal{D}_\parallel} \sum_{n=1}^{\infty} \frac{(-1)^n (L_n(2x) - L_{n-1}(2x) + 2L_{n-1}^{(1)}(2x) + 4\theta_{n-2} \cdot L_{n-2}^{(2)}(2x) - 2L_{n-1}^{(1)}(2x) - 4\theta_{n-3} \cdot L_{n-3}^{(2)}(2x))}{1 - 2\frac{neB}{\mathcal{D}_\parallel}} \end{aligned} \quad (63)$$

With these expressions, we evaluate

$$\mathcal{D}(k) = \mathcal{A}_3^2 \mathbf{k}_\perp^2 - (\mathcal{A}_1^2 - \mathcal{A}_2^2) (k_\parallel^2 - m^2), \quad (64)$$

and finally evaluate z and z_3 with Eq. (30a) and Eq. (30b), respectively. These results can be appreciated in Figs. 5–10, as a function of the energy scale q_0/m , as well as the magnitude of the average background magnetic field eB/m^2 , respectively.

C. Ultra-intense (LLL) field $eB/m^2 \gg 1$

Let us now analyze the asymptotic behaviour of the quasi-particle renormalization parameters z , z_3 , and v'/c , respectively, in the ultra-intense magnetic field regime $eB/m^2 \gg 1$. Here, we obtain the corresponding asymptotic expression for $\mathcal{A}_1(q)$ by considering only the lowest-Landau level (LLL) $n = 0$ in Eq. (46). Therefore, we have

$$\mathcal{A}_1(q) \sim i \frac{e^{-\mathbf{q}_\perp^2/eB}}{q_\parallel^2 - m^2}, \quad (65)$$

and the corresponding expressions for its derivatives are

$$\mathcal{A}_2(q) = \frac{e^{-\mathbf{q}_\perp^2/eB}}{q_\parallel^2 - m^2}, \quad (66)$$

and

$$\mathcal{A}_3(q) = 0. \quad (67)$$

Similarly, we also have

$$\mathcal{D}(q) = 2 \frac{e^{-2\mathbf{q}_\perp^2/eB}}{q_\parallel^2 - m^2}. \quad (68)$$

Finally, the integrals of $\mathcal{A}_1(q)$ are given, in this approximation, by the expressions

$$\tilde{\mathcal{A}}_1(q) = \frac{\pi^2 eB}{\sqrt{q_0^2 - m^2}}, \quad (69a)$$

$$\tilde{\mathcal{A}}_2(q_0) = \frac{\pi}{\sqrt{q_0^2 - m^2}}. \quad (69b)$$

Applying these asymptotic results for the ultra-strong field regime, and substituting into the general definitions Eq. (30a) and Eq. (30b), we obtain explicit analytical expressions for the renormalization factors z and z_3 , respectively, as follows

$$z = 1 + \frac{3}{4} \frac{\Delta(eB) e^{-\mathbf{q}_\perp^2/eB}}{\pi \sqrt{q_0^2 - m^2}}, \quad (70)$$

and

$$\begin{aligned} z_3 &= \left(1 + \frac{\Delta(eB) e^{-\mathbf{q}_\perp^2/eB}}{4\pi \sqrt{q_0^2 - m^2}} \right) z^{-1} \\ &= \frac{1 + \frac{\Delta(eB) e^{-\mathbf{q}_\perp^2/eB}}{4\pi \sqrt{q_0^2 - m^2}}}{1 + \frac{3}{4} \frac{\Delta(eB) e^{-\mathbf{q}_\perp^2/eB}}{\pi \sqrt{q_0^2 - m^2}}} \end{aligned} \quad (71)$$

Remarkably, while for very large magnetic fields $z \sim eB/m^2$ grows linearly, z_3 instead converges asymptotically to the constant limit

$$\lim_{\frac{eB}{m^2} \rightarrow \infty} z_3 = \frac{1}{3}. \quad (72)$$




FIG. 5. Wavefunction renormalization factor z as a function of the dimensionless energy scale q_0/m . Here $\mathbf{p}_\perp^2 = \mathbf{p}_\parallel^2 - m^2$.




FIG. 6. Wavefunction renormalization factor z as a function of the dimensionless magnetic field scale eB/m^2 . Here $\mathbf{p}_\perp^2 = \mathbf{p}_\parallel^2 - m^2$, and $eB/m^2 \in [10, 500]$




FIG. 7. Charge renormalization factor z_3 as a function of the dimensionless energy scale q_0/m . Here $\mathbf{p}_\perp^2 = p_\parallel^2 - m^2$.




FIG. 9. Effective refraction index v'/c as a function of the dimensionless energy scale q_0/m . Here $\mathbf{p}_\perp^2 = p_\parallel^2 - m^2$.




FIG. 8. Charge renormalization factor z_3 as a function of the dimensionless magnetic field scale eB/m^2 . Here $\mathbf{p}_\perp^2 = p_\parallel^2 - m^2$, and $eB/m^2 \in [10, 500]$




FIG. 10. Effective refraction index v'/c as a function of the dimensionless magnetic field scale eB/m^2 . Here $\mathbf{p}_\perp^2 = p_\parallel^2 - m^2$, and $eB/m^2 \in [10, 500]$

Let us now summarize the behavior of the renormalization factors z and z_3 as a function of the energy scale q_0/m , and the average background magnetic field eB/m^2 , respectively, in the whole range of both parameters, as displayed in Figs. 5–8, respectively. As can be appreciated in Fig. 5, z presents a monotonically decreasing behaviour as a function of the energy q_0/m , that asymptotically reaches the limit

$z \rightarrow 1$ as $q_0/m \gg 1$, for all values of the average background magnetic field eB/m^2 . In physical terms, this shows that the quasi-particle renormalization due to the random magnetic field fluctuations tends to be negligible as the energy of the propagating fermions becomes very large, but in contrast it can be quite significant at low energy scales. This trend is also consistent with the effective refraction index $v'/c = z^{-1}$,

as shown in Figs. 9,10. For low energy scales, $v'/c < 1$, indicating a strong renormalization of the effective group velocity of the propagating quasi-particles due to the presence of the magnetic background fluctuations. In contrast, for larger energy scales the effect becomes weaker, thus recovering the asymptotic limit $v'/c \rightarrow 1$ as $q_0/m \gg 1$. Since low energy and momentum components in the Fourier representation of the propagator correspond to long-wavelength components in the space of configurations, our results are consistent with the fact that such long-wavelength components are more sensitive to the spatial distribution of the magnetic fluctuations, and hence experience a higher degree of decoherence, thus reducing the corresponding group velocity. In contrast, the high-energy Fourier modes of the propagator, that correspond to short-wavelength components in the configuration space, are less sensitive to the presence of spatial fluctuations of the background magnetic field.

Concerning the charge renormalization factor z_3 , as can be appreciated in Fig. 7 it experiences a strong effect $z_3 < 1$ at low quasi-particle energies q_0/m , but this effect becomes negligible at large energy scales $q_0/m \gg 1$, since $z_3 \rightarrow 1$ as an asymptotic limit. This behavior, that can be interpreted physically as a charge screening due to the spatial magnetic fluctuations in the background, is consistent with the aforementioned interpretation for the effective index of refraction as a function of energy. On the other hand, as can be appreciated in Fig. 8, z_3 tends to decrease as a function of the average background magnetic field intensity, achieving an asymptotic limit $z_3 \rightarrow 1/3$ as shown in Eq. (72).

VI. VERTEX CORRECTIONS AT $O(\Delta^2)$

Let us now consider the renormalization of the effective interaction term $\Delta \rightarrow \tilde{\Delta}$, that characterizes the strength of the effective interaction vertex in the effective, averaged action for the replica system, Eq. (10). Following the skeleton diagrams for the perturbation theory, as depicted in Fig. 2, the diagrams contributing at order Δ^2 to the 4-point vertex $\hat{\Gamma}$ are depicted in Fig. 11.

Therefore, the matrix elements corresponding to each diagram are given by the following integral expressions

$$\hat{\Gamma}_{(a)} = \int \frac{d^3q}{(2\pi)^3} \gamma^i S_F(p-q) \gamma^j \otimes \gamma_i S_F(p'-q) \gamma_j, \quad (73a)$$

$$\hat{\Gamma}_{(b)} = \int \frac{d^3q}{(2\pi)^3} \gamma^i S_F(p-q) \gamma^j \otimes \gamma_i S_F(p'+q) \gamma_j, \quad (73b)$$

and

$$\hat{\Gamma}_{(c)} = \int \frac{d^3q}{(2\pi)^3} \gamma^i S_F(p+q) \gamma^j \otimes \gamma_i S_F(p'-q) \gamma_j. \quad (73c)$$

In order to compute the expressions above, it is convenient to introduce the notation

$$\hat{\Gamma}^{(\lambda,\sigma)} = \int \frac{d^3q}{(2\pi)^3} \gamma^i S_F(p+\lambda q) \gamma^j \otimes \gamma_i S_F(p'+\sigma q) \gamma_j, \quad (74)$$




FIG. 11. Diagrams contributing to the 4-point vertex function $\hat{\Gamma}$ at order Δ^2 .

where $\lambda, \sigma = \pm 1$. Then, we have the correspondence $\hat{\Gamma}_{(a)} = \hat{\Gamma}^{(-,-)}$, $\hat{\Gamma}_{(b)} = \hat{\Gamma}^{(-,+)}$, and $\hat{\Gamma}_{(c)} = \hat{\Gamma}^{(+,-)}$, respectively.

By considering the tensor structure of the propagator, it is straightforward to realize that the full vertex, taking into account the multiplicity and symmetry factors for each diagram, is given by

$$\hat{\Gamma} = 2\hat{\Gamma}^{(-,-)} + 2\hat{\Gamma}^{(-,+)} + 4\hat{\Gamma}^{(+,-)}. \quad (75)$$

The former leads to an effective interaction of the form

$$\hat{\Gamma} = \tilde{\Delta}(\bar{\psi}\gamma^i\psi)(\bar{\psi}\gamma^i\psi) + \text{other tensor structures}, \quad (76)$$

where the renormalized coefficient $\tilde{\Delta}$ is given, up to second order in Δ (see Appendix D for details) by the expression

$$\begin{aligned} \tilde{\Delta} = & \Delta + 2\Delta^2 \left(\mathcal{J}_2^{(-,-)} + \mathcal{J}_2^{(-,+)} + 2\mathcal{J}_2^{(+,-)} \right) \\ & + (1 - \partial_x^2)(1 - \partial_y^2)\mathcal{J}_3^{(-,-)} + (1 - \partial_x^2)(1 - \partial_y^2)\mathcal{J}_3^{(-,+)} \\ & + 2(1 - \partial_x^2)(1 - \partial_y^2)\mathcal{J}_3^{(+,-)}. \end{aligned} \quad (77)$$

Here, we defined the integrals

$$\mathcal{J}_1^{(\lambda,\sigma)}(p,p') \equiv \int \frac{d^3q}{(2\pi)^3} \mathcal{A}_1(p+\lambda q) \mathcal{A}_1(p'+\sigma q), \quad (78a)$$

$$\mathcal{J}_2^{(\lambda,\sigma)}(p,p') \equiv \int \frac{d^3q}{(2\pi)^3} q_{\parallel}^2 \mathcal{A}_1(p+\lambda q) \mathcal{A}_1(p'+\sigma q), \quad (78b)$$

and

$$\mathcal{J}_3^{(\lambda,\sigma)}(p,p') \equiv \int \frac{d^3q}{(2\pi)^3} \mathbf{q}_{\perp}^2 \mathcal{A}_1(p+\lambda q) \mathcal{A}_1(p'+\sigma q). \quad (78c)$$




FIG. 12. real and imaginary parts of the correction coefficient from Eq. (77) as a function of the fermion's energy. Here, we take $Q = 0$ and $\mathbf{p}_\perp^2 = p_0^2 - m^2$.

In order to calculate the integrals \mathcal{J}_i , we shall use the analytical expression for $\mathcal{A}_1(k)$ Eq. (A24) (details in Appendix A):

$$\mathcal{A}_1(k) = \frac{i e^{-\mathbf{k}_\perp^2/eB}}{2eB} \exp\left[-\frac{i\pi(k_\parallel^2 - m^2)}{2eB}\right] \times \Gamma\left(-\frac{k_\parallel^2 - m^2}{2eB}\right) U\left(-\frac{k_\parallel^2 - m^2}{2eB}, 0, \frac{2\mathbf{k}_\perp^2}{eB}\right). \quad (79)$$

A. The integral \mathcal{J}_1

Let us first consider the integral

$$\mathcal{J}_1^{(\lambda, \sigma)}(p, p') = \int \frac{d^3q}{(2\pi)^3} \mathcal{A}_1(p + \lambda q) \mathcal{A}_1(p' + \sigma q). \quad (80)$$

For the case $(\lambda, \sigma) = (-1, -1)$ we change the integration variables as follows

$$\begin{aligned} p' - q &= q' + Q, \\ p - q &= q' - Q. \end{aligned} \quad (81)$$

For notational simplicity, in what follows we shall use q

instead of q' , and we shall define the parameters

$$\begin{aligned} a &= -\frac{\mathcal{D}_\parallel(q_3 + Q_\parallel)}{2eB}, \\ a' &= -\frac{\mathcal{D}_\parallel(q_3 - Q_\parallel)}{2eB}. \end{aligned} \quad (82)$$

Furthermore, we shall use the identity (for $z = 2\mathbf{q}_\perp^2/eB$)

$$\Gamma(a)U(a, \epsilon, z) = \frac{1}{a}M(a, \epsilon, z) + \Gamma(-1 + \epsilon)zM(1 + a, 2, z), \quad (83)$$

along with the singular expansion for $\epsilon \rightarrow 0^+$

$$\Gamma(-1 + \epsilon) = \frac{-1}{\epsilon} + \gamma_e - 1 + O(\epsilon), \quad (84)$$

where $\gamma_e = 0.577$ is the Euler-Mascheroni constant. In addition, given that there is a strong exponential damping in the integral, we consider the expansion of the Kummer function for small values of its argument, that is given by

$$M(a, b, z) = 1 + \frac{a}{b}z + O(z^2), \quad (85)$$

so that, after regularization by removing the divergences in $1/\epsilon$, we end up with the integral

$$\begin{aligned} \mathcal{J}_1^{(-, -)}(p, p') &= \left(\frac{i}{2eB}\right)^2 e^{-\frac{2\mathbf{Q}_\perp^2}{eB}} \int_{-\infty}^{\infty} \frac{dq_3}{2\pi} e^{i\pi(a+a')} \int_0^{\infty} \frac{d^2q_\perp}{(2\pi)^2} e^{-\frac{2\mathbf{q}_\perp^2}{eB}} \Gamma(a)U\left(a, \epsilon, \frac{(q+Q)_\perp^2}{eB}\right) \Gamma(a')U\left(a', \epsilon, \frac{(q-Q)_\perp^2}{eB}\right) \\ &= \frac{1}{(2\pi)^3} \left(\frac{i}{2eB}\right)^2 e^{-\frac{2\mathbf{Q}_\perp^2}{eB}} \int_{-\infty}^{\infty} dq_3 e^{i\pi(a+a')} \\ &\quad \times \int_0^{\infty} d^2q_\perp e^{-\frac{2\mathbf{q}_\perp^2}{eB}} \left[\frac{1}{a} + \frac{\gamma_e - 1}{eB} (\mathbf{q}_\perp^2 + 2\mathbf{Q}_\perp \cdot \mathbf{q}_\perp + \mathbf{Q}_\perp^2)\right] \left[\frac{1}{a'} + \frac{\gamma_e - 1}{eB} (\mathbf{q}_\perp^2 - 2\mathbf{Q}_\perp \cdot \mathbf{q}_\perp + \mathbf{Q}_\perp^2)\right]. \end{aligned} \quad (86)$$

Furthermore, we shall set the external 3-momenta to zero,

except for the presence of the Q_\perp factors, that we shall keep as

finite in order to use this expression as a generating function. Therefore, the integral reduces to the simpler expression

$$\begin{aligned} \mathcal{J}_1^{(-,-)}(p, p') &= -\frac{eB}{4\pi^2} e^{-\frac{2\mathbf{Q}_\perp^2}{eB}} \\ &\times \int_{-\infty}^{\infty} \frac{dq_3}{(q_3^2 + m^2 + i\epsilon)^2} \exp\left[\frac{i\pi}{2eB} (q_3^2 + m^2)\right] \\ &\times \left[1 + \frac{(\gamma_e - 1)\mathbf{Q}_\perp^2 (q_3^2 + m^2)}{(eB)^2} + \frac{(\gamma_e - 1)(q_3^2 + m^2)}{eB}\right] \end{aligned} \quad (87)$$

Performing the last integral explicitly, we obtain (details in Appendix D)

$$\begin{aligned} \mathcal{J}_1^{(-,-)}(p, p') &= -\frac{eB}{4\pi^2} e^{-\frac{2\mathbf{Q}_\perp^2}{eB}} \left\{ \frac{(1-i)\pi}{2\sqrt{eB}m^2} e^{\frac{i\pi m^2}{2eB}} \right. \\ &+ \frac{(eB + i\pi m^2)\pi}{2(eB)m^3} \left[1 - \sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\frac{1-i}{\sqrt{2}} \frac{m}{\sqrt{eB}}\right)\right] \\ &\left. + \frac{\pi(\gamma_e - 1)}{(eB)m} \left(1 + \frac{\mathbf{Q}_\perp^2}{eB}\right) \left[1 - \sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\frac{1-i}{\sqrt{2}} \frac{m}{\sqrt{eB}}\right)\right] \right\}, \end{aligned} \quad (88)$$

where $\operatorname{erf}(x)$ is the error function.

B. The integral \mathcal{J}_2

For this second integral, we notice that $q_\parallel^2 = -q_3^2$, so that after integration over $z = 2\mathbf{Q}_\perp^2/(eB)$ we obtain (details in Appendix D)

$$\begin{aligned} \mathcal{J}_2^{(-,-)}(p, p') &= \frac{eB}{4\pi^2} e^{-\frac{2\mathbf{Q}_\perp^2}{eB}} \\ &\times \int_{-\infty}^{\infty} dq_3 \frac{q_3^2}{(q_3^2 + m^2 + i\epsilon)^2} \exp\left[\frac{i\pi}{2eB} (q_3^2 + m^2)\right] \\ &\times \left[1 + \frac{(\gamma_e - 1)\mathbf{Q}_\perp^2 (q_3^2 + m^2)}{(eB)^2} + \frac{(\gamma_e - 1)(q_3^2 + m^2)}{eB}\right] \end{aligned} \quad (89)$$

After performing the remaining momentum integral over q_3 , as shown in detail in Appendix D, we finally obtain

$$\begin{aligned} \mathcal{J}_2^{(-,-)}(p, p') &= \frac{eB}{4\pi^2} e^{-\frac{2\mathbf{Q}_\perp^2}{eB}} \\ &\times \left\{ \frac{(i-1)\pi}{\sqrt{eB}} e^{\frac{i\pi m^2}{2eB}} \right. \\ &+ \frac{(eB - i\pi m^2)\pi}{2(eB)m} \left[1 - \sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\frac{1-i}{\sqrt{2}} \frac{m}{\sqrt{eB}}\right)\right] \\ &\left. - \frac{m\pi(\gamma_e - 1)}{(eB)} \left(1 + \frac{\mathbf{Q}_\perp^2}{eB}\right) \left[1 - \sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\frac{1-i}{\sqrt{2}} \frac{m}{\sqrt{eB}}\right)\right] \right\}. \end{aligned}$$

C. The integral \mathcal{J}_3

By following the same procedure as in the previous two cases, as shown in Appendix D, it is straightforward to obtain the analytical expression

$$\begin{aligned} \mathcal{J}_3^{(-,-)}(p, p') &= -\frac{eB^2}{8\pi^2} e^{-\frac{2\mathbf{Q}_\perp^2}{eB}} \left\{ \frac{(1-i)\pi}{2\sqrt{eB}m^2} e^{\frac{i\pi m^2}{2eB}} \right. \\ &+ \frac{(eB + i\pi m^2)\pi}{2(eB)m^3} \left[1 - \sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\frac{1-i}{\sqrt{2}} \frac{m}{\sqrt{eB}}\right)\right] \\ &\left. + \frac{\pi(\gamma_e - 1)\mathbf{Q}_\perp^2}{(eB)^2 m} \left[1 - \sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\frac{1-i}{\sqrt{2}} \frac{m}{\sqrt{eB}}\right)\right] \right\}. \end{aligned} \quad (90)$$

Moreover, it is straightforward to verify the following relations

$$\begin{aligned} \mathcal{J}_n^{(+,+)}(p, p') &= \mathcal{J}_n^{(-,-)}(p, p') \\ \mathcal{J}_n^{(+,-)}(p, p') &= \mathcal{J}_n^{(-,+)}(p, p') = \mathcal{J}_n^{(-,-)}(p, p') \big|_{Q \rightarrow P}, \end{aligned} \quad (91)$$

for $n = 1, 2, 3$, that allows us to generate all the remaining expressions from these three explicit analytical results.

An explicit numerical evaluation of our analytical expressions for the renormalized effective interaction, expressed by the combination $(\hat{\Delta} - \Delta)/(2\Delta^2)$, is displayed in Fig. 12. Clearly, this effective coupling develops both a real as well as an imaginary part (left and right panels in Fig. 12, respectively). In particular, the emergence of an imaginary component implies an imaginary contribution to the self-energy, corresponding to a relaxation time (spectral broadening) of the quasi-particle spectrum. This is a natural consequence of the decoherence mechanism induced by the random fluctuating magnetic environment. On the other hand, as can be appreciated in Fig. 12, both the real and imaginary contributions to the effective interaction $\hat{\Delta}$ display a large enhancement (in absolute value) at low energy scales $p_0/m < 1$, while asymptotically $\hat{\Delta} \rightarrow \Delta$ at higher energies $p_0/m \gg 1$. This strong renormalization effect at low-energies is consistent with the effect observed in the previous section for the charge z_3 and refraction index v'/c , respectively, and can be explained in similar terms due to the short-range spatial distribution of the magnetic noise, that therefore renormalizes mainly the long-wavelength components of the propagator, corresponding to the small energy-momentum components in Fourier space.

VII. CONCLUSIONS

We have studied the effects of quenched, white noise spatial fluctuations in an otherwise uniform background magnetic field, over the properties of the QED fermion propagator. This configuration is important in different physical scenarios, including heavy-ion collisions and the quark-gluon plasma, where spatial anisotropies of the background magnetic field may be present. We developed explicit results, that

we carried over by combining the replica method to average over spatial fluctuations, with a perturbation theory based on the Schwinger propagator for the average background field. Upon averaging over magnetic fluctuations, we obtained an effective action in the replica fields, with an effective particle-particle interaction proportional to the strength Δ of the spatial auto-correlation function of the background noise. Our perturbative results show that, up to first order in Δ , the propagator retains its form, thus representing renormalized quasi-particles with the same mass $m' = m$, but propagating in the medium with a magnetic field and noise-dependent index of refraction $v'/c = z^{-1}$, and effective charge $e' = z_3 e$, where z and z_3 are renormalization factors. We showed that z presents a monotonically decreasing behaviour as a function of the energy q_0/m , that reaches the asymptotic limit $z \rightarrow 1$ as $q_0/m \gg 1$, for all values of the average background magnetic field eB/m^2 . In physical terms, this shows that the quasi-particle renormalization due to the random magnetic field fluctuations, while being quite significant at low energy scales, tends to be negligible as the energy of the propagating fermions becomes very large. This trend is also observed in the effective refraction index $v'/c = z^{-1}$, since at low energy scales $v'/c < 1$, indicating a strong renormalization of the effective group velocity of the propagating quasi-particles due to the presence of the magnetic background fluctuations. In contrast, for larger energy scales the effect becomes weaker, thus recovering the asymptotic limit $v'/c \rightarrow 1$ as $q_0/m \gg 1$.

Our results show that the effective quasi-particle charge experiences a strong renormalization $z_3 < 1$ at low energies q_0/m , while the effect becomes negligible a large energy scales $q_0/m \gg 1$, since $z_3 \rightarrow 1$ as an asymptotic limit. We interpret this as a charge screening due to the spatial magnetic fluctuations in the background. On the other hand, z_3 tends to decrease as a function of the average background magnetic field intensity, achieving an asymptotic limit $z_3 \rightarrow 1/3$ as shown in Eq. (72).

We remark that both the effective refraction index $v'/c = z^{-1}$ and charge screening z_3 display a similar, and therefore consistent, renormalization behaviour of the quasi-particle properties in the magnetically fluctuating environment. In order to understand such effects in physical terms, we remark that the low energy and momentum components in the Fourier representation of the propagator correspond to long-wavelength components in the space of configurations. Therefore, our results are consistent with the fact that such long-wavelength components are more sensitive to the spatial dis-

tribution of the background magnetic noise, and hence experience a higher degree of decoherence, thus reducing the corresponding group velocity and enhancing the charge screening. In contrast, the high-energy Fourier components of the propagator, that correspond to short-wavelength components in the configuration space, are less sensitive to the presence of spatial fluctuations of the background magnetic field. In addition, we remark that the intensity of the average background magnetic field defines a characteristic length-scale known as the Landau radius $l_B = 1/\sqrt{eB}$, that determines the support of the quasi-particle propagator in configuration space. Moreover, in the semi-classical picture this length-scale represents the typical size of the “cyclotron radius” of the helycoidal trajectories that propagate along the magnetic field axis. Therefore, the stronger the magnetic field, the smaller the Landau radius, and hence the quasi-particle propagator is modulated towards higher momentum and energy components that, as previously discussed, are more sensitive to the magnetic noise renormalization effects, as is verified by the trend observed both in z_3 and in v'/c , that strongly decrease as the average magnetic field intensity increases $eB/m^2 \gg 1$.

Moreover, we also showed that 4-point vertex corrections at the second order in Δ^2 lead to a renormalized $\hat{\Delta} = \Delta + O(\Delta^2)$, whose relative magnitude grows with the average magnetic field intensity eB/m^2 , and tend to decrease with the quasi-particle energy scale p_0/m , in agreement with the behavior of v'/c and z_3 and the physical interpretation previously discussed.

The analysis and results presented in this work only concern the study of the quasi-particle fermion propagator in the noisy magnetic field background. However, the effective model obtained via the replica method and its consequences can be extended towards the study of other physical quantities, such as the photon polarization tensor. We are currently investigating this and it will be communicated in a separate article.

ACKNOWLEDGMENTS

J.D.C.-Y. and E.M. acknowledge financial support from ANID PIA Anillo ACT/192023. E.M. also acknowledges financial support from Fondecyt 1190361. M. L. acknowledges support from ANID/CONICYT FONDECYT Regular (Chile) under grants No. 1200483, 1190192 and 1220035. M. L. also acknowledges support from ANID/PIA/APOYO AFB180002 (Chile).

-
- [1] Sk Noor Alam, Victor Roy, Shakeel Ahmad, and Subhasis Chattopadhyay, “Electromagnetic field fluctuation and its correlation with the participant plane in Au + Au and isobaric collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. D* **104**, 114031 (2021).
 - [2] Alejandro Ayala, Jorge David Castaño-Yepes, LA Hernández, Ana Julia Mizher, María Elena Tejeda-Yeomans, and R Zamora, “Anisotropic photon emission from gluon fusion and splitting in a strong magnetic background I: The two-gluon one-

- photon vertex,” (2022), [arXiv:2209.09364 \[hep-ph\]](https://arxiv.org/abs/2209.09364).
- [3] Inghirami, Gabriele, Mace, Mark, Hirono, Yuji, Del Zanna, Luca, Kharzeev, Dmitri E., and Bleicher, Marcus, “Magnetic fields in heavy ion collisions: flow and charge transport,” *Eur. Phys. J. C* **80**, 293 (2020).
- [4] Alejandro Ayala, Jorge David Castaño Yepes, Isabel Domínguez Jiménez, Jordi Salinas San Martín, and María Elena Tejeda-Yeomans, “Centrality dependence of photon yield and elliptic flow from gluon fusion and splitting

- induced by magnetic fields in relativistic heavy-ion collisions,” *Eur. Phys. J. A* **56**, 53 (2020).
- [5] Alejandro Ayala, Jorge David Castano-Yepes, Cesareo A. Dominguez, Luis A. Hernandez, Saul Hernandez-Ortiz, and Maria Elena Tejada-Yeomans, “Prompt photon yield and elliptic flow from gluon fusion induced by magnetic fields in relativistic heavy-ion collisions,” *Phys. Rev. D* **96**, 014023 (2017), [Erratum: *Phys. Rev. D* 96, 119901 (2017)].
- [6] Wit Busza, Krishna Rajagopal, and Wilke van der Schee, “Heavy ion collisions: The big picture and the big questions,” *Annual Review of Nuclear and Particle Science* **68**, 339–376 (2018), <https://doi.org/10.1146/annurev-nucl-101917-020852>.
- [7] Koichi Hattori and Daisuke Satow, “Electrical conductivity of quark-gluon plasma in strong magnetic fields,” *Phys. Rev. D* **94**, 114032 (2016).
- [8] Koichi Hattori and Daisuke Satow, “Gluon spectrum in a quark-gluon plasma under strong magnetic fields,” *Phys. Rev. D* **97**, 014023 (2018).
- [9] Michael Buballa, “Njl-model analysis of dense quark matter,” *Physics Reports* **407**, 205–376 (2005).
- [10] D. Blaschke, A. Ayriyan, and A. (Editors) Friesen, “Compact stars in the QCD phase diagram,” (Universe MppiaG, 2020).
- [11] Julian Schwinger, “On gauge invariance and vacuum polarization,” *Phys. Rev.* **82**, 664–679 (1951).
- [12] W. Dittrich and M. Reuter, “Effective Lagrangians in Quantum Electrodynamics, lecture notes in physics,” (Springer-Verlag, Berlin-Heidelberg, 1985).
- [13] W. Dittrich and H. Gies, “Probing the quantum vacuum: Perturbative effective action approach in quantum electrodynamics and its application, springer tracts in modern physics,” (Springer-Verlag, Berlin-Heidelberg, 2000).
- [14] C. A. Dominguez, Luis A. Hernández, Marcelo Loewe, Cristian Villavicencio, and R. Zamora, “Magnetic field dependence of nucleon parameters from qcd sum rules,” *Phys. Rev. D* **102**, 094007 (2020).
- [15] Koichi Hattori and Kazunori Itakura, “Vacuum birefringence in strong magnetic fields: (i) photon polarization tensor with all the landau levels,” *Annals of Physics* **330**, 23–54 (2013).
- [16] Alejandro Ayala, Jorge David Castaño Yepes, M. Loewe, and Enrique Muñoz, “Gluon polarization tensor in a magnetized medium: Analytic approach starting from the sum over landau levels,” *Phys. Rev. D* **101**, 036016 (2020).
- [17] Ayala, Alejandro, Hernández, Luis A., Loewe, Marcelo, and Villavicencio, Cristian, “Qcd phase diagram in a magnetized medium from the chiral symmetry perspective: the linear sigma model with quarks and the nambu-jona-lasinio model effective descriptions,” *Eur. Phys. J. A* **57**, 234 (2021).
- [18] Alejandro Ayala, Jorge David Castaño Yepes, M. Loewe, and Enrique Muñoz, “Fermion mass and width in qed in a magnetic field,” *Phys. Rev. D* **104**, 016006 (2021).
- [19] Vladimir A. Miransky and Igor A. Shovkovy, “Quantum field theory in a magnetic field: From quantum chromodynamics to graphene and dirac semimetals,” *Physics Reports* **576**, 1–209 (2015), quantum field theory in a magnetic field: From quantum chromodynamics to graphene and Dirac semimetals.
- [20] Holger Gies and Lars Roessler, “Vacuum polarization tensor in inhomogeneous magnetic fields,” *Phys. Rev. D* **84**, 065035 (2011).
- [21] P.-L. Zhao, A.-M. Wang, and G.-Z. Liu, “Effects of random potentials in three-dimensional quantum electrodynamics,” *Phys. Rev. B* **95**, 235144 (2017).
- [22] Valery Gusynin, Anthony Hams, and Manuel Reenders, “Non-perturbative infrared dynamics of three-dimensional qed with a four-fermion interaction,” *Phys. Rev. D* **63**, 045025 (2001).
- [23] M. Mèzard and G. Parisi, “Replica field theory for random manifolds,” *J. Phys. I France* **1**, 809–836 (1991).
- [24] M. Kardar, G. Parisi, and Y.-C. Zhang, “Dynamic scaling of growing interfaces,” *Phys. Rev. Lett.* **56**, 889–892 (1986).
- [25] I. S. Gradshteyn and I. M. Ryzik, “Table of Integrals, Series and Products, 6th ed.” (Academic Press, San Diego-CA, 2000) pp. 992–8.975–1.
- [26] I. S. Gradshteyn and I. M. Ryzik, “Table of Integrals, Series and Products, 6th ed.” (Academic Press, San Diego-CA, 2000) pp. 803–7.414–6.

Appendix A: The coefficient \mathcal{A}_1

We can calculate the integral \mathcal{A}_1 in terms of Landau levels by means of the generating function of the Laguerre polynomials

$$e^{-\frac{x}{2} \frac{1-t}{1+t}} = (1+t)e^{-x/2} \sum_{n=0}^{\infty} (-t)^n L_n^0(x), \quad (\text{A1})$$

since

$$\begin{aligned} e^{-ix \tan v} &= e^{-x(1-e^{-2iv})/(1+e^{-2iv})} \\ &= (1+e^{-2iv}) e^{-x} \sum_{n=0}^{\infty} (-1)^n e^{-2in v} L_n^0(2x) \end{aligned} \quad (\text{A2})$$

Therefore, we have (for $x = \mathbf{k}_{\perp}^2/eB$)

$$\begin{aligned} \mathcal{A}_1 &= e^{-x} \left(\sum_{n=0}^{\infty} (-1)^n L_n^0(2x) \int_0^{\infty} d\tau e^{i(\mathcal{D}_{\parallel} - 2(n+1)eB)\tau} \right. \\ &\quad \left. + \sum_{n=0}^{\infty} (-1)^n L_n^0(2x) \int_0^{\infty} d\tau e^{i(\mathcal{D}_{\parallel} - 2neB)\tau} \right) \end{aligned} \quad (\text{A3})$$

Evaluating the exponential integrals,

$$\begin{aligned} \mathcal{A}_1 &= ie^{-x} \left(\sum_{n=0}^{\infty} (-1)^n \frac{L_n^0(2x)}{\mathcal{D}_{\parallel} - 2(n+1)eB} \right. \\ &\quad \left. + \sum_{n=0}^{\infty} (-1)^n \frac{L_n^0(2x)}{\mathcal{D}_{\parallel} - 2neB} \right) \\ &= i \frac{e^{-x}}{\mathcal{D}_{\parallel}} \left[1 + \sum_{n=1}^{\infty} \frac{(-1)^n [L_n^0(2x) - L_{n-1}^0(2x)]}{1 - 2n \frac{eB}{\mathcal{D}_{\parallel}}} \right] \end{aligned} \quad (\text{A4})$$

This expansion seems is fair for $eB > 0$. However, we would like to inspect if it is possible to use it to generate a valid expansion near $eB = 0$.

1. Expansion for low magnetic fields

Notice that, from Eq. (44) we have the identities

$$e^{-x} \sum_{n=0}^{\infty} (-1)^n e^{-2in v} L_n^0(2x) = \frac{e^{-ix \tan v}}{1 + e^{-2iv}} \quad (\text{A5})$$

$$e^{-x} \sum_{n=0}^{\infty} (-1)^n e^{-2i(n+1)v} L_n^0(2x) = \frac{e^{-2iv} e^{-ix \tan v}}{1 + e^{-2iv}} \quad (\text{A6})$$

Therefore, we can calculate the following expansions

$$\begin{aligned} e^{-x} \sum_{n=0}^{\infty} \frac{(-1)^n L_n^0(2x)}{\mathcal{D}_{\parallel} - 2(n+1)eB} &= \frac{1}{\mathcal{D}_{\parallel}} \sum_{k=0}^{\infty} \left(\frac{2eB}{\mathcal{D}_{\parallel}} \right)^k \\ &\times \left[e^{-x} \sum_{n=0}^{\infty} (-1)^n (n+1)^k L_n^0(2x) \right] \\ &= \frac{1}{\mathcal{D}_{\parallel}} \sum_{k=0}^{\infty} \left(\frac{2eB}{\mathcal{D}_{\parallel}} \right)^k \frac{1}{(-2i)^k} \frac{\partial^k}{\partial v^k} \left(\frac{e^{-2iv} e^{-ix \tan v}}{1 + e^{-2iv}} \right) \Big|_{v \rightarrow 0} \quad (\text{A7}) \end{aligned}$$

and similarly

$$\begin{aligned} e^{-x} \sum_{n=0}^{\infty} \frac{(-1)^n L_n^0(2x)}{\mathcal{D}_{\parallel} - 2neB} &= \frac{1}{\mathcal{D}_{\parallel}} \sum_{k=0}^{\infty} \left(\frac{2eB}{\mathcal{D}_{\parallel}} \right)^k \\ &\times \left[e^{-x} \sum_{n=0}^{\infty} (-1)^n n^k L_n^0(2x) \right] \\ &= \frac{1}{\mathcal{D}_{\parallel}} \sum_{k=0}^{\infty} \left(\frac{2eB}{\mathcal{D}_{\parallel}} \right)^k \frac{1}{(-2i)^k} \frac{\partial^k}{\partial v^k} \left(\frac{e^{-ix \tan v}}{1 + e^{-2iv}} \right) \Big|_{v \rightarrow 0} \quad (\text{A8}) \end{aligned}$$

Substituting both expressions into Eq.(A4), we obtain the infinite series

$$\mathcal{A}_1 = \frac{i}{\mathcal{D}_{\parallel}} \left(1 + \sum_{k=1}^{\infty} \left(\frac{ieB}{\mathcal{D}_{\parallel}} \right)^k \mathcal{E}_k(x) \right) \quad (\text{A9})$$

Here, we have defined the polynomials $\mathcal{E}_k(x)$ (for $x = k_{\perp}^2/eB$) as generated by the function $e^{-ix \tan v}$,

$$\mathcal{E}_k(x) = \lim_{v \rightarrow 0} \frac{\partial^k}{\partial v^k} (e^{-ix \tan v}) \quad (\text{A10})$$

The first few cases

$$\begin{aligned} \mathcal{E}_1(x) &= -ix \\ \mathcal{E}_2(x) &= -x^2 \\ \mathcal{E}_3(x) &= -2ix + ix^3 \\ \mathcal{E}_4(x) &= x^4 - 8x^2 \\ \mathcal{E}_5(x) &= -ix^5 + 20ix^3 - 16ix \\ \mathcal{E}_6(x) &= -x^6 + 40x^4 - 136x^2 \\ &\vdots \end{aligned} \quad (\text{A11})$$

Substituting these expressions into the series Eq. (34), it can be reorganized as an expansion in terms of the variable

$y = k_{\perp}^2/\mathcal{D}_{\parallel}$, as follows

$$\begin{aligned} \mathcal{A}_1 &= \frac{i}{\mathcal{D}_{\parallel}} \left[(1 + y + y^2 + y^3 + \dots) \right. \\ &\quad \left. - 2 \left(\frac{eB}{\mathcal{D}_{\parallel}} \right)^2 y (1 + 4y + 10y^2 + 20y^4 + \dots) \right. \\ &\quad \left. + 8 \left(\frac{eB}{\mathcal{D}_{\parallel}} \right)^4 y (2 + 17y + 77y^2 + \dots) \right] + O \left(\frac{eB}{\mathcal{D}_{\parallel}} \right)^6 \\ &= \frac{i}{\mathcal{D}_{\parallel}} \left[\frac{1}{1-y} - 2 \left(\frac{eB}{\mathcal{D}_{\parallel}} \right)^2 \frac{y}{(1-y)^4} \right. \\ &\quad \left. + 8 \left(\frac{eB}{\mathcal{D}_{\parallel}} \right)^4 \frac{y(3y+2)}{(1-y)^7} \right] + O \left(\frac{eB}{\mathcal{D}_{\parallel}} \right)^6 \quad (\text{A13}) \end{aligned}$$

Finally, using the simple identity

$$\mathcal{D}_{\parallel}(1-y) = \mathcal{D}_{\parallel}(1 - \frac{k_{\perp}^2}{\mathcal{D}_{\parallel}}) = k^2 - m^2 + i\epsilon \quad (\text{A14})$$

we obtain the final form

$$\begin{aligned} \mathcal{A}_1 &= \frac{i}{k^2 - m^2 + i\epsilon} \left[1 - 2 \left(\frac{eB}{\mathcal{D}_{\parallel}} \right)^2 \frac{y}{(1-y)^3} \right. \\ &\quad \left. + 8 \left(\frac{eB}{\mathcal{D}_{\parallel}} \right)^4 \frac{y(3y+2)}{(1-y)^6} \right] + O \left(\frac{eB}{\mathcal{D}_{\parallel}} \right)^6 \\ &= \frac{i}{k^2 - m^2 + i\epsilon} + \frac{-2i(eB)^2 k_{\perp}^2}{[k^2 - m^2 + i\epsilon]^4} + O((eB)^4). \quad (\text{A15}) \end{aligned}$$

2. A closed expression in terms of Hypergeometric functions

In order to simplify the integrals \mathcal{J}_i , let us to provide an analytical expression for $\mathcal{A}_1(k)$. From the generating function of the Laguerre's polynomials:

$$\sum_{n=0}^{\infty} (-1)^n t^n L_n^{\alpha}(x) = \frac{1}{(1+t)^{\alpha}} \exp \left(\frac{t}{1+t} x \right), \quad (\text{A16})$$

then, by defining:

$$b \equiv \frac{t}{1+t}, \quad (\text{A17})$$

we get:

$$(1-b)^{t+\alpha} e^{bx} = \sum_{n=0}^{\infty} (-1)^n \frac{b^n}{(1-b)^n} L_n^{\alpha}(x). \quad (\text{A18})$$

Multiplying by $b^{\beta}/(1-b)^{\beta+2}$:

$$b^{\beta} (1+b)^{\alpha-\beta-1} e^{-bx} = \sum_{n=0}^{\infty} (-1)^n b^{n+\beta} (1-b)^{-n-\beta-2} L_n^{\alpha}(x), \quad (\text{A19})$$

so that:

$$\int_0^{-\infty} db b^\beta (1+b)^{\alpha-\beta-1} e^{-bx} = - \sum_{n=0}^{\infty} \frac{(-1)^n L_n^\alpha(x)}{n+\beta+1}. \quad (\text{A20})$$

Now by setting $b \rightarrow -b$:

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(-1)^n L_n^\alpha(x)}{n+\beta+1} &= (-1)^\beta \int_0^{\infty} db b^\beta (1+b)^{\alpha-\beta-1} e^{-bx} \\ &= e^{i\pi\beta} \Gamma(1+\beta) U(1+\beta, 1+\alpha, x), \end{aligned} \quad (\text{A21})$$

where $\Gamma(z)$, and $U(a, b, z)$ are the gamma-function and the confluent hypergeometric function, respectively.

Now, from Eq. (A4):

$$\begin{aligned} \mathcal{A}_1 &= ie^{-x} \sum_{n=0}^{\infty} \left(\frac{(-1)^n L_n^0(2x)}{\mathcal{D}_\parallel - 2(n+1)eB} + \frac{(-1)^n L_n^0(2x)}{\mathcal{D}_\parallel - 2neB} \right) \\ &= -\frac{ie^{-x}}{2eB} \sum_{n=0}^{\infty} \left[\frac{(-1)^n L_n^0(2x)}{n + \left(1 - \frac{\mathcal{D}_\parallel}{2eB}\right)} + \frac{(-1)^n L_n^0(2x)}{n - \frac{\mathcal{D}_\parallel}{2eB}} \right]. \end{aligned} \quad (\text{A22})$$

Therefore, by using Eq. (A21):

$$\begin{aligned} \mathcal{A}_1 &= -\frac{ie^{-x}}{2eB} \\ &\times \left[e^{-i\pi \frac{\mathcal{D}_\parallel}{2eB}} \Gamma\left(1 - \frac{\mathcal{D}_\parallel}{2eB}\right) U\left(1 - \frac{\mathcal{D}_\parallel}{2eB}, 1, 2x\right) \right. \\ &\left. + e^{-i\pi \left(1 + \frac{\mathcal{D}_\parallel}{2eB}\right)} \Gamma\left(-\frac{\mathcal{D}_\parallel}{2eB}\right) U\left(-\frac{\mathcal{D}_\parallel}{2eB}, 1, 2x\right) \right]. \end{aligned} \quad (\text{A23})$$

The latter can be simplified with the properties of the gamma and the hypergeometric functions. First, from the identity $\Gamma(z+1) = z\Gamma(z)$:

$$\begin{aligned} \mathcal{A}_1 &= \frac{ie^{-x}}{2eB} \exp\left(-\frac{i\pi\mathcal{D}_\parallel}{2eB}\right) \Gamma\left(-\frac{\mathcal{D}_\parallel}{2eB}\right) \\ &\times \left[\frac{\mathcal{D}_\parallel}{2eB} U\left(1 - \frac{\mathcal{D}_\parallel}{2eB}, 1, 2x\right) + U\left(-\frac{\mathcal{D}_\parallel}{2eB}, 1, 2x\right) \right]. \end{aligned} \quad (\text{A24})$$

Moreover, given that:

$$U(a, b, z) - aU(a+1, b, z) = U(a, b-1, z), \quad (\text{A25})$$

we finally arrive to:

$$\begin{aligned} \mathcal{A}_1(k) &= \frac{ie^{-\mathbf{k}_\perp^2/eB}}{2eB} \exp\left[-\frac{i\pi\left(k_\parallel^2 - m^2\right)}{2eB}\right] \\ &\times \Gamma\left(-\frac{k_\parallel^2 - m^2}{2eB}\right) U\left(-\frac{k_\parallel^2 - m^2}{2eB}, 0, \frac{2\mathbf{k}_\perp^2}{eB}\right). \end{aligned} \quad (\text{A26})$$

Appendix B: The density of states $\rho(E)$

In this appendix, we show the details for the calculation of the density of states for the Landau level spectrum $\rho(E)$ defined in the main text. We start from the definition of the density of states

$$\begin{aligned} \rho(E) &= \int_{-\infty}^{\infty} \frac{dp_3}{2\pi} \sum_{n=0}^{\infty} \delta\left(E - \sqrt{p_3^2 + m^2 + 2(n+1)eB}\right) \\ &= 2 \sum_{n=0}^{\infty} \int_0^{\infty} \frac{dp_3}{2\pi} \frac{\delta\left(p^3 - \sqrt{E^2 - m^2 - 2(n+1)eB}\right)}{p^3/E} \\ &= \frac{E}{\pi} \sum_{n=0}^{\infty} \frac{\Theta\left(E - \sqrt{m^2 + 2(n+1)eB}\right)}{\sqrt{E^2 - m^2 - 2(n+1)eB}} \end{aligned} \quad (\text{B1})$$

Since this function, and hence the corresponding sum, is defined for each fixed value of the energy E , there is a maximum integer $n = N_{max}(E)$ at which the sum is truncated by the condition imposed on the Heaviside step function

$$E - \sqrt{m^2 + 2(N_{max} + 1)eB} = 0, \quad (\text{B2})$$

that leads to the definition (with $[z]$ the lowest integer part)

$$N_{max}(E) = \lfloor \frac{E^2 - m^2}{2eB} - 1 \rfloor. \quad (\text{B3})$$

Hence, we have

$$\begin{aligned} \rho(E) &= \Theta(E - \sqrt{m^2 + 2eB}) \frac{E}{\pi} \\ &\times \sum_{n=0}^{N_{max}(E)} \frac{1}{\sqrt{E^2 - m^2 - 2(n+1)eB}} \end{aligned} \quad (\text{B4})$$

In this finite sum, $0 \leq n \leq N_{max}(E)$, we can redefine the index by

$$\ell \equiv N_{max}(E) - n \implies 0 \leq \ell \leq N_{max}(E), \quad (\text{B5})$$

and hence we have the equivalent expression

$$\begin{aligned} \rho(E) &= \Theta(E - \sqrt{m^2 + 2eB}) \frac{E}{\pi\sqrt{eB}} \\ &\times \sum_{\ell=0}^{N_{max}(E)} \frac{1}{\sqrt{\frac{E^2 - m^2 - 2(N_{max}(E)+1)eB}{2eB} + \ell}} \end{aligned} \quad (\text{B6})$$

Finally, using the property of the Riemann Zeta function

$$\sum_{\ell=0}^N (z + \ell)^{-s} = \zeta(s, z) - \zeta(s, z + N + 1), \quad (\text{B7})$$

we obtain

$$\begin{aligned} \rho(E) &= \Theta(E - \sqrt{m^2 + 2eB}) \frac{E}{\pi\sqrt{eB}} \\ &\times \left[\zeta\left(\frac{1}{2}, \frac{E^2 - m^2 - 2eB}{2eB} - N_{max}(E)\right) \right. \\ &\left. - \zeta\left(\frac{1}{2}, \frac{E^2 - m^2}{2eB}\right) \right] \end{aligned} \quad (\text{B8})$$

Appendix C: Computing \tilde{A}_1 and \tilde{A}_2

1. Weak magnetic field limit

We need to compute:

$$\tilde{\mathcal{A}}_1(q_0) = -2i(eB)^2 \int d^3p \frac{p_\perp^2}{(q_0^2 - p_3^2 - p_\perp^2 - m^2 + i\epsilon)^4} \quad (C1)$$

so that in spherical coordinates, with $p_3 = p \cos \theta$ and $p_\perp = p \sin \theta$:

$$\begin{aligned} \tilde{\mathcal{A}}_1(q_0) &= -4\pi i(eB)^2 \int_0^\pi d\theta \sin^3 \theta \int_0^\infty dp \frac{p^4}{(q_0^2 - p^2 - m^2 + i\epsilon)^4} \\ &= -4\pi i(eB)^2 \frac{4}{3} \int_0^\infty dp \frac{p^4}{(q_0^2 - p^2 - m^2 + i\epsilon)^4} \\ &= -4\pi i(eB)^2 \frac{4}{3} \int_0^\infty dp \frac{p^4}{(q_0^2 - p^2 - m^2 + i\epsilon)^4} \end{aligned} \quad (C2)$$

By defining

$$a^2 \equiv q_0^2 - m^2 + i\epsilon, \quad (C3)$$

and

$$a_\pm = \sqrt{q_0^2 - m^2} \pm i\epsilon, \quad (C4)$$

we can use complex integration:

$$\begin{aligned} \tilde{\mathcal{A}}_1(q_0) &= -4\pi i(eB)^2 \frac{4}{3} \frac{1}{2} \int_{-\infty}^\infty dz \frac{z^4}{(z^2 - a^2)^4} \\ &= -4\pi i(eB)^2 \frac{4}{3} \frac{1}{2} \frac{2\pi i}{3!} \lim_{z \rightarrow a_+} \frac{d^3}{dz^3} \left[\frac{z^4}{(z - a_-)^4} \right] \\ &= -4\pi i(eB)^2 \frac{4}{3} \frac{1}{2} \frac{2\pi i}{3!} \frac{(-3)}{16(q_0^2 - m^2)^{3/2}} \\ &= -\frac{\pi^2}{6} \frac{(eB)^2}{(q_0^2 - m^2)^{3/2}}. \end{aligned} \quad (C5)$$

On the other hand, at order $\mathcal{O}((eB)^2)$:

$$\begin{aligned} \tilde{\mathcal{A}}_2(q_0) &= i \int_{-\infty}^{+\infty} \frac{dp_3}{q_0^2 - (p^3)^2 - m^2 + i\epsilon} \\ &= -i \int_{-\infty}^{+\infty} \frac{dz}{z^2 - a^2} \\ &= -i \int_{-\infty}^{+\infty} \frac{dz}{(z - a_+)(z + a_-)} \end{aligned} \quad (C6)$$

By choosing a contour closing upside, we get from the residue theorem:

$$-i \int_{-\infty}^{+\infty} \frac{dz}{(z - a_+)(z + a_-)} = \frac{\pi}{a_+}. \quad (C7)$$

Then:

$$\tilde{\mathcal{A}}_2(q_0) = \frac{\pi}{\sqrt{q_0^2 - m^2}} \quad (C8)$$

2. Arbitrary magnetic field

From the definition of $\tilde{\mathcal{A}}_1$:

$$\begin{aligned} \tilde{\mathcal{A}}_1(q_0) &= \int d^3p \mathcal{A}_1(q_0, p_3; \mathbf{p}_\perp) \\ &= i \int d^3p \left[\frac{e^{-\mathbf{p}_\perp^2/eB}}{q_0^2 - p_3^2 - m^2 + i\epsilon} \right. \\ &\quad \left. + \sum_{n=1}^{\infty} (-1)^n e^{-\mathbf{p}_\perp^2/eB} \frac{L_n\left(\frac{2\mathbf{p}_\perp^2}{eB}\right) - L_{n-1}\left(\frac{2\mathbf{p}_\perp^2}{eB}\right)}{q_0^2 - p_3^2 - m^2 - 2neB + i\epsilon} \right] \\ &= \mathcal{I}_1 + \sum_{n=1}^{\infty} (-1)^n \mathcal{I}_{2,n} \end{aligned} \quad (C9)$$

Here, we defined

$$\begin{aligned} \mathcal{I}_1 &= i \int d^3p \frac{e^{-\mathbf{p}_\perp^2/eB}}{q_0^2 - p_3^2 - m^2 + i\epsilon} \\ &= -i \int d^2p_\perp e^{-\mathbf{p}_\perp^2/eB} \int_{-\infty}^\infty \frac{dz}{(z - a - i\epsilon)(z + a + i\epsilon)} \\ &= -i\pi(eB) \frac{2\pi i}{2(a + i\epsilon)} = \frac{\pi^2 eB}{\sqrt{q_0^2 - m^2 + i\epsilon}}, \end{aligned} \quad (C10)$$

and

$$\mathcal{I}_{2,n} = i \int d^3p e^{-\mathbf{p}_\perp^2/eB} \frac{L_n\left(\frac{2\mathbf{p}_\perp^2}{eB}\right) - L_{n-1}\left(\frac{2\mathbf{p}_\perp^2}{eB}\right)}{q_0^2 - p_3^2 - m^2 - 2neB + i\epsilon} \quad (C11)$$

In cylindrical coordinates, with azimuthal symmetry, $d^3p = dp_3 \pi d(\mathbf{p}_\perp^2)$. Moreover, in the integral over \mathbf{p}_\perp , define $x = \frac{2\mathbf{p}_\perp^2}{eB}$, such that

$$\begin{aligned} \mathcal{I}_{2,n} &= \frac{\pi qB}{2} \int_{-\infty}^\infty \frac{dp_3}{q_0^2 - m^2 - p_3^2 - 2nqB + i\epsilon} \\ &\quad \times \int_0^\infty dx e^{-x/2} [L_n(x) - L_{n-1}(x)] \\ &= 2\pi qB (-1)^n \int_{-\infty}^\infty \frac{dp_3}{q_0^2 - m^2 - p_3^2 - 2nqB + i\epsilon} \end{aligned} \quad (C12)$$

where we used the identity (Gradshteyn-Ryzhik)

$$\int_0^\infty dx e^{-bx} L_n(x) = (b-1)^n b^{-n-1} \quad \text{Re } b > 0 \quad (C13)$$

Inserting Eq. (47) and Eq. (49) into Eq. (46), we obtain (after shifting the index $n \rightarrow n+1$)

$$\begin{aligned} \tilde{\mathcal{A}}_1(q_0) &= \frac{\pi^2(eB)}{\sqrt{q_0^2 - m^2 + i\epsilon}} \\ &\quad + 2\pi qB \int_{-\infty}^{+\infty} dp_3 \sum_{n=0}^{\infty} \frac{1}{q_0^2 - m^2 - p_3^2 - 2(n+1)eB + i\epsilon} \end{aligned} \quad (C14)$$

Let us introduce the density of states for Landau levels

$$\rho(E) = \int_{-\infty}^\infty \frac{dp_3}{2\pi} \sum_{n=0}^{\infty} \delta(E - E_n(p_3)), \quad (C15)$$

with the dispersion relation for the spectrum

$$E_n(p_3) = \sqrt{p_3^2 + m^2 + 2(n+1)eB}. \quad (\text{C16})$$

With these definitions, we obtain from Eq. (52) the exact expression

$$\tilde{\mathcal{A}}_1(q_0) = \frac{\pi^2 eB}{\sqrt{q_0^2 - m^2 + i\epsilon}} + 4\pi^2 eB \int_{-\infty}^{+\infty} dE \frac{\rho(E)}{q_0^2 - E^2 + i\epsilon}, \quad (\text{C17})$$

where, as shown in detail in Appendix B

$$\begin{aligned} \rho(E) &= \Theta(E - \sqrt{m^2 + 2eB}) \frac{E}{\pi\sqrt{eB}} \\ &\times \left[\zeta\left(\frac{1}{2}, \frac{E^2 - m^2 - 2eB}{2eB}\right) - N_{max}(E) \right] \\ &- \zeta\left(\frac{1}{2}, \frac{E^2 - m^2}{2eB}\right), \end{aligned} \quad (\text{C18})$$

where we defined

$$N_{max}(E) = \lfloor \frac{E^2 - m^2}{2eB} - 1 \rfloor, \quad (\text{C19})$$

with $\lfloor x \rfloor$ the integer part of x , and $\zeta(n, z)$ is the Riemann zeta function.

On the other hand:

$$\begin{aligned} \tilde{\mathcal{A}}_2(q_0) &= \int_{-\infty}^{\infty} dp_3 p \mathcal{A}_1(q_0, p_3; \mathbf{p}_\perp = 0) \\ &= i \int_{-\infty}^{\infty} dp_3 \frac{1}{q_0^2 - p_3^2 - m^2 + i\epsilon} \\ &+ i \int_{-\infty}^{\infty} dp_3 \sum_{n=1}^{\infty} (-1)^n \frac{L_n(0) - L_{n-1}(0)}{q_0^2 - p_3^2 - m^2 - 2neB + i\epsilon}, \end{aligned} \quad (\text{C20})$$

where the second term vanishes, given that $L_n(0) = 1 \ \forall n$. Hence

$$\tilde{\mathcal{A}}_2(q_0) = \mathcal{I}_1 = \frac{\pi^2 eB}{\sqrt{q_0^2 - m^2 + i\epsilon}}. \quad (\text{C21})$$

3. Strong magnetic field limit

In this limit:

$$\mathcal{A}_1(q) = i \frac{e^{-\mathbf{q}_\perp^2/eB}}{q_\parallel^2 - m^2}, \quad (\text{C22a})$$

$$\mathcal{A}_2(q) = \frac{e^{-\mathbf{q}_\perp^2/eB}}{q_\parallel^2 - m^2}, \quad (\text{C22b})$$

$$\mathcal{A}_3(q) = 0, \quad (\text{C22c})$$

$$\mathcal{D}(q) = 2 \frac{e^{-2\mathbf{q}_\perp^2/eB}}{q_\parallel^2 - m^2} \quad (\text{C22d})$$

and

$$\tilde{\mathcal{A}}_1(q) = \frac{\pi^2 eB}{\sqrt{q_0^2 - m^2}} \quad (\text{C23a})$$

$$\tilde{\mathcal{A}}_2(q_0) = \frac{\pi}{\sqrt{q_0^2 - m^2}}. \quad (\text{C23b})$$

Appendix D: Vertex corrections at $O(\Delta^2)$

The diagrams contributing at order Δ^2 to the 4-point vertex are depicted in Fig. 11, and hence their corresponding matrix elements are given by the following integral expressions

$$\hat{\Gamma}_{(a)} = \int \frac{d^3 q}{(2\pi)^3} \gamma^i S_F(p - q) \gamma^j \otimes \gamma_i S_F(p' - q) \gamma_j, \quad (\text{D1a})$$

$$\hat{\Gamma}_{(b)} = \int \frac{d^3 q}{(2\pi)^3} \gamma^i S_F(p - q) \gamma^j \otimes \gamma_i S_F(p' + q) \gamma_j, \quad (\text{D1b})$$

and

$$\hat{\Gamma}_{(c)} = \int \frac{d^3 q}{(2\pi)^3} \gamma^i S_F(p + q) \gamma^j \otimes \gamma_i S_F(p' - q) \gamma_j \quad (\text{D1c})$$

In order to compute the former expressions, it is convenient to introduce the notation

$$\hat{\Gamma}^{(\lambda, \sigma)} = \int \frac{d^3 q}{(2\pi)^3} \gamma^i S_F(p + \lambda q) \gamma^j \otimes \gamma_i S_F(p' + \sigma q) \gamma_j. \quad (\text{D2})$$

where $\lambda, \sigma = \pm 1$. Then, we have the correspondence $\hat{\Gamma}_{(a)} = \hat{\Gamma}^{(-, -)}$, $\hat{\Gamma}_{(b)} = \hat{\Gamma}^{(-, +)}$, and $\hat{\Gamma}_{(c)} = \hat{\Gamma}^{(+, -)}$, respectively. By considering the tensor structure of the propagator, it is straightforward to realize that the full vertex, taking into account the multiplicity factors for each diagram,

$$\hat{\Gamma} = 2\hat{\Gamma}^{(-, -)} + 2\hat{\Gamma}^{(-, +)} + 4\hat{\Gamma}^{(+, -)}, \quad (\text{D3})$$

leads to an effective interaction of the form

$$\hat{\Gamma} = \tilde{\Delta}(\bar{\psi} \gamma^i \psi)(\bar{\psi} \gamma^i \psi) + \text{other tensor structures}, \quad (\text{D4})$$

where the renormalized coefficient $\tilde{\Delta}$ is given up to second order in Δ by

$$\begin{aligned} \tilde{\Delta} &= \Delta + 2(\Delta)^2 \left(\mathcal{J}_2^{(-, -)} + \mathcal{J}_2^{(-, +)} + 2\mathcal{J}_2^{(+, -)} \right. \\ &+ (1 - \partial_x^2)(1 - \partial_y^2) \mathcal{J}_3^{(-, -)} + (1 - \partial_x^2)(1 - \partial_y^2) \mathcal{J}_3^{(-, +)} \\ &+ 2(1 - \partial_x^2)(1 - \partial_y^2) \mathcal{J}_3^{(+, -)} \left. \right) \end{aligned} \quad (\text{D5})$$

Now, from Eq. (16):

$$\begin{aligned} \hat{\Gamma}^{(\lambda, \sigma)} &= - \left(\hat{\Gamma}_{11}^{(\lambda, \sigma)} + \hat{\Gamma}_{12}^{(\lambda, \sigma)} + \hat{\Gamma}_{13}^{(\lambda, \sigma)} + \hat{\Gamma}_{21}^{(\lambda, \sigma)} \right. \\ &+ \hat{\Gamma}_{22}^{(\lambda, \sigma)} + \hat{\Gamma}_{23}^{(\lambda, \sigma)} + \hat{\Gamma}_{31}^{(\lambda, \sigma)} + \hat{\Gamma}_{32}^{(\lambda, \sigma)} + \hat{\Gamma}_{33}^{(\lambda, \sigma)} \left. \right), \end{aligned} \quad (\text{D6})$$

where

$$\hat{\Gamma}_{11}^{(\lambda, \sigma)} = \int \frac{d^3 q}{(2\pi)^3} (m + \not{p}_{\parallel} + \lambda \not{q}_{\parallel}) (m + \not{p}'_{\parallel} + \sigma \not{q}_{\parallel}) \times \mathcal{A}_1(p + \lambda q) \mathcal{A}_1(p' + \sigma q), \quad (\text{D7a})$$

$$\hat{\Gamma}_{12}^{(\lambda, \sigma)} = i\gamma^1 \gamma^2 \int \frac{d^3 q}{(2\pi)^3} (m + \not{p}_{\parallel} + \lambda \not{q}_{\parallel}) (m + \not{p}'_{\parallel} + \sigma \not{q}_{\parallel}) \times \mathcal{A}_1(p + \lambda q) \mathcal{A}_2(p' + \sigma q), \quad (\text{D7b})$$

$$\hat{\Gamma}_{13}^{(\lambda, \sigma)} = \int \frac{d^3 q}{(2\pi)^3} (m + \not{p}_{\parallel} + \lambda \not{q}_{\parallel}) (\not{p}'_{\perp} + \sigma \not{q}_{\perp}) \times \mathcal{A}_1(p + \lambda q) \mathcal{A}_3(p' + \sigma q), \quad (\text{D7c})$$

$$\hat{\Gamma}_{21}^{(\lambda, \sigma)} = i\gamma^1 \gamma^2 \int \frac{d^3 q}{(2\pi)^3} (m + \not{p}_{\parallel} + \lambda \not{q}_{\parallel}) (m + \not{p}'_{\parallel} + \sigma \not{q}_{\parallel}) \times \mathcal{A}_2(p + \lambda q) \mathcal{A}_1(p' + \sigma q), \quad (\text{D7d})$$

$$\hat{\Gamma}_{22}^{(\lambda, \sigma)} = - \int \frac{d^3 q}{(2\pi)^3} (m + \not{p}_{\parallel} + \lambda \not{q}_{\parallel}) (m + \not{p}'_{\parallel} + \sigma \not{q}_{\parallel}) \times \mathcal{A}_2(p + \lambda q) \mathcal{A}_2(p' + \sigma q), \quad (\text{D7e})$$

$$\hat{\Gamma}_{23}^{(\lambda, \sigma)} = i\gamma^1 \gamma^2 \int \frac{d^3 q}{(2\pi)^3} (m + \not{p}_{\parallel} + \lambda \not{q}_{\parallel}) (\not{p}'_{\perp} + \sigma \not{q}_{\perp}) \times \mathcal{A}_2(p + \lambda q) \mathcal{A}_3(p' + \sigma q), \quad (\text{D7f})$$

$$\hat{\Gamma}_{31}^{(\lambda, \sigma)} = \int \frac{d^3 q}{(2\pi)^3} (\not{p}_{\perp} + \lambda \not{q}_{\perp}) (m + \not{p}'_{\parallel} + \sigma \not{q}_{\parallel}) \times \mathcal{A}_3(p + \lambda q) \mathcal{A}_1(p' + \sigma q), \quad (\text{D7g})$$

$$\hat{\Gamma}_{32}^{(\lambda, \sigma)} = -i\gamma^1 \gamma^2 \int \frac{d^3 q}{(2\pi)^3} (\not{p}_{\perp} + \lambda \not{q}_{\perp}) (m + \not{p}'_{\parallel} + \sigma \not{q}_{\parallel}) \times \mathcal{A}_3(p + \lambda q) \mathcal{A}_2(p' + \sigma q), \quad (\text{D7h})$$

$$\hat{\Gamma}_{33}^{(\lambda, \sigma)} = \int \frac{d^3 q}{(2\pi)^3} (\not{p}_{\perp} + \lambda \not{q}_{\perp}) (\not{p}'_{\perp} + \sigma \not{q}_{\perp}) \times \mathcal{A}_3(p + \lambda q) \mathcal{A}_3(p' + \sigma q). \quad (\text{D7i})$$

The latter equations can be condensed by defining a single master integral in terms of \mathcal{A}_1 and its derivatives. To do so, note that:

$$\begin{aligned} & (m + \not{p}_{\parallel} + \lambda \not{q}_{\parallel}) (m + \not{p}'_{\parallel} + \sigma \not{q}_{\parallel}) \\ &= m^2 + m [\not{p}_{\parallel} + \not{p}'_{\parallel} + (\sigma + \lambda) \not{q}_{\parallel}] + \not{p}_{\parallel} \not{p}'_{\parallel} \\ &+ \sigma \not{p}_{\parallel} \not{q}_{\parallel} + \lambda \not{q}_{\parallel} \not{p}'_{\parallel} + \lambda \sigma (\not{q}_{\parallel})^2, \end{aligned} \quad (\text{D8})$$

and given that \mathcal{A}_i are even functions of q , the linear terms can be ignored. Then:

$$\begin{aligned} & (m + \not{p}_{\parallel} + \lambda \not{q}_{\parallel}) (m + \not{p}'_{\parallel} + \sigma \not{q}_{\parallel}) \\ & \rightarrow m^2 + m (\not{p}_{\parallel} + \not{p}'_{\parallel}) + \not{p}_{\parallel} \not{p}'_{\parallel} + \lambda \sigma q_{\parallel}^2. \end{aligned} \quad (\text{D9})$$

Now, it is convenient to define:

$$\begin{aligned} P &\equiv \frac{p' + p}{2}, \\ Q &\equiv \frac{p' - p}{2}, \end{aligned} \quad (\text{D10})$$

from which:

$$\begin{aligned} & (m + \not{p}_{\parallel} + \lambda \not{q}_{\parallel}) (m + \not{p}'_{\parallel} + \sigma \not{q}_{\parallel}) \\ & \rightarrow m^2 + 2m \not{P}_{\parallel} + (\not{P}_{\parallel} - \not{Q}_{\parallel}) (\not{P}_{\parallel} + \not{Q}_{\parallel}) + \lambda \sigma q_{\parallel}^2. \end{aligned} \quad (\text{D11})$$

Similarly:

$$\begin{aligned} & (m + \not{p}_{\parallel} + \lambda \not{q}_{\parallel}) (\not{p}'_{\perp} + \sigma \not{q}_{\perp}) \\ & \rightarrow m (\not{P}_{\perp} + \not{Q}_{\perp}) + (\not{P}_{\parallel} - \not{Q}_{\parallel}) (\not{P}_{\perp} + \not{Q}_{\perp}), \end{aligned} \quad (\text{D12})$$

$$\begin{aligned} & (\not{p}_{\perp} + \lambda \not{q}_{\perp}) (m + \not{p}'_{\parallel} + \sigma \not{q}_{\parallel}) \\ & \rightarrow m (\not{P}_{\perp} - \not{Q}_{\perp}) + (\not{P}_{\perp} - \not{Q}_{\perp}) (\not{P}_{\parallel} + \not{Q}_{\parallel}), \end{aligned} \quad (\text{D13})$$

and

$$\begin{aligned} & (\not{p}_{\perp} + \lambda \not{q}_{\perp}) (\not{p}'_{\perp} + \sigma \not{q}_{\perp}) \\ & \rightarrow (\not{P}_{\perp} - \not{Q}_{\perp}) (\not{P}_{\perp} + \not{Q}_{\perp}) - \lambda \sigma q_{\perp}^2. \end{aligned} \quad (\text{D14})$$

Moreover, Eqs. (20) provide relations between \mathcal{A}_1 and \mathcal{A}_2 and \mathcal{A}_3 so that by introducing the variables

$$x = \frac{\mathbf{p}_{\perp}^2}{eB}, \quad y = \frac{\mathbf{p}'_{\perp}^2}{eB}, \quad (\text{D15})$$

so that

$$\begin{aligned}
\hat{\Gamma}^{(\lambda, \sigma)} = & - \left[(\not{P}_{\parallel} - \not{Q}_{\parallel}) (\not{P}_{\parallel} + \not{Q}_{\parallel}) + 2m\not{P}_{\parallel} + m^2 \right] [1 + \partial_x \partial_y - \gamma^1 \gamma^2 (\partial_x - \partial_y)] \mathcal{J}_1^{(\lambda, \sigma)}(p, p') \\
& - \left[(\not{P}_{\parallel} - \not{Q}_{\parallel}) (\not{P}_{\perp} + \not{Q}_{\perp}) + m (\not{P}_{\perp} + \not{Q}_{\perp}) \right] (1 - \gamma^1 \gamma^2 \partial_x) (1 - \partial_y^2) \mathcal{J}_1^{(\lambda, \sigma)}(p, p') \\
& - \left[(\not{P}_{\perp} - \not{Q}_{\perp}) (\not{P}_{\parallel} + \not{Q}_{\parallel}) + m (\not{P}_{\perp} - \not{Q}_{\perp}) \right] (1 + \gamma^1 \gamma^2 \partial_y) (1 - \partial_x^2) \mathcal{J}_1^{(\lambda, \sigma)}(p, p') \\
& - (\not{P}_{\perp} - \not{Q}_{\perp}) (\not{P}_{\perp} + \not{Q}_{\perp}) (1 - \partial_x^2) (1 - \partial_y^2) \mathcal{J}_1^{(\lambda, \sigma)}(p, p') \\
& - \lambda \sigma [1 + \partial_x \partial_y - \gamma^1 \gamma^2 (\partial_x - \partial_y)] \mathcal{J}_2^{(\lambda, \sigma)}(p, p') - \lambda \sigma (1 - \partial_x^2) (1 - \partial_y^2) \mathcal{J}_3^{(\lambda, \sigma)}(p, p'), \tag{D16}
\end{aligned}$$

where

$$\mathcal{J}_1^{(\lambda, \sigma)}(p, p') \equiv \int \frac{d^3 q}{(2\pi)^3} \mathcal{A}_1(p + \lambda q) \mathcal{A}_1(p' + \sigma q), \tag{D17a}$$

$$\mathcal{J}_2^{(\lambda, \sigma)}(p, p') \equiv \int \frac{d^3 q}{(2\pi)^3} q_{\parallel}^2 \mathcal{A}_1(p + \lambda q) \mathcal{A}_1(p' + \sigma q), \tag{D17b}$$

and

$$\mathcal{J}_3^{(\lambda, \sigma)}(p, p') \equiv \int \frac{d^3 q}{(2\pi)^3} \mathbf{q}_{\perp}^2 \mathcal{A}_1(p + \lambda q) \mathcal{A}_1(p' + \sigma q), \tag{D17c}$$

In order to simplify the integrals \mathcal{J}_i , we shall use the analytical expression for $\mathcal{A}_1(k)$ Eq. (A24) (details in Appendix A):

$$\begin{aligned}
\mathcal{A}_1(k) = & \frac{i e^{-\mathbf{k}_{\perp}^2 / eB}}{2eB} \exp \left[-\frac{i\pi (k_{\parallel}^2 - m^2)}{2eB} \right] \\
& \times \Gamma \left(-\frac{k_{\parallel}^2 - m^2}{2eB} \right) U \left(-\frac{k_{\parallel}^2 - m^2}{2eB}, 0, \frac{2\mathbf{k}_{\perp}^2}{eB} \right). \tag{D18}
\end{aligned}$$

1. The integral \mathcal{J}_1

We shall consider the integral:

$$\mathcal{J}_1^{(\lambda, \sigma)}(p, p') = \int \frac{d^3 q}{(2\pi)^3} \mathcal{A}_1(p + \lambda q) \mathcal{A}_1(p' + \sigma q) \tag{D19}$$

For the case $(\lambda, \sigma) = (-1, -1)$ we change the integration variables as follows

$$\begin{aligned}
p' - q &= q' + Q \\
p - q &= q' - Q \tag{D20}
\end{aligned}$$

in what follows, we shall use q instead of q' . For notation simplicity, we shall define the parameters

$$\begin{aligned}
a &= -\frac{\mathcal{D}_{\parallel}(q_3 + Q_{\parallel})}{2eB}, \\
a' &= -\frac{\mathcal{D}_{\parallel}(q_3 - Q_{\parallel})}{2eB}, \tag{D21}
\end{aligned}$$

and we shall use the identity

$$\Gamma(a)U(a, \epsilon, z) = \frac{1}{a}M(a, \epsilon, z) + \Gamma(-1 + \epsilon)zM(1 + a, 2, z), \tag{D22}$$

together with for $\epsilon \rightarrow 0^+$

$$\Gamma(-1 + \epsilon) = \frac{-1}{\epsilon} + \gamma_e - 1 + O(\epsilon), \tag{D23}$$

where $\gamma_e = 0.577$ is the Euler-Mascheroni constant. Also, given that there is a strong exponential damping in the integral, we consider the expansion of the Kummer function for small values of its argument, that is given by

$$M(a, b, z) = 1 + \frac{a}{b}z + O(z^2), \tag{D24}$$

so that, after removing the divergences, we end up with the integral

$$\begin{aligned}
\mathcal{J}_1^{(-, -)}(p, p') &= \left(\frac{i}{2eB} \right)^2 e^{-\frac{2\mathbf{Q}_{\perp}^2}{eB}} \int_{-\infty}^{\infty} \frac{dq_3}{2\pi} e^{i\pi(a+a')} \int_0^{\infty} \frac{d^2 q_{\perp}}{(2\pi)^2} e^{-\frac{2\mathbf{q}_{\perp}^2}{eB}} \Gamma(a)U \left(a, \epsilon, \frac{(q + Q)_{\perp}^2}{eB} \right) \Gamma(a')U \left(a', \epsilon, \frac{(q - Q)_{\perp}^2}{eB} \right) \\
&= \frac{1}{(2\pi)^3} \left(\frac{i}{2eB} \right)^2 e^{-\frac{2\mathbf{Q}_{\perp}^2}{eB}} \int_{-\infty}^{\infty} dq_3 e^{i\pi(a+a')} \\
&\times \int_0^{\infty} d^2 q_{\perp} e^{-\frac{2\mathbf{q}_{\perp}^2}{eB}} \left[\frac{1}{a} + \frac{\gamma_e - 1}{eB} (\mathbf{q}_{\perp}^2 + 2\mathbf{Q}_{\perp} \cdot \mathbf{q}_{\perp} + \mathbf{Q}_{\perp}^2) \right] \left[\frac{1}{a'} + \frac{\gamma_e - 1}{eB} (\mathbf{q}_{\perp}^2 - 2\mathbf{Q}_{\perp} \cdot \mathbf{q}_{\perp} + \mathbf{Q}_{\perp}^2) \right] \tag{D25}
\end{aligned}$$

Let us focus into the integrand. At order $\mathcal{O}(\mathbf{q}_\perp^2)$, we have:

$$\begin{aligned} & \left[\frac{1}{a} + \frac{\gamma_e - 1}{eB} (\mathbf{q}_\perp^2 + 2\mathbf{Q}_\perp \cdot \mathbf{q}_\perp + \mathbf{Q}_\perp^2) \right] \left[\frac{1}{a'} + \frac{\gamma_e - 1}{eB} (\mathbf{q}_\perp^2 - 2\mathbf{Q}_\perp \cdot \mathbf{q}_\perp + \mathbf{Q}_\perp^2) \right] \\ & \simeq \frac{1}{aa'} + \frac{(\gamma_e - 1)\mathbf{q}_\perp^2}{eB} \left(\frac{1}{a} + \frac{1}{a'} + \frac{2(\gamma_e - 1)\mathbf{Q}_\perp^2}{eB} \right) + \frac{(\gamma_e - 1)(2\mathbf{Q}_\perp \cdot \mathbf{q}_\perp)}{eB} \left(\frac{1}{a'} - \frac{1}{a} \right) \\ & - \frac{(\gamma_e - 1)^2(2\mathbf{Q}_\perp \cdot \mathbf{q}_\perp)^2}{eB^2} + \frac{(\gamma_e - 1)\mathbf{Q}_\perp^2}{eB} \left(\frac{1}{a} + \frac{1}{a'} \right). \end{aligned} \quad (\text{D26})$$

Then, by defining:

$$z \equiv \frac{2\mathbf{q}_\perp^2}{eB}, \quad (\text{D27})$$

such that

$$\begin{aligned} \mathbf{Q}_\perp \cdot \mathbf{q}_\perp &= |\mathbf{Q}_\perp| |\mathbf{q}_\perp| \cos \theta = \sqrt{\frac{eB}{2}} |\mathbf{Q}_\perp| z^{1/2} \cos \theta, \\ d^3q &= dq_3 d^2q_\perp = \frac{eB}{4} dz d\theta dq_3, \end{aligned} \quad (\text{D28})$$

where after angular integration we obtain:

$$\begin{aligned} \mathcal{J}_1^{(-,-)}(p, p') &= \frac{1}{(2\pi)^3} \frac{\pi eB}{2} \left(\frac{i}{2eB} \right)^2 e^{-\frac{2\mathbf{Q}_\perp^2}{eB}} \\ &\times \int_{-\infty}^{\infty} dq_3 e^{-\frac{i\pi}{2eB} (\mathcal{D}_\parallel(q_3 + Q_\parallel) + \mathcal{D}_\parallel(q_3 - Q_\parallel))} \\ &\times \int_0^\infty dz e^{-z} \left\{ \frac{1}{aa'} + \frac{(\gamma_e - 1)}{2} \left(\frac{1}{a} + \frac{1}{a'} \right) z \right. \\ &\left. + \frac{(\gamma_e - 1)\mathbf{Q}_\perp^2}{eB} \left(\frac{1}{a} + \frac{1}{a'} \right) \right\}. \end{aligned} \quad (\text{D29})$$

Performing the integration over z :

$$\begin{aligned} \mathcal{J}_1^{(-,-)}(p, p') &= -\frac{eB}{4\pi^2} e^{-\frac{2\mathbf{Q}_\perp^2}{eB}} e^{-\frac{i\pi(\mathbf{Q}_\parallel^2 - m^2)}{eB}} \\ &\times \int_{-\infty}^{\infty} dq_3 \frac{\exp\left(\frac{i\pi}{2eB} q_3^2\right)}{\left(Q_\parallel^2 - q_3^2 - m^2 + i\epsilon\right)^2 - 4Q_3^2 q_3^2} \\ &\times \left[1 - \frac{(\gamma_e - 1)\mathbf{Q}_\perp^2 (Q_\parallel^2 - q_3^2 - m^2)}{(eB)^2} \right. \\ &\left. - \frac{(\gamma_e - 1)(Q_\parallel^2 - q_3^2 - m^2)}{eB} \right] \end{aligned} \quad (\text{D30})$$

In what follows, we shall set the external 3-momenta to zero, except for the presence of the Q_\perp factors, those we shall keep in order to use this expression as a generating function.

Then:

$$\begin{aligned} \mathcal{J}_1^{(-,-)}(p, p') &= -\frac{eB}{4\pi^2} e^{-\frac{2\mathbf{Q}_\perp^2}{eB}} \\ &\times \int_{-\infty}^{\infty} \frac{dq_3}{(q_3^2 + m^2 + i\epsilon)^2} \exp\left[\frac{i\pi}{2eB} (q_3^2 + m^2)\right] \\ &\times \left[1 + \frac{(\gamma_e - 1)\mathbf{Q}_\perp^2 (q_3^2 + m^2)}{(eB)^2} + \frac{(\gamma_e - 1)(q_3^2 + m^2)}{eB} \right] \end{aligned} \quad (\text{D31})$$

Now, from the well-known results

$$\begin{aligned} &\int_{-\infty}^{\infty} \frac{dx}{x^2 + m^2} \exp\left[\frac{i\pi}{2b} (x^2 + m^2)\right] \\ &= \frac{\pi}{m} \left[1 - (1 - i)C\left(\frac{m}{\sqrt{b}}\right) - (1 + i)S\left(\frac{m}{\sqrt{b}}\right) \right], \end{aligned} \quad (\text{D32a})$$

and

$$\begin{aligned} &\int_{-\infty}^{\infty} \frac{dx}{(x^2 + m^2)^2} \exp\left[\frac{i\pi}{2b} (x^2 + m^2)\right] \\ &= \frac{(1 - i)\pi}{2\sqrt{b}m^2} e^{\frac{i\pi m^2}{2b}} \\ &+ \frac{(b + i\pi m^2)\pi}{2bm^3} \left[1 - (1 - i)C\left(\frac{m}{\sqrt{b}}\right) - (1 + i)S\left(\frac{m}{\sqrt{b}}\right) \right], \end{aligned} \quad (\text{D32b})$$

where $C(x)$ and $S(x)$ are the cosine and sine Fresnel integrals, respectively. From the property:

$$C(x) + iS(x) = \sqrt{\frac{\pi}{2}} \frac{1 + i}{2} \text{erf}\left(\frac{1 - i}{\sqrt{2}} x\right), \quad (\text{D33})$$

we have:

$$\begin{aligned} \mathcal{J}_1^{(-,-)}(p, p') &= -\frac{eB}{4\pi^2} e^{-\frac{2\mathbf{Q}_\perp^2}{eB}} \left\{ \frac{(1 - i)\pi}{2\sqrt{eB}m^2} e^{\frac{i\pi m^2}{2eB}} \right. \\ &+ \frac{(eB + i\pi m^2)\pi}{2(eB)m^3} \left[1 - \sqrt{\frac{\pi}{2}} \text{erf}\left(\frac{1 - i}{\sqrt{2}} \frac{m}{\sqrt{eB}}\right) \right] \\ &\left. + \frac{\pi(\gamma_e - 1)}{(eB)m} \left(1 + \frac{\mathbf{Q}_\perp^2}{eB} \right) \left[1 - \sqrt{\frac{\pi}{2}} \text{erf}\left(\frac{1 - i}{\sqrt{2}} \frac{m}{\sqrt{eB}}\right) \right] \right\}, \end{aligned} \quad (\text{D34})$$

where $\text{erf}(x)$ is the error function.

2. The integral \mathcal{J}_2

For this integral, note that $q_{\parallel}^2 = -q_3^2$, so that after integration over z we get:

$$\begin{aligned} \mathcal{J}_2^{(-,-)}(p, p') &= \frac{eB}{4\pi^2} e^{-\frac{2\mathbf{Q}_{\perp}^2}{eB}} \\ &\times \int_{-\infty}^{\infty} dq_3 \frac{q_3^2}{(q_3^2 + m^2 + i\epsilon)^2} \exp \left[\frac{i\pi}{2eB} (q_3^2 + m^2) \right] \\ &\times \left[1 + \frac{(\gamma_e - 1)\mathbf{Q}_{\perp}^2 (q_3^2 + m^2)}{(eB)^2} + \frac{(\gamma_e - 1)(q_3^2 + m^2)}{eB} \right] \end{aligned} \quad (\text{D35})$$

By using:

$$\begin{aligned} &\int_{-\infty}^{\infty} dx \frac{x^2}{x^2 + m^2} \exp \left[\frac{i\pi}{2b} (x^2 + m^2) \right] \\ &= (1 + i)e^{\frac{i\pi m^2}{2b}} \sqrt{b} \\ &- m\pi \left[1 - (1 - i)C \left(\frac{m}{\sqrt{b}} \right) - (1 + i)S \left(\frac{m}{\sqrt{b}} \right) \right], \end{aligned} \quad (\text{D36a})$$

and

$$\begin{aligned} &\int_{-\infty}^{\infty} dx \frac{x^2}{(x^2 + m^2)^2} \exp \left[\frac{i\pi}{2b} (x^2 + m^2) \right] \\ &= \frac{(i - 1)\pi}{\sqrt{b}} e^{\frac{i\pi m^2}{2b}} \\ &+ \frac{(ib + \pi m^2)\pi}{2bm} \left[-i + (1 + i)C \left(\frac{m}{\sqrt{b}} \right) - (1 - i)S \left(\frac{m}{\sqrt{b}} \right) \right], \end{aligned} \quad (\text{D36b}) \quad \text{for } n = 1, 2, 3.$$

we get:

$$\begin{aligned} \mathcal{J}_2^{(-,-)}(p, p') &= \frac{eB}{4\pi^2} e^{-\frac{2\mathbf{Q}_{\perp}^2}{eB}} \\ &\times \left\{ \frac{(i - 1)\pi}{\sqrt{eB}} e^{\frac{i\pi m^2}{2eB}} \right. \\ &+ \frac{(eB - i\pi m^2)\pi}{2(eB)m} \left[1 - \sqrt{\frac{\pi}{2}} \operatorname{erf} \left(\frac{1 - i}{\sqrt{2}} \frac{m}{\sqrt{eB}} \right) \right] \\ &\left. - \frac{m\pi(\gamma_e - 1)}{(eB)} \left(1 + \frac{\mathbf{Q}_{\perp}^2}{eB} \right) \left[1 - \sqrt{\frac{\pi}{2}} \operatorname{erf} \left(\frac{1 - i}{\sqrt{2}} \frac{m}{\sqrt{eB}} \right) \right] \right\} \end{aligned} \quad (\text{D37})$$

3. The integral \mathcal{J}_3

By following the same procedure, it is straightforward to obtain at order $\mathcal{O}(z)$:

$$\begin{aligned} \mathcal{J}_3^{(-,-)}(p, p') &= -\frac{eB^2}{8\pi^2} e^{-\frac{2\mathbf{Q}_{\perp}^2}{eB}} \left\{ \frac{(1 - i)\pi}{2\sqrt{eB}m^2} e^{\frac{i\pi m^2}{2eB}} \right. \\ &+ \frac{(eB + i\pi m^2)\pi}{2(eB)m^3} \left[1 - \sqrt{\frac{\pi}{2}} \operatorname{erf} \left(\frac{1 - i}{\sqrt{2}} \frac{m}{\sqrt{eB}} \right) \right] \\ &\left. + \frac{\pi(\gamma_e - 1)\mathbf{Q}_{\perp}^2}{(eB)^2 m} \left[1 - \sqrt{\frac{\pi}{2}} \operatorname{erf} \left(\frac{1 - i}{\sqrt{2}} \frac{m}{\sqrt{eB}} \right) \right] \right\}, \end{aligned} \quad (\text{D37})$$

Moreover, it is easy to check that

$$\begin{aligned} \mathcal{J}_n^{(+,+)}(p, p') &= \mathcal{J}_n^{(-,-)}(p, p') \\ \mathcal{J}_n^{(+,-)}(p, p') &= \mathcal{J}_n^{(-,+)}(p, p') = \mathcal{J}_n^{(-,-)}(p, p') \mid_{Q \rightarrow P}, \end{aligned} \quad (\text{D38})$$