

On the Possibility of Demonstrating Confinement in Non-Supersymmetric Theories by Deforming Confining Supersymmetric Theories

Michael Dine,

*Santa Cruz Institute for Particle Physics and Department of Physics,
University of California, Santa Cruz,
Santa Cruz, CA, USA*

E-mail: mdine@ucsc.edu

ABSTRACT: We review arguments that supersymmetric QCD with N colors and $N_f \geq 0$ massive quarks confines. We explain that this remains the case for small soft breakings and small supersymmetric masses for any quarks. These results continue to hold at large N . We then consider these theories with large soft breakings – real QCD – describing the problem in terms of the direct computation of the linear potential between a heavy quark and antiquark. We provide naive arguments that confinement persists, and then give a more general treatment. The features required for the naive arguments are generally believed to hold at large N . Extending them to finite N requires only mild assumptions.

Contents

1	Introduction: Arguing for Confinement in Non-Supersymmetric Theories Obtained by Deformation of Confining Supersymmetric Theories	1
2	Confinement in Supersymmetric Gauge Theories	2
3	Lessons from the CP^N Models in Two Dimensions	3
4	The Essence of the Argument for Confinement in Four Dimensions	6
5	Confinement in the Large N Expansion With Broken Supersymmetry	9
6	Theories with N_f Light Quarks At Large and Finite N	9
7	The QCD Description of the η' at Large N	9
8	θ Dependence	10
9	Conclusion: Even With Large Soft Breakings, SUSY QCD is Confining	11

1 Introduction: Arguing for Confinement in Non-Supersymmetric Theories Obtained by Deformation of Confining Supersymmetric Theories

Confinement and the existence of a mass gap are readily established features of many $N = 1$ supersymmetric gauge theories. This is the case, in particular, for gauge theories without matter fields and gauge theories with only massive matter. Exploiting tools such as holomorphy, it is possible to study many aspects of the problem, such as the role of monopole condensation[1, 2]. It is natural to ask whether these results can be extended to demonstrate confinement in non-supersymmetric theories. One could start, for example, by taking a supersymmetric gauge theory without matter, and then introducing a gaugino mass term. Due to the mass gap, for small gaugino mass both the mass gap and confinement persist. The question is: is there a deconfining phase transition as m_λ becomes large compared to the QCD scale?

The analogous problem can be addressed in CP^N theories in two dimensions at large N [3], where one can readily check that there is no phase transition as one passes from the supersymmetric to the non-supersymmetric theory. These theories are confining and gapped in both limits. Confinement is a phenomenon of the far infrared theory, and the infrared theory is the same in both limits, up to a rescaling of lengths. In four dimensional gauge theories, we don't have a simple (continuum) picture of confinement, so the problem is more challenging. Again, if we deform the supersymmetric theories with small soft

breaking terms, confinement will survive, as readily follows from the existence of a mass gap. But for large soft breakings, i.e. to describe real QCD, more is required to establish confinement. In [4], arguments were put forward that if one considered theories with anomaly mediated supersymmetry breaking, there would be no phase transition. These arguments relied on working only to linear order in soft breakings, and considering the monopole condensate, but as explained in [5], non-linear effects (in the low energy dynamics) become important (and difficult to control) as we approach the QCD scale. So while the result is plausible, its foundations are shaky.

But focussing on the heavy quark-antiquark potential (Wilson line), we will argue that a deconfining transition can be ruled out. Framed this way, confinement follows from rather mild assumptions about the theory in the far infrared. In large N , these assumptions reduce to properties which are generally assumed to be true of the planar diagram expansion. In this paper, we give simple arguments for confinement from this perspective. Possible loopholes are enumerated, but are not very plausible.

The rest of this note is organized as follows. We briefly review confinement in supersymmetric theories in the next section. In section 3, we then remark on confinement in the two-dimensional CP^N models at large N , with and without supersymmetry, noting certain features which we expect to be relevant to four dimensions. Turning to four dimensions, we give naive arguments for confinement and then a more general argument for confinement for large m_λ in section 4. We explore the question of large N theories in four dimensions with small and large soft breakings in section 5. Limiting consideration to CP conserving values of parameters reduces the problem of confinement to standard assumptions about large N . In section 6, we extend the analysis to theories with light quarks. In section 7, we compare these assumptions with those governing the η' at large N [6]. In section 8, we remark on possible behaviors with CP violating parameters. Our conclusion, that confinement holds in the theories with large soft breakings, is presented in section 9.

2 Confinement in Supersymmetric Gauge Theories

Proving confinement in non-abelian gauge theories without supersymmetry remains one of the great challenges of theoretical physics. What is remarkable is that, with $N = 1$ supersymmetry, one can establish confinement in a range of gauge theories, in a variety of ways. These include:

1. Following Seiberg and Witten[1, 2], one can prove confinement by exhibiting monopole condensation in theories which can be obtained by deformation of $N = 2$ theories.
2. Starting with $SU(N)$ supersymmetric QCD with $N - 1$ massive flavors, one can perform a systematic computation, for small masses, of the spectrum and other features. Using holomorphy, one can study the theory in various regimes of quark mass, establishing, among other results, gaugino condensation and the existence of a mass gap in the $N_F = 0$ theory[7].

In the case of the $SU(N)$ $N = 2$ theory, before one adds a mass term for the adjoint field, ϕ , there are N points on the moduli space of vacua where monopoles become

massless:

$$\langle u \rangle \equiv \langle \text{Tr } \phi^2 \rangle = u_0 e^{\frac{2\pi i}{N}}. \quad (2.1)$$

These transform into one another under a Z_N symmetry. Adding a large mass for the adjoint, the theory at low energies is supersymmetric $SU(N)$ QCD without matter, and these states correspond to the N possible values of the gaugino condensate, $\langle \lambda\lambda \rangle \propto e^{\frac{2\pi i}{N}}$. With a gaugino mass, the degeneracy is lifted, and all but one of the states are unstable. For large enough gaugino mass, we would expect that all but the ground state are highly unstable.

So confinement and the existence of a mass gap are common in supersymmetric field theories. Notably, they are features of certain large N theories, such as the $SU(N)$ supersymmetric gauge theory without matter, or with number of flavors, N_f , much less than N , providing additional tools to study non-supersymmetric theories.

3 Lessons from the CP^N Models in Two Dimensions

The CP^N models at large N in two dimensions[3] are instructive for the case of four dimensional QCD. These theories are confining and gapped. While confinement itself is much simpler in two than in four dimensions, other aspects of the problem, particularly the scaling of terms in a four dimensional effective action, have close parallels. What is similar about the two theories is that confinement is a problem of the far infrared. As a result, it is natural to focus on a Wilsonian action with cutoff, Λ_w , well below the scale of the theory, Λ (the dynamically generated scale in the two dimensional case, Λ_{QCD} , the QCD scale in the four dimensional case). Confinement is a feature of this very low energy effective theory.

For the non-supersymmetric CP^N models, the microscopic degrees of freedom are a set of N complex scalars on the CP^N manifold; more conveniently, they are a set of scalar fields along with a scalar lagrange multiplier and a vector field with no kinetic term at the classical level. The scalars gain mass of order Λ , the dynamically generated scale of the theory. In the far infrared, the the scalar degrees of freedom are integrated out and the Wilsonian action is written in terms of a gauge field, A_μ , and its field strength $F_{\mu\nu}$. In the non-supersymmetric case, the low scale effective action can be computed in the loop expansion, with successive loops suppressed by by powers of $1/N$. At one loop, one has a series of operators of the form $F_{\mu\nu}^{2n}$, with coefficients scaling as $N\Lambda^{-4n+2}$. But setting aside the explicit one loop computation, we can understand this scaling by the requirement that if one perform a one loop computation with the effective action, it yields corrections to the effective action smaller than, or at least not parameterically larger than, the action itself. Here we can take the ultraviolet cutoff to be (at most) of order Λ . So, for example, writing the coefficient of $F_{\mu\nu}^4$ as $\frac{1}{M^6}$, the one loop correction to the $F_{\mu\nu}^2$ coupling behaves as

$$N^0 \frac{1}{M^6} \int^\Lambda d^2 k \frac{k^4}{k^2} \sim N^0 \frac{\Lambda^4}{M^6}. \quad (3.1)$$

This is (up to a factor of N) of order the leading term in the action, provided $M \sim \Lambda$. We can place a sharper bound if we insist on a particular behavior of the $1/N$ expansion; the result, needless to say, is in agreement with the explicit computation.

The hierarchy of operators is also consistent with confinement. F^2 is the most relevant operator of the low energy theory and the F^2 term in the action makes the dominant contribution to the potential at large distances. For $\Lambda_w \ll \Lambda$, the coefficient of F^2 does not evolve with decreasing Λ_w . It is crucial that the infrared theory is confining for any value of the coefficient of the $F_{\mu\nu}^2$ term, $\frac{1}{e_0^2}$. Changing e_0 , the coefficient of the most relevant operator in the infrared theory, just rescales lengths in the large distance theory. In four dimensions, we will argue for similar behavior.

The theory exhibits non-trivial θ dependence[3]. We will consider θ -dependence later in four dimensions, where our arguments for confinement will hold on the CP conserving subspace of couplings.

The supersymmetric version of the theory microscopically includes fermionic partners of the scalar fields. It is convenient to work with the theory in the form described in [3], in which form it is relatively easy to explicitly break the supersymmetry. In terms of the scalars, n_i fermions ψ^i , bosonic auxiliary fields $\lambda, \sigma, \pi, A_\mu$ and fermionic auxiliary fields $\chi, \bar{\chi}$, the lagrangian is:

$$\mathcal{L} = \frac{N}{g^2} [n_i^* (\partial_\mu - iA_\mu - i\pi)^2 n^i + \bar{\psi}_i (i\partial - \mathcal{A}) \psi^i - \bar{\psi}_i m_\psi \psi^i - \frac{1}{2} (\sigma^2 + \pi^2) - \alpha m_\psi \sigma \quad (3.2)$$

$$- \frac{1}{\sqrt{2}} \bar{\psi}^i (\sigma + i\pi\gamma_5) \psi^i + \lambda (n_i^* n^i - 1) + \bar{\chi} n_i \psi^i + \bar{\psi}_i n^i \chi^*].$$

Here we have included explicit supersymmetry violating terms, proportional to m_ψ . The term linear in σ is necessary for the renormalizability of the theory. Including m_ψ , we can perform a one loop computation at large N (the leading effect) of the effective action. We can look for stationary points of this action. We can obtain, as in [3], the values of σ, π , and λ at large N by computing the effective action at one loop. Setting $\pi = A_\mu = 0$, and differentiating the action with respect to λ and σ yields two equations:

$$N \int \frac{d^2k}{(2\pi)^2} \frac{1}{k^2 + \lambda} = \frac{N}{g^2}; \quad (3.3)$$

$$N \int \frac{d^2k}{(2\pi)^2} \frac{\sqrt{2}(\frac{\sigma}{\sqrt{2}} + m_\psi)}{k^2 + (\frac{\sigma}{\sqrt{2}} + m_\psi)^2} = N(\frac{\sigma}{g^2}). \quad (3.4)$$

Here we have noted that while in general there is a different renormalization of the fermion and boson kinetic terms, at large N they are the same, so the same factor of $1/g^2$ appears in both 3.3 and 3.4. The first equation is solved by

$$\lambda \equiv \Lambda^2 = M^2 e^{-\frac{4\pi}{g^2}}. \quad (3.5)$$

The second is solved, for small m_ψ , by $\sigma^2 = 2\lambda$. We can think of the scale, Λ , as a fixed physical scale, analogous to Λ_{QCD} , measuring $m_F = m_\psi + \frac{\sigma}{\sqrt{2}}$ and other scales relative to this one. For large m_ψ , $\frac{\sigma}{\sqrt{2}} + m_\psi \sim m_\psi$. We can understand this if we think of integrating

out the fermion, with mass much greater than Λ , In this case, the effects on the dynamics at low energies are small .

At scales below the scale Λ , the system, in either limit, is described by the gauge field A_μ . This field acquires a non-trivial kinetic term at one loop. It does not create physical excitations, but it does give rise to a potential between static charged sources, which grows linearly at large distances; equivalently, it yields an area law for large Wilson loops. The vacuum polarization is given by:

$$\pi_{\mu\nu}(q) = (\pi_{\mu\nu})_B(q) + (\pi_{\mu\nu})_F(q) \quad (3.6)$$

where

$$(\pi_{\mu\nu})_B(q) = (g_{\mu\nu}q^2 - q_\mu q_\nu)\pi_B(q^2); \quad \pi_B(q^2) = \frac{1}{4\pi} \frac{1}{m_B^2} \quad q^2 \ll m_B^2. \quad (3.7)$$

$$(\pi_{\mu\nu})_F(q) = (g_{\mu\nu}q^2 - q_\mu q_\nu)\pi_F(q^2); \quad \pi_F(q^2) = \frac{1}{4\pi} \frac{1}{m_F^2} \quad q^2 \ll m_F^2. \quad (3.8)$$

Each of these terms has the structure of a local interaction,

$$-\frac{1}{4e_0^2}F_{\mu\nu}^2, \quad (3.9)$$

with the bosonic term dominating for large m_F^2 .

In either limit, the low energy theory is the $U(1)$ gauge theory. We can think of a Wilsonian effective action, obtained by integrating out physics at scales above Λ_w . Λ_w can be arbitrarily small; we still find a linear potential. This potential arises from the most relevant operator, $F_{\mu\nu}^2$, with coefficient $-\frac{1}{4e_0^2}$. This is a reflection of the fact that the linear potential is a phenomena of the far infrared in this theory – arbitrarily far. Less relevant operators yield contributions to the potential suppressed by powers of $(\Lambda R)^{-1}$. The coupling e_0^2 is proportional to $\Lambda = \Lambda(\Lambda_w)$, which we can now think of as a coupling in the Wilsonian theory. For small Λ_w , examining equation 3.3, we see that Λ goes to a constant as $\Lambda_w \rightarrow 0$ as $(\frac{\Lambda_w}{\Lambda})^2$. As a result, the coefficient of the separation R in the linear potential tends to a constant. We will argue for similar behavior in four dimensional gauge theory.

We might have anticipated this result by noting, first, that the supersymmetric theory is confining. Because the theory is gapped, we expect this to remain true in the presence of a small m_ψ . For small $\Lambda_w \ll m_F$, $\Lambda_w \ll \Lambda$, the theory is a $U(1)$ gauge theory, characterized by the coefficient of the most relevant operator. As a result, the theory is confining for all Λ_w , and therefore for all m_F . The argument could have failed if there were two most relevant operators of the same dimension in the small Λ_w theory. Also, if the most relevant operator was marginal, it could conceivably have a fixed point for some value of the coupling. In four dimensions, we don't know the structure of the theory in the far infrared, but we can attempt to make an argument for confinement along these lines.

It is worth noting that confinement occurs in a regime where there are no physical excitations of the system, so Greens functions of gauge-invariant local operators fall to zero exponentially rapidly with growing separation. Such correlators can be represented by contact terms in the low energy theory; the value of this contact term can be thought

of as information which supplements the Wilsonian action. The Wilson loop is different; not being a correlation function of a local operator, it does not satisfy a simple spectral relation. Its large distance behavior can be computed from the Wilsonian action with arbitrarily small Λ_w . There are lessons here for four dimensions. In particular, for small m_λ , the theory with $N_f = 0$ is gapped, confining, and has non-trivial θ dependence, implying a non-vanishing contact term in the $F\tilde{F}$ two-point function. This is necessary to give the correct θ -dependence, and should be thought of as a feature which supplements the Wilsonian action. Similarly, the large radius Wilson loop obeys an area law, which can be computed from the small Λ_w Wilsonian action. The question is: do these features change as we take m_λ large? What might change in this case as the theory flows to small Λ_w ?

4 The Essence of the Argument for Confinement in Four Dimensions

Turning now to QCD, to address confinement, rather than attacking the question of the fate of the magnetic monopole condensate at large m_λ , we focus on the heavy quark-antiquark potential. Taking R as the quark anti-quark separation (and q the corresponding momentum transfer), for confinement we can study arbitrarily large R (small q) for general m_λ . In particular, we can take $R \gg m_\lambda^{-1}$. In the broken supersymmetric gauge theory without matter fields, for sufficiently small m_λ , the potential is linear.

Because confinement is a feature of the far infrared, it makes sense to consider a Wilsonian effective action with scale Λ_w . Typically in thinking about this action in the continuum, one takes $\Lambda_w \gg \Lambda_{QCD}$, so that the action can be computed in a weak coupling approximation. This would be appropriate to the problem of large m_λ , where integrating out the gaugino would yield a theory well described in this way. But the case of which we have knowledge is that of small m_λ , so we'll be interested in considering a range of Λ_w , including Λ_w well below the QCD scale. The problem is then not a weak coupling one.

Consider, then, the softly broken $N = 1$ gauge theory without matter fields as a function of $x = \frac{m_\lambda}{\Lambda_{QCD}}$. We will restrict, first, to $\theta = 0$. More precisely we will take the θ parameter in the langrangian to be zero, m_λ to be real, and the phase of the gaugino condensate or of the field u of equation 2.1 to be zero. For small values of x , $x < x_0$, for some x_0 , we know that the low energy theory confines and is gapped. Confinement is the result of some dynamics at scales well below the mass gap. The CP^N model illustrates this, as we have seen in section 3. There it is the gauge field in the far infrared which is responsible for confinement. This field, at low energies, does not create any of the particles of the theory, which are massive.

Integrating out physics above a scale Λ_w slightly below the gluino mass, we are left with an effective theory with Wilsonian cutoff, Λ_w slightly less than m_λ . The pure gauge theory is confining for some $\frac{\Lambda_w}{\Lambda_{QCD}} < y_0$, where y_0 is slightly less than x_0 . Now provided that the theory flows smoothly from larger y to smaller, confinement is established.

For this to fail, the theory obtained by flow from large Λ_w to small Λ_w , i.e. the theory obtained in the far infrared by integrating out a heavy gluino must be different than that

obtained by integrating out a light gluino. This does not seem particularly plausible, and we will now argue that the flow is smooth.

We first give two somewhat naive arguments for confinement for large m_λ . Each relies on the fact of confinement for small gluino mass. The first starts by examining the diagrams of perturbation theory, and noting that a substantial enhancement of the individual contributions at large distances is required to obtain a linear potential. Such enhancements take the form of (powers of) logarithms of q . For $q \ll \Lambda_w$ and $m_\lambda \sim \Lambda_w$, large logs only arise from diagrams with gluons.¹ So only gauge bosons play a role in confinement. The confining potential, taken to be the sum of these diagrams, is then a feature of the pure gauge theory, with Wilsonian cutoff Λ_w and with a coupling $g(\Lambda_w)$ in the effective action. The coefficient of the linear term in the potential, k , depends only on the scale parameter of the effective theory with small Λ_w , $\tilde{\Lambda}_{QCD}$, $k = a\tilde{\Lambda}_{QCD}^2$. Here

$$\tilde{\Lambda}_{QCD} = \Lambda_w e^{-\int_{g_{ref}}^{g(\Lambda_w)} \frac{dg'}{\beta(g')}}. \quad (4.1)$$

In this equation, g_{ref} is some fixed reference value of the coupling. The form of this result will not change as we increase Λ_w in the bosonic theory, as the effect is just a change in the overall scale. So confinement survives at large m_λ . One might question the reliability of the guidance provided by Feynman diagrams, but other known non-perturbative effects, such as instantons, are also dominated at large distances by the purely gluonic contributions. The usual large N picture, as we will discuss further below, assumes that features of QCD, at least at large N , are reflected in the Feynman diagram expansion. As for Λ in the CP^N model, $\tilde{\Lambda}$ tends to a constant as $\Lambda_w \rightarrow 0$, as follows from the fact of confinement in the small m_λ theory.

We can reframe this argument, referring to a naive understanding of the operators of QCD, an understanding which holds more rigorously at large N . For this argument for confinement, we will make three assumptions:

1. We can write an effective action, at large distances, in terms of gauge fields $A_\mu (F_{\mu\nu})$. This is plausible, and, at large N , a basic feature.
2. Operators have roughly their naive dimensions. In particular, there is a single marginal operator, $F_{\mu\nu}^2$, and associated dimensionless coupling, $g^2(\Lambda_w)$.
3. The theory with small m_λ and $\Lambda_w < m_\lambda$, defines the infrared sector of actual QCD, for some values of the parameters of the effective lagrangian. This is the case for large m_λ , where integrating out the gaugino generates a change in the marginal coupling and leaves over some irrelevant operators. In computing the potential at very large distances. With our assumption above the situation is similar, in that the contributions of irrelevant operators to the potential at large distances will be suppressed by powers of $(R\Lambda_w)^{-1}$.

¹Here we are working, as suggested above, with an effective theory where the gluinos have been integrated out, giving contact terms in the action.

4. There are no zeros of the beta function. As will see, and we can state more sharply in this formulation, we can actually rule out this possibility.

With these assumptions, the theory in the far infrared has only one parameter, a mass or inverse length scale, $\tilde{\Lambda}_{QCD}(\Lambda_w)$. To define this, we proceed somewhat heuristically, writing an effective action which governs very low scale phenomena ($\mu = m_\lambda, \Lambda_w$) as

$$\mathcal{L} = -\frac{1}{4g^2(\mu)}F_{\mu\nu}^2 + \dots, \quad (4.2)$$

and we have dropped higher dimension terms suppressed by powers of μ . Arguments of the type we gave for the CP^N models bound these higher dimension terms by powers of $1/\Lambda_w$ or $\frac{1}{\tilde{\Lambda}_{QCD}}$. Indeed we can again define the scale $\tilde{\Lambda}_{QCD}$ as in equation 4.1.

We know that confinement occurs for small m_λ . We know, as well, that the coefficient of the linear potential tends to a constant as $\Lambda_w \rightarrow 0$. As a consequence, $\tilde{\Lambda}_{QCD}$ also tends to a constant as $\Lambda_w \rightarrow 0$. This is similar to the behavior of the CP^N theory. Confinement, then, depends only on the length parameter $\tilde{\Lambda}_{QCD}$. As we increase m_λ , $\tilde{\Lambda}_{QCD}$ will change, but confinement will occur, simply associated with this different scale.

We can argue against an obstruction as we vary $g(\Lambda_w)$ by noting that a break in the flow would correspond to an infrared fixed point at some intermediate value of Λ_w . If this were all there is, then the theory in this region, starting with some Λ_w would flow to this fixed point, which is inconsistent with the fact that it is gapped and confining. So there must be at least two such points (and in general an even number). But now the theory would have an ultraviolet fixed point starting at small Λ_w , which is not consistent with the fact that the theory with small m_λ is asymptotically free. At the level of planar Feynman diagrams, we can make diagrammatic versions of these statements, and we will do that in the next section.

The argument for confinement can be stated more generally. Confinement follows if:

1. In the low Λ_w theory, there is a single, most relevant operator. Correspondingly, for a given Λ_w , there is a single parameter which characterizes the flow, $g(\Lambda_w)$. $g(\Lambda_w)$ flows with Λ_w according to the renormalization group.
2. There are no fixed points in $g(\Lambda_w)$.

If these statements are true, then the system flows from large Λ_w to smaller Λ_w where the system is known to be confining. Equivalently, we can define a parameter $\tilde{\Lambda}_{QCD}$, and again the confining parameter is independent of Λ_w .

If there were two most relevant operators, then as one increased m_λ , it is conceivable that one could follow a trajectory which at some point becomes non-confining. For small values of $g(\Lambda_w)$, there is only one such operator. Suppose *there are* two or more most relevant operators (degenerate in dimension) at larger $g(\Lambda_w)$, i.e. for $g(\Lambda_w) > g_0$. This would appear to imply that actual QCD is a multiparameter theory, both in the supersymmetric case and the non-supersymmetric one. Given all we know about the supersymmetric theory, the former possibility appears highly unlikely. Alternatively, two operators might

cross in dimension at a particular value of the coupling. But in order that one theory track the higher point, one the lower point, the system would have to be finely tuned.

In this more general framework, the possibility of a fixed point (or multiple fixed points) can again be ruled out by facts we know about the theory. In particular, if there were a single fixed point, then starting with small m_λ and $\Lambda_w < m_\lambda$, the theory would flow at large distances to the fixed point. But this is inconsistent with the fact of confinement in the small m_λ theory. If there were a pair of fixed points (more generally an even number of fixed points), we would also encounter inconsistency with known facts of the small Λ_w theory.

5 Confinement in the Large N Expansion With Broken Supersymmetry

These arguments for confinement are particularly compelling in the large N limit. The main reason one might question the Feynman diagram argument is that Feynman diagrams possibly might not capture the essence of the theory, but at large N , the planar diagrams *should* capture the theory's essential features. Also at large N , the assumptions underlying the second argument are believed to hold: the system is described, in particular, in terms of gluons and gluinos, and the arguments used to bound the coefficients of operators in the effective action (and correspondingly the dimensions of these operators) should hold.

6 Theories with N_f Light Quarks At Large and Finite N

We can extend these result to theories with N_f light quarks. Add to the action of this theory mass terms for the squarks and a mass for the gluino. For small values of the soft breaking parameters, one can find the ground state and calculate an effective action and the spectrum of the theory. The theory is gapped and confined. One can again ask what happens when the soft breakings are increased, eventually to values larger than the QCD scale. Once again, we can focus on the heavy quark-antiquark potential. At large distances, this is an infrared phenomenon, as before. Taking the fermion masses to be non-zero but small, then as we did for the light gluino, we can consider the far infrared theory, and argue that this theory is the one we have encountered in the theory with $N_f = 0$. Again this theory is described by a single parameter and is necessarily confining. At large N with $N_f \ll N$, we can simply repeat the arguments of the $N_f = 0$ theory.

7 The QCD Description of the η' at Large N

While often unstated, the assumption that the underlying degrees of freedom at large N in QCD in the far infrared are the gluon fields (and the dominance of planar diagrams) underlies the usual treatment of the η' at large N [6]. The analysis relies heavily on the anomaly equation,

$$\partial_\mu j^{\mu 5} = \frac{2N_f}{16\pi^2} F\tilde{F}, \quad (7.1)$$

or, in terms of the η' ,

$$f_\pi \partial^2 \eta' = \frac{2N_f}{16\pi^2} F\tilde{F}. \quad (7.2)$$

From this, one can compute the η' mass in terms of the leading contribution in $1/N$ to the two point function of the η' at distances of order a power of N times Λ_{QCD}^{-1} but smaller than a power of N times the QCD scale[6]. The result is expressed in terms of the correlator

$$U(k) = \langle (F(x)\tilde{F}(x) F(0)\tilde{F}(0))_k \rangle \quad (7.3)$$

at zero (small, by a power of N) momentum. All of this is written explicitly in terms of the degrees of freedom arising from A^μ .

Because the correlator is evaluated at zero momentum, and because of the observational fact that the theory is gapped, it is, as in the case of CP^N , controlled by a contact term

$$U(x) = K\delta(x). \quad (7.4)$$

This contact term can be computed, for example, by computing the θ^2 term in the action for a given Λ_w . Because of the infrared sensitivity of the computation and because the system is gapped, this term is invariant under changes of Λ_w . Contributions to $U(k)$ from operators of higher dimension than $F_{\mu\nu}^2$ in the Wilsonian action are suppressed by powers of $\frac{m_{\eta'}^2}{\Lambda^2} \sim N^{-1}$.

This is all quite parallel to the questions which arise for confinement: one can study the deep infrared (small Λ_w) and work with the degrees of freedom of the original continuum action. The dominant term in the action is again the naive leading term.

8 θ Dependence

Once we allow a non-zero θ , or equivalently treat m_λ as complex, a richer phase structure is possible. In particular, if we take m_λ and $\langle\lambda\lambda\rangle$ as real, we can introduce a $\theta F\tilde{F}$ term, and our lagrangian now has two couplings of dimension four:

$$\mathcal{L} = -\frac{1}{4g_{eff}^2(\Lambda_w)}F_{\mu\nu}^2 + \frac{\theta}{16\pi^2}F\tilde{F}. \quad (8.1)$$

Our discussion of the last section applies to the theory along the line $\theta = 0$. It established that there is confinement in this space. With non-zero θ , a more intricate phase structure arises, and there are trajectories which are sensitive to these phase changes. The supersymmetric gauge theory with small gaugino mass exhibits a branched structure, with phase transitions as one varies θ . Recall in particular that:

$$m_\lambda\langle\lambda\lambda\rangle \propto e^{i(\theta + \frac{2\pi k}{N})}. \quad (8.2)$$

As a result, the theory undergoes phase transitions as θ varies. It is quite possible that there are regions in the space $(g_{eff}(\Lambda_w), \theta)$ with non-zero θ which evolve under the renormalization group to unconfined regions[8].

9 Conclusion: Even With Large Soft Breakings, SUSY QCD is Confining

From these rather simple considerations, it appears that, by deforming supersymmetric QCD with large gaugino and squark masses, one obtains a theory which is gapped and confined. To summarize the basic argument:

1. The linear confining potential of the supersymmetric theories survives with small soft breakings, due to the presence of a mass gap.
2. The linear confining potential of the slightly broken supersymmetric theories can be computed at arbitrarily large distance.
3. One can work with a Wilsonian action with scale Λ_w small compared to the soft breakings and large compared to R^{-1} .
4. For small Λ_w and large R , only the gluon fields contribute to the potential.
5. The potential depends only on $\tilde{\Lambda}_{QCD}$, a parameter which depends in the usual renormalization group fashion on the value of Λ_w and the most relevant coupling. Increasing m_λ just results in a rescaling of lengths.

The last two points are basic features of the perturbative expansion, but it is hard to see how they could fail to hold more generally – at long enough distances, the gaugino mass should be irrelevant in the technical sense.

For large N , the analysis is rigorous to the extent that large N is defined as a sum of an infinite set of planar diagrams, resummed in some way. For finite N , a breakdown of the components of the basic argument would be quite bizarre. Confinement in the theory with large breaking of supersymmetry, then, seems an inevitable consequence of the fact that the supersymmetric theory is confining and that confinement is a feature of the (arbitrarily) far infrared.

Acknowledgments

We thank P. Draper, G. Festuccia, N. Seiberg and E. Witten for conversations and critical comments. This work was supported in part by U.S. Department of Energy grant No. DE-FG02-04ER41286.

References

- [1] N. Seiberg and E. Witten, *Electric - magnetic duality, monopole condensation, and confinement in $N=2$ supersymmetric Yang-Mills theory*, *Nucl. Phys. B* **426** (1994) 19–52, [[hep-th/9407087](#)]. [Erratum: *Nucl.Phys.B* 430, 485–486 (1994)].
- [2] N. Seiberg and E. Witten, *Monopoles, duality and chiral symmetry breaking in $N=2$ supersymmetric QCD*, *Nucl. Phys. B* **431** (1994) 484–550, [[hep-th/9408099](#)].
- [3] E. Witten, *Instantons, the Quark Model, and the $1/n$ Expansion*, *Nucl. Phys. B* **149** (1979) 285–320.

- [4] C. Csáki, A. Gomes, H. Murayama, and O. Telem, *Demonstration of Confinement and Chiral Symmetry Breaking in $SO(Nc)$ Gauge Theories*, *Phys. Rev. Lett.* **127** (2021), no. 25 251602, [[arXiv:2106.10288](#)].
- [5] M. Dine and Y. Yu, *Challenges to Obtaining Results for Real QCD from SUSY QCD*, [arXiv:2205.00115](#).
- [6] E. Witten, *Current Algebra Theorems for the $U(1)$ Goldstone Boson*, *Nucl. Phys. B* **156** (1979) 269–283.
- [7] N. Seiberg, *The Power of holomorphy: Exact results in 4-D SUSY field theories*, in *Particles, Strings, and Cosmology (PASCOS 94)*, pp. 0357–369, 5, 1994. [hep-th/9408013](#).
- [8] D. Gaiotto, A. Kapustin, Z. Komargodski, and N. Seiberg, *Theta, Time Reversal, and Temperature*, *JHEP* **05** (2017) 091, [[arXiv:1703.00501](#)].