

NON-PRINCIPAL T -DUALITY, GENERALIZED COMPLEX GEOMETRY AND BLOW-UPS

GIL R. CAVALCANTI AND ALDO WITTE

ABSTRACT. We extend the notion of T -duality to manifolds endowed with non-principal torus actions. The singularities of the torus action are controlled by a certain Lie algebroid, called the elliptic tangent bundle. Using this Lie algebroid, we explain how certain invariant generalized complex structures can be transported via T -duality. Along the way, we use the elliptic tangent bundle to define connections for these torus action, and give new insight to the classification of such actions by Haefliger-Salem.

Dedicated to the memory of Gilles Castel

CONTENTS

1. Introduction	2
1.1. T -duality	2
1.2. Blow-ups	3
1.3. Generalized complex structures	4
1.4. Organisation of the paper	4
1.5. Acknowledgements	5
2. Divisors and Lie algebroids; generalized complex structures and their lifts	5
2.1. Divisors	5
2.2. Lie algebroids and residue maps	6
2.3. Generalized complex structures	7
2.4. Lifting generalized complex structures	8
3. Classification of torus actions and connections	10
3.1. Haefliger–Salem classification	10
3.2. Torus actions, divisors and connections	11
3.3. Smoothness of the curvature	14
3.4. Čech–de Rham isomorphism	15
3.5. Connection approach to Haefliger-Salem class	16
4. The singular differential geometry of the correspondence space	17
5. Non principal T -duality	18
5.1. T -duality	18
5.2. Courant algebroids	19
6. Blow-ups	23
6.1. Blow-ups in generalized complex geometry	23
6.2. Blowing up for the elliptic tangent bundle	24
6.3. T -duality and blow-up	26
7. Examples	27
7.1. Basic examples	27
7.2. Non-zero residue elliptic symplectic structures	29
References	30

1. INTRODUCTION

In this paper we will study three topics: T -duality for torus actions with fixed points, blow-ups of elliptic divisors and will apply those two to the study of generalized complex structures. While each one of these has a life on its own, the interplay between them makes them all the more interesting.

1.1. T -duality. T -duality is an equivalence between quantum field theories with very different properties. Its origins trace back to a paper by Sathiapalan [19] where it was noted that different compactifications with S^1 -symmetry were related by, among other things, inversion of the radius of the circles. This relation is frequently referred to as taking “dual circles”, “exchange of tangent and cotangent directions” or “swap of winding and momentum”. The precise relationship between (local) T -dual Riemannian structures was first understood by Buscher in [6] and was developed further by Roček and Verlinde in [18]. A global notion of T -duality was formulated in [3, 4] as an equivalence between *principal torus bundles*. In [7] it was observed that T -duality can be understood as an isomorphism of Courant algebroids, therefore formalizing the operation “exchange of tangent and cotangent directions” and allowing for a systematic transport of geometric structures between T -dual spaces. Because this well established notion of T -duality only holds for principal torus bundles, we refer to it as *principal T -duality*.

Once we move away from principal circle bundles and start considering circle actions with fixed points problems quickly pop up. Namely, the inversion of radii, a hallmark of T -duality, implies that every fixed point on one side is matched by an open end of a manifold of the T -dual side. Performing T -duality this way amounts to removing the fixed points and only considering the resulting open manifold where the action is free. For example, taking the 2-sphere with rotation along the z -axis, the dual becomes an open cylinder with an incomplete metric. Versions of this phenomenon appear in several different places, most notably, when trying to produce SYZ-mirrors of Fano varieties [1, 11, 14]. Further, the strategy of “simply ignore fixed points” may leave one with the impression that we are missing important behaviour that takes place “at infinity”.

The main goal of the current article is to develop a framework for T -duality, which allows for torus actions which locally decompose as standard rotations of the plane. We call such actions *standard*. The new input that allows us to study these actions without compromising on compactness of the T -dual are *elliptic divisors* and *elliptic tangent bundles*. An elliptic divisor is an ideal, $I_{|D|}$, of smooth functions, which comes naturally associated to a standard torus actions. The elliptic tangent bundle is a Lie algebroid, $\mathcal{A}_{|D|}$, defined as the vector fields which preserve the elliptic divisor.

As with principal T -duality, T -duality for these actions will provide an isomorphism of Courant algebroids, but now using the elliptic tangent bundle:

Theorem 5.13. *Let M, \hat{M} be manifolds with standard T^k actions and let $\mathcal{A}_{|D|} \rightarrow M, \mathcal{A}_{|\hat{D}|} \rightarrow \hat{M}$ denote the induced elliptic tangent bundles. Let $H \in \Omega_{T^k}^3(\mathcal{A}_{|D|}), \hat{H} \in \Omega^3(\mathcal{A}_{|\hat{D}|})$ be closed invariant forms. If (M, H) and (\hat{M}, \hat{H}) are T -dual then there exists an isomorphism of Courant algebroids:*

$$(1.1) \quad (\mathcal{A}_{|D|} \oplus \mathcal{A}_{|D|}^*)/T^k \xrightarrow{\cong} (\mathcal{A}_{|\hat{D}|} \oplus \mathcal{A}_{|\hat{D}|}^*)/T^k.$$

This isomorphism of Courant algebroids allows us to transport geometric structures which are invariant under the torus action between on $\mathcal{A}_{|D|}$ over M and $\mathcal{A}_{|\hat{D}|}$ over \hat{M} . As we will see later, this has implications for T -duality of complex manifolds as well as further special classes of generalized complex manifolds.

Already in the case of principal torus bundles, T -duals may not exist [4]. In that case, the proof of existence is constructive and involves choices of connections and a switch between the Chern class of the torus bundle with a collection of cohomology classes determined by the background 3-form. To carry out a similar construction of T -duals for torus actions with fixed points, one first needs to develop the theory of connections and curvature for those actions.

1.1.1. Torus actions and connections. An integral part of this paper is the classification of torus actions on manifolds. For principal torus actions, the relevant datum is its characteristic class. For general torus actions, this question is studied by Haefliger and Salem in [13]. They show that the action is classified by its local model and a class in $\check{H}^2(M; \mathbb{Z}^k)$. The class in $\check{H}^2(M; \mathbb{Z}^k)$ has a geometric interpretation: it is the obstruction to finding a global cross-section for the quotient map $M \rightarrow M/T^k$.

Although the class of [13] resembles a Chern class, no connection approach exists. In this article we will show that one may use the elliptic tangent bundle to give a definition of connections for standard T^k -actions. The curvature of these connections then induces the class of Haefliger-Salem in the following way:

Proposition 3.25. *Let $\pi : M \rightarrow B$ be a manifold endowed with a standard T^k -action and let $\pi' : M' \rightarrow B$ be a manifold endowed with a locally equivalent action. Let Θ, Θ' be connection one-forms on M respectively M' . Then $[d\Theta - d\Theta'] \in H^2(A_{\partial B}; \mathfrak{t}^k)$ is representable by a smooth form, and descends to the Haefliger-Salem class of M' in $\check{H}^2(B; \mathfrak{t}^k)$.*

With the notions of connection and curvature at hand we can move on to produce a construction of a T -duals, therefore establishing conditions that ensure their existence (c.f. Theorem 5.8). The T -dual we construct has a T^k -action which is locally equivalent to original one. Yet we also observe that there are other T -duals to (M, H) which do not satisfy this property, hence the fixed point set gives rise to a new source of non-uniqueness of T -duals.

1.2. Blow-ups. Blow-ups come from complex/algebraic geometry as a method to resolve singularities, but have encountered applications in the study other geometric structures specially as a way to construct new examples [15, 10, 2].

The minimum amount of data needed to construct canonically the blow-up of a space at a point is a holomorphic ideal for that point [2]. As it is, holomorphic ideals, \mathcal{I}_D , are the heart and soul of complex log tangent bundles. Further, there is a close connection between complex log and elliptic tangent bundles: complex log tangent bundles are in correspondence with elliptic tangent bundles together with coorientations [8]. So, after a finite set of choices, there is a canonical blow-up construction for manifolds with elliptic divisors. Not only that but the blow-up itself inherits canonically an elliptic divisor and in many ways the blow-up manifold is indistinguishable from the original one.

Theorem 6.9. *Let $I_{|D|}$ be an elliptic divisor, and let $\mathcal{A}_{|D|}$ be the associated elliptic tangent bundle. Let $Y \subset D$ be a submanifold compatible with D in a suitable sense. Then after a choice of coorientations there is a canonical blow-up \tilde{M} of M along Y and there exists a natural elliptic divisor $I_{|\tilde{D}|}$ on \tilde{M} such that the blow-down map $p : \tilde{M} \rightarrow M$ provides a Lie algebroid submersion*

$$p : \mathcal{A}_{|\tilde{D}|} \rightarrow \mathcal{A}_{|D|}.$$

This means in particular that any structure on $\mathcal{A}_{|D|}$ can be pulled back to $\tilde{\mathcal{A}}_{|\tilde{D}|}$.

When M comes equipped with a torus action and $Y = \{*\}$ is a fixed point of this action, the blow-up \tilde{M} naturally inherits a torus action and we can compare the constructions of T -duality and blow-up:

Proposition 6.15. *T-duality and blowing up commute.*

This procedure will allow us to construct many examples of T -dual manifolds and transport geometric structures between them.

1.3. Generalized complex structures. As an illustration of how to transport structures between T -dual spaces, we will study how generalized complex structures behave in this framework. The easy result here is that Theorem 5.13 allows us to transport invariant structures between $\mathcal{A}_{|D|} \oplus \mathcal{A}_{|D|}^*$ and $\mathcal{A}_{|\hat{D}|} \oplus \mathcal{A}_{|\hat{D}|}^*$, which is directly analogous to what happens with principal T -duality. The new insight here is that this transport in some cases may happen in disguise and associated to a *lifting* question.

One interesting phenomenon we encounter is the following: Given a manifold with elliptic divisor $(M, \mathcal{I}_{|D|})$ and a generalized complex structure \mathcal{J} on M , under favourable conditions we can lift \mathcal{J} canonically to a generalized complex structure \mathcal{J}_A on $\mathcal{A}_{|D|}$. In some cases, this switch can be subtle enough that goes unnoticed. For example, if (M, I) is a complex manifold and $\mathcal{I}_{|D|}$ is the elliptic divisor associated to a complex hypersurface $D \subset M$, then I lifts to a complex structure on $\mathcal{A}_{|D|}$. At the other extreme, the type of structure that lifts to a symplectic structure on the elliptic tangent bundle is a stable generalized complex structure on M [8, 9] and this switch is very visible because stable structures would normally be recognised as singular symplectic structures on M .

Pairing this remark with the framework we developed for T -duality, we produce examples with the following characteristics: we look for a manifold with a torus action compatible elliptic divisor $(M, \mathcal{I}_{|D|})$ together with a generalized complex structure \mathcal{J} on M such that \mathcal{J} lifts to a generalized complex structure, $\mathcal{J}_{\mathcal{A}_{|D|}}$ on $\mathcal{A}_{|D|}$. We then transport $\mathcal{J}_{\mathcal{A}_{|D|}}$ to a generalized complex structure $\hat{\mathcal{J}}_{\mathcal{A}_{|\hat{D}|}}$ on $\mathcal{A}_{|\hat{D}|}$ via T -duality and study when $\hat{\mathcal{J}}_{\mathcal{A}_{|\hat{D}|}}$ is itself a lift of a generalized complex structure $\hat{\mathcal{J}}$ on \hat{M} . When this happens, it makes sense to call $\hat{\mathcal{J}}$ the structure T -dual to \mathcal{J} . We also notice that the condition that a structure can be lifted to $\mathcal{A}_{|D|}$ is a very restrictive one and therefore examples satisfying all the hypotheses above are rare. Yet, once one example is found, we can use the theory of blow-ups within generalized complex geometry [2] to produce infinite families of T -dual structures satisfying the lifting conditions described above.

Stable generalized complex structures can be viewed as symplectic structures on $\mathcal{A}_{|D|}$, and can hence be transported in this manner. There is however an intriguing difference with T -duality for principal torus actions: A generalized complex structure on M which lifts to $\mathcal{A}_{|D|}$ will always induce a generalized complex structure on the T -dual Lie algebroid $\mathcal{A}_{|\hat{D}|} \rightarrow \hat{M}$, however it might not induce a generalized complex structure on M . We will, for instance, encounter complex structures on the elliptic tangent bundle on manifolds which are not almost complex.

1.4. Organisation of the paper. This paper is organized as follows. In Section 2 we will recall the notions of elliptic divisors, and their associated Lie algebroids. Furthermore, we recall how stable generalized complex structures can be viewed as symplectic structures on Lie algebroids. More generally, we discuss lifting generalized complex structures to Lie algebroids. In Section 3 we recall the classification of torus actions by Heafiger-Salem. Afterwards we introduce the notion of a connection for a standard torus action, and show how these can be used to understand this classification. In Section 4 we study singular fibre products, which we will encounter in the description of T -duality. In Section 5 we introduce the the notion of non-principal T -duality, using the Lie algebroids from Section 2. In Section 6 we show that the elliptic tangent bundle interacts very well with (generalized) complex blow-ups, and show that T -dualizing commutes with blowing-up. Finally, in Section 7 we discuss concrete examples.

1.5. Acknowledgements. The second author was supported by the NWO through the UGC Graduate Programme (Netherlands), and through the FWO and FNRS under EOS project G0H4518N, and FWO Project G083118N (Belgium). The authors would like to thank Maarten Mol and Álvaro del Pino for useful discussions. The authors are grateful to the organizers of the workshop “Generalized Geometry in Interaction”, Madrid 2022, where a preliminary version of this article was presented.

2. DIVISORS AND LIE ALGEBROIDS; GENERALIZED COMPLEX STRUCTURES AND THEIR LIFTS

As we mentioned in the introduction, the main objective of this paper is to extend T -duality to torus actions with fixed points without giving up on compactness of the spaces involved. The object that will allow us to accomplish this task are Lie algebroids, specifically Lie algebroids associated to divisors as introduced in [8, 9]. Not only will these allow us to lift torus actions, but also we will arrive at the very natural problem of lifting geometric structures from the underlying manifold to the Lie algebroid.

In this section give the first step towards our goal and recall the theory of divisors and Lie algebroids (for more details, see [8, 9]). We also introduce several relevant geometric structures, including generalized complex structures, and their lifts to these Lie algebroids

2.1. Divisors. Inspired by divisors in algebraic geometry, we will make use of the following notion to deal with singularities of geometric structures:

Definition 2.1. A **real (resp. complex) divisor** on M is a locally principal ideal $I \subset C^\infty(M; \mathbb{R})$ (respectively $C^\infty(M; \mathbb{C})$) which is locally generated by functions with a nowhere dense zero-set. \diamond

Definition 2.2. Let (M, I_M) and (N, I_N) be manifolds together with divisors. A smooth map $\varphi : M \rightarrow N$ is a **morphism** of divisors if $\varphi^* I_N = I_M$. It is said to be a **diffeomorphism** of divisors if φ is moreover a diffeomorphism. \diamond

We will encounter three types of divisors in this article:

Definition 2.3. Let M^n be a smooth manifold.

- A **real log divisor** on M is a real divisor, I_Z , such that for each $p \in M$ there are coordinates $(x_1, \dots, x_n) : U \rightarrow \mathbb{R}^n$ centred at p and an integer k such that

$$I_Z|_U = \langle x_1 \cdot \dots \cdot x_k \rangle.$$

- A **complex log divisor** on M is a complex divisor I_D , such that for each $p \in M$ there are coordinates $(z_1, \dots, z_k, x_{2k+1}, \dots, x_n) : U \rightarrow \mathbb{C}^k \times \mathbb{R}^{n-2k}$ centred at p such that

$$I_D|_U = \langle z_1 \cdot \dots \cdot z_k \rangle.$$

- An **elliptic divisor** on M is a real divisor $I_{|D|}$, such that for each $p \in M$ there are coordinates $(x_1, \dots, x_n) : U \rightarrow \mathbb{R}^n$ centred at p and an integer k such that

$$I_{|D|}|_U = \langle r_1^2 \cdot \dots \cdot r_k^2 \rangle,$$

$$\text{with } r_i^2 = x_{2i-1}^2 + x_{2i}^2.$$

In these three cases, the number k is the **multiplicity** of p . The vanishing set of the ideal is called the **vanishing locus**. We call each of these divisors **smooth** if their vanishing locus is an embedded smooth submanifold, that is, if the multiplicity of each point is either 0 or 1. \diamond

In general, the vanishing locus is only (the image of) an immersed submanifold with transverse intersections, and is naturally stratified by the multiplicity of its points. It is convenient to introduce some notation to refer to the different strata of the vanishing locus, so if we let, say, D , denote the vanishing locus of any of the divisors above, the set of points with multiplicity

at least k will be denoted by $D(k)$ and the set of points with multiplicity at precisely k will be denoted by $D[k]$. Each of the $D(k)$ is itself an immersed submanifold, and hence we may speak of:

Definition 2.4. Let $I_{|D|}$ be an elliptic divisor, and let $\iota : N \rightarrow D(k)$ be an immersion. The image of a connected component of N is called an **irreducible component** of $D(k)$. \diamond

Given a complex log divisor I_D , one may consider its **associated elliptic divisor** defined by the condition $I_{|D|} \otimes \mathbb{C} = I_D \otimes \bar{I}_D$. In this case, $D[1]$ is automatically co-oriented by the differential of any local generator of I_D . Conversely, given an elliptic divisor $I_{|D|}$ for which $D[1]$ is co-oriented, there exists a unique complex log divisor I_D inducing $I_{|D|}$ and the given co-orientation.

2.2. Lie algebroids and residue maps. Using the Serre-Swann theorem, one may show that vector fields preserving the divisors from Definition 2.3 define vector bundles and the natural identification of sections of these vector bundles with vector fields makes these vector bundles into Lie algebroids:

Definition 2.5. Let M be a manifold, and $I_Z, I_D, I_{|D|}$ a real log, complex log, respectively elliptic divisor on M . Vector fields preserving these ideals define Lie algebroids:

- $\mathcal{A}_Z \rightarrow TM$, the **real log-tangent bundle**.
- $\mathcal{A}_D \rightarrow T_{\mathbb{C}}M$, the **complex log-tangent bundle**.
- $\mathcal{A}_{|D|} \rightarrow TM$, the **elliptic tangent bundle**. \diamond

As the divisors under consideration have local normal forms, so do have the associated Lie algebroids:

Remark 2.6. In the coordinates of Definition 2.3, we have:

- $\Gamma(\mathcal{A}_Z) = \langle x_1 \partial_{x_1}, \dots, x_k \partial_{x_k}, \partial_{x_{k+1}}, \dots, \partial_{x_n} \rangle$.
- $\Gamma(\mathcal{A}_D) = \langle z_1 \partial_{z_1}, \partial_{\bar{z}_1}, \dots, z_k \partial_{z_k}, \partial_{\bar{z}_k}, \partial_{x_{2k+1}}, \dots, \partial_{x_n} \rangle$.
- $\Gamma(\mathcal{A}_{|D|}) = \langle r_1 \partial_{r_1}, \partial_{\theta_1}, \dots, r_k \partial_{r_k}, \partial_{\theta_k}, \partial_{x_{2k+1}}, \dots, \partial_{x_n} \rangle$,

with $r_i \partial_{r_i} = x_{2i-1} \partial_{x_{2i-1}} + x_{2i} \partial_{x_{2i}}$ and $\partial_{\theta_i} = x_{2i-1} \partial_{x_{2i}} - x_{2i} \partial_{x_{2i-1}}$. \diamond

For each of the Lie algebroids above, we obtain a corresponding de Rham complex on the exterior power of the dual vector bundle. The relation between complex and elliptic tangent bundles translates into the existence of maps between their corresponding de Rham complexes. Indeed, given a complex log divisor, I_D , and associated elliptic divisor, $I_{|D|}$, taking the imaginary part of a complex log form is a map $\Im^* : (\Omega^\bullet(\mathcal{A}_D), d_{\mathcal{A}_D}) \rightarrow (\Omega^\bullet(\mathcal{A}_{|D|}), d_{\mathcal{A}_{|D|}})$.

There are several residue maps associated to these Lie algebroids, obtained by taking the coefficients of certain singular generators. We quickly recall the ones we need in this article. Given a smooth elliptic divisor, with D co-orientable, it is explained in [8] that there are residue maps:

$$\begin{aligned} \text{Res}_q : (\Omega^\bullet(\mathcal{A}_{|D|}), d_{\mathcal{A}_{|D|}}) &\rightarrow (\Omega^{\bullet-2}(D), d), & \alpha &\mapsto \iota_D^*(\iota_{r\partial_r \wedge \partial_\theta} \alpha) \\ \text{Res}_r : (\Omega^\bullet(\mathcal{A}_{|D|}), d_{\mathcal{A}_{|D|}}) &\rightarrow (\Omega^{\bullet-1}(S^1 ND), d), & \alpha &\mapsto \iota_{r\partial_r} \alpha, \end{aligned}$$

where $S^1 ND$ is the sphere bundle of the normal bundle of D .

Let $I_{|D|}$ be a general elliptic divisor with $D[1]$ co-orientable. The inclusion $\iota_{M \setminus D(2)} : M \setminus D(2) \rightarrow M$, pulls the elliptic divisor $I_{|D|}$ back to a smooth elliptic divisor with vanishing locus $D[1]$. Hence we may define residue maps by composition:

$$\begin{aligned} \text{Res}_q : \Omega^\bullet(\mathcal{A}_{|D|}) &\rightarrow \Omega^{\bullet-2}(D[1]), & \alpha &\mapsto \iota_{D[1]}^*(\iota_{r\partial_r \wedge \partial_\theta}(\iota_{M \setminus D(2)}^* \alpha)) \\ \text{Res}_r : \Omega^\bullet(\mathcal{A}_{|D|}) &\rightarrow \Omega^{\bullet-1}(S^1 ND[1]), & \alpha &\mapsto \iota_{r\partial_r}(\iota_{M \setminus D(2)}^* \alpha) \end{aligned}$$

There exists several higher order residue maps, however defining these very quickly becomes a combinatorial task. As we are mainly interested in two-forms, we will define:

Definition 2.7. Let $I_{|D|}$ be an elliptic divisor, $\omega \in \Omega^2(\mathcal{A}_{|D|})$ and let $p \in D[2]$. Using the coordinates from Remark 2.6 we define

$$\text{Res}_{r_i r_j} \omega(p) = \omega_p(r_i \partial_{r_i}, r_j \partial_{r_j}), \quad \text{Res}_{r_i \theta_j} \omega(p) = \omega_p(r_i \partial_{r_i}, \partial_{\theta_j}), \quad \text{Res}_{\theta_i \theta_j}(\omega)(p) = \omega_p(\partial_{\theta_i}, \partial_{\theta_j}). \quad \diamond$$

Note that these residues are only well-defined up to sign. Similarly, one may define residue maps for the real log tangent bundle. If I_Z is a smooth real log divisor, we may define

$$\text{Res} : \Omega^\bullet(\mathcal{A}_Z) \rightarrow \Omega^{\bullet-1}(Z), \quad \alpha \mapsto \iota_Z^*(\iota_{x \partial_x} \alpha).$$

Again, for a general real log divisor we may define $\text{Res}^1 : \Omega^\bullet(\mathcal{A}_Z) \rightarrow \Omega^{\bullet-1}(Z[1])$ by precomposing with the pullback $M \setminus D(2) \hookrightarrow M$. The higher residues are again somewhat involved to define. For two-forms we define:

$$(2.1) \quad \text{Res}_{x_i x_j}^2(\omega)(p) = \omega_p(x_i \partial_{x_i}, x_j \partial_{x_j}), \quad p \in Z[2].$$

2.3. Generalized complex structures. In this section we recall the definition of generalized complex structures and focus our attention on the class of stable generalized complex structures called stable. We will recall that these structures are in correspondence with symplectic structures on the elliptic tangent bundle.

Given a manifold M , we form $\mathbb{T}M = TM \oplus T^*M$. This bundle is endowed with the natural symmetric pairing given by evaluation of forms on vectors.

Definition 2.8. A **generalized complex structure** on a manifold M , is a pair (\mathbb{J}, H) , where $H \in \Omega^3(M)$ is a closed real 3-form and \mathbb{J} is a complex structure on $\mathbb{T}M$ orthogonal with respect to the natural pairing and whose $+i$ -eigenbundle, $L \subset \mathbb{T}_{\mathbb{C}}M$ is involutive with respect to the Courant–Dorfmann bracket:

$$[[X + \xi, Y + \eta]]_H = [X, Y] + \mathcal{L}_X \eta - \iota_Y d\xi + \iota_X \iota_Y H, \quad X + \xi, Y + \eta \in \mathbb{T}M. \quad \diamond$$

Given a generalized complex structure one may decompose it as a block matrix with respect to the splitting of $\mathbb{T}M$. The skew-symmetry of \mathbb{J} ensures this decomposition is of the form

$$\mathbb{J} = \begin{pmatrix} J & \pi_{\mathbb{J}}^\sharp \\ \sigma^\flat & -J^* \end{pmatrix}, \quad J \in \text{End}(TM), \sigma \in \Omega^2(M), \pi_{\mathbb{J}} \in \mathfrak{X}^2(M).$$

The bivector $\pi_{\mathbb{J}}$ is Poisson. Two generalized complex structures $(\mathbb{J}_1, H_1), (\mathbb{J}_2, H_2)$ are said to be **gauge equivalent**, if there exists a two-form $B \in \Omega^2(M)$ such that $dB = H_1 - H_2$ and

$$\mathbb{J}_2 = \begin{pmatrix} 1 & 0 \\ B^\flat & 1 \end{pmatrix} \mathbb{J}_1 \begin{pmatrix} 1 & 0 \\ -B^\flat & 1 \end{pmatrix}.$$

If two generalized complex structures $\mathbb{J}_1, \mathbb{J}_2$ are gauge equivalent, then $\pi_{\mathbb{J}_1} = \pi_{\mathbb{J}_2}$.

The Clifford action of $X + \xi \in \mathbb{T}_{\mathbb{C}}M$ on $\rho \in \wedge^n T_{\mathbb{C}}M$ is defined via $(X + \xi) \cdot \rho = \iota_X \rho + \xi \wedge \rho$. Given a generalized complex structure, one may define its **canonical bundle** $K \subset \wedge^{\text{top}} T_{\mathbb{C}}M$ via

$$L = \{u \in T_{\mathbb{C}}M : u \cdot K = 0\}.$$

The dual bundle K^* has a natural section s , called the **anticanonical** section which is defined via $s : \Gamma(K) \rightarrow C^\infty(M; \mathbb{C}), \rho \mapsto \rho_0$.

Definition 2.9. A generalized complex structure (\mathbb{J}, H) on M is called **stable** if $s(\Gamma(K)) \subset C^\infty(M; \mathbb{C})$ defines a complex log divisor. \diamond

Example 2.10. Let (M, J, π) be a holomorphic Poisson manifold with $\pi = \pi_R + i\pi_I$ and denote by $\pi_I^\sharp: T^*M \rightarrow TM$ the map associated to π_I . Then

$$\mathbb{J}_{J,\pi} := \begin{pmatrix} -J & 4\pi_I^\sharp \\ 0 & J^* \end{pmatrix}$$

defines a generalized complex structure. If furthermore $\wedge^n \pi(\Gamma(\det(T^*M))) \subset C^\infty(M; \mathbb{C})$ defines a complex log divisor, $\mathbb{J}_{J,\pi}$ defines a stable generalized complex structure. Holomorphic Poisson structures with these properties are called *log symplectic*. \triangle

These generalized complex structures can be described as symplectic structures on Lie algebroids:

Theorem 2.11 ([9]). *There is a one-to-one correspondence between gauge equivalence classes $[(\mathbb{J}, H)]$ of stable generalized complex structures on M and symplectic forms $\omega \in \Omega^2(\mathcal{A}_{|D|})$ on the elliptic tangent bundle with $D[1]$ co-orientable and satisfying the following equations:*

- $\text{Res}^q(\omega) = 0$
- $\text{Res}_{\theta_i r_j} \omega = \text{Res}_{r_i \theta_j} \omega$
- $\text{Res}_{r_i r_j} \omega = -\text{Res}_{\theta_i \theta_j} \omega$

Given (\mathbb{J}, H) , the associated $\omega \in \Omega^2(\mathcal{A}_{|D|})$ is given by $\pi_{\mathbb{J}}^{-1}$.

2.4. Lifting generalized complex structures. Lifting Poisson structures to Lie algebroids is a well-established field of study. However, lifts of generalized complex and Dirac structures to Lie algebroids have not yet received much attention. As we will see, the questions of which structures lift to Lie algebroids and which structures descend from Lie algebroids arise naturally.

2.4.1. Severa class. Let us first shortly study Courant algebroids where we replace the role of the tangent bundle by a general Lie algebroid. Given any Lie algebroid \mathcal{A} and $H \in \Omega^3(\mathcal{A})$ closed one may define the standard \mathcal{A} -Courant algebroid, in the usual manner. We set $\mathbb{A} = \mathcal{A} \oplus \mathcal{A}^*$ and for $X + \xi, Y + \eta \in \Gamma(\mathcal{A} \oplus \mathcal{A}^*)$.

$$\begin{aligned} \langle X + \xi, Y + \eta \rangle &= \frac{1}{2}(\eta(X) + \xi(Y)), \\ \llbracket X + \xi, Y + \eta \rrbracket_H &= [X, Y] + \mathcal{L}_X \eta - \iota_Y d\xi + \iota_X \iota_Y H. \end{aligned}$$

This notion, which is a particular case of a Lie bialgebroid, can be axiomatised as follows:

Definition 2.12. An **exact \mathcal{A} -Courant algebroid** is a vector bundle extension

$$(2.2) \quad 0 \rightarrow \mathcal{A}^* \xrightarrow{\pi^*} \mathbb{A} \xrightarrow{\pi} \mathcal{A} \rightarrow 0,$$

together with a non-degenerate pairing $\langle \cdot, \cdot \rangle$ on \mathbb{A} , and a bracket $\llbracket \cdot, \cdot \rrbracket$ on $\Gamma(\mathbb{A})$ satisfying:

- $\llbracket e_1, \llbracket e_2, e_3 \rrbracket \rrbracket = \llbracket \llbracket e_1, e_2 \rrbracket, e_3 \rrbracket + \llbracket e_2, \llbracket e_1, e_3 \rrbracket \rrbracket$
- $\pi(\llbracket e_1, e_2 \rrbracket) = [\pi(e_1), \pi(e_2)]_{\mathcal{A}}$
- $\llbracket e_1, f e_2 \rrbracket = f \llbracket e_1, e_2 \rrbracket + \mathcal{L}_{\rho(\pi(e_1))}(f) e_2$
- $\mathcal{L}_{\rho(\pi(e_1))}(\langle e_2, e_3 \rangle) = \langle \llbracket e_1, e_2 \rrbracket, e_3 \rangle + \langle e_2, \llbracket e_1, e_3 \rrbracket \rangle$
- $\llbracket e_1, e_1 \rrbracket = \pi^* d_{\mathcal{A}} \langle e_1, e_1 \rangle$.

\diamond

Given an exact \mathcal{A} -Courant algebroid, the composition $\rho \circ \pi : \mathbb{A} \rightarrow TM$ together with $\langle \cdot, \cdot \rangle$ and $\llbracket \cdot, \cdot \rrbracket$ will give \mathbb{A} the structure of a non-exact Courant algebroid.

Just as for ordinary exact courant algebroids, isotropic splittings $s : \mathcal{A} \rightarrow \mathbb{A}$ of (2.2) are classified by closed three-forms $H \in \Omega^3(\mathcal{A})$ appearing as the curvature

$$\iota_X \iota_Y H = s^*[s(X), s(Y)].$$

The difference between two splittings $s - s'$, defines a two-form $B \in \Omega^2(\mathcal{A})$ via $(s - s')(X) = \iota_X B$. Consequently $[H] \in H^3(\mathcal{A})$ is independent of the isotropic splitting and determines the \mathcal{A} -Courant algebroid structure completely. Hence, this cohomology class is the natural extension of the **Severa-class** of exact Courant algebroids.

2.4.2. Lifting Dirac structures. The answer to the question on how to lift structures takes different forms depending on what one is trying to lift. Here we will focus on Dirac and generalized complex structures, whose definitions on \mathbb{A} is completely analogous to the definition on TM :

Definition 2.13. Let $\mathcal{A} \rightarrow M$ be a Lie algebroid. An **\mathcal{A} -Dirac structure** is a subbundle $E_{\mathcal{A}} \subset \mathbb{A}$ together with a three-form $H \in \Omega_{\text{cl}}^3(\mathcal{A})$, such that $E_{\mathcal{A}}$ is isotropic with respect to the pairing and involutive with respect to the bracket on \mathbb{A} .

An **\mathcal{A} -generalized complex structure** is a pair $(\mathcal{J}_{\mathcal{A}}, H_{\mathcal{A}})$ where $H_{\mathcal{A}} \in \Omega_{\text{cl}}^3(\mathcal{A})$ and $\mathcal{J}_{\mathcal{A}}$ is an orthogonal complex structure $\mathcal{J}_{\mathcal{A}}$ on \mathbb{A} for which $\mathbb{A}^{1,0}$, the $+i$ -eigenbundle, defines an $\mathbb{A}_{\mathbb{C}}$ -Dirac structure. \diamond

Definition 2.14. Let $\mathcal{A} \rightarrow M$ be a Lie algebroid. An \mathcal{A} -Dirac structure $(E_{\mathcal{A}}, H_{\mathcal{A}})$ is a **lift** of a Dirac structure (E, H) on M if $H_{\mathcal{A}} = \rho_{\mathcal{A}}^* H$ and $\mathfrak{F}_{\rho}(E_{\mathcal{A}}) = E$. \diamond

Here \mathfrak{F}_{ρ} denotes the Dirac pushforward via the anchor map.

This notion matches the notion of a lift of a Poisson structure, when that is considered as a Dirac structure. Indeed, given a Lie algebroid $\mathcal{A} \rightarrow M$, a Poisson structure $\pi_{\mathcal{A}} \in \Gamma(\wedge^2 \mathcal{A})$ and a Poisson structure $\pi \in \mathfrak{X}^2(M)$ one can readily see that $\rho_* \pi_{\mathcal{A}} = \pi$ if and only if $E_{\pi_{\mathcal{A}}} = \text{graph}(\pi_{\mathcal{A}}) \subset \mathbb{A}$ is a lift of $E_{\pi} = \text{graph}(\pi)$.

For generalized complex structures, however, working only with the push-forward map is too restrictive, so the notion of lift requires some tweaking.

Definition 2.15. Let $\rho: \mathcal{A} \rightarrow TM$ be a Lie algebroid over M

- $\tilde{X} + \tilde{\xi} \in \mathbb{A}$ is **related** to $X + \xi \in TM$ if $\rho(\tilde{X}) = X$ and $\rho^* \xi = \tilde{\xi}$.
- A generalized complex structure $(\mathcal{J}_{\mathcal{A}}, H_{\mathcal{A}})$ on \mathbb{A} is a **lift** of a generalized complex structure (\mathcal{J}, H) on M if whenever $\tilde{X} + \tilde{\xi} \in \mathbb{A}$ is related to $X + \xi \in TM$, $\mathcal{J}_{\mathcal{A}}(\tilde{X} + \tilde{\xi})$ is related to $\mathcal{J}(X + \xi)$. \diamond

Example 2.16. In the situation above, if $J: TM \rightarrow TM$ and $J_{\mathcal{A}}: \mathcal{A} \rightarrow \mathcal{A}$ are complex structures we can promote them to generalized complex structures and ask when $J_{\mathcal{A}}$ is a lift of J . It turns out this happens if and only if the following diagram commutes

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{J_{\mathcal{A}}} & \mathcal{A} \\ \downarrow \rho & & \downarrow \rho \\ TM & \xrightarrow{J} & TM \end{array}$$

which agrees with what one would like to call the lift of a complex structure. \triangle

Remark 2.17. In this paper we will work with Lie algebroids \mathcal{A} with dense isomorphism locus, U . In this case, we see that $\mathcal{J}_{\mathcal{A}}$ is a lift of \mathcal{J} on U if and only if $\mathcal{J}_{\mathcal{A}}|_U = (\rho|_U) \circ \mathcal{J} \circ (\rho|_U)^{-1}$. If that holds, by continuity $\mathcal{J}_{\mathcal{A}}$ is related to \mathcal{J} in all of M . Therefore we see that the question of whether \mathcal{J} admits a lift to a Lie algebroid with dense isomorphism locus becomes the question of existence of a smooth extension of $(\rho|_U) \circ \mathcal{J} \circ (\rho|_U)^{-1}$ beyond the isomorphism locus. \diamond

Example 2.18. It is also good to keep in mind that there are (generalized) complex structures on Lie algebroids, which do arise as lifts of structures on M . Consider, for example, the elliptic

tangent bundle $\mathcal{A}_{|D|}$ on \mathbb{C}^2 , associated to the ideal $\langle r_1^2 r_2^2 \rangle$. The following defines a complex structure on $\mathcal{A}_{|D|}$:

$$\begin{aligned} z_1 \partial_{z_1} &\mapsto z_2 \partial_{z_2}, & \bar{z}_1 \partial_{\bar{z}_1} &\mapsto \bar{z}_2 \partial_{\bar{z}_2}, \\ z_2 \partial_{z_2} &\mapsto -z_1 \partial_{z_1}, & \bar{z}_2 \partial_{\bar{z}_2} &\mapsto -\bar{z}_1 \partial_{\bar{z}_1}. \end{aligned}$$

However, the induced complex structure on $\mathbb{C}^2 \setminus D$ does not extend over D , as we have, for example,

$$\partial_{z_1} \mapsto \frac{z_2}{z_1} \partial_{z_2}. \quad \triangle$$

3. CLASSIFICATION OF TORUS ACTIONS AND CONNECTIONS

Having introduced the geometric structures of interest last section, now we move on to study manifolds with the relevant torus actions. We begin this section by recalling the classification of certain torus actions by Haefliger and Salem in [13]. The relevant cohomology class for this classification is $\check{H}^2(B; \mathbb{Z}^n)$, where B is the quotient space of the torus action. Although the Haefliger–Salem class greatly resembles the Chern class of a principal S^1 -bundle, no such description exists at the moment in terms of connections and curvature. We will use the elliptic tangent bundle to provide a connection point of view on this class.

3.1. Haefliger–Salem classification. The first ingredient of the classification is the following result:

Theorem 3.1 (Theorem 3.1 in [13]). *Let M be a manifold with a T^k -action, and let $\pi: M \rightarrow B$ denote the quotient map. Any diffeomorphism $h: M \rightarrow M$ commuting with the action of T^k and preserving all orbits is of the form $h(x) = f(\pi(x)) \cdot x$, where $f \in C^\infty(B; T^k)$.*

Here we say that $f \in C^\infty(B)$ if $\pi^* f$ is smooth on M . Similarly, we define $\Omega^\bullet(B)$ to correspond to $\Omega_{\text{bas}}^\bullet(M)$.

Remark 3.2. The discussion in [13] concerns itself with the more general setting in which M is also allowed to be an orbifold. As our case of interest is when the total space is a smooth manifold we will immediately restrict to that case. \diamond

Definition 3.3. Let M, M' be two manifolds endowed with T^k -actions and let $\pi: M \rightarrow B, \pi': M' \rightarrow B$ denote the quotient maps. Then M and M' are said to be **locally equivalent** if each point in B has a neighbourhood U such that there exists a T^k -equivariant diffeomorphism from $\pi^{-1}(U)$ to $\pi'^{-1}(U)$, covering the identity on B . \diamond

When fixing an effective T^k -action on a manifold M it is possible to classify all other such actions which are locally equivalent to M :

Proposition 3.4 (Proposition 4.2 in [13]). *Let $\pi: M \rightarrow B$ be a manifold endowed with an effective torus action with connected isotropy groups. The equivalence classes of manifolds endowed with locally equivalent torus actions form an affine space modelled on $\check{H}^2(B; \mathbb{Z}^k)$.*

Since we will need to get our hands on the cohomology classes prescribed by the proposition, we give a sketch of the proof. Notice that there is no preferred manifold endowed with a torus action locally equivalent to that of M and the class in question measures how different from M another such manifold is.

Proof. Let $\pi': M' \rightarrow B$ be a manifold endowed with a locally equivalent torus action. Therefore, there exists a cover $\mathcal{U} = \{U_i\}_{i \in I}$ of B together with equivariant diffeomorphisms $h_i: \pi^{-1}(U_i) \rightarrow \pi'^{-1}(U_i)$. The transition maps $h_j^{-1} \circ h_i: \pi^{-1}(U_i \cap U_j) \rightarrow \pi^{-1}(U_i \cap U_j)$ satisfy the assumptions of

Theorem 3.1 and therefore there exists $f_{ji} \in C^\infty(U_i \cap U_j; T^k)$ such that $h_j^{-1} \circ h_i(x) = f_{ji}(\pi(x)) \cdot x$. These functions combine to give a cocycle $\{f_{ji}\} \in \check{C}^1(\mathcal{U}; \mathcal{T}^k)$, where \mathcal{T}^k is the sheaf of smooth functions with values in T^k . Using the exponential sequence one obtains the desired class in $\check{H}^2(\mathcal{U}; \mathbb{Z}^k)$.

Conversely, given a class in $\check{H}^2(\mathcal{U}; \mathbb{Z}^k)$, one uses again the exponential sequence to obtain a cocycle $\{f_{ji}\} \in \check{C}^1(\mathcal{U}; \mathcal{T}^k)$. Using this cocycle one may glue the disjoint unions of the $\pi^{-1}(U_i)$ to obtain a manifold M' endowed with a locally equivalent torus action. One readily checks that cohomologous cocycles result in equivalent T^k manifolds. \square

Definition 3.5. Let $\pi : M \rightarrow B$ be a manifold endowed with an effective torus action with connected isotropy groups, and $\pi' : M' \rightarrow B$ be a locally equivalent action. We will refer to the corresponding class in $\check{H}^2(B, \mathbb{Z}^k)$ as the **Haefliger-Salem class** of M' with respect to M . \diamond

Remark 3.6. A key difference with the classification of principal torus bundles in terms of their Chern class, is that the classification in Proposition 3.4 is affine in nature. Given a base B , there is no canonical torus actions playing the role $B \times T^k$ has in the principal case. \diamond

Remark 3.7. Note that the results in [13] are about orbifolds, but as we assume that M is a smooth manifold, any locally equivalent spaces will be smooth manifolds as well. \diamond

3.2. Torus actions, divisors and connections. We will lift torus actions to the elliptic tangent bundle in such a way that the associated infinitesimal action becomes free. The required notion we need is the following:

Definition 3.8. A T^k action and an elliptic divisor, $I_{|D|}$ on a manifold M are **compatible** if

- the action is effective,
- the isotropy group of each point is connected,
- the points with non-trivial isotropy are contained in D ,
- the action preserves the ideal $I_{|D|}$.

\diamond

These assumptions allow us to lift to the elliptic tangent bundle:

Proposition 3.9. Let $I_{|D|}$ be an elliptic divisor compatible with a T^k -action on M . Then there exists an injective Lie algebroid homomorphism $\tilde{a} : \mathfrak{t}^k \rightarrow \mathcal{A}_{|D|}$ such that the following diagram commutes:

$$\begin{array}{ccc} & & \mathcal{A}_{|D|} \\ & \nearrow \tilde{a} & \downarrow \\ \mathfrak{t}^k & \xrightarrow{a} & TM \end{array}$$

Proof. The existence of the Lie algebroid map is a local statement, so we can work in neighbourhoods of points.

Given a point p where action is free, there are two possibilities, either $p \notin D$ or $p \in D$. In the first case, \mathcal{A} isomorphic to TM via the anchor map in a neighbourhood of p and the action has a unique lift to $\mathcal{A}_{|D|}$. If $p \in D$, because the action preserves the ideal $\mathcal{A}_{|D|}$, then the orbit of p lies within the same stratum as p , say $D[i]$ and because the action is free, we get a rank k subbundle $a(\mathfrak{t}^k) \subset TD[i] \hookrightarrow \mathcal{A}_{|D|}$, and therefore a has a lift, which is unique by continuity.

Given a point where the action has nontrivial isotropy, say $T^l \subset T^k$, we can split $T^k = T^l \times T^{k-l}$. By the previous argument the action of T^{k-l} lifts to $\mathcal{A}_{|D|}$ and the lift is made of a subspace that maps injectively to TM via the anchor. Now we focus on the action of T^l . Since

p is fixed by the action of T^l and the action is effective we can linearise it and assume that, in appropriate coordinates (with values in $\mathbb{C}^m \times \mathbb{R}^{n-2m}$), the action is given by

$$(\theta_1, \dots, \theta_l) \cdot (z_1, \dots, z_m, x_1, \dots, x_{n-2m}) \mapsto (e^{i(\sum_j a_{1,j}\theta_j)} z_1, \dots, e^{i(\sum_j a_{m,j}\theta_j)} z_m, x_1, \dots, x_{n-2m}),$$

where $m \geq l$ and the matrix $A = (a_{i,j})_{ij}$ has integral coefficients. Because all isotropy groups are connected A has an $l \times l$ submatrix with determinant ± 1 .

Suppose f is a local generator of the elliptic divisor $I_{|D|}$. Since D has codimension 2, the locus where f is nonzero is connected and hence we may take f to be non-negative, vanishing only at D . Since f is non-negative, by taking its average over the T^l action, we obtain another nonnegative generator of the ideal, still denoted by f , which is invariant under the T^l action. The invariant polynomials of the T^l -action are generated by $(r_1^2, \dots, r_m^2, x_1, \dots, x_{n-2m})$, and hence a result by Schwarz [20] implies that f must be of the form $f = g(r_1^2, \dots, r_m^2, x_1, \dots, x_{n-2m})$, for some smooth g . Because f has this form, it is immediate that the vector fields $\partial_{\vartheta_1}, \dots, \partial_{\vartheta_m}$ preserve the ideal generated by f , where we use polar coordinates for z_i : $z_i = r_i e^{i\vartheta_i}$. Hence ∂_{ϑ_i} give rise to (non-vanishing) elements of $\mathcal{A}_{|D|}$. Because the vector fields generating the T^k -action are given by $A(\partial_{\vartheta_1}), \dots, A(\partial_{\vartheta_l}) \in \text{span}\langle \partial_{\vartheta_1}, \dots, \partial_{\vartheta_m} \rangle$, and A has maximal rank, we see that these vector fields span a rank l -subspace of $\mathcal{A}_{|D|}$ which over D are in the kernel of the anchor map and hence are independent from the generators corresponding to the T^{k-l} action. \square

Studying general torus actions goes beyond the scope of this article. For instance, one problem which occurs is that there is too much freedom in choosing a compatible elliptic divisor:

Example 3.10. Consider \mathbb{C}^2 , endowed with the S^1 -action $\theta_1 \cdot (z_1, z_2) = (e^{i\theta_1} z_1, z_2)$ and the compatible elliptic divisor generated by $|z_1|^2 |z_2|^2$. Using coordinates (t, z_2) for the quotient space $B = \mathbb{R}_{\geq 0} \times \mathbb{C}$, we have that B inherits a quotient divisor $I_{\text{red}} = \langle t |z_2|^2 \rangle$. Which is a mixture between an elliptic and real log divisor. More generally, for a T^k -action on \mathbb{C}^n the induced divisor on the base can have any kind of mixture between elliptic and real log divisors. \triangle

We restrict ourselves to a very well-behaved class of singular torus actions, but still interesting enough to allow for many examples:

Definition 3.11. We say that a T^k -action on a manifold M^n is **standard** if each orbit has a neighbourhood U such that

$$U \simeq T^{k-l} \times \mathbb{C}^l \times \mathbb{R}^m \quad \text{for some } k, l, m \text{ such that } k + l + m = n,$$

in which the action is given by

$$\begin{aligned} (e^{i\theta_1}, \dots, e^{i\theta_k}) \cdot (\varphi_1, \dots, \varphi_{k-l}, z_1, \dots, z_l, x_1, \dots, x_m) \\ = (e^{i\theta_1} \varphi_1, \dots, e^{i\theta_{k-l}} \varphi_{k-l}, e^{i\theta_{k-l+1}} z_1, \dots, e^{i\theta_k} z_l, x_1, \dots, x_m), \end{aligned}$$

possibly after changing the generators of T^k . \diamond

For general T^k -actions, deriving conditions when they are standard is not simple. However, for full torus actions one can show the following:

Lemma 3.12. *A T^n -action on a manifold M^{2n} which is effective and has connected isotropy groups is standard.*

The local normal form of a standard T^k -action immediately implies the following:

Proposition 3.13. *Let $\pi : M \rightarrow B := M/T^k$ be a manifold endowed with a standard T^k -action. Then*

- B is a manifold with corners.
- The action is compatible with the elliptic divisor $I_{|D|} := \pi^* I_{\partial B}$.
- π induces a Lie algebroid submersion $\mathcal{A}_{|D|} \rightarrow \mathcal{A}_{\partial B}$.

In particular there is a canonical compatible elliptic divisor associated to a standard T^k -action. Therefore, in what follows when given a standard T^k -action on M , $I_{|D|}$ will always denote the canonical associated elliptic divisor.

Example 3.14. Consider $S^n \subset \mathbb{R}^{n+1}$, and the circle action given by rotating the first two coordinates. This action is standard with quotient space diffeomorphic to D^n and elliptic divisor generated by $x_1^2 + x_2^2$. \triangle

Because a standard T^k -action lifts to the Lie algebroids so does the projection map, and we obtain a diagram as follows:

$$\begin{array}{ccc} \mathcal{A}_{|D|} & \xrightarrow{\tilde{\pi}} & \mathcal{A}_{\partial B} \\ \downarrow & & \downarrow \\ M & \xrightarrow{\pi} & B \end{array}$$

Hence, we obtain a short exact sequence of vector bundles over M :

$$0 \rightarrow \ker d\tilde{\pi} \rightarrow \mathcal{A}_{|D|} \rightarrow \pi^* \mathcal{A}_{\partial B} \rightarrow 0$$

We may quotient by T^k , to obtain the sequence of vector bundles over B :

$$0 \rightarrow B \times \mathfrak{t}^k \rightarrow \mathcal{A}_{|D|}/T^k \rightarrow \mathcal{A}_{\partial B} \rightarrow 0$$

One may think of this sequence as an $\mathcal{A}_{|D|}$ version of the Atiyah sequence for principal bundles. With this in mind, vector bundle splittings of this sequence give rise to the notion of connections for the T^k -action on M . Concretely, in terms of differential forms we have:

Definition 3.15. Let M be a manifold endowed with a T^k action, and let $I_{|D|}$ be a compatible elliptic divisor. A **connection-one form** on M is a Lie algebroid one-form $\Theta \in \Omega^1(\mathcal{A}_{|D|}; \mathfrak{t}^k)$ such that

- $m_\lambda^* \Theta = \Theta$ for all $\lambda \in T^k$
- $\iota_{a(v)} \Theta = v$, for all $v \in \mathfrak{t}^k$.

\diamond

3.2.1. Complex line bundles. Let $L \rightarrow M$ be a complex line bundle. Then $\Gamma(L^{1,0*}) \subset C^\infty(M; \mathbb{C})$ naturally defines a smooth complex log divisor. The associated elliptic divisor $I_{|D|}$ is naturally compatible with the standard S^1 -action on L .

Connections for the S^1 -action arise from linear connections on L :

Lemma 3.16. *Let ∇ be a complex linear connection on a complex line bundle $L \rightarrow M$, and let $\zeta \in \Omega^1(L \setminus M; \mathbb{C})$ denote the corresponding connection one-form on the associated \mathbb{C}^* -bundle. Then ζ extends over the zero-section as a complex logarithmic form, and $\Theta := \Im(\zeta) \in \Omega^2(\mathcal{A}_{|D|})$ defines a connection one-form for the associated S^1 -action*

Proof. Let $z_\alpha : L|_{U_\alpha} \rightarrow \mathbb{C}$ be local fibre coordinates of L . Write $z_\alpha = r_\alpha e^{i\theta_\alpha}$, and consider the complex log forms $d \log z_\alpha = d \log r_\alpha + i d\theta_\alpha$. If $z_\alpha = g_\alpha^\beta z_\beta$, with g_β^α transition functions, we find that $d \log z_\alpha - d \log z_\beta = d \log g_\beta^\alpha$. Given a complex linear connection ∇ on L , we may consider the corresponding connection one-forms $\xi_\alpha \in \Omega^1(U_\alpha; \mathbb{C})$, which satisfy $\xi_\alpha - \xi_\beta = d \log g_\beta^\alpha$. We conclude that $\zeta := d \log z_\alpha + \xi_\alpha$ defines a global complex logarithmic form \square

If we let $K = K_1 + iK_2$ denote the curvature of the connection ∇ on L , we find that $d\Theta = K_2$ is actually a smooth form.

Remark 3.17. The form Θ , restricted to the complement of the zero-section, also goes by the name of the global angular form. The above discussion shows that the global angular form can be extended over the zero-section as an elliptic differential form. \diamond

3.3. Smoothness of the curvature. Given a standard T^k -action and a corresponding connection one-form $\Theta \in \Omega^1(\mathcal{A}_{|D|}; \mathfrak{t}^k)$, the curvature, $d\Theta$, is basic and descends to a form in $\Omega^2(\mathcal{A}_{\partial B}; \mathfrak{t}^k)$. We saw for complex line bundles that there are connections for which the curvature was in fact a smooth form. This is a general phenomenon for standard torus actions. To proof this, we need to understand the cohomology of the log-tangent bundle on the base:

Proposition 3.18 ([16],[12]). *Let (M, I_Z) be a manifold endowed with real log divisor. Then there exists an isomorphism*

$$(3.1) \quad H^\bullet(\mathcal{A}_Z) \simeq H^\bullet(M) \oplus H^{\bullet-1}(Z[1]) \oplus H^{\bullet-2}(Z[2]) \oplus \cdots,$$

where $Z[n]$ denotes the intersection stratification of Z . The maps $H^\bullet(\mathcal{A}_Z) \rightarrow H^{\bullet-i}(Z[i])$ are induced by the residue maps Res^i , and the map $H^\bullet(M) \rightarrow H^\bullet(\mathcal{A}_Z)$ is the pullback via the anchor.

Remark 3.19. The map Res^2 is defined combining the local residues $\text{Res}_{x_i x_j}$, as in (2.1). We refer to ([16],[12]) for the details. \diamond

Using (3.1) we see that we can associate a well-defined smooth cohomology class to a class in $H^\bullet(\mathcal{A}_Z)$. Explicitly we may also phrase this as follows:

Lemma 3.20. *Let (M, Z) be a manifold endowed with a immersed hypersurface, with self-transverse intersections. Let $\alpha \in \Omega^\bullet(M)$ be such that $[\rho^*\alpha] = 0 \in H^\bullet(\mathcal{A}_Z)$. Then $[\alpha] = 0 \in H^\bullet(M)$.*

Recall from Lemma 3.13 that the quotient map of a standard T^k -action induces a Lie algebroid submersion $\pi : \mathcal{A}_{|D|} \rightarrow \mathcal{A}_{\partial B}$. This map is compatible with the residues in the following manner.

Lemma 3.21. *Let $\pi : M \rightarrow B$ be endowed with a standard T^k -action. Then for $i = 1, 2$ the following diagram commutes:*

$$\begin{array}{ccc} \Omega^2(\mathcal{A}_{|D|}) & \xrightarrow{\text{Res}_r^i} & \Omega^{2-i}(T^i ND[i]) \\ \pi^* \uparrow & & \uparrow \pi^* \\ \Omega^2(\mathcal{A}_{\partial B}) & \xrightarrow{\text{Res}^i} & \Omega^{2-i}(\partial B[i]) \end{array}$$

As $D[2]$ is a discrete set of points, $\text{Res}_r^2 \omega$ is defined to be $\text{Res}_{r_1 r_2}(\omega)(p)$. A similar result holds for higher order residues, but is more involved to state.

Using the description of the cohomology of $\mathcal{A}_{\partial B}$, we can show that the curvature of a connection, $d\Theta$, is cohomologically smooth:

Lemma 3.22. *Let M be endowed with a standard T^k -action and let $\Theta \in \Omega^1(\mathcal{A}_{|D|}; \mathfrak{t}^k)$ be a connection-one form with curvature $K \in \Omega^2(\mathcal{A}_{\partial B}; \mathfrak{t}^k)$. Then there exists a smooth form $\tilde{K} \in \Omega^2(B; \mathfrak{t}^k)$ with $[\rho^* \tilde{K}] = [K] \in H^2(\mathcal{A}_{\partial B}; \mathfrak{t}^k)$.*

Proof. Using (3.1) we have to show that the residues Res^i of K vanish. By rank reasons Res^i vanishes for $i \geq 3$. We have that $p^* \text{Res}^1(K) = \text{Res}_r^1(d\Theta) = d\text{Res}_r^1(\Theta)$. By T^n -invariance of Θ we must have that $\text{Res}_r^1(\Theta)$ is T^n -invariant as well, and thus of the form p^*f , with $f \in$

$C^\infty(B)$. Therefore $p^* \text{Res}^1(K) = p^* df$, and we conclude that $\text{Res}^1(K)$ is exact. We also have that $p^* \text{Res}^2(K) = \text{Res}_r^2(d\Theta)$, but the latter is necessarily zero by smoothness of the coefficient functions of elliptic one-forms. We conclude that all the residues of $[K]$ vanish, as these are precisely the maps $H^\bullet(\mathcal{A}_{\partial B}) \rightarrow H^{\bullet-i}(\partial B[i])$ in Equation (3.1), we conclude that $[K] = [\rho^* \tilde{K}]$, for some smooth $\tilde{K} \in \Omega^2(B; \mathfrak{t}^k)$. \square

The following example shows that Lemma 3.20 is not a given when the elliptic divisor and action are independent:

Example 3.23. Let M be a manifold endowed with a smooth elliptic divisor $I_{|D|}$, with D co-orientable, and consider the trivial S^1 -bundle, $\pi : M \times S^1 \rightarrow M$. The elliptic divisor is trivially compatible with the S^1 -action, and hence we can choose a connection one-form for this action. Any such form will be given by $\Theta = \alpha + d\theta$, with $\alpha \in \Omega^1(\mathcal{A}_{|D|})$ and $d\theta \in \Omega^1(S^1)$.

We may pick $\alpha = 0$, and then the curvature of Θ simply vanishes. However, we may change α in such a manner that $d\Theta$ remains smooth but is no longer exact as an element in $H^2(M)$, contradicting the conclusion of Lemma 3.20.

The construction goes as follows: Consider a metric on a tubular neighbourhood \mathcal{U} of D in M , and let $r^2 : \mathcal{U} \rightarrow \mathbb{R}$ denote the associated squared distance function. Let $\Theta_M \in \Omega^1(\mathcal{A}_{|D|})$ be a connection one-form on \mathcal{U} , obtained as in Lemma 3.16 by viewing \mathcal{U} as an (open) in a complex line bundle.

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function with $\text{supp} f \subset [-1, 1]$, and $f|_{[-\epsilon, \epsilon]} \equiv 1$ for some $\epsilon < 1$. Define an elliptic form $\alpha = f(r^2)\Theta_M$. The cohomology class $d\alpha$ is concentrated in \mathcal{U} , where it is precisely the curvature of the normal bundle of D . Consequently, we see that $\Theta = \alpha + d\theta$ has smooth curvature, which is not cohomologous to zero. \triangle

3.3.1. Local description of connection one-forms. For future reference we recall the local description for connection one-forms on principal bundles:

Proposition 3.24. *Let $\pi : P \rightarrow B$ be a principal G -bundle, and let $h_i : \pi^{-1}(U_i) \rightarrow U_i \times G$ be local trivialisations for $i = 1, 2$. Let $\Theta \in \Omega^1(P; \mathfrak{g})$ be a connection one-form on P . Then*

$$h_i^* \Theta = \omega_{\text{MC}} + \pi^* \alpha_i,$$

with $\omega_{\text{MC}} \in \Omega^1(G; \mathfrak{g})$ the Maurer-Cartan form, and $\alpha_i \in \Omega^1(U_i; \mathfrak{g})$. Moreover, if we let $f_{ji} \in C^\infty(U_i \cap U_j; G)$ be the transition functions given by $h_j^{-1} \circ h_i(x, g) = (x, f_{ji}(x) \cdot g)$, we have

$$\alpha_i - \text{Ad}_{f_{ji}}^* \alpha_j = \pi^* d \log f_{ji}.$$

3.4. Čech-de Rham isomorphism. To continue, we need to recall a basic version of the Čech-de Rham isomorphism.

Let M be endowed with a proper group action by some group G . Let Ω_{bas}^n denote the sheaf of basic differential forms and let Z_{bas}^n denote the sheaf of closed basic differential forms. Both are sheaves of $C^\infty(M)_{\text{bas}}$ -modules and fit into the short exact sequence

$$(3.2) \quad 0 \rightarrow Z_{\text{bas}}^n \rightarrow \Omega_{\text{bas}}^n \rightarrow Z_{\text{bas}}^{n+1} \rightarrow 0.$$

Consider the basic Čech cocycle groups $C_{\text{bas}}^k(\mathcal{U}, \mathcal{F})$, for which the maps $U_{i_1 \dots i_k} \rightarrow \mathcal{F}$ are invariant with respect to the G -action on $U_{i_1 \dots i_k}$.

By existence of a G -invariant partition of unity on M , the sheaves Ω_{bas}^n are fine, and hence $H_{\text{bas}}^i(M, \Omega_{\text{bas}}^n) = \{0\}$ for all $i \geq 1$. Therefore the connecting morphism of the above exact sequence induces an isomorphism $H_{\text{bas}}^i(M, Z_{\text{bas}}^{n+1}) \rightarrow H_{\text{bas}}^i(M, Z_{\text{bas}}^n)$. Applying this consecutively, one obtains the isomorphism between $H_{\text{bas}}^{i+1}(M, \mathbb{R}) \rightarrow H_{\text{bas}}^1(M, Z_{\text{bas}}^i)$. As the latter corresponds to $H_{dR, \text{bas}}^{i+1}(M)$ one obtains the desired isomorphism.

We also consider a basic version of the exponential sequence:

$$(3.3) \quad 0 \rightarrow \mathbb{Z}^n \rightarrow C_{\text{bas}}^\infty(M; \mathbb{R}^n) \rightarrow C_{\text{bas}}^\infty(M; T^n) \rightarrow 0.$$

The long exact sequence will induce an isomorphism $\check{H}_{\text{bas}}^1(M, C_{\text{bas}}^\infty(M; T^n)) \rightarrow \check{H}_{\text{bas}}^2(M; \mathbb{Z}^n)$. Combining this with the Čech-de Rham isomorphism we obtain a map to $H_{\text{bas}}^2(M; \mathbb{Z}^n)$.

3.5. Connection approach to Haefliger-Salem class. We now have all the theory in place to compare the curvature of a connection for a standard T^k -action with the Haefliger-Salem class:

Proposition 3.25. *Let $\pi : M \rightarrow B$ be a manifold endowed with a standard T^k -action and let $\pi' : M' \rightarrow B$ be a manifold endowed with a locally equivalent action. Let Θ, Θ' be connection one-forms on M respectively M' . Then $[d\Theta - d\Theta'] \in H^2(A_{\partial B}; \mathfrak{t}^k)$ is representable by a smooth form, and descends to the Haefliger-Salem class of M' in $\check{H}^2(B; \mathfrak{t}^k)$.*

Proof. Let $I_{|D|}, I_{|D'|}$ denote the induced elliptic divisors on M respectively M' . By Lemma 3.22, we have that both $[d\Theta]$ and $[d\Theta']$ are representable by smooth forms, therefore we can modify Θ and Θ' such that $d\Theta$ and $d\Theta'$ are smooth forms. We will still denote the modified connections by Θ and Θ' .

Let $\mathcal{U} = \{U_i\}$ be an open cover of B , together with equivariant diffeomorphism $h_i : \pi^{-1}(U_i) \rightarrow \pi'^{-1}(U_i)$ provided by the fact that the actions on M and M' are locally equivalent. Furthermore, assume that over $U_i \setminus \partial B$ both π and π' are trivialisable as principal T^k -bundles.

Because the h_i are equivariant, they are morphisms of divisors, and hence we may consider $h_i^* \Theta' \in \Omega^1(\mathcal{A}_{|D|}|_{\pi^{-1}(U_i)}; \mathfrak{t}^k)$. One readily shows that $h_i^* \Theta'$ is a connection on $\pi^{-1}(U_i)$ and thus $\eta_i := \Theta - h_i^* \Theta' \in \Omega_{\text{bas}}^1(\mathcal{A}_{|D|}|_{\pi^{-1}(U_i)}; \mathfrak{t}^k)$ is a basic form.

Let $f_{ji} \in C^\infty(U_i \cap U_j; T^k)$ be the transition functions given by $h_j \circ h_i(x) = (x, f_{ji}(x))$. By Lemma 3.24, $\eta_i - \eta_j = h_i^* \Theta' - h_j^* \Theta'$ coincides with $\pi^* d \log f_{ji}$ on $\pi^{-1}((U_i \cap U_j) \setminus \partial B)$. By density, we have that $\eta_i - \eta_j = \pi^* d \log f_{ji}$ on the entirety of $\pi^{-1}(U_i \cap U_j)$. As the η_i are closed, we conclude that $\{\eta_i - \eta_j\}_{ij}$ defines a cocycle in $\check{C}_{\text{bas}}^1(M, Z_{\text{bas}}^1)$.

The Haefliger-Salem class is defined by applying the exponential sequence to the class $\{f_{ji}\}$ in $\check{H}^1(\mathcal{U}; T^k)$. Inspecting the Čech-de Rham isomorphism, we conclude that this class corresponds to the class defined by $\{\eta_i - \eta_j\}_{ij}$. We are left to show that $\{\eta_i - \eta_j\}_{ij} \in \check{H}^1(\mathcal{U}, Z_{\text{bas}}^1)$ corresponds to $[d\Theta - d\Theta']$. By exactness of (3.2), we know that there exist a refinement $\tilde{\mathcal{U}} = \{\tilde{U}_i\}$ of \mathcal{U} and $\tilde{\eta}_i \in \Omega_{\text{bas}}^1(\pi^{-1}(\tilde{U}_i); \mathfrak{g})$ such that $\tilde{\eta}_i - \tilde{\eta}_j = \eta_i - \eta_j$. The class in $H_{dR, \text{bas}}^2(M, \mathfrak{t}^k)$ corresponding to $\{\eta_i - \eta_j\}_{ij}$ is given by $d\tilde{\eta}_i$.

Note that $\tilde{\eta}_i - \eta_i = \tilde{\eta}_j - \eta_j$, and hence these forms glue to give a globally defined form $\eta \in \Omega_{\text{bas}}^1(M; \mathfrak{g})$. Because $d\eta = d\tilde{\eta}_i - d\eta_i$, and $d\eta_i = d\Theta - h_i^* d\Theta'$, we have that $d\eta$ is smooth. Hence, by Lemma 3.20 we have that $d\eta$ is exact in $H^2(B; \mathfrak{t}^k)$. Consequently $[d\tilde{\eta}_i] = [d\eta_i] \in H^2(B; \mathfrak{t}^k)$. But as $d\eta_i = d\Theta - h_i^* d\Theta'$, we conclude that the class in $H_{dR, \text{bas}}^2(M, \mathfrak{t}^k)$ corresponding to $\{\eta_i - \eta_j\}_{ij}$ is given by $d\Theta - d\Theta'$, which finishes the proof. \square

The description of the Haefliger-Salem class in terms of connections has the following consequence:

Lemma 3.26. *Let $\pi : M \rightarrow B$ be a standard T^k -action. Then there exists a locally equivalent action $\pi' : M' \rightarrow B$ which admits a flat connection.*

Proof. Let Θ be a connection on M , with integral curvature. The class $[d\Theta] \in H^2(B; \mathbb{Z}^k)$ corresponds to a locally equivalent torus action $M' \rightarrow B$. Given a connection Θ' on M' , we have by Proposition 3.25 that $[d\Theta - d\Theta']$ defines the Haefliger-Salem class of M' . But as this class is given by $[d\Theta]$, we conclude that $[d\Theta'] = 0$ which finishes the proof. \square

The importance of this lemma is that in the classification of standard T^k -actions, it is now possible to pick as a reference an action $M_0 \rightarrow B$ with a flat connection. Using this reference, and a locally equivalent action $M' \rightarrow B$, we find that the Haefliger-Salem class of M_0 with respect to M' is given exactly by $[d\Theta']$ for some connection Θ' on M' .

Definition 3.27. Let $M \rightarrow B$ be a standard T^k -action. The locally equivalent action $M_0 \rightarrow B$ which admits a flat connection is called the **reference action**. \diamond

When $M \rightarrow B$ is a principal T^k -action, the reference action is simply the trivial bundle $B \times T^k$. Contrary to in the principal case, there might be several non locally equivalent standard T^k -actions over the same base with flat connections. We will see examples of this in Section 5.2.1.

4. THE SINGULAR DIFFERENTIAL GEOMETRY OF THE CORRESPONDENCE SPACE

Let $M \rightarrow B, \hat{M} \rightarrow B$ be manifolds endowed with standard torus actions. The fibre-product $M \times_B \hat{M} \subset M \times \hat{M}$ fails to be a smooth submanifold. This space plays an important role in T -duality, and therefore we will shortly study notions for smooth objects on this space.

Let us start with the simplest example:

Example 4.1. Let M, \hat{M} both denote \mathbb{C} endowed with the standard S^1 -action, and $B = \mathbb{R}_{\geq 0}$ the quotient. Then

$$M \times_B \hat{M} = \{(z, \tilde{z}) : |z| = |\tilde{z}|\},$$

and hence we may consider singular coordinates $(r, \theta, \hat{\theta}) \in \mathbb{R}_{\geq 0} \times T^2$. These coordinates are well-defined except at $r = 0$. We see that $M \times_B \hat{M}$ is a cone over a torus. \triangle

The space $\mathbb{C} \times_{\mathbb{R}_{\geq 0}} \mathbb{C}$ will serve as local model for the spaces we want to consider, so we will describe smoothness in terms of it:

Definition 4.2. A function $f : \mathbb{C} \times_{\mathbb{R}_{\geq 0}} \mathbb{C} \rightarrow \mathbb{R}$ is smooth, if it is the restriction of a smooth function $\tilde{f} \in C^\infty(\mathbb{C} \times \mathbb{C})$. \diamond

Definition 4.3. A map $\Phi : \mathbb{C} \times_{\mathbb{R}_{\geq 0}} \mathbb{C} \rightarrow \mathbb{C} \times_{\mathbb{R}_{\geq 0}} \mathbb{C}$ is said to be smooth, if for every $f \in C^\infty(\mathbb{C} \times_{\mathbb{R}_{\geq 0}} \mathbb{C})$, $f \circ \Phi$ is smooth. \diamond

Taking products, we can talk about smooth maps and diffeomorphisms of $(\mathbb{C} \times_{\mathbb{R}_{\geq 0}} \mathbb{C})^l \times \mathbb{R}^k$, and charts and atlases of this form. From the local form of standard T^k -actions it immediately follows that:

Lemma 4.4. Let M, \hat{M} be standard T^k -actions over the same base B . Then $M \times_B \hat{M}$ admits an atlas of coordinate charts of the form $(\mathbb{C} \times_{\mathbb{R}_{\geq 0}} \mathbb{C})^l \times \mathbb{R}^k$.

A point in $M \times_B \hat{M}$ has singular coordinates

$$(4.1) \quad (r_1, \dots, r_l, \theta_1, \dots, \theta_l, \hat{\theta}_1, \dots, \hat{\theta}_l, \psi_1, \dots, \psi_{n-l}, x_1, \dots, x_{n-l}, \hat{x}_1, \dots, \hat{x}_{n-l}).$$

Here the $\{r_i\}$ are one set of radial coordinates on \mathbb{C}^l , where $\{\theta_i\}, \{\hat{\theta}_i\}$ are two sets of polar coordinates on \mathbb{C}^l . The $\{\psi_i\}$ are honest coordinates on T^{n-l} and the $\{x_i\}$ and $\{\hat{x}_i\}$ are coordinates on the two copies of \mathbb{R}^{n-l} .

Remark 4.5 (Induced divisor). Let M, \hat{M} be manifolds endowed with standard T^k -actions and same base B and let $I_{|D|}$ and $I_{|\hat{D}|}$ denote the corresponding elliptic divisors. Then the restriction of $I_{|D|}$ and $I_{|\hat{D}|}$ to $M \times_B \hat{M}$ coincides. In the coordinates of (4.1) the induced elliptic divisor on $M \times_B \hat{M}$ is given by $\langle r_1, \dots, r_l \rangle$. \diamond

Rather than trying to find a notion of tangent bundle for $M \times_B \hat{M}$, we use the ambient elliptic tangent bundle on $M \times \hat{M}$ to define a Lie algebroid over $M \times_B \hat{M}$.

The first observation is that the way $M \times_B \hat{M}$ sits inside the smooth manifold $M \times \hat{M}$ is compatible with the Lie algebroid $\mathcal{A}_{|D|} \boxtimes \mathcal{A}_{|\hat{D}|}$ in the following manner:

Definition 4.6. Let $\mathcal{A} \rightarrow M$ be a Lie algebroid and $N \subset M$ a subset, and let $\Gamma_N(\mathcal{A})$ denote the sections of \mathcal{A} tangent to N . Then N is **adapted** to \mathcal{A} if $\text{Sing}(N) \subseteq \{x \in M : \rho_x \text{ is not of full rank}\}$ and the pull-back sheaf $\iota_N^* \Gamma_N(\mathcal{A})$ is locally free over $\iota_N^* C^\infty(M)$. \diamond

As $\iota_N^* \Gamma(\mathcal{A})$ is locally free, we may view it as a topological vector bundle over N . By declaring sections to be smooth if they are restrictions of smooth sections of \mathcal{A} , we can view \mathcal{A} as a smooth Lie algebroid over N .

Example 4.7. A subspace $N \subset M$ is adapted to TM if and only if it is an embedded submanifold. In this case the pull-back sheaf is simply given by vector fields on N . \triangle

The rationale behind this definition is as follows. We allow subspaces to have singularities, but they need to occur precisely when the anchor of \mathcal{A} is not an isomorphism.

Example 4.8. Let $M = \mathbb{R}^2$, and let \mathcal{A} be the Lie algebroid generated by $x\partial_x$ and $y\partial_y$. Let N denote the union of the diagonal and anti-diagonal. If $X = fx\partial_x + gy\partial_y$ is tangent to the diagonal, it implies it must preserve the ideal generated by $x - y$. Therefore, $X = f(x\partial_x + y\partial_y)$. As these vector fields are also tangent to the anti-diagonal we conclude that N is an \mathcal{A} -adepted subspace, and the restricted Lie algebroid is generated by $x\partial_x + y\partial_y$. \triangle

From the local form of standard T^k -actions, we immediately obtain the following:

Lemma 4.9. Let M, \hat{M} be standard T^k -actions over the same base B . Then $M \times_B \hat{M}$ is an $\mathcal{A}_{|D|} \boxtimes \mathcal{A}_{|\hat{D}|}$ -adapted subspace.

We denote the induced Lie algebroid on $M \times_B \hat{M}$ by $\mathcal{A}_{|D|} \times_B \mathcal{A}_{|\hat{D}|}$.

5. NON PRINCIPAL T -DUALITY

Using the Lie algebroids from Section 2 we will extend the notions of T -duality to standard torus actions. To proof the existence of T -duals we will use the classification of torus actions, and the connection point of view thereof, as described in Section 3. Moreover, we will stress the differences with T -duality for principal torus bundles; in particular the new source of topology change arising from the singularities of the torus action.

5.1. T -duality. In this subsection we will recall the notions of T -duality for principal torus actions. We will be brief and refer to [3] and [7] for more details.

Definition 5.1. Let M and \hat{M} be principal T^k -bundles with base B and let $H \in \Omega_{T^k}^3(M)$, $\hat{H} \in \Omega_{T^k}^3(\hat{M})$ be T^k -invariant closed three-forms. Consider the fibre-product, $M \times_B \hat{M}$, and the following diagram:

$$\begin{array}{ccccc}
 & (M \times_B \hat{M}, p^*H - \hat{p}^*\hat{H}) & & & \\
 & \swarrow p & & \searrow \hat{p} & \\
 (M, H) & & & & (\hat{M}, \hat{H}) \\
 & \searrow \pi & & \swarrow \hat{\pi} & \\
 & B & & &
 \end{array}$$

We say that (M, H) and (\hat{M}, \hat{H}) are T -**dual** if there exists a T^{2k} -invariant two-form $F \in \Omega_{T^{2k}}^2(M \times_B \hat{M})$ such that

$$dF = p^*H - \hat{p}^*\hat{H},$$

and the restriction

$$F|_{\mathfrak{t}_M \times \mathfrak{t}_{\hat{M}}} : \mathfrak{t}_M \times \mathfrak{t}_{\hat{M}} \rightarrow \mathbb{R},$$

where $\mathfrak{t}_M, \mathfrak{t}_{\hat{M}}$ denotes the tangent space to the fibres of p, \hat{p} respectively, is non-degenerate. We call $M \times_B \hat{M}$ the **correspondence space**. \diamond

T -duals always exist, provided the three-form is integral:

Theorem 5.2 ([3]). *Let $M \rightarrow B$ be a principal T^k -bundle, and $H \in \Omega_{T^k}^3(M)$ with integral cohomology class. Then there exists a principal T^k -bundle, \hat{M} T -dual to M .*

T -dual spaces have isomorphic complexes of invariant differential forms:

Theorem 5.3 ([3]). *Let (M, H) and (\hat{M}, \hat{H}) be T -dual with $dF = p^*H - \hat{p}^*\hat{H}$. Then*

$$\tau : (\Omega_{T^k}^\bullet(M), d_H) \rightarrow (\Omega_{T^k}^\bullet(\hat{M}), d_{\hat{H}}), \quad \rho \mapsto \int_{T^k} e^F \wedge \rho,$$

is an isomorphism of cochain complexes. Here the integral is over the fibres of \tilde{p} .

5.2. Courant algebroids. T -duality induces an isomorphism of invariant sections of the standard Courant algebroid:

Theorem 5.4 ([7]). *If (M, H) and (\hat{M}, \hat{H}) are T -dual, then there is an isomorphism of Courant algebroids*

$$\varphi : (TM \oplus T^*M)/T^k \rightarrow (T\hat{M} \oplus T^*\hat{M})/T^k,$$

such that $\tau(v \cdot \rho) = \varphi(v) \cdot \tau(\rho)$, for all $v \in (TM \oplus T^*M)/T^k$ and $\rho \in \Omega^\bullet(M)$.

The strength of this description is that it immediately allows one to transport structures:

Theorem 5.5 ([7]). *Let (M, H) and (\hat{M}, \hat{H}) be T -dual spaces. Then any Dirac, generalized complex, generalized Kähler or SKT structure on M which is invariant under the torus action is transformed via φ into a structure of the same kind on \hat{M} .*

We are now ready to introduce the notion of T -duality for standard torus actions: **Do we want to add a remark that, at least the definition makes sense for more general compatible torus actions?**

Definition 5.6. Let M, \hat{M} be two manifolds endowed with standard T^k -actions over the same base B and let $I_{|D|}, I_{|\hat{D}|}$ be the induced elliptic divisors. Furthermore, suppose we are given $H \in \Omega_{\text{cl}}^3(\mathcal{A}_{|D|}), \hat{H} \in \Omega_{\text{cl}}^3(\mathcal{A}_{|\hat{D}|})$ closed and T^k -invariant. We consider the following diagram:

$$\begin{array}{ccccc} & & (\mathcal{A}_{|D|} \times_B \mathcal{A}_{|\hat{D}|} \rightarrow M \times_B \hat{M}, p^*H - \hat{p}^*\hat{H}) & & \\ & \swarrow p & & \searrow \hat{p} & \\ (\mathcal{A}_{|D|} \rightarrow M, H) & & & & (\mathcal{A}_{|\hat{D}|} \rightarrow \hat{M}, \hat{H}) \\ & \searrow \pi & & \swarrow \hat{\pi} & \\ & & \mathcal{A}_{\partial B} \rightarrow B & & \end{array}$$

We say that (M, H) and (\hat{M}, \hat{H}) are **T -dual** if there exists a T^{2k} -invariant two-form $F \in \Omega^2(\mathcal{A}_{|D|} \times_B \mathcal{A}_{|\hat{D}|})$ such that $p^*H - \hat{p}^*\hat{H} = dF$ and

$$(5.1) \quad F : \mathfrak{t}_M \otimes \mathfrak{t}_{\hat{M}} \rightarrow \mathbb{R},$$

is non-degenerate. \diamond

Let us expand on this definition. First $\mathcal{A}_{|D|} \times_B \mathcal{A}_{|\hat{D}|}$ is the Lie algebroid induced by viewing $M \times_B \hat{M}$ as an $(\mathcal{A}_{|D|} \boxtimes \mathcal{A}_{|\hat{D}|})$ -adapted subspace of $M \times \hat{M}$. The two-form F is a form for this Lie algebroid, and can in practise be thought of as the restriction of a two-form $\tilde{F} \in \Omega^2(\mathcal{A}_{|D|} \boxtimes \mathcal{A}_{|\hat{D}|})$. Here \mathfrak{t}_{M_i} denotes the kernel of the Lie algebroid submersion $\mathcal{A}_{|D|} \times_B \mathcal{A}_{|\hat{D}|} \rightarrow \mathcal{A}_{|D_i|}$ induced by p_i .

Example 5.7. Consider \mathbb{C} with the standard S^1 -action, and $H = 0$. Then $F = d\theta_1 \wedge d\theta_2$ provides a T -dual from $(\mathbb{C}, 0)$ to itself. \triangle

A connection one-form $\Theta \in \Omega^1(\mathcal{A}_{|D|}; \mathfrak{t})$ induces an isomorphism $\mathcal{A}_{|D|}/T^k \simeq \mathfrak{t}^k \oplus \mathcal{A}_{\partial B}$, therefore we obtain a decomposition

$$\Omega_{T^k}^3(\mathcal{A}_{|D|}) = \sum_{i=0}^3 \Omega^i(\mathcal{A}_{\partial B}, \wedge^{3-i}\mathfrak{t}^*).$$

The T -duality conditions forces the component of H in $\Omega^0(\mathcal{A}_{\partial B}; \wedge^3\mathfrak{t}^*)$ to vanish, and the component in $\Omega^1(\mathcal{A}_{\partial B}; \wedge^2\mathfrak{t}^*)$ to be exact. When considering geometric structures (up to gauge equivalence), it is the cohomology class $[H]$, rather than the form itself, which is of importance. Hence we can impose that we consider H which only have two terms in this decomposition:

$$(5.2) \quad H = \langle \hat{c}, \Theta \rangle + h,$$

for some $\hat{c} \in \Omega^2(\mathcal{A}_{\partial B}; \mathfrak{t}^*)$ and $h \in \Omega^3(\mathcal{A}_{\partial B})$. Using this decomposition, and the relation of the curvature with the Haefliger-Salem class, we can now address the question of existence of T -duals.

Theorem 5.8. *Given a manifold $M \rightarrow B$ endowed with a standard T^k and an invariant closed three-form $H \in \Omega^3(\mathcal{A}_{|D|})$. If $[\hat{c}]$ as in Equation (5.2) defines a class in $H^2(B; \mathbb{Z}^k)$, then there a T -dual $\hat{M} \rightarrow B$.*

Proof. Let $M_0 \rightarrow B$, be the reference torus action of M . Using Proposition 3.4, with respect to M_0 and \hat{M} , the class $[\hat{c}] \in H^2(B; \mathbb{Z}^k)$ corresponds to a manifold $\hat{M} \rightarrow B$ endowed with a locally equivalent action. Moreover, there exists a connection $\hat{\Theta} \in \Omega^1(\mathcal{A}_{|\hat{D}|}; \mathfrak{t}^k)$ for which $d\hat{\Theta} = \hat{c}$. Now define

$$\hat{H} = \langle c, \hat{\Theta} \rangle + h \in \Omega^3(\mathcal{A}_{|\hat{D}|}),$$

with $c = d\Theta$, for some connection $\Theta \in \Omega^1(M; \mathfrak{t}^k)$. Now (M, H) and (\hat{M}, \hat{H}) are T -dual because

$$p^*H - \hat{p}^*\hat{H} = \langle \hat{c}, \Theta \rangle - \langle c, \hat{\Theta} \rangle = \langle d\hat{\Theta}, \Theta \rangle - \langle d\Theta, \hat{\Theta} \rangle = -d\langle \Theta, \hat{\Theta} \rangle,$$

and as Θ and $\hat{\Theta}$ are connection-one-forms, we obtain that $F = -\langle \Theta, \hat{\Theta} \rangle$ satisfies the requirements of T -duality. \square

Remark 5.9. Note that it might be preferable to put a direct condition on H , rather than the $[\hat{c}]$ in the decomposition. However, this will also impose further conditions on the h term which are not strictly necessary. Still, in practice we will assume that H is smooth and defines an integral cohomology class, which implies the assumptions in Proposition 5.8 \diamond

Remark 5.10 (Non-uniqueness). This proposition provides the existence of T -duals, but these are by no means unique. Already in the case without fixed points, the construction does not take any torsion of the torus bundle into account. But also changing the specific F used may change the topology of the resulting T -dual, see also Bunke-Schick [5]. Now the non uniqueness goes even further: because we made use of the classification of Haefliger-Salem, any T -dual produced using Proposition 5.8 will necessarily be locally equivalent to M . However, that needn't be the case as we will see momentarily. \diamond

5.2.1. Topology change from singularities. The singularities of the torus actions provide a new source of topology change for T -duals:

Proposition 5.11. *Let M, \hat{M} be manifolds with standard T^k -actions over the same contractible base B . Then $(M, H = 0)$ and $(\hat{M}, \hat{H} = 0)$ are T -dual.*

Proof. Because the base of M and \hat{M} is contractible there exists flat connections $\Theta, \hat{\Theta}$ on M respectively \hat{M} . Then, as in the proof of Proposition 5.8 (the restriction of) $F = -\langle \Theta, \hat{\Theta} \rangle$ provides a T -dual between $(M, 0)$ and $(\hat{M}, 0)$. \square

In [17] a classification of four-manifolds with effective T^2 -actions and connected isotropy groups appears. All such manifolds are of the following form:

- $\#n(S^2 \times S^2)$, $n \in \mathbb{N}$
- $\#n(\mathbb{C}P^2) \# m(\overline{\mathbb{C}P^2})$, $n, m \in \mathbb{N}$
- S^4

All of these manifolds have a contractible base, so we conclude by the above Theorem that if two have the same base they are T -dual. The only invariant of the base is the number of corners, which can be read off from the Euler characteristic of the total space. For instance $2\mathbb{C}P^2$, $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$ and $S^2 \times S^2$ are all T -dual. The common base of these manifolds is contractible, and hence the actions are necessarily not locally equivalent.

Just as in principal T -duality, we obtain an isomorphism of differential complexes. The proof is omitted as it is identical:

Proposition 5.12. *Let M, \hat{M} be manifolds endowed with standard T^k -actions over the same base B . Let $H \in \Omega_{\text{cl}}^3(\mathcal{A}_{|D|})$, $\hat{H} \in \Omega_{\text{cl}}^3(\mathcal{A}_{|\hat{D}|})$, and let $F \in \Omega^2(\mathcal{A}_{|D|} \times_B \mathcal{A}_{|\hat{D}|})$ provide a T -dual. Then*

$$(5.3) \quad \tau : \Omega_{T^k}^\bullet(\mathcal{A}_{|D|}) \rightarrow \Omega_{T^k}^\bullet(\mathcal{A}_{|\hat{D}|}), \quad \tau(p) = \iota_{X_1 \wedge \dots \wedge X_n}(e^F \wedge \rho),$$

is an isomorphism of differential complexes $(\Omega_{T^k}^\bullet(M; \mathcal{A}_{|D|}), d_H)$ and $(\Omega_{T^k}^\bullet(\hat{M}; \mathcal{A}_{|\hat{D}|}), d_{\hat{H}})$. Here $X_1, \dots, X_n \in \Gamma(\mathcal{A}_{|D|})$ are the infinitesimal generators of the torus action on M .

Moreover, we may consider the standard $\mathcal{A}_{|D|}$ -Courant algebroid, $\mathcal{A}_{|D|} \oplus \mathcal{A}_{|D|}^*$, and obtain Theorem 5.4 in our setting. Again, the proof is identical to the principal case and thus omitted.

Theorem 5.13. *Let M, \hat{M} be manifolds endowed with standard T^k -actions over the same base B . Let $H \in \Omega_{\text{cl}}^3(\mathcal{A}_{|D|})$, $\hat{H} \in \Omega_{\text{cl}}^3(\mathcal{A}_{|\hat{D}|})$, and assume that (M, H) and (\hat{M}, \hat{H}) are T -dual. Then there is an isomorphism between Courant algebroids $(\mathcal{A}_{|D|} \oplus \mathcal{A}_{|D|}^*)/T^k \rightarrow M$ and $(\mathcal{A}_{|\hat{D}|} \oplus \mathcal{A}_{|\hat{D}|}^*)/T^k \rightarrow \hat{M}$.*

Using this isomorphism we are able to transport geometric structures on $\mathcal{A}_{|D|} \rightarrow M$ to geometric structures on $\mathcal{A}_{|\hat{D}|} \rightarrow \hat{M}$:

Theorem 5.14. *Let (M, H) and (\hat{M}, \hat{H}) be T -dual. Then any Dirac, generalized complex or SKT structure on $\mathcal{A}_{|D|} \rightarrow M$ invariant under the torus action will be transported to a structure of the same kind on $\mathcal{A}_{|\hat{D}|} \rightarrow \hat{M}$.*

However, there is one key difference compared to the case of principal T -duality. Given a T^k -invariant generalized complex structure on M which lifts to a generalized complex structure on $\mathcal{A}_{|D|} \rightarrow M$, we can transport it to a generalized complex structure on $\mathcal{A}_{|\hat{D}|} \rightarrow \hat{M}$. However, a priori there is no guarantee that it again descends to a generalized complex structure on M . We illustrate this phenomenon by the following examples:

Example 5.15 (\mathbb{C}). Consider $M = \mathbb{C}$ with the standard S^1 -action, and endow it with the elliptic symplectic form $d \log r \wedge d\theta = \frac{dx \wedge dy}{x^2 + y^2}$. If we let $\hat{M} = \mathbb{C}$ denote another copy of \mathbb{C} , we have that $F = d\theta \wedge d\hat{\theta}$ provides a T -dual. Under this T -duality, the elliptic symplectic structures $e^{id \log r \wedge d\theta}$ is sent to spinor $d \log \hat{z}$. This spinor corresponds to the complex structure on $\mathcal{A}_{|D|} = \{r\partial_r, \partial_\theta\}$ defined by $r\partial_r \mapsto \partial_\theta$. It is immediate to see that this complex structure is a lift of the standard complex structure on \mathbb{C} . Hence we conclude that the elliptic symplectic form, which is not a generalized complex structure, is T -dual to the standard complex structure. \triangle

The next example will illustrate that T -duality for standard torus actions is very sensitive to the specific $F \in \Omega^2(\mathcal{A}_{|D|} \times_B \mathcal{A}_{|\hat{D}|})$ chosen:

Example 5.16 (\mathbb{C}^2). Consider $M = \mathbb{C}^2$ with coordinates (z_1, z_2) and $\hat{M} = \mathbb{C}^2$ with coordinates (\hat{z}_1, \hat{z}_2) . Endow both with the standard T^2 -action. There are ample choices for a T -duality between $(M, 0)$ and $(\hat{M}, 0)$. Indeed, any of the following two-forms will suffice:

- $F_{1,\pm} = d\theta_1 \wedge d\hat{\theta}_1 \pm d\theta_2 \wedge d\hat{\theta}_2$.
- $F_{2,\pm} = -d\theta_1 \wedge d\hat{\theta}_2 \pm d\theta_2 \wedge d\hat{\theta}_1$.

Although each of these forms satisfy the conditions necessary of T -duality, the behaviour with respect to specific generalized complex structures is quite different.

We consider the stable generalized complex structure given by $\rho = z_1 z_2 + dz_1 \wedge dz_2$. The corresponding elliptic symplectic form is $\omega = d \log r_1 \wedge d\theta_2 + d\theta_1 \wedge d \log r_2$ with spinor $\rho = 1 + id \log r_1 \wedge d\theta_2 + id\theta_1 \wedge d \log r_2 - d \log r_1 \wedge d\theta_1 \wedge d \log r_2 \wedge d\theta_2$.

By Theorem 5.14 we know that T -duality with any of these forms will provide generalized complex structures on the elliptic tangent bundle. However, a direct computation will show that only $F_{2,+}$ provides a generalized complex structure on M . In that case we have that the T -dual generalized complex structure is given by:

$$\rho_{2,+} = d \log z_1 \wedge d \log z_2.$$

One immediately checks that $\rho_{2,+}$ is a lift of the standard complex structure on \mathbb{C} . In conclusion, this elliptic symplectic structure on \mathbb{C}^2 is T -dual to the standard complex structure on \mathbb{C}^2 .

The story for the real part of ρ is quite different. Consider the elliptic symplectic form $\omega = d \log r_1 \wedge d \log r_2 - d\theta_1 \wedge d\theta_2$, with spinor

$$\rho = 1 + id \log r_1 \wedge d \log r_2 - id\theta_1 \wedge d\theta_2 + d\theta_1 \wedge d\theta_2 \wedge d \log r_1 \wedge d \log r_2$$

. Again, one can perform a direct computation showing that only $F_{1,-}$ will provide a generalized complex structure on M . In that case the T -dual is given by:

$$-ie^{-i(d \log r_1 \wedge d \log r_2 + d\theta_1 \wedge d\theta_2)}$$

In conclusion we see that ρ is T -dual to itself. \triangle

6. BLOW-UPS

In this section we will study the interaction between elliptic geometry and blow-ups. We will work with the formalism of generalized complex blow-ups introduced in [2]. We will show that blow-ups of submanifolds of the degeneracy locus of an elliptic divisor induce Lie algebroid submersions between the respective elliptic tangent bundles. Torus actions are preserved by blow-ups at fixed points, and hence we may consider the relation with T -duality. We will show that T -duality and blowing up commutes.

6.1. Blow-ups in generalized complex geometry. The contents of this subsection appear in [2], we recall them for completeness.

As we do not assume any holomorphicity of our manifolds, more data is necessary in constructing a blow-up:

Definition 6.1 ([2]). Let M^n be a smooth manifold, and let Y be a closed embedded submanifold of codimension $2l$. A **holomorphic ideal** for Y is a sheaf of ideals $I_Y \subset C^\infty(M; \mathbb{C})$ such that:

- $I_Y|_{M \setminus Y} = C^\infty(M; \mathbb{C})|_{M \setminus Y}$.
- For every $y \in Y$, there is a neighbourhood U with coordinates $z_1, \dots, z_l, x_{2l+1}, \dots, x_n : U \rightarrow \mathbb{C}^l \times \mathbb{R}^{n-2l}$ such that $Y \cap U = \{z_1 = \dots = z_l = 0\}$ and $I_Y|_U = \langle z_1, \dots, z_l \rangle$. \diamond

As shown in [2], a submanifold Y admits a holomorphic ideal if and only if the normal bundle NY admits a complex structure.

Theorem 6.2 ([2]). *Given a submanifold Y and a holomorphic ideal I_Y , there exists a, unique up to isomorphism, manifold \tilde{M} together with a map $p : \tilde{M} \rightarrow M$ such that p^*I_Y is a divisor satisfying the following universal property: For any smooth map $f : X \rightarrow M$, such that f^*I_Y is a divisor, there exists a unique $\tilde{f} : X \rightarrow \tilde{M}$ such that the following diagram commutes:*

$$\begin{array}{ccc} X & \xrightarrow{\tilde{f}} & \tilde{M} \\ & \searrow f & \downarrow p \\ & & M \end{array}$$

We call \tilde{M} the **blow-up of I_Y in M** , and $p : \tilde{M} \rightarrow M$ the blow-down map. The natural manifolds one can blow-up in generalized complex geometry are the following:

Definition 6.3 ([2]). Let (M, \mathbb{J}, H) be a generalized complex manifold. A submanifold Y is called **generalized Poisson** if

$$\mathbb{J}N^*Y = N^*Y. \quad \diamond$$

An immediate consequence of this definition is that Y is a Poisson submanifold of the underlying Poisson structure $\pi_{\mathbb{J}}$. Moreover, if Q is a holomorphic Poisson submanifold and \mathbb{J}_Q is the associated generalized complex structure, a submanifold Y is a generalized Poisson submanifold if and only if it is a holomorphic Poisson submanifold.

By definition, the generalized complex structure induces a complex structure on the normal bundle of a generalized Poisson submanifold. Hence:

Lemma 6.4 ([2]). *Let (M, \mathbb{J}, H) be a generalized complex manifold and Y a generalized Poisson submanifold. Then there exist a canonical holomorphic ideal I_Y for Y for which the induced complex structure on NY coincides with the one induced by \mathbb{J} .*

Given that a generalized Poisson submanifold gives rise to a holomorphic ideal, and hence a canonical blow-up, the next step is to determine when the generalized complex structure lifts to the blow-up. For this we need the following notion:

Definition 6.5. A Lie algebra \mathfrak{g} is said to be degenerate if the map $\Lambda^3(\mathfrak{g}) \rightarrow \text{Sym}^2(\mathfrak{g})$

$$x \wedge y \wedge z \mapsto [x, y] \wedge z,$$

vanishes. ◇

Given a generalized Poisson submanifold $Y \subset (M, \mathbb{J})$, it is in particular a Poisson submanifold and hence one obtains a fibre-wise Lie algebra structure on N^*Y . This Lie algebra structure captures whether the generalized complex structure lifts to the blow-up:

Theorem 6.6 ([2]). *Let (M, \mathbb{J}, H) be a generalized complex manifold and Y a generalized Poisson submanifold. Then there exists a generalized complex structure $\tilde{\mathbb{J}}$ on the blow-up $p : \tilde{M} \rightarrow M$, with respect to the canonical holomorphic ideal I_Y , for which p is a generalized holomorphic map, if and only if the Lie algebra structure on N^*Y is degenerate.*

We may blow-up components of the degeneracy locus of a stable generalized complex structure:

Lemma 6.7. *Let (M, \mathbb{J}) be a stable generalized complex manifold, and let $Y \subset D(l)$ be an irreducible component. Then Y is a generalized Poisson submanifold for which the fibre-wise Lie algebra structure on N^*Y is trivial.*

Proof. First we note that Y is a union of symplectic leaves of the underlying Poisson structure $\pi_{\mathbb{J}}$, and hence a Poisson submanifold. For every $p \in Y$, there exists a neighbourhood U , with a complex structure and a holomorphic Poisson structure Q inducing the generalized complex structure (up to gauge equivalence). We have that $Y \cap U$, can be written as the zero-set of some of the complex coordinates, and hence it is a complex submanifold of U . Therefore, $Y \cap U$ is a holomorphic Poisson submanifold of Q , and consequently a generalized Poisson submanifold of \mathbb{J} on U . As being a generalized Poisson submanifold is a point-wise condition, this shows that Y is a generalized Poisson submanifold of M .

Because $\pi_{\mathbb{J}}$ is an elliptic symplectic structure it lifts to an element of $\Gamma(\wedge^2 \mathcal{A}_{|D|})$, in particular the bivector is tangent to D . Because of the geometry of D , it follows that $\pi_{\mathbb{J}}$ is tangent to each stratum $D[l]$, and therefore in particular to Y . Consequently, the Lie algebra structure induced on N^*Y is trivial. □

We conclude that irreducible components of the degeneracy locus of a stable generalized complex structure may be blown-up in a generalized complex fashion.

6.2. Blowing up for the elliptic tangent bundle. There is yet another explanation why one can blow up irreducible components of the degeneracy locus of a stable generalized complex structure. Namely, the elliptic tangent bundle is very well-behaved under blow-ups. To state a result independent of a generalized complex structure, we need that the holomorphic ideal is adapted to the elliptic ideal:

Definition 6.8. Let M^n be endowed with an elliptic divisor $I_{|D|}$, and Y^{n-2l} a closed embedded submanifold. We say that a holomorphic ideal I_Y for Y is **adapted** to $I_{|D|}$ if there exists local coordinates $z_1, \dots, z_k, x_{2k+1}, \dots, x_{n-2k}$ for which z_1, \dots, z_l are as in Definition 6.1 and $I_{|D|}|_U = \langle |z_1|^2 \cdot \dots \cdot |z_k|^2 \rangle$. ◇

Note that this definition in particular implies that $Y \subset D$. Given such a holomorphic ideal, we may blow-up the elliptic divisor to obtain:

Theorem 6.9. *Let $(M, I_{|D|})$ be a manifold endowed with an elliptic divisor, and let $Y \subset D(k)$ be a irreducible component. Assume that we are given a holomorphic ideal I_Y for Y adapted*

to $I_{|D|}$. Then there exists an elliptic ideal $I_{|\tilde{D}|}$ on the blow-up \tilde{M} such that the blow-down map induces a fibre-wise isomorphism:

$$p_* : (\mathcal{A}_{|\tilde{D}|})_x \xrightarrow{\cong} (\mathcal{A}_{|D|})_{p(x)},$$

for all $x \in \tilde{M}$.

Proof. To study the blow-down map we need to use local coordinates describing it. Given $l \geq k$ let $x \in Y \cap D[l]$, let $U = \mathbb{C}^l \times \mathbb{R}^m$ be a local chart with coordinates $(z_1, \dots, z_l, x_1, \dots, x_m)$, with the z_i as in Definition 6.1. Let $\tilde{U} = \tilde{\mathbb{C}}^l \times \mathbb{R}^m$, and $p = (p', \text{Id}) : \tilde{U} \rightarrow U$ be the blow-down map. The manifold $\tilde{\mathbb{C}}^l$ has a cover of l -coordinate charts given by

$$(v_1, \dots, v_{i-1}, z_i, v_{i+1}, \dots, v_l) \leftrightarrow (z_i \cdot (v_1, \dots, v_{i-1}, 1, v_{i+1}, \dots, v_l), \\ [v_1 : \dots : v_{i-1} : 1 : v_{i+1} : \dots : v_l]).$$

In these coordinate charts the blow-down map is given by

$$p' : (v_1, \dots, v_{i-1}, z_i, v_{i+1}, \dots, v_l) \mapsto (z_i v_1, \dots, z_i v_{i-1}, z_i, z_i v_{i+1}, \dots, z_i v_l).$$

With these coordinates at hand, we can consider the local form of $p^* I_{|D|}$. Note that this does not provide an elliptic divisor yet, however $I_{|D|} := \sqrt{p^* I_{|D|}}$ does. Because a vector field preserves $p^* I_{|D|}$ if and only if it preserves $\sqrt{p^* I_{|D|}}$ we find that the blow-down map induces a Lie algebroid map $p_* : \mathcal{A}_{|\tilde{D}|} \rightarrow \mathcal{A}_{|D|}$.

To check that the blow-down map p induces a fibre-wise isomorphism, it suffices to show that it pulls-back a volume element $(\mathcal{A}_{|D|})_{p(x)}$ to a volume element of $(\mathcal{A}_{|\tilde{D}|})_x$. In the above coordinates, the following would define a local volume form on $(\mathcal{A}_{|D|})_x$:

$$\Omega = d \log r_1 \wedge d\theta_1 \wedge \dots \wedge d \log r_l \wedge d\theta_l \wedge dx_1 \wedge \dots \wedge dx_m.$$

Therefore

$$p^* \Omega = d \log \tilde{r}_1 \wedge d\hat{\theta}_1 \wedge \dots \wedge d \log \tilde{r}_l \wedge d\hat{\theta}_l \wedge d \log r_{l+1} \wedge d\theta_{l+1} \wedge \dots \wedge d \log r_k \wedge d\theta_k \wedge dx_1 \wedge \dots \wedge dx_m,$$

where the $(\tilde{r}_j, \hat{\theta}_j)$ are the polar coordinates corresponding to $(v_1, \dots, v_{i-1}, z_i, v_{i+1}, \dots, v_l)$. We conclude that p indeed induces the fibrewise isomorphism as required. \square

Example 6.10 (Blowing up points). Let M be an orientable manifold and let $Y = \{p\}$ be a point. Then we can always find complex coordinates around p , z_1, \dots, z_l , such that $I_Y = \langle z_1, \dots, z_l \rangle$ is a holomorphic ideal. The topology of the blow-up will depend on the parity of l , and whether the coordinates z_1, \dots, z_l are orientation preserving. If l is odd, the blow-up will be $M \# \mathbb{C}P^l$, if l is even it will be $M \# \overline{\mathbb{C}P^l}$ if the coordinates are orientated and $M \# \mathbb{C}P^l$ if they are not. \triangle

Example 6.11 (Global normal crossing). Let $I_{|D|}$ be a global normal crossing divisor, and let $Y \subset D(l)$ be an irreducible component. We have that Y must then necessarily be of the form $D_1 \cap \dots \cap D_l$, for some of the components of D . If $D[1]$ is co-orientable, it follows that each D_i is co-orientable as well, and one can show that there exists a complex structure on ND_i , inducing the elliptic divisor (see [9]). Combining these complex structures to a complex structure on NY , gives a holomorphic ideal for Y which is compatible with $I_{|D|}$. \triangle

Given a generalized complex structure on any of the Lie algebroids $\mathcal{A}_{|D|}$ may lift it to a structure of the same kind on the blown-up Lie algebroid. But similarly as for T -duality, it might happen that said structure does not descend to give a generalized complex structure on \tilde{M} . However, when the holomorphic ideal is induced by the generalized complex structure, this will always be the case:

Lemma 6.12. *Let (M, \mathbb{J}) be a stable generalized complex manifold, and let $Y \subset D(k)$ be an irreducible component. Let $\omega \in \Omega^2(\mathcal{A}_{|D|})$ denote the elliptic symplectic structure corresponding to \mathbb{J} . Let I_Y be the canonical holomorphic ideal for Y . Then this ideal is adapted to $I_{|D|}$, and the blow-up $p^*\omega$ will correspond to the generalized complex blow-up of \mathbb{J} as in Lemma 6.7.*

We conclude that the point of view of blowing-up using the elliptic tangent bundle results in the same blow-up as in [2]. However, Theorem 6.9 can be used in a larger context: for instance, elliptic symplectic forms not corresponding to generalized complex structures may be blown-up using this result.

6.2.1. Complex structures on the blow-up. Let $(M, I_{|D|})$ is a manifold endowed with an elliptic divisor, and \mathbb{J} is a generalized complex structure on M , which lifts to $\mathcal{A}_{|D|}$. As pointed out before, if we are given $Y \subset D(k)$ an irreducible component, and I_Y a holomorphic ideal, then the blow-up of \mathbb{J} induces a generalized complex structure on $\mathcal{A}_{|\tilde{D}|}$, however this needn't descend to a generalized complex structure on \tilde{M} . The following example shows this can already happen with complex structures:

Example 6.13 (Complex structures on $n\mathbb{CP}^2 \# m\overline{\mathbb{CP}^2}$). Consider the standard complex structure J on \mathbb{CP}^2 , and $\mathcal{O}(3)$ and the section s corresponding to the polynomial $z_0 z_1 z_2$. Then $I_D := s^*(\Gamma(\mathcal{O}(3))) \subset C^\infty(\mathbb{CP}^2, \mathbb{C})$ defines a complex log divisor with degeneracy locus $D = \{[0 : z_1 : z_2]\} \cup \{[z_0 : 0 : z_2]\} \cup \{[z_0 : z_1 : 0]\}$, and let $I_{|D|}$ be the associated elliptic divisor. As D is a complex submanifold of \mathbb{CP}^2 , J will preserve the log tangent bundle A_D , and consequently induce a complex structure $J_{\mathcal{A}_{|D|}}$ on $\mathcal{A}_{|D|} \rightarrow \mathbb{CP}^2$.

We may also consider $\overline{\mathbb{CP}^2}$, with the same elliptic divisor $I_{|D|}$. Although this is not an almost complex manifold, it does admit a complex structure on $\mathcal{A}_{|D|}$. Indeed, one may show that in affine coordinates (w_1, w_2) , the spinor $\rho = d \log w_1 \wedge d \log \overline{w}_2$ gives rise to a global complex structure on $\mathcal{A}_{|D|} \rightarrow \overline{\mathbb{CP}^2}$.

Now, let $Y = \{\text{pt}\} \subset \mathbb{CP}^2$ be any of the vertices of D , and pick a holomorphic ideal for Y . As described in Example 6.10, the corresponding total space will be either $\mathbb{CP}^2 \# \mathbb{CP}^2$ or $\mathbb{CP}^2 \# \overline{\mathbb{CP}^2}$. In either case, we obtain a complex structure $J_{\mathcal{A}_{|\tilde{D}|}}$ on the blown-up Lie algebroid $\mathcal{A}_{|\tilde{D}|}$.

On $\mathbb{CP}^2 \# \overline{\mathbb{CP}^2}$, the blow-up coincides with the usual complex blow-up and $J_{\mathcal{A}_{|\tilde{D}|}}$ is simply a lift of the complex structure on $\mathbb{CP}^2 \# \overline{\mathbb{CP}^2}$. However, $\mathbb{CP}^2 \# \mathbb{CP}^2$ is not even an almost complex manifold, and hence the induced complex structure on $J_{\mathcal{A}_{|\tilde{D}|}}$ does not descend to a (generalized) complex structure on $\mathbb{CP}^2 \# \mathbb{CP}^2$. Continuing inductively, we obtain complex structures on elliptic tangent bundles over $n\mathbb{CP}^2 \# m\mathbb{CP}^2$, for all $n, m \in \mathbb{N}$. \triangle

6.3. T -duality and blow-up. We now study how T -duality interacts with blow-ups. First we recall that toric actions are preserved under blow-ups:

Lemma 6.14. *Let M^{2n} admit a standard T^n action, and let $p \in M$ be a fixed point. Then, both $M \# \mathbb{CP}^n$ and $M \# \overline{\mathbb{CP}^n}$ admits a standard T^n -action for which the blow-down map is equivariant.*

The fact that the blow-down map induces a fibre-wise isomorphism between elliptic tangent bundles immediately implies the following:

Proposition 6.15 (T -duality commutes with blow-up). *Let M^{2n} and \hat{M}^{2n} be endowed with standard T^n -actions over the same base B , and let $I_{|D|}, I_{|\hat{D}|}$ denote the induced elliptic divisors.*

Let $H \in \Omega_{\text{cl}}^3(\mathcal{A}_{|D|})$, $\hat{H} \in \Omega_{\text{cl}}^3(\mathcal{A}_{|\hat{D}|})$, and $F \in \Omega^2(\mathcal{A}_{|D|} \times_B \mathcal{A}_{|\hat{D}|})$ provide a T -dual between (M, H) and (\hat{M}, \hat{H}) .

Let $x \in M, \hat{x} \in \hat{M}$ be fixed points for the action, and let X, Y denote either $\mathbb{C}P^n$ or $\overline{\mathbb{C}P^n}$. Let $q : M \#_x X \rightarrow M$, $\hat{q} : \hat{M} \#_{\hat{x}} Y \rightarrow \hat{M}$ denote the blow-down maps. Then we have a canonical T -duality between the blow-ups:

$$\begin{array}{ccc}
 ((M \# X) \times_{\tilde{B}} (\hat{M} \# Y), p^* q^* H - \hat{p}^* \hat{q}^* \hat{H}) & & \\
 \swarrow p & & \searrow \hat{p} \\
 (M \# X, q^* H) & & (\hat{M} \# Y, \hat{q}^* \hat{H}) \\
 \searrow \pi & & \swarrow \hat{\pi} \\
 & \tilde{B} &
 \end{array}$$

Consequently, the following diagram commutes:

$$\begin{array}{ccc}
 \Omega^\bullet(\mathcal{A}_{|\tilde{D}|}) & \xrightarrow{\tilde{\tau}} & \Omega^\bullet(\mathcal{A}_{|\hat{\tilde{D}}|}) \\
 q^* \uparrow & & \hat{q}^* \uparrow \\
 \Omega^\bullet(\mathcal{A}_{|D|}) & \xrightarrow{\tau} & \Omega^\bullet(\mathcal{A}_{|\hat{D}|})
 \end{array}$$

Proof. By Theorem 6.9 q and \hat{q} induce fibre-wise isomorphisms between elliptic tangent bundles. Let $Q : (M \# X) \times_{\tilde{B}} (\hat{M} \# Y) \rightarrow M \times_B \hat{M}$ denote the induced map between the correspondence spaces. Consequently Q induces a fibre-wise isomorphism between the elliptic tangent bundles on the correspondence spaces. Therefore $Q^* F$ will satisfy the equation for T -duality. \square

Corollary 6.16. Let (M, H) , (\hat{M}, \hat{H}) be as in the setting of Proposition 6.15, and assume there is a generalized complex structure \mathbb{J} on (M, H) T -dual to a generalized complex structure $\hat{\mathbb{J}}$ on (\hat{M}, \hat{H}) . Then the T -duality from the previous Proposition 6.15 sends $\tilde{\mathbb{J}}$ to $\hat{\tilde{\mathbb{J}}}$.

Proof. As $\tilde{M} \setminus E \simeq \hat{M} \simeq \{\hat{x}\}$, we have that the T -dual of $\tilde{\mathbb{J}}$ coincides with $\hat{\tilde{\mathbb{J}}}$ outside of E . By continuity they must coincide over the entirety of \tilde{M}_2 . \square

7. EXAMPLES

In this final section we will provide explicit examples of T -dual geometric structures on manifolds endowed with standard torus actions. We will define some basic examples, and use these to produce more examples by blowing up.

7.1. Basic examples. We start by consider simple examples of generalized complex structures. The analysis heavily depends on the local description in Example 5.16. So recall from there that any of the following two-forms:

- $F_{1,\pm} = d\theta_1 \wedge d\hat{\theta}_1 \pm d\theta_2 \wedge d\hat{\theta}_2$.
- $F_{2,\pm} = -d\theta_1 \wedge d\hat{\theta}_2 \pm d\theta_2 \wedge d\hat{\theta}_1$.

provides a T -duality from $(\mathbb{C}, 0)$ to itself.

Example 7.1 ($S^1 \times S^3$). Let T^2 act on $S^3 \subset \mathbb{C}^2$, and extend the action trivially to an action on $S^1 \times S^3$. The quotient space B is an annulus. Viewing $S^1 \times S^3$ as the Hopf surface $(\mathbb{C}^2 \setminus \{0\})/\mathbb{Z}$, the stable generalized complex structure $\tilde{\rho} = z_1 z_2 + dz_1 \wedge dz_2$ on $\mathbb{C}^2 \setminus \{0\}$ descends to a stable generalized complex structure ρ on $S^1 \times S^3$. All the two forms $F_{i,\pm}$ are \mathbb{Z} -invariant and thus

descend to two-forms on $S^1 \times S^3$. Therefore, as in Example 5.16, we obtain that T -duality induced by the two-form $F_{2,+}$ sends ρ to the standard complex structure on $S^1 \times S^3$.

If we instead consider the generalized complex structure induced by $\tilde{\rho}_2 = z_1 z_2 + idz_1 \wedge dz_2$, and use the two-form induced by $F_{1,-}$, we find that ρ_2 is T -dual to itself. \triangle

Example 7.2 (\mathbb{CP}^2). We consider \mathbb{CP}^2 in the usual fashion as $(\mathbb{C}^3 \setminus \{0\})/\mathbb{C}$ with torus action $(\lambda_1, \lambda_2) \cdot [z_0 : z_1 : z_2] = [e^{i\lambda_1} z_0 : e^{i\lambda_2} z_1 : z_2]$. Consider the complex log forms $\tilde{\zeta}_1 = d \log z_0 - d \log z_2$ and $\tilde{\zeta}_2 = d \log z_1 - d \log z_2$. As these are basic with respect to the \mathbb{C}^* -action, they descend to complex log forms ζ_1, ζ_2 . We let $\Theta_1 = \Im^* \zeta_1, \Theta_2 = \Im^* \zeta_2$ denote the imaginary parts. Then (Θ_1, Θ_2) provides a connection-one form for the T^2 -action on \mathbb{CP}^2 .

Consider another copy of \mathbb{CP}^2 , and denote the same connection one-form by $(\hat{\Theta}_1, \hat{\Theta}_2)$. Similarly to Example 5.16 we now have that any of the following two-forms provides a self T -dual for $(\mathbb{CP}^2, 0)$.

$$\begin{aligned} F_{1,\pm} &= \Theta_1 \wedge \hat{\Theta}_1 \pm \Theta_2 \wedge \hat{\Theta}_2, \\ F_{2,\pm} &= -\Theta_1 \wedge \hat{\Theta}_2 \pm \Theta_2 \wedge \hat{\Theta}_1. \end{aligned}$$

We will now the action of this T -duality to the generalized complex structures on \mathbb{CP}^2 .

Consider the holomorphic Poisson structure $\tilde{\pi} = z_0 z_1 \partial_{z_0} \wedge \partial_{z_1}$. Because this is \mathbb{C}^* -invariant it induces a holomorphic Poisson structure π on \mathbb{CP}^2 , which is moreover log symplectic. One can show, as in Example 5.16, that $F_{2,+}$ provides a T -dual between the corresponding stable generalized complex structure and the standard complex structure on \mathbb{CP}^2 . Similarly, consider the holomorphic Poisson structure $\tilde{\pi}_2 = iz_0 z_1 \partial_{z_0} \wedge \partial_{z_1}$, and the induced generalized complex structure ρ_2 . Then $F_{1,-}$, as in Example 5.16, will provide a T -dual between ρ_2 and itself. \triangle

Example 7.3 ($S^2 \times S^2$). Identify $M = S^2 \times S^2$ with $\mathbb{CP}^1 \times \mathbb{CP}^1$, and consider the torus action given by $(\lambda_1, \lambda_2) \cdot ([z_0 : z_1], [w_0 : w_1]) = ([e^{i\lambda_1} z_0 : z_1], [e^{i\lambda_2} w_0 : w_1])$. Let $\tilde{\zeta}_1 = d \log z_0 - d \log z_1$ and $\tilde{\zeta}_2 = d \log w_0 - d \log w_1$ be complex log-forms on $(\mathbb{C}^2 \setminus \{0\})^2$. As these are basic they induce complex log-forms ζ_1, ζ_2 on M , and we may consider their imaginary parts $\Theta_1 = \Im^* \zeta_1, \Theta_2 = \Im^* \zeta_2$. The form $(\Theta_1, \Theta_2) \in \Omega^2(\mathcal{A}_{|D|}; \mathfrak{t}^2)$ provides a connection one-form for the torus action on M . Let \hat{M} denote another copy of $S^2 \times S^2$, and consider the same connection one-form $(\hat{\Theta}_1, \hat{\Theta}_2)$. As in the previous example, the form $F = -\Theta_1 \wedge \hat{\Theta}_2 + \Theta_2 \wedge \hat{\Theta}_1$ provides a self T -dual on $(M, 0)$.

In affine coordinates $t = z_0/z_1, s = w_0/w_1$ we consider the holomorphic Poisson structure $\pi = tw \partial_t \partial_w$. In any other set of affine coordinates, this Poisson structure has the same form and hence defines a global holomorphic Poisson structure on M , which is moreover log symplectic. As in the previous example, the corresponding stable generalized complex structure is T -dual to the standard complex structure on $S^2 \times S^2$. And the stable generalized complex structure associated to $\pi = itw \partial_t \partial_w$ is T -dual to itself via $F = \Theta_1 \wedge \hat{\Theta}_1 - \Theta_2 \wedge \hat{\Theta}_2$. \triangle

We may now blow-up \mathbb{CP}^2 to obtain more examples of stable generalized complex structures, which are T -dual:

Example 7.4 ($n\mathbb{CP}^2 \# m\overline{\mathbb{CP}^2}$). Let $\tilde{\pi}_1 = z_0 z_1 \partial_{z_0} \wedge \partial_{z_1}$, and $\tilde{\pi}_2 = iz_0 z_1 \partial_{z_0} \wedge \partial_{z_1}$ be holomorphic Poisson structures on $\mathbb{C}^3 \setminus \{0\}$, and ρ_1, ρ_2 the induced stable generalized complex structures on \mathbb{CP}^2 , as in Example 7.2. We may perform consecutive blow-ups of either of these structures, to obtain families of generalized complex structures $\rho_{1,n,m}, \rho_{2,n,m}$ on $\mathcal{A}_{|D|} \rightarrow n\mathbb{CP}^2 \# m\overline{\mathbb{CP}^2}$. One readily shows that these generalized complex structures on $\mathcal{A}_{|D|}$, induced generalized complex structures on $n\mathbb{CP}^2 \# m\overline{\mathbb{CP}^2}$ if and only if n is odd.

We have seen in Example 7.2 that ρ_1 can be made T -dual to the standard complex structure, and ρ_2 dual to itself. By Proposition 6.15 it then follows that each $\rho_{2,n,m}$ is T -dual to itself, and $\rho_{1,n,m}$ is T -dual to the complex structures on $\mathcal{A}_{|D|}$ described in Example 6.13. \triangle

Remark 7.5. We obtained the families $\rho_{2,n,m}$ also in Theorem 7.5 in [9] via a connected sum procedure. But not the families $\rho_{1,n,m}$. This is of relevance, as we see that their behaviour under T -duality is very different. \diamond

Example 7.6 ($n\mathbb{C}P^{2k} \# m\overline{\mathbb{C}P}^{2k}$). The previous example is in no way restrictive to four dimensions. One may as easily start with the Poisson structure $z_0 z_1 \partial_{z_0} \wedge \partial_{z_1} + \cdots + z_{2(k-1)} z_{2k} \partial_{z_{2(k-1)}} \wedge \partial_{z_{2k}}$ on \mathbb{C}^{2k+1} , and repeat the above to obtain T -dual generalized complex structures on $n\mathbb{C}P^{2k}$. After blowing up at fixed points one again obtains many families of T -dual generalized complex structures on $n\mathbb{C}P^{2k} \# m\overline{\mathbb{C}P}^{2k}$. \triangle

7.2. Non-zero residue elliptic symplectic structures. Another class of structures which behave well with respect to T -duality are elliptic symplectic forms with non-zero elliptic residue. These structures admit a \mathbb{C}^* -invariant normal form:

Lemma 7.7 ([21]). *Let $(M, I_{|D|})$ be a manifold with a smooth elliptic divisor, and let $\omega \in \Omega^2(\mathcal{A}_{|D|})$ be an elliptic symplectic form with $\lambda := \text{Res}_q(\omega) \neq 0$. Then there exists a canonical flat connection on ν_D and a tubular neighbourhood of D such that*

$$(7.1) \quad \omega = \lambda \rho \wedge \Theta + p^* \omega_D.$$

Here ρ, Θ are the connection one-forms corresponding to the flat connection, as defined in 3.16 and $\omega_D = \iota_D^* \omega \in \Omega^2(D)$.

Conversely, given any complex line bundle $L \rightarrow D$, ∇ a flat connection on L , $\lambda \neq 0$ and $\omega_D \in \Omega^2(D)$ symplectic, (7.1) defines an elliptic symplectic form on L .

The form ω is S^1 -invariant, and hence we can consider its T -dual. It will appear via the following lemma:

Lemma 7.8. *Let (M, J) be a generalized complex manifold, and $L \rightarrow M$ a complex vector bundle, endowed with a flat connection ∇ . Then $\text{tot}(L)$ inherits a natural generalized complex structure.*

Proof. We define \tilde{J} on $\text{tot}(L)$ by specifying it on linear and fibrewise constant vector fields. Given $X \in T_p M$, we define $\tilde{J}(\text{hor}^\nabla(X)) = \text{hor}^\nabla(J(X))$. And given $v \in T_p^{\text{vert}} L$, we define $\tilde{J}(v) = iv$, under the identification of $T_p^{\text{vert}} L \simeq L_p$. Because the connection is flat, \tilde{J} will be integrable. \square

Lemma 7.9. *Given (D, ω_D) a symplectic manifold, $p: L \rightarrow M$ a complex line bundle with flat connection ∇ , and $\lambda \in \mathbb{R} \setminus \{0\}$ then ω as in (7.1) is T -dual to the type 1 generalized complex structure induced by Lemma 7.8*

Proof. Let $\hat{\Theta}$ denote the same connection one-form, on another copy of $L \rightarrow M$. Then $F = \Theta \wedge \hat{\Theta}$ provides the desired T -dual. We have that, using τ from Proposition 5.12 that

$$\tau(e^{i\omega}) = i\lambda(\rho + i\Theta) \wedge e^{i\omega_D}.$$

Locally, this corresponds to the spinor given by $i\lambda dz \wedge e^{i\omega_D}$, which is precisely the generalized complex structure form Lemma 7.8. \square

Remark 7.10. Given a smooth elliptic divisor $I_{|D|}$, elliptic symplectic forms with zero elliptic residue have a similar S^1 -invariant normal form as in (7.1). Unfortunately, the corresponding T -dual generalized complex structure on $\mathcal{A}_{|D|}$ does not descend to a generalized complex structure on M . \diamond

REFERENCES

- [1] D. Auroux, L. Katzarkov, and D. Orlov, *Mirror symmetry for del Pezzo surfaces: vanishing cycles and coherent sheaves*, Invent. Math. **166** (2006), no. 3, 537–582.
- [2] M. Bailey, G. Cavalcanti, and J. Duran, *Blow-ups in generalized complex geometry*, Transactions of the American Mathematical Society **371** (2016), 2109–2131.
- [3] P. Bouwknegt, J. Evslin, and V. Mathai, *T-Duality: Topology Change from H-Flux*, Communications in Mathematical Physics **249** (2004), 383–416.
- [4] P. Bouwknegt, K. Hannabuss, and V. Mathai, *T-duality for principal torus bundles*, Journal of High Energy Physics **2004** (2004), no. 03, 018.
- [5] U. BUNKE and T. SCHICK, *ON THE TOPOLOGY OF T-DUALITY*, Reviews in Mathematical Physics **17** (2005), no. 01, 77–112, <https://doi.org/10.1142/S0129055X05002315>.
- [6] T. Buscher, *A symmetry of the string background field equations*, Physics Letters B **194** (1987), no. 1, 59–62.
- [7] G. Cavalcanti and M. Gualtieri, *Generalized complex geometry and T-duality*, A Celebration of the Mathematical Legacy of Raoul Bott (CRM Proceedings and Lecture Notes) (2010), 341–366.
- [8] G. R. Cavalcanti and M. Gualtieri, *Stable generalized complex structures*, Proc. Lond. Math. Soc. **116** (2018), no. 5, 1075–1111.
- [9] G. R. Cavalcanti, R. L. Klaasse, and A. Witte, *Self-crossing stable generalized complex structures*, Journal of symplectic geometry **20** (2022), 761–811.
- [10] A. Fino and A. Tomassini, *Blow-ups and resolutions of strong Kähler with torsion metrics*, Adv. Math. **221** (2009), no. 3, 914–935.
- [11] M. Gross, P. Hacking, and S. Keel, *Mirror symmetry for log Calabi-Yau surfaces I*, Publ. Math. Inst. Hautes Études Sci. **122** (2015), 65–168.
- [12] M. Gualtieri, S. Li, A. Pelayo, and T. S. Ratiu, *The tropical momentum map: a classification of toric log symplectic manifolds*, Math. Ann. **367** (2017), no. 3–4, 1217–1258.
- [13] A. Haefliger and E. Salem, *Actions of Tori on Orbifolds*, Annals of Global Analysis and Geometry (1991), 37–59.
- [14] K. Hori and C. Vafa, *Mirror Symmetry*, 2000. [arXiv:hep-th/0002222](https://arxiv.org/abs/hep-th/0002222) [hep-th].
- [15] D. McDuff, *Examples of simply-connected symplectic non-Kählerian manifolds*, J. Differential Geom. **20** (1984), no. 1, 267–277.
- [16] E. Miranda and G. Scott, *The geometry of E-manifolds*, 2018. [arXiv:1802.02959](https://arxiv.org/abs/1802.02959) [math.SG].
- [17] P. Orlik and F. Raymond, *Actions of the Torus on 4-Manifolds. I*, Transactions of the American Mathematical Society **152** (1970), no. 2, 531–559.
- [18] M. Roček and E. Verlinde, *Duality, quotients, and currents*, Nuclear Physics B **373** (1992), no. 3, 630–646.
- [19] B. Sathiapalan, *Duality in statistical mechanics and string theory*, Phys. Rev. Lett. **58** (1987), 1597–1599.
- [20] G. W. Schwarz, *Smooth functions invariant under the action of a compact Lie group*, Topology **14** (1975), 63–68.
- [21] A. Witte, *Between generalized complex and Poisson geometry*, PhD thesis, Utrecht University (2021), 1–211.

DEPARTMENT OF MATHEMATICS, UTRECHT UNIVERSITY, 3508 TA UTRECHT, THE NETHERLANDS
 Email address: g.r.cavalcanti@uu.nl

DEPARTEMENT OF MATHEMATICS, CELESTIJNENLAAN 200B, B-3001 LEUVEN (HEVERLEE), BELGIUM
 Email address: aldowitte@hotmail.nl