

TTK-22-43, P3H-22-120
December 6, 2022

$t\bar{t}Z$ in the 4ℓ channel at NLO in QCD

JASMINA NASUFI¹

*Institute for Theoretical Particle Physics and Cosmology RWTH-Aachen University,
D-52056 Aachen, Germany*

NLO QCD corrections to the process $pp \rightarrow e^+ \nu_e \mu^- \bar{\nu}_\mu \tau^+ \tau^- b \bar{b} + X$ are presented with the full off-shell effects included. The calculation includes all resonant and non-resonant Feynman diagrams, photon and Z -gauge boson contributions, as well as interference effects, all incorporated at the matrix element level. Furthermore all heavy intermediate particles are described via Breit-Wigner propagators. Theoretical uncertainties related to the scale variation and choice of PDF sets were investigated at the integrated and differential level. The modelling is studied by a direct comparison of the full off-shell process to the NWA, where tops, Z - and W -gauge bosons are on-shell. Moreover, motivated by experimental cuts, we also investigate the impact of imposing a window cut around the mass of the Z -gauge boson on the full off-shell predictions.

PRESENTED AT

15th International Workshop on Top Quark Physics
Durham, UK, 4–9 September, 2022

¹The work of Jasmina Nasufi was supported by the Deutsche Forschungsgemeinschaft (DFG) under grant 396021762 - TRR 257: P3H - Particle Physics Phenomenology after the Higgs Discovery and under grant 400140256 - GRK 2497: The physics of the heaviest particles at the Large Hadron Collider.

1 Introduction

The associated production of a top pair and a Z -gauge boson is a very important process to study at the LHC. It features some of the heaviest particles known to-date and its signature can therefore be modified by various new physics scenarios (e.g. ref. [1, 2]). Furthermore, $t\bar{t}Z$ comprises an important background to the SM process $t\bar{t}H$ (e.g. ref. [3]) and other BSM processes (e.g. ref. [4]) in the multi-lepton channels. It has been measured at the LHC by both ATLAS and CMS (e.g. ref [5, 6]). In this work, we focus on $pp \rightarrow e^+\nu_e\mu^-\bar{\nu}_\mu\tau^+\tau^-b\bar{b} + X$ at NLO in QCD with all full off-shell effects included. All leptons and b -jets are treated as massless. The default fiducial phase space region is defined by applying transverse momentum, rapidity and ΔR cuts on all final states. For details, please see ref. [7].

2 Results

The integrated fiducial cross sections at LO and NLO in QCD, with a dynamic scale setting and the NNPDF3.1 PDF set are:

$$\sigma_{\text{full off-shell}}^{\text{LO}} = 80.32^{+32\%}_{-22\%} \text{ ab} \quad \sigma_{\text{full off-shell}}^{\text{NLO}} = 98.88^{+1\%}_{-6\%} \text{ ab} \quad (1)$$

Thus the integrated cross section is increased by +23% once NLO QCD corrections are applied. The size of the scale uncertainty is reduced to 6% maximally at NLO. On the left of fig. 1 the integrated cross section is computed for different re-scalings of the renormalization μ_R and factorization μ_F scales by a factor ξ , with respect to the central dynamic scale choice. We note that the variation in the integrated cross section is driven primarily by μ_R . Yet another systematic source of theory uncertainties are intrinsic PDF uncertainties. For the NLO cross section in eq. (1) the PDF uncertainty is about $\pm 1\%$. Another way of evaluating PDF uncertainties is to compare the predictions computed with different central PDF sets. This is




Figure 1: (Left) Scale dependance plot for the dynamic scale. (Right) Various PDF sets compared against each other and the scale uncertainty.




Figure 2: Differential fiducial cross sections for $t\bar{t}Z$ with various modelling approaches.

shown on the right of fig. 1, where we note that for most PDF sets the biggest difference to the default NNPDF3.1 PDF set as well as their corresponding intrinsic PDF uncertainties are around 3% or smaller. The difference to NNPDF3.1 is larger only for the ABMP16 PDF set, where it amounts to about 5%. In all cases, scale variation dominates the theoretical uncertainties.

Furthermore, we also investigate the modelling in the presence of the full off-shell effects by comparing to the full NWA_{full} and the NWA with LO decays $\text{NWA}_{\text{LOdec}}$. In the NWA, $\tau^+\tau^-$ can only originate from an on-shell Z -gauge boson, where the latter is emitted in the production stage. On the other hand, full off-shell predictions allow for off-mass-shell and $\gamma^* \rightarrow \tau^+\tau^-$ contributions. This leads to sizable full off-shell effects of about 11% at the integrated level. To investigate them further, we impose an additional window cut $|M_{\tau^+\tau^-} - m_Z| < X$, that aims to remove photon induced contributions. By varying X between 25 GeV and 10 GeV full off-shell effects are reduced below 3%.

The impact of this cut can be visualized even better at the differential level in fig. 2. For Z -boson related observables such as $p_{T\tau^+\tau^-}$, imposing this cut with $X = 10, 20$ GeV to the full off-shell predictions leads to better agreement with the NWA_{full} even in the high p_T regions, where full off-shell effects are sizable. On the other hand, for top-quark related observables, such as $p_{Tb_1b_2}$, the window cut does not remove the single-resonant contributions, visible at the end of the plotted p_T spectrum. Due to the photon induced contributions, full off-shell effects are also visible for angular observables.

3 Conclusion

In conclusion, NLO QCD corrections to $t\bar{t}Z$ in the 4ℓ decay channel are large (23%) and they reduce the dominant scale uncertainties to at most 6%. Full off-shell effects for this process are sizable (11%) due to the photon contribution and interference effects. We find that by imposing a window cut with $X = 10, 20$ GeV one can effectively remove the full off-shell effects related to the Z -gauge boson.

References

- [1] N. Greiner, K. Kong, J.-C. Park, S. C. Park and J.-C. Winter, *Model-Independent Production of a Top-Philic Resonance at the LHC*, *JHEP* **04** (2015) 029 [[1410.6099](#)].
- [2] U. Baur, A. Juste, L. H. Orr and D. Rainwater, *Probing electroweak top quark couplings at hadron colliders*, *Phys. Rev. D* **71** (2005) 054013 [[hep-ph/0412021](#)].
- [3] ATLAS collaboration, *Evidence for the associated production of the Higgs boson and a top quark pair with the ATLAS detector*, *Phys. Rev. D* **97** (2018) 072003 [[1712.08891](#)].
- [4] ATLAS collaboration, *Search for supersymmetry at $\sqrt{s} = 13$ TeV in final states with jets and two same-sign leptons or three leptons with the ATLAS detector*, *Eur. Phys. J. C* **76** (2016) 259 [[1602.09058](#)].
- [5] CMS collaboration, *Measurement of top quark pair production in association with a Z boson in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **03** (2020) 056 [[1907.11270](#)], copyright terms of the CC BY 4.0 license apply.
- [6] ATLAS collaboration, *Measurements of the inclusive and differential production cross sections of a top-quark–antiquark pair in association with a Z boson at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **81** (2021) 737 [[2103.12603](#)].
- [7] G. Bevilacqua, H. B. Hartanto, M. Kraus, J. Nasufi and M. Worek, *NLO QCD corrections to full off-shell production of $t\bar{t}Z$ including leptonic decays*, *JHEP* **08** (2022) 060 [[2203.15688](#)], copyright terms of the CC BY 4.0 license apply.