# $3n + 3^k$ : New Perspective on Collatz Conjecture

Naouel Boulkaboul\*

Independent researcher, Algiers, Algeria

October 2022

#### Abstract

Collatz conjecture is generalized to  $3n + 3^k$   $(k \in N)$ . Operating as usual, every sequence seems to reach  $3^k$  and end up in the loop  $3^k, 4.3^k, 2.3^k, 3^k$ . The usual 3n + 1 conjecture is recovered for k = 0. For k > 0, we noticed the existence of a sequence of period 3, namely,  $3^{k-1}, 2.3^k, 3^k$ , alongside the cycle  $4.3^k, 2.3^k, 3^k$  encountered in the 3n + 1(k = 0) sequence. A term formula of the  $3n + 3^k$  conjecture has been derived, and hence the *total stopping time*.

#### **1** Introduction

Credited to the mathematician Lothar Collatz, the Collatz conundrum (see for instance Ref. [1] and references therein), which was brought forward in the 1930s, is one of the simplest yet unsolved conjectures in mathematics. "This is a really dangerous problem. People become obsessed with it and it really is impossible," as stated by Jeffrey Lagarias. Such a conjecture asks whether repeating two simple arithmetic operations will eventually reach 1. That is to say, let  $N := \{0, 1, 2...\}$  denote the natural numbers, so that  $N + 1 = \{1, 2, 3...\}$  are the positive integers. Now, pick an arbitrary positive integer  $n \in N + 1$  and apply the following operation on it: If the number is even, divide it by two; while if the number is odd, triple it and add one. Let the process be denoted by f(n). That is

$$f(n) = \begin{cases} \frac{n}{2} & \text{if } n \equiv 0 \pmod{2} \\ 3n+1 & \text{if } n \equiv 1 \pmod{2} \end{cases}$$
(1)

The main task is to form a sequence by performing this operation repeatedly, and take the output at each step as the input at the next. Consequently, one may notice that such a process will eventually lead to the number 1, regardless of which positive integer is chosen initially.

If we set  $f^0(n) = n$  and  $f^l(n) = f(f^{l-1}(n))$  for  $l \in N$ . Then, the Collatz sequence for n reads

$$C(n) = \{f^l(n)\}_{l=0}^{\infty},\tag{2}$$

Strictly speaking, one may note that every Collatz sequence ends up in the loop 1, 4, 2, 1. To disprove Collatz conjecture, one has to show that there exists some starting number which yields a sequence that does not include 1. Such a sequence would either enter a repeating cycle that excludes 1, or increase infinitely. No such sequence has been found so far [2].

In the present paper, we show that Collatz conjecture 3n + 1 may possibly be a part of a more generalized case of the form  $3n + 3^k$  with  $k \in N$ .

# 2 $3n+3^k$ Conjecture

We provide the following operation, pick an arbitrary positive integer  $n \in N + 1$ , if the number is even, divide it by two; otherwise if the number is odd, triple it and add  $3^k$  with  $k \in N$ .

$$g_k(n) = \begin{cases} \frac{n}{2} & \text{if } n \equiv 0 \pmod{2} \\ 3n+3^k & \text{if } n \equiv 1 \pmod{2} \\ k \in N \end{cases}$$
(3)

<sup>\*</sup>Email: nwl\_qcd@yahoo.com

Repeating the two operations will eventually lead to  $3^k$   $(k \in N)$ . Hence, one may argue that the original Collatz conjecture 3n + 1 is nothing more than  $3n + 3^k$  conjecture with k = 0, i. e.  $f(n) = g_0(n)$ . Apparently, each positive integer, upon repetitive application of  $g_k(n)$ , will end in a repeating sequence of the form  $3^k, 4.3^k, 2.3^k, 3^k$ . In the 3n + 1 (or equivalently  $3n + 3^k$  with k = 0) sequence, the only cycle of period 3 is known to be the sequence  $4.3^k, 2.3^k, 3^k$  [3]. For k > 0 though, along with the aforementioned cycle, one may distinguish another sequence of period 3, that is  $3^{k-1}, 2.3^k, 3^k$ . Although such a generalized conjecture seems to be true, it has yet to be numerically verified for at least the largest set of numbers that has been checked in the case of 3n + 1 so far, i. e.  $n \leq 2^{100000} - 1$  [4], even though a general proof of it is still lacking. Furthermore, up to this point it is not clear whether there exists a certain value of k for or beyond which the conjecture cease to be hold.

## **3** Term formula for $3n + 3^k$ sequence

Let  $C_k^l(n)$  include the first l terms of the  $3n + 3^k$  sequence for n. We set m as the number of odd terms in  $C_k^l(n)$ ,  $d_i$  as the number of consecutive even terms immediately following the *i*th odd term and  $d_0$  as the number of even terms preceding the first odd term. It follows that the next term in the  $3n + 3^k$  sequence for n is <sup>1</sup>.

$$g_k^l(n) = \frac{3^m}{2^{l-m}}n + 3^k \epsilon \sum_{j=1}^m \frac{3^{m-j}}{2^{\sum_{i=j}^m d_i}}$$
(4)

where  $\epsilon = 0$  if m = 0 and  $\epsilon = 1$  if  $m \neq 0$ , note that  $l - m = d_0 + d_1 + ... + d_m$ . It is worth noting that, for the original 3n + 1 Collatz sequence, the above relation reduces to

$$g_0^l(n) = \frac{3^m}{2^{l-m}}n + \epsilon \sum_{j=1}^m \frac{3^{m-j}}{2^{\sum_{i=j}^m d_i}}$$
(5)

 $3n + 3^k$  sequences of integers ranging from 1 to 17 for k = 0, 1, 2, 3 and 4 are listed in Table. 1. In the k-sequences chosen, it is evident that, after consecutive application of  $g_k^l$ , the aforementioned integers eventually reach  $3^k$ .

### 4 Total stopping time

We call the *total stopping time* of n the smallest t such that  $n_t = 3^k$ , with  $n_t$  being the value of  $g_k$  applied to n recursively t times. It is worth mentioning that t is nothing but the number of integers in the  $3n + 3^k$  sequence just preceding the  $3^k$  term, i. e.  $n_t = g_k^t(n)$ . Thus, using the relation (4), the *total stopping time* can be written as

$$t = \log_2 \left[ \frac{2^m 3^m}{3^k (1 - \epsilon \sum_{j=1}^m \frac{3^{m-j}}{2^{\sum_{i=j}^m d_i}})} n \right]$$
(6)

If t does not exist we say that the *total stopping time* is infinite. It follows that, one has to prove that all positive integers yield a finite total stopping time in order to prove the  $3n + 3^k$  conjecture. In an endeavor to prove the 3n + 1 conjecture, mathematicians used to inspect what it is called the *stopping time*, namely the least positive l for which  $f^l(n) = g_0^l(n) < n$  [6–9]. If they can prove that all positive integers have a finite stopping time, they can prove by induction that the Collatz conjecture is true. It is nothing short of reasonable since for n > 1,  $f^{l}(n) = g_{0}^{l}(n) = 1$  cannot occur without the occurrence of some  $f^{l}(n) = g_{0}^{l}(n) < n$  [10]. Indeed, in the 1970s, it has been shown that almost all Collatz sequences eventually reach a number that is smaller than where it started [11]. In 2019, Terence Tao proved that for almost all numbers the Collatz sequence of n leads to a lower value, showing that the Collatz conjecture holds true for almost all numbers [12]. But the question to be asked is whether it is the right (or sufficient) way to prove the Collatz conjecture. As obvious as it seems, if the  $3n + 3^k$ conjecture turns out to be true for all  $n \in N+1$ , then the statement that, for each  $n \in N+1$  there exists  $l \in N+1$ such that  $q_k^l(n) < n$ , does not hold for  $n < 3^k$ . This can be easily seen in the 3n + 27 sequence for n = 5, 9, 13 (see for instance Table. 1). Alternatively, we must focus on showing that each  $n \in N + 1$  has a finite total stopping time and hence proving that the  $3n + 3^k$  sequence inclines toward  $3^k$  no matter what integer we start with. That is to say, one has to test the validity of the following statement: "for each  $n \in N+1$ , there exists  $t \in N+1$  such that  $g_k^t(n) = 3^k$ .", without making use of the stopping time.

<sup>&</sup>lt;sup>1</sup>Note that this formula is different from the one derived by L E. Garner [5]

Now, suppose that two integers  $n_1$  and  $n_2$  have the same total stopping time t within the sequence  $3n + 3^k$ , then from Eq. (6) we have

$$n_{2} = \frac{2^{m_{1}}3^{m_{1}}}{2^{m_{2}}3^{m_{2}}} \frac{1 - \epsilon \sum_{j=1}^{m_{2}} \frac{3^{m_{2}-j}}{2\sum_{i=j}^{m_{2}}d_{2,i}}}{1 - \epsilon \sum_{j=1}^{m_{1}} \frac{3^{m_{1}-j}}{2\sum_{i=j}^{m_{1}}d_{1,i}}} n_{1}$$

$$\tag{7}$$

where  $m_1(m_2)$  and  $d_{1,i}(d_{2,i})$  are, respectively, the number of odd terms and the number of consecutive even terms immediately following the *i*th odd term in  $C_k^t(n_1)$  ( $C_k^t(n_2)$ ), namely the set of *t* terms of the  $3n + 3^k$  sequence for  $n_1(n_2)$ .

It is obvious that for  $m_1 = m_2 = 0$ , we have  $n_2 = n_1$ . Hence, no two integers, for which the corresponding sequences contain even terms only, have the same *total stopping time*. Sequences that have the same *total stopping time* and the same cycle of period 3 have the same number of odd terms, a feature that can be figured out by checking Figs. 1–3 which depict the *total stopping time* and the number of odd terms in the set  $C_k^{t+1}(n)(k = 0, 1, 2)$  for different values of n. In the 3n + 9 sequence for example (upper plot in Fig. 3), there exist two cycles of period 3: 36, 18, 9 and 3, 18, 9, the integers 32, 33 and 35 have the same *total stopping time*, i.e. t = 10, but only 33 and 35 share the same number of odd terms m + 1 = 4, which is due to the fact that both of them possesses the same cycle of period 3, i.e. 36, 18, 9, whereas 32 has the cycle 3, 18, 9.

Remarkably, It is worth noting that sequences that possess the same number of odd terms in the set  $C_k^{t+1}(n)$  do not necessarily yield the same *total stopping time*, even if they share the same cycle of period 3, take the example of 1 and 2 in the 3n + 9 sequence, both of them have the same number of odd terms in the set  $C_9^{t+1}(n)$ , i.e. m + 1 = 3, and the same cycle 3, 18, 9 (check Table. 1) meanwhile they have different *total stopping time*, t = 5 and t = 6, respectively.

### References

- [1] Jeffrey C Lagarias. The 3x+ 1 problem: An annotated bibliography (1963–1999). The ultimate challenge: the 3x, 1:267–341, 2003.
- [2] David Barina. Convergence verification of the collatz problem. The Journal of Supercomputing, 77(3):2681– 2688, 2021.
- [3] Roger E Zarnowski. Generalized inverses and the total stopping times of collatz sequences. Linear and Multilinear Algebra, 49(2):115–130, 2001.
- [4] Wei Ren, Simin Li, Ruiyang Xiao, and Wei Bi. Collatz conjecture for 2<sup>^</sup>100000-1 is true-algorithms for verifying extremely large numbers. In 2018 IEEE SmartWorld, Ubiquitous Intelligence & Computing, Advanced & Trusted Computing, Scalable Computing & Communications, Cloud & Big Data Computing, Internet of People and Smart City Innovation (SmartWorld/SCALCOM/UIC/ATC/CBDCom/IOP/SCI), pages 411–416. IEEE, 2018.
- [5] Lynn E Garner. On the collatz 3n+1 algorithm. Proceedings of the American Mathematical Society, 82(1):19-22, 1981.
- [6] Ivan Korec. A density estimate for the 3x + 1 problem. Mathematica Slovaca, 44(1):85–89, 1994.
- [7] H Nelson Crooks Jr and Chigozie Nwoke. Collatz conjecture: Patterns within. arXiv preprint arXiv:2209.05995, 2022.
- [8] Michael A Idowu. A novel theoretical framework formulated for information discovery from number system and collatz conjecture data. *Proceedia Computer Science*, 61:105–111, 2015.
- [9] Rafael Ruggiero. The relationship between stopping time and number of odd terms in collatz sequences. arXiv preprint arXiv:1911.01229, 2019.
- [10] Jeffrey C Lagarias. The ultimate challenge: The 3x + 1 problem. American Mathematical Soc., 2010.
- [11] Riho Terras. A stopping time problem on the positive integers. Acta Arithmetica, 3(30):241–252, 1976.
- [12] T Tao. Almost all orbits of the collatz map attain almost bounded values. arXiv preprint arXiv:1909.03562, 2019.




Figure 1: Behavior of the *total stopping time* and the number of odd terms in the set  $C_1^{t+1}(n)$  of the 3n + 1 sequence for  $n \in [1, 100]$  (top),  $n \in [500, 600]$  (middle) and  $n \in [900, 1000]$  (bottom), the blue bars represent the *total stopping time* while the red bars depict the number of odd terms in the sequence.




Figure 2: Behavior of the *total stopping time* and the number of odd terms in the set  $C_3^{t+1}(n)$  of the 3n + 3 sequence for  $n \in [1, 100]$  (top),  $n \in [500, 600]$  (middle) and  $n \in [900, 1000]$  (bottom), the blue bars represent the *total stopping time* while the red bars depict the number of odd terms in the sequence.




Figure 3: Behavior of the *total stopping time* and the number of odd terms in the set  $C_9^{t+1}(n)$  of the 3n + 9 sequence for  $n \in [1, 100]$  (top),  $n \in [500, 600]$  (middle) and  $n \in [900, 1000]$  (bottom), the blue bars represent the *total stopping time* while the red bars depict the number of odd terms in the sequence.

								3n + 1									
n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
	4	1	10	2	16	3	22	4	28	5	34	6	40	7	46	8	52
	2	4	5	1	8	10	11	2	14	16	17	3	20	22	23	4	26
	1	2	16	4	4	5	34	1	7	8	52	10	10	11	70	2	13
		1	8	2	2	16	17	4	22	4	26	5	5	34	35	1	40
			4	1	1	8	52	2	11	2	13	16	16	17	106	4	20
			2		4	4	26	1	34	1	40	8	8	52	53	2	10
			1		2	2	13		17	4	20	4	4	26	160	1	5
			4		1	1	40		52 96	2	10	2	2	13	80		16
			2			4	20		26	1	5	1	1	40	40		8
			1			2	10		13		16	4	4	20 10	20		4
						1	5 16		$\frac{40}{20}$		8	$\frac{2}{1}$	$\frac{2}{1}$	10 E	10		2
							$\frac{16}{8}$		20 10		4	1	1	$\frac{5}{16}$	5 16		1
											2				16		$\frac{4}{2}$
							$\frac{4}{2}$		$\frac{5}{16}$		$\frac{1}{4}$			8	$\frac{8}{4}$		2 1
							$\frac{2}{1}$		8		$\frac{4}{2}$			$\frac{4}{2}$	$\frac{4}{2}$		T
							4		4		1			1	1		
							$\frac{4}{2}$		2		1			4	4		
							1		1					2	$\frac{1}{2}$		
							T		4					$\frac{2}{1}$	1		
									2					1	1		
									1								
								3n + 3									
$\overline{n}$	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
	6	1	12	2	18	3	24	4	30	5	36	6	42	$\overline{7}$	48	8	54
	3	6	6	1	9	12	12	2	15	18	18	3	21	24	24	4	27
	12	3	3	6	30	6	6	1	48	9	9	12	66	12	12	2	84
	6	12		3	15	3	3	6	24	30	30	6	33	6	6	1	42
	3	6		12	48		12	3	12	15	15	3	102	3	3	6	21
		3		6	24		6	12	6	48	48		51	12	12	3	66
				3	12		3	6	3	24	24		156	6	6	12	33
					6			3	12	12	12		78	3	3	6	102
					3				6	6	6		39			3	51
					12				3	3	3		120				156
					6					12	12		60				78
					3					6	6		30				39
										3	3		15				120
													48				60
													24				30
													12 C				15
													6				48
													3				24
													12 C				$\frac{12}{6}$
													6				6
																	0
													3				3
																	$\frac{3}{12}$
																	3

								3n + 9									
$\overline{n}$	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
	12	1	18	2	24	3	30	4	36	5	42	6	48	7	54	8	60
	6	12	9	1	12	18	15	2	18	24	21	3	24	30	27	4	30
	3	6	36	12	6	9	54	1	9	12	72	18	12	15	90	2	15
	18	3	18	6	3	36	27	12	36	6	36	9	6	54	45	1	54
	9	18	9	3	18	18	90	6	18	3	18	36	3	27	144	12	27
	36	9		18	9	9	45	3	9	18	9	18	18	90	72	6	90
	18	36		9	36		144	18		9	36	9	9	45	36	3	45
	9	18		36	18		72	9		36	18		36	144	18	18	144
		9		18	9		36	36		18	9		18	72	9	9	72
				9			18	18		9			9	36	36	36	36
							9	9						18	18	18	18
							36							9	9	9	9
							18							36			36
							9							18			18
														9			9
								3n + 27									
n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
	30	1	36	2	42	3	48	4	54	5	60	6	66	7	72	8	78
	15	30	18	1	21	36	24	2	27	42	30	3	33	48	36	4	39
	72	15	9	30	90	18	12	1	108	21	15	36	126	24	18	2	144
	36	72	54	15	45	9	6	30	54	90	72	18	63	12	9	1	72
	18	36	27	72	162	54	3	15	27	45	36	9	216	6	54	30	36
	9	18	108	36	81	27	36	72	108	162	18	54	108	3	27	15	18
	54	9	54	18	270	108	18	36	54	81	9	27	54	36	108	72	9
	27	54	27	9	135	54	9	18	27	270	54	108	27	18	54	36	54
	108	27		54	432	27	54	9		135	27	54	108	9	27	18	27
	54	108		27	216		27	54		432	108	27	54	54		9	108
	27	54		108	108		108	27		216	54		27	27		54	54
		27		54	54		54	108		108	27			108		27	27
				27	27		27	54		54				54		108	
					108			27		27				27		54	
					54					108						27	
					27					54							
										27							

								3n + 81									
$\overline{n}$	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
	84	1	90	2	96	3	102	4	108	5	114	6	120	7	126	8	132
	42	84	45	1	48	90	51	2	54	96	57	3	60	102	63	4	66
	21	42	216	84	24	45	234	1	27	48	252	90	30	51	270	2	33
	144	21	108	42	12	216	117	84	162	24	126	45	15	234	135	1	180
	72	144	54	21	6	108	432	42	81	12	63	216	126	117	486	84	90
	36	72	27	144	3	54	216	21	324	6	270	108	63	432	243	42	45
	18	36	162	72	90	27	108	144	162	3	135	54	270	216	810	21	216
	9	18	81	36	45	162	54	72	81	90	486	27	135	108	405	144	108
	108	9	324	18	216	81	27	36		45	243	162	486	54	1296	72	54
	54	108	162	9	108	324	162	18		216	810	81	243	27	648	36	27
	27	54	81	108	54	162	81	9		108	405	324	810	162	324	18	162
	162	27		54	27	81	324	108		54	1296	162	405	81	162	9	81
	81	162		27	162		162	54		27	648	81	1296	324	81	108	324
	324	81		162	81		81	27		162	324		648	162	324	54	162
	162	324		81	324			162		81	162		324	81	162	27	81
	81	162		324	162			81		324	81		162		81	162	
		81		162	81			324		162	324		81			81	
				81				162		81	162		324			324	
								81			81		162			162	
													81			81	

Table 1:  $3n + 3^k$  sequence with k = 0, 1, 2, 3 and 4 for n ranging from 1 to 17.