## INTRODUCTION TO TOPOLOGICAL SYMMETRY IN QFT

DANIEL S. FREED

ABSTRACT. This brief note publicizes the quantum framework for symmetry that is developed in our joint paper [FMT] with Greg Moore and Constantin Teleman. We include additional motivation and an application to a selection rule for line defects in 4-dimensional gauge theories.

This note<sup>1</sup> is a preview (appearing in arrears) of joint work with Greg Moore and Constantin Teleman [FMT], to which we refer for further exposition, further development, more examples, and for extensive references to the literature, since we omit citations here.

As a first introduction to our framework for topological symmetries, consider the simpler setting of isometries in Riemannian geometry. Let M be a Riemannian manifold. In differential geometry we define the notions of (i) a Lie group G, and (ii) the action of G on M by isometries. These are both fairly straightforward definitions. What is more difficult (Myers-Steenrod theorem) is to prove that the group of *all* isometries of M has a canonical Lie group structure. Our main definitions for topological symmetries in quantum field theory are analogous to (i) and (ii); we do not attempt to give structure to the collection of all topological symmetries of a quantum field theory.<sup>2</sup>

Here are the main points:

- (1) We define separately (i) abstract symmetry data, and (ii) a concrete realization of abstract symmetry data on a quantum field theory. The data (i) is a quiche<sup>3</sup> ( $\sigma, \rho$ ) in which  $\sigma$  is a topological field theory and  $\rho$  is a topological boundary theory. The data of the action (ii) of ( $\sigma, \rho$ ) on a quantum field theory F belongs to nontopological field theory if F is not topological.
- (2) Topological symmetry is treated as a *quantum* object: the pair  $(\sigma, \rho)$ . This goes beyond the usual coupling of a theory to a background gauge field, which treats symmetry in a *classical* context. We illustrate the power of this perspective by means of an example at the end of this note.

Date: July 5, 2023.

This material is based upon work supported by the National Science Foundation under Grant Number DMS-2005286 and by the Simons Foundation Award 888988 as part of the Simons Collaboration on Global Categorical Symmetries. It is also based upon work supported by the National Science Foundation under Grant No. 1440140, while the author was in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the fall semester of 2022.

<sup>&</sup>lt;sup>1</sup>which will appear in the conference proceedings of String Math 2022

<sup>&</sup>lt;sup>2</sup>There are well-developed mathematical foundations for *topological* field theory, but the geometric development of general quantum field theories is at a much more nascent phase. So while we treat topological field theories rigorously, we take a more heuristic approach to nontopological quantum field theories. Therefore, we do not even attempt a definition of topological symmetries in a nontopological quantum field theory.

<sup>&</sup>lt;sup>3</sup>This strange terminology for an object that encodes abstract quantum symmetries evokes a pair  $(\sigma, \rho)$  in which  $\rho$  is the crust and  $\sigma$  the filling; imagine the sandwich on the left of Figure 3 made open-faced by omitting the right "piece of bread"  $\tilde{F}$ . One can (and one did!) point out that a quiche is not an open-faced sandwich. (Google 'tartine')

(3) Our definitions are inspired by an analogy that we explicate below:

(1) the pair (A, R) of an algebra A and a right module  $R \iff a$  quiche  $(\sigma, \rho)$ element of  $A \iff b$  defect in  $(\sigma, \rho)$ 

Our framework applies to internal topological symmetries of quantum field theories. It includes homotopical symmetries: higher groups, 2-groups, etc.

To reinforce the power of separating abstract symmetry from its concrete action, we begin with an illustration from representation theory. Then we recount the key definitions from [FMT]. We end with an application of our framework to 4-dimensional gauge theory, or more to properly 4dimensional quantum field theories with a certain symmetry: the derivation of a selection rule for line defects.

I warmly thank Greg Moore and Constantin Teleman for their collaboration and their feedback on this note.

## Motivation: Computations in representations of $\mathfrak{sl}_2\mathbb{R}$

# 2-dimensional representation. Set

(2) 
$$h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \qquad e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \qquad f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

These matrices form a basis of the Lie algebra  $\mathfrak{sl}_2\mathbb{R}$ . Straightforward addition and multiplication of  $2 \times 2$  matrices verifies the identity

(3) 
$$\frac{1}{2}h^2 + ef + fe = \frac{1}{2}h^2 + h + 2fe$$

In fact, both sides equal the scalar matrix  $\begin{pmatrix} 3/2 & 0 \\ 0 & 3/2 \end{pmatrix}$ .

**3-dimensional representation.** Here the matrices that represent the elements (2) of  $\mathfrak{sl}_2\mathbb{R}$  are

(4) 
$$h' = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -2 \end{pmatrix} \qquad e' = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{pmatrix} \qquad f' = \begin{pmatrix} 0 & 0 & 0 \\ 2 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

One can compute—perhaps not quite as easily—that the identity (3) holds:

(5) 
$$\frac{1}{2}(h')^2 + e'f' + f'e' = \frac{1}{2}(h')^2 + h' + 2f'e'$$

Again both sides are scalar matrices, namely 4 times the identity matrix.

 $\mathbf{2}$ 

Infinite dimensional representations. The Lie group  $\mathrm{SL}_2\mathbb{R}$  acts on the projective line  $\mathbb{RP}^1$  as fractional linear transformations, effectively through its quotient  $\mathrm{PSL}_2\mathbb{R}$ . For each complex number  $\lambda \in \mathbb{C}$ , there is an induced action on  $\lambda$ -differentials  $\phi(x)(dx)^{\lambda}$ . The infinitesimal action of the Lie algebra  $\mathfrak{sl}_2\mathbb{R}$  is by the first-order differential operators

(6)  
$$\begin{aligned}
\tilde{h}: \phi \longmapsto -2x\phi' - 2\lambda\phi \\
\tilde{e}: \phi \longmapsto -\phi' \\
\tilde{f}: \phi \longmapsto x^2\phi' + 2\lambda x\phi
\end{aligned}$$

Simple calculus manipulations verify the identity (3), which here says that the ostensibly secondorder differential operator on each side of

(7) 
$$\frac{1}{2}\tilde{h}^2 + \tilde{e}\tilde{f} + \tilde{f}\tilde{e} = \frac{1}{2}\tilde{h}^2 + \tilde{h} + 2\tilde{f}\tilde{e}$$

is the scalar operator that multiplies  $\phi$  by  $4\lambda^2 - 2\lambda$ .

The universal enveloping algebra. Let  $\mathcal{U}(\mathfrak{sl}_2\mathbb{R})$  be the unital associative algebra generated by  $\mathfrak{sl}_2\mathbb{R}$  subject to the relation generated by equating commutators in  $\mathcal{U}(\mathfrak{sl}_2\mathbb{R})$  with the Lie bracket in  $\mathfrak{sl}_2\mathbb{R}$ :

(8) 
$$xy - yx = [x, y], \quad x, y \in \mathfrak{sl}_2\mathbb{R}.$$

Any linear representation of the Lie algebra  $\mathfrak{sl}_2\mathbb{R}$  extends to a left module for this *universal enveloping algebra*  $\mathcal{U}(\mathfrak{sl}_2\mathbb{R})$ . A special case of (8) is the relation

$$(9) ef = fe + h$$

in  $\mathcal{U}(\mathfrak{sl}_2\mathbb{R})$ , where  $e, f, h \in \mathfrak{sl}_2\mathbb{R}$  are the matrices (2). Substitute (9) into the left hand side of (3) to immediately obtain the right hand side of (3). That's it! This simple manipulation in the abstract algebra  $\mathcal{U}(\mathfrak{sl}_2\mathbb{R})$  proves that the relation (3) holds in *every* linear representation of  $\mathfrak{sl}_2\mathbb{R}$ , and so in particular immediately proves (5) and (7). Observe that the expression  $h^2/2 + ef + fe$  lies in the center of  $\mathcal{U}(\mathfrak{sl}_2\mathbb{R})$ , and therefore necessarily acts as a scalar in every irreducible representation, as we also saw in the three representations considered above. (However, this centrality is not the main point we are making here.)

The framework we introduce for symmetry in quantum field theory enables similar universal computations in an abstract quiche  $(\sigma, \rho)$ , and these computations imply relations in every quantum field theory on which  $(\sigma, \rho)$  acts. Computations are made with defects in the topological field theory  $(\sigma, \rho)$ .

# Motivation: Algebras

We first remark that the word 'symmetry' is usually used for a transformation of a mathematical object that preserves structure, so in particular is invertible. But if, say, a finite group G acts linearly on a complex vector space V, then one can take linear combinations of the linear operators corresponding to elements of G to endow V with the structure of a module over the group algebra  $\mathbb{C}[G]$ . In this way one can talk about an "algebra of symmetries", though the algebra contains many non-units (noninvertible elements). As an extreme case, we might now call the action of  $0 \in \mathbb{C}[G]$  a "symmetry" of V. Whether or not such "noninvertible symmetries" merit this nomenclature, it is these algebras of symmetries that motivate our definitions in field theory.

Abstract algebras of symmetries. Let A be an algebra, and for simplicity suppose that the ground field is  $\mathbb{C}$ . For our motivational exposition it suffices to assume that A and the modules that follow are finite dimensional. We do not use a topology on A. Let R be the right regular A-module, i.e., the vector space underlying the algebra A together with the right module structure  $R \times A \to R$ given by multiplication. Then an abstract algebra of symmetries is the pair (A, R).

**Concrete algebras of symmetries.** Let V be a complex vector space. The action of (A, R) on V is specified by the data of a pair  $(L, \theta)$ : L is a left A-module and  $\theta$  is an isomorphism of vector spaces

(10) 
$$\theta \colon R \otimes_A L \xrightarrow{\cong} V$$

Since R is the right regular module, the tensor product in the domain of (10) is  $A \otimes_A L$ , and there is a natural isomorphism of this mixing of A and L with the vector space that underlies the left module L. Passing from a module to its underlying vector space is such a natural process that we do not usually bother to articulate the isomorphism (10). But in the field theory context there is no operation analogous to 'underlying vector space'; Example 17 below suggests such an operation is not possible. Elements of  $A = \text{End}_A(R)$  act as linear operators on V via the isomorphism  $\theta$ .

A generalization. We can relax the condition that R be the *regular* module and consider (A, R) with any right module. Then this very general notion of abstract symmetry allows many possibilities. For example, if  $A = \mathbb{C}$  is the ground field as an algebra and R is any vector space, then an (A, R)action on a vector space V decomposes V as a tensor product  $R \otimes L$  for some vector space L. For general (A, R), the algebra  $\operatorname{End}_A(R)$  acts on  $R \otimes_A L$  and so too on V via the isomorphism  $\theta$ .

## Quiches and their action on quantum field theories

**Nomenclature:** from algebra to field theory. A Wick-rotated field theory is a linear representation of a bordism category: this is the elevator speech version of Graeme Segal's axiom system. In this sense a Wick-rotated field theory is analogous to a linear representation of a Lie group, or to a left module over an algebra.<sup>4</sup> We adopt terminology from these analogs. A field theory has a (spacetime) dimension and a collection of background fields that determine the domain bordism

<sup>&</sup>lt;sup>4</sup>Suppose A is an algebra and M is a right A-module. Then one can contemplate an algebra A' that acts on M on the left and commutes with the A-action. For example, one often considers the commutant  $A' = \text{End}_A(M)$ . In the analogy with field theory, A is the bordism category, the module M is the Wick-rotated field theory, and A' is the quiche, defined in the next few sections.




FIGURE 1. A domain wall, a right boundary theory, and a left boundary theory

category; specifying these is analogous to specifying which Lie group or algebra we are representing. But in this note we often leave these important specifications implicit. Suppose  $\sigma, \sigma_1, \sigma_2$  are field theories (of the same dimension on the same background fields). We use the following nomenclature for the domain walls and boundaries in Figure 1:

(11) domain wall 
$$\delta: \sigma_1 \to \sigma_2 \iff (\sigma_2, \sigma_1)$$
-bimodule  
(11) right boundary theory  $\rho: \sigma \to \mathbb{1} \iff$  right  $\sigma$ -module  
left boundary theory  $\widetilde{F}: \mathbb{1} \to \sigma \implies$  left  $\sigma$ -module

Here 1 is the tensor unit theory:  $1 \otimes F \cong F$  for all theories F. Furthermore, we use the notation

(12) 
$$\rho \otimes_{\sigma} \widetilde{F}$$

for the dimensional reduction F of  $\sigma$  along the interval pictured in Figure 2, a presentation of F that we sometimes call a *sandwich*: the bulk theory  $\sigma$  is sandwiched between the right boundary theory  $\rho$  and the left boundary theory  $\widetilde{F}$ .




FIGURE 2. The interval used in the dimensional reduction (12)

Abstract field theoretic symmetry data: the quiche. The nonnegative integer n in the following is the dimension of field theories on which the quiche  $(\sigma, \rho)$  acts.

**Definition 13.** Fix  $n \in \mathbb{Z}^{\geq 0}$ . Then an *n*-dimensional quicke is a pair  $(\sigma, \rho)$  in which  $\sigma$  is an (n+1)-dimensional topological field theory and  $\rho$  is a right topological  $\sigma$ -module.

It is the topological nature of the field theory  $\sigma$  and its right boundary theory  $\rho$  that make this definition appropriate for topological symmetries. One can relax the definition and only require that  $\sigma$  be a once-categorified *n*-dimensional topological field theory; in its role as symmetries of an *n*-dimensional field theory, it is not necessary to evaluate  $\sigma$  on arbitrary (n + 1)-manifolds.

Remark 14. In the motivational case of algebras, there is a distinguished right module: the regular module. Similarly, there are classes of topological field theories  $\sigma$  for which we can define the notion of a regular  $\sigma$ -module. The examples in this note are of this type.

The most basic quiche is associated to a finite group G and any nonnegative integer n. Let  $\sigma$  be free (n + 1)-dimensional gauge theory with gauge group G, and let  $\rho$  be the right boundary theory which sums over sections of the restriction of a principal G-bundle to the boundary. (This is sometimes called a Dirichlet boundary theory: the fluctuating principal G-bundle is trivialized on the boundary.) The detailed example at the end of this note is a homotopical variant in which G is the homotopical abelian group BA, the classifying space of a finite abelian group A.

Action of a quicke on a quantum field theory. The following definition is illustrated in Figure 3.




FIGURE 3.  $(\sigma, \rho)$ -module data on a field theory F

**Definition 15.** Let  $(\sigma, \rho)$  be an *n*-dimensional quiche, and let *F* be an *n*-dimensional field theory. A  $(\sigma, \rho)$ -module structure on *F* is a pair  $(\tilde{F}, \theta)$  in which  $\tilde{F}$  is a left  $\sigma$ -module and  $\theta$  is an isomorphism

(16) 
$$\theta \colon \rho \otimes_{\sigma} \widetilde{F} \xrightarrow{\cong} F$$

of absolute n-dimensional theories.

Neither F nor  $\widetilde{F}$  is assumed to be topological. (See footnote <sup>2</sup> for the mathematical status that we adopt in this note and in [FMT] for nontopological field theories.)

**Example 17** (center symmetry in gauge theory). Let H be a Lie group and suppose  $A \subset H$  is a finite subgroup of the center of H. Then A is abelian. Set  $\overline{H} = H/A$ . (As an example, take  $H = SU_2$  and  $A = \mu_2 = \{\pm 1\} \subset SU_2$  the center; then  $\overline{H} \cong SO_3$ .) Suppose F is an H-gauge theory in some dimension n, and assume that F carries the homotopical symmetry of the group BA, the classifying space of A. For example, the theory may have matter fields that are neutral under the central subgroup  $A \subset H$ , in which case it has BA-symmetry. A map into BA classifies a principal A-bundle, and there is a bilinear pairing {principal A-bundles} × {H-connections}  $\rightarrow$ {H-connections}: multiplication  $A \times H \rightarrow H$  is a homomorphism of Lie groups since A is central. We assume that the quantum field theory F has this action as a symmetry.) In the physics literature

### TOPOLOGICAL SYMMETRY IN QFT

this is termed a "1-form symmetry". It is usually expressed by introducing a new background field an A-gerbe<sup>5</sup>—and extending the theory F to a theory with this background/classical field. In [FMT] we advocate for the quantum quiche picture:  $\sigma$  is the quantum theory that sums over A-gerbes,  $\rho$  is the boundary theory that trivializes an A-gerbe, and  $\tilde{F}$  is an  $\overline{H}$ -gauge theory. Observe that no details about the quantum field theory F are used to describe the symmetry; we only have to know that it carries a BA-symmetry. In particular, our characterization of F as "an H-gauge theory" is not necessary, and furthermore this characterization only pertains to the description of the theory, not to the abstract theory: the possibility of isomorphisms between H-gauge theories and quantum field theories that lack a fluctuating H-connection obstructs the statement "The theory F is an H-gauge theory".

## Quotients

Augmentations of algebras. We begin as before with motivation from algebra.

**Definition 18.** Let A be an algebra over  $\mathbb{C}$ . An *augmentation* of A is an algebra homomorphism  $\epsilon_A \colon A \to \mathbb{C}$  from A to the ground field.

An augmentation gives the ground field  $\mathbb{C}$  the structure of an A-module  $\mathbb{C}_{\epsilon_A}$ , which we take to be a right module: set  $\lambda \cdot a = \lambda \epsilon_A(a)$  for  $\lambda \in \mathbb{C}_{\epsilon_A}$ ,  $a \in A$ . Let R be the right regular module, and let V be a vector space equipped with an (A, R)-action, including the structure isomorphism (10) that recovers the underlying vector space V. Define the vector space

(19) 
$$Q := \mathbb{C}_{\epsilon_A} \otimes_A L$$

Then Q is by definition the quotient of V by the action of (A, R) using the augmentation  $\epsilon_A$ .

**Example 20.** Let G be a finite group and  $A = \mathbb{C}[G]$  its complex group algebra. A character  $\chi$  of A determines the augmentation that sends  $g \mapsto \chi(g)$ ,  $g \in G$ . In particular, there is a canonical augmentation from the unit character:  $\chi(g) = 1$  for all  $g \in G$ . Let S be a set with a (left) G-action, and let  $V = \mathbb{C}\langle S \rangle$  be the free vector space on S. Then for the unit character, Q in (19) can be identified with  $\mathbb{C}\langle S/G \rangle$ , the free vector space on the quotient of S by the G-action. Any character  $\chi$  determines a complex line bundle  $\mathcal{L} \to S//G$  over the groupoid (stack) quotient S//G. In this case Q can be identified with the vector space of sections of  $\mathcal{L} \to S//G$ . Note that at each  $s \in S$  the stabilizer subgroup  $G_s \subset G$  acts on the fiber  $\mathcal{L}_s$  by a character. If that character is nontrivial, then all sections vanish at s.

Augmentations do not always exist and, as in Example 20, they may not be unique.

<sup>&</sup>lt;sup>5</sup>An A-gerbe is a geometric representative of a cohomology class in  $H^2(-; A)$ ; it arises here as the obstruction to lifting a principal  $\overline{H}$ -bundle to a principal H-bundle. For  $H = SU_2$  and  $A = \mu_2$  this obstruction is the second Stiefel-Whitney class.

Quotients in field theory. This often goes under the name 'gauging a symmetry'. Let  $(\sigma, \rho)$  be an n-dimensional quiche for some nonnegative integer n. Just as the notion of a *regular* module only makes sense in special contexts (Remark 14), so too does the notion of an *augmentation* of  $(\sigma, \rho)$ . In fully local topological field theory, one special context occurs when the codomain (n + 1)-category of the theory  $\sigma$  is the Morita category of algebras in some n-category. In that case we have the notions of both the regular module and augmentations: an augmentation is an algebra map to the tensor unit. We realize the augmentation  $\epsilon$  as a right  $\sigma$ -module. Then if  $\epsilon^L$  is its left adjoint—a particular dual left  $\sigma$ -module—the sandwich  $\rho \otimes_{\sigma} \epsilon^L$  is the trivial theory. Augmentations may not exist, whereas the regular module of an algebra always exists. For the gauge theory of a finite group G introduced after Remark 14, an augmentation is the right boundary theory  $\epsilon$  which does not have any additional fluctuating fields: there is no constraint on the bulk fluctuating G-bundle, hence the quantization sums over all G-bundles. As mentioned, this is the process of quotienting, or gauging, (by) the G-symmetry. The quotient theory along the augmentation  $\epsilon$  is by definition

(21) 
$$F/\sigma := \epsilon \otimes_{\sigma} \widetilde{F}$$

as illustrated in Figure 4.




FIGURE 4. On the left, the structure isomorphism of the  $(\sigma, \rho)$ -module structure on F; on the right, the definition of the quotient theory  $F/\sigma$ 

# Line defects in 4-dimensional theories with BA-symmetry

What follows is an extended example to illustrate the power of the quantum quiche framework. Our exposition is inspired by the papers [GMN, AST]. We first state the result, which is a selection rule for line<sup>6</sup> defects. Then we briefly introduce *finite homotopy theories*, an important general class of topological field theories. We conclude with a derivation of the selection rule.

Setup and the selection rule. As in Example 17, let H be a Lie group and suppose  $A \subset H$  is a finite subgroup of its center. Suppose F is a 4-dimensional field theory with BA-symmetry; we refer to it as an H-gauge theory. (But see the comment at the end of Example 17.) For every pair (A', q) consisting of a subgroup  $A' \subset A$  and a quadratic function  $q: A' \to \mathbb{C}^{\times}$  there is an associated<sup>7</sup>

<sup>&</sup>lt;sup>6</sup>Affine geometry: point, line, plane. Differential geometry: point, curve, surface. 'Curve defects' anyone?

<sup>&</sup>lt;sup>7</sup>Below we define the theory  $F_{A',q}$  from F using only the BA-symmetry of F; see (34) and Figure 5. The descriptions as gauge theories are not only superfluous but also do not have any intrinsic meaning. Furthermore, an important feature of our approach, as opposed to that in [GMN, AST], is that we give a *local* construction of  $F_{A',q}$  from F. Characterizing a quantum field theory by cataloging its line defects is not a local procedure. Here we compute the selection rule on line defects from the local description of the theory, and moreover our computation takes place entirely in the topological field theory that encodes the BA-symmetry.

H/A'-gauge theory  $F_{A',q}$ . We compute a selection rule for line defects in this theory. As we explain below, the category of line defects is a sum of categories indexed by

$$(22) (m,e) \in A \times A^{\vee},$$

where  $A^{\vee} = \text{Hom}(A, \mathbb{C}^{\times})$  is the group of characters—the Pontrjagin dual group—to A. Now the quadratic function  $q: A' \to \mathbb{C}^{\times}$  has an associated bicharacter

(23) 
$$b_q \colon A' \times A' \to \mathbb{C}^{\times},$$

and so by transposition a homomorphism

(24) 
$$\epsilon_q \colon A' \to (A')^{\vee}.$$

The selection rule for line defects in the theory  $F_{A',q}$  is:

(25) 
$$\begin{aligned} m \in A' \\ e \big|_{A'} &= \epsilon_q(m)^{-1} \end{aligned}$$

For later use: The quadratic function q gives rise to a *Pontrjagin square* cohomology operation

(26) 
$$\mathfrak{P}_q \colon H^2(X; A') \longrightarrow H^4(X; \mathbb{C}^{\times})$$

on any topological space X, and it too is a quadratic function:

(27) 
$$\mathfrak{P}_q(x_1 + x_2) = \mathfrak{P}_q(x_1) \cdot \mathfrak{P}_q(x_2) \cdot \beta_q(x_1, x_2), \qquad x_1, x_2 \in H^2(X; A'),$$

where  $\beta_q \colon H^2(X; A') \times H^2(X; A') \to H^4(X; \mathbb{C}^{\times})$  is the symmetric bihomomorphism defined by the cup product and the bicharacter (23) on coefficients.

Finite homotopy theories. These topological field theories are constructed by a finite, homotopical version of the Feynman path integral, hence are amenable to computations using techniques from algebraic topology. Some boundaries and defects in these theories have semiclassical descriptions, so are similarly susceptible to straightforward computation. The data that defines a finite homotopy theory is a pair  $(\mathcal{X}, \lambda)$  consisting of a  $\pi$ -finite space<sup>8</sup>  $\mathcal{X}$  and a cocycle  $\lambda$  on  $\mathcal{X}$ ; the theory exists in any dimension. In our example we use cocycles for singular cohomology with coefficients in  $\mathbb{C}^{\times}$ .

Set  $(\mathfrak{X}, \lambda) = (B^2A, 1)$ , where  $B^2A$  is an Eilenberg-MacLane space K(A, 2) and 1 is the identity cocycle that represents the identity element in  $H^4(B^2A; \mathbb{C}^{\times})$ . We construct a 5-dimensional

<sup>&</sup>lt;sup>8</sup>The  $\pi$ -finiteness condition— $\mathfrak{X}$  has finitely many path components; there exists  $Q \in \mathbb{Z}^{\geq 0}$  such that  $\pi_q(\mathfrak{X}, x) = 0$  for all q > Q,  $x \in \mathfrak{X}$ ; and  $\pi_q(\mathfrak{X}, x)$  is a finite group for all  $q \in \mathbb{Z}^{\geq 0}$ ,  $x \in \mathfrak{X}$ —is only needed to define the partition function in the top dimension. Without that assumption we obtain a once-categorified field theory. Our example  $\mathfrak{X} = B^2 A$  is  $\pi$ -finite, so defines a full topological field theory.

topological field theory  $\sigma$ . Let M be a closed manifold of dimension  $\leq 4$ ; we briefly describe the quantization  $\sigma(M)$  for dim M = 4, 3, 2. Consider the mapping space

(28) 
$$\mathfrak{X}^M = \operatorname{Map}(M, B^2 A).$$

This mapping space is the space of "fluctuating fields" of the theory on M; its quantizations (30)–(32) proceed via a homotopical form of integration. Eilenberg-MacLane spaces classify singular cohomology, from which it follows that

(29)  
$$\pi_0(\mathfrak{X}^M) = H^2(M; A)$$
$$\pi_1(\mathfrak{X}^M) = H^1(M; A)$$
$$\pi_2(\mathfrak{X}^M) = H^0(M; A)$$

Also, the mapping space  $\mathfrak{X}^M = \operatorname{Map}(M, B^2 A)$  is a product of Eilenberg-MacLane spaces: all k-invariants vanish. For dim M = 4 the vector space  $\sigma(M)$  is the space of locally constant complex functions on  $\mathfrak{X}^M$ :

(30) 
$$\sigma(M) = \operatorname{Fun}_{\operatorname{flat}}(\mathfrak{X}^M) = \operatorname{Fun}(\pi_0 \mathfrak{X}^M), \quad \dim M = 4.$$

Notice that this quantization only uses the set  $\pi_0 \mathfrak{X}^M$ . For dim M = 3 the linear category  $\sigma(M)$  is the category of flat complex vector bundles on  $\mathfrak{X}^M$ :

(31) 
$$\sigma(M) = \operatorname{Vect}_{\operatorname{flat}}(\mathfrak{X}^M) = \operatorname{Vect}(\pi_{\leq 1}\mathfrak{X}^M), \quad \dim M = 3.$$

Let  $\mathbb{V}$  denote the linear category of finite dimensional complex vector spaces and linear maps. Notice that the quantization (31) only uses the fundamental groupoid  $\pi_{\leq 1} \mathcal{X}^M$ ; the category  $\sigma(M)$  is equivalent to the category of complex vector bundles over that groupoid, i.e., the category of functors  $\pi_{\leq 1} \mathcal{X}^M \to \mathbb{V}$ . Now a  $\mathbb{V}$ -module is a complex linear category, and a flat  $\mathbb{V}$ -bundle over a topological space is a local system of  $\mathbb{V}$ -modules. For dim M = 2 the linear 2-category  $\sigma(M)$  is the 2-category of flat  $\mathbb{V}$ -bundles on  $\mathcal{X}^M$ :

(32) 
$$\sigma(M) = \operatorname{Cat}_{\operatorname{flat}}(\mathfrak{X}^M) = \operatorname{Cat}(\pi_{\leq 2}\mathfrak{X}^M), \quad \dim M = 2.$$

Let C denote the linear 2-category of suitable complex linear categories. Again the quantization (32) depends only on a truncation of the mapping space, here the fundamental 2-groupoid  $\pi_{\leq 2} \mathfrak{X}^M$ , and (32) is the 2-category of functors  $\pi_{\leq 2} \mathfrak{X}^M \to \mathbb{C}$ . Thus to each object of  $\pi_{\leq 2} \mathfrak{X}^M$  is assigned a linear category, to each 1-morphism is assigned a linear functor, and to each 2-morphism is assigned a natural transformation of functors. In particular, an elements of  $\pi_2(\mathfrak{X}^M, \phi)$  acts as an automorphism of the identity functor of the category attached to  $\phi \in \mathfrak{X}^M$ .

There are semiclassical data that specify some special topological boundary theories and topological defects in a finite homotopy theory. The semiclassical data of a topological boundary theory is a ( $\pi$ -finite) map  $f: \mathcal{Y} \to \mathcal{X}$  together with a suitable cochain on  $\mathcal{Y}$ ; for a topological defect the codomain is an iterated free loop space of  $\mathcal{X}$ . A regular boundary theory has  $\mathcal{Y} = *$  a point; an augmentation has  $f = \mathrm{id}_{\mathcal{X}}$ . We defer to [FMT, Appendix A] for details. **Proof of the selection rule.** Recall the setup in Example 17. Namely, A is a finite abelian group,  $\sigma$  is the 5-dimensional finite homotopy theory constructed from  $(B^2A, 1)$ , and  $\rho$  is the right regular boundary theory that quantizes a basepoint  $* \to B^2A$ . Furthermore, F is a 4-dimensional quantum field theory with BA-symmetry, and  $(\tilde{F}, \theta)$  is the data that expresses the BA-symmetry. A special case already mentioned is gauge theory: H is a Lie group that contains A as a subgroup of its center, then set  $\overline{H} = H/A$ , and suppose F as an H-gauge theory with BA-symmetry, in which case  $\widetilde{F}$  is an  $\overline{H}$ -gauge theory (obtained by coupling H-connections to A-gerbes).

As stated in the text following (32), semiclassical data for a right boundary theory is a  $\pi$ -finite map  $\mathcal{Y} \to B^2 A$  together with a cocycle  $\mu$  on  $\mathcal{Y}$  for a class in  $H^4(\mathcal{Y}; \mathbb{C}^{\times})$ . If  $A' \subset A$  is a subgroup, then  $\mathcal{Y} = B^2 A' \to B^2 A$  is a  $\pi$ -finite map. There is an isomorphism

(33) 
$$H^4(B^2A'; \mathbb{C}^{\times}) \cong \{ \text{quadratic functions } q \colon A' \longrightarrow \mathbb{C}^{\times} \}$$

by a classical computation of Eilenberg-MacLane: the quadratic function q maps to the Pontrjagin square  $\mathcal{P}_q(\iota) \in H^4(B^2A'; \mathbb{C}^{\times})$  of the tautological class  $\iota \in H^2(B^2A'; A')$ . Hence a pair (A', q)determines a right topological boundary theory  $R_{A',q}$  for  $\sigma$ . Furthermore, the sandwich

is the 4-dimensional quantum field theory which is a twisted quotient by BA', as illustrated in Figure 5. In the language of gauge theories,  $R_{A',q} \otimes_{\sigma} \widetilde{F}$  is a theory with gauge group H/A'; the quadratic form determines topological terms in the action of that gauge theory.




FIGURE 5. The q-twisted quotient of F by BA'

Fix A', q. We investigate line defects in the theory  $R_{A',q} \otimes_{\sigma} \widetilde{F}$ . The computation is local, but for ease of language assume given a 4-manifold M and a 1-dimensional submanifold  $C \subset M$  on which we "wrap" defects. In the sandwich picture, these are defects supported on the 2-dimensional submanifold  $[0,1] \times C \subset [0,1] \times M$ ; see Figure 6. The 1-category of line defects<sup>9</sup> supported on  $C \subset M$ is the inverse limit  $\lim_{\epsilon \to 0} \operatorname{Hom}(1, \sigma(L_{\epsilon}))$ , where in the sandwich picture  $L_{\epsilon}$  is the Cartesian product of [0,1] and the linking 2-sphere of size  $\epsilon$  of  $C \subset M$  at some  $p \in C$ . To compute  $\sigma$  on  $L_{\epsilon} \approx [0,1] \times S^2$ , cut along  $\{1/2\} \times S^2$  and so express  $\sigma(L_{\epsilon})$  as the 1-category

(35) 
$$\operatorname{Hom}(\mathcal{V}_{A',q},\mathcal{V}_{\widetilde{F}}(\epsilon))$$

<sup>&</sup>lt;sup>9</sup>The global statement on C is correct if the normal bundle to  $C \subset M$  is framed. In any case, it is the local statement at a point of C that is our focus here.

between objects in the 2-category  $\sigma(S^2)$ .<sup>10</sup> Observe that  $\mathcal{V}_{A',q}$  is topological—the top half-cylinder  $\Gamma$ in Figure 6 lies in the domain of the topological field theory  $(\sigma, R_{A',q})$ —so no  $\lim_{\epsilon \to 0}$  is required. The limit is required for  $\mathcal{V}_{\widetilde{F}}(\epsilon)$ , but the derivation of the selection rule (25) uses only the top halfcylinder  $\Gamma$ , so we do not encounter the limit in the sequel.




FIGURE 6. The 1-category of line defects

Remark 36. This slicing of  $L_{\epsilon}$  is the maneuver that sequesters the topological computation from the nontopological part of the theory. Such sequestration maneuvers on defects are what the sandwich presentation of the theory make possible.

Next, we compute the 2-category  $\sigma(S^2)$  along which we factor in (35). From (29) we have

(37)  

$$\pi_0 (\operatorname{Map}(S^2, B^2 A)) = H^2(S^2; A) \cong A$$

$$\pi_1 (\operatorname{Map}(S^2, B^2 A)) = H^1(S^2; A) = 0$$

$$\pi_2 (\operatorname{Map}(S^2, B^2 A)) = H^0(S^2; A) \cong A$$

Therefore, according to (32) the quantization  $\sigma(S^2)$  of  $\mathcal{M}$  is the 2-category of flat  $\mathbb{V}$ -bundles  $\mathcal{V} \to \pi_{\leq 2} \mathcal{M}$ . The computation (37) implies that for each  $m \in \pi_0 \mathcal{M}$  we have a linear category  $\mathcal{V}^{(m)}$  equipped with an action of  $\pi_2 \cong A$  by automorphisms of the identity functor, hence  $\mathcal{V}^{(m)}$  decomposes as

(38) 
$$\mathcal{V}^{(m)} = \bigoplus_{e} \mathcal{V}^{(m,e)} \cdot e, \qquad e \in H^0(S^2; A)^{\vee} \cong A^{\vee}.$$

In the gauge theory context, m and e are called discrete magnetic and electric fluxes.

As preparation for the final step of the computation, fix a cocycle  $\mu_q$  that represents the class in  $H^4(B^2A'; \mathbb{C}^{\times})$  that corresponds to q under the isomorphism (33). Suppose  $f_1, f_2: S^2 \to B^2A'$  are maps whose homotopy classes are  $m_1, m_2 \in A'$ , respectively. Then

(39) 
$$\langle (f_1 \times f_2)^*(\mu_q), [S^2 \times S^2] \rangle = \beta_q(m_1, m_2),$$





FIGURE 7. The map  $\mathfrak{G}' \longrightarrow \mathfrak{G}$  of 2-groupoids; see (42)

follows from an application of (27) with  $x_i = f_i^*(\iota)$ , where  $\iota \in H^2(B^2A'; A')$  is the tautological class.

Finally, we compute the value of  $(\sigma, R_{A',q})$  on the half-cylinder  $\Gamma$  in Figure 6; the result is an object  $\mathcal{V}_{A',q} = \bigoplus_{m,e} \mathcal{V}_{A',q}^{(m,e)} \cdot e$  in the 2-category  $\sigma(S^2)$ . Note that  $\Gamma$  is a morphism  $\emptyset^2 \to S^2$  in the bordism category; the  $R_{A',q}$ -coloring of the left boundary has the effect of "coning off" that boundary component. Intuitively,  $\Gamma: \emptyset^2 \to S^2$  is the bordism that computes the value of the boundary theory  $R_{A',q}$  on  $S^2$ . The semiclassical value of this bordism is the correspondence



in which the cocycle  $\tau^2(\mu_q)$  is the transgression of  $\mu_q$ ; its cohomology class lies in  $H^2(\operatorname{Map}(S^2, B^2A'); \mathbb{C}^{\times})$ . Represent  $\tau^2(\mu_q)$  as a flat  $\mathbb{V}$ -line bundle  $\mathcal{K} \to \operatorname{Map}(S^2, B^2A')$ . Formally, the quantization of  $\Gamma$  is

(41) 
$$\mathcal{V}_{A',q} = (p_1)_* \big( \mathcal{K} \otimes p_0^*(\underline{\mathbb{V}}) \big) = (p_1)_* (\mathcal{K}),$$

where  $\underline{\mathbb{V}} \to *$  is the trivial  $\mathbb{V}$ -line bundle. We compute  $(p_1)_*(\mathcal{K})$  using the map

(42) 
$$\mathcal{G}' := \pi_{<2} \operatorname{Map}(S^2, B^2 A') \xrightarrow{\jmath} \pi_{<2} \operatorname{Map}(S^2, B^2 A) =: \mathcal{G}$$

of fundamental 2-groupoids, as depicted in Figure 7. (Recall the homotopy groups (37) and the fact that all k-invariants vanish. Hence we have strict models in which (42) is an inclusion.) Our task is to compute the pushforward  $j_*(\mathcal{K}) \to \mathcal{G}$  of the V-line bundle  $\mathcal{K} \to \mathcal{G}'$ . First, it follows from (39) that for  $m \in A'$ , the action of  $\pi_2(\mathcal{G}', m) \cong A'$  on  $\mathcal{K}_m$  is via the character  $\epsilon_q(m)$ , where recall the definition (24) of  $\epsilon_q$ . Now since j induces an inclusion  $A' \hookrightarrow A$  on  $\pi_0$ , it follows that  $\mathcal{V}_{A',q}^{(m,e)} = 0$  if  $m \notin A'$ . This proves the first of the selection rules (25). For the second, observe that for  $m \in A'$ , the pushforward of  $\mathcal{K}_m$  yields a V-module over  $\mathcal{G}$  supported at  $m \in A' \subset A$  on which

<sup>&</sup>lt;sup>10</sup>To glue we must reverse the depicted arrow of time at the boundary of the top half-cylinder  $\Gamma$ . The finite semisimplicity of  $\sigma(S^2)$  makes this a straightforward operation.

the action of  $\pi_2(\mathfrak{G}, m) \cong A$  is by the induced representation  $\operatorname{Ind}_{A'}^A(\epsilon_q(m))$ . The latter is the sum of 1-dimensional representations: characters of A that restrict to  $\epsilon_q(m)$  on A' appear with multiplicity one. Therefore, the  $\mathbb{V}$ -bundle  $\mathcal{V}_{A',q} \to \mathfrak{G}$  has support on  $A' \subset A$  and the fiber at  $m \in A'$  is a sum of  $\mathbb{V}$ -lines indexed by characters of A whose restriction to A' is  $\epsilon_q(m)$ . It only remains to observe that the inverse in the second selection rule in (25) is due to the fact that  $\mathcal{V}_{A',q}$  is in the domain of (35), so appears dualized.

# References

- [AST] Ofer Aharony, Nathan Seiberg, and Yuji Tachikawa, Reading between the lines of four-dimensional gauge theories, JHEP 08 (2013), 115, arXiv:1305.0318 [hep-th]. 8
- [FMT] Daniel S. Freed, Gregory W. Moore, and Constantin Teleman, Topological symmetry in quantum field theory, arXiv:2209.07471.1, 2, 6, 7, 10
- [GMN] Davide Gaiotto, Gregory W. Moore, and Andrew Neitzke, Framed BPS States, Adv. Theor. Math. Phys. 17 (2013), no. 2, 241–397, arXiv:1006.0146 [hep-th]. 8

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS, AUSTIN, TX 78712 Email address: dafr@math.utexas.edu