APPROXIMATION OF NONLINEAR FILTERING FOR MULTISCALE MCKEAN-VLASOV STOCHASTIC DIFFERENTIAL EQUATIONS*

HUIJIE QIAO AND WANLIN WEI

School of Mathematics, Southeast University, Nanjing, Jiangsu 211189, P.R.China

ABSTRACT. The work concerns approximation of nonlinear filtering for multiscale McKean-Vlasov stochastic differential equations. First of all, by a Poisson equation we prove an average principle. Then we define nonlinear filtering of the origin multiscale equations and the average equation, and again by the Poisson equation show approximation between nonlinear filtering of the slow part for the origin multiscale equations and that of the average equation.

1. Introduction

Let $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{t\in[0,T]}, \mathbb{P})$ be a complete filtered probability space. B, W are n-dimensional and m-dimensional standard Brownian motions defined on $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{t\in[0,T]}, \mathbb{P})$, respectively. Moreover, B and W are mutually independent. Consider the following system on $\mathbb{R}^n \times \mathbb{R}^m$:

$$\begin{cases}
dX_{t}^{\varepsilon} = b_{1}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon})dt + \sigma_{1}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon})dB_{t}, \\
X_{0}^{\varepsilon} = x_{0}, \quad 0 \leqslant t \leqslant T, \\
dZ_{t}^{\varepsilon} = \frac{1}{\varepsilon}b_{2}(\mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon})dt + \frac{1}{\sqrt{\varepsilon}}\sigma_{2}(\mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon})dW_{t}, \\
Z_{0}^{\varepsilon} = z_{0}, \quad 0 \leqslant t \leqslant T,
\end{cases} \tag{1}$$

where these mappings $b_1: \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \to \mathbb{R}^n$, $\sigma_1: \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \to \mathbb{R}^{n \times n}$, $b_2: \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \to \mathbb{R}^m$, $\sigma_2: \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \to \mathbb{R}^{m \times m}$ are all Borel measurable, $\mathscr{L}_{X_{\varepsilon}}^{\mathbb{P}}$ denotes the distribution of $X_{\varepsilon}^{\varepsilon}$ under the probability measure \mathbb{P} , and $\mathcal{P}_2(\mathbb{R}^n)$ denotes the collection of probability measures on \mathbb{R}^n with finite second moments.

If $b_1, \sigma_1, b_2, \sigma_2$ don't depend on the distribution $\mathscr{L}_{X_t^\varepsilon}^{\mathbb{P}}$, and b_2, σ_2 also depend on the state X_t^ε , the system (1) is usually called multiscale stochastic differential equations (SDEs for short) or slow-fast systems. And $X_t^\varepsilon, Z_t^\varepsilon$ are called the slow and fast parts, respectively. These systems are often used to model phenomena from many fields, such as biology, chemistry and engineering. Moreover, there are many results about them. We only mention the averaging principle for them. In [9], Khasminskii first studied the averaging principle. Later, many authors generalized the result (See e.g. [15, 16, 17, 18, 19, 20] and the references therein). In particular, Pardoux and Veretennikov [18, 19, 20] systematically investigated approximation between X_t^ε and the solution of the average equation through Poisson equations.

AMS Subject Classification(2020): 60G35; 35K55.

Keywords: Multiscale McKean-Vlasov stochastic differential equations; the Poisson equation; the average principle; approximation of nonlinear filtering.

^{*}This work was partly supported by NSF of China (No.12071071).

Corresponding author: Huijie Qiao, hjqiaogean@seu.edu.cn.

If $b_1, \sigma_1, b_2, \sigma_2$ depend on distributions, the system (1) is generally called multiscale McKean-Vlasov SDEs. Since McKean-Vlasov SDEs are more complex than SDEs, there are only a few results about the averaging principle. Let us review them. When σ_1 doesn't depend on Z_t^{ε} and b_2, σ_2 also depend on the state X_t^{ε} , Röckner, Sun and Xie [25] showed that the slow part X_t^{ε} of the system (1) converges to an average system in the L^2 sense. Recently, Xu, Liu, Liu and Miao [29] extended the result in [25] to the case where b_2, σ_2 depend on $(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}, Z_t^{\varepsilon}, \mathscr{L}_{Z_t^{\varepsilon}}^{\mathbb{P}})$. Very recently, we [22] considered the same system to that in [29] and established the L^p $(p \geq 2)$ convergence. In the paper, we observe the system (1), that is, σ_1 does depend on Z_t^{ε} . The price to pay is that b_2, σ_2 don't depend on the state X_t^{ε} and the distribution $\mathscr{L}_{Z_t^{\varepsilon}}^{\mathbb{P}}$.

Next, we take an observation process Y_t^{ε} , i.e.

$$Y_t^{\varepsilon} = V_t + \int_0^t h(X_s^{\varepsilon}, \mathcal{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) \mathrm{d}s, \tag{2}$$

where V is a l-dimensional Brownian motion independent of B, W, and $h: \mathbb{R}^n \times \mathcal{P}(\mathbb{R}^n) \times \mathbb{R}^m \to \mathbb{R}^l$ is Borel measurable. Thus, the system (1) and the process (2) form a signal-observation system. Here we as well as study nonlinear filtering problems of this signal-observation system. That is, we estimate and predict $(X_{\cdot}^{\varepsilon}, \mathscr{L}_{X_{\cdot}^{\varepsilon}}^{\mathbb{P}}, Z_{\cdot}^{\varepsilon})$ based on the information of Y_{\cdot}^{ε} . In fact, these problems have appeared in many fields, such as stochastic control, financial modeling, speech and image processing, and Bayesian networks. Furthermore, if $b_1, \sigma_1, b_2, \sigma_2, h$ don't depend on the distribution $\mathscr{L}_{X_{\epsilon}^{\varepsilon}}^{\mathbb{P}}$, these problems have been widely studied (See [1, 5, 6, 8, 17, 21, 23] and references therein). If σ_1, h don't depend on $Z_{\epsilon}^{\varepsilon}$ and b_2, σ_2 also depend on the state $X_{\epsilon}^{\varepsilon}$ and the distribution $\mathscr{L}_{Z_{\epsilon}^{\varepsilon}}^{\mathbb{P}}$, for the signal-observation system (1)+(2) we [22] showed the convergence of the nonlinear filtering for $(X_{\epsilon}^{\varepsilon}, \mathscr{L}_{X_{\epsilon}^{\varepsilon}}^{\mathbb{P}})$ in the L^p $(p \geqslant 1)$ sense. In the paper, we require that σ_1, h depend on $Z_{\epsilon}^{\varepsilon}$ and obtain the weak convergence.

As a whole, our contribution are two-folded:

- By a Poisson equation we prove that the slow part of the original system weakly converges to the average system.
- By the same Poisson equation we establish that the nonlinear filtering of the slow part weakly converges to that of the average system.

It is worthwhile to mentioning our results. In [13], three authors studied the following multiscale McKean-Vlasov SDEs:

$$\begin{cases}
dX_{t}^{\varepsilon} = b_{1}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon}, \mathcal{L}_{Z_{t}^{\varepsilon}}^{\mathbb{P}})dt + \sigma_{1}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}})dB_{t}, \\
X_{0}^{\varepsilon} = x_{0}, \quad 0 \leqslant t \leqslant T, \\
dZ_{t}^{\varepsilon} = \frac{1}{\varepsilon}b_{2}(\mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon}, \mathcal{L}_{Z_{t}^{\varepsilon}}^{\mathbb{P}})dt + \frac{1}{\sqrt{\varepsilon}}\sigma_{2}(\mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon}, \mathcal{L}_{Z_{t}^{\varepsilon}}^{\mathbb{P}})dW_{t}, \\
Z_{0}^{\varepsilon} = z_{0}, \quad 0 \leqslant t \leqslant T.
\end{cases} \tag{3}$$

Then they defined

$$\bar{b}_1(x,\mu) = \int_{\mathbb{R}^m} b_1(x,\mu,z,\nu^{\mu}) \nu^{\mu}(\mathrm{d}z), \quad \bar{\sigma}_1 \bar{\sigma}_1^*(x,\mu) = \int_{\mathbb{R}^m} (\sigma_1 \sigma_1^*)(x,\mu,z,\nu^{\mu}) \nu^{\mu}(\mathrm{d}z), \quad (4)$$

where ν^{μ} is the unique invariant probability measure for the fast equation (See Section 3), constructed the average equation (11) and presented a similar average principle under strong conditions. It is obvious that the system (3) is more general than the system (1). However, comparing (4) with (10), we think that our definitions of $\bar{b}_1, \bar{\sigma}_1$ are more natural.

Moreover, our proof of the average principle is more succinct. Therefore, we don't simply seek generality.

The paper proceeds as follows. In Section 2, we introduce some related notations and the definition of L-derivatives. Then we state two main theorems with full assumptions in Section 3. The proofs of two main theorems are placed in Section 4 and 5, respectively. In Section 6, we prove (39).

The following convention will be used throughout the paper: C with or without indices will denote different positive constants whose values may change from one place to another.

2. Preliminary

In the section, we will recall some notations, and the definition of L-derivative for functions on $\mathcal{P}_2(\mathbb{R}^n)$ and list all the assumptions.

2.1. **Notations.** In this subsection, we introduce some notations used in the sequel.

 $|\cdot|$, $||\cdot||$ represent the norm of a vector and a matrix, respectively. $\langle\cdot,\cdot\rangle$ stands for the inner product of vectors on \mathbb{R}^n . A^* denotes the transpose of the matrix A.

Let $\mathcal{B}_b(\mathbb{R}^n)$ be the set of all bounded Borel measurable functions on \mathbb{R}^n . Let $C(\mathbb{R}^n)$ be the set of all functions which are continuous on \mathbb{R}^n . $C^k(\mathbb{R}^n)$ represents the collection of all functions which are continuous differentiable up to k-order. Define the Banach space:

$$C_{lip}(\mathbb{R}^n) := \left\{ \varphi \in C(\mathbb{R}^n) : \|\varphi\|_{C_{lip}(\mathbb{R}^n)} = \sup_{x \in \mathbb{R}^n} \frac{|\varphi(x)|}{1 + |x|^2} + \sup_{x \neq y} \frac{|\varphi(x) - \varphi(y)|}{|x - y|} < \infty \right\}.$$

Let $\mathscr{B}(\mathbb{R}^n)$ denote the Borel σ -field on \mathbb{R}^n . $\mathcal{P}(\mathbb{R}^n)$ consists of all probability measures on $\mathscr{B}(\mathbb{R}^n)$ carrying the usual topology of weak convergence. Let $\mathscr{B}(\mathbb{R}^n)$ satisfying:

$$\|\mu\|^2 := \int_{\mathbb{R}^n} |x|^2 \mu(dx) < \infty.$$

We put on $\mathcal{P}_2(\mathbb{R}^n)$ a topology induced by the following metric:

$$\rho(\mu,\nu) = \sup_{\|\varphi\|_{C_{lin}(\mathbb{R}^n)} \le 1} \left| \int_{\mathbb{R}^n} \varphi(x) \mu(dx) - \int_{\mathbb{R}^n} \varphi(x) \nu(dx) \right|.$$

Thus, $(\mathcal{P}_2(\mathbb{R}^n), \rho)$ is a Polish space.

2.2. L-derivative for functions on $\mathcal{P}_2(\mathbb{R}^n)$. In the subsection we recall the definition of L-derivative for functions on $\mathcal{P}_2(\mathbb{R}^n)$.

The definition was first introduced by Lions (c.f. [2]). Moreover, he used some abstract probability spaces to describe the L-derivatives. Here, for the convenience to understand the definition, we apply a straight way to state it (c.f. [24]). Let I be the identity map on \mathbb{R}^n . For $\mu \in \mathcal{P}_2(\mathbb{R}^n)$ and $\phi \in L^2(\mathbb{R}^n, \mathscr{B}(\mathbb{R}^n), \mu; \mathbb{R}^n), <\mu, \phi>:= \int_{\mathbb{R}^n} \phi(x)\mu(\mathrm{d}x)$. Moreover, by simple calculation, it holds that $\mu \circ (I + \phi)^{-1} \in \mathcal{P}_2(\mathbb{R}^n)$.

Definition 2.1. (i) A function $f : \mathcal{P}_2(\mathbb{R}^n) \to \mathbb{R}$ is called L-differentiable at $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, if the functional

$$L^{2}(\mathbb{R}^{n}, \mathscr{B}(\mathbb{R}^{n}), \mu; \mathbb{R}^{n}) \ni \phi \mapsto f(\mu \circ (I + \phi)^{-1})$$

is Fréchet differentiable at $\phi = 0$; that is, there exists a unique $\gamma \in L^2(\mathbb{R}^n, \mathscr{B}(\mathbb{R}^n), \mu; \mathbb{R}^n)$ such that

$$\lim_{\langle \mu, |\phi|^2 \rangle \to 0} \frac{f(\mu \circ (I + \phi)^{-1}) - f(\mu) - \langle \mu, \gamma \cdot \phi \rangle}{\sqrt{\langle \mu, |\phi|^2 \rangle}} = 0.$$

In the case, we denote $\partial_{\mu} f(\mu) = \gamma$ and call it the L-derivative of f at μ .

(ii) A function $f: \mathcal{P}_2(\mathbb{R}^n) \to \mathbb{R}$ is called L-differentiable on $\mathcal{P}_2(\mathbb{R}^n)$ if L-derivative $\partial_{\mu} f(\mu)$ exists for all $\mu \in \mathcal{P}_2(\mathbb{R}^n)$.

Definition 2.2. The function f is said to be in $C^k(\mathcal{P}_2(\mathbb{R}^n))$, if $\mu \mapsto f(\mu)$ is k-times continuous L-differentiable on $\mathcal{P}_2(\mathbb{R}^n)$.

Definition 2.3. The function $F: \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \mapsto \mathbb{R}$ is said to be in $C^{k,k}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$, if for $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, $F(\cdot,\mu)$ is in $C^k(\mathbb{R}^n)$, and for $x \in \mathbb{R}^n$, $F(x,\cdot)$ is in $C^k(\mathcal{P}_2(\mathbb{R}^n))$. If $F \in C^{k,k}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$, and itself and all its derivatives are bounded and jointly continuous in the corresponding variable family, we say $F \in C_b^{k,k}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$.

Definition 2.4. The function $\Psi: \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \mapsto \mathbb{R}$ is said to be in $C^{k,k,k}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m)$, if for $(x, z) \in \mathbb{R}^n \times \mathbb{R}^m$, $\Psi(x, \cdot, z) \in C^k(\mathcal{P}_2(\mathbb{R}^n))$ and for $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, $\Psi(\cdot, \mu, \cdot) \in C^k(\mathbb{R}^n \times \mathbb{R}^m)$. If $\Psi \in C^{k,k,k}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m)$, and itself and all its derivatives are bounded and jointly continuous in the corresponding variable family, we say $\Psi \in C^{k,k,k}_b(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m)$.

- 2.3. **Assumptions.** In the subsection, we list all the assumptions used in the sequel.
- $(\mathbf{H}_{b_1,\sigma_1}^1)$ There exists a constant $L_{b_1,\sigma_1} > 0$ such that for $x_i \in \mathbb{R}^n$, $\mu_i \in \mathcal{P}_2(\mathbb{R}^n)$, $z_i \in \mathbb{R}^m$, i = 1, 2,

$$|b_1(x_1, \mu_1, z_1) - b_1(x_2, \mu_2, z_2)|^2 + ||\sigma_1(x_1, \mu_1, z_1) - \sigma_1(x_2, \mu_2, z_2)||^2$$

$$\leqslant L_{b_1, \sigma_1} \Big(|x_1 - x_2|^2 + \rho^2(\mu_1, \mu_2) + |z_1 - z_2|^2 \Big).$$

 $(\mathbf{H}_{\sigma_1}^2)$ There exists a constant l > 0 such that for $x \in \mathbb{R}^n$, $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, $z \in \mathbb{R}^m$, $h \in \mathbb{R}^n$,

$$\langle \sigma_1(x,\mu,z)h,h\rangle \geqslant l|h|^2.$$

 $(\mathbf{H}_{b_1,\sigma_1}^3) \ \partial_x b_1(x,\mu,z), \partial_\mu b_1(x,\mu,z)(y), \partial_z b_1(x,\mu,z), \partial_{xx} b_1(x,\mu,z), \partial_{xz} b_1(x,\mu,z), \partial_{zz} b_1(x,\mu,z), \\ \partial_y \partial_\mu b_1(x,\mu,z)(y) \text{ exist for any } (x,\mu,y,z) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^n \times \mathbb{R}^m \text{ and are uniformly bounded. And there exists a constant } \gamma_1 \in (0,1] \text{ such that for } z_i \in \mathbb{R}^m, \\ i = 1,2,$

$$\sup_{x \in \mathbb{R}^{n}, \mu \in \mathcal{P}_{2}(\mathbb{R}^{n})} \|\partial_{x}b_{1}(x, \mu, z_{1}) - \partial_{x}b_{1}(x, \mu, z_{2})\| \leqslant C|z_{1} - z_{2}|^{\gamma_{1}},$$

$$\sup_{x \in \mathbb{R}^{n}, \mu \in \mathcal{P}_{2}(\mathbb{R}^{n})} \|\partial_{\mu}b_{1}(x, \mu, z_{1}) - \partial_{\mu}b_{1}(x, \mu, z_{2})\|_{L^{2}(\mu)} \leqslant C|z_{1} - z_{2}|^{\gamma_{1}},$$

$$\sup_{x \in \mathbb{R}^{n}, \mu \in \mathcal{P}_{2}(\mathbb{R}^{n})} \|\partial_{z}b_{1}(x, \mu, z_{1}) - \partial_{z}b_{1}(x, \mu, z_{2})\| \leqslant C|z_{1} - z_{2}|^{\gamma_{1}},$$

$$\sup_{x \in \mathbb{R}^{n}, \mu \in \mathcal{P}_{2}(\mathbb{R}^{n})} \|\partial_{xx}b_{1}(x, \mu, z_{1}) - \partial_{xx}b_{1}(x, \mu, z_{2})\| \leqslant C|z_{1} - z_{2}|^{\gamma_{1}},$$

$$\sup_{x \in \mathbb{R}^{n}, \mu \in \mathcal{P}_{2}(\mathbb{R}^{n})} \|\partial_{xz}b_{1}(x, \mu, z_{1}) - \partial_{xz}b_{1}(x, \mu, z_{2})\| \leqslant C|z_{1} - z_{2}|^{\gamma_{1}},$$

$$\sup_{x \in \mathbb{R}^{n}, \mu \in \mathcal{P}_{2}(\mathbb{R}^{n})} \|\partial_{zz}b_{1}(x, \mu, z_{1}) - \partial_{zz}b_{1}(x, \mu, z_{2})\| \leqslant C|z_{1} - z_{2}|^{\gamma_{1}},$$

$$\sup_{x \in \mathbb{R}^{n}, \mu \in \mathcal{P}_{2}(\mathbb{R}^{n})} \|\partial_{y}\partial_{\mu}b_{1}(x, \mu, z_{1}) - \partial_{y}\partial_{\mu}b_{1}(x, \mu, z_{2})\|_{L^{2}(\mu)} \leqslant C|z_{1} - z_{2}|^{\gamma_{1}}.$$

Moreover, the above conditions also hold for $\sigma_1 \sigma_1^*$.

 $(\mathbf{H}_{b_2,\sigma_2}^1)$ There exists a constant $L_{b_2,\sigma_2} > 0$ such that for $\mu_i \in \mathcal{P}_2(\mathbb{R}^n)$, $z_i \in \mathbb{R}^m$, i = 1, 2,

$$|b_2(\mu_1, z_1) - b_2(\mu_2, z_2)|^2 + ||\sigma_2(\mu_1, z_1) - \sigma_2(\mu_2, z_2)||^2 \leqslant L_{b_2, \sigma_2} \Big(\rho^2(\mu_1, \mu_2) + |z_1 - z_2|^2\Big).$$

 $(\mathbf{H}_{b_2,\sigma_2}^2)$ There exists a constant $\beta > 0$ with $\frac{\beta}{2} > L_{b_2,\sigma_2}$ such that for $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, $z_i \in \mathbb{R}^m$, i = 1, 2,

$$2\langle z_1 - z_2, b_2(\mu, z_1) - b_2(\mu, z_2) \rangle + \|\sigma_2(\mu, z_1) - \sigma_2(\mu, z_2)\|^2 \leqslant -\beta |z_1 - z_2|^2.$$

 $(\mathbf{H}_{b_2,\sigma_2}^{2'})$ For any $p \geqslant 1$, there exists a constant $\beta' > 0$ with $\frac{\beta'}{2} > (2p-1)L_{b_2,\sigma_2}$ such that for $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, $z_i \in \mathbb{R}^m$, i = 1, 2,

$$2\langle z_1 - z_2, b_2(\mu, z_1) - b_2(\mu, z_2) \rangle + (2p - 1) \|\sigma_2(\mu, z_1) - \sigma_2(\mu, z_2)\|^2 \leqslant -\beta' |z_1 - z_2|^2.$$

 $(\mathbf{H}_{b_2,\sigma_2}^3)$ $\partial_{\mu}b_2(\mu,z)(y), \partial_z b_2(\mu,z), \partial_{zz}b_2(\mu,z), \partial_y \partial_{\mu}b_2(\mu,z)(y)$ exist for any $(\mu,y,z) \in \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^n \times \mathbb{R}^m$ and are uniformly bounded. And there exists a constant $\gamma_2 \in (0,1]$ such that for $z_i \in \mathbb{R}^m$, i = 1, 2,

$$\begin{split} \sup_{\mu \in \mathcal{P}_{2}(\mathbb{R}^{n})} \| \partial_{\mu} b_{2}(\mu, z_{1}) - \partial_{\mu} b_{2}(\mu, z_{2}) \|_{L^{2}(\mu)} & \leq C |z_{1} - z_{2}|^{\gamma_{2}}, \\ \sup_{\mu \in \mathcal{P}_{2}(\mathbb{R}^{n})} \| \partial_{z} b_{2}(\mu, z_{1}) - \partial_{z} b_{2}(\mu, z_{2}) \| & \leq C |z_{1} - z_{2}|^{\gamma_{2}}, \\ \sup_{\mu \in \mathcal{P}_{2}(\mathbb{R}^{n})} \| \partial_{zz} b_{2}(\mu, z_{1}) - \partial_{zz} b_{2}(\mu, z_{2}) \| & \leq C |z_{1} - z_{2}|^{\gamma_{2}}, \\ \sup_{\mu \in \mathcal{P}_{2}(\mathbb{R}^{n})} \| \partial_{y} \partial_{\mu} b_{2}(\mu, z_{1}) - \partial_{y} \partial_{\mu} b_{2}(\mu, z_{2}) \|_{L^{2}(\mu)} & \leq C |z_{1} - z_{2}|^{\gamma_{2}}. \end{split}$$

Moreover, the above conditions also hold for σ_2 .

 (\mathbf{H}_h) h is bounded, and there is a constant $L_h > 0$ such that

$$|h(x_1, \mu_1, z_1) - h(x_2, \mu_2, z_2)|^2 \le L_h(|x_1 - x_2|^2 + \rho^2(\mu_1, \mu_2) + |z_1 - z_2|^2)$$

Remark 2.5. (i) $(\mathbf{H}_{b_1,\sigma_1}^1)$ yields that there exists a constant $\bar{L}_{b_1,\sigma_1} > 0$ such that for $x \in \mathbb{R}^n$, $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, $z \in \mathbb{R}^m$,

$$|b_1(x,\mu,z)|^2 + ||\sigma_1(x,\mu,z)||^2 \leqslant \bar{L}_{b_1,\sigma_1}(1+|x|^2+||\mu||^2+|z|^2).$$
 (5)

(ii) $(\mathbf{H}_{b_2,\sigma_2}^1)$ implies that there exists a constant $\bar{L}_{b_2,\sigma_2} > 0$ such that for $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, $z \in \mathbb{R}^m$,

$$|b_2(\mu, z)|^2 + ||\sigma_2(\mu, z)||^2 \leqslant \bar{L}_{b_2, \sigma_2}(1 + ||\mu||^2 + |z|^2).$$
(6)

(iii) $(\mathbf{H}_{b_2,\sigma_2}^1)$ and $(\mathbf{H}_{b_2,\sigma_2}^2)$ yield that there exists a constant C > 0 such that for $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, $z \in \mathbb{R}^m$

$$2\langle z, b_2(\mu, z) \rangle + \|\sigma_2(\mu, z)\|^2 \leqslant -\alpha |z|^2 + C(1 + \|\mu\|^2), \tag{7}$$

where $\alpha := \frac{\beta}{2} - L_{b_2,\sigma_2}$.

(iv) $(\mathbf{H}_{b_2,\sigma_2}^1)$ and $(\mathbf{H}_{b_2,\sigma_2}^{2'})$ imply that there exists a constant C > 0 such that for $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, $z \in \mathbb{R}^m$

$$2\langle z, b_2(\mu, z) \rangle + (2p - 1) \|\sigma_2(\mu, z)\|^2 \leqslant -\alpha' |z|^2 + C(1 + \|\mu\|^2), \tag{8}$$

where $\alpha' := \frac{\beta'}{2} - (2p-1)L_{b_2,\sigma_2}$.

Here we mention that $(\mathbf{H}_{b_2,\sigma_2}^{2'})$ is stronger than $(\mathbf{H}_{b_2,\sigma_2}^{2})$, and $(\mathbf{H}_{b_2,\sigma_2}^{2})$ and $(\mathbf{H}_{b_2,\sigma_2}^{2'})$ are used in Theorem 3.1 and 3.2, respectively.

3. Main results

In the section, we sate our main results in this paper.

3.1. The average principle. Under these assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$ $(\mathbf{H}_{b_2,\sigma_2}^1)$, by [28, Theorem 2.1] or [3, Theorem 3.1], we know that the slow-fast system (1) has a unique strong solution $(X_{\cdot}^{\varepsilon}, Z_{\cdot}^{\varepsilon})$.

Next, we take any $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, and fix it. Consider the following SDE:

$$\begin{cases} dZ_t^{\mu,z_0} = b_2(\mu, Z_t^{\mu,z_0}) dt + \sigma_2(\mu, Z_t^{\mu,z_0}) dW_t, \\ Z_0^{\mu,z_0} = z_0, \quad 0 \leqslant t \leqslant T. \end{cases}$$
(9)

Based on [7], it holds that under the assumption $(\mathbf{H}_{b_2,\sigma_2}^1)$, the above equation has a unique strong solution Z^{μ,z_0} . Besides, Z^{μ,z_0} is a Markov process. Let $p_t(\mu;z_0,A)$ denote the transition probability of Z^{μ,z_0} for $t \geq 0$ and $A \in \mathcal{B}(\mathbb{R}^m)$. The associated transition semigroup $\{P_t^{\mu}\}_{t\geq 0}$ is given by

$$(P_t^{\mu}\phi)(z_0) = \int_{\mathbb{R}^m} \phi(z') p_t(\mu; z_0, dz'), \quad \phi \in \mathcal{B}_b(\mathbb{R}^m).$$

Under the assumption $(\mathbf{H}_{b_2,\sigma_2}^2)$, by [28, Theorem 3.1], one could obtain that there exists a unique invariant probability measure ν^{μ} for Z_{\cdot}^{μ,z_0} .

In the following, set for $(x, \mu) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)$

$$\bar{b}_1(x,\mu) := \int_{\mathbb{R}^m} b_1(x,\mu,z) \nu^{\mu}(\mathrm{d}z), \quad \Sigma(x,\mu) := \int_{\mathbb{R}^m} (\sigma_1 \sigma_1^*)(x,\mu,z) \nu^{\mu}(\mathrm{d}z), \tag{10}$$

and then $\Sigma(x,\mu)$ is a positive definite symmetric matrix. Thus, we construct an average equation on $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{t\in[0,T]}, \mathbb{P})$ as follows:

$$\begin{cases}
d\bar{X}_t = \bar{b}_1(\bar{X}_t, \mathscr{L}_{\bar{X}_t}^{\mathbb{P}})dt + \bar{\sigma}_1(\bar{X}_t, \mathscr{L}_{\bar{X}_t}^{\mathbb{P}})dB_t, \\
\bar{X}_0 = x_0,
\end{cases}$$
(11)

where $\bar{\sigma}_1(x,\mu)$ is the square root of $\Sigma(x,\mu)$, i.e. $\bar{\sigma}_1(x,\mu)$ is a positive definite symmetric matrix satisfying $\Sigma(x,\mu) = (\bar{\sigma}_1\bar{\sigma}_1)(x,\mu)$.

Now, it is the position to state the first main result of this paper.

Theorem 3.1. Suppose that these assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$ $(\mathbf{H}_{\sigma_1}^2)$ $(\mathbf{H}_{b_1,\sigma_1}^3)$ $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^3)$ -hold. Then $\{X_t^{\varepsilon}, t \in [0,T]\}$ converges weakly to $\{\bar{X}_t, t \in [0,T]\}$ in $C([0,T],\mathbb{R}^n)$, where \bar{X}_t is a strong solution of Eq.(11).

The proof of the above theorem is placed in the near section.

3.2. Convergence for nonlinear filtering. Set

$$(\Lambda_t^{\varepsilon})^{-1} := \exp\left\{-\int_0^t h^i(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) dV_s^i - \frac{1}{2} \int_0^t |h(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon})|^2 ds\right\}.$$

Here and hereafter, we use the convention that repeated indices imply the summation. Under (\mathbf{H}_h) , we get that

$$\mathbb{E}\left(\exp\left\{\frac{1}{2}\int_0^T |h(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon})|^2 \mathrm{d}s\right\}\right) < \infty,$$

and furthermore $(\Lambda_t^{\varepsilon})^{-1}$ is an exponential martingale under the measure \mathbb{P} . Define a probability measure \mathbb{P}^{ε} via

$$\frac{\mathrm{d}\mathbb{P}^{\varepsilon}}{\mathrm{d}\mathbb{P}} = (\Lambda_T^{\varepsilon})^{-1}.$$

Then by the Girsanov theorem, it holds that Y^{ε} is a Brownian motion under the probability measure \mathbb{P}^{ε} .

Define the nonlinear filtering for $(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}, Z_t^{\varepsilon})$: for any $\Psi \in \mathcal{B}_b(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m)$

$$\rho_t^\varepsilon(\Psi) := \mathbb{E}^{\mathbb{P}^\varepsilon}[\Psi(X_t^\varepsilon, \mathscr{L}_{X_t^\varepsilon}^\mathbb{P}, Z_t^\varepsilon) \Lambda_t^\varepsilon | \mathscr{F}_t^{Y^\varepsilon}], \quad \pi_t^\varepsilon(\Psi) := \mathbb{E}[\Psi(X_t^\varepsilon, \mathscr{L}_{X_t^\varepsilon}^\mathbb{P}, Z_t^\varepsilon) | \mathscr{F}_t^{Y^\varepsilon}],$$

where $\mathscr{F}_t^{Y^{\varepsilon}} := \sigma\{Y_s^{\varepsilon}, 0 \leqslant s \leqslant t\} \vee \mathcal{N}$, and \mathcal{N} denotes the collection of all zero sets under the \mathbb{P} -measure. Here ρ_t^{ε} , π_t^{ε} are called the unnormalized and normalized filtering of $(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}, Z_t^{\varepsilon})$ with respect to $\mathscr{F}_t^{Y^{\varepsilon}}$, respectively. By the Kallianpur-Striebel formula, we get the following relationship between $\rho_t^{\varepsilon}(\Psi)$ and $\pi_t^{\varepsilon}(\Psi)$:

$$\pi_t^{\varepsilon}(\Psi) = \frac{\rho_t^{\varepsilon}(\Psi)}{\rho_t^{\varepsilon}(1)}.$$

Define the (x,μ) -marginal of ρ_t^{ε} as follows:

$$\rho_t^{\varepsilon,x,\mu}(F) = \int_{\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m} F(x,\mu) \rho_t^{\varepsilon}(\mathrm{d}x,\mathrm{d}\mu,\mathrm{d}z), \quad F \in \mathcal{B}_b(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)),$$

and then it holds that

$$\pi_t^{\varepsilon,x,\mu}(F) = \frac{\rho_t^{\varepsilon,x,\mu}(F)}{\rho_t^{\varepsilon,x,\mu}(1)},$$

where $\pi_t^{\varepsilon,x,\mu}$ is the (x,μ) -marginal of π_t^{ε} .

Next, set

$$\bar{\Lambda}_t := \exp \left\{ \int_0^t \bar{h}^i(\bar{X}_s, \mathscr{L}_{\bar{X}_s}^{\mathbb{P}}) dY_s^{\varepsilon, i} - \frac{1}{2} \int_0^t |\bar{h}(\bar{X}_s, \mathscr{L}_{\bar{X}_s}^{\mathbb{P}})|^2 ds \right\},
\bar{\rho}_t(F) := \mathbb{E}^{\mathbb{P}^{\varepsilon}} [F(\bar{X}_t, \mathscr{L}_{\bar{X}_t}^{\mathbb{P}}) \bar{\Lambda}_t | \mathscr{F}_t^{Y^{\varepsilon}}],
\bar{\pi}_t(F) := \frac{\bar{\rho}_t(F)}{\bar{\rho}_t(1)},$$

where $\bar{h}(x,\mu) := \int_{\mathbb{R}^m} h(x,\mu,z) \nu^{\mu}(\mathrm{d}z)$. Then about the relationship between $\pi_t^{\varepsilon,x,\mu}$ and $\bar{\pi}_t$, we have the following result which is the second main result for this paper.

Theorem 3.2. Assume that $(\mathbf{H}_{b_1,\sigma_1}^1)$ $(\mathbf{H}_{\sigma_1}^2)$ $(\mathbf{H}_{b_1,\sigma_1}^3)$ $(\mathbf{H}_{b_2,\sigma_2}^1)$ $(\mathbf{H}_{b_2,\sigma_2}^{2'})$ $(\mathbf{H}_{b_2,\sigma_2}^3)$ and (\mathbf{H}_h) hold. Then for any $t \in [0,T]$, $\pi_t^{\varepsilon,x,\mu}$ converges weakly to $\bar{\pi}_t$ as $\varepsilon \to 0$.

We will prove the above theorem in Section 5.

4. Proof of Theorem 3.1

In the section, we prove Theorem 3.1. First of all, we make some estimates.

4.1. Some estimates for $X^{\varepsilon}, Z^{\varepsilon}$.

Lemma 4.1. Under assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$ $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^2)$, there exists a constant C > 0such that

$$\sup_{t \in [0,T]} \mathbb{E}|X_t^{\varepsilon}|^2 \leqslant C(1+|x_0|^2+|z_0|^2), \quad \sup_{t \in [0,T]} \mathbb{E}|Z_t^{\varepsilon}|^2 \leqslant C(1+|x_0|^2+|z_0|^2).$$

Proof. For X_t^{ε} , based on the Hölder inequality, the isometric formula and (5), we can get

$$\mathbb{E}|X_{t}^{\varepsilon}|^{2} \leqslant 3|x_{0}|^{2} + 3\mathbb{E}\left|\int_{0}^{t}b_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon})\mathrm{d}s\right|^{2} + 3\mathbb{E}\left|\int_{0}^{t}\sigma_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon})\mathrm{d}B_{s}\right|^{2} \\
\leqslant 3|x_{0}|^{2} + 3t\int_{0}^{t}\mathbb{E}|b_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon})|^{2}\mathrm{d}s + 3\int_{0}^{t}\mathbb{E}\|\sigma_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon})\|^{2}\mathrm{d}s \\
\leqslant 3|x_{0}|^{2} + 3(T+1)\int_{0}^{t}\bar{L}_{b_{1},\sigma_{1}}(1+\mathbb{E}|X_{s}^{\varepsilon}|^{2} + \|\mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}\|^{2} + \mathbb{E}|Z_{s}^{\varepsilon}|^{2})\mathrm{d}s \\
\leqslant 3|x_{0}|^{2} + 3(T+1)\int_{0}^{t}\bar{L}_{b_{1},\sigma_{1}}(1+2\mathbb{E}|X_{s}^{\varepsilon}|^{2} + \mathbb{E}|Z_{s}^{\varepsilon}|^{2})\mathrm{d}s, \tag{12}$$

where $\|\mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}\|^2 = \mathbb{E}|X_s^{\varepsilon}|^2$.

For Z_t^{ε} , applying the Itô formula to $|Z_t^{\varepsilon}|^2$ and taking the expectation, one could obtain that

$$\mathbb{E}|Z_t^{\varepsilon}|^2 = |z_0|^2 + \frac{2}{\varepsilon} \mathbb{E} \int_0^t \langle Z_s^{\varepsilon}, b_2(\mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) \rangle \mathrm{d}s + \frac{1}{\varepsilon} \mathbb{E} \int_0^t \|\sigma_2(\mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon})\|^2 \mathrm{d}s,$$

and then

$$\frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E}|Z_{t}^{\varepsilon}|^{2} = \frac{2}{\varepsilon} \mathbb{E}\langle Z_{t}^{\varepsilon}, b_{2}(\mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon}) \rangle + \frac{1}{\varepsilon} \mathbb{E}\|\sigma_{2}(\mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon})\|^{2} \\
\leqslant \frac{1}{\varepsilon} \mathbb{E}\left(-\alpha|Z_{t}^{\varepsilon}|^{2} + C(1 + \|\mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}\|^{2})\right) \\
\leqslant \frac{-\alpha}{\varepsilon} \mathbb{E}|Z_{t}^{\varepsilon}|^{2} + \frac{C}{\varepsilon}\left(1 + \mathbb{E}|X_{t}^{\varepsilon}|^{2}\right),$$

where the above inequality is based on (7) and $\|\mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}\|^2 = \mathbb{E}|X_t^{\varepsilon}|^2$. By the comparison theorem, we have that

$$\mathbb{E}|Z_{t}^{\varepsilon}|^{2} \leqslant |z_{0}|^{2}e^{-\frac{\alpha}{\varepsilon}t} + \frac{C}{\varepsilon} \int_{0}^{t} e^{-\frac{\alpha}{\varepsilon}(t-s)} \left(1 + \mathbb{E}|X_{s}^{\varepsilon}|^{2}\right) ds$$

$$\leqslant |z_{0}|^{2} + C\left(1 + \sup_{s \in [0,t]} \mathbb{E}|X_{s}^{\varepsilon}|^{2}\right). \tag{13}$$

Finally, inserting (13) in (12), by the Gronwall inequality one can get that

$$\sup_{t \in [0,T]} \mathbb{E}|X_t^{\varepsilon}|^2 \leqslant C(1+|x_0|^2+|z_0|^2),$$

which is just the required first result. Moreover, this together with (13) implies the second estimate.

Lemma 4.2. Suppose that the assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$ $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^2)$ hold. Then it holds that for any $\delta > 0$ and any $\{\mathscr{F}_t\}$ -stopping time $\tau \leqslant T$,

$$\mathbb{E}|X_{\tau+\delta}^{\varepsilon} - X_{\tau}^{\varepsilon}|^{2} \leqslant C(1 + |x_{0}|^{2} + |z_{0}|^{2})(\delta^{2} + \delta).$$

Proof. From the Hölder inequality, the isometric formula and Lemma 4.1, it follows that

$$\begin{split} \mathbb{E}|X_{\tau+\delta}^{\varepsilon} - X_{\tau}^{\varepsilon}|^{2} &\leqslant 2\mathbb{E}\Big|\int_{\tau}^{\tau+\delta}b_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon})\mathrm{d}s\Big|^{2} + 2\mathbb{E}\Big|\int_{\tau}^{\tau+\delta}\sigma_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon})\mathrm{d}B_{s}\Big|^{2} \\ &\leqslant 2\delta\mathbb{E}\int_{\tau}^{\tau+\delta}|b_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon})|^{2}\mathrm{d}s + 2\mathbb{E}\int_{\tau}^{\tau+\delta}\|\sigma_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon})\|^{2}\mathrm{d}s \\ &\leqslant (2\delta+2)C\mathbb{E}\int_{\tau}^{\tau+\delta}(1+|X_{s}^{\varepsilon}|^{2}+\|\mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}\|^{2}+|Z_{s}^{\varepsilon}|^{2})\mathrm{d}s \\ &= (2\delta+2)C\mathbb{E}\int_{\tau}^{\tau+\delta}(1+|X_{s}^{\varepsilon}|^{2}+\mathbb{E}|X_{s}^{\varepsilon}\|^{2}+|Z_{s}^{\varepsilon}|^{2})\mathrm{d}s \\ &\leqslant (2\delta+2)C\mathbb{E}\int_{\tau}^{\tau+\delta}(1+|X_{s}^{\varepsilon}|^{2}+C(1+|x_{0}|^{2}+|z_{0}|^{2})+|Z_{s}^{\varepsilon}|^{2})\mathrm{d}s \\ &= (2\delta+2)C\mathbb{E}\int_{0}^{\delta}(1+|X_{\tau+u}^{\varepsilon}|^{2}+C(1+|x_{0}|^{2}+|z_{0}|^{2})+|Z_{\tau+u}^{\varepsilon}|^{2})\mathrm{d}u \\ &\leqslant C(1+|x_{0}|^{2}+|z_{0}|^{2})(\delta^{2}+\delta), \end{split}$$

which completes the proof.

Now, combining Lemma 4.1 and 4.2 with [10, Theorem 2.7], we draw the following conclusion.

Lemma 4.3. Under assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^2)$, $\{X_t^{\varepsilon}, t \in [0,T]\}$ is relatively weakly compact in $C([0,T],\mathbb{R}^n)$.

Here, we mention that the slow-fast system (1) is a Markov process and its infinitesimal generator is as follows: for $\Psi \in C^{2,2,2}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m)$,

$$(\mathcal{L}^{X^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}},Z^{\varepsilon}}\Psi)(x,\mu,z):=(\mathcal{L}^{X^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}}\Psi)(x,\mu,z)+(\mathcal{L}^{Z^{\varepsilon}}\Psi)(x,\mu,z),$$

where

$$(\mathcal{L}^{X^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}}\Psi)(x,\mu,z): = \partial_{x_{i}}\Psi(x,\mu,z)b_{1}^{i}(x,\mu,z) + \frac{1}{2}\partial_{x_{i}x_{j}}\Psi(x,\mu,z)(\sigma_{1}\sigma_{1}^{*})^{ij}(x,\mu,z)$$

$$+ \int_{\mathbb{R}^{n}}(\partial_{\mu}\Psi)_{i}(x,\mu,z)(y)b_{1}^{i}(y,\mu,z)\mu(\mathrm{d}y)$$

$$+ \frac{1}{2}\int_{\mathbb{R}^{n}}\partial_{y_{i}}(\partial_{\mu}\Psi)_{j}(x,\mu,z)(y)(\sigma_{1}\sigma_{1}^{*})^{ij}(y,\mu,z)\mu(\mathrm{d}y),$$

$$1 \qquad 1$$

$$(\mathcal{L}^{Z^{\varepsilon}}\Psi)(x,\mu,z): = \frac{1}{\varepsilon}\partial_{z_{i}}\Psi(x,\mu,z)b_{2}^{i}(\mu,z) + \frac{1}{2\varepsilon}\partial_{z_{i}z_{k}}\Psi(x,\mu,z)(\sigma_{2}\sigma_{2}^{*})^{ik}(\mu,z).$$

4.2. Some estimates for the frozen equation (9).

Lemma 4.4. Assume that $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^2)$ hold. Then we have for $\mu, \mu_1, \mu_2 \in \mathcal{P}_2(\mathbb{R}^n)$,

$$\mathbb{E}|Z_t^{\mu,z_0}|^2 \leqslant C(1+\|\mu\|^2+|z_0|^2), \quad t \geqslant 0,$$

$$\mathbb{E}|Z_t^{\mu_1,z_0}-Z_t^{\mu_2,z_0}|^2 \leqslant L_{b_2,\sigma_2}Te^{(L_{b_2,\sigma_2}+1)t}\rho^2(\mu_1,\mu_2), \quad t \geqslant 0.$$

Proof. First of all, by the same deduction to that for Z^{ε} in Lemma 4.1, one can get the first estimate.

For the second estimate, take T > 0 and note that for any $t \in [0, T]$

$$Z_t^{\mu_1,z_0} - Z_t^{\mu_2,z_0} = \int_0^t \left(b_2(\mu_1, Z_s^{\mu_1,z_0}) - b_2(\mu_2, Z_s^{\mu_2,z_0}) \right) ds$$
$$+ \int_0^t \left(\sigma_2(\mu_1, Z_s^{\mu_1,z_0}) - \sigma_2(\mu_2, Z_s^{\mu_2,z_0}) \right) dW_s.$$

Thus, applying the Itô formula to $|Z_t^{\mu_1,z_0} - Z_t^{\mu_2,z_0}|^2$ and taking expectation on two sides, we obtain that

$$\mathbb{E}|Z_{t}^{\mu_{1},z_{0}} - Z_{t}^{\mu_{2},z_{0}}|^{2}
= 2\mathbb{E}\int_{0}^{t} \langle Z_{s}^{\mu_{1},z_{0}} - Z_{s}^{\mu_{2},z_{0}}, b_{2}(\mu_{1}, Z_{s}^{\mu_{1},z_{0}}) - b_{2}(\mu_{2}, Z_{s}^{\mu_{2},z_{0}})\rangle ds
+ \mathbb{E}\int_{0}^{t} \|\sigma_{2}(\mu_{1}, Z_{s}^{\mu_{1},z_{0}}) - \sigma_{2}(\mu_{2}, Z_{s}^{\mu_{2},z_{0}})\|^{2} ds
\leq \mathbb{E}\int_{0}^{t} (|Z_{s}^{\mu_{1},z_{0}} - Z_{s}^{\mu_{2},z_{0}}|^{2} + |b_{2}(\mu_{1}, Z_{s}^{\mu_{1},z_{0}}) - b_{2}(\mu_{2}, Z_{s}^{\mu_{2},z_{0}})|^{2}) ds
+ \mathbb{E}\int_{0}^{t} \|\sigma_{2}(\mu_{1}, Z_{s}^{\mu_{1},z_{0}}) - \sigma_{2}(\mu_{2}, Z_{s}^{\mu_{2},z_{0}})\|^{2} ds
\leq (L_{b_{2},\sigma_{2}} + 1) \int_{0}^{t} \mathbb{E}|Z_{s}^{\mu_{1},z_{0}} - Z_{s}^{\mu_{2},z_{0}}|^{2} ds + L_{b_{2},\sigma_{2}} T \rho^{2}(\mu_{1}, \mu_{2}).$$

By the Gronwall inequality, it holds that

$$\mathbb{E}|Z_t^{\mu_1,z_0} - Z_t^{\mu_2,z_0}|^2 \leqslant L_{b_2,\sigma_2} T e^{(L_{b_2,\sigma_2}+1)t} \rho^2(\mu_1,\mu_2).$$

For $t \in [T, 2T], [2T, 3T], \dots$, the similar deduction to that for $t \in [0, T]$ implies the required result.

Now, it is necessary to point out that the infinitesimal generator of Z^{μ,z_0} or the transition semigroup $\{P^{\mu}_t\}_{t\geqslant 0}$ is the operator $\varepsilon \mathcal{L}^{Z^{\varepsilon}}$.

4.3. Some results for the average equation (11).

Lemma 4.5. Suppose that $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{\sigma_1}^2)$ and $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^2)$ hold. Then there exists a constant C > 0 such that

$$\|\bar{\sigma}_1(x_1,\mu_1) - \bar{\sigma}_1(x_2,\mu_2)\| \le C(|x_1 - x_2| + \rho(\mu_1,\mu_2)).$$

Moreover, Eq.(11) has a unique strong solution \bar{X} satisfying for any $q \geqslant 1$

$$\sup_{t \in [0,T]} \mathbb{E}|\bar{X}_t|^{2q} \leqslant C(1+|x_0|^{2q}).$$

Proof. First of all, note that

$$\|\bar{\sigma}_{1}(x_{1}, \mu_{1}) - \bar{\sigma}_{1}(x_{2}, \mu_{2})\| = \|(\Sigma(x_{1}, \mu_{1}) - \Sigma(x_{2}, \mu_{2}))(\bar{\sigma}_{1}(x_{1}, \mu_{1}) + \bar{\sigma}_{1}(x_{2}, \mu_{2}))^{-1}\|$$

$$\leq \|\Sigma(x_{1}, \mu_{1}) - \Sigma(x_{2}, \mu_{2})\|\|(\bar{\sigma}_{1}(x_{1}, \mu_{1}) + \bar{\sigma}_{1}(x_{2}, \mu_{2}))^{-1}\|$$

$$\leq C\|\Sigma(x_{1}, \mu_{1}) - \Sigma(x_{2}, \mu_{2})\|,$$
(14)

where the last inequality is based on $(\mathbf{H}_{\sigma_1}^2)$. Therefore, we only need to estimate $\|\Sigma(x_1, \mu_1) - \Sigma(x_2, \mu_2)\|$. From the definition of $\Sigma(x, \mu)$, the Hölder inequality and Lemma 4.4, it follows

that

$$\begin{split} &\|\Sigma(x_1,\mu_1) - \Sigma(x_2,\mu_2)\| \\ &= \left\| \int_{\mathbb{R}^m} (\sigma_1 \sigma_1^*)(x_1,\mu_1,z) \nu^{\mu_1}(\mathrm{d}z) - \int_{\mathbb{R}^m} (\sigma_1 \sigma_1^*)(x_2,\mu_2,z) \nu^{\mu_2}(\mathrm{d}z) \right\| \\ &= \left\| \lim_{S \to \infty} \frac{1}{S} \int_0^S \mathbb{E}(\sigma_1 \sigma_1^*)(x_1,\mu_1,Z_t^{\mu_1,z_0}) \mathrm{d}t - \lim_{S \to \infty} \frac{1}{S} \int_0^S \mathbb{E}(\sigma_1 \sigma_1^*)(x_2,\mu_2,Z_t^{\mu_2,z_0}) \mathrm{d}t \right\| \\ &\leqslant \lim_{S \to \infty} \frac{1}{S} \int_0^S \mathbb{E} \|(\sigma_1 \sigma_1^*)(x_1,\mu_1,Z_t^{\mu_1,z_0}) - (\sigma_1 \sigma_1^*)(x_2,\mu_2,Z_t^{\mu_2,z_0}) \| \mathrm{d}t \\ &\leqslant \lim_{S \to \infty} \frac{1}{S} \int_0^S \mathbb{E} \|\sigma_1(x_1,\mu_1,Z_t^{\mu_1,z_0}) \sigma_1^*(x_1,\mu_1,Z_t^{\mu_1,z_0}) \\ &- \sigma_1(x_1,\mu_1,Z_t^{\mu_1,z_0}) \sigma_1^*(x_2,\mu_2,Z_t^{\mu_2,z_0}) \| \mathrm{d}t \\ &+ \lim_{S \to \infty} \frac{1}{S} \int_0^S \mathbb{E} \|\sigma_1(x_1,\mu_1,Z_t^{\mu_1,z_0}) \sigma_1^*(x_2,\mu_2,Z_t^{\mu_2,z_0}) \| \mathrm{d}t \\ &\leqslant \lim_{S \to \infty} \frac{1}{S} \int_0^S \left(\mathbb{E} \|\sigma_1(x_1,\mu_1,Z_t^{\mu_1,z_0}) - \sigma_1^*(x_2,\mu_2,Z_t^{\mu_2,z_0}) \|^2 \right)^{1/2} \right. \\ &\leqslant \lim_{S \to \infty} \frac{1}{S} \int_0^S \left(\mathbb{E} \|\sigma_1^*(x_1,\mu_1,Z_t^{\mu_1,z_0}) - \sigma_1(x_2,\mu_2,Z_t^{\mu_2,z_0}) \|^2 \right)^{1/2} \right. \\ &\left. \mathbb{E} \|\sigma_1(x_1,\mu_1,Z_t^{\mu_1,z_0}) - \sigma_1(x_2,\mu_2,Z_t^{\mu_2,z_0}) \|^2 \right)^{1/2} \right. \\ &\leqslant C(|x_1 - x_2| + \rho(\mu_1,\mu_2)) + C \lim_{S \to \infty} \frac{1}{S} \int_0^S \left(\mathbb{E} |Z_t^{\mu_1,z_0} - Z_t^{\mu_2,z_0}|^2 \right)^{1/2} \, \mathrm{d}t \\ &\leqslant C(|x_1 - x_2| + \rho(\mu_1,\mu_2)). \end{split}$$

Finally, inserting the above inequality in (14), one could obtain that

$$\|\bar{\sigma}_1(x_1,\mu_1) - \bar{\sigma}_1(x_2,\mu_2)\| \le C(|x_1 - x_2| + \rho(\mu_1,\mu_2)).$$

Besides, by $(\mathbf{H}_{b_1,\sigma_1}^1)$ and [25, Lemma 3.8], it holds that

$$|\bar{b}_1(x_1, \mu_1) - \bar{b}_1(x_2, \mu_2)|^2 \leqslant L_{\bar{b}_1} \Big(|x_1 - x_2|^2 + \rho^2(\mu_1, \mu_2) \Big),$$
 (15)

where $L_{\bar{b}_1} > 0$ is a constant. Thus, Eq.(11) has a unique strong solution \bar{X} . (c.f. [3, Theorem 3.1]). Then, by the same deduction to that in Lemma 4.1, we also obtain the required estimate. This proof is complete.

Here, we remind that the solution \bar{X} of the average equation (11) is a Markov process and its infinitesimal generator is given by

$$(\mathcal{L}^{\bar{X},\mathcal{L}^{\mathbb{P}}_{\bar{X}}}F)(x,\mu): = \partial_{x_{i}}F(x,\mu)\bar{b}_{1}^{i}(x,\mu) + \frac{1}{2}\partial_{x_{i}x_{j}}F(x,\mu)(\bar{\sigma}_{1}\bar{\sigma}_{1}^{*})^{ij}(x,\mu)$$

$$+ \int_{\mathbb{R}^{n}}(\partial_{\mu}F)_{i}(x,\mu)(y)\bar{b}_{1}^{i}(y,\mu)\mu(\mathrm{d}y)$$

$$+ \frac{1}{2}\int_{\mathbb{R}^{n}}\partial_{y_{i}}(\partial_{\mu}F)_{j}(x,\mu)(y)(\bar{\sigma}_{1}\bar{\sigma}_{1}^{*})^{ij}(y,\mu)\mu(\mathrm{d}y), \quad F \in C^{2,2}(\mathbb{R}^{n}\times\mathcal{P}_{2}(\mathbb{R}^{n})).$$

4.4. **Some estimates for a Poisson equation.** Consider the following Poisson equation:

$$(\varepsilon \mathcal{L}^{Z^{\varepsilon}} \chi)(x, \mu, z) = -\left[(\mathcal{L}^{X^{\varepsilon}, \mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}} - \mathcal{L}^{\bar{X}, \mathcal{L}_{\bar{X}}^{\mathbb{P}}}) F \right] (x, \mu, z), \tag{16}$$

where $F \in C^{2,2}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$. Set

$$\Phi(x,\mu,z) := (\mathcal{L}^{X^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}}F)(x,\mu,z),$$

and then by the simple calculation, it holds that

$$\bar{\Phi}(x,\mu) := \int_{\mathbb{R}^m} \Phi(x,\mu,z) \nu^{\mu}(\mathrm{d}z) = (\mathcal{L}^{\bar{X},\mathscr{L}_{\bar{X}}^{\mathbb{P}}} F)(x,\mu).$$

Proposition 4.6. Suppose that assumptions $(\mathbf{H}^1_{b_1,\sigma_1})$, $(\mathbf{H}^2_{\sigma_1})$, $(\mathbf{H}^1_{b_2,\sigma_2})$ - $(\mathbf{H}^3_{b_2,\sigma_2})$ hold. Set

$$\chi_F(x,\mu,z) := \int_0^\infty P_t^\mu \left[\Phi(x,\mu,\cdot) - \bar{\Phi}(x,\mu) \right](z) \mathrm{d}t. \tag{17}$$

Then $\chi_F(x,\mu,z)$ is the unique solution for Eq.(16). Assume more $(\mathbf{H}_{b_1,\sigma_1}^3)$ and $F(\cdot,\cdot) \in C_b^{4,4}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$. Then $\chi_F(\cdot,\cdot,\cdot) \in C^{2,2,2}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m)$ and the following results hold for $x \in \mathbb{R}^n, \mu \in \mathcal{P}_2(\mathbb{R}^n), z \in \mathbb{R}^m$

$$\max\{|\chi_{F}(x,\mu,z)|, |\partial_{x}\chi_{F}(x,\mu,z)|, \|\partial_{\mu}\chi_{F}(x,\mu,z)(\cdot)\|_{L^{2}(\mu)}, |\partial_{z}\chi_{F}(x,\mu,z)|\}$$

$$\leqslant C(1+|x|+\|\mu\|+|z|),$$

$$\max\{\|\partial_{xx}\chi_{F}(x,\mu,z)\|, \|\partial_{y}\partial_{\mu}\chi_{F}(x,\mu,z)(\cdot)\|_{L^{2}(\mu)}\}$$

$$\leqslant C(1+|x|+\|\mu\|+|z|).$$

Proof. First of all, by the same deduction to that of [22, Lemma 4.5], it holds that

$$\begin{split} \int_0^\infty \left| P_t^\mu \left[\Phi(x,\mu,\cdot) - \bar{\Phi}(x,\mu) \right](z) \right| \mathrm{d}t &= \int_0^\infty |\mathbb{E}\Phi(x,\mu,Z_t^{\mu,z}) - \bar{\Phi}(x,\mu)| \mathrm{d}t \\ &\leqslant \int_0^\infty C e^{-\frac{\beta}{2}t} (1 + |x| + \|\mu\| + |z|) \mathrm{d}t \\ &= \frac{2C}{\beta} (1 + |x| + \|\mu\| + |z|). \end{split}$$

So the right side of (17) is well-defined. Then, acting the generator $\varepsilon \mathcal{L}^{Z^{\varepsilon}}$ on $\chi_F(x,\mu,z)$, we obtain that

$$\varepsilon(\mathcal{L}^{Z^{\varepsilon}}\chi_{F})(x,\mu,z) = \int_{0}^{\infty} (\varepsilon\mathcal{L}^{Z^{\varepsilon}}P_{t}^{\mu}) \left[\Phi(x,\mu,\cdot) - \bar{\Phi}(x,\mu)\right](z) dt
= \int_{0}^{\infty} \frac{dP_{t}^{\mu} \left[\Phi(x,\mu,\cdot) - \bar{\Phi}(x,\mu)\right](z)}{dt} dt
= \lim_{t \to \infty} P_{t}^{\mu} \left[\Phi(x,\mu,\cdot) - \bar{\Phi}(x,\mu)\right](z) - \left[\Phi(x,\mu,\cdot) - \bar{\Phi}(x,\mu)\right](z)
= -\left[\Phi(x,\mu,\cdot) - \bar{\Phi}(x,\mu)\right](z),$$

which yields that χ_F is a solution for Eq.(16). Besides, based on $(\mathbf{H}_{b_2,\sigma_2}^3)$, we know that the solutions of Eq.(16) are unique up to an additive constant. Thus, $\chi_F(x,\mu,z)$ is the unique solution for Eq.(16).

Under $(\mathbf{H}_{b_1,\sigma_1}^3)$ and $F(\cdot,\cdot) \in C_b^{4,4}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$, some computation with [25, Proposition 4.1] implies that $\chi_F(\cdot,\cdot,\cdot) \in C^{2,2,2}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m)$ and for $x \in \mathbb{R}^n$, $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, $z \in \mathbb{R}^m$

$$\max\{|\chi_{F}(x,\mu,z)|, |\partial_{x}\chi_{F}(x,\mu,z)|, \|\partial_{\mu}\chi_{F}(x,\mu,z)(\cdot)\|_{L^{2}(\mu)}, |\partial_{z}\chi_{F}(x,\mu,z)|\}$$

$$\leqslant C(1+|x|+\|\mu\|+|z|),$$

$$\max\{\|\partial_{xx}\chi_{F}(x,\mu,z)\|, \|\partial_{y}\partial_{\mu}\chi_{F}(x,\mu,z)(\cdot)\|_{L^{2}(\mu)}\}$$

$$\leqslant C(1+|x|+\|\mu\|+|z|).$$

The proof is complete.

Now, it is the position to prove Theorem 3.1.

Proof of Theorem 3.1.

First of all, applying the Itô formula to $F(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}})$ for $F \in C_b^{4,4}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$, one could obtain that for $0 \leq s < t \leq T$

$$F(X_{t}^{\varepsilon}, \mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}) = F(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}) + \int_{s}^{t} (\mathcal{L}^{X^{\varepsilon}, \mathscr{L}_{X^{\varepsilon}}^{\mathbb{P}}} F)(X_{r}^{\varepsilon}, \mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}, Z_{r}^{\varepsilon}) dr$$

$$+ \int_{s}^{t} \frac{\partial F}{\partial x_{i}} (X_{r}^{\varepsilon}, \mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}) \sigma_{1}^{ij} (X_{r}^{\varepsilon}, \mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}, Z_{r}^{\varepsilon}) dB_{r}^{j},$$

and furthermore

$$F(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}) - F(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}) - \int_{s}^{t} (\mathcal{L}^{\bar{X}, \mathcal{L}_{\bar{X}}^{\mathbb{P}}} F)(X_{r}^{\varepsilon}, \mathcal{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}) dr$$

$$= \int_{s}^{t} (\mathcal{L}^{X^{\varepsilon}, \mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}} F)(X_{r}^{\varepsilon}, \mathcal{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}, Z_{r}^{\varepsilon}) dr - \int_{s}^{t} (\mathcal{L}^{\bar{X}, \mathcal{L}_{\bar{X}}^{\mathbb{P}}} F)(X_{r}^{\varepsilon}, \mathcal{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}) dr$$

$$+ \int_{s}^{t} \frac{\partial F}{\partial x_{i}} (X_{r}^{\varepsilon}, \mathcal{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}) \sigma_{1}^{ij}(X_{r}^{\varepsilon}, \mathcal{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}, Z_{r}^{\varepsilon}) dB_{r}^{j}. \tag{18}$$

Thus, multiplying a bounded \mathscr{F}_s -measurable functional Γ_s of the process $\{\bar{X}_t, t \in [0, T]\}$ and taking the expectation under the measure \mathbb{P} on both sides of (18), we have that

$$\mathbb{E}\left[\Gamma_{s}(\bar{X}_{.})\left(F(X_{t}^{\varepsilon},\mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}})-F(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}})-\int_{s}^{t}(\mathcal{L}^{\bar{X}_{.}}\mathcal{L}_{\bar{X}}^{\mathbb{P}}F)(X_{r}^{\varepsilon},\mathcal{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}})\mathrm{d}r\right)\right]$$

$$=\mathbb{E}\left[\Gamma_{s}(\bar{X}_{.})\int_{s}^{t}\left[(\mathcal{L}^{X^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}}F)(X_{r}^{\varepsilon},\mathcal{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}},Z_{r}^{\varepsilon})-(\mathcal{L}^{\bar{X}_{.}}\mathcal{L}_{\bar{X}}^{\mathbb{P}}F)(X_{r}^{\varepsilon},\mathcal{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}})\right]\mathrm{d}r\right]$$

$$=-\mathbb{E}\left[\Gamma_{s}(\bar{X}_{.})\int_{s}^{t}\varepsilon(\mathcal{L}^{Z^{\varepsilon}}\chi_{F})(X_{r}^{\varepsilon},\mathcal{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}},Z_{r}^{\varepsilon})\mathrm{d}r\right],$$
(19)

where Proposition 4.6 is used in the last equality.

Next, we observe the right hand side of (19). By applying the Itô formula to $\varepsilon \chi_F(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}, Z_t^{\varepsilon})$, it holds that

$$\varepsilon \chi_{F}(X_{t}^{\varepsilon}, \mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon}) - \varepsilon \chi_{F}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) - \varepsilon \int_{s}^{t} (\mathcal{L}^{X^{\varepsilon}, \mathscr{L}_{X^{\varepsilon}}^{\mathbb{P}}} \chi_{F}) (X_{r}^{\varepsilon}, \mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}, Z_{r}^{\varepsilon}) dr
= \varepsilon \int_{s}^{t} (\mathcal{L}^{Z^{\varepsilon}} \chi_{F}) (X_{r}^{\varepsilon}, \mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}, Z_{r}^{\varepsilon}) dr + \int_{s}^{t} \frac{\varepsilon \partial \chi_{F}}{\partial x_{i}} (X_{r}^{\varepsilon}, \mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}, Z_{r}^{\varepsilon}) \sigma_{1}^{ij} (X_{r}^{\varepsilon}, \mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}, Z_{r}^{\varepsilon}) dB_{r}^{j}$$

$$+ \int_{s}^{t} \frac{\varepsilon \partial \chi_{F}}{\partial z_{i}} (X_{r}^{\varepsilon}, \mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}, Z_{r}^{\varepsilon}) \sigma_{2}^{ik} (X_{r}^{\varepsilon}, \mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}, Z_{r}^{\varepsilon}) dW_{r}^{k}.$$

So, multiplying Γ_s and taking the expectation on both sides of the above equality, we know that

$$\mathbb{E}\left[\Gamma_{s}(\bar{X}_{\cdot})\int_{s}^{t}\varepsilon(\mathcal{L}^{Z^{\varepsilon}}\chi_{F})(X_{r}^{\varepsilon},\mathcal{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}},Z_{r}^{\varepsilon})\mathrm{d}r\right]$$

$$= \mathbb{E}\left[\Gamma_{s}(\bar{X}_{\cdot})\left(\varepsilon\chi_{F}(X_{t}^{\varepsilon},\mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}},Z_{t}^{\varepsilon})-\varepsilon\chi_{F}(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}},Z_{s}^{\varepsilon})-\varepsilon\int_{s}^{t}(\mathcal{L}^{X^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}}\chi_{F})(X_{r}^{\varepsilon},\mathcal{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}},Z_{r}^{\varepsilon})\mathrm{d}r\right)\right],$$

which together with (19) yields that

$$\mathbb{E}\Big[\Gamma_{s}(\bar{X}_{\cdot})\Big(F(X_{t}^{\varepsilon},\mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}) - F(X_{s}^{\varepsilon},\mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}) - \int_{s}^{t} (\mathcal{L}^{\bar{X},\mathscr{L}_{\bar{X}}^{\mathbb{P}}}F)(X_{r}^{\varepsilon},\mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}) dr\Big)\Big]$$

$$= -\varepsilon \mathbb{E}\Big[\Gamma_{s}(\bar{X}_{\cdot})\Big(\chi_{F}(X_{t}^{\varepsilon},\mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}},Z_{t}^{\varepsilon}) - \chi_{F}(X_{s}^{\varepsilon},\mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}},Z_{s}^{\varepsilon}) - \int_{s}^{t} (\mathcal{L}^{X^{\varepsilon},\mathscr{L}_{X_{\varepsilon}}^{\mathbb{P}}}\chi_{F})(X_{r}^{\varepsilon},\mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}},Z_{r}^{\varepsilon}) dr\Big)\Big].$$

Taking the limits on two sides of the above equality, by the boundedness of $\Gamma_s(\bar{X}_{\cdot})$ and Proposition 4.6 we obtain that

$$\lim_{\varepsilon \to 0} \mathbb{E} \Big[\Gamma_s(\bar{X}_{\cdot}) \Big(F(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}) - F(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}) - \int_s^t (\mathcal{L}^{\bar{X}, \mathscr{L}_{\bar{X}}^{\mathbb{P}}} F)(X_r^{\varepsilon}, \mathscr{L}_{X_r^{\varepsilon}}^{\mathbb{P}}) \mathrm{d}r \Big) \Big] = 0,$$

which implies that the weak limit of $\{X_t^{\varepsilon}, t \in [0, T]\}$ is a solution of the martingale problem associated with $(\mathcal{L}^{\bar{X}, \mathcal{L}_{\bar{X}}^{\mathbb{P}}}, \delta_{x_0})$. Since the solution of the martingale problem associated with $(\mathcal{L}^{\bar{X}, \mathcal{L}_{\bar{X}}^{\mathbb{P}}}, \delta_{x_0})$ is unique, $\{X_t^{\varepsilon}, t \in [0, T]\}$ converges weakly to $\{\bar{X}_t, t \in [0, T]\}$, which completes the proof.

5. Proof of Theorem 3.2

In the section, we prove Theorem 3.2. To do this, we prepare some important estimates.

5.1. Some estimates for $\Lambda^{\varepsilon}_{\cdot}, \bar{\Lambda}_{\cdot}$

Lemma 5.1. Suppose that $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{b_2,\sigma_2}^1)$ and (\mathbf{H}_h) hold. Then we have that for any $q \geqslant 1$ and $s, t \in [0,T]$,

$$\mathbb{E}^{\mathbb{P}^\varepsilon} |\Lambda^\varepsilon_t|^{2q} \leqslant C, \qquad \mathbb{E}^{\mathbb{P}^\varepsilon} |\Lambda^\varepsilon_s - \Lambda^\varepsilon_t|^{2q} \leqslant C |s-t|^q,$$

where the constant C is independent of ε .

Proof. First of all, note that

$$\Lambda_t^{\varepsilon} = \exp\left\{ \int_0^t h^i(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) dY_s^{\varepsilon, i} - \frac{1}{2} \int_0^t |h(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon})|^2 ds \right\}.$$

Thus, by the Itô formula, it holds that

$$\Lambda_t^{\varepsilon} = 1 + \int_0^t \Lambda_s^{\varepsilon} h^i(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) dY_s^{\varepsilon, i}.$$
 (20)

Moreover, based on the BDG inequality and (\mathbf{H}_h) , one could obtain that

$$\begin{split} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Lambda_{t}^{\varepsilon}|^{2q} &\leqslant 2^{2q-1} + 2^{2q-1}l^{2q-1} \sum_{i=1}^{l} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \Big(\int_{0}^{t} \Lambda_{s}^{\varepsilon} h^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \mathrm{d}Y_{s}^{\varepsilon, i} \Big)^{2q} \\ &\leqslant 2^{2q-1} + C \sum_{i=1}^{l} \int_{0}^{t} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Lambda_{s}^{\varepsilon} h^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon})|^{2q} \mathrm{d}s \\ &\leqslant 2^{2q-1} + C \int_{0}^{t} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Lambda_{s}^{\varepsilon}|^{2q} \mathrm{d}s, \end{split}$$

which together with the Gronwall inequality yields that

$$\mathbb{E}^{\mathbb{P}^{\varepsilon}}(\Lambda_{t}^{\varepsilon})^{2q} \leqslant C. \tag{21}$$

Finally, we investigate that for $0 \le t < s \le T$,

$$\Lambda_s^{\varepsilon} - \Lambda_t^{\varepsilon} = \int_t^s \Lambda_r^{\varepsilon} h^i(X_r^{\varepsilon}, \mathscr{L}_{X_r^{\varepsilon}}^{\mathbb{P}}, Z_r^{\varepsilon}) dY_r^{\varepsilon, i},$$

and furthermore

$$\mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Lambda_{s}^{\varepsilon} - \Lambda_{t}^{\varepsilon}|^{2q} \leqslant l^{2q-1} \sum_{i=1}^{l} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left(\int_{t}^{s} \Lambda_{r}^{\varepsilon} h^{i}(X_{r}^{\varepsilon}, \mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}, Z_{r}^{\varepsilon}) \mathrm{d}Y_{r}^{\varepsilon, i} \right)^{2q}$$

$$\leqslant C(s-t)^{q-1} \sum_{i=1}^{l} \int_{t}^{s} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Lambda_{r}^{\varepsilon} h^{i}(X_{r}^{\varepsilon}, \mathscr{L}_{X_{r}^{\varepsilon}}^{\mathbb{P}}, Z_{r}^{\varepsilon})|^{2q} \mathrm{d}r$$

$$\leqslant C(s-t)^{q-1} \sum_{i=1}^{l} \int_{t}^{s} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Lambda_{r}^{\varepsilon}|^{2q} \mathrm{d}r$$

$$\leqslant C(s-t)^{q},$$

where the last inequality is based on (21).

 $\textbf{Lemma 5.2.} \ \ \textit{Under these assumptions} \ (\textbf{H}^1_{b_1,\sigma_1}), \ (\textbf{H}^1_{b_2,\sigma_2}) \ \ \textit{and} \ (\textbf{H}_h), \ \textit{it holds that}$

$$\sup_{t \in [0,T]} \mathbb{E} |\Lambda_t^{\varepsilon}|^{-2} \leqslant C.$$

Proof. Note that

$$\begin{split} \mathbb{E}(\Lambda_t^{\varepsilon})^{-2} &= \mathbb{E}\left[\left(\exp\left\{-\int_0^t h^i(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) \mathrm{d}V_s^i - \frac{1}{2}\int_0^t |h(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon})|^2 \mathrm{d}s\right\}\right)^2\right] \\ &= \mathbb{E}\left[\exp\left\{-\int_0^t 2h^i(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) \mathrm{d}V_s^i - \frac{1}{2}\int_0^t |2h(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon})|^2 \mathrm{d}s\right\} \\ & \cdot \exp\left\{\frac{1}{2}\int_0^t |2h(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon})|^2 \mathrm{d}s - \int_0^t |h(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon})|^2 \mathrm{d}s\right\}\right] \\ & \leqslant \exp\{CT\}\mathbb{E}\left[\exp\left\{-\int_0^t 2h^i(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) \mathrm{d}V_s^i - \frac{1}{2}\int_0^t |2h(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon})|^2 \mathrm{d}s\right\}\right] \\ &= \exp\{CT\}, \end{split}$$

where the last equality is based on the fact that $\exp\left\{-\int_0^t 2h^i(X_s^\varepsilon, \mathscr{L}_{X_s^\varepsilon}^\mathbb{P}, Z_s^\varepsilon) dV_s^i - \frac{1}{2} \int_0^t |2h(X_s^\varepsilon, \mathscr{L}_{X_s^\varepsilon}^\mathbb{P}, Z_s^\varepsilon)|^2 ds\right\}$ is an exponential martingale. The proof is complete. \square

Lemma 5.3. Under these assumptions $(\mathbf{H}^1_{b_1,\sigma_1})$, $(\mathbf{H}^2_{\sigma_1})$, $(\mathbf{H}^1_{b_2,\sigma_2})$ - $(\mathbf{H}^2_{b_2,\sigma_2})$, (\mathbf{H}_h) , there exists a constant C>0 such that for any $q\geqslant 1$,

$$\sup_{t \in [0,T]} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\bar{\Lambda}_t|^{2q} \leqslant C.$$

Since the proof of the above lemma is similar to that for Λ^{ε} in Lemma 5.1, we omit it.

5.2. Some estimates for $\rho_t^{\varepsilon}(1)$, $\bar{\rho}_t(1)$. Let us firstly establish the Zakai equations about Eq.(1) and Eq.(11).

Lemma 5.4. (The Zakai equation)

(i) For $\Psi \in C_b^{2,2,2}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m)$, the Zakai equation of Eq.(1) is given by

$$\rho_t^{\varepsilon}(\Psi) = \rho_0^{\varepsilon}(\Psi) + \int_0^t \rho_s^{\varepsilon}(\mathcal{L}^{X^{\varepsilon}, \mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}, Z^{\varepsilon}}\Psi) ds + \int_0^t \rho_s^{\varepsilon}(\Psi h^i) dY_s^{\varepsilon, i}, \quad \rho_0^{\varepsilon}(\Psi) = \Psi(x_0, \delta_{x_0}, z_0), (22)$$

(ii) For $F \in C_b^{2,2}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$, the Zakai equation of Eq.(11) is given by

$$\bar{\rho}_t(F) = \bar{\rho}_0(F) + \int_0^t \bar{\rho}_s(\mathcal{L}^{\bar{X},\mathcal{L}_{\bar{X}}^{\mathbb{P}}} F) \mathrm{d}s + \int_0^t \bar{\rho}_s(F\bar{h}^i) \mathrm{d}Y_s^{\varepsilon,i}, \quad \bar{\rho}_0(F) = F(x_0, \delta_{x_0}). \tag{23}$$

Proof. Applying the Itô formula to $\Psi(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}, Z_t^{\varepsilon})$, one could obtain that

$$\Psi(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}, Z_t^{\varepsilon}) = \Psi(x_0, \delta_{x_0}, z_0) + \int_0^t (\mathcal{L}^{X^{\varepsilon}, \mathscr{L}_{X^{\varepsilon}}^{\mathbb{P}}, Z^{\varepsilon}} \Psi)(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) ds
+ \int_0^t \frac{\partial \Psi}{\partial x_i} (X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) \sigma_1^{ij} (X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) dB_s^j
+ \int_0^t \frac{\partial \Psi}{\partial z_i} (X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) \sigma_2^{ik} (X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, Z_s^{\varepsilon}) dW_s^k.$$

So, by combining the above deduction with (20), the Itô formula implies that

$$\Psi(X_{t}^{\varepsilon}, \mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon})\Lambda_{t}^{\varepsilon} = \Psi(x_{0}, \delta_{x_{0}}, z_{0}) + \int_{0}^{t} \Lambda_{s}^{\varepsilon} (\mathcal{L}^{X^{\varepsilon}, \mathscr{L}_{X^{\varepsilon}}^{\mathbb{P}}, Z^{\varepsilon}} \Psi)(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) ds
+ \int_{0}^{t} \Psi(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} h^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}) dY_{s}^{\varepsilon, i}
+ \int_{0}^{t} \Lambda_{s}^{\varepsilon} \frac{\partial \Psi}{\partial x_{i}} (X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \sigma_{1}^{ij}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) dB_{s}^{j}
+ \int_{0}^{t} \Lambda_{s}^{\varepsilon} \frac{\partial \Psi}{\partial z_{i}} (X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \sigma_{2}^{ik}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) dW_{s}^{k}.$$

Taking the conditional expectation about $\mathscr{F}_t^{Y^{\varepsilon}}$ on both sides of the above equality under the measure \mathbb{P}^{ε} , one can get that

$$\mathbb{E}^{\mathbb{P}^{\varepsilon}}[\Psi(X_{t}^{\varepsilon}, \mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon})\Lambda_{t}^{\varepsilon}|\mathscr{F}_{t}^{Y^{\varepsilon}}]$$

$$= \Psi(x_{0}, \delta_{x_{0}}, z_{0}) + \int_{0}^{t} \mathbb{E}^{\mathbb{P}^{\varepsilon}}[\Lambda_{s}^{\varepsilon}(\mathcal{L}^{X^{\varepsilon}, \mathscr{L}_{X^{\varepsilon}}^{\mathbb{P}}, Z^{\varepsilon}}\Psi)(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon})|\mathscr{F}_{s}^{Y^{\varepsilon}}] ds$$

$$\begin{split} &+ \int_{0}^{t} \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} h^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}) |\mathscr{F}_{s}^{Y^{\varepsilon}}] \mathrm{d} Y_{s}^{\varepsilon, i} \\ &= \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi(x_{0}, \delta_{x_{0}}, z_{0}) |\mathscr{F}_{0}^{Y^{\varepsilon}}] + \int_{0}^{t} \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Lambda_{s}^{\varepsilon} (\mathcal{L}^{X^{\varepsilon}, \mathscr{L}_{X^{\varepsilon}}^{\mathbb{P}}, Z^{\varepsilon}} \Psi) (X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) |\mathscr{F}_{s}^{Y^{\varepsilon}}] \mathrm{d} s \\ &+ \int_{0}^{t} \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} h^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}) |\mathscr{F}_{s}^{Y^{\varepsilon}}] \mathrm{d} Y_{s}^{\varepsilon, i}, \end{split}$$

which together with the definition of $\rho_t^{\varepsilon}(\Psi)$ yields that

$$\rho_t^{\varepsilon}(\Psi) = \rho_0^{\varepsilon}(\Psi) + \int_0^t \rho_s^{\varepsilon}(\mathcal{L}^{X^{\varepsilon}, \mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}, Z^{\varepsilon}}\Psi) ds + \int_0^t \rho_s^{\varepsilon}(\Psi h^i) dY_s^{\varepsilon, i}, \quad \rho_0^{\varepsilon}(\Psi) = \Psi(x_0, \delta_{x_0}, z_0).$$

By the same deduction to that of (i), we obtain (ii). The proof is complete.

Lemma 5.5. Under these assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{\sigma_1}^2)$, $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^2)$, (\mathbf{H}_h) , there exists a constant C > 0 such that for any $q \geqslant 1$,

$$\mathbb{E}\left(\sup_{t\in[0,T]}|\rho_t^{\varepsilon}(1)|^q\right)\leqslant C,\quad \mathbb{E}\left(\sup_{t\in[0,T]}|\bar{\rho}_t(1)|^q\right)\leqslant C.$$

Proof. First of all, from the Hölder inequality, it follows that

$$\mathbb{E}\left(\sup_{t\in[0,T]}|\rho_t^{\varepsilon}(1)|^q\right) = \mathbb{E}^{\mathbb{P}^{\varepsilon}}\left[\Lambda_T^{\varepsilon}\left(\sup_{t\in[0,T]}|\rho_t^{\varepsilon}(1)|^q\right)\right] = \left[\mathbb{E}^{\mathbb{P}^{\varepsilon}}(\Lambda_T^{\varepsilon})^2\right]^{\frac{1}{2}}\left[\mathbb{E}^{\mathbb{P}^{\varepsilon}}\left(\sup_{t\in[0,T]}|\rho_t^{\varepsilon}(1)|^{2q}\right)\right]^{\frac{1}{2}}.(24)$$

Then we estimate $|\rho_t^{\varepsilon}(1)|^{2q}$. Since $\rho_t^{\varepsilon}(1)$ satisfies Eq.(22) with $\Psi(x,\mu,z)=1$, it holds that

$$\begin{split} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left(\sup_{t \in [0,T]} |\rho_{t}^{\varepsilon}(1)|^{2q} \right) & \leqslant \quad 2^{2q-1} + 2^{2q-1} l^{2q-1} \sum_{i=1}^{l} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left(\sup_{t \in [0,T]} \left| \int_{0}^{t} \rho_{s}^{\varepsilon}(h^{i}) \mathrm{d}Y_{s}^{\varepsilon,i} \right|^{2q} \right) \\ & \leqslant \quad 2^{2q-1} + 2^{2q-1} l^{2q-1} \left[\frac{2q(2q-1)}{2} \right]^{q} \sum_{i=1}^{l} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left(\int_{0}^{T} |\rho_{s}^{\varepsilon}(h^{i})|^{2} \mathrm{d}s \right)^{q} \\ & \leqslant \quad 2^{2q-1} + 2^{2q-1} l^{2q-1} \left[\frac{2q(2q-1)}{2} \right]^{q} T^{q-1} \sum_{i=1}^{l} \int_{0}^{T} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\rho_{s}^{\varepsilon}(h^{i})|^{2q} \mathrm{d}s \\ & \leqslant \quad 2^{2q-1} + C \sum_{i=1}^{l} \int_{0}^{T} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [h^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} | \mathscr{F}_{s}^{Y^{\varepsilon}} \right] \right|^{2q} \mathrm{d}s \\ & \leqslant \quad 2^{2q-1} + C \sum_{i=1}^{l} \int_{0}^{T} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left| h^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} \right|^{2q} \mathrm{d}s \\ & \leqslant \quad 2^{2q-1} + C \int_{0}^{T} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Lambda_{s}^{\varepsilon}|^{2q} \mathrm{d}s \\ & \leqslant \quad 2^{2q-1} + C \int_{0}^{T} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Lambda_{s}^{\varepsilon}|^{2q} \mathrm{d}s \\ & \leqslant \quad C \end{split}$$

where the last step is based on (21). So, inserting the above inequality in (24), by (21) we obtain that

$$\mathbb{E}\left(\sup_{t\in[0,T]}|\rho_t^\varepsilon(1)|^q\right)\leqslant C.$$

Finally, the same deduction to that for the above inequality implies the second estimate. The proof is complete. \Box

Lemma 5.6. Under these assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{\sigma_1}^2)$, $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^2)$, (\mathbf{H}_h) , there exists a constant C > 0 such that for $t \in [0,T]$

$$(\bar{\rho}_t(1))^{-1} \leqslant C$$
, a.s..

Since the proof of the above lemma is similar to that of [21, Lemma 5.1], we omit it.

5.3. Some estimates for $\xi^{\varepsilon} := \rho^{\varepsilon,x,\mu} - \bar{\rho}$. In order to investigate $\xi^{\varepsilon} := \rho^{\varepsilon,x,\mu} - \bar{\rho}$, we prepare some following stronger moment estimates.

Lemma 5.7. Under assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$ $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^{2'})$, there exists a constant C > 0 such that

$$\sup_{t \in [0,T]} \mathbb{E}|X_t^{\varepsilon}|^{2p} \leqslant C(1+|x_0|^{2p}+|z_0|^{2p}), \quad \sup_{t \in [0,T]} \mathbb{E}|Z_t^{\varepsilon}|^{2p} \leqslant C(1+|x_0|^{2p}+|z_0|^{2p}),$$

where p is the same to that in $(\mathbf{H}_{b_2,\sigma_2}^{2'})$.

We don't prove the lemma above, since its proof is similar to that for Lemma 4.1.

Lemma 5.8. Under these assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$ $(\mathbf{H}_{\sigma_1}^2)$ $(\mathbf{H}_{b_2,\sigma_2}^1)$, $(\mathbf{H}_{b_2,\sigma_2}^2)$, and (\mathbf{H}_h) , $\{\xi_t^{\varepsilon}, t \in [0,T]\}$ is relatively weakly compact in $C([0,T], \mathcal{M}(\mathbb{R}^n \times \mathscr{P}_2(\mathbb{R}^n)))$, where $\mathcal{M}(\mathbb{R}^n \times \mathscr{P}_2(\mathbb{R}^n))$ stands for the collection of all the finite measures on $\mathbb{R}^n \times \mathscr{P}_2(\mathbb{R}^n)$.

Proof. First of all, from Lemma 5.5, it follows that for $F \in C_b^{2,2}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$,

$$\mathbb{E}|\xi_t^{\varepsilon}(F)| = \mathbb{E}|\rho_t^{\varepsilon,x,\mu}(F) - \bar{\rho}_t(F)| = \mathbb{E}|\rho_t^{\varepsilon}(F) - \bar{\rho}_t(F)| \leqslant \mathbb{E}|\rho_t^{\varepsilon}(F)| + \mathbb{E}|\bar{\rho}_t(F)|$$

$$\leqslant \|F\|_{C_b^{2,2}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))} \mathbb{E}|\rho_t^{\varepsilon}(1)| + \|F\|_{C_b^{2,2}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))} \mathbb{E}|\bar{\rho}_t(1)|$$

$$\leqslant C\|F\|_{C_b^{2,2}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))},$$

which yields that

$$\sup_{\varepsilon} \sup_{t \in [0,T]} \mathbb{E}|\xi_t^{\varepsilon}(F)| < \infty. \tag{25}$$

Besides, by Lemma 5.4, it holds that

$$\rho_t^{\varepsilon,x,\mu}(F) = F(x_0, \delta_{x_0}) + \int_0^t \rho_s^{\varepsilon,x,\mu} \left((\mathcal{L}^{X^{\varepsilon}, \mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}} F)(\cdot, \cdot, Z_s^{\varepsilon}) \right) ds + \int_0^t \rho_s^{\varepsilon,x,\mu} \left(F h^i(\cdot, \cdot, Z_s^{\varepsilon}) \right) dY_s^{\varepsilon,i},$$

$$\bar{\rho}_t(F) = F(x_0, \delta_{x_0}) + \int_0^t \bar{\rho}_s (\mathcal{L}^{\bar{X}, \mathcal{L}_{\bar{X}}^{\mathbb{P}}} F) ds + \int_0^t \bar{\rho}_s (F \bar{h}^i) dY_s^{\varepsilon,i}.$$

Based on the above equation and the isometric formula, we can obtain that for any $\delta > 0$ and any $\{\mathscr{F}_t\}$ -stopping time $0 \leq \tau < \tau + \delta \leq T$,

$$\mathbb{E}|\xi_{\tau+\delta}^{\varepsilon}(F) - \xi_{\tau}^{\varepsilon}(F)|^{2} = \mathbb{E}^{\mathbb{P}^{\varepsilon}}\left[|\xi_{\tau+\delta}^{\varepsilon}(F) - \xi_{\tau}^{\varepsilon}(F)|^{2}\Lambda_{T}^{\varepsilon}\right] \\ \leqslant (\mathbb{E}^{\mathbb{P}^{\varepsilon}}|\xi_{\tau+\delta}^{\varepsilon}(F) - \xi_{\tau}^{\varepsilon}(F)|^{4})^{1/2}(\mathbb{E}^{\mathbb{P}^{\varepsilon}}(\Lambda_{T}^{\varepsilon})^{2})^{1/2},$$

and

$$\mathbb{E}^{\mathbb{P}^{\varepsilon}} |\xi_{\tau+\delta}^{\varepsilon}(F) - \xi_{\tau}^{\varepsilon}(F)|^{4} \leqslant \mathbb{E}^{\mathbb{P}^{\varepsilon}} |(\rho_{\tau+\delta}^{\varepsilon,x,\mu}(F) - \bar{\rho}_{\tau+\delta}(F)) - (\rho_{\tau}^{\varepsilon,x,\mu}(F) - \bar{\rho}_{\tau}(F))|^{4}$$

$$\leqslant 2^{3} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\rho_{\tau+\delta}^{\varepsilon,x,\mu}(F) - \rho_{\tau}^{\varepsilon,x,\mu}(F)|^{4} + 2^{3} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\bar{\rho}_{\tau+\delta}(F) - \bar{\rho}_{\tau}(F)|^{4}$$

$$\leqslant 4^{3}\mathbb{E}^{\mathbb{P}^{\varepsilon}} \Big| \int_{\tau}^{\tau+\delta} \rho_{s}^{\varepsilon,x,\mu} \left((\mathcal{L}^{X^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}} F)(\cdot,\cdot,Z_{s}^{\varepsilon}) \right) ds \Big|^{4} \\
+4^{3}\mathbb{E}^{\mathbb{P}^{\varepsilon}} \Big| \int_{\tau}^{\tau+\delta} \rho_{s}^{\varepsilon,x,\mu} \left(Fh^{i}(\cdot,\cdot,Z_{s}^{\varepsilon}) \right) dY_{s}^{\varepsilon,i} \Big|^{4} \\
+4^{3}\mathbb{E}^{\mathbb{P}^{\varepsilon}} \Big| \int_{\tau}^{\tau+\delta} \bar{\rho}_{s} (\mathcal{L}^{\bar{X},\mathcal{L}_{\bar{X}}^{\mathbb{P}}} F) ds \Big|^{4} + 4^{3}\mathbb{E}^{\mathbb{P}^{\varepsilon}} \Big| \int_{\tau}^{\tau+\delta} \bar{\rho}_{s} (F\bar{h}^{i}) dY_{s}^{\varepsilon,i} \Big|^{4} \\
\leqslant 4^{3}\delta^{3}\mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{\tau}^{\tau+\delta} \Big| \rho_{s}^{\varepsilon,x,\mu} \left((\mathcal{L}^{X^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}} F)(\cdot,\cdot,Z_{s}^{\varepsilon}) \right) \Big|^{4} ds \\
+4^{3}\delta\mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{\tau}^{\tau+\delta} \Big| \rho_{s}^{\varepsilon,x,\mu} \left(Fh^{i}(\cdot,\cdot,Z_{s}^{\varepsilon}) \right) \Big|^{4} ds \\
+4^{3}\delta^{3}\mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{\tau}^{\tau+\delta} \Big| \bar{\rho}_{s} (\mathcal{L}^{\bar{X},\mathcal{L}_{\bar{X}}^{\mathbb{P}}} F) \Big|^{4} ds + 4^{3}\delta\mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{\tau}^{\tau+\delta} |\bar{\rho}_{s}(F\bar{h}^{i})|^{4} ds \\
=: I_{1} + I_{2} + I_{3} + I_{4}.$$

For $I_1 + I_3$, it holds that

$$\begin{split} I_{1} + I_{3} &\leqslant 4^{3} \delta^{3} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{\tau}^{\tau + \delta} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \Big[|(\mathcal{L}^{X^{\varepsilon}, \mathcal{L}^{\mathbb{P}}_{X^{\varepsilon}}} F)(X_{s}^{\varepsilon}, \mathcal{L}^{\mathbb{P}}_{X_{s}^{\varepsilon}}, Z_{s}^{\varepsilon})|^{4} |\Lambda_{s}^{\varepsilon}|^{4} |\mathcal{F}^{Y^{\varepsilon}}_{S}| \, \mathrm{d}s \\ &+ 4^{3} \delta^{3} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{\tau}^{\tau + \delta} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \Big[|(\mathcal{L}^{\bar{X}, \mathcal{L}^{\mathbb{P}}_{\bar{X}}} F)(\bar{X}_{s}, \mathcal{L}^{\mathbb{P}}_{\bar{X}_{s}})|^{4} |\bar{\Lambda}_{s}|^{4} |\mathcal{F}^{Y^{\varepsilon}}_{S}| \, \mathrm{d}s \\ &\leqslant 4^{3} \delta^{3} C \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{\tau}^{\tau + \delta} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \Big[(1 + |X_{s}^{\varepsilon}|^{8} + \mathbb{E}|X_{s}^{\varepsilon}|^{8} + |Z_{s}^{\varepsilon}|^{8}) |\Lambda_{s}^{\varepsilon}|^{4} |\mathcal{F}^{Y^{\varepsilon}}_{S}| \, \mathrm{d}s \\ &+ 4^{3} \delta^{3} C \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{\tau}^{\tau + \delta} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \Big[(1 + |X_{s}^{\varepsilon}|^{8} + \mathbb{E}|X_{s}^{\varepsilon}|^{8}) |\bar{\Lambda}_{s}|^{4} |\mathcal{F}^{Y^{\varepsilon}}_{S}| \, \mathrm{d}s \\ &\leqslant 4^{3} \delta^{3} C \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{0}^{\delta} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \Big[(1 + |X_{\tau + u}^{\varepsilon}|^{8} + C + |Z_{\tau + u}^{\varepsilon}|^{8}) |\Lambda_{\tau + u}^{\varepsilon}|^{4} |\mathcal{F}^{Y^{\varepsilon}}_{\tau + u}| \, \mathrm{d}u \\ &+ 4^{3} \delta^{3} C \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{0}^{\delta} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \Big[(1 + |\bar{X}_{\tau + u}|^{8} + C) |\bar{\Lambda}_{\tau + u}|^{4} |\mathcal{F}^{Y^{\varepsilon}}_{\tau + u}| \, \mathrm{d}u \\ &\leqslant 4^{3} \delta^{3} C \int_{0}^{\delta} \left(1 + \mathbb{E}^{\mathbb{P}^{\varepsilon}} |X_{\tau + u}^{\varepsilon}|^{16} + C + \mathbb{E}^{\mathbb{P}^{\varepsilon}} |Z_{\tau + u}^{\varepsilon}|^{16} \right)^{1/2} \left(\mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Lambda_{\tau + u}^{\varepsilon}|^{8} \right)^{1/2} \, \mathrm{d}u \\ &\leqslant C \delta^{4}, \end{split}$$

where the last inequality is based on Lemma 5.7, 5.1, 4.5 and 5.3. Moreover, by the boundedness of F, h, we have that $I_2 + I_4 \leq C\delta^2$. Thus, these estimates imply that

$$\mathbb{E}|\xi_{\tau+\delta}^{\varepsilon}(F) - \xi_{\tau}^{\varepsilon}(F)|^{2} \leqslant C(\delta^{2} + \delta),$$

and then

$$\lim_{\delta \to 0} \limsup_{\epsilon \downarrow 0} \sup_{\tau \leqslant T} \mathbb{E} |\xi_{\tau+\delta}^{\epsilon}(F) - \xi_{\tau}^{\epsilon}(F)|^2 = 0.$$
 (26)

Finally, combining (25) (26) with [10, Theorem 2.7], we obtain that $\{\xi_t^{\varepsilon}(F), t \in [0, T]\}$ is relatively weakly compact in $C([0, T], \mathbb{R})$, which together with [11, Theorem 6.2] yields that $\{\xi_t^{\varepsilon}, t \in [0, T]\}$ is relatively weakly compact.

Here, in order to prove that $\{\xi^{\varepsilon_k}\}$ converges weakly to 0, we need the following estimate.

Lemma 5.9. Suppose that the assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$ $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^{2'})$ hold. Then it holds that for any $\delta > 0$ and any $t \in [0,T]$,

$$\mathbb{E}|X_{t+\delta}^{\varepsilon} - X_t^{\varepsilon}|^{2p} \leqslant C(1 + |x_0|^{2p} + |z_0|^{2p})(\delta^{2p} + \delta^p),$$

where p is the same to that in $(\mathbf{H}_{b_2,\sigma_2}^{2'})$.

We don't prove the lemma above, since its proof is similar to that for Lemma 4.2.

Proposition 5.10. Under these assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$ $(\mathbf{H}_{\sigma_1}^2)$ $(\mathbf{H}_{b_1,\sigma_1}^3)$ $(\mathbf{H}_{b_2,\sigma_2}^1)$ $(\mathbf{H}_{b_2,\sigma_2}^3)$ and (\mathbf{H}_h) , there exists a subsequence $\{\xi^{\varepsilon_k}\}$ which converges weakly to 0 in $C([0,T], \mathcal{M}(\mathbb{R}^n \times \mathscr{P}_2(\mathbb{R}^n)))$.

Proof. Step 1. We prove that for any $t \in [0,T]$ and $F \in C_b^{4,4}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$

$$\lim_{\varepsilon \to 0} \mathbb{E} \left| \xi_t^{\varepsilon}(F) - \int_0^t \xi_s^{\varepsilon} (\mathcal{L}^{\bar{X}, \mathscr{L}_{\bar{X}}^{\mathbb{P}}} F) \mathrm{d}s - \int_0^t \xi_s^{\varepsilon} (F\bar{h}^i) \mathrm{d}Y_s^{\varepsilon, i} \right|^2 = 0.$$

First of all, we define a perturbed test function $F^{\varepsilon}(x,\mu,z)$ as follows:

$$F^{\varepsilon}(x,\mu,z) = F(x,\mu) + \varepsilon \chi_F(x,\mu,z),$$

where $\chi_F(x,\mu,z)$ is the unique solution of the Poisson equation (16). Note that

$$\rho_t^\varepsilon(F^\varepsilon) = \rho_t^\varepsilon(F) + \rho_t^\varepsilon(\varepsilon\chi_F) = \rho_t^{\varepsilon,x,\mu}(F) + \varepsilon\rho_t^\varepsilon(\chi_F).$$

Thus, based on Lemma 5.4, we get that

$$\begin{split} \xi_t^{\varepsilon}(F) &= \rho_t^{\varepsilon,x,\mu}(F) - \bar{\rho}_t(F) = \rho_t^{\varepsilon}(F^{\varepsilon}) - \varepsilon \rho_t^{\varepsilon}(\chi_F) - \bar{\rho}_t(F) \\ &= -\varepsilon \rho_t^{\varepsilon}(\chi_F) + \rho_0^{\varepsilon}(F^{\varepsilon}) + \int_0^t \rho_s^{\varepsilon}(\mathcal{L}^{X^{\varepsilon},\mathcal{L}^{\mathcal{P}}_{X^{\varepsilon}},\mathcal{Z}^{\varepsilon}}F^{\varepsilon}) \mathrm{d}s + \int_0^t \rho_s^{\varepsilon}(F^{\varepsilon}h^i) \mathrm{d}Y_s^{\varepsilon,i} \\ &- \left(\bar{\rho}_0(F) + \int_0^t \bar{\rho}_s(\mathcal{L}^{\bar{X},\mathcal{L}^{\mathcal{P}}_{X}}F) \mathrm{d}s + \int_0^t \bar{\rho}_s(F\bar{h}^i) \mathrm{d}Y_s^{\varepsilon,i}\right) \\ &= &- \varepsilon \rho_t^{\varepsilon}(\chi_F) + \rho_0^{\varepsilon,x,\mu}(F) + \varepsilon \rho_0^{\varepsilon}(\chi_F) + \int_0^t \rho_s^{\varepsilon}(\mathcal{L}^{X^{\varepsilon},\mathcal{L}^{\mathcal{P}}_{X^{\varepsilon}}}F + \varepsilon \mathcal{L}^{X^{\varepsilon},\mathcal{L}^{\mathcal{P}}_{X^{\varepsilon}}}\chi_F + \varepsilon \mathcal{L}^{Z^{\varepsilon}}\chi_F) \mathrm{d}s \\ &+ \int_0^t \rho_s^{\varepsilon}(Fh^i) \mathrm{d}Y_s^{\varepsilon,i} + \varepsilon \int_0^t \rho_s^{\varepsilon}(\chi_F h^i) \mathrm{d}Y_s^{\varepsilon,i} - \bar{\rho}_0(F) - \int_0^t \bar{\rho}_s(\mathcal{L}^{\bar{X},\mathcal{L}^{\mathcal{P}}_{X}}F) \mathrm{d}s \\ &- \int_0^t \bar{\rho}_s(F\bar{h}^i) \mathrm{d}Y_s^{\varepsilon,i} \\ &= &- \varepsilon \rho_t^{\varepsilon}(\chi_F) + \varepsilon \rho_0^{\varepsilon}(\chi_F) + \int_0^t \rho_s^{\varepsilon}(\mathcal{L}^{\bar{X},\mathcal{L}^{\mathcal{P}}_{X}}F) \mathrm{d}s + \varepsilon \int_0^t \rho_s^{\varepsilon}(\mathcal{L}^{X^{\varepsilon},\mathcal{L}^{\mathcal{P}}_{X^{\varepsilon}}}\chi_F) \mathrm{d}s \\ &+ \int_0^t \rho_s^{\varepsilon}(Fh^i) \mathrm{d}Y_s^{\varepsilon,i} + \varepsilon \int_0^t \rho_s^{\varepsilon}(\chi_F h^i) \mathrm{d}Y_s^{\varepsilon,i} - \int_0^t \bar{\rho}_s(\mathcal{L}^{\bar{X},\mathcal{L}^{\mathcal{P}}_{X}}F) \mathrm{d}s \\ &- \int_0^t \bar{\rho}_s(F\bar{h}^i) \mathrm{d}Y_s^{\varepsilon,i} + \varepsilon \int_0^t \rho_s^{\varepsilon}(\chi_F h^i) \mathrm{d}Y_s^{\varepsilon,i} - \int_0^t \bar{\rho}_s(\mathcal{L}^{\bar{X},\mathcal{L}^{\mathcal{P}}_{X}}F) \mathrm{d}s \\ &- \int_0^t \bar{\rho}_s(F\bar{h}^i) \mathrm{d}Y_s^{\varepsilon,i} + \varepsilon \int_0^t \rho_s^{\varepsilon}(\chi_F h^i) \mathrm{d}Y_s^{\varepsilon,i} - \int_0^t \bar{\rho}_s(\mathcal{L}^{\bar{X},\mathcal{L}^{\mathcal{P}}_{X}}F) \mathrm{d}s \\ &- \int_0^t \bar{\rho}_s(F\bar{h}^i) \mathrm{d}Y_s^{\varepsilon,i} + \varepsilon \int_0^t \rho_s^{\varepsilon}(\chi_F h^i) \mathrm{d}Y_s^{\varepsilon,i} - \int_0^t \bar{\rho}_s(\mathcal{L}^{\bar{X},\mathcal{L}^{\mathcal{P}}_{X}}F) \mathrm{d}s \\ &- \int_0^t \bar{\rho}_s(F\bar{h}^i) \mathrm{d}Y_s^{\varepsilon,i} + \varepsilon \int_0^t \rho_s^{\varepsilon}(\chi_F h^i) \mathrm{d}Y_s^{\varepsilon,i} - \int_0^t \bar{\rho}_s(\mathcal{L}^{\bar{X},\mathcal{L}^{\mathcal{P}}_{X}}F) \mathrm{d}s \\ &- \int_0^t \bar{\rho}_s(F\bar{h}^i) \mathrm{d}Y_s^{\varepsilon,i} + \varepsilon \int_0^t \rho_s^{\varepsilon}(\chi_F h^i) \mathrm{d}Y_s^{\varepsilon,i} - \varepsilon \int_0^t \bar{\rho}_s(\chi_F h^i) \mathrm{d}Y_s^{\varepsilon,i} - \varepsilon$$

$$= -\varepsilon \rho_{t}^{\varepsilon}(\chi_{F}) + \varepsilon \rho_{0}^{\varepsilon}(\chi_{F}) + \int_{0}^{t} \rho_{s}^{\varepsilon,x,\mu}(\mathcal{L}^{\bar{X},\mathcal{L}^{\mathbb{P}}_{\bar{X}}}F) ds + \varepsilon \int_{0}^{t} \rho_{s}^{\varepsilon}(\mathcal{L}^{X^{\varepsilon},\mathcal{L}^{\mathbb{P}}_{X^{\varepsilon}}}\chi_{F}) ds$$

$$+ \int_{0}^{t} \rho_{s}^{\varepsilon}(Fh^{i}) dY_{s}^{\varepsilon,i} + \varepsilon \int_{0}^{t} \rho_{s}^{\varepsilon}(\chi_{F}h^{i}) dY_{s}^{\varepsilon,i} - \int_{0}^{t} \bar{\rho}_{s}(\mathcal{L}^{\bar{X},\mathcal{L}^{\mathbb{P}}_{\bar{X}}}F) ds - \int_{0}^{t} \bar{\rho}_{s}(F\bar{h}^{i}) dY_{s}^{\varepsilon,i}$$

$$- \int_{0}^{t} \rho_{s}^{\varepsilon,x,\mu}(F\bar{h}^{i}) dY_{s}^{\varepsilon,i} + \int_{0}^{t} \rho_{s}^{\varepsilon,x,\mu}(F\bar{h}^{i}) dY_{s}^{\varepsilon,i}$$

$$= -\varepsilon \rho_{t}^{\varepsilon}(\chi_{F}) + \varepsilon \rho_{0}^{\varepsilon}(\chi_{F}) + \int_{0}^{t} \xi_{s}^{\varepsilon}(\mathcal{L}^{\bar{X},\mathcal{L}^{\mathbb{P}}_{\bar{X}}}F) ds + \int_{0}^{t} \xi_{s}^{\varepsilon}(F\bar{h}^{i}) dY_{s}^{\varepsilon,i}$$

$$+ \varepsilon \int_{0}^{t} \rho_{s}^{\varepsilon}(\mathcal{L}^{X^{\varepsilon},\mathcal{L}^{\mathbb{P}}_{X^{\varepsilon}}}\chi_{F}) ds + \varepsilon \int_{0}^{t} \rho_{s}^{\varepsilon}(\chi_{F}h^{i}) dY_{s}^{\varepsilon,i} + \int_{0}^{t} \rho_{s}^{\varepsilon}(Fh^{i} - F\bar{h}^{i}) dY_{s}^{\varepsilon,i},$$

where the fact that $\rho_0^{\varepsilon,x,\mu}(F) = \bar{\rho}_0(F) = F(x_0, \delta_{x_0})$ is used in the fourth equality. Moreover, it holds that

$$\mathbb{E} \left| \xi_{t}^{\varepsilon}(F) - \int_{0}^{t} \xi_{s}^{\varepsilon}(\mathcal{L}^{\bar{X},\mathcal{L}_{\bar{X}}^{\mathbb{P}}}F) ds - \int_{0}^{t} \xi_{s}^{\varepsilon}(F\bar{h}^{i}) dY_{s}^{\varepsilon,i} \right|^{2} \\
= \mathbb{E} \left| -\varepsilon \rho_{t}^{\varepsilon}(\chi_{F}) + \varepsilon \rho_{0}^{\varepsilon}(\chi_{F}) + \varepsilon \int_{0}^{t} \rho_{s}^{\varepsilon}(\mathcal{L}^{X^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}}\chi_{F}) ds + \varepsilon \int_{0}^{t} \rho_{s}^{\varepsilon}(\chi_{F}h^{i}) dY_{s}^{\varepsilon,i} \right|^{2} \\
+ \int_{0}^{t} \rho_{s}^{\varepsilon}(Fh^{i} - F\bar{h}^{i}) dY_{s}^{\varepsilon,i} \right|^{2} \\
\leq 5\varepsilon^{2} \mathbb{E} |\rho_{t}^{\varepsilon}(\chi_{F})|^{2} + 5\varepsilon^{2} \mathbb{E} |\rho_{0}^{\varepsilon}(\chi_{F})|^{2} + 5\varepsilon^{2} \mathbb{E} \left| \int_{0}^{t} \rho_{s}^{\varepsilon}(\mathcal{L}^{X^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}}\chi_{F}) ds \right|^{2} \\
+ 5\varepsilon^{2} \mathbb{E} \left| \int_{0}^{t} \rho_{s}^{\varepsilon}(\chi_{F}h^{i}) dY_{s}^{\varepsilon,i} \right|^{2} + 5\mathbb{E} \left| \int_{0}^{t} \rho_{s}^{\varepsilon}(Fh^{i} - F\bar{h}^{i}) dY_{s}^{\varepsilon,i} \right|^{2} \\
=: J_{1} + J_{2} + J_{3} + J_{4} + J_{5}. \tag{27}$$

For J_1 , based on the Hölder inequality, the Jensen inequality, Lemma 5.1, 5.7 and Proposition 4.6, it holds that

$$J_{1} = 5\varepsilon^{2}\mathbb{E}^{\mathbb{P}^{\varepsilon}}[|\rho_{t}^{\varepsilon}(\chi_{F})|^{2}\Lambda_{T}^{\varepsilon}] \leq 5\varepsilon^{2}(\mathbb{E}^{\mathbb{P}^{\varepsilon}}|\rho_{t}^{\varepsilon}(\chi_{F})|^{4})^{1/2}(\mathbb{E}^{\mathbb{P}^{\varepsilon}}|\Lambda_{T}^{\varepsilon}|^{2})^{1/2}$$

$$\leq 5\varepsilon^{2}C(\mathbb{E}^{\mathbb{P}^{\varepsilon}}[\mathbb{E}^{\mathbb{P}^{\varepsilon}}[|\chi_{F}(X_{t}^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}},Z_{t}^{\varepsilon})|^{4}|\Lambda_{t}^{\varepsilon}|^{4}|\mathcal{F}_{t}^{Y^{\varepsilon}}]])^{1/2}$$

$$\leq 5\varepsilon^{2}C(\mathbb{E}^{\mathbb{P}^{\varepsilon}}|\chi_{F}(X_{t}^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}},Z_{t}^{\varepsilon})|^{8})^{1/4}(\mathbb{E}^{\mathbb{P}^{\varepsilon}}|\Lambda_{t}^{\varepsilon}|^{8})^{1/4}$$

$$\leq 5\varepsilon^{2}C(\mathbb{E}^{\mathbb{P}^{\varepsilon}}(1+|X_{t}^{\varepsilon}|^{8}+\|\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}\|^{8}+|Z_{t}^{\varepsilon}|^{8}))^{1/4}$$

$$\leq C\varepsilon^{2}. \tag{28}$$

We treat J_2 . It is easy to see that

$$J_2 = 5\varepsilon^2 |\chi_F(x_0, \delta_{x_0}, z_0)|^2.$$
 (29)

For J_3 , Proposition 4.6 and (5) implies

$$|(\mathcal{L}^{X^{\varepsilon},\mathcal{L}_{X^{\varepsilon}}^{\mathbb{P}}}\chi_{F})(x,\mu,z)| \leqslant C(1+|x|+||\mu||+|z|).$$

And based on the Hölder inequality and the same deduction to that for J_1 , we know that

$$J_3 \leqslant 5T\varepsilon^2 \int_0^T \mathbb{E}|\rho_s^{\varepsilon}(\mathcal{L}^{X^{\varepsilon},\mathscr{L}_{X^{\varepsilon}}^{\mathbb{P}}}\chi_F)|^2 ds \leqslant C\varepsilon^2.$$
 (30)

For J_4 , by the boundedness of h, the BDG inequality and the similar deduction to that for J_1 , it holds that

$$J_{4} = 5\varepsilon^{2} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\Lambda_{T}^{\varepsilon} \left| \int_{0}^{t} \rho_{s}^{\varepsilon}(\chi_{F}h^{i}) dY_{s}^{\varepsilon,i} \right|^{2} \right]$$

$$\leq 5\varepsilon^{2} \left[\mathbb{E}^{\mathbb{P}^{\varepsilon}} (\Lambda_{T}^{\varepsilon})^{2} \right]^{\frac{1}{2}} \left[\mathbb{E}^{\mathbb{P}^{\varepsilon}} \left| \int_{0}^{t} \rho_{s}^{\varepsilon}(\chi_{F}h^{i}) dY_{s}^{\varepsilon,i} \right|^{4} \right]^{\frac{1}{2}}$$

$$\leq C\varepsilon^{2} \left[\sum_{i=1}^{l} \int_{0}^{T} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\rho_{s}^{\varepsilon}(\chi_{F}h^{i})|^{4} ds \right]^{\frac{1}{2}} \leq C\varepsilon^{2}. \tag{31}$$

Next, we deal with J_5 . Set $\Psi(x,\mu,z) := F(x,\mu)h(x,\mu,z) - F(x,\mu)\bar{h}(x,\mu)$, and then by the boundedness of F and (\mathbf{H}_h) , it holds that $\Psi(x,\mu,z)$ is bounded and Lipschitz continuous. Then, applying the Hölder inequality and the BDG inequality, one could obtain that

$$J_{5} = 5\mathbb{E} \left| \int_{0}^{t} \rho_{s}^{\varepsilon}(\Psi^{i}) dY_{s}^{\varepsilon,i} \right|^{2} = 5\mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\Lambda_{T}^{\varepsilon} \left| \int_{0}^{t} \rho_{s}^{\varepsilon}(\Psi^{i}) dY_{s}^{\varepsilon,i} \right|^{2} \right]$$

$$\leq 5[\mathbb{E}^{\mathbb{P}^{\varepsilon}} (\Lambda_{T}^{\varepsilon})^{2}]^{\frac{1}{2}} \left[\mathbb{E}^{\mathbb{P}^{\varepsilon}} \left| \int_{0}^{t} \rho_{s}^{\varepsilon}(\Psi^{i}) dY_{s}^{\varepsilon,i} \right|^{4} \right]^{\frac{1}{2}} \leq C \left[\sum_{i=1}^{l} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{0}^{T} \left| \rho_{s}^{\varepsilon}(\Psi^{i}) \right|^{4} ds \right]^{\frac{1}{2}}$$

$$\leq C \left[\sum_{i=1}^{l} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{0}^{T} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} |\mathcal{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right]^{\frac{1}{2}}.$$

$$(32)$$

In the following, we are devoted to estimating the last term of the above inequality. First, we divide the time interval [0, T] into parts of size δ_{ε} : $0 = t_0 < t_1 < t_2 < \cdots < t_N \leq T, N = \left[\frac{T}{\delta_{\varepsilon}}\right], t_{j+1} - t_j = \delta_{\varepsilon}, j = 0, 1, 2, \cdots, N-1$. Then it holds that

$$\mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\int_{0}^{T} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} | \mathscr{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right]$$

$$= \sum_{j=0}^{N-1} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\int_{t_{j}}^{t_{j+1}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} | \mathscr{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right]$$

$$+ \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\int_{N\delta_{\varepsilon}}^{T} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} | \mathscr{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right].$$

For clarity and simplicity, we use $[t, t + \delta_{\varepsilon}]$ instead of $[t_j, t_{j+1}]$, and the analysis for the remainder term over the interval follows from the same argument. So, we know that

$$\mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\int_{0}^{T} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} | \mathscr{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right]$$

$$\leqslant (N+1) \sup_{t \in [0,T]} \left[\mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{t}^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} | \mathscr{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right]$$

$$\leqslant \left(\frac{T}{\delta_{\varepsilon}} + 1\right) \sup_{t \in [0,T]} \left[\mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{t}^{t + \delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\Psi^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} | \mathscr{F}_{s}^{Y^{\varepsilon}} \right] \right|^{4} \mathrm{d}s \right]. \tag{33}$$

Next, we study

$$\mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{t}^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} | \mathscr{F}_{s}^{Y^{\varepsilon}}] \right|^{4} \mathrm{d}s.$$

From the Hölder inequality and the Jensen inequality, it follows that

$$\mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\int_{t}^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} | \mathcal{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right] \\
= \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\int_{t}^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} - \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} \right. \\
\left. + \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} - \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} + \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} | \mathcal{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right] \\
\leqslant 3^{3} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\int_{t}^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} - \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} | \mathcal{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right] \\
+ 3^{3} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\int_{t}^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} - \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} | \mathcal{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right] \\
\leqslant 3^{3} \int_{t}^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} | \mathcal{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right] \\
\leqslant 3^{3} \int_{t}^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left| \left(\Psi^{i}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) - \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \right) \Lambda_{s}^{\varepsilon} \right|^{4} ds \right] \\
+ 3^{3} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\int_{t}^{t+\delta_{\varepsilon}} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left| \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} | \mathcal{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right] \\
+ 3^{3} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\int_{t}^{t+\delta_{\varepsilon}} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left| \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} | \mathcal{F}_{s}^{Y^{\varepsilon}}] \right|^{4} ds \right] \\
=: K_{1} + K_{2} + K_{3}. \tag{34}$$

For K_1 , by the Hölder inequality, and Lemma 5.1 and 5.9, it holds that

$$K_{1} \leqslant 3^{3} \int_{t}^{t+\delta_{\varepsilon}} \left(\mathbb{E}^{\mathbb{P}^{\varepsilon}} | \Psi^{i}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) - \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) |^{8} \right)^{\frac{1}{2}} [\mathbb{E}^{\mathbb{P}^{\varepsilon}} | \Lambda_{s}^{\varepsilon} |^{8}]^{\frac{1}{2}} ds$$

$$\leqslant C \int_{t}^{t+\delta_{\varepsilon}} \left(\mathbb{E}^{\mathbb{P}^{\varepsilon}} | \Psi^{i}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) - \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) |^{8} \right)^{\frac{1}{2}} ds$$

$$\leqslant C \int_{t}^{t+\delta_{\varepsilon}} \left(\mathbb{E}^{\mathbb{P}^{\varepsilon}} \left(|X_{s}^{\varepsilon} - X_{t}^{\varepsilon}|^{8} + \rho^{8} (\mathcal{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}) \right) \right)^{\frac{1}{2}} ds$$

$$\leqslant C \int_{t}^{t+\delta_{\varepsilon}} \left(\mathbb{E}^{\mathbb{P}^{\varepsilon}} |X_{s}^{\varepsilon} - X_{t}^{\varepsilon}|^{8} + \mathbb{E} |X_{s}^{\varepsilon} - X_{t}^{\varepsilon}|^{8} \right)^{\frac{1}{2}} ds$$

$$= C \int_{t}^{t+\delta_{\varepsilon}} \left(\mathbb{E}(\Lambda_{T}^{\varepsilon})^{-1} |X_{s}^{\varepsilon} - X_{t}^{\varepsilon}|^{8} + \mathbb{E} |X_{s}^{\varepsilon} - X_{t}^{\varepsilon}|^{8} \right)^{\frac{1}{2}} ds$$

$$\leqslant C \int_{t}^{t+\delta_{\varepsilon}} \left((\mathbb{E}|X_{s}^{\varepsilon} - X_{t}^{\varepsilon}|^{16})^{\frac{1}{2}} + \mathbb{E}|X_{s}^{\varepsilon} - X_{t}^{\varepsilon}|^{8} \right)^{\frac{1}{2}} ds$$

$$\leqslant C \int_{t}^{t+\delta_{\varepsilon}} (\delta_{\varepsilon}^{4} + \delta_{\varepsilon}^{2}) ds \leqslant C(\delta_{\varepsilon}^{5} + \delta_{\varepsilon}^{3}), \tag{35}$$

where $\rho^2(\mathscr{L}_{X_s^{\varepsilon}}^{\mathbb{P}}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}) \leqslant \mathbb{E}|X_s^{\varepsilon} - X_t^{\varepsilon}|^2$.

From the boundedness of Ψ and Lemma 5.1, it follows that

$$K_2 \leqslant C \int_{t}^{t+\delta_{\varepsilon}} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Lambda_s^{\varepsilon} - \Lambda_t^{\varepsilon}|^4 \mathrm{d}s \leqslant C \delta_{\varepsilon}^3.$$
 (36)

To treat K_3 , we introduce an auxiliary process as follows: for $s \in [k\delta_{\varepsilon}, (k+1)\delta_{\varepsilon})$,

$$\begin{cases}
d\hat{Z}_{s}^{\varepsilon} = \frac{1}{\varepsilon} b_{2}(\mathscr{L}_{X_{k\delta_{\varepsilon}}^{\varepsilon}}^{\mathbb{P}}, \hat{Z}_{s}^{\varepsilon}) ds + \frac{1}{\sqrt{\varepsilon}} \sigma_{2}(\mathscr{L}_{X_{k\delta_{\varepsilon}}^{\varepsilon}}^{\mathbb{P}}, \hat{Z}_{s}^{\varepsilon}) dW_{s}, \\
\hat{Z}_{k\delta_{\varepsilon}}^{\varepsilon} = Z_{k\delta_{\varepsilon}}^{\varepsilon}.
\end{cases} (37)$$

Then it holds that

$$K_{3} = 3^{3} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{t}^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{t}^{\varepsilon}, \mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} - \Psi^{i}(X_{t}^{\varepsilon}, \mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, \hat{Z}_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} \right| + \Psi^{i}(X_{t}^{\varepsilon}, \mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, \hat{Z}_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} |\mathscr{F}_{s}^{Y^{\varepsilon}}|^{4} ds$$

$$\leq 6^{3} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{t}^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{t}^{\varepsilon}, \mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} - \Psi^{i}(X_{t}^{\varepsilon}, \mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, \hat{Z}_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} |\mathscr{F}_{s}^{Y^{\varepsilon}}|^{4} ds$$

$$+ 6^{3} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_{t}^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^{i}(X_{t}^{\varepsilon}, \mathscr{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, \hat{Z}_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} |\mathscr{F}_{s}^{Y^{\varepsilon}}|^{4} ds$$

$$=: D_{1} + D_{2}.$$

For D_1 , based on the Jensen inequality, the Hölder inequality and (21), it holds that

$$D_{1} \leqslant C \int_{t}^{t+\delta_{\varepsilon}} \mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon} - \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, \hat{Z}_{s}^{\varepsilon}) \Lambda_{t}^{\varepsilon}|^{4} ds$$

$$\leqslant C \int_{t}^{t+\delta_{\varepsilon}} \left(\mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) - \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, \hat{Z}_{s}^{\varepsilon})|^{8} \right)^{\frac{1}{2}} (\mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Lambda_{t}^{\varepsilon}|^{8})^{\frac{1}{2}} ds$$

$$\leqslant C \int_{t}^{t+\delta_{\varepsilon}} \left(\mathbb{E}^{\mathbb{P}^{\varepsilon}} |\Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) - \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, \hat{Z}_{s}^{\varepsilon})|^{8} \right)^{\frac{1}{2}} ds$$

$$= C \int_{t}^{t+\delta_{\varepsilon}} \left(\mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[|\Psi^{i}(x, \delta_{x}, Z_{s}^{\varepsilon, \delta_{x}, z}) - \Psi^{i}(x, \delta_{x}, \hat{Z}_{s}^{\varepsilon, \delta_{x}, z})|^{8} \right] ds$$

$$\left| (x, \delta_{x}, z) = (X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon}) \right] ds, \tag{38}$$

where $Z_s^{\varepsilon,\delta_x,z}$ represents the solution to the fast part of Eq.(1), but $\mathscr{L}_{X^{\varepsilon}}^{\mathbb{P}}$ starts in δ_x at the time t and Z^{ε} starts in z at the time t, and $\hat{Z}_s^{\varepsilon,\delta_x,z}$ stands for the solution to the auxiliary equation (37), but $\mathscr{L}_{X^{\varepsilon}}^{\mathbb{P}}$ starts in δ_x at the time t and Z^{ε} starts in z at the time t. By Lemma 5.2, it holds that for $s \in [t, t + \delta_{\varepsilon}]$

$$\mathbb{E}^{\mathbb{P}^{\varepsilon}}[|\Psi^{i}(x,\delta_{x},Z_{s}^{\varepsilon,\delta_{x},z}) - \Psi^{i}(x,\delta_{x},\hat{Z}_{s}^{\varepsilon,\delta_{x},z})|^{8}] \leqslant C\mathbb{E}^{\mathbb{P}^{\varepsilon}}[|Z_{s}^{\varepsilon,\delta_{x},z} - \hat{Z}_{s}^{\varepsilon,\delta_{x},z}|^{8}]$$

$$\leqslant C \left(\mathbb{E}[|Z_s^{\varepsilon,\delta_x,z} - \hat{Z}_s^{\varepsilon,\delta_x,z}|^{16}] \right)^{\frac{1}{2}} \left(\mathbb{E}[|\Lambda_T^{\varepsilon}|^{-2}] \right)^{\frac{1}{2}}$$

$$\leqslant C \left(\mathbb{E}[|Z_s^{\varepsilon,\delta_x,z} - \hat{Z}_s^{\varepsilon,\delta_x,z}|^{16}] \right)^{\frac{1}{2}}.$$

Note that

$$\mathbb{E}[|Z_s^{\varepsilon,\delta_x,z} - \hat{Z}_s^{\varepsilon,\delta_x,z}|^{16}] \leqslant C(1+|x|^{16}+|z|^{16})(\frac{\delta_{\varepsilon}^{15}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^7}{\varepsilon^8})(\delta_{\varepsilon}^{17} + \delta_{\varepsilon}^9)e^{C(\frac{\delta_{\varepsilon}^{16}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^8}{\varepsilon^8})}, \tag{39}$$

where the above inequality is proved in the Appendix. Thus, we have that

$$\mathbb{E}^{\mathbb{P}^{\varepsilon}}[|\Psi^{i}(x,\delta_{x},Z_{s}^{\varepsilon,\delta_{x},z}) - \Psi^{i}(x,\delta_{x},\hat{Z}_{s}^{\varepsilon,\delta_{x},z})|^{8}] \\ \leqslant C(1+|x|^{8}+|z|^{8})\left(\left(\frac{\delta_{\varepsilon}^{15}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^{7}}{\varepsilon^{8}}\right)(\delta_{\varepsilon}^{17} + \delta_{\varepsilon}^{9})e^{C(\frac{\delta_{\varepsilon}^{16}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^{8}}{\varepsilon^{8}})}\right)^{\frac{1}{2}}.$$

Inserting the above inequality into (38), by Lemma 5.7 one could obtain that

$$D_1 \leqslant C\delta_{\varepsilon} \left(\left(\frac{\delta_{\varepsilon}^{15}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^{7}}{\varepsilon^{8}} \right) (\delta_{\varepsilon}^{17} + \delta_{\varepsilon}^{9}) e^{C\left(\frac{\delta_{\varepsilon}^{16}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^{8}}{\varepsilon^{8}} \right)} \right)^{\frac{1}{4}}. \tag{40}$$

Next, we calculate D_2 . Set

$$\mathcal{H}_s := \mathscr{F}_s^{Y^{\varepsilon}} \vee \mathscr{F}_t^{X^{\varepsilon}} \vee \mathscr{F}_t^{Z^{\varepsilon}}, \quad s \in [t, t + \delta_{\varepsilon}),$$

where $\mathscr{F}_t^{X^{\varepsilon}}$, $\mathscr{F}_t^{Z^{\varepsilon}}$ denote the usual augmentation of $\sigma\{X_r^{\varepsilon}, 0 \leqslant r \leqslant t\}$, $\sigma\{Z_r^{\varepsilon}, 0 \leqslant r \leqslant t\}$, respectively. Then from the tower property of the conditional expectation, the Jensen inequality, Lemma 5.1 and the Hölder inequality, it follows that

$$\begin{split} D_2 &= 6^3 \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_t^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^i(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}^{\varepsilon}}, \hat{Z}_s^{\varepsilon}) \Lambda_t^{\varepsilon} | \mathscr{F}_s^{Y^{\varepsilon}}] \right|^4 \mathrm{d}s \\ &= 6^3 \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_t^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^i(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}^{\varepsilon}}, \hat{Z}_s^{\varepsilon}) \Lambda_t^{\varepsilon} | \mathcal{F}_s^{Y^{\varepsilon}}] \right|^4 \mathrm{d}s \\ &= 6^3 \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_t^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\Lambda_t^{\varepsilon} \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^i(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}^{\varepsilon}}, \hat{Z}_s^{\varepsilon}) | \mathcal{H}_s] | \mathscr{F}_s^{Y^{\varepsilon}} \right] \right|^4 \mathrm{d}s \\ &= 6^3 \mathbb{E}^{\mathbb{P}^{\varepsilon}} \int_t^{t+\delta_{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[\Lambda_t^{\varepsilon} \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^i(x, \delta_x, \hat{Z}_s^{\varepsilon, \delta_x, z}) | (x, \delta_x, z) = (X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}^{\varepsilon}}, Z_t^{\varepsilon})] | \mathscr{F}_s^{Y^{\varepsilon}} \right] \right|^4 \mathrm{d}s \\ &\leq 6^3 \int_t^{t+\delta_{\varepsilon}} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left| \Lambda_t^{\varepsilon} \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^i(x, \delta_x, \hat{Z}_s^{\varepsilon, \delta_x, z}) | (x, \delta_x, z) = (X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}^{\varepsilon}}, Z_t^{\varepsilon})] \right|^4 \mathrm{d}s \\ &\leq 6^3 \int_t^{t+\delta_{\varepsilon}} \left(\mathbb{E}^{\mathbb{P}^{\varepsilon}} | \Lambda_t^{\varepsilon} |^8 \right)^{\frac{1}{2}} \left(\mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^i(x, \delta_x, \hat{Z}_s^{\varepsilon, \delta_x, z}) | (x, \delta_x, z) = (X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}^{\varepsilon}}, Z_t^{\varepsilon})] \right|^4 \mathrm{d}s \\ &\leq C \int_t^{t+\delta_{\varepsilon}} \left(\mathbb{E}^{\mathbb{P}^{\varepsilon}} | \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^i(x, \delta_x, \hat{Z}_s^{\varepsilon, \delta_x, z}) | (x, \delta_x, z) = (X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}^{\varepsilon}}, Z_t^{\varepsilon})] \right|^8 \mathrm{d}s \\ &\leq C \delta_{\varepsilon}^{\frac{1}{2}} \left(\int_t^{t+\delta_{\varepsilon}} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^i(x, \delta_x, \hat{Z}_s^{\varepsilon, \delta_x, z}) | (x, \delta_x, z) = (X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}^{\varepsilon}}, Z_t^{\varepsilon})] \right|^8 \mathrm{d}s \right)^{\frac{1}{2}} \\ &= C \delta_{\varepsilon}^{\frac{1}{2}} \left(\int_t^{t+\delta_{\varepsilon}} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left| \mathbb{E}^{\mathbb{P}^{\varepsilon}} [\Psi^i(x, \delta_x, \hat{Z}_s^{\varepsilon, \delta_x, z}) | (x, \delta_x, z) = (X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}^{\varepsilon}}, Z_t^{\varepsilon})] \right|^8 \mathrm{d}s \right)^{\frac{1}{2}} \end{aligned}$$

$$= C\delta_{\varepsilon}^{\frac{1}{2}} \varepsilon^{\frac{1}{2}} \left(\int_{0}^{\frac{\delta_{\varepsilon}}{\varepsilon}} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[P_{r}^{\mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}} \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon}) \right]^{8} dr \right)^{\frac{1}{2}}$$

$$\leq C\delta_{\varepsilon}^{\frac{1}{2}} \varepsilon^{\frac{1}{2}} \left(\int_{0}^{\infty} \mathbb{E}^{\mathbb{P}^{\varepsilon}} \left[P_{r}^{\mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}} \Psi^{i}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}, Z_{t}^{\varepsilon}) \right]^{8} dr \right)^{\frac{1}{2}}. \tag{41}$$

In the following, we deal with $|P_r^{\mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}}\Psi^i(X_t^{\varepsilon},\mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}},Z_t^{\varepsilon})|^8$. Note that

$$P_r^{\mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}} \Psi^i(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}, Z_t^{\varepsilon}) = \mathbb{E}\left[\Psi^i(x, \delta_x, Z_r^{\delta_x, z}) \middle| (x, \delta_x, z) = (X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}, Z_t^{\varepsilon})\right].$$

Thus, by the same deduction as Lemma 4.5 in [22], we get that

$$\begin{split} & \left| P_r^{\mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}} \Psi^i(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}, Z_t^{\varepsilon}) - \bar{\Psi}^i(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}) \right|^2 \\ = & \left| \mathbb{E} \left[\Psi^i(x, \delta_x, Z_r^{\delta_x, z}) - \bar{\Psi}^i(x, \delta_x) \middle| (x, \delta_x, z) = (X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}, Z_t^{\varepsilon}) \right] \right|^2 \\ \leqslant & C e^{-\beta' r} (1 + |X_t^{\varepsilon}|^2 + \|\mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}\|^2 + |Z_t^{\varepsilon}|^2), \end{split}$$

where $\bar{\Psi}(x,\mu) := \int_{\mathbb{R}^m} \Psi(x,\mu,z) \nu^{\mu}(\mathrm{d}z)$. Since $\bar{\Psi}(x,\mu) = 0$, one could obtain that

$$\left| P_r^{\mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}} \Psi^i(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}, Z_t^{\varepsilon}) \right|^2 \leqslant C e^{-\beta' r} (1 + |X_t^{\varepsilon}|^2 + \|\mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}\|^2 + |Z_t^{\varepsilon}|^2),$$

and

$$\left| P_r^{\mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}} \Psi^i(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}, Z_t^{\varepsilon}) \right|^8 \leqslant C e^{-4\beta' r} (1 + |X_t^{\varepsilon}|^8 + \|\mathscr{L}_{X_t^{\varepsilon}}^{\mathbb{P}}\|^8 + |Z_t^{\varepsilon}|^8).$$

Next, by inserting the above inequality in (41), it holds that

$$D_{2} \leqslant C\delta_{\varepsilon}^{\frac{1}{2}} \varepsilon^{\frac{1}{2}} \left(\int_{0}^{\infty} e^{-4\beta' r} \mathbb{E}^{\mathbb{P}^{\varepsilon}} (1 + |X_{t}^{\varepsilon}|^{8} + \|\mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}\|^{8} + |Z_{t}^{\varepsilon}|^{8}) dr \right)^{\frac{1}{2}}$$

$$\leqslant C\delta_{\varepsilon}^{\frac{1}{2}} \varepsilon^{\frac{1}{2}} \left(\mathbb{E}^{\mathbb{P}^{\varepsilon}} (1 + |X_{t}^{\varepsilon}|^{8} + \|\mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}\|^{8} + |Z_{t}^{\varepsilon}|^{8}) \right)^{\frac{1}{2}}$$

$$\leqslant C\delta_{\varepsilon}^{\frac{1}{2}} \varepsilon^{\frac{1}{2}} \left(\mathbb{E} (1 + |X_{t}^{\varepsilon}|^{16} + \|\mathcal{L}_{X_{t}^{\varepsilon}}^{\mathbb{P}}\|^{16} + |Z_{t}^{\varepsilon}|^{16}) \right)^{\frac{1}{4}}$$

$$\leqslant C\delta_{\varepsilon}^{\frac{1}{2}} \varepsilon^{\frac{1}{2}}, \tag{42}$$

where Lemma 5.7 is used in the last inequality.

So, combining (42) with (40), we know that

$$K_3 \leqslant C\delta_{\varepsilon} \left(\left(\frac{\delta_{\varepsilon}^{15}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^{7}}{\varepsilon^{8}} \right) (\delta_{\varepsilon}^{17} + \delta_{\varepsilon}^{9}) e^{C\left(\frac{\delta_{\varepsilon}^{16}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^{8}}{\varepsilon^{8}} \right)} \right)^{\frac{1}{4}} + C\delta_{\varepsilon}^{\frac{1}{2}} \varepsilon^{\frac{1}{2}}. \tag{43}$$

Inserting (35), (36) and (43) into (34), one can get that

$$\begin{split} & \mathbb{E}^{\mathbb{P}^{\varepsilon}} \Bigg[\int_{t}^{t+\delta_{\varepsilon}} \Big| \mathbb{E}^{\mathbb{P}^{\varepsilon}} \big[\Psi^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} \big| \mathscr{F}_{s}^{Y^{\varepsilon}} \big] \Big|^{4} \mathrm{d}s \Bigg] \\ \leqslant & C(\delta_{\varepsilon}^{5} + \delta_{\varepsilon}^{3}) + C\delta_{\varepsilon}^{3} + C\delta_{\varepsilon} \Big((\frac{\delta_{\varepsilon}^{15}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^{7}}{\varepsilon^{8}}) (\delta_{\varepsilon}^{17} + \delta_{\varepsilon}^{9}) e^{C(\frac{\delta_{\varepsilon}^{16}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^{8}}{\varepsilon^{8}})} \Big)^{\frac{1}{4}} + C\delta_{\varepsilon}^{\frac{1}{2}} \varepsilon^{\frac{1}{2}}, \end{split}$$

which together with (33) implies that

$$\begin{split} & \mathbb{E}^{\mathbb{P}^{\varepsilon}} \Bigg[\int_{0}^{T} \Big| \mathbb{E}^{\mathbb{P}^{\varepsilon}} \big[\Psi^{i}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}^{\mathbb{P}}, Z_{s}^{\varepsilon}) \Lambda_{s}^{\varepsilon} \big| \mathscr{F}_{s}^{Y^{\varepsilon}} \big] \Big|^{4} \mathrm{d}s \Bigg] \\ & \leqslant \quad (\frac{T}{\delta_{\varepsilon}} + 1) \Big[C(\delta_{\varepsilon}^{5} + \delta_{\varepsilon}^{3}) + C\delta_{\varepsilon}^{3} + C\delta_{\varepsilon} \Big((\frac{\delta_{\varepsilon}^{15}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^{7}}{\varepsilon^{8}}) (\delta_{\varepsilon}^{17} + \delta_{\varepsilon}^{9}) e^{C(\frac{\delta_{\varepsilon}^{16}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^{8}}{\varepsilon^{8}})} \Big)^{\frac{1}{4}} + C\delta_{\varepsilon}^{\frac{1}{2}} \varepsilon^{\frac{1}{2}} \Big] \\ & \leqslant \quad C(T + \delta_{\varepsilon}) \Big[(\delta_{\varepsilon}^{4} + \delta_{\varepsilon}^{2}) + \delta_{\varepsilon}^{2} + \Big((\frac{\delta_{\varepsilon}^{15}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^{7}}{\varepsilon^{8}}) (\delta_{\varepsilon}^{17} + \delta_{\varepsilon}^{9}) e^{C(\frac{\delta_{\varepsilon}^{16}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^{8}}{\varepsilon^{8}})} \Big)^{\frac{1}{4}} + \frac{\varepsilon^{\frac{1}{2}}}{\delta_{\varepsilon}^{\frac{1}{2}}} \Big]. \end{split}$$

From the above inequality and (32), it follows that

$$J_5 \leqslant C(T + \delta_{\varepsilon})^{\frac{1}{2}} \left[(\delta_{\varepsilon}^4 + \delta_{\varepsilon}^2) + \delta_{\varepsilon}^2 + \left(\left(\frac{\delta_{\varepsilon}^{15}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^7}{\varepsilon^8} \right) (\delta_{\varepsilon}^{17} + \delta_{\varepsilon}^9) e^{C(\frac{\delta_{\varepsilon}^{16}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^8}{\varepsilon^8})} \right)^{\frac{1}{4}} + \frac{\varepsilon^{\frac{1}{2}}}{\delta_{\varepsilon}^{\frac{1}{2}}} \right]^{\frac{1}{2}}. \tag{44}$$

Finally, inserting (28)-(31) and (44) into (27), we have

$$\mathbb{E} \left| \xi_t^{\varepsilon}(F) - \int_0^t \xi_s^{\varepsilon} (\mathcal{L}^{\bar{X}, \mathcal{L}_{\bar{X}}^{\mathbb{P}}} F) \mathrm{d}s - \int_0^t \xi_s^{\varepsilon} (F\bar{h}^i) \mathrm{d}Y_s^{\varepsilon, i} \right|^2 \\ \leqslant C\varepsilon^2 + C(T + \delta_{\varepsilon})^{\frac{1}{2}} \left[(\delta_{\varepsilon}^4 + \delta_{\varepsilon}^2) + \delta_{\varepsilon}^2 + \left((\frac{\delta_{\varepsilon}^{15}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^7}{\varepsilon^8}) (\delta_{\varepsilon}^{17} + \delta_{\varepsilon}^9) e^{C(\frac{\delta_{\varepsilon}^{16}}{\varepsilon^{16}} + \frac{\delta_{\varepsilon}^8}{\varepsilon^8})} \right)^{\frac{1}{4}} + \frac{\varepsilon^{\frac{1}{2}}}{\delta_{\varepsilon}^{\frac{1}{2}}} \right]^{\frac{1}{2}}.$$

Choosing $\delta_{\varepsilon} = \varepsilon (-ln\varepsilon)^{\frac{1}{32}}$ and taking the limit on both sides of the above inequality, it holds that

$$\lim_{\varepsilon \to 0} \mathbb{E} \left| \xi_t^{\varepsilon}(F) - \int_0^t \xi_s^{\varepsilon}(\mathcal{L}^{\bar{X}, \mathscr{L}_{\bar{X}}^{\mathbb{P}}} F) \mathrm{d}s - \int_0^t \xi_s^{\varepsilon}(F\bar{h}^i) \mathrm{d}Y_s^{\varepsilon, i} \right|^2 = 0.$$

Step 2. We prove that there exists a subsequence $\{\xi^{\varepsilon_k}\}$ which converges weakly to 0. By Lemma 5.8 and Step 1, we know that there exists a subsequence $\{\xi^{\varepsilon_k}\}$ which converges weakly to ξ in $C([0,T],\mathcal{M}(\mathbb{R}^n\times\mathscr{P}_2(\mathbb{R}^n))$ as $k\to\infty$, and furthermore for $t\in[0,T]$ and $F\in C_b^{4,4}(\mathbb{R}^n\times\mathscr{P}_2(\mathbb{R}^n))$, $\xi_t(F)$ satisfies the equation

$$\xi_t(F) - \int_0^t \xi_s(\mathcal{L}^{\bar{X},\mathcal{L}_{\bar{X}}^{\mathbb{P}}} F) \mathrm{d}s - \int_0^t \xi_s(F\bar{h}^i) \mathrm{d}Y_s = 0, \tag{45}$$

where Y is a l-dimensional Brownian motion. For Eq.(45), by [14, Theorem 4.9], it holds that its solutions are unique. Besides, note that 0 is a solution to Eq.(45). Thus, $\xi = 0$, that is, $\{\xi^{\varepsilon_k}\}$ converges weakly to 0. The proof is complete.

Now, it is the position to prove Theorem 3.2.

Proof of Theorem 3.2. For $t \in [0,T]$ and $F \in C_b^{4,4}(\mathbb{R}^n \times \mathscr{P}_2(\mathbb{R}^n))$, it holds that

$$\pi_t^{\varepsilon,x,\mu}(F) - \bar{\pi}_t(F) = \frac{\rho_t^{\varepsilon,x,\mu}(F) - \bar{\rho}_t(F)}{\bar{\rho}_t(1)} - \pi_t^{\varepsilon,x,\mu}(F) \frac{\rho_t^{\varepsilon,x,\mu}(1) - \bar{\rho}_t(1)}{\bar{\rho}_t(1)} = \frac{\xi_t^{\varepsilon}(F)}{\bar{\rho}_t(1)} - \pi_t^{\varepsilon,x,\mu}(F) \frac{\xi_t^{\varepsilon}(1)}{\bar{\rho}_t(1)}.$$

So, based on Lemma 5.6 and 5.10, and the boundedness of $\pi_t^{\varepsilon,x,\mu}(F)$, we obtain that $\pi_t^{\varepsilon,x,\mu}$ converges weakly to $\bar{\pi}_t$ as $\varepsilon \to 0$. The proof is complete.

6. Appendix

In the section, we prove (39).

Proof of (39).

By the Hölder inequality, the BDG inequality, $(\mathbf{H}_{b_2,\sigma_2}^1)$ and Lemma 5.9, we get that for $s \in [t, t + \delta_{\varepsilon}]$

$$\begin{split} & \mathbb{E}[|Z_{\varepsilon}^{s,\delta_{x},z}-\hat{Z}_{\varepsilon}^{s,\delta_{x},z}|^{16}] \\ & = \mathbb{E}\left[\left|\frac{1}{\varepsilon}\int_{t}^{s}\left(b_{2}(\mathcal{L}_{X_{\varepsilon}^{s,\delta_{x},z}}^{p},Z_{\varepsilon}^{s,\delta_{x},z}) - b_{2}(\delta_{x},\hat{Z}_{r}^{s,\delta_{x},z})\right)\mathrm{d}r \right. \\ & + \frac{1}{\sqrt{\varepsilon}}\int_{t}^{s}\left(\sigma_{2}(\mathcal{L}_{X_{\varepsilon}^{s,\delta_{x},z}}^{p},Z_{\varepsilon}^{s,\delta_{x},z}) - b_{2}(\delta_{x},\hat{Z}_{r}^{s,\delta_{x},z})\right)\mathrm{d}V \\ & + \frac{1}{\sqrt{\varepsilon}}\int_{t}^{s}\left(\sigma_{2}(\mathcal{L}_{X_{\varepsilon}^{s,x,\delta_{x},z}}^{p},Z_{\varepsilon}^{s,\delta_{x},z}) - \sigma_{2}(\delta_{x},\hat{Z}_{r}^{s,\delta_{x},z})\right)\mathrm{d}V \Big|^{16} \right] \\ & \leq \frac{2^{15}}{\varepsilon^{16}}\mathbb{E}\left[\left|\int_{t}^{s}\left(\sigma_{2}(\mathcal{L}_{X_{\varepsilon}^{s,x,\delta_{x},z}}^{p},Z_{r}^{s,\delta_{x},z}) - b_{2}(\delta_{x},\hat{Z}_{r}^{s,\delta_{x},z})\right)\mathrm{d}V \Big|^{16} \right] \\ & \leq C_{\varepsilon}^{\frac{15}{\varepsilon^{16}}}\int_{s}^{s}\mathbb{E}\left|b_{2}(\mathcal{L}_{X_{\varepsilon}^{p,x,\delta_{x},z}}^{p},Z_{r}^{s,\delta_{x},z}) - \sigma_{2}(\delta_{x},\hat{Z}_{r}^{s,\delta_{x},z})\right|^{16}\mathrm{d}r \\ & + C_{\varepsilon}^{\frac{5}{\varepsilon^{16}}}\int_{t}^{s}\mathbb{E}\left|b_{2}(\mathcal{L}_{X_{\varepsilon}^{p,x,\delta_{x},z}}^{p},Z_{r}^{s,\delta_{x},z}) - b_{2}(\delta_{x},\hat{Z}_{r}^{s,\delta_{x},z})\right|^{16}\mathrm{d}r \\ & = C_{\varepsilon}^{\frac{5}{\varepsilon^{16}}}\int_{t}^{s}\mathbb{E}\left|b_{2}(\mathcal{L}_{X_{\varepsilon}^{p,x,\delta_{x},z}}^{p},Z_{r}^{s,\delta_{x},z}) - b_{2}(\delta_{x},Z_{r}^{s,\delta_{x},z}) + b_{2}(\delta_{x},Z_{r}^{s,\delta_{x},z}) - b_{2}(\delta_{x},\hat{Z}_{r}^{s,\delta_{x},z})\right|^{16}\mathrm{d}r \\ & + C_{\varepsilon}^{\frac{5}{\varepsilon^{16}}}\int_{t}^{s}\mathbb{E}\left|b_{2}(\mathcal{L}_{X_{\varepsilon}^{p,x,\delta_{x},z}}^{p},Z_{r}^{s,\delta_{x},z}) - b_{2}(\delta_{x},Z_{r}^{s,\delta_{x},z}) + \sigma_{2}(\delta_{x},Z_{r}^{s,\delta_{x},z}) - \sigma_{2}(\delta_{x},\hat{Z}_{r}^{s,\delta_{x},z})\right|^{16}\mathrm{d}r \\ & + C_{\varepsilon}^{\frac{5}{\varepsilon^{16}}}\int_{t}^{s}\mathbb{E}\left|b_{2}(\mathcal{L}_{X_{\varepsilon}^{p,x,\delta_{x},z}}^{p},Z_{r}^{s,\delta_{x},z}) - b_{2}(\delta_{x},Z_{r}^{s,\delta_{x},z})\right|^{16}\mathrm{d}r \\ & + C_{\varepsilon}^{\frac{5}{\varepsilon^{16}}}\int_{t}^{s}\mathbb{E}\left|b_{2}(\mathcal{L}_{X_{\varepsilon}^{p,x,\delta_{x},z}}^{p},Z_{r}^{s,\delta_{x},z}) - b_{2}(\delta_{x},Z_{r}^{s,\delta_{x},z})\right|^{16}\mathrm{d}r \\ & + C_{\varepsilon}^{\frac{5}{\varepsilon^{16}}}\int_{t}^{s}\mathbb{E}\left|b_{2}(\sigma_{x},Z_{r}^{s,\delta_{x},z},Z_{r}^{s,\delta_{x},z}) - \sigma_{2}(\delta_{x},Z_{r}^{s,\delta_{x},z})\right|^{16}\mathrm{d}r \\ & + C_{\varepsilon}^{\frac{5}{\varepsilon^{16}}}\int_{t}^{s}\mathbb{E}\left|b_{2}(\sigma_{x},Z_{r}^{s,\delta_{x},z},Z_{r}^{s,\delta_{x},z}) - \sigma_{2}(\delta_{x},Z_{r}^{s,\delta_{x},z})\right|^{16}\mathrm{d}r \\ & + C_{\varepsilon}^{\frac{5}{\varepsilon^{16}}}\int_{t}^{s}\mathbb{E}\left|\delta_{x}(z,Z_{r}^{s,s},z_{x},Z_{r}^{s,s},z_{x},z_{x},z_{x},z_{x},z_{x},z_{x},z_{x},z_{x},z_{x},z_{x},z_{x},z_{x},z_{x},z_{x},z_{x},z_{x},z$$

Then the Gronwall inequality gives us

$$\mathbb{E}[|Z_s^{\varepsilon,\delta_x,z} - \hat{Z}_s^{\varepsilon,\delta_x,z}|^{16}] \leqslant C(1+|x|^{16}+|z|^{16})(\frac{\delta_\varepsilon^{15}}{\varepsilon^{16}} + \frac{\delta_\varepsilon^7}{\varepsilon^8})(\delta_\varepsilon^{17} + \delta_\varepsilon^9)e^{C(\frac{\delta_\varepsilon^{16}}{\varepsilon^{16}} + \frac{\delta_\varepsilon^8}{\varepsilon^8})}.$$

References

- [1] R. Beeson, N. Namachchivaya and N. Perkowski: Approximation of the filter equation for multiple timescale, correlated, nonlinear systems, SIAM J. Math. Anal., 54(2022)3054-3090.
- [2] P. Cardaliaguet: Notes on mean field games (from P.L. Lion's lectures at College de France). https://www.ceremade.dauphine.fr/cardalia/MFG100629.pdf, 2013.
- [3] X. J. Ding and H. J. Qiao: Euler-Maruyama approximations for stochastic McKean-Vlasov equations with non-Lipschitz coefficients, *Journal of Theoretical Probability*, 34(2021)1408-1425.
- [4] X. J. Ding and H. J. Qiao: Stability for stochastic McKean-Vlasov equations with non-Lipschitz coefficients, SIAM J. Control Optim., 59(2021)887-905.
- [5] E. Goggin: Convergence in distribution of conditional expectations, Ann. Probab., 22(1994)1097-1114.
- [6] E. Goggin: Convergence of filters with applications to the Kalman-Bucy case, IEEE Trans. Inform. Theory., 38(1992)1091-1100.
- [7] Z. Huang: Basis of Stochastic Analysis (in Chinese). Second ed., Science Press, Beijing, 2001.
- [8] P. Imkeller, N. S. Namachchivaya, N. Perkowski and H. C. Yeong: Dimensional reduction in nonlinear filtering: a homogenization approach, *The Annals of Applied Probability*, 23(2013), 2290-2326.
- [9] R.Z. Khasminskii: On an averaging principle for Itô stochastic differential equations, Kibernetica, 4(1968)260-279.
- [10] T. G. Kurtz: Approximation of population processes. In: Vol. 36 of CBMS-NSF Regional Conf. Series in Appl. Math. Philadelphia: SIAM, 1981.
- [11] H. J. Kushner: Weak Convergence Methods and Singularly Perturbed Stochastic Control and Filtering Problems, Systems & Control: Foundations & Applications 3. Birkhäuser, Boston, 1990.
- [12] J. Li and H. Min: Weak solutions of mean-field stochastic differential equations, Stochastic Analysis and Applications, 35(2017)542-568.
- [13] Y. Li, F. Wu and L. Xie: Poisson equation on Wasserstein space and diffusion approximations for McKean-Vlasov equation, http://arxiv.org/abs/2203.12796.
- [14] M. Q. Liu and H. J. Qiao: Uniqueness and superposition of the space-distribution dependent Zakai equations, http://arxiv.org/abs/2008.01596.
- [15] W. Liu, M. Röckner, X. B. Sun, Y. C. Xie: Averaging principle for slow-fast stochastic differential equations with time dependent locally Lipschitz coefficients, *Journal of Differential Equations*, 268(2020)2910-2948.
- [16] D. Liu: Strong convergence of principle of averaging for multiscale stochastic dynamical systems, Commun. Math. Sci., 8(2010)999-1020.
- [17] V. M. Lucic and A. J. Heunis: Convergence of nonlinear filters for randomlyperturbed dynamical systems, *Appl. Math. Optim.*, 48 (2003)93-128.
- [18] E. Pardoux and A. Y. Veretennikov: On poisson equation and diffusion approximation 1, Ann. Probab., 29(2001)1061-1085.
- [19] E. Pardoux and A. Y. Veretennikov: On poisson equation and diffusion approximation 2, Ann. Probab., 31(2003)1166-1192.
- [20] E. Pardoux and A. Y. Veretennikov: On poisson equation and diffusion approximation 3, Ann. Probab., 33(2005)1111-1133.
- [21] H. J. Qiao: Convergence of nonlinear filterings for stochastic dynamical systems with Lévy noises, Stochastic Analysis and Applications, 40(2022)352-378.
- [22] H. J. Qiao and W. L. Wei: Efficient filtering for multiscale McKean-Vlasov Stochastic differential equations, http://arxiv.org/abs/2206.05037.
- [23] H. J. Qiao, Y. J. Zhang and J. Q. Duan: Effective filtering on a random slow manifold, Nonlinearity, 31(2018)4649-4666.
- [24] P. Ren and F.-Y. Wang: Space-distribution PDEs for path independent additive functionals of McKean-Vlasov SDEs, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 23(2020)2050018.

- [25] M. Röckner, X. B. Sun and Y. C. Xie: Strong convergence order for slow-fast McKean-Vlasov stochastic differential equations, *Annales de l'I.H.P.Probabilités et statistiques*, 57(2021)547-576.
- [26] A. S. Sznitman: Topics in propagation of chaos, École d'Été de Probabilitiés de Saint-Flour XIX-1989, Lecture Notes in Mathematics Vol. 1464. Berlin, Springer, 1991.
- [27] N. Sen and P. E. Caines: Nonlinear filtering theory for McKean-Vlasov type stochastic differential equations, SIAM J. Control Optim., 54(2016)153-174.
- [28] F.-Y. Wang: Distribution dependent SDEs for Landau type equations, *Stochastic Process. Appl.*, 128(2018)595-621.
- [29] J. Xu, J. F. Liu, J. C. Liu and Y. Miao: Strong averaging principle for two-time-scale stochastic McKean-Vlasov equations, *Applied Mathematics and Optimization*, 84(2021)837-867.