

Tetrahedron of flavors: One Three to rule them all

Aharon Davidson*

Physics Department, Ben-Gurion University of the Negev, Beer-Sheva 84105, Israel

(Dated: October 10, 2022)

Under the conception that the total number three of fermion families must have the one and the same gauge theoretical origin as all other threes which accompany the single family grand unifiable group structure, we trade the trinification $SU(3)$ symmetry building block for its semi-simple Vertical-Horizontal symmetric $SU(3)_V \otimes SU(3)_H$ extension. The anomaly free flavor-chiral fermion representation is then constructed, solely out of threes and without any superfluous replication, by treating each standard V -triplet (and anti triplet) as an H -singlet, and conversely, by letting each standard V -singlet transform as an H -triplet (or anti triplet). The model can be schematically described by a tetrahedron of flavors. In its bi-trinification phase, the model exhibits two coupled trinification cycles, sharing the same color group, and is furthermore accompanied by a built-in dark sector. In its isomorphic quartification phase, the model presents a novel non-Pati-Salam version of quark/lepton correspondence, where quarks are paired with anti-leptons to cancel horizontal anomalies, and all fermion masses stem from one and the same Yukawa source.

INTRODUCTION

Decoding the mass spectrum of quarks and leptons is without any doubt the holy grail of theoretical particle physics. While the mass generating Higgs mechanism, mediated by Yukawa couplings, is rooted in the heart of the standard electro/nuclear theory, we still do not have the slightest idea what physics actually determines the eigenmasses and mixings. At this stage, even an empirical formula to account for the observed mass hierarchy is certainly welcome. The structure of the Fermi mass matrix is however just the tip of the flavor puzzle iceberg.

The various fermionic pieces which make a single family (or a generation) are well organized, up to some electro/nuclear neutral members, within the framework of the by now standard $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$ gauge theory. Anomaly cancellation is then a necessary self consistency condition, but obviously not a sufficient one for shedding light on the full picture. The consequent grand unification procedure to combine the otherwise separated fermionic pieces into a single representation has been exclusively realized for the $SU(5) \subset SO(10) \subset E(6)$ group sequence, and is most welcome, but leaves a fundamental question unanswered. Namely, why can/must grand unification be realized (at least partially) already at the single family level, while apparently leaving the observed multiplicity of fermion families field/group theoretically out of the game.

At the time, embedding the single family GUT within a larger multi family GUT seemed to be a logical step forwards. Unfortunately, such a potentially promising direction has been practically exhausted, and so far failed to deliver. On the short list of theoretical (on top of experimental) obstacles encountered one can find:

- (i) The magic yet theoretically challenging number of exactly *three* standard fermion families.
- (ii) The restrictive chiral flavor structure required at the

low energy group theoretical reduction level.

(iii) The ever growing sizes (counting generators) of the candidate GUT groups involved.

(iv) Heavy field theoretical artillery, such as supersymmetry and dimensional reduction, while being attractive from various other reasons, has not contributed so far its part to the flavor puzzle. The more so superstring theory, including its string phenomenology outer branch, has not been proven too useful either.

It may well be that the tempting strategy of passing through the single family grand unification stage may have counter intuitively blocked us from unveiling the full flavor picture.

VERTICAL \leftrightarrow HORIZONTAL SYMMETRY

Equipped with no compelling answers, the flavor puzzle has sourced plenty of imaginative ideas, ranging from atomic style isotopes all the way to compositeness, involving a variety of field theoretical techniques. In this paper, however, with the focus solely on group theory, we humbly follow the hypothesis that the total number *three* of fermion families shares the one and the same gauge theoretical origin as all other *threes* which govern the single family unifiable (not grand unified) group structure. Adopting this line of thought, let our starting point be trinification [1, 2], based on the semi-simple gauge group

$$G = SU(3)_C \otimes SU(3)_L \otimes SU(3)_R . \quad (1)$$

An accompanying C_3 discrete symmetry is optional, and becomes mandatory if a common gauge coupling constant is in order. Associated with is the familiar anomaly free

flavor chiral representation of left handed fermions

$$\psi_L = \begin{array}{|c|c|c|c|} \hline q & 3 & 3^* & 1 \\ \hline q^c & 3^* & 1 & 3 \\ \hline \ell + \ell^c & 1 & 3 & 3^* \\ \hline \end{array} \quad (2)$$

which can always be supplemented, if so required, by a bunch of dark G -singlets

$$\chi = \begin{array}{|c|c|c|c|} \hline \chi & 1 & 1 & 1 \\ \hline \end{array} \quad (3)$$

Schematically, the model can be neatly represented by a triangle, as depicted in Fig.1, where the vertices represent fermions and the edges stand for the various $SU(3)$ group factors involved.




FIG. 1: The trinification model: The vertices are associated with the various fermions involved., and the three edges stand for the three $SU(3)$ group factors involved. An incoming/outgoing arrow signals a triplet/anti-triplet of the respective $SU(3)$. Schematically, blue and red circles represent triplets and anti-triplets respectively

Already at the trinification level, prior to $E(6)$ unification [3], there is a heavy price (not everyone is willing) to pay, namely extending the standard single fermion family, composed of 16 members, to include extra $27-16=11$ non-standard members. The latter supplement, easily classified by means of the electric charge operator

$$Q = T_{3L} + T_{3R} + \frac{1}{2}(Y_L + Y_R) , \quad (4)$$

decouples however once $SU(3)_L \otimes SU(3)_R$ parent symmetry breaks down to its left-right symmetric sub-group $SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$. Furthermore, notice that the full quark/lepton correspondence, i.e. treating lepton number as the forth color, [4] is not restored until (and if) $E(6)$ embedding is realized.

Multiplying the trinification group G by extra $SU(3)$ factors is then quite a conservative step forwards. $SU(3)^N$ models, with $N > 3$, mostly single family models, have been extensively studied by Ma [5] and by others [6]. Horizontal symmetries [7], primarily invoked to classify the otherwise electro/nuclear degenerate fermionic families, start from minimal $U(1)$, local or else global (incorporating the Peccei-Quinn mechanism [8]) and in

particular include an $SU(3)$ example [9]. Here, we revive the idea of Vertical \leftrightarrow Horizontal symmetry [10], and trade the single family group, now referred to as G_V , for its $G_V \otimes G_H$ semi-simple extension. Note that models incorporating semi-simple group structure, for example Pati-Salam's $SU(4)^4$ [4], first ever unification scheme (originally designed for two families), the hybrid single family left-right symmetric $SU(5)_L \otimes SU(5)_R$ model [11] hosting chiral color [12], and multi family models of the $SO(10) \otimes SO(10)$ type [13], have already been discussed in the literature.

To construct the tenable $G_V \otimes G_H$ fermion representation, we adopt the following prescription:

Treat each standard V -triplet (and anti-triplet) as an H -singlet, and conversely, let each standard V -singlet transform as an H -triplet (or anti-triplet).

This prescription generalizes Eq.(2) into

$$\psi_L = \begin{array}{|c|c|c|c|c|c|} \hline & \text{V} & & \text{H} & & \\ \hline 3 & 3^* & 1 & 1 & 1 & x \\ \hline 3^* & 1 & 3 & 1 & y & 1 \\ \hline 1 & 3 & 3^* & z & 1 & 1 \\ \hline 1 & 1 & 1 & z^* & y^* & x^* \\ \hline \end{array} \quad (5)$$

The cancelation of ABJ anomalies is done pairwise, while maintaining overall flavor chirality, so that each of the $SU(3)$ representations x, y, z can be either a 3 or else a 3^* . While the particular choice seems irrelevant from any individual $SU(3)$ point of view, a matter of definition, it does become relevant in the presence of an accompanying discrete symmetry. Altogether, with only $SU(3)$ triplets and anti-triplets at our disposal, only two independent configurations exist.

The manifestly symmetric configuration

$$x = y = z \quad , \quad (6)$$

could have been our naive preference. But unfortunately, as demonstrated in Fig.(2), its associated discrete symmetry does not really go much beyond the original trinity model, offering the same three ways to identify $\{q, q^c, \ell + \ell^c\}$, and this without involving $V \leftrightarrow H$ interplay, and without reviving quark/lepton correspondence.




FIG. 2: With blue and red circles representing triplets and anti-triplets respectively, the three trinification ways to identify $\{q, q^c, \ell + \ell^c\}$ are illustrated by $\{a, b, c\}$ cyclic permutations accompanied by suitable re-arrangements of the six $SU(3)$ group factors.

TETRAHEDRON OF FLAVORS

In this paper, however, from reasons to be specified, we construct the so-called bi-trinification model (along with its isomorphic quartification phase) based on the alternative configuration

$$x = y = z^* \quad \text{up to permutations.} \quad (7)$$

It offers six variant ways to identify $\{q, q^c, \ell + \ell^c\}$, each of which comes with its own (practically equivalent) permuted and/or conjugate horizontal assignments. To fully appreciate this point, as demonstrated in Fig.3, one may rearrange rows as well as columns, identify an alternative trinification V-sector (three left columns), and then verify how one H-sector (three right columns) gets systematically replaced by an equivalent one, such that $(x, y, z) \rightarrow (x', y', z')$, where the latter is a permuted version of the former, with or without conjugation.




FIG. 3: Horizontal bi-trinification: By rearranging rows (fermion assignments) and columns ($SU(3)$ group factors), while recasting the exact structure of the trinification V-sector (marked with lines), one can easily switch from one H-sector (three right columns) to another. The six variants are thus equivalent to each other. Note that triplets and anti-triplets are represented here by blue and red circles, respectively.

The fermion representation furnishes a tetrahedron. While the edges, numbered $k = 1, 2, \dots, 6$, correspond to the individual $SU(3)_k$ group factors, the vertices are associated with the four fermion classes $q, q^c, \ell + \ell^c, \chi$. The various arrows involved, making the edges directed, stand for triplets (outgoing arrows) and anti-triplets (ingoing arrows). Notice that there are in fact two different ways to consistently identify the V-sector. Sticking to Fig.(3), it is either the 231-cycle (three upper configurations) or alternatively the 246-cycle (three lower configurations). It is this characteristic feature which justifies using the terminology bi-trinification. The two cycles correspond to the trinifications $SU(3)_C \otimes [SU(3)_L \otimes SU(3)_R]_{V,H}$, respectively. This in turn gives rise to two electric charges

$$Q_{V,H} = \left[T_{3L} + T_{3R} + \frac{1}{2} Y_{L+R} \right]_{V,H}, \quad (8)$$




FIG. 4: The Tetrahedron model (Bi-Trinification phases): The two coupled 3-cycles (red arrows), namely 231 (left plot) and 246 (right plot), describe two different trinification V-sectors. Attached to each one of them is a corresponding H-sector (blue arrows). In this phase, ℓ and ℓ^c conventionally share a common leptonic vertex, and associated with the fourth vertex χ is a novel dark (= electro/nuclear neutral) sector.

one of which, that is Q_H , must be spontaneously broken, thereby opening the door for the Holdom effect [14], also known as photon mixing.

The χ -fermions are required on anomaly cancelation grounds. They are V-sector singlets by construction, and as such, have no standard electro/nuclear interactions. This makes them, by definition, candidates for dark matter particles of the WIMP kind. They do interact with ordinary matter though, with the H -sector serving as the tenable portal. The mechanism involved, as demonstrated in Fig.(5) for the special lepto/dark case, is the exchange of super heavy horizontal gauge bosons.




FIG. 5: Dark matter portal: Dark matter particles (V-sector singlets) interact with standard model particles (V-sector non singlets) by exchanging heavy horizontal gauge bosons. Depicted in this figure is the lepto/dark case.

BI-TRI/QUART ISOMORPHISM

Alternative rearrangements of rows and columns can make the single family group flavor assignments depart from their built-in trinification (and thus bi-trinification) construction, and at the expense of a smaller H-sector, extend the V-sector into a quartification phase. Depicted in Fig.(6) are the four associated quartification

variant configurations. Indeed, they share a common extended V-sector (left four columns connected with lines) and admit shrank (only two, rather than three, right columns) horizontal structures. They are equivalent to each other. And most importantly, the existence of these four variants serves to establish that, as far as our model is concerned, bi-trinification and quartification are two faces of the same coin.




FIG. 6: Horizontal quartification: Alternative rearrangements of particle assignments (rows) and $SU(3)$ group factors (columns) depart from trinification, and extend the V-sector (marked with lines) into quartification at the expense of a smaller (two right columns) H-sector. The equivalence of these four variants to each other is interpreted as a novel version of quark/lepton correspondence. As before, triplets and anti-triplets are represented by blue and red circles, respectively.

Back to the tetrahedron of flavors, the quartification phase calls for some reshuffling of fermion assignments. To be specific, the leptonic sub-representation bifurcates, such that ℓ and ℓ^c belong now to two different vertices. To classify the fermions under the extended single family group $SU(3)^4$, with the individual subscripts being C, L, R, N (with N denoting the new comer), we introduce the fully symmetric electric charge formula

$$Q = T_{3L} + T_{3R} + T_{3N} + \frac{1}{2}(Y_L + Y_R + Y_N) . \quad (9)$$

Apart from the manifest LRN symmetry, it has the advantage that all individual electric charges involved are standard. The electric charge Q , which generalizes Eq.(4), should be contrasted with Eq.(8), and with a different electric charge formula recently advocated for a similar $SU(3)^4$ model. To be more explicit, and expose the extra nine leptons per family, introduced on top of the trinification scheme, we piecewise specify the $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ content of the

fermion representation, namely

$$\begin{aligned} (3, 3^*, 1, 1) &= (3, 2, 1)_{\frac{1}{3}} + (3, 1, 1)_{-\frac{2}{3}} , \\ (3^*, 1, 3, 1) &= (3^*, 1, 2)_{-\frac{1}{3}} + (3^*, 1, 1)_{\frac{2}{3}} , \\ (1, 3, 1, 3^*) &= (1, 2, 1)_{-1,1} + (1, 2, 1)_{-1} + \\ &\quad + (1, 1, 1)_{0,2} + (1, 1, 1)_0 , \\ (1, 1, 3^*, 3) &= (1, 1, 2)_{-1,1} + (1, 1, 2)_1 + \\ &\quad + (1, 1, 1)_{-2,0} + (1, 1, 1)_0 . \end{aligned} \quad (10)$$

Of special interest are those vector-like fermions which carry non-standard $B - L$ charges. From a group theory point of view, they are primarily responsible for the small value

$$\sin^2 \theta_W \rightarrow \frac{\sum T_{L3}^2}{\sum Q^2} = \frac{1}{4} \quad (11)$$

of the Weinberg angle at the symmetry limit, which is smaller than the conventional value of $\frac{3}{8}$. However, the vector-like surplus of quarks and leptons is not protected by the $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ symmetry subgroup, and decouples at some presumably very heavy mass scale. Under the latter residual symmetry, we are left with the exact flavor chiral family of the Left-Right symmetric model, that is

$$\psi = (3, 2, 1)_{\frac{1}{3}} + (3^*, 1, 2)_{-\frac{1}{3}} + (1, 2, 1)_{-1} + (1, 1, 2)_1 . \quad (12)$$




FIG. 7: The Tetrahedron model (quartification phase): The focus now is on the 4-cycle 2451 (red arrows) which describes a single family $SU(3)^4$ fermionic representation. It accounts for the revival of a full $q, q^c \leftrightarrow \ell, \ell^c$ quark/lepton correspondence, as expressed by the four variants shown in Fig.(6). The blue arrows account for the threefold family structure under horizontal $SU(3)_I \otimes SU(3)_{II}$.

QUARK/LEPTON CORRESPONDENCE RESURRECTION

$B - L$ does not serve in our model as the forth color. The reason is obvious: There is no Pati-Salam $SU(4)$ to

the rescue. Quark/lepton correspondence is absent from the trinification model as well. Unlike q and q^c which are assigned to two different vertices, see Fig.(1), ℓ and ℓ^c share a common third vertex. It is only in Ma's $SU(3)^4$ model that quark/lepton correspondence has been revived, at least in the sense that ℓ and ℓ^c split vertices, see Fig.(7). But there is more to it.

The H-sector is primarily in charge of the threefold family replication. Denoting the quartification horizontal group by $SU(3)_I \otimes SU(3)_{II}$, and choosing for definiteness one of the four variants depicted in Fig.(6), we face the horizontal assignments

$$q \sim (1, 3), \quad q^c \sim (3^*, 1), \quad \ell \sim (3, 1), \quad \ell^c \sim (1, 3^*). \quad (13)$$

It is the pair $\{q^c, \ell\}$ which cancels the $SU(3)_I$ anomalies, while $\{q, \ell^c\}$ takes care of the $SU(3)_{II}$ anomalies. Such an unprecedented horizontal pairing, which seems as a unified generalization of Foot-Lew quark/lepton symmetry [15], is fully dictated by the group theoretical structure of our tetrahedron model, and gives a novel perspective to the inter relations between quarks and leptons.

Another attractive aspect of quartification has to do with the Higgs sector. The standard model has already taught us that quarks and leptons alike should acquire their masses via Yukawa couplings with the one and the same Higgs scalar. One complex doublet suffices to govern all masses (and may even be the only source of spontaneous symmetry breaking). Truly, this feature is forcefully shared by the original $SU(3)^3$ trinification model as well. Starting from the fermionic representation Eq.(2), the mass generating job is carried out by the scalar ϕ which exhibits the fermion bilinear quantum numbers

$$\phi \sim qq^c \sim \ell\ell^c \sim (1, 3^*, 3). \quad (14)$$

This is strikingly not the case for the tetrahedron model in the bi-trinification phase, for which

$$\begin{aligned} qq^c &\sim (1, 3^*, 3 || 1, 3, 3^*) \\ \ell\ell^c &\sim (1, 3^*, 3 || 3^*, 1, 1) \end{aligned} \quad (15)$$

The more so, once χ has entered the game, we cannot avoid having $(\ell + \ell^c)\chi \sim (1, 3, 3^* || 1, 3^*, 3)$ as well. This seems to suggest that ℓ and ℓ^c better not share a common tetrahedron vertex. Indeed, one is back on safe (and quite attractive) grounds provided the tetrahedron model in adopted in its the quartification phase.

Choosing for the sake of definiteness a particular horizontal variant, say

q	3	3^*	1	1	1	3
q^c	3^*	1	3	1	3^*	1
ℓ	1	3	1	3^*	3	1
ℓ^c	1	1	3^*	3	1	3^*

(16)

quartification naturally offers the conjugate fermion bilinear quantum numbers

$$\begin{aligned} qq^c &\sim \phi \sim (1, 3^*, 3, 1 || 3^*, 3) \\ \ell\ell^c &\sim \phi^* \sim (1, 3, 3^*, 1 || 3, 3^*) \end{aligned} \quad (17)$$

In turn, as dictated by their revived correspondence, quarks and leptons do share now a common Yukawa coupling origin. The associated VEV pattern needs not be simple, to say the least, reflecting the double spinorial structure (expressed by $i, j = 1, 2, 3$) of $\langle \phi \rangle = v_{L,R}^{ij}$ under $SU(3)_{I,II}$ of each of the two (LR symmetric) Weinberg-Salam doublets. To add a complication to the list, notice that generically the two VEV matrices v_L^{ij} and v_R^{ij} cannot be diagonalized simultaneously. This is apparently the reason underlying the fermion mixing phenomenon, but unfortunately not even a tentative mixing formula can be derived at this preliminary stage. It is highly suggestive that $SU(3)_I \otimes SU(3)_{II}$ gets broken at a certain stage down to $SU(3)_{I+II}$, under which ϕ transforms as a singlet, with all horizontal fine details ripped off.

EPILOGUE

In this paper, unfortunately and admittedly, we have offered no detailed insight into the structure of the Fermi mass matrix. While this is undoubtedly a drawback, it should be gracefully criticized, recalling that after forty years of flavor puzzle frustration, and after so many imaginative yet unripe theoretical ideas per year, no one can so far claim even the slightest of victories. The major clues, so we would like to believe, rowing against the conventional stream, are already with us. Namely, the challenging total number three of fermion families, and the striking failure of all grand unification trails (string theory and extra dimensions included) to account for it. The naive yet powerful idea that all threes which accompany flavor physics must have the one and the same gauge theoretical origin then simply paves the way for Vertical-Horizontal symmetric trinification. We can only hope that the resulting hereby presented tetrahedron of flavors, stemming from trinification but eventually focused on its quartification (rather than bi-trinification) phase, would shed some light on the quark/lepton correspondence and elegantly account for the threefold family replication, at least at the group theoretical level.

* Electronic address: davidson@bgu.ac.il; URL: <https://physics.bgu.ac.il/~davidson>

- [1] A. de Rujula, H. Georgi and S.L. Glashow, in *Fifth Workshop on Grand Unification*, edited by K. Kang, H. Fried and P. Frampton, World Scientific (1984).
- [2] K.S. Babu, X.G. He and S. Pakvasa, Phys. Rev. D33, 763 (1986); J.E. Kim, Phys. Lett. B591, 119 (2004); J.

- Sayre, S. Wiesenfeldt and S. Willenbrock, Phys. Rev. D73, 035013 (2006); J. Hetzel and B. Stech, Phys. Rev. D91, 055026 (2015); G.M. Pelaggi, A. Strumia and S. Vignali, JHEP 08, 130 (2015); K.S. Babu, B. Bajc, M. Nemevšek and Z. Tavartkiladze, AIP Conf. Proc. 1900, 020002 (2017); Z.W. Wang, A. Al Balushi, R. Mann and H.M. Jiang, Phys. Rev. D99, 115017 (2019); K.S. Babu, S. Jana and A. Thapa, JHEP 02, 051 (2022).
- [3] F. Gursev, P. Ramond and P. Sikivie, Phys. Lett. 60B, 177 (1976); Y. Achiman and B. Stech, Phys. Lett. 77B, 389 (1978); Q. Shafi, Phys. Lett. B79, 301 (1978); R. Barbieri and D.V. Nanopoulos, Phys. Lett. 91B, 369 (1980).
- [4] J.C. Pati and A. Salam, Phys. Rev. D10, 275 (1974).
- [5] E. Ma, Phys. Lett. B593, 198 (2004); E. Ma, Mod. Phys. Lett. A20, 1953 (2005); E. Ma, M. Mondragon and G. Zoupanos, JHEP 12, 026 (2005); Q.H. Cao, S.L. Chen and E. Ma, Phys. Rev. D73, 015009 (2006).
- [6] K.S. Babu, I. Thomas, W. Kephart and H. Pas, Phys. Rev. D77, 116006 (2008); T. Banks and J.F. Fortin, JHEP 07, 046 (2009); C. Kownacki, E. Ma, N. Pollard, O. Popov and M. Zakeri, Phys. Lett. B769, 267 (2017); *ibid* Nucl. Phys. B928, 520 (2018); *ibid* Phys. Lett. B777, 121 (2018); E. Ma, Phys. Lett. B820, 136561 (2021).
- [7] F. Wilczek and A. Zee, Phys. Rev. Lett. 42, 421 (1979); A. Davidson, Phys. Rev. Lett. 43, 92 (1979).
- [8] A. Davidson and K.C. Wali, Phys. Rev. Lett. 48, 11 (1982); F. Wilczek and A. Zee, Phys. Rev. Lett., 49:1549 (1982).
- [9] T. Yanagida, Phys. Rev. D20, 2986 (1979); Z.G. Berezhiani, Phys. Lett. B150, 177 (1985).
- [10] A. Davidson and K.C. Wali, Phys. Lett. B94, 359 (1980).
- [11] A. Davidson and K.C. Wali, Phys. Rev. Lett. 58, 2623 (1987).
- [12] P.H. Frampton and S.L. Glashow, Phys. Lett. B190, 157(1987); A. Davidson and K.C. Wali, Phys. Lett. B194, 247 (1987).
- [13] A. Davidson, P.D. Mannheim and K.C. Wali, Phys. Rev. Lett. 47, 149 (1981).
- [14] B. Holdom, Phys. Lett. 166B, 196 (1986).
- [15] R. Foot, and H. Lew, Phys. Rev. D41, 3502 (1990); R. Foot, H. Lew and R.R. Volkas, Phys. Rev. D44, 1531 (1991).