ON THE TAKAI DUALITY FOR L^p OPERATOR CROSSED PRODUCTS

ZHEN WANG AND SEN ZHU

ABSTRACT. The aim of this paper is to study a problem raised by N. C. Phillips concerning the existence of Takai duality for L^p operator crossed products $F^p(G,A,\alpha)$, where G is a locally compact Abelian group, A is an L^p operator algebra and α is an isometric action of G on A. Inspired by D. Williams' proof for the Takai duality theorem for crossed products of C^* -algebras, we construct a homomorphism Φ from $F^p(\hat{G}, F^p(G,A,\alpha), \hat{\alpha})$ to $\mathcal{K}(l^p(G)) \otimes_p A$ which is a natural L^p -analog of D. Williams' map. For countable discrete Abelian groups G and separable unital L^p operator algebras A which have unique L^p operator matrix norms, we show that Φ is an isomorphism if and only if either G is finite or p=2; in particular, Φ is an isometric isomorphism in the case that p=2. Moreover, it is proved that Φ is equivariant for the double dual action $\hat{\alpha}$ of G on $F^p(\hat{G},F^p(G,A,\alpha),\hat{\alpha})$ and the action $Ad\rho\otimes\alpha$ of G on $\mathcal{K}(l^p(G))\otimes_p A$.

1. Introduction

The Takai duality theorem is an important result concerning crossed products of C^* algebras. Recall that a C^* -dynamical system is a triple (A, G, α) consisting of a C^* -algebra A, a locally compact group G and a continuous homomorphism $\alpha: G \to \operatorname{Aut}(A)$, where $\operatorname{Aut}(A)$ is the group of automorphisms of A. Let $A \rtimes_{\alpha} G$ be the full crossed product of A by G. If G is an Abelian group with dual group \hat{G} , then there is a dual action $\hat{\alpha}$ of \hat{G} on $A \rtimes_{\alpha} G$ which is given by $\hat{\alpha}_{\gamma}: C_c(G,A) \to C_c(G,A), \ \hat{\alpha}_{\gamma}(f)(s) := \overline{\gamma(s)}f(s)$ for all $\gamma \in \hat{G}, s \in G$ and $f \in C_c(G, A)$, and $\hat{\alpha}_{\gamma}$ can extend to $A \rtimes_{\alpha} G$ by continuity. Therefore $(A \rtimes_{\alpha} G, \hat{G}, \hat{\alpha})$ is a C^* -dynamical system called the dual system. The Takai duality theorem [22] says that the iterated crossed product $(A \rtimes_{\alpha} G) \rtimes_{\hat{\alpha}} \hat{G}$ is *-isomorphic to $A \otimes \mathcal{K}(L^{2}(G))$, where $\mathcal{K}(L^{2}(G))$ denotes the algebra of compact operators on $L^2(G)$. Moreover, this isomorphism is also equivariant for the double dual action $\hat{\alpha}$ of G on $(A \rtimes_{\alpha} G) \rtimes_{\hat{\alpha}} \hat{G}$ and the action $\alpha \otimes \operatorname{Ad} \rho$ of G on $A \otimes \mathcal{K}(L^2(G))$. This result has many applications in the K-theory of operator algebras. For example, it can be used to prove Connes' Thom isomorphism theorem (see [1, Theorem 10.2.2] or [4]). In 1986, I. Raeburn [20] gave another proof of Takai's theorem by exploiting the universal properties of crossed products of C^* -algebras. D. Williams provided a variation of I. Raeburn's proof, with an extra contribution by S. Echterhoff (see [23, Theorem 7.1]).

The aim of this paper is to study a problem raised by N. C. Phillips [17, Problem 8.7] concerning the Takai duality for L^p operator crossed products. To proceed, we first introduce some notations and terminology. The reader is referred to [11] or [16] for more details.

Date: December 13, 2022.

²⁰¹⁰ Mathematics Subject Classification. Primary 22D35, 47L65; Secondary 43A25, 47L10.

 $Key\ words\ and\ phrases.\ L^p$ operator crossed products, Takai duality, Locally compact Abelian groups, L^p operator algebras.

The class of L^p operator algebras is a natural generalization of operator algebras on Hilbert spaces, and was first studied in Herz's influential paper [13] on harmonic analysis on L^p spaces. People's renewed interests in L^p operator algebras were inspired by N. C. Phillips [16], who showed that a few of the important constructions of C^* -algebras can be generalized to the setting of L^p operator algebras. Recently the study of L^p operator algebras has received much attention, and a number of authors have made significant contributions to it (see [2, 6, 7, 8, 9, 10, 11, 12, 14, 16]).

The L^p operator crossed products were introduced by N. C. Phillips [16] with the aim to compute the K-theory groups of L^p Cuntz algebras. Recall that a Banach algebra A is an L^p operator algebra if it can be isometrically represented on an L^p space $(p \in [1, \infty))$. Let $\pi: A \to \mathcal{B}(L^p(X, \nu))$ be a representation of A. We say that π is σ -finite if ν is σ -finite. Let G be a locally compact group. Then there exists a left Haar measure μ on G. An L^p operator algebra dynamical system is a triple (G, A, α) consisting of a locally compact group G, an L^p operator algebra A and a continuous homomorphism $\alpha: G \to \operatorname{Aut}(A)$, where $\operatorname{Aut}(A)$ is the group of isometric automorphisms of A. A contractive covariant representation of (G, A, α) on an L^p space E is a pair (π, ν) consisting of a nondegenerate contractive homomorphism $\pi: A \to \mathcal{B}(E)$ and an isometric group representation $\nu: G \to \mathcal{B}(E)$, satisfying the covariance condition

$$v_t \pi(a) v_{t-1} = \pi(\alpha_t(a))$$

for all $t \in G$ and $a \in A$. A covariant representation (π, v) of (G, A, α) is σ -finite if π is σ -finite.

Denote by $C_c(G, A, \alpha)$ the vector space of continuous compactly supported functions $G \to A$, made into an algebra over \mathbb{C} with product given by twisted convolution, that is,

$$(f * g)(t) := \int_G f(s)\alpha_s(g(s^{-1}t))d\mu(s)$$

for all $f, g \in C_c(G, A, \alpha)$ and $t \in G$. The integrated form of (π, v) is the nondegenerate contractive homomorphism $\pi \rtimes v : C_c(G, A, \alpha) \to \mathcal{B}(E)$ given by

$$(\pi \rtimes v)(f)(\xi) := \int_{G} \pi(f(t))v_{t}(\xi)d\mu(t)$$

for all $f \in C_c(G, A, \alpha)$ and for all $\xi \in E$. Denote by $\operatorname{Rep}_p(G, A, \alpha)$ the class of all nondegenerate σ -finite contractive covariant representations of (G, A, α) on L^p -spaces. The full L^p operator crossed product $F^p(G, A, \alpha)$ is defined as the completion of $C_c(G, A, \alpha)$ in the norm

$$||f||_{F^p(G,A,\alpha)} := \sup \{||(\pi \rtimes v)(f)|| : (\pi,v) \in \operatorname{Rep}_p(G,A,\alpha)\}.$$

Let (G, A, α) be an L^p operator algebra dynamical system. Given a nondegenerate σ finite contractive representation $\pi_0 : A \to \mathcal{B}(E_0)$ on an L^p space E_0 , its associated regular
covariant representation is the pair $(\pi, \lambda_p^{E_0})$ on $L^p(G) \otimes_p E_0 \cong L^p(G, E_0)$ given by

$$\pi(a)(\xi)(s) := \pi_0(\alpha_{s^{-1}}(a))(\xi(s))$$

and

$$\lambda_p^{E_0}(s)(\xi)(t) := \xi(s^{-1}t)$$

for all $a \in A$, $\xi \in L^p(G, E_0)$, and $s, t \in G$. We denote by $\operatorname{RegRep}_p(G, A, \alpha)$ the class consisting of nondegenerate σ -finite contractive regular covariant representations of (G, A, α) , which

is clearly a subclass of $\operatorname{Rep}_p(G, A, \alpha)$. The reduced L^p operator crossed product $F_{\lambda}^p(G, A, \alpha)$ is defined as the completion of $C_c(G, A, \alpha)$ in the norm

$$||f||_{F^p_{\lambda}(G,A,\alpha)} := \sup\{||(\pi \rtimes v)(f)|| : (\pi,v) \in \operatorname{RegRep}_p(G,A,\alpha)\}.$$

By [18, Theorem 7.1], if G is amenable, then the identity map on $C_c(G, A, \alpha)$ can be extended to an isometric isomorphism from $F^p(G, A, \alpha)$ onto $F^p_{\lambda}(G, A, \alpha)$. If $A = \mathbb{C}$, then it is easy to see that $F^p(G, A, \mathrm{id})$ is the full group L^p operator algebra $F^p(G)$ and $F^p_{\lambda}(G, A, \mathrm{id})$ is the reduced group L^p operator algebra $F^p_{\lambda}(G)$, where id is the trivial action of G on \mathbb{C} .

If, in addition, G is Abelian, then we let \hat{G} denote the dual group of G. For each $\gamma \in \hat{G}$, N. C. Phillips defined an isomorphism $\hat{\alpha}_{\gamma} : C_c(G, A, \alpha) \to C_c(G, A, \alpha)$ by $\hat{\alpha}_{\gamma}(f)(s) := \overline{\gamma(s)}f(s)$ for all $f \in C_c(G, A, \alpha)$ and $s \in G$ (see [16, Definition 3.15]). Thus $\hat{\alpha}_{\gamma}$ can extend to an isometry on $F^p(G, A, \alpha)$ by continuity (see [16, Theorem 3.18]). Hence there is a dual system $(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$. So we obtain the iterated L^p operator crossed product $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$.

In [17, Problem 8.7], N. C. Phillips raised the following natural problem concerning the existence of the Takai duality theorem for L^p operator crossed products.

Problem 1.1. Let $p \in [1, \infty)$, A be an L^p operator algebra, G be a locally compact Abelian group, and $\alpha : G \to \operatorname{Aut}(A)$ be an isometric action of G on A. Then is $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$ isomorphic to $\mathcal{K}(L^p(G)) \otimes_p A$? Here $\mathcal{K}(L^p(G))$ denotes the algebra of compact operators on $L^p(G)$ and \otimes_p denotes the spatial L^p operator tensor product of L^p operator algebras.

The reader is referred to [16] for the details of spatial L^p operator tensor products.

The aim of this paper is to consider Problem 1.1 following D. Williams' proof for the Takai duality theorem. We first briefly recall D. Williams' approach (see [23, Theorem 7.1]). Suppose that G is a locally compact Abelian group and (A, G, α) is a C^* -dynamical system. D. Williams constructed a map Ψ from $(A \rtimes_{\alpha} G) \rtimes_{\hat{\alpha}} \hat{G}$ to $\mathcal{K}(L^2(G)) \otimes A$ as a composition of four *-homomorphisms

$$(A \rtimes_{\alpha} G) \rtimes_{\hat{\alpha}} \hat{G} \xrightarrow{\Psi_{1}} (A \rtimes_{\operatorname{id}} \hat{G}) \rtimes_{\widehat{\operatorname{id}}^{-1} \otimes \alpha} G \xrightarrow{\Psi_{2}} C_{0}(G, A) \rtimes_{\operatorname{lt} \otimes \alpha} G$$

$$\downarrow_{\Psi_{3}} , \qquad (1)$$

$$\mathcal{K}(L^{2}(G)) \otimes A \qquad \stackrel{\Psi_{4}}{\longleftarrow} C_{0}(G, A) \rtimes_{\operatorname{lt} \otimes \operatorname{id}} G$$

and proved that Ψ_1, Ψ_2, Ψ_3 and Ψ_4 are isomorphisms. It follows readily that $\Psi = \Psi_4 \circ \Psi_3 \circ \Psi_2 \circ \Psi_1$ is an isomorphism.

Our approach to the Takai duality for L^p operator crossed products is inspired by the preceding construction. Unless otherwise stated, we assume throughout the following that (G, A, α) is an L^p operator algebra dynamical system, where G is a countable discrete Abelian group, and A is a separable unital L^p operator algebra which has unique L^p operator matrix

norms (see Definition 2.3). We construct L^p -analogues of the construction (1) as follows:

$$F^{p}(\hat{G}, F^{p}(G, A, \alpha), \hat{\alpha}) \xrightarrow{\Phi_{1}} F^{p}(G, F^{p}(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha) \xrightarrow{\Phi_{2}} F^{p}(G, C_{0}(G, A), \operatorname{lt} \otimes \alpha)$$

$$\downarrow^{\Phi_{3}}$$

$$\mathcal{K}(l^{p}(G)) \otimes_{p} A \xrightarrow{\Phi_{4}} F^{p}(G, C_{0}(G, A), \operatorname{lt} \otimes \operatorname{id})$$

Next we will explain the constructions of Φ_1, Φ_2, Φ_3 and Φ_4 .

The construction of Φ_1 . Let β be the trivial action of \hat{G} on A. For each $s \in G$, we define an isometric isomorphism $\hat{\beta}_s : C(\hat{G}, A, \beta) \to C(\hat{G}, A, \beta)$ by $\hat{\beta}_s(\varphi)(\gamma) := \gamma(s)\varphi(\gamma)$ for all $\varphi \in C(\hat{G}, A, \beta)$ and $\gamma \in \hat{G}$. Then we can form the tensor product action of G on $F^p(\hat{G}, A, \beta)$ by $(\hat{\beta} \otimes \alpha)_s(\varphi)(\gamma) := \gamma(s)\alpha_s(\varphi(\gamma))$. Thus $(\hat{\beta} \otimes \alpha)_s$ extends to an element of $\operatorname{Aut}(F^p(\hat{G}, A, \beta))$. For each $F \in C_c(\hat{G} \times G, A)$, we define $\Phi_1(F)$ as

$$\Phi_1(F)(s,\gamma) := \gamma(s)F(\gamma,s), \quad s \in G, \gamma \in \hat{G}.$$
(2)

It will be shown that Φ_1 is an isometric homomorphism with dense range (see Proposition 3.1). Then Φ_1 extends to an isometric isomorphism from $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$ onto $F^p(G, F^p(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha)$.

The construction of Φ_2 . Let lt: $G \to \operatorname{Aut}(C_0(G))$ be the isometric action induced by left translation of G on itself. Then we can form the tensor product action of G on $C_0(G,A)$ by $(\operatorname{lt} \otimes \alpha)_t f(s) := \alpha_t(f(t^{-1}s))$ for all $f \in C_0(G,A)$ and $s,t \in G$. Let $\Gamma_p: F_{\lambda}^p(\hat{G}) \to C_0(G)$ is the Gelfand transformation of $F_{\lambda}^p(\hat{G})$. Since the Gelfand transformation coincides with the Fourier transformation on $L^1(\hat{G})$, we have $\Gamma_p(\varphi)(t) = \hat{\varphi}(t) = \int_{\hat{G}} \varphi(\gamma) \overline{\gamma(t)} d\mu(\gamma)$ for all $\varphi \in L^1(\hat{G})$. For $\varphi \in C(\hat{G})$ and $a \in A$, the linear span of the elementary tensors $\varphi \otimes a$ are dense in $F^p(\hat{G},A,\beta)$. Since β is the trivial action of \hat{G} on A, by the universal property of $F^p(\hat{G},A,\beta)$ (see [16, Theorem 3.6]), there exists a contractive homomorphism from $F^p(\hat{G},A,\beta)$ to $C_0(G,A)$ by sending $\varphi \otimes a$ to $\Gamma_p(\varphi) \otimes a$. We denote this homomorphism by φ_2 . It will be shown that φ_2 is equivariant for the action $\hat{\beta} \otimes \alpha$ of G on $F^p(\hat{G},A,\beta)$ and the action lt $\otimes \alpha$ of G on $C_0(G,A)$. By the universal property of $F^p(G,F^p(\hat{G},A,\beta),\hat{\beta})$ (see [16, Theorem 3.6]), there is a homomorphism $\Phi_2 := \varphi_2 \rtimes \operatorname{id}: F^p(G,F^p(\hat{G},A,\beta),\hat{\beta} \otimes \alpha) \to F^p(G,C_0(G,A),\operatorname{lt} \otimes \alpha)$ (see Proposition 3.3) which maps $F \in C_c(G \times \hat{G},A)$ into $C_c(G,C_0(G,A))$ by the formula

$$\Phi_2(F)(s,t) := \int_{\hat{G}} F(s,\gamma) \overline{\gamma(t)} d\mu(\gamma), \qquad s,t \in G.$$
 (3)

We remark that Φ_2 is essentially induced by the Gelfand transformation Γ_p of $F_{\lambda}^p(\hat{G})$.

The construction of Φ_3 . Let id be the trivial action of G on A. Then we can form the tensor product action of G on $C_0(G,A)$ by $(\operatorname{lt} \otimes \operatorname{id})_t f(s) := f(t^{-1}s)$ for all $s,t \in G$ and $f \in C_0(G,A)$. For $F \in C_c(G,C_0(G,A))$, let $\Phi_3(F)$ be defined as

$$\Phi_3(F)(s,t) := \alpha_t^{-1}(F(s,t)), \quad s, t \in G.$$
(4)

It will be also shown that Φ_3 is isometric (see Proposition 3.7). Hence Φ_3 extends to an isometric isomorphism from $F^p(G, C_0(G, A), lt \otimes \alpha)$ onto $F^p(G, C_0(G, A), lt \otimes id)$.

The construction of Φ_4 . E. Gardella and H. Thiel [8, Theorem 4.3] have showed that $F^p(G, C_0(G), lt)$ is isometrically isomorphic to \overline{M}_G^p (see [16, Example 1.6] for the definition of \overline{M}_G^p). \overline{M}_G^p equals $\mathcal{K}(l^p(G))$ when p > 1, and is strictly smaller than $\mathcal{K}(l^1(G))$ when p = 1 and G is infinite. Since A has unique L^p operator matrix norms, the norm on spatial L^p operator tensor product $\mathcal{K}(l^p(G)) \otimes_p A$ is independent of the choice of isometric representation of A (see Lemma 2.4). We assume that $A \subseteq \mathcal{B}(E)$ is a closed subalgebra on some separable L^p space E. By the universal property of full L^p operator crossed product $F^p(G, C_0(G, A), lt \otimes id)$ (see [16, Theorem 3.6]), there exists a homomorphism $\Phi_4 : F^p(G, C_0(G, A), lt \otimes id) \to \overline{M}_G^p \otimes_p A \subseteq \mathcal{K}(l^p(G)) \otimes_p A$ which maps $F \in C_c(G \times G, A) \subseteq F^p(G, C_0(G, A), lt \otimes id)$ to $\mathcal{K}(l^p(G)) \otimes_p A$ by the formula

$$\Phi_4(F)h(t) := \sum_{s \in G} F(s, t)h(s^{-1}t), \tag{5}$$

where $h \in C_c(G, E) \subseteq l^p(G, E)$.

We will prove that the map $\Phi = \Phi_4 \circ \Phi_3 \circ \Phi_2 \circ \Phi_1$ is an equivariant homomorphism for the double dual action $\hat{\alpha}$ of G on $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$ and the action $\operatorname{Ad}\rho \otimes \alpha$ of G on $\mathcal{K}(l^p(G)) \otimes_p A$. The Pontryagin Duality Theorem allows us to identify G with the dual of \hat{G} where $s \in G$ is associated to the character $\gamma \mapsto \gamma(s)$ on \hat{G} . Therefore there is a double dual action $\hat{\alpha}$ of G on the iterated crossed product $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$. The double dual action $\hat{\alpha}$ of G on $C_c(\hat{G} \times G, A) \subseteq F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$ is given by $\hat{\alpha}_t(F)(\gamma, s) := \overline{\gamma(t)}F(\gamma, s)$ for all $t \in G$. Next let us introduce the action of G on $\mathcal{K}(l^p(G)) \otimes_p A$. Let $\rho : G \to \mathcal{B}(l^p(G))$ be the right-regular representation and form the dynamical system $\operatorname{Ad}\rho : G \to \operatorname{Aut}(\mathcal{K}(l^p(G)))$, where $(\operatorname{Ad}\rho)_s T := \rho(s)T\rho(s^{-1})$. Then we have the tensor product dynamical system $\operatorname{Ad}\rho \otimes \alpha : G \to \operatorname{Aut}(\mathcal{K}(l^p(G))) \otimes_p A$, where $(\operatorname{Ad}\rho \otimes \alpha)_s (T \otimes a) = (\operatorname{Ad}\rho)_s (T) \otimes \alpha_s(a)$.

The main result of this paper is the following theorem.

Theorem 1.2. Let (G, A, α) be an L^p operator algebra dynamical system, where G is a countable discrete Abelian group, and A is a separable unital L^p operator algebra which has unique L^p operator matrix norms. Let Φ_1 , Φ_2 , Φ_3 , Φ_4 be defined as in (2), (3), (4), (5) and $\Phi = \Phi_4 \circ \Phi_3 \circ \Phi_2 \circ \Phi_1$. Then

- (i) Φ_1 and Φ_3 are two isometric isomorphisms for $p \in [1, \infty)$;
- (ii) Φ_2 is an isomorphism if and only if either G is finite or p=2; in particular, $F^2(\hat{G}, F^2(G, A, \alpha), \hat{\alpha})$ is isometrically isomorphic to $F^p(G, C_0(G, A), \operatorname{lt} \otimes \alpha)$;
- (iii) Φ_4 is an isometric homomorphism for $p \in [1, \infty)$, and Φ_4 is an isometric isomorphism if and only if $p \in (1, \infty)$;
- (iv) Φ is equivariant for the double dual action $\hat{\alpha}$ of G on $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$ and the action $Ad\rho \otimes \alpha$ of G on $\mathcal{K}(l^p(G)) \otimes_p A$.

Corollary 1.3. Under the same assumption in Theorem 1.2, the map $\Phi : F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha}) \to \mathcal{K}(l^p(G)) \otimes_p A$ is an isomorphism if and only if either G is finite or p = 2. In particular, $F^2(\hat{G}, F^2(G, A, \alpha), \hat{\alpha})$ is isometrically isomorphic to $\mathcal{K}(l^2(G)) \otimes_2 A$.

Remark 1.4. By Theorem 1.2, Problem 1.1 is reduced to studying when $F^p(G, F^p(G, A, \beta), \hat{\beta} \otimes \alpha)$ is isomorphic to $F^p(G, C_0(G, A), lt \otimes \alpha)$. Our result shows that the natural homomorphism Φ_2 between them, induced by the Gelfand transformation of $F^p_{\lambda}(\hat{G})$, is not an isomorphism.

This suggests a negative answer to N. C. Phillips' problem on the L^p Takai duality in the case that $p \in [1, \infty) \setminus \{2\}$ and G is infinite.

Remark 1.5. Usually the norm on spatial L^p operator tensor product depends on how an L^p operator algebra is represented (see [16, page 8]). This differs significantly from the C^* -algebra case. To ensure the "uniqueness" of the norm on the spatial L^p operator tensor product, we only consider L^p operator algebras with unique L^p operator matrix norms. Furthermore, the integrated form of a nondegenerate σ -finite isometric regular covariant representation of (G, A, α) is an isometric representation of $F^p_{\lambda}(G, A, \alpha)$ (see Lemma 2.5). We do not know whether the result of Theorem 1.2 still holds without the assumption that A has unique L^p operator matrix norms.

The paper is organized as follows. In Section 2, we make some preparation for the proof of Theorem 1.2. Subsection 2.1 is devoted to the description of nondegenerate contractive representation of $F^p(G, A, \alpha)$, showing that each such representation is the integrated form for some nondegenerate contractive covariant representation of (G, A, α) (see Theorem 2.1), which will be useful in the proof of Theorem 1.2 (i). In Subsection 2.2, we prove two properties of L^p operator algebras with unique L^p operator matrix norms (see Lemmas 2.4 and 2.5). Section 3 is devoted to the proof of Theorem 1.2.

2. Preparation

In this section we make some preparation for the proof of Theorem 1.2.

2.1. Contractive representations of $F^p(G, A, \alpha)$.

This subsection is devoted to the description of nondegenerate contractive representation of $F^p(G, A, \alpha)$. The following theorem is the main result, which will be useful to prove Theorem 1.2 (i).

Theorem 2.1. Let (G, A, α) be an L^p operator algebra dynamical system, where G is a locally compact group, and A is an L^p operator algebra with a two-sided contractive approximate identity. Then every nondegenerate σ -finite contractive representation of $F^p(G, A, \alpha)$ has the form $\pi \rtimes v$ for some nondegenerate σ -finite contractive covariant representation (π, v) of (G, A, α) .

Proof. We let $L: F^p(G, A, \alpha) \to \mathcal{B}(E)$ be a nondegenerate σ -finite contractive representation of $F^p(G, A, \alpha)$ on an L^p space E. Next we shall construct a nondegenerate σ -finite contractive covariant representation of (G, A, α) on E admitting L as its integrated form. The full L^p operator crossed product $F^p(G, A, \alpha)$ in general does not contain a copy of either A or G. However, the left multiplier algebra of $F^p(G, A, \alpha)$ does. Our main approach is to construct maps from A and G into the left multiplier algebra of $F^p(G, A, \alpha)$. Let $M_L(F^p(G, A, \alpha))$ be the left multiplier algebra of $F^p(G, A, \alpha)$, and we write $\iota: F^p(G, A, \alpha) \to M_L(F^p(G, A, \alpha))$ for the canonical inclusion. Then, by [11, Theorem 4.5], there exists a canonical extension $\overline{L}: M_L(F^p(G, A, \alpha)) \to \mathcal{B}(E)$ such that $L = \overline{L} \circ \iota$.

Step 1. The construction of an injective homomorphism $i_A: A \to M_L(F^p(G, A, \alpha))$.

For each $f \in C_c(G, A, \alpha)$, we define $i_A(a)f(s) := af(s)$. Clearly, if (π_1, u) is a contractive covariant representation of (G, A, α) on an L^p space E_0 , then we have

$$\pi_1 \rtimes u(i_A(a)f) = \pi_1(a) \circ \pi_1 \rtimes u(f). \tag{6}$$

It follows that

$$||i_A(a)f|| \le ||a||||f||.$$

Therefore $i_A(a)$ extends to a map from $F^p(G, A, \alpha)$ to itself. For $f, g \in C_c(G, A, \alpha)$, note that

$$(i_A(a)f) * g(t) = \int_G i_A(a)f(s)\alpha_s(g(s^{-1}t))d\mu(s)$$
$$= a\int_G f(s)\alpha_s(g(s^{-1}t))d\mu(s)$$
$$= i_A(a)(f * g)(t).$$

Since $i_A(a)$ is bounded and $C_c(G, A, \alpha)$ is dense in $F^p(G, A, \alpha)$, it follows that $i_A(a) \in M_L(F^p(G, A, \alpha))$. Obviously, i_A is a homomorphism from A to $M_L(F^p(G, A, \alpha))$.

To see that i_A is injective, let $\rho_0: A \to \mathcal{B}(E_1)$ be an faithful nondegenerate representation of A on an L^p space E_1 , and let $(\rho, \lambda_p^{E_1})$ be its associated regular covariant representation on $L^p(G, E_1)$. Then $\rho: A \to \mathcal{B}(L^p(G, E_1))$ is faithful and nondegenerate (see [16, Lemma 2.11(7)]. Let $\rho \rtimes \lambda_p^{E_1}: M_L(F^p(G, A, \alpha)) \to \mathcal{B}(L^p(G, E_1))$ be the canonical extension of $\rho \rtimes \lambda_p^{E_1}$. Since A has a two-sided contractive approximate identity, by [11, Theorem 5.5], there is a two-sided contractive approximate identity in $F^p(G, A, \alpha)$. Since $(\rho, \lambda_p^{E_1})$ is nondegenerate, by (6), we have $\rho(a) = \overline{\rho \rtimes \lambda_p^{E_1}} \circ i_A(a)$. It follows that i_A is injective.

Step 2. The construction of an injective homomorphism from G into the group of invertible isometries in $M_L(F^p(A, G, \alpha))$.

For each $s \in G$, we define $i_G(s) : C_c(G, A, \alpha) \to C_c(G, A, \alpha)$ as $i_G(s)f(t) := \alpha_s(f(s^{-1}t))$ for $f \in C_c(G, A, \alpha)$ and $t \in G$. Note that if (π_2, w) is a contractive covariant representation of (G, A, α) on an L^p -space E_2 , then for $\xi \in E_2$ we have

$$\pi_2 \rtimes w(i_G(s)f))(\xi) = \int_G \pi_2(i_G(s)f(t))w_t(\xi)d\mu(t)$$

$$= \int_G \pi_2(\alpha_s(f(s^{-1}t)))w_t(\xi)d\mu(t)$$

$$= \int_G \pi_2(\alpha_s(f(t)))w_{st}(\xi)d\mu(t)$$

$$= \int_G w_s\pi_2(f(t))w_{s^{-1}}w_{st}(\xi)d\mu(t)$$

$$= w_s \circ \pi_2 \rtimes w(f)(\xi).$$

Hence we have

$$\pi_2 \rtimes w(i_G(s)f)) = w_s \circ \pi_2 \rtimes w(f). \tag{7}$$

Since w_s is an isometry, by the universal property of $F^p(G, A, \alpha)$ (see [16, Theorem 3.6]), we have $||i_G(s)f|| = ||f||$. Thus we can extend $i_G(s)$ to an isometric map from $F^p(G, A, \alpha)$ to itself.

For $f, g \in C_c(G, A, \alpha)$ and $r, t \in G$, we have

$$(i_{G}(r)f) * g(t) = \int_{G} i_{G}(r)f(s)\alpha_{s}(g(s^{-1}t))d\mu(s)$$

$$= \int_{G} \alpha_{r}(f(r^{-1}s))\alpha_{s}(g(s^{-1}t))d\mu(s)$$

$$= \int_{G} \alpha_{r}(f(s))\alpha_{rs}(g(s^{-1}r^{-1}t))d\mu(s)$$

$$= \alpha_{r}(\int_{G} f(s)\alpha_{s}(g(s^{-1}r^{-1}t))d\mu(s)$$

$$= \alpha_{r}(f * g(r^{-1}t))$$

$$= i_{G}(r)(f * g)(t).$$

Since $i_G(r)$ is isometric on $F^p(G, A, \alpha)$ and $C_c(G, A, \alpha)$ is dense in $F^p(G, A, \alpha)$, it follows that $i_G(r) \in M_L(F^p(A, G, \alpha))$. It is easy to check $i_G(rs) = i_G(r)i_G(s)$ and $i_G(r)^{-1} = i_G(r^{-1})$. Then i_G is an invertible isometry-valued homomorphism.

To show that i_G is injective, let $\pi_0: A \to \mathcal{B}(E_0)$ be a faithful nondegenerate representation of A on an L^p space E_0 , and let $(\pi_3, \lambda_p^{E_0})$ be its associated regular covariant representation on $L^p(G, E_0)$. Let $\overline{\pi_3} \rtimes \lambda_p^{E_0}: M_L(F^p(G, A, \alpha)) \to \mathcal{B}(L^p(G, E_0))$ be the canonical extension of $\pi_3 \rtimes \lambda_p^{E_0}$. By (7), for each $f \in C_c(G, A, \alpha)$, we have $\pi_3 \rtimes \lambda_p^{E_0}(i_G(s)f)) = \lambda_p^{E_0}(s) \circ \pi_3 \rtimes \lambda_p^{E_0}(f)$. Since there is a two-sided contractive approximate identity in $F^p(G, A, \alpha)$ (see [11, Theorem 5.5]) and $(\pi_3, \lambda_p^{E_0})$ is nondegenerate, we have $\lambda_p^{E_0}(s) = \overline{\pi_3} \rtimes \lambda_p^{E_0} \circ i_G(s)$. So, if $s \neq e$, then we certainly have $\lambda_p^{E_0}(s) \neq I_{L^p(G,E_0)}$, where $I_{L^p(G,E_0)}$ is the identity operator on $L^p(G,E_0)$. This shows that $i_G(s) \neq I$ if $s \neq e$, where I is the identity element in $M_L(F^p(G,A,\alpha))$.

It is clear that $\overline{L} \circ i_G$ is a representation of G and $\overline{L} \circ i_A$ is a representation of A. Let $(\pi, v) = (\overline{L} \circ i_A, \overline{L} \circ i_G)$. Now it suffices to check that (π, v) is the nondegenerate σ -finite contractive covariant representation of (G, A, α) and $L = \pi \rtimes v$.

Step 3. Verification.

For $f \in C_c(G, A, \alpha)$, $s, t \in G$ and $a \in A$, note that

$$i_G(s)i_A(a)i_G(s^{-1})f(t) = \alpha_s(i_A(a)i_G(s^{-1})f(s^{-1}t)) = \alpha_s(ai_G(s^{-1})f(s^{-1}t))$$
$$= \alpha_s(a)\alpha_s(\alpha_{s^{-1}}(f(ss^{-1}t))) = i_A(\alpha_s(a))f(t)$$

and

$$(\overline{L} \circ i_G)(s)(\overline{L} \circ i_A)(a)(\overline{L} \circ i_G)(s^{-1}) = \overline{L} \circ i_G(s)i_A(a)i_G(s^{-1}) = \overline{L} \circ i_A(\alpha_s(a)).$$

Since L is σ -finite and contractive, it is easy to check that $\overline{L} \circ i_A$ is also σ -finite and contractive. Now it remains to show that $\overline{L} \circ i_A$ is nondegenerate. If $\{e_i\}$ is a left contractive approximate identity of A, then $\{i_A(e_i)\}$ is a left contractive approximate identity in $M_L(F^p(G, A, \alpha))$. For $f \in C_c(G, A, \alpha)$ and $\xi \in E$, we have

$$\lim_{i} (\overline{L} \circ i_{A}(e_{i})) L(f) \xi = \lim_{i} (\overline{L}(e_{i}f) \xi = \lim_{i} L(e_{i}f) \xi = L(f) \xi.$$

Since L is nondegenerate, it follows that $\{\overline{L} \circ i_A(e_i)\}$ converges to the identity operator I_E in $\mathcal{B}(E)$. Hence $\pi = \overline{L} \circ i_A$ is nondegenerate. Thus (π, v) is a nondegenerate covariant representation of (G, A, α) on the L^p space E.

For $a \in A$ and $g \in C_c(G)$, note that

$$(\overline{L} \circ i_A) \rtimes (\overline{L} \circ i_G)(i_A(a)i_G(g)) = \int_G \overline{L} \circ i_A(i_A(a)g(s))\overline{L} \circ i_G(s)d\mu(s)$$

$$= \int_G \overline{L} \circ i_A(ag(s))\overline{L} \circ i_G(s)d\mu(s)$$

$$= \overline{L} \circ i_A(a) \int_G g(s)\overline{L} \circ i_G(s)d\mu(s)$$

$$= \overline{L} \circ i_A(a)\overline{L} \circ i_G(g) = \overline{L}(i_A(a)i_G(g))$$

$$= L(i_A(a)i_G(g)).$$

By [23, Lemma 1.87], the span of elements of the form $i_A(a)i_G(g)$ are dense in $F^p(G, A, \alpha)$ (in fact, $i_A(a)i_G(g)(s) = g(s)a$). It follows that $L = (\overline{L} \circ i_A) \rtimes (\overline{L} \circ i_G) = \pi \rtimes v$.

2.2. Unique L^p operator matrix norms.

In this subsection, we shall prove two properties of L^p operator algebras with unique L^p operator matrix norms.

Definition 2.2 ([19, Definition 3.3]). Let (X, \mathcal{B}, μ) be a measure space, and let $A \subseteq \mathcal{B}(L^p(X, \mu))$ be a closed subalgebra. We equip $M_n(A)$ with the matrix norms coming from the identification of $M_n(A)$ with a closed subalgebra of $\mathcal{B}(L^p(\{1, 2, ..., n\} \times X, \nu \times \mu))$, where ν is the counting measure on $\{1, 2, ..., n\}$.

Definition 2.3 ([19, Definition 4.1]). Let $p \in [1, \infty)$ and A be a separable L^p operator algebra. We say that A has unique L^p operator matrix norms if whenever (X, \mathcal{B}, μ) and (Y, \mathcal{C}, ν) are σ -finite measure spaces such that $L^p(X, \mu)$ and $L^p(Y, \nu)$ are separable, $\pi : A \to \mathcal{B}(L^p(X, \mu))$ and $\sigma : A \to \mathcal{B}(L^p(Y, \nu))$ are isometric representations, and $\pi(A)$ and $\sigma(A)$ are given the matrix normed structure of Definition 2.2, then $\sigma \circ \pi^{-1} : \pi(A) \to \sigma(A)$ is completely isometric.

 M_n^p and C(X) are basic examples with unique L^p operator matrix norms (see [19, Corollary 4.4 and Proposition 4.6]), where X is compact metrizable space.

Given a countable set X endowed with the counting measure, we denote by $l^p(X)$ the corresponding L^p space. When $X = \mathbb{N}$, we simply write l^p for $l^p(\mathbb{N})$.

Lemma 2.4. Let $B \subseteq \mathcal{B}(l^p)$ be a closed subalgebra, and let A be a separable L^p operator algebra with unique L^p operator matrix norms. Then the norm on spatial L^p operator tensor product $B \otimes_p A$ is independent of the choice of isometric representation of A.

Proof. For each $n \geq 0$, we let P_n be the canonical projection on l^p defined by $P_n(\sum_{i=0}^{\infty} \alpha_i e_i) = \sum_{i=0}^{i=n} \alpha_i e_i$, where $\{e_i\}$ is the standard basis in l^p and $\alpha_i \in \mathbb{C}$. Then we have $||P_n x - x|| \to 0$ for all $x \in l^p$. It is not hard to show that for every $T \in \mathcal{B}(l^p \otimes_p E)$, where E is a separable L^p space, we have

$$||T|| = \sup_{n} \{||(P_n \otimes I_E)T(P_n \otimes I_E)||\},$$

where I_E is the identity operator on E. Since A is separable, by [16, Proposition 1.25] and [19, Lemma 2.7], we only need separable L^p spaces and the measure is σ -finite. Let $\pi_1: A \to \mathcal{B}(L^p(X,\mu))$ and $\pi_2: A \to \mathcal{B}(L^p(Y,\nu))$ be two isometric representations of A on two separable L^p spaces and the measures μ , ν are σ -finite, respectively. Thus, if $\sum b_i \otimes a_i \in B \otimes_{alg} A$ (\otimes_{alg} is the algebraic tensor product), then

$$\|\sum b_i \otimes \pi_1(a_i)\| = \sup_n \{\|\sum (P_n b_i P_n) \otimes \pi_1(a_i)\|\}$$

and

$$\|\sum b_i \otimes \pi_2(a_i)\| = \sup_n \{\|\sum (P_n b_i P_n) \otimes \pi_2(a_i)\|\}.$$

Since $P_n\mathcal{B}(l^p)P_n$ is isomorphic to M_n^p , and A has unique L^p operator matrix norms, we have $\|\sum (P_nb_iP_n)\otimes \pi_1(a_i)\| = \|\sum (P_nb_iP_n)\otimes \pi_2(a_i)\|$ for each n. Therefore $\|\sum b_i\otimes \pi_1(a_i)\| = \|\sum b_i\otimes \pi_2(a_i)\|$.

The preceding lemma shows that if A is a separable L^p operator algebra with unique L^p operator matrix norms, and G is a countable discrete group, then the norm on spatial L^p operator tensor product $\mathcal{K}(l^p(G))\otimes_p A$ is independent of the choice of isometric representation of A.

If A is a C^* -algebra, then it is well known that the reduced crossed product $A \rtimes_{\alpha,r} G$ does not depend on the choice of faithful representation of $A \subseteq \mathcal{B}(H)$, since there is a unique C^* -norm on $M_n(A)$ (see [3, Proposition 8.4]). For L^p operator algebras with unique L^p operator matrix norms, we have an analogous result.

Lemma 2.5. Let (G, A, α) be an L^p operator algebra dynamical system, where G is a countable discrete group and A is a separable L^p operator algebra with unique L^p operator matrix norms. Let $\pi_0 : A \to \mathcal{B}(E_0)$ be a nondegenerate σ -finite isometric representation of A on an L^p space E_0 , and let $(\pi, \lambda_p^{E_0})$ be its associated regular covariant representation. Then the integrated form $\pi \rtimes \lambda_p^{E_0}$ is an isometric representation of $F_\lambda^p(G, A, \alpha)$.

Proof. We give the proof by proving two claims.

Claim 1. For any $f \in C_c(G, A, \alpha)$, it holds that

$$||f||_{F^p_{\lambda}(G,A,\alpha)} = \sup\{||(\pi \rtimes v(f))|| : (\pi,v) \in \operatorname{IsoRegRep}_p(G,A,\alpha)\},$$

where IsoRegRep_p(G, A, α) denotes the class consisting of nondegenerate σ -finite isometric regular covariant representation of (G, A, α) .

By [16, Lemma 3.19], there is a nondegenerate σ -finite isometric representation $\rho_0: A \to \mathcal{B}(L^p(X,\mu))$ such that, with $(\rho, \lambda_p^{L^p(X,\mu)})$ being its associated regular covariant representation, the representation $\rho \rtimes \lambda_p^{L^p(X,\mu)}$ is nondegenerate and isometric on $F_{\lambda}^p(G,A,\alpha)$. Then one can see Claim 1.

Claim 2. The norm of $\pi \rtimes \lambda_p^{E_0}(f)$ is independent of the choice of nondegenerate σ -finite isometric representation π_0 .

Since A is separable, by [16, Proposition 1.25] and [19, Lemma 2.7], we may assume that E_0 is a separable L^p space and the measure from E_0 is σ -finite. Arbitrarily choose a nonempty finite set $F \subseteq G$. Let $P \in \mathcal{B}(l^p(G))$ be the finite rank projection onto the span of $\{\delta_t : t \in F\}$, where $\{\delta_t\}_{t \in G}$ is the canonical basis of $l^p(G)$. Let I be the identity operator in $\mathcal{B}(E_0)$. Rather than compute the norm of $f \in C_c(G, A, \alpha)$ under the representation $\pi \rtimes \lambda_p^{E_0} : C_c(G, A, \alpha) \to \mathcal{B}(l^p(G) \otimes_p E_0)$, we will cut by the projection $P \otimes I$ and show that the norm of $(P \otimes I)\pi \rtimes \lambda_p^{E_0}(f)(P \otimes I)$ is independent of the isometric representation $\pi_0 : A \to \mathcal{B}(E_0)$. By taking a limit over finite sets in G, we conclude that so is the norm of $\pi \rtimes \lambda_p^{E_0}(f)$.

Since F is a finite subset of G, let M_F^p be the collection of those $T \in \mathcal{B}(l^p(G))$ satisfying $T(l^p(G \setminus F)) = \{0\}$ and $T(l^p(F)) \subseteq l^p(F)$ (see [16, Example 1.6]). Let $\{e_{s,t}\}_{s,t\in F}$ be the canonical matrix units of $P\mathcal{B}(l^p(G))P \cong M_F^p$ and fix arbitrary $a \in A$. We claim that

$$(P \otimes I)\pi(a) = (P \otimes I)\pi(a)(P \otimes I) = \sum_{t \in F} e_{t,t} \otimes \pi_0(\alpha_{t^{-1}}(a)).$$

This is clear if $\pi(a)$ is considered as a diagonal matrix in $\mathcal{B}(\bigoplus_{t\in G} E_0)$. Identifying $l^p(G)\otimes_p E_0$ with the L^p direct sum $\bigoplus_{t\in G} E_0$, we may simply take the L^p direct sum representation

$$\pi(a) = \bigoplus_{t \in G} \pi_0(\alpha_{t^{-1}}(a)) \in \mathcal{B}(\bigoplus_{t \in G} E_0).$$

In the spatial L^p operator tensor product picture, we have

$$\pi(a) = \sum_{t \in G} e_{t,t} \otimes \pi_0(\alpha_{t-1}(a)),$$

where the convergence is in the strong operator topology.

Let $\lambda_p: G \to \mathcal{B}(l^p(G))$ be the left-regular representation. For each $s \in G$, we have $\lambda_p^{E_0}(s) = \lambda_p(s) \otimes I$. Thus one can see that

$$(P \otimes I)\pi(a)(\lambda_p^{E_0}(s))(P \otimes I) = (P \otimes I)\pi(a)(\lambda_p(s) \otimes I)(P \otimes I)$$

$$= (\sum_{t \in F} e_{t,t} \otimes \pi_0(\alpha_{t^{-1}}(a)))(P\lambda_p(s)P \otimes I)$$

$$= (\sum_{t \in F} e_{t,t} \otimes \pi_0(\alpha_{t^{-1}}(a)))(\sum_{r \in F \cap sF} e_{r,s^{-1}r} \otimes I)$$

$$= \sum_{r \in F \cap sF} e_{r,s^{-1}r \otimes \pi_0(\alpha_{r^{-1}}(a))} \in M_F^p \otimes \pi_0(A).$$

Now if $f = \sum_{s \in G} a_s \delta_s \in C_c(G, A, \alpha)$, then we have

$$(P \otimes I)\pi \rtimes \lambda_p^{E_0}(f)(P \otimes I) = \sum_{s \in G} \sum_{r \in F \cap sF} e_{r,s^{-1}r} \otimes \pi_0(\alpha_{r^{-1}}(a_s)) \in M_F^p \otimes \pi_0(A).$$

Since A has unique L^p operator norms, the norm of $(P \otimes I)\pi \rtimes \lambda_p^{E_0}(f)(P \otimes I)$ is independent of the choice of nondegenerate σ -finite isometric representation $\pi_0: A \to \mathcal{B}(E_0)$. This proves Claim 2.

By Claims 1 and 2, $\pi \rtimes \lambda_p^{E_0}$ is an isometric representation of $F_{\lambda}^p(G,A,\alpha)$.

3. Proof of Theorem 1.2

In the rest of this paper, let G be a countable discrete Abelian group. Thus we may assume that the Haar measure on G is the counting measure. Since the dual group \hat{G} is compact, we also denote by μ the unique normalized positive Haar measure on \hat{G} .

3.1. The isometry property of Φ_1 .

The aim of this subsection is to prove the following result.

Proposition 3.1. Let $\Phi_1: F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha}) \to F^p(G, F^p(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha)$ be the map defined in (2) under the assumption of Theorem 1.2. Then Φ_1 is an isometric isomorphism.

We first make some preparation. For the reader's convenience, we recall the definitions of $\hat{\alpha}$ and $\hat{\beta} \otimes \alpha$. For each $\gamma \in \hat{G}$, $\hat{\alpha}_{\gamma} : C_c(G, A, \alpha) \to C_c(G, A, \alpha)$ is defined by $\hat{\alpha}_{\gamma}(f)(s) := \overline{\gamma(s)}f(s)$ for all $f \in C_c(G, A, \alpha)$ and $s \in G$. Then $\hat{\alpha}_{\gamma}$ extends to an isometry on $F^p(G, A, \alpha)$. Let β be the trivial action of \hat{G} on A. For each $s \in G$, we define an isometric isomorphism $\hat{\beta}_s : C(\hat{G}, A, \beta) \to C(\hat{G}, A, \beta)$ by $\hat{\beta}_s(\varphi)(\gamma) := \gamma(s)\varphi(\gamma)$ for all $\varphi \in C(\hat{G}, A, \beta)$ and $\gamma \in \hat{G}$. Then we can form the tensor product action of G on $F^p(\hat{G}, A, \beta)$ by $(\hat{\beta} \otimes \alpha)_s(\varphi)(\gamma) := \gamma(s)\alpha_s(\varphi(\gamma))$. Hence $(\hat{\beta} \otimes \alpha)_s$ extends to an element of $\operatorname{Aut}(F^p(\hat{G}, A, \beta))$.

Lemma 3.2. Let (G, A, α) be an L^p operator algebra dynamical system, where G is a countable discrete Abelian group, and A is an L^p operator algebra. Then

- (i) $C_c(\hat{G} \times G, A)$ is a dense subalgebra of $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$.
- (ii) $C_c(G \times \hat{G}, A)$ is a dense subalgebra of $F^p(G, F^p(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha)$.

Proof. (i) If $F \in C_c(\hat{G} \times G, A)$, then $\lambda_F(\gamma)(s) := F(\gamma, s)$ defines an element $\lambda_F \in C_c(\hat{G}, C_c(G, A))$ $\subseteq C_c(\hat{G}, F^p(G, A, \alpha))$ for all $\gamma \in \hat{G}$ and $s \in G$. That is, $\lambda_F(\gamma) \in C_c(G, A)$ for all $\gamma \in \hat{G}$. For $\sigma \in \hat{G}$ and $F_1, F_2 \in C_c(\hat{G} \times G, A)$, note that

$$\lambda_{F_1} * \lambda_{F_2}(\sigma) = \int_{\hat{G}} \lambda_{F_1}(\gamma) * \hat{\alpha}_{\gamma}(\lambda_{F_2}(\gamma^{-1}\sigma)) d\mu(\gamma)$$

takes value in $F^p(G, A, \alpha)$; moreover, by [23, Lemma 1.108], the former integral takes value in $C_c(G, A) \subseteq F^p(G, A, \alpha)$. For each $t \in G$, we have

$$\lambda_{F_1} * \lambda_{F_2}(\sigma)(t) = \int_{\hat{G}} \sum_{s \in G} \lambda_{F_1}(\gamma)(s) \alpha_s(\hat{\alpha}_{\gamma}(\lambda_{F_2}(\gamma^{-1}\sigma))(s^{-1}t)) d\mu(\gamma)$$
$$= \int_{\hat{G}} \sum_{s \in G} \lambda_{F_1}(\gamma)(s) \overline{\gamma(s^{-1}t)} \alpha_s(\lambda_{F_2}(\gamma^{-1}\sigma)(s^{-1}t)) d\mu(\gamma).$$

For convenience, we can rewrite this equality as

$$F_1 * F_2(\sigma, t) = \int_{\hat{G}} \sum_{s \in G} F_1(\gamma, s) \overline{\gamma(s^{-1}t)} \alpha_s(F_2(\gamma^{-1}\sigma, s^{-1}t)) d\mu(\gamma).$$

Therefore we can view $C_c(\hat{G} \times G, A)$ as a subalgebra of $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$.

Since $C_c(G, A, \alpha)$ is dense in $F^p(G, A, \alpha)$, by [23, Lemma 1.87], $C(\hat{G}, C_c(G, A, \alpha), \hat{\alpha})$ is dense in $C(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$. Thus $C_c(\hat{G} \times G, A)$ is dense in $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$.

Using a similar argument (i) as in the proof of Lemma 3.2, one can prove (ii) by exchanging G with \hat{G} .

Proof of Proposition 3.1. The proof consists of four claims.

Claim 1. Φ_1 is a homomorphism from $C_c(\hat{G} \times G, A)$ onto $C_c(G \times \hat{G}, A)$.

Obviously, Φ_1 is a surjective map. Next, we will show that Φ_1 is a homomorphism. For each $F_1, F_2 \in C_c(\hat{G} \times G, A)$, we have

$$\lambda_{\Phi_1(F_1)} * \lambda_{\Phi_1(F_2)}(t) = \sum_{s \in G} \lambda_{\Phi_1(F_1)}(s) * (\hat{\beta} \otimes \alpha)_s (\lambda_{\Phi_1(F_2)}(s^{-1}t)).$$

Then we have

$$\begin{split} \Phi_1(F_1) * \Phi_1(F_2)(t,\sigma) &= \lambda_{\Phi_1(F_1)} * \lambda_{\Phi_1(F_2)}(t)(\sigma) \\ &= \sum_{s \in G} \int_{\hat{G}} \lambda_{\Phi_1(F_1)}(s)(\gamma) \beta_{\gamma}((\hat{\beta} \otimes \alpha)_s (\lambda_{\Phi_1(F_2)}(s^{-1}t))(\gamma^{-1}\sigma)) d\mu(\gamma) \\ &= \sum_{s \in G} \int_{\hat{G}} \lambda_{\Phi_1(F_1)}(s)(\gamma)(\gamma^{-1}\sigma)(s) \alpha_s (\lambda_{\Phi_1(F_2)}(s^{-1}t)(\gamma^{-1}\sigma)) d\mu(\gamma) \\ &= \sum_{s \in G} \int_{\hat{G}} \gamma(s) F_1(\gamma, s)(\gamma^{-1}\sigma)(s)(\gamma^{-1}\sigma)(s^{-1}t) \alpha_s (F_2(\gamma^{-1}\sigma, s^{-1}t)) d\mu(\gamma) \\ &= \sum_{s \in G} \int_{\hat{G}} F_1(\gamma, s) \overline{\gamma(s^{-1}t)} \sigma(t) \alpha_s (F_2(\gamma^{-1}\sigma, s^{-1}t)) d\mu(\gamma) \\ &= \int_{\hat{G}} \sum_{s \in G} F_1(\gamma, s) \overline{\gamma(s^{-1}t)} \sigma(t) \alpha_s (F_2(\gamma^{-1}\sigma, s^{-1}t)) d\mu(\gamma) \\ &= \sigma(t) \lambda_{F_1} * \lambda_{F_2}(\sigma)(t) = \Phi_1(F_1 * F_2)(t, \sigma). \end{split}$$

This shows that $\Phi_1(F_1 * F_2) = \Phi_1(F_1) * \Phi_1(F_2)$. Hence Φ_1 is a homomorphism from $C_c(\hat{G} \times G, A)$ onto $C_c(G \times \hat{G}, A)$.

Since Φ_1 maps a dense subalgebra onto a dense subalgebra, it suffices to show that Φ_1 is isometric for the universal norms.

Let L be a nondegenerate σ -finite contractive representation of $F^p(G, F^p(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha)$. By Theorem 2.1, there exists a nondegenerate σ -finite contractive covariant representation (R, U) of $(G, F^p(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha)$ such that $L = R \rtimes U$, and there exists a nondegenerate σ -finite contractive covariant representation (π, V) of (\hat{G}, A, β) such that $R = \pi \rtimes V$. Claim 2. $U_sV_{\gamma} = \gamma(s)V_{\gamma}U_s$ for each $\gamma \in \hat{G}$ and each $s \in G$.

Let $i_{\hat{G}}$ be the canonical map of \hat{G} into $M_L(F^p(\hat{G}))$. Then $i_{\hat{G}}(\gamma)\varphi(\sigma) = \beta_{\gamma}(\varphi(\gamma^{-1}\sigma)) = \varphi(\gamma^{-1}\sigma)$ for all $\gamma, \sigma \in \hat{G}$ and $\varphi \in C(\hat{G})$. For each $s \in G$,

$$\hat{\beta}_s \circ i_{\hat{G}}(\gamma)\varphi(\sigma) = \sigma(s)i_{\hat{G}}(\gamma)\varphi(\sigma) = \sigma(s)\varphi(\gamma^{-1}\sigma) = \gamma(s)(\gamma^{-1}\sigma)(s)\varphi(\gamma^{-1}\sigma)$$
$$= \gamma(s)\hat{\beta}_s\varphi(\gamma^{-1}\sigma) = \gamma(s)i_{\hat{G}}(\gamma)\circ\hat{\beta}_s\varphi(\sigma),$$

it follows that $\hat{\beta}_s \circ i_{\hat{G}}(\gamma) = \gamma(s)i_{\hat{G}}(\gamma) \circ \hat{\beta}_s$.

Now if $a \in A$, $\varphi \in C(\hat{G})$ and $f \in C_c(G)$, then the functions which are elementary tensors of the form $\varphi \otimes f \otimes a$ span a dense subset of $C_c(\hat{G} \times G, A)$. Since

$$U_{s}V_{\gamma}L(\varphi \otimes f \otimes a) = U_{s}V_{\gamma}R \rtimes U(\varphi \otimes f \otimes a) = U_{s}V_{\gamma}\pi(a)V(\varphi)U(f)$$

$$= U_{s}\pi(a)V_{\gamma}V(\varphi)U(f) = U_{s}\pi(a)V(i_{\hat{G}}(\gamma)\varphi)U(f)$$

$$= U_{s}\pi \rtimes V(i_{\hat{G}}(\gamma)(\varphi) \otimes a)U(f) = \pi \rtimes V((\hat{\beta} \otimes \alpha)_{s}(i_{\hat{G}}(\gamma)(\varphi) \otimes a))U_{s}U(f)$$

$$= \pi(\alpha_{s}(a))V(\hat{\beta}_{s} \circ i_{\hat{G}}(\gamma)\varphi)U_{s}U(f) = \pi(\alpha_{s}(a))V(\gamma(s)i_{\hat{G}}(\gamma) \circ \hat{\beta}_{s}(\varphi))U_{s}U(f)$$

$$= \gamma(s)\pi(\alpha_{s}(a))V_{\gamma}V(\hat{\beta}_{s}(\varphi))U_{s}U(f) = \gamma(s)V_{\gamma}\pi(\alpha_{s}(a))V(\hat{\beta}_{s}(\varphi))U_{s}U(f)$$

$$= \gamma(s)V_{\gamma}U_{s}\pi \rtimes V((\hat{\beta} \otimes \alpha)_{s^{-1}}(\hat{\beta}_{s}(\varphi) \otimes \alpha_{s}(a))U(f)$$

$$= \gamma(s)V_{\gamma}U_{s}\pi(a)V(\varphi)U(f) = \gamma(s)V_{\gamma}U_{s}L(\varphi \otimes f \otimes a),$$

and L is nondegenerate, it follows that

$$U_s V_{\gamma} = \gamma(s) V_{\gamma} U_s. \tag{8}$$

Claim 3. (π, U) is a σ -finite covariant representation of (G, A, α) and $(\pi \rtimes U, V)$ is a σ -finite covariant representation of $(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$.

For each $b \in A$, a similar computation as in the proof of (8) shows that

$$U_{s}\pi(b)\pi(a)V(\varphi)U(f) = U_{s}\pi(ba)V(\varphi)U(f)$$

$$= U_{s}\pi \rtimes V(\varphi \otimes ba)U(f)$$

$$= \pi \rtimes V((\hat{\beta} \otimes \alpha)_{s}(\varphi \otimes ba))U_{s}U(f)$$

$$= \pi(\alpha_{s}(b))\pi \rtimes V(\hat{\beta}_{s}(\varphi) \otimes \alpha_{s}(a))U_{s}U(f)$$

$$= \pi(\alpha_{s}(b))U_{s}\pi \rtimes V(\varphi \otimes a)U(f)$$

$$= \pi(\alpha_{s}(b))U_{s}\pi(a)V(\varphi)U(f).$$

Thus (π, U) is a covariant representation of (G, A, α) . Obviously, the representation π is σ -finite.

Since the action of \hat{G} on A is trivial, by (8), we have

$$V_{\gamma}\pi \rtimes U(a\otimes f) = \pi(a)V_{\gamma}\sum_{s\in G}f(s)U_{s} = \pi(a)\sum_{s\in G}f(s)\overline{\gamma(s)}U_{s}V_{\gamma} = \pi \rtimes U(\hat{\alpha}_{\gamma}(a\otimes f))V_{\gamma}.$$

Hence $(\pi \rtimes U, V)$ is a covariant representation of $(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$. Since U is σ -finite, it follows that $\pi \rtimes U$ is σ -finite.

Claim 4. $||\Phi_1(F)|| \le ||F||$.

From Claim 3, $L' = (\pi \rtimes U) \rtimes V$ is a contractive representation of $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$. By the Fubini theorem and (8), we have

$$L(\Phi_1(F)) = \sum_{s \in G} \pi \rtimes V(\lambda_{\Phi_1(F)}(s))U_s = \sum_{s \in G} \int_{\hat{G}} \pi(\Phi_1(F)(s,\gamma))V_{\gamma}U_s d\mu(\gamma)$$
$$= \sum_{s \in G} \int_{\hat{G}} \pi(F(\gamma,s))\gamma(s)V_{\gamma}U_s d\mu(\gamma) = \int_{\hat{G}} \sum_{s \in G} \pi(F(\gamma,s))U_s V_{\gamma} d\mu(\gamma) = L'(F).$$

It follows that $||\Phi_1(F)|| \le ||F||$. Reversing the above argument gives $||F|| \le ||\Phi_1(F)||$. This completes the proof.

3.2. A characterization of Φ_2 being an isomorphism.

The aim of this subsection is to prove the following.

Proposition 3.3. Let $\Phi_2 : F^p(G, F^p(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha) \to F^p(G, C_0(G, A), \operatorname{lt} \otimes \alpha)$ be defined in (3) under the assumption of Theorem 1.2. Then Φ_2 is an isomorphism if and only if either G is finite or p = 2; in particular, Φ_2 is an isometric isomorphism for p = 2.

Clearly, Φ_2 is defined in terms of the Gelfand transformation of reduced group L^p operator algebra. The following theorem concerning the Gelfand transformation of $F_{\lambda}^p(G)$ will be useful later.

Theorem 3.4 ([24, Theorem 2.7] or [8, Proposition 3.22 and Corollary 3.20]). Let G be a locally compact Abelian group and $p \in [1, \infty) \setminus \{2\}$. Then

- (i) the Gelfand transformation $\Gamma_p: F_{\lambda}^p(G) \to C_0(\hat{G})$ is an injective contractive map with dense range;
- (ii) the Gelfand transformation Γ_p is surjective if and only if G is finite.

Lemma 3.5. Let A be a separable L^p operator algebra with unique L^p operator matrix norms, G be a countable discrete Abelian group, and $\varphi_2 : F^p(\hat{G}, A, \beta) \to C_0(G, A)$ be defined in (3).

- (i) If $p \in [1, \infty) \setminus \{2\}$, then φ_2 is an isomorphism if and only if G is finite.
- (ii) If p = 2, then φ_2 is an isometric isomorphism.

Proof. We first prove two key claims.

Claim 1. There exists a nondegenerate σ -finite isometric representation π_0 of A on some L^p space such that $F^p(\hat{G}, A, \beta)$ is isometrically isomorphic to $F^p_{\lambda}(\hat{G}) \otimes_p \pi_0(A)$.

Since \hat{G} is amenable, it follows that $F^p(\hat{G}, A, \beta)$ is isometrically isomorphic to $F^p_{\lambda}(\hat{G}, A, \beta)$ (see [18, Theorem 7.1]). By [16, Lemma 3.19], there exists a a nondegenerate σ -finite isometric representation $\pi_0: A \to \mathcal{B}(L^p(X, \nu))$ such that, with $(\pi, \lambda_p^{L^p(X, \nu)})$ being its associated regular covariant representation, the representation $\pi \rtimes \lambda_p^{L^p(X, \nu)}$ is nondegenerate and isometric on $F^p_{\lambda}(\hat{G}, A, \beta)$. Since β is the trivial action of \hat{G} on A, it follows that $F^p_{\lambda}(\hat{G}, A, \beta)$

is isometrically isomorphic to $F_{\lambda}^{p}(\hat{G}) \otimes_{p} \pi_{0}(A)$. We denote by ρ the isometric isomorphism from $F^{p}(\hat{G}, A, \beta)$ onto $F_{\lambda}^{p}(\hat{G}) \otimes_{p} \pi_{0}(A)$.

Claim 2. $\varphi_2: F^p(\hat{G}, A, \beta) \to C_0(G, A)$ is an injective homomorphism.

Note that the norm on $C_0(G) \otimes_p A \cong C_0(G,A)$ is independent of the choice of isometric representation of A, then we let $\mathrm{Id}_{C_0(G)} \otimes \pi_0$ be the isometric isomorphism from $C_0(G) \otimes_p A$ onto $C_0(G) \otimes_p \pi_0(A)$ by sending $f \otimes a$ to $f \otimes \pi_0(a)$ for all $f \in C_0(G)$ and $a \in A$. Hence, the homomorphism $(\mathrm{Id}_{C_0(G)} \otimes \pi_0) \circ \varphi_2 \circ \rho^{-1} : F_\lambda^p(\hat{G}) \otimes_p \pi_0(A) \to C_0(G) \otimes_p \pi_0(A)$ sends $\varphi \otimes \pi_0(a)$ to $\Gamma_p(\varphi) \otimes \pi_0(a)$ for all $\varphi \in F_\lambda^p(\hat{G})$. Hence $(\mathrm{Id}_{C_0(G)} \otimes \pi_0) \circ \varphi_2 \circ \rho^{-1} = \Gamma_p \otimes \mathrm{id}_{\pi_0(A)}$. To show φ_2 is an injective homomorphism, it suffices to show that $\Gamma_p \otimes \mathrm{id}_{\pi_0(A)}$ is an injective homomorphism.

If $\xi \in L^p(\hat{G} \times X, \mu \times \nu)$, then $\xi(x)(\gamma) := \xi(\gamma, x)$ defines an element $\xi(x) \in L^p(\hat{G}, \mu)$. Since \hat{G} is a compact Abelian group and $\mu(\hat{G}) = 1$, by [21, Exercise 5 (a)], it follows that $\xi(x) \in L^1(G, \mu) \subseteq F_{\lambda}^p(\hat{G})$. Let $\hat{\xi}(t, x) := \Gamma_p(\xi(x))(t)$. If $F = \sum_{i=1}^m f_i \otimes \pi_0(a_i)$ and $\xi = \sum_{j=1}^n g_j \otimes h_j$, where $f_i \in C(\hat{G})$, $a_i \in A$, $g_j \in L^p(\hat{G}, \mu)$ and $h_j \in L^p(X, \nu)$, then a direct computation shows that

$$\Gamma_p \otimes \mathrm{id}_{\pi_0(A)}(F)(\widehat{\xi}) = (\sum_{i=1}^m \Gamma_p(f_i) \otimes \pi_0(a_i))(\sum_{j=1}^n \Gamma_p(g_j) \otimes h_j)$$
$$= \sum_{i,j} \Gamma_p(f_i * g_j) \otimes \pi_0(a_i)h_j = \widehat{F(\xi)}.$$

Note that the elements of the form $\sum_{i=1}^{m} f_i \otimes \pi_0(a_i)$ are dense in $F_{\lambda}^p(\hat{G}) \otimes_p \pi_0(A)$ and the elements of form $\sum_{j=1}^{n} g_j \otimes h_j$ are dense in $L^p(\hat{G} \times X, \mu \times \nu)$. Since $\Gamma_p \otimes \mathrm{id}_{\pi_0(A)}$ is contractive, it follows that

$$(\Gamma_p \otimes \mathrm{id}_A)(F)(\widehat{\xi}) = \widehat{F(\xi)}$$
 for all $F \in F_\lambda^p(\widehat{G}) \otimes_p A$, $\xi \in L^p(\widehat{G} \times X, \mu \times \nu)$. (9)

For a proof of Claim 2 by contradiction, we assume that $\Gamma_p \otimes \operatorname{id}_{\pi_0(A)}(F) = 0$ for some nonzero $F \in F_{\lambda}^p(\hat{G}) \otimes_p \pi_0(A)$. Then there exists $\xi \in L^p(\hat{G} \times X)$ such that $F(\xi) \neq 0$. By (9), we have $\Gamma_p \otimes \operatorname{id}_{\pi_0(A)}(F)(\widehat{\xi}) = 0$. This implies $\widehat{F(\xi)} = 0$. So, for $x \in X$ and $t \in G$, we have $\widehat{F(\xi)}(t,x) = \Gamma_p(F(\xi)(x))(t) = 0$. Then $\Gamma_p(F(\xi)(x)) = 0$. Since Γ_p is injective, we have $F(\xi) = 0$, which is a contradiction. This proves Claim 2.

If G is a locally compact Abelian group, then G is finite if and only if its dual group \hat{G} is finite (see [5, Theorem 9.15]). By Theorem 3.4, the Gelfand transformation $\Gamma_p: F_{\lambda}^p(\hat{G}) \to C_0(G)$ is an isomorphism if and only if either p=2 or G is finite. If $p \in [1,\infty) \setminus \{2\}$, then φ_2 is an isomorphism if and only if G is finite.

Now assume that p=2. Since $\Gamma_2: F_{\lambda}^2(\hat{G}) \to C_0(G)$ is an isometric isomorphism, by [15, Corollary 1.12], $\Gamma_2 \otimes \operatorname{id}_{\pi_0(A)}: F_{\lambda}^2(\hat{G}) \otimes_2 \pi_0(A) \to C_0(G) \otimes_2 \pi_0(A)$ is an isometric isomorphism. \square

The following proposition will be useful in the proof for the injectivity of Φ_2 .

Proposition 3.6 ([16, Propositions 4.8 and 4.9]). Let $p \in [1, \infty)$ and G be a countable discrete group. Then there exist contractive linear maps $\{E_s : s \in G\}$ from $F^p_{\lambda}(G, A, \alpha)$ to A satisfying

(i) for any $a \in A$, $s, t \in G$,

$$E_s(a\delta_t) = \begin{cases} a, & s = t, \\ 0, & s \neq t, \end{cases}$$

where δ_s is the function in $C_c(G)$ which is 1 at s and zero elsewhere;

(ii) for any $f \in F_{\lambda}^{p}(G, A, \alpha)$, f = 0 if and only if $E_{s}(f) = 0$ for all $s \in G$.

Proof of Proposition 3.3. We first prove two key claims.

Claim 1. $\Phi_2 = \varphi_2 \rtimes id$.

The Gelfand transformation $\Gamma_p: F^p_{\lambda}(\hat{G}) \to C_0(G)$ sends $\varphi \in L^1(\hat{G})$ to its Fourier transformation $\hat{\varphi} \in C_0(G)$ which is given by $\hat{\varphi}(t) = \int_{\hat{G}} \varphi(\gamma) \overline{\gamma(t)} d\mu(\gamma)$. Hence the homomorphism $\varphi_2: F^p(\hat{G}, A, \beta) \to C_0(G, A)$ satisfies

$$\varphi_2(\psi)(t) = \int_{\hat{G}} \psi(\gamma) \overline{\gamma(t)} d\mu(\gamma)$$

for all $\psi \in C(\hat{G}, A)$. Since

$$\varphi_{2}((\hat{\beta} \otimes \alpha)_{s}(\psi))(t) = \int_{\hat{G}} (\hat{\beta} \otimes \alpha)_{s}(\psi)(\gamma) \overline{\gamma(t)} d\mu(\gamma)$$

$$= \int_{\hat{G}} \gamma(s) \alpha_{s}(\psi(\gamma)) \overline{\gamma(t)} d\mu(\gamma)$$

$$= \alpha_{s}(\int_{\hat{G}} \psi(\gamma) \overline{\gamma(s^{-1}t)} d\mu(\gamma))$$

$$= (\operatorname{lt} \otimes \alpha)_{s}(\varphi_{2}(\psi))(t),$$

it follows that φ_2 is equivariant.

Let L be a nondegenerate σ -finite contractive representation of $F^p(G, C_0(G, A), \operatorname{lt} \otimes \alpha)$. Then, by Theorem 2.1, there exists a nondegenerate σ -finite contractive covariant representation (π, v) of $(G, C_0(G, A), \operatorname{lt} \otimes \alpha)$ such that $L = \pi \rtimes v$. Since $\varphi_2(F^p(\hat{G}, A, \beta))$ is dense in $C_0(G, A)$, it follows that $(\pi \circ \varphi_2, v)$ is a nondegenerate σ -finite contractive covariant representation of $(G, F^p(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha)$. By the universal property of $F^p(G, F^p(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha)$ (see [16, Theorem 3.6]), there exists a contractive homomorphism $\varphi_2 \rtimes \operatorname{id} : F^p(G, F^p(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha) \to F^p(G, C_0(G, A), \operatorname{lt} \otimes \alpha)$. For each $F \in C_c(G \times \hat{G}, A)$, we have $\lambda_F(s) \in C(\hat{G}, A, \beta) \subseteq F^p(\hat{G}, A, \beta)$ and $(\varphi_2 \rtimes \operatorname{id})(F)(s) = \varphi_2(\lambda_F(s))$. Hence

$$(\varphi_2 \rtimes \mathrm{id})(F)(s,t) = \int_{\hat{G}} \lambda_F(s)(\gamma) \overline{\gamma(t)} d\mu(\gamma) = \int_{\hat{G}} F(s,\gamma) \overline{\gamma(t)} d\mu(\gamma).$$

This implies that $\Phi_2 = \varphi_2 \rtimes id$.

Claim 2. Φ_2 is an injective homomorphism.

Since G is amenable, by [18, Theorem 7.1], there are two isometric isomorphisms π_1 : $F^p(G, F^p(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha) \to F^p_{\lambda}(G, F^p(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha)$ and $\pi_2 : F^p(G, C_0(G, A), \operatorname{lt} \otimes \alpha) \to F^p_{\lambda}(G, C_0(G, A), \operatorname{lt} \otimes \alpha)$.

For each $s \in G$, let E_s be the linear map as in Proposition 3.6. Then we have the following commutative diagram

$$F^{p}(G, F^{p}(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha) \xrightarrow{\Phi_{2}} F^{p}(G, C_{0}(G, A), \operatorname{lt} \otimes \alpha)$$

$$\downarrow^{\pi_{1}} \qquad \qquad \downarrow^{\pi_{2}}$$

$$F^{p}_{\lambda}(G, F^{p}(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha) \qquad F^{p}_{\lambda}(G, C_{0}(G, A), \operatorname{lt} \otimes \alpha)$$

$$\downarrow^{E_{s}} \qquad \qquad \downarrow^{E_{s}}$$

$$F^{p}(\hat{G}, A, \beta) \xrightarrow{\varphi_{2}} C_{0}(G, A).$$

Choose $F \in F^p(G, F^p(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha)$ such that $\Phi_2(F) = 0$. Then we have $\varphi_2 \circ E_s(\pi_1(F)) = 0$. Since φ_2 is injective, we get that $E_s(\pi_1(F)) = 0$ for all $s \in G$. By Proposition 3.6, we have $\pi_1(F) = 0$. This will imply that F = 0. This proves Claim 2.

If $p \in [1, \infty) \setminus \{2\}$, then, by Lemma 3.5, φ_2 is an isomorphism if and only if G is finite. Hence Φ_2 is an isomorphism if and only if G is finite.

Now assume that p=2. We have proved that Φ_2 is a contractive homomorphism. Since φ_2 is an equivariant isometric isomorphism, it follows that φ_2^{-1} is also an equivariant isometric isomorphism. By the universal property of $F^2(G, C_0(G, A), \operatorname{lt} \otimes \alpha)$ (see [16, Theorem 3.6]), there exists a contractive homomorphism $\varphi_2^{-1} \rtimes \operatorname{id} : F^2(G, C_0(G, A), \operatorname{lt} \otimes \alpha) \to F^2(G, F^2(\hat{G}, A, \beta), \hat{\beta} \otimes \alpha)$. Then, by the Fourier inversion formula and a direct calculation similar to that in the proof of Claim 1, one can check that Φ_2 is the inverse of $\varphi_2^{-1} \rtimes \operatorname{id}$. Hence Φ_2 is an isometric isomorphism.

3.3. The isometry property of Φ_3 .

The aim of this subsection is to prove Φ_3 is an isometric isomorphism and $\Phi_3 \circ \Phi_2 \circ \Phi_1$ is equivariant for the double dual action $\hat{\alpha}$ of G on $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$ and the action $(\operatorname{rt} \otimes \alpha) \otimes \operatorname{id}$ of G on $F^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$.

Lemma 3.7. Let $\Phi_3: F^p(G, C_0(G, A), \operatorname{lt} \otimes \alpha) \to F^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$ be defined in (4) under the assumption of Theorem 1.2. Then Φ_3 is an isometric isomorphism.

Proof. We can define an isometric isomorphism $\varphi_3: C_0(G,A) \to C_0(G,A)$ by $\varphi_3(f)(t) := \alpha_t^{-1}(f(t))$. Then

$$\varphi_3((\operatorname{lt} \otimes \alpha)_s(f))(t)) = \alpha_t^{-1}(\alpha_s(f(s^{-1}t)))$$

$$= \alpha_{s^{-1}t}^{-1}(f(s^{-1}t))$$

$$= (\operatorname{lt} \otimes \operatorname{id})_s(\varphi_3(f))(t).$$

So φ_3 is equivariant. By the universal property of $F^p(G, C_0(G, A), lt \otimes \alpha)$ and $F^p(G, C_0(G, A), lt \otimes id)$ (see [16, Theorem 3.6]), using a similar argument in Lemma 3.3, one can show that $\Phi_3 = \varphi_3 \rtimes id$ is an isometric isomorphism.

For the reader's convenience, we recall the double dual action $\hat{\alpha}$ of G on $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$ and the action $(\operatorname{rt}\otimes\alpha)\otimes\operatorname{id}$ of G on $F^p(G, C_0(G, A), \operatorname{lt}\otimes\operatorname{id})$. The double dual action $\hat{\alpha}$ of G on $C_c(\hat{G}\times G, A)\subseteq F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$ is given by $\hat{\alpha}_t(F)(\gamma, s):=\overline{\gamma(t)}F(\gamma, s)$ for all $t\in G$. Let rt denote the right-translation of G on $C_0(G)$, that is, $(\operatorname{rt})_t f(s):=f(st)$ for all $f\in C_0(G)$. Then we get an L^p operator algebra dynamical system $\operatorname{rt}\otimes\alpha:G\to\operatorname{Aut}(C_0(G,A))$, where $(\operatorname{rt}\otimes\alpha)_t f(s):=\alpha_t(f(st))$ for all $f\in C_0(G,A)$ and $t\in G$. Thus we get an L^p operator algebra dynamical system $(\operatorname{rt}\otimes\alpha)\otimes\operatorname{id}:G\to\operatorname{Aut}(F^p(G,C_0(G,A),\operatorname{lt}\otimes\operatorname{id}))$, where $((\operatorname{rt}\otimes\alpha)\otimes\operatorname{id})_r F(s,t):=\alpha_r(F(s,tr))$ for all $F\in C_c(G\times G,A)\subseteq F^p(G,C_0(G,A),\operatorname{lt}\otimes\operatorname{id})$ and $f\in G$.

Lemma 3.8. The homomorphism

$$\Phi_3 \circ \Phi_2 \circ \Phi_1 : F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha}) \to F^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$$

is equivariant for the double dual action $\hat{\alpha}$ of G on $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$ and the action $(\operatorname{rt} \otimes \alpha) \otimes \operatorname{id}$ of G on $F^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$.

Proof. For $F \in C_c(\hat{G} \times G, A)$ and $s, t \in G$, we have

$$\begin{split} \Phi_3 \circ \Phi_2 \circ \Phi_1(F)(s,t) &= \alpha_t^{-1}(\Phi_2 \circ \Phi_1(F)(s,t)) \\ &= \alpha_t^{-1}(\int_{\hat{G}} \Phi_1(F)(s,\gamma) \overline{\gamma(t)} d\mu(\gamma)) \\ &= \int_{\hat{G}} \alpha_t^{-1}(F(\gamma,s)) \overline{\gamma(s^{-1}t)} d\mu(\gamma). \end{split}$$

Then, for each $r \in G$, we have

$$\Phi_{3} \circ \Phi_{2} \circ \Phi_{1}(\hat{\alpha}_{r}(F))(s,t) = \int_{\hat{G}} \alpha_{t}^{-1}(\hat{\alpha}_{r}(F)(\gamma,s)) \overline{\gamma(s^{-1}t)} d\mu(\gamma)
= \int_{\hat{G}} \alpha_{t}^{-1}(F(\gamma,s)) \overline{\gamma(s^{-1}tr)} d\mu(\gamma)
= \alpha_{r}(\int_{\hat{G}} \alpha_{tr}^{-1}(F(\gamma,s) \overline{\gamma(s^{-1}tr)} d\mu(\gamma))
= \alpha_{r}(\Phi_{3} \circ \Phi_{2} \circ \Phi_{1}(F)(s,tr)).$$

Hence $(\Phi_3 \circ \Phi_2 \circ \Phi_1) \circ \hat{\hat{\alpha}}_r = ((\operatorname{rt} \otimes \alpha) \otimes \operatorname{id})_r \circ (\Phi_3 \circ \Phi_2 \circ \Phi_1).$

3.4. The isometry property of Φ_4 .

The aim of this subsection is to prove the following.

Proposition 3.9. Let $\Phi_4: F^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id}) \to \overline{M}_G^p \otimes_p A$ be defined in (5) under the assumption of Theorem 1.2. Then Φ_4 is an isometric isomorphism which is equivariant for the action $(\operatorname{rt} \otimes \alpha) \otimes \operatorname{id}$ of G on $F^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$ and the action $\operatorname{Ad}\rho \otimes \alpha$ of G on

 $\overline{M}_G^p \otimes_p A$. Moreover, the right-hand side equals $\mathcal{K}(l^p(G)) \otimes_p A$ if and only if p > 1 or G is finite.

To prove Proposition 3.9, we make some preparation.

Lemma 3.10. Let G be a countable discrete group and A be a separable unital L^p operator algebra. If A has unique L^p operator matrix norms, then so does $C_0(G, A)$.

Proof. Let $\pi: C_0(G, A) \to \mathcal{B}(L^p(X, \mu))$ and $\sigma: C_0(G, A) \to \mathcal{B}(L^p(Y, \nu))$ be two isometric representations. By [19, Lemma 5.15], $C_c(G, A)$ has unique L^p operator norms. For any $F \in M_n(C_c(G, A))$, we have $\|\mathrm{id}_{M_n} \otimes \pi(F)\| = \|\mathrm{id}_{M_n} \otimes \sigma(F)\|$. Note that $C_c(G, A)$ is dense in $C_0(G, A)$. Then $C_0(G, A)$ has unique L^p operator matrix norms.

The following result concerning compact operators on L^p spaces will be useful in the proof of Proposition 3.9.

Theorem 3.11 ([8, Theorem 4.3]). Let $p \in [1, \infty)$, G be a discrete group, and $\operatorname{lt}: G \to \operatorname{Aut}(C_0(G))$ be the isometric action induced by the left translation of G on itself. Then there are natural isometric isomorphisms:

$$F^p(G, C_0(G), \mathrm{lt}) \xrightarrow{\kappa} F_{\lambda}^p(G, C_0(G), \mathrm{lt}) \xrightarrow{\iota} \overline{M}_G^p$$

Moreover, the right-hand side equals $\mathcal{K}(l^p(G))$ when p > 1, and is strictly smaller than $\mathcal{K}(l^1(G))$ when p = 1 and G is infinite.

Proof of Proposition 3.9. We give the proof by proving two key claims.

Claim 1. $\Phi_4: F^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id}) \to \overline{M}_G^p \otimes_p A$ is an isometric isomorphism.

Since G is amenable, by [18, Theorem 7.1]), $F^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$ is isometrically isomorphic to $F^p_{\lambda}(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$; we denote by π this isomorphism. It suffices to show that $F^p_{\lambda}(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$ is isometrically isomorphic to $\overline{M}^p_G \otimes_p A$.

Firstly, we construct an isometric representation of $\overline{M}_G^p \otimes_p A$ and $F_\lambda^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$ by using the property of unique L^p operator matrix norms. Since A has unique L^p operator matrix norms, by Lemma 2.4, the norm on spatial L^p operator tensor product $\overline{M}_G^p \otimes_p A$ is independent of the choice of isometric representation of A. Let $\pi_0 : A \to \mathcal{B}(E)$ be a nondegenerate σ -finite isometric representation of A. Then there is a natural isometric representation of $\overline{M}_G^p \otimes_p A$ on $l^p(G, E)$.

Furthermore, there exists a canonical nondegenerate σ -finite isometric representation π_1 of $C_0(G,A)$ on $l^p(G,E)$ which maps $f\otimes a$ to $M_f\otimes \pi_0(a)$, where $f\in C_0(G), a\in A$, and M_f is the multiplication operator on $l^p(G)$. Let $(\pi,\lambda_p^{l^p(G,E)})$ be its associated regular covariant representation. Then, by Lemmas 2.5 and 3.10, the integrated form $\pi\rtimes\lambda_p^{l^p(G,E)}$ is an isometric representation of $F_{\lambda}^p(G,C_0(G,A),\operatorname{lt}\otimes\operatorname{id})$.

Now we are going to construct an isometric isomorphism from $F^p_{\lambda}(G, C_0(G, A), lt \otimes id)$ to $\overline{M}^p_G \otimes_p A$.

We define an invertible isometry $V: l^p(G) \otimes_p l^p(G) \otimes_p E \to l^p(G) \otimes_p l^p(G) \otimes_p E$ as $\delta_s \otimes \delta_t \otimes \xi \mapsto \delta_{st} \otimes \delta_t \otimes \xi$, where δ_s is the canonical basis of $l^p(G)$ for all $s \in G$ and $\xi \in E$.

We denote by $I_{l^p(G)}$ and I_E the identity operator on $l^p(G)$ and E, respectively. A direct computation shows that

$$V\pi(f\otimes a)(\delta_s\otimes \delta_t\otimes \xi) = V(\delta_s\otimes \pi_0((\operatorname{lt}\otimes \operatorname{id})_{s^{-1}}(f\otimes a))(\delta_t\otimes \xi))$$

$$= V(\delta_s\otimes M_{(\operatorname{lt})_{s^{-1}}(f)}\delta_t\otimes a\xi))$$

$$= V(\delta_s\otimes f(st)\delta_t\otimes a\xi)$$

$$= f(st)\delta_{st}\otimes \delta_t\otimes a\xi$$

$$= (M_f\otimes I_{l^p(G)}\otimes a)(\delta_{st}\otimes \delta_t\otimes \xi)$$

$$= (M_f\otimes I_{l^p(G)}\otimes a)(V(\delta_s\otimes \delta_t\otimes \xi)).$$

It follows that

$$V\pi(f\otimes a)V^{-1} = M_f\otimes I_{l^p(G)}\otimes a. \tag{10}$$

It is easy to check that $\lambda_p^{l^p(G,E)} = \lambda_p \otimes I_{l^p(G)} \otimes I_E$, where $\lambda_p : G \to \mathcal{B}(l^p(G))$ is the left-regular representation of G on $l^p(G)$. For each $r \in G$, one can check that

$$V(\lambda_{p}(r) \otimes I_{l^{p}(G)} \otimes I_{E})V^{-1}(\delta_{s} \otimes \delta_{t} \otimes \xi) = V(\lambda_{p}(r) \otimes I_{l^{p}(G)} \otimes I_{E})(\delta_{st^{-1}} \otimes \delta_{t} \otimes \xi)$$

$$= V(\delta_{rst^{-1}} \otimes \delta_{t} \otimes \xi)$$

$$= \delta_{rs} \otimes \delta_{t} \otimes \xi$$

$$= (\lambda_{p}(r) \otimes I_{l^{p}(G)} \otimes I_{E})(\delta_{s} \otimes \delta_{t} \otimes \xi).$$

It follows that

$$V(\lambda_p(r) \otimes I_{l^p(G)} \otimes I_E)V^{-1} = \lambda_p(r) \otimes I_{l^p(G)} \otimes I_E.$$
(11)

Since $f \in C_0(G)$, by (10) and (11), we have $V(F_{\lambda}^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id}))V^{-1} \cong \overline{M}_G^p \otimes_p \mathbb{C}I_{l^p(G)} \otimes_p A$. Obviously, $\overline{M}_G^p \otimes_p \mathbb{C}I_{l^p(G)} \otimes_p A$ is isometrically isomorphic to $\overline{M}_G^p \otimes_p A$. Since V is an isometry, it follows that $F_{\lambda}^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$ is isometrically isomorphic to $\overline{M}_G^p \otimes_p A$. We denote this isomorphism by φ_4 . It is easy to check that $\Phi_4 = \varphi_4 \circ \pi$. Hence Φ_4 is an isometric isomorphism from $F^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$ onto $\overline{M}_G^p \otimes_p A$. This proves Claim 1.

By Theorem 3.11, $\overline{M}_G^p \otimes_p A$ equals $\mathcal{K}(l^p(G)) \otimes_p A$ if and only if either p > 1 or G is finite.

Claim 2. Φ_4 is equivariant for the action $(\operatorname{rt} \otimes \alpha) \otimes \operatorname{id}$ of G on $F^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$ and the action $\operatorname{Ad}\rho \otimes \alpha$ of G on $\overline{M}_G^p \otimes_p A$.

For $F \in C_c(G \times G, A) \subseteq F^p(G, C_0(G, A), \operatorname{lt} \otimes \operatorname{id})$ and $h \in C_c(G, E) \subseteq l^p(G, E)$, we have $\Phi_4(F)h(t) = \sum_{s \in G} F(s, t)h(s^{-1}t)$. For $r \in G$, $T \in \mathcal{K}(l^p(G))$ and $a \in A$, we have

$$(\operatorname{Ad}\rho \otimes \alpha)_r(T \otimes a) = (\operatorname{Ad}\rho)_r(T) \otimes \alpha_r(a) = (\rho_r \otimes I_E)((I_{l^pG}) \otimes \alpha_r)(T \otimes a))(\rho_{r^{-1}} \otimes I_E). \text{ Hence}$$

$$(\operatorname{Ad}\rho \otimes \alpha)_r(\Phi_4(F))h(t) = (\rho_r \otimes I_E)((I_{l^pG}) \otimes \alpha_r)(\Phi_4(F)))(\rho_{r^{-1}} \otimes I_E)h(t)$$

$$= ((I_{l^pG}) \otimes \alpha_r)(\Phi_4(F)))(\rho_{r^{-1}} \otimes I_E)h(tr)$$

$$= \sum_{s \in G} \alpha_r(F(s, tr))(\rho_{r^{-1}} \otimes I_E)h(s^{-1}tr)$$

$$= \sum_{s \in G} \alpha_r(F(s, tr))h(s^{-1}t)$$

$$= \sum_{s \in G} ((\operatorname{rt} \otimes \alpha) \otimes \operatorname{id})_r(F)(s, t)h(s^{-1}t)$$

$$= \Phi_4(((\operatorname{rt} \otimes \alpha) \otimes \operatorname{id})_r(F))h(t).$$

Therefore, $(\mathrm{Ad}\rho\otimes\alpha)_r\circ\Phi_4=\Phi_4\circ((\mathrm{rt}\otimes\alpha)\otimes\mathrm{id})_r$ as required.

Proof of Theorem 1.2. Statements (i), (ii) and (iii) in Theorem 1.2 follow from Propositions 3.1, 3.3, 3.7 and 3.9. By Lemma 3.8 and Proposition 3.9, the map Φ is equivariant for the double dual action $\hat{\alpha}$ of G on $F^p(\hat{G}, F^p(G, A, \alpha), \hat{\alpha})$, and the action $Ad\rho \otimes \alpha$ of G on $\mathcal{K}(l^p(G)) \otimes_p A$.

ACKNOWLEDGEMENTS

The first author is supported by the National Natural Science Foundation of China (grant numbers 12201240, 11971253), the China Postdoctoral Science Foundation (grant number 2022M711310), the Natural Science Foundation of Fujian Province (grant number 2020J05206) and the Scientific Research Project of Putian University (grant number 2020001). The second author is supported by the National Natural Science Foundation of China (grant number 12171195).

References

- [1] B. Blackadar, K-Theory for Operator Algebras, Second edition, Mathematical Sciences Research Institute Publications, 5, Cambridge University Press, Cambridge, 1998.
- [2] D. Blecher and N. C. Phillips, L^p operator algebras with approximate identities I, Pacific J. Math. 303 (2019), no. 2, pp. 401-457.
- [3] N. P. Brown and N. Ozawa, C^* -algebras and finite-dimensional approximations, Graduate Studies in Mathematics, 88, American Mathematical Society, Providence, RI, 2008.
- [4] A. Connes, An analogue of the Thom isomorphism for crossed products of a C^* -algebra by an action of \mathbb{R} , Adv. Math. 39 (1981), pp. 31-55.
- J. B. Conway, A Course in Functional Analysis, Graduate Texts in Mathematics, 96, Springer, New York, 1990.
- [6] Y. Choi, E. Gardella and H. Thiel, Rigidity results for L^p -operator algebras and applications, Preprint (arxiv 1909.03612), 2019.
- [7] E. Gardella and M. Lupini, Representations of étale groupoids on L^p spaces, Adv. Math. 318 (2017), pp. 233-278.
- [8] E. Gardella and H. Thiel, Group algebras acting on L^p spaces, J. Fourier Anal. Appl. 21 (2015), no. 6, pp. 1310-1343.
- [9] E. Gardella and H. Thiel, Quotients of Banach algebras acting on L^p spaces, Adv. Math. 296 (2016), pp. 85-92.

- [10] E. Gardella, H. Thiel, Isomorphisms of algebras of convolution operators, To appear in Ann. Sci. École Nor. Sup, Preprint (arxiv 1809.01585), 2018.
- [11] E. Gardella and H. Thiel, Representations of p-convolution algebras on L^q -spaces. Trans. Amer. Math. Soc. 371 (2019), no. 3, pp. 2207-2236.
- [12] E. Gardella, A modern look at algebras of operators on L^p spaces, Exo. Math. 39 (2021), no. 3, pp. 420-453.
- [13] C. Herz, The theory of *p*-spaces with an application to convolution operators, Trans. Amer. Math. Soc. 154 (1971), pp. 69-82.
- [14] S. Hejazian and S. Pooya, Simple reduced L^p -operator crossed products with unique trace, J. Operator Theory 74 (2015), no. 1, pp. 133-147.
- [15] G. Pisier, Tensor products of C^* -algebras and operator spaces—the Connes-Kirchberg problem, London Mathematical Society Student Texts, 96, Cambridge University Press, Cambridge, 2020.
- [16] N. C. Phillips, Crossed products of L^p operator algebras and the K-theory of Cuntz algebras on L^p spaces, Preprint (arXiv:1309.6406), 2013.
- [17] N. C. Phillips, "Open problems related to operator algebras on L^p spaces", preprint, 2014, https://pdfs.semanticscholar.org/0823/5038ec45079e7721a59021a4492da2c2b1a3.pdf.
- [18] N. C. Phillips, Operator algebras on L^p spaces which "look like" C^* -algebras, 2014, https://pages.uoregon.edu/ncp/Talks/20140527_GPOTS/LpOpAlgs_TalkSummary.pdf.
- [19] N. C. Phillips and M. G. Viola, Classification of L^p AF algebras, Internat. J. Math. 31 (2020), no. 13, 2050088, 41 pp.
- [20] I. Raeburn, On crossed products and Takai duality, Proc. Edinburgh Math. Soc. (2) 31 (1988), pp. 321-330
- [21] W. Rudin, Real and complex analysis, Third edition, McGraw-Hill Book Company, New York, 1987.
- [22] H. Takai, On a duality for crossed products of C^* -algebras, J. Funct. Anal. 19 (1975), pp. 25-39.
- [23] D. Williams, Crossed Products of C*-Algebras, Mathematical Surveys and Monographs, 134, American Mathematical Society, Providence, RI, 2007.
- [24] Z. Wang and Y. Zeng, Gelfand theory of reduced group L^p operator algebra, Ann. Funct. Anal. 13, 14 (2022).

Current address: Department of Mathematics, Jilin University, Changchun 130012, P. R. China

KEY LABORATORY OF APPLIED MATHEMATICS OF FUJIAN PROVINCE UNIVERSITY, SCHOOL OF MATHEMATICS AND FINANCE, PUTIAN UNIVERSITY, PUTIAN 351100, P. R. CHINA

Email address: zwangmath@jlu.edu.cn

DEPARTMENT OF MATHEMATICS, JILIN UNIVERSITY, CHANGCHUN 130012, P. R. CHINA

Email address: zhusen@jlu.edu.cn