

LONG TIME H^1 -STABILITY OF FAST L2-1 $_{\sigma}$ METHOD ON GENERAL NONUNIFORM MESHES FOR SUBDIFFUSION EQUATIONS*

CHAOYU QUAN [†], XU WU [‡], AND JIANG YANG [§]

Abstract. In this work, the global-in-time H^1 -stability of a fast L2-1 $_{\sigma}$ method on general nonuniform meshes is studied for subdiffusion equations, where the convolution kernel in the Caputo fractional derivative is approximated by sum of exponentials. Under some mild restrictions on time stepsize, a bilinear form associated with the fast L2-1 $_{\sigma}$ formula is proved to be positive semidefinite for all time. As a consequence, the uniform global-in-time H^1 -stability of the fast L2-1 $_{\sigma}$ schemes can be derived for both linear and semilinear subdiffusion equations, in the sense that the H^1 -norm is uniformly bounded as the time tends to infinity. To the best of our knowledge, this appears to be the first work for the global-in-time H^1 -stability of fast L2-1 $_{\sigma}$ scheme on general nonuniform meshes for subdiffusion equations. Moreover, the sharp finite time H^1 -error estimate for the fast L2-1 $_{\sigma}$ schemes is reproved based on more delicate analysis of coefficients where the restriction on time step ratios is relaxed comparing to existing works.

Key words. subdiffusion equations, fast L2-1 $_{\sigma}$ method, global-in-time H^1 -stability, nonuniform meshes, sharp error estimate

AMS subject classifications. 35R11, 65M12

1. Introduction. In the past decade, the time-fractional diffusion equation [29, 7] attract much attention from researchers in theoretical and numerical analysis. Developing stable and accurate numerical method has been a core issue in this longstanding and active research area. Many well-known schemes have been developed and analyzed, including the piecewise polynomial interpolation methods (such as L1, L2-1 $_{\sigma}$ and L2 schemes), discontinuous Galerkin (dG) methods, and convolution quadrature (CQ) methods under the assumption that the solution is sufficiently smooth; see [17, 39, 2, 6, 28, 9, 30]. However, the solutions to time-fractional problems typically admit weak singularities. This inspires researchers to design numerical schemes to overcome this singularity problem. For example, the L1, L2-1 $_{\sigma}$, L2 and dG methods on the graded meshes or general nonuniform meshes have been developed to overcome the singularities of time-fractional diffusion equation in [38, 14, 4, 16, 15, 20, 21, 22, 30, 31]. The L1, L2 and CQ schemes with proper initial correction using uniform step size can regain the desired high-order accuracy for linear subdiffusion problems with singularities; see [44, 42, 9, 11]. Recently, there are also plenty of works on the stability and convergence of numerical methods for semilinear time-fractional equations; see [10, 1, 41, 23, 5, 18]. Particularly, the energy dissipation property for a class of fractional phase field models has been a core issue [23, 5, 12], which essentially amounts to uniform global-in-time H^1 -stability. This motivates us to study the long time H^1 -stability of various numerical schemes for

*Submitted in Nov 2022.

[†]SUSTech International Center for Mathematics, Southern University of Science and Technology, Shenzhen, P.R. China (quancy@sustech.edu.cn).

[‡]Department of Mathematics, Harbin Institute of Technology, Harbin 150001, China; Department of Mathematics, Southern University of Science and Technology, Shenzhen, China (11849596@mail.sustech.edu.cn).

[§]Department of Mathematics, SUSTech International Center for Mathematics & National Center for Applied Mathematics Shenzhen (NCAMS), Guangdong Provincial Key Laboratory of Computational Science and Material Design, Southern University of Science and Technology, Shenzhen, China (yangj7@sustech.edu.cn).

semilinear time-fractional equations.

In addition to the weak singularity problem the accuracy of numerical solutions, another important issue is the storage problem due to the nonlocality of the fractional derivatives. Precisely speaking, all the aforementioned works require $\mathcal{O}(N)$ storage and $\mathcal{O}(N^2)$ computational cost for N time steps, which are too expensive. Then the fast algorithms to reduce computational storage and cost have been developed. Lubich-Schädle [27] propose a new algorithm for the evaluation of convolution integrals for wave problems with non-reflecting boundary conditions. Based on the block triangular Toeplitz matrix or block triangular Toeplitz-like matrix, Ke-Ng-Sun [13] and Lu-Peng-Sun [26] propose a fast approximation to time-fractional derivative. Baffet-Hesthaven [3] compress the kernel in the Laplace domain and obtained a sum-of-poles approximation for the Laplace transform of the kernel. Jiang-Zhang-Zhang-Zhang [8] propose the sum-of-exponentials (SOE) approximation to speed up the evaluation of weakly singular Caputo derivative kernel, which significantly reduces the computational storage and cost. Along this way, the fast L1 and L2-1 $_{\sigma}$ schemes are proposed and analyzed on both uniform and nonuniform meshes in [24, 43, 19, 25, 36, 37], and Zhu-Xu [45] study the fast L2 method on uniform meshes based on the SOE approximation. The SOE approximation allows us to reduce the storage and overall computational cost from $\mathcal{O}(N)$ and $\mathcal{O}(N^2)$ for the L1, L2-1 $_{\sigma}$, and L2 scheme to $\mathcal{O}(N_q)$ and $\mathcal{O}(N_q N)$ for the fast method, respectively, with N being the number of the time steps and N_q being the number of the quadrature nodes.

In this work, we apply the fast L2-1 $_{\sigma}$ scheme on general nonuniform meshes for subdiffusion equation with homogeneous Dirichlet boundary condition, where the L2-1 $_{\sigma}$ method [2] and the SOE technique [8, 43] are combined to approximate the Caputo fractional derivative. Given a fractional order $\alpha \in (0, 1)$, an absolute tolerance error $\varepsilon \ll 1$, a cut-off time $\Delta t > 0$ and a fixed time $T_{\text{soe}} > \Delta t$, it is well-known that there exists an SOE approximation to the convolution kernel, see Lemma 2.1 for details. Denote the fast L2-1 $_{\sigma}$ operator based on this SOE approximation by $F_k^{\alpha,*}$. In [23], the authors state that the positive semidefiniteness of the bilinear given as below form with the standard L2-1 $_{\sigma}$ formula is an open problem,

$$\mathcal{B}_n(v, w) = \sum_{k=1}^n \langle F_k^{\alpha,*} v, \delta_k w \rangle, \quad \delta_k w := w^k - w^{k-1}, \quad n \geq 1,$$

which is solved recently in [34] by some of us. In this work, we generalize this positive semidefiniteness to the fast L2-1 $_{\sigma}$ formula. We prove that above bilinear form is positive semidefinite if the time steps $\{\tau_k\}_{k \geq 1}$ satisfy (see Theorem 3.2)

$$\begin{aligned} \rho_k := \tau_k / \tau_{k-1} &\geq 0.475329, \quad \varepsilon \leq \min_{k \geq 1} \frac{1}{5(1-\alpha)(\sigma\tau_k)^{\alpha}}, \quad \Delta t \leq \min_{k \geq 2} \sigma\tau_k, \\ T_{\text{soe}} &\geq \max_{k \geq 2} (\sigma\tau_{k+1} + \tau_k). \end{aligned}$$

Based on this result, we prove that the H^1 -stability of the fast L2-1 $_{\sigma}$ scheme for linear subdiffusion equation holds true even when the time t is larger than T_{soe} and tends to infinity, (see Theorem 4.1), while for semilinear subdiffusion equations, an additional restriction on τ_k should be satisfied to ensure similar stability (see Theorem 4.2). To the best of our knowledge, this positive semidefiniteness result as well as the corresponding global-in-time H^1 -stability is new. On the other hand, the H^1 -error estimate of the fast L2-1 $_{\sigma}$ method have been well studied for subdiffusion equations on general nonuniform meshes in [19, 25]. But we still repeat the proof of the finite

time optimal convergence rate in H^1 -norm and reduce the restriction on time step ratios from $\rho_k \geq 2/3$ to $\rho_k \geq 0.475329$.

This work is organized as follows. In Section 2, the L2-1 $_{\sigma}$ formula and the SOE approximation are recalled, and the fast L2-1 $_{\sigma}$ formula is then derived. In Section 3, we prove the positive semidefiniteness of the bilinear form \mathcal{B}_n under some mild restrictions on the time steps. In Section 4, we establish the global-in-time H^1 -stability of the L2-1 $_{\sigma}$ scheme for both the linear and semilinear subdiffusion equations, based on the above positive semidefiniteness result. In Section 5, the sharp finite time H^1 -error estimate of the fast L2-1 $_{\sigma}$ scheme for the subdiffusion equation is provided. In Section 6, we provide some numerical results to verify our theorems. Some conclusions are provided in Section 7.

2. Fast L2-1 $_{\sigma}$ formula. We first recall the L2-1 $_{\sigma}$ method and then the fast L2-1 $_{\sigma}$ method on an arbitrary nonuniform mesh for approximating the Caputo fractional derivative defined by

$$\partial_t^{\alpha} u = \frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{u'(s)}{(t-s)^{\alpha}} ds.$$

In the following content, we consider a nonuniform time mesh $0 = t_0 < t_1 < \dots < t_k \dots$ with time step $\tau_k = t_k - t_{k-1}$.

2.1. L2-1 $_{\sigma}$ formula. We first briefly recall the L2-1 $_{\sigma}$ scheme, which is constructed in [2]. The fractional derivative $\partial_t^{\alpha} u(t_k^*)$ is approximated by the L2-1 $_{\sigma}$ formula

$$(2.1) \quad L_k^{\alpha,*} u = \frac{1}{\Gamma(1-\alpha)} \left(\sum_{j=1}^{k-1} \int_{t_{j-1}}^{t_j} \frac{\partial_s \Pi_{2,j}(s)}{(t_k^* - s)^{\alpha}} ds + \int_{t_{k-1}}^{t_k^*} \frac{\partial_s \Pi_{1,k}(s)}{(t_k^* - s)^{\alpha}} ds \right) \\ = \frac{1}{\Gamma(1-\alpha)} \left(\sum_{j=1}^{k-1} (a_j^{(k)} u^{j-1} + b_j^{(k)} u^j + c_j^{(k)} u^{j+1}) \right) + \frac{\sigma^{1-\alpha} (u^k - u^{k-1})}{\Gamma(2-\alpha) \tau_k^{\alpha}},$$

where $t_k^* := t_{k-1} + \sigma \tau_k$, $\sigma = 1 - \alpha/2$,

$$\Pi_{2,j}(t) := \frac{(t-t_j)(t-t_{j+1})}{(t_{j-1}-t_j)(t_{j-1}-t_{j+1})} u^{j-1} + \frac{(t-t_{j-1})(t-t_{j+1})}{(t_j-t_{j-1})(t_j-t_{j+1})} u^j \\ + \frac{(t-t_{j-1})(t-t_j)}{(t_{j+1}-t_{j-1})(t_{j+1}-t_j)} u^{j+1}, \\ \Pi_{1,k}(t) := \frac{t-t_k}{t_{k-1}-t_k} u^{k-1} + \frac{t-t_{k-1}}{t_k-t_{k-1}} u^k,$$

for $1 \leq j \leq k-1$, and

$$(2.2) \quad a_j^{(k)} = \int_{t_{j-1}}^{t_j} \frac{2s-t_j-t_{j+1}}{\tau_j(\tau_j+\tau_{j+1})} \frac{1}{(t_k^*-s)^{\alpha}} ds, \quad b_j^{(k)} = \int_{t_{j-1}}^{t_j} \frac{2s-t_{j-1}-t_{j+1}}{-\tau_j\tau_{j+1}} \frac{1}{(t_k^*-s)^{\alpha}} ds, \\ c_j^{(k)} = \int_{t_{j-1}}^{t_j} \frac{2s-t_{j-1}-t_j}{\tau_{j+1}(\tau_j+\tau_{j+1})} \frac{1}{(t_k^*-s)^{\alpha}} ds.$$

The coefficients satisfy that $a_j^{(k)} + b_j^{(k)} + c_j^{(k)} = 0$, for $1 \leq j \leq k-1$. Then the discrete fractional derivative in (2.1) can be reformulated as

$$(2.3) \quad L_k^{\alpha,*} u = \frac{1}{\Gamma(1-\alpha)} \left(c_{k-1}^{(k)} \delta_k u - a_1^{(k)} \delta_1 u + \sum_{j=2}^{k-1} d_j^{(k)} \delta_j u \right) + \frac{\sigma^{1-\alpha}}{\Gamma(2-\alpha) \tau_k^{\alpha}} \delta_k u,$$

where $\delta_j u = u^j - u^{j-1}$ and $d_j^{(k)} := c_{j-1}^{(k)} - a_j^{(k)}$. Here we make a convention that $a_1^{(1)} = 0$ and $c_0^{(1)} = 0$; see [34] for details.

2.2. Fast L2-1 $_\sigma$ formula. Note that the right-hand side of (2.3) involves a sum of all previous solutions $\{u^j\}_{j=0}^k$, which reflects the memory effect of the non-local fractional operator. This leads to expensive computational cost as the number of time steps increases. To improve the efficiency, one can use the following SOE approximation result:

LEMMA 2.1 ([8, 24]). *For the given $\alpha \in (0, 1)$, an absolute tolerance error $\varepsilon \ll 1$, a cut-off time $\Delta t > 0$ and a fixed time $T_{\text{soe}} > \Delta t$, there exists a positive integer N_q , positive quadrature nodes θ^ℓ and corresponding positive weights ϖ^ℓ ($1 \leq \ell \leq N_q$) such that*

$$\left| t^{-\alpha} - \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell t} \right| \leq \varepsilon, \quad \forall t \in [\Delta t, T_{\text{soe}}],$$

where the number N_q of quadrature nodes satisfies

$$N_q = \mathcal{O}\left(\log \frac{1}{\varepsilon} \left(\log \log \frac{1}{\varepsilon} + \log \frac{T_{\text{soe}}}{\Delta t}\right) + \log \frac{1}{\Delta t} \left(\log \log \frac{1}{\varepsilon} + \log \frac{1}{\Delta t}\right)\right).$$

REMARK 2.1. *Note that in this work, T_{soe} is not necessarily the total time. In the later H^1 -stability analysis, the total time can go to infinity for fixed T_{soe} .*

The L2-1 $_\sigma$ formula $L_k^{\alpha,*} u$ is split into a history part $H(t_k^*)$ and a local part $L(t_k^*)$ as:

$$H(t_k^*) := \frac{1}{\Gamma(1-\alpha)} \int_0^{t_{k-1}^*} \frac{\partial_s \Pi_2(s)}{(t_k^* - s)^\alpha} ds, \quad L(t_k^*) := \frac{1}{\Gamma(1-\alpha)} \int_{t_{k-1}^*}^{t_k^*} \frac{\partial_s \Pi_{1,k}(s)}{(t_k^* - s)^\alpha} ds,$$

where $\Pi_2(s) = \Pi_{2,j}(s)$ on the interval (t_{j-1}, t_j) , $1 \leq j \leq k-1$. The fast L2-1 $_\sigma$ formula is obtained by replacing $(t_k^* - s)^{-\alpha}$ in $H(t_k^*)$ with the SOE approximation in Lemma 2.1:

$$\begin{aligned} (2.4) \quad F_k^{\alpha,*} u &= \frac{1}{\Gamma(1-\alpha)} \left(\int_0^{t_{k-1}^*} \partial_s \Pi_2(s) \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell (t_k^* - s)} ds + \int_{t_{k-1}^*}^{t_k^*} \frac{\partial_s \Pi_{1,k}(s)}{(t_k^* - s)^\alpha} ds \right) \\ &= \frac{\sum_{\ell=1}^{N_q} \varpi^\ell H^\ell(t_k^*)}{\Gamma(1-\alpha)} + \frac{\sigma^{1-\alpha} (u^k - u^{k-1})}{\Gamma(2-\alpha) \tau_k^\alpha}, \end{aligned}$$

where $H^\ell(t_k^*) = \int_0^{t_{k-1}^*} \partial_s \Pi_2(s) e^{-\theta^\ell (t_k^* - s)} ds$ for $1 \leq \ell \leq N_q$. $H^\ell(t_k^*)$ can be computed by the following recurrence formulation:

$$\begin{aligned} H^\ell(t_k^*) &= \int_0^{t_{k-2}^*} \partial_s \Pi_2(s) e^{-\theta^\ell (t_k^* - s)} ds + \int_{t_{k-2}^*}^{t_{k-1}^*} \partial_s \Pi_{2,k-1}(s) e^{-\theta^\ell (t_k^* - s)} ds \\ &= e^{-\theta^\ell [(1-\sigma)\tau_{k-1} + \sigma\tau_k]} H^\ell(t_{k-1}^*) + \hat{a}_{k-1}^{k,\ell} u^{k-2} + \hat{b}_{k-1}^{k,\ell} u^{k-1} + \hat{c}_{k-1}^{k,\ell} u^k, \end{aligned}$$

with $H^\ell(t_1^*) = 0$ and for $k \geq 2$

$$(2.5) \quad \hat{a}_{k-1}^{k,\ell} = \int_{t_{k-2}^*}^{t_{k-1}^*} \frac{2s - t_{k-1} - t_k}{\tau_{k-1}(\tau_{k-1} + \tau_k)} e^{-\theta^\ell (t_k^* - s)} ds, \quad \hat{b}_{k-1}^{k,\ell} = \int_{t_{k-2}^*}^{t_{k-1}^*} \frac{2s - t_{k-2} - t_k}{-\tau_{k-1}\tau_k} e^{-\theta^\ell (t_k^* - s)} ds,$$

$$\hat{c}_{k-1}^{k,\ell} = \int_{t_{k-2}}^{t_{k-1}} \frac{2s - t_{k-2} - t_{k-1}}{\tau_k(\tau_{k-1} + \tau_k)} e^{-\theta^\ell(t_k^* - s)} ds.$$

Comparing $L_k^{\alpha,*}u$ with $F_k^{\alpha,*}u$, we see that the former requires all the previous time step value $\{u^j\}_{j=1}^k$, while the latter only needs u^k , u^{k-1} , u^{k-2} , and $H^\ell(t_k^*)$, $\ell = 1, \dots, N_q$. This means that approximating $\partial_t^\alpha u(t_k^*)$ by $F_k^{\alpha,*}u$ rather than $L_k^{\alpha,*}u$ could reduce the storage and computational cost, when k is large. Roughly speaking, the storage cost can be reduced from $\mathcal{O}(N)$ to $\mathcal{O}(N_q)$ and the computational cost can be reduced from $\mathcal{O}(N^2)$ to $\mathcal{O}(N_q N)$ by replacing $L_k^{\alpha,*}u$ with $F_k^{\alpha,*}u$.

The main purpose of this work is to establish global-in-time H^1 stability of the fast L2-1 $_\sigma$ scheme for subdiffusion equation.

3. Positive semidefiniteness of bilinear form \mathcal{B}_n . In this part, we state and prove the main results on positive semidefiniteness of a bilinear form associated with $F_k^{\alpha,*}u$, that will be used to establish the global-in-time H^1 -stability of the fast L2-1 $_\sigma$ scheme for subdiffusion equations.

Firstly, we shall reformulate the formula (2.4) as

$$\begin{aligned} (3.1) \quad F_k^{\alpha,*}u &= \frac{1}{\Gamma(1-\alpha)} \left(\sum_{j=1}^{k-1} \int_{t_{j-1}}^{t_j} \partial_s \Pi_{2,j}(s) \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell(t_k^* - s)} ds + \int_{t_{k-1}}^{t_k^*} \frac{\partial_s \Pi_{1,k}(s)}{(t_k^* - s)^\alpha} ds \right) \\ &= \frac{1}{\Gamma(1-\alpha)} \sum_{j=1}^{k-1} (\hat{a}_j^{(k)} u^{j-1} + \hat{b}_j^{(k)} u^j + \hat{c}_j^{(k)} u^{j+1}) + \frac{\sigma^{1-\alpha}}{\Gamma(2-\alpha)\tau_k^\alpha} \delta_k u \\ &= \frac{1}{\Gamma(1-\alpha)} \sum_{j=1}^{k-1} (\hat{c}_j^{(k)} \delta_{j+1} u - \hat{a}_j^{(k)} \delta_j u) + \frac{\sigma^{1-\alpha}}{\Gamma(2-\alpha)\tau_k^\alpha} \delta_k u \\ &= \frac{1}{\Gamma(1-\alpha)} \left(\hat{c}_{k-1}^{(k)} \delta_k u + \sum_{j=1}^{k-1} \hat{d}_j^{(k)} \delta_j u \right) + \frac{\sigma^{1-\alpha}}{\Gamma(2-\alpha)\tau_k^\alpha} \delta_k u, \end{aligned}$$

where $\delta_j u = u^j - u^{j-1}$,

$$\begin{aligned} (3.2) \quad \hat{a}_j^{(k)} &= \int_{t_{j-1}}^{t_j} \frac{2s - t_j - t_{j+1}}{\tau_j(\tau_j + \tau_{j+1})} \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell(t_k^* - s)} ds, \\ \hat{b}_j^{(k)} &= \int_{t_{j-1}}^{t_j} \frac{2s - t_{j-1} - t_{j+1}}{-\tau_j \tau_{j+1}} \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell(t_k^* - s)} ds, \\ \hat{c}_j^{(k)} &= \int_{t_{j-1}}^{t_j} \frac{2s - t_{j-1} - t_j}{\tau_{j+1}(\tau_j + \tau_{j+1})} \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell(t_k^* - s)} ds, \\ \hat{d}_j^{(k)} &= \hat{c}_{j-1}^{(k)} - \hat{a}_j^{(k)}, \end{aligned}$$

for $1 \leq j \leq k-1$. In the reformulation (3.1), we use the fact $\hat{a}_j^{(k)} + \hat{b}_j^{(k)} + \hat{c}_j^{(k)} = 0$. Here we make a convention that $\hat{c}_0^{(k)} = 0$, i.e., $\hat{d}_1^{(k)} = -\hat{a}_1^{(k)}$.

We now give some properties of the fast L2-1 $_\sigma$ coefficients $\hat{d}_j^{(k)}$ in (3.2).

LEMMA 3.1 (Properties of $\hat{d}_j^{(k)}$). *For the fast L2-1 $_\sigma$ coefficients $\hat{d}_j^{(k)}$ on a nonuniform mesh $\{\tau_j\}_{j \geq 1}$ defined in (3.2), the following properties hold:*

- (P1) $\hat{d}_j^{(k)} > 0$, $1 \leq j \leq k-1$, $k \geq 2$;
- (P2) $\hat{d}_j^{(k+1)} - \hat{d}_j^{(k)} < 0$, $1 \leq j \leq k-1$, $k \geq 2$.

Furthermore, if the nonuniform mesh $\{\tau_j\}_{j \geq 1}$ with $\rho_j := \tau_j/\tau_{j-1}$ satisfies

$$(3.3) \quad \rho_j \geq \eta \approx 0.475329, \quad \forall j \geq 2,$$

where η is the unique positive root of $1 - 3\rho^2(1 + \rho) = 0$, then

$$(P3) \quad \hat{d}_{j+1}^{(k)} - \hat{d}_j^{(k)} > 0, \quad 1 \leq j \leq k-2, \quad k \geq 3;$$

$$(P4) \quad \hat{d}_{j+1}^{(k)} - \hat{d}_j^{(k)} > \hat{d}_{j+1}^{(k+1)} - \hat{d}_j^{(k+1)}, \quad 1 \leq j \leq k-2, \quad k \geq 3.$$

Proof. We first provide two equivalent reformulations of $\hat{a}_j^{(k)}$ according to (3.2): $\forall 1 \leq j \leq k-1$,

$$(3.4) \quad \begin{aligned} \hat{a}_j^{(k)} &= \sum_{\ell=1}^{N_q} \int_0^1 \frac{-2\tau_j(1-s) - \tau_{j+1}}{\tau_j + \tau_{j+1}} \varpi^\ell e^{-\theta^\ell(t_k^* - (t_{j-1} + s\tau_j))} ds \\ &= \sum_{\ell=1}^{N_q} \frac{1}{\tau_j + \tau_{j+1}} \int_0^1 \varpi^\ell e^{-\theta^\ell(t_k^* - (t_{j-1} + s\tau_j))} d(\tau_j s^2 - (2\tau_j + \tau_{j+1})s) \\ &= \sum_{\ell=1}^{N_q} \varpi^\ell \left(-e^{-\theta^\ell(t_k^* - t_j)} + \frac{\theta^\ell \tau_j}{\tau_j + \tau_{j+1}} \int_0^1 (\tau_j + \tau_{j+1} + s\tau_j)(1-s) e^{-\theta^\ell(t_k^* - t_j + s\tau_j)} ds \right) \end{aligned}$$

and

$$(3.5) \quad \begin{aligned} \hat{a}_j^{(k)} &= \sum_{\ell=1}^{N_q} \int_0^1 \frac{-2\tau_j(1-s) - \tau_{j+1}}{\tau_j + \tau_{j+1}} \varpi^\ell e^{-\theta^\ell(t_k^* - (t_{j-1} + s\tau_j))} ds \\ &= \sum_{\ell=1}^{N_q} \int_0^1 \frac{-2\tau_j s - \tau_{j+1}}{(\tau_j + \tau_{j+1})} \varpi^\ell e^{-\theta^\ell(t_k^* - t_j + s\tau_j)} ds \\ &= \sum_{\ell=1}^{N_q} \frac{1}{\tau_j + \tau_{j+1}} \int_0^1 \varpi^\ell e^{-\theta^\ell(t_k^* - t_j + s\tau_j)} d(-\tau_j s^2 - \tau_{j+1}s) \\ &= - \sum_{\ell=1}^{N_q} \varpi^\ell \left(e^{-\theta^\ell(t_k^* - t_{j-1})} + \frac{\theta^\ell \tau_j}{\tau_j + \tau_{j+1}} \int_0^1 (\tau_j + \tau_{j+1} - s\tau_j)(1-s) e^{-\theta^\ell(t_k^* - t_{j-1} - s\tau_j)} ds \right). \blacksquare \end{aligned}$$

Furthermore, we also reformulate $\hat{c}_j^{(k)}$ in (3.2) as: $\forall 1 \leq j \leq k-1$,

$$(3.6) \quad \begin{aligned} \hat{c}_j^{(k)} &= \sum_{\ell=1}^{N_q} \int_0^1 \frac{\tau_j^2(2s-1)}{\tau_{j+1}(\tau_j + \tau_{j+1})} \varpi^\ell e^{-\theta^\ell(t_k^* - (t_{j-1} + s\tau_j))} ds \\ &= \sum_{\ell=1}^{N_q} \frac{\tau_j^2}{\tau_{j+1}(\tau_j + \tau_{j+1})} \int_0^1 \varpi^\ell e^{-\theta^\ell(t_k^* - (t_{j-1} + s\tau_j))} d(s^2 - s) \\ &= \sum_{\ell=1}^{N_q} \frac{\varpi^\ell \theta^\ell \tau_j^3}{\tau_{j+1}(\tau_j + \tau_{j+1})} \int_0^1 s(1-s) e^{-\theta^\ell(t_k^* - t_j + s\tau_j)} ds. \end{aligned}$$

From equations (3.5)–(3.6), it is easy to verify $\hat{a}_j^{(k)} < 0$ and $\hat{c}_j^{(k)} > 0$, which imply $\hat{d}_j^{(k)} = \hat{c}_{j-1}^{(k)} - \hat{a}_j^{(k)} > 0$, i.e., (P1) holds (note that $\hat{c}_0^{(k)} = 0$). Moreover, for any fixed s , we know that $e^{-\theta^\ell(t_k^* - t_{j-1})}$, $e^{-\theta^\ell(t_k^* - t_{j-1} - s\tau_j)}$, and $e^{-\theta^\ell(t_k^* - t_j + s\tau_j)}$ decrease w.r.t. k .

Combining this with $\theta^\ell > 0$ and $\varpi^\ell > 0$ stated in Lemma 2.1, we can claim that $\hat{a}_j^{(k+1)} - \hat{a}_j^{(k)} > 0$, $\hat{c}_j^{(k+1)} - \hat{c}_j^{(k)} < 0$ and $\hat{d}_j^{(k+1)} - \hat{d}_j^{(k)} < 0$, i.e., (P2) holds.

We now turn to prove the properties (P3) and (P4). From equations (3.4)–(3.6), we can derive

$$(3.7) \quad \begin{aligned} \hat{d}_{j+1}^{(k)} - \hat{d}_j^{(k)} &= (\hat{c}_j^{(k)} - \hat{c}_{j-1}^{(k)}) - (\hat{a}_{j+1}^{(k)} - \hat{a}_j^{(k)}) \\ &= \sum_{\ell=1}^{N_q} \varpi^\ell \theta^\ell \left(G_{j,k}^\ell + \frac{\tau_{j+1}}{\tau_{j+1} + \tau_{j+2}} \int_0^1 (\tau_{j+1} + \tau_{j+2} - s\tau_{j+1})(1-s)e^{-\theta^\ell(t_k^* - t_j - s\tau_{j+1})} ds \right), \end{aligned}$$

where

$$(3.8) \quad \begin{aligned} G_{j,k}^\ell &= \frac{\tau_j^3}{\tau_{j+1}(\tau_j + \tau_{j+1})} \int_0^1 s(1-s)e^{-\theta^\ell(t_k^* - t_j + s\tau_j)} ds \\ &\quad - \frac{\tau_{j-1}^3}{\tau_j(\tau_{j-1} + \tau_j)} \int_0^1 s(1-s)e^{-\theta^\ell(t_k^* - t_{j-1} + s\tau_{j-1})} ds \\ &\quad + \frac{\tau_j}{\tau_j + \tau_{j+1}} \int_0^1 (\tau_j + \tau_{j+1} + s\tau_j)(1-s)e^{-\theta^\ell(t_k^* - t_j + s\tau_j)} ds \\ &= \left(\frac{\tau_j^3}{\tau_{j+1}(\tau_j + \tau_{j+1})} + \frac{(4\tau_j + 3\tau_{j+1})\tau_j}{\tau_j + \tau_{j+1}} \right) \int_0^1 s(1-s)e^{-\theta^\ell(t_k^* - t_j + s\tau_j)} ds \\ &\quad - \frac{\tau_{j-1}^3}{\tau_j(\tau_{j-1} + \tau_j)} \int_0^1 s(1-s)e^{-\theta^\ell(t_k^* - t_{j-1} + s\tau_{j-1})} ds \\ &\quad + e^{-\theta^\ell(t_k^* - t_j)} \tau_j \int_0^1 (1-3s)(1-s)e^{-\theta^\ell s\tau_j} ds. \end{aligned}$$

and we use the forms (3.4) for $\hat{a}_j^{(k)}$ and (3.5) for $\hat{a}_{j+1}^{(k)}$. Here we make a convention $\tau_0 = 0$, so that $\hat{d}_1^{(k)} = -\hat{a}_1^{(k)}$. Under the condition (3.3), we have

$$\frac{(4\tau_j + 3\tau_{j+1})\tau_j}{\tau_j + \tau_{j+1}} - \frac{\tau_{j-1}^3}{\tau_j(\tau_{j-1} + \tau_j)} = \tau_j \left(\frac{1}{1 + \rho_{j+1}} + \frac{3\rho_j^2(1 + \rho_j) - 1}{\rho_j^2(1 + \rho_j)} \right) \geq 0.$$

Combining this with the fact $e^{-\theta^\ell(t_k^* - t_j + s\tau_j)} > e^{-\theta^\ell(t_k^* - t_{j-1} + s\tau_{j-1})}$ and

$$\begin{aligned} \theta^\ell \tau_j \int_0^1 (1-3s)(1-s)e^{-\theta^\ell s\tau_j} ds &= \int_0^1 (1-3s)(1-s) \left(-e^{-\theta^\ell s\tau_j} \right)' ds \\ &= 1 + \int_0^1 (6s-4)e^{-\theta^\ell s\tau_j} ds = 1 - e^{-\theta^\ell \xi_j^\ell \tau_j}, \end{aligned}$$

where ξ_j^ℓ is a constant in $(0, 1)$, we can find that

$$(3.9) \quad G_{j,k}^\ell > (\theta^\ell)^{-1} e^{-\theta^\ell(t_k^* - t_j)} \left(1 - e^{-\theta^\ell \xi_j^\ell \tau_j} \right) > 0, \quad \forall j < k.$$

Then we can derive from (3.7) that

$$(3.10) \quad \hat{d}_{j+1}^{(k)} - \hat{d}_j^{(k)} > \sum_{\ell=1}^{N_q} \frac{\varpi^\ell \theta^\ell \tau_{j+1}}{\tau_{j+1} + \tau_{j+2}} \int_0^1 (\tau_{j+1} + \tau_{j+2} - s\tau_{j+1})(1-s)e^{-\theta^\ell(t_k^* - t_j - s\tau_{j+1})} ds.$$

Therefore, (P3) holds. Moreover, the property (P4) holds follows from

$$\begin{aligned} & (\hat{d}_{j+1}^{(k)} - \hat{d}_j^{(k)}) - (\hat{d}_{j+1}^{(k+1)} - \hat{d}_j^{(k+1)}) \\ &= \sum_{\ell=1}^{N_q} \varpi^\ell \theta^\ell \left(1 - e^{\theta^\ell (t_k^* - t_{k+1}^*)} \right) \\ & \quad \left(G_{j,k}^\ell + \frac{\tau_{j+1}}{\tau_{j+1} + \tau_{j+2}} \int_0^1 (\tau_{j+1} + \tau_{j+2} - s\tau_{j+1})(1-s)e^{-\theta^\ell (t_k^* - t_j - s\tau_{j+1})} ds \right) > 0, \end{aligned}$$

according to (3.7). \square

Now we can state and prove the following main theorem.

THEOREM 3.2. *Consider a nonuniform mesh $\{\tau_k\}_{k \geq 1}$ satisfying that for $k \geq 2$*

$$(3.11) \quad \begin{aligned} \rho_k &\geq \eta \approx 0.475329, \quad \varepsilon \leq \min_{k \geq 1} \frac{1}{5(1-\alpha)(\sigma\tau_k)^\alpha}, \\ \Delta t &\leq \min_{k \geq 2} \sigma\tau_k, \quad T_{\text{soe}} \geq \max_{k \geq 2} (\sigma\tau_{k+1} + \tau_k), \end{aligned}$$

where ε , Δt , and T_{soe} are fixed in Lemma 2.1, and $\sigma = 1 - \alpha/2$. Then for any function u defined on $[0, \infty) \times \Omega$ and $n \geq 1$,

$$\mathcal{B}_n(u, u) = \sum_{k=1}^n \langle F_k^{\alpha,*} u, \delta_k u \rangle \geq \sum_{k=1}^n \frac{[\mathbf{B}]_{kk}}{\Gamma(1-\alpha)} \|\delta_k u\|^2 \geq 0,$$

where $\|\cdot\|$ is the L^2 -norm and \mathbf{B} is defined in (3.13).

Proof. According to (3.1), we can rewrite $\mathcal{B}_n(u, u)$ in the following matrix form

$$(3.12) \quad \mathcal{B}_n(u, u) = \sum_{k=1}^n \langle F_k^{\alpha,*} u, \delta_k u \rangle = \frac{1}{\Gamma(1-\alpha)} \int_{\Omega} \psi \mathbf{M} \psi^T dx,$$

where $\psi = [\delta_1 u, \delta_2 u, \dots, \delta_n u]$ and $\mathbf{M} = \mathbf{A} + \mathbf{B}$ with

$$\mathbf{A} = \begin{pmatrix} \beta_1 & & & & \\ \hat{d}_1^{(2)} & \beta_2 & & & \\ \hat{d}_1^{(3)} & \hat{d}_2^{(3)} & \beta_3 & & \\ \vdots & \vdots & \ddots & \ddots & \\ \hat{d}_1^{(n)} & \hat{d}_2^{(n)} & \dots & \hat{d}_{n-1}^{(n)} & \beta_n \end{pmatrix}$$

and

$$(3.13) \quad \mathbf{B} = \text{diag} \left(\frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_1^\alpha} - \beta_1, \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_2^\alpha} + \hat{c}_1^{(2)} - \beta_2, \dots, \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_n^\alpha} + \hat{c}_{n-1}^{(n)} - \beta_n \right)$$

with

$$(3.14) \quad 2\beta_1 = \hat{d}_1^{(2)}, \quad 2\beta_k - \hat{d}_k^{(k+1)} = \hat{d}_{k-1}^{(k)} - \hat{d}_{k-1}^{(k+1)}, \quad 2 \leq k \leq n.$$

Consider the following symmetric matrix $\mathbf{S} = \mathbf{A} + \mathbf{A}^T$. According to the properties (P1)–(P4) in Lemma 3.1, if the condition (3.11) holds, all the elements of \mathbf{S} are positive, and \mathbf{S} satisfies the following three properties:

- (1) $\forall 1 \leq j < i \leq n$, $[\mathbf{S}]_{i-1,j} \geq [\mathbf{S}]_{i,j}$;

- (2) $\forall 1 < j \leq i \leq n$, $[\mathbf{S}]_{i,j-1} < [\mathbf{S}]_{i,j}$;
 (3) $\forall 1 < j < i \leq n$, $[\mathbf{S}]_{i-1,j-1} - [\mathbf{S}]_{i,j-1} \leq [\mathbf{S}]_{i-1,j} - [\mathbf{S}]_{i,j}$.

From [32, Lemma 2.1], $\mathbf{S} = \mathbf{A} + \mathbf{A}^T$ is positive semidefinite.

Next we will prove $[\mathbf{B}]_{kk} \geq 0$, $k \geq 1$, under the condition (3.11). When $k = 1$, from the second equality in (3.5), $2\beta_1 = \hat{d}_1^{(2)} = -\hat{a}_1^{(2)}$ in (3.14), and Lemma 2.1, we have

(3.15)

$$\begin{aligned} [\mathbf{B}]_{11} &= \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_1^\alpha} - \frac{1}{2} \sum_{\ell=1}^{N_q} \int_0^1 \frac{2s + \rho_2}{1 + \rho_2} \varpi^\ell e^{-\theta^\ell(\sigma\tau_2 + s\tau_1)} ds \\ &> \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_1^\alpha} - \frac{1}{2} \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell\sigma\tau_2} \int_0^1 \frac{2s + \rho_2}{1 + \rho_2} ds = \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_1^\alpha} - \frac{1}{2} \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell\sigma\tau_2} \\ &\geq \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_1^\alpha} - \frac{1}{2}(\sigma\tau_2)^{-\alpha} - \frac{1}{2}\varepsilon \geq \frac{1}{2(1-\alpha)(\sigma\tau_1)^\alpha} (2 - \alpha - (1-\alpha)\eta^{-\alpha}) - \frac{1}{2}\varepsilon. \end{aligned}$$

When $k \geq 2$, from $2\beta_k - \hat{d}_k^{(k+1)} = \hat{d}_{k-1}^{(k)} - \hat{d}_{k-1}^{(k+1)}$ in (3.14), we have


$$(3.16) \quad [\mathbf{B}]_{kk} = \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_k^\alpha} + \frac{1}{2}\hat{c}_{k-1}^{(k)} + \frac{1}{2}[\hat{c}_{k-1}^{(k)} - (\hat{d}_{k-1}^{(k)} - \hat{d}_{k-1}^{(k+1)} + \hat{d}_k^{(k+1)})].$$

From the equations (3.4)–(3.6) and $\hat{d}_j^{(k)} = \hat{c}_{j-1}^{(k)} - \hat{a}_j^{(k)}$, when the condition (3.11) holds, we can derive for $k \geq 2$

$$\begin{aligned} &\hat{c}_{k-1}^{(k)} - (\hat{d}_{k-1}^{(k)} - \hat{d}_{k-1}^{(k+1)} + \hat{d}_k^{(k+1)}) \\ &= (\hat{c}_{k-1}^{(k)} - \hat{c}_{k-1}^{(k+1)}) - (\hat{c}_{k-2}^{(k)} - \hat{c}_{k-2}^{(k+1)}) + (\hat{a}_{k-1}^{(k)} - \hat{a}_{k-1}^{(k+1)} + \hat{a}_k^{(k+1)}) \\ &= \sum_{\ell=1}^{N_q} \varpi^\ell \theta^\ell (1 - e^{\theta^\ell(t_k^* - t_{k+1}^*)}) G_{k-1,k}^\ell - \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell(\sigma\tau_k)} \\ &\quad - \sum_{\ell=1}^{N_q} \frac{\varpi^\ell \theta^\ell \tau_k}{\tau_k + \tau_{k+1}} \int_0^1 (\tau_k + \tau_{k+1} - s\tau_k)(1-s) e^{-\theta^\ell(t_{k+1}^* - t_{k-1} - s\tau_k)} ds \\ &\geq - \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell(\sigma\tau_k)} - \sum_{\ell=1}^{N_q} \frac{\varpi^\ell \theta^\ell \tau_k}{\tau_k + \tau_{k+1}} \int_0^1 (\tau_{k+1} + s\tau_k) s e^{-\theta^\ell(\sigma\tau_{k+1} + s\tau_k)} ds, \end{aligned} \quad (3.17)$$

where $G_{k-1,k}^\ell > 0$ is defined in (3.8) and we use (3.4) for $\hat{a}_{k-1}^{(k)}$ and $\hat{a}_{k-1}^{(k+1)}$, and (3.5) for $\hat{a}_k^{(k+1)}$. Moreover, from Lemma 2.1, we have

$$\begin{aligned} (3.18) \quad &\sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell\sigma\tau_k} + \sum_{\ell=1}^{N_q} \frac{\varpi^\ell \theta^\ell \tau_k}{\tau_k + \tau_{k+1}} \int_0^1 (\tau_{k+1} + s\tau_k) s e^{-\theta^\ell(\sigma\tau_{k+1} + s\tau_k)} ds \\ &\leq \sum_{\ell=1}^{N_q} \varpi^\ell \left(e^{-\theta^\ell\sigma\tau_k} + \theta^\ell \tau_k \int_0^1 e^{-\theta^\ell(\sigma\tau_{k+1} + s\tau_k)} ds \right) \\ &= \sum_{\ell=1}^{N_q} \varpi^\ell \left(e^{-\theta^\ell\sigma\tau_k} + \int_0^1 (-e^{-\theta^\ell(\sigma\tau_{k+1} + s\tau_k)})' ds \right) \end{aligned}$$

FIG. 1. $f_1(\alpha)$ and $f_2(\alpha)$ w.r.t. $\alpha \in (0, 1)$.

$$\begin{aligned}
&= \sum_{\ell=1}^{N_q} \varpi^\ell \left(e^{-\theta^\ell \sigma \tau_k} + e^{-\theta^\ell \sigma \tau_{k+1}} - e^{-\theta^\ell (\sigma \tau_{k+1} + \tau_k)} \right) \\
&\leq (\sigma \tau_k)^{-\alpha} + (\sigma \tau_{k+1})^{-\alpha} - (\sigma \tau_{k+1} + \tau_k)^{-\alpha} + 3\varepsilon
\end{aligned}$$

Combining (3.16)–(3.18), $\sigma = 1 - \alpha/2$ and the fact $\hat{c}_{k-1}^{(k)} > 0$, we can derive

$$\begin{aligned}
[\mathbf{B}]_{kk} &\geq \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_k^\alpha} - \frac{1}{2}(\sigma \tau_k)^{-\alpha} - \frac{1}{2}(\sigma \tau_{k+1})^{-\alpha} + \frac{1}{2}(\sigma \tau_{k+1} + \tau_k)^{-\alpha} - \frac{3}{2}\varepsilon \\
(3.19) \quad &= \frac{1}{2(1-\alpha)(\sigma \tau_k)^\alpha} \left(1 - (1-\alpha)\sigma^\alpha [(\sigma \tau_{k+1})^{-\alpha} - (\sigma \tau_{k+1} + \tau_k)^{-\alpha}] \right) - \frac{3}{2}\varepsilon \\
&\geq \frac{1}{2(1-\alpha)(\sigma \tau_k)^\alpha} \left(1 - (1-\alpha)\sigma^\alpha [(\sigma \eta)^{-\alpha} - (\sigma \eta + 1)^{-\alpha}] \right) - \frac{3}{2}\varepsilon.
\end{aligned}$$

We now plot the following functions in Figure 1

$$f_1(\alpha) = 2 - \alpha - (1 - \alpha)\eta^{-\alpha}, \quad f_2(\alpha) = 1 - (1 - \alpha)\sigma^\alpha [(\sigma \eta)^{-\alpha} - (\sigma \eta + 1)^{-\alpha}],$$

where $\eta \approx 0.475329$ from (3.3). We can see that $f_i(\alpha) \geq 0.6$, for $i = 1, 2$. Thus $[\mathbf{B}]_{kk} \geq 0$, $k \geq 1$, under the condition (3.11). □

4. Long time H^1 -stability of the fast L2-1 $_\sigma$ scheme.

4.1. Linear subdiffusion equation. For simplicity, we first consider the following linear subdiffusion equation:

$$\begin{aligned}
(4.1) \quad &\partial_t^\alpha u(t, x) = \Delta u(t, x) + f(t, x), \quad (t, x) \in (0, \infty) \times \Omega, \\
&u(t, x) = 0, \quad (t, x) \in (0, \infty) \times \partial\Omega, \\
&u(0, x) = u^0(x), \quad x \in \Omega,
\end{aligned}$$

where Ω is a bounded Lipschitz domain in \mathbb{R}^d . Given an arbitrary nonuniform mesh $\{\tau_k\}_{k \geq 1}$, the fast L2-1 $_\sigma$ scheme of this subdiffusion equation is written as

$$\begin{aligned}
(4.2) \quad &F_k^{\alpha,*} u = (1 - \alpha/2)\Delta u^k + \alpha/2\Delta u^{k-1} + f(t_k^*, \cdot), \quad \text{in } \Omega, \\
&u^k = 0, \quad \text{on } \partial\Omega.
\end{aligned}$$

THEOREM 4.1. *Assume that $f(t, x) \in L^{\infty}([0, \infty); L^2(\Omega)) \cap BV([0, \infty); L^2(\Omega))$ is a bounded variation function in time and $u^0 \in H_0^1(\Omega)$. Fix ε , Δt , and T_{soe} in Lemma 2.1. For an arbitrary nonuniform mesh $\{\tau_k\}_{k \geq 1}$ satisfying (3.11), then the numerical solution u^n of the fast L2-1 $_{\sigma}$ scheme (4.2) satisfies the following global-in-time H^1 -stability*

$$\|\nabla u^n\| \leq \|\nabla u^0\| + 2C_f C_{\Omega}, \quad \forall n \geq 1,$$

where $C_f = 2\|f\|_{L^{\infty}([0, \infty); L^2(\Omega))} + \|f\|_{BV([0, \infty); L^2(\Omega))}$, and C_{Ω} is the Sobolev embedding constant depending on Ω and the spatial dimension d .

Proof. Due to the positive semidefiniteness of bilinear form \mathcal{B}_n established in Theorem 3.2, the proof is quite similar as the one of [33, Theorem 4.1]. We omit it here. \square

4.2. Semilinear subdiffusion equation. In this part, we consider the semilinear subdiffusion equation

$$(4.3) \quad \begin{aligned} \partial_t^{\alpha} u(t, x) &= \nu^2 \Delta u(t, x) - f(u(t, x)), & (t, x) \in (0, \infty) \times \Omega, \\ u(0, x) &= u^0(x), & x \in \Omega, \end{aligned}$$

where $\nu > 0$ is the diffusion coefficient, f is some nonlinear functional of $u(t, x)$, and Ω is a bounded Lipschitz domain in \mathbb{R}^d . For simplicity, we consider the periodic boundary condition or the homogeneous Dirichlet/Neumann boundary condition and assume that

$$(4.4) \quad \sup_{u \in \mathbb{R}} |f'(u)| \leq L$$

for some constant $L > 0$. One of such models is the time fractional phase-field model, which is revealed to admit different dynamical scales [40].

We consider the following scheme for the semilinear subdiffusion equation (4.3) in Ω , using the fast L2-1 $_{\sigma}$ formula for Caputo derivative and the Newton linearization for nonlinearity:

$$(4.5) \quad F_k^{\alpha, *} u = (1 - \alpha/2) \nu^2 \Delta u^k + (\alpha/2) \nu^2 \Delta u^{k-1} - f(u^{k-1}) - (1 - \alpha/2) f'(u^{k-1}) \delta_k u.$$

THEOREM 4.2. *Assume that f in (4.3) satisfies (4.4). Fix ε , Δt , and T_{soe} in Lemma 2.1. For an arbitrary nonuniform mesh $\{\tau_k\}_{k \geq 1}$ satisfying (3.11) and*

$$(4.6) \quad \begin{aligned} \tau_1^{\alpha} &\leq \frac{2 - \alpha - (1 - \alpha)\eta^{-\alpha}}{\sigma^{\alpha}[(3 - \alpha)\Gamma(2 - \alpha)L + (1 - \alpha)\varepsilon]}, \\ \tau_k^{\alpha} &\leq \frac{1 - (1 - \alpha)\sigma^{\alpha}[(\sigma\eta)^{-\alpha} - (\sigma\eta + 1)^{-\alpha}]}{\sigma^{\alpha}[(3 - \alpha)\Gamma(2 - \alpha)L + 3(1 - \alpha)\varepsilon]}, \quad k \geq 2, \end{aligned}$$

where $\sigma = 1 - \alpha/2$, then the numerical solution u^n of the fast L2-1 $_{\sigma}$ scheme (4.5) satisfies the following energy stability: $\forall n \geq 1$,

$$\int_{\Omega} \left(\frac{\nu^2}{2} |\nabla u^n|^2 + F(u^n) \right) dx \leq \int_{\Omega} \left(\frac{\nu^2}{2} |\nabla u^0|^2 + F(u^0) \right) dx,$$

where F is a primitive integral of f , i.e., $F'(u) = f(u)$.

Proof. Fix n and let $E^n := \int_{\Omega} \left(\frac{\nu^2}{2} |\nabla u^n|^2 + F(u^n) \right) dx$. Multiplying (4.5) by $\delta_k u$ and integrating over Ω yield

$$\begin{aligned} & \langle F_k^{\alpha,*} u, \delta_k u \rangle \\ &= -\frac{\nu^2}{2} (\|\nabla u^k\|^2 - \|\nabla u^{k-1}\|^2 + (1-\alpha)\|\nabla \delta_k u\|^2) \\ & \quad - \langle F(u^k) - F(u^{k-1}), 1 \rangle + \frac{1}{2} \langle f'(\omega) \delta_k u, \delta_k u \rangle - (1-\alpha/2) \langle f'(u^{k-1}) \delta_k u, \delta_k u \rangle \\ &= -E^k + E^{k-1} - \frac{\nu^2(1-\alpha)}{2} \|\nabla \delta_k u\|^2 + \frac{1}{2} \langle (f'(\omega) - 2(1-\alpha/2)f'(u^{k-1})) \delta_k u, \delta_k u \rangle \\ &\leq -E^k + E^{k-1} + \frac{(3-\alpha)L}{2} \|\delta_k u\|^2, \end{aligned}$$

where ω is between u^k and u^{k-1} . Summing up the above inequality over k yields

$$\sum_{k=1}^n \langle F_k^{\alpha} u, \delta_k u \rangle \leq -E^n + E^0 + \frac{(3-\alpha)L}{2} \sum_{k=1}^n \|\delta_k u\|^2.$$

From Theorem 3.2, we have

$$E^n \leq E^0 - \sum_{k=1}^n \left(\frac{[\mathbf{B}]_{kk}}{\Gamma(1-\alpha)} - \frac{(3-\alpha)L}{2} \right) \|\delta_k u\|^2.$$

According to (3.15) and (3.19), under the condition (4.6), we have

$$\frac{[\mathbf{B}]_{kk}}{\Gamma(1-\alpha)} - \frac{(3-\alpha)L}{2} \geq 0, \quad 1 \leq k \leq n$$

and consequently $E^n \leq E^0$ for any $n \geq 1$. \square

5. Sharp finite time H^1 -error estimate. In this part, we present the sharp H^1 -error estimate of the fast L2-1 $_{\sigma}$ scheme (4.2) for the linear subdiffusion equation (4.1) in finite time $t \in (0, T_{\text{soe}}]$ with some constraints on the time step. We consider time meshes $\{\tau_k\}_{k=1}^N$:

$$(5.1) \quad 0 = t_0 < t_1 < \dots < t_k \dots < t_N = T_{\text{soe}}, \quad \text{with} \quad \tau_k = t_k - t_{k-1},$$

where T_{soe} is given in Lemma 2.1. We first reformulate the discrete formula (3.1):

$$F_k^{\alpha,*} u = \frac{1}{\Gamma(1-\alpha)} \left([\mathbf{M}]_{k,k} u^k - \sum_{j=2}^k ([\mathbf{M}]_{k,j} - [\mathbf{M}]_{k,j-1}) u^{j-1} - [\mathbf{M}]_{k,1} u^0 \right),$$

where \mathbf{M} is given by (3.12). We now give some properties on $[\mathbf{M}]_{k,j}$.

LEMMA 5.1. *Given a nonuniform mesh $\{\tau_k\}_{k=1}^N$ in (5.1) satisfying*

$$(5.2) \quad \rho_k \geq \eta \approx 0.475329, \quad \Delta t \leq \min_{2 \leq k \leq N} \sigma \tau_k, \quad \varepsilon \leq \min_{2 \leq k \leq N} \frac{1-\alpha}{\sigma} \hat{c}_{k-1}^{(k)},$$

where $\varepsilon, \Delta t$ are fixed in Lemma 2.1 and $\sigma = 1 - \alpha/2$, then following properties of $[\mathbf{M}]_{k,j}$ given by (3.12) hold:

(Q1) for $1 \leq j \leq k-1$,

$$[\mathbf{M}]_{k,j} \geq \frac{\eta}{(1+\eta)\tau_j} \sum_{\ell=1}^{N_q} \int_{t_{j-1}}^{t_j} \varpi^\ell e^{-\theta^\ell(t_k^*-s)} ds, \quad [\mathbf{M}]_{k,k} \geq \frac{1}{\tau_k} \int_{t_{k-1}}^{t_k^*} (t_k^* - s)^{-\alpha} ds;$$

(Q2) for $2 \leq j \leq k-1$,

$$\begin{aligned} [\mathbf{M}]_{k,j} - [\mathbf{M}]_{k,j-1} &\geq \sum_{\ell=1}^{N_q} \frac{\varpi^\ell \theta^\ell \tau_j}{\tau_j + \tau_{j+1}} \int_0^1 (\tau_j + \tau_{j+1} - s\tau_j)(1-s)e^{-\theta^\ell(t_k^*-t_{j-1}-s\tau_j)} ds, \\ [\mathbf{M}]_{k,k} - [\mathbf{M}]_{k,k-1} &\geq \frac{\alpha}{2(1-\alpha)(\sigma\tau_k)^\alpha} - \varepsilon; \end{aligned}$$

(Q3)

$$\frac{1-\alpha}{\sigma} [\mathbf{M}]_{k,k} - [\mathbf{M}]_{k,k-1} \geq 0, \quad k \geq 2.$$

Proof. From (3.12) and the first equation in (3.4), for $1 \leq j \leq k-1$,

$$\begin{aligned} [\mathbf{M}]_{k,j} &= \hat{d}_j^{(k)} \geq -\hat{a}_j^{(k)} \geq \sum_{\ell=1}^{N_q} \int_0^1 \frac{\tau_{j+1}}{\tau_j + \tau_{j+1}} \varpi^\ell e^{-\theta^\ell(t_k^*-(t_{j-1}+s\tau_j))} ds \\ &\geq \frac{\eta}{(1+\eta)\tau_j} \sum_{\ell=1}^{N_q} \int_{t_{j-1}}^{t_j} \varpi^\ell e^{-\theta^\ell(t_k^*-s)} ds, \end{aligned}$$

and

$$[\mathbf{M}]_{k,k} = \hat{c}_{k-1}^{(k)} + \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_k^\alpha} \geq \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_k^\alpha} = \frac{1}{\tau_k} \int_{t_{k-1}}^{t_k^*} (t_k^* - s)^{-\alpha} ds.$$

For $2 \leq j \leq k-1$, $[\mathbf{M}]_{k,j} - [\mathbf{M}]_{k,j-1} = \hat{d}_j^{(k)} - \hat{d}_{j-1}^{(k)}$. Then the first the result of (Q2) is from (3.10). Moreover from Lemma 2.1, if (3.3) holds for $j = k-1$, we have

$$\begin{aligned} [\mathbf{M}]_{k,k} - [\mathbf{M}]_{k,k-1} &= \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_k^\alpha} + (\hat{c}_{k-1}^{(k)} - \hat{c}_{k-2}^{(k)} + \hat{a}_{k-1}^{(k)}) \\ &= \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_k^\alpha} + \sum_{\ell=1}^{N_q} \varpi^\ell \theta^\ell G_{k-1,k}^\ell - \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell \sigma \tau_k} \\ &> \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_k^\alpha} - \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell \sigma \tau_k} \geq \frac{\sigma^{1-\alpha}}{(1-\alpha)\tau_k^\alpha} - (\sigma\tau_k)^{-\alpha} - \varepsilon = \frac{\alpha}{2(1-\alpha)(\sigma\tau_k)^\alpha} - \varepsilon. \end{aligned}$$

where $G_{k-1,k}^\ell > 0$ is defined in (3.8). For the property (Q3), from (3.4) and (3.6) and Lemma 2.1 we have

$$\begin{aligned} &\frac{1-\alpha}{\sigma} [\mathbf{M}]_{k,k} - [\mathbf{M}]_{k,k-1} \\ &= (\sigma\tau_k)^{-\alpha} + \frac{1-\alpha}{\sigma} \hat{c}_{k-1}^{(k)} - \sum_{\ell=1}^{N_q} \varpi^\ell \theta^\ell \left(\frac{\tau_{k-2}^3}{\tau_{k-1}(\tau_{k-2} + \tau_{k-1})} \int_0^1 s(1-s)e^{-\theta^\ell(t_k^*-t_{k-2}+s\tau_{k-2})} ds \right. \\ &\quad \left. - \frac{\tau_{k-1}}{\tau_{k-1} + \tau_k} \int_0^1 (\tau_{k-1} + \tau_k + s\tau_{k-1})(1-s)e^{-\theta^\ell(t_k^*-t_{k-1}+s\tau_{k-1})} ds \right) - \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell \sigma \tau_k} \end{aligned}$$

$$\geq (\sigma\tau_k)^{-\alpha} + \frac{1-\alpha}{\sigma} \hat{c}_{k-1}^{(k)} - \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell \sigma \tau_k} \geq \frac{1-\alpha}{\sigma} \hat{c}_{k-1}^{(k)} - \varepsilon \geq 0,$$

where we use the same technique as the proof of (3.9) to obtain the first inequality. \square

We now analyze the approximation error of the fast L2-1 $_\sigma$ formula in the following lemma.

LEMMA 5.2. *Given a function u satisfying $|\partial_t^m u(t)| \leq C_m(1+t^{\alpha-m})$ for $m = 1, 3$ and the nonuniform mesh $\{\tau_k\}_{k=1}^N$ in (5.1) satisfying (5.2). The approximation error is defined by*

(5.3)

$$r_k := \frac{1}{\Gamma(1-\alpha)} \left(\int_0^{t_{k-1}^*} (t_k^* - s)^{-\alpha} \partial_s u(s) \, ds - \int_0^{t_{k-1}^*} \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell (t_k^* - s)} \partial_s I_2 u(s) \, ds \right. \\ \left. + \int_{t_{k-1}}^{t_k^*} (t_k^* - s)^{-\alpha} \partial_s [u(s) - \Pi_{1,k} u(s)] \, ds \right) \quad k \geq 1,$$

where $I_2 u = \Pi_{2,j} u$ on (t_{j-1}, t_j) for $j < k$, and $\Pi_{1,j}$ and $\Pi_{2,j}$ are two standard Lagrange interpolation operators with the interpolation points: $\{t_{j-1}, t_j\}$, $\{t_{j-1}, t_j, t_{j+1}\}$, respectively. Then for $k \geq 1$,

$$|r_k| \leq \frac{C}{\Gamma(1-\alpha)} \left\{ [\mathbf{M}]_{k,1} (t_2^\alpha / \alpha + t_2) + \sum_{j=2}^k ([\mathbf{M}]_{k,j} - [\mathbf{M}]_{k,j-1}) (1 + \rho_{j+1}) (1 + t_{j-1}^{\alpha-3}) \tau_j^3 \right. \\ \left. + \varepsilon (t_{k-1} + t_{k-1}^\alpha / \alpha + 2\sigma(1 + t_{k-1}^{\alpha-3}) \tau_k^3) \mathbf{1}_{k \geq 2} \right\},$$

where $C > 0$ is a constant depending only on C_m for $m = 1, 3$ and $\rho_{k+1} = 1$.

Proof. The case of $k = 1$ can be verified easily. We now consider the case of $k \geq 2$. From (5.3) and Lemma 2.1, it is not difficult to derive

(5.4)

$$|r_k| \leq \frac{\varepsilon}{\Gamma(1-\alpha)} \int_0^{t_{k-1}^*} |\partial_s u(s)| \, ds + \frac{1}{\Gamma(1-\alpha)} \left| \int_{t_{k-1}}^{t_k^*} (t_k^* - s)^{-\alpha} \partial_s [u(s) - \Pi_{1,k} u(s)] \, ds \right| \\ + \frac{1}{\Gamma(1-\alpha)} \left| \int_0^{t_{k-1}^*} \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell (t_k^* - s)} [\partial_s u(s) - \partial_s I_2 u(s)] \, ds \right|.$$

Using $|\partial_s u(s)| \leq C_1(1 + s^{\alpha-1})$ gives

$$\int_0^{t_{k-1}^*} |\partial_s u(s)| \, ds \leq C_1 (t_{k-1} + t_{k-1}^\alpha / \alpha).$$

Similar to [34, Lemma 4.5], we can derive the following inequalities based on Lemma 5.1:

$$\left| \int_0^{t_1} \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell (t_k^* - s)} [\partial_s u(s) - \partial_s I_2 u(s)] \, ds \right| \leq C [\mathbf{M}]_{k,1} (t_2^\alpha / \alpha + t_2), \\ \left| \int_{t_{j-1}}^{t_j} \sum_{\ell=1}^{N_q} \varpi^\ell e^{-\theta^\ell (t_k^* - s)} [\partial_s u(s) - \partial_s I_2 u(s)] \, ds \right|$$

$$\leq C_3([\mathbf{M}]_{k,j} - [\mathbf{M}]_{k,j-1})(1 + \rho_{j+1})(1 + t_{j-1}^{\alpha-3})\tau_j^3, \quad 2 \leq j \leq k-1,$$

and

$$\left| \int_{t_{k-1}}^{t_k^*} (t_k^* - s)^{-\alpha} \partial_s [u(s) - \Pi_{1,k} u(s)] ds \right| \leq 2C_3([\mathbf{M}]_{k,k} - [\mathbf{M}]_{k,k-1})\sigma(1 + t_{k-1}^{\alpha-3})\tau_k^3 + 2\varepsilon C_3\sigma(1 + t_{k-1}^{\alpha-3})\tau_k^3.$$

Combining the above inequalities, we can obtain the desired result. \square

THEOREM 5.3 (Sharp finite time H^1 -error estimate). *Assume that $u \in C^3((0, T_{\text{soe}}], H_0^1(\Omega) \cap H^2(\Omega))$ is a solution to (4.1) in finite time $(0, T_{\text{soe}}]$ and $|\partial_t^m u(t)| \leq C_m(1 + t^{\alpha-m})$, for $m = 1, 2, 3$, $0 < t \leq T_{\text{soe}}$. If the nonuniform mesh $\{\tau_k\}_{k=1}^N$ in (5.1) satisfies (5.2) and $\varepsilon \leq \frac{1}{2}T_{\text{soe}}^{-\alpha}$, then the numerical solutions of fast L2-1 $_{\sigma}$ scheme (4.2) satisfies the following error estimate*

$$\begin{aligned} \max_{1 \leq k \leq N} \|e^k\|_{H^1(\Omega)} \leq C \bigg\{ & \left(\frac{t_2^\alpha/\alpha + t_2}{1 - \alpha} \max_{2 \leq k \leq N} (1 + \rho_{k+1})(1 + t_{k-1}^{\alpha-3})(t_{k-1}^*)^\alpha \tau_k^3 \tau_{k-1}^{-\alpha} \right. \\ & + (\tau_1^\alpha/\alpha + \tau_1)\tau_1^{\alpha/2} + \sqrt{\Gamma(1 - \alpha)} \max_{2 \leq k \leq N} (t_k^*)^{\alpha/2} (1 + t_{k-1}^{\alpha-2})\tau_k^2 \\ & \left. + \varepsilon \max_{2 \leq k \leq N} (t_k^*)^\alpha (t_{k-1} + t_{k-1}^\alpha/\alpha + 2\sigma(1 + t_{k-1}^{\alpha-3})\tau_k^3) \right\}, \end{aligned}$$

where $e^k := u(t_k) - u^k$ and C is a constant depending only on C_m , $m = 1, 2, 3$ and Ω . Moreover, for the graded mesh with grading parameter $r \geq 1$, i.e.,

$$(5.5) \quad t_j = \left(\frac{j}{N}\right)^r T_{\text{soe}}, \quad \tau_j = t_j - t_{j-1} = \left[\left(\frac{j}{N}\right)^r - \left(\frac{j-1}{N}\right)^r\right] T_{\text{soe}},$$

we have

$$(5.6) \quad \max_{1 \leq k \leq N} \|e^k\|_{H^1(\Omega)} \leq \tilde{C} \left(N^{-\min\{r\alpha, 2\}} + \varepsilon \right).$$

where \tilde{C} depends on α , Ω , T_{soe} , and C_m with $m = 1, 2, 3$.

Proof. Based on Lemma 5.2, the proof is similar to [34, Theorem 4.6] except replacing the test function $e_k^* = (1 - \alpha/2)e^k + \alpha/2e^{k-1}$ with $-\Delta e_k^*$. We omit it here and leave it for readers. \square

REMARK 5.1. The sharp finite time H^1 -error estimate of the fast L2-1 $_{\sigma}$ scheme for subdiffusion equations has also been well studied in [19, 25]. In Theorem 5.3, we reduce the restriction $\rho_k \geq 2/3$ in [19, 25] to $\rho_k \geq \eta \approx 0.475329$.

6. Numerical results. In this section, we provide some numerical results for the fast L2-1 $_{\sigma}$ schemes (4.2) and (4.5). As in [22, 4], all the discrete coefficients $a_j^{(k)}$, $c_j^{(k)}$, $\hat{a}_{k-1}^{k,\ell}$ and $\hat{c}_{k-1}^{k,\ell}$ in (2.2) and (2.5) are computed by adaptive Gauss-Kronrod quadrature, to avoid roundoff error problems.

EXAMPLE 6.1. Consider the subdiffusion equation (4.1) with $f(t, x, y) = \Gamma(1 + \alpha)(x^2 - 1)(y^2 - 1) - 2t^\alpha(x^2 + y^2 - 2)$, thus the exact solution $u(t, x, y) = t^\alpha(x^2 - 1)(y^2 - 1)$ in $\Omega = [-1, 1]^2$.

In this example, we use the graded mesh (5.5). We first compare the computational cost of the fast L2-1 $_{\sigma}$ scheme (4.2) with the standard L2-1 $_{\sigma}$ scheme in [2]. Here we

TABLE 1

(Example 6.1) $\max_{1 \leq k \leq N} \|u(t_k) - u^k\|$ for graded meshes with different grading parameters r and time step numbers N where $\alpha = 0.3$.

	$N = 100$	$N = 200$	$N = 400$	$N = 800$	$N = 1600$
$r = 1/\alpha$	1.8477e-3	9.5337e-4	4.8429e-4	2.4407e-4	1.2252e-4
order	—	0.9546	0.9772	0.9885	0.9943
$r = 2/\alpha$	5.2619e-5	1.3355e-5	3.3696e-6	8.4709e-7	2.1248e-7
order	—	1.9782	1.9868	1.9920	1.9951
$r = 3/\alpha$	1.1576e-4	2.9563e-5	7.4851e-6	1.8855e-6	4.7355e-7
order	—	1.9693	1.9817	1.9890	1.9934

TABLE 2

(Example 6.1) $\max_{1 \leq k \leq N} \|u(t_k) - u^k\|$ for graded meshes with different grading parameters r and time step numbers N where $\alpha = 0.5$.

	$N = 100$	$N = 200$	$N = 400$	$N = 800$	$N = 1600$
$r = 1/\alpha$	2.1627e-3	1.1124e-3	5.6416e-4	2.8409e-4	1.4255e-4
order	—	0.9592	0.9795	0.9897	0.9949
$r = 2/\alpha$	3.4309e-5	8.6705e-6	2.1836e-6	5.4867e-7	1.3764e-7
order	—	1.9844	1.9894	1.9927	1.9950
$r = 3/\alpha$	7.5018e-5	1.9013e-5	4.7967e-6	1.2065e-6	3.0290e-7
order	—	1.9802	1.9869	1.9912	1.9940

use spectral collocation method in space with 10^2 Chebyshev–Gauss–Lobatto points. We plot the CPU time w.r.t. the total number of time steps N for both schemes with $\alpha = 0.5$ on the left-hand side of Figure 2. Here we set $T_{\text{soe}} = 1$, $\varepsilon = 1e-12$, $\Delta t = \sigma\tau_2$ in Lemma 2.1 and the grading parameter $r = 1.5$ in (5.5). We observe that the CPU time of the fast L2- 1_σ scheme increases linearly w.r.t. N , while the cost of the original L2- 1_σ scheme increases quadratically.

To better understand the relationship between the number of quadrature nodes N_q and the parameters $(\varepsilon, \Delta t)$ in Lemma 2.1, we plot N_q w.r.t. ε for fixed $\Delta t = 1e-5$ and N_q w.r.t. Δt for fixed $\varepsilon = 1e-13$, respectively in the middle and right subfigures of Figure 2 where $T_{\text{soe}} = 1$. We can see that N_q increases almost linearly w.r.t. both $\log(\varepsilon)$ and $\log(\Delta t)$.




FIG. 2. (Example 6.1) Left: CPU time in seconds w.r.t. number of time steps N with $T_{\text{soe}} = 1$, $\varepsilon = 1e-12$, $\Delta t = \sigma\tau_2$ for the SOE approximation in Lemma 2.1 and $\alpha = 0.5$, $r = 1.5$ for the graded mesh (5.5). Middle: N_q w.r.t. ε with $T_{\text{soe}} = 1$, $\Delta t = 1e-5$. Right: N_q w.r.t. Δt with $T_{\text{soe}} = 1$, $\varepsilon = 1e-13$.

TABLE 3

(Example 6.1) $\max_{1 \leq k \leq N} \|u(t_k) - u^k\|$ for graded meshes with different grading parameters r and time step numbers N where $\alpha = 0.7$.

	$N = 100$	$N = 200$	$N = 400$	$N = 800$	$N = 1600$
$r = 1/\alpha$	1.8710e-3	9.5944e-4	4.8584e-4	2.4447e-4	1.2262e-4
order	—	0.9636	0.9817	0.9908	0.9954
$r = 2/\alpha$	2.0342e-5	5.1591e-6	1.2961e-6	3.2443e-7	8.1133e-8
order	—	1.9793	1.9929	1.9982	1.9996
$r = 3/\alpha$	3.9451e-5	9.9647e-6	2.5108e-6	6.3158e-7	1.5866e-7
order	—	1.9852	1.9887	1.9911	1.9930




FIG. 3. (Example 6.1) Maximum H^1 -error w.r.t. N for $\alpha = 0.3, 0.5, 0.7$ from left to right with different r for graded meshes (5.5).

We now test the convergence of the fast L2-1 $_{\sigma}$ scheme (4.2). Let $T_{\text{soe}} = 1$, $\varepsilon = 1e-12$, $\Delta t = \sigma\tau_2$ in the Lemma 2.1, and apply the spectral collocation method in the space with 25^2 Chebyshev–Gauss–Lobatto points. Table 1–3 present the maximum L^2 -errors with $\alpha = 0.3, 0.5, 0.7$ and $r = 1/\alpha, 2/\alpha, 3/\alpha$ for graded meshes respectively. It can be observed that the convergence rates in L^2 -norm are $N^{-\min\{r\alpha, 2\}}$ for the fast L2-1 $_{\sigma}$ scheme (4.2) on graded meshes. The maximum L^2 -error is computed by $\max_{1 \leq k \leq N} \|u(t_k) - u^k\|$. We show the maximum H^1 -error in Figure 3 with $\alpha = 0.3, 0.5, 0.7$ and $r = 1/\alpha, 2/\alpha, 3/\alpha$ for graded meshes respectively. The maximum H^1 -error is computed by $\max_{1 \leq k \leq N} \|\nabla(u(t_k) - u^k)\|$. We can also find that the convergence rates in H^1 -norm for the fast L2-1 $_{\sigma}$ scheme (4.2) on graded meshes are $N^{-\min\{r\alpha, 2\}}$, which is consistent with Theorem 5.3.

EXAMPLE 6.2 (Long time simulation). Consider the subdiffusion equation (4.1), with two different sources terms $f_1(t, x, y) = t \sin(0.2t)$, $f_2(t, x, y) = 5 \exp(-0.0005t) \sin(0.005t)$. The initial condition is set to $u^0 = \sin(\pi x) \sin(\pi y)$ in $\Omega = [-1, 1]^2$.

We test the global-in-time H^1 -stability of the fast L2-1 $_{\sigma}$ scheme (4.2) with source term f_1 and f_2 respectively. The spectral collocation method is used in space with 10^2 Chebyshev–Gauss–Lobatto points. We set $T_{\text{soe}} = 1$, $\varepsilon = 1e-12$, $\Delta t = \sigma\tau_2$ in Lemma 2.1. In addition, we use the following nonuniform mesh:

$$\tau_j = \begin{cases} \left(\frac{j}{100}\right)^r - \left(\frac{j-1}{100}\right)^r & j \leq 100 \\ 1.005\tau_{j-1} & j > 100, 1.005\tau_{j-1} < \tau_{\max} := 0.2, \\ \tau_{\max} & \text{otherwise,} \end{cases}$$




FIG. 4. (Example 6.2) Long time H^1 -instability. Left: $\|\nabla u\|$ w.r.t. time t . Right: Time step w.r.t. time t .




FIG. 5. (Example 6.2) Long time H^1 -stability. Left: $\|\nabla u\|$ w.r.t. time: t . Right: Time step w.r.t. time: t .

where $r = 2/\alpha$.

We plot $\|\nabla u\|$ w.r.t. time t in Figure 4 and Figure 5 for f_1 and f_2 respectively, computed by the fast L2-1 $_\sigma$ scheme (4.2) with $\alpha = 0.5$. Note that the time t_n can be much larger than T_{soe} . We can see that $\|\nabla u\|$ is bounded in the case of f_2 , but not in the case of f_1 . One explanation is that $f_1 \notin L^\infty([0, \infty); L^2(\Omega)) \cap BV([0, \infty); L^2(\Omega))$ so that the assumption on source term in Theorem 4.1 is not satisfied, while $f_2 \in L^\infty([0, \infty); L^2(\Omega)) \cap BV([0, \infty); L^2(\Omega))$ holds true.

EXAMPLE 6.3. (a) Consider the semilinear subdiffusion equation

$$\partial_t^\alpha u = \Delta u + \sin(u) + g(t, x, y), \quad (t, x, y) \in (0, 1] \times (-1, 1)^2$$

with homogeneous Dirichlet boundary condition. To investigate the convergence orders in time, the initial condition u^0 and source term g are chosen such that the exact solution is $u(t, x, y) = t^\alpha \sin(\pi x) \sin(\pi y)$.

(b) Consider the semilinear subdiffusion equation

$$\partial_t^\alpha u = \nu^2 \Delta u + \sin(u), \quad (t, x, y) \in (0, \infty) \times (-1, 1)^2$$

with homogeneous Dirichlet boundary condition, $\nu^2 = 0.01$, and initial condition $u^0 = \sin(\pi x) \sin(\pi y)$.

For the problem (a), we use the graded meshes (5.5). We set $T_{\text{soe}} = 1$, $\varepsilon = 1e - 12$, $\Delta t = \sigma \tau_2$ in Lemma 2.1, and use spectral collocation method in space with 25^2

Chebyshev–Gauss–Lobatto points. Figure 6 shows the maximum L_2 -errors and maximum H^1 -error for $\alpha = 0.3, 0.5, 0.7$ with $r = 1/\alpha, 2/\alpha, 3/\alpha$ for graded meshes (5.5). Again, it can be observed that the convergence rates in both L^2 -norm and H^1 -norm are $N^{-\min\{r\alpha, 2\}}$ for graded meshes, which agree with the results in [25].




FIG. 6. (Example 6.3) Maximum L^2 -error (top) and H^1 -error (bottom) w.r.t. N for $\alpha = 0.3, 0.5, 0.7$ from left to right, with different r for graded mesh (5.5).

We now turn to show the energy evolution for the problem (b) of Example 6.3 for $\alpha = 0.3, 0.5, 0.7$. The energy is $E = \int_{\Omega} \left(\frac{\nu^2}{2} |\nabla u|^2 + \cos(u) \right) dx$. The spectral collocation method is applied in space with 10^2 Chebyshev–Gauss–Lobatto points. Let $T_{\text{soe}} = 300$, $\varepsilon = 1e-12$, $\Delta t = \sigma \tau_2$ in Lemma 2.1. We use the following nonuniform time mesh:

$$(6.1) \quad \tau_j = \begin{cases} 0.1 \left[\left(\frac{j}{100} \right)^r - \left(\frac{j-1}{100} \right)^r \right] & j \leq 100 \\ 1.01\tau_{j-1} & j > 100, 1.01\tau_{j-1} < \tau_{\max} := 0.02, \\ \tau_{\max} & \text{otherwise,} \end{cases}$$

where $r = 3$. This mesh satisfies all constraints in Theorem 4.2. The energy evolution and time steps are illustrated in Figure 7, which is consistent with Theorem 4.2.

EXAMPLE 6.4. Consider the time-fractional Allen–Cahn (TFAC) equation

$$\partial_t^\alpha u = \nu^2 \Delta u - f(u), \quad (t, x, y) \in (0, \infty) \times (-1, 1)^2$$

with homogeneous Dirichlet boundary condition, $\nu^2 = 0.01$, $u^0 = 0.05(2 * \text{rand}(x, y) - 1)$, and $f(u) = u^3 - u$, where $\text{rand}(\cdot)$ stands for the random values in $(0, 1)$.

Note that the global Lipschitz property fails for the TFAC equation. Fortunately, it is well studied in [5] that the TFAC equation satisfies the maximum principle, i.e.,




FIG. 7. (Example 6.3) Left: Energy evolution w.r.t. time t with $\alpha = 0.3, 0.5, 0.7$. Right: Time step w.r.t. time t .

$\|u\|_\infty \leq 1$ if $\|u^0\|_\infty \leq 1$. We use the following truncation technique for $f(u)$ (see for example [35])

$$\hat{f}(u) = \begin{cases} (3M^2 - 1)u - 2M^3, & u > M, \\ u^3 - u, & |u| \leq M, \\ (3M^2 - 1)u + 2M^3, & u < -M, \end{cases}$$

one primitive integral of which is

$$\hat{F}(u) = \begin{cases} \frac{3M^2 - 1}{2}u^2 - 2M^3u + \frac{3M^4 + 1}{4}, & u > M, \\ \frac{1}{4}(u^2 - 1)^2, & |u| \leq M, \\ \frac{3M^2 - 1}{2}u^2 + 2M^3u + \frac{3M^4 + 1}{4}, & u < -M. \end{cases}$$

Here $M \geq 1$ is some positive constant. In particular, we set $M = 1$ so that $\sup_{u \in \mathbb{R}} |\hat{f}'(u)| \leq 2$, i.e., the Lipschitz condition is satisfied. Let $T_{\text{soe}} = 300$, $\varepsilon = 1e-12$, $\Delta t = \sigma \tau_2$ in Lemma 2.1. The space is discretized by the spectral collocation method with 10^2 Chebyshev–Gauss–Lobatto points. For the time discretization, we still use the time mesh (6.1).

Firstly we plot the energy evolution for the fast L2-1 $_\sigma$ scheme (4.5) using the aforementioned truncation technique with $\alpha = 0.4, 0.6, 0.8$. In this case, the energy is computed by $\hat{E} = \int_\Omega \left(\frac{\nu^2}{2} |\nabla u|^2 + \hat{F}(u) \right) dx$. It can be verified that the time mesh (6.1) satisfies all the conditions in Theorem 4.2 for $\alpha = 0.6, 0.8$. The energy stability result can be observed on the left-hand side of Figure 8 which agrees with Theorem 4.2. However, the energy stability is still observed for $\alpha = 0.4$ despite that the conditions in Theorem 4.2 are not satisfied.

Secondly, we compare the original energies of the fast L2-1 $_\sigma$ schemes (4.5) with $f(u)$ and $\hat{f}(u)$, on the right-hand side of Figure 8, where we take the time mesh (6.1) and $\alpha = 0.8$. Here the original energy is calculated by $E = \int_\Omega \left(\frac{\nu^2}{2} |\nabla u|^2 + \frac{1}{4}(u^2 - 1)^2 \right) dx$. It is observed that the corresponding original energies almost coincide and both of them decay monotonically in time, although we can not provide a rigorous proof of this energy dissipation.




FIG. 8. (Example 6.4) Left: Energy evolution w.r.t. time t for $\alpha = 0.4, 0.6, 0.8$. Right: Comparison of original energies w.r.t. time t with $\alpha = 0.8$ for $f(u)$ and $\hat{f}(u)$.

7. Conclusion. We have proved a new global-in-time H^1 -stability of the fast L2-1 $_{\sigma}$ scheme for linear and semilinear subdiffusion equations. The proposed scheme is constructed by combining the L2-1 $_{\sigma}$ scheme with the SOE approximation to the convolution kernel involved in the Caputo fractional derivative. A crucial bilinear form \mathcal{B}_n , associated with the fast L2-1 $_{\sigma}$ formula, is proved to be positive semidefinite under some mild constraints of time steps for fixed SOE parameters ε , Δt , and T_{soe} . Note that this positive semidefiniteness holds not only when $t \leq T_{\text{soe}}$ but also for any time $t > T_{\text{soe}}$. Therefore, the uniform long time H^1 -stability still holds as the time goes to infinity.

We also revisit the sharp H^1 -error estimate in finite time for the fast L2-1 $_{\sigma}$ scheme, and we relax the constraint on the stepsize ratios from $\rho_k \geq 2/3$ in [19, 25] to $\rho_k \geq \eta \approx 0.475329$. So far, the long time optimal H^1 -error estimate is still open for the fast L2-1 $_{\sigma}$ scheme with general nonuniform meshes. This might be solved with the help of positive semidefiniteness of the bilinear form associated with $F_k^{\alpha,*}$.

Acknowledgements. C. Quan is supported by NSFC Grant 12271241, the Stable Support Plan Program of Shenzhen Natural Science Fund (Program Contract No. 20200925160747003), and Shenzhen Science and Technology Program (Grant No. RCYX20210609104358076). J. Yang is supported by the National Natural Science Foundation of China (NSFC) Grant No. 12271240, the NSFC/Hong Kong RGC Joint Research Scheme (NSFC/RGC 11961160718), the fund of the Guangdong Provincial Key Laboratory of Computational Science and Material Design, China (No.2019B030301001), and the Shenzhen Natural Science Fund (RCJC20210609103819018).

REFERENCES

- [1] Mariam Al-Maskari and Samir Karaa, *Numerical approximation of semilinear subdiffusion equations with nonsmooth initial data*, SIAM Journal on Numerical Analysis **57** (2019), no. 3, 1524–1544.
- [2] Anatoly A Alikhanov, *A new difference scheme for the time fractional diffusion equation*, Journal of Computational Physics **280** (2015), 424–438.
- [3] Daniel Baffet and Jan S Hesthaven, *A kernel compression scheme for fractional differential equations*, SIAM Journal on Numerical Analysis **55** (2017), no. 2, 496–520.
- [4] Hu Chen and Martin Stynes, *Error analysis of a second-order method on fitted meshes for a time-fractional diffusion problem*, Journal of Scientific Computing **79** (2019), no. 1, 624–647.

- [5] Qiang Du, Jiang Yang, and Zhi Zhou, *Time-fractional Allen–Cahn equations: analysis and numerical methods*, Journal of Scientific Computing **85** (2020), no. 2, 1–30.
- [6] Guang-hua Gao, Zhi-zhong Sun, and Hong-wei Zhang, *A new fractional numerical differentiation formula to approximate the Caputo fractional derivative and its applications*, Journal of Computational Physics **259** (2014), 33–50.
- [7] Rudolf Gorenflo, Francesco Mainardi, Daniele Moretti, and Paolo Paradisi, *Time fractional diffusion: a discrete random walk approach*, Nonlinear Dynamics **29** (2002), no. 1, 129–143.
- [8] Shidong Jiang, Jiwei Zhang, Qian Zhang, and Zhimin Zhang, *Fast evaluation of the Caputo fractional derivative and its applications to fractional diffusion equations*, Communications in Computational Physics **21** (2017), no. 3, 650–678.
- [9] Bangti Jin, Buyang Li, and Zhi Zhou, *Correction of high-order BDF convolution quadrature for fractional evolution equations*, SIAM Journal on Scientific Computing **39** (2017), no. 6, A3129–A3152.
- [10] ———, *Numerical analysis of nonlinear subdiffusion equations*, SIAM Journal on Numerical Analysis **56** (2018), no. 1, 1–23.
- [11] ———, *Subdiffusion with time-dependent coefficients: improved regularity and second-order time stepping*, Numerische Mathematik **145** (2020), no. 4, 883–913.
- [12] Samir Karaa, *Positivity of discrete time-fractional operators with applications to phase-field equations*, SIAM Journal on Numerical Analysis **59** (2021), no. 4, 2040–2053.
- [13] Rihuan Ke, Michael K Ng, and Hai-Wei Sun, *A fast direct method for block triangular Toeplitz-like with tri-diagonal block systems from time-fractional partial differential equations*, Journal of Computational Physics **303** (2015), 203–211.
- [14] Natalia Kopteva, *Error analysis of the L1 method on graded and uniform meshes for a fractional-derivative problem in two and three dimensions*, Mathematics of Computation **88** (2019), no. 319, 2135–2155.
- [15] ———, *Error analysis of an L2-type method on graded meshes for a fractional-order parabolic problem*, Mathematics of Computation **90** (2021), no. 327, 19–40.
- [16] Natalia Kopteva and Xiangyun Meng, *Error analysis for a fractional-derivative parabolic problem on quasi-graded meshes using barrier functions*, SIAM Journal on Numerical Analysis **58** (2020), no. 2, 1217–1238.
- [17] TAM Langlands and Bruce I Henry, *The accuracy and stability of an implicit solution method for the fractional diffusion equation*, Journal of Computational Physics **205** (2005), no. 2, 719–736.
- [18] Buyang Li and Shu Ma, *Exponential convolution quadrature for nonlinear subdiffusion equations with nonsmooth initial data*, SIAM Journal on Numerical Analysis **60** (2022), no. 2, 503–528.
- [19] Xin Li, Hong-lin Liao, and Luming Zhang, *A second-order fast compact scheme with unequal time-steps for subdiffusion problems*, Numerical Algorithms **86** (2021), no. 3, 1011–1039.
- [20] Hong-lin Liao, Dongfang Li, and Jiwei Zhang, *Sharp error estimate of the nonuniform L1 formula for linear reaction-subdiffusion equations*, SIAM Journal on Numerical Analysis **56** (2018), no. 2, 1112–1133.
- [21] Hong-lin Liao, William McLean, and Jiwei Zhang, *A Discrete Grönwall Inequality with Applications to Numerical Schemes for Subdiffusion Problems*, SIAM Journal on Numerical Analysis **57** (2019), no. 1, 218–237.
- [22] Hong-Lin Liao, William McLean, and Jiwei Zhang, *A second-order scheme with nonuniform time steps for a linear reaction-subdiffusion problem*, Communications in Computational Physics **30** (2021), no. 2, 567–601.
- [23] Hong-lin Liao, Tao Tang, and Tao Zhou, *A second-order and nonuniform time-stepping maximum-principle preserving scheme for time-fractional Allen–Cahn equations*, Journal of Computational Physics **414** (2020), 109473.
- [24] Hong-lin Liao, Yonggui Yan, and Jiwei Zhang, *Unconditional convergence of a fast two-level linearized algorithm for semilinear subdiffusion equations*, Journal of Scientific Computing **80** (2019), no. 1, 1–25.
- [25] Nan Liu, Yanping Chen, Jiwei Zhang, and Yanmin Zhao, *Unconditionally optimal H^1 -error estimate of a fast nonuniform L2-1 σ scheme for nonlinear subdiffusion equations*, Numerical Algorithms (2022), 1–23.
- [26] Xin Lu, Hong-Kui Pang, and Hai-Wei Sun, *Fast approximate inversion of a block triangular Toeplitz matrix with applications to fractional sub-diffusion equations*, Numerical Linear Algebra with Applications **22** (2015), no. 5, 866–882.
- [27] Christian Lubich and Achim Schädle, *Fast convolution for nonreflecting boundary conditions*, SIAM Journal on Scientific Computing **24** (2002), no. 1, 161–182.

- [28] Chunwan Lv and Chuanju Xu, *Error analysis of a high order method for time-fractional diffusion equations*, SIAM Journal on Scientific Computing **38** (2016), no. 5, A2699–A2724.
- [29] Ralf Metzler and Joseph Klafter, *The random walk's guide to anomalous diffusion: a fractional dynamics approach*, Physics reports **339** (2000), no. 1, 1–77.
- [30] Kassem Mustapha, Basheer Abdallah, and Khaled M Furati, *A discontinuous Petrov–Galerkin method for time-fractional diffusion equations*, SIAM Journal on Numerical Analysis **52** (2014), no. 5, 2512–2529.
- [31] Kassem Mustapha and William McLean, *Uniform convergence for a discontinuous galerkin, time-stepping method applied to a fractional diffusion equation*, IMA Journal of Numerical Analysis **32** (2012), no. 3, 906–925.
- [32] Chaoyu Quan, Tao Tang, and Jiang Yang, *How to define dissipation-preserving energy for time-fractional phase-field equations*, CSIAM Transactions on Applied Mathematics **1** (2020), no. 3, 478–490.
- [33] Chaoyu Quan and Boyi Wang, *Energy stable $L2$ schemes for time-fractional phase-field equations*, Journal of Computational Physics **458** (2022), 111085.
- [34] Chaoyu Quan and Xu Wu, *On stability and convergence of $L2-1_\sigma$ method on general nonuniform meshes for subdiffusion equation*, arXiv preprint arXiv:2208.01384 (2022).
- [35] Jie Shen and Xiaofeng Yang, *Numerical approximations of Allen–Cahn and Cahn–Hilliard equations*, Discrete & Continuous Dynamical Systems-A **28** (2010), no. 4, 1669.
- [36] Jin-ye Shen, Zhi-zhong Sun, and Rui Du, *Fast finite difference schemes for time-fractional diffusion equations with a weak singularity at initial time*, East Asian J. Appl. Math. **8** (2018), no. 4, 834–858.
- [37] Kerui Song and Pin Lyu, *A high-order and fast scheme with variable time steps for the time-fractional black-scholes equation*, Mathematical Methods in the Applied Sciences (2021).
- [38] Martin Stynes, Eugene O’Riordan, and José Luis Gracia, *Error analysis of a finite difference method on graded meshes for a time-fractional diffusion equation*, SIAM Journal on Numerical Analysis **55** (2017), no. 2, 1057–1079.
- [39] Zhi-zhong Sun and Xiaonan Wu, *A fully discrete difference scheme for a diffusion-wave system*, Applied Numerical Mathematics **56** (2006), no. 2, 193–209.
- [40] Tao Tang, Boyi Wang, and Jiang Yang, *Asymptotic analysis on the sharp interface limit of the time-fractional cahn-hilliard equation*, SIAM J. Appl. Math. **82** (2022), no. 3, 773–792.
- [41] Kai Wang and Zhi Zhou, *High-order time stepping schemes for semilinear subdiffusion equations*, SIAM Journal on Numerical Analysis **58** (2020), no. 6, 3226–3250.
- [42] Yanyuan Xing and Yubin Yan, *A higher order numerical method for time fractional partial differential equations with nonsmooth data*, Journal of Computational Physics **357** (2018), 305–323.
- [43] Yonggui Yan, Zhi-Zhong Sun, and Jiwei Zhang, *Fast evaluation of the Caputo fractional derivative and its applications to fractional diffusion equations: a second-order scheme*, Communications in Computational Physics **22** (2017), no. 4, 1028–1048.
- [44] Yubin Yan, Monzorul Khan, and Neville J Ford, *An analysis of the modified $L1$ scheme for time-fractional partial differential equations with nonsmooth data*, SIAM Journal on Numerical Analysis **56** (2018), no. 1, 210–227.
- [45] Hongyi Zhu and Chuanju Xu, *A fast high order method for the time-fractional diffusion equation*, SIAM Journal on Numerical Analysis **57** (2019), no. 6, 2829–2849.