

MULTILINEAR SINGULAR INTEGRALS WITH HOMOGENEOUS KERNELS NEAR L^1

GEORGIOS DOSIDIS AND LENKA SLAVÍKOVÁ

ABSTRACT. We obtain the optimal open range of $L^{p_1}(\mathbb{R}^n) \times \dots \times L^{p_m}(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)$ bounds for multilinear singular integral operators with homogeneous kernels of the form $\Omega(\frac{x}{|x|})|x|^{-mn}$, where Ω is a function in $L^q(\mathbb{S}^{mn-1})$ with vanishing integral and $q > 1$.

1. INTRODUCTION

Singular integral operators of the form

$$(1) \quad Tf(x) := pv \int_{\mathbb{R}^n} f(y)K(x-y)dy$$

have been extensively studied ever since their introduction by Calderón and Zygmund in [1]. In the special case of homogeneous kernels $K(x) = \frac{\Omega(x/|x|)}{|x|^n}$, which includes the Hilbert and the Riesz transforms, boundedness of the operator T from $L^p(\mathbb{R}^n)$ to itself can be obtained for all $1 < p < \infty$ without any smoothness assumption on Ω . This was first proven in [2] under the assumption that Ω has vanishing integral on the unit sphere and belongs to the Orlicz space $L \log L(\mathbb{S}^{n-1})$, and subsequently improved in [8, 9] by making use of Hardy spaces. Endpoint bounds were obtained in [5, 6, 20, 24].

In this paper, we investigate the multilinear variant of the operator (1). Namely, given $m \in \mathbb{N}$ and a function Ω with zero mean on the unit sphere which, in addition, belongs to $L^q(\mathbb{S}^{mn-1})$ for some $q > 1$, we set $K(x) = \frac{\Omega(x')}{|x|^{mn}}$ for $x \in \mathbb{R}^{mn} \setminus \{0\}$, where $x' = \frac{x}{|x|}$, and consider the multilinear singular integral operator

$$(2) \quad T_{\Omega}^m(f_1, f_2, \dots, f_m)(x) := pv \int_{\mathbb{R}^{mn}} K(x-y_1, x-y_2, \dots, x-y_m) \prod_{i=1}^m f_i(y_i)dy.$$

Throughout the paper, we assume that exponents p_1, \dots, p_m, p satisfy the Hölder scaling $\frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i}$, and we study the question of $L^{p_1}(\mathbb{R}^n) \times \dots \times L^{p_m}(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)$ boundedness of T_{Ω}^m .

The $L^{p_1}(\mathbb{R}^n) \times L^{p_2}(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)$ boundedness of the bilinear operator T_{Ω}^2 was proven by Coifman and Meyer in [7] under the assumption that Ω is a function of bounded variation and $1 < p_1, p_2, p < \infty$. This result led, in particular, to an alternative proof of L^p estimates for Calderón's first commutator, an operator introduced by Calderón in [4] in connection with the study of elliptic partial differential equations. A yet another proof of the boundedness of Calderón's commutator relevant for our present approach was given by Muscalu in [22]. An extension of the Coifman–Meyer theorem [7] to arbitrary exponents satisfying $1 < p_1, p_2 < \infty$, with p being possibly less than 1, was obtained by Kenig and Stein in [21] and by Grafakos and Torres in [18].

The authors were supported by the Primus research programme PRIMUS/21/SCI/002 of Charles University.

The first result concerning bounds for T_Ω^2 in the case when Ω does not possess any smoothness was obtained in [10]. In that paper, the authors considered the one-dimensional case $n = 1$ and proved boundedness of the operator T_Ω^2 in a certain range of exponents under the assumption that the even part of Ω belongs to the Hardy space $H^1(\mathbb{S}^1)$. The higher-dimensional case was discussed in [13], where boundedness of T_Ω^2 in the full range $1 < p_1, p_2 < \infty$ was proven under the assumption that $\Omega \in L^\infty(\mathbb{S}^{2n-1})$, and boundedness in the “local L^2 case”, that is $2 \leq p_1, p_2 \leq \infty$ and $1 \leq p \leq 2$, was proven for $\Omega \in L^2(\mathbb{S}^{2n-1})$. The local L^2 case bound was later improved in [17], weakening the L^2 assumption on Ω to the requirement that $\Omega \in L^q(\mathbb{S}^{2n-1})$ with $q > 4/3$. Boundedness in the extended range of exponents $\frac{1}{p} + \frac{1}{q} < 2$, when $q > 4/3$, was proven recently in [19].

In this work, we extend the boundedness results for T_Ω^2 to the case $1 < q \leq 4/3$. Let $d\sigma$ denote the surface measure on the sphere.

Theorem 1. *Let $1 < p_1, p_2 < \infty$, $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$, and suppose that $\Omega \in L^q(\mathbb{S}^{2n-1})$, $q > 1$, with $\int_{\mathbb{S}^{2n-1}} \Omega(\theta) d\sigma(\theta) = 0$. If*

$$(3) \quad \frac{1}{p} + \frac{1}{q} < 2,$$

then there exists a constant $C = C(n, p_1, p_2, q)$ such that

$$(4) \quad \|T_\Omega^2(f_1, f_2)\|_{L^p(\mathbb{R}^n)} \leq C \|\Omega\|_{L^q(\mathbb{S}^{2n-1})} \|f_1\|_{L^{p_1}(\mathbb{R}^n)} \|f_2\|_{L^{p_2}(\mathbb{R}^n)}.$$

The proof of Theorem 1 relies on estimates obtained in [13, 19] combined with a new estimate of minimal blow-up when $\Omega \in L^1(\mathbb{S}^{2n-1})$.

Remark. *The question of weak-type boundedness of T_Ω^2 in the limiting case $\frac{1}{p} + \frac{1}{q} = 2$ remains open. In addition, no bounds are known for the operator T_Ω^2 when Ω belongs merely to $L \log L(\mathbb{S}^{2n-1})$ except in the one-dimensional setting, in which a partial result was obtained in [10].*

We graph the range of boundedness of T_Ω^2 . The dotted line corresponds to $\frac{1}{p} + \frac{1}{q} = 2$. Here and in what follows, q' stands for the exponent satisfying $\frac{1}{q} + \frac{1}{q'} = 1$.




FIGURE 1. Optimal range of $L^{p_1}(\mathbb{R}^n) \times L^{p_2}(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)$ boundedness of T_Ω^2 when $\Omega \in L^q(\mathbb{S}^{2n-1})$, $q > 1$.

The boundedness of the multilinear operator T_Ω^m with a rough kernel was first treated in [14], where initial $L^2(\mathbb{R}^n) \times \dots \times L^2(\mathbb{R}^n) \rightarrow L^{2/m}(\mathbb{R}^n)$ bounds were established. In order to discuss the boundedness region for this operator, we introduce some notation.

For any multiindex $\alpha \in \{0, 1\}^m$, we write $|\alpha| = \sum_{i=1}^m \alpha_i$. If $\vec{\frac{1}{p}} = \left(\frac{1}{p_1}, \dots, \frac{1}{p_m}\right)$, we set $\frac{1}{p_\alpha} = \alpha \cdot \vec{\frac{1}{p}} = \sum_{i=1}^m \frac{\alpha_i}{p_i}$. Then $\frac{1}{p} = \frac{1}{p_{(1, \dots, 1)}} = \sum_{i=1}^m \frac{1}{p_i}$. For any $q > 1$, we write $\vec{\frac{1}{p}} \in \mathcal{H}^m(q)$ if for all $\alpha \in \{0, 1\}^m$,

$$(5) \quad \frac{1}{p_\alpha} + \frac{|\alpha| - 1}{q} < |\alpha|.$$

When $|\alpha| = 0$ or 1 , Equation (5) is satisfied for all $1 < p_1, \dots, p_m < \infty$, while the condition for $|\alpha| = m$ is

$$\frac{1}{p} + \frac{m-1}{q} < m.$$

We graph $\mathcal{H}^3(q)$ in the Figure 2 below. The equation for $|\alpha| = 3$ corresponds to the red triangle, while the 3 equations for $|\alpha| = 2$, which have the form

$$\frac{1}{p_1} + \frac{1}{p_2} + \frac{1}{q} < 2, \quad \frac{1}{p_1} + \frac{1}{p_3} + \frac{1}{q} < 2, \quad \frac{1}{p_2} + \frac{1}{p_3} + \frac{1}{q} < 2,$$

correspond to the three red rectangles.

Recently in [15], the bounds for T_Ω^m were extended from the middle point $(\frac{1}{2}, \dots, \frac{1}{2})$ to the whole region $\mathcal{H}^m(q)$, when $q \geq 2$. Similar to the bilinear case, we extend these boundedness results to the case $1 < q < 2$.

Theorem 2. *Let $1 < p_1, p_2, \dots, p_m < \infty$, $\frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i}$ be such that $(\frac{1}{p_1}, \dots, \frac{1}{p_m}) \in \mathcal{H}^m(q)$ and suppose that $\Omega \in L^q(\mathbb{S}^{mn-1})$, $q > 1$, with $\int_{\mathbb{S}^{mn-1}} \Omega(\theta) d\sigma(\theta) = 0$. Then there exists a constant $C = C(n, p_1, p_2, \dots, p_m, q)$ such that*

$$(6) \quad \|T_\Omega^m(f_1, \dots, f_m)\|_{L^p(\mathbb{R}^n)} \leq C \|\Omega\|_{L^q(\mathbb{S}^{mn-1})} \prod_{i=1}^m \|f_i\|_{L^{p_i}(\mathbb{R}^n)}.$$

In particular, if $1 < p, p_1, p_2, \dots, p_m < \infty$, then Equation (6) holds for all $q > 1$. Note that for $m = 2$, the range $\mathcal{H}^2(q)$ corresponds to the condition $\frac{1}{p} + \frac{1}{q} < 2$, so Theorem 1 is a special case of Theorem 2. Moreover, for $m = 1$, the condition in Equation (5) becomes $p > 1$, which is the range of boundedness for the linear operator when $\Omega \in L^q(\mathbb{S}^{n-1})$ for some $q > 1$.




FIGURE 2. Optimal range of $L^{p_1}(\mathbb{R}^n) \times L^{p_2}(\mathbb{R}^n) \times L^{p_3}(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)$ boundedness of T_Ω^3 when $\Omega \in L^q(\mathbb{S}^{3n-1})$, $q > 1$.

Theorem 2 provides the optimal range of exponents such that the strong-type boundedness holds. Building on a counterexample introduced in [10] and refined in [16], it was proven in [15] that $\mathcal{H}^m(q)$ is the largest open set of exponents in which T_Ω^m is bounded. The counterexample in the next proposition also deals with the endpoints in $\mathcal{H}^m(q)$ for which we have equality in one or more of the equations in (5).

Proposition 3. *If $q > 1$, $1 < p_1, \dots, p_m < \infty$, $\frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i}$ and $(\frac{1}{p_1}, \dots, \frac{1}{p_m}) \notin \mathcal{H}^m(q)$, then there exist an odd function $\Omega \in L^q(\mathbb{S}^{mn-1})$ and $f_i \in L^{p_i}(\mathbb{R}^n)$, $i = 1, \dots, m$, such that $T_\Omega^m(f_1, \dots, f_m) \notin L^p(\mathbb{R}^n)$.*

In Section 2, we use the bootstrap argument introduced in [15] to reduce Theorem 2 to proving an estimate of minimal blow-up in the Banach range of exponents when $\Omega \in L^1(\mathbb{S}^{mn-1})$. This estimate is stated in Proposition 4 below and will be proved in Section 3. Section 4 contains the proof of Proposition 3.

Acknowledgment. We thank the anonymous referee for their valuable comments that helped to improve the exposition. We also thank Bae Jun Park for pointing out to us an error in the statement of Proposition 4 in an earlier version of the manuscript.

2. REDUCTIONS

We fix $m \in \mathbb{N}$, $q \geq 1$ and $\Omega \in L^q(\mathbb{S}^{mn-1})$ with mean value zero and in the following we simplify notation writing T instead of T_Ω^m . Let η be a Schwartz function in \mathbb{R}^{mn} such that $\widehat{\eta}(\xi) = 1$ when $|\xi| \leq 1$ and $\widehat{\eta}(\xi) = 0$ when $|\xi| \geq 2$, and let $\widehat{\beta} = \widehat{\eta} - \widehat{\eta}(2\cdot)$. We decompose the kernel K by setting $K^i(y) = K(y)\widehat{\beta}(2^{-i}y)$ for $i \in \mathbb{Z}$ and

$$K_j^i(y) = \left(\widehat{K}^i(\cdot)\widehat{\beta}(2^{-j+i}\cdot) \right)^\vee(y), \quad K_j = \sum_{i=-\infty}^{\infty} K_j^i, \quad j \in \mathbb{Z}.$$

We decompose $T = \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} T_j^i = \sum_{j=-\infty}^{\infty} T_j$ accordingly, where K_j^i is the kernel of T_j^i and K_j the kernel of T_j . We note that

$$\widehat{K}_j = \sum_{i=-\infty}^{\infty} \widehat{K}_j^i(2^i\cdot),$$

by the homogeneity of K .

Let $1 < p_1, p_2, \dots, p_m < \infty$ and let p satisfy $\frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i}$. We obtain Theorem 2 by proving bounds of the form

$$(7) \quad \|T_j\|_{L^{p_1}(\mathbb{R}^n) \times \dots \times L^{p_m}(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)} \lesssim Q(j) \|\Omega\|_{L^q(\mathbb{S}^{mn-1})}$$

that are summable in j . Here and in the following we are using the usual notation $A \lesssim B$ to denote that there exists a constant c such that $A \leq cB$ and we use $A \lesssim_\gamma B$ to imply that c depends on the parameter γ .

We first recall two estimates of the type (7) which are available in the literature. The following proposition follows from the multilinear Calderón-Zygmund theory [18]. Its specific version stated below was proved in [13, Proposition 3] in the bilinear case, and the same argument translates also to the multilinear setting, as observed in [15, Equation (3.3)].

Proposition A ([13, 15]). *Let $q > 1$ and $0 < \delta < 1/q'$. Then inequality (7) holds with $Q(j) = 2^{(mn-\delta)j}$ if $j \geq 0$ and $Q(j) = 2^{-|j|(1-\delta)}$ if $j < 0$.*

Let $\mathcal{C}^m(q) = (0, \frac{1}{q}]^m$. The following proposition follows from the results in [15, Section 3].

Proposition B ([15]). *Let $q \geq 2$ and $\vec{\frac{1}{p}} \in \mathcal{C}^m(q)$. Then there exist $r = r(\vec{\frac{1}{p}}, n) > 0$ and $j_0 = j_0(m, n) \in \mathbb{Z}$ such that inequality (7) holds with $Q(j) = 2^{-rj}$ if $j \geq j_0$.*

In what follows, we fix j_0 from Proposition B and we only consider $j \geq j_0$ in light of Proposition A. The following proposition will be proved in Section 3.

Proposition 4. *Let $j \geq j_0$, $p > 1$, $q = 1$ and $\varepsilon > 0$. Then inequality (7) holds with $Q(j) \lesssim_\varepsilon 2^{j\varepsilon}$.*

We assume Proposition 4 for now and we state and prove several claims that will gradually improve the dependence of the bound in estimate (7) on j . Theorem 2 will then be a direct consequence of Proposition A and Claim 7 below.

Claim 5. *Let $j \geq j_0$, $q > 1$ and $\vec{\frac{1}{p}} \in \mathcal{C}^m(q)$. Then there exists an $r = r(\vec{\frac{1}{p}}, q, n) > 0$ such that inequality (7) holds with $Q(j) = 2^{-rj}$.*

Proof. If $q \geq 2$ then the claim follows from Proposition B. To prove the claim in the case $1 < q < 2$, we will use complex interpolation between the estimates in Proposition B and Proposition 4. To be able to proceed with the interpolation argument, we view T_j as an $(m+1)$ -linear operator. More precisely, let

$$\tilde{T}(\Omega, f_1, \dots, f_m)(x) := pv \int_{\mathbb{R}^{mn}} (\Omega - \text{Avg}(\Omega))(y') |y|^{-mn} \prod_{i=1}^m f_i(x - y_i) dy,$$

where $\text{Avg}(\Omega) = \int_{\mathbb{S}^{mn-1}} \Omega d\sigma$ is the mean value of Ω . Note that \tilde{T} is $(m+1)$ -linear and for every Ω with mean zero and for any Schwartz functions f_1, \dots, f_m , it agrees with the singular integral defined in (2). Therefore, it is given by an m -linear convolution with a distribution W_Ω and $W_{\Omega - \text{Avg}(\Omega)} = W_\Omega$, with equality holding in the sense of distributions.

Similar to T_j , we define \tilde{T}_j by setting $\tilde{K}_\Omega = \frac{\Omega(y') - \text{Avg}(\Omega)}{|y|^{mn}}$, $\tilde{K}_\Omega^i(y) = \tilde{K}_\Omega(y) \widehat{\beta}(2^{-i}y)$,

$$\left(\tilde{K}_\Omega\right)_j^i(y) = \left(\widehat{\tilde{K}_\Omega^i(\cdot)} \widehat{\beta}(2^{-j+i}\cdot)\right)^\vee(y), \quad \left(\tilde{K}_\Omega\right)_j = \sum_{i=-\infty}^{\infty} \left(\tilde{K}_\Omega\right)_j^i,$$

and

$$\tilde{T}_j(\Omega, f_1, \dots, f_m)(x) = pv \int_{\mathbb{R}^{mn}} \left(\tilde{K}_\Omega\right)_j(y) \prod_{i=1}^m f_i(x - y_i) dy.$$

It is immediate that the bounds described above for T_j also hold for \tilde{T}_j and we can use $(m+1)$ -linear interpolation (see [3, Lemma 2.1 and Proposition 2.2] for the multilinear interpolation result used here) between Banach space $(m+1)$ -tuplets of the form $L^q(\mathbb{S}^{mn-1})$, $L^{p_1}(\mathbb{R}^n)$, \dots , $L^{p_m}(\mathbb{R}^n)$.

Fix a $1 < q < 2$ and a $\vec{\frac{1}{p}} \in \mathcal{C}^m(q)$ and assume that $p_1 = \min\{p_1, \dots, p_m\}$ (without a loss of generality, since the other cases are symmetric) and note that $p_1 > 2$ since $q < 2$. Then $\frac{p_1}{2} \vec{\frac{1}{p}} = \left(\frac{1}{2}, \frac{1}{2} \frac{p_1}{p_2}, \dots, \frac{1}{2} \frac{p_1}{p_m}\right) \in \mathcal{C}^m(2)$ and Proposition B yields

$$(8) \quad \|\tilde{T}_j\|_{L^2(\mathbb{S}^{mn-1}) \times L^2(\mathbb{R}^n) \times L^{\frac{2p_2}{p_1}}(\mathbb{R}^n) \times \dots \times L^{\frac{2p_m}{p_1}}(\mathbb{R}^n) \rightarrow L^{\frac{2p}{p_1}}(\mathbb{R}^n)} \lesssim 2^{-r_0 j}$$

for some positive exponent r_0 .

Let $\varepsilon > 0$ be small, to be determined momentarily. If $\delta \in (0, 1)$ satisfies $\delta < p$, then Proposition 4 and the embedding $L^{\frac{1}{1-\delta}}(\mathbb{S}^{mn-1}) \hookrightarrow L^1(\mathbb{S}^{mn-1})$ imply that there is a constant C_ε such that

$$(9) \quad \|\tilde{T}_j\|_{L^{\frac{1}{1-\delta}}(\mathbb{S}^{mn-1}) \times L^{\frac{p_1}{\delta}}(\mathbb{R}^n) \times L^{\frac{p_2}{\delta}}(\mathbb{R}^n) \times \dots \times L^{\frac{p_m}{\delta}}(\mathbb{R}^n) \rightarrow L^{\frac{p}{\delta}}(\mathbb{R}^n)} \lesssim C_\varepsilon 2^{\varepsilon j}$$

Interpolating between the estimates (8) and (9) we obtain

$$(10) \quad \|\tilde{T}_j\|_{L^q(\mathbb{S}^{mn-1}) \times L^{p_1}(\mathbb{R}^n) \times \dots \times L^{p_m}(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)} \lesssim C_\varepsilon^{\frac{p_1-2}{p_1-2\delta}} (2^j)^\varepsilon \frac{p_1-2}{p_1-2\delta} - r_0 \frac{2-2\delta}{p_1-2\delta}.$$

Finally, choosing $\varepsilon < r_0 \frac{2-2\delta}{p_1-2}$ completes the proof of the claim. \square

Claim 6. *Let $j \geq j_0$, $q > 1$, $\frac{\vec{1}}{p} \in \mathcal{H}^m(q)$ and $\varepsilon > 0$. Then inequality (7) holds with $Q(j) \lesssim_\varepsilon 2^{j\varepsilon}$.*

Proof. Claim 5 yields that (7) holds with $Q(j) \lesssim_\varepsilon 2^{j\varepsilon}$ whenever $\frac{\vec{1}}{p} \in \mathcal{C}^m(q)$. The proof then follows from the extension result of [15, Proposition 6.2]. \square

Claim 7. *Let $j \geq j_0$, $\frac{\vec{1}}{p} \in \mathcal{H}^m(q)$ and $q > 1$. Then there exists an $r = r(\frac{\vec{1}}{p}, q, n) > 0$ such that inequality (7) holds with $Q(j) \lesssim 2^{-rj}$.*

Proof. We observe that the point $(\frac{1}{q}, \frac{1}{q}, \dots, \frac{1}{q})$ is in the interior of $\mathcal{H}^m(q)$ and interpolate between Claim 5 and Claim 6 using [12, Theorem 7.2.2]. \square

3. PROOF OF PROPOSITION 4

A critical tool for the proof of Proposition 4 is a bound for the shifted maximal and square functions. Let φ and ψ be Schwartz functions on \mathbb{R}^n with compact Fourier supports such that $\widehat{\psi}$ is supported away from the origin. For $t > 0$ and $v \in \mathbb{R}^n$, we denote $\varphi_t^v(x) = t^n \varphi(tx - v)$ and similarly for $\psi_t^v(x)$.

With this notation, we have

Proposition C ([12, 22]). *Let $1 < p < \infty$. Then*

$$\left\| \sup_{r \in \mathbb{Z}} |f * \varphi_{2^r}^v| \right\|_{L^p(\mathbb{R}^n)} \lesssim \log(2 + |v|) \|f\|_{L^p(\mathbb{R}^n)}$$

and

$$\left\| \left(\sum_{r \in \mathbb{Z}} |f * \psi_{2^r}^v|^2 \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^n)} \lesssim \log(2 + |v|) \|f\|_{L^p(\mathbb{R}^n)}.$$

In the one-dimensional setting, Proposition C is proven in [22, Theorems 4.1, 5.1]. The first of the two inequalities is also contained in [25, Chapter II, 5.10]. The proof in the n -dimensional case can be found in [12, Proposition 7.5.1].

We next apply Proposition C to prove Proposition 4. Our argument follows the ideas from [22, 23].

Proof of Proposition 4. Given $j \in \mathbb{Z}$, let Λ_j be the $(m+1)$ -linear form given by

$$\Lambda_j(f_1, \dots, f_{m+1}) = \int_{\mathbb{R}^n} T_j(f_1, \dots, f_m)(x) f_{m+1}(x) dx.$$

Since $p > 1$, we observe that Proposition 4 will follow if we prove that

$$|\Lambda_j(f_1, \dots, f_{m+1})| \lesssim j^m \|\Omega\|_{L^1(\mathbb{S}^{mn-1})} \|f_{m+1}\|_{L^{p'}(\mathbb{R}^n)} \prod_{i=1}^m \|f_i\|_{L^{p_i}(\mathbb{R}^n)}$$

for $j \geq j_0$. To simplify notation, we write $\vec{x} = (x_1, \dots, x_m) \in \mathbb{R}^{mn}$ for $x_i \in \mathbb{R}^n$. We rewrite the form Λ_j as follows:

$$\begin{aligned} \Lambda_j(f_1, \dots, f_{m+1}) &= \sum_{k \in \mathbb{Z}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^{mn}} \widehat{K}_j^0(2^k \vec{\xi}) \left(\prod_{i=1}^m \widehat{f}_i(\xi_i) \right) f_{m+1}(x) e^{2\pi i x \cdot (\xi_1 + \dots + \xi_m)} d\vec{\xi} dx \\ &= \sum_{k \in \mathbb{Z}} \int_{\mathbb{R}^{mn}} \widehat{K}_j^0(2^k \vec{\xi}) \left(\prod_{i=1}^m \widehat{f}_i(\xi_i) \right) \widehat{f_{m+1}}(-\xi_1 - \dots - \xi_m) d\vec{\xi}. \end{aligned}$$

Notice that $\widehat{K}_j^0(2^k \cdot)$ is supported in the mn -dimensional annulus where $|\vec{\xi}| \sim 2^{j-k}$. Therefore, at least two of the vectors $\xi_1, \dots, \xi_m, (-\xi_1 - \dots - \xi_m)$ have to belong to an n -dimensional annulus of radius about 2^{j-k} . By a smooth partition of unity, it suffices to consider pieces of the form

$$\sum_{k \in \mathbb{Z}} \int_{\mathbb{R}^{mn}} \widehat{K}_j^0(2^k \vec{\xi}) \left(\prod_{i=1}^m \widehat{f}_i(\xi_i) \widehat{\phi}_i(2^{k-j} \xi_i) \right) \widehat{f_{m+1}}(-\xi_1 - \dots - \xi_m) \widehat{\phi_{m+1}}(2^{k-j}(-\xi_1 - \dots - \xi_m)) d\vec{\xi},$$

where ϕ_i , $i = 1, \dots, m+1$, are Schwartz functions with compact Fourier support such that at least two of them are Fourier supported away from the origin. We denote these two functions by ϕ_l and $\phi_{l'}$. Using Fourier inversion, the form can then be written as

$$\begin{aligned} &\sum_{k \in \mathbb{Z}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^{mn}} 2^{-kmn} K_j^0(2^{-k} \vec{y}) \left(\prod_{i=1}^m f_i * (\phi_i)_{2^{j-k}}(x - y_i) \right) f_{m+1} * (\phi_{m+1})_{2^{j-k}}(x) d\vec{y} dx \\ &= \sum_{k \in \mathbb{Z}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^{mn}} K_j^0(\vec{y}) \left(\prod_{i=1}^m f_i * (\phi_i)_{2^{j-k}}(x - 2^k y_i) \right) f_{m+1} * (\phi_{m+1})_{2^{j-k}}(x) d\vec{y} dx \\ &= \sum_{k \in \mathbb{Z}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^{mn}} K_j^0(\vec{y}) \prod_{i=1}^{m+1} f_i * (\phi_i)_{2^{j-k}}^{2^j y_i}(x) d\vec{y} dx, \end{aligned}$$

where we have set $y_{m+1} = 0$. We also formally set $p_{m+1} = p'$. For a fixed $\vec{y} \in \mathbb{R}^{mn}$, we estimate

$$\begin{aligned} &\left| \sum_{k \in \mathbb{Z}} \int_{\mathbb{R}^n} \prod_{i=1}^{m+1} f_i * (\phi_i)_{2^{j-k}}^{2^j y_i}(x) dx \right| \\ &\leq \int_{\mathbb{R}^n} \prod_{i \neq l, l'} \sup_{k \in \mathbb{Z}} |f_i * (\phi_i)_{2^{j-k}}^{2^j y_i}(x)| \prod_{r \in \{l, l'\}} \left(\sum_{k \in \mathbb{Z}} |f_r * (\phi_r)_{2^{j-k}}^{2^j y_r}(x)|^2 \right)^{1/2} dx \\ &\lesssim \prod_{i \neq l, l'} \left\| \sup_{k \in \mathbb{Z}} |f_i * (\phi_i)_{2^{j-k}}^{2^j y_i}| \right\|_{L^{p_i}(\mathbb{R}^n)} \prod_{r \in \{l, l'\}} \left\| \left(\sum_{k \in \mathbb{Z}} |f_r * (\phi_r)_{2^{j-k}}^{2^j y_r}|^2 \right)^{1/2} \right\|_{L^{p_r}(\mathbb{R}^n)} \\ &\lesssim \left(\prod_{i=1}^m \log(2 + 2^j |y_i|) \|f_i\|_{L^{p_i}(\mathbb{R}^n)} \right) \|f_{m+1}\|_{L^{p'}(\mathbb{R}^n)} \\ &\lesssim j^m \left(\prod_{i=1}^m \log(2 + |y_i|) \|f_i\|_{L^{p_i}(\mathbb{R}^n)} \right) \|f_{m+1}\|_{L^{p'}(\mathbb{R}^n)}, \end{aligned}$$

using Proposition C. Therefore, it suffices to show that

$$\int_{\mathbb{R}^{mn}} |K_j^0(\vec{y})| \prod_{i=1}^m \log(2 + |y_i|) d\vec{y} \lesssim \|\Omega\|_{L^1(\mathbb{S}^{mn-1})}.$$

We recall that $K_j^0 = K^0 * \beta_{2^j}$. Since β is a Schwartz function, we may estimate it as

$$|\beta(x)| \lesssim \sum_{k=1}^{\infty} 2^{-2mnk} \chi_{\mathbb{B}^{mn}(0, 2^k)}(x).$$

We further notice that the function $|K^0| * \chi_{\mathbb{B}^{mn}(0, 2^{k-j})}$ is supported in the set $\mathbb{B}^{mn}(0, 2^{k+1})$. Therefore,

$$\begin{aligned} & \int_{\mathbb{R}^{mn}} |K_j^0(\vec{y})| \prod_{i=1}^m \log(2 + |y_i|) d\vec{y} \\ & \lesssim \sum_{k=1}^{\infty} 2^{mnj-2mnk} \int_{\mathbb{R}^{mn}} |K^0| * \chi_{\mathbb{B}^{mn}(0, 2^{k-j})}(\vec{y}) \prod_{i=1}^m \log(2 + |y_i|) d\vec{y} \\ & \lesssim \sum_{k=1}^{\infty} k^m 2^{mnj-2mnk} \| |K^0| * \chi_{\mathbb{B}^{mn}(0, 2^{k-j})} \|_{L^1(\mathbb{R}^{mn})} \lesssim \|K^0\|_{L^1(\mathbb{R}^{mn})} \sum_{k=1}^{\infty} k^m 2^{-mnk} \\ & \lesssim \|K^0\|_{L^1(\mathbb{R}^{mn})} \lesssim \|\Omega\|_{L^1(\mathbb{S}^{mn-1})}, \end{aligned}$$

as desired. \square

4. COUNTEREXAMPLE

Proof of Proposition 3. Since $(\frac{1}{p_1}, \dots, \frac{1}{p_m}) \notin \mathcal{H}^m(q)$, there is a multiindex α such that (5) is not satisfied. Without a loss of generality, we can assume that $\alpha = (1, \dots, 1, 0, \dots, 0)$, with $|\alpha| = \kappa \geq 2$, since the cases $|\alpha| = 0$ and $|\alpha| = 1$ are trivial, and the operator is unaffected by permutations of the functions f_i .

For $i = 1, \dots, m$, let

$$\begin{cases} f_i(x) &= |x|^{-n/p_i} |\log|x||^{-\gamma/p_i} \chi_{|x| < \frac{1}{2}}(x), & \text{for } i = 1, \dots, \kappa, \\ f_i(x) &= |x|^{-n/p_i} |\log|x||^{-\gamma/p_i} \chi_{|x| > 2}(x), & \text{for } i = \kappa + 1, \dots, m, \end{cases}$$

for some $\gamma > 1$ to be determined, and note that $f_i \in L^{p_i}(\mathbb{R}^n)$. For $\theta = (\theta_1, \dots, \theta_m) \in \mathbb{S}^{mn-1}$, we define

$$\Omega(\theta) = \text{sgn}(\theta_1^1) \chi_{\mathbb{B}^{\kappa n}(0, \frac{1}{4\sqrt{\kappa}})}(\theta_1, \dots, \theta_\kappa) \left(\sum_{i=2}^{\kappa} |\theta_1 - \theta_i|^2 \right)^{-\frac{(\kappa-1)n}{2q}} \left| \log \sum_{i=2}^{\kappa} |\theta_1 - \theta_i|^2 \right|^{-\frac{\gamma}{q}},$$

where θ_1^1 denotes the first coordinate of θ_1 . Then Ω is an odd function in $L^q(\mathbb{S}^{mn-1})$. Indeed, using the co-area formula, [11, Appendix D], and setting $r_{\theta, \kappa} = \sqrt{1 - \sum_{i=1}^{\kappa} |\theta_i|^2}$, we have

$$\int_{\mathbb{S}^{mn-1}} |\Omega(\theta)|^q d\theta = \int_{\mathbb{B}^{\kappa n}} \int_{r_{\theta, \kappa} \mathbb{S}^{(m-\kappa)n-1}} |\Omega(\theta)|^q \frac{d\sigma(\theta_{\kappa+1}, \dots, \theta_m)}{r_{\theta, \kappa}} d\theta_1 \dots d\theta_\kappa.$$

Since $1 - \sum_{i=1}^k |\theta_i|^2 \approx 1$ when θ belongs to the support of Ω , the previous expression is bounded by

$$\begin{aligned} & \int_{\mathbb{B}^{\kappa n}(0, \frac{1}{4\sqrt{\kappa}})} \left(\sum_{i=2}^{\kappa} |\theta_1 - \theta_i|^2 \right)^{-\frac{(\kappa-1)n}{2}} \left| \log \sum_{i=2}^{\kappa} |\theta_1 - \theta_i|^2 \right|^{-\gamma} d\theta_1 \cdots d\theta_{\kappa} \\ & \lesssim \int_{\mathbb{B}^{(\kappa-1)n}(0, \frac{1}{4\sqrt{\kappa}})} |u|^{-(\kappa-1)n} |\log |u||^{-\gamma} du_2 \cdots du_{\kappa} < \infty, \end{aligned}$$

where the first inequality follows by the change of coordinates to $u_i = \theta_1 - \theta_i$ for $i = 2, \dots, \kappa$.

Let $x \in \mathbb{R}^n$ be a vector such that $x^1 > 10$. We denote $\bar{x} = (x, x, \dots, x) \in (\mathbb{R}^n)^m$. Then

$$T_{\Omega}^m(f_1, \dots, f_m)(x) = \int_{|y_1|, \dots, |y_{\kappa}| < \frac{1}{2}} \cdots \int_{|y_{\kappa+1}|, \dots, |y_m| > 2} \left(\prod_{i=1}^m |y_i|^{-n/p_i} |\log |y_i||^{-\gamma/p_i} \right) \frac{\Omega\left(\frac{\bar{x}-y}{|\bar{x}-y|}\right)}{|\bar{x}-y|^{mn}} dy.$$

Note that, since $|y_1| < \frac{1}{2}$, the first coordinate of $(\bar{x} - y)_1$ is positive, and thus the integrand in the above integral is positive. Therefore, we can estimate it from below by the integral over the smaller region $A \times B$, where

$$A = \left\{ (y_1, \dots, y_{\kappa}) \in \mathbb{B}^{\kappa n} \left(0, \frac{1}{|x|}\right) : |y_1| \approx \cdots \approx |y_{\kappa}| \approx |y_1 - y_2| \approx \cdots \approx |y_1 - y_{\kappa}| \right\}$$

and

$$B = \mathbb{B}^{n(m-\kappa)} \left(\underbrace{(x, \dots, x)}_{m-\kappa}, \frac{|x|}{2} \right).$$

We write $y = (z, z')$, where $z = (y_1, \dots, y_{\kappa})$ and $z' = (y_{\kappa+1}, \dots, y_m)$. In $A \times B$, we have for $i = 2, \dots, \kappa$,

$$|y_1 - y_i| \approx |z| < \frac{1}{|x|} \quad \text{and} \quad |\bar{x} - y| \approx |z'| \approx |x|,$$

and therefore, for $y \in A \times B$,

$$\begin{aligned} \Omega\left(\frac{\bar{x}-y}{|\bar{x}-y|}\right) &= \left(\sum_{i=2}^{\kappa} \left| \frac{y_1 - y_i}{|\bar{x}-y|} \right|^2 \right)^{-\frac{(\kappa-1)n}{2q}} \left| \log \sum_{i=2}^{\kappa} \left| \frac{y_1 - y_i}{|\bar{x}-y|} \right|^2 \right|^{-\frac{\gamma}{q}} \\ &\approx \left(\frac{|z|}{|x|} \right)^{-\frac{(\kappa-1)n}{q}} \left| \log \frac{|z|}{|x|} \right|^{-\frac{\gamma}{q}} \\ &\gtrsim |x|^{\frac{(\kappa-1)n}{q}} |z|^{-\frac{(\kappa-1)n}{q}} |\log |z||^{-\frac{\gamma}{q}}. \end{aligned}$$

Setting $\frac{1}{p_{\alpha}} = \sum_{i=1}^{\kappa} \frac{1}{p_i}$ and $\frac{1}{\tilde{p}_{\alpha}} = \sum_{i=\kappa+1}^m \frac{1}{p_i}$, we estimate the absolute value of $T_{\Omega}^m(f_1, \dots, f_m)(x)$ from below by

$$\begin{aligned} & \int_B \int_A \left(\prod_{i=1}^m |y_i|^{-n/p_i} |\log |y_i||^{-\gamma/p_i} \right) \frac{\Omega\left(\frac{\bar{x}-y}{|\bar{x}-y|}\right)}{|\bar{x}-y|^{mn}} dy \\ & \gtrsim |x|^{-mn + \frac{(\kappa-1)n}{q}} \int_B |z'|^{-\frac{n}{\tilde{p}_{\alpha}}} (\log |z'|)^{-\frac{\gamma}{\tilde{p}_{\alpha}}} dz' \int_A |z|^{-n\left(\frac{1}{p_{\alpha}} + \frac{\kappa-1}{q}\right)} |\log |z||^{-\frac{\gamma}{p_{\alpha}} - \frac{\gamma}{q}} dz \\ & \gtrsim (\log |x|)^{-\frac{\gamma}{\tilde{p}_{\alpha}}} |x|^{-mn + \frac{(\kappa-1)n}{q} - \frac{n}{\tilde{p}_{\alpha}} + (m-\kappa)n} \int_A |z|^{-n\left(\frac{1}{p_{\alpha}} + \frac{\kappa-1}{q}\right)} |\log |z||^{-\frac{\gamma}{p_{\alpha}} - \frac{\gamma}{q}} dz, \end{aligned}$$

which diverges when $\frac{1}{p_\alpha} + \frac{\kappa-1}{q} > \kappa = |\alpha|$.

When $\frac{1}{p_\alpha} + \frac{\kappa-1}{q} = \kappa$, the exponents in the previous expression simplify to

$$(\log |x|)^{-\frac{\gamma}{p_\alpha}} |x|^{-\frac{n}{p}} \int_A |z|^{-\kappa n} |\log |z||^{-\frac{\gamma}{p_\alpha} - \frac{\gamma}{q}} dz,$$

which is equivalent to

$$(\log |x|)^{1 - \frac{\gamma}{p} - \frac{\gamma}{q}} |x|^{-\frac{n}{p}}.$$

Therefore,

$$\int_{\mathbb{R}^n} |T_\Omega^m(f_1, \dots, f_m)(x)|^p dx \gtrsim \int_{\{x \in \mathbb{R}^n: x^1 > 10\}} |x|^{-n} (\log |x|)^{p - \gamma - \frac{\gamma p}{q}} dx,$$

which is finite if and only if $p - \gamma - \frac{\gamma p}{q} < -1 \Leftrightarrow \gamma > \frac{p+1}{\frac{p}{q}+1}$. Since $q > 1$, we can choose $1 < \gamma < \frac{p+1}{\frac{p}{q}+1}$ in the definitions of f_1, \dots, f_m , and Ω to obtain the counterexample. \square

STATEMENTS AND DECLARATIONS

Conflict of interest. On behalf of all authors, the corresponding author states that there is no conflict of interest.

Data availability. Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

REFERENCES

- [1] A. P. Calderón, A. Zygmund. *On the existence of certain singular integrals*, Acta Math. **88** (1952), no. 1, 85–139.
- [2] A. P. Calderón, A. Zygmund. *On singular integrals*, Amer. J. Math. **78** (1956), 289–309.
- [3] J.-K. Bak, D. M. Oberlin, A. Seeger. *Restriction of Fourier transforms to curves and related oscillatory integrals*, American Journal of Mathematics **131**(2) (2009), 277–311.
- [4] A. P. Calderón. *Commutators of singular integral operators*, Proc. Nat. Acad. Sci. U.S.A. **53** (1965), 1092–1099.
- [5] M. Christ. *Weak type (1, 1) bounds for rough operators I*, Ann. Math. **128** (1988), 19–42.
- [6] M. Christ, J. L. Rubio de Francia. *Weak type (1, 1) bounds for rough operators II*, Invent. Math. **93** (1988), 225–237.
- [7] R. R. Coifman, Y. Meyer. *On commutators of singular integrals and bilinear singular integrals*, Trans. Amer. Math. Soc. **212** (1975), 315–331.
- [8] R. R. Coifman, G. Weiss. *Extensions of Hardy spaces and their use in analysis*, Bull. Amer. Math. Soc. **83** (1977), 569–645.
- [9] W. C. Connett. *Singular integrals near L^1 , in Harmonic analysis in Euclidean spaces, Part 1*, (Williamstown 1978), Proc. Sympos. Pure Math. 35, Amer. Math. Soc., Providence (1979), 163–165.
- [10] G. Diestel, L. Grafakos, P. Honzik, Z. Si, M. E. Terwilleger. *Method of rotations for bilinear singular integrals*, Commun. Math. Anal. 2011, Conference 3, 99–107.
- [11] L. Grafakos. *Classical Fourier Analysis*. 3rd Ed., GTM **249**, Springer NY, 2014.
- [12] L. Grafakos. *Modern Fourier Analysis*. 3rd Ed., GTM **250**, Springer NY, 2014.
- [13] L. Grafakos, D. He, P. Honzik. *Rough bilinear singular integrals*, Adv. Math. **326** (2018), 54–78.
- [14] L. Grafakos, D. He, P. Honzik, B. J. Park. *Initial $L^2 \times \dots \times L^2$ bounds for multilinear operators*, Preprint, <https://arxiv.org/abs/2010.15312>
- [15] L. Grafakos, D. He, P. Honzik, B. J. Park. *Multilinear rough singular integrals operators*, Preprint, <https://arxiv.org/abs/2207.00764>
- [16] L. Grafakos, D. He, L. Slavíková. *Failure of the Hörmander kernel condition for multilinear Calderón–Zygmund operators*, C. R. Math. Acad. Sci. Paris **357** (2019), no. 4, 382388.

- [17] L. Grafakos, D. He, L. Slavíková. $L^2 \times L^2 \rightarrow L^1$ boundedness criteria, *Math. Ann.* **376** (2020), no. 1–2, 431–455.
- [18] L. Grafakos, R. H. Torres. *Multilinear Calderón-Zygmund Theory*, *Adv. Math.* **165** (2002), 124–164.
- [19] D. He, B. J. Park. *Improved estimates for bilinear rough singular integrals*, *Math. Ann.* **386** (2023), no. 3–4, 1951–1978.
- [20] S. Hofmann. *Weak type (1, 1) boundedness of singular integrals with nonsmooth kernels*, *Math. Soc.* **103** (1988), 260–264.
- [21] C. Kenig, E. M. Stein. *Multilinear estimates and fractional integration*, *Math. Res. Lett.* **6** (1999), no. 1, 1–15.
- [22] C. Muscalu. *Calderón commutators and the Cauchy integral on Lipschitz curves revisited: I. First commutator and generalizations*, *Rev. Mat. Iberoam.* **30** (2014), no. 2, 727–750.
- [23] C. Muscalu. *Calderón commutators and the Cauchy integral on Lipschitz curves revisited II. The Cauchy integral and its generalizations*, *Rev. Mat. Iberoam.* **30** (2014), no. 3, 1089–1122.
- [24] A. Seeger. *Singular integral operators with rough convolution kernels*, *J. Amer. Math. Soc.* **9** (1996), 9–105.
- [25] E. M. Stein. *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*. Princeton Mathematical Series, **43**, Springer, New Jersey, 1993.

GEORGIOS DOSIDIS. DEPARTMENT OF MATHEMATICAL ANALYSIS, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY, SOKOLOVSKÁ 83, 186 75 PRAHA 8, CZECH REPUBLIC

E-mail address: `Dosidis@karlin.mff.cuni.cz`

LENKA SLAVÍKOVÁ. DEPARTMENT OF MATHEMATICAL ANALYSIS, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY, SOKOLOVSKÁ 83, 186 75 PRAHA 8, CZECH REPUBLIC

E-mail address: `slavikova@karlin.mff.cuni.cz`