

Unpolarized QED parton distribution functions in NLO

A.B. Arbuzov^{a,b*}, U.E. Voznaya^{a,b†}

December 22, 2022

^a*Joint Institute for Nuclear Research, Joliot-Curie str. 6, Dubna, 141980, Moscow region, Russia*

^b*Dubna state university, Universitetskaya str. 19, Dubna, 141980, Moscow region, Russia*

Abstract

Perturbative solutions for unpolarized QED parton distribution and fragmentation functions are presented explicitly in the next-to-leading logarithmic approximation. The scheme of iterative solution of QED evolution equations is described in detail. Terms up to $\mathcal{O}(\alpha^3 L^2)$ are calculated analytically, where $L = \ln(\mu^2/m_e^2)$ is the large logarithm which depends on the factorization energy scale $\mu \gg m_e$. The results are process independent, they are relevant for future high-precision experiments.

PACS: 12.20.-m, 13.40.-f

1 Introduction

The studies of high-energy physics processes at future high-luminosity electron-positron colliders such as the FCC-ee [1] and CEPC [2] require very precise calculations of QED radiative corrections for construction of sufficiently accurate theoretical predictions of these processes [3]. In spite of the existence of powerful methods for calculations in perturbative QED, it is still very difficult to compute complete radiative corrections in $\mathcal{O}(\alpha^3)$. On the other hand, we know that higher-order radiative corrections enhanced by the so-called large logarithms typically provide the bulk of the effect. For sufficiently inclusive observables, all terms enhanced by the large logs can be calculated with the help of QED parton distribution function (PDF) approach which is based on the factorization theorem.

The parton distribution functions approach in QED was developed analogously to the QCD one by E.A. Kuraev and V.S. Fadin [4] in 1980-s as a step in preparation for high-precision measurements at LEP. Actually even earlier, it has been already applied to quasi-elastic neutrino scattering in Ref. [5]. The approach is widely used for calculation of QED radiative corrections in high energy physics. QED evolution equation are just a reduction of the QCD DGLAP (Dokshitzer-Gribov-Lipatov-Altarelli-Parisi) evolution equations [6, 7, 8] to the abelian case of QED. The equations are based on the renormalization group and scale

*arbuzov@theor.jinr.ru

†voznaya@theor.jinr.ru

invariance. They allow to effectively account the logarithmic dependence on the factorization scale.

The possibility to exponentiate a part of the QED parton distribution functions was already considered in [4], see more details in [9]. The exponentiation is natural because of the Yennie-Frautschi-Suura theorem [10]. And it can be done based on the known exact solution of the evolution equations in the limit of soft radiation [6]. But in the present paper we will consider only perturbative order-by-order representations of QED PDFs, while exponentiation of our results will be presented elsewhere.

Perturbative solutions of QED evolution equations in the leading logarithmic approximation are known up to the fifth order for non-singlet electron distribution functions [11, 12]. Singlet higher-order leading log contributions to this function were computed in [13]. One can see that the existing results for electron PDFs in the collinear leading logarithmic approximation are well cross-checked and used for estimates of radiative corrections to a wide class of processes.

As concerns QED PDFs in the next-to-leading logarithmic approximation, it is much less elaborated and exploited. In the $\mathcal{O}(\alpha^2 L)$ spacelike electron PDFs were first considered in Ref. [14]. Later also timelike electron PDFs in the same order were used to describe radiative corrections to muon decay spectrum [15] and deep inelastic scattering [16]. The resummation in the leading order in the unpolarized case was considered in the work [17]. Relatively recently in the series of papers [18, 19, 20], the results for the leading and next-to-leading radiative corrections to electron-positron annihilation into a virtual photon or Z boson up to the $\mathcal{O}(\alpha^6 L^5)$ order were presented. Those results were obtained with the help of spacelike QED PDFs, but explicit expressions for the functions were not given. Moreover, those results are somewhat incomplete since the transitions from electrons into positrons were missed.

In some recent works the evolution was considered with alternative factorization (Delta, which is DIS-like) and renormalization ($\alpha(m_Z)$ and G_μ) schemes [21, 22, 23, 24]. Electrons, and then all "QED flavors", i.e., other charged leptons and quarks were considered.

In this article, we describe in detail the iterative solution of evolution equations of parton distribution functions in QED. The results up to the third iteration are shown explicitly. We consider process-independent PDFs which describe the probability density of finding massless partons (electron, positron or photon) inside electrons and photons. Both spacelike and timelike QED PDFs are evaluated.

2 Master Formula

The cross-section of a high-energy $ab \rightarrow cd$ process with charged particles in the initial and final states in the next-to-leading order (NLO) approximation can be represented in the following form [25]

$$d\sigma_{ab \rightarrow cd}^{\text{NLO}} = \sum_{i,j,k,l} \int_{\bar{z}_1}^1 dz_1 \int_{\bar{z}_2}^1 dz_2 D_{ia}^{\text{str}}(z_1, \frac{\mu_0^2}{\mu^2}) D_{jb}^{\text{str}}(z_2, \frac{\mu_0^2}{\mu^2}) \\ \times \left(d\sigma_{ij \rightarrow kl}^{\text{Born}}(z_1, z_2) + d\bar{\sigma}_{ij \rightarrow kl}^{(1)}(z_1, z_2) + \mathcal{O}(\alpha^2 L^0) \right)$$

$$\times \int_{\bar{y}_1}^1 \frac{dy_1}{Y_1} \int_{\bar{y}_2}^1 \frac{dy_2}{Y_2} D_{ck}^{\text{frg}}\left(\frac{y_1}{Y_1}, \frac{\mu_0^2}{\mu^2}\right) D_{dl}^{\text{frg}}\left(\frac{y_2}{Y_2}, \frac{\mu_0^2}{\mu^2}\right) + \mathcal{O}\left(\frac{\mu_0^2}{\mu^2}\right), \quad (1)$$

where \bar{z}_i and \bar{y}_i are some minimal energy fractions defined by experimental conditions; z_i are the energy fractions of the incoming partons; Y_i are the energy fractions of the outgoing particles; $d\bar{\sigma}_{ij \rightarrow kl}^{(0)}(z_1, z_2)$ and $d\bar{\sigma}_{ij \rightarrow kl}^{(1)}(z_1, z_2)$ are the Born massless parton cross-section and the $\mathcal{O}(\alpha)$ contribution to it. The bar in $d\bar{\sigma}_{ij \rightarrow kl}^{(1)}(z_1, z_2)$ denotes application of a subtraction scheme to exclude mass singularities. Here we will use the standard $\overline{\text{MS}}$ scheme. The spacelike (marked "str"), i.e., structure, and timelike (marked "frg"), i.e., fragmentation, functions D_{ij} depend on energy fractions and the ratio $\frac{\mu_0^2}{\mu^2}$, which is the argument of the large logarithm, respectively. Note that in QED the renormalization scale μ_0 is typically chosen to be equal to the electron mass. The factorization scale is usually chosen to be equal to the energy scale of the hard sub-process. For example, for the process of electron positron annihilation into a Z boson, we have $\mu = M_Z$ and $\mu_0 = m_e$ and the large logarithm is really large numerically, $L = \ln(M_Z^2/m_e^2) \approx 24$.

As the result, the cross-section in the NLO approximation takes into account the QED radiative corrections enhanced by the large logarithms and reads

$$d\sigma_{ab \rightarrow cd}^{\text{NLO}} = d\sigma_{ab \rightarrow cd}^{(0)} \left\{ 1 + \sum_{k=1}^{\infty} \left(\frac{\alpha}{2\pi}\right)^k \sum_{l=k-1}^k c_{k,l} L^l + \mathcal{O}(\alpha^k L^{k-2}) \right\}, \quad (2)$$

where $c_{k,l}$ are the coefficients to be computed. The terms of the type $\alpha^k L^k$ provide the leading order (LO) logarithmic approximation, and the ones of the type $\alpha^k L^{k-1}$ yield the NLO contribution.

3 QED Evolution Equations

The main goal of the article is to describe how to compute the QED parton distribution functions. Let us consider QED evolution equations for PDFs in the spacelike region. The equations are induced by the renormalization group and have the following form, see, e.g., Ref. [26]:

$$D_{ba}(x, \mu^2, \mu_0^2) = \delta(1-x)\delta_{ba} + \sum_{i=e, \bar{e}, \gamma} \int_{\mu_0^2}^{\mu^2} \frac{dt\alpha(t)}{2\pi t} \int_x^1 \frac{dy}{y} D_{ia}(y, t, \mu_0) P_{bi}\left(\frac{x}{y}\right), \quad (3)$$

where index a corresponds to the initial particle, e.g., an electron; and indices b and i mark QED partons which can be photons(γ) or massless electrons (e) and positrons (\bar{e}). The parton distribution function $D_{ba}(x, \mu^2)$ describes the probability density to find the massless parton b in the initial (massive) particle a with the of energy fraction x of the initial particle energy. Here μ is the factorization scale which is the characteristic energy of a hard sub-process and μ_0 is the renormalization scale. $P_{ji}(x)$ are called splitting functions and describe a perturbative transformation of parton i into parton j which takes the energy fraction x . They can be expanded in a series in α

$$P_{ji}(x) = P_{ji}^{(0)}(x) + \frac{\alpha}{2\pi} P_{ji}^{(1)}(x) + \mathcal{O}(\alpha^2). \quad (4)$$

The splitting function $P_{ee}^{(1)}(x)$ includes contributions of different sub-processes. Expression for this function can be derived from the analogous expression for quark-quark function in QCD $P_{qq}^{(1)}(x)$ [27] with singlet and non-singlet parts:

$$P_{q_i q_k} = \delta_{ik} P_{qq}^{NS} + P_{qq}^S, \quad (5)$$

where indices i and k denote quark flavors. In reduction to QED we have only one flavor¹. The singlet part of our function corresponds to Feynman diagrams with discontinuous electron lines between the initial and final electrons. The non-singlet part corresponds to the Feynman diagrams with continuous electron line. The non-singlet part contains contributions from initial state photon radiation, final state fermion pair radiation, where the fermions are neither electron nor positron and interference. The singlet part corresponds to electron-positron pair final state radiation [14]. The splitting function of electron in positron type $P_{e\bar{e}}^{(1)}(x)$ also have singlet and non-singlet parts, so the functions $P_{ee}^{(1)}(x)$ and $P_{e\bar{e}}^{(1)}(x)$ are

$$P_{ee}^{(1)} = P_{ee}^{NS} + P_{ee}^S, \quad P_{e\bar{e}}^{(1)} = P_{e\bar{e}}^{NS} + P_{e\bar{e}}^S, \quad (6)$$

There are obvious equalities [27] connecting splitting functions which involve electrons and positrons because of the C parity conservation in QED:

$$P_{ee} = P_{\bar{e}\bar{e}}, \quad (7)$$

$$P_{e\bar{e}} = P_{\bar{e}e}, \quad (8)$$

$$P_{e\gamma} = P_{\bar{e}\gamma}, \quad (9)$$

$$P_{\gamma e} = P_{\gamma\bar{e}}, \quad (10)$$

and at the two loop level there is a relation between singlet contributions [27]:

$$P_{e\bar{e}}^S = P_{ee}^S. \quad (11)$$

We consider spacelike and timelike parton distribution functions (or, in another notation, structure and fragmentation functions). Spacelike functions are denoted $[P^{(i)}]_S$, and timelike $[P^{(i)}]_T$. The spacelike functions correspond to transitions from massive particles into massless ones, and the timelike ones correspond to transition from massless particles into massive ones. The difference between timelike and spacelike functions appear in QED only in NLO [26] and only in the Θ parts. In the leading order spacelike and timelike functions are equal. Δ parts of timelike and spacelike functions are also equal. There is the relation between spacelike and timelike functions in the leading order:

$$-x[P_{ee}^{(0)}]_T\left(\frac{1}{x}\right) = [P_{ee}^{(0)}]_S(x), \quad (12)$$

which is called Gribov-Lipatov relation [28]. It is violated in the higher orders:

$$[P^{(i)}]_T(x) = [P^{(i)}]_S(x) + \Delta(x). \quad (13)$$

¹The possibility to include in QED, e.g., muons and then have $N_f = 2$ is straightforward.

The $\Delta(x)$ term can be calculated using analytical continuation of the timelike function to the nonphysical region $x > 1$ [28]. For electron-electron function, the correction to the Gribov-Lipatov relation reads

$$\begin{aligned} \Delta_{ee}(x) = & \left(\frac{\alpha}{2\pi}\right)^2 C_f \left[4 \frac{1+x^2}{1-x} \ln x \ln(1-x) \right. \\ & \left. + \left(\frac{6}{1-x} - 5 - 5x \right) \ln x + \left(1+x - 2 \frac{1+x^2}{1-x} \right) \ln^2 x \right]. \end{aligned} \quad (14)$$

3.1 Running coupling constant

The expression for the QED running coupling constant in the $\overline{\text{MS}}$ scheme can be found, e.g., in [29, 30],

$$\alpha(\mu^2) = \frac{\alpha(\mu_0)}{1 + \overline{\Pi}(\mu, \mu_0, \alpha(0))}, \quad (15)$$

with

$$\begin{aligned} \overline{\Pi}(\mu, \mu_0, \alpha(0)) = & \frac{\alpha(0)}{\pi} \left(\frac{5}{9} - \frac{L}{3} \right) + \left(\frac{\alpha(0)}{\pi} \right)^2 \left(\frac{55}{48} - \zeta_3 - \frac{L}{4} \right) \\ & + \left(\frac{\alpha(0)}{\pi} \right)^3 \left(\frac{-L^2}{24} \right) + \dots \end{aligned} \quad (16)$$

where $L = \ln(\mu^2/\mu_0^2)$ is again the large logarithm. After expansion, we get

$$\begin{aligned} \alpha(\mu^2) = & \frac{\alpha(\mu_0)}{2\pi} \left\{ 1 + \frac{\alpha(0)}{2\pi} \left(-\frac{10}{9} + \frac{2}{3}L \right) + \left(\frac{\alpha(0)}{2\pi} \right)^2 \left(-\frac{1085}{324} + 4\zeta_3 \right. \right. \\ & \left. \left. - \frac{13}{27}L + \frac{4}{9}L^2 \right) + \mathcal{O}(\alpha^3(0)) \right\}, \end{aligned} \quad (17)$$

where $\zeta_n \equiv \zeta(n)$ is the Riemann zeta function. Here we put $\mu_0 = m_e$ and assume $\alpha(\mu_0^2) \approx \alpha(0) \equiv \alpha$.

3.2 Convolution and plus prescription

The evolution equations involve the convolution operation

$$(f \otimes g)(x) \equiv \int_0^1 dz \int_0^1 dy f(z)g(y)\delta(x-yz) = \int_0^1 \frac{dz}{z} f(z)g\left(\frac{x}{z}\right). \quad (18)$$

Many of relevant functions have a regularization at $x \rightarrow 1$ and can be represented as a sum of the so-called Θ and Δ parts:

$$f(x) = \lim_{\Delta \rightarrow 0} \left(f_{\Theta}(x)\Theta(1-x-\Delta) + f_{\Delta}\delta(1-x) \right), \quad (19)$$

where $\Theta(x)$ is the standard Heaviside step function. In many cases, such functions can be regularized with the help of the *plus prescription* which acts in an integral with a regular

function as follows

$$\int_z^1 dx [f(x)]_+ g(x) = \int_0^1 dx f(x) [g(x)\Theta(x-z) - g(1)]. \quad (20)$$

If function $f(x)$ is regularized by the plus prescription, it satisfies the sum rule

$$f_\Delta = - \int_0^{1-\Delta} f_\Theta(z) dz. \quad (21)$$

For the convolution of two functions with Δ parts we get

$$(f \otimes g)_\Theta(z) = \lim_{\Delta \rightarrow 0} \left\{ \int_{z/(1-\Delta)}^{1-\Delta} \frac{dx}{x} f_\Theta(x) g_\Theta\left(\frac{z}{x}\right) + f_\Delta g_\Theta(z) + f_\Theta(z) g_\Delta \right\}. \quad (22)$$

Note that in this way we get only the Θ part of the convolution. If the result satisfies sum rule (21), its Δ part is easily restored. If the sum rule is not applicable, the Δ part of a convolution can be calculated as

$$(f \otimes g)_\Delta = f_\Delta g_\Delta - \int_{1-\Delta}^1 dy f(y) \int_{1-\Delta}^{\frac{1-\Delta}{y}} g(x) dx, \quad (23)$$

as follows from the definitions (18) and (19).

3.3 Initial conditions

We are going to solve evolution equations (3) by iterations. So we need some initial conditions. In the NLO approximation in the $\overline{\text{MS}}$ scheme, they are [31, 18]

$$D_{ee}^{(0)}(x, \mu^2) = \delta(1-x) + \frac{\alpha}{2\pi} d_{ee}^{(1)}(x), \quad (24)$$

$$D_{\gamma e}^{(0)}(x, \mu^2) = \frac{\alpha}{2\pi} d_{\gamma e}^{(1)}(x), \quad (25)$$

$$D_{e\gamma}^{(0)}(x, \mu^2) = \frac{\alpha}{2\pi} d_{e\gamma}^{(1)}(x), \quad (26)$$

$$D_{e\bar{e}}^{(0)}(x, \mu^2) = \frac{\alpha}{2\pi} d_{e\bar{e}}^{(1)}(x) = 0, \quad (27)$$

$$\begin{aligned} d_{ee}^{(1)}(x) &= \left[\frac{1+x^2}{1-x} \left(\ln \frac{\mu_0^2}{m_e^2} - 1 - 2 \ln(1-x) \right) \right]_+ \\ &= \left[\frac{1+x^2}{1-x} (-1 - 2 \ln(1-x)) \right]_+, \end{aligned} \quad (28)$$

note that here and in what follows we apply the natural choice of the QED renormalization constant $\mu_0 = m_e$ [18],

$$d_{\gamma e}^{(1)}(x) = -\frac{1 + (1-x)^2}{2x}(2 \ln x + 1), \quad (29)$$

$$d_{e\gamma}^{(1)}(x) = 0. \quad (30)$$

They are fixed with the help of the known results of perturbative calculations. Using the complete result in first order $\mathcal{O}(\alpha)$ we can fix functions $d_{ba}^{(1)}(x)$ in a chosen subtraction scheme. In any case, the following equality should be held

$$\sigma_{e\bar{e}\rightarrow\gamma^*}^{(1)}(x) = \bar{\sigma}_{e\bar{e}\rightarrow\gamma^*}^{(1)}(x) + 2\frac{\alpha}{2\pi} [P_{ee}^{(0)}(x)L + d_{ee}^{(1)}(x)] + \mathcal{O}\left(\frac{m_e^2}{\mu^2}\right), \quad (31)$$

where $\sigma^{(1)}$ is the complete one-loop contribution to the electron-positron annihilation process into a virtual photon $\bar{e} + e \rightarrow \gamma^*$ in the standard QED with massive electrons. Here x is the energy fraction of the produced virtual photon with respect to the total c.m.s. energy of the initial particles. To restore the mass dependence in the results of calculations within a massless theory (up to $\mathcal{O}(m_e^2/\mu^2)$ terms) one can use the so-called *massification* procedure. The above equation is an example of this procedure.

We can fix the next-to-leading contributions to the splitting functions and the subtracted one-loop cross-sections $\bar{\sigma}_{ij\rightarrow kl}^{(1)}$ with the help of a complete $\mathcal{O}(\alpha^2)$ result, e.g., for the same process of electron-positron annihilation with initial state radiation corrections [14, 18]. Note that separation of term of the right-hand side of Eq. (31) depends on the chosen scheme and factorization scale.

3.4 Iterative solution

Evolution equations can be solved using the iterative method with the initial conditions (24). On the first step, we substitute the initial conditions for $D_{ia}(e, s')$ in the evolution equations (3) under the integral on the right-hand side. The result of the first iteration reads

$$\begin{aligned} D_{ee}^{(1)}(x, \mu) &= D_{ee}^{(0)} + \frac{\alpha}{2\pi} \left(P_{ee} \otimes D_{ee}^{(0)} + P_{e\gamma} \otimes D_{\gamma e}^{(0)} + P_{e\bar{e}} \otimes D_{\bar{e}e}^{(0)} \right) \\ &= \delta(1-x) + d_{ee}^{(1)}(x) + \frac{\alpha}{2\pi} \left(P_{ee}^{(0)} + \frac{\alpha}{2\pi} P_{ee}^{(1)} \right) \otimes (\delta(1-x) + d_{ee}^{(1)}(x)) \\ &\quad + \left(P_{eg}^{(0)} + \frac{\alpha}{2\pi} P_{eg}^{(1)} \right) \otimes d_{\gamma e}^{(1)}(x) = 1 + \frac{\alpha}{2\pi} d_{ee}^{(1)}(x) + \frac{\alpha}{2\pi} L P_{ee}^{(0)}, \end{aligned} \quad (32)$$

$$\begin{aligned} D_{\gamma e}^{(1)}(x, \mu) &= D_{\gamma e}^{(0)} + \frac{\alpha}{2\pi} (P_{\gamma\gamma} \otimes D_{\gamma e}^{(0)} + P_{\gamma e} \otimes D_{ee}^{(0)} + P_{\gamma\bar{e}} \otimes D_{\bar{e}e}^{(0)}) = d_{\gamma e}^{(1)}(x) \\ &\quad + \frac{\alpha}{2\pi} \left(P_{\gamma\gamma}^{(0)} + \frac{\alpha}{2\pi} P_{\gamma\gamma}^{(1)} \right) \otimes d_{\gamma e}^{(1)}(x) + \frac{\alpha}{2\pi} \left(P_{\gamma e}^{(0)} + \frac{\alpha}{2\pi} P_{\gamma e}^{(1)} \right) \otimes (\delta(1-x) \\ &\quad + d_{ee}^{(1)}(x)) = \frac{\alpha}{2\pi} d_{\gamma e}^{(1)}(x) + \frac{\alpha}{2\pi} L P_{\gamma e}^{(0)}, \end{aligned} \quad (33)$$

$$D_{e\bar{e}}^{(1)}(x, \mu) = D_{e\bar{e}}^{(0)} + \frac{\alpha}{2\pi} (P_{ee}^{(0)} \otimes D_{e\bar{e}}^{(0)} + P_{e\bar{e}}^{(0)} \otimes D_{ee}^{(0)} + P_{e\gamma}^{(0)} \otimes D_{e\gamma}^{(0)}) = 0. \quad (34)$$

On the second iteration step, we put the result of the first step under the integral (3). The results of the second and the third iterations are presented in the Appendix B.

We also show the results of iterations as functions of z . The difference between spacelike and timelike functions is only in $P_{ij}^{(1)}$, so the parton distributions $D_{ij}^{(1)}$ are equal for spacelike and timelike cases. As we are interested only in LO and NLO contributions, we can omit terms proportional to $(\frac{\alpha}{2\pi})^{k+2} L^k$. The results are presented in the Appendix C. We used splitting functions and massless Wilson coefficients [18, 32, 27, 14] which are given in the Appendix A.

4 Conclusions

In this way, we described the iterative method of solving QED evolution equations. Extension of the presented calculations to higher orders is straightforward. But as concerns next-to-leading contributions, e.g. $\mathcal{O}(\alpha^4 L^3)$, they do not seem numerically relevant for the near future collider experiments. Note that our results are scheme and scale dependent. Application of other schemes and variation of renormalization and factorization scales are possible within the same line of calculations. Here, we presented only the electron parton distribution functions, photon PDFs will be presented elsewhere.

The calculated QED parton distribution functions can be applied to various high-energy processes with participation of electrons and/or positrons. Note that the PDFs here are treated in the collinear approximation which should match the experimental conditions of particle registration of the process under study. In other words, the physical observable should not be sensitive to transverse momenta of initial and/or final state radiation. In any case, the results obtained within the collinear approximation will serve as benchmarks for Monte Carlo simulations which describe the complete kinematics of events.

As we mentioned in the introduction, our results partially intersect with the ones given in Refs. [18, 19, 20]. Results a detailed comparison will be presented later.

Acknowledgments

A.A. is grateful for support to the RFBR grant N 20-02-00441.

Appendix A

Process-independent parton splitting functions read

$$\begin{aligned}
P_{ee}^{(0)}(x) &= \left[\frac{1+x^2}{1-x} \right]_+, & P_{e\gamma}^{(0)}(x) &= x^2 + (1-x)^2, & P_{e\bar{e}}^{(0)}(x) &= 0, & (A.1) \\
P_{\gamma e}^{(0)}(x) &= \frac{1+(1-x)^2}{x}, & P_{\gamma\gamma}^{(0)}(x) &= \frac{\beta_0}{2} \delta(1-x), \\
P_{e\gamma}^{(1)}(x) &= C_f T_f (4 - 9x - (1-4x) \ln x - (1-2x) \ln^2 x + 4 \ln(1-x) + \\
&\quad + (2(\ln(1-x) - \ln x)^2 - 4(\ln(1-x) - \ln x) - 4\zeta(2) + 10) P_{e\gamma}^{(0)}(x)), \\
P_{\gamma e}^{(1)}(x) &= C_f^2 \left(-\frac{5}{2} - \frac{7}{2}x + \left(2 + \frac{7}{2}x \right) \ln x - \left(1 - \frac{1}{2}x \right) \ln x^2 - 2x \ln(1-x) \right. \\
&\quad \left. - (3 \ln(1-x) + \ln^2(1-x)) P_{\gamma e}^{(0)}(x) \right) + C_f \left(-\frac{4x}{3} - \left(\frac{20}{9} + \frac{4 \ln(1-x)}{3} \right) P_{\gamma e}^{(0)}(x) \right),
\end{aligned}$$

$$\begin{aligned}
P_{ee}^{NS}(x) &= C_f^2 \left(\left(-2 \ln x \ln(1-x) - \frac{3}{2} \ln x \right) \frac{1+x^2}{1-x} - \left(\frac{3}{2} + \frac{7}{2}x \right) \ln x - \right. \\
&\quad \left. - \frac{1}{2}(1+x) \ln^2 x - 5(1-x) \right) + C_f T_f \left(-\frac{2}{3} \ln x - \frac{10}{9} \frac{1+x^2}{1-x} - \frac{4}{3}(1-x) \right) + \\
&\quad + \delta(1-x) \left(C_f^2 \left(\frac{3}{8} - \frac{\pi}{2} + 6\zeta_3 \right) - C_f T_f \left(\frac{1}{6} + \frac{2\pi^2}{9} \right) \right), \tag{A.2}
\end{aligned}$$

$$\begin{aligned}
P_{ee}^S(x) &= C_f T_f \left(\frac{20}{9x} - 2 + 6x - \frac{56}{9}x^2 + (1 + 5x + \frac{8}{3}x^2) \ln x - (1+x) \ln^2 x \right), \\
[P_{ee}^{(1)}(x)]_S &= P_{ee}^{NS}(x) + P_{ee}^S(x), \\
[P_{ee}^{(1)}(x)]_T &= C_f^2 \left(-5 + 5x - \left(\frac{3}{2} + \frac{7}{2}x \right) \ln x + \frac{1}{2}(1+x) \ln^2 x + \right. \\
&\quad \left. \left(\frac{3}{2} \ln x - 2 \ln^2 x + 2 \ln x \ln(1-x) \right) \frac{1+x^2}{1-x} \right) + C_f T_f \left(-\frac{52}{3} + \frac{28}{3}x + \frac{112}{9}x^2 - \frac{40}{9x} + \right. \\
&\quad \left. + \left(10 - 18x - \frac{16}{3}x^2 \right) \ln x + 2(1+x) \ln^2 x + \left(-\frac{2}{3} \ln x - \frac{10}{9} \right) \frac{1+x^2}{1-x} \right), \tag{A.3}
\end{aligned}$$

$$\begin{aligned}
[P_{\gamma e}^{(1)}(x)]_T &= C_f^2 \left[-\frac{1}{2} + \frac{9}{2}x \left(-8 + \frac{1}{2}x \right) \ln x + 2x \ln(1-x) + \left(1 - \frac{1}{2} \right) \ln^2 x + \right. \\
&\quad \left. + \left(\ln^2(1-x) + 4 \ln x \ln(1-x) - 8S_1(x) - \frac{4}{3}\pi^2 \right) P_{\gamma e^{(0)}} \right], \tag{A.4}
\end{aligned}$$

$$\begin{aligned}
P_{e\bar{e}}^{(1)}(x) &= C_f^2 \left(2 \frac{1+x^2}{1+x} S_2(x) + 2(1+x) \ln x + 4(1-x) \right), \\
S_2(x) &= \int_{\frac{x}{1+x}}^{\frac{1}{1+x}} \frac{dz}{z} \ln \frac{1-z}{z} = -3\zeta_2 - \frac{\ln^2 x}{2} + 2 \ln x \ln(1+x) + 2\text{Li}_2(1+x); \\
S_1(x) &= \int_0^{1-x} \frac{dz}{z} \ln(1-z), \tag{A.5}
\end{aligned}$$

where $P_{ee}^{NS}(x)$ and $P_{ee}^S(x)$ are non-singlet and singlet contributions. Note that in QED the QCD constants are reduced to $C_f = 1$ and $T_f = 1$.

Appendix B

$$\begin{aligned}
D_{ee}^{(\text{II})}(x, \mu) &= \delta(1-x) + \frac{\alpha}{2\pi} d_{ee}^{(1)}(x) + \frac{\alpha}{2\pi} L P_{ee}^{(0)} + \left(\frac{\alpha}{2\pi} \right)^2 L \left(d_{\gamma e}^{(1)}(x) \otimes P_{e\gamma}^{(0)} \right. \\
&\quad \left. + P_{ee}^{(1)} - \frac{10}{9} P_{ee}^{(0)} + P_{ee}^{(0)} \otimes d_{ee}^{(1)}(x) \right) + \left(\frac{\alpha}{2\pi} \right)^2 L^2 \left(\frac{1}{2} P_{\gamma e}^{(0)} \otimes P_{e\gamma}^{(0)} + \frac{1}{3} P_{ee}^{(0)} \right. \\
&\quad \left. + \frac{1}{2} P_{ee}^{(0)} \otimes P_{ee}^{(0)} \right), \tag{B.1}
\end{aligned}$$

$$D_{\gamma e}^{(\text{II})}(x, \mu) = \frac{\alpha}{2\pi} d_{\gamma e}^{(1)}(x) + \frac{\alpha}{2\pi} L P_{\gamma e}^{(0)} + \left(\frac{\alpha}{2\pi} \right)^2 L \left(P_{\gamma e}^{(1)} + d_{\gamma e}^{(0)} \otimes P_{\gamma\gamma}^{(0)} \right)$$

$$\begin{aligned}
& -\frac{10}{9}P_{\gamma e}^{(0)} + d_{ee}^{(1)}(x) \otimes P_{\gamma e}^{(0)} + \left(\frac{\alpha}{2\pi}\right)^2 L^2 \left(\frac{1}{3}P_{\gamma e}^{(0)} + \frac{1}{2}P_{\gamma e}^{(0)} \otimes P_{\gamma\gamma}^{(0)} \right. \\
& \left. + \frac{1}{2}P_{ee}^{(0)} \otimes P_{\gamma e}^{(0)} \right), \tag{B.2}
\end{aligned}$$

$$\begin{aligned}
D_{ee}^{(\text{II})}(x, \mu) &= \left(\frac{\alpha}{2\pi}\right)^2 L \left(P_{e\bar{e}}^{(1)} + d_{\gamma e}^{(1)} \otimes P_{e\gamma}^{(0)} \right) \\
&+ \left(\frac{\alpha}{2\pi}\right)^2 L^2 \left(\frac{1}{2}P_{e\gamma}^{(0)} \otimes P_{\gamma\bar{e}}^{(0)} \right). \tag{B.3}
\end{aligned}$$

$$\begin{aligned}
D_{ee}^{(\text{III})}(x, \mu) &= 1 + \frac{\alpha}{2\pi}d_{ee}^{(1)} + \frac{\alpha}{2\pi}LP_{ee}^{(0)} + \left(\frac{\alpha}{2\pi}\right)^2 L \left(d_{\gamma e}^{(1)} \otimes P_{e\gamma}^{(0)} + P_{ee}^{(1)} \right. \\
&- \left. \frac{10}{9}P_{ee}^{(0)} + P_{ee}^{(0)} \otimes d_{ee}^{(1)} \right) + \left(\frac{\alpha}{2\pi}\right)^2 L^2 \left(\frac{1}{2}P_{\gamma e}^{(0)} \otimes P_{e\gamma}^{(0)} + \frac{1}{3}P_{ee}^{(0)} \right) \\
&+ \frac{1}{2}P_{ee}^{(0)} \otimes P_{ee}^{(0)} + \left(\frac{\alpha}{2\pi}\right)^3 L^2 \left(\frac{1}{2}P_{\gamma e}^{(1)} \otimes P_{e\gamma}^{(0)} + \frac{1}{2}P_{\bar{e}e}^{(0)} \otimes P_{e\bar{e}}^{(1)} + \frac{1}{3}d_{\gamma e}^{(1)} \otimes P_{e\gamma}^{(0)} \right. \\
&+ \frac{1}{2}d_{\gamma e}^{(1)} \otimes P_{\gamma\gamma}^{(0)} \otimes P_{e\gamma}^{(0)} + \frac{1}{2}P_{\gamma e}^{(0)} \otimes P_{e\gamma}^{(1)} - \frac{10}{9}P_{\gamma e}^{(0)} \otimes P_{e\gamma}^{(0)} + \frac{2}{3}P_{ee}^{(1)} \\
&+ \frac{1}{2}d_{ee}^{(1)} \otimes P_{\gamma e}^{(0)} \otimes P_{e\gamma}^{(0)} - \frac{13}{54}P_{ee}^{(0)} + \frac{1}{2}P_{ee}^{(0)} \otimes d_{\gamma e}^{(1)} \otimes P_{e\gamma}^{(0)} + P_{ee}^{(0)} \otimes P_{ee}^{(1)} \\
&+ \frac{1}{3}P_{ee}^{(0)} \otimes d_{ee}^{(1)}s - \frac{10}{9}P_{ee}^{(0)} \otimes P_{ee}^{(0)} + \frac{1}{2}P_{ee}^{(0)} \otimes P_{ee}^{(0)} \otimes d_{ee}^{(1)} \left. \right) + \left(\frac{\alpha}{2\pi}\right)^3 L^3 \left(\frac{4}{27}P_{ee}^{(0)} \right. \\
&+ \frac{1}{6}P_{\gamma\bar{e}}^{(0)} \otimes P_{\bar{e}e}^{(0)} \otimes P_{e\gamma}^{(0)} + \frac{1}{3}P_{\gamma e}^{(0)} \otimes P_{e\gamma}^{(0)} + \frac{1}{6}P_{\gamma e}^{(0)} \otimes P_{\gamma\gamma}^{(0)} \otimes P_{e\gamma}^{(0)} \\
&\left. + \frac{1}{3}P_{ee}^{(0)} \otimes P_{\gamma e}^{(0)} \otimes P_{e\gamma}^{(0)} + \frac{1}{3}P_{ee}^{(0)} \otimes P_{ee}^{(0)} + \frac{1}{6}P_{ee}^{(0)} \otimes P_{ee}^{(0)} \otimes P_{ee}^{(0)} \right), \tag{B.4}
\end{aligned}$$

$$\begin{aligned}
D_{ge}^{(\text{III})}(x, \mu) &= \frac{\alpha}{2\pi}d_{\gamma e}^{(1)} + \frac{\alpha}{2\pi}LP_{\gamma e}^{(0)} + \left(\frac{\alpha}{2\pi}\right)^2 L \left(P_{\gamma e}^{(1)} + d_{\gamma e}^{(1)} \otimes P_{\gamma\gamma}^{(0)} \right. \\
&+ \left. d_{ee}^{(1)} \otimes P_{\gamma e}^{(0)} \right) + \left(\frac{\alpha}{2\pi}\right)^2 L^2 \left(\frac{1}{2}P_{\gamma\bar{e}}^{(0)} \otimes P_{\bar{e}e}^{(0)} + \frac{1}{3}P_{\gamma e}^{(0)} + \frac{1}{2}P_{\gamma e}^{(0)} \otimes P_{\gamma\gamma}^{(0)} + \frac{1}{2}P_{ee}^{(0)} \otimes P_{\gamma e}^{(0)} \right) \\
&+ \left(\frac{\alpha}{2\pi}\right)^3 L^2 \left(\frac{2}{3}P_{\gamma e}^{(1)} + \frac{1}{2}P_{\gamma e}^{(1)} \otimes P_{\gamma\gamma}^{(0)} + \frac{1}{2}P_{\gamma\bar{e}}^{(0)} \otimes P_{\bar{e}e}^{(1)} - \frac{10}{9}P_{\gamma\bar{e}}^{(0)} \otimes P_{\bar{e}e}^{(0)} \right. \\
&+ \frac{1}{2}P_{\gamma\bar{e}}^{(1)} \otimes P_{\bar{e}e}^{(0)} + \frac{1}{3}d_{\gamma e}^{(1)} \otimes P_{\gamma\gamma}^{(0)} + \frac{1}{2}d_{\gamma e}^{(1)} \otimes (P_{\gamma\gamma}^{(0)})^2 + \frac{1}{2}d_{\gamma e}^{(1)} \otimes P_{\gamma\bar{e}}^{(0)} \otimes P_{\bar{e}\gamma}^{(0)} - \frac{13}{54}P_{\gamma e}^{(0)} \\
&+ \frac{1}{2}P_{\gamma e}^{(0)} \otimes P_{\gamma\gamma}^{(1)} - \frac{10}{9}P_{\gamma e}^{(0)} \otimes P_{\gamma\gamma}^{(0)} + \frac{1}{2}P_{\gamma e}^{(0)} \otimes d_{\gamma e}^{(1)} \otimes P_{e\gamma}^{(0)} + \frac{1}{2}P_{ee}^{(1)} \otimes P_{\gamma e}^{(0)} \\
&+ \frac{1}{2}d_{ee}^{(1)} \otimes P_{\gamma\bar{e}}^{(0)} \otimes P_{\bar{e}e}^{(0)} + \frac{1}{3}d_{ee}^{(1)} \otimes P_{\gamma e}^{(0)} + \frac{1}{2}d_{ee}^{(1)} \otimes P_{\gamma e}^{(0)} \otimes P_{\gamma\gamma}^{(0)} + \frac{1}{2}P_{ee}^{(0)} \otimes P_{\gamma e}^{(1)} \\
&- \left. \frac{10}{9} \otimes P_{ee}^{(0)} \otimes P_{\gamma e}^{(0)} + \frac{1}{2}P_{ee}^{(0)} \otimes d_{ee}^{(1)} \otimes P_{\gamma e}^{(0)} \right) + \left(\frac{\alpha}{2\pi}\right)^3 L^3 \left(\frac{1}{3}P_{\gamma\bar{e}}^{(0)} \otimes P_{\bar{e}e}^{(0)} \right. \\
&+ \frac{1}{6}P_{\gamma\bar{e}}^{(0)} \otimes P_{\bar{e}e}^{(0)} \otimes P_{\gamma\gamma}^{(0)} + \frac{4}{27}P_{\gamma e}^{(0)} + \frac{1}{3}P_{\gamma e}^{(0)} \otimes P_{\gamma\gamma}^{(0)} + \frac{1}{6}P_{\gamma e}^{(0)} \otimes P_{\gamma\gamma}^{(0)} \otimes P_{\gamma\gamma}^{(0)} \\
&+ \frac{1}{6}P_{\gamma e}^{(0)} \otimes P_{\gamma\bar{e}}^{(0)} \otimes P_{\bar{e}\gamma}^{(0)} + \frac{1}{6}P_{\gamma e}^{(0)} \otimes P_{\gamma e}^{(0)} \otimes P_{e\gamma}^{(0)} + \frac{1}{6}P_{ee}^{(0)} \otimes P_{\gamma\bar{e}}^{(0)} \otimes P_{\bar{e}e}^{(0)} \\
&\left. + \frac{1}{3}P_{ee}^{(0)} \otimes P_{\gamma e}^{(0)} + \frac{1}{6}P_{ee}^{(0)} \otimes P_{\gamma e}^{(0)} \otimes P_{\gamma\gamma}^{(0)} + \frac{1}{6}P_{ee}^{(0)} \otimes P_{ee}^{(0)} \otimes P_{\gamma e}^{(0)} \right), \tag{B.5}
\end{aligned}$$

$$\begin{aligned}
D_{e\bar{e}}^{(\text{III})}(x, \mu) &= \left(\frac{\alpha}{2\pi}\right)^2 L \left(P_{e\bar{e}}^{(1)} + d_{\gamma e}^{(1)} \otimes P_{e\gamma}^{(0)} \right) \\
&+ \left(\frac{\alpha}{2\pi}\right)^2 L^2 \left(\frac{1}{2} P_{e\gamma}^{(0)} \otimes P_{\gamma\bar{e}}^{(0)} \right) + \left(\frac{\alpha}{2\pi}\right)^3 L^2 \left(\frac{2}{3} P_{e\bar{e}}^{(1)} + \frac{1}{2} P_{\gamma\bar{e}}^{(0)} \otimes P_{e\gamma}^{(1)} \right. \\
&- \frac{10}{9} P_{\gamma\bar{e}}^{(0)} \otimes P_{e\gamma}^{(0)} + \frac{1}{2} P_{\gamma\bar{e}}^{(1)} \otimes P_{e\gamma}^{(0)} + \frac{1}{3} d_{\gamma e}^{(1)} \otimes P_{e\gamma}^{(0)} \\
&+ \frac{1}{2} d_{\gamma e}^{(1)} \otimes P_{\gamma\gamma}^{(0)} \otimes P_{e\gamma}^{(0)} + \frac{1}{2} d_{ee}^{(1)} \otimes P_{\gamma\bar{e}}^{(0)} \otimes P_{e\gamma}^{(0)} + \frac{1}{2} P_{e\bar{e}}^{(0)} \otimes P_{e\bar{e}}^{(1)} \\
&+ \left. \frac{1}{2} P_{ee}^{(0)} \otimes P_{e\bar{e}}^{(1)} + \frac{1}{2} P_{ee}^{(0)} \otimes d_{\gamma e}^{(1)} \otimes P_{e\gamma}^{(0)} \right) + \left(\frac{\alpha}{2\pi}\right)^3 L^3 \left(\frac{1}{3} P_{e\gamma}^{(0)} \otimes P_{\gamma\bar{e}}^{(0)} \right. \\
&+ \frac{1}{6} P_{e\gamma}^{(0)} \otimes P_{\gamma\bar{e}}^{(0)} \otimes P_{\gamma\gamma}^{(0)} \\
&+ \left. \frac{1}{6} P_{e\bar{e}}^{(0)} \otimes P_{\gamma\bar{e}}^{(0)} \otimes P_{e\gamma}^{(0)} + \frac{1}{6} P_{ee}^{(0)} \otimes P_{\gamma\bar{e}}^{(0)} \otimes P_{e\gamma}^{(0)} \right). \tag{B.6}
\end{aligned}$$

Because of the charge conjugation parity, these functions obey the equalities (7,8,9,10).

Appendix C

$$[D_{ee}^{(\text{I})}(z)]_S = \frac{\alpha}{2\pi} (-1 - 2 \ln(1-z)) \frac{1+z^2}{1-z} + \frac{\alpha}{2\pi} L \frac{1+z^2}{1-z} + 1 \tag{C.1}$$

$$[D_{e\bar{e}}^{(\text{I})}(z)]_S = 0, \tag{C.2}$$

$$[D_{\gamma e}^{(\text{I})}(z)]_S = \frac{\alpha}{2\pi} (-2 - 4 \ln z) \frac{(1-z)^2 + 1}{z} + \frac{\alpha}{2\pi} L \frac{(1-z)^2 + 1}{z}; \tag{C.3}$$

$$\begin{aligned}
[D_{ee}^{(\text{II})}(z)]_S &= \frac{\alpha}{2\pi} (-1 - 2 \ln(1-z)) \frac{1+z^2}{1-z} + \frac{\alpha}{2\pi} L \frac{1+z^2}{1-z} \\
&+ \left(\frac{\alpha}{2\pi}\right)^2 L \left[-\frac{137}{18} - \frac{2}{z} - \frac{31}{9(1-z)} + \frac{271}{18} z - 2z^2 + 4\zeta_2 \frac{1+z^2}{1-z} \right. \\
&- (2+2z)\text{Li}_2(1-z) + \ln z \left(-\frac{8}{3z} - \frac{1}{3(1-z)} - \frac{22}{3} - \frac{4}{3}z + \frac{8}{3}z^2 \right) \\
&+ \ln(1-z) \left(11 - \frac{14}{1-z} + 3z \right) - 6 \ln^2(1-z) \frac{1+z^2}{1-z} \\
&+ \left. \ln z \ln(1-z) \left(-4 + \frac{4}{1-z} - 4z \right) + \ln^2 z \left(-\frac{7}{2} - \frac{7}{2}z \right) \right] \\
&+ \left(\frac{\alpha}{2\pi}\right)^2 L^2 \left[-\frac{7}{3} + \frac{2}{3z} + \frac{11}{3(1-z)} - \frac{4}{3}z - \frac{2}{3}z^2 + 2 \ln(1-z) \frac{1+z^2}{1-z} \right. \\
&+ \left. \ln z \left(\frac{5}{2} - \frac{2}{1-z} + \frac{5}{2}z \right) \right] + 1; \tag{C.4}
\end{aligned}$$

$$\begin{aligned}
[D_{\gamma e}^{(\text{II})}(z)]_S &= \left(\frac{\alpha}{2\pi}\right) (-1 - 2 \ln z) \frac{(1-z)^2 + 1}{z} \\
&+ \left(\frac{\alpha}{2\pi}\right) L \frac{(1-z)^2 + 1}{z} + \left(\frac{\alpha}{2\pi}\right)^2 L \left[\frac{29}{6} - \frac{22}{3z} - \frac{2}{1-z} - \frac{17}{2}z \right.
\end{aligned}$$

$$\begin{aligned}
& +z^2 + \ln(1-z) \left(\frac{38}{3} - \frac{38}{3z} - \frac{4}{1-z} - \frac{25}{3}z + 2z^2 \right) - \ln^2(1-z) \\
& \times \frac{(1-z)^2 + 1}{z} + \ln z \left(-\frac{2}{3} + \frac{8}{3z} + \frac{29}{6}z \right) + \ln^2 z \left(-1 + \frac{z}{2} \right) \\
& + \left(\frac{\alpha}{2\pi} \right)^2 L^2 \left[1 - \frac{1}{4}z + \ln(1-z) \frac{(1-z)^2 + 1}{z} + \ln z \left(1 - \frac{z}{2} \right) \right]; \tag{C.5}
\end{aligned}$$

$$\begin{aligned}
[D_{e\bar{e}}^{(\text{III})}(z)]_S &= \left(\frac{\alpha}{2\pi} \right)^2 L \left[1 - \frac{2}{z} + 3z - 2z^2 - 6\zeta_2 \frac{1+z^2}{1+z} \right. \\
& + 4\text{Li}_2(1+z) \frac{1+z^2}{1+z} + \ln z \left(-3 - \frac{8}{3z} + 3z + \frac{8}{3}z^2 \right) \\
& \left. + \ln^2 z \left(-4 + \frac{2}{1+z} - 2z \right) \right] + \left(\frac{\alpha}{2\pi} \right)^2 L^2 \left[\frac{1}{2} + \frac{2}{3z} - \frac{1}{2}z \right. \\
& \left. - \frac{2}{3}z^2 + (1+z) \ln z \right], \tag{C.6}
\end{aligned}$$

The third iteration:

$$\begin{aligned}
[D_{e\bar{e}}^{(\text{III})}(z)]_S &= \frac{\alpha}{2\pi} (-1 - 2\ln(1-z)) \frac{1+z^2}{1-z} + \frac{\alpha}{2\pi} L \frac{1+z^2}{1-z} \\
& + \left(\frac{\alpha}{2\pi} \right)^2 L \left[-\frac{137}{18} - \frac{2}{z} - \frac{31}{9(1-z)} + \frac{271}{18}z - 2z^2 + 4\zeta_2 \frac{1+z^2}{1-z} \right. \\
& + \text{Li}_2(1-z)(-2-2z) + \ln(1-z) \left(11 - \frac{14}{1-z} + 3z \right) \\
& - 6\ln^2(1-z) \frac{1+z^2}{1-z} + \ln z \left(-\frac{22}{3} - \frac{8}{3z} - \frac{1}{3(1-z)} - \frac{4z}{3} \right. \\
& \left. + \frac{8}{3}z^2 \right) + 4\ln z \ln(1-z) \frac{z^2}{1-z} - \frac{7}{2}(1+z)\ln^2 z \left. \right] \\
& + \left(\frac{\alpha}{2\pi} \right)^2 L^2 \left[-\frac{7}{3} + \frac{2}{3z} + \frac{11}{3(1-z)} - \frac{4}{3}z - \frac{2}{3}z^2 \right. \\
& + 2\ln(1-z) \frac{1+z^2}{1-z} + \ln z \left(-\frac{2}{1-z} + \frac{5}{2} + \frac{5}{2}z \right) \left. \right] + \left(\frac{\alpha}{2\pi} \right)^3 L^2 \left[\frac{679}{27} + \frac{59}{27z} \right. \\
& - \frac{403}{54(1-z)} - \frac{847}{54}z - \frac{113}{27}z^2 + \zeta_2 \left(\frac{18}{1-z} - 19 + z + \frac{8}{3}z^2 - \frac{8}{3z} \right) - 10\zeta_3 \frac{1+z^2}{1-z} \\
& - (7+7z)\text{S}_{1,2}(1-z) + \text{Li}_2(1-z) \left(-12 - 10z + \frac{4}{3}z^2 + \ln(1-z)(-12-12z) \right. \\
& \left. + \ln z(-9-9z) \right) + 12(1+z)\text{Li}_3(1-z) + \ln(1-z) \left(-\frac{49}{36} - \frac{2}{3z} - \frac{557}{18(1-z)} \right. \\
& \left. + \frac{1235}{36}z - \frac{4}{3}z^2 + 20\zeta_2 \frac{1+z^2}{1-z} \right) + \ln^2(1-z) \left(\frac{53}{2} - \frac{2}{3z} - \frac{34}{1-z} + \frac{15}{2}z + \frac{2}{3}z^2 \right) \\
& - 8\ln^3(1-z) \frac{1+z^2}{1-z} + \ln z \left(\frac{65}{4} + \frac{17}{6(1-z)} - \frac{61}{4}z + \frac{28}{9}z^2 - 4\zeta_2 \frac{1+z^2}{1-z} \right) \\
& \left. + \ln z \ln(1-z) \left(-\frac{68}{3} - \frac{8}{3z} + \frac{28}{3(1-z)} - \frac{26}{3}z + 4z^2 \right) \right]
\end{aligned}$$

$$\begin{aligned}
& +12 \ln z \ln^2(1-z) \frac{z^2}{1-z} + \ln^2 z \left(-\frac{1}{2} + \frac{7}{3(1-z)} - \frac{7}{2}z - 4z^2 \right) \\
& + \ln^2 z \ln(1-z) \left(-\frac{11}{2} - \frac{11}{2}z \right) + \ln^3 z \left(\frac{4}{3} + \frac{4}{3}z \right) \Big] \\
& + \left(\frac{\alpha}{2\pi} \right)^3 L^3 \left[-\frac{971}{216} + \frac{8}{27z} + \frac{491}{108(1-z)} - 2\zeta_2 \frac{1+z^2}{1-z} - \frac{11}{216}z - \frac{8}{27}z^2 \right. \\
& + \text{Li}_2(1-z) \left(\frac{7}{3} + \frac{7}{3}z \right) + \ln(1-z) \left(-\frac{17}{3} + \frac{8}{9z} + \frac{26}{3(1-z)} - 3z - \frac{8}{9}z^2 \right) \\
& + 2 \ln^2(1-z) \frac{1+z^2}{1-z} + \ln z \left(\frac{151}{36} - \frac{13}{3(1-z)} + \frac{103}{36}z + \frac{8}{9}z^2 \right) \\
& + \ln z \ln(1-z) \left(\frac{13}{3} - \frac{4}{1-z} + \frac{13}{3}z \right) + \ln^2 z \left(-\frac{11}{12} + \frac{2}{3(1-z)} \right. \\
& \left. \left. - \frac{11}{12}z \right) \right] + 1, \tag{C.7}
\end{aligned}$$

$$\begin{aligned}
[D_{\gamma_e}^{(\text{III})}(z)]_S &= \frac{\alpha}{2\pi^2} (-1 - 2 \ln z) \frac{(1-z)^2 + 1}{z} + \frac{\alpha}{2\pi^2} L \frac{(1-z)^2 + 1}{z} \\
& + \left(\frac{\alpha}{2\pi^2} \right)^2 L \left[\frac{29}{6} - \frac{22}{3z} - \frac{2}{1-z} - \frac{17}{2}z + z^2 + \ln(1-z) \left(\frac{38}{3} - \frac{38}{3z} \right. \right. \\
& \left. \left. - \frac{4}{1-z} - \frac{25}{3}z + 2z^2 \right) - \ln^2(1-z) \frac{(1-z)^2 + 1}{z} + \ln z \left(-\frac{2}{3} + \frac{8}{3z} + \frac{29}{6}z \right) \right. \\
& \left. + \ln^2 z \left(-1 + \frac{1}{2}z \right) \right] + \left(\frac{\alpha}{2\pi^2} \right)^2 L^2 \left[1 - \frac{z}{4} + \ln(1-z) \frac{(1-z)^2 + 1}{z} + \ln z \left(1 - \frac{z}{2} \right) \right] \\
& + \left(\frac{\alpha}{2\pi^2} \right)^3 L^2 \left[-\frac{6877}{216} + \frac{5}{54z} - \frac{92}{9(1-z)} + \frac{15595}{432}z + \frac{529}{108}z^2 + \left(-\frac{14}{3} - \frac{46}{36}z - \frac{61}{6z} \right) \zeta_2 \right. \\
& + \left(10 + 3z + \frac{4}{z} \right) \zeta_3 + \left(-8 - \frac{4}{z} \right) \text{S}_{1,2}(1-z) + \text{Li}_2(1+z) \left(10 + \frac{6}{z} + 4z \right) \\
& + \text{Li}_2(1-z) \left(-3 + \frac{10}{3z} - z + \ln(1-z) \left(-8 - \frac{4}{z} + 4z \right) + \ln z \left(-6 + \frac{4}{z} - z \right) \right) \\
& + \text{Li}_3(1-z) \left(8 + \frac{4}{z} - 4z \right) + \text{Li}_3(z^2) \left(-1 + \frac{1}{z} - \frac{z}{2} \right) + \text{Li}_3(1+z) \left(-8 - \frac{8}{z} - 4z \right) \\
& + \ln(1+z) \zeta_2 \left(12 + \frac{12}{z} + 6z \right) + \ln(1-z) \left(\frac{445}{18} - \frac{331}{18z} - \frac{23}{3(1-z)} - \frac{94}{9}z + \frac{23}{6}z^2 \right. \\
& \left. + 4 \frac{(z-1)^2 + 1}{z} \zeta_2 \right) + \ln^2(1-z) \left(-\frac{23}{3} + \frac{5}{3z} - \frac{1}{1-z} + \frac{7}{3}z + \frac{1}{2}z^2 \right) \\
& - 2 \ln^3(1-z) \frac{(1-z)^2 + 1}{z} + \ln z \left(-\frac{275}{6} + \frac{20}{3z} + \frac{11}{2(1-z)} - \frac{23}{12}z \right. \\
& \left. - \frac{127}{36}z^2 + \left(6 - \frac{8}{z} - z \right) \zeta_2 + \frac{4}{z} \text{Li}_2(1+z) \right) + \ln z \ln(1-z) \left(\frac{7}{3} - \frac{19}{6}z \right) \\
& + \ln z \ln^2(1-z) (-6 + 3z) + \ln^2 z \left(-\frac{35}{12} + \frac{8}{3z} - \frac{1}{2(1-z)} + \frac{1}{3}z - \frac{1}{4}z^2 \right)
\end{aligned}$$

$$\begin{aligned}
& + \ln^2 z (\ln(1+z) + \ln(1-z)) \left(-2 + \frac{2}{z} - z \right) + \ln^3 z \left(-\frac{5}{2} + \frac{19}{12}z \right) \Big] \\
& + \left(\frac{\alpha}{2\pi^2} \right)^3 L^3 \left[\left(\frac{193}{36} - \frac{62}{27z} - \frac{191}{72}z + \frac{8}{27}z^2 \right) - \frac{2}{3}\zeta_2 \frac{(z-1)^2 + 1}{z} + \frac{4}{3z} \text{Li}_2(1-z) \right. \\
& + \ln(1-z) \left(\frac{4}{9} + \frac{8}{9z} + \frac{1}{9}z \right) + \frac{2}{3} \ln^2(1-z) \frac{(z-1)^2 + 1}{z} + \ln z \left(-\frac{7}{18} - \frac{8}{9z} + \frac{10}{9}z \right) \\
& \left. + \ln z \ln(1-z) \left(\frac{4}{3} - \frac{2}{3}z \right) + \ln^2 z \left(\frac{1}{2} - \frac{1}{4}z \right) \right], \tag{C.8}
\end{aligned}$$

$$\begin{aligned}
[D_{e\bar{e}}^{(\text{III})}(z)]_S &= s \left(\frac{\alpha}{2\pi^2} \right)^2 L \left[1 - \frac{2}{z} + 3z - 2z^2 - 6\zeta_2 \frac{1+z^2}{1+z} + 4\text{Li}_2(1+z) \frac{1+z^2}{1+z} \right. \\
& + \ln z \left(-3 - \frac{8}{3z} + 3z + \frac{8}{3}z^2 \right) + \ln^2 z \left(-4 + \frac{2}{1+z} - 2z \right) \Big] \\
& + \left(\frac{\alpha}{2\pi^2} \right)^2 L^2 \left[\frac{1}{2} + \frac{2}{3z} - \frac{1}{2}z - \frac{2}{3}z^2 + (1+z) \ln z \right] + \left(\frac{\alpha}{2\pi^2} \right)^3 L^2 \left[\frac{97}{36} + \frac{59}{27z} \right. \\
& - \frac{25}{36}z - \frac{113}{27}z^2 + \text{Li}_2(1+z) \left(\frac{52}{3(1+z)} - \frac{50}{3} + \frac{2}{3}z \right) + \zeta_3 \left(8 - \frac{12}{1+z} - 8z \right) \\
& + \zeta_2 \left(23 - \frac{8}{3z} - \frac{26}{1+z} + z + \frac{8}{3}z^2 \right) + (-10 + 6z) \text{S}_{1,2}(1-z) \\
& + \text{Li}_2(1-z) \left\{ -\frac{4}{3} + \frac{2}{3}z + \frac{4}{3}z^2 - 8 \ln(1+z) \frac{1+z^2}{1+z} + \ln(1-z) \left(-10 + \frac{16}{1+z} \right. \right. \\
& \left. \left. + 6z \right) + (-14 + 2z) \ln z \right\} - 8 \left(\text{Li}_2 \left(\frac{1+z}{2} \right) - \text{Li}_2 \left(\frac{1-z}{1+z} \right) \right) (\ln(1+z) \\
& - \ln(1-z)) \frac{1+z^2}{1+z} + \text{Li}_3(1-z) \left(-6 + \frac{16}{1+z} + 10z \right) + 16\text{Li}_3(1+z) \frac{1+z^2}{1+z} \\
& - 4 \text{Li}_3(1-z^2) \frac{1+z^2}{1+z} + \text{Li}_3(z^2) \left(-1 - \frac{2}{1+z} + z \right) - 8\text{Li}_3 \left(\frac{z}{1+z} \right) \frac{1+z^2}{1+z} \\
& + \ln(1+z) \left(-32\zeta_2 - 4 \ln^2(2) + 8\text{Li}_2(1+z) \right) \frac{1+z^2}{1+z} + \frac{16}{3} \ln(1+z)^3 \frac{1+z^2}{1+z} \\
& + \ln(1-z) \left(\frac{1}{2} - \frac{2}{3z} + \frac{3}{2}z - \frac{4}{3}z^2 + (-8\zeta_2 + 4 \ln^2(2)) \frac{1+z^2}{1+z} \right) \\
& + 8 \ln(1-z) \ln(1+z) \ln(2) \frac{1+z^2}{1+z} - 12 \ln(1-z) \ln^2(1+z) \frac{1+z^2}{1+z} \\
& + \ln^2(1-z) \left(-\frac{1}{2} - \frac{2}{3z} + \frac{1}{2}z + \frac{2}{3}z^2 - 8 \ln(2) \frac{1+z^2}{1+z} \right) \\
& + 8 \ln^2(1-z) \ln(1+z) \frac{1+z^2}{1+z} + \ln z \left(\frac{353}{36} + \frac{71}{36}z + \frac{28}{9}z^2 + \left(-6 + \frac{24}{1+z} - 2z \right) \zeta_2 \right. \\
& \left. - 8\text{Li}_2(1+z) \right) - 4 \ln z \ln^2(1+z) \frac{1+z^2}{1+z} + \ln z \ln(1-z) \\
& \times \left(-\frac{10}{3} - \frac{8}{3z} + \frac{8}{3}z + 4z^2 \right) + (-1-z) \ln z \ln^2(1-z) + \ln^2 z \left(-\frac{10}{3} + \frac{13}{3(1+z)} \right)
\end{aligned}$$

$$\begin{aligned}
& -\frac{9}{2}z - 4z^2) + \ln^2 z \ln(1+z) \left(-6 + \frac{4}{1+z} + 6z\right) + \ln^2 z \ln(1-z) (-9-z) \\
& + \ln^3 z \left(\frac{13}{6} - \frac{4}{3(1+z)} + \frac{1}{6}z\right) + \left(\frac{\alpha}{2\pi^2}\right)^3 L^3 \left[-\frac{19}{18} + \frac{8}{27} + \frac{19}{18}z - \frac{8}{27}z^2\right. \\
& + \left(\frac{4}{3}z + \frac{4}{3}\right) \text{Li}_2(1-z) + \ln(1-z) \left(\frac{8}{9z} + \frac{2}{3} - \frac{2}{3}z - \frac{8}{9}z^2\right) \\
& \left. + \ln z \left(\frac{4}{9} + \frac{10}{9}z + \frac{8}{9}z^2\right) + \ln z \ln(1-z) \left(\frac{4}{3} + \frac{4}{3}z\right) + \left(-\frac{1}{3} - \frac{1}{3}z\right) \ln^2 z\right]. \quad (\text{C.9})
\end{aligned}$$

And for timelike functions:

$$[D_{ee}^{(\text{I})}(z)]_T = [D_{ee}^{(\text{I})}(z)]_S, \quad (\text{C.10})$$

$$[D_{e\bar{e}}^{(\text{I})}(z)]_T = 0 = [D_{e\bar{e}}^{(\text{I})}(z)]_S, \quad (\text{C.11})$$

$$[D_{\gamma e}^{(\text{I})}(z)]_T = [D_{\gamma e}^{(\text{I})}(z)]_S, \quad (\text{C.12})$$

$$\begin{aligned}
[D_{ee}^{(\text{II})}(z)]_T &= \frac{\alpha}{2\pi}(-1 - 2\ln(1-z))\frac{1+z^2}{1-z} \\
&+ \frac{\alpha}{2\pi}L\frac{1+z^2}{1-z} + \left(\frac{\alpha}{2\pi}\right)^2 L \left[-\frac{389}{18} - \frac{26}{3z} - \frac{31}{9(1-z)} + \frac{307}{18}z + \frac{50}{3}z^2 + 4\frac{1+z^2}{1-z}\zeta_2\right. \\
&+ \text{Li}_2(1-z)(-2-2z) + \ln(1-z) \left(11 - \frac{14}{1-z} + 3z\right) - 6\ln^2(1-z)\frac{1+z^2}{1-z} \\
&+ \ln z \left(-\frac{10}{3} - \frac{8}{3z} + \frac{17}{3(1-z)} - \frac{76}{3}z - \frac{16}{3}z^2\right) + \ln z \ln(1-z) \left(-8 + \frac{12}{1-z} - 8z\right) \\
&+ \ln^2 z \left(\frac{5}{2} - \frac{4}{1-z} + \frac{5}{2}z\right) + \left(\frac{\alpha}{2\pi}\right)^2 L^2 \left[-\frac{7}{3} + \frac{2}{3z} + \frac{11}{3(1-z)} - \frac{4}{3}z - \frac{2}{3}z^2\right. \\
&\left. + 2\ln(1-z)\frac{1+z^2}{1-z} + \ln z \left(\frac{5}{2} - \frac{2}{1-z} + \frac{5}{2}z\right)\right] + 1, \quad (\text{C.13})
\end{aligned}$$

$$\begin{aligned}
[D_{\gamma e}^{(\text{II})}(z)]_T &= \frac{\alpha}{2\pi}(-1 - 2\ln z)\frac{(z-1)^2+1}{z} + \left(\frac{\alpha}{2\pi}\right)L\frac{(z-1)^2+1}{z} \\
&+ \left(\frac{\alpha}{2\pi}\right)^2 L \left[\frac{43}{18} - \frac{26}{9z} - \frac{2}{1-z} + \frac{55}{18}z + z^2 - 8\zeta_2\frac{(z-1)^2+1}{z} - 8\text{Li}_2(1-z)\frac{(z-1)^2+1}{z}\right. \\
&+ \ln(1-z) \left(4 - \frac{4}{z} - \frac{4}{1-z} + 2z^2\right) + \ln^2(1-z)\frac{(z-1)^2+1}{z} + \ln z \left(-\frac{32}{3} + \frac{8}{3z} + \frac{11}{6}z\right) \\
&+ 4\ln z \ln(1-z)\frac{(z-1)^2+1}{z} + \ln^2 z \left(1 - \frac{1}{2}z\right) \left. \right] \\
&+ \left(\frac{\alpha}{2\pi}\right)^2 L^2 \left[1 - \frac{1}{4}z + \ln z \left(1 - \frac{1}{2}z\right) + \ln(1-z)\frac{(z-1)^2+1}{z}\right], \quad (\text{C.14})
\end{aligned}$$

$$\begin{aligned}
[D_{e\bar{e}}^{(\text{II})}(z)]_T &= \left(\frac{\alpha}{2\pi^2}\right)^2 L \left[1 - \frac{2}{z} + 3z - 2z^2 - 6\zeta_2\frac{1+z^2}{1+z} + 4\frac{1+z^2}{1+z}\text{Li}_2(1+z)\right. \\
&+ \ln z \left(-3 - \frac{8}{3z} + 3z + \frac{8}{3}z^2\right) + \ln^2 z \left(-4 + \frac{2}{1+z} - 2z\right) \left. \right] \\
&+ \left(\frac{\alpha}{2\pi^2}\right)^2 L^2 \left[\frac{1}{2} + \frac{2}{3z} - \frac{1}{2}z - \frac{2}{3}z^2 + (1+z)\ln z\right], \quad (\text{C.15})
\end{aligned}$$

$$\begin{aligned}
[D_{ee}^{(\text{III})}(z)]_T &= \frac{\alpha}{2\pi^2} (-1 - 2\ln(1-z)) \frac{1+z^2}{1-z} \\
&+ \frac{\alpha}{2\pi^2} L \frac{1+z^2}{1-z} + \left(\frac{\alpha}{2\pi^2}\right)^2 L \left[-\frac{389}{18} - \frac{26}{3z} - \frac{31}{9(1-z)} + \frac{307}{18}z + \frac{50}{3}z^2 + 4\zeta_2 \frac{1+z^2}{1-z} \right. \\
&+ \text{Li}_2(1-z)(-2-2z) + \ln(1-z) \left(11 - \frac{14}{1-z} + 3z \right) - 6\ln^2(1-z) \frac{1+z^2}{1-z} \\
&+ \ln z \left(-\frac{10}{3} - \frac{8}{3z} + \frac{17}{3(1-z)} - \frac{76}{3}z - \frac{16}{3}z^2 \right) + \ln z \ln(1-z) \left(-8 + \frac{12}{1-z} - 8z \right) \\
&+ \ln^2 z \left(\frac{5}{2} - \frac{4}{1-z} + \frac{5}{2}z \right) \left. \right] + \left(\frac{\alpha}{2\pi^2}\right)^2 L^2 \left[-\frac{7}{3} + \frac{2}{3z} + \frac{11}{3(1-z)} - \frac{4}{3}z - \frac{2}{3}z^2 \right. \\
&+ 2\ln(1-z) \frac{1+z^2}{1-z} + \ln z \left(\frac{5}{2} - \frac{2}{1-z} + \frac{5}{2}z \right) \left. \right] + \left(\frac{\alpha}{2\pi^2}\right)^3 L^2 \left[\frac{823}{27} - \frac{34}{9z} - \frac{403}{54(1-z)} \right. \\
&- \frac{1045}{54}z + \frac{1}{9}z^2 - 10\zeta_3 \frac{1+z^2}{1-z} + \zeta_2 \left(\frac{8}{3z} + \frac{18}{1-z} - 15 - 3z - \frac{8}{3}z^2 \right) \\
&+ \text{S}_{1,2}(1-z)(-47-47z) + \text{Li}_2(1-z) \left(-\frac{13}{3} - \frac{16}{z} - \frac{94}{3}z + \frac{20}{3}z^2 \right. \\
&- (14\ln(1-z) + 17\ln z)(1+z) + \text{Li}_3(1-z)(14+14z) \\
&+ \ln(1-z) \left(-\frac{943}{36} - \frac{89}{9z} - \frac{557}{18(1-z)} + \frac{1661}{36}z + \frac{188}{9}z^2 \right. \\
&+ 20\zeta_2 \frac{1+z^2}{1-z} \left. \right) + \ln^2(1-z) \left(27 - \frac{2}{z} - \frac{34}{1-z} + 7z + 2z^2 \right) - 8\ln^3(1-z) \frac{1+z^2}{1-z} \\
&+ \ln z \left(\frac{757}{36} + \frac{95}{6(1-z)} - \frac{1115}{36}z - \frac{188}{9}z^2 + \zeta_2 \left(20 - \frac{24}{1-z} + 20z \right) \right) \\
&+ \ln z \ln(1-z) \left(-\frac{71}{3} + \frac{116}{3(1-z)} - \frac{152}{3}z - \frac{28}{3}z^2 \right) \\
&+ \ln z \ln^2(1-z) \left(-21 + \frac{28}{1-z} - 21z \right) + \ln^2 z \left(\frac{91}{6} - \frac{37}{3(1-z)} + \frac{145}{6}z + \frac{32}{3}z^2 \right) \\
&+ \ln^2 z \ln(1-z) \left(\frac{29}{2} - \frac{16}{1-z} + \frac{29}{2}z \right) + \ln^3 z \left(-\frac{10}{3} + \frac{8}{3(1-z)} - \frac{10}{3}z \right) \left. \right] \\
&+ \left(\frac{\alpha}{2\pi^2}\right)^3 L^3 \left[-\frac{971}{216} + \frac{8}{27z} + \frac{491}{108(1-z)} - \frac{11}{216}z - \frac{8}{27}z^2 - 2\zeta_2 \frac{1+z^2}{1-z} \right. \\
&+ \text{Li}_2(1-z) \left(\frac{7}{3} + \frac{7}{3}z \right) + \ln(1-z) \left(-\frac{17}{3} + \frac{8}{9z} + \frac{26}{3(1-z)} - 3z - \frac{8}{9}z^2 \right) \\
&+ 2\ln^2(1-z) \frac{1+z^2}{1-z} + \ln z \left(\frac{151}{36} - \frac{13}{3(1-z)} + \frac{103}{36}z + \frac{8}{9}z^2 \right) \\
&+ \ln z \ln(1-z) \left(\frac{13}{3} - \frac{4}{1-z} + \frac{13}{3}z \right) + \ln^2 z \left(-\frac{11}{12} + \frac{2}{3(1-z)} - \frac{11}{12}z \right) \left. \right] + 1, \quad (\text{C.16})
\end{aligned}$$

$$[D_{\gamma e}^{(\text{III})}(z)]_T = \frac{\alpha}{2\pi^2} (-1 - 2\ln z) \frac{(z-1)^2 + 1}{z} + \frac{\alpha}{2\pi^2} L \frac{(z-1)^2 + 1}{z}$$

$$\begin{aligned}
& + \left(\frac{\alpha}{2\pi^2}\right)^2 L \left(\frac{43}{18} - \frac{26}{9z} - \frac{2}{1-z} + \frac{55}{18}z + z^2 - 8\frac{(z-1)^2+1}{z} \zeta_2 \right. \\
& - 8\text{Li}_2(1-z)\frac{(z-1)^2+1}{z} + \ln(1-z) \left(4 - \frac{4}{z} - \frac{4}{1-z} + 2z^2 \right) + \frac{(z-1)^2+1}{z} \ln^2(1-z) \\
& \left. + \ln z \left(-\frac{32}{3} + \frac{8}{3z} + \frac{11}{6}z \right) + 4\frac{(z-1)^2+1}{z} \ln z \ln(1-z) + \ln^2 z \left(1 - \frac{1}{2}z \right) \right] \\
& + \left(\frac{\alpha}{2\pi^2}\right)^2 L^2 \left[1 - \frac{1}{4}z + \frac{(z-1)^2+1}{z} \ln(1-z) + \ln z \left(1 - \frac{1}{2}z \right) \right] \\
& + \left(\frac{\alpha}{2\pi^2}\right)^3 L^2 \left[-\frac{529}{24} - \frac{346}{54z} + \frac{10}{3(1-z)} + \frac{1267}{48}z + \frac{25}{108}z^2 + \zeta_3 \left(10 + \frac{4}{z} + 3z \right) \right. \\
& + \zeta_2 \left(-\frac{70}{3z} - \frac{8}{1-z} + \frac{10}{3} - \frac{85}{6}z + 4z^2 \right) + \text{S}_{1,2}(1-z) \left(-4 - \frac{4}{z} - 2z \right) \\
& + \text{Li}_2(1+z) \left(\frac{6}{z} + 10 + 4z \right) + \text{Li}_2(1-z) \left(7 - \frac{14}{3z} - \frac{8}{1-z} - 6z + 4z^2 + \ln z \left(-6 + \frac{4}{z} - z \right) \right. \\
& \left. - \frac{12}{z} \ln(1-z) \right) + \frac{12}{z} \text{Li}_3(1-z) + \text{Li}_3(1+z) \left(-8 - \frac{8}{z} - 4z \right) + \text{Li}_3(z^2) \left(-1 + \frac{1}{z} - \frac{1}{2}z \right) \\
& + \ln(1+z)\zeta_2 \left(12 + \frac{12}{z} + 6z \right) + \ln(1-z) \left(\frac{15}{2} - \frac{37}{6z} + \frac{2}{3(1-z)} - \frac{43}{6}z - \frac{1}{3}z^2 \right. \\
& \left. + 4\zeta_2 \frac{(z-1)^2+1}{z} \right) + \ln^2(1-z) \left(-\frac{31}{3} + \frac{13}{3z} + \frac{1}{1-z} + \frac{11}{3}z - \frac{1}{2}z^2 \right) \\
& - 2\ln^3(1-z)\frac{(z-1)^2+1}{z} + \ln z \left(-\frac{55}{2} - \frac{4}{9z} - \frac{15}{2(1-z)} - \frac{115}{12}z - \frac{41}{36}z^2 + \zeta_2 \left(6 - \frac{8}{z} - z \right) \right) \\
& + \ln z \ln(1-z) \left(\frac{13}{3} + \frac{4}{z} + \frac{4}{1-z} - \frac{19}{6}z - 2z^2 \right) + \ln z \ln^2(1-z)(-6+3z) \\
& + \ln^2 z \left(\frac{41}{12} + \frac{1}{2(1-z)} + \frac{25}{6}z + \frac{1}{4}z^2 + \ln(1+z) \left(-2 + \frac{2}{z} - z \right) + \ln(1-z) \left(\frac{2}{z} - 2z \right) \right) \\
& \left. + \ln^3 z \left(-\frac{1}{2} + \frac{7}{12}z \right) \right] + \left(\frac{\alpha}{2\pi^2}\right)^3 L^3 \left[\frac{193}{36} - \frac{62}{27z} - \frac{191}{72}z + \frac{8}{27}z^2 - \frac{2}{3} \frac{(z-1)^2+1}{z} \zeta_2 \right. \\
& + \frac{4}{3z} \text{Li}_2(1-z) + \ln(1-z) \left(\frac{4}{9} + \frac{8}{9z} + \frac{1}{9}z \right) + \frac{2}{3} \ln^2(1-z)\frac{(z-1)^2+1}{z} \\
& \left. + \ln z \left(-\frac{7}{18} - \frac{8}{9z} + \frac{10}{9}z \right) + \frac{2}{3} \ln z \ln(1-z)(2-z) + \frac{1}{4} \ln^2 z (2-z) \right], \tag{C.17}
\end{aligned}$$

$$\begin{aligned}
[D_{e\bar{e}}^{(\text{III})}(z)]_T & = \left(\frac{\alpha}{2\pi^2}\right)^2 L \left[1 - \frac{2}{z} + 3z - 2z^2 - 6\zeta_2 \frac{1+z^2}{1+z} + 4\text{Li}_2(1+z) \frac{1+z^2}{1+z} \right. \\
& \left. + \ln z \left(-3 - \frac{8}{3z} + 3z + \frac{8}{3}z^2 \right) + \ln^2 z \left(-4 + \frac{2}{1+z} - 2z \right) \right] \\
& + \left(\frac{\alpha}{2\pi^2}\right)^2 L^2 \left[\left(\frac{1}{2} + \frac{2}{3z} - \frac{1}{2}z - \frac{2}{3}z^2 \right) + \ln z(1+z) \right] + \left(\frac{\alpha}{2\pi^2}\right)^3 L^2 \left[\left(\frac{109}{36} - \frac{2}{27z} \right. \right. \\
& \left. \left. - \frac{97}{36}z - \frac{7}{27}z^2 + \zeta_3 \left(8 - \frac{12}{1+z} - 8z \right) + \zeta_2 \left(25 + \frac{8}{3z} - \frac{22}{1+z} - z - \frac{8}{3}z^2 \right) \right) \right]
\end{aligned}$$

$$\begin{aligned}
& +S_{1,2}(1-z)(-62-46z) + \text{Li}_2(1+z) \left(-\frac{46}{3} + \frac{44}{3(1+z)} - \frac{2}{3}z \right) \\
& + \text{Li}_2(1-z) \left(-\frac{35}{3} - \frac{16}{z} + \frac{76}{3}z + \frac{68}{3}z^2 - 8\ln(1+z)\frac{1+z^2}{1+z} \right. \\
& + \ln(1-z) \left(-12 + \frac{16}{1+z} + 4z \right) + \ln z(-38-22z) \\
& - 8 \left(\text{Li}_2\left(\frac{1+z}{2}\right) - \text{Li}_2\left(\frac{1-z}{1+z}\right) \right) (\ln(1+z) - \ln(1-z))\frac{1+z^2}{1+z} \\
& + \text{Li}_3(1-z)(-4 + \frac{16}{1+z} + 12z) + 16 \text{Li}_3(1+z)\frac{1+z^2}{1-z} - 4 \text{Li}_3(1-z^2)\frac{1+z^2}{1+z} \\
& + \text{Li}_3(z^2) \left(-1 - \frac{2}{1+z} + z \right) - 8\text{Li}_3\left(\frac{z}{1+z}\right)\frac{1+z^2}{1+z} + \ln(1+z)(8\text{Li}_2(1+z) - 32\zeta_2 \\
& - 4\ln^2(2))\frac{1+z^2}{1+z} + \frac{16}{3}\ln(1+z)^3\frac{1+z^2}{1+z} + \ln(1-z) \left(\frac{11}{3} + \frac{31}{9z} + \frac{28}{3}z - \frac{148}{9}z^2 \right. \\
& + \left. (-8\zeta_2 + 4\ln^2(2))\frac{1+z^2}{1+z} \right) + 8\ln(1-z)\ln(1+z)\ln(2)\frac{1+z^2}{1+z} \\
& - 12\ln(1-z)\ln^2(1+z)\frac{1+z^2}{1+z} - 12\ln(1-z)\ln^2(1+z)\frac{1+z^2}{1+z} \\
& + \ln^2(1-z) \left(-\frac{2}{z} + 2z^2 - 8\frac{1+z^2}{1+z}\ln(2) \right) + 8\ln^2(1-z)\ln(1+z)\frac{1+z^2}{1+z} \\
& + \ln z \left(\frac{59}{12} + \frac{119}{12}z + \frac{188}{9}z^2 + \zeta_2 \left(2 + \frac{24}{1+z} + 6z \right) - \frac{8\text{Li}_2}{1+z} \right) - 4\ln z \ln^2(1+z)\frac{1+z^2}{1+z} \\
& + \ln z \ln(1-z) \left(\frac{1}{3} + \frac{40}{3}z + \frac{20}{3}z^2 \right) - 2(1+z)\ln z \ln^2(1-z) + \ln^2 z \left(1 + \frac{11}{3}z + \frac{19}{2}z \right. \\
& + \left. \frac{8}{3}z^2 + \ln(1+z) \left(-6 + \frac{4}{1+z} + 6z \right) + \ln(1-z)(-7+z) \right) \\
& - \ln^3 z \left(\frac{5}{6} - \frac{4}{3(1+z)} + \frac{7}{6}z \right) \Big] + \left(\frac{\alpha}{2\pi^2} \right)^3 L^3 \left[-\frac{19}{18} + \frac{8}{27z} + \frac{19}{18}z - \frac{8}{27}z^2 \right. \\
& + \left. \frac{4}{3}\text{Li}_2(1-z)(1+z) + \ln(1-z) \left(\frac{2}{3} + \frac{8}{9z} - \frac{2}{3}z - \frac{8}{9}z^2 \right) \right. \\
& + \left. \ln z \left(\frac{4}{9} + \frac{10}{9}z + \frac{8}{9}z^2 \right) + \frac{4}{3}(1+z)\ln z \ln(1-z) - \frac{1}{3}\ln^2 z(1+z) \right]. \tag{C.18}
\end{aligned}$$

References

- [1] A. Abada, et al., FCC-ee: The Lepton Collider: Future Circular Collider Conceptual Design Report Volume 2, Eur. Phys. J. ST 228 (2) (2019) 261–623. doi:10.1140/epjst/e2019-900045-4.
- [2] M. Dong, et al., CEPC Conceptual Design Report: Volume 2 - Physics & Detector (11 2018). arXiv:1811.10545.

- [3] S. Jadach, M. Skrzypek, QED challenges at FCC-ee precision measurements, *Eur. Phys. J. C* 79 (9) (2019) 756. [arXiv:1903.09895](#), [doi:10.1140/epjc/s10052-019-7255-9](#).
- [4] E. A. Kuraev, V. S. Fadin, On Radiative Corrections to $e^+ e^-$ Single Photon Annihilation at High-Energy, *Sov. J. Nucl. Phys.* 41 (1985) 466–472.
- [5] A. De Rujula, R. Petronzio, A. Savoy-Navarro, Radiative Corrections to High-Energy Neutrino Scattering, *Nucl. Phys. B* 154 (1979) 394–426. [doi:10.1016/0550-3213\(79\)90039-7](#).
- [6] V. N. Gribov, L. N. Lipatov, Deep inelastic $e p$ scattering in perturbation theory, *Sov. J. Nucl. Phys.* 15 (1972) 438–450.
- [7] G. Altarelli, G. Parisi, Asymptotic Freedom in Parton Language, *Nucl. Phys. B* 126 (1977) 298–318. [doi:10.1016/0550-3213\(77\)90384-4](#).
- [8] Y. L. Dokshitzer, Calculation of the Structure Functions for Deep Inelastic Scattering and $e^+ e^-$ Annihilation by Perturbation Theory in Quantum Chromodynamics., *Sov. Phys. JETP* 46 (1977) 641–653.
- [9] M. Cacciari, A. Deandrea, G. Montagna, O. Nicosini, QED structure functions: A Systematic approach, *EPL* 17 (1992) 123–128. [doi:10.1209/0295-5075/17/2/007](#).
- [10] D. R. Yennie, S. C. Frautschi, H. Suura, The infrared divergence phenomena and high-energy processes, *Annals Phys.* 13 (1961) 379–452. [doi:10.1016/0003-4916\(61\)90151-8](#).
- [11] M. Przybycien, A Fifth order perturbative solution to the Gribov-Lipatov equation, *Acta Phys. Polon. B* 24 (1993) 1105–1114. [arXiv:hep-th/9511029](#).
- [12] A. B. Arbuzov, Nonsinglet splitting functions in QED, *Phys. Lett. B* 470 (1999) 252–258. [arXiv:hep-ph/9908361](#), [doi:10.1016/S0370-2693\(99\)01290-3](#).
- [13] J. Blumlein, H. Kawamura, Universal higher order QED corrections to polarized lepton scattering, *Nucl. Phys. B* 708 (2005) 467–510. [arXiv:hep-ph/0409289](#), [doi:10.1016/j.nuclphysb.2004.12.001](#).
- [14] F. A. Berends, W. L. van Neerven, G. J. H. Burgers, Higher Order Radiative Corrections at LEP Energies, *Nucl. Phys. B* 297 (1988) 429, [Erratum: *Nucl.Phys.B* 304, 921 (1988)]. [doi:10.1016/0550-3213\(88\)90313-6](#).
- [15] A. Arbuzov, K. Melnikov, $O(\alpha^2 \ln(m(\mu) / m(e)))$ corrections to electron energy spectrum in muon decay, *Phys. Rev. D* 66 (2002) 093003. [arXiv:hep-ph/0205172](#), [doi:10.1103/PhysRevD.66.093003](#).
- [16] J. Blumlein, H. Kawamura, $O(\alpha^2 L)$ radiative corrections to deep inelastic ep scattering, *Phys. Lett. B* 553 (2003) 242–250. [arXiv:hep-ph/0211191](#), [doi:10.1016/S0370-2693\(02\)03194-5](#).

- [17] J. Blumlein, H. Kawamura, Universal higher order singlet QED corrections to unpolarized lepton scattering, *Eur. Phys. J. C* 51 (2007) 317–333. [arXiv:hep-ph/0701019](#), [doi:10.1140/epjc/s10052-007-0300-0](#).
- [18] J. Blumlein, A. De Freitas, W. van Neerven, Two-loop QED Operator Matrix Elements with Massive External Fermion Lines, *Nucl. Phys. B* 855 (2012) 508–569. [arXiv:1107.4638](#), [doi:10.1016/j.nuclphysb.2011.10.009](#).
- [19] J. Ablinger, J. Blümlein, A. De Freitas, K. Schönwald, Subleading Logarithmic QED Initial State Corrections to $e^+e^- \rightarrow \gamma^*/Z^{0*}$ to $O(\alpha^6 L^5)$, *Nucl. Phys. B* 955 (2020) 115045. [arXiv:2004.04287](#), [doi:10.1016/j.nuclphysb.2020.115045](#).
- [20] J. Blümlein, A. De Freitas, K. Schönwald, The QED initial state corrections to the forward-backward asymmetry of $e^+e^- \rightarrow \gamma^*/Z^{0*}$ to higher orders, *Phys. Lett. B* 816 (2021) 136250. [arXiv:2102.12237](#), [doi:10.1016/j.physletb.2021.136250](#).
- [21] S. Frixione, Initial conditions for electron and photon structure and fragmentation functions, *JHEP* 11 (2019) 158. [arXiv:1909.03886](#), [doi:10.1007/JHEP11\(2019\)158](#).
- [22] V. Bertone, M. Cacciari, S. Frixione, G. Stagnitto, The partonic structure of the electron at the next-to-leading logarithmic accuracy in QED, *JHEP* 03 (2020) 135, [Erratum: *JHEP* 08, 108 (2022)]. [arXiv:1911.12040](#), [doi:10.1007/JHEP03\(2020\)135](#).
- [23] S. Frixione, On factorisation schemes for the electron parton distribution functions in QED, *JHEP* 07 (2021) 180, [Erratum: *JHEP* 12, 196 (2012)]. [arXiv:2105.06688](#), [doi:10.1007/JHEP07\(2021\)180](#).
- [24] V. Bertone, M. Cacciari, S. Frixione, G. Stagnitto, M. Zaro, X. Zhao, Improving methods and predictions at high-energy e+e- colliders within collinear factorisation, *JHEP* 10 (2022) 089. [arXiv:2207.03265](#), [doi:10.1007/JHEP10\(2022\)089](#).
- [25] A. B. Arbuzov, E. S. Scherbakova, Next-to-leading order corrections to Bhabha scattering in renormalization group approach. I. Soft and virtual photonic contributions, *JETP Lett.* 83 (2006) 427–432. [arXiv:hep-ph/0602119](#), [doi:10.1134/S0021364006100018](#).
- [26] A. B. Arbuzov, Leading and Next-to-Leading Logarithmic Approximations in Quantum Electrodynamics, *Phys. Part. Nucl.* 50 (6) (2019) 721–825. [doi:10.1134/S1063779619060029](#).
- [27] R. K. Ellis, W. Vogelsang, The Evolution of parton distributions beyond leading order: The Singlet case (2 1996). [arXiv:hep-ph/9602356](#).
- [28] G. Curci, W. Furmanski, R. Petronzio, Evolution of Parton Densities Beyond Leading Order: The Nonsinglet Case, *Nucl. Phys. B* 175 (1980) 27–92. [doi:10.1016/0550-3213\(80\)90003-6](#).
- [29] P. A. Baikov, K. G. Chetyrkin, J. H. Kuhn, C. Sturm, The relation between the QED charge renormalized in MSbar and on-shell schemes at four loops, the QED on-shell beta-function at five loops and asymptotic contributions to the muon anomaly at five and six loops, *Nucl. Phys. B* 867 (2013) 182–202. [arXiv:1207.2199](#), [doi:10.1016/j.nuclphysb.2012.09.018](#).

- [30] S. G. Gorishnii, A. L. Kataev, S. A. Larin, The three loop QED contributions to the photon vacuum polarization function in the $\overline{\text{MS}}$ scheme and the four loop corrections to the QED beta function in the on-shell scheme, *Phys. Lett. B* 273 (1991) 141–144, [Erratum: *Phys.Lett.B* 275, 512 (1992), Erratum: *Phys.Lett.B* 341, 448 (1995)]. doi:10.1016/0370-2693(91)90568-B.
- [31] A. Arbuzov, Higher order QED corrections to muon decay spectrum, *JHEP* 03 (2003) 063. arXiv:hep-ph/0206036, doi:10.1088/1126-6708/2003/03/063.
- [32] W. Furmanski, R. Petronzio, Singlet Parton Densities Beyond Leading Order, *Phys. Lett. B* 97 (1980) 437–442. doi:10.1016/0370-2693(80)90636-X.