

Renormalization of antiferromagnetic magnons by superconducting condensate and quasiparticles

A. M. Bobkov,¹ S. A. Sorokin,^{2,1} and I. V. Bobkova^{1,2}

¹*Moscow Institute of Physics and Technology, Dolgoprudny, 141700 Moscow region, Russia*

²*National Research University Higher School of Economics, 101000 Moscow, Russia*

(Dated: December 6, 2022)

The ability to modify and tune the spin-wave dispersion is one of the most important requirements for engineering of magnonic networks. In this study we demonstrate the promise of synthetic thin-film hybrids composed of an antiferromagnetic insulator (AF) and a normal (N) or superconducting (S) metal for tuning and modifying the spin-wave dispersion in antiferromagnetic insulators. The key ingredient is the uncompensated magnetic moment at the AF/S(N) interface, which induces an effective exchange field in the adjacent metal via the interface exchange interaction. The exchange field spin polarizes quasiparticles in the metal and induces spinful triplet Cooper pairs screening the magnon. The quasiparticle and Cooper pair polarization renormalizes the magnon dispersion. The renormalization results in the splitting of the otherwise degenerate AF magnon modes with no need of applying a magnetic field. It is also proposed that measurements of the renormalized dispersion relations can provide the amplitude of the effective exchange field induced by the AF in the adjacent metal.

I. INTRODUCTION

Magnonics is devoted to the exploration of spin waves in magnetic structures. Now it is very rapidly developing mainly due to the impressive advance of nanotechnology, the development of new experimental techniques and the promise of a new generation of devices in which magnons would be employed¹⁻³. In particular, the interconversion between magnonic spin signals and electron spin signals has been investigated⁴⁻⁹.

The other important question is a renormalization of magnon characteristics in hybrid structures, and, in particular, the influence of the adjacent metal on the magnon characteristics in thin film ferromagnet/normal metal (F/N) or ferromagnet superconductor (F/S) heterostructures. The ability to modify and tune the spin-wave dispersion is one of the most important requirements for engineering of magnonic networks. In particular, it was found that the adjacent metal works as a spin sink strongly influencing Gilbert damping of the magnon modes¹⁰⁻²⁰. Also, the adjacent superconducting layer can result in shifting of $k = 0$ magnon frequencies (Kittel mode)^{15,18-20}. The exact mechanism is still under debate now^{21,22}, but it is probably of electromagnetic nature and is not related to the exchange-based proximity effects between the ferromagnet and the superconductor. The electromagnetic interaction between the ferromagnet and superconductor also results in appearance of magnon-fluxon excitations²³ and efficient gating of magnons²⁴. Further it was reported that the proximity effect in thin film F/S hybrids, that is the generation of an effective exchange field in S by the proximity to F, results in appearing of composite particles of different physical nature. They are composed of a magnon in F and an accompanying cloud of spinful triplet pairs in S, which was termed magnon-cooparon²⁵. The cloud of triplet pairs results in the essential renormalization of the magnon stiffness and its spin. Furthermore, for thin film

S/F bilayers with an unconventional spinful triplet superconductor, a conversion of Cooper pair supercurrents to magnon spin currents has been suggested²⁶.

On the other hand, antiferromagnets (AFs) have recently gathered interest as alternatives to ferromagnets as active components in spintronics applications due to their robustness against perturbation by magnetic fields, the absence of parasitic stray fields and ultrafast dynamics²⁷⁻²⁹. In particular, antiferromagnet/normal metal hybrids have been actively studied in the context of spin transport experiments³⁰⁻³³. Antiferromagnetic magnons have been predicted to be effective mediators of the superconducting pairing in antiferromagnet/metal hybrid structures³⁴⁻³⁷.

The influence of superconductivity on the magnon spectra in antiferromagnetic superconductors has been studied in Ref. 38. It was reported that the superconductivity enhances $k = 0$ magnon frequency thus making the magnon spectrum nonmonotonic function of the magnon wave number. However, much less is known about the back action of the adjacent metal on the magnon characteristics in thin film AF/N or AF/S heterostructures. Here we address this question for antiferromagnetic insulator/normal (superconducting) metal interface if the corresponding interface is uncompensated, that is, there is a nonzero average magnetization at the interface. Due to the interface exchange coupling an effective exchange field is induced in the adjacent metal producing an electron spin polarization there³⁹. This makes the thin conducting film proximitized by the uncompensated antiferromagnet very much like a conductor in a contact with a ferromagnetic insulator. The effective exchange field generates singlet-triplet conversion in the superconducting film⁴⁰, superconductivity suppression, spin-splitting of the superconducting DOS⁴¹ and giant spin-dependent Seebeck effect⁴².

We study the back action of the spin polarization induced in the superconductor or normal metal on the

magnon spectrum of the antiferromagnetic insulator. We consider an easy-axis antiferromagnet. It is known that in this case there are two oppositely polarized magnon modes, which are degenerate unless an external magnetic field is applied. We demonstrate that the back action of the metal removes the degeneracy with no need for application of the field. Moreover, the group velocities of the oppositely polarized modes become essentially different. The general reason is that the sublattice symmetry-breaking exchange at the uncompensated interface results in the symmetry-breaking between the two opposite-spin magnon modes. The dynamical polarization of the conduction electrons interacts with the oppositely polarized modes asymmetrically. This effect takes place both for AF/N and AF/S heterostructures. It has two contributions. The first one is from quasiparticles, which dominate at small magnon wave numbers k . The second contribution is caused by the interaction of magnons with superconducting condensate and is explained by the appearance of the magnon-cooparon composite particles, analogously to the case of F/S hybrids²⁵. Further we show that damping of the magnon modes can also be renormalized by the back action of the metal, that is by magnon-assisted processes of the electronic quasiparticles spin flip. Finally, we demonstrate the appearance of the additional modes in the magnon spectrum due to the hybridization between the magnons and electron paramagnetic resonance mode, which is caused by the exchange field induced in the metal by the antiferromagnet itself.

The paper is organized as follows. In Sec. II the system under consideration and the model under study are described. In Sec. III we formulate the theoretical framework to treat the problem. In Sec. IV the results for the magnon spectrum renormalization are presented and discussed. Short summary of our research is given in Sec. V.

II. SYSTEM AND MODEL

The model system that we consider is shown in Fig. 1. It is a thin film bilayer consisting of an antiferromagnetic insulator with homogeneous Néel order interfaced to a conventional spin-singlet superconductor. AF/N interface is considered as a limiting case, corresponding to temperatures above the superconducting critical temperature. We assume uncompensated magnetic moments at the AF/S interface, that is the interface possesses finite magnetization. Under these conditions, we expect the uncompensated AF to induce a spin-splitting field in the superconductor³⁹ via an interfacial exchange interaction. It is assumed that the thickness of the S film d_S is smaller than the superconducting coherence length ξ_S . Then the induced spin-splitting can be considered as uniform along the y -direction. In general the magnetic proximity effect at an AF/S uncompensated interface is not reduced to the effective exchange only³⁹, analogously to the case of the ferromagnetic insulator/superconductor

interface^{43,44}. However, in the framework of the present study we neglect other terms which can be viewed as additional magnetic impurities in the superconductor and focus on the effect of the exchange field.




FIG. 1. Sketch of the AF/S bilayer under consideration and a schematic illustration of the circularly-polarized magnon modes, which are energetically degenerate without a coupling to the S(N) layer and an applied magnetic field.

We treat S within the quasiclassical framework and solve the Eilenberger equation to obtain Green's function and, consequently, the electron spin polarization, which is expressed in terms of the Green's function. The AF insulator is treated via the two-sublattice Landau-Lifshitz-Gilbert (LLG) description. The sublattices A and B are marked by the corresponding superscripts. The two subsystems are coupled due to the interfacial exchange, which is assumed nonzero between S and the sublattice A. The AF film is also assumed to be thin with uniform Néel order along the y -direction. The interface exchange leads to a spin-splitting term in the Eilenberger equation describing S and a spin torque term describing the magnetization dynamics for the AF sublattice A. The overall system dynamics is determined by solving the coupled equations self-consistently.

We wish to examine wavevector-resolved excitations of the hybrid. Thus we can obtain the complete information needed for examining arbitrary wavepackets generated by a given experimental method. To this end, we assume existence of a spin wave with wavevector \mathbf{k} in the AF [Fig. 1] such that the magnetization unit vector $\mathbf{m}^{A,B}(\mathbf{r}, t) = \mathbf{m}_0^{A,B} + \delta\mathbf{m}^{A,B}(\mathbf{r}, t)$ consists of the equilibrium part $\mathbf{m}_0^{A,B} = \pm\mathbf{e}_z$ and the excitation part $\delta\mathbf{m}_{\pm}^{A,B}(\mathbf{r}, t) = \delta m [\cos(\mathbf{k}\mathbf{r} + \omega t)\mathbf{e}_x \pm \sin(\mathbf{k}\mathbf{r} + \omega t)\mathbf{e}_y] \exp(-\kappa t)$. Unlike the ferromagnetic case, in AFs two magnon modes⁴⁵ are possible for the same wavevector \mathbf{k} . They are described by the excitation parts $\delta\mathbf{m}_{\pm}^{A,B}(\mathbf{r}, t)$ and are called by ω_{\pm} modes. The magnon modes are schematically shown in Fig. 1. In these modes the two sublattice magnetizations

are nearly opposite to each other. They precess circularly, counterclockwise in one of the modes and clockwise in the other. On the quantum language they correspond to magnons carrying the opposite spins. In the absence of the applied field and proximity to the superconductor (or normal metal) these modes are degenerate. The effect of the own dipolar fields can also remove the degeneracy and strongly hybridizes the spin carried by the modes⁴⁶. But here we neglect the effect of dipolar fields because according to our numerical estimates the coupling to the S(N) film generates larger splitting of the modes, of the order of tens of GHz, while the splitting due to the dipolar interaction was estimated as $\sim \text{GHz}$ ⁴⁶. In addition the proximity-induced splitting manifest very strong temperature dependence at low (superconducting) temperatures, what makes possible to distinguish it. We assume magnon wavevector \mathbf{k} to be in plane of the AF/S interface.

III. FORMALISM

A. Description of magnons in AF

The magnetization dynamics is described within the Landau-Lifshitz-Gilbert framework. The LLG equation can be written for each sublattice separately⁴⁷:

$$\frac{\partial \mathbf{m}^i}{\partial t} = -\gamma \mathbf{m}^i \times \mathbf{H}_{\text{eff}}^i + \sum_j \alpha_{ij} \mathbf{m}^i \times \frac{\partial \mathbf{m}^j}{\partial t} + \mathbf{N}^i, \quad (1)$$

where $i = A, B$ is the sublattice index and $-\gamma$ with $\gamma > 0$ is the AF gyromagnetic ratio, $\mathbf{H}_{\text{eff}}^i = Km_z^i \mathbf{e}_z + A \nabla^2 \mathbf{m}^i - J_{\text{AF}} \mathbf{m}^{\bar{i}} + \mathbf{H}$ is the effective magnetic field in the AF containing an external magnetic field \mathbf{H} , easy-axis anisotropy with easy axis along z -direction and anisotropy constant K , exchange stiffness A and antiferromagnetic exchange coupling between two sublattices with coupling constant $J_{\text{AF}} > 0$. $\bar{i} = B(A)$ for $i = A(B)$. α_{ij} is the 2×2 Gilbert dissipation matrix^{47,48}, which can be characterized by two real positive numbers α and α_c as follows: $\alpha_{AA} = \alpha_{BB} = \alpha$ and $\alpha_{AB} = \alpha_{BA} = \alpha_c$. The last term in Eq. (1) represents the torque experienced by the sublattice.

The torque \mathbf{N}^i can be calculated starting from the effective exchange interaction between the spin densities

on the two sides of the AF/S interface:

$$H_{\text{int}} = - \int d^2 \mathbf{r} J \mathbf{S}^A \cdot \mathbf{s}, \quad (2)$$

where \mathbf{s} is the electronic spin density operator in the S film, \mathbf{S}^A is the localized spin operator in the AF film, belonging to the sublattice A. J parameterizes interfacial exchange interaction between AF and S and the integration is performed over the 2D interface. It has been shown³⁹ that this exchange interaction hamiltonian results in the appearance of the exchange field $\mathbf{h}(\mathbf{r}) = JM\mathbf{m}^A(\mathbf{r})/(2\gamma d_S)$ in the S film, where M is the sublattice saturation magnetization.

From Eq. (2) one obtains the additional contribution to the Landau-Lifshitz-Gilbert equation written in the form of an interface torque acting on the magnetization:

$$\mathbf{N}_{\text{int}}^A = J\delta(y - y_I)\mathbf{m}^A(y) \times \mathbf{s}, \quad \mathbf{N}^B = 0, \quad (3)$$

where the interface is located at $y = y_I$. We assume that the antiferromagnetic film is thin and its magnetization for a given sublattice is homogeneous in the y -direction. In this case Eqs. (1) and (3) can be averaged over the thickness d_{AF} of the AF film. For the averaged torque we thus obtain

$$\mathbf{N}^A = \frac{J\mathbf{m}^A \times \mathbf{s}}{d_{\text{AF}}} = \tilde{J}\mathbf{m}^A \times \mathbf{s}, \quad \mathbf{N}^B = 0. \quad (4)$$

Here $\tilde{J} \equiv J/d_{\text{AF}}$. We expand the electronic spin polarization over the three perpendicular vectors as

$$\mathbf{s} = s_0 \mathbf{e}_z + \delta s_{\parallel} \delta \mathbf{m}^A + \delta s_{\perp} (\delta \mathbf{m}^A \times \mathbf{e}_z), \quad (5)$$

where s_0 is the equilibrium value of the electron spin polarization in the superconductor, corresponding to the absence of the magnon. δs_{\parallel} and δs_{\perp} describe the dynamic corrections to the spin polarization due to the magnon. The components s_0 , δs_{\parallel} and δs_{\perp} are in general functions of (ω, \mathbf{k}) and are calculated in the framework of the Usadel equations below. Substituting the expressions for \mathbf{s} and $\mathbf{m}^{A,B}(\mathbf{r}, t) = \pm \mathbf{e}_z + \delta \mathbf{m}^{A,B}(\mathbf{r}, t)$ in Eq. (1) above and linearizing this equation with respect to $\delta \mathbf{m}^{A,B}$, we obtain

$$\begin{pmatrix} -\omega - i\kappa \pm B_{\pm} \pm \delta\omega_{\pm} \pm i\omega\alpha - i\chi_{\pm} \\ \mp \gamma J_{\text{AF}} \mp i\omega\alpha_c \end{pmatrix} \begin{pmatrix} \delta m_{\pm}^A \\ \delta m_{\pm}^B \end{pmatrix} = 0, \quad (6)$$

where $B_{\pm} = \gamma(K + J_{\text{AF}} + Ak^2 \pm H) = B \pm \gamma H$. Eq. (6) decouples giving two independent circularly polarized magnon modes $\delta m_{\pm}^{A(B)} = \delta m_x^{A(B)} \pm i\delta m_y^{A(B)}$. Terms $\delta\omega_{\pm}$ and χ_{\pm} result from the torque acting on the AF from the

superconductor. $\delta\omega_{\pm} = \tilde{J}[\delta s_{\parallel}(\pm\omega, \pm\mathbf{k}) - s_0(\pm\omega, \pm\mathbf{k})]$ and $\chi_{\pm} = \tilde{J}\delta s_{\perp}(\pm\omega, \pm\mathbf{k})$. Neglecting the small terms resulting from $(\omega\alpha)^2$, $(\omega\alpha_c)^2$ and χ_{\pm}^2 in Eq. (6) we obtain the following expression for the mode frequencies,

which are determined by the real part ω of the solution $\omega_{\pm} + i\kappa_{\pm}$, providing zero determinant of the matrix in Eq. (6):

$$\omega_{\pm} = \pm \left(\frac{\delta\omega_{\pm}}{2} + \gamma H \right) + \sqrt{\left(\frac{\delta\omega_{\pm}}{2} \right)^2 + \omega_0^2 + \delta\omega_{\pm} B}, \quad (7)$$

where $\omega_0 = \gamma \sqrt{(K + J_{\text{AF}} + Ak^2)^2 - J_{\text{AF}}^2}$ is the AF magnon frequency at $H = 0$ and without any contact with S(N) layer. It is seen from Eq. (7) that the interaction of the AF magnons with polarized electrons in the adjacent metal layer removes degeneracy of the magnon modes even at zero magnetic field.

The magnon decay rate κ is also renormalized by the influence of the metal layer. The renormalized κ is determined by the imaginary part κ of the solution $\omega_{\pm} + i\kappa_{\pm}$ providing zero determinant of the matrix in Eq. (6) and up to the linear order with respect to α , α_c and χ_{\pm} takes the form:

$$\kappa_{\pm} = \frac{\alpha \omega \left(B + \frac{\delta\omega_{\pm}}{2} \right) - J_{\text{AF}} \omega \alpha_c \mp \frac{\chi_{\pm}}{2} (B_{-} \pm \omega)}{\sqrt{\left(\frac{\delta\omega_{\pm}}{2} \right)^2 + \omega_0^2 + \delta\omega_{\pm} B}}. \quad (8)$$

B. Electron spin polarization in S(N)

Now our goal is to calculate the electron spin polarization \mathbf{s} in the superconductor. The approach closely follows Ref. 25, where it has been used for treating S/F bilayers. Here we provide the corresponding equations for consistency and generalize the calculations for the case of two magnon modes. \mathbf{s} can be calculated via the quasiclassical Green's function as

$$\mathbf{s} = -\frac{N_F}{16} \int d\varepsilon \text{Tr}_4 \left[\boldsymbol{\sigma} \tau_z \check{g}^K \right], \quad (9)$$

where \check{g}^K is the Keldysh component of the quasiclassical Green's function \check{g} , which is a 8×8 matrix in the direct product of Keldysh, spin and particle-hole spaces. It obeys the Usadel equation:

$$iD \nabla (\check{g} \otimes \nabla \check{g}) = [\varepsilon \tau_z - \mathbf{h} \boldsymbol{\sigma} \tau_z + \Delta i \tau_y, \check{g}]_{\otimes}, \quad (10)$$

where $[A, B]_{\otimes} = A \otimes B - B \otimes A$ and we work in the mixed (ε, t) representation with $A \otimes B = \exp[(i/2)(\partial_{\varepsilon_1} \partial_{t_2} - \partial_{\varepsilon_2} \partial_{t_1})] A(\varepsilon_1, t_1) B(\varepsilon_2, t_2)|_{\varepsilon_1=\varepsilon_2=\varepsilon; t_1=t_2=t}$. In case $A[B](\varepsilon, t) = A[B](\varepsilon) \exp[i\Omega_{A[B]} t]$ the \otimes -product is reduced to $A(\varepsilon, t) \otimes B(\varepsilon, t) = A(\varepsilon - \Omega_{A/2}, t) \exp[i(\Omega_A + \Omega_B)t]$. $\tau_{x,y,z}$ and $\sigma_{x,y,z}$ are Pauli matrices in particle-hole and spin spaces, respectively. Δ is the superconducting order parameter. The explicit structure of the Green's function in the Keldysh space takes the form:

$$\check{g} = \begin{pmatrix} \check{g}^R & \check{g}^K \\ 0 & \check{g}^A \end{pmatrix}, \quad (11)$$

where $\check{g}^{R(A)}$ is the retarded (advanced) component of the Green's function and \check{g}^K is the Keldysh component. Further we express the Keldysh part of the Green's function

via the retarded, advanced Green's functions and the distribution function $\check{\varphi}$ as follows: $\check{g}^K = \check{g}^R \otimes \check{\varphi} - \check{\varphi} \otimes \check{g}^A$.

The exchange field is taken in the form of a time-independent component and a circularly polarized magnon:

$$\mathbf{h}_{\pm} = h_0 \mathbf{e}_z + \delta h \cos(\mathbf{k}\mathbf{r} + \omega t) \mathbf{e}_x \pm \delta h \sin(\mathbf{k}\mathbf{r} + \omega t) \mathbf{e}_y \quad (12)$$

Please note that \mathbf{h}_{\pm} , respectively induced by the ω_{\pm} magnon modes, does not have a sublattice index. This is because the superconductor is only coupled to the A-sublattice of the antiferromagnet and the induced exchange field is just an imprint of this sublattice $\mathbf{h}_{\pm} = JM \mathbf{m}_{\pm}^A / (2\gamma d_S)$. Then

$$\delta \mathbf{h}_{\pm} \boldsymbol{\sigma} = \delta h e^{\mp i(\mathbf{k}\mathbf{r} + \omega t) \sigma_z} \boldsymbol{\sigma}_x. \quad (13)$$

The quasiclassical Green's function is to be found in the form: $\check{g}_{\pm} = \check{g}_0 + \delta \check{g}_{\pm}$, where \check{g}_0 is the Green's function in the absence of the magnon and $\delta \check{g}_{\pm}$ is the first order correction with respect to $\delta \mathbf{h}_{\pm}$. Taking into account that $\nabla \check{g}_0 = 0$ (we assume that in the absence of the magnon the bilayer is spatially homogeneous along the interface) from Eq. (10) we obtain the following equation for $\delta \check{g}_{\pm}$:

$$iD \check{g}_0 \otimes \nabla^2 \delta \check{g}_{\pm} = [\varepsilon \tau_z - h_0 \mathbf{e}_z \sigma_z \tau_z + \tau_z \hat{\Delta}, \delta \check{g}_{\pm}]_{\otimes} - [\delta h e^{-i(\mathbf{k}\mathbf{r} + \omega t) \sigma_z} \boldsymbol{\sigma}_x \tau_z, \check{g}_0]_{\otimes} \quad (14)$$

Introducing the unitary operator $\hat{U} = e^{-i(\mathbf{k}\mathbf{r} + \omega t) \sigma_z / 2}$ we can transform the Green's function as follows:

$$\delta \check{g}_{\pm} = U \otimes \delta \check{g}_m \otimes U^{\dagger}. \quad (15)$$

In case if the system is spatially homogeneous except for the magnon, $\delta \check{g}_m$ does not depend on coordinates. Then

$$\nabla^2 \delta \check{g}_{\pm} = -\frac{k^2}{2} \check{U} \otimes (\delta \check{g}_m - \sigma_z \delta \check{g}_m \sigma_z) \otimes U^{\dagger} = -k^2 \delta \check{g}_{\pm} \quad (16)$$

where when passing to the second equality it is used that $\delta \check{g}_{\pm} = \delta \hat{g}_x \sigma_x + \delta \hat{g}_y \sigma_y$ and has no z -component in the spin space according to the spin structure of the magnon exchange field $\delta \mathbf{h}_{\pm}$.

From the normalization condition $\check{g} \otimes \check{g} = 1$ it follows that $\check{g}_0 \otimes \delta \check{g}_{\pm} = -\delta \check{g}_{\pm} \otimes \check{g}_0$. It gives us $\check{g}_0 \otimes \delta \check{g}_{\pm} = (1/2)[\check{g}_0, \delta \check{g}_{\pm}]_{\otimes}$. Eq. (14) takes the form:

$$\begin{aligned} & [\varepsilon \tau_z + i \frac{Dk^2}{2} \check{g}_0 - h_0 \mathbf{e}_z \sigma_z \tau_z + \tau_z \hat{\Delta}, \delta \check{g}_{\pm}]_{\otimes} - \\ & [\delta h e^{-i(\mathbf{k}\mathbf{r} + \omega t) \sigma_z} \boldsymbol{\sigma}_x \tau_z, \check{g}_0]_{\otimes} = 0. \end{aligned} \quad (17)$$

From Eq. (17) the following equation for $\delta \check{g}_m$ is obtained:

$$\begin{aligned} & [\hat{\Lambda}_d \tau_z + \hat{\Lambda}_{od} i \tau_y, \delta \check{g}_m] = \delta h [\sigma_x \tau_z, \\ & \frac{1}{2} ((g_{0,s} + g_{0,a} \sigma_z) \tau_z + (f_{0,s} + f_{0,a} \sigma_z) i \tau_y)]. \end{aligned} \quad (18)$$

It does not contain time dependence and \otimes -products. In Eq. (18) we use the following definitions

$$g_{0,s(a)} = g_{0,\uparrow}(\varepsilon + \frac{\omega}{2}) \pm g_{0,\downarrow}(\varepsilon - \frac{\omega}{2}), \quad (19)$$

$$f_{0,s(a)} = f_{0,\uparrow}(\varepsilon + \frac{\omega}{2}) \pm f_{0,\downarrow}(\varepsilon - \frac{\omega}{2}), \quad (20)$$

where $g_{0,\uparrow(\downarrow)}$ represent the bulk Green's functions for the superconductor in the exchange field h_0 :

$$g_{0,\uparrow(\downarrow)}^R = \frac{|\varepsilon \mp h_0|}{\sqrt{(\varepsilon + i\delta \mp h_0)^2 - \Delta^2}} \quad (21)$$

$$f_{0,\uparrow(\downarrow)}^R = \frac{\Delta \text{sgn}(\varepsilon \mp h_0)}{\sqrt{(\varepsilon + i\delta \mp h_0)^2 - \Delta^2}}, \quad (22)$$

and $g(f)_{0,\uparrow(\downarrow)}^A = -g(f)_{0,\uparrow(\downarrow)}^{R*}$, $g(f)_{0,\uparrow(\downarrow)}^K = [g(f)_{0,\uparrow(\downarrow)}^R - g(f)_{0,\uparrow(\downarrow)}^A] \tanh[\varepsilon/2T]$.

$$\hat{\Lambda}_d = \Lambda_d^0 + \Lambda_d^z \sigma_z = \varepsilon + \frac{iDk^2}{4} g_{0,s} + \left(\frac{\omega}{2} - h_0 + \frac{iDk^2}{4} g_{0,a}\right) \sigma_z, \quad (23)$$

$$\hat{\Lambda}_{od} = \Lambda_{od}^0 + \Lambda_{od}^z \sigma_z = \Delta + \frac{iDk^2}{4} f_{0,s} + \frac{iDk^2}{4} f_{0,a} \sigma_z. \quad (24)$$

Solving Eq. (18) we obtain:

$$\delta \check{g}_m = \delta g_{mx} \sigma_x \tau_z + \delta f_{mx} \sigma_x i \tau_y, \quad (25)$$

where

$$\delta g_{mx} = \frac{\delta h [f_{0,s} \Lambda_{od}^z - g_{0,a} \Lambda_d^0]}{2[\Lambda_d^0 \Lambda_d^z - \Lambda_{od}^0 \Lambda_{od}^z]} \quad (26)$$

The distribution function also acquires a correction due to the magnon: $\check{\varphi}_+ = \check{\varphi}_0 + \delta \check{\varphi}_+$. It is convenient to work with the transformed distribution function $\hat{U}^\dagger \check{\varphi}_+ \hat{U} = \check{\varphi}_m + \delta \varphi_m$, which does not depend on time and spatial coordinates. Here $\check{\varphi}_m = (1/2)[\varphi_{m,s} + \varphi_{m,a} \sigma_z]$ with $\varphi_{m,s(a)} = \tanh[(\varepsilon + \omega/2)/2T] \pm \tanh[(\varepsilon - \omega/2)/2T]$ is the result of the unitary transformation of the equilibrium distribution function $\check{\varphi}_0 = \tanh[\varepsilon/2T]$. From the Keldysh part of the Usadel equation (18) we can obtain the following equation for the first order correction to the distribution function $\delta \check{\varphi}_m$:

$$iDk^2 [\delta \check{\varphi}_m - \check{g}_{m0}^R \delta \check{\varphi}_m \check{g}_{m0}^A] + \check{g}_{m0}^R [\check{K}, \delta \check{\varphi}_m] - [\check{K}, \delta \check{\varphi}_m] \check{g}_{m0}^A + \check{g}_{m0}^R [\check{\varphi}_m, \delta h \sigma_x \tau_z] - [\check{\varphi}_m, \delta h \sigma_x \tau_z] \check{g}_{m0}^A = 0, \quad (27)$$

where

$$\check{K} = (\varepsilon + (\omega/2 - h_0) \sigma_z) \tau_z + \Delta i \tau_y, \quad (28)$$

$$\check{g}_{m0}^{R,A} = \hat{U}^\dagger \otimes \check{g}_0^{R,A} \otimes \hat{U} = (1/2)[(g_{0,s}^{R,A} + g_{0,a}^{R,A} \sigma_z) \tau_z + (f_{0,s}^{R,A} + f_{0,a}^{R,A} \sigma_z) i \tau_y], \quad (29)$$

Structure of Eq. (27) dictates that

$$\delta \check{\varphi}_m = \begin{pmatrix} 0 & \delta \varphi_m^\uparrow \\ \delta \varphi_m^\downarrow & 0 \end{pmatrix}. \quad (30)$$

Substituting Eq. (30) into Eq. (27) we obtain the following result:

$$\delta \varphi_m^\sigma = -\delta h \varphi_{m,a} \times \frac{2h_{0\omega} G_{-, \sigma} + i\sigma Dk^2 (g_{m\sigma}^R - g_{m\bar{\sigma}}^A)}{4h_{0\omega}^2 G_{-, \sigma} + 4h_{0\omega} i\sigma Dk^2 (g_{m\sigma}^R - g_{m\bar{\sigma}}^A) - (Dk^2)^2 G_{+, \sigma}}. \quad (31)$$

where we introduce the spin subband index $\sigma = \uparrow (\downarrow)$ in the subscripts/superscripts and $\sigma = \pm 1$ for spin-up(down) subbands, respectively, if it is as a factor. $\bar{\sigma} = -\sigma$, $h_{0\omega} = \omega/2 - h_0$, $g(f)_{m\sigma}^{R,A} = g(f)_{0,\sigma}^{R,A}(\varepsilon + \sigma\omega/2)$ and $G_{\pm, \sigma} = 1 - g_{m\sigma}^R g_{m\bar{\sigma}}^A \pm f_{m\sigma}^R f_{m\bar{\sigma}}^A$.

The electron spin polarization \mathbf{s}_+ , induced in the superconductor by the equilibrium magnetization \mathbf{m}_0^A and the magnon mode $\delta \mathbf{m}_+^A$ is calculated from Eq. (9). It can be decomposed into three orthogonal components according to Eq. (5). The corresponding components take the form:

$$s_0 = -\frac{N_F}{4} \int_{-\infty}^{\infty} d\varepsilon \tanh \frac{\varepsilon}{2T} \text{Re}[g_{0,\uparrow}^R - g_{0,\downarrow}^R], \quad (32)$$

$$\delta s_{\parallel}(\omega, \mathbf{k}) = -\frac{N_F h_0}{8\delta h} \int_{-\infty}^{\infty} d\varepsilon \left\{ 2\varphi_{m,s} \text{Re}[\delta g_{mx}^R] + \sum_{\sigma} (g_{m\sigma}^R - g_{m\bar{\sigma}}^A) \delta \varphi_m^{\sigma} \right\}, \quad (33)$$

$$\delta s_{\perp}(\omega, \mathbf{k}) = \frac{N_F h_0}{8\delta h} \int_{-\infty}^{\infty} d\varepsilon \left\{ 2\varphi_{m,a} \text{Im}[\delta g_{mx}^R] + i \sum_{\sigma} \sigma (g_{m\sigma}^R - g_{m\bar{\sigma}}^A) \delta \varphi_m^{\sigma} \right\}. \quad (34)$$

The electron spin polarization induced by the mode $\delta \mathbf{m}_+^A$ can be obtained from Eqs. (33) and (34) by the substitution $\omega \rightarrow -\omega$ and $\mathbf{k} \rightarrow -\mathbf{k}$.

IV. RESULTS AND DISCUSSION

The dispersion relations for the both AF magnon modes $\omega_{\pm}(\mathbf{k})$ at $H = 0$ are presented in Fig. 2. Panel (a) represents the general view of the mode frequencies on a large scale of the order of Δ_0 , where Δ_0 is a value of the superconducting gap in the bulk superconductor at $T = 0$ and without a proximity to the antiferromagnet. The numerical parameters of the AF/S bilayer are taken to be close to the material parameters of MnF_2 (AF)⁴⁵ and Nb (S). The upper mode is the original AF mode, split and renormalized by superconductivity. While the characteristic AF frequencies $\omega_0 \sim \Delta_0$ are of the order of THz, all the effects of splitting and renormalization are controlled by the parameter $\beta = \tilde{J} N_F \Delta_0$. Taking $d_{\text{AF}} = d_S$, $M \sim 230\text{G}$, $\gamma = 2\mu_B/\hbar$ and $N_F \approx 1.3 \times 10^{35} \text{erg}^{-1} \text{cm}^{-3}$ for Nb, we obtain for chosen materials $\beta \sim 0.05h_0$. The equilibrium effective exchange field in the S(N) layer for the results presented in Fig. 2 is assumed to be $h_0 = 0.2\Delta_0$. This value corresponds to the available estimates of the effective field at the uncompensated AF/N interfaces of real materials³⁹.

For the chosen parameters we obtain that the superconductivity-induced splitting of the AF magnon

mode is of the order of tens of GHz. For this reason panel (b) represents the upper mode on the appropriate scale. First of all, it is seen from Fig. 2(a) that the magnon modes are split even at zero magnetic field. Both $\omega_{\pm}(\mathbf{k} = 0)$ and the magnon velocity $\mathbf{v}_{m,\pm} = d\omega_{\pm}/d\mathbf{k}$ are renormalized by the proximity to the adjacent metal. The physical reason for this splitting is that only the magnetization of A -sublattice \mathbf{m}^A is coupled to the conductor. The amplitudes of magnon magnetizations at A -sublattice are different for \mathbf{m}_{\pm} modes, see Fig. 1. This results in different effective exchange fields, induced in the conductor by these modes, what, in its turn, results in nonequivalent back action of the electron polarization on the magnon.




FIG. 2. Magnon dispersion $\omega(k)$ for $h_0 = 0.2\Delta_0$. ω is measured in units of Δ_0 and k is measured in units of $2c^{-1}$, where c is a length scale of the order of the interatomic distance. We take $c = 3.3\text{\AA}$ in accordance with the lattice period along \hat{c} -axis of MnF_2 ⁴⁹. Panel (a) represents the general view of the magnon modes dispersion in the AF/S bilayer. Panel (b) is the zoom of the rectangular region in panel (a). Different colors correspond to different temperatures. ω_+ -mode is plotted by solid lines and ω_- -mode is plotted by dashed lines. The original AF dispersion $\omega_0(k)$ is shown by grey line in panel (b). Panel (c) is the zoom of ω_{2h_0} -mode, which is seen as a black point in panel (a). Different colors are again different temperatures. The decay rate is encoded by the brightness of the corresponding line. We take $\beta = 0.05h_0$, $\alpha = 2 \times 10^{-4}$, $\alpha_c = 10^{-4}$, $\xi_S = 10\text{nm}$, $\Delta_0 = 18\text{K}$, $\gamma K = 0.061\Delta_0$, $\gamma J_{AF} = 3.92\Delta_0$, $\gamma A(c/2\xi_S)^2 = 2.26\Delta_0$ throughout the paper.

Moreover, due to the resonance shape of $\delta\omega_{\pm}(\omega_{\pm})$ [see Eq.(37) below], the nonlinear equation (7) gives additional solutions for magnon frequencies, which do not occur in the absence of the adjacent conductor. The corresponding magnon frequencies are located at $\omega_{2h_0} \approx 2h_0$. Their physical origin is from the hybridization of the magnon modes and the electron paramagnetic resonance mode in the conductor. However, the important point is that in the considered case the electron paramagnetic res-

onance mode does not occur in the S(N) layer by itself and appears due to the interaction with the antiferromagnet. They are represented in panel (c) of Fig. 2. We discuss the additional modes in more details below.

Let us first concentrate on the renormalization of the AF magnon mode by quasiparticles or normal electrons. In panel (b) of Fig. 2 the corresponding frequencies are plotted for several temperatures. Different colors correspond to different temperatures. Solid and dashed lines of the same color represent ω_{\pm} for the same temperature. The renormalization of the magnon modes is originated from two different physical sources. One of them is the influence of quasiparticle polarization and the other one is the effect of accompanying cloud of Cooper pairs. To study both contributions in more detail, in Fig. 3 we plot the dependence of the mode frequencies ω_{\pm} on temperature for three representative values of the magnon wave number k .




FIG. 3. Dependence of the mode frequencies ω_+ (blue) and ω_- (red) on temperature for three representative values of the magnon wave number k : (a) $k = 0$, (b) $kc/2 = 0.009$, (c) $kc/2 = 0.03$. For all panels black dashed line is ω_0 . Vertical grey line marks the superconducting critical temperature of the AF/S bilayer. $h_0 = 0.2\Delta_0$.

From panel (a) of Fig. 3 one can see that at $k = 0$ (Kittel mode) the renormalization is fully suppressed at $T = 0$. That means the quasiparticle nature of the effect. Due to the presence of the superconducting gap the quasiparticles in the S layer are suppressed at $T \rightarrow 0$. Physically the renormalization of the $k = 0$ AF frequency can be understood as follows. The AF magnetization, including the equilibrium magnetization and a magnon, induces an exchange field \mathbf{h} in the superconductor. This exchange field polarizes quasiparticles. If the AF magnetization were static, the quasiparticle polarization would be aligned with it and would not generate a torque on the AF magnetization. However, due to the dynamical nature of the AF magnetization with finite frequency ω_0 the instantaneous quasiparticle polarization in S is not aligned with it and works as an additional contribution to the anisotropy field, but different for the both AF magnon modes. The perpendicular component of the polarization is determined by the difference $\delta s_{\parallel}(\pm\omega, k = 0) - s_0(\pm\omega, k = 0)$, which disappears at $\omega_{\pm}/\Delta_0 \rightarrow 0$. This clearly indicates that the asymmetric shift of the $k = 0$ magnons originates from the retardation effects. As it is seen from Fig. 3(a) the $k = 0$ frequency shift is the most pronounced in the normal state when the superconducting gap disappears. In this limit the induced electron polarization in N can be found analytically:

$$\delta s_{\parallel,N} = N_F \frac{\omega h_0 (\omega - 2h_0)}{(\omega - 2h_0)^2 + (Dk^2)^2}, \quad (35)$$

$$\delta s_{\perp,N} = -N_F \frac{\omega h_0 Dk^2}{(\omega - 2h_0)^2 + (Dk^2)^2}, \quad (36)$$

while $s_{0,N} = 0$. For the ω_+ mode this results in the following expressions for $\delta\omega$ and χ :

$$\delta\omega_{+,N} = N_F \tilde{J} \frac{\omega h_0 (\omega - 2h_0)}{(\omega - 2h_0)^2 + (Dk^2)^2}, \quad (37)$$

$$\chi_{+,N} = -N_F \tilde{J} \frac{\omega h_0 Dk^2}{(\omega - 2h_0)^2 + (Dk^2)^2}. \quad (38)$$

$\delta\omega_{-,N}(\omega) = \delta\omega_{+,N}(-\omega)$, the same is valid for $\chi_{-,N}$. It is seen that $\delta\omega_+ \neq \delta\omega_-$. The most important contribution of the coupling $\delta\omega_{\pm}$ to the AF mode splitting in Eq. (7) is via the last term $\delta\omega_{\pm}B$. Disregarding the smaller terms containing $\delta\omega_{\pm}$ in this equation, at $H = 0$ and $k = 0$ we obtain that $\omega_{\pm} \approx \sqrt{\omega_0^2 + \delta\omega_{\pm}B} \approx \sqrt{2\gamma J_{\text{AF}}(\gamma K + \delta\omega_{\pm}/2)}$. That is, $\delta\omega_{\pm}$ acts on the AF magnetization at $k = 0$ in the same way as an anisotropy field does, but the contribution to the anisotropy field is different for the both magnon modes.

The similar shift of the Kittel mode $k = 0$ also takes place for S/F bilayers^{20,25}, but in that case the characteristic magnon frequencies are of the order of GHz and, consequently, $\omega_0/\Delta_0 \ll 1$. This leads to the fact that the effective dynamical contribution to the anisotropy field is small. Therefore, the $k = 0$ shift is negligible for ferromagnetic magnons and the main renormalization there is due to the Cooper pairs²⁵. Here, for the AF problem,

the both sources of the renormalization are essential. For small magnon wave numbers k the main contribution is due to quasiparticles. This contribution is suppressed at $k \sim \sqrt{|\omega - 2h_0|/D}$, as it is seen from Eq. (35).

To effectively generate equal-spin pairs screening the magnon, the spatial inhomogeneity of the exchange field in the superconductor is required²⁵. Consequently, this process is more efficient at large k , comparable to the inverse superconducting coherence length $\xi_S^{-1} = \sqrt{D/\Delta_0}$, which in our case corresponds to $k_S c/2 = \xi_S^{-1} c/2 = 0.0165$. Panel (c) of Fig. 3 demonstrates the temperature dependence of the magnon frequencies renormalization at large $k > k_S$. It is seen that the temperature dependence is opposite to the $k = 0$ limit, presented in panel (a). This is because the quasiparticle renormalization is already not effective at such wave numbers and the main effect is due to the screening of the magnons by the Cooper pairs. Panel (b) shows an intermediate case when the pair contribution only starts to come into play.




FIG. 4. Magnon dispersion in the AF/S bilayer at (a) $h_0 = 0.35\Delta_0$ and (b) $h_0 = 0.5\Delta_0$. Inset to panel (b) demonstrates ω_{2h_0} mode. Temperature $T = 0.2\Delta_0$ for the both panels.

Fig. 4 demonstrates how the renormalization of the magnons depends on the value of the induced exchange field h_0 . While our basic consideration presented above corresponds to $2h_0 = 0.4\Delta_0 < \omega_0(k = 0)$, panels (a) and (b) of Fig. 4 illustrate two other possible cases $2h_0 \approx \omega_0$ and $2h_0 > \omega_0$, respectively. It is worth mentioning that the amplitude of the effective exchange $\sim d_S^{-1}$ and, therefore, can be tuned by varying the metal layer thickness. It is seen from Eq. (37) that $\delta\omega_+$ has a resonance shape peaked at $2h_0$ and originated from the electron

paramagnetic resonance in the S(N) layer. That results in additional solutions of nonlinear Eq. (7) located at $\omega_{2h_0} \approx 2h_0$. When $\omega_0(k=0)$ and ω_{2h_0} are well separated, the corresponding additional branches are above or below the main AF splitted modes and do not disturb them, as it is shown in Fig. 2(a) and Fig. 4(b). However, for the case $\omega_{2h_0} \approx \omega_0$ the additional mode is strongly coupled to ω_+ resulting in very essential modifications of the dispersion curves, as it is seen from Fig. 4(a). In principle, the additional mode, originated from the electron paramagnetic resonance can serve as an experimental probe of the effective exchange field induced in the metal layer.




FIG. 5. (a) Dependence of the modes $\omega_{\pm}(k=0)$ and ω_{2h_0} on the external magnetic field, applied along the easy axis. Grey lines correspond to $\omega_{0,\pm}(k=0)$. While $\omega_{0,+}$ is seen in panel (a), $\omega_{0,-}$ cannot be distinguished from the renormalized red line ω_- . (b) Difference between the renormalized magnon frequencies of the AF/S(N) bilayer and ω_0 as a function of the applied field. $k=0$. Solid lines represent the difference $\omega_+(H) - \omega_{0,+}(H)$ (blue) and $\omega_-(H) - \omega_{0,-}(H)$ (red) at $T = 0.4\Delta_0$, and the dotted lines correspond to the same differences but plotted at $T = T_c$. $h_0 = 0.2\Delta_0$ for the both panels.

Further, in Fig. 5 we present the dependence of the split magnon frequencies at $k=0$ on the external field H , applied along the easy-axis of the antiferromagnet. The typical linear field-induced splitting of the both modes is shown in panel (a). As it was already discussed above, the renormalization of the AF frequencies by the adjacent conductor is not clearly seen on the scale Δ_0 . For this reason in panel (b) we plot the difference between the magnon frequencies in the AF/S(N) bilayer and the corresponding magnon frequency of the bare an-

tiferromagnet $\omega_{\pm}(k=0) - \omega_{0,\pm}(k=0)$ as a function of H . Here $\omega_{0,\pm}$ means the corresponding mode without the proximity to the metal. The solid lines represent low-temperature results calculated in the superconducting state at $T = 0.4\Delta_0$, and the dotted lines are the high-temperature results at $T = T_c$, which actually correspond to the normal state. It is seen that the difference $\omega_{\pm} - \omega_{0,\pm}$ grows with temperature supporting the quasi-particle origin of the $k=0$ renormalization, discussed above. It is also important that $\omega_{\pm} - \omega_{0,\pm}$ is an asymmetric function of the applied field. This is obviously due to the presence of the effective exchange h_0 in the metal layer. In this case the applied field also contributes to the effective exchange as $h_0 \rightarrow h_0 + \mu_B H$. The difference $\omega_{\pm} - \omega_{0,\pm}$ goes to zero at $\gamma H = -2h_0$. This provides another method of experimental measurement of h_0 : the induced exchange is determined by the applied field when the difference between $\omega_+(k=0)$ in the superconducting and normal states of the metal layer disappears.

V. CONCLUSIONS

In the present paper we have investigated the renormalization of the magnon modes in a thin-film antiferromagnetic insulator caused by the proximity to a thin metal layer, which can be normal or superconducting. The key point is the presence of an uncompensated magnetic moment at the AF/S interface, which induces an effective exchange field in the adjacent metal via the interface exchange interaction. The exchange field spin polarizes quasiparticles in the metal and induces spinful triplet Cooper pairs screening the magnon, which provide an additional dynamical contribution to the magnetic anisotropy and influence the magnon velocity. The renormalization results in the splitting of the AF magnon modes with no need of applying a magnetic field. The physical reason for this splitting is asymmetric coupling of two antiferromagnet sublattices to the metal. The proximity effect also leads to appearance of additional modes in the magnon spectrum due to the hybridization between the magnons and electron paramagnetic resonance mode, which is caused by the exchange field induced in the metal by the antiferromagnet itself.

The dependence of the renormalized magnon modes in the AF/S(N) bilayer on the applied magnetic field is also studied. It is proposed that measurements of the renormalized dispersion relations can provide direct information about the amplitude of the effective exchange field induced by the AF in the adjacent metal. Our study demonstrates the promise of synthetic thin-film hybrids for tuning and modifying the spin-wave dispersion in antiferromagnets by manipulating the temperature or thickness of the metallic layer.

ACKNOWLEDGMENTS

We are grateful to Akashdeep Kamra for valuable discussions. The financial support from the Russian Sci-

ence Foundation via the RSF project No.22-42-04408 is acknowledged.

-
- ¹ S. O. Demokritov and A. N. Slavin, *Magnonics from Fundamentals to Applications* (Berlin: Springer, 2013).
- ² S.M. Rezende, *Fundamentals of Magnonics* (Berlin: Springer, 2020).
- ³ Anjan Barman, Gianluca Gubbiotti, S. Ladak, A. O. Adeyeye, M. Krawczyk, J. Gräfe, C. Adelman, S. Coto-fana, A. Naeemi, V. I. Vasyuchka, B. Hillebrands, S. A. Nikitov, H. Yu, D. Grundler, A. V. Sadovnikov, A. A. Grachev, S. E. Sheshukova, J.-Y. Duquesne, M. Marangolo, G. Csaba, W. Porod, V. E. Demidov, S. Urazhdin, S. O. Demokritov, E. Albisetti, D. Petti, R. Bertacco, H. Schultheiss, V. V. Kruglyak, V. D. Poimanov, S. Sahoo, J. Sinha, H. Yang, M. Münzenberg, T. Moriyama, S. Mizukami, P. Landeros, R. A. Gallardo, G. Carlotti, J.-V. Kim, R. L. Stamps, R. E. Camley, B. Rana, Y. Otani, W. Yu, T. Yu, G. E. W. Bauer, C. Back, G. S. Uhrig, O. V. Dobrovolskiy, B. Budinska, H. Qin, S. van Dijken, A. V. Chumak, A. Khitun, D. E. Nikonov, I. A. Young, B. W. Zingsem, and M. Winklhofer, “The 2021 magnonics roadmap,” *Journal of Physics: Condensed Matter* **33**, 413001 (2021).
- ⁴ A. V. Chumak, V. I. Vasyuchka, A. A. Serga, and B. Hillebrands, “Magnon spintronics,” *Nature Physics* **11**, 453–461 (2015).
- ⁵ Y. Kajiwara, K. Harii, S. Takahashi, J. Ohe, K. Uchida, M. Mizuguchi, H. Umezawa, H. Kawai, K. Ando, K. Takanashi, S. Maekawa, and E. Saitoh, “Transmission of electrical signals by spin-wave interconversion in a magnetic insulator,” *Nature* **464**, 262–266 (2010).
- ⁶ Akashdeep Kamra and Wolfgang Belzig, “Magnon-mediated spin current noise in ferromagnet | nonmagnetic conductor hybrids,” *Phys. Rev. B* **94**, 014419 (2016).
- ⁷ Mathias Weiler, Matthias Althammer, Michael Schreier, Johannes Lotze, Matthias Pernpeintner, Sibylle Meyer, Hans Huebl, Rudolf Gross, Akashdeep Kamra, Jiang Xiao, Yan-Ting Chen, HuJun Jiao, Gerrit E. W. Bauer, and Sebastian T. B. Goennenwein, “Experimental test of the spin mixing interface conductivity concept,” *Phys. Rev. Lett.* **111**, 176601 (2013).
- ⁸ T. Kato, Y. Ohnuma, M. Matsuo, J. Rech, T. Jonckheere, and T. Martin, “Microscopic theory of spin transport at the interface between a superconductor and a ferromagnetic insulator,” *Phys. Rev. B* **99**, 144411 (2019).
- ⁹ Morten Amundsen, Irina V. Bobkova, and Akashdeep Kamra, “Magnonic spin joule heating and rectification effects,” *Phys. Rev. B* **106**, 144411 (2022).
- ¹⁰ Yaroslav Tserkovnyak, Arne Brataas, Gerrit E. W. Bauer, and Bertrand I. Halperin, “Nonlocal magnetization dynamics in ferromagnetic heterostructures,” *Rev. Mod. Phys.* **77**, 1375–1421 (2005).
- ¹¹ Yuichi Ohnuma, Hiroto Adachi, Eiji Saitoh, and Sadamichi Maekawa, “Enhanced dc spin pumping into a fluctuating ferromagnet near T_C ,” *Phys. Rev. B* **89**, 174417 (2014).
- ¹² C. Bell, S. Milikisyants, M. Huber, and J. Aarts, “Spin dynamics in a superconductor-ferromagnet proximity system,” *Phys. Rev. Lett.* **100**, 047002 (2008).
- ¹³ Kun-Rok Jeon, Chiara Ciccarelli, Hidekazu Kurebayashi, Lesley F. Cohen, Sachio Komori, Jason W. A. Robinson, and Mark G. Blamire, “Abrikosov vortex nucleation and its detrimental effect on superconducting spin pumping in pt/nb/ni₈₀fe₂₀/Nb/Pt proximity structures,” *Phys. Rev. B* **99**, 144503 (2019).
- ¹⁴ Kun-Rok Jeon, Chiara Ciccarelli, Hidekazu Kurebayashi, Lesley F. Cohen, Xavier Montiel, Matthias Eschrig, Sachio Komori, Jason W. A. Robinson, and Mark G. Blamire, “Exchange-field enhancement of superconducting spin pumping,” *Phys. Rev. B* **99**, 024507 (2019).
- ¹⁵ Kun-Rok Jeon, Chiara Ciccarelli, Hidekazu Kurebayashi, Lesley F. Cohen, Xavier Montiel, Matthias Eschrig, Thomas Wagner, Sachio Komori, Anand Srivastava, Jason W.A. Robinson, and Mark G. Blamire, “Effect of meissner screening and trapped magnetic flux on magnetization dynamics in thick Nb/ni₈₀fe₂₀/Nb trilayers,” *Phys. Rev. Applied* **11**, 014061 (2019).
- ¹⁶ Kun-Rok Jeon, Chiara Ciccarelli, Andrew J. Ferguson, Hidekazu Kurebayashi, Lesley F. Cohen, Xavier Montiel, Matthias Eschrig, Jason W. A. Robinson, and Mark G. Blamire, “Enhanced spin pumping into superconductors provides evidence for superconducting pure spin currents,” *Nature Materials* **17**, 499–503 (2018).
- ¹⁷ Yunyan Yao, Qi Song, Yota Takamura, Juan Pedro Cascales, Wei Yuan, Yang Ma, Yu Yun, X. C. Xie, Jagadeesh S. Moodera, and Wei Han, “Probe of spin dynamics in superconducting nbn thin films via spin pumping,” *Phys. Rev. B* **97**, 224414 (2018).
- ¹⁸ L. L. Li, Y. L. Zhao, X. X. Zhang, and Y. Sun, “Possible evidence for spin-transfer torque induced by spin-triplet supercurrents,” *Chin. Phys. Lett.* **35**, 077401 (2018).
- ¹⁹ I.A. Golovchanskiy, N.N. Abramov, V.S. Stolyarov, V.I. Chichkov, M. Silaev, I.V. Shchetinin, A.A. Golubov, V.V. Ryazanov, A.V. Ustinov, and M.Yu. Kupriyanov, “Magnetization dynamics in proximity-coupled superconductor-ferromagnet-superconductor multilayers,” *Phys. Rev. Applied* **14**, 024086 (2020).
- ²⁰ M. A. Silaev, “Finite-frequency spin susceptibility and spin pumping in superconductors with spin-orbit relaxation,” *Phys. Rev. B* **102**, 144521 (2020).
- ²¹ M.A. Silaev, “Anderson-higgs mass of magnons in superconductor/ferromagnet/superconductor systems,” (2022), arXiv:2207.13201 [cond-mat.supr-con].
- ²² S. V. Mironov and A. I. Buzdin, “Giant demagnetization effects induced by superconducting films,” *Applied Physics Letters* **119**, 102601 (2021).
- ²³ O. V. Dobrovolskiy, R. Sachser, T. Brächer, T. Böttcher, V. V. Kruglyak, R. V. Vovk, V. A. Shklovskij, M. Huth, B. Hillebrands, and A. V. Chumak, “Magnon-fluxon interaction in a ferromagnet/superconductor heterostructure,” *Nature Physics* **15**, 477–482 (2019).

- ²⁴ Tao Yu and Gerrit E. W. Bauer, “Efficient gating of magnons by proximity superconductors,” *Phys. Rev. Lett.* **129**, 117201 (2022).
- ²⁵ Irina V. Bobkova, Alexander M. Bobkov, Akashdeep Kamra, and Wolfgang Belzig, “Magnon-cooperons in magnet-superconductor hybrids,” *Communications Materials* **3**, 95 (2022).
- ²⁶ Lina G. Johnsen, Haakon T. Simensen, Arne Brataas, and Jacob Linder, “Magnon spin current induced by triplet cooper pair supercurrents,” *Phys. Rev. Lett.* **127**, 207001 (2021).
- ²⁷ V. Baltz, A. Manchon, M. Tsoi, T. Moriyama, T. Ono, and Y. Tserkovnyak, “Antiferromagnetic spintronics,” *Rev. Mod. Phys.* **90**, 015005 (2018).
- ²⁸ T. Jungwirth, X. Marti, P. Wadley, and J. Wunderlich, “Antiferromagnetic spintronics,” *Nature Nanotechnology* **11**, 231–241 (2016).
- ²⁹ Arne Brataas, Bart van Wees, Olivier Klein, Grégoire de Loubens, and Michel Viret, “Spin insulatronics,” *Physics Reports* **885**, 1–27 (2020).
- ³⁰ T. Wimmer, A. Kamra, J. Gückelhorn, M. Opel, S. Geprägs, R. Gross, H. Huebl, and M. Althammer, “Observation of antiferromagnetic magnon pseudospin dynamics and the Hanle effect,” *Phys. Rev. Lett.* **125**, 247204 (2020).
- ³¹ Akashdeep Kamra and Wolfgang Belzig, “Spin pumping and shot noise in ferrimagnets: Bridging ferro- and antiferromagnets,” *Phys. Rev. Lett.* **119**, 197201 (2017).
- ³² Ran Cheng, Jiang Xiao, Qian Niu, and Arne Brataas, “Spin pumping and spin-transfer torques in antiferromagnets,” *Phys. Rev. Lett.* **113**, 057601 (2014).
- ³³ Eirik Erlandsen and Asle Sudbø, “Magnon drag in a metal-insulating antiferromagnet bilayer,” *Phys. Rev. B* **105**, 184434 (2022).
- ³⁴ Eirik Erlandsen, Akashdeep Kamra, Arne Brataas, and Asle Sudbø, “Enhancement of superconductivity mediated by antiferromagnetic squeezed magnons,” *Phys. Rev. B* **100**, 100503 (2019).
- ³⁵ Eirik Erlandsen, Arne Brataas, and Asle Sudbø, “Magnon-mediated superconductivity on the surface of a topological insulator,” *Phys. Rev. B* **101**, 094503 (2020).
- ³⁶ Eirik Erlandsen and Asle Sudbø, “Schwinger boson study of superconductivity mediated by antiferromagnetic spin fluctuations,” *Phys. Rev. B* **102**, 214502 (2020).
- ³⁷ Even Thingstad, Eirik Erlandsen, and Asle Sudbø, “Eliashberg study of superconductivity induced by interfacial coupling to antiferromagnets,” *Phys. Rev. B* **104**, 014508 (2021).
- ³⁸ Aleksandr I. Buzdin and L. N. Bulaevskii, “Antiferromagnetic superconductors,” *Soviet Physics Uspekhi* **29**, 412 (1986).
- ³⁹ Akashdeep Kamra, Ali Rezaei, and Wolfgang Belzig, “Spin splitting induced in a superconductor by an antiferromagnetic insulator,” *Phys. Rev. Lett.* **121**, 247702 (2018).
- ⁴⁰ A. I. Buzdin, “Proximity effects in superconductor-ferromagnet heterostructures,” *Rev. Mod. Phys.* **77**, 935–976 (2005).
- ⁴¹ F. Sebastian Bergeret, Mikhail Silaev, Pauli Virtanen, and Tero T. Heikkilä, “Colloquium: Nonequilibrium effects in superconductors with a spin-splitting field,” *Rev. Mod. Phys.* **90**, 041001 (2018).
- ⁴² G. A. Bobkov, I. V. Bobkova, A. M. Bobkov, and Akashdeep Kamra, “Thermally induced spin torque and domain-wall motion in superconductor/antiferromagnetic-insulator bilayers,” *Phys. Rev. B* **103**, 094506 (2021).
- ⁴³ Audrey Cottet, Daniel Huertas-Hernando, Wolfgang Belzig, and Yuli V. Nazarov, “Spin-dependent boundary conditions for isotropic superconducting green’s functions,” *Phys. Rev. B* **80**, 184511 (2009).
- ⁴⁴ M. Eschrig, A. Cottet, W. Belzig, and J. Linder, “General boundary conditions for quasiclassical theory of superconductivity in the diffusive limit: application to strongly spin-polarized systems,” *New Journal of Physics* **17**, 083037 (2015).
- ⁴⁵ Sergio M. Rezende, Antonio Azevedo, and Roberto L. Rodríguez-Suárez, “Introduction to antiferromagnetic magnons,” *Journal of Applied Physics* **126**, 151101 (2019).
- ⁴⁶ Akashdeep Kamra, Utkarsh Agrawal, and Wolfgang Belzig, “Noninteger-spin magnonic excitations in untextured magnets,” *Phys. Rev. B* **96**, 020411 (2017).
- ⁴⁷ Akashdeep Kamra, Roberto E. Troncoso, Wolfgang Belzig, and Arne Brataas, “Gilbert damping phenomenology for two-sublattice magnets,” *Phys. Rev. B* **98**, 184402 (2018).
- ⁴⁸ H. Y. Yuan, Qian Liu, Ke Xia, Zhe Yuan, and X. R. Wang, “Proper dissipative torques in antiferromagnetic dynamics,” *Europhysics Letters* **126**, 67006 (2019).
- ⁴⁹ Maurice Griffel and J. W. Stout, “Preparation of single crystals of manganous fluoride. the crystal structure from x-ray diffraction. the melting point and density,” *Journal of the American Chemical Society* **72**, 4351–4353 (1950).