

Pair correlation function of the one-dimensional Riesz gas

C. W. J. Beenakker

Instituut-Lorentz, Universiteit Leiden, P.O. Box 9506, 2300 RA Leiden, The Netherlands

(Dated: December 2022)

A method from random-matrix theory is used to calculate the pair correlation function of a one-dimensional gas of $N \gg 1$ classical particles with a power law repulsive interaction potential $u(x) \propto |x|^{-s}$ (a so-called Riesz gas). An integral formula for the covariance of single-particle operators is obtained which generalizes known results in the limits $s \rightarrow -1$ (Coulomb gas) and $s \rightarrow 0$ (log-gas). As an application, we calculate the variance of the center of mass of the Riesz gas, which has a universal large- N limit that does not depend on the shape of the confining potential.

I. INTRODUCTION

The one-dimensional (1D) Riesz gas [1–4], describes N classical particles that move on a line (the x -axis) with a repulsive interaction potential $u(x)$ of the form

$$u(x) = \begin{cases} \text{sign}(s)|x|^{-s} & \text{for } s > -2, \\ -\ln|x| & \text{for } s = 0. \end{cases} \quad (1.1)$$

The particles are prevented from moving off to infinity by a confining potential $V(x)$.

The cases $s = 0$ and $s = -1$ are also referred to as log-gas and Coulomb gas, respectively. The log-gas plays a central role in random-matrix theory (RMT) [5]. Experimentally, an interaction potential with an adjustable exponent $s \lesssim 1$ has been realized in a chain of trapped ion spins [6].

Thermal averages $\langle \dots \rangle$, at inverse temperature β , are defined with respect to the Gibbs measure

$$P(x_1, x_2, \dots, x_N) = Z^{-1} \times \exp\left(-\beta \left[J \sum_{i < j=1}^N u(x_i - x_j) + \sum_{i=1}^N V(x_i) \right]\right), \quad (1.2)$$

where Z normalizes the distribution to unity. The interaction strength is parameterized by $J > 0$. The average density is

$$\rho(x) = \left\langle \sum_{i=1}^N \delta(x - x_i) \right\rangle, \quad (1.3)$$

normalized to the particle number,

$$\int_{-\infty}^{\infty} \rho(x) dx = N. \quad (1.4)$$

Much is known about the dependence of $\rho(x)$ on the range of the interaction [7–10], in the three regimes $s > 1$ (short-range repulsion), $-1 < s < 1$ (weakly long-range repulsion), and $-2 < s < -1$ (strongly long-range repulsion). In what follows we consider the second regime, $-1 < s < 1$, which includes the log-gas at $s = 0$ and the Coulomb gas in the limit $s \rightarrow -1$.

For $N \gg 1$ the density has a compact support, which may consist of multiple disjoint intervals. Considering a

single interval (a, b) , an end point may be N -independent, fixed by a hard wall, or it may be N -dependent, freely adjustable in a smooth potential. The density vanishes as a powerlaw at a free end point, while it diverges (with an integrable singularity) at a fixed end point.

For example, in the case of a quadratic confinement $V(x) \propto x^2$, there are two free N -dependent end points at $b = -a \propto N^{1/(2+s)}$ and the density profile is [9]

$$\rho(x) \propto (b^2 - x^2)^{(s+1)/2}, \quad |s| < 1. \quad (1.5)$$

The density profile becomes flat in the Coulomb gas limit, while the log-gas has the Wigner semicircle law [11]. Alternatively, if we set $V \equiv 0$ and confine the Riesz gas by a hard wall at $x = a$ and $x = b$, then the density diverges on approaching a fixed end point [10],

$$\rho(x) \propto (x - a)^{(s-1)/2} (b - x)^{(s-1)/2}, \quad |s| < 1. \quad (1.6)$$

Knowledge of the density allows us to calculate by integration the average $\langle F \rangle$ of a single-particle observable $F = \sum_{i=1}^N f(x_i)$ (also known as a “linear statistic”). For $N \gg 1$ we can restrict the integral to the interval (a, b) ,

$$\langle F \rangle = \int_a^b \rho(x) f(x) dx. \quad (1.7)$$

Going beyond the first moment, Flack, Majumdar, and Schehr recently obtained [12] a strikingly simple formula for the large- N limit of the variance of F in the Coulomb case [13],

$$\text{Var } F = \frac{1}{2\beta J} \int_a^b [df(x)/dx]^2 dx \quad \text{for } s = -1. \quad (1.8)$$

The corresponding formula in the RMT case $s = 0$ is known [14, 15], but results for other values of the interaction parameter s are not known. The aim of this paper is to provide that information for the entire range $-1 < s < 1$.

For that purpose one needs the connected pair correlation function

$$R(x, y) = \left\langle \sum_{i,j=1}^N \delta(x - x_i) \delta(y - x_j) \right\rangle - \rho(x)\rho(y), \quad (1.9)$$

which gives the variance upon integration,

$$\text{Var } F = \int_a^b dx \int_a^b dy R(x, y) f(x) f(y). \quad (1.10)$$

In RMT there are basically two methods to compute $R(x, y)$. The method of orthogonal polynomials [16] applies to specific confining potentials (typically linear or quadratic) and then gives results for any N . The alternative method of functional derivatives [17] (equivalently, the method of loop equations [18]) takes the large- N limit, but then works generically for any form of confinement. Since the latter method does not assume a logarithmic repulsion, it is the method of choice in what follows.

II. PAIR CORRELATION FUNCTION

For $N \gg 1$ the pair correlation function oscillates rapidly on the scale of the interparticle spacing $\delta x \simeq (b-a)/N$. These oscillations are irrelevant for the computation of the variance of an observable that varies smoothly on the scale of δx , so that in the large- N limit it is sufficient to know the smoothed correlation function.

The method of functional derivatives starts from the exact representation

$$R(x, y) = -\frac{1}{\beta} \frac{\delta \rho(x)}{\delta V(y)}. \quad (2.1)$$

The variation of the density is to be carried out at con-

stant particle number,

$$\int_a^b \delta \rho(x) dx = 0. \quad (2.2)$$

The integration interval (a, b) is the support of the smoothed particle density $\rho(x)$ in the large- N limit. (We assume that the confining potential produces a support in a single interval.)

In the regime $-1 < s < 1$ of a weakly long-ranged repulsion, and for $J \gtrsim 1/\beta$, variations in the smoothed density ρ and in the confining potential V are related by the condition of mechanical equilibrium [9, 19],

$$J \int_a^b u(x-y) \delta \rho(y) dy + \delta V(x) = \text{constant}, \quad a < x < b. \quad (2.3)$$

Taking the derivative with respect to x we have a singular integral equation,

$$J(|s| + \delta_{0,s}) \mathcal{P} \int_a^b dy \delta \rho(y) \frac{\text{sign}(x-y)}{|x-y|^{s+1}} = \frac{d}{dx} \delta V(x), \quad a < x < b, \quad (2.4)$$

which we need to invert in order to obtain the functional derivative (2.1). (The symbol \mathcal{P} indicates the principal value of the integral.)

For $-1 < s < 1$ the general solution to Eq. (2.4) is given by the Sonin inversion formula [20–22] (see App. A),

$$J(|s| + \delta_{0,s}) \delta \rho(x) = C_{\delta V} [(x-a)(b-x)]^{s-} - C_1 \mathcal{S}_{\delta V}(x), \quad (2.5a)$$

$$\mathcal{S}_{\delta V}(x) = -(x-a)^{s+} \frac{d}{dx} \int_x^b dt \frac{(t-x)^{s-}}{(t-a)^s} \frac{d}{dt} \int_a^t dy (y-a)^{s-} (t-y)^{s+} \frac{d}{dy} \delta V(y), \quad (2.5b)$$

$$C_1 = \frac{\sin(\pi s_+) \Gamma(s+1)}{\pi s_+ \Gamma(s_+)^2}, \quad s_{\pm} = (s \pm 1)/2. \quad (2.5c)$$

The coefficient $C_{\delta V}$ is fixed by the constraint (2.2),

$$C_{\delta V} = \frac{C_1}{C_2} \int_a^b dx \mathcal{S}_{\delta V}(x), \quad (2.6a)$$

$$C_2 = \int_a^b dx [(x-a)(b-x)]^{s-} = \frac{(b-a)^s \sqrt{\pi} \Gamma(s_+)}{2^s \Gamma(1+s/2)}. \quad (2.6b)$$

The function $\Gamma(x)$ is the usual gamma function.

The pair correlation function $R(x, y)$ is obtained from Eq. (2.1) as a distribution, defined by its action on a test function $g(y)$ upon integration over y . Using the

functional-derivative identities

$$\int_a^b dy \frac{\delta \mathcal{S}_{\delta V}(x)}{\delta V(y)} g(y) = \mathcal{S}_g(x), \quad (2.7a)$$

$$\int_a^b dy \frac{\delta C_{\delta V}}{\delta V(y)} g(y) = \frac{C_1}{C_2} \int_a^b dx \mathcal{S}_g(x), \quad (2.7b)$$

we find the following expression:

$$J(|s| + \delta_{0,s}) \int_a^b dy R(x, y) g(y) = C_1 \mathcal{S}_g(x) - [(x-a)(b-x)]^{s-} \frac{C_1}{C_2} \left(\int_a^b dx \mathcal{S}_g(x) \right). \quad (2.8)$$

III. COVARIANCE OF SINGLE-PARTICLE OBSERVABLES

Eq. (2.8) provides a formula for the covariance of the two observables $F = \sum_{i=1}^N f(x_i)$ and $G = \sum_{i=1}^N g(x_i)$,

$$\begin{aligned} \text{CoVar}(F, G) &= \langle FG \rangle - \langle F \rangle \langle G \rangle \\ &= \int_a^b dx \int_a^b dy R(x, y) f(x) g(y). \end{aligned} \quad (3.1)$$

The covariance is given by integrals over $f(x)$ and $\mathcal{S}_g(x)$,

$$\begin{aligned} J\beta(|s| + \delta_{0,s}) \text{CoVar}(F, G) &= C_1 \int_a^b dx f(x) \mathcal{S}_g(x) \\ &- \frac{C_1}{C_2} \left(\int_a^b dx \mathcal{S}_g(x) \right) \int_a^b dx [(x-a)(b-x)]^{s-} f(x). \end{aligned} \quad (3.2)$$

The variance of a single observable then follows from $\text{Var} F = \text{CoVar}(F, F)$.

For general functions f, g the integrals in Eq. (3.2) may be carried out numerically (see App. A). Closed-form expressions can be obtained for polynomial functions. It is convenient to shift the origin of the coordinate system so that $(a, b) \rightarrow (0, L)$. We define $X_p = \sum_i x_i^p$, $p \geq 1$, and obtain the covariance

$$\begin{aligned} J\beta(|s| + \delta_{0,s}) \text{CoVar}(X_p, X_q) &= \frac{2\pi L^{p+q+s} \Gamma(s)}{\Gamma(-p-s_-) \Gamma(p+s+1) \Gamma(-q-s_-) \Gamma(q+s+1)} \\ &\times \frac{pqs \sin(\pi s_+)}{(p+q+s) (\cos[\pi(p+q+s)] + \cos[\pi(p-q)])}. \end{aligned} \quad (3.3)$$

For the variance this reduces to

$$J\beta(|s| + \delta_{0,s}) \text{Var} X_p = \frac{2\pi L^{2p+s} p^2 s \Gamma(s) \sin(\pi s_+)}{(2p+s) \Gamma(-p-s_-)^2 \Gamma(p+s+1)^2 (1 + \cos[\pi(2p+s)])}. \quad (3.4)$$

We have checked that the general formula (3.3) agrees with the known formulas in the Coulomb gas limit [12] (see also App. B),

$$\begin{aligned} J\beta \text{CoVar}(F, G) &= \frac{1}{2} \int_0^L \frac{df(x)}{dx} \frac{dg(x)}{dx} dx, \quad \text{for } s = -1, \\ \Rightarrow J\beta \text{CoVar}(X_p, X_q) &= \frac{pqL^{p+q-1}}{2(p+q-1)}, \end{aligned} \quad (3.5)$$

and in the log-gas limit [23, 24],

$$\begin{aligned} J\beta \text{CoVar}(F, G) &= \frac{1}{\pi^2} \mathcal{P} \int_0^L dx \int_0^L dy \frac{g(y) df(x)/dx}{y-x} \sqrt{\frac{x(L-x)}{y(L-y)}}, \quad \text{for } s = 0, \\ \Rightarrow J\beta \text{CoVar}(X_p, X_q) &= \frac{\pi L^{p+q}}{(p+q) \Gamma(\frac{1}{2}-p) \Gamma(p) \Gamma(\frac{1}{2}-q) \Gamma(q) \cos(\pi p) \cos(\pi q)}. \end{aligned} \quad (3.6)$$

IV. VARIANCE OF THE CENTER OF MASS

By way of illustration, we compute the variance of the center of mass $M = N^{-1} \sum_{i=1}^N x_i$ of the Riesz gas. Eq.

(3.4) for $p = 1$ gives

$$\begin{aligned} \text{Var} M &= \frac{C_s \sqrt{\pi}}{2^{s+3} \Gamma(\frac{1}{2}-s/2) \Gamma(2+s/2)}, \\ C_s &= \frac{L^{s+2}}{N^2 J\beta |s| + \delta_{0,s}}. \end{aligned} \quad (4.1)$$




FIG. 1. Variance of the center of mass of the Riesz gas, computed from Eq. (4.1). The calculation applies to the interval $|s| < 1$, the dashed curve is the analytical continuation of Eq. (4.1) to smaller s .

The dimensionless coefficient C_s is N -independent if the system is scaled at constant interparticle spacing $\delta x = L/N$ with interaction strength $J \propto N^s$.

The dependence of $\text{Var } M$ on the interaction exponent s is plotted in Fig. 1. At the upper limit $s \rightarrow 1$ we find $C_s^{-1} \text{Var } M \rightarrow 0$, to leading order in $1/N$. This is consistent with the fact that the repulsion is short-range for $s > 1$, so we would expect the positions of the particles to fluctuate independently. The variance of the center of mass (rescaled by C_s) would then be of order $1/N$.

The theory applies to the interval $-1 < s < 1$ of a weakly long-range repulsion. The dashed curve in Fig. 1 is the analytical continuation of Eq. (4.1) to the interval $-2 < s < -1$ of a strongly long-range repulsion. We surmise that the formula remains valid in that regime.

V. CONCLUSION

In conclusion, we have computed the pair correlation function of a 1D system of classical particles with a long-range power law repulsion. In the thermodynamic limit (particle number N and system size L to infinity at fixed interparticle spacing $\delta x = L/N$), and upon smoothing over δx , the pair correlation function becomes a *universal* function of the power law exponent $s \in (-1, 1)$ —

$$S_-(x) = -C_1(x-a)^{s-} \frac{d}{dx} \int_x^b dt \frac{(t-x)^{s+}}{(t-a)^s} \frac{d}{dt} \int_a^t dy (y-a)^{s+} (t-y)^{s-} g'(y), \quad (\text{A3})$$

$$S_+(x) = C_1(x-a)^{s+} \frac{d}{dx} \int_x^b dt \frac{(t-x)^{s-}}{(t-a)^s} \frac{d}{dt} \int_a^t dy (y-a)^{s-} (t-y)^{s+} g'(y). \quad (\text{A4})$$

For the general solution we take either one of these two particular solutions and add it to an arbitrary multiple

independent of the shape of the confining potential $V(x)$ for a given single-interval support (a, b) of the average density. So it does not matter if the Riesz gas is confined to the interval $(0, L)$ by a soft parabolic potential or by a hard-wall confinement — the density fluctuations are the same even though the average density profile $\rho(x)$ is very different in the two cases [10].

Our result (3.2) for the covariance of a pair of single-particle observables generalizes old results for a logarithmic repulsion [14, 15] and a very recent result for a linear repulsion [12]. We rely on a solution of an integral equation that requires $|s| < 1$ (weakly long-range regime), but the method of functional derivatives that we have used can be applied also outside of this interval. The independence of the pair correlation function $R(x, y) = -\beta^{-1} \delta \rho(x) / \delta V(y)$ on the shape of the confining potential follows directly from the linearity of the relation between ρ and V , so we expect a universal result also for $-2 < s < -1$. In the short-range regime $s > 1$, in contrast, the ρ - V relation is nonlinear [9] and no universal answer is expected.

ACKNOWLEDGMENTS

The author receives funding from the European Research Council (Advanced Grant 832256).

Appendix A: Sonin inversion formula

We summarize results from Refs. 20–22 on the solution of the singular integral equation

$$\mathcal{P} \int_a^b dy S(y) \frac{\text{sign}(x-y)}{|x-y|^{s+1}} = g'(x), \quad x \in (a, b), \quad |s| < 1. \quad (\text{A1})$$

The homogenous integral equation, with zero on the right-hand-side, has the solution

$$S_0(x) = [(x-a)(b-x)]^{s-}. \quad (\text{A2})$$

The Sonin inversion formula gives two particular solutions to the inhomogeneous integral equation,

of the homogeneous solution,

$$S(x) = S_{\pm}(x) + \text{constant} \times S_0(x), \quad (\text{A5})$$

where “constant” means independent of x . Since, by construction, the difference of two particular solutions solves the homogeneous integral equation, it does not matter for the general solution which particular solution we choose.

In the main text we chose $S_+(x)$. This has the benefit over $S_-(x)$ that the derivatives can be eliminated upon partial integration,

$$\int_a^b dx f(x) S_+(x) = -C_1 \int_a^b dx [f'(x)(x-a) + s_+ f(x)] (x-a)^{s_-} \times \int_x^b dt \frac{(t-x)^{s_-}}{(t-a)^s} \int_a^t dy s_+(y-a)^{s_-} (t-y)^{s_-} g'(y). \quad (\text{A6})$$

The three definite integrals of Eq. (A6) are in a form that can be evaluated numerically, without the need to take derivatives.

Appendix B: Coulomb gas limit

The Coulomb gas limit $s = -1$ can be obtained from the general formulas for $|s| < 1$ by means of the identities [25]

$$\lim_{\epsilon \searrow 0} \int_x^b \frac{\epsilon f(y)}{(y-x)^{1-\epsilon}} dy = f(x), \quad (\text{B1a})$$

$$\lim_{\epsilon \searrow 0} \int_a^b \frac{\epsilon f(x)}{[(b-x)(x-a)]^{1-\epsilon}} dx = \frac{f(a) + f(b)}{b-a}. \quad (\text{B1b})$$

We start from the derivative-free representation (A6) of the integrals in Eq. (3.2), and apply Eq. (B1) first to the integral over y ,

$$\begin{aligned} I_1(t) &= \lim_{s \searrow -1} \int_a^t dy s_+(y-a)^{s_-} (t-y)^{s_-} g'(y) \\ &= \frac{g'(t) + g'(a)}{t-a}, \end{aligned} \quad (\text{B2})$$

then to the integral over t ,

$$\begin{aligned} I_2(x) &= \lim_{s \searrow -1} \int_x^b dt \frac{(t-x)^{s_-}}{(t-a)^s} I_1(t) \\ &= \frac{1}{s_+} [g'(x) + g'(a)], \end{aligned} \quad (\text{B3})$$

and finally to the integral over x ,

$$\begin{aligned} I_3 &= \lim_{s \searrow -1} \int_a^b dx [f'(x)(x-a) + s_+ f(x)] (x-a)^{s_-} I_2(x) \\ &= \frac{1}{s_+} \int_a^b dx f'(x) [g'(x) + g'(a)] + \frac{2}{s_+} f(a) g'(a). \end{aligned} \quad (\text{B4})$$

Moreover, since $1/C_2 \rightarrow \frac{1}{2} s_+(b-a)$ for $s \rightarrow -1$, we have

$$\lim_{s \searrow -1} \frac{1}{C_2} \int_a^b dx [(x-a)(b-x)]^{s_-} f(x) = \frac{1}{2} [f(b) + f(a)]. \quad (\text{B5})$$

Using also $C_1 = \frac{1}{2} s_+ + \mathcal{O}(s+1)^2$ we thus arrive at

$$\begin{aligned} \lim_{s \searrow -1} C_1 \int_a^b dx f(x) \mathcal{S}_g(x) &= \frac{1}{2} \int_a^b dx f'(x) [g'(x) + g'(a)] + f(a) g'(a), \end{aligned} \quad (\text{B6a})$$

$$\begin{aligned} \lim_{s \searrow -1} \frac{C_1}{C_2} \left(\int_a^b dx \mathcal{S}_g(x) \right) \int_a^b dx [(x-a)(b-x)]^{s_-} f(x) &= \frac{1}{2} g'(a) [f(b) + f(a)]. \end{aligned} \quad (\text{B6b})$$

Substitution of Eq. (B6) into Eq. (3.2) gives

$$\lim_{s \searrow -1} \text{CoVar}(F, G) = \frac{1}{2J\beta} \int_a^b dx f'(x) g'(x), \quad (\text{B7})$$

in accord with Ref. [12].

-
- [1] M. Riesz, *Intégrales de Riemann-Liouville et potentiels*, Acta Sci. Math. Szeged **9**, 1 (1938).
 [2] M. Riesz, *L'intégrale de Riemann-Liouville et le problème de Cauchy*, Acta Math. **81**, 1 (1949) doi: 10.1007/BF02395016.
 [3] S. Serfaty, *Microscopic description of Log*

- and Coulomb gases*, arXiv:1709.04089 doi: <https://doi.org/10.48550/arXiv.1709.04089>.
 [4] M. Levin, *Coulomb and Riesz gases: The known and the unknown*, J. Math. Phys. **63**, 061101 (2022) doi: 10.1063/5.0086835.
 [5] P. J. Forrester, *Log-Gases and Random Matrices* (Prince-

- ton University Press, Princeton, 2010).
- [6] J. Zhang, G. Pagano, P. W. Hess, A. Kyprianidis, P. Becker, H. Kaplan, A. V. Gorshkov, Z.-X. Gong, and C. Monroe, *Observation of a many-body dynamical phase transition with a 53-qubit quantum simulator*, Nature **551**, 601 (2017) doi:10.1038/nature24654.
- [7] T. Leblé and S. Serfaty, *Large deviation principle for empirical fields of Log and Riesz gases*, Invent. Math. **210**, 645 (2017) doi:10.1007/s00222-017-0738-0.
- [8] D. P. Hardin, T. Leblé, E. B. Saff, and S. Serfaty, *Large deviation principles for hypersingular Riesz gases*, Constr. Approx. **48**, 61 (2018) doi:10.1007/s00365-018-9431-9.
- [9] S. Agarwal, A. Dhar, M. Kulkarni, A. Kundu, S. N. Majumdar, D. Mukamel, and G. Schehr, *Harmonically confined particles with long-range repulsive interactions*, Phys. Rev. Lett. **123**, 100603 (2019) doi:10.1103/PhysRevLett.123.100603.
- [10] J. Kethepalli, M. Kulkarni, A. Kundu, S. N. Majumdar, D. Mukamel, and G. Schehr, *Harmonically confined long-ranged interacting gas in the presence of a hard wall*, J. Stat. Mech. 103209 (2021) doi:10.1088/1742-5468/ac2896.
- [11] E. P. Wigner, *On the statistical distribution of the widths and spacings of nuclear resonance levels*, Math. Proc. Cambridge Philos. Soc. **47**, 790 (1951) doi:10.1017/S0305004100027237.
- [12] A. Flack, S. N. Majumdar, and G. Schehr, *An exact formula for the variance of linear statistics in the one-dimensional jellium model*, arXiv:2211.11850 doi:10.48550/arXiv.2211.11850.
- [13] To compare with Ref. [12] substitute $\beta J \mapsto 2\alpha N$.
- [14] M. L. Mehta, *Random Matrices* (Elsevier, Amsterdam, 2004).
- [15] C. W. J. Beenakker, *Random-matrix theory of quantum transport*, Rev. Mod. Phys. **69**, 731 (1997) doi:10.1103/RevModPhys.69.731.
- [16] M. L. Mehta, *On the statistical properties of the level spacings in nuclear spectra*, Nucl. Phys. **18**, 395 (1960) doi:10.1016/0029-5582(60)90413-2.
- [17] C. W. J. Beenakker, *Universality in the random-matrix theory of quantum transport*, Phys. Rev. Lett. **70**, 1155 (1993) doi:10.1103/PhysRevLett.70.1155.
- [18] J. Ambjørn, J. Jurkiewicz, and Yu. M. Makeenko, *Multiloop correlators for two-dimensional quantum gravity*, Phys. Lett. B **251**, 517 (1990) doi:10.1016/0370-2693(90)90790-D.
- [19] The variational equation (2.3) does not contain a contribution $\propto \rho(b)\delta b$ from a variation of the end point b , for two reasons: Either $\delta b = 0$, for a fixed end point, or $\rho(b) = 0$, for a free end point. Similarly for the end point a .
- [20] N. Ya. Sonin, *Studies on Cylinder Functions and Special Polynomials* (Gostekhizdat, Moscow, 1954; in Russian).
- [21] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional integrals and derivatives: theory and applications* (Gordon and Breach, Philadelphia, 1993).
- [22] S. V. Buldyrev, S. Havlin, A. Ya. Kazakov, M. G. E. da Luz, E. P. Raposo, H. E. Stanley, and G. M. Viswanathan, *Average time spent by Lévy flights and walks on an interval with absorbing boundaries*, Phys. Rev. E **64**, 041108 (2001) doi:10.1103/PhysRevE.64.041108.
- [23] C. W. J. Beenakker, *Universality of Brézin and Zee's spectral correlator*, Nucl. Phys. B **422**, 515 (1994) doi:10.1016/0550-3213(94)2894-8.
- [24] F. D. Cunden and P. Vivo, *Universal covariance formula for linear statistics on random matrices*, Phys. Rev. Lett. **113**, 070202 (2014) doi:PhysRevLett.113.070202.
- [25] Eq. (B1) is a special case of Ramanujan's master theorem, for a proof see <https://mathoverflow.net/q/435723/11260>