

Hawking effect can generate physically inaccessible genuine tripartite nonlocality

Tinggui Zhang^{1,†} Xin Wang¹ and Shao-Ming Fei^{2,3,1}

¹ School of Mathematics and Statistics,
Hainan Normal University, Haikou, 571158, China

² School of Mathematical Sciences,
Capital Normal University, Beijing 100048, China

³ Max-Planck-Institute for Mathematics in the Sciences,
Leipzig 04103, Germany

[†] Correspondence to tinggui333@163.com

We explore the acceleration effect on the genuine tripartite nonlocality (GTN) for one or two accelerated detector(s) coupled to the vacuum field with initial mixed tripartite states. We show that the Hawking radiation degrades the physically accessible GTN, which suffers from “sudden death” at certain critical Hawking temperature. An novel phenomenon has been observed first time that the Hawking effect can generate the physically inaccessible GTN for fermion fields in curved spacetime, the “sudden birth” of the physically inaccessible GTN. This result shows that the GTN can pass through the event horizon of black hole for certain mixed initial states. We also derived analytically the tradeoff relations of genuine tripartite entanglement (GTE) and quantum coherence under the influence of Hawking effect.

PACS numbers: 04.70.Dy, 03.65.Ud, 04.62.+v

INTRODUCTION

Raised by Einstein et al. [1] in discussions on the incompleteness of quantum mechanics in 1935, quantum nonlocality [2] has been extensively studied and played significant roles in a variety of quantum tasks [3]. Recent studies have shown that quantum nonlocality is also a unique quantum resource for some device-independent quantum tasks such as the key distribution [4], random expansion [5], random amplification [6] and the related experiments [7–9]. In particular, in 1987 a special kind of nonlocality called genuine tripartite nonlocality (GTN) was introduced by Svetlichny for tripartite systems [10]. Such multipartite nonlocal correlations have not only foundational implications [11] but also novel applications in quantum communication and quantum computation [12–15], as well as in phase transitions and criticality in many-body systems [14]. GTN can be detected by the violation of Svetlichny inequality. Let $A_i = \vec{a}_i \cdot \vec{\sigma}$, $B_i = \vec{b}_i \cdot \vec{\sigma}$ and $C_i = \vec{c}_i \cdot \vec{\sigma}$, $i = 0, 1$, be the measurement observables associated with the first, second and third qubit, respectively, with \vec{a}_i , \vec{b}_i and \vec{c}_i the real unit vectors, $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ the vector given by the standard Pauli matrices. The Svetlichny operator is defined by

$$\begin{aligned} \mathbb{S} = & A_0(B_0 + B_1)C_0 + A_0(B_0 - B_1)C_1 \\ & + A_1(B_0 - B_1)C_0 - A_1(B_0 + B_1)C_1. \end{aligned} \quad (1)$$

If a state ρ admits bi-local hidden variable model, then the expectation value of the Svetlichny operator satisfies the following Svetlichny’s inequality,

$$\mathbb{S}(\rho) = |\text{Tr}(\mathbb{S}\rho)| \leq 4. \quad (2)$$

If the inequality (2) is violated, ρ must be a genuine three-qubit nonlocally correlated state.

In 1974, Hawking discovered black hole evaporation caused by thermal radiation emitted by the black hole [16]. Soon after, Unruh in 1976 proposed a very important prediction based on the effect of Hawking radiation, which says that a vacuum state observed by an observer who stays in flat Minkowski spacetime would be detected as a thermal state from another observer who hovers near the event horizon of the black hole with uniform acceleration [17, 18]. In recent years, as the intersection of classical informatics, quantum mechanics, special and general relativity and quantum field theory, the problem of quantum information in non-inertial system and curved spacetime has become a new hotspot in the research of quantum information science [19–45]. Undoubtedly, the study of quantum information under the framework of relativity, especially in the non-inertial frame and curved spacetime, not only is of great significance to the development and perfectness of quantum information theory, but also plays a positive role in understanding the entropy and information paradox in black holes. For examples, Fuentes et al. [19] show that the entanglement in noninertial frames is characterized by the observer-dependent property. Martín-Martínez et al. [22] studied the entanglement degradation affected by the Hawking effect of Schwarzschild black hole. Mann et al. [30] pointed out that incorporating concepts from quantum theory into relativity can yield novel and interesting effects. More recently, Wu et al. [43] studied the GTN and the genuine tripartite entanglement (GTE) of Dirac fields in the background of a Schwarzschild black hole. Li et al. [44] explored the tripartite entropic uncertainty and genuine tripartite quantum coherence of Dirac fields in the background of the Garfinkle-Horowitz-Strominger (GHS) dilatation space-time.

March attention has been paid to the multipar-

tite Greenberger-Horne-Zeilinger (GHZ) state $|GHZ\rangle = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle)$ or the generalized GHZ states under the Hawking radiation. It is recently found in Refs. [43] and [44] that the GTN is not redistributable, but GTE can be redistributed through Hawking effect. With the growth of the Hawking temperature, the physically accessible GTN decreases, but physically inaccessible GTN is not generated. The GTE however behaves differently: with the loss of the physically accessible GTE, the physically inaccessible GTE is generated continually through Hawking effect. This is inconsistent with intuition since quantum entanglement and quantum nonlocality are in consistent for pure states [46–48]. To clarify such problems, in this article we study the GTN of multipartite mixed states for Dirac fields in the background of a Schwarzschild black hole. The paper is organized as follows. In the second section, we briefly recall the quantization of Dirac fields in the background of Schwarzschild black hole, and the GTN, GTE and quantum coherence for three-qubit X-states. In the third section, we calculate and analyze the GTN of a detailed mixed state after one or bipartite acceleration. A novel phenomenon that is basically different from that of pure states is illustrated: the Hawking effect may not only degrades the physically accessible GTN, but also enhances the physically inaccessible GTN. The “sudden death” of physically accessible GTN takes place at some critical Hawking temperature, while “sudden birth” happens to the physically inaccessible GTN. We show that such phenomenon may be not observed for other mixed states. In the fourth section, we consider the quantum correlation GTE and quantum coherence. Analytical tradeoff relations among the distributions of GTE and coherence are derived. The last section is the conclusions and discussions.

QUANTIZATION OF DIRAC FIELDS IN A SCHWARZSCHILD BLACK HOLE

We first recall the evolution process of vacuum state of Dirac fields in a Schwarzschild black hole [43, 44]. The Dirac equation under a general background spacetime can be written as [49],

$$[\gamma^a e_a^\mu (\partial_\mu + \Gamma_\mu)]\Phi = 0, \quad (3)$$

where γ^a are the Dirac matrices, e_a^μ is the inverse of the tetrad e_μ^a , $\Gamma_\mu = \frac{1}{8}[\gamma^a, \gamma^b]e_a^\nu e_{b\nu;\mu}$ are the spin connection coefficients. The metric of the Schwarzschild black hole can be written as [24, 50]

$$ds^2 = - \left(\frac{r-2M}{r-2D}\right)dt^2 + \left(\frac{r-2M}{r-2D}\right)^{-1}dr^2 + r(r-2D)(d\theta^2 + \sin^2\theta d\phi^2), \quad (4)$$

where M denotes the mass of the black hole, and D denotes the parameter of the dilation field. The Hawking

temperature is given by $T = \frac{1}{8\pi(M-D)}$. The thermal Fermi-Dirac distribution of particles with Hawking temperature T was observed in Refs. [51, 52].

Solving the Dirac equation (3) near the event horizon of black hole, one gets a set of positive frequency outgoing solutions for the inside and outside regions of the event horizon, [22, 23]

$$\phi_k^{I+} = \xi e^{-i\omega\mu}, \quad \phi_k^{II+} = \xi e^{i\omega\mu}, \quad (5)$$

where k represents the field mode, ξ denotes four-component Dirac spinor, ω is a monochromatic frequency and $\mu = t - r_*$, where the tortoise coordinate $r_* = r + 2M \ln \frac{r-2M}{2M}$. ϕ_k^{I+} and ϕ_k^{II+} are usually called Schwarzschild modes. Particles and antiparticles are classified according to future-directed timelike Killing vector under each region.

An analytic continuation of (5) [44] gives rise to a complete basis of positive energy modes, i.e., the Kruskal modes [52]. Then one can use the Schwarzschild mode and the Kruskal mode to expand the Dirac field, respectively, leading to the Bogoliubov transformations between annihilation operator and creation operator under the Schwarzschild and Kruskal coordinates [42]. After properly normalizing the state vector, the vacuum state and excited state of the Kruskal particle in the single-mode approximation are given by

$$|0\rangle = \cos\beta |0\rangle_I^+ |0\rangle_{II}^- + \sin\beta |1\rangle_I^+ |1\rangle_{II}^-, \quad (6)$$

$$|1\rangle = |1\rangle_I^+ |0\rangle_{II}^-, \quad (7)$$

where $|n\rangle_I$ and $|n\rangle_{II}$ are the number states for the particle outside the region (physically accessible) and the antiparticle inside the region of the event horizon (physically inaccessible), respectively, the superscripts + and – indicate the particle and antiparticle, respectively, $\cos\beta = (e^{-\frac{\omega}{T}})^{-\frac{1}{2}}$ and $\sin\beta = (e^{\frac{\omega}{T}})^{-\frac{1}{2}}$.

In the following we will consider three-qubit X-state give by density matrix,

$$\rho_X = \begin{pmatrix} d_1 & 0 & 0 & 0 & 0 & 0 & 0 & f_1 \\ 0 & d_2 & 0 & 0 & 0 & 0 & f_2 & 0 \\ 0 & 0 & d_3 & 0 & 0 & f_3 & 0 & 0 \\ 0 & 0 & 0 & d_4 & f_4 & 0 & 0 & 0 \\ 0 & 0 & 0 & f_4^* & e_4 & 0 & 0 & 0 \\ 0 & 0 & f_3^* & 0 & 0 & e_3 & 0 & 0 \\ 0 & f_2^* & 0 & 0 & 0 & 0 & e_2 & 0 \\ f_1^* & 0 & 0 & 0 & 0 & 0 & 0 & e_1 \end{pmatrix}$$

in the orthogonal basis $\{|000\rangle, |001\rangle, \dots, |111\rangle\}$. The GTN (Svetlichny value) of ρ_X is simply given by [53]

$$S(\rho_X) = \max\{8\sqrt{2}|f_i|, 4|N|\}, \quad (8)$$

where $N = d_1 - d_2 - d_3 + d_4 - e_4 + e_3 + e_2 - e_1$. The GTE of ρ_X is given by [54]

$$E(\rho_X) = 2 \max\{0, |f_i| - m_i\}, \quad i = 1, 2, 3, 4, \quad (9)$$

where $m_i = \sum_{j \neq i}^4 \sqrt{d_j e_j}$. The quantum coherence of ρ_X is given by [55]

$$C(\rho_X) = C_{l_1}(\rho_X) = \sum_{i \neq j} |\rho_{ij}| = 2 \sum_{i=1}^4 |f_i|. \quad (10)$$

EVOLUTION OF GTN FOR MIXED STATES IN SCHWARZSCHILD BLACK HOLE

We first consider the initial tripartite mixed state $\rho_1 = p|GHZ\rangle\langle GHZ| + (1-p)|000\rangle\langle 000|$ of the dirac fields shared by Alice, Bob and Charlie in the asymptotically flat region, where $p \in [0, 1]$.

Case(1): Alice and Bob stay stationarily, while Charlie moves with a uniform acceleration in his vacuum cavity. Applying the transformation of Eqs. (6) and (7) for detector C, we obtain a 4-partite state $\rho_{ABC_I C_{II}}$ associated with subsystems A, B and C_I observed by Alice, Bob and Charlie outside the event horizon of black hole, respectively, and the subsystem C_{II} observed by anti-Charlie inside the event horizon. Since the interior region of black hole is causally disconnected from the exterior region, Alice, Bob and Charlie cannot access the modes inside the event horizon. The mode C_I outside the event horizon is called the accessible modes, and the mode C_{II} inside the event horizon the inaccessible modes. By tracing over the inaccessible mode on $\rho_{ABC_I C_{II}}$, we obtain the reduced density matrix,

$$\rho_{ABC_I} = \begin{pmatrix} \left(\frac{2-p}{2}\right) \cos^2 r_c & 0 & 0 & 0 & 0 & 0 & \frac{p}{2} \cos r_c \\ 0 & \left(\frac{2-p}{2}\right) \sin^2 r_c & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \frac{p}{2} \cos r_c & 0 & 0 & 0 & 0 & 0 & \frac{p}{2} \end{pmatrix},$$

where $\cos r_c = (e^{-\frac{\omega}{T_c}})^{-\frac{1}{2}}$ and $\sin r_c = (e^{\frac{\omega}{T_c}})^{-\frac{1}{2}}$ are functions of the Hawking temperature T_c . For simplicity, we set $\omega = 1$ and instead of T_c , we use r_c in the following analysis.

From Eq. (8), we obtain the GTN of the state ρ_{ABC_I} ,

$$S(\rho_{ABC_I}) = \max\{8\sqrt{2}|\frac{p}{2} \cos r_c|, 4|(1 - \frac{p}{2}) \cos 2r_c - \frac{p}{2}|\}.$$

In FIG. 1 (a) we plot the Svetlichny value $S(\rho_{ABC_I})$ as function of r_c (the Hawking temperature T) and p . We find that, with the increase of Hawking temperature T and the decreasing of parameter p from 1 to 0, $S(\rho_{ABC_I})$ is larger than 4 at first and then smaller than 4. This implies two interesting aspects. One is that the thermal noise introduced by Hawking temperature destroys the




FIG. 1: Case(1): Alice and Bob stay stationarily, while Charlie moves with a uniform acceleration in his vacuum cavity. (a) the Svetlichny value $S(\rho_{ABC_I})$ as function of r_c (the Hawking temperature T) and p . (b) the Svetlichny value $S(\rho_{ABC_{II}})$ as function of r_c and p .

physically accessible GTN between Alice, Bob and Charlie. And the “sudden death” takes place at a critical temperature T_c . Another phenomenon is that with the decrease of the parameter p , the GTN transition from the maximally entangled initial state to the complete separable one is also influenced by Hawking effect.

Similarly, by tracing over the accessible mode C_I in $\rho_{ABC_I C_{II}}$ we get the reduced density matrix,

$$\rho_{ABC_{II}} = \begin{pmatrix} \left(\frac{2-p}{2}\right) \cos^2 r_c & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \left(\frac{2-p}{2}\right) \sin^2 r_c & 0 & 0 & 0 & \frac{p}{2} \sin r_c & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{p}{2} \sin r_c & 0 & 0 & 0 & \frac{p}{2} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

and the GTN of the state $\rho_{ABC_{II}}$,

$$S(\rho_{ABC_{II}}) = \max\{8\sqrt{2}|\frac{p}{2} \sin r_c|, 4|(\frac{2-p}{2}) \cos 2r_c + \frac{p}{2}|\}.$$

From FIG. 1 (b) we see that $S(\rho_{ABC_{II}})$ is always small than 4 for any r_c and p . Thus the physically inaccessible GTN between modes A, B and C_{II} cannot be produced.

Case(2): Alice still stays at an asymptotically flat region, while Bob and Charlie hover outside the event horizon of the black hole. Then in terms of the Kruskal modes for Alice and Schwarzschild modes for Bob and Charlie, the initial state ρ_1 can be rewritten as a 5-partite entangled state $\rho_{AB_I B_{II} C_I C_{II}}$ consisted by subsystem A observed by Alice, subsystems B_I and C_I observed by Bob and Charlie outside the event horizon of black hole, and subsystems B_{II} and C_{II} observed by anti-Bob and anti-Charlie inside the event horizon, respectively. We call the modes B_I and C_I outside the event horizon the accessible modes, and the modes B_{II} and C_{II} inside the event horizon the inaccessible modes. Tracing over the inaccessible modes in state $\rho_{AB_I B_{II} C_I C_{II}}$, we obtain the reduced density operator,

$$\rho_{AB_I C_I} = \begin{pmatrix} d_1 & 0 & 0 & 0 & 0 & 0 & 0 & f_1 \\ 0 & d_2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & d_3 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & d_4 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ f_1 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{p}{2} \end{pmatrix},$$

where $d_1 = (\frac{2-p}{2}) \cos^2 r_b \cos^2 r_c$, $d_2 = (\frac{2-p}{2}) \cos^2 r_b \sin^2 r_c$, $d_3 = (\frac{2-p}{2}) \sin^2 r_b \cos^2 r_c$, $d_4 = (\frac{2-p}{2}) \sin^2 r_b \sin^2 r_c$ and $f_1 = \frac{p}{2} \cos r_b \cos r_c$. Here, $\cos r_b = (e^{-\frac{\alpha}{T_b}})^{-\frac{1}{2}}$ and $\sin r_b = (e^{\frac{\alpha}{T_b}})^{-\frac{1}{2}}$ are functions of the Hawking temperature T_b . We obtain the GTN for $\rho_{AB_I C_I}$,

$$S(\rho_{AB_I C_I}) = \max\{4\sqrt{2}p|\cos r_b \cos r_c|, 4|(\frac{2-p}{2}) \cos 2r_b \cos 2r_c - \frac{p}{2}|\}.$$

In FIG. 2, we plot the Svetlichny value $S(\rho_{AB_I C_I})$ as function of $r_b = r_c$ and p . It is shown that $S(\rho_{AB_I C_I})$ is larger than 4 at first and then smaller than 4 with the increase of Hawking temperature T and the decrease of p . This implies that the thermal noise introduced by Hawking temperature destroys the physically accessible GTN between Alice, Bob and Charlie. The ‘‘sudden death’’ of such physically accessible GTN takes place at some critical temperature T_b .

The reduced density matrix $\rho_{AB_I B_{II}}$ can be expressed




FIG. 2: Case(2): The Svetlichny value $S(\rho_{AB_I C_I})$ as a function of $r_c = r_b$ and p for the case that Alice stays stationarily, while Bob and Charlie move with a uniform acceleration in their vacuum cavities.

as

$$\rho_{AB_I B_{II}} = \begin{pmatrix} d_1 & 0 & 0 & 0 & 0 & 0 & 0 & f_1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{p}{2} & 0 \\ f_1 & 0 & 0 & 0 & 0 & 0 & 0 & e_1 \end{pmatrix},$$

with the matrix elements give by $d_1 = (\frac{2-p}{2}) \cos^2 r_b$, $f_1 = (\frac{2-p}{2}) \cos r_b \sin r_b$ and $e_1 = (\frac{2-p}{2}) \sin^2 r_b$. In FIG. 3, we plot the Svetlichny value $S(\rho_{AB_I B_{II}})$ as function of $r_b = r_c$ and p . We find that $S(\rho_{AB_I B_{II}})$ is smaller than 4 at first and then larger than 4 with the increase of Hawking temperature T and the decrease of p . This phenomenon is quite novel as the birth of the physically inaccessible GTN for $\rho_{AB_I B_{II}}$ has never been observed in previous literature with pure initial states [43, 44]. To see this new phenomenon more clearly, we draw $S(\rho_{AB_I B_{II}})$ corresponding to different values of p in FIG. 3(b). We see that when $p = 1$, the birth of GTN does not show up. As the value of p becomes smaller, GTN becomes more than 4 when $p = \frac{1}{2}, \frac{1}{4}$ and 0. This result shows that the GTN can pass through the event horizon of black hole for the case of mixed states.

Due the symmetry under the interchange of Bob and Charlie, we have $S(\rho_{AC_I C_{II}}) = S(\rho_{AB_I B_{II}})$ and $S(\rho_{AB_I C_{II}}) = S(\rho_{AB_{II} C_I})$. Therefore, the GTN of $\rho_{AC_I C_{II}}$ has similar properties to $\rho_{AB_I B_{II}}$ under the influence of Hawking effect. We only need to consider the GTN for the rest two types of reduced states $\rho_{AB_I C_{II}}$ and $\rho_{AB_{II} C_I}$ of $\rho_{AB_I B_{II} C_I C_{II}}$. By direct calculation, we




FIG. 3: Case(2): Alice stays stationarily, while Bob and Charlie move with a uniform acceleration in their vacuum cavities. (a) the Svetlichny value $S(\rho_{AB|B_{II}})$ as function of r_b and p . (b) the Svetlichny value $S(\rho_{AB|B_{II}})$ as function of r_b . The dot-dashed gray, the dashed blue, the dotted red and the solid blue lines are for $p = 1, \frac{1}{2}, \frac{1}{4}, 0$, respectively.

have

$$\rho_{AB|C_{II}} = \begin{pmatrix} d_1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & d_2 & 0 & 0 & 0 & 0 & f_2 & 0 \\ 0 & 0 & d_3 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & d_4 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & f_2 & 0 & 0 & 0 & 0 & \frac{p}{2} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

where $d_1 = \left(\frac{2-p}{2}\right) \cos^2 r_b \cos^2 r_c$, $d_2 = \left(\frac{2-p}{2}\right) \cos^2 r_b \sin^2 r_c$, $d_3 = \left(\frac{2-p}{2}\right) \sin^2 r_b \cos^2 r_c$, $d_4 = \left(\frac{2-p}{2}\right) \sin^2 r_b \sin^2 r_c$ and $f_2 = \frac{p}{2} \cos r_b \cos r_c$,




FIG. 4: Case(2): Alice stays stationarily, while Bob and Charlie move with a uniform acceleration in their vacuum cavities. (a) the Svetlichny value $S(\rho_{AB|C_{II}})$ as function of $r_c = r_b$ and p . (b) The Svetlichny value $S(\rho_{AB_{II}|C_I})$ as function of $r_c = r_b$ and p .

and

$$\rho_{AB_{II}C_{II}} = \begin{pmatrix} d_1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & d_2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & d_3 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & d_4 & f_4 & 0 & 0 & 0 \\ 0 & 0 & 0 & f_4 & \frac{p}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

where $d_1 = \left(\frac{2-p}{2}\right) \cos^2 r_b \cos^2 r_c$, $d_2 = \left(\frac{2-p}{2}\right) \cos^2 r_b \sin^2 r_c$, $d_3 = \left(\frac{2-p}{2}\right) \sin^2 r_b \cos^2 r_c$, $d_4 = \left(\frac{2-p}{2}\right) \sin^2 r_b \sin^2 r_c$ and $f_4 = \frac{p}{2} \sin r_b \sin r_c$.

Using (8) we obtain $S(\rho_{AB|C_{II}})$ and $S(\rho_{AB_{II}|C_I})$, respectively. FIG. 4 shows that the GTN cannot be generated between A, B_I and C_{II} or between A, B_{II} and C_{II} .




FIG. 5: Alice stays stationarily, while Bob and Charlie move with a uniform acceleration in their vacuum cavities. The Svetlichny value $S(\rho_{AB_I B_{II}})$ with respect to the initial state ρ_2 as function of r_b and p .

We have used the initial mixed state $\rho_1 = p|GHZ\rangle\langle GHZ| + (1-p)|000\rangle\langle 000|$ to display the novel phenomenon of sudden birth of GTN. However, such phenomenon may not appear for other mixed states. Let us consider the mixed state $\rho_2 = p|GHZ\rangle\langle GHZ| + (1-p)|111\rangle\langle 111|$. For ρ_2 the GTN for other reduced states are similar to ρ_1 , except for $S(\rho_{AB_I B_{II}})$ which is always less than 4, namely, no GTN is generated, See FIG. 5. The only difference between the states ρ_1 and ρ_2 is that the vector $|000\rangle$ is replaced with $|111\rangle$. Their different behaviors on the new generation of physically inaccessible GTN are due to that the evolution process of vacuum state of Dirac fields in a Schwarzschild black hole has different effects on $|0\rangle$ and $|1\rangle$.

EVOLUTION OF THE GTE AND QUANTUM COHERENCE IN SCHWARZSCHILD BLACK HOLE

In this section, we explore the GTE and quantum coherence under evolution of ρ_1 after the acceleration effect for one or two accelerated detector(s) coupled to the vacuum field.

It is very coincidental that according to Eqs. (9) and (10), we find that for all the tripartite reduced states in the previous section, the GTE is equal to the quantum coherence. For the case (1), we have

$$\begin{aligned} E(\rho_{ABC_I}) &= C(\rho_{ABC_I}) = p \cos r_c, \\ E(\rho_{ABC_{II}}) &= C(\rho_{ABC_{II}}) = p \sin r_c. \end{aligned}$$

For the case (2), we obtain

$$\begin{aligned} E(\rho_{AB_I C_I}) &= C(\rho_{AB_I C_I}) = p |\cos r_b \cos r_c|, \\ E(\rho_{AB_I C_{II}}) &= C(\rho_{AB_I C_{II}}) = p |\cos r_b \sin r_c|, \\ E(\rho_{AB_I B_{II}}) &= C(\rho_{AB_I B_{II}}) = (2-p) |\cos r_b \sin r_b|, \\ E(\rho_{AB_{II} C_{II}}) &= C(\rho_{AB_{II} C_{II}}) = p |\sin r_b \sin r_c|. \end{aligned}$$

Due to the symmetry between Bob and Charlie, it is easy to find that

$$\begin{aligned} E(\rho_{AC_I C_{II}}) &= E(\rho_{AB_I B_{II}}) = (2-P) |\sin r_c \cos r_c|, \\ E(\rho_{AB_{II} C_I}) &= E(\rho_{AB_I C_{II}}) = p |\cos r_c \sin r_b|. \end{aligned}$$

The tradeoff relationship and the monogamy relations are important issues in characterizing the quantum correlations in quantum information theory, which are deeply related to quantum uncertainty and incompatibility principle. Noted that the acceleration of Bob and Charlie is completed under the same Hawking radiation. Hence, the corresponding changes of the Hawking temperatures T_b and T_c are the same. Therefore, the functions $\cos r_b$ and $\cos r_c$ corresponding to T are independent individually, namely, $\cos r_b = \cos r_c$ and $\sin r_b = \sin r_c$. Therefore, we have the following tradeoff relations between the physically accessible GTE (quantum coherence) and the physically inaccessible GTE (quantum coherence),

$$E^2(\rho_{ABC_I}) + E^2(\rho_{ABC_{II}}) = p^2 = E^2(\rho_1), \quad (11)$$

$$E(\rho_{AB_I C_I}) + E(\rho_{AB_{II} C_{II}}) = p = E(\rho_1), \quad (12)$$

$$\begin{aligned} E^2(\rho_{AB_I C_I}) + E^2(\rho_{AB_{II} C_I}) + E^2(\rho_{AB_I C_{II}}) \\ + E^2(\rho_{AB_{II} C_{II}}) = p^2. \end{aligned} \quad (13)$$

The above trade-off relations give restrictions on the redistribution of entanglement from physically accessible to physically inaccessible patterns. For instance, (12) shows that the total physically accessible GTE $E(\rho_{AB_I C_I})$ and the physically inaccessible GTE $E(\rho_{AB_{II} C_{II}})$ is equal to the initial GTE of ρ_1 . Therefore, under the time evolution how much the physically accessible GTE decreases implies how much the physically inaccessible GTE increases. The Eq. (13) poses a strong restrictive relationship among the four genuine tripartite entanglement. These tradeoff relations shed new lights on understanding the quantum information theory in the framework of relativity.

CONCLUSIONS AND DISCUSSIONS

Quantum nonlocality is a distinctive feature of quantum mechanics. We have investigated the effect of Hawking radiation on GTN for Dirac fields in Schwarzschild spacetime. We first time show that the Hawking effect can not only degrades the physically accessible GTN, but also enhance the physically inaccessible GTN. The ‘‘sudden death’’ of the physically accessible GTN at certain critical Hawking temperature and the ‘‘sudden birth’’ of

the physically inaccessible GTN imply that, with the loss of the physically accessible GTN, the physically inaccessible GTN is generated continually through Hawking effect. This phenomenon is observed first time for GTN. If we regard the redistribution of nonlocality as a kind of phenomenon of information tunnelling, this phenomenon shows that not only the flow of quantum entanglement can pass through the black hole perspective, but also the nonlocality can either. This phenomenon also enriches further understanding of relativistic quantum information.

We have also investigated the problem for quantum correlation GTE and quantum coherence. Interesting trade-off relations are derived among the GTEs and the quantum coherence, which restrict the distribution of GTE and coherence in the subsystems during the evolution. With regard to the related theories of entanglement measurement under the effect of Hawking radiation for Dirac fields in Schwarzschild spacetime, the previous literatures mainly focused on pure initial states. Based on initial mixed states, we have shown the new phenomenon that the physically inaccessible GTN is generated continually through Hawking effect. From the view of resource theory in quantum mechanics, the quantum correlations including entanglement and nonlocality, as well as the coherence may have universal redistribution properties. Our results may highlight new understanding of relativistic quantum information.

Acknowledgments

This work is supported by the National Natural Science Foundation of China (NSFC) under Grant Nos.11861031, 12075159 and 12171044; the Hainan Provincial Natural Science Foundation of China under Grant Nos.121RC539, the specific research fund of the Innovation Platform for Academicians of Hainan Province under Grant No. YSPTZX202215, Beijing Natural Science Foundation (Grant No. Z190005).

[1] A. Einstein, B. Podolsky and N. Rosen, *Phys. Rev.* **47**, 777 (1935)
 [2] H. Cao, Z. Guo, *Sci. China Phys. Mech. Astron.* **62**, 30311 (2019)
 [3] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani and S. Wehner, *Rev. Mod. Phys.* **86**, 419 (2014)
 [4] A. Acin, N. Brunner, N. Gisin, S. Massar, S. Pironio and V. Scarani, *Phys. Rev. Lett.* **98**, 230501 (2007)
 [5] S. Pironio, et al., *Nature*, **464**, 1021 (2010)
 [6] R. Colbeck and R. Renner, *Nature Physics*, **8**, 450 (2012)
 [7] M. H. Li, et al., *Phys. Rev. Lett.* **126**, 050503 (2021)
 [8] W. Z. Liu, et al., *Nature Physics*, **17**, 448 (2021)
 [9] L. K. Shalm, et al., *Nature Physics*, **17**, 452 (2021)

[10] G. Svetlichny, *Phys. Rev. D* **35**, 3066 (1987)
 [11] N. Gisin, *Quantum Reality, Relativistic Causality, and Closing the Epistemic Circle. The Western Ontario Series in Philosophy of Science*, **73**, Springer, Dordrecht. https://doi.org/10.1007/978-1-4020-9107-0_9 (2009).
 [12] V. Scarani and N. Gisin, *Phys. Rev. Lett.* **87**, 117901 (2001).
 [13] M. Seevinck, Ph.D. thesis, Utrecht University, arXiv:0811.1027v2 [quant-ph].
 [14] P. Zoller et al., *Eur. Phys. J. D* **36**, 203 (2005).
 [15] C. Y. Lu et al., *Nature Phys.* **3**, 91 (2007).
 [16] S.W. Hawking, *Nature* **248**, 30 (1974)
 [17] W.G. Unruh, *Phys. Rev. D* **14** 870 (1976)
 [18] L.C.B. Crispino, A. Higuchi and G.E.A. Matsas, *Rev. Mod. Phys.* **80**, 787 (2008)
 [19] I. Fuentes-Schuller, R.B. Mann, *Phys. Rev. Lett.* **95**, 120404 (2005)
 [20] P.M. Alsing, I. Fuentes-Schuller, R.B. Mann, T.E. Tessier, *Phys. Rev. A* **74**, 032326 (2006)
 [21] J. Wang, Q. Pan, S. Chen, J. Jing, *Phys. Lett. B* **677**, 186 (2009)
 [22] E. Martín-Martínez, L.J. Garay, J. León, *Phys. Rev. D* **82**, 064006 (2010)
 [23] J. Wang, Q. Pan, J. Jing, *Phys. Lett. B* **692**, 202 (2010)
 [24] J. Wang, Q. Pan, J. Jing, *Ann. Phys.* **325**, 1190 (2010)
 [25] E. Martín-Martínez, L.J. Garay, J. León, *Phys. Rev. D* **82**, 064028 (2010)
 [26] J. Wang, J. Jing, *Phys. Rev. A* **82**, 032324 (2010)
 [27] S. Khan, M.K. Khan, *J. Phys. A Math. Theor.* **44**, 045305 (2011)
 [28] J. Wang, J. Jing, *Phys. Rev. A* **83**, 022314 (2011)
 [29] B. Nasr Esfahani, M. Shamirzaie, M. Soltani, *Phys. Rev. D* **84**, 025024 (2011)
 [30] R.B. Mann, T.C. Ralph, *Class. Quantum Gravity* **29**, 220301(2012)
 [31] Z. Tian, J. Jing, *J. High Energy Phys.* **04**, 109 (2013)
 [32] S. Xu, X.K. Song, J.D. Shi, L. Ye, *Phys. Rev. D* **89**, 065022 (2014)
 [33] Z. Tian, J. Jing, *J. High Energy Phys.* **07**, 089 (2014)
 [34] S. Kanno, J.P. Shock, J. Soda, *Phys. Rev. D* **94**, 125014 (2016)
 [35] Y. Dai, Z. Shen, Y. Shi, *Phys. Rev. D* **94**, 025012 (2016)
 [36] N. Friis, *New J. Phys.* **18**, 033014 (2016)
 [37] X. Liu, Z. Tian, J. Wang, J. Jing, *Phys. Rev. D* **97**, 105030 (2018)
 [38] W.C. Qiang, G.H. Sun, Q. Dong, and S.H. Dong, *Phys. Rev. A* **98**, 022320 (2018)
 [39] A.J. Torres-Arenas, Q. Dong, G.H. Sun, W.C. Qiang, S.H. Dong, *Phys. Lett. B* **789**, 93 (2019)
 [40] S.M. Wu, H.S. Zeng, *Int. J. Theor. Phys.* **59**, 861 (2020)
 [41] S.M. Wu, H.S. Zeng, *Class. Quantum Gravity* **37**, 115003 (2020)
 [42] S.M. Wu, Z.C. Li and H.S. Zeng, *EPL*, **129**, 40002 (2020)
 [43] S.M. Wu, H.S. Zeng, *Eur. Phys. J. C* **82**, 4 (2022)
 [44] L.J. Li, F. Ming, X.K. Song, L. Ye, D. Wang, *Eur. Phys. J. C* **82**, 726 (2022)
 [45] S.M. Wu, H.S. Zeng and T. Liu, *New J. Phys.* **24**, 073004 (2022)
 [46] N. Gisin, *Phys. Lett. A* **154**, 201(1991)
 [47] S.X. Yu, Q. Chen, C.J. Zhang, C.H. Lai, and C.H. Oh, *Phys. Rev. Lett.* **109**, 120402 (2012)
 [48] S.H. Jiang, Z.P. Xu, H.Y. Su, A. K. Pati and J.L. Chen, *Phys. Rev. Lett.* **120**, 050403 (2018)
 [49] D.R. Brill, J.A. Wheeler, *Rev. Mod. Phys.* **29**, 465 (1957)

- [50] D. Garfinkle, G.T. Horowitz, A. Strominger, Phys. Rev. D **43**, 3140 (1991);
A. Gareia, D. Galtsov, O. Kechkin, Phys. Rev. Lett. **74**, 1276 (1995)
- [51] S.W. Hawking, Commun. Math. Phys. **43**, 199 (1975);
G.W. Gibbons, S.W. Hawking, Phys. Rev. D **15**, 2738 (1977)
- [52] T. Damoar, R. Ruffini, Phys. Rev. D **14**, 332 (1976)
- [53] K. Wang, Z.J. Zheng, Sci. Rep. **10**, 6621 (2020);
K. Wang, Y. Liang, Z.J. Zheng, Quantum Inf. Process. **19**, 140 (2020)
- [54] Z.H. Ma, Z.H. Chen, J.L. Chen, C. Spengler, A. Gabriel, M. Huber, Phys. Rev. A **83**, 062325 (2011);
S.M. Hashemi Rafsanjani, M. Huber, C.J. Broadbent, J.H. Eberly, Phys. Rev. A **86**, 062303 (2012)
- [55] T. Baumgratz, M. Cramer, M.B. Plenio, Phys. Rev. Lett. **113**, 140401 (2014);
A. Streltsov, G. Adesso, M.B. Plenio, Rev. Mod. Phys. **89**, 041003 (2017);
M.L. Hu, X. Hu, J.C. Wang, Y. Peng, Y.R. Zhang, H. Fan, Phys. Rep. **762**, 1 (2018)