

Wrapping Cycles in Delaunay Complexes: Bridging Persistent Homology and Discrete Morse Theory

Ulrich Bauer   

Department of Mathematics and Munich Data Science Institute,
Technische Universität München, Germany

Fabian Roll   

Department of Mathematics, Technische Universität München, Germany

Abstract

We study the connection between discrete Morse theory and persistent homology in the context of shape reconstruction methods. Specifically, we consider the construction of Wrap complexes, introduced by Edelsbrunner as a subcomplex of the Delaunay complex, and the construction of lexicographic optimal homologous cycles, also considered by Cohen–Steiner, Lieutier, and Vuillamy in a similar setting. We show that for any cycle in a Delaunay complex for a given radius parameter, the lexicographically optimal homologous cycle is supported on the Wrap complex for the same parameter, thereby establishing a close connection between the two methods. We obtain this result by establishing a fundamental connection between reduction of cycles in the computation of persistent homology and gradient flows in the algebraic generalization of discrete Morse theory.

2012 ACM Subject Classification Mathematics of computing → Geometric topology; Theory of computation → Computational geometry

Keywords and phrases persistent homology, discrete Morse theory, apparent pairs, Wrap complex, lexicographic optimal chains, shape reconstruction

Supplementary Material Source code publicly available under MIT license.

Software (Source code): <https://github.com/fabian-roll/wrappingcycles> [10]


archived at `swb:1:dir:5edd13ee3b341318256c442f703d84a0b1b27c6a`

Funding This research has been supported by the Deutsche Forschungsgemeinschaft (DFG – German Research Foundation) – Project-ID 195170736 – TRR109 *Discretization in Geometry and Dynamics*.

Acknowledgements We thank Herbert Edelsbrunner and Marian Mrozek for interesting discussions about the connection between algebraic gradient flow in discrete Morse theory and matrix reduction in persistent homology.

1 Introduction

Reconstructing shapes and submanifolds from point clouds is a classical topic in computational geometry. Starting in the 2000s, several key results have been achieved [1, 2, 12, 14, 21, 23, 24, 37, 39], culminating in theoretical homeomorphic reconstruction guarantees for a method based on the Delaunay triangulation [14]. The method is theoretical in nature, and their practical applicability is hindered by complexity and robustness issues. A major challenge is caused by *slivers* [13], which are simplices in the Delaunay triangulation with small volume but no short edges, and which have to be handled explicitly. In contrast, several related Delaunay-based methods have proven highly robust and successful in practice, in particular, Morse-theory based methods such as *Wrap* and related constructions [6, 22, 24, 38–40] and homological methods based on minimal cycles [4, 17–19, 42]. The latter methods produce *water-tight* surfaces (that is, boundaries of solids) by construction, gracefully circumventing




■ **Figure 1** Left: Delaunay triangulation of a point cloud, with critical simplices highlighted. Middle: Wrap complex for a small radius parameter. Right: lexicographically minimal cycle for the most persistent feature (black contour), shown together with its bounding chain (shaded blue).

the issue of slivers. On the other hand, the *Wrap complex* [6] is always homotopy-equivalent to a union of balls of a given radius, but may still contain critical sliver simplices. It is therefore desirable to identify a subcomplex that is free from slivers.

Even though the Morse-theoretic and the homological method seem closely related in spirit, up to now the connection between them has not been fully explored. While the development of Wrap preceded the introduction of Forman’s discrete Morse theory [28], it has since been redeveloped within this framework [6], which provides powerful tools for proving geometric and topological properties of this construction. A bridge from Wrap to homology-based methods advances our understanding of their geometric and topological properties, potentially paving the way for reconstruction guarantees that extend beyond surfaces. Indeed, a synthesis of Wrap, discrete Morse theory, and persistent homology in the context of surface reconstruction has been envisioned already in Edelsbrunner’s original paper describing the Wrap algorithm [24].

Contributions The objective of this paper is twofold. First, we provide a tight link between persistent homology and discrete Morse theory, unifying the central notions of persistence pairs and of gradient pairs in a common framework. Despite various results connecting both theories (see, e.g., [5, 8, 35]), this kind of interface has been missing from the literature. Using a specific refinement of the sublevel set filtration of a discrete Morse function, we demonstrate how the persistence pairs yield an algebraic gradient that contains as a subset the gradient pairs of the discrete Morse function, and such that the corresponding algebraic gradient flow can be viewed as a variant of the reduction algorithm for computing persistent homology (see Section 4). Second, we use these insights to establish a strong connection between the aforementioned Morse-theoretic and homological approaches to shape reconstruction. Our main result (specializing Theorem 29 to the Delaunay radius function, see Example 6), shows that the lexicographically minimal cycles (see Definition 4) in a Delaunay complex are all supported on the corresponding Wrap complex (see Definition 8), as illustrated in Figure 1. In the statement of the theorem, the order on simplices follows the convention outlined in Section 2.3, and homology is considered with coefficients in a specified field.

► **Theorem A.** *Let $X \subset \mathbb{R}^d$ be a finite subset in general position, let $r \in \mathbb{R}$, and let $h \in H_*(\text{Del}_r(X))$ be a homology class of the Delaunay complex $\text{Del}_r(X)$. Then the lexicographically minimal cycle of h , with respect to the Delaunay-lexicographic order on the simplices, is supported on the Wrap complex $\text{Wrap}_r(X)$.*



■ **Figure 2** The lexicographically minimal cycle corresponding to the most persistent feature of the Delaunay filtration for 3D scan point clouds [32] yields an accurate reconstruction of the surface.

As a consequence of our main result, we obtain the following connection between Wrap complexes and persistent homology (see Section 2.1) of the Delaunay filtration, specializing Corollary 30 to the Delaunay radius function. Our result states that the minimal cycle homologous to the boundary of a simplex killing homology in the Delaunay filtration – obtained by a variant of the familiar matrix reduction algorithm for computing persistence – is supported on the Wrap complex for the birth value of the corresponding feature.

► **Corollary B.** *Let $X \subset \mathbb{R}^d$ be a finite subset in general position and (σ, τ) a non-zero persistence pair of the lexicographically refined Delaunay filtration. Let $r = r_X(\sigma)$ and $s = r_X(\tau)$ be the radius of the smallest empty circumsphere of σ and of τ , respectively. Then the lexicographically minimal cycle of $[\partial\tau]$ in the Delaunay complex $\text{Del}_s^{\leq}(X)$, given as the column R_τ of the totally reduced filtration boundary matrix, is supported on the Wrap complex $\text{Wrap}_r(X)$.*

The totally reduced filtration boundary matrix can be computed using Algorithm 2. The connection between lexicographically minimal cycles and matrix reduction algorithms used in the context of persistent homology has already been discussed in [18].

For a sufficiently good sample of a compact d -submanifold of Euclidean space, the union of closed balls centered at the sample points deformation retracts onto the submanifold by a closest point projection [3, 37]. As the Delaunay complex is naturally homotopy equivalent to the union of closed balls [7], this implies that the fundamental class of the manifold is captured directly in the d -dimensional persistent homology of the Delaunay filtration through a natural isomorphism. Similar observations that are based on the induced maps in persistent homology and with weaker assumptions have also been made before [15] [11, Section 11.4]. Moreover, in the preprint [17, Theorem 9.1] it is argued that the unique non-trivial lexicographically optimal 2-cycle in the Čech complex of a sample of a 2-submanifold of Euclidean space yields a homeomorphic reconstruction of the submanifold. Note that the lexicographically optimal cycles considered in [17–19, 42] are based on a slightly different total order on simplices, refining the minimum enclosing radius function. Relating these particular choices and results to our results and those in [6] remains an interesting open problem.

Together with Corollary B, the discussion above suggests the following heuristic for a simple and robust algorithm for shape reconstruction from a point cloud, combining Wrap and lexicographically minimal cycles with persistent homology: Take the most persistent d -dimensional feature of the Delaunay filtration, i.e., the interval in the barcode with the largest death/birth ratio. Intuitively, this feature is born at a small scale and only gets filled in at a large scale. By Corollary B, the corresponding lexicographically minimal cycle is guaranteed to be supported on the Wrap complex for a small scale parameter. See Figure 2 for an illustration, which can readily be reproduced using the code provided in [10] by executing `docker build -o output github.com/fabian-roll/wrappingcycles` on any machine with Docker installed and configured with sufficient memory (16GB recommended).

2 Preliminaries

A finite *simplicial complex* K is a collection of finite nonempty sets such that for any set $\sigma \in K$ and any nonempty subset $\rho \subseteq \sigma$ one has $\rho \in K$. A set $\sigma \in K$ is called a *simplex*, and $\dim \sigma = \text{card } \sigma - 1$ is its *dimension*. Moreover, ρ is said to be a *face of* σ and σ a *coface of* ρ . If $\dim \rho = \dim \sigma - 1$, then we call ρ a *facet of* σ , σ a *cofacet of* ρ , and (ρ, σ) a *facet pair*.

2.1 Persistent homology and apparent pairs

Based chain complexes and filtrations We assume the reader to be familiar with the basics of homological algebra (see, e.g., [36, 43]). By a *based chain complex* (C_*, Σ_*) (sometimes also called a *Lefschetz complex* [34]) we mean a bounded chain complex $C_* = (C_n, \partial)_{n \in \mathbb{N}}$ of finite dimensional vector spaces over a field \mathbb{F} together with a basis Σ_n for each C_n . Consider the canonical bilinear form $\langle \cdot, \cdot \rangle$ on C_* for the given basis Σ_* , i.e., for $a, b \in \Sigma_*$ we have $\langle a, b \rangle = 0$ if $a \neq b$ and $\langle a, a \rangle = 1$. Given two basis elements $c \in \Sigma_n$ and $e \in \Sigma_{n+1}$ such that $\langle \partial e, c \rangle \neq 0$, we call c a *facet of* e and e a *cofacet of* c , and we call (c, e) a *facet pair*.

For a poset P and any element $p \in P$ we denote by $\downarrow p = \{q \in P \mid q \leq p\}$ the *down set* of p . A *filtration* of (C_*, Σ_*) is a collection of based chain complexes $(C_*^i, \Sigma_*^i)_{i \in I}$, where I is a totally ordered indexing set, such that $\Sigma_*^i \subseteq \Sigma_*$ spans the subcomplex C_*^i of C_* for all $i \in I$, and $i \leq j$ implies $\Sigma_*^i \subseteq \Sigma_*^j$. We call the filtration an *elementwise filtration* if for any j with immediate predecessor i we have that $\Sigma_*^j \setminus \Sigma_*^i$ contains exactly one basis element σ_j . Thus, elementwise filtrations of (C_*, Σ_*) correspond bijectively to total orders $<$ on Σ_* such that the down sets $\downarrow \sigma_j = \{\sigma_i \mid \sigma_i \leq \sigma_j\}$ span subcomplexes.

Our main example for a based chain complex is the simplicial chain complex $C_*(K)$ of a simplicial complex K with coefficients in a field. If the vertices of K are totally ordered, then there is a canonical basis of $C_n(K)$ consisting of the n -dimensional simplices of K oriented according to the given vertex order, and a simplexwise filtration of K induces a canonical elementwise filtration of $C_*(K)$.

Matrix reduction For a based chain complex $(C_*, \Sigma_* = \sigma_1 < \dots < \sigma_l)$ with an elementwise filtration, we often identify an element of C_* with its coordinate vector in \mathbb{F}^l . The *filtration boundary matrix* D of an elementwise filtration is the matrix that represents the boundary map ∂ with respect to the total order on the basis elements induced by the filtration.

For a matrix R , we denote by R_j the j th column of R , and by $R_{i,j}$ the entry of R in row i and column j . The *pivot* of a column R_j , denoted by $\text{PivInd } R_j$, is the maximal row index i with $R_{i,j} \neq 0$, taken to be 0 if all entries are 0. Otherwise, the non-zero entry is called the *pivot entry*, denoted by $\text{PivEnt } R_i$. We define $\text{PivInds } R = \{i \mid i = \text{PivInd } R_j \neq 0\}$ to be the collection of all non-zero column pivots. Moreover, we call a column R_j *reduced* if

Input: $D = \partial$ an $l \times l$ filtration boundary matrix
Result: $R = D \cdot S$ with R reduced and S full rank upper triangular
 $R = D$; $S = \text{Id}$;
for $j = 1$ **to** l **do**
 while *there exists* $i < j$ *with* $\text{PivInd } R_i = \text{PivInd } R_j > 0$ **do**
 $\mu = -\text{PivEnt } R_j / \text{PivEnt } R_i$;
 $R_j = R_j + \mu \cdot R_i$; $S_j = S_j + \mu \cdot S_i$;
return R, S

■ **Algorithm 1** Standard matrix reduction

Input: $D = \partial$ an $l \times l$ filtration boundary matrix
Result: $R = D \cdot S$ with R totally reduced and S full rank upper triangular
 $R = D$; $S = \text{Id}$;
for $j = 1$ **to** l **do**
 while *there exist* $s, i < j$ *with* $\text{PivInd } R_i = s$ *and* $R_{s,j} \neq 0$ **do**
 $\mu = -R_{s,j} / R_{s,i}$;
 $R_j = R_j + \mu \cdot R_i$; $S_j = S_j + \mu \cdot S_i$;
return R, S

■ **Algorithm 2** Exhaustive matrix reduction

its pivot cannot be decreased by adding a linear combination of the columns R_i with $i < j$, and we call the matrix R reduced if all its columns are reduced. Finally, we call the matrix R *totally reduced* if for each $i < j$ we have $R_{s,j} = 0$, where $s = \text{PivInd } R_i$.

We call a matrix S a *reduction matrix* if it is a full rank upper triangular matrix such that $R = D \cdot S$ is reduced and S is *homogeneous*, meaning that respects the degrees in the chain complex. Any such reduction $R = D \cdot S$ of the filtration boundary matrix induces a direct sum decomposition (see, e.g., [5, 20]) of C_* into elementary chain complexes in the following way: If $R_j \neq 0$, then we have the summand

$$\cdots \rightarrow 0 \rightarrow \langle S_j \rangle \xrightarrow{\partial} \langle R_j \rangle \rightarrow 0 \rightarrow \cdots ,$$

in which case we call j a *death index*, $i = \text{PivInd } R_j$ a *birth index*, and (i, j) an *index persistence pair*. If $R_i = 0$ and $i \notin \text{PivInds } R$, then we have the summand

$$\cdots \rightarrow 0 \rightarrow \langle S_i \rangle \rightarrow 0 \rightarrow \cdots ,$$

in which case we call i an *essential index*. Moreover, we call the element σ_i a *birth*, *death*, or *essential element*, if its index is a birth, death, or essential index. Similarly, we call a pair of elements (σ_i, σ_j) a *persistence pair*, if the pair (i, j) is an index persistence pair. Note that this is independent of the specific reduction of the filtration boundary matrix. By taking the intersection with the filtration, one obtains elementary filtered chain complexes, in which R_j is a cycle appearing in the filtration at index $i = \text{PivInd } R_j$ and becoming a boundary when S_j enters the filtration at index j , and in which an essential cycle S_i enters the filtration at index i . Thus, the *barcode* of the *persistent homology* [25] of the elementwise filtration is given by the collection of intervals $\{[i, j) \mid (i, j) \text{ index persistence pair}\} \cup \{[i, \infty) \mid i \text{ essential index}\}$.

Such a reduction $R = D \cdot S$ can be computed by a variant of Gaussian elimination [16], as in Algorithm 1. A slight modification is Algorithm 2, which computes a totally reduced filtration boundary matrix, as used in Corollary B. This is also known as *exhaustive reduction*, and appears in various forms in the literature [18, 26, 27, 44].

Apparent pairs Many optimization schemes have been developed in order to speed up the computation of persistent homology. One of them is based on *apparent pairs* [5], a concept which lies at the interface of persistence and discrete Morse theory.

► **Definition 1.** Let $(C_*, \Sigma_* = \sigma_1 < \dots < \sigma_l)$ be a based chain complex with an elementwise filtration. We call a pair of basis elements (σ_i, σ_j) an *apparent pair* if σ_i is the maximal facet of σ_j and σ_j is the minimal cofacet of σ_i .

In the context of persistence, the interest in apparent pairs stems from the following observation [5], immediate from the definitions.

► **Lemma 2.** For any apparent pair (σ, τ) of an elementwise filtration, the column of τ in the filtration boundary matrix is reduced, and (σ, τ) is a persistence pair.

2.2 Lexicographic optimality

In this section, we introduce the lexicographic order on chains for a based chain complex $(C_*, \Sigma_* = \sigma_1 < \dots < \sigma_l)$ with an elementwise filtration, extending the definitions in [18]. For any chain $c = \sum_i \lambda_i \sigma_i \in C_n$, we define its *support* $\text{supp}_{\Sigma_*} c$ to be the set of basis elements $\sigma_i \in \Sigma_n$ with $\lambda_i \neq 0$. Note that this is not to be confused with the supporting subcomplex, which also contains the faces of these basis elements. Given a totally ordered set (X, \leq) , we consider the lexicographic order \preceq on the power set 2^X given by identifying any subset $A \subseteq X$ with its characteristic function and considering the lexicographic order on the set of characteristic functions. Explicitly, for $A, B \subseteq X$ we have $A \preceq B$ if and only if $A = B$ or the maximal element of the symmetric difference $(A \setminus B) \cup (B \setminus A)$ is contained in B .

► **Definition 3.** The lexicographic preorder \sqsubseteq on the collection of chains C_n is given by $c_1 \sqsubseteq c_2$ if and only if $\text{supp}_{\Sigma_*} c_1 \preceq \text{supp}_{\Sigma_*} c_2$ in the lexicographic order on subsets of Σ_n . We write \sqsubset for the corresponding strict preorder.

If we consider a simplicial chain complex with coefficients in $\mathbb{Z}/2\mathbb{Z}$, then this preorder is a total order, and it coincides with the one considered in [18, Definition 2.1].

► **Definition 4.** We call a chain $c \in C_n$ *lexicographically minimal*, or *irreducible*, if there exists no strictly smaller homologous chain $(c + \partial e) \sqsubset c$ in the lexicographic preorder, where $e \in C_{n+1}$. Otherwise, we call the chain c *reducible*.

It follows from Proposition 39 that each homology class of the chain complex C_* has a unique lexicographically minimal representative cycle, regardless of the coefficient field.

2.3 Discrete Morse theory and apparent pairs

Following closely the exposition in [9], we use the following generalization of a discrete Morse function, originally due to Forman [6, 28, 29]. Let K be a simplicial complex.

► **Definition 5.** A function $f: K \rightarrow \mathbb{R}$ is a *generalized discrete Morse function* if

- f is *monotonic*, i.e., for any $\sigma, \tau \in K$ with $\sigma \subseteq \tau$ we have $f(\sigma) \leq f(\tau)$, and
- there exists a (unique) partition \hat{V} of K into intervals $[\rho, \phi] = \{\psi \in K \mid \rho \subseteq \psi \subseteq \phi\}$ in the face poset such that any pair of simplices $\sigma \subseteq \tau$ satisfies $f(\sigma) = f(\tau)$ if and only if σ and τ belong to a common interval in the partition.

The collection of regular intervals, $[\rho, \phi]$ with $\rho \neq \phi$, is the discrete gradient V of f on K , and any singleton interval $[\sigma, \sigma] = \{\sigma\}$, as well as the corresponding simplex σ , is *critical*.

If W is another discrete gradient on K , then we say that V is a *refinement* of W if each interval in the gradient partition \hat{W} is a disjoint union of intervals in \hat{V} . If the refinement preserves the set of critical simplices, we call it a *regular refinement*. Moreover, if each regular interval in V only consists of a pair of simplices, we simply call f a *discrete Morse function*. We often refer to a discrete gradient without explicit mention of the function f , noting that different functions can have the same gradient.

For a monotonic function $g: K \rightarrow \mathbb{R}$ we write $S_r(g) = g^{-1}(-\infty, r] \subseteq K$ for the sublevel set and $S_r^<(g) = g^{-1}(-\infty, r) \subseteq K$ for the open sublevel set of g at scale $r \in \mathbb{R}$.

► **Example 6.** For a finite subset $X \subset \mathbb{R}^d$ in general position, the *Čech radius function* $r_\emptyset: \check{\text{Cech}}(X) \rightarrow \mathbb{R}$ as well as the *Delaunay radius function* $r_X: \text{Del}(X) \rightarrow \mathbb{R}$, which assign to a simplex the radius of its smallest enclosing sphere and smallest empty circumsphere, respectively, are both generalized discrete Morse functions [6]. Moreover, for $r \in \mathbb{R}$ their sublevel sets at scale r are the *Čech complex* $\check{\text{Cech}}_r(X)$ and the *Delaunay complex* $\text{Del}_r(X)$, respectively. Similarly, their open sublevel sets at scale r are the *Čech complex* $\check{\text{Cech}}_r^<(X)$ and the *Delaunay complex* $\text{Del}_r^<(X)$, respectively.

We now explain the connection between discrete Morse theory and apparent pairs (Definition 1). Let $f: K \rightarrow \mathbb{R}$ be a monotonic function and assume that the vertices of K are totally ordered. The *f-lexicographic order* is the total order \leq_f on K given by ordering the simplices by their value under f , then by dimension, then by the lexicographic order induced by the total vertex order.

A persistence pair (σ, τ) of the elementwise filtration determined by the f -lexicographic order is a *zero persistence pair* if $f(\sigma) = f(\tau)$. The collection of apparent pairs of a simplexwise filtration forms a discrete gradient [5, Lemma 3.5], the *apparent pairs gradient*. There is a further connection between apparent pairs and discrete Morse functions: Let $f: K \rightarrow \mathbb{R}$ be a generalized discrete Morse function with discrete gradient V . Consider the regular refinement of V obtained by applying a *minimal vertex refinement* to each interval:

$$\tilde{V} = \{(\psi \setminus \{v\}, \psi \cup \{v\}) \mid \psi \in [\rho, \phi] \in V, v = \min(\phi \setminus \rho)\}.$$

By the following proposition from [9, Lemma 9], this regular refinement is induced by the apparent pairs gradient for the simplexwise filtration determined by the f -lexicographic order. In particular, the zero persistence pairs of this simplexwise filtration are precisely the zero persistence apparent pairs, and the V -critical simplices of K are precisely the simplices that are either essential or contained in a non-zero persistence pair.

► **Proposition 7.** *The zero persistence apparent pairs with respect to the f -lexicographic order are precisely the gradient pairs of \tilde{V} .*

3 Descending complexes and gradient refinements

We extend the definition of the *Wrap complex* from [6] to an arbitrary subset C of the set of critical simplices with respect to a discrete gradient V and study its behavior under gradient refinements. We further extend it to monotonic functions $g: K \rightarrow \mathbb{R}$ that are *compatible* with V : all simplices within an interval $I \in \hat{V}$ have the same function value, $g(I) \in \mathbb{R}$.

The gradient partition \hat{V} has a canonical poset structure $\leq_{\hat{V}}$ given by the transitive closure of the relation $I \sim J$ if and only if there exists a face $\sigma \in I$ of a simplex $\tau \in J$. The *down set* of a subset $A \subseteq \hat{V}$ is the set of intervals $\downarrow A = \{I \in \hat{V} \mid \exists J \in A : I \leq_{\hat{V}} J\}$, and for $r \in \mathbb{R}$ we denote the discrete gradient V restricted to the sublevel set $S_r(g)$ by

$V_r = \{I \in V \mid g(I) \leq r\}$. Note that if $I \leq_{\hat{V}} J$, then $g(I) \leq g(J)$, and hence for a subset $A \subseteq \hat{V}_r \subseteq \hat{V}$ the down sets with respect to the canonical poset structure on \hat{V}_r and with respect to the canonical poset structure on \hat{V} coincide.

► **Definition 8.** For a discrete gradient V on K , the descending complex is the subcomplex

$$D(V) = \bigcup \downarrow \{\{\sigma\} \mid \sigma \in K \text{ critical}\}$$

of K given by the union of intervals in the down sets of the critical intervals. More generally, if C is a subset of the set of critical simplices, the descending complex $D(V, C)$ is the subcomplex

$$D(V, C) = \bigcup \downarrow \{\{\sigma\} \mid \sigma \in C\}$$

of K . Moreover, for a monotonic function $g: K \rightarrow \mathbb{R}$ that is compatible with V , the descending complex $D_r(V, g)$ at scale $r \in \mathbb{R}$ is the subcomplex

$$D_r(V, g) = D(V_r) = D(V, \text{Crit}_r(V, g)) = \bigcup \downarrow \{\{\sigma\} \mid \sigma \in K \text{ critical}, g(\sigma) \leq r\}$$

of $S_r(g)$, where $\text{Crit}_r(V, g) = \{\sigma \in K \text{ critical} \mid g(\sigma) \leq r\}$. If V is the discrete gradient of a generalized discrete Morse function f , we simply write $D_r(f)$ for $D_r(V, f)$.

The descending complex $D(V)$ in the context of discrete Morse theory is motivated by the concept of a *descending* or *stable manifold* of a critical point from smooth Morse theory, which is central in the original definition of the Wrap complex [24]. Note that the descending complex $D_r(r_X) \subseteq \text{Del}_r(X)$ of the Delaunay radius function r_X (see Example 6) is precisely the *Wrap complex*, $\text{Wrap}_r(X)$, from [6]. It has been shown that a variant of the Wrap complex can be used to topologically reconstruct a submanifold by choosing a suitable subset of critical simplices [22, 39, 40], which also motivates our definition of $D(V, C)$.

We now study the behavior of the descending complex under gradient refinements.

► **Proposition 9.** Let V be a discrete gradient on K , let C be a subset of V -critical simplices, and let W be a refinement of V . Then the descending complex $D(W, C)$ is a subcomplex of the descending complex $D(V, C)$.

Proof. Note first that every V -critical simplex is also W -critical, as W is a refinement of V . By the same reason, there exists a set map $\varphi: \hat{W} \rightarrow \hat{V}$ between the gradient partitions such that for every $B \in \hat{W}$ we have $B \subseteq \varphi(B)$. It follows straightforwardly from the definition of the canonical poset structures on \hat{W} and \hat{V} that φ is a poset map. Thus, for every W -critical simplex $\sigma \in C$ and interval $A \in \hat{W}$ with $A \leq_{\hat{W}} \{\sigma\}$, we have $\varphi(A) \leq_{\hat{V}} \varphi(\{\sigma\}) = \{\sigma\} \in \hat{V}$, as σ is also V -critical. It now follows directly from the construction of the descending complexes, that $A \subseteq \varphi(A) \subseteq D(V, C)$ and $D(W, C) \subseteq D(V, C)$. ◀

► **Remark 10.** If L is a subcomplex of K and the complement $K \setminus L$ is the disjoint union of regular intervals in V , then V induces a collapse $K \searrow L$ [6, Theorem 2.2]. It follows directly from this and the construction of $D(V)$, that $D(V)$ is the smallest subcomplex of K such that V induces a collapse $K \searrow D(V)$. Moreover, if W is a regular refinement of V , implying $D(W) \subseteq D(V)$ by Proposition 9, the complement $D(V) \setminus D(W)$ is the disjoint union of regular intervals in W . Similar to before, it follows that W induces a collapse $D(V) \searrow D(W)$. In particular, the inclusion $D(W) \hookrightarrow D(V)$ is a homotopy equivalence (see Figure 3).



■ **Figure 3** Left: Generalized discrete gradient (blue) with corresponding descending complex (green). Right: lexicographic gradient refinement (blue) with corresponding descending complex (green).

4 Algebraic Morse theory and persistence

We saw that the apparent pairs are closely related to persistent homology and discrete Morse theory. In this section, we show how all the persistence pairs are related to algebraic Morse theory [30,31,41], also called algebraic discrete Morse theory. We also show how this approach connects to lexicographically minimal cycles. Let (C_*, Σ_*) be a based chain complex.

- **Definition 11.** A function $f: \Sigma_* \rightarrow \mathbb{R}$ is an algebraic Morse function if
- f is monotonic, i.e., for any facet $\sigma \in \Sigma_n$ of $\tau \in \Sigma_{n+1}$ we have $f(\sigma) \leq f(\tau)$, and
 - there exists a (unique) disjoint collection V of facet pairs such that every facet pair (μ, η) satisfies $f(\mu) = f(\eta)$ if and only if $(\mu, \eta) \in V$.

We call V the algebraic gradient of f on Σ_* , and a basis element critical if it is not contained in any pair of V . Moreover, for $(\sigma, \tau) \in V$ we call σ a gradient facet and τ a gradient cofacet.

We often refer to an algebraic gradient without explicit mention of the associated function.

► **Remark 12.** The definitions of algebraic Morse function and algebraic gradient generalize those from discrete Morse theory. Let $f: K \rightarrow \mathbb{R}$ be a discrete Morse function with discrete gradient V . Recall that the simplicial chain complex $C_*(K)$ has a basis Σ_* consisting of the simplices of K with some chosen orientation. We can now interpret f as an algebraic Morse function on this basis Σ_* and the discrete gradient V as an algebraic gradient.

4.1 Gradient pairs from persistence pairs

We now explain how all persistence pairs determine an algebraic gradient that relates to discrete Morse theory through apparent pairs (Lemma 2 and Proposition 7). This establishes the framework for a key step in our proof of Theorem A. Let $(C_*, \Sigma_* = \sigma_1 < \dots < \sigma_l)$ be a based chain complex with an elementwise filtration, and let $R = D \cdot S$ be a reduction of the filtration boundary matrix. For any chain $c \in C_n$, we denote by $\text{Pivot}_{\Sigma_*} c = \max \text{supp}_{\Sigma_*} c$ the maximal basis element in the basis representation of c with respect to Σ_* . If v is the coordinate vector in \mathbb{R}^l representing c , we also write $\text{Pivot}_{\Sigma_*} v$ for $\text{Pivot}_{\Sigma_*} c = \sigma_{\text{PivInd } v}$.

The direct sum decomposition of filtered chain complexes explained in Section 2.1 yields a straightforward interpretation of persistence pairs as an algebraic gradient, which we discuss in Appendix A.1. However, this gradient is not suitable for our purposes as it neither relates directly to apparent pairs nor lexicographically optimal cycles. Hence, we consider an alternative approach that uses the distinctness of non-zero pivot elements in the reduced matrix $R = D \cdot S$. To this end, we equip the chain complex C_* with the new ordered basis $\Omega_* = \tau_1 < \dots < \tau_l$ given by

$$\tau_i = \begin{cases} \sigma_i & \text{if } i \text{ is a birth or essential index,} \\ S_i & \text{if } i \text{ is a death index.} \end{cases}$$

We call this basis Ω_* the *reduction basis*. Note that with respect to the original basis we have $\text{Pivot}_{\Sigma_*} \tau_i = \sigma_i$ for all i . Moreover, note that for every death index j and $R_j = D \cdot S_j$ we have $\text{Pivot}_{\Omega_*} R_j = \text{Pivot}_{\Sigma_*} R_j$. By pairing the death columns S_j with the pivot elements $\text{Pivot}_{\Omega_*} R_j$ of their boundaries $R_j = D \cdot S_j$, we obtain a set of disjoint pairs, which we call the *reduction gradient* of S :

$$\{(\text{Pivot}_{\Omega_*} R_j, S_j) \mid j \text{ is a death index}\}.$$

► **Proposition 13.** *The reduction gradient is an algebraic gradient on Ω_* .*

Proof. Consider the function $f: \Omega_* \rightarrow \mathbb{N}$ with values $f(\text{Pivot}_{\Omega_*} R_j) = f(S_j) = \text{PivInd } R_j$ for every death index j and $f(\tau_i) = i$ for every essential index i . Note that, as S is a full rank upper triangular matrix, the ordered basis $\Omega_* = \tau_1 < \dots < \tau_l$ is compatible with the given elementwise filtration, in the sense that $\downarrow \tau_j$ induces the same subcomplex of C_* as $\downarrow \sigma_j$ for every j . In particular, whenever τ_i is a facet of τ_j , we have $i < j$. The function f assigns to any basis element of the form $\tau_j = S_j$, for j a death index, the index $i = \text{PivInd } R_j$ of its maximal facet $\text{Pivot}_{\Omega_*} R_j = \text{Pivot}_{\Sigma_*} R_j = \tau_i \in \Omega_*$, and to all other basis elements τ_i their index i . This implies that f is monotonic with algebraic gradient the reduction gradient. ◀

The reduction gradient is closely related to apparent pairs (Definition 1) in the following sense. For any apparent pair (σ_i, σ_j) of the elementwise filtration, the column of σ_j in the filtration boundary matrix is reduced (Lemma 2). Therefore, we are led to define a reduction matrix S to be *apparent pairs compatible* if the column S_j contains only one non-zero entry $S_{j,j} = 1$ for every apparent pair (σ_i, σ_j) . Note that the collection of apparent pairs of the elementwise filtration forms an algebraic gradient on Σ_* [33, Lemma 2.2; 5, Lemma 3.5] that we call the *apparent pairs gradient*. The following is a direct consequence of the definitions.

► **Lemma 14.** *If the reduction matrix S is apparent pairs compatible, then the apparent pairs gradient of the elementwise filtration on Σ_* is a subset of the reduction gradient of S on Ω_* .*

► **Remark 15.** Both of the reduction algorithms in Section 2.1 compute a reduction $R = D \cdot S$ of the filtration boundary matrix, noting explicitly that S is homogeneous. Furthermore, Algorithm 1 computes a reduction matrix S that is also apparent pairs compatible.

4.2 The flow of an algebraic gradient

We now introduce the flow determined by an algebraic gradient, study its behavior under gradient containment, and analyze in the subsequent sections its relation to lexicographically minimal cycles (Proposition 26), as well as the descending complex (Proposition 28). While the remainder of the paper is mainly focused on cycles, in this section we present results that hold more generally for chains, and which may be of independent interest. Moreover, in Appendix A.4 we demonstrate how the algebraic flow on a cycle can be interpreted as a variant of Gaussian elimination, tying it closely to the exhaustive reduction (Algorithm 2).

Let (C_*, Σ_*) be a based chain complex and V an algebraic gradient on Σ_* . The following definition, originally for discrete gradients [28], carries over naturally to the algebraic setting.

► **Definition 16.** *The flow $\Phi: C_* \rightarrow C_*$ determined by V is the chain map given by*

$$\Phi(c) = c + \partial F(c) + F(\partial c),$$

where $F: C_* \rightarrow C_{*+1}$ is the unique linear map defined on the basis elements $\sigma \in \Sigma_*$ as

$$F(\sigma) = \begin{cases} -\frac{1}{\langle \partial \tau, \sigma \rangle} \cdot \tau & \text{if } \sigma \text{ is contained in a pair } (\sigma, \tau) \in V, \\ 0 & \text{otherwise.} \end{cases}$$

Note that, by construction, the map F is a chain homotopy between the identity and the flow Φ . In particular, if c is a cycle, then the flow reduces to $\Phi(c) = c + \partial F(c)$ and therefore acts on each homology class of the chain complex by a change of representative cycle. Moreover, if $f: \Sigma_* \rightarrow \mathbb{R}$ is an algebraic Morse function, then, by construction, the associated flow Φ acts for any $r \in \mathbb{R}$ on the subcomplex of C_* spanned by the sublevel set $S_r(f) = f^{-1}(-\infty, r]$.

Forman [28] proved that the sequence $(\Phi^n)_n$ stabilizes for the cellular chain complex of a finite CW-complex. This generalizes to our setting of arbitrary finite chain complexes, and we denote the *stabilized flow* by $\Phi^\infty = \Phi^r$, where r is a large enough natural number:

► **Proposition 17.** *There exists an $r \in \mathbb{N}$ such that for all $s \geq r$ we have $\Phi^s = \Phi^r$.*

The proof of this proposition uses the following lemma, which is a straightforward generalization of the corresponding statement in [28], and is of independent interest to us.

► **Lemma 18.** *Let $\sigma \in \Sigma_n$ be critical. Then for all $r \in \mathbb{N}$ the iterated flow $\Phi^{r+1}(\sigma)$ is given by*

$$\Phi^{r+1}(\sigma) = \sigma + w + \Phi(w) + \cdots + \Phi^r(w),$$

where $w = F(\partial\sigma)$. Moreover, we have $\Phi^s(w) \in \text{im } F$ for all $s \in \mathbb{N}$, and $F(\Phi^{r+1}(\sigma)) = 0$.

Based on the concept of a discrete flow, Forman [28] considered the subcomplex of the cellular chain complex given by the flow invariant chains and proved the following proposition, which generalizes to our setting. Consider the subcomplex of Φ -invariant chains,

$$C_*^\Phi = \{c \in C_* \mid \Phi(c) = c\}.$$

► **Proposition 19.** *The Φ -invariant chains are spanned by the image of the critical basis elements under the stabilized flow $C_*^\Phi = \text{span}\{\Phi^\infty(\sigma) \mid \sigma \in \Sigma_n \text{ critical}\}$.*

We now relate the flow invariant chains of an algebraic gradient to those of its subgradients. A proof of the following statement can be found in Appendix A.2.

► **Proposition 20.** *Let (C_*, Σ_*) be a based chain complex and $W \subseteq V$ two algebraic gradients on Σ_* . Consider the flows $\Psi, \Phi: C_* \rightarrow C_*$ determined by W and V , respectively. Then any Φ -invariant chain is also Ψ -invariant, i.e., we have $C_*^\Phi \subseteq C_*^\Psi$.*

Note that the flow Φ can be written as a sum of flows $\Phi = \sum_{(a,b) \in V} \Phi^{(a,b)} - (\text{card } V - 1) \cdot \text{id}$, where $\Phi^{(a,b)}$ is the flow determined by the algebraic gradient $\{(a,b)\}$ on Σ_* . Together with Proposition 20, this proves the following.

► **Corollary 21.** *Let V be an algebraic gradient on Σ_* with associated flow $\Phi: C_* \rightarrow C_*$. Then a chain is Φ -invariant if and only if it is $\Phi^{(a,b)}$ -invariant for every pair $(a,b) \in V$.*

For a based chain complex (C_*, Σ_*) and algebraic gradient V on Σ_* , we denote by $\partial_n V = \{\partial b \mid \exists (a,b) \in V \text{ with } a \in \Sigma_n\}$ the set of gradient cofacet boundaries in degree n . We say that two cycles $z, z' \in Z_n$ are V -homologous if there exists an element $\partial e \in \text{span } \partial_n V$ such that $z - z' = \partial e$. Observe that for any cycle $z \in Z_n$, the cycle $\Phi(z) = z + \partial F(z)$ is V -homologous to z . A proof of the following statement can be found in Appendix A.2.

► **Proposition 22.** *Let V be an algebraic gradient on Σ_* with associated flow $\Phi: C_* \rightarrow C_*$. Then a cycle $z \in Z_n$ is Φ -invariant if and only if it contains no gradient facets of V . Moreover, if $z' \in Z_n$ is any Φ -invariant cycle that is V -homologous to z , then $z = z'$.*

Now let $(C_*, \Sigma_* = \sigma_1 < \dots < \sigma_l)$ be a based chain complex with an elementwise filtration, and let $R = D \cdot S$ be a reduction of the filtration boundary matrix. In order to relate the flow determined by the reduction gradient to the flow determined by the zero persistence apparent pairs gradient, and therefore to discrete Morse theory (Propositions 7 and 28), we call a reduction matrix S *death-compatible* if for every death index j and non-zero entry $S_{i,j} \neq 0$ we also have that i is a death index.

► **Remark 23.** Both algorithms in Section 2.1 compute a death-compatible reduction matrix S .

The following is a direct consequence of Lemma 14 and the definitions.

► **Lemma 24.** *If the reduction matrix S is apparent pairs and death-compatible, then the flows determined by the apparent pairs gradient of the elementwise filtration as an algebraic gradient on Σ_* and as an algebraic gradient on Ω_* , respectively, coincide.*

4.3 Relating the algebraic flow and lexicographically minimal cycles

We now relate the flow invariant cycles determined by a reduction gradient to lexicographically minimal homologous cycles. Let $(C_*, \Sigma_* = \sigma_1 < \dots < \sigma_l)$ be a based chain complex with an elementwise filtration, and let $R = D \cdot S$ be a reduction of the filtration boundary matrix. The following result [18, Lemmas 3.3 and 3.5] provides an equivalent condition for minimality.

► **Proposition 25.** *A cycle is lexicographically minimal with respect to the elementwise filtration if and only if its support contains only death elements and essential elements.*

We provide another characterization in terms of the algebraic flow determined by the reduction gradient of S (defined on the reduction basis Ω_*), which we call the *reduction flow*.

► **Proposition 26.** *Let $(C_*, \Sigma_* = \sigma_1 < \dots < \sigma_l)$ be a based chain complex with an elementwise filtration, and let $R = D \cdot S$ be a reduction of the filtration boundary matrix by a death-compatible reduction matrix S . Then for a cycle $z \in Z_n$ the following are equivalent:*


- z is lexicographically minimal with respect to the ordered basis Σ_* ;
- z is invariant under the reduction flow.

We defer the proofs to Appendix A.3.

5 Relating algebraic reduction gradients and discrete gradients

We are now ready to relate our results for the algebraic flow determined by the reduction gradient to discrete Morse theory. In particular, we show that for a generalized discrete Morse function $f: K \rightarrow \mathbb{R}$, the lexicographically minimal cycles with respect to the f -lexicographic order on the simplices are supported on the descending complexes of f . In Appendix B, we show that the reduction chains, given as the columns of a suitable reduction matrix, are also supported on the descending complexes of f . In particular, this is true of the columns of the totally reduced filtration boundary matrix, considered as cycles in the sublevel set corresponding to the pivot index. This summarizes the relations between reductions of the filtration boundary matrix and the descending complexes of f . Recall that a discrete gradient V on K gives rise to a flow $\Phi: C_*(K) \rightarrow C_*(K)$.

► **Lemma 27.** *The flow Φ restricts to a chain map on the descending complex $D(V) \subseteq K$, i.e., for $c \in C_*(D(V))$ we also have $\Phi(c) \in C_*(D(V))$.*



■ **Figure 4** Discrete gradient (blue) with corresponding descending complex (green). Left: Cycle c (red). Right: Stabilized cycle $\Phi(c) = \Phi^\infty(c)$ (red), supported on the descending complex (green).

Proof. Recall from Definition 16 that Φ is given by $\Phi(c) = c + \partial F(c) + F(\partial c)$, where $F: C_*(K) \rightarrow C_{*+1}(K)$ is the linear map with $F(\sigma) = -\langle \partial\tau, \sigma \rangle^{-1} \cdot \tau$ if $(\sigma, \tau) \in V$ and 0 on all other simplices. Thus, if $\eta \in D(V)$ is any simplex, then $\partial\eta$ and $F(\eta)$ are both contained in $C_*(D(V))$, by definition of the descending complex. Therefore, if $c \in C_*(D(V))$ is any chain, then the chains ∂c , $F(c)$, and $F(\partial c)$ are also contained in $C_*(D(V))$. This shows that the chain $\Phi(c) = c + \partial F(c) + F(\partial c)$ is contained in $C_*(D(V))$, proving the claim. ◀

The Φ -invariant chains are supported on the descending complex, as illustrated in Figure 4.

► **Proposition 28.** *The Φ -invariant chains of $C_*(K)$ are supported on the descending complex $D(V)$, i.e., we have $C_*^\Phi(K) \subseteq C_*(D(V))$.*

Proof. Let $c \in C_*^\Phi(K)$ be any Φ -invariant chain; we show that c is contained in $C_*(D(V))$. By Proposition 19, we can assume without loss of generality that c is of the form $\Phi^\infty(\sigma)$ for a critical simplex σ . By definition, $\Phi^\infty = \Phi^r$ for a large enough r , and by Lemma 18, $\Phi^r(\sigma)$ is given by $\Phi^r(\sigma) = \sigma + w + \Phi(w) + \dots + \Phi^{r-1}(w)$, where $w = F(\partial\sigma)$. It follows directly from the definition of the descending complex (Definition 8) that σ , $\partial\sigma$, and $w = F(\partial\sigma)$ are contained in $C_*(D(V))$. By Lemma 27, we have $\Phi^k(w) \in C_*(D(V))$ for every k , and therefore $c = \Phi^r(\sigma) = \sigma + w + \Phi(w) + \dots + \Phi^{r-1}(w) \in C_*(D(V))$, proving the claim. ◀

The following, together with Example 6, directly implies Theorem A.

► **Theorem 29.** *Let f be a generalized discrete Morse function, let $r \in \mathbb{R}$, and let $h \in H_*(S_r(f))$ be a homology class of the sublevel set $S_r(f)$. Then the lexicographically minimal cycle of h , with respect to the f -lexicographic order, is supported on the descending complex $D_r(f)$.*

Proof. Let V be the discrete gradient of f , and let W be the zero persistence apparent pairs gradient induced by the f -lexicographic order, which is a regular refinement of V by Proposition 7. Note that f is compatible with W .

Recall from Remarks 15 and 23 that Algorithm 1 computes a reduction $R = D \cdot S$ of the filtration boundary matrix, corresponding to the simplexwise filtration of $S_r(f)$ induced by the f -lexicographic order, such that the reduction matrix S is apparent pairs compatible and also death-compatible. Consider the corresponding reduction gradient on the corresponding reduction basis Ω_* with associated reduction flow $\Psi: C_*(S_r(f)) \rightarrow C_*(S_r(f))$.

The lexicographically minimal cycle $z \in Z_*(S_r(f))$ of h is a Ψ -invariant cycle according to Proposition 26. We show that z is contained in $C_*(D_r(f))$, which proves the claim: As S is apparent pairs compatible, it follows from Lemma 14 and Proposition 20 that z is also invariant under the algebraic flow determined by the zero persistence apparent pairs gradient W_r on Ω_* . Lemma 24 implies that this flow coincides with the algebraic flow $\Phi: C_*(S_r(f)) \rightarrow C_*(S_r(f))$ determined by the zero persistence apparent pairs gradient W_r .

on the standard basis given by the simplices of $S_r(f)$. It now follows from Proposition 28, the definition of descending complex (Definition 8), and Proposition 9 that z is contained in

$$C_*^\Phi(S_r(f)) \subseteq C_*(D(W_r)) = C_*(D_r(W, f)) \subseteq C_*(D_r(V, f)) = C_*(D_r(f)). \quad \blacktriangleleft$$

Let D be the filtration boundary matrix of the simplexwise filtration of K induced by the f -lexicographic order, and let $R = D \cdot S$ be a totally reduced reduction of D . The following, together with Example 6, directly implies Corollary B.

► **Corollary 30.** *Let f be a generalized discrete Morse function and (σ, τ) a non-zero persistence pair of the simplexwise filtration induced by the f -lexicographic order. Let $r = f(\sigma)$ and $s = f(\tau)$ be the function values of σ and of τ , respectively. Then the lexicographically minimal cycle of $[\partial\tau]$ in the open sublevel set $S_s^<(f)$, given as the column R_τ of the totally reduced filtration boundary matrix, is supported on the descending complex $D_r(f)$.*

Proof. As (σ, τ) is a non-zero persistence pair, we know that $f(\sigma) < f(\tau) = s$ and that τ is a critical simplex. As R is a reduction of D , this implies that R_τ and $\partial\tau$ are homologous cycles in $S_s^<(f)$. Since R is totally reduced, the cycle R_τ does not contain a (non-essential) birth simplex of the (smaller) simplexwise filtration of $S_s^<(f)$ induced by the f -lexicographic order. Thus, Proposition 25 implies that R_τ is the lexicographically minimal cycle of $[\partial\tau]$ in $S_s^<(f)$. As $r = f(\sigma)$ and R is a reduction of D , the cycle R_τ is supported on the subcomplex $S_r(f)$ of $S_s^<(f)$, implying that it is also a lexicographically minimal cycle in $S_r(f)$. It now follows from Theorem 29 that the cycle R_τ is supported on the descending complex $D_r(f)$. ◀

References

- 1 Nina Amenta, Sunghee Choi, Tamal K. Dey, and Naveen Leekha. A simple algorithm for homeomorphic surface reconstruction. *Internat. J. Comput. Geom. Appl.*, 12(1-2):125–141, 2002. 16th Annual Symposium on Computational Geometry (Kowloon, 2000). doi:10.1142/S0218195902000773.
- 2 Nina Amenta, Sunghee Choi, and Ravi Krishna Kolluri. The power crust, unions of balls, and the medial axis transform. *Comput. Geom.*, 19(2-3):127–153, 2001. Combinatorial curves and surfaces. doi:10.1016/S0925-7721(01)00017-7.
- 3 Dominique Attali, Hana Dal Poz Kouřimská, Christopher Fillmore, Ishika Ghosh, André Lieutier, Elizabeth Stephenson, and Mathijs Wintraecken. Tight bounds for the learning of homotopy à la Niyogi, Smale, and Weinberger for subsets of Euclidean spaces and of Riemannian manifolds. In *40th International Symposium on Computational Geometry (SoCG 2024)*, volume 224 of *Leibniz International Proceedings in Informatics (LIPIcs)*, 2024. arXiv:2206.10485.
- 4 Dominique Attali and André Lieutier. Delaunay-like triangulation of smooth orientable submanifolds by ℓ_1 -norm minimization. In *38th International Symposium on Computational Geometry*, volume 224 of *LIPIcs. Leibniz Int. Proc. Inform.*, pages Art. No. 8, 16. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2022. doi:10.4230/lipics.socg.2022.8.
- 5 Ulrich Bauer. Ripser: efficient computation of Vietoris-Rips persistence barcodes. *J. Appl. Comput. Topol.*, 5(3):391–423, 2021. doi:10.1007/s41468-021-00071-5.
- 6 Ulrich Bauer and Herbert Edelsbrunner. The Morse theory of Čech and Delaunay complexes. *Trans. Amer. Math. Soc.*, 369(5):3741–3762, 2017. doi:10.1090/tran/6991.
- 7 Ulrich Bauer, Michael Kerber, Fabian Roll, and Alexander Rolle. A unified view on the functorial nerve theorem and its variations. *Expo. Math.*, 41(4):125503, 2023. doi:10.1016/j.exmath.2023.04.005.
- 8 Ulrich Bauer, Carsten Lange, and Max Wardetzky. Optimal topological simplification of discrete functions on surfaces. *Discrete Comput. Geom.*, 47(2):347–377, 2012. doi:10.1007/s00454-011-9350-z.

- 9 Ulrich Bauer and Fabian Roll. Gromov Hyperbolicity, Geodesic Defect, and Apparent Pairs in Vietoris-Rips Filtrations. In *38th International Symposium on Computational Geometry (SoCG 2022)*, volume 224 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 15:1–15:15, 2022. doi:10.4230/LIPIcs.SoCG.2022.15.
- 10 Ulrich Bauer and Fabian Roll. WrappingCycles, 2023. Executable directly by running `docker build -o output github.com/fabian-roll/wrappingcycles` on any machine with Docker installed and configured with sufficient memory (16GB recommended). URL: <https://github.com/fabian-roll/wrappingcycles>.
- 11 Jean-Daniel Boissonnat, Frédéric Chazal, and Mariette Yvinec. *Geometric and topological inference*. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 2018. doi:10.1017/9781108297806.
- 12 Jean-Daniel Boissonnat and Arijit Ghosh. Manifold reconstruction using tangential Delaunay complexes. *Discrete Comput. Geom.*, 51(1):221–267, 2014. doi:10.1007/s00454-013-9557-2.
- 13 Siu-Wing Cheng, Tamal K. Dey, Herbert Edelsbrunner, Michael A. Facello, and Shang-Hua Teng. Sliver exudation. *J. ACM*, 47(5):883–904, 2000. doi:10.1145/355483.355487.
- 14 Siu-Wing Cheng, Tamal K. Dey, and Edgar A. Ramos. Manifold reconstruction from point samples. In *Proceedings of the Sixteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1018–1027. ACM, New York, 2005. doi:10.5555/1070432.1070579.
- 15 David Cohen-Steiner, Herbert Edelsbrunner, and John Harer. Stability of persistence diagrams. *Discrete Comput. Geom.*, 37(1), 2007. doi:10.1007/s00454-006-1276-5.
- 16 David Cohen-Steiner, Herbert Edelsbrunner, and Dmitriy Morozov. Vines and vineyards by updating persistence in linear time. In *Computational geometry (SCG’06)*, pages 119–126. ACM, New York, 2006. doi:10.1145/1137856.1137877.
- 17 David Cohen-Steiner, André Lieutier, and Julien Vuillamy. Lexicographic optimal chains and manifold triangulations. Research report, INRIA Sophia-Antipolis Méditerranée, December 2019. URL: <https://hal.science/hal-02391190>.
- 18 David Cohen-Steiner, André Lieutier, and Julien Vuillamy. Lexicographic optimal homologous chains and applications to point cloud triangulations. *Discrete Comput. Geom.*, 68(4):1155–1174, 2022. Special Issue: 36th Annual Symposium on Computational Geometry (SoCG 2020). doi:10.1007/s00454-022-00432-6.
- 19 David Cohen-Steiner, André Lieutier, and Julien Vuillamy. Delaunay and regular triangulations as lexicographic optimal chains. *Discrete Comput. Geom.*, 70(1):1–50, 2023. doi:10.1007/s00454-023-00485-1.
- 20 Vin de Silva, Dmitriy Morozov, and Mikael Vejdemo-Johansson. Dualities in persistent (co)homology. *Inverse Problems*, 27(12):124003, 17, 2011. doi:10.1088/0266-5611/27/12/124003.
- 21 Tamal K. Dey. *Curve and surface reconstruction: algorithms with mathematical analysis*, volume 23 of *Cambridge Monographs on Applied and Computational Mathematics*. Cambridge University Press, Cambridge, 2007. doi:10.1017/CB09780511546860.
- 22 Tamal K. Dey, Joachim Giesen, Edgar A. Ramos, and Bardia Sadri. Critical points of the distance to an epsilon-sampling of a surface and flow-complex-based surface reconstruction. In *Computational geometry (SCG’05)*, pages 218–227. ACM, New York, 2005. doi:10.1145/1064092.1064126.
- 23 Herbert Edelsbrunner. The union of balls and its dual shape. *Discrete Comput. Geom.*, 13(3-4):415–440, 1995. doi:10.1007/BF02574053.
- 24 Herbert Edelsbrunner. Surface reconstruction by wrapping finite sets in space. In *Discrete and computational geometry*, volume 25 of *Algorithms Combin.*, pages 379–404. Springer, Berlin, 2003. doi:10.1007/978-3-642-55566-4_17.
- 25 Herbert Edelsbrunner and John L. Harer. *Computational topology*. American Mathematical Society, Providence, RI, 2010. An introduction. doi:10.1090/mbk/069.
- 26 Herbert Edelsbrunner and Katharina Ölsböck. Tri-partitions and bases of an ordered complex. *Discrete Comput. Geom.*, 64(3):759–775, 2020. doi:10.1007/s00454-020-00188-x.

- 27 Herbert Edelsbrunner and Afra Zomorodian. Computing linking numbers of a filtration. *Homology Homotopy Appl.*, 5(2):19–37, 2003. Algebraic topological methods in computer science (Stanford, CA, 2001). doi:10.4310/HHA.2003.v5.n2.a2.
- 28 Robin Forman. Morse theory for cell complexes. *Adv. Math.*, 134(1):90–145, 1998. doi:10.1006/aima.1997.1650.
- 29 Ragnar Freij. Equivariant discrete Morse theory. *Discrete Math.*, 309(12):3821–3829, 2009. doi:10.1016/j.disc.2008.10.029.
- 30 Michael Jöllenbeck and Volkmar Welker. Minimal resolutions via algebraic discrete Morse theory. *Mem. Amer. Math. Soc.*, 197(923):vi+74, 2009. doi:10.1090/memo/0923.
- 31 Dmitry Kozlov. *Combinatorial algebraic topology*, volume 21 of *Algorithms and Computation in Mathematics*. Springer, Berlin, 2008. doi:10.1007/978-3-540-71962-5.
- 32 Stanford Computer Graphics Laboratory. The Stanford 3D Scanning Repository. URL: <http://graphics.stanford.edu/data/3Dscanrep/>.
- 33 Leon Lampret. Chain complex reduction via fast digraph traversal. Preprint, 2020. arXiv:1903.00783.
- 34 S. Lefschetz. *Algebraic Topology*. AMS books online. American Mathematical Society, 1942.
- 35 Konstantin Mischaikow and Vidit Nanda. Morse theory for filtrations and efficient computation of persistent homology. *Discrete Comput. Geom.*, 50(2):330–353, 2013. doi:10.1007/s00454-013-9529-6.
- 36 James R. Munkres. *Elements of algebraic topology*. Addison-Wesley Publishing Company, Menlo Park, CA, 1984.
- 37 Partha Niyogi, Stephen Smale, and Shmuel Weinberger. Finding the homology of submanifolds with high confidence from random samples. *Discrete Comput. Geom.*, 39(1-3):419–441, 2008. doi:10.1007/s00454-008-9053-2.
- 38 Cédric Portaneri, Mael Rouxel-Labbé, Michael Hemmer, David Cohen-Steiner, and Pierre Alliez. Alpha wrapping with an offset. *ACM Trans. Graph.*, 41(4):127:1–127:22, 2022. doi:10.1145/3528223.3530152.
- 39 Edgar A. Ramos and Bardia Sadri. Geometric and topological guarantees for the WRAP reconstruction algorithm. In *Proceedings of the Eighteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1086–1095. ACM, New York, 2007.
- 40 Bardia Sadri. Manifold homotopy via the flow complex. *Comput. Graph. Forum*, 28(5):1361–1370, 2009. URL: <https://doi.org/10.1111/j.1467-8659.2009.01513.x>, doi:10.1111/J.1467-8659.2009.01513.X.
- 41 Emil Sköldberg. Morse theory from an algebraic viewpoint. *Trans. Amer. Math. Soc.*, 358(1):115–129, 2006. doi:10.1090/S0002-9947-05-04079-1.
- 42 Julien Vuillamy. *Planimetric simplification and lexicographic optimal chains for 3D urban scene reconstruction*. Theses, Université Côte d’Azur, September 2021. URL: <https://hal.science/tel-03339931>.
- 43 Charles A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1994. doi:10.1017/CB09781139644136.
- 44 Afra Zomorodian and Gunnar Carlsson. Computing persistent homology. *Discrete Comput. Geom.*, 33(2):249–274, 2005. doi:10.1007/s00454-004-1146-y.

A Algebraic Morse theory and persistence

A.1 Gradient pairs from persistence pairs

In this section, we discuss how the direct sum decomposition of filtered chain complexes yields a straightforward interpretation of persistence pairs as an algebraic gradient, which was mentioned in Section 4.1.

Consider the direct sum decomposition explained in Section 2.1 and equip the chain complex C_* with the new ordered basis $E_* = \eta_1 < \dots < \eta_l$ given by

$$\eta_i = \begin{cases} R_j & \text{if there exists an index persistence pair } (i, j), \\ S_i & \text{if } i \text{ is a death or essential index.} \end{cases}$$

We call E_* the *decomposition basis*, noting that it induces a decomposition of the filtered chain complex as described in Section 2.1. Note that with respect to the original basis we have $\text{Pivot}_{\Sigma_*} \eta_i = \sigma_i$ for all i . By pairing the death columns S_j with their boundaries $R_j = D \cdot S_j$, we obtain a set of disjoint pairs that we call the *decomposition gradient* of S :

$$\{(R_j, S_j) \mid j \text{ is a death index}\}.$$

► **Proposition 31.** *The decomposition gradient is an algebraic gradient on the basis E_* .*

Proof. Consider the function $f: E_* \rightarrow \mathbb{N}$ with values $f(R_j) = f(S_j) = j$ for every death index j and $f(S_i) = i$ for every essential index i . Note first that for any death index j , the basis element R_j is the only facet of S_j , and R_j has no facets as it is a cycle. Similarly, for any essential index i , the basis element S_i has no facets, as well. Thus, it is immediate that f is a monotonic function whose algebraic gradient is the decomposition gradient. ◀

A.2 The flow of an algebraic gradient

In this section, we provide the relegated proofs for the statements in Section 4.2.

Proof of Proposition 20. Let $c \in C_n$ be a Φ -invariant chain. Recall from Definition 16 that Φ is given by $\Phi(c) = c + \partial F(c) + F(\partial c)$, where $F: C_* \rightarrow C_{*+1}$ is the linear map with $F(\sigma) = -\langle \partial \tau, \sigma \rangle^{-1} \cdot \tau$ if $(\sigma, \tau) \in V$ and 0 on all other basis elements of Σ_* . By Proposition 19 and linearity, we can assume without loss of generality that c is of the form $\Phi^\infty(\eta)$ for a V -critical basis element $\eta \in \Sigma_n$. By definition, $\Phi^\infty = \Phi^{r+1}$ for a large enough $r \in \mathbb{N}$, and by Lemma 18, the chain $c = \Phi^{r+1}(\eta)$ satisfies $F(c) = F(\Phi^{r+1}(\eta)) = 0$. Hence, we have

$$c = \Phi(c) = c + F(\partial c) + \partial F(c) = c + F(\partial c)$$

from which we conclude that $F(\partial c) = 0$ holds. By construction of F , this implies that c and ∂c are the sums of critical elements and gradient cofacets of V .

Similarly to before, Ψ is given by $\Psi(c) = c + \partial E(c) + E(\partial c)$, where $E: C_* \rightarrow C_{*+1}$ is the linear map with $E(\sigma) = -\langle \partial \tau, \sigma \rangle^{-1} \cdot \tau$ if $(\sigma, \tau) \in W$ and 0 on all other basis elements of Σ_* . As $W \subseteq V$, it holds true that c and ∂c are the sums of critical elements and gradient cofacets of W , as well. By construction of E , this implies $E(c) = E(\partial c) = 0$, and hence

$$\Psi(c) = c + \partial E(c) + E(\partial c) = c + 0 + 0 = c,$$

meaning that c is Ψ -invariant, proving the claim. ◀

The following statement is a direct consequence of the definitions.

► **Lemma 32.** *Let (C_*, Σ_*) be a based chain complex and let $\sigma_1 < \dots < \sigma_l$ be any total order on the basis Σ_* corresponding to an elementwise filtration that refines the sublevel set filtration induced by an algebraic Morse function f . Then, with respect to this total order, for any gradient pair $(a, b) \in V$ of f the pivot of ∂b is equal to a , $\text{Pivot}_{\Sigma_*} \partial b = a$.*

Let (C_*, Σ_*) be a based chain complex and V an algebraic gradient on Σ_* . Let L_n be the linear subspace of C_n spanned by the gradient facets of V and consider the canonical linear projection $\pi_n: \text{span } \partial_n V \rightarrow L_n$ with respect to the basis Σ_n .

► **Lemma 33.** *The subset $\partial_n V = \{\partial b \mid \exists (a, b) \in V \text{ with } a \in \Sigma_n\} \subseteq C_n$ is linearly independent, and the canonical linear projection $\pi_n: \text{span } \partial_n V \rightarrow L_n$ is an isomorphism.*

Proof. By Lemma 32, we can take a total order $\sigma_1 < \dots < \sigma_l$ on the basis Σ_* such that for any pair $(a, b) \in V$ the pivot of ∂b is equal to a , $\text{Pivot}_{\Sigma_*} \partial b = a$. As the pairs in V are disjoint, we know that the pivots $\text{Pivot}_{\Sigma_*} \partial b = a$ for $(a, b) \in V$ are distinct. In particular, this implies that the subset $\partial_n V$ is linearly independent.

To prove the second claim, note that, similarly to before, the pivots $\text{Pivot}_{\Sigma_*} \pi_n(\partial b) = a$ for $(a, b) \in V$ with $a \in \Sigma_n$ are distinct. This implies that the linearly independent vectors $\partial_n V$ are sent to linearly independent vectors in L_n . As $\text{span } \partial_n V$ and L_n have the same dimension $\text{card}(\{(a, b) \in V \mid a \in \Sigma_n\})$, this shows that π_n is an isomorphism. ◀

Proof of Proposition 22. We start by proving the first claim. On the one hand, we know from Proposition 19 and Lemma 18, that a Φ -invariant chain contains no gradient facets of V . On the other hand, if a cycle contains no gradient facets of V , then it is $\Phi^{(a,b)}$ -invariant for every pair $(a, b) \in V$, and it follows from Corollary 21 that it is also Φ -invariant.

To prove the second claim, let $z' \in Z_n$ be any Φ -invariant cycle that is V -homologous to z . Then by definition, there exists a $\partial e \in \text{span } \partial_n V$ with $z - z' = \partial e$. Recall from Lemma 33, that the projection $\pi_n: \text{span } \partial_n V \rightarrow L_n$ onto the subspace of C_n spanned by the gradient facets of V is an isomorphism. By assumption, the difference $z - z' = \partial e$ is Φ -invariant, and therefore it contains no gradient facets of V , by our first claim. Thus, we have $\pi_n(\partial e) = 0$, implying $z - z' = \partial e = 0$, and proving the second claim $z = z'$. ◀

Let $(C_*, \Sigma_* = \sigma_1 < \dots < \sigma_l)$ be a based chain complex with an elementwise filtration. Motivated by Lemma 32, we make the following definition.

► **Definition 34.** *We call an algebraic gradient V on Σ_* reduced if for all pairs $(a, b) \in V$ we have $\text{Pivot}_{\Sigma_*} \partial b = a$, and we call it reduced in degree n , or n -reduced, if this holds for all pairs $(a, b) \in V$ with $a \in \Sigma_n$.*

Motivated by the notion of two cycles being V -homologous, and by Proposition 22, we make the following definition.

► **Definition 35.** *We say that an algebraic gradient V on Σ_* generates the n -boundaries if the set of gradient cofacet boundaries $\partial_n V$ generates the entire subspace of boundaries $B_n \subseteq C_n$.*

► **Remark 36.** By construction, any reduction gradient is n -reduced and generates the n -boundaries for all n .

A.3 Relating the algebraic flow and lexicographically minimal cycles

In this section, we prove Propositions 26 and 39 through a sequence of general statements. As we deal with two different bases in Proposition 26, we first introduce the *n-reduction gradient*, which provides an intermediate between lexicographically minimal cycles with respect to Σ_* and the reduction gradient defined on the reduction basis Ω_* .

Let $(C_*, \Sigma_* = \sigma_1 < \dots < \sigma_l)$ be a based chain complex with an elementwise filtration, and let $R = D \cdot S$ be a reduction of the filtration boundary matrix. We equip the chain complex C_* with the new ordered basis $\Pi_* = \tau_1 < \dots < \tau_l$ given by

$$\tau_i = \begin{cases} S_i & \text{if } i \text{ is a death index and } \sigma_i \in C_{n+1}, \\ \sigma_i & \text{otherwise.} \end{cases}$$

We call Π_* the *n-reduction basis*. Note that Σ_* and Π_* induce the same lexicographic preorder on C_n . In particular, note that for every death index j with $\sigma_j \in C_{n+1}$ we have $\text{Pivot}_{\Pi_*} R_j = \text{Pivot}_{\Sigma_*} R_j$, where $R_j = D \cdot S_j$. Similarly to Proposition 13, the collection of disjoint pairs

$$\{(\text{Pivot}_{\Pi_*} R_j, S_j) \mid j \text{ is a death index and } \sigma_j \in C_{n+1}\}$$

is an algebraic gradient on Π_* , the *n-reduction gradient* of S . By construction, the *n-reduction gradient* is *n-reduced* and generates the *n-boundaries*.

We now prove some general statements about algebraic gradients that are *n-reduced* and generate the *n-boundaries*. Let $(C_*, \Pi_* = \tau_1 < \dots < \tau_l)$ be any based chain complex with an elementwise filtration.

► **Lemma 37.** *If V is an n -reduced algebraic gradient on Π_* that generates the n -boundaries, then a cycle $z \in Z_n$ is reducible if and only if its support in the basis Π_* contains a gradient facet of V .*

Proof. Assume that the cycle z is reducible. By definition, there exists a cycle $z' \sqsubset z$ and a chain $e \in C_{n+1}$ such that $z - z' = \partial e$. As V generates the *n-boundaries*, the boundary $\partial e \in B_n = \text{span } \partial_n V$ can be written as a linear combination $\partial e = \sum \lambda_i \partial b_i$ for some $(a_i, b_i) \in V$ with $\lambda_i \neq 0$ and $a_i \in \Pi_n$. Since the chains ∂b_i have distinct pivots, as V is *n-reduced* and the pairs in V are disjoint, there exists a unique chain ∂b_j with $\text{Pivot}_{\Pi_*} \partial e = \text{Pivot}_{\Pi_*} \partial b_j = a_j$. As $\lambda_j \neq 0$, the gradient facet a_j must be contained in z , as otherwise $z \sqsubseteq z'$, contradicting the assumption.

For the converse, note that if the cycle z contains some gradient facet a with $(a, b) \in V$, then $z' = \Phi^{(a,b)}(z) = z + \langle z, a \rangle \cdot \partial F(a)$, where $F(a) = -\langle \partial b, a \rangle^{-1} \cdot b$, is a homologous cycle with $z' \sqsubset z$, since $\text{Pivot}_{\Pi_*} \partial b = a$ and a is contained in $\text{supp}_{\Pi_*} z$ but not in $\text{supp}_{\Pi_*} z'$. ◀

► **Proposition 38.** *If V is an n -reduced algebraic gradient on a basis Π_* that generates the n -boundaries, then for a cycle $z \in Z_n$ the following are equivalent:*

- z is lexicographically minimal with respect to the basis Π_* ;
- z is invariant under the algebraic flow Φ determined by V , i.e., it satisfies $\Phi(z) = z$.

Proof. Let $z \in Z_n$ be a Φ -invariant cycle. By Proposition 22, this is equivalent to z not containing a gradient facet, and that, in turn is equivalent by Lemma 37 to z being irreducible, which is the same as being lexicographically minimal. ◀

In particular, note that Proposition 38 implies that every *n-reduced* algebraic gradient on Π_* that generates the *n-boundaries* has the same set of invariant cycles.

Proof of Proposition 26. By construction of the algebraic flow and under the assumption that S is death-compatible, the reduction flow agrees on cycles with the flow determined by the n -reduction gradient on the n -reduction basis Π_* . Thus, together with Proposition 38, this implies that the cycle z is invariant under the reduction flow if and only if z is lexicographically minimal with respect to the ordered basis Π_* . As mentioned before, the ordered bases Π_* and Σ_* induce the same lexicographic preorder on C_n , and hence the claim follows. ◀

Next, we show the uniqueness of optimal homologous cycles:

► **Proposition 39.** *Let $(C_*, \Sigma_* = \sigma_1 < \dots < \sigma_l)$ be a based chain complex with an elementwise filtration, and let z be lexicographically minimal cycle with respect to the basis Σ_* . If $c \in Z_n$ is homologous to z and $c \sqsubseteq z$ with respect to Σ_* , then we must have $c = z$.*

Proof. Let $R = D \cdot S$ be a reduction of the filtration boundary matrix, and let V be the n -reduction gradient on the n -reduction basis Π_* . Recall that V is n -reduced and generates the n -boundaries by construction. As mentioned before, the ordered bases Π_* and Σ_* induce the same lexicographic preorder on C_n , and thus, both cycles z and c are also lexicographically minimal with respect to the basis Π_* . By Proposition 38, this implies that both cycles are invariant under the flow determined by V . As V generates the n -boundaries, for z and c being homologous is the same as being V -homologous, and hence Proposition 22 implies $c = z$. ◀

Finally, we give a proof of Proposition 25 using these results.

Proof of Proposition 25. By Remark 23, we know that there exists a death-compatible reduction matrix S . Thus, by Proposition 26, a cycle z is lexicographically minimal with respect to the elementwise filtration if and only if z is invariant under the reduction flow. As in the proof of Proposition 26, this is equivalent to z being invariant under the flow of the n -reduction gradient. This in turn is, by Proposition 22, equivalent to z not containing any gradient facets of the n -reduction gradient. Finally, by the construction of the n -reduction gradient, this is equivalent to z not containing any (non-essential) birth elements, meaning that it only contains death elements and essential elements. ◀

A.4 Algebraic flow as matrix reduction

We demonstrate how the algebraic flow on a cycle can be interpreted as a variant of Gaussian elimination, tying it closely to the exhaustive reduction introduced in Section 2.1.

Let $(C_*, \Sigma_* = \sigma_1 < \dots < \sigma_l)$ be a based chain complex with an elementwise filtration, and let V be an algebraic gradient on Σ_* .

Input: $D = \partial$ the $l \times l$ filtration boundary matrix, c a cycle

for $i = 1, \dots, l$ **do**

if $c_i \neq 0$ **and** $(\sigma_i, \sigma_j) \in V$ **a gradient pair then**

$\mu = -c_i / D_{i,j};$

$c = c + \mu \cdot D_j;$

return c

■ **Algorithm 3** Gradient flow reduction

► **Proposition 40.** *If V is n -reduced and $c \in Z_n$ a cycle, then Algorithm 3 computes $\Phi(c)$.*

Proof. By definition, for any cycle c the algebraic flow is given by $\Phi(c) = c + \partial F(c)$. We denote by L the set of pairs (i, j) with $c_i \neq 0$ and $(\sigma_i, \sigma_j) \in V$. Recall from Definition 16 that $F: C_* \rightarrow C_{*+1}$ is the linear map with $F(\sigma_i) = -\langle \partial\sigma_j, \sigma_i \rangle^{-1} \cdot \sigma_j$ if $(\sigma_i, \sigma_j) \in V$ and 0 on all other basis elements. Note that $\partial\sigma_j$ is represented by the column D_j and $\langle \partial\sigma_j, \sigma_i \rangle = D_{i,j}$. Thus, we have

$$\partial F(c) = \sum_{(i,j) \in L} c_i \cdot \partial F(\sigma_i) = \sum_{(i,j) \in L} (-c_i \langle \partial\sigma_j, \sigma_i \rangle^{-1}) \cdot \partial\sigma_j = \sum_{(i,j) \in L} (-c_i / D_{i,j}) \cdot D_j.$$

By assumption of V being n -reduced, these columns D_j have distinct pivots and thus, traversing and updating the cycle c from small to large index has the same effect as adding a summand from the sum above to the cycle c one after the other. Hence, Algorithm 3 computes $\Phi(c)$. ◀

To compute the image of a cycle under the stabilized flow, one can use Algorithm 4, which resembles the exhaustive reduction from Section 2.1, and that does not require any additional specific choices besides the algebraic gradient, in contrast to Algorithm 3.

Input: $D = \partial$ the filtration boundary matrix, c a cycle
while *there exists i with $c_i \neq 0$ and $(\sigma_i, \sigma_j) \in V$ a gradient pair* **do**
 $\mu = -c_i / D_{i,j};$
 $c = c + \mu \cdot D_j;$
return c
■ **Algorithm 4** Stabilized flow reduction

► **Proposition 41.** *For any cycle $c \in Z_n$, Algorithm 4 computes $\Phi^\infty(c)$.*


Proof. Note that we can choose a different elementwise filtration by the σ_i without crucially affecting Algorithm 4. Moreover, by Lemma 32, we can assume that V is n -reduced. Thus, the algorithm is essentially the same as Algorithm 2, implying that it terminates.

Now note that Algorithm 4 computes a V -homologous cycle c' of c that contains no gradient facets, and thus it is Φ -invariant by Proposition 22. By definition, the stabilized cycle $\Phi^\infty(c)$ is also a Φ -invariant cycle and V -homologous to c . Therefore, both c' and $\Phi^\infty(c)$ are Φ -invariant cycles that are V -homologous. Proposition 22 now implies $c' = \Phi^\infty(c)$. ◀

B Reduction chains and descending complexes

For a generalized discrete Morse function $f: K \rightarrow \mathbb{R}$, we know from Corollary 30 that the columns of the totally reduced filtration boundary matrix corresponding to non-zero persistence pairs are supported on the descending complex. In this section, we show that the reduction chains, i.e., the chains given as the columns of the reduction matrix, that correspond to essential and non-zero persistence pairs are also supported on the descending complex.

Let D be the filtration boundary matrix of the simplexwise filtration of K induced by the f -lexicographic order, and let S be a death-compatible reduction matrix with $R = D \cdot S$ totally reduced.



■ **Figure 5** Discrete gradient (blue) together with the unique critical simplex (orange).

► **Proposition 42.** *Let $f: K \rightarrow \mathbb{R}$ be a generalized discrete Morse function and let $\tau \in K$ be a critical simplex, i.e., a simplex that is either essential or contained in a non-zero persistence pair of the simplexwise filtration induced by the f -lexicographic order. Let $r = f(\tau)$ be the function value of τ . Then the chain given as the column S_τ of the reduction matrix is supported on the descending complex $D_r(f)$.*

Proof. Note that if τ is an essential simplex or a birth simplex contained in a non-zero persistence pair, then it is an essential simplex of the (smaller) simplexwise filtration of $S_r(f)$ induced by the f -lexicographic order. Let W_r be the zero persistence apparent pairs gradient of this simplexwise filtration and denote by $\Phi: C_*(S_r(f)) \rightarrow C_*(S_r(f))$, with $\Phi(c) = c + F(\partial c) + \partial F(c)$, the associated flow.

Assume first that τ is an essential simplex of the simplexwise filtration of $S_r(f)$. Then, the chain c given as the column S_τ of the reduction matrix is a cycle, i.e., $\partial c = 0$. Moreover, since S is death-compatible, we also have $F(c) = 0$. Hence, we get

$$\Phi(c) = c + F(\partial c) + \partial F(c) = c + 0 + 0 = c$$

and the chain c is Φ -invariant. Analogously as in the proof of Theorem 29, we conclude that c is supported on the descending complex $D_r(f)$.

Now assume that τ is a death simplex of the simplexwise filtration of $S_r(f)$ that is contained in a non-zero persistence pair (σ, τ) , implying that τ is not contained in a zero-persistence apparent pair. By Definition 1, and as the matrix R is totally reduced, the column $R_\tau = D \cdot S_\tau$ does not contain any apparent facet of the simplexwise filtration. Therefore, the chain c given as the column S_τ of the reduction matrix satisfies $F(\partial c) = 0$. As before, since S is death-compatible, we also have $F(c) = 0$. Hence, we get

$$\Phi(c) = c + F(\partial c) + \partial F(c) = c + 0 + 0 = c$$

and the chain c is Φ -invariant. Analogously to before, we conclude that c is supported on the descending complex $D_r(f)$, proving the claim. ◀

► **Remark 43.** Note that the assumption in Corollary 30 and Proposition 42 on the pair (σ, τ) to be a non-zero persistence pair can not be dropped. Consider, for example, the simplicial complex in Figure 5 and the discrete Morse function $f: K \rightarrow \mathbb{R}$ that assigns to the simplices σ and τ the value 3 and to all other simplices the value as indicated. Then the pair (σ, τ) is a zero persistence apparent pair, but neither $S_\tau = \tau$ nor $S_\sigma = R_\tau = \partial \tau$ is supported on the descending complex $D_3(f)$, which only consists of the orange vertex with function value 0.