

Antipodal Self-Duality for a Four-Particle Form Factor

Lance J. Dixon¹, Ömer Gürdoğan², Yu-Ting Liu^{1,3}, Andrew J. McLeod^{4,5} and Matthias Wilhelm⁶

¹ *SLAC National Accelerator Laboratory, Stanford University, Stanford, CA 94309, USA*

² *School of Physics & Astronomy, University of Southampton, Southampton, SO17 1BJ, UK*

³ *Kavli Institute for Theoretical Physics, UC Santa Barbara, Santa Barbara, CA 93106, USA*

⁴ *CERN, Theoretical Physics Department, 1211 Geneva 23, Switzerland*

⁵ *Mani L. Bhaumik Institute for Theoretical Physics,*

Department of Physics and Astronomy, UCLA, Los Angeles, CA 90095, USA and

⁶ *Niels Bohr International Academy, Niels Bohr Institute, Blegdamsvej 17, 2100 Copenhagen Ø, Denmark*

We bootstrap the symbol of the maximal-helicity-violating four-particle form factor for the chiral part of the stress-tensor supermultiplet in planar $\mathcal{N} = 4$ super-Yang-Mills theory at two loops. When minimally normalized, this symbol involves only 34 letters and obeys the extended Steinmann relations in all partially-overlapping three-particle momentum channels. In addition, the remainder function for this form factor exhibits an antipodal self-duality: it is invariant under the combined operation of the antipodal map defined on multiple polylogarithms—which reverses the order of the symbol letters—and a simple kinematic map. This self-duality holds on a four-dimensional parity-preserving kinematic hypersurface. It implies the antipodal duality recently noticed between the three-particle form factor and the six-particle amplitude in this theory.

Introduction

Symmetries play a central role in modern formulations of fundamental physics, where they reflect simple facts about the world such as conservation laws and how different types of particles interact. However, sometimes new symmetries emerge in our theories whose physical implications aren't immediately clear. These discoveries often lead to the development of more powerful mathematical techniques for making predictions, and have the potential to guide us to new physical principles.

The planar limit of $\mathcal{N} = 4$ super-Yang-Mills (SYM) theory has proven to be an especially rewarding place to look for novel symmetries in particle physics. Most famously, scattering amplitudes and form factors in this theory are dual to Wilson loops with lightlike edges [1–13] and as such respect a dual conformal symmetry [1, 14–17]. These quantities also exhibit interesting number-theoretic symmetries [18], as well as intriguing connections to cluster algebras, tropical fans, and positive geometries [19–43]. While most of these special properties don't directly generalize to non-supersymmetric theories, their study has still led to significant improvements in our understanding of more general classes of amplitudes, form factors, and Feynman integrals (see for instance [36, 44–50]).

Much of this recent progress has been fuelled by the in-depth study of amplitudes that evaluate to multiple polylogarithms (MPLs) [51–56], which are endowed with a Hopf algebra structure [57–62]. In particular, one part of this structure—the symbol map—greatly simplifies the study of what sequences of discontinuities appear in polylogarithmic functions. This fact has been leveraged to bootstrap certain amplitudes and form factors in planar $\mathcal{N} = 4$ SYM theory to extremely high loop orders [27, 45, 63–74].

In an unexpected recent development, these bootstrap results have revealed a mysterious new duality between the maximally-helicity-violating (MHV) six-particle amplitude in parity-preserving kinematics, and the MHV three-particle form factor that involves a single insertion of the chiral stress tensor multiplet, which includes the BPS operator $\text{tr}(\phi^2)$ [75]. Namely, these two quantities are related to each other by another one of the maps that appears in the Hopf algebra structure of MPLs: the antipode map. At the level of the symbol, the antipode map simply reverses the order of discontinuities in MPLs. Thus, this duality can be loosely understood as the observation that the three-particle form factor and the MHV six-particle amplitude (in parity-preserving kinematics) encode exactly the same sequences of discontinuities, but in the opposite order—after a suitable map between the kinematic variables that describe the two processes. At the moment, no physical argument is known for why this duality should hold, but it has been checked explicitly through at least seven loops. Interestingly, two-loop MHV amplitudes have also been shown to exhibit a different type of antipodal symmetry in parity-even kinematics, which is conjectured to hold to all particle multiplicity [76].

In this paper, we describe a more general antipodal duality, which applies to four-particle form factors. Namely, the four-particle form factor is *self*-dual under the action of the antipode in a four-dimensional subspace of its kinematics. In fact, as we will show below, this new duality *implies* the duality between the three-particle form factor and the six-particle amplitude, due to the fact that these quantities respectively appear in the double and triple collinear limits of the four-particle form factor.

In order to substantiate this claim, we first bootstrap the symbol of the four-particle form factor at two loops. We do this by constructing the space of symbols that in-

volve the letters that appear in the Feynman integrals that contribute to this form factor [48, 77–79]. Amazingly, the form factor is uniquely identified in this space by just the first-entry condition, invariance under a standard set of discrete symmetries, and the strict (leading-power) double collinear limits. (This rigidity is reminiscent of the seven-particle amplitude, whose three-loop MHV symbol could be bootstrapped with only mild collinear information [80].) Three other limits—triple-collinear limits, the recently-computed limit of light-like operator momenta [81], and the near-collinear limit—serve as cross-checks on our result.

Multiple normalizations are needed to expose the different properties of this four-particle form factor. In one normalization, the symbol of the form factor obeys the extended Steinmann relations (defined below) in all partially-overlapping three-particle momentum channels. In another normalization, this form factor is antipodally self-dual. We describe these normalizations, as well as the self-duality map, in more detail below. We provide the symbol in both normalizations in ancillary files, which also describe its various discrete symmetries and kinematic limits.

The Bootstrap

We begin our bootstrap by removing the infrared divergences and the MHV tree-level prefactor from the four-particle MHV form factor [82] $\mathcal{F}_4^{\text{MHV}}$,

$$\mathcal{F}_4^{\text{MHV}} = \mathcal{F}_4^{\text{min}} \times F_4, \quad (1)$$

where

$$\mathcal{F}_4^{\text{min}} = \mathcal{F}_4^{\text{MHV, tree}} \times \exp \left[-\frac{g^2}{\epsilon^2} \sum_{i=1}^4 \left(\frac{\mu^2}{-s_{i,i+1}} \right)^\epsilon \right]. \quad (2)$$

Here, the 't Hooft coupling is $g^2 = N_c g_{\text{YM}}^2 / (16\pi^2)$ and we have omitted contributions proportional to transcendental constants. Our main objective is to calculate F_4 , which depends on eight dimensionless ratios:

$$u_i = \frac{(p_i + p_{i+1})^2}{q^2}, \quad v_i = \frac{(p_i + p_{i+1} + p_{i+2})^2}{q^2}, \quad (3)$$

where $i = 1, 2, 3, 4$ and $q = \sum_{i=1}^4 p_i$ is the (generically off-shell) momentum of the operator insertion. All indices should be understood to be mod 4. Due to momentum conservation and the masslessness of the four particles, $p_i^2 = 0$, these variables satisfy three constraints:

$$-u_1 + u_3 + v_4 + v_1 = 1, \quad (4)$$

$$-u_2 + u_4 + v_1 + v_2 = 1, \quad (5)$$

$$-u_3 + u_1 + v_2 + v_3 = 1. \quad (6)$$

Correspondingly, the four-particle form factor depends on five independent variables.

When expanded perturbatively in the coupling,

$$F_4 = 1 + \sum_{L=1}^{\infty} g^{2L} F_4^{(L)}, \quad (7)$$

the L -loop contribution $F_4^{(L)}$ is expected to be expressible as a linear combination of MPLs of weight $2L$ with rational coefficients. The symbol of an MPL can be defined iteratively by its total differential [58]:

$$dG = \sum_{x \in \mathcal{L}} G^x d \ln x \quad \Rightarrow \quad \mathcal{S}(G) = \sum_{x \in \mathcal{L}} \mathcal{S}(G^x) \otimes x, \quad (8)$$

where G^x are MPLs of one lower weight. The $d \ln$ arguments x are referred to as symbol letters, while the total multiplicative span of the letters appearing in an MPL is referred to as its symbol alphabet. For more background on MPLs and the symbol map, see for instance [83].

The symbol of the one-loop form factor is [7]

$$\begin{aligned} \mathcal{S}(F_4^{(1)}) &= 2v_1 \otimes (1 - v_1) + \frac{u_1}{u_2 u_4} \otimes u_1 \\ &+ \frac{u_1}{v_4 v_1} \otimes \frac{u_1 - v_4 v_1}{u_1} + \text{cyclic}, \end{aligned} \quad (9)$$

where the cyclic transformation maps $u_i \mapsto u_{i+1}$ and $v_i \mapsto v_{i+1}$. We recall that the two-loop remainder function is related to the form factor itself by

$$R_4^{(2)} = F_4^{(2)} - \frac{1}{2} [F_4^{(1)}]^2, \quad (10)$$

and $R_4^{(2)}$ has smooth behavior in factorization limits.

In order to bootstrap $F_4^{(2)}$ or $R_4^{(2)}$, we first assemble the alphabet of symbol letters that can appear in the four-particle form factor. Since the Feynman integrals that contribute to this form factor are all known [48, 77, 78], this can be done easily.¹ Altogether, the relevant Feynman integrals depend on 113 independent letters. Five different square roots appear in this alphabet, one of which involves the dihedrally-invariant argument

$$\text{tr}_5 = 4i\epsilon_{\alpha\beta\gamma\delta} p_1^\alpha p_2^\beta p_3^\gamma p_4^\delta, \quad (11)$$

and four of which are organized into pairs of two-orbits under the action of the dihedral group.² The dihedral group D_4 is generated by the order-four cyclic transformation and a flip transformation, for example $u_1 \leftrightarrow u_4$, $u_2 \leftrightarrow u_3$, $v_1 \leftrightarrow v_3$. Spacetime parity acts by flipping the

¹ While the nonplanar double pentagon Feynman integrals that contribute to this form factor have not yet been published [78], they do not give rise to letters beyond those that appear in the planar pentabox and nonplanar hexabox topologies [79].

² In the notation of [48], these four additional roots correspond to two orientations of Δ_3 and two orientations of Σ_5 . Further orientations of these roots don't show up because they correspond to different planar orderings.

sign in front of tr_5 , while leaving the sign in front of the other square roots intact.

We next construct the space of integrable weight-four symbols that draws upon this alphabet. We are only interested in symbols that have the correct branch cuts (satisfy the first-entry condition [84]), which in this context states that only the 8 letters $\{u_i, v_i\}$ can occur in the first entry. We also impose invariance under the dihedral symmetry group D_4 . There are 522 independent symbols s_i satisfying these conditions, which we use to formulate our initial ansatz for the symbol of the remainder function $R_4^{(2)}$:

$$\mathcal{S}(R_4^{(2)}) = \sum_{i=1}^{522} c_i s_i, \quad (12)$$

where the c_i are undetermined rational coefficients.

To fix the values of the coefficients in (12), we first require that our ansatz is even under all elements of the algebraic Galois group, which flip the signs in front of each of the 5 square roots separately. (Note that one of the elements is parity.) This imposes 148 independent conditions on the coefficients in our ansatz. Next, we require that our ansatz for $R_4^{(2)}$ reduces to the three-particle form-factor remainder $R_3^{(2)}$ [65] when two of the external particles become collinear. In this limit, the 113-letter alphabet involves 25 spurious letters, in addition to the 6 letters describing the three-particle form factor F_3 . Matching our ansatz onto the correct expression completely fixes the remaining 374 coefficients, and thus uniquely determines the symbol of $R_4^{(2)}$, or equivalently of $F_4^{(2)}$. The numbers of free parameters at each stage in the calculation are collected in Table I.

Although we started with an initial ansatz of over one hundred letters, our result for the symbol of $F_4^{(2)}$ only involves 34 letters (notably, all five square roots still appear). As expected, it also obeys the Steinmann relations [85, 86], which forbid sequential discontinuities in partially-overlapping momentum channels; in the case of massless scattering amplitudes, these relations apply when both channels contain at least three particles [70]. However, while the Steinmann relations only put constraints on the first two entries of the symbol, many amplitudes have been found to obey an *extended* set of Steinmann relations, in which the same constraints hold for all adjacent entries in the symbol [18, 72, 87, 88]. In processes involving one massive and four massless external legs, the (extended) Steinmann relations forbid the letter v_i from appearing next to v_j in the symbol when $j \neq i$. Notably, the two-loop master integrals that contribute to $F_4^{(2)}$ only obey the Steinmann relations—not the extended Steinmann relations—between v_i and v_{i+2} [48]. However, the extended Steinmann relations are obeyed in *all* channels by $F_4^{(2)}$. Similarly, higher-point amplitudes in this theory obey all extended Steinmann relations when normalized minimally (although amplitudes

Constraints	Parameters
first entry, integrability, D_4 invariance	522
Galois symmetry	374
strict double collinear limit $\rightarrow R_3^{(2)}$	0
strict triple collinear limit $\rightarrow \hat{R}_6^{(2)}$	0
light-like limit	0
FFOPE	0

TABLE I. Number of free parameters that remain after imposing each constraint in the bootstrap procedure, starting with the 113-letter alphabet. The limits are taken at leading power, except for the FFOPE.

do not respect dual conformal invariance in this normalization) [28, 72].

Special Kinematic Limits

We can check our results in a number of different kinematic limits. First we consider the light-like limit, where the operator momentum $q^2 \rightarrow 0$. In this limit, the 34 letters that appear in the symbol of $R_4^{(2)}$ reduce to 13 independent multiplicative combinations. Nine of these combinations match the light-like letters reported in [81], while four are spurious and must drop out of R_4 . We have confirmed that our $R_4^{(2)}$ symbol correctly reproduces the symbol of the light-like form factor remainder reported in [81].

Although our bootstrap procedure made use of information about the strict, or leading power, collinear limit, we can still make nontrivial predictions for the subleading powers in expansion of $F_4^{(2)}$ around this limit. Such terms are predicted by the recently-developed Form Factor Operator Product Expansion (FFOPE) [89–91]. To carry out this cross-check, we rewrite our kinematic variables in terms of the OPE variables T , T_2 , S , S_2 , and f_2 [89, 92]:

$$\begin{aligned} u_1 &= \frac{T^2 T_2^2}{(T^2 + 1)(S^2 + T^2 + T_2^2 + 1)}, \\ u_2 &= \left[(1+T^2) + \frac{S^2(S_2 T_2(1+f_2^2) + f_2(1+S_2^2+T^2+T_2^2))}{f_2 S_2^2} \right]^{-1}, \\ u_3 &= \frac{S^2}{(T^2 + 1)(S^2 + T^2 + T_2^2 + 1)}, \\ u_4 &= \frac{S^2 T^2}{S_2^2} u_2, \\ v_1 &= \frac{T_2^2 + 1}{S^2 + T^2 + T_2^2 + 1}, \end{aligned} \quad (13)$$

while v_2 , v_3 , and v_4 are fixed by the relations (4)–(6). The near-collinear limit in these variables corresponds to an expansion around small values of T_2 , which we have computed by expanding our symbol to $\mathcal{O}(T^2 T_2^2)$. We checked this expansion against the predictions made by




FIG. 1. The four-particle form factor is antipodally self-dual in parity-preserving kinematics. This duality maps the double and triple collinear limits of the form factor to each other. Because the four-particle form factor reduces to the three-particle form factor and the six-particle amplitude in these limits, the self-duality of the four-particle form factor implies the duality observed in [75].

the FFOPE (using the procedure explained in [90]) for the $T^2 T_2 \ln(T)$, $T^2 T_2 \ln(T_2)$, $T^2 T_2$, and $T^2 T_2^2 \ln(T_2)$ contributions.³ Each of these checks was carried out as a series in S and S_2 and to all available powers of f_2 .

Finally, it can be seen from the OPE [89–104], as well as from arguments based on dual conformal invariance and factorization [9], that the four-particle form factor remainder R_4 must reduce to the six-particle MHV amplitude’s remainder \hat{R}_6 as $T \rightarrow 0$ in the parametrization introduced in (13). In both cases, the limit can be interpreted as a triple-collinear limit; in the six-particle case, the limit covers all of the dual-conformally invariant phase space, allowing the triple-collinear splitting amplitude’s finite part to be identified with \hat{R}_6 . We have checked that this limit is indeed obeyed.

Antipodal Self-Duality

As discussed in the last section, the four-particle form factor possesses kinematic limits in which it reduces to the three-particle form factor and to the six-particle amplitude. These two quantities were recently discovered to be antipodally dual [75], making it tempting to investigate whether the four-particle form factor could be dual to itself in parity-preserving kinematics, where tr_5 vanishes. In the OPE variables, this hypersurface simply corresponds to setting $f_2 = 1$. In the u_i and v_i variables,

it requires setting the Gram determinant

$$u_2^2 [u_1^2 - 2u_1(1 + u_3) + (1 - u_3)^2] + [u_1 v_2 - v_1(v_2 - u_3)]^2 - 2u_2 [u_1^2 v_2 + u_1(u_3(2 - v_1 - v_2) - v_2(1 + v_1)) + v_1(1 - u_3)(v_2 - u_3)] \quad (14)$$

to zero. Note that none of the other four square roots rationalize on this $\text{tr}_5 = 0$ surface.

Surprisingly, we find that an antipodal self-duality does in fact hold on this parity-preserving hypersurface:

$$R_4|_{\text{tr}_5=0} = S(R_4|_{\text{tr}_5=0})|_{u_i, v_i \rightarrow g(u_i), g(v_i)} \quad (15)$$

where $S(F)$ denotes the polylogarithmic antipode of F , which acts at the level of the symbol as [105, 106]

$$S(x_1 \otimes x_2 \otimes \cdots \otimes x_m) = (-1)^m x_m \otimes \cdots \otimes x_2 \otimes x_1, \quad (16)$$

and the kinematic map is defined by

$$g(u_1) = u_1 \sqrt{\frac{u_2 u_4}{(u_2 - v_1 v_2)(u_4 - v_3 v_4)}}, \quad (17)$$

$$g(v_1) = (1 - v_1) \sqrt{\frac{u_1 u_2}{(u_1 - v_4 v_1)(u_2 - v_1 v_2)}}, \quad (18)$$

plus cyclic images. In the OPE variables, this mapping takes an even simpler form:

$$g(T) = \sqrt{\frac{T_2}{S_2}}, \quad g(S) = \sqrt{\frac{1}{T_2 S_2}}, \quad (19)$$

$$g(T_2) = \frac{T}{S}, \quad g(S_2) = \frac{1}{TS},$$

and it is clear that $g^2 = 1$. Notably, this map reduces to the duality map described in [75] upon identifying T_2 and S_2 with the OPE parameters that describe the six-particle, which were denoted \hat{T} and \hat{S} in [75]. This identification naturally arises in the triple-collinear limit of R_4 , where the form factor reduces to the six-particle remainder \hat{R}_6 , which is mapped to the double collinear limit of R_4 by (19). However, the self-duality of R_4 in (15) holds more generally in the *full* four-dimensional space of parity-preserving kinematics.⁴

While antipodal self-duality (15) holds for the remainder function R_4 , there is an obstruction to it holding for F_4 . Namely, the one-loop form factor $F_4^{(1)}$ contains final entries that are not in the set $\{g(u_i), g(v_i)\}$, $i = 1, 2, 3, 4$, dictated by antipodal self-duality. From the symbol (9) of $F_4^{(1)}$, it is easy to see that the final entry sitting behind the first entry v_1 is just $g(v_1)^2$. However, the final

³ We note that the contributions at $\mathcal{O}(T^0)$ simply reproduce the OPE expansion of \hat{R}_6 , while the contributions at $\mathcal{O}(T_2^0)$ reproduce the OPE expansion of R_3 .

⁴ Note from eqs. (17) and (18) that the light-like limit $u_i, v_i \rightarrow \infty$ maps to finite u_i, v_i under the kinematic map g , which implies that the light-like form factor does not exhibit the antipodal self-duality we have found for $q^2 \neq 0$.

entry sitting behind u_1 is $(u_1 - v_4 v_1)/(u_4 u_2)$, and its logarithm is not a linear combination of $\{\ln g(u_i), \ln g(v_i)\}$. Furthermore, it does not seem possible to repair this obstruction by any simple adjustment of the normalization $F_4^{(1)}$ that is consistent with both the first-entry condition and the Steinmann relations.

Discussion and Conclusions

In this letter, we have bootstrapped the two-loop four-particle MHV form factor of the chiral part of the stress-tensor supermultiplet in planar $\mathcal{N} = 4$ SYM theory, and have found that it possesses an antipodal self-duality in parity-preserving kinematics. While we can only check this duality at two loops, the previously-identified antipodal duality between R_3 and \hat{R}_6 that it implies has been checked through seven loops (and even eight loops [107]), which strongly indicates that the self-duality of R_4 will hold to all loop orders.

The two-loop four-particle form factor is a highly constrained quantity, and we have only needed to impose invariance under its Galois and dihedral symmetries, as well as knowledge of its double-collinear limit, in order to uniquely fix all coefficients in our ansatz. This has left us with a wealth of cross-checks on our result, including the comparison of the near-collinear limit to the FFOPE, triple collinear limits, and the light-like limit. This also raises the question of whether the form factor can be bootstrapped to higher loop orders. The main unknown is whether new letters appear at three loops, beyond the 113 letters that appear in the two-loop integrals contributing to this quantity. While great strides have been made recently towards better characterizing the analytic structure of Feynman integrals from first principles (see for instance [108–117]), this question remains hard to address directly, and we leave it to future work.

It is natural to wonder if further antipodal (self-)dualities hold between form factors and amplitudes at higher particle multiplicity. Notably, as one increases the number of scattering particles, form factors contain an increasingly rich pattern of form factors and amplitudes in their (multi-)collinear limits, giving rise to many intriguing possibilities. In a similar vein, it would be interesting to calculate the two-loop NMHV form factor for the chiral part of the stress-tensor supermultiplet, and to search for further antipodal dualities that involve this quantity. More generally, it would be interesting to understand what the antipodal dualities we have already observed tell us about the physics underlying the three- and four-particle form factors.

The two-loop three-particle form factor is known to satisfy the principle of maximal transcendentality (PMT) [118–121], meaning that the $\mathcal{N} = 4$ SYM result for $\text{tr}(\phi^2)$ matches the maximally-transcendental part of

the Higgs-to-three-gluon amplitude in pure Yang-Mills theory (or QCD) in the leading large-top-mass limit (operator $\text{tr}(F^2)$) [62, 65, 122–126], for all gluon helicity configurations. The PMT has recently been extended (in a different way) to the four-gluon form factor of $\text{tr}(F^3)$ [126]. Here we ask: can the PMT for $\text{tr}(\phi^2)$ be extended to any Higgs-to-four-gluon helicity amplitudes? At one loop, the PMT already fails for the color-ordered helicity configurations $(----)$ and $(-+-+)$, and their parity conjugates [127, 128], but it works for $(-++)$ and $(----)$ and their parity conjugates [129, 130].

At two loops and leading-color, we cannot say much about $(-++)$ currently, because our starting point (12) was dihedrally invariant, and the $(-++)$ form factor (divided by the tree) need not be invariant, although its leading-transcendental part happens to be at one loop. That leaves $(----)$, which is dihedrally invariant. The leading transcendental, weight-four, parts of its double collinear limits match those in planar $\mathcal{N} = 4$ SYM, thanks to the PMT holding for the Higgs-to-three-gluon amplitude and for the $g \rightarrow gg$ splitting amplitude [131]. We found earlier that there is a unique weight-four, dihedrally- and parity-invariant function with specified double-collinear limits. Thus $R_4^{(2)}$ should also provide the parity-even part of the two-loop remainder for $A_4^{(2)}(\phi, 1^-, 2^-, 3^-, 4^-)$, where $\phi = H + iA$, with H the Higgs boson and A a pseudoscalar coupling to $\text{tr}(F\tilde{F})$ [124]. We also find a unique weight-four parity-odd dihedrally invariant function that vanishes in both the double and triple collinear limits. We leave to future work the tantalizing questions of whether the coefficient of this parity-odd function vanishes, as suggested by the PMT, and more generally, how much of the two-loop Higgs-to-four-gluon amplitude in QCD can be bootstrapped.

We would like to thank S. Abreu and V. Sotnikov for stimulating discussions, and Gang Yang for helpful comments on the manuscript. This research was supported by the US Department of Energy under contract DE-AC02-76SF00515, by the National Science Foundation under Grant No. NSF PHY-1748958, and by the Munich Institute for Astro-, Particle and BioPhysics (MIAPbP) which is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2094 – 390783311. LD and YL thank the KITP, and LD, AM and MW thank MIAPbP, for support and hospitality. YL was also supported in part by the Heising-Simons Foundation and the Simons Foundation. MW was supported in part by the ERC starting grant 757978 and grant 00025445 from Villum Fonden. ÖG is supported by the UKRI/EPSCRC Stephen Hawking Fellowship EP/T016396/1.

-
- [1] L. F. Alday and J. M. Maldacena, Gluon scattering amplitudes at strong coupling, *JHEP* **06**, 064, arXiv:0705.0303 [hep-th].
- [2] J. Drummond, G. Korchemsky, and E. Sokatchev, Conformal properties of four-gluon planar amplitudes and Wilson loops, *Nucl. Phys. B* **795**, 385 (2008), arXiv:0707.0243 [hep-th].
- [3] A. Brandhuber, P. Heslop, and G. Travaglini, MHV amplitudes in $\mathcal{N} = 4$ super Yang-Mills and Wilson loops, *Nucl. Phys. B* **794**, 231 (2008), arXiv:0707.1153 [hep-th].
- [4] J. Drummond, J. Henn, G. Korchemsky, and E. Sokatchev, Conformal Ward identities for Wilson loops and a test of the duality with gluon amplitudes, *Nucl. Phys. B* **826**, 337 (2010), arXiv:0712.1223 [hep-th].
- [5] L. F. Alday and R. Roiban, Scattering Amplitudes, Wilson Loops and the String/Gauge Theory Correspondence, *Phys. Rept.* **468**, 153 (2008), arXiv:0807.1889 [hep-th].
- [6] T. Adamo, M. Bullimore, L. Mason, and D. Skinner, Scattering Amplitudes and Wilson Loops in Twistor Space, *J. Phys. A* **44**, 454008 (2011), arXiv:1104.2890 [hep-th].
- [7] A. Brandhuber, B. Spence, G. Travaglini, and G. Yang, Form Factors in $\mathcal{N} = 4$ Super Yang-Mills and Periodic Wilson Loops, *JHEP* **01**, 134, arXiv:1011.1899 [hep-th].
- [8] L. F. Alday and J. Maldacena, Comments on gluon scattering amplitudes via AdS/CFT, *JHEP* **0711**, 068, arXiv:0710.1060 [hep-th].
- [9] Z. Bern, L. J. Dixon, D. A. Kosower, R. Roiban, M. Spradlin, C. Vergu, and A. Volovich, The Two-Loop Six-Gluon MHV Amplitude in Maximally Supersymmetric Yang-Mills Theory, *Phys. Rev.* **D78**, 045007 (2008), arXiv:0803.1465 [hep-th].
- [10] J. Drummond, J. Henn, G. Korchemsky, and E. Sokatchev, Hexagon Wilson loop = six-gluon MHV amplitude, *Nucl. Phys. B* **815**, 142 (2009), arXiv:0803.1466 [hep-th].
- [11] J. Maldacena and A. Zhiboedov, Form factors at strong coupling via a Y-system, *JHEP* **11**, 104, arXiv:1009.1139 [hep-th].
- [12] R. Ben-Israel, A. G. Tumanov, and A. Sever, Scattering amplitudes — Wilson loops duality for the first non-planar correction, *JHEP* **08**, 122, arXiv:1802.09395 [hep-th].
- [13] L. Bianchi, A. Brandhuber, R. Panerai, and G. Travaglini, Dual conformal invariance for form factors, *JHEP* **02**, 134, arXiv:1812.10468 [hep-th].
- [14] J. M. Drummond, J. Henn, V. A. Smirnov, and E. Sokatchev, Magic identities for conformal four-point integrals, *JHEP* **01**, 064, arXiv:hep-th/0607160 [hep-th].
- [15] Z. Bern, M. Czakon, L. J. Dixon, D. A. Kosower, and V. A. Smirnov, The Four-Loop Planar Amplitude and Cusp Anomalous Dimension in Maximally Supersymmetric Yang-Mills Theory, *Phys. Rev.* **D75**, 085010 (2007), arXiv:hep-th/0610248 [hep-th].
- [16] Z. Bern, J. Carrasco, H. Johansson, and D. Kosower, Maximally supersymmetric planar Yang-Mills amplitudes at five loops, *Phys. Rev.* **D76**, 125020 (2007), arXiv:0705.1864 [hep-th].
- [17] J. M. Drummond, J. Henn, G. P. Korchemsky, and E. Sokatchev, Dual superconformal symmetry of scattering amplitudes in $\mathcal{N} = 4$ super-Yang-Mills theory, *Nucl. Phys. B* **828**, 317 (2010), arXiv:0807.1095 [hep-th].
- [18] S. Caron-Huot, L. J. Dixon, F. Dulat, M. Von Hippel, A. J. McLeod, and G. Papathanasiou, The Cosmic Galois Group and Extended Steinmann Relations for Planar $\mathcal{N} = 4$ SYM Amplitudes, *JHEP* **09**, 061, arXiv:1906.07116 [hep-th].
- [19] J. Golden, A. B. Goncharov, M. Spradlin, C. Vergu, and A. Volovich, Motivic Amplitudes and Cluster Coordinates, *JHEP* **1401**, 091, arXiv:1305.1617 [hep-th].
- [20] J. Golden and M. Spradlin, A Cluster Bootstrap for Two-Loop MHV Amplitudes, *JHEP* **02**, 002, arXiv:1411.3289 [hep-th].
- [21] J. Golden, M. F. Paulos, M. Spradlin, and A. Volovich, Cluster Polylogarithms for Scattering Amplitudes, *J. Phys. A* **47**, 474005 (2014), arXiv:1401.6446 [hep-th].
- [22] J. Golden and M. Spradlin, An analytic result for the two-loop seven-point MHV amplitude in $\mathcal{N} = 4$ SYM, *JHEP* **1408**, 154, arXiv:1406.2055 [hep-th].
- [23] J. Drummond, J. Foster, and Ö. Gürdoğan, Cluster Adjacency Properties of Scattering Amplitudes in $\mathcal{N} = 4$ Supersymmetric Yang-Mills Theory, *Phys. Rev. Lett.* **120**, 161601 (2018), arXiv:1710.10953 [hep-th].
- [24] J. L. Bourjaily, A. J. McLeod, M. von Hippel, and M. Wilhelm, Rationalizing Loop Integration, *JHEP* **08**, 184, arXiv:1805.10281 [hep-th].
- [25] J. Drummond, J. Foster, and Ö. Gürdoğan, Cluster adjacency beyond MHV, *JHEP* **03**, 086, arXiv:1810.08149 [hep-th].
- [26] J. Golden and A. J. McLeod, Cluster Algebras and the Subalgebra Constructibility of the Seven-Particle Remainder Function, *JHEP* **01**, 017, arXiv:1810.12181 [hep-th].
- [27] J. Drummond, J. Foster, Ö. Gürdoğan, and G. Papathanasiou, Cluster adjacency and the four-loop NMHV heptagon, *JHEP* **03**, 087, arXiv:1812.04640 [hep-th].
- [28] J. Golden, A. J. McLeod, M. Spradlin, and A. Volovich, The Sklyanin Bracket and Cluster Adjacency at All Multiplicity, *JHEP* **03**, 195, arXiv:1902.11286 [hep-th].
- [29] J. Golden and A. J. McLeod, The two-loop remainder function for eight and nine particles, *JHEP* **06**, 142, arXiv:2104.14194 [hep-th].
- [30] J. Drummond, J. Foster, Ö. Gürdoğan, and C. Kalousios, Tropical Grassmannians, cluster algebras and scattering amplitudes, *JHEP* **04**, 146, arXiv:1907.01053 [hep-th].
- [31] J. Drummond, J. Foster, Ö. Gürdoğan, and C. Kalousios, Algebraic singularities of scattering amplitudes from tropical geometry, *JHEP* **04**, 002, arXiv:1912.08217 [hep-th].
- [32] N. Arkani-Hamed, T. Lam, and M. Spradlin, Non-perturbative geometries for planar $\mathcal{N} = 4$ SYM amplitudes, *JHEP* **03**, 065, arXiv:1912.08222 [hep-th].
- [33] N. Henke and G. Papathanasiou, How tropical are seven- and eight-particle amplitudes?, *JHEP* **08**, 005, arXiv:1912.08254 [hep-th].
- [34] J. Drummond, J. Foster, Ö. Gürdoğan, and C. Kalousios, Tropical fans, scattering equations and amplitudes, *JHEP* **11**, 071, arXiv:2002.04624 [hep-th].
- [35] J. Mago, A. Schreiber, M. Spradlin, and A. Volovich, Symbol alphabets from plabic graphs, *JHEP* **10**, 128, arXiv:2007.00646 [hep-th].

- [36] D. Chicherin, J. M. Henn, and G. Papathanasiou, Cluster algebras for Feynman integrals, *Phys. Rev. Lett.* **126**, 091603 (2021), arXiv:2012.12285 [hep-th].
- [37] J. Mago, A. Schreiber, M. Spradlin, A. Yelleshpur Srikant, and A. Volovich, Symbol alphabets from plabic graphs II: rational letters, *JHEP* **04**, 056, arXiv:2012.15812 [hep-th].
- [38] A. Herderschee, Algebraic branch points at all loop orders from positive kinematics and wall crossing, *JHEP* **07**, 049, arXiv:2102.03611 [hep-th].
- [39] S. He, Z. Li, and Q. Yang, Notes on cluster algebras and some all-loop Feynman integrals, *JHEP* **06**, 119, arXiv:2103.02796 [hep-th].
- [40] J. Mago, A. Schreiber, M. Spradlin, A. Yelleshpur Srikant, and A. Volovich, Symbol alphabets from plabic graphs III: $n = 9$, *JHEP* **09**, 002, arXiv:2106.01406 [hep-th].
- [41] N. Henke and G. Papathanasiou, Singularities of eight- and nine-particle amplitudes from cluster algebras and tropical geometry, *JHEP* **10**, 007, arXiv:2106.01392 [hep-th].
- [42] L. Ren, M. Spradlin, and A. Volovich, Symbol alphabets from tensor diagrams, *JHEP* **12**, 079, arXiv:2106.01405 [hep-th].
- [43] Q. Yang, Schubert problems, positivity and symbol letters, *JHEP* **08**, 168, arXiv:2203.16112 [hep-th].
- [44] J. L. Bourjaily, E. Gardi, A. J. McLeod, and C. Vergu, All-mass n -gon integrals in n dimensions, *JHEP* **08** (08), 029, arXiv:1912.11067 [hep-th].
- [45] L. J. Dixon, A. J. McLeod, and M. Wilhelm, A Three-Point Form Factor Through Five Loops, *JHEP* **04**, 147, arXiv:2012.12286 [hep-th].
- [46] S. He, Z. Li, Q. Yang, and C. Zhang, Feynman Integrals and Scattering Amplitudes from Wilson Loops, *Phys. Rev. Lett.* **126**, 231601 (2021), arXiv:2012.15042 [hep-th].
- [47] S. He, Z. Li, and Q. Yang, Kinematics, cluster algebras and Feynman integrals, (2021), arXiv:2112.11842 [hep-th].
- [48] S. Abreu, H. Ita, B. Page, and W. Tschernow, Two-loop hexa-box integrals for non-planar five-point one-mass processes, *JHEP* **03**, 182, arXiv:2107.14180 [hep-ph].
- [49] S. He, J. Liu, Y. Tang, and Q. Yang, The symbology of Feynman integrals from twistor geometries, (2022), arXiv:2207.13482 [hep-th].
- [50] J. M. Henn, A. Matijasić, and J. Miczajka, One-loop hexagon integral to higher orders in the dimensional regulator, (2022), arXiv:2210.13505 [hep-th].
- [51] K.-T. Chen, Iterated path integrals, *Bull. Amer. Math. Soc.* **83**, 831 (1977).
- [52] A. B. Goncharov, Geometry of configurations, polylogarithms, and motivic cohomology, *Adv. Math.* **114**, 197 (1995).
- [53] A. B. Goncharov, Multiple polylogarithms, cyclotomy and modular complexes, *Math. Res. Lett.* **5**, 497 (1998), arXiv:1105.2076 [math.AG].
- [54] E. Remiddi and J. Vermaseren, Harmonic polylogarithms, *Int. J. Mod. Phys. A* **15**, 725 (2000), arXiv:hep-ph/9905237.
- [55] J. M. Borwein, D. M. Bradley, D. J. Broadhurst, and P. Lisonek, Special values of multiple polylogarithms, *Trans. Am. Math. Soc.* **353**, 907 (2001), arXiv:math/9910045 [math-ca].
- [56] S. Moch, P. Uwer, and S. Weinzierl, Nested sums, expansion of transcendental functions and multiscale multiloop integrals, *J. Math. Phys.* **43**, 3363 (2002), arXiv:hep-ph/0110083 [hep-ph].
- [57] A. B. Goncharov, Galois symmetries of fundamental groupoids and noncommutative geometry, *Duke Math. J.* **128**, 209 (2005), arXiv:math/0208144 [math.AG].
- [58] A. B. Goncharov, M. Spradlin, C. Vergu, and A. Volovich, Classical Polylogarithms for Amplitudes and Wilson Loops, *Phys. Rev. Lett.* **105**, 151605 (2010), arXiv:1006.5703 [hep-th].
- [59] F. Brown, On the decomposition of motivic multiple zeta values, *Adv. Studies in Pure Math.* **63**, 31 (2012), arXiv:1102.1310 [math.NT].
- [60] F. Brown, Mixed Tate motives over \mathbb{Z} , *Ann. of Math.* (2) **175**, 949 (2012), arXiv:1102.1312 [math.AG].
- [61] C. Duhr, H. Gangl, and J. R. Rhodes, From polygons and symbols to polylogarithmic functions, *JHEP* **10**, 075, arXiv:1110.0458 [math-ph].
- [62] C. Duhr, Hopf algebras, coproducts and symbols: an application to Higgs boson amplitudes, *JHEP* **1208**, 043, arXiv:1203.0454 [hep-ph].
- [63] L. J. Dixon, J. M. Drummond, and J. M. Henn, Bootstrapping the three-loop hexagon, *JHEP* **1111**, 023, arXiv:1108.4461 [hep-th].
- [64] L. J. Dixon, J. M. Drummond, and J. M. Henn, Analytic result for the two-loop six-point NMHV amplitude in $\mathcal{N} = 4$ super Yang-Mills theory, *JHEP* **1201**, 024, arXiv:1111.1704 [hep-th].
- [65] A. Brandhuber, G. Travaglini, and G. Yang, Analytic two-loop form factors in $\mathcal{N} = 4$ SYM, *JHEP* **05**, 082, arXiv:1201.4170 [hep-th].
- [66] L. J. Dixon, J. M. Drummond, M. von Hippel, and J. Pennington, Hexagon functions and the three-loop remainder function, *JHEP* **1312**, 049, arXiv:1308.2276 [hep-th].
- [67] L. J. Dixon and M. von Hippel, Bootstrapping an NMHV amplitude through three loops, *JHEP* **1410**, 65, arXiv:1408.1505 [hep-th].
- [68] L. J. Dixon, J. M. Drummond, C. Duhr, and J. Pennington, The four-loop remainder function and multi-Regge behavior at NNLLA in planar $\mathcal{N} = 4$ super-Yang-Mills theory, *JHEP* **1406**, 116, arXiv:1402.3300 [hep-th].
- [69] L. J. Dixon, M. von Hippel, and A. J. McLeod, The four-loop six-gluon NMHV ratio function, *JHEP* **01**, 053, arXiv:1509.08127 [hep-th].
- [70] S. Caron-Huot, L. J. Dixon, A. McLeod, and M. von Hippel, Bootstrapping a Five-Loop Amplitude Using Steinmann Relations, *Phys. Rev. Lett.* **117**, 241601 (2016), arXiv:1609.00669 [hep-th].
- [71] L. J. Dixon, J. Drummond, T. Harrington, A. J. McLeod, G. Papathanasiou, and M. Spradlin, Heptagons from the Steinmann Cluster Bootstrap, *JHEP* **02**, 137, arXiv:1612.08976 [hep-th].
- [72] S. Caron-Huot, L. J. Dixon, F. Dulat, M. von Hippel, A. J. McLeod, and G. Papathanasiou, Six-Gluon amplitudes in planar $\mathcal{N} = 4$ super-Yang-Mills theory at six and seven loops, *JHEP* **08**, 016, arXiv:1903.10890 [hep-th].
- [73] L. J. Dixon and Y.-T. Liu, Lifting Heptagon Symbols to Functions, *JHEP* **10**, 031, arXiv:2007.12966 [hep-th].
- [74] L. J. Dixon, Ö. Gürdoğan, A. J. McLeod, and M. Wilhelm, Bootstrapping a stress-tensor form factor through eight loops, *JHEP* **07**, 153, arXiv:2204.11901 [hep-th].

- [75] L. J. Dixon, Ö. Gürdoğan, A. J. McLeod, and M. Wilhelm, Folding Amplitudes into Form Factors: An Antipodal Duality, *Phys. Rev. Lett.* **128**, 111602 (2022), arXiv:2112.06243 [hep-th].
- [76] Y.-T. Liu, Antipodal symmetry of two-loop MHV amplitudes, *JHEP* **09**, 131, arXiv:2207.11815 [hep-th].
- [77] S. Abreu, H. Ita, F. Moriello, B. Page, W. Tschernow, and M. Zeng, Two-Loop Integrals for Planar Five-Point One-Mass Processes, *JHEP* **11**, 117, arXiv:2005.04195 [hep-ph].
- [78] S. Abreu, D. Chicherin, H. Ita, B. Page, V. Sotnikov, W. Tschernow, and S. Zoia, to appear.
- [79] S. Abreu, private communication.
- [80] J. M. Drummond, G. Papathanasiou, and M. Spradlin, A Symbol of Uniqueness: The Cluster Bootstrap for the 3-Loop MHV Heptagon, *JHEP* **03**, 072, arXiv:1412.3763 [hep-th].
- [81] Y. Guo, L. Wang, and G. Yang, Analytic Four-Point Lightlike Form Factors and OPE of Null-Wrapped Polygons, (2022), arXiv:2209.06816 [hep-th].
- [82] A. Brandhuber, Ö. Gürdoğan, R. Mooney, G. Travaglini, and G. Yang, Harmony of Super Form Factors, *JHEP* **10**, 046, arXiv:1107.5067 [hep-th].
- [83] C. Duhr, Mathematical aspects of scattering amplitudes, in *Theoretical Advanced Study Institute in Elementary Particle Physics: Journeys Through the Precision Frontier: Amplitudes for Colliders* (2015) pp. 419–476, arXiv:1411.7538 [hep-ph].
- [84] D. Gaiotto, J. Maldacena, A. Sever, and P. Vieira, Pulling the straps of polygons, *JHEP* **12**, 011, arXiv:1102.0062 [hep-th].
- [85] O. Steinmann, Über den Zusammenhang zwischen den Wightmanfunktionen und der retardierten Kommutatoren, *Helv. Physica Acta* **33**, 257 (1960).
- [86] O. Steinmann, Wightman-Funktionen und retardierten Kommutatoren. II, *Helv. Physica Acta* **33**, 347 (1960).
- [87] S. Caron-Huot, L. J. Dixon, M. von Hippel, A. J. McLeod, and G. Papathanasiou, The Double Pentadladder Integral to All Orders, *JHEP* **07**, 170, arXiv:1806.01361 [hep-th].
- [88] S. He, Z. Li, and Q. Yang, Comments on all-loop constraints for scattering amplitudes and Feynman integrals, *JHEP* **01**, 073, [Erratum: *JHEP* **05**, 076 (2022)], arXiv:2108.07959 [hep-th].
- [89] A. Sever, A. G. Tumanov, and M. Wilhelm, Operator Product Expansion for Form Factors, *Phys. Rev. Lett.* **126**, 031602 (2021), arXiv:2009.11297 [hep-th].
- [90] A. Sever, A. G. Tumanov, and M. Wilhelm, An Operator Product Expansion for Form Factors II. Born level, *JHEP* **10**, 071, arXiv:2105.13367 [hep-th].
- [91] A. Sever, A. G. Tumanov, and M. Wilhelm, An Operator Product Expansion for Form Factors III. Finite Coupling and Multi-Particle Contributions, *JHEP* **03**, 128, arXiv:2112.10569 [hep-th].
- [92] B. Basso, A. Sever, and P. Vieira, Spacetime and Flux Tube S-Matrices at Finite Coupling for $\mathcal{N} = 4$ Supersymmetric Yang-Mills Theory, *Phys. Rev. Lett.* **111**, 091602 (2013), arXiv:1303.1396 [hep-th].
- [93] L. F. Alday, D. Gaiotto, J. Maldacena, A. Sever, and P. Vieira, An Operator Product Expansion for Polygonal null Wilson Loops, *JHEP* **1104**, 088, arXiv:1006.2788 [hep-th].
- [94] B. Basso, A. Sever, and P. Vieira, Space-time S-matrix and Flux tube S-matrix II. Extracting and Matching Data, *JHEP* **1401**, 008, arXiv:1306.2058 [hep-th].
- [95] B. Basso, A. Sever, and P. Vieira, Space-time S-matrix and Flux-tube S-matrix III. The two-particle contributions, *JHEP* **08**, 085, arXiv:1402.3307 [hep-th].
- [96] B. Basso, A. Sever, and P. Vieira, Collinear Limit of Scattering Amplitudes at Strong Coupling, *Phys. Rev. Lett.* **113**, 261604 (2014), arXiv:1405.6350 [hep-th].
- [97] B. Basso, A. Sever, and P. Vieira, Space-time S-matrix and Flux-tube S-matrix IV. Gluons and Fusion, *JHEP* **09**, 149, arXiv:1407.1736 [hep-th].
- [98] A. Belitsky, Nonsinglet pentagons and NMHV amplitudes, *Nucl. Phys. B* **896**, 493 (2015), arXiv:1407.2853 [hep-th].
- [99] A. Belitsky, Fermionic pentagons and NMHV hexagon, *Nucl. Phys. B* **894**, 108 (2015), arXiv:1410.2534 [hep-th].
- [100] B. Basso, J. Caetano, L. Cordova, A. Sever, and P. Vieira, OPE for all Helicity Amplitudes, *JHEP* **08**, 018, arXiv:1412.1132 [hep-th].
- [101] A. V. Belitsky, On factorization of multiparticle pentagons, *Nucl. Phys. B* **897**, 346 (2015), arXiv:1501.06860 [hep-th].
- [102] B. Basso, J. Caetano, L. Cordova, A. Sever, and P. Vieira, OPE for all Helicity Amplitudes II. Form Factors and Data Analysis, *JHEP* **12**, 088, arXiv:1508.02987 [hep-th].
- [103] B. Basso, A. Sever, and P. Vieira, Hexagonal Wilson loops in planar $\mathcal{N} = 4$ SYM theory at finite coupling, *J. Phys. A* **49**, 41LT01 (2016), arXiv:1508.03045 [hep-th].
- [104] A. Belitsky, Matrix pentagons, *Nucl. Phys. B* **923**, 588 (2017), arXiv:1607.06555 [hep-th].
- [105] A. Goncharov, Multiple polylogarithms and mixed Tate motives, (2001), arXiv:math/0103059 [math.AG].
- [106] F. Brown, Single-Valued Motivic Periods and Multiple Zeta Values, *SIGMA* **2**, e25 (2014), arXiv:1309.5309 [math.NT].
- [107] L. J. Dixon and Y.-T. Liu, An Eight Loop Amplitude via Antipodal Duality, to appear.
- [108] S. Abreu, R. Britto, C. Duhr, and E. Gardi, From multiple unitarity cuts to the coproduct of Feynman integrals, *JHEP* **10**, 125, arXiv:1401.3546 [hep-th].
- [109] S. Bloch and D. Kreimer, Cutkosky Rules and Outer Space, (2015), arXiv:1512.01705 [hep-th].
- [110] S. Abreu, R. Britto, C. Duhr, and E. Gardi, Algebraic Structure of Cut Feynman Integrals and the Diagrammatic Coaction, *Phys. Rev. Lett.* **119**, 051601 (2017), arXiv:1703.05064 [hep-th].
- [111] J. L. Bourjaily, H. Hannesdottir, A. J. McLeod, M. D. Schwartz, and C. Vergu, Sequential Discontinuities of Feynman Integrals and the Monodromy Group, *JHEP* **01**, 205, arXiv:2007.13747 [hep-th].
- [112] H. S. Hannesdottir, A. J. McLeod, M. D. Schwartz, and C. Vergu, Implications of the Landau equations for iterated integrals, *Phys. Rev. D* **105**, L061701 (2022), arXiv:2109.09744 [hep-th].
- [113] H. S. Hannesdottir and S. Mizera, What is the $i\epsilon$ for the S-matrix?, (2022), arXiv:2204.02988 [hep-th].
- [114] M. Mühlbauer, Cutkosky's Theorem for Massive One-Loop Feynman Integrals - Part 1, (2022), arXiv:2206.08402 [math-ph].
- [115] H. S. Hannesdottir, A. J. McLeod, M. D. Schwartz, and C. Vergu, Constraints on Sequential Discontinuities from the Geometry of On-shell Spaces, (2022), arXiv:2211.07633 [hep-th].

- [116] J. L. Bourjaily, C. Vergu, and M. von Hippel, Landau Singularities and Higher-Order Roots, (2022), arXiv:2208.12765 [hep-th].
- [117] M. Mühlbauer, On the Homology of Unions of Certain Non-Degenerate Quadrics in General Position, (2022), arXiv:2211.06683 [math-ph].
- [118] A. Kotikov and L. Lipatov, DGLAP and BFKL evolution equations in the $\mathcal{N} = 4$ supersymmetric gauge theory, in *35th Annual Winter School on Nuclear and Particle Physics* (2001) arXiv:hep-ph/0112346.
- [119] A. Kotikov and L. Lipatov, DGLAP and BFKL equations in the $\mathcal{N} = 4$ supersymmetric gauge theory, Nucl. Phys. B **661**, 19 (2003), [Erratum: Nucl.Phys.B 685, 405–407 (2004)], arXiv:hep-ph/0208220.
- [120] A. Kotikov, L. Lipatov, A. Onishchenko, and V. Velizhanin, Three loop universal anomalous dimension of the Wilson operators in $\mathcal{N} = 4$ SUSY Yang-Mills model, Phys. Lett. B **595**, 521 (2004), [Erratum: Phys.Lett.B 632, 754–756 (2006)], arXiv:hep-th/0404092.
- [121] A. Kotikov, L. Lipatov, A. Rej, M. Staudacher, and V. Velizhanin, Dressing and wrapping, J. Stat. Mech. **0710**, P10003 (2007), arXiv:0704.3586 [hep-th].
- [122] F. Wilczek, Decays of Heavy Vector Mesons Into Higgs Particles, Phys. Rev. Lett. **39**, 1304 (1977).
- [123] M. A. Shifman, A. Vainshtein, and V. I. Zakharov, Remarks on Higgs Boson Interactions with Nucleons, Phys. Lett. B **78**, 443 (1978).
- [124] L. J. Dixon, E. Glover, and V. V. Khoze, MHV rules for Higgs plus multi-gluon amplitudes, JHEP **12**, 015, arXiv:hep-th/0411092.
- [125] T. Gehrmann, M. Jaquier, E. Glover, and A. Koukoutsakis, Two-Loop QCD Corrections to the Helicity Amplitudes for $H \rightarrow 3$ partons, JHEP **02**, 056, arXiv:1112.3554 [hep-ph].
- [126] Y. Guo, Q. Jin, L. Wang, and G. Yang, Deciphering the maximal transcendentality principle via bootstrap, JHEP **09**, 161, arXiv:2205.12969 [hep-th].
- [127] E. W. N. Glover, P. Mastrolia, and C. Williams, One-loop phi-MHV amplitudes using the unitarity bootstrap: The General helicity case, JHEP **08**, 017, arXiv:0804.4149 [hep-ph].
- [128] S. Badger, E. W. Nigel Glover, P. Mastrolia, and C. Williams, One-loop Higgs plus four gluon amplitudes: Full analytic results, JHEP **01**, 036, arXiv:0909.4475 [hep-ph].
- [129] S. D. Badger, E. W. N. Glover, and K. Risager, One-loop phi-MHV amplitudes using the unitarity bootstrap, JHEP **07**, 066, arXiv:0704.3914 [hep-ph].
- [130] S. D. Badger and E. W. N. Glover, One-loop helicity amplitudes for $H \rightarrow$ gluons: The All-minus configuration, Nucl. Phys. B Proc. Suppl. **160**, 71 (2006), arXiv:hep-ph/0607139.
- [131] Z. Bern, L. J. Dixon, and D. A. Kosower, Two-loop $g \rightarrow gg$ splitting amplitudes in QCD, JHEP **08**, 012, arXiv:hep-ph/0404293.