Wave turbulence and the kinetic equation beyond leading order

Vladimir Rosenhaus¹ and Michael Smolkin²

¹ Initiative for the Theoretical Sciences The Graduate Center, CUNY 365 Fifth Ave, New York, NY 10016, USA

² The Racah Institute of Physics The Hebrew University of Jerusalem Jerusalem 91904, Israel

We derive a scheme by which to solve the Liouville equation perturbatively in the nonlinearity, which we apply to weakly nonlinear classical field theories. Our solution is a variant of the Prigogine diagrammatic method, and is based on an analogy between the Liouville equation in infinite volume and scattering in quantum mechanics, described by the Lippmann-Schwinger equation. The motivation for our work is wave turbulence: a broad class of nonlinear classical field theories are believed to have a stationary turbulent state — a far-from-equilibrium state, even at weak coupling. Our method provides an efficient way to derive properties of the weak wave turbulent state. A central object in these studies, which is a reduction of the Liouville equation, is the kinetic equation, which governs the occupation numbers of the modes. All properties of wave turbulence to date are based on the kinetic equation found at leading order in the weak nonlinearity. We explicitly obtain the kinetic equation to next-to-leading order.

Contents

1. Introduction	1
2. A perturbative solution of the Liouville equation	2
3. Quartic field theory: leading order	7
4. Quartic field theory: next-to-leading order	10
5. Discussion	16
A. From the master equation to the kinetic equation	n 17
B. Frequency renormalization	19

1. Introduction

This paper will present a scheme by which to solve the Liouville equation for a classical weakly nonlinear system in infinite volume, perturbatively in the nonlinearity. Our expansion will be analogous to the construction of scattering states in quantum mechanics. We will apply our method to weakly nonlinear field theories and derive new results for the kinetic equation governing the occupation numbers of the modes, which is central to studies of weak wave turbulence [1–3]. Wave turbulence has been shown to occur in an incredible range of contexts, from surface gravity waves to waves on vibrating elastic plates, see [4–23] for some recent work.

The Liouville equation is perhaps most familiar in the context of kinetic theory of a large number of interacting particles. Under a range of assumptions, primarily that the gas is dilute, the Liouville equation becomes the Boltzmann equation, governing the single particle phase space density. The corrections to the Boltzmann equation away from the dilute gas limit can be large and give qualitatively new effects [24–26], and have been studied in a number of works [27–35].

The Liouville equation can just as well be studied for interacting waves, rather than interacting particles. In the weak interaction limit it reduces to the wave kinetic equation, the wave analog of the Boltzmann equation [36]. In addition to the thermal state there is, surprisingly, a second stationary solution – a far-from-equilibrium turbulent state (the Kolmogorov-Zakharov state). The properties of this turbulent state, such as the occupation numbers of the modes and the correlations between modes, are largely unknown, except at leading order in the weak interaction. However, experience from other contexts, such as the kinetic theory of gases or perturbative quantum field theory, suggests that going beyond leading order is possible, valuable, and can lead to qualitatively

new effects. There has been a recent push to study the wave turbulent state beyond leading order [4,5], and this work is in line with this motivation.

Long ago Prigogine observed that the Liouville equation can be viewed as a Schrödinger equation, and used time-dependent perturbation theory to study the approach to equilibrium, going beyond the Boltzmann equation [34]. We build on Prigogine's work, with the distinction that we draw an analogy between stationary solutions of the Liouville equation and solutions of the time-independent Schrödinger equation. More specifically, with a continuum of modes, the Liouville equation is analogous to the Lippmann-Schwinger equation, used in quantum mechanics to describe scattering states. We proceed to solve the Liouville equation, formulated as a Lippmann-Schwinger equation, perturbatively in the nonlinearity. We do this explicitly for field theories with a weak quartic interaction, to the first two orders in the nonlinearity.

The paper is organized as follows. In Sec. 2 we view the Liouville equation as a Schrödinger equation and set up the perturbation theory for the solution. In Sec. 3 we specialize to Hamiltonians for waves with a weak quartic interaction and work out the perturbative solution of the Liouville equation at leading order in the quartic interaction, reproducing the well-known kinetic equation for waves. In Sec. 4 we extend this to next-to-leading order in the interaction and, in particular, derive the kinetic equation to next-to-leading order. We conclude in Sec. 5.

2. A perturbative solution of the Liouville equation

The statistical mechanics of N degrees of freedom is concerned with solving for the evolution of the phase space density $\rho(J_i, \alpha_i, t)$, a function of the action, J_i , and angle, α_i , variables, with i running from 1 to N. The evolution is governed by the Liouville equation,

$$\frac{\partial \rho}{\partial t} = \{H, \rho\} \ . \tag{2.1}$$

One is often interested in stationary solutions, which describe the late-time behavior. For almost any interacting system, the expectation is that this stationary solution is the thermal state. Surprisingly, for many field theories there is an additional stationary solution, which is the "wave turbulent" state. Concretely, what is known is this: for a field theory, at weak coupling, the stationary solution of the Liouville equation is of the form,

$$\rho(J) = \exp\left(-\sum_{p} \frac{J_p}{n_p}\right), \qquad \langle J_p \rangle \equiv \int dJ J_p \, \rho(J) = n_p, \qquad (2.2)$$

where the sum is over all Fourier modes p. The thermal solution has the occupation numbers,

$$n_p = \frac{T}{\omega_p} \,\,, \tag{2.3}$$

which is the Rayleigh-Jeans solution, where T is the temperature (this is simply the Bose-Einstein occupation number at large temperatures) and ω_p is the energy of mode p. For certain Hamiltonians, there is a second solution – the wave turbulent, or Kolmogorov-Zakharov, solution – with ρ of the form (2.2), but with an n_p that scales as a power of p,

$$n_p \sim p^{-\gamma} \,\,\,(2.4)$$

where γ depends on the form of the interaction. This is a stationary solution which is very different from the thermal solution; it is a state that is far from equilibrium.

The thermal state at finite coupling is similar to the one at weak coupling: it is of the form (2.2), but the ω_p appearing in n_p in (2.3) is replaced with the full energy of mode p. On the other hand, the density $\rho(J_i, \alpha_i)$ for the turbulent state beyond leading order in the coupling is far more interesting and elaborate: it will be a complicated function of the J_i and the α_i . Finding $\rho(J_i, \alpha_i)$ is a major problem, and this will be our goal: to systematically solve the Liouville equation, perturbatively in the coupling, for a time-independent ρ .

A way of organizing the perturbation series is to notice that the Liouville equation can be viewed as a Schrödinger equation. Explicitly, after multiplication by i, (2.1) becomes,

$$i\frac{\partial \rho}{\partial t} = L\rho , \qquad (2.5)$$

where

$$L = i \sum_{j} \left(\frac{\partial H}{\partial \alpha_{j}} \frac{\partial}{\partial J_{j}} - \frac{\partial H}{\partial J_{j}} \frac{\partial}{\partial \alpha_{j}} \right) . \tag{2.6}$$

A notable difference with the Schrödinger equation is that the phase space density $\rho(J_i, \alpha_i)$ depends on both the action and the angle variables, unlike the wavefunction which depends on either the positions or the momenta.

We are interested in stationary solutions of the Liouville equation. This amounts to finding ρ for which $L\rho = 0$; or, in the language of the Schrödinger equation, solving the time-independent Schrödinger equation for eigenfunctions with zero eigenvalue. There are two kinds of eigenfunctions of the Schrödinger equation: bound states and scattering states. It is the scattering states that we are interested in. In quantum mechanics one finds the scattering state eigenfunctions by solving the Lippmann-Schwinger equation. Let us recall how this works; we will then do something analogous for the Liouville equation.

Lippmann-Schwinger equation

The Hamiltonian in quantum mechanics is split into a free part and an interacting part, $H = H_0 + V$, so that the Schrödinger equation is,

$$(H_0 + V)|\psi\rangle = E|\psi\rangle . (2.7)$$

The solution is formally [37],

$$|\psi\rangle = |\psi_0\rangle + G(E)V|\psi\rangle , \quad G(E) = \frac{1}{E - H_0 + i\epsilon} ,$$
 (2.8)

where $|\psi_0\rangle$ is the solution of the free Schrödinger equation, $H_0|\psi\rangle=E|\psi\rangle$. ¹ Eq. 2.8 is the Lippmann-Schwinger equation. Iterating the equation gives a perturbative expansion for the wavefunction,

$$|\psi\rangle = |\psi_0\rangle + G(E)V|\psi_0\rangle + G(E)VG(E)V|\psi_0\rangle + \dots$$
 (2.9)

It is useful to write the Lippmann-Schwinger equation in position space,

$$\psi(x) = \psi_0(x) + \int dx' G(x, x') V(x') \psi(x') , \quad G(x, x') = \langle x | G(E) | x' \rangle , \qquad (2.10)$$

where the position-space wavefunction is $\psi(x) = \langle x|\psi\rangle$, and the Green's function is $G(x,x') = \langle x|G(E)|x'\rangle$, while the potential is taken to be local, $\langle x|V|x'\rangle = V(x)\delta(x-x')$. For H_0 corresponding to a free particle, $H_0 = p^2/2m$, the propagator is,

$$G_{\pm}(x,x') = \int \frac{d^3p}{(2\pi)^3} \frac{e^{i\vec{p}\cdot(\vec{x}-\vec{x}')}}{E - \frac{p^2}{2m} \pm i\epsilon} = -\frac{m}{2\pi} \frac{e^{ik|\vec{x}-\vec{x}'|}}{|\vec{x}-\vec{x}'|} , \quad E = \frac{k^2}{2m} .$$
 (2.11)

Finally, we note that the Lippmann-Schwinger equation applies to scattering states, which have a continuum of energy levels. This was important to us when we identified the energy of the eigenstates of both the free Hamiltonian and the interacting Hamiltonian as E. This is to be contrasted with bound states: the bound state energy levels are discretely spaced and change as the Hamiltonian changes.

¹The $i\epsilon$ that we added sets the boundary conditions. In particular, we can view the $i\epsilon$ as being added to the energy and making it slightly complex, $E \to E + i\epsilon$. The eigenfunctions have the time evolution e^{-iEt} , which becomes $e^{-iEt}e^{\epsilon t}$. This vanishes in the far past, which is what we want: we want the far past to have eigenfunctions of the free Schrödinger equation. The choice of $+i\epsilon$ which we made corresponds to a retarded propagator. It means that a plane wave is ingoing and a scattered wave is outgoing. The choice of $-i\epsilon$ corresponds to an advanced propagator. Notice that the $i\epsilon$ is absent once we Fourier transform to position space in (2.11). If we had picked $-i\epsilon$ in (2.8), instead of $+i\epsilon$, the exponential in (2.11) would have had a minus sign, leading to a scattered wave that is ingoing.

Liouville equation

We now return to the Liouville equation (2.5) and split the Hamiltonian into a free part, which depends only on the action variables, and an interacting part,

$$H = H_0(J_i) + V(J_i, \alpha_i)$$
 (2.12)

The Liouville operator is correspondingly $L = L_0 + \delta L$ where,

$$L_0 = -i \sum_{j} \omega(J_j) \frac{\partial}{\partial \alpha_j} , \qquad \omega(J_j) \equiv \frac{\partial H_0}{\partial J_j} , \qquad (2.13)$$

$$\delta L = i \sum_{j} \left(\frac{\partial V}{\partial \alpha_{j}} \frac{\partial}{\partial J_{j}} - \frac{\partial V}{\partial J_{j}} \frac{\partial}{\partial \alpha_{j}} \right) . \tag{2.14}$$

Our goal is to solve the Liouville equation for a time-independent phase space density ρ . We denote such a state in ket notation as $|\Phi\rangle$, so that $\rho(J_i, \alpha_i) = \langle J_i, \alpha_i | \Phi \rangle$. As a result of the Liouville equation (2.5) $|\Phi\rangle$ must satisfy,

$$(L_0 + \delta L)|\Phi\rangle = 0. (2.15)$$

In analogy with the Lippmann-Schwinger equation (2.8), the solution of (2.15) is,

$$|\Phi\rangle = |\Phi_0\rangle + G\delta L|\Phi\rangle , \quad G = \frac{1}{-L_0 + i\epsilon} ,$$
 (2.16)

where $|\Phi_0\rangle$ solves $L_0|\Phi_0\rangle = 0$. This is just like (2.8) with L_0 instead of H_0 and with E = 0. Here the propagator (or Green's function) G is the inverse of the free Liouville operator L_0 . Let us look at the Green's function in more detail. Writing its matrix element as, $\langle \vec{\alpha} | G | \vec{\alpha}' \rangle = G(\vec{\alpha} - \vec{\alpha}')$, we have,

$$L_0G(\vec{\alpha}) = -\delta(\vec{\alpha}) , \qquad G(\vec{\alpha}) = \sum_{\vec{\alpha}} \frac{e^{i\vec{n}\cdot\vec{\alpha}}}{-\vec{n}\cdot\vec{\omega} + i\epsilon} ,$$
 (2.17)

where $\delta(\vec{\alpha}) \equiv \prod_j \delta(\alpha_j)$. Here \vec{n} is a vectors of integers, and we have switched to vector notation, $\vec{J}, \vec{\alpha}$ to denote the J_i, α_i . We denote the Fourier transform of the Green's function by $G(\vec{n})$,

$$G(\vec{n}) = \frac{1}{-\vec{n} \cdot \vec{\omega} + i\epsilon} \ . \tag{2.18}$$

Notice that the eigenfunctions $|\vec{n}\rangle$ of the free Liouville operator L_0 are plane waves,

$$\langle \vec{\alpha} | \vec{n} \rangle = \exp(i\vec{n} \cdot \vec{\alpha}) , \qquad L_0 | \vec{n} \rangle = \vec{n} \cdot \vec{\omega} | \vec{n} \rangle .$$
 (2.19)

We could multiply this by any function $\rho(\vec{J})$ of the action variables and it would continue to be an eigenfunction of L_0 . We are particularly interested in the eigenfunctions of L_0 with zero eigenvalue,

as they correspond to the solutions $|\Phi_0\rangle$ of the free Liouville equation,

$$\langle \vec{\alpha} | \Phi_0 \rangle = \exp(i\vec{n} \cdot \vec{\alpha}) \, \rho(\vec{J}) \,, \quad \vec{n} \cdot \vec{\omega} = 0 \,,$$
 (2.20)

where $\rho(\vec{J})$ is any function of the action variables. Even though this a function of the \vec{J} variables as well, for notational simplicity we omit writing an explicit \vec{J} on the left-hand side. Also, to be clear, we wrote the sum in (2.13) and (2.14) to be over a seemingly discrete index j, however we really have in mind that j is momentum, which is continuous. For instance, the true meaning of $\vec{n} \cdot \vec{\omega}$ is,

$$\vec{n} \cdot \vec{\omega} \equiv \sum_{i} n_i \omega_i = \sum_{p} n_p \omega_p = \int d^d p \, n_p \, \omega_p \ . \tag{2.21}$$

We will continue to use the $\vec{n} \cdot \vec{\omega}$ notation due to its simplicity. The $\vec{n} \cdot \vec{\omega}$ is evidently analogous to energy in the discussion of the Lippmann-Schwinger equation.

The Liouville equation (2.16), written in the plane wave basis, is,

$$\langle \vec{n} | \Phi \rangle = \langle \vec{n} | \Phi_0 \rangle + \sum_{\vec{n}'} G(\vec{n}) \langle \vec{n} | \delta L | \vec{n}' \rangle \langle \vec{n}' | \Phi \rangle , \qquad (2.22)$$

where we used that $\langle \vec{n}|G|\vec{n}'\rangle = \delta_{n,n'}G(\vec{n})$. The transformation between the state in the two bases, $|\vec{\alpha}\rangle$ and $|\vec{n}\rangle$, is as usual through a Fourier transform,

$$\langle \vec{\alpha} | \Phi \rangle = \sum_{\vec{n}} \langle \vec{\alpha} | \vec{n} \rangle \langle \vec{n} | \Phi \rangle = \sum_{\vec{n}} e^{i\vec{n}\cdot\vec{\alpha}} \langle \vec{n} | \Phi \rangle . \tag{2.23}$$

Just as we did for the Lippmann-Schwinger equation (2.9), we may perturbatively expand $|\Phi\rangle$ around $|\Phi_0\rangle$,

$$\langle \vec{n} | \Phi \rangle = \langle \vec{n} | \Phi_0 \rangle + \sum_{\vec{n}'} G(\vec{n}) \langle \vec{n} | \delta L | \vec{n}' \rangle \langle n' | \Phi_0 \rangle + \sum_{\vec{n}', \vec{n}''} G(\vec{n}) \langle \vec{n} | \delta L | \vec{n}' \rangle G(\vec{n}') \langle \vec{n}' | \delta L | \vec{n}'' \rangle \langle \vec{n}'' | \Phi_0 \rangle + \dots (2.24)$$

This is an expression that builds the solution $|\Phi\rangle$ of the full Liouville equation in terms of a solution $|\Phi_0\rangle$ of the free Liouville equation. Let us take a special case: $\langle \vec{\alpha} | \Phi_0 \rangle$ which is independent of $\vec{\alpha}$, ²

$$\langle \vec{\alpha} | \Phi_0 \rangle = \rho(\vec{J}) \ . \tag{2.25}$$

The Fourier transform $\langle \vec{n} | \Phi_0 \rangle$ is correspondingly only nonzero for $\vec{n} = 0$, with $\langle \vec{n} = 0 | \Phi_0 \rangle = \rho(J)$.

²We do this because later on we will be interested in $\langle J_r \rangle = \int dJ d\alpha J_r \langle \alpha | \Phi_0 \rangle$, which will vanish unless $\langle \alpha | \Phi_0 \rangle$ (2.20) is independent of α .

We therefore set $\vec{n} = 0$ in (2.24) so that it becomes,

$$\langle 0|\Phi\rangle = \langle 0|\Phi_0\rangle + G(0)\langle 0|\delta L|0\rangle\langle 0|\Phi_0\rangle + \sum_{\vec{n}'} G(0)\langle 0|\delta L|\vec{n}'\rangle G(\vec{n}')\langle \vec{n}'|\delta L|0\rangle\langle 0|\Phi_0\rangle + \dots$$
 (2.26)

In the next section we will specialize these equations to a field theory with a quartic interaction.

3. Quartic field theory: leading order

We now turn to our Hamiltonian of interest, for a field theory with a quartic interaction. Rather than working with real fields ϕ_p and momentum conjugate π_p , it is common to work with complex fields a_p and their complex conjugate a_p^* , which we will denote by a_p^{\dagger} . The Hamiltonian is taken to be [2],

$$H = \sum_{p} \omega_{p} a_{p}^{\dagger} a_{p} + \sum_{p_{1}, p_{2}, p_{3}, p_{4}} \lambda_{p_{1} p_{2} p_{3} p_{4}} a_{p_{1}}^{\dagger} a_{p_{2}}^{\dagger} a_{p_{3}} a_{p_{4}} , \qquad (3.1)$$

where ω_p is an arbitrary real function of p and the coupling $\lambda_{p_1p_2p_3p_4}$ is any complex function of the momenta p_1, p_2, p_3, p_4 which has the symmetry properties $\lambda_{p_1p_2p_3p_4} = \lambda_{p_2p_1p_3p_4} = \lambda_{p_1p_2p_4p_3} = \lambda_{p_3p_4p_1p_2}^*$. Phase space in these variables is a function of the a_p and a_p^{\dagger} . It is convenient to switch to action-angle variables, J_p, α_p ,

$$a_p = \sqrt{J_p}e^{-i\alpha_p} , \quad a_p^{\dagger} = \sqrt{J_p}e^{i\alpha_p} , \qquad (3.2)$$

so that the Hamiltonian becomes,

$$H = \sum_{i} \omega_{i} J_{i} + \sum_{i,j,k,l} \lambda_{ijkl} \sqrt{J_{i} J_{j} J_{k} J_{l}} \exp\left(i(\alpha_{i} + \alpha_{j} - \alpha_{k} - \alpha_{l})\right) , \qquad (3.3)$$

where we have switched to labeling the momenta p by the discrete index i, for notational convenience. The advantage of action-angle variables is that the free part of the Hamiltonian depends only on the action variables.

Computing the Liouville operator, we have that L_0 is (2.13) with $\omega(J_i) = \omega_i$, while δL in (2.14) is,

$$\delta L = -\sum_{i,j,k,l} \lambda_{ijkl} \sqrt{J_i J_j J_k J_l} e^{-i\vec{e}_{ij;kl} \cdot \vec{\alpha}} \left[\left(\partial_i + \partial_j - \partial_k - \partial_l \right) + \frac{i}{2} \left(\frac{\partial_{\alpha_i}}{J_i} + \frac{\partial_{\alpha_j}}{J_j} + \frac{\partial_{\alpha_k}}{J_k} + \frac{\partial_{\alpha_l}}{J_l} \right) \right] , \quad (3.4)$$

where we are using the shorthand $\partial_i \equiv \partial_{J_i}$. In addition, we have defined $\vec{e}_{ij;kl}$, which is a vector that has a -1 in the *i*'th and *j*'th entry and a +1 in the *k*'th and *l*'th entry, and a zero elsewhere;

³The relation between them is, as usual, $\phi_k = \frac{1}{2\sqrt{\omega_k}}(a_k + a_k^{\dagger})$ and $\pi_k = \frac{i}{2\sqrt{\omega_k}}(a_k^{\dagger} - a_k)$.

in other words,

$$-\vec{e}_{ij:kl} \cdot \vec{\alpha} = \alpha_i + \alpha_j - \alpha_k - \alpha_l . \tag{3.5}$$

The matrix element of δL trivially follows,

$$\langle \vec{n} | \delta L | \vec{n}' \rangle = -\sum_{i,j,k,l} \lambda_{ijkl} \sqrt{J_i J_j J_k J_l} \, \delta_{\vec{n} + \vec{e}_{ij;kl}, \vec{n}'} \left[\left(\partial_i + \partial_j - \partial_k - \partial_l \right) - \frac{1}{2} \left(\frac{n_i'}{J_i} + \frac{n_j'}{J_j} + \frac{n_k'}{J_k} + \frac{n_l'}{J_l} \right) \right] . \quad (3.6)$$

An equivalent way of writing this, which will also be useful is,

$$\langle \vec{n} | \delta L | \vec{n}' \rangle = -\sum_{i,j,k,l} \lambda_{ijkl} \, \delta_{\vec{n} + \vec{e}_{ij;kl},\vec{n}'} \left[\left(\partial_i + \partial_j - \partial_k - \partial_l \right) - \frac{1}{2} \left(\frac{n_i}{J_i} + \frac{n_j}{J_j} + \frac{n_k}{J_k} + \frac{n_l}{J_l} \right) \right] \sqrt{J_i J_j J_k J_l} , \quad (3.7)$$

where we used the commutation relation $\left[\partial, \sqrt{J}\right] = \frac{1}{2J}\sqrt{J}$ to go between the two expressions. Now, using these matrix elements in (2.26) gives,

$$\langle 0|\Phi\rangle = \langle 0|\Phi_0\rangle + G(0)\sum_{ijkl}|\lambda_{ijkl}|^2G(\vec{e}_{ij;kl})\left(\partial_i + \partial_j - \partial_k - \partial_l\right)J_iJ_jJ_kJ_l\left(\partial_k + \partial_l - \partial_i - \partial_j\right)\langle 0|\Phi_0\rangle + \dots ,$$
(3.8)

where we used that, in order for the matrix element $\langle \vec{0}|\delta L|\vec{n}'\rangle$ to be nonzero, the intermediate state $|\vec{n}\rangle$ has to take the form $|\vec{ij};kl\rangle$, by which we mean the mode occupation numbers are: $n_i = n_j = -1$ and $n_k = n_l = 1$, and all other $n_a = 0$.

Notice that the second term in (3.8) has a G(0). From the form of $G(\vec{n})$ given in (2.18), we see that G(0) diverges in the limit that ϵ goes to zero. This means that the quantity multiplying G(0) must vanish,

$$\sum_{ijkl} |\lambda_{ijkl}|^2 \frac{1}{\omega_i + \omega_j - \omega_k - \omega_l + i\epsilon} \left(\partial_i + \partial_j - \partial_k - \partial_l \right) J_i J_j J_k J_l \left(\partial_k + \partial_l - \partial_i - \partial_j \right) \rho(\vec{J}) = 0 , \qquad (3.9)$$

where we used the explicit form of $G(\vec{e}_{ij;kl})$ and replaced $\langle 0|\Phi_0\rangle$ with $\rho(\vec{J})$. Next, we note that,

$$\frac{1}{x+i\epsilon} = P\frac{1}{x} - i\pi\delta(x) , \qquad (3.10)$$

where P denotes principal value. Since the principal part of $1/(\omega_k+\omega_l-\omega_i-\omega_j)$ is odd under exchange of i, j with k, l, whereas the other terms in (3.9) are even, we are left with the delta function piece,

$$\sum_{ijkl} |\lambda_{ijkl}|^2 \delta(\omega_i + \omega_j - \omega_k - \omega_l) \left(\partial_i + \partial_j - \partial_k - \partial_l\right) J_i J_j J_k J_l \left(\partial_k + \partial_l - \partial_i - \partial_j\right) \rho(\vec{J}) = 0 . \tag{3.11}$$

Before proceeding further with analyzing this equation, let us take stock of where we are. Our problem was to solve the Liouville equation. We wrote this in the form (2.16) resembling the Lippmann-Schwinger equation, which gives the solution as a series built off of the solution of the free Liouville equation. We took a special solution of the free Liouville equation $\langle \vec{\alpha} | \Phi_0 \rangle = \rho(\vec{J})$ (2.25) which is independent of the angles $\vec{\alpha}$. However we found that, rather than (2.16) giving us corrections to this, perturbatively in the coupling, as one is used to in quantum mechanics, it gave us a constraint on $\rho(\vec{J})$: at leading order it must satisfy (3.11).

The equation (3.11) is not new: it was found by Prigogine [34]. He studied what one might call the analog of time-dependent perturbation theory for the Liouville equation, whereas we are studying the analog of time-independent perturbation theory. Prigogine found,

$$\frac{d\rho(\vec{J})}{\partial t} = -4\pi \sum_{ijkl} |\lambda_{ijkl}|^2 \delta(\omega_i + \omega_j - \omega_j - \omega_l) \left(\partial_i + \partial_j - \partial_k - \partial_l\right) J_i J_j J_k J_l \left(\partial_k + \partial_l - \partial_i - \partial_j\right) \rho(\vec{J}) , \quad (3.12)$$

which of course agrees with (3.11) if one is looking for stationary $\rho(\vec{J})$, as we are. Let us see how (3.12) reduces to the kinetic equation. Defining the expectation values,

$$\langle J_i \rangle = \int dJ J_i \, \rho(J) \;, \quad \langle J_i J_j J_k \rangle = \int dJ J_i \, J_j \, J_k \, \rho(J) \;,$$
 (3.13)

and multiplying (3.12) by J_r and integrating with respect to \vec{J} we get,

$$\frac{\partial \langle J_r \rangle}{\partial t} = -4\pi \int dJ J_r \sum_{i,j,k,l} |\lambda_{ijkl}|^2 \delta(\omega_i + \omega_j - \omega_k - \omega_l) \left(\partial_i + \partial_j - \partial_k - \partial_l\right) J_i J_j J_k J_l \left(\partial_k + \partial_l - \partial_i - \partial_j\right) \rho(J) . \tag{3.14}$$

We integrate by parts to the left (twice) and identify the resulting integrals as expectation values,

$$\frac{\partial \langle J_r \rangle}{\partial t} = 4\pi \sum_{i,j,k,l} \left(\delta_{ir} + \delta_{jr} - \delta_{kr} - \delta_{lr} \right) |\lambda_{ijkl}|^2 \delta(\omega_i + \omega_j - \omega_k - \omega_l) \left(\langle J_j J_k J_l \rangle + \langle J_i J_k J_l \rangle - \langle J_i J_j J_l \rangle - \langle J_i J_j J_k \rangle \right) ,$$
(3.15)

where δ_{ij} is the Kronecker delta function. Now, if we take $\rho(J)$ to be an exponential,

$$\rho(J) = \frac{1}{\prod_{i} n_{i}} \exp\left(-\sum_{i} \frac{J_{i}}{n_{i}}\right) , \qquad \langle J_{i} \rangle = n_{i} , \qquad (3.16)$$

then all the correlation functions factorize into one-point functions and we are left with,

$$\frac{\partial n_r}{\partial t} = 4\pi \sum_{i,i,k,l} \left(\delta_{ir} + \delta_{jr} - \delta_{kr} - \delta_{lr} \right) \left| \lambda_{ijkl} \right|^2 \delta(\omega_i + \omega_j - \omega_k - \omega_l) \left(\frac{1}{n_i} + \frac{1}{n_j} - \frac{1}{n_k} - \frac{1}{n_l} \right) n_i n_j n_k n_l . \quad (3.17)$$

This is the standard (leading order) kinetic equation for waves [2].

Using our perturbative (in the coupling) solution of the Liouville equation analog of the Lippmann-Schwinger equation, we can go beyond the leading order kinetic equation, and compute

the kinetic equation to any order in the coupling. Generalizing what we just did, from (2.22) we have that,

$$\langle \vec{0} | \Phi \rangle = \langle \vec{0} | \Phi_0 \rangle + G(\vec{0}) \sum_{\vec{n}'} \langle \vec{0} | \delta L | \vec{n}' \rangle \langle n' | \Phi \rangle . \tag{3.18}$$

Since $G(\vec{0})$ is divergent, it must be the case that,

$$\sum_{\vec{n}'} \langle \vec{0} | \delta L | \vec{n}' \rangle \langle n' | \Phi \rangle = 0 . \tag{3.19}$$

We will refer to this as the master equation. We may multiply this by J_r and integrate over \vec{J} , and identify the resulting quantity as proportional to $\frac{\partial n_r}{\partial t}$,

$$\frac{\partial n_r}{\partial t} \propto \int dJ J_r \sum_{\vec{n}'} \langle \vec{0} | \delta L | \vec{n}' \rangle \langle n' | \Phi \rangle . \tag{3.20}$$

This is the kinetic equation: for a stationary state, both sides vanish, as should be the case. We therefore have a clear procedure for computing the kinetic equation to any order in the interaction strength: we perturbatively solve (2.16) for $|\Phi\rangle$ with the initial condition that $|\Phi_0\rangle$ is angle independent i.e, $\langle \vec{n} | \Phi_0 \rangle$ is nonzero for $\vec{n} = 0$, and insert the result into (3.20). This is what we will do in the next section.

We end this section with a comment. Perturbatively solving the Liouville equation is equivalent to perturbatively computing integrals of motion, see [38–40]. The connection with the Lippmann-Schwinger equation gives a systematic and unambiguous scheme by which to compute the kinetic equation to any order in the nonlinearity. It is important that we are studying field theories in the continuum, which have an infinite number of degrees of freedom. One might have worried that once one goes away from the free limit, integrability will be broken (perturbation theory will break down) and there will be no conserved quantities. However, since we have a field theory, any function $\rho(\vec{J})$ is conserved for the free theory; once interactions are turned on most functions indeed stop being conserved, but some remain conserved: those that satisfy the equations, such as (3.9), that our perturbation theory forces on us.

4. Quartic field theory: next-to-leading order

In this section we continue the discussion in the previous section, but go to next-to-leading order in the nonlinearity. To recap: our goal is to solve the Liouville equation for the phase space distribution $\Phi(\vec{J}, \vec{\alpha})$, which is a function of the action J_i and angle α_i variables of all the modes, which have been grouped into vectors, \vec{J} and $\vec{\alpha}$. We work perturbatively in the interaction, starting with the solution $\Phi_0(\vec{J}, \vec{\alpha})$ of the Liouville equation for a free Hamiltonian.

We are particularly interested in the case in which Φ_0 is independent of the angle. It is convenient to change basis, from angles $\vec{\alpha}$ to the Fourier transform variables of occupation numbers \vec{n} . The Φ_0 that is $\vec{\alpha}$ independent then only depends on the zero mode, $\vec{n} = 0$. Writing Φ_0 in ket notation, it is $\langle \vec{n} = 0 | \Phi_0 \rangle$. Going back to our perturbative expansion (2.26) for $\langle 0 | \Phi \rangle$, we add the next-to-leading order term,

$$\langle 0|\Phi\rangle = \langle 0|\Phi_0\rangle + G(0)((\delta\mathcal{L})_{\text{first}} + (\delta\mathcal{L})_{\text{second}} + \dots), \qquad (4.1)$$

where we have dropped the $\langle 0|\delta L|0\rangle$ term, since it vanishes for our δL , and we defined,

$$(\delta \mathcal{L})_{\text{first}} = \sum_{\vec{n}'} \langle \vec{0} | \delta L | \vec{n}' \rangle G(\vec{n}') \langle \vec{n}' | \delta L | 0 \rangle \langle 0 | \Phi_0 \rangle$$

$$(4.2)$$

$$(\delta \mathcal{L})_{\text{second}} = \sum_{\vec{n}', \vec{n}''} \langle \vec{0} | \delta L | \vec{n}' \rangle G(\vec{n}') \langle \vec{n}' | \delta L | \vec{n}'' \rangle G(\vec{n}'') \langle \vec{n}'' | \delta L | 0 \rangle \langle 0 | \Phi_0 \rangle . \tag{4.3}$$

As argued in the previous section, since G(0) diverges we must have the master equation,

$$(\delta \mathcal{L})_{\text{first}} + (\delta \mathcal{L})_{\text{second}} = 0$$
, (4.4)

to third order in the coupling.

In the previous section we computed the first term $(\delta \mathcal{L})_{\text{first}}$, see (3.11). Here we will compute the next order term, $(\delta \mathcal{L})_{\text{second}}$. Before proceeding, it is useful to streamline our notation and give graphical representations of the terms that appear. We will label modes by $1, 2, 3, \ldots$ instead of i, j, k, \ldots . In addition, we can make use of the specific form of δL for a field theory with a quartic interaction in order to specify which states can appear in the perturbative expansion. Namely, δL acts by "creating" two particles and "destroying" two other particles, and its matrix element was given earlier in (3.6). We have that $(\delta \mathcal{L})_{\text{first}}$ (4.2) reduces to,

$$(\delta \mathcal{L})_{\text{first}} = \sum_{1,\dots,4} G(\vec{e}_{12;34}) \langle \vec{0} | \delta L | \overline{12}; 34 \rangle \langle \overline{12}; 34 | \delta L | 0 \rangle \langle 0 | \Phi_0 \rangle . \tag{4.5}$$

Our notation is such that when we write $|\overline{12};34\rangle$ we mean a state in which modes 1 and 2 have occupation number -1 and modes 3 and 4 have occupation number 1. We may represent (4.5) graphically, as shown in Fig. 1. One can imagine a vertical dashed line running through the center of the figure which represents the intermediate state $|\overline{12};34\rangle$. The arrows on the lines indicate the direction of flow of occupation number: an arrow pointing to the right corresponds to occupation number plus one, while an arrow pointing to the left corresponds to occupation number minus one.




Figure 1: A diagrammatic representation of $(\delta \mathcal{L})_{\text{first}}$ in (4.5). We start with vacuum $|0\rangle$. The interaction δL , which occurs at the left vertex, turns this into the state $|\overline{12};34\rangle$. The two "particles" and two "holes" each propagate to the right. The interaction δL then transforms the state back into the vacuum $|0\rangle$.

In this new notation (4.5) in explicit form is

$$(\delta \mathcal{L})_{\text{first}} = \sum_{1,\dots,4} G(\vec{e}_{12;34}) \lambda_{1234} \lambda_{3412} \left(\partial_1 + \partial_2 - \partial_3 - \partial_4 \right) J_1 J_2 J_3 J_4 \left(\partial_3 + \partial_4 - \partial_1 - \partial_2 \right) \left\langle 0 | \Phi_0 \right\rangle . \tag{4.6}$$

Let us now look at the next order term $(\delta \mathcal{L})_{\text{second}}$ (4.3). To compute this we need to enumerate all the possible intermediate states $|\vec{n}'\rangle$ and $|\vec{n}''\rangle$. The most efficient way to do this is with the aid of diagrams describing the process. The diagrams will have three vertices, one for each δL . There are four diagrams to consider. Two of the diagrams are shown below, in Fig. 2. We also need to




Figure 2: A diagrammatic representation of the contributions to $(\delta \mathcal{L})_{\text{second}}$.

add the same diagrams, with all the arrows reversed. These are shown below, in Fig. 3. We split




Figure 3: The same diagrams as in Fig. 2, but with all arrows reversed, also contribute to $(\delta \mathcal{L})_{\text{second}}$.

$$(\delta \mathcal{L})_{\text{second}}$$
 into
$$(\delta \mathcal{L})_{\text{second}} = (\delta \mathcal{L})_{\text{second}}^a + (\delta \mathcal{L})_{\text{second}}^b , \qquad (4.7)$$

where $(\delta \mathcal{L})_{\text{second}}^a$ denotes the contribution that is the sum of the diagrams Fig. 2(a) and Fig. 3(a), and $(\delta \mathcal{L})_{\text{second}}^b$ denotes the contribution that is the sum of the diagrams Fig. 2(b) and Fig. 3(b). Let us start with Fig. 2(a). This diagram can be cut in two different ways, producing two distinct intermediate states, see Fig 4.




Figure 4: The diagram shown in Fig. 2(a). We use a vertical dashed line to represent an insertion of an intermediate state. Depending on the two ways in which we rotate Fig. 2(a), we produce different intermediate states, as shown here in (i) and (ii). In (i) the first (leftmost) intermediate state that the vertical dashed line crosses is $|\overline{12};34\rangle$ and the second is $|\overline{56};34\rangle$. In (ii) the first state is also $|\overline{12};34\rangle$, but the second is $|\overline{12};56\rangle$.

We start with Fig. 4(i), which corresponds to a sequence of states $|0\rangle \rightarrow |\overline{12};34\rangle \rightarrow |\overline{56};34\rangle \rightarrow |0\rangle$. Its contribution is,

$$\sum_{1,\dots,6} G(0)G(\vec{e}_{12;34})G(\vec{e}_{56;34}) \langle 0|\delta L|\overline{12};34\rangle\langle \overline{12};34|\delta L|\overline{56};34\rangle\langle \overline{56};34|\delta L|0\rangle\langle 0|\Phi_0\rangle . \tag{4.8}$$

Let us evaluate this explicitly, making use of the matrix elements (3.6). We get,

$$-\sum_{1,\dots,6} G(0)G(\vec{e}_{12;34})G(\vec{e}_{56;34})\lambda_{1234}\lambda_{5612}\lambda_{3456} \left(\partial_{1}+\partial_{2}-\partial_{3}-\partial_{4}\right)J_{1}J_{2}\sqrt{J_{3}J_{4}J_{5}J_{6}}$$

$$\left[\left(\partial_{5}+\partial_{6}-\partial_{1}-\partial_{2}\right)+\frac{1}{2}\left(\frac{1}{J_{5}}+\frac{1}{J_{6}}\right)\right]\sqrt{J_{5}J_{6}J_{3}J_{4}}\left(\partial_{3}+\partial_{4}-\partial_{5}-\partial_{6}\right)\left\langle 0|\Phi_{0}\right\rangle ,\quad (4.9)$$

where we used the matrix element in the form of (3.7) for $\langle \vec{0}|\delta L|\overline{12};34\rangle$, and in the form of (3.6) for $\langle \overline{12};34|\delta L|\overline{56};34\rangle$ and $\langle \overline{56};34|\delta L|0\rangle$. It is convenient to commute through the J_5 and J_6 so as

to combine them. Using $\left[\partial, \sqrt{J}\right] = \frac{1}{2J}\sqrt{J}$ this gives,

$$\begin{split} -\sum_{1,\dots,6} G(0)G(\vec{e}_{12;34})G(\vec{e}_{56;34})\lambda_{1234}\lambda_{5612}\lambda_{3456} \left(\partial_{1}+\partial_{2}-\partial_{3}-\partial_{4}\right)J_{1}J_{2}J_{3}J_{4} \\ \left(\partial_{5}+\partial_{6}-\partial_{1}-\partial_{2}\right)J_{5}J_{6} \left(\partial_{3}+\partial_{4}-\partial_{5}-\partial_{6}\right)\left\langle 0|\Phi_{0}\right\rangle \;. \quad (4.10) \end{split}$$

Next, we note that from Fig. 3(a) we will have an identical contribution, but with all the arrows reversed. Rather than redo the computation, it is useful to understand how to get the result by a symmetry transformation.

Particle-hole symmetry

Let us understand what it means to have the arrows run in the opposite direction. For the couplings, it means $\lambda_{ijkl} \to \lambda_{klij} = \lambda_{ijkl}^*$. It also means that the sign of the angles is reversed, $\vec{\alpha} \to -\vec{\alpha}$. In terms of $G(\vec{\alpha})$, reversing the sign of $\vec{\alpha}$ corresponds in Fourier space to sending $G(\vec{n})$ to $G(-\vec{n})$. Since with our notation our occupation numbers are either plus or minus one, reversing arrows means $G(\vec{e}_{ij;kl}) \to G(\vec{e}_{kl:ij}) = -G^*(\vec{e}_{ij:kl})$.

Now, to see the effect of reversing the arrows on the matrix elements: we are swapping particles and holes, i.e $|ij;kl\rangle \to |kl;ij\rangle$. We also need to understand the effect of swapping the arrows on the explicit form of the matrix elements. In terms of the variables, we see from (3.6) that flipping the sign of $\vec{\alpha}$ corresponds to $\vec{J} \to -\vec{J}$.

There is an easier way of understanding the mapping: it follows from a symmetry of the Hamiltonian (3.3). In particular, the Hamiltonian is invariant under 4

$$\omega_i \to -\omega_i \;, \quad J_i \to -J_i \;, \quad \alpha_i \to -\alpha_i \;, \quad \lambda_{1234} \to \lambda_{-1-2-3-4} \equiv \lambda_{1234}^* \;, \quad i = 1, 2, 3, 4 \;.$$
 (4.11)

It is also invariant under a flip in sign of just two of the indices,

$$\omega_i \to -\omega_i$$
, $J_i \to -J_i$, $\alpha_i \to -\alpha_i$, $\lambda_{1234} \to \lambda_{1-2-34} \equiv \lambda_{1324}$, $i = 2, 3$. (4.12)

Now, returning to (4.10) and performing the described transformation gives the complex conjugate of (4.10), with a minus sign. Summing the two gives what (with foresight) we denote as

⁴We have introduced the notation that if one of the indices of the coupling appears with a minus sign in the first two slots, it means the same as the index appearing without a minus sign in one of the last two slots, and likewise if an index appears with a minus sign in the last two slots, it means the same as the index appearing without a minus sign in one of the first two slots, i.e. $\lambda_{1-2-34} \equiv \lambda_{1324}$ and $\lambda_{-1-2-3-4} = \lambda_{3412}$.

 $\frac{1}{2} \left(\delta \mathcal{L} \right)_{\text{second}}^{a} \text{ where,}$

$$(\delta \mathcal{L})_{\text{second}}^{a} = -4i \sum_{1,\dots,6} \text{Im} \left(G(\vec{e}_{12;34}) G(\vec{e}_{56;34}) \lambda_{1234} \lambda_{5612} \lambda_{3456} \right) (\partial_{1} + \partial_{2} - \partial_{3} - \partial_{4}) J_{1} J_{2} J_{3} J_{4}$$

$$(\partial_{5} + \partial_{6} - \partial_{1} - \partial_{2}) J_{5} J_{6} (\partial_{3} + \partial_{4} - \partial_{5} - \partial_{6}) \langle 0 | \Phi_{0} \rangle .$$
 (4.13)

We also need to include the diagram in Fig. 4(ii). We notice that it is the same as Fig. 4(i) with $1, 2 \leftrightarrow 3, 4$ and the arrows going in the opposite direction. But 1, 2, 3, 4 are just dummy variables, so for that contribution we can simply change variables $1, 2 \leftrightarrow 3, 4$. So we just need to have the arrows go in the opposite direction. But to get (4.13) we already took the arrows to go in both directions, so this gives back the same thing: $\frac{1}{2} (\delta \mathcal{L})_{\text{second}}^a$. In total we thus have $(\delta \mathcal{L})_{\text{second}}^a$.

Next we turn to the diagram in Fig. 2(b). We notice Fig. 2(b) can be obtained from Fig. 2(a) by swapping 2 and 3 and then reversing the arrows on 2, 3, 6. We must also multiply by a combinatorial factor of 4 (because 1 and 2 can be exchanged, and 3 and 4 can be exchanged). Performing this transformation, $2 \to -3$, $3 \to -2$, and $6 \to -6$, on $(\delta \mathcal{L})^a_{\text{second}}$ we get $(\delta \mathcal{L})^b_{\text{second}}$,

$$(\delta \mathcal{L})_{\text{second}}^{b} = 16i \sum_{1,\dots,6} \text{Im} \left(G(\vec{e}_{12;34}) G(\vec{e}_{52;46}) \lambda_{1234} \lambda_{5316} \lambda_{4625} \right) (\partial_{1} + \partial_{2} - \partial_{3} - \partial_{4}) J_{1} J_{2} J_{3} J_{4}$$

$$(\partial_{5} + \partial_{3} - \partial_{1} - \partial_{6}) J_{5} J_{6} \left(\partial_{4} + \partial_{6} - \partial_{2} - \partial_{5} \right) \langle 0 | \Phi_{0} \rangle .$$
 (4.14)

We have now computed all the terms appearing in the master equation (4.4) to third order in the coupling.

Kinetic equation

As we established in the previous section, the kinetic equation is found by integrating the master equation,

$$\frac{\partial n_r}{\partial t} = -4i \int dJ J_r \left((\delta \mathcal{L})_{\text{first}} + (\delta \mathcal{L})_{\text{second}} \right) . \tag{4.15}$$

This is straightforward to carry out; the details are in Appendix A. The result for the first order term is,

$$\int dJ J_r (\delta \mathcal{L})_{\text{first}} = -\sum_{1,\dots,4} G(\vec{e}_{12;34}) \lambda_{1234} \lambda_{3412} (\delta_{1r} + \delta_{2r} - \delta_{3r} - \delta_{4r}) \prod_{i=1}^4 n_i \left(\frac{1}{n_1} + \frac{1}{n_2} - \frac{1}{n_3} - \frac{1}{n_4} \right) , \quad (4.16)$$

and for the second order term is,

$$\int dJ J_r \left(\delta \mathcal{L}\right)_{\text{second}}^a = -4i \sum_{1,\dots,6} \text{Im} \left(G(\vec{e}_{12;34}) G(\vec{e}_{56;34}) \lambda_{1234} \lambda_{5612} \lambda_{3456} \right) \left(\delta_{1r} + \delta_{2r} - \delta_{3r} - \delta_{4r} \right)$$

$$\prod_{i=1}^{6} n_i \left(\frac{1}{n_1} + \frac{1}{n_2} \right) \left(\frac{1}{n_3} + \frac{1}{n_4} - \frac{1}{n_5} - \frac{1}{n_6} \right) , \quad (4.17)$$

and something similar for $\int dJ J_r \left(\delta \mathcal{L}\right)_{\text{second}}^b$ which is obtained by an index interchange. Now, for $G(\vec{e}_{12;34})$ recall its definition (2.18),

$$G(\vec{e}_{12;34}) = \frac{1}{\omega_1 + \omega_2 - \omega_3 - \omega_4 + i\epsilon} = P \frac{1}{\omega_1 + \omega_2 - \omega_3 - \omega_4} - i\pi\delta(\omega_3 + \omega_4 - \omega_1 - \omega_2) , \qquad (4.18)$$

where we split the term into a principal part and a delta function, using (3.10). A potentially more useful form of the kinetic equation is one in which one makes use of (4.18) to write the terms, grouping them by the number of delta functions. Doing this, along with some permutations of the indices to combine and simplify terms (see Appendix A for details) we get for the kinetic equation,

$$\frac{dn_1}{dt} = 16\pi \sum_{2,3,4} \delta(\omega_{12;34}) \lambda_{1234}^2 \prod_{i=1}^4 n_i \left(\frac{1}{n_1} + \frac{1}{n_2} - \frac{1}{n_3} - \frac{1}{n_4} \right) \\
\left[1 + 4 \sum_{5,6} \frac{\lambda_{1256} \lambda_{5634}}{\lambda_{1234}} \frac{n_5 + n_6}{\omega_{12;56}} + 16 \sum_{5,6} \frac{\lambda_{1635} \lambda_{2546}}{\lambda_{1234}} \frac{n_6 - n_5}{\omega_{46;25}} \right] , \quad (4.19)$$

where here we have taken the couplings to be real and we defined $\omega_{ij;kl} \equiv \omega_i + \omega_j - \omega_k - \omega_l$, and when we write $1/\omega$ we really mean the principal part of $1/\omega$. This matches the next-to-leading order kinetic equation we found in [5].

5. Discussion

We have studied classical field theories with weak nonlinearity. A fundamental object is the kinetic equation, describing the time evolution of the occupation number n_k of a mode of momentum k. We presented a scheme to find the kinetic equation perturbatively in the nonlinearity, and found it explicitly to next-to-leading order for a field theory with a quartic interaction. Our method amounts to perturbatively solving the Liouville equation, which we did by analogy with the Lippmann-Schwinger equation used in scattering in quantum mechanics.

The expansion we found is analogous to perturbation theory in many-body quantum mechanics or quantum field theory. In particular, the phase space density is a function of the action and angle variables of every mode p. The effect of a quartic interaction on the Liouville operator is that it adds one unit of angular dependence for some modes p_1 and p_2 , and subtracts one unit for modes

 p_3 and p_4 ; it is as if the interaction creates particles of momenta p_1 and p_2 and destroys particles of momenta p_3 and p_4 . In this way, each term in the perturbative expansion is represented by a Feynman diagram. Our seed for the perturbative expansion was a phase space density that only depends on the action variables. Order by order in perturbation theory, angular dependence for an increasing number of modes is acquired.

More broadly, the standard approach to solving the Liouville equation is through timedependent perturbation theory [34]. In contrast, our method is like time-independent perturbation theory and works by analogy with scattering, which is far simpler. This approach may be useful in other contexts.

Rather than doing a phase space average, an alternative way to derive the kinetic equation is to introduce external Gaussian-random forcing and dissipation, perform an average over the forcing, and at the end take both the forcing and dissipation to zero while maintaining a fixed ratio. This is what we did in earlier work [5]. The method in this paper is technically far simpler, although the method of [5] is perhaps conceptually simpler.

One motivation for studying weakly nonlinear field theories is the existence of turbulent cascades (the Kolmogorov-Zakharov state). The state is found by looking for stationary solutions of the wave kinetic equation. This is always done for the part of the kinetic equation that is leading order in the nonlinearity. It will be interesting to see how accounting for the next-to-leading order term in the kinetic equation modifies the turbulent state.

Although we focused on the occupation numbers of modes, the method presented here allows one to compute much more: it systematically gives the entire state, as a function of all the action and angle variables, perturbatively in the coupling of the nonlinear term. An important problem is to decide which quantity in particular provides an insightful characterization of the turbulent state, and to then compute it.

Acknowledgments

We are indebted to G. Falkovich for making us aware of the Prigogine diagrammatic method and for valuable discussions. We thank M. Srednicki for helpful discussions. The work of VR is supported in part by NSF grant 2209116 and by the ITS through a Simons grant. The work of MS is supported by the Israeli Science Foundation Center of Excellence (grant No. 2289/18). This work was also supported in part by BSF grant 2022113.

A. From the master equation to the kinetic equation

In this appendix we fill in some of details involved in getting the kinetic equation to next-to-leading order. In particular, we start with the kinetic equation (4.15), make use of $(\delta \mathcal{L})_{\text{first}}$ in (4.6) and $(\delta \mathcal{L})_{\text{second}}$ in (4.7), (4.13), and (4.14) to obtain the final form of the kinetic equation in (4.19).

We first look at the integral of $(\delta \mathcal{L})_{\text{first}}$, given in (4.6),

$$\int dJ J_r (\delta \mathcal{L})_{\text{first}} = \int dJ J_r \sum_{1,\dots,4} G(\vec{e}_{12;34}) \lambda_{1234} \lambda_{3412} (\partial_1 + \partial_2 - \partial_3 - \partial_4) J_1 J_2 J_3 J_4 (\partial_3 + \partial_4 - \partial_1 - \partial_2) \rho(\vec{J})$$
(A.1)

where $\langle 0|\Phi_0\rangle \equiv \rho(\vec{J})$. As shown in Sec. 3, we repeatedly integrate by parts to the left and then use the exponential distribution for $\rho(\vec{J})$, as given in (3.16), so that this turns into,

$$\int dJ J_r \left(\delta \mathcal{L}\right)_{\text{first}} = -\sum_{1,\dots,4} G(\vec{e}_{12;34}) \lambda_{1234} \lambda_{3412} \left(\delta_{1r} + \delta_{2r} - \delta_{3r} - \delta_{4r}\right) \prod_{i=1}^4 n_i \left(\frac{1}{n_1} + \frac{1}{n_2} - \frac{1}{n_3} - \frac{1}{n_4}\right) . \quad (A.2)$$

Next we look at the integral of $(\delta \mathcal{L})_{\text{second}}^a$, given in (4.13),

$$\int dJ J_r (\delta \mathcal{L})_{\text{second}}^a = -4i \int d\vec{J} J_r \sum_{1,\dots,6} \text{Im} \left(G(\vec{e}_{12;34}) G(\vec{e}_{56;34}) \lambda_{1234} \lambda_{5612} \lambda_{3456} \right) (\partial_1 + \partial_2 - \partial_3 - \partial_4)$$

$$J_1 J_2 J_3 J_4 (\partial_5 + \partial_6 - \partial_1 - \partial_2) J_5 J_6 (\partial_3 + \partial_4 - \partial_5 - \partial_6) \rho(\vec{J}) . \quad (A.3)$$

We repeatedly integrate by parts to the left so that this turns into,

$$\int dJ J_r (\delta \mathcal{L})_{\text{second}}^a = -4i \int d\vec{J} \sum_{1,\dots,6} \text{Im} \left(G(\vec{e}_{12;34}) G(\vec{e}_{56;34}) \lambda_{1234} \lambda_{5612} \lambda_{3456} \right) (\delta_{1r} + \delta_{2r} - \delta_{3r} - \delta_{4r})$$

$$J_1 J_2 J_3 J_4 J_5 J_6 \left(\frac{1}{J_1} + \frac{1}{J_2} \right) \left(\frac{1}{J_3} + \frac{1}{J_4} - \frac{1}{J_5} - \frac{1}{J_6} \right) \rho(\vec{J}) . \quad (A.4)$$

Evaluating using an exponential distribution for $\rho(\vec{J})$, as given in (3.16), turns all the J_i into n_i ,

$$\int dJ J_r (\delta \mathcal{L})_{\text{second}}^a = -4i \sum_{1,\dots,6} \text{Im} \left(G(\vec{e}_{12;34}) G(\vec{e}_{56;34}) \lambda_{1234} \lambda_{5612} \lambda_{3456} \right) (\delta_{1r} + \delta_{2r} - \delta_{3r} - \delta_{4r})$$

$$\prod_{i=1}^{6} n_i \left(\frac{1}{n_1} + \frac{1}{n_2} \right) \left(\frac{1}{n_3} + \frac{1}{n_4} - \frac{1}{n_5} - \frac{1}{n_6} \right) . \quad (A.5)$$

The other term, $\int dJ J_r (\delta \mathcal{L})_{\text{second}}^b$, is obtained by the transformation discussed previously: $2 \to -3$, $3 \to -2$, and $6 \to -6$, and multiplying by a combinatorial factor of 4, giving,

$$\int dJ J_r (\delta \mathcal{L})_{\text{second}}^b = 16i \sum_{1,\dots,6} \text{Im} \left(G(\vec{e}_{12;34}) G(\vec{e}_{52;46}) \lambda_{1234} \lambda_{5316} \lambda_{4625} \right) \left(\delta_{1r} + \delta_{2r} - \delta_{3r} - \delta_{4r} \right)$$

$$\prod_{i=1}^{6} n_i \left(\frac{1}{n_1} - \frac{1}{n_3} \right) \left(\frac{1}{n_4} + \frac{1}{n_6} - \frac{1}{n_2} - \frac{1}{n_5} \right) . \quad (A.6)$$

Combining all these terms gives, via (4.15), the next-to-leading order kinetic equation. We

would like to make sure that it matches the result we found previously in [5]. Focusing on (A.5), we see that we can replace $(\delta_{1r}+\delta_{2r}-\delta_{3r}-\delta_{4r}) \to 2(\delta_{1r}-\delta_{3r})$ because of the symmetry of everything else under $1 \leftrightarrow 2$ or $3 \leftrightarrow 4$. For the term coming with δ_{3r} we do a change of variables, $(1,2) \leftrightarrow (3,4)$,

$$\int dJ J_r \left(\delta \mathcal{L}\right)_{\text{second}}^a \\
= -8i \sum_{1,\dots,6} \delta_{1r} \prod_{i=1}^6 n_i \left\{ \text{Im} \left(G(\vec{e}_{12;34}) G(\vec{e}_{56;34}) \lambda_{1234} \lambda_{5612} \lambda_{3456} \right) \left(\frac{1}{n_1} + \frac{1}{n_2} \right) \left(\frac{1}{n_3} + \frac{1}{n_4} - \frac{1}{n_5} - \frac{1}{n_6} \right) \\
+ \text{Im} \left(G(\vec{e}_{12;34}) G(\vec{e}_{12;56}) \lambda_{1234} \lambda_{5612} \lambda_{3456} \right) \left(\frac{1}{n_3} + \frac{1}{n_4} \right) \left(\frac{1}{n_1} + \frac{1}{n_2} - \frac{1}{n_5} - \frac{1}{n_6} \right) \right\} , \quad (A.7)$$

where we used that $G(-\vec{e}) = -G^*(\vec{e})$. Next, we insert the following relation,

$$G(\vec{e}_{12;34})(G(\vec{e}_{56;34}) + G(\vec{e}_{12;56})) = G(\vec{e}_{56;34})G(\vec{e}_{12;56})$$
(A.8)

into (A.7). Writing the $G(\vec{e}_{ij;kl})$ in (A.7) explicitly, we have,

$$\int dJ J_r \left(\delta \mathcal{L}\right)_{\text{second}}^a = -8i \sum_{1,\dots,6} \delta_{1r} \prod_{i=1}^6 n_i \text{Im} \left[\lambda_{1234} \lambda_{5612} \lambda_{3456} \left\{ \left(\frac{1}{n_1} + \frac{1}{n_2} \right) \left(\frac{1}{n_3} + \frac{1}{n_4} \right) \frac{1}{\omega_{34;56} - i\epsilon} \frac{1}{\omega_{56;12} - i\epsilon} \right. \\ \left. - \left(\frac{1}{n_5} + \frac{1}{n_6} \right) \left(\frac{1}{n_1} + \frac{1}{n_2} \right) \frac{1}{\omega_{34;12} - i\epsilon} \frac{1}{\omega_{34;56} - i\epsilon} - \left(\frac{1}{n_5} + \frac{1}{n_6} \right) \left(\frac{1}{n_3} + \frac{1}{n_4} \right) \frac{1}{\omega_{34;12} - i\epsilon} \frac{1}{\omega_{56;12} - i\epsilon} \right\} \right], \quad (A.9)$$

where we defined the shorthand $\omega_{ij;kl} \equiv \omega_i + \omega_j - \omega_k - \omega_l$. This precisely reproduces what we had in the kinetic equation in [5]. ⁵ There we proceeded by using (4.18) on all the $G(\vec{e}_{ij;kl})$ terms, grouping terms together based on the number of delta functions of the ω_i , and performing the change of variables (3, 4) \leftrightarrow (5, 6) on some of the terms in order to show that they are either zero due to antisymmetry, or to combine them with other terms. The end result is the kinetic equation (4.19).

B. Frequency renormalization

In the Hamiltonian (3.3),

$$H = \sum_{i} \omega_{i} J_{i} + \sum_{i,j,k,l} \lambda_{ijkl} \sqrt{J_{i} J_{j} J_{k} J_{l}} \exp\left(i(\alpha_{i} + \alpha_{j} - \alpha_{k} - \alpha_{l})\right) , \qquad (B.1)$$

⁵In particular, in [5] insert (4.25) into (4.32) and use that the $Im(x^*) = -Im(x)$.

it is useful to split off the term in the interaction in which some pair of indices are equal,

$$H = \sum_{i} \omega_{i} J_{i} + \sum_{(i,j)\neq(k,l)} \lambda_{ijkl} \sqrt{J_{i} J_{j} J_{k} J_{l}} \exp\left(i(\alpha_{i} + \alpha_{j} - \alpha_{k} - \alpha_{l})\right) + H_{ren} , \qquad (B.2)$$

$$H_{ren} = 2 \sum_{i\neq k} \lambda_{ikik} J_{i} J_{k} + \sum_{i} \lambda_{iiii} J_{i}^{2} .$$

The corresponding matrix element of $(\delta L)_{ren}$ for H_{ren} is, from (2.14),

$$\langle \vec{n} | (\delta L)_{ren} | \vec{n}' \rangle = 4 \, \delta_{\vec{n}, \vec{n}'} \sum_{i \neq k} \lambda_{ikik} J_k n_i + 2 \, \delta_{\vec{n}, \vec{n}'} \sum_i \lambda_{iiii} J_i n_i . \tag{B.3}$$

Switching to notation in which we call mode i to be 1 and mode k to be 5, we have that the contribution at order λ^3 which involves one insertion of $(\delta L)_{ren}$ is,

$$(\delta \mathcal{L})_{\text{ren}} = -4 \sum_{1,\dots,5} G(\vec{e}_{12;34})^2 |\lambda_{1234}|^2 (\partial_1 + \partial_2 - \partial_3 - \partial_4) J_1 J_2 J_3 J_4$$

$$J_5 (\lambda_{1515} + \lambda_{2525} - \lambda_{3535} - \lambda_{4545}) (\partial_3 + \partial_4 - \partial_1 - \partial_2) \langle 0 | \Phi_0 \rangle , \quad (B.4)$$

where we have accounted for the fact that $(\delta L)_{ren}$ can be inserted along any of the lines 1, 2, 3, 4. The contribution to the kinetic equation is,

$$\int dJ J_r (\delta \mathcal{L})_{\text{ren}} = i \sum_{1,\dots,4} \text{Im}(G(\vec{e}_{12;34})^2) |\lambda_{1234}|^2 (\delta_{1r} + \delta_{2r} - \delta_{3r} - \delta_{4r}) \prod_{i=1}^4 n_i \left(\frac{1}{n_1} + \frac{1}{n_2} - \frac{1}{n_3} - \frac{1}{n_4}\right)$$

$$(\delta \omega_1 + \delta \omega_2 - \delta \omega_3 - \delta \omega_4) \quad (B.5)$$

where we used that the term with $\text{Re}(G(\vec{e}_{12;34})^2)$ gives zero due to the symmetry under $1, 2 \leftrightarrow 3, 4$, and where

$$\delta\omega_a = 4\sum_5 \lambda_{a5a5} n_5 \ . \tag{B.6}$$

We note that,

$$G(\vec{e}_{12;34}) = \frac{1}{\omega_{12;34} + i\epsilon} , \quad \Rightarrow \operatorname{Im}(G(\vec{e}_{12;34})^2) = -\frac{\partial}{\partial \omega_{12;34}} \operatorname{Im} \frac{1}{\omega_{12;34} + i\epsilon} = \pi \delta'(\omega_{12;34})$$
(B.7)

The contribution to the kinetic equation of (B.5) is therefore,

$$\left(\frac{dn_r}{\partial t}\right)_{\text{ren}} = 4\sum_{1,\dots,4} \pi \delta'(\omega_{12;34}) |\lambda_{1234}|^2 (\delta_{1r} + \delta_{2r} - \delta_{3r} - \delta_{4r}) \prod_{i=1}^4 n_i \left(\frac{1}{n_1} + \frac{1}{n_2} - \frac{1}{n_3} - \frac{1}{n_4}\right) (\delta\omega_1 + \delta\omega_2 - \delta\omega_3 - \delta\omega_4)$$
(B.8)

The reason we interpreted this as a frequency renormalization is because if one takes the leading order kinetic equation,

$$\frac{\partial n_r}{\partial t} = 4 \sum_{1,\dots,4} (\delta_{1r} + \delta_{2r} - \delta_{3r} - \delta_{4r}) \pi \delta(\omega_{12;34}) |\lambda_{1234}|^2 \prod_{i=1}^4 n_i \left(\frac{1}{n_1} + \frac{1}{n_2} - \frac{1}{n_3} - \frac{1}{n_4} \right)$$
(B.9)

and sets $\omega_a \to \omega_a + \delta \omega_a$, the change is precisely (B.8).

References

- [1] V. Zakharov, "Weak turbulence in media with a decay spectrum," J Appl Mech Tech Phys 6 (1965) 22–24.
- [2] V. E. Zakharov, V. S. L'vov, and G. Falkovich, Kolmogorov Spectra of Turbulence I: Wave Turbulence. Springer-Verlag, 1992.
- [3] S. Nazarenko, Wave Turbulence. Springer-Verlag Berlin Heidelberg, 2011.
- [4] M. Shavit and G. Falkovich, "Singular Measures and Information Capacity of Turbulent Cascades," *Phys. Rev. Lett.* **125** (2020) 104501, arXiv:1911.12670 [physics.flu-dyn].
- [5] V. Rosenhaus and M. Smolkin, "Feynman rules for forced wave turbulence," arXiv:2203.08168 [cond-mat.stat-mech].
- [6] C. Rodda, C. Savaro, G. Davis, J. Reneuve, P. Augier, J. Sommeria, T. Valran, S. Viboud, and N. Mordant, "Experimental observations of internal wave turbulence transition in a stratified fluid," *Physical Review Fluids* 7 (2022) no. 9, 094802, arXiv:2209.03616 [physics.flu-dyn].
- [7] E. Falcon and N. Mordant, "Experiments in Surface Gravity Capillary Wave Turbulence," *Annual Review of Fluid Mechanics* **54** (Jan, 2022) 1–25, arXiv:2107.04015 [physics.flu-dyn].
- [8] E. Kochurin, G. Ricard, N. Zubarev, and E. Falcon, "Three-dimensional direct numerical simulation of free-surface magnetohydrodynamic wave turbulence," *Physical Review E* **105** (2022) no. 6, L063101, arXiv:2205.11516 [physics.plasm-ph].
- [9] G. Düring, C. Josserand, and S. Rica, "Wave turbulence theory of elastic plates," *Physica D: Nonlinear Phenomena* **347** (2017) 42–73.
- [10] A. I. Dyachenko, A. O. Korotkevich, and V. E. Zakharov, "Weak Turbulent Kolmogorov Spectrum for Surface Gravity Waves," *Physical Review Letters* 92 (Apr. 2004).

- [11] J. Orosco, W. Connacher, and J. Friend, "Identification of weakly-to strongly-turbulent three-wave processes in a micro-scale system," arXiv:2205.01803 [physics.flu-dyn].
- [12] J. Guioth, F. Bouchet, and G. L. Eyink, "Path large deviations for the kinetic theory of weak turbulence," arXiv:2203.11737 [cond-mat.stat-mech].
- [13] M. Gałka, P. Christodoulou, M. Gazo, A. Karailiev, N. Dogra, J. Schmitt, and Z. Hadzibabic, "Emergence of isotropy and dynamic self-similarity in the birth of two-dimensional wave turbulence," arXiv:2203.09514 [cond-mat.quant-gas].
- [14] N. Vladimirova, M. Shavit, and G. Falkovich, "Fibonacci Turbulence," *Physical Review X* 11 (Jun, 2021), arXiv:2101.10418 [nlin.CD].
- [15] G. Falkovich, Y. Kadish, and N. Vladimirova, "Multi-mode correlations and the entropy of turbulence," arXiv:2209.05816 [nlin.CD].
- [16] J. W. Banks, T. Buckmaster, A. O. Korotkevich, G. Kovacic, and J. Shatah, "Direct verification of the kinetic description of wave turbulence," arXiv:2109.02477 [physics.flu-dyn].
- [17] S. Schlichting and D. Teaney, "The First fm/c of Heavy-Ion Collisions," Ann. Rev. Nucl. Part. Sci. 69 (2019) 447–476, arXiv:1908.02113 [nucl-th].
- [18] A. Chatrchyan, K. T. Geier, M. K. Oberthaler, J. Berges, and P. Hauke, "Analog cosmological reheating in an ultracold Bose gas," Phys. Rev. A 104 (2021) no. 2, 023302, arXiv:2008.02290 [cond-mat.quant-gas].
- [19] Y. Zhu, B. Semisalov, G. Krstulovic, and S. Nazarenko, "Direct and inverse cascades in turbulent Bose-Einstein condensate," arXiv:2208.09279 [cond-mat.quant-gas].
- [20] G. Dematteis and Y. V. Lvov, "The Structure of Energy Fluxes in Wave Turbulence," arXiv:2205.12899 [physics.flu-dyn].
- [21] V. David and S. Galtier, "Wave turbulence in inertial electron magnetohydrodynamics," *Journal of Plasma Physics* 88 (2022) no. 5, 905880509, arXiv:2209.08577 [physics.plasm-ph].
- [22] Y. Sano, N. Navon, and M. Tsubota, "Emergent isotropy of a wave-turbulent cascade in the Gross-Pitaevskii model," arXiv:2209.08973 [cond-mat.quant-gas].
- [23] N. Berti, K. Baudin, A. Fusaro, G. Millot, A. Picozzi, and J. Garnier, "Interplay of thermalization and strong disorder: Wave turbulence theory, numerical simulations, and experiments in multimode optical fibers," *Phys. Rev. Lett.* 129 (2022) no. 6, 063901, arXiv:2207.08680 [physics.optics].

- [24] K. Kawasaki and I. Oppenheim, "Logarithmic term in the density expansion of transport coefficients," *Phys. Rev.* **139** (Sep. 1965) A1763–A1768.
- [25] J. R. Dorfman and E. G. D. Cohen, "Velocity correlation functions in two and three dimensions," *Phys. Rev. Lett.* **25** (Nov, 1970) 1257–1260.
- [26] J. R. Dorfman, T. R. Kirkpatrick, and J. V. Sengers, "Why Non-equilibrium is Different," arXiv:1512.02679 [cond-mat.stat-mech].
- [27] J. R. Dorfman, H. van Beijeren, and T. R. Kirkpatrick, Contemporary Kinetic Theory of Matter. Cambridge University Press, 2021.
- [28] M. S. Green, "Boltzmann equation from the statistical mechanical point of view," *The Journal of Chemical Physics* **25** (1956) no. 5, 836–855.
- [29] E. Cohen, "On the generalization of the boltzmann equation to general order in the density," *Physica* **28** (1962) no. 10, 1025–1044.
- [30] R. Zwanzig, "Method for finding the density expansion of transport coefficients of gases," *Physical Review* **129** (1963) no. 1, 486.
- [31] N. Bogoliubov, Problems of Dynamic Theory in Statistical Physics, 1946.
- [32] J. Brocas, "On the comparison between two generalized boltzmann equations," Advances in Chemical Physics (1967) 317–381.
- [33] R. Balescu, "Irreversible processes in ionized gases," The Physics of Fluids 3 (1960) no. 1, 52–63.
- [34] I. Prigogine, *Non-equilibrium Statistical Mechanics*. Interscience Publishers, 1962. Reprint: Dover Publications, 2017.
- [35] R. L. Liboff, Kinetic Theory: Classical, Quantum, and Relativistic Descriptions. Springer, 2003.
- [36] R. Peierls, "On the kinetic theory of heat conduction in crystals," Ann. Phys. **395** (1929) no. 8, 1055.
- [37] J. J. Sakurai, Modern Quantum Mechanics. Addison-Wesley, 1994.
- [38] A. Polyakov unpublished
- [39] V. Gurarie, Statistics without thermodynamic equilibrium. PhD thesis, Princeton University, 1996. https://scholar.google.com/citations?view_op=view_citation&hl=en&

- $user=1X40ds8AAAAJ\&cstart=100\&pagesize=100\&sortby=pubdate\&citation_for_view=1X40ds8AAAAJ:f2IySw72cVMC.$
- [40] V. Gurarie, "Probability density, diagrammatic technique, and epsilon expansion in the theory of wave turbulence," *Nucl. Phys. B* **441** (1995) 569–594, arXiv:hep-th/9405077.