

Testing axion couplings to leptons in Z decays at future e^+e^- colliders

Lorenzo Calibbi, Zijie Huang, Shaoyang Qin,
Yiming Yang,¹ and Xiaoyue Yin

School of Physics, Nankai University, Tianjin 300071, China

Abstract

We study the possibility of probing the existence of a light, invisible, axion-like particle (ALP) a in leptonic decays of the Z boson at the proposed high-energy e^+e^- colliders, CEPC and FCC-ee. Both projects plan to run at the Z pole, collecting $10^{12} - 10^{13}$ visible Z decays. We show that, searching for the emission of an invisible ALP from leptons in leptonic Z decays, this enormous statistics could allow to constrain the ALP couplings to leptons at an unprecedented level for laboratory experiments. In particular, within a Monte Carlo simulation framework, we estimate that CEPC/FCC-ee can be sensitive to the coupling of an invisible ALP to muons up to $f_a/C_{\mu\mu}^A \approx 1$ TeV — where f_a is the ALP decay constant — corresponding to $\text{BR}(Z \rightarrow \mu^+\mu^- a) \approx 2 \times 10^{-11}$.

arXiv:2212.02818v1 [hep-ph] 6 Dec 2022

¹Corresponding author, *e-mail*: yang_yiming@mail.nankai.edu.cn

Contents

1	Introduction	2
2	ALP couplings to leptons and decays	3
3	ALP emission in leptonic Z decay	5
4	Collider simulation and discussion	6
5	Summary and conclusions	10

1 Introduction

Despite the undeniable success of the Standard Model (SM) and the absence at present of conclusive evidence for new physics (NP) at the Large Hadron Collider (LHC) and low-energy experiments, a number of experimental observations and open problems call for its extension — among others: dark matter, neutrino masses, matter-antimatter asymmetry, the flavour puzzle.

Interestingly, a wide class of models that have been proposed to address these questions predicts the existence of so-called axion-like particles (ALPs), light weakly-coupled pseudoscalar fields that are the pseudo-Nambu-Goldstone bosons (PNGBs) of spontaneously broken global symmetries. The prototypical example is the axion, which arises from the breaking of the Peccei-Quinn $U(1)$ symmetry that was introduced to solve the strong CP problem [1–3]. Similarly, other well-motivated ALPs are light remnants of high-energy global symmetries associated with further outstanding problems of the SM, such as the lepton number within neutrino mass models [4–6], or flavour symmetries [7–9], e.g. in theories addressing the observed hierarchical pattern of fermion masses and mixing and the strong CP problem at the same time [10, 11]. In these latter examples, ALP couplings to charged leptons, which are the focus of the present work, are unavoidable, although model-dependent — for more instances of ‘leptonic’ axion and ALP models, see [12]. As all ALP couplings, these are inversely proportional to the scale f_a at which the symmetry is broken and thus suppressed if this occurs at energies much larger than the lepton mass scale.

ALP interactions with charged leptons can be tested at laboratory experiments either directly or indirectly via the coupling to photons that they unavoidably induce at the loop level. Such laboratory constraints come from measurements of electric [13] and magnetic dipole moments — in certain regions of the parameter space, ALP contributions can be large enough as to account for the observed muon $g - 2$ discrepancy, see e.g. [14–18] — from beam dump experiments [19, 20], and from low-energy e^+e^- colliders such as B factories [21].¹ However, most of these probes tend to lose sensitivity if the ALP is light and weakly-coupled enough to be long-lived, that is, for large values of f_a .²

The latter situation is best constrained by astrophysical observations. Emission of light particles would, in fact, contribute to the energy loss of stellar systems altering their evolution [25]. Consequently, very stringent bounds to ALP-electron couplings can be derived from observations of populations of stars such as white dwarfs (WD) and red giants (RG) [26, 27], as long as the ALPs are light enough to be produced inside the star ($m_a \lesssim \mathcal{O}(10)$ keV) and weakly-coupled enough to avoid to be reabsorbed. Hence, these limits do not apply to heavier ALPs and/or ALPs characterised by medium-to-low values of the breaking scale f_a , while

¹In presence of flavour-violating couplings, severe constraints are also provided by searches for lepton flavour violation [12, 22, 23].

²An exception is provided by the search for light dark matter particles in e^+e^- annihilations at the NA64 experiment [24], which is also sensitive to an invisible ALP coupling to electrons.

still long-lived for what concerns laboratory experiments. Similarly, the observation of the supernova explosion SN1987A can be used to set limits on ALP couplings to electrons [28] that are somewhat weaker than those from WD and RG but extend to $m_a \lesssim \mathcal{O}(100)$ MeV given the much hotter environment. Furthermore, the SN1987A data were recently employed to set bounds to the coupling to muons as well [29, 30], the latter ones being sensitive also to the low f_a regime. These constraints are however affected by our uncertainty about the supernova explosion mechanism and could even disappear in certain scenarios thereof [31].

The above discussion shows that it would be desirable to have independent direct laboratory probes of the ALP couplings to leptons, sensitive to the case of a long-lived (and thus invisible) ALP. Recent proposals along this direction have focused on the emission of ALPs in meson decays, showing how charged pion [32] and kaon [33] decays can be sensitive to, respectively, the coupling to electrons and muons of a light invisible particle.

The aim of the present note is to show that the proposed high-energy e^+e^- accelerators, the Circular Electron-Positron Collider (CEPC) [34, 35] and the e^+e^- stage of the Future Circular Collider (FCC-ee) [36, 37], can provide a unique opportunity to discover or constrain invisible ALPs through their couplings to charged leptons. Both proposals plan to run for several years at a centre of mass energy around the Z pole, $\sqrt{s} \simeq 91$ GeV, hence operating as “Tera Z ” factories, that is, producing more than 10^{12} visible Z boson decays at two interaction points. As the Z boson has a probability of about 10% of decaying into two charged leptons, the CEPC and FCC-ee would produce more than 10^{11} boosted lepton pairs, $\ell^+\ell^-$ ($\ell = e, \mu, \tau$), thus offering the opportunity to test the emission of an ALP a from the final state leptons in the process $Z \rightarrow \ell^+\ell^-a$. In this paper, we focus on the parameter space where a would decay outside the detector hence appearing as missing energy and we simulate the signal and background for $e^+e^- \rightarrow Z^{(*)} \rightarrow \ell^+\ell^-a$, in order to estimate the sensitivity of CEPC/FCC-ee to the ALP couplings to leptons.

A number of recent works studied possible searches for ALPs at future e^+e^- colliders [18, 38–41], focusing on other ALP-SM interactions and production modes and/or on promptly-decaying ALPs—which is a consequence of the relatively strong couplings necessary e.g. for an explanation of the muon $g - 2$ anomaly. As mentioned above, here we are instead concerned with lighter and more weakly-coupled ALPs that would escape the detector unseen.

The rest of the paper is organised as follows. In Section 2, we set the notation, review the effective Lagrangian describing ALP-lepton interactions, and discuss the ALP decay modes and lifetime. In Section 3, we discuss ALP emission in leptonic Z decays and present an analytical estimate of the number of ALP events to be expected at future e^+e^- colliders. In Section 4, we show the result of our simulation and its impact on the ALP parameter space. Finally, in Section 5, we summarise and conclude.

2 ALP couplings to leptons and decays

In the following, we review the simplified-model approach to lepton-ALP couplings that we employ in our analysis. The interactions we are interested in can be written as a dimension-five operator involving a derivative ALP coupling with a leptonic axial current:

$$\mathcal{L}_{\text{eff}} \supset \sum_{\ell=e,\mu,\tau} \frac{C_{\ell\ell}^A}{2f_a} \partial_\mu a (\bar{\ell}\gamma^\mu\gamma_5\ell), \quad (1)$$

where a represents the ALP field, ℓ denotes charged leptons (that is, $\ell = e, \mu, \tau$), and $C_{\ell\ell}^A$ are model-dependent dimensionless coefficients that we will be treating as free parameters throughout the paper. The ALP decay constant f_a is a scale related to the spontaneous breaking of the global $U(1)$ symmetry the ALP is associated to. Notice that in this work we restrict the discussion to flavour-conserving couplings to leptons. For the very tight constraints that arise in

presence of flavour-violating ALP-lepton interactions, in particular from lepton-flavour-violating decays $\ell \rightarrow \ell' a$ into an invisible ALP, see [12].

Upon integrating by parts and inserting the equations of motion of the lepton fields, the above Lagrangian is equivalent — up to a shift of the anomalous coupling to photons — to the following dimension-four interaction terms involving leptonic pseudoscalar currents, see e.g. [42]:

$$\mathcal{L}_{\text{eff}} \supset -i \sum_{\ell} g_{\ell} a (\bar{\ell} \gamma_5 \ell), \quad g_{\ell} \equiv C_{\ell\ell}^A \frac{m_{\ell}}{f_a}, \quad (2)$$

which shows that the ALP-lepton couplings are proportional to the lepton mass m_{ℓ} and thus larger for heavier generations.

The above lepton-ALP interactions induce, when kinematically allowed, ALP decays into lepton pairs, whose width reads:

$$\Gamma(a \rightarrow \ell^+ \ell^-) = \frac{m_a}{8\pi} m_{\ell}^2 \left(\frac{C_{\ell\ell}^A}{f_a} \right)^2 \sqrt{1 - \frac{4m_{\ell}^2}{m_a^2}}. \quad (3)$$

The same interactions unavoidably contribute to the effective ALP coupling into photons E_{eff} , which we define as:

$$\mathcal{L}_{\text{eff}} \supset E_{\text{eff}} \frac{\alpha_{\text{em}}}{4\pi} \frac{a}{f_a} F \tilde{F}. \quad (4)$$

This quantity depends on a model-dependent UV contribution and, through lepton loops, on the ALP couplings to leptons in Eq. (1):

$$E_{\text{eff}} = E_{\text{UV}} + \sum_{\ell} C_{\ell\ell}^A B(z), \quad (5)$$

where $z = 4m_{\ell}^2/m_a^2$ and the loop function can be written as [43]:

$$B(z) = 1 - z f^2(z), \quad f(z) = \begin{cases} \arcsin \frac{1}{\sqrt{z}}, & z \geq 1, \\ \frac{\pi}{2} + \frac{i}{2} \ln \frac{1+\sqrt{1-z}}{1-\sqrt{1-z}}, & z < 1. \end{cases} \quad (6)$$

Notice that the latter contribution correctly decouples in the limit $m_a \ll m_{\ell}$, as $B(z) \rightarrow 0$ for $z \rightarrow \infty$. In terms of the above-defined effective coupling, the decay rate into photons reads

$$\Gamma(a \rightarrow \gamma\gamma) = \frac{\alpha_{\text{em}}^2 E_{\text{eff}}^2 m_a^3}{64\pi^3 f_a^2}. \quad (7)$$

Finally, ALP decays into neutrinos $a \rightarrow \nu_i \bar{\nu}_i$ are typically suppressed by the small neutrino masses, hence we set $\Gamma(a \rightarrow \nu_i \bar{\nu}_i) = 0$ throughout the paper.³ Thus we use the following expression for the total decay width of the ALP:

$$\Gamma_a = \Gamma(a \rightarrow \gamma\gamma) + \sum_{\ell} \theta(m_a - 2m_{\ell}) \times \Gamma(a \rightarrow \ell^+ \ell^-). \quad (8)$$

The above expression enables us to assess under which conditions the emitted ALPs preferably decay outside of the detector, thus giving rise to the missing energy signature that we are going to study in the following sections. In Figure 1, we highlight in grey the region of the parameter space where the proper decay length $c\tau_a \equiv c/\Gamma_a$ of a ‘purely muonic’ ALP ($C_{\mu\mu}^A = 1$, $C_{ee}^A = 0$, $E_{\text{UV}} = 0$)⁴ is of the order of the size of typical collider experiment detectors or smaller,

³Furthermore, ALPs decaying into neutrinos inside a collider detector would still appear as missing energy. Hence, these decay modes would not change the following discussion.

⁴Notice that the coupling to tau leptons $C_{\tau\tau}^A$ has no effect on the ALP lifetime as long as $m_a \ll m_{\tau}$.




Figure 1: ALP decay length in the (m_a, f_a) plane: in the gray region an ALP coupling only to muons mostly decays inside the detector ($c\tau_a < 10$ m). The coloured lines show how the boundary of the region is modified for different choices of the couplings to leptons and photons.

$c\tau_a < 10$ m. Outside this region, the ALP is typically long-lived. The figure also shows how the boundary of the region varies for different choices of the coupling to electrons C_{ee}^A and the UV-dependent coupling to photons E_{UV} . As one can see, the ALP is mostly long-lived as long as its mass is below the kinematic threshold of the decay into a muon pair, $m_a < 2m_\mu \simeq 210$ MeV. This is also true even in presence of ALP couplings to electrons and photons, at least if they feature a certain hierarchical structure such that $C_{ee}^A \ll C_{\mu\mu}^A$ — otherwise the ALP would mostly decay as $a \rightarrow e^+e^-$ for low values of the ALP decay constant, $f_a \lesssim 1$ TeV, and $m_a > 2m_e$. For reasons that we will be discussed in the following sections, we will mostly focus on ALPs coupling to muons and set $C_{ee}^A = 0$. For definiteness, we will also assume $E_{UV} = 0$, which corresponds to the case of a PNGB of a $U(1)$ symmetry free of electromagnetic anomalies, an example being provided by the majoron, the PNGB associated to the spontaneous breaking of the lepton number [4–6]. Furthermore, Figure 1 shows that $E_{UV} \neq 0$ would not affect much the region we are interested in where a is long-lived (unless $|E_{UV}| \gg 1$). In Section 4, we will comment about the impact of relaxing the above assumptions ($C_{ee}^A = 0, E_{UV} = 0$) on the prospected CEPC/FCC-ee constraints on the ALP parameter space.

3 ALP emission in leptonic Z decay

As we aim at estimating the sensitivity of CEPC/FCC-ee on ALPs emitted by one of the final state leptons in leptonic Z decays, we present here the analytical expression for the decay width of $Z \rightarrow \ell^+\ell^-a$. Following from the diagrams in Figure 2 and the interaction in the Lagrangian given in Eq. (1), the partial Z decay width we are interested in reads

$$\Gamma(Z \rightarrow \ell^+\ell^-a) = \frac{G_F m_\ell^2}{192\sqrt{2}\pi^3 m_Z^3} \left(\frac{C_{\ell\ell}^A}{f_a}\right)^2 \int dm_{13}^2 dm_{23}^2 \left[\frac{m_Z^2(m_{23}^4 + m_{13}^4)}{m_{23}^2 m_{13}^2} (2c_{2w}^2 - 2c_{2w} + 1) - (1 - 2c_{2w})^2 (m_{13}^2 + m_{23}^2) \right], \quad (9)$$

where m_Z is the Z boson mass, G_F is the Fermi constant, $c_{2w} \equiv \cos 2\theta_w$ with θ_w denoting the weak mixing angle, and $m_{13}^2 \equiv (p_1 + p)^2$, $m_{23}^2 \equiv (p_2 + p)^2$, with the 4-momenta defined as in




Figure 2: Feynman diagrams of leptonic Z decay with ALP emission.

Figure 2. The integration is performed over the usual phase space available to a three-body decay, see e.g. [44].

Although the process we are going to simulate is $e^+e^- \rightarrow Z^* \rightarrow \ell^+\ell^-a$, the above analytical expression of the width of on-shell Z decays already allows us to obtain a first approximate assessment of the capability of future colliders. Taking into account the total width of the Z boson, $\Gamma_Z \simeq 2.5$ GeV [44], Eq. (9) numerically gives for $m_a \ll m_\ell$:

$$\text{BR}(Z \rightarrow e^+e^-a) \approx 1.2 \times 10^{-13} \left(\frac{100 \text{ GeV}}{f_a/C_{ee}^A} \right)^2, \quad (10)$$

$$\text{BR}(Z \rightarrow \mu^+\mu^-a) \approx 2.5 \times 10^{-9} \left(\frac{100 \text{ GeV}}{f_a/C_{\mu\mu}^A} \right)^2, \quad (11)$$

$$\text{BR}(Z \rightarrow \tau^+\tau^-a) \approx 3.3 \times 10^{-7} \left(\frac{100 \text{ GeV}}{f_a/C_{\tau\tau}^A} \right)^2. \quad (12)$$

The above values correspond to the largest possible branching ratios that we can arguably expect within viable UV completions of the effective Lagrangian in Eq. (1), as it may be challenging to formulate such theories for $f_a \lesssim 100$ GeV.

According to the CEPC conceptual design report (CDR) [34,35], in the course of the Z -pole run, two detectors should collect data amounting to an integrated luminosity of $\mathcal{L} = 50 \text{ ab}^{-1}$, equivalent to approximately 2×10^{12} visible Z decays. More recent assessments of the expected accelerator performance have updated this figure to $\mathcal{L} = 96 \text{ ab}^{-1}$, that is, more than 4×10^{12} Z decays [45]. Similarly, the FCC-ee CDR [36,37] estimates that the collider should deliver in total $\mathcal{L} = 150 \text{ ab}^{-1}$ ($\approx 7 \times 10^{12}$ Z decays) at two interaction points. Comparing these planned luminosities with the estimates for the branching ratios in Eqs. (10)-(12), one can see that no signal events are to be expected from $Z \rightarrow e^+e^-a$, as a consequence of the process being suppressed by a factor $\sim (m_e/f_a)^2$, see Eq. (9). On the other hand, we can foresee that CEPC/FCC-ee would produce up to $\mathcal{O}(10^4)$ $Z \rightarrow \mu^+\mu^-a$ events and $\mathcal{O}(10^6)$ $Z \rightarrow \tau^+\tau^-a$ events. For this reason, in the following, we will mostly focus on $Z \rightarrow \mu^+\mu^-a$ and comment about $Z \rightarrow \tau^+\tau^-a$ that, despite the larger probability, is much more difficult to disentangle from the background as a consequence of the missing energy due to the neutrinos from tau decays.

4 Collider simulation and discussion

In order to simulate the process we are interested in— $e^+e^- \rightarrow \ell^+\ell^-a$ with a centre of mass energy $\sqrt{s} \simeq 91$ GeV—we implement the Lagrangian in Eq. (1) in FEYNRULES [46] and we employ the resulting model files [47] within the MADGRAPH framework [48,49] in order to

	$\sigma(e^+e^- \rightarrow \ell^+\ell^-a)$	$\sigma(e^+e^- \rightarrow \ell^+\ell^-\nu\bar{\nu})$	$\sigma(e^+e^- \rightarrow \ell^+\ell^-(\gamma))$
$\ell = e$	7.1×10^{-9} pb	1.9×10^{-3} pb	4490 pb
$\ell = \mu$	7.6×10^{-5} pb	2.9×10^{-4} pb	2024 pb
$\ell = \tau$	1.1×10^{-2} pb	2.8×10^{-4} pb	2020 pb

Table 1: LO cross sections of signal (for $f_a/C_{\ell\ell}^A = 100$ GeV, $m_a = 10^{-6}$ GeV) and background processes for e^+e^- collisions with $\sqrt{s} = 91.2$ GeV, cf. the main text for details.

compute cross sections and generate events for the signal and the relevant SM backgrounds. As a cross check of the model's implementation, we calculated $\text{BR}(Z \rightarrow \ell^+\ell^+a)$ with MADGRAPH finding a very good agreement with the results shown in Eqs. (10)-(12). The simulation we present below makes also use of the parton shower generator PYTHIA [50] and the fast detector simulator DELPHES [51] adopting the parameters of a typical CEPC detector [52]. Finally, we analyse the resulting events by means of ROOT [53] and MADANALYSIS [54].

Having set up the above computation framework, we proceed as follows: (i) We perform a full fast simulation (from event generation to detector response) assuming a given value of $f_a/C_{\ell\ell}^A$ and different values of m_a (although the signal acceptance will turn out to be insensitive to the latter parameter); (ii) Then, we analyse the results of the simulation, in particular the event distributions, to find the suitable kinematical variables to cut on, in order to reduce the background and retain the signal; (iii) Finally, we use the estimated efficiencies of our kinematical cuts to calculate the minimum signal cross section corresponding to a detectable number of events and obtain from it to expected limits on the model's parameters.

The ALP appears as missing energy inside the detector, since we focus on the parameter space where it is characterised by a long lifetime, as shown in Section 2. Therefore, the main physical background for our processes is given by $e^+e^- \rightarrow \ell^+\ell^-\nu\bar{\nu}$, typically stemming from the rare four-body decay $Z \rightarrow \ell^+\ell^-\nu\bar{\nu}$. The cross sections of signal (for representative choices of the ALP parameters) and background processes, as calculated at leading order (LO) by MADGRAPH, are displayed in Table 1. In addition, there is a sizeable probability of radiation from the initial or final state leptons of one or more photons, $e^+e^- \rightarrow \ell^+\ell^-\gamma$. These processes can fake our signal if the photons, especially the soft ones, get undetected. In order to account for such an instrumental background, we generate large samples of $e^+e^- \rightarrow \ell^+\ell^-$ events and let PYTHIA and DELPHES simulate, respectively, the electromagnetic showers and the detector response. As a reference, the LO cross sections for $e^+e^- \rightarrow \ell^+\ell^-$ are also shown in Table 1.

The cross section of $e^+e^- \rightarrow e^+e^-a$ displayed in Table 1 confirms that, as argued by the analytical estimates in Section 3, constraining ALP-electron interactions is beyond the sensitivity of Tera Z factories. Hence we only need to simulate the processes $e^+e^- \rightarrow \mu^+\mu^-a$ and $e^+e^- \rightarrow \tau^+\tau^-a$. Table 1 shows that the cross section of the latter process is relatively sizeable, in fact larger than the SM process $\sigma(e^+e^- \rightarrow \tau^+\tau^-\nu\bar{\nu})$ as long as $f_a \lesssim 600$ GeV, as follows from the scaling $\sigma(e^+e^- \rightarrow \tau^+\tau^-a) \sim (m_\tau/f_a)^2$. The challenge of this mode is rather related to the large amount of missing energy coming from the neutrinos in the tau decays that makes it difficult to detect the missing energy associated to possible ALP emission. This is illustrated in the second row of Figure 3, where the distribution of missing transverse energy (MET) and transverse momentum of the most energetic lepton (PT) are shown for events with taus decaying leptonically into different flavour leptons, $\tau^+\tau^- \rightarrow e^+\mu^- + \text{MET}$ or $e^-\mu^+ + \text{MET}$. As we can see, both the MET and PT distributions from $e^+e^- \rightarrow \tau^+\tau^-a$ and $e^+e^- \rightarrow \tau^+\tau^-$ are similar. Therefore, in order to search for an ALP signal, one should rather rely on hadronically decaying taus, which requires modelling the tau tagging capabilities of the future detectors and the related uncertainty. We postpone such an analysis to future work and focus in the following on the muon mode only.




Figure 3: Distribution, after the basic selection cuts, of missing transverse energy (MET) and transverse momentum of the most energetic lepton (PT) for $e^+e^- \rightarrow \ell^+\ell^-a$ (signal), $e^+e^- \rightarrow \ell^+\ell^-\nu\bar{\nu}$ and $e^+e^- \rightarrow \ell^+\ell^-(\gamma)$ (background), with $\ell = \mu$ (first row), $\ell = \tau$ with $\tau^+\tau^-$ decaying leptonically to different flavour leptons (second row). Note that (unlike $e^+e^- \rightarrow \ell^+\ell^-a$ and $e^+e^- \rightarrow \ell^+\ell^-\nu\bar{\nu}$) the $e^+e^- \rightarrow \ell^+\ell^-$ is not normalised according to its cross section, otherwise its large rate would not let its distribution be clearly displayed.

We select $\mu^+\mu^- + \text{MET}$ events imposing the following basic angular and isolation requirements: $|\eta_{\mu_i}| < 2.5$ for the pseudorapidity of both muons, and $\theta_{\mu_1\mu_2} > 0.2$ for the separation angle between them. We label as μ_1 (μ_2) the most (least) energetic particle of the $\mu^+\mu^-$ pair. The first row of Figure 3 clearly shows that the MET and PT distributions provide an handle to disentangle the signal events $\mu^+\mu^-a$ from the background stemming from, respectively, $e^+e^- \rightarrow \mu^+\mu^-(\gamma)$ (where the photons are missed) and $e^+e^- \rightarrow \mu^+\mu^-\nu\bar{\nu}$. In particular, a cut MET > 7 GeV along with the following PT cut can eliminate entirely $e^+e^- \rightarrow \mu^+\mu^-(\gamma)$ with undetected photons entirely. On the other hand, an upper cut MET < 35 GeV can help reduce the $\mu^+\mu^-\nu\bar{\nu}$ background retaining most of the signal events, cf. Figure 3(a). Secondly, a cut on the momentum of the most energetic muon μ_1 , PT > 43 GeV, enables us to efficiently distinguish the signal from the neutrino background, as shown in Figure 3(b). To summarise, we impose the following requirements on our simulated events:

- **Basic selection:** $\theta_{\mu_1\mu_2} > 0.2$, $|\eta_{\mu_i}| < 2.5$;
- **Cut 1:** missing energy, $7 \text{ GeV} < \text{MET} < 35 \text{ GeV}$;
- **Cut 2:** μ_1 momentum, PT $> 43 \text{ GeV}$.

		Basic cuts	Cut 1	Cut 2
ALP Signal	acceptance	0.88	0.77	0.23
	σ [$\times 10^{-5}$ pb]	6.7	5.9	1.7
$\nu\bar{\nu}$ background	acceptance	0.91	0.77	0.0028
	σ [$\times 10^{-4}$ pb]	2.6	2.2	0.0081
γ background	acceptance	0.94	5×10^{-6}	0

Table 2: Signal and background cross section and acceptance after the cuts described in the main text. The signal cross section was calculated assuming $f_a/C_{\mu\mu}^A = 100$ GeV.

	$\mathcal{L} = 50 \text{ ab}^{-1}$	$\mathcal{L} = 100 \text{ ab}^{-1}$	$\mathcal{L} = 150 \text{ ab}^{-1}$
σ_{95} ($\sigma_{5\sigma}$) [$\times 10^{-6}$ pb]	1.3 (4.1)	0.88 (2.6)	0.71 (2.0)
f_{95} ($f_{5\sigma}$) [GeV]	770 (430)	930 (540)	1040 (620)
BR_{95} ($\text{BR}_{5\sigma}$) [$\times 10^{-11}$]	4.0 (13)	2.7 (8.5)	2.1 (6.6)

Table 3: Projected 95% CL limit (5σ -discovery sensitivity) on the signal cross section, the ALP-muon coupling $f_a/C_{\mu\mu}^A$, and $\text{BR}(Z \rightarrow \mu^+\mu^- a)$. See the text for details.

The efficiency of the above cuts is displayed in Table 2. As we can see, they can effectively remove the background while retaining about 23% of the signal events. We can use the resulting acceptance of signal and background to estimate the significance of the proposed search for a given value of the integrated luminosity \mathcal{L} . As customary, we define the effective significance of the signal in terms of the number of signal (n_s) and background (n_b) events after the cuts as

$$s = \frac{n_s}{\sqrt{n_b + n_s}}. \quad (13)$$

Setting $s = 2$ in this expression, we can derive the 95% confidence level (CL) upper limit on the number of signal events that, for a given \mathcal{L} , the future Tera Z factories can set. Similarly, setting $s = 5$, we obtain the value of n_s corresponding to a 5σ discovery. This information can be translated into limits on the signal cross section (and equivalently on the decay rate of $Z \rightarrow \mu^+\mu^- a$) and consequently on the ALP coupling to muons. In Table 3, we show the resulting limits for three benchmark values for the integrated luminosity that CEPC and FCC-ee expect to collect at two interaction points from the operation of the Z-pole run, as reported in [34–37, 45]. In particular, the table displays the 95% CL limits we obtain on the signal cross section ($\equiv \sigma_{95}$), on $f_a/C_{\mu\mu}^A$ ($\equiv f_{95}$), and on $\text{BR}(Z \rightarrow \mu^+\mu^- a)$ ($\equiv \text{BR}_{95}$), as well as the corresponding 5σ discovery values ($\sigma_{5\sigma}$, $f_{5\sigma}$, $\text{BR}_{5\sigma}$). As we can see, a search of this kind at a Tera Z factory can be sensitive to ALP decay constants up to about 1 TeV and yield $\text{BR}(Z \rightarrow \mu^+\mu^- a) \lesssim 2 \times 10^{-12}$ as the upper bound on the branching ratio for the Z decay into a muonic ALP.

The results of Table 3 can be employed to show the sensitivity reach of CEPC/FCC-ee on the ALP parameter space. In Figure 4, we plot the estimated 95% CL exclusion lines on the $(m_a, f_a/C_{\mu\mu}^A)$ plane for the case of an anomaly-free ALP ($E_{\text{UV}} = 0$) whose coupling to electron vanishes ($C_{ee}^A = 0$), showing three representative values of \mathcal{L} as in Table 3. Notice that the limits in the figure can be also read in terms of the dimensionless coupling g_μ , as defined in Eq. (2), that can be excluded down to $g_\mu \approx 10^{-4}$. In Figure 4, the number of signal events have been convoluted with the probability that the ALP decays inside a typical detector (to be concrete, we assumed a flight distance $L = 10$ m), taking into account the proper decay length $c\tau_a$ calculated from Eq. (8). As expected from the discussion in Section 2, this cuts off




Figure 4: Prospected CEPC/FCC-ee 95% CL exclusion on the $(m_a, f_a/C_{\mu\mu}^A)$ plane for a muonic ALP ($C_{ee}^A = 0$, $E_{UV} = 0$) with different assumptions for the integrated luminosity \mathcal{L} , as indicated. On the right side of the dashed grey line the proper decay length of the ALP, as calculated using Eq. (8), is $c\tau_a < 10$ m. The region to the left of the red dashed line is excluded by SN1987A data, according to the analysis in [30].

the sensitivity of our search for an invisible ALP around the kinematical threshold of the decay $a \rightarrow \mu^+\mu^-$. For $m_a \gtrsim 2m_\mu$, the model can be tested through searches for $Z \rightarrow \mu^+\mu^-\mu^+\mu^-$ — with the invariant mass $m_{\mu\mu}^2$ of one of the muon pairs featuring a resonance in correspondence of $m_{\mu\mu}^2 = m_a^2$ — as discussed in [18]. We note that Figure 1 shows how the parameter space that our search is sensitive to would be reduced if we switch on C_{ee}^A and E_{UV} . In particular, for a coupling to electrons of the same order as the one to muons, $C_{ee}^A \approx C_{\mu\mu}^A$, our search would lose sensitivity for $m_a \gtrsim 2m_e \approx 1$ MeV, where ALPs could be sought through the search for a di-electron resonance in $Z \rightarrow \mu^+\mu^-e^+e^-$.

Finally, Figure 4 also shows the SN1987A constraint on light particles coupling to muons, as calculated in [30], which excludes the displayed parameter space for $f_a/C_{\mu\mu}^A \gtrsim 50$ GeV, $m_a \lesssim \mathcal{O}(10)$ MeV. For models with $C_{ee}^A \neq 0$, supernova constraints and other star cooling bounds associated to ALP emission from electrons instead lose sensitivity for $f_a/C_{ee}^A \lesssim 10^5$ GeV (see e.g. [12]) leaving the range of parameters accessible at colliders unconstrained. This observation, as well as the interplay between our exclusion lines and the SN1987A bound in Figure 4, shows how searches for long-lived ALPs in $e^+e^- \rightarrow Z^{(*)} \rightarrow \ell^+\ell^-a$ at future colliders may be nicely complementary to astrophysical probes of such light particles, in particular in view of the uncertainty of the supernova constraint [31].

5 Summary and conclusions

Axion-like particles are common by-products of new physics models addressing outstanding problems of the Standard Model. In this note, motivated by the fact that interactions with SM leptons are unavoidable for a wide range of well-motivated ALP scenarios, we have discussed the capability of the proposed high-energy e^+e^- colliders CEPC and FCC-ee of probing this class of particles through their couplings with charged leptons. We considered as production process the ALP emission from leptons in the large samples of $Z \rightarrow \ell^+\ell^-$ decays that CEPC and FCC-ee will collect operating as Tera Z factories, focusing in particular on the case of long-lived

ALPs that would appear as missing energy in the detector.

We first calculated the rate of $Z \rightarrow \ell^+\ell^-a$, in order to analytically estimate the number of events that one can expect at Tera Z factories. Then we employed a fast simulation framework (MADGRAPH+PYTHIA+DELPHES) to compute the LO cross section of $e^+e^- \rightarrow \ell^+\ell^-a$ and that of the relevant backgrounds, as well as to generate the corresponding events. The main findings of our work can be summarised as follows.

- As is already clear from the analytical estimates in Section 3, the number of $e^+e^- \rightarrow e^+e^-a$ events that a Tera Z can yield is too low to constrain the ALP coupling to electrons, unless $f_a/C_{ee}^A \ll 100$ GeV—see Eq. (10)—a situation that may be difficult to map to a viable UV-complete model.
- Concerning the muon mode, the signal $e^+e^- \rightarrow \mu^+\mu^-a$ can emerge from the background constituted by $\mu^+\mu^-\nu\bar{\nu}$ and $\mu^+\mu^-\gamma$ (where γ gets undetected), if one applies simple cuts on the missing energy of the event and the momentum of the most energetic muon, see Table 2.
- We estimate that such a crude “cut and count” search—that could be certainly improved and optimised—may be sensitive to ALP-muon interactions characterised by $f_a/C_{\mu\mu}^A \lesssim 1$ TeV, or equivalently $g_\mu \gtrsim 10^{-4}$, corresponding to $\text{BR}(Z \rightarrow \mu^+\mu^-a) \gtrsim 2 \times 10^{-12}$ —see Table 3 and Figure 4. Our summary plot in Figure 4 also shows that our search would lose sensitivity for $m_a \gtrsim 2m_\mu$ —where Tera Z factories may instead search for $Z \rightarrow \mu^+\mu^-a \rightarrow \mu^+\mu^-\mu^+\mu^-$ —and highlights the interplay between collider searches and astrophysical constraints (from SN1987A).
- The tau channel $e^+e^- \rightarrow \tau^+\tau^-a$ enjoys a two order of magnitude enhancement compared to the muon one, $\sim (m_\tau/m_\mu)^2$, such that $\sigma(e^+e^- \rightarrow \tau^+\tau^-a) > \sigma(e^+e^- \rightarrow \tau^+\tau^-\nu\bar{\nu})$ for $f_a/C_{\tau\tau}^A \lesssim 600$ GeV. Nevertheless, the missing energy associated to the ALP is way more difficult to detect due to that carried by neutrinos from $\tau^+\tau^-$ decays. As a consequence, a dedicated study focusing on hadronic taus and the expected tagging efficiencies of the CEPC/FCC-ee detectors would be necessary to assess the Tera Z sensitivity on $f_a/C_{\tau\tau}^A$.

In conclusion, we believe that the above results reinforce the physics case of a Z -pole run of the proposed e^+e^- colliders. Indeed, our study provides a further example of how Tera Z factories, besides performing very precise electroweak measurements and conducting an excellent particle physics programme (see [55] for a recent overview), could also effectively contribute to searches for new physics particles.

Acknowledgements

We would like to thank Prof. Chen Ning for valuable discussions and Wang Bin for technical support. This work was supported by the Innovation and Entrepreneurship Training Program for College Students of Tianjin (No. 202110055325). LC is partially supported by the National Natural Science Foundation of China under the Grant No. 12035008.

References

- [1] R. D. Peccei and H. R. Quinn, *CP Conservation in the Presence of Instantons*, *Phys. Rev. Lett.* **38** (1977) 1440–1443.
- [2] F. Wilczek, *Problem of Strong P and T Invariance in the Presence of Instantons*, *Phys. Rev. Lett.* **40** (1978) 279–282.
- [3] S. Weinberg, *A New Light Boson?*, *Phys. Rev. Lett.* **40** (1978) 223–226.

- [4] Y. Chikashige, R. N. Mohapatra and R. Peccei, *Are There Real Goldstone Bosons Associated with Broken Lepton Number?*, *Phys. Lett. B* **98** (1981) 265–268.
- [5] G. Gelmini and M. Roncadelli, *Left-Handed Neutrino Mass Scale and Spontaneously Broken Lepton Number*, *Phys. Lett. B* **99** (1981) 411–415.
- [6] H. M. Georgi, S. L. Glashow and S. Nussinov, *Unconventional Model of Neutrino Masses*, *Nucl. Phys. B* **193** (1981) 297–316.
- [7] F. Wilczek, *Axions and Family Symmetry Breaking*, *Phys. Rev. Lett.* **49** (1982) 1549–1552.
- [8] D. B. Reiss, *Can the Family Group Be a Global Symmetry?*, *Phys. Lett. B* **115** (1982) 217–220.
- [9] D. Chang and G. Senjanovic, *On Axion and Familons*, *Phys. Lett. B* **188** (1987) 231.
- [10] Y. Ema, K. Hamaguchi, T. Moroi and K. Nakayama, *Flaxion: a minimal extension to solve puzzles in the standard model*, *JHEP* **01** (2017) 096, [[1612.05492](#)].
- [11] L. Calibbi, F. Goertz, D. Redigolo, R. Ziegler and J. Zupan, *Minimal axion model from flavor*, *Phys. Rev. D* **95** (2017) 095009, [[1612.08040](#)].
- [12] L. Calibbi, D. Redigolo, R. Ziegler and J. Zupan, *Looking forward to lepton-flavor-violating ALPs*, *JHEP* **09** (2021) 173, [[2006.04795](#)].
- [13] L. Di Luzio, R. Gröber and P. Paradisi, *Hunting for CP-violating axionlike particle interactions*, *Phys. Rev. D* **104** (2021) 095027, [[2010.13760](#)].
- [14] W. J. Marciano, A. Masiero, P. Paradisi and M. Passera, *Contributions of axionlike particles to lepton dipole moments*, *Phys. Rev. D* **94** (2016) 115033, [[1607.01022](#)].
- [15] M. Bauer, M. Neubert, S. Renner, M. Schnubel and A. Thamm, *Axionlike Particles, Lepton-Flavor Violation, and a New Explanation of a_μ and a_e* , *Phys. Rev. Lett.* **124** (2020) 211803, [[1908.00008](#)].
- [16] D. Buttazzo, P. Panci, D. Teresi and R. Ziegler, *Xenon1T excess from electron recoils of non-relativistic Dark Matter*, *Phys. Lett. B* **817** (2021) 136310, [[2011.08919](#)].
- [17] M. A. Buen-Abad, J. Fan, M. Reece and C. Sun, *Challenges for an axion explanation of the muon $g - 2$ measurement*, *JHEP* **09** (2021) 101, [[2104.03267](#)].
- [18] J. Liu, X. Ma, L.-T. Wang and X.-P. Wang, *The ALP explanation to muon $(g - 2)$ and its test at future Tera-Z and Higgs factories*, [2210.09335](#).
- [19] R. Essig, R. Harnik, J. Kaplan and N. Toro, *Discovering New Light States at Neutrino Experiments*, *Phys. Rev. D* **82** (2010) 113008, [[1008.0636](#)].
- [20] B. Döbrich, J. Jaeckel, F. Kahlhoefer, A. Ringwald and K. Schmidt-Hoberg, *ALPtraum: ALP production in proton beam dump experiments*, *JHEP* **02** (2016) 018, [[1512.03069](#)].
- [21] M. J. Dolan, T. Ferber, C. Hearty, F. Kahlhoefer and K. Schmidt-Hoberg, *Revised constraints and Belle II sensitivity for visible and invisible axion-like particles*, *JHEP* **12** (2017) 094, [[1709.00009](#)].
- [22] C. Cornella, P. Paradisi and O. Sumensari, *Hunting for ALPs with Lepton Flavor Violation*, *JHEP* **01** (2020) 158, [[1911.06279](#)].

- [23] M. Bauer, M. Neubert, S. Renner, M. Schnubel and A. Thamm, *Flavor probes of axion-like particles*, *JHEP* **09** (2022) 056, [[2110.10698](#)].
- [24] Y. M. Andreev et al., *Improved exclusion limit for light dark matter from $e+e-$ annihilation in NA64*, *Phys. Rev. D* **104** (2021) L091701, [[2108.04195](#)].
- [25] G. G. Raffelt, *Stars as laboratories for fundamental physics*. 1996.
- [26] N. Viaux, M. Catelan, P. B. Stetson, G. Raffelt, J. Redondo, A. A. R. Valcarce et al., *Neutrino and axion bounds from the globular cluster M5 (NGC 5904)*, *Phys. Rev. Lett.* **111** (2013) 231301, [[1311.1669](#)].
- [27] M. M. Miller Bertolami, B. E. Melendez, L. G. Althaus and J. Isern, *Revisiting the axion bounds from the Galactic white dwarf luminosity function*, *JCAP* **1410** (2014) 069, [[1406.7712](#)].
- [28] G. G. Raffelt, *Astrophysical methods to constrain axions and other novel particle phenomena*, *Phys. Rept.* **198** (1990) 1–113.
- [29] R. Bollig, W. DeRocco, P. W. Graham and H.-T. Janka, *Muons in Supernovae: Implications for the Axion-Muon Coupling*, *Phys. Rev. Lett.* **125** (2020) 051104, [[2005.07141](#)].
- [30] D. Croon, G. Elor, R. K. Leane and S. D. McDermott, *Supernova Muons: New Constraints on Z' Bosons, Axions and ALPs*, *JHEP* **01** (2021) 107, [[2006.13942](#)].
- [31] N. Bar, K. Blum and G. D’Amico, *Is there a supernova bound on axions?*, *Phys. Rev. D* **101** (2020) 123025, [[1907.05020](#)].
- [32] W. Altmannshofer, J. A. Dror and S. Gori, *New Insights Into Axion-Lepton Interactions*, [2209.00665](#).
- [33] G. Krnjaic, G. Marques-Tavares, D. Redigolo and K. Tobioka, *Probing Muonphilic Force Carriers and Dark Matter at Kaon Factories*, *Phys. Rev. Lett.* **124** (2020) 041802, [[1902.07715](#)].
- [34] CEPC STUDY GROUP collaboration, *CEPC Conceptual Design Report: Volume 1 - Accelerator*, [1809.00285](#).
- [35] CEPC STUDY GROUP collaboration, M. Dong et al., *CEPC Conceptual Design Report: Volume 2 - Physics & Detector*, [1811.10545](#).
- [36] FCC collaboration, A. Abada et al., *FCC Physics Opportunities: Future Circular Collider Conceptual Design Report Volume 1*, *Eur. Phys. J. C* **79** (2019) 474.
- [37] FCC collaboration, A. Abada et al., *FCC-ee: The Lepton Collider: Future Circular Collider Conceptual Design Report Volume 2*, *Eur. Phys. J. ST* **228** (2019) 261–623.
- [38] M. Bauer, M. Heiles, M. Neubert and A. Thamm, *Axion-Like Particles at Future Colliders*, *Eur. Phys. J. C* **79** (2019) 74, [[1808.10323](#)].
- [39] H.-Y. Zhang, C.-X. Yue, Y.-C. Guo and S. Yang, *Searching for axionlike particles at future electron-positron colliders*, *Phys. Rev. D* **104** (2021) 096008, [[2103.05218](#)].
- [40] C.-X. Yue, S. Yang, H. Wang and N. Zhang, *Prospects for detecting axionlike particles via the decay $Z \rightarrow af\bar{f}$ at future Z factories*, *Phys. Rev. D* **105** (2022) 115027, [[2204.04702](#)].

- [41] C.-T. Lu, *Lighting Electroweak-Violating ALP-Lepton Interactions at e^+e^- and ep Colliders*, [2210.15648](#).
- [42] M. Bauer, M. Neubert, S. Renner, M. Schnubel and A. Thamm, *The Low-Energy Effective Theory of Axions and ALPs*, *JHEP* **04** (2021) 063, [[2012.12272](#)].
- [43] M. Bauer, M. Neubert and A. Thamm, *Collider Probes of Axion-Like Particles*, *JHEP* **12** (2017) 044, [[1708.00443](#)].
- [44] P. D. Group, R. L. Workman, V. D. Burkert, V. Crede, E. Klempt, U. Thoma et al., *Review of Particle Physics*, *Progress of Theoretical and Experimental Physics* **2022** (08, 2022) .
- [45] J. Guimaraes da Costa, “*Status of the Circular Electron Positron Collider.*” <https://indico.ihep.ac.cn/event/17020/contributions/117889/attachments/64239/74992/20221024-CEPC-Workshop-goodR.pdf>, 2022.
- [46] A. Alloul, N. D. Christensen, C. Degrande, C. Duhr and B. Fuks, *FeynRules 2.0 - A complete toolbox for tree-level phenomenology*, *Comput. Phys. Commun.* **185** (2014) 2250–2300, [[1310.1921](#)].
- [47] C. Degrande, C. Duhr, B. Fuks, D. Grellscheid, O. Mattelaer and T. Reiter, *UFO - The Universal FeynRules Output*, *Comput. Phys. Commun.* **183** (2012) 1201–1214, [[1108.2040](#)].
- [48] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer and T. Stelzer, *MadGraph 5 : Going Beyond*, *JHEP* **06** (2011) 128, [[1106.0522](#)].
- [49] F. Maltoni and T. Stelzer, *MadEvent: Automatic event generation with MadGraph*, *JHEP* **02** (2003) 027, [[hep-ph/0208156](#)].
- [50] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159–177, [[1410.3012](#)].
- [51] DELPHES 3 collaboration, J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaître, A. Mertens et al., *DELPHES 3, A modular framework for fast simulation of a generic collider experiment*, *JHEP* **02** (2014) 057, [[1307.6346](#)].
- [52] C. Chen, X. Mo, M. Selvaggi, Q. Li, G. Li, M. Ruan et al., *Fast simulation of the CEPC detector with Delphes*, [1712.09517](#).
- [53] R. Brun and F. Rademakers, *ROOT: An object oriented data analysis framework*, *Nucl. Instrum. Meth. A* **389** (1997) 81–86.
- [54] E. Conte, B. Fuks and G. Serret, *MadAnalysis 5, A User-Friendly Framework for Collider Phenomenology*, *Comput. Phys. Commun.* **184** (2013) 222–256, [[1206.1599](#)].
- [55] CEPC PHYSICS STUDY GROUP collaboration, H. Cheng et al., *The Physics potential of the CEPC. Prepared for the US Snowmass Community Planning Exercise (Snowmass 2021)*, in *2022 Snowmass Summer Study*, 5, 2022, [2205.08553](#).