

# Response to Comment arXiv:2210.10539v1

Ivan Horváth<sup>1,2,\*</sup> and Peter Markoš<sup>3,†</sup>

<sup>1</sup>Nuclear Physics Institute CAS, 25068 Řež (Prague), Czech Republic

<sup>2</sup>University of Kentucky, Lexington, KY 40506, USA

<sup>3</sup>Dept. of Experimental Physics, Faculty of Mathematics, Physics and Informatics, Comenius University in Bratislava, Mlynská Dolina 2, 842 28 Bratislava, Slovakia

(Dated: Dec 5, 2022)

In recent work [1], we calculated the infrared (IR) effective counting dimension  $d_{\text{IR}}$  [2–4] for critical states of 3D Anderson models in universality classes O, U, S and AIII. The results entailed two new messages. (m1) Space effectively occupied by critical electron is of dimension  $d_{\text{IR}} \approx 8/3$ . Demonstrated properties of effective counting imply that the meaning and relevance of this is fully analogous to e.g. Minkowski dimension of ternary Cantor set being  $d_{\text{UV}} = \log_3(2)$ . Indeed, both statements express the scaling of properly defined physical volume; both dimensions are measure-based. (m2) The values of  $d_{\text{IR}}$  in studied classes coincide to better than two parts per mill with comparable errors. We dubbed this finding superuniversality of  $d_{\text{IR}}$  since other critical indices tend to differ to a notably larger degree. Exact superuniversality offers itself as a possibility, but obviously cannot be demonstrated in a finite numerical computation, only ascertained with better bounds or disproved.

In Ref. [5] Burmistrov claims that he “proved” inexact nature of (m2) by invoking the multifractal (MF) formalism. He derived MF representation for average effective count  $\langle \mathcal{N}_* \rangle$ , involving  $f(3)$  and  $\alpha_0$  ( $f(\alpha_0) = 3$ ), namely

$$\langle \mathcal{N}_* \rangle \simeq 4c_0 \frac{L^{f(3)}}{\sqrt{\ln(L)}} \quad , \quad c_0 = \sqrt{|f''(\alpha_0)|/(2\pi)} \quad (1)$$

The asymptotically equal sign ( $\simeq$ ) suggests that (1) conveys an exact  $L \rightarrow \infty$  leading term of  $\langle \mathcal{N}_* \rangle$ . However, in the ensuing discussion,  $4c_0$  appears to be treated as approximation to proportionality constant, which is what we will assume. Burmistrov then concludes that (1) “proves the absence of “superuniversality” of  $\langle \mathcal{N}_* \rangle$ ” due to numerically known class-dependent values of  $f(3)$  and  $c_0$ . We note in passing that superuniversality of  $\langle \mathcal{N}_* \rangle$ , exact or not, was in fact not claimed or invoked in [1].

To put formula (1) in context, recall that MF formalism was created to describe UV measure singularities arising e.g. in strange attractors [6, 7]. The method identifies sets  $A_\alpha$  of local singularities with Hölder strength  $\alpha > 0$ , treating their Hausdorff dimensions  $f_H(\alpha) = \dim_H(A_\alpha)$  as characteristics of interest. Note that  $f_H(\alpha)$  is a proper measure-based dimension of spatial set  $A_\alpha$ .

Common variation is the moment method which avoids computing  $\alpha(x)$  at each point for the price of coarsening

the singularity information. Evaluation of the associated  $f_m(\alpha)$  proceeds by computing continuum of generalized dimensions which are not measure-based and do not represent dimensions of space in themselves. Nevertheless,  $f_H(\alpha)$  and  $f_m(\alpha)$  coincide under certain analytic assumptions by virtue of associated transformations. However, the moment method does not identify the singularity set  $A_\alpha$ . What it does provide is a trick to evaluate  $f_H(\alpha)$  valid in certain circumstances. Without explicit check that  $f_H(\alpha) = f_m(\alpha)$  or explicit verification that needed analytic properties are met, one has no choice but to treat  $f_m(\alpha)$  as approximation to  $f_H(\alpha)$ .

In Anderson criticality, the situation is somewhat different. This is an IR problem which means that the notion of local singularity is absent and the very concept of  $A_\alpha$  becomes fuzzy. The moment method is then normally taken to *define* MF spectrum  $f(\alpha) \equiv f_m(\alpha)$  upon formally replacing the UV cutoff  $a$  with  $1/L$ . However, one is left with little regarding the information on populations (subsets of space) whose dimensions is  $f_m(\alpha)$  meant to represent. Unlike in UV problem of a strange attractor, there is no readily available analogue of  $A_\alpha$  and  $f_H(\alpha)$  to independently check upon.

Formula (1) may prove valuable in this regard since it represents MF *prediction* ( $f \equiv f_m$ ) for  $\langle \mathcal{N}_* \rangle$  which has independent definition and well-defined spatial meaning. Degree of consistency between true  $\langle \mathcal{N}_* \rangle$  and (1) can thus test assumptions associated with the MF approach. We perform such comparison to  $\langle \mathcal{N}_* \rangle$  data of Ref. [1]. To minimize finite- $L$  systematics, only the upper segment of all data ( $64 \leq L \leq 128$ ) is used to fit for  $c_0$  and  $f(3)$ . We find that instead of improving, fit form (1) worsens  $\chi^2/\text{DOF}$  relative to pure power ( $1.5 \rightarrow 2.9$ ). More importantly, resulting  $f(3)$  is sharply inconsistent with the MF-computed value  $f(3) = 2.733(3)$  [8], used in Burmistrov argument. Described results refer to O class but situation in other classes is the same.

The above is conveyed by Fig. 1(a) where  $1/L$  behavior of  $f(3)$ , expressed from Eq. (1), is shown together with fitted  $1/L=0$  value (with error) and the associated MF prediction. Analogous plot for pure power is also shown (full circles). Fitted  $d_{\text{IR}}$  here is consistent with one obtained from the extrapolated pair method [1]. Approach of data to fitted values in both cases is of course by design, similarly to plots in Fig. 1 of Ref. [5]. Note also that the obtained  $c_0$  differs roughly by factor of two from its approximate MF prediction as shown in Fig. 1(b).

The above discussion warrants the following points.

\* [ihorv2@g.uky.edu](mailto:ihorv2@g.uky.edu)

† [peter.markos@fmph.uniba.sk](mailto:peter.markos@fmph.uniba.sk)




FIG. 1. MF-computed indices  $f(3)$  [8] and  $c_0$  compared to those obtained from Eq. (1) and  $\langle N_* \rangle$  data of Ref. [1].

(i) Given the scale of inconsistencies found in the MF prediction (1), we conclude that arguments of Ref. [5] do not have tangible bearing on our (m1) and (m2), including the possibility of exact superuniversality in  $d_{\text{IR}}$ . One should also keep in mind that superuniversality and  $d_{\text{IR}} \approx 8/3$  are two distinct conclusions.

(ii) Our findings suggest that derivation of (1) may involve assumptions on  $\alpha$ -populations that are not justified. For example, setting  $p(x) \equiv L^{-\alpha(x)}$  ( $p(x)$  = probability at  $x$ ), which forces certain structure on  $\alpha$ -populations without knowing what they actually are, may be too cavalier. Showing that this is harmless in  $L \rightarrow \infty$  limit is non-trivial even for moderately complex populations. Similar goes for setting  $\nu(\alpha) \equiv L^{f(\alpha)}$  where  $\nu(\alpha)$  is the  $\alpha$ -population count. Assumptions on analyticity of  $f(\alpha)$ ,  $f_m(\alpha)$  and quadratic local nature of maximum at  $\alpha_0$  are also present. Characterizations such as “exact” and “proof” in this context are perhaps too strong and premature.

(iii) Arguments of [5], leading to claim of systematic effect in the analysis of [1], use numerical results of [8]. But systematics involved in the latter work is not discussed

although smaller systems and smaller statistics were used. (Systematics in [1] was also studied in Ref. [4] albeit from a different angle.) Are there logs affecting standard MF analyses, appearing e.g. via arbitrary replacement  $p \equiv L^{-\alpha} \rightarrow p \equiv \text{const } L^{-\alpha}$ . Are they accounted for?

(iv) The presence of  $\log(L)$  powers in dimension estimates is always possible since they leave it intact. MF arguments are not necessary to invoke the possibility. But in the absence of a firm prediction, establishing their presence is a delicate numerical issue. Indeed, it is difficult to show that the influence of an unknown log is larger than that of subleading powers. Statistical strength of available data is frequently not sufficient to do that. Occam’s razor approach is then an accepted resolution.

(v) We strongly disagree with the comment “ $d_{\text{IR}}$  is nothing but  $f(d)$ ”. There was also a calculus student who once declared “ $\pi$  is nothing but  $22/7 - \int_0^1 dx x^4(1-x)^4/(1+x^2)$ ”. Perhaps there is a bit more to  $d_{\text{IR}}$  than the currently hypothetical  $d_{\text{IR}} = f_m(3)$ . Its meaning is expressed in (m1), being worked out in requisite detail by Refs. [2–4].

(vi) Regarding comments on prospects for superuniversality in various dimensions, we repeat that our claim is only for  $d_{\text{IR}}$  at level better than two parts per mill in  $D=3$ . The gist of our message is that properties of space in Anderson transitions may be special relative to other characteristics. Scenario where superuniversality in  $d_{\text{IR}}$  is violated in  $D=2 + \epsilon$  but very accurate or exact in  $D=3$  and higher dimensions is not contradictory in our opinion.

## ACKNOWLEDGMENTS

P.M. was supported by Slovak Grant Agency VEGA, Project n. 1/0101/20.

- 
- [1] I. Horváth and P. Markoš, *Phys. Rev. Lett.* **129**, 106601 (2022), [arXiv:2110.11266 \[cond-mat.dis-nn\]](#).
  - [2] I. Horváth and R. Mendris, *Entropy* **22**, 1273 (2020), [arXiv:1807.03995 \[quant-ph\]](#).
  - [3] A. Alexandru and I. Horváth, *Phys. Rev. Lett.* **127**, 052303 (2021), [arXiv:2103.05607 \[hep-lat\]](#).
  - [4] I. Horváth, P. Markoš, and R. Mendris, (2022), [arXiv:2205.11520 \[hep-lat\]](#).
  - [5] I. S. Burmistrov, arXiv e-prints (2022), [arXiv:2210.10539 \[cond-mat.dis-nn\]](#).
  - [6] K. Falconer, *Fractal Geometry: Mathematical Foundations and Applications*, 3rd ed. (Wiley, 2014).
  - [7] T. C. Halsey, M. H. Jensen, L. P. Kadanoff, I. Procaccia, and B. I. Shraiman, *Phys. Rev. A* **33**, 1141 (1986).
  - [8] L. Ujfalusi and I. Varga, *Phys. Rev. B* **91**, 184206 (2015).