# On the Proof of Chiral Symmetry Breaking from Anomaly Matching in QCD-like Theories

Luca Ciambriello,  $^{1,\,*}$ Roberto Contino,  $^{2,\,3,\,\dagger}$  Andrea Luzio,  $^{4,\,5,\,\ddagger}$ 

Marcello Romano,<sup>6,§</sup> and Ling-Xiao Xu<sup>7,8,9,¶</sup>

<sup>1</sup>Interdisciplinary Laboratories for Advanced Materials Physics (i-LAMP) and Dipartimento di Matematica e Fisica, Università Cattolica del Sacro Cuore, Brescia, Italy

<sup>2</sup>Dipartimento di Fisica, Sapienza Università di Roma, Italy

<sup>3</sup>Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Roma, Italy

<sup>4</sup>Scuola Normale Superiore, Pisa, Italy

<sup>5</sup>Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Pisa, Italy

<sup>6</sup> Université Paris-Saclay, CNRS, CEA, Institut de Physique Théorique, 91191, Gif-sur-Yvette, France

<sup>7</sup>Dipartimento di Fisica e Astronomia 'G. Galilei', Università di Padova, Italy

<sup>8</sup>Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Padova, Italy

<sup>9</sup>Abdus Salam International Centre for Theoretical Physics, Trieste, Italy

We critically reconsider the argument based on 't Hooft anomaly matching that aims at proving chiral symmetry breaking in 4d QCD-like theories with  $N_c > 2$  colors and  $N_f$  flavors. The main line of reasoning relies on a property of the solutions of the anomaly matching and persistent mass equations called  $N_f$ -independence. In previous works, the validity of  $N_f$ -independence was assumed based on qualitative arguments, but it was never proven rigorously. We provide a detailed proof and clarify under which (dynamical) conditions it holds. Our results are valid for a generic spectrum of massless composite fermions including baryons and exotics.

<sup>\*</sup> luca.ciambriello@unicatt.it

 $<sup>^{\</sup>dagger}$ roberto.contino@uniroma1.it

 $<sup>^{\</sup>ddagger}$  and rea.luzio@sns.it

<sup>&</sup>lt;sup>§</sup> marcello.romano@ipht.fr

<sup>¶</sup> lxu@ictp.it

#### CONTENTS

I. Introduction	2
II. Characterizing Massless Bound States	6
A. Massless Bound States and Composite Operators	6
B. Equivalent Tensors	9
III. Anomaly Matching and Persistent Mass Conditions	10
A. Anomaly Matching Conditions	11
B. Persistent Mass Conditions	12
IV. Proof of Chiral Symmetry Breaking from $N_f$ -independence	15
A. Proof of $N_f$ -independence	17
1. PMC $[N_f]$ are $N_f$ -dependent in general	18
2. Condition for $N_f$ -independence of PMC $[N_f, 1]$	19
3. Condition for $N_f$ -independence of AMC $[N_f]$	23
4. Proof of Eq. (IV.23): downlifting is the inverse of uplifting	25
V. Conclusions	27
Acknowledgments	28
A. Technicalities on Anomalies	29
B. Equivalent Tensors	30
C. PMC with equal masses	34
D. General Proof of Eq. (IV.23)	37
References	39

### I. INTRODUCTION

Confinement and chiral symmetry breaking are the two phenomena exhibited by QCD at low energy. Their full theoretical understanding still remains elusive due to the fact that QCD gets strongly coupled in the infrared (IR). One particular question is whether confinement enforces chiral symmetry breaking *inevitably* in QCD. In this work, we address this question in QCD-like theories with  $SU(N_c)$  gauge group and  $N_f$  flavors of vectorlike quarks in the fundamental representation. A QCD-like theory is expected to confine when the number of massless flavors is smaller than a critical value  $N_f^{CFT}$ , above which it is either conformal (for  $N_f^{CFT} \leq N_f < 11N_c/2$ ) or IR free (for  $N_f \geq 11N_c/2$ ). Therefore, we consider  $N_c > 2$  and  $2 \leq N_f < N_f^{CFT}$ . For  $N_c = 2$ , quarks and antiquarks transform in equivalent color representations, and this case needs to be addressed separately.

Among the available theoretical tools, 't Hooft anomaly matching [1] stands out as one of the few that can constrain the IR spectrum of confining gauge theories. A QCD-like theory with  $N_f$  massless flavors has a global chiral symmetry <sup>1</sup>

$$\mathcal{G}[N_f] = SU(N_f)_L \times SU(N_f)_R \times U(1)_B, \qquad (I.1)$$

where  $U(1)_B$  is baryon number. Confinement implies that the low-energy spectrum consists of color-singlet bound states.<sup>2</sup> If chiral symmetry is unbroken by the vacuum, then such bound states transform in non-trivial irreducible representations (irreps) of  $\mathcal{G}[N_f]$ . Furthermore, massless composite fermions must exist to reproduce in the infrared (IR) the anomalies of  $\mathcal{G}[N_f]$  induced by the quarks in the ultraviolet (UV). In other words, the UV and IR anomalies of the chiral symmetry group  $\mathcal{G}[N_f]$  need to be matched. This implies a set of equations called Anomaly Matching Conditions (AMC). They were first derived by 't Hooft in Ref. [1] by coupling background gauge fields to the anomalous quark currents and introducing color-neutral spectator fermions. A derivation based on analyticity and unitarity was given in Refs. [2, 3], see also Ref. [4] for implications in the large- $N_c$  limit.

Besides 't Hooft anomaly matching, the low-energy spectrum of QCD-like theories also needs to satisfy Persistent Mass Conditions (PMC) [5], originally formulated by 't Hooft as decoupling conditions [1]. As we will see, PMC are an implication of the Vafa-Witten theorem valid for vectorlike gauge theories with  $\theta = 0$  [6]. Their physical significance can be summarized as follows: when *i* flavors are given finite and unequal masses, with  $1 \le i \le N_f - 2$ , the symmetry  $\mathcal{G}[N_f]$  is explicitly broken to

$$\mathcal{G}[N_f, i] = SU(N_f - i)_L \times SU(N_f - i)_R \times U(1)_{H_1} \times \dots \times U(1)_{H_i} \times U(1)_B, \qquad (I.2)$$

where each vector factor  $U(1)_{H_k}$  is defined such that the k-th massive quark has charge  $H_k = 1$ , while the other quarks have  $H_k = 0$ . Bound states that are charged under  $U(1)_{H_1} \times \cdots \times U(1)_{H_i}$ must contain massive flavors of quarks or antiquarks as their microscopic constituents, and the

<sup>&</sup>lt;sup>1</sup> In this paper, we neglect the discrete factors that appear in the global symmetry group. We define  $U(1)_B$  in a standard way and assign baryon number  $1/N_c$  to quarks.

<sup>&</sup>lt;sup>2</sup> There exists no rigorous definition of confinement in QCD with dynamical quarks in the fundamental representation. In this paper, by confinement we mean a phase of the theory where the only asymptotic states are color-singlet particles. We assume, in particular, that asymptotic states do not include extended objects interpolated by nonlocal operators.

unbroken chiral symmetry cannot prevent them from acquiring a mass. In other words, such composite fermions with non-zero charge under  $U(1)_{H_1} \times \cdots \times U(1)_{H_i}$  must furnish vectorlike representations under  $\mathcal{G}[N_f, i]$ .<sup>3</sup> Mathematically, this translates into the PMC. In particular, PMC equations with more than one massive flavors were not fully exploited in the previous literature but, as it turns out, they are key in the approach of Ref. [7].

One possible strategy to prove chiral symmetry breaking consists in making use of AMC and PMC. These form a system of linear equations in the indices of massless composite fermions. The index of a given irrep of  $\mathcal{G}[N_f]$  is defined as the number of left-handed minus right-handed helicity states in the spectrum transforming as such irrep. If there exist no integral values of these indices that satify both AMC and PMC, then chiral symmetry breaking must occur. This line of reasoning was first proposed by 't Hooft [1] and afterwards pursued by many other authors [2, 8–13] under specific assumptions. In his seminal paper, 't Hooft worked out the cases  $N_c = 3$  and  $N_c = 5$ , showing that no integral solution exists for  $N_f > 2$ . He assumed that the spectrum of massless fermions include only baryons with baryon number b = 1, and that mixed representations have vanishing index for  $N_c = 5$ . Frishman et al. [2] extended the analysis to generic  $N_c$  by working under the same assumptions. They found no integral solution. (The real solution found in Ref. [2] is one where 'elbow'-shape Young Tableaux have indices  $\pm 1/N_c^2$ , while all the other indices are zero. For  $N_c = 5$ , this coincides with the real solution found by 't Hooft.) Considering only massless baryons, Cohen and Frishman [12] further showed that integer solutions do exist for  $N_f = 2$  and arbitrary  $N_c$  besides the  $\sigma$ -model solution already found by 't Hooft.

An important step forward towards a general proof was made by Farrar in Ref. [9]. She considered the possible existence of exotic massless bound states in the spectrum, i.e. bound states with antiquark constituents, and was able to prove chiral symmetry breaking by assuming ' $N_f$ independence'. According to the latter, the solutions of AMC and PMC are the same for any number of flavors. This property was first proposed by 't Hooft in Ref. [1] and justified by him through qualitative dynamical considerations. It was later suggested by Ref. [2] that it comes as a consequence of the PMC, and similar arguments were presented in a more quantitative fashion in [10, 11]. On the other hand, Cohen and Frishman pointed out in Ref. [12] that, if only minimal baryons are considered as massless bound states, it is necessary to separate the regimes with  $N_f > N_c$  and  $N_f \leq N_c$ . Their work suggests that  $N_f$ -independence cannot hold in general, but only for sufficiently large  $N_f$  and according to the spectrum of putative massless composite

<sup>&</sup>lt;sup>3</sup> Here we focus on bound states with non-zero baryon number, which therefore transform as complex representations of  $\mathcal{G}[N_f, i]$ . Only Dirac mass terms are allowed for these states.

fermions.

A different argument to prove chiral symmetry breaking using 't Hooft anomaly matching was pursued independently by Schwimmer in Ref. [8]. He considered an  $SU(N_f|N_f)$  superalgebra whose grading acts so that odd supergenerators mix massless states with opposite helicity. Although strictly speaking  $SU(N_f|N_f)$  is not a symmetry of a QCD-like theory, it can be used as a classification group. Schwimmer argued that, for a purely baryonic massless spectrum, irreducible representations of  $SU(N_f|N_f)$  satisfy PMC trivially, but they cannot furnish an integral solution of AMC. In order to elevate this simple and elegant argument to the level of a proof of chiral symmetry breaking, one should demonstrate that it works for a generic spectrum with exotics and that it gives the most general solution of PMC, i.e. that every PMC solution is a collection of superalgebra irreps. This evidence is still missing.

Despite many papers on the subject and the various approaches explored in the literature, proving chiral symmetry breaking in a generic QCD-like confining theory is still an open issue. In particular, one would like a proof valid for the most generic spectrum of massless bound states and for arbitrary  $N_c$  and  $N_f$ . In this work, we reconsider the argument based on 't Hooft anomaly matching and  $N_f$ -independence with the aim of formulating it in a rigorous fashion. We derive a proof of  $N_f$ -independence as a mathematical property of AMC and PMC equations and clarify under which assumptions it holds. The latter turn out to be dynamical requirements on the spectrum of massless bound states. Our study extends previous analyses and corrects a few statements on the form of the solution that were made in the literature. We report in a companion paper [14] the explicit solutions of the system of AMC and PMC equations that we derived for specific values of  $N_c$  and  $N_f$ , and focus here on the more formal arguments. In a third paper, we further provide two novel arguments to prove chiral symmetry breaking which apply to a generic spectrum of massless bound states [7].

There exist other arguments, not based on 't Hooft anomaly matching, that can support chiral symmetry breaking. One attempt makes use of the mass inequality  $m_B \ge m_{\pi}$ , which has been proven by Weingarten to hold in QCD-like theories [15], see also [16]. Here  $m_B$  is the mass of the lightest baryon and  $m_{\pi}$  is the mass of the lightest state interpolated by the operator  $\bar{q}\gamma^5 q$ . If one assumes that chiral symmetry is not spontaneously broken, then matching 't Hooft anomalies requires  $m_B = 0$ , and the mass inequality implies  $m_{\pi} = 0$ . Although this proves that there must exist at least one massless scalar in the spectrum (something which is in fact rather interesting per se), it does not contradict the initial assumption of unbroken chiral symmetry, since the axial current  $J^{\mu}_A$  and the state  $\pi$  transform differently under  $SU(N_f)_L \times SU(N_f)_R$ , hence  $\langle 0|J^{\mu}_A|\pi\rangle = 0$ . Therefore, this argument does not seem conclusive. A different approach aims at deducing the properties of real QCD from those of  $\mathcal{N} = 1$  SUSY QCD [17, 18] by adding small supersymmetrybreaking terms [19–25]. Although the IR dynamics of near-SUSY QCD can be characterized rather precisely and various exotic phases can be identified, extrapolating to real QCD by sending the supersymmetry-breaking scale to infinity is challenging due to the possible occurence of phase transitions [25, 26]. In this work, we do not rely on supersymmetry.

The paper is organized as follows.

- In Section II we characterize the most general spectrum of massless composite fermions using tensor notation. Massless fermions can be classified by their representations under the chiral symmetry group  $\mathcal{G}[N_f]$ , and are interpolated from the vacuum by composite operators which transform as tensors of  $\mathcal{G}[N_f]$ .
- Section III gives a concise review on AMC and PMC equations. In particular, we justify PMC from the Vafa-Witten theorem [6], and make some original considerations regarding PMC with more than one massive flavors.
- We enunciate and prove our theorem on  $N_f$ -independence in Section IV. There we derive several features and important details that were overlooked in the previous literature, in particular the conditions under which  $N_f$ -independence holds true.

Let us define 'class A' tensors as composite operators made of  $n_q$  quarks and  $n_{\bar{q}}$  antiquarks (where  $n_q = bN_c + n_{\bar{q}}$ , and b is the baryon number) satisfying  $N_f > bN_c + 2 n_{\bar{q}}$ . We find that  $N_f$ -independence is valid when the only massless bound states to have non-vanishing indices are those interpolated by 'class A' tensors. For the baryonic case, our result is consistent with Ref. [12] when b = 1 and  $n_{\bar{q}} = 0$ .

• Finally, we conclude in Section V and clarify the most technical aspects of our analysis in Appendices A, B, C and D.

Throughout the paper, several examples are provided which we hope can be useful for the reader.

### II. CHARACTERIZING MASSLESS BOUND STATES

### A. Massless Bound States and Composite Operators

We focus on values of  $N_c$  and  $N_f$  for which the theory is in the confining phase, and assume that chiral symmetry is unbroken by the strong dynamics. This requires the existence of massless fermionic bound states to reproduce 't Hooft anomalies in the IR. According to the Weinberg-Witten theorem [27], a theory with Lorentz-covariant conserved currents cannot contain massless particles of spin j > 1/2 that have a non-vanishing corresponding charge. Therefore, we only consider fermionic bound states of spin j = 1/2. Massless bound states in the spectrum can be classified in terms of irreducible representations (irreps) of the flavor symmetry group  $\mathcal{G}[N_f]$ , cf. Eq. (I.1). In particular, one can study the gauge-invariant composite local operators that interpolate massless fermions from the vacuum. Being color singlets, such operators are characterized by the sum rule

$$n_q - n_{\bar{q}} = bN_c , \qquad (\text{II.1})$$

where  $n_q$ ,  $n_{\bar{q}}$  are the numbers of quark and antiquark fields forming the operator, and b is its baryon number. Since the bound state is a fermion, b and  $N_c$  must be both odd integers. As explained in the next section, one can focus on bound states with positive b without loss of generality.

We define *baryons* those bound states that can be interpolated by operators with only quarks (i.e. with  $n_{\bar{q}} = 0$ ), and call *exotics* all the other bound states. Exotics thus transform in representations of  $\mathcal{G}[N_f]$  that cannot be obtained through operators with only quarks.

As a consequence of their quark and antiquark content, composite operators transform as tensors of  $SU(N_f)_L \times SU(N_f)_R$  with  $n_q$  contravariant upper indices and  $n_{\bar{q}}$  covariant lower indices. One can thus establish a correspondence between the space of interpolating operators and the abstract space of tensors of  $SU(N_f)_L \times SU(N_f)_R$ . Notice that, while a given operator corresponds to a single tensor, the converse is not true: different operators with the same quark and antiquark (hence flavor) content will correspond to the same tensor. Traceless tensors <sup>4</sup> transform as irreducible representations of the global symmetry group, and thus play an important role. When representing traceless tensors, we will distinguish between upper (lower) indices of  $SU(N_f)_L$ , denoted as  $n_L$  ( $\bar{n}_L$ ), and upper (lower) indices of  $SU(N_f)_R$ , denoted as  $n_R$  ( $\bar{n}_R$ ). For each set of n indices, one can consider an associated Young Tableau (YT), denoted as  $\{n\}$  in the following, with n boxes. We will use the compact notation  $T_{\{\bar{n}_L\};\{\bar{n}_R\}}^{\{n_R\}}$  to indicate traceless tensors of total upper rank  $n = n_L + n_R$ and total lower rank  $\bar{n} = \bar{n}_L + \bar{n}_R$ . The corresponding operators will have  $n_q = n + \Delta$  quarks and  $n_{\bar{q}} = \bar{n} + \Delta$  antiquarks, where  $\Delta$  is the number of quark and antiquark indices contracted in a flavor singlet. Each traceless tensor satisfying the sum rule (II.1) corresponds to a unique massless

<sup>&</sup>lt;sup>4</sup> A tensor is defined to be traceless if the  $(SU(N_f)_L \times SU(N_f)_R)$ -invariant contraction of any upper index with any lower index identically vanishes. In general, any tensor can be decomposed as a direct sum of terms, each being a traceless tensor or the product of one (lower-rank) traceless tensor times a certain number of invariant tensors  $\delta_j^i$ (see for example Ref. [28], chapter 13).

bound state (although, similarly to the case of operators, the converse is not true). For a given theory with  $N_c$  colors and  $N_f$  flavors, the set of all possible traceless tensors that satisfy Eq. (II.1) forms a tensor space denoted as  $\mathcal{T}[N_f]$ :

Composite Operators ~ 
$$T^{\{n_L\};\{n_R\}}_{\{\bar{n}_L\};\{\bar{n}_R\}} \in \mathcal{T}[N_f]$$
. (II.2)

For convenience, we will denote by  $\mathcal{R}[N_f]$  the corresponding space of representations of  $\mathcal{G}[N_f]$ . When no confusion arises, we will use the shorthand notation

$$T_{\{\bar{n}\}}^{\{n\}} \equiv T_{\{\bar{n}_L\};\{\bar{n}_R\}}^{\{n_L\};\{n_R\}}, \qquad (\text{II.3})$$

where each of the symbols  $\{n\}$  and  $\{\bar{n}\}$  collectively denotes a pair of YTs with  $n = n_L + n_R$  and  $\bar{n} = \bar{n}_L + \bar{n}_R$ .

Indices of SU(N) can be raised (lowered) by means of the fully antisymmetric invariant tensor  $\epsilon^{i_1i_2\cdots i_N}$  ( $\epsilon_{i_1i_2\cdots i_N}$ ). To find the representation characterizing a given tensor one can proceed as follows [28]: for each kind of indices, starting from the (antiquark) YT { $\bar{n}$ } one finds the dual tableau  $\widetilde{\{\bar{n}\}}$  obtained by replacing each column of length  $\zeta$  with a column of length  $N_f - \zeta$  (columns of  $\widetilde{\{\bar{n}\}}$  must of course be arranged in reverse order compared to those in { $\bar{n}$ }); then, the representation to be found is that given by the 'combined' YT (CYT) obtained as the union of the columns of {n} and  $\widetilde{\{\bar{n}\}}$ . <sup>5</sup> For brevity, the CYT obtained from {n} and { $\bar{n}$ } will be denoted as

$$\{n;\bar{n}\} \equiv \{n\} + \{\bar{n}\} , \qquad (\text{II.4})$$

where the 'sum' on the right-hand side means adding together the columns of  $\{n\}$  and  $\{\overline{n}\}$  to make a new YT.

A massless bound state interpolated by a traceless tensor  $T_{\{\bar{n}_L\},\{\bar{n}_R\}}^{\{n_L\},\{\bar{n}_R\}}$  (i.e. by an operator corresponding to that tensor <sup>6</sup>) transforms as an irrep r of  $\mathcal{G}[N_f]$  characterized by a pair of CYT and baryon number b given by Eq. (II.1):

$$r = (\{n_L; \bar{n}_L\}, \{n_R; \bar{n}_R\}, b) \quad . \tag{II.5}$$

**Example 2.1.** Consider  $T_{\{0\};\{0\}}^{\{3_a\};\{0\}}$ , where  $\{3_a\} = \square$ . Such tensor exists for  $N_f \ge 3$  and satisfies the sum rule of Eq. (II.1) with b = 1 for  $N_c = 3$ . It interpolates a baryon transforming as  $r = (\square, s, 1)$ , where s stands for the singlet representation of  $SU(N_f)$ .

<sup>&</sup>lt;sup>5</sup> For example, the adjoint representation of  $SU(N_f)$  can be thought of as the traceless part of the tensor product of a fundamental and an anti-fundamental representation, which can be written as the tensor  $T_{\{1\}}^{\{1\}}$ . The CYT has two columns, one with  $N_f - 1$  boxes and one with a single box. According to the procedure outlined above, the CYT is obtained by first replacing  $\{1\}$  from the lower index by a column of  $N_f - 1$  boxes, and then joining it with the box from the upper index.

<sup>&</sup>lt;sup>6</sup> In the following, when we say that a tensor interpolates a bound state we mean that the bound state is interpolated by some local operator which corresponds to that tensor.

#### **B.** Equivalent Tensors

In general, different traceless tensors of  $\mathcal{T}(N_f)$  interpolate states in different representations of  $\mathcal{G}[N_f]$ . There are some important exceptions however, which we call equivalent tensors.

First of all, tensors interpolating the same representation of  $\mathcal{G}[N_f]$  can have YTs differing by the presence of one (or more) set of  $N_f$  fully antisymmetrized indices, i.e. flavor singlets

$$\epsilon_{i_1i_2\cdots i_{N_f}}q^{i_1}q^{i_2}\cdots q^{i_{N_f}} . \tag{II.6}$$

Tensors of this kind can still obey the sum rule (II.1) with the same b, and thus be equivalent under  $\mathcal{G}[N_f]$ , if such flavor singlets are located in different YTs.

**Example 2.2.** Consider  $T_{\{1\};\{0\}}^{\{N_f\};\{0\}}$  and  $T_{\{1\};\{0\}}^{\{0\};\{N_f\}}$ , for  $bN_c = N_f - 1$ , where  $\{N_f\}$  and  $\{1\}$  are respectively a YT with a single column of length  $N_f$  and a YT with one box. Both these tensors transform as an anti-fundamental of  $SU(N_f)_L$ , a singlet representation of  $SU(N_f)_R$ , and have baryon number  $b = (N_f - 1)/N_c$ .

There is another, less trivial, way in which equivalent tensors can arise. Notice that two different pairs of upper and lower YTs (for either left or right indices) can correspond to the same representation of  $SU(N_f)$ . Two tensors are equivalent under  $\mathcal{G}[N_f]$ , i.e. correspond to the same representation, if their CYTs for both left and right indices are identical and their baryon number is the same.

**Example 2.3.** Consider  $T_{\{0\};\{m\}}^{\{0\};\{n\}}$  and  $T_{\{0\};\{N_f-m\}}^{\{0\};\{N_f-m\}}$  with  $n, m < N_f$  and  $bN_c = n - m$ , where all the indices are fully antisymmetrized. These two equivalent tensors transform trivially under  $SU(N_f)_L$  and as an irrep of  $SU(N_f)_R$  characterized by a CYT with two columns, one with n boxes and one with  $N_f - m$  boxes. They have the same baryon number  $b = (n - m)/N_c$ . The case with m = 0 gives an example of a tensor with lower (antiquark) indices interpolating a baryonic bound state:  $T_{\{0\};\{N_f-n\}}^{\{0\};\{N_f\}}$  is equivalent to the purely baryonic operator  $T_{\{0\};\{0\}}^{\{0\};\{n\}}$ .

As it will become clear in the following, understanding the properties of equivalent tensors helps us to understand the validity of  $N_f$ -independence and of the superalgebra approach. It is useful to classify traceless tensors into three classes according to their baryon number b and their ranks  $\bar{n} = \bar{n}_L + \bar{n}_R$  and  $n = n_L + n_R$ , as depicted in Fig. 1. For fixed  $N_c$  and  $N_f$ , the three classes are defined as follows:

• Class A: tensors with  $n + \bar{n} = 2\bar{n} + bN_c < N_f$ .




FIG. 1. Different classes of tensors that interpolate massless bound states in a confining QCD-like theory with  $N_c$  colors and  $N_f$  flavors, under the assumption of unbroken chiral symmetry. Tensors are defined in terms of their baryon number b (c.f. Eq. (II.1)) and antiquark number  $\bar{n} = \bar{n}_L + \bar{n}_R$ .

- Class B: tensors with  $n = \bar{n} + bN_c \leq N_f \leq 2\bar{n} + bN_c = n + \bar{n}$ ; when  $n = N_f$  a flavor singlet can appear in the traceless part (i.e.  $\epsilon_{i_1i_2\cdots i_{N_f}}q^{i_1}q^{i_2}\cdots q^{i_{N_f}}$ ), otherwise they contain no flavor singlets; since  $\bar{n} < n$ , given that b > 0, no flavor singlets made of antiquark indices can appear.
- Class C: tensors with  $n = \bar{n} + bN_c > N_f$ .

The following result holds true:

Lemma 1 (Absence of Equivalent Tensors). Any two class-A tensors cannot transform in the same representations under  $\mathcal{G}[N_f]$ , i.e. they are not equivalent. Any two terms in their tensor decomposition under  $\mathcal{G}[N_f, 1]$  (cf. Eq. (I.2)) which have non-zero  $U(1)_{H_1}$  charge are not equivalent either.

The proof is given in Appendix **B**.

### **III. ANOMALY MATCHING AND PERSISTENT MASS CONDITIONS**

The main purpose of this section is to give a review on AMC and PMC (see also [29]). Some important properties of the PMC overlooked in the previous literature will be also discussed.

In order to write the AMC and PMC, we first introduce the index  $\ell(r)$  to denote the multiplicity of the (massless) bound states in the irrep r, defined as the number of times these appear in the spectrum with helicity +1/2 minus the number of times they appear with helicity -1/2 [1]:

$$\ell(r) \equiv \mathcal{N}(r)_{+1/2} - \mathcal{N}(r)_{-1/2} .$$
 (III.1)

Since  $\mathcal{N}(r)_{\pm 1/2}$  are positive integers, the indices  $\ell$  are also integers. The invariance of the theory under CPT implies that

$$\mathcal{N}(r)_{-1/2} = \mathcal{N}(r^*)_{+1/2}$$
 (III.2)

Here  $r^*$  is the complex conjugate of r, in particular it has opposite baryon number. This property allows us to consider only bound states with positive baryon number without loss of generality: the multiplicity of a massless bound state with negative baryon number is equal and opposite to the multiplicity of its antiparticle, whose baryon number is positive.

Parity is unbroken in the confining phase of vectorlike gauge theories with massive fermions and  $\theta = 0$  [30]. Continuity implies that the parity-preserving vacuum is a global minimum also in the limit of vanishing quark masses, although degenerate parity-breaking minima might appear in principle. If the vacuum preserves parity, then it follows, for any  $r = (r_L, r_R, b)$ , that

$$\mathcal{N}(r)_{-1/2} = \mathcal{N}(r_P)_{+1/2} ,$$
 (III.3)

where  $r_P = (r_R, r_L, b)$  is the parity conjugate of r. Notice however that our results do not rely on the assumption of unbroken parity.

### A. Anomaly Matching Conditions

There are four 't Hooft anomalies that must be matched in the IR by composite massless fermions: those of  $[SU(N_f)_L]^2 U(1)_B$ ,  $[SU(N_f)_L]^3$ ,  $[SU(N_f)_R]^2 U(1)_B$  and  $[SU(N_f)_R]^3$ . <sup>7</sup> The AMC of  $[SU(N_f)_L]^2 U(1)_B$  and  $[SU(N_f)_L]^3$  can be written explicitly as

$$\sum_{r \in \mathcal{R}[N_f]} \ell(r) A_i(r) = N_c A_i(r_{q_L}), \qquad (\text{III.4})$$

where  $A_i(r)$  is the anomaly contribution of the representation r of  $\mathcal{G}[N_f]$ , and where i = 2 for  $[SU(N_f)_L]^2 U(1)_B$  and i = 3 for  $[SU(N_f)_L]^3$ . The left-hand side of this equation encodes the anomaly of the bound states, expressed as a sum over the contributions of all possible representations r in the spectrum multiplied by their index  $\ell$ ; the right-hand side encodes instead the anomaly

<sup>&</sup>lt;sup>7</sup> If parity is spontaneously broken by the vacuum, the spectrum of massless bound states charged under  $U(1)_B$  can be chiral and the  $[U(1)_B]^3$  anomaly must be also matched. In the following, we will not make use of such additional anomaly matching equation, as we want our approach to be as general as possible.

12

of the left-handed quarks  $q_L$ , which transform in the representation  $r_{q_L} = (\Box, s, 1/N_c)$ . Here and in the following we assume  $N_f > 2$ . For  $N_f = 2$ , since all the representations of SU(2) are real or pseudo-real, there is only one AMC, that of  $[SU(2)_L]^2 U(1)_B$ , which has integer solutions [1, 12].

The anomaly of a representation  $r = (r_L, r_R, b)$  can be computed as

$$A_i(r) = d(r_R) D_i(r_L) , \qquad (\text{III.5})$$

where  $d(r_R)$  is the dimension of  $r_R$ , and  $D_{i=2,3}(r_L)$  are the anomaly traces of respectively  $[SU(N_f)_L]^2 U(1)_B$  and  $[SU(N_f)_L]^3$ , see Appendix A for their definitions. Both quantities depend on  $N_f$  explicitly. For the quarks, in particular,  $A_i(r_{q_L}) = D_i(\Box)$ . By conveniently defining the following ratios of anomaly traces

$$\bar{D}_i(r_L) \equiv \frac{D_i(r_L)}{D_i(\Box)} , \qquad i = 2,3$$
(III.6)

the AMC can be recast into the form

$$\sum_{r \in \mathcal{R}[N_f]} \ell(r) \, d(r_R) \, \bar{D}_i(r_L) = N_c \quad . \tag{III.7}$$

The AMC for  $[SU(N_f)_R]^2 U(1)_B$  and  $[SU(N_f)_R]^3$  can be derived in a similar way. One thus has a system of four linear, non-homogeneous equations for the variables  $\ell$ , that will be denoted as  $AMC[N_f]$  in the following to stress its dependence on the number of flavors  $N_f$ .

If  $N_c$  is even, any color-singlet bound state is necessarily a boson. This directly implies that 't Hooft anomalies cannot be matched unless chiral symmetry is spontaneously broken and massless Nambu-Goldstone bosons are present in the spectrum. In theories with  $N_c$  odd, proving chiral symmetry breaking is not equally easy. It is possible to show that  $AMC[N_f]$  do not have integer solutions when  $N_f$  is an integral multiple of a prime factor p > 1 of  $N_c$ , see Ref. [7]. For such special values of  $N_f$  chiral symmetry breaking is therefore proven. A first application of this argument was made by Preskill and Weinberg in Ref. [5] in the context of three-flavor QCD and for the case of baryons with b = 1, see also [29]. In the following, we will discuss the case of theories with  $N_c$ odd and  $N_f$  generic.

### **B.** Persistent Mass Conditions

In its original formulation, the Persistent Mass Condition asserts that any bound state containing massive constituents must also be massive [5]. It is however possible to give the following more precise statement that does not make reference to internal constituents of the bound states and relies instead only on their representation under the chiral symmetry group. <sup>8</sup>

When one flavor is given a mass, each massless fermionic bound state transforming as an irrep of  $\mathcal{G}[N_f]$  breaks up into irreps of the unbroken flavor group  $\mathcal{G}[N_f, 1]$ . The Persistent Mass Condition requires that, after such decomposition, the bound states with nonzero  $U(1)_{H_1}$  charge must be massive, i.e. come in vectorlike pairs with vanishing index. More in general, when *i* flavors are given unequal masses, all the bound states that are charged under  $U(1)_{H_1} \times \cdots \times U(1)_{H_i} \subset \mathcal{G}[N_f, i]$  must be massive, i.e. have vanishing index.

As formulated above, the Persistent Mass Condition can be proven by using the arguments articulated by Vafa and Witten in Ref. [6]. Let us analyze the case of one massive flavor, the generalization to additional massive flavors being straightforward. We consider a regularized version of our theory with finite cutoff  $\Lambda$ . <sup>9</sup> Let m be the bare mass of the massive flavor and  $\epsilon$  that of all the remaining flavors. At the end we will take the limit  $\epsilon \to 0$  and then remove the cutoff. We stress that m and  $\epsilon$  are bare and not renormalized masses. Consider a local operator T(x)that interpolates bound states with  $U(1)_{H_1}$  charge  $H_1 \neq 0$ . The results of Ref. [6] imply that the two-point function of T(x) satisfies the bound

$$|\langle T^{\dagger}(x)T(y)\rangle| \lesssim C e^{-(Qm+q\epsilon)|x-y|}$$
(III.8)

where  $Q = |H_1|$ ,  $q = |bN_c - H_1|$  and C is a prefactor that does not depend on x and y. Such exponential fall-off means that any bound state excited by T(x) has mass  $M \ge Qm + q\epsilon$ , where of course M depends on  $\Lambda$ . If we now let  $\epsilon \to 0$  we obtain the lower bound  $M \ge |H_1|m > 0$ . This result holds in the regularized theory with finite cutoff. It states that for any finite value of  $\Lambda$ , the spectrum of the bound states excited by T(x) must be vectorlike, i.e. the corresponding index is 0. Since the theory is defined as the limit for  $\Lambda \to \infty$  of the regularized one, this means that the index will vanish also after the cutoff is removed. <sup>10</sup> This proves the PMC.

<sup>&</sup>lt;sup>8</sup> Although intuitive, the notion of internal constituents can be made precise only in the framework of a given phenomenological model for the bound states and, as such, it is not convenient.

<sup>&</sup>lt;sup>9</sup> Our argument, as well as those of [6], is formulated for a theory on the continuum. One might want to resort to lattice regularization for a fully rigorous definition of the theory. Notice that the assumptions on which the Vafa-Witten theorem relies are fullfilled only on specific lattice realizations, like that given by staggered fermions. See for example [31–33].

<sup>&</sup>lt;sup>10</sup> If one uses the renormalization group to resum the leading logs, one finds that the bare mass m goes to zero if the limit  $\Lambda \to \infty$  is taken while keeping the renormalized mass fixed and finite. We thank M. Bochicchio for pointing this to us. While this seems to nullify the bound on M, the spectrum of bound states excited by T(x) remains vectorlike and this is sufficient to prove that the index is zero after removing the cutoff.

Notice that it was crucial to assume  $H_1 \neq 0$  for the operator T(x). No firm statement can be made on bound states with  $H_1 = 0$ , even though they can be interpolated by operators featuring massive quarks and antiquarks. In fact, since  $H_1 = 0$ , there will necessarily exist at least one operator without heavy flavors that can interpolate those states. This also means that decoupling cannot be used to forbid massless states with  $H_1 = 0$ . Indeed, decoupling is just the statement that one can reproduce physical results at low energy by means of an effective theory defined in terms of degrees of freedom with  $H_1 = 0$ . To the best of our knowledge, the proof of PMC and the importance of the condition  $H_1 \neq 0$  were not explicitly discussed in the previous literature. Going beyond a naive formulation of PMC is especially required when dealing with exotic states.

On the mathematical level, the Persistent Mass Condition implies a series of equations which must be satisfied by the indices  $\ell$ . To discuss this, it is useful to introduce the following notation: we will denote by

- $\mathcal{R}_0[N_f, i]$  the space of irreps of  $\mathcal{G}[N_f, i]$  with all vanishing charges  $(H_k = 0 \text{ with } k = 1, \dots, i);$
- $\hat{\mathcal{R}}[N_f, i]$  the space of irreps of  $\mathcal{G}[N_f, i]$  with  $H_i \neq 0$  and  $H_k = 0$  for  $k = 1, \ldots, i 1$ .

In the case of one massive flavor, the index of any irrep  $\hat{r}_1$  of  $\mathcal{G}[N_f, 1]$  with  $H_1 \neq 0$  must vanish:

$$0 = \ell(\hat{r}_1) = \sum_{r \in \mathcal{R}[N_f]} k(r \to \hat{r}_1) \, \ell(r) \,, \qquad \forall \; \hat{r}_1 \in \hat{\mathcal{R}}[N_f, 1] \,. \tag{III.9}$$

The coefficient  $k(r \to \hat{r}_1)$  equals the number of times  $\hat{r}_1$  appears in the decomposition of r. Let  $\hat{r}_1 = (\{m_L; \bar{m}_L\}, \{m_R; \bar{m}_R\}, H_1, b)$ . It is easy to see that

$$H_1 = bN_c - (m_L + m_R) + (\bar{m}_L + \bar{m}_R), \qquad (\text{III.10})$$

while the baryon number b is the same as that of the parent irrep r. The system of all PMC (one equation for each irrep  $\hat{r}_1 \in \hat{\mathcal{R}}[N_f, 1]$ ) that are obtained by giving one of the  $N_f$  flavors a mass will be denoted as  $PMC[N_f, 1]$  in the following. Notice that  $PMC[N_f, 1]$  depend implicitly on  $N_f$ , i.e. in general they have different expressions in confining theories with different number of flavors. This point will be elaborated further in Section. IV A.

Additional PMC arise when more than one flavor is given a mass. For example, when a second flavor becomes massive, those states which had  $H_1 = 0$  in the first step will now break up into multiplets of the unbroken flavor group  $\mathcal{G}[N_f, 2]$ , cf. Eq. (I.2). After this second step of decomposition, the states which are charged nontrivially under  $U(1)_{H_2}$  are subject to  $PMC[N_f, 2]$ , i.e. equations of the form

$$0 = \ell(\hat{r}_2) = \sum_{r_1 \in \mathcal{R}_0[N_f, 1]} k(r_1 \to \hat{r}_2) \,\ell(r_1) \ , \qquad \forall \ \hat{r}_2 \in \hat{\mathcal{R}}[N_f, 2] \,, \tag{III.11}$$

where the sum runs over all irreps  $r_1$  of  $\mathcal{G}[N_f, 1]$  with  $H_1 = 0$ . Let  $\hat{r}_2 = (\hat{r}_{2L}, \hat{r}_{2R}, H_1 = 0, H_2, b)$ , with  $\hat{r}_{2L} = \{m'_L; \bar{m}'_L\}, \hat{r}_{2R} = \{m'_R; \bar{m}'_R\}$ ; then it follows that

$$H_2 = bN_c - (m'_L + m'_R) + (\bar{m}'_L + \bar{m}'_R).$$
(III.12)

Notice that  $PMC[N_f, 2]$  also depend implicitly on  $N_f$ . More in general, in a theory with  $N_f$  flavors there exist  $N_f - 2$  sets of PMC, denoted by  $PMC[N_f, i]$  with  $1 \le i \le N_f - 2$ , obtained by considering all possible numbers of massive flavors and giving them unequal masses. Giving the same mass to different flavors does not lead to new equations, as shown in Appendix C. Since there are no PMC in a theory with 2 flavors, the largest number of flavors that can be made massive to obtain new equations is  $N_f - 2$ . <sup>11</sup> The collection of all the PMC in a theory with  $N_f$  flavors will be denoted by  $PMC[N_f]$  in the following. To the best of our knowledge, previous works only considered PMC with one massive flavor [1, 2, 8–13]; these are denoted as  $PMC[N_f, 1]$  in this paper. On the other hand, PMC with more than one massive flavors play an important role in the proof of chiral symmetry breaking (see Ref. [7]).

It is useful to notice that  $PMC[N_f, 2]$  are related to  $PMC[N_f - 1, 1]$ . Indeed, each of the bound states that remain massless after giving one flavor a mass is in one-to-one correspondence with a massless bound state of a theory with  $N_f - 1$  flavors. As a consequence,  $PMC[N_f, 2]$  have the same form as  $PMC[N_f - 1, 1]$ . Indeed, the latter are written as

$$0 = \ell(\hat{r}'_1) = \sum_{r' \in \mathcal{R}[N_f - 1]} k(r' \to \hat{r}'_1) \ell(r') , \qquad \forall \; \hat{r}'_1 \in \hat{\mathcal{R}}[N_f - 1, 1], \qquad (\text{III.13})$$

which is identical to Eq. (III.11) provided one identifies r' with  $r_1$  and sets  $\ell(r') = \ell(r_1)$ . The identification is possible because  $\mathcal{R}_0[N_f, 1] = \mathcal{R}[N_f - 1]$ , i.e. any irrep of  $\mathcal{G}[N_f, 1]$  with  $H_1 = 0$  can be seen as an irrep of  $\mathcal{G}[N_f - 1]$ . More in general, it is simple to see that  $\text{PMC}[N_f, i]$  have the same form as  $\text{PMC}[N_f - 1, i - 1]$ , for  $1 < i \leq N_f - 2$ . The general structure of PMC when  $N_f$  is varied is therefore the one sketched in Fig. 2.

### IV. PROOF OF CHIRAL SYMMETRY BREAKING FROM N<sub>f</sub>-INDEPENDENCE

The AMC[ $N_f$ ] and PMC[ $N_f$ ] give a linear system of equations on the indices  $\ell$ . As first suggested by 't Hooft [1], it is useful to consider such system and its solutions as functions of  $N_f$ . This

<sup>&</sup>lt;sup>11</sup> If  $N_f = 2$  and one flavor is given a mass, the unbroken group  $\mathcal{G}[2, 1] = U(1)_H \times U(1)_B$  is vectorlike. If parity is unbroken in the vacuum, this means that all the indices of irreps of  $\mathcal{G}[2, 1]$  vanish automatically. On the other hand, if parity is broken by the vacuum, then non-trivial PMC equations arise. In the following, we will not make use of such additional PMC equations for  $N_f = 2$ , as we want our approach to be as general as possible.




FIG. 2. General structure of persistent mass conditions  $PMC[N_f]$  for different values of  $N_f$ . The set of PMC obtained by giving different masses to *i* flavors, with  $1 \le i \le N_f - 2$ , is denoted as  $PMC[N_f, i]$ . Sets of PMC equations connected by a line have the same form.

is independent of whether the theory with  $N_f$  flavors confines or not; one can indeed consider Eq. (III.4) as a rule to construct the AMC[ $N_f$ ] for any  $N_f$ , and similarly Eqs. (III.9) and (III.11) as rules to construct the PMC[ $N_f$ ]. Under particular assumptions that will be discussed in detail, the following property then holds and is key to prove chiral symmetry breaking:

 $N_f$ -independence: Any finite set of real indices  $\{\ell\}$  that solves the system of AMC and PMC for a confining theory with  $N_f$  flavors is also a solution of the same equations for any  $N'_f \ge N_f$ .

In order to properly state  $N_f$ -independence one should better qualify what it exactly means that the same set of indices  $\{\ell\}$  is a solution of AMC and PMC for different numbers of flavors. As we will show, one needs to establish a map between irreps of the flavor groups  $\mathcal{G}[N_f]$  and  $\mathcal{G}[N'_f]$ , i.e. a map between states of different theories. For the moment we will content ourselves with the naive definition given above and explore its consequences.

The validity of  $N_f$ -independence implies that chiral symmetry must be broken in the theory with  $N_f$  flavors. There are at least two ways to see this. The first makes use of the following simple argument: let us suppose that for  $N_f$  flavors and  $N_c$  colors the theory confines and chiral symmetry is not spontaneously broken by the vacuum. This requires that  $AMC[N_f]$  have integer solutions. By  $N_f$ -independence, this in turn implies that  $AMC[N'_f]$  must also have integer solutions for any  $N'_f \geq N_f$ . However, we know that  $AMC[p\,m]$  do not admit integer solutions for any positive integer m, where p is any prime factor of  $N_c$ . This contradicts the initial assumption and thus proves (as long as  $N_f$ -independence is valid) that chiral symmetry is spontaneously broken in the theory with  $N_f$  flavors. Notice that it is not necessary to assume that the theory with  $N_c$  colors and  $N'_f > N_f$  flavors confines.

A second way to see that chiral symmetry breaking is implied by  $N_f$ -independence is by means of the following argument. The anomaly coefficients  $A_i(r)$  have polynomial dependence on  $N_f$ . Therefore, as long as  $N_f$ -independence holds, the AMC can be recast, for any  $N'_f \ge N_f$ , as

$$a_h(\{\ell\}) (N'_f)^h + a_{h-1}(\{\ell\}) (N'_f)^{h-1} + \dots + a_1(\{\ell\}) N'_f + a_0(\{\ell\}) = 0 , \qquad (IV.1)$$

where the coefficients  $a_i(\{\ell\})$  depend on the indices  $\ell$ . In other words, the polynom in  $N'_f$  of Eq. (IV.1) generates the AMC for any  $N'_f \geq N_f$ . Tensors with n upper and  $\bar{n}$  lower indices contribute to the AMC with terms of order  $n + \bar{n} - 1$  or smaller in  $N_f$ . The highest degree of the polynom, h, is thus finite if the spectrum of representations, hence of interpolating tensor operators, is finite:

$$h = n_{max} + \bar{n}_{max} - 1 = bN_c + 2\bar{n}_{max} - 1 .$$
 (IV.2)

 $N_f$ -independence implies that Eq. (IV.1) has a solution for every integer  $N'_f \ge N_f$  once its coefficients are evaluated on the set of indices  $\{\ell\}$  which solves  $AMC[N_f] \cup PMC[N_f]$ . But a polynom can have a number of solutions bigger than its degree only if it vanishes identically. This implies that its coefficients must individually vanish:

$$a_i(\{\ell\}) = 0, \quad \text{for all} \quad 0 \le i \le h \ . \tag{IV.3}$$

As pointed out in Ref. [9] by Farrar, the equation  $a_0(\{\ell\}) = 0$  does not admit integral solutions. Hence, no integral solution of the AMC[ $N_f$ ] exists, and this means that the assumption of unbroken chiral symmetry is not consistent: chiral symmetry must be spontaneously broken in the theory with  $N_f$  flavors.

### A. Proof of $N_f$ -independence

In the past literature  $N_f$ -independence was assumed to hold based on empirical arguments. Given its centrality to prove chiral symmetry breaking, we want to derive a rigorous proof and spell out explicitly the assumptions which underlie its validity. Following Refs. [2, 10, 11], we can first check whether  $PMC[N'_f]$  can be solved, for any  $N'_f \geq N_f$ , by the same indices that solve  $PMC[N_f]$ , and then check whether a solution of  $AMC[N_f]$  is also a solution of  $AMC[N'_f]$ , up to equations implied by  $PMC[N'_f]$ . If the solution of  $AMC[N_f] \cup PMC[N_f]$  is also a solution of  $AMC[N'_f] \cup PMC[N'_f]$ , then  $N_f$ -independence follows. We find that this is possible only for a spectrum of massless bound states that are interpolated by class-A tensors.

# 1. $PMC[N_f]$ are $N_f$ -dependent in general

In order to see if a real solution of  $PMC[N_f]$  is also a real solution of  $PMC[N'_f]$  for  $N'_f > N_f$  we start by considering  $N'_f = N_f + 1$  and then proceed iteratively. The relation between  $PMC[N_f]$  and  $PMC[N_f + 1]$  is clarified by recalling that  $PMC[N_f + 1, i]$  have the same form as  $PMC[N_f, i - 1]$ , for  $1 < i \leq N_f - 1$ , see Section IIIB. This means that  $PMC[N_f + 1]$  contain  $PMC[N_f]$  plus the additional set of equations  $PMC[N_f + 1, 1]$ . It is thus crucial to understand whether the latter are truly linearly-independent equations. We will now show that  $PMC[N_f + 1, 1]$  are identical to  $PMC[N_f, 1]$  under specific conditions.

The form of  $PMC[N_f, 1]$  is in general different in theories with different numbers of flavors because of the following three reasons:

- 1. New tensor structures arise for  $N'_f > N_f$ . In general, such new tensor structures imply new irreps in the theory with  $N'_f$  flavors. For example, tensors with  $bN_c$  fully antisymmetrized upper indices exist only for  $N_f \ge bN_c$  but not for  $N_f < bN_c$ . Indeed, the  $bN_c$  boxes can only be arranged such that the YT has at most  $N_f$  rows. Clearly, there are more irreps allowed for  $N_f \ge bN_c$  than for  $N_f < bN_c$ . This point was emphasized in Ref. [12] for a spectrum of massless baryons with b = 1.
- 2. Different tensors of  $\mathcal{G}[N_f]$  can become equivalent or cease to be so when the number of flavors is increased. In other words, two different tensors can interpolate the same irrep of  $\mathcal{G}[N'_f]$ but different irreps of  $\mathcal{G}[N_f]$ , or viceversa.
- 3. Different tensors of  $\mathcal{G}[N_f, 1]$  can become equivalent or cease to be so when the number of flavors is increased. Accordingly, a single PMC equation splits into different PMC equations as tensors become inequivalent, while different PMC equations collapse into a single equation when tensors become equivalent.

The last point is best clarified by looking at an explicit example.

**Example 4.1**. Let us consider a theory with  $N_c = 3$  and the following two tensors

$$T_1 = T_{\{1\};\{0\}}^{\{1\};\{3_s\}}, \qquad T_2 = T_{\{2_a\};\{0\}}^{\{3_a\};\{2_s\}}, \qquad (IV.4)$$

where  $\{3_s\} = \Box \Box$ ,  $\{3_a\} = \Box$ ,  $\{2_s\} = \Box \Box$ ,  $\{2_a\} = \Box$ . The decomposition of  $T_1$  and  $T_2$  in tensors of  $\mathcal{G}[N_f, 1]$  contains the following two terms, both with  $H_1 = 3$ :

$$T_{1} = T'_{1} + \cdots , \quad T'_{1} = T^{\{1\};\{0\}}_{\{1\};\{0\}}$$

$$T_{2} = T'_{2} + \cdots , \quad T'_{2} = T^{\{2_{a}\};\{0\}}_{\{2_{a}\};\{0\}}$$
(IV.5)

where the dots indicate other terms in the decomposition. For  $N_f = 4$ ,  $T_1$  and  $T_2$  transform respectively as the irreps

$$r_1 = \left( \square, \square, 1 \right), \quad r_2 = \left( \square, \square, 1 \right)$$
(IV.6)

of  $\mathcal{G}[4]$  and are thus inequivalent. The tensors  $T'_1$  and  $T'_2$ , on the other hand, both transform as the representation ( $\square$ , s, 3, 1) of  $\mathcal{G}[4, 1]$  and are thus equivalent. The PMC[4, 1] equation corresponding to that representation reads

$$0 = \ell(r_1) + \ell(r_2) + \cdots$$
 (IV.7)

where the dots indicate contributions from operators other than  $T_1$  and  $T_2$ . For  $N_f = 5$ ,  $T_1$  and  $T_2$  transform as the following irreps

$$r_1 = \left( \square, \square, 1 \right), \quad r_2 = \left( \square, \square, 1 \right)$$
 (IV.8)

of  $\mathcal{G}[5]$  while  $T'_1$  and  $T'_2$  transform respectively as

$$\left(\begin{array}{c} \hline \\ \hline \\ \end{array}, s, 3, 1\right)$$
 and  $\left(\begin{array}{c} \hline \\ \hline \\ \end{array}, s, 3, 1\right)$ . (IV.9)

The latter are different irreps of  $\mathcal{G}[5, 1]$ , which means that  $T'_1$  and  $T'_2$  are inequivalent for  $N_f = 5$ . As a result, Eq. (IV.7) splits into two equations of PMC[5, 1]:

$$0 = \ell(r_1) + \cdots, \quad 0 = \ell(r_2) + \cdots.$$
 (IV.10)

In a similar fashion, one can construct examples where, as  $N_f$  is increased, different PMC equations collapse into a single PMC equation because tensors of  $\mathcal{G}[N_f, 1]$  become equivalent.

# 2. Condition for $N_f$ -independence of $PMC[N_f, 1]$

We have shown that the form of  $PMC[N_f, 1]$  is  $N_f$ -dependent in general, which implies that  $PMC[N_f+1, 1]$  are linearly independent with respect to  $PMC[N_f, 1]$ . This is in turn an obstruction to the  $N_f$ -independence of the solution of PMC. However,  $N_f$ -independence can be still obtained if

one imposes suitable restrictions on the spectrum of possible massless states. In order to determine such restrictions, we introduce the notions of uplifting of tensors.

We define *uplifting* as a map between traceless tensors of  $\mathcal{T}[N_f]$  and  $\mathcal{T}[N_f + 1]$ . Given any tensor  $T \in \mathcal{T}[N_f]$ , the uplifted one is defined as the tensor in  $\mathcal{T}[N_f + 1]$  which is identical to T, but with indices running from 1 to  $N_f + 1$ :

$$T^{\{n_L\};\{n_R\}}_{\{\bar{n}_L\};\{\bar{n}_R\}} \in \mathcal{T}[N_f] \to T^{\{n_L\};\{n_R\}}_{\{\bar{n}_L\};\{\bar{n}_R\}} \in \mathcal{T}[N_f+1].$$
(IV.11)

We notice that:

- For any tensor  $T_{\{\bar{n}_L\};\{\bar{n}_R\}}^{\{n_L\};\{\bar{n}_R\}} \in \mathcal{T}[N_f]$  there exists a corresponding well defined tensor  $T_{\{\bar{n}_L\};\{\bar{n}_R\}}^{\{n_L\};\{\bar{n}_R\}} \in \mathcal{T}[N_f+1]$ . This is because the YTs  $\{n_L\}, \{n_R\}, \{\bar{n}_L\}$  and  $\{\bar{n}_R\}$  have at most  $N_f$  rows, hence are well defined for a theory with  $N_f + 1$  flavors as well.
- Uplifting is an injective map: Any two different tensors of  $\mathcal{T}[N_f]$  are mapped to different tensors of  $\mathcal{T}[N_f + 1]$ .
- Uplifting is not surjective, since there are tensors of  $\mathcal{T}[N_f + 1]$  which cannot be obtained from the uplifting of any tensor of  $\mathcal{T}[N_f]$ . In particular, uplifting cannot reach tensors that have  $N_f + 1$  fully antisymmetrized upper (or lower) indices forming a singlet, since such configuration does not exist for tensors of  $\mathcal{T}[N_f]$ .

The cartoon of Fig. 3 summarizes the properties of tensor uplifting.

Uplifting has been defined as a map between tensors, but it can be used to induce a corresponding map between representations of  $\mathcal{G}[N_f]$  and  $\mathcal{G}[N_f + 1]$ . In order to do so, one has to fix the redundancy implied by the tensor notation and choose one representative tensor for each representation. A potential obstacle to a meaningful definition is however posed by the existence of equivalent tensors. It may happen, indeed, that two different tensors that are equivalent in a theory with  $N_f$  flavors become inequivalent when the number of flavors is increased to  $N_f + 1$ ; in this case the same irrep of  $\mathcal{G}[N_f]$  could be mapped to two different irreps of  $\mathcal{G}[N_f + 1]$ , which is not acceptable. We know, on the other hand, that two class-A tensors cannot be equivalent, see Lemma 1 at the end of Section II. If we thus consider only representations of  $\mathcal{G}[N_f]$  and  $\mathcal{G}[N_f + 1]$ that are interpolated by class-A tensors, then a meaninful definition of uplifting is possible also for irreps. More explicitly: given any representation r interpolated by a class-A tensor of  $\mathcal{T}[N_f]$ , the uplifted representation U[r] is defined to be the one interpolated by the class-A tensor of  $\mathcal{T}[N_f + 1]$ which is identical to T. Since Lemma 1 ensures that two class-A tensors cannot be equivalent, there




FIG. 3. This cartoon tries to capture some of the properties of tensor uplifting. The two planes represent the space of tensors  $\mathcal{T}(N_f)$  and  $\mathcal{T}(N_f + 1)$ , the dots (whose color corresponds to a particular property) being the individual tensors. The uplifting map is represented by the upward black arrows, while the curved gray lines represent the equivalence relation between equivalent tensors. The green box in the lower plane represents the subset of class-A tensors in  $\mathcal{T}(N_f)$ ; its image through uplifting is represented by the green box in  $\mathcal{T}(N_f+1)$ . The latter is a proper subset of the set of class-A tensors in  $\mathcal{T}(N_f+1)$ , pictured by the red box. In this cartoon one can notice some of the properties explained in the text, in particular: *i*) two class-A tensors cannot be equivalent; *ii*) outside class-A tensors, the equivalence classes are not respected under uplifting (see the yellow dots); *iii*) the image of  $\mathcal{T}(N_f)$  through uplifting does not coincide with  $\mathcal{T}(N_f+1)$ (the black dot is not the uplifted of any tensor in  $\mathcal{T}(N_f)$ ).

exists at most one class-A tensor interpolating any given representation, i.e. any given state. <sup>12</sup> This means that the above definition is unique and gives an injective mapping.

**Example 4.2**. Let us consider a theory with  $N_c = 3$  and the following two tensors

$$T_a = T_{\{0\};\{0\}}^{\{3\};\{0\}}, \qquad T_b = T_{\{1\};\{0\}}^{\{4\};\{0\}}$$
(IV.12)

where {3} and {4} are YT with a single column with respectively 3 and 4 rows. For  $N_f = 4$ ,  $T_a$  is class A and  $T_b$  is class B; they transform under the same representation of  $\mathcal{G}[4]$ ,

$$r_a = \left( \bigsqcup, s, 1 \right) = \left( \bigsqcup, s, 1 \right) = r_b, \qquad (IV.13)$$

<sup>&</sup>lt;sup>12</sup> Class-A tensors interpolating a baryon will necessarily have  $\bar{n} = 0$ , while exotics will be interpolated by class-A tensors with  $\bar{n} \neq 0$ .

and are therefore equivalent. For  $N_f = 5$ ,  $T_a$  and  $T_b$  are still respectively of class A and B, but are now inequivalent since

$$r_a = \left(\square, s, 1\right) \neq \left(\square, s, 1\right) = r_b,$$
 (IV.14)

Establishing an uplifting map between the representations of  $N_f = 4$  and  $N_f = 5$  theories is thus possible only by focusing on representations interpolated by class-A tensors, i.e. by considering  $r_a$ and discarding  $r_b$ .

Notice that although every class-A irrep in  $\mathcal{R}[N_f]$  can be uplifted to a class-A irrep in  $\mathcal{R}[N_f+1]$ , the converse is not true.<sup>13</sup> In particular, irreps in  $\mathcal{R}[N_f+1]$  that are interpolated by tensors with  $n + \bar{n} = N_f$  are not the uplift of any class-A irrep in  $\mathcal{R}[N_f]$ .

Taking one step further, the map between representations corresponds to a map between the physical massless states of a theory with  $N_f$  flavors and those of a theory with  $N_f + 1$  flavors, provided that the latter theory is confining. Such last step is however not necessary to establish  $N_f$ -independence, since  $PMC[N_f]$  can be considered as mathematical equations rather than physical conditions.

Uplifting can be used to establish  $N_f$ -independence of  $PMC[N_f, 1]$ . It is easy to see that  $N_f$ independence holds if it is possible to establish an injective map between representations of  $\mathcal{G}[N_f]$ and  $\mathcal{G}[N_f + 1]$ , as well as between terms in their decompositions when one flavor is given a mass. The definition of uplifting given above has such property, because Lemma 1 ensures that within
class A no equivalent tensors exist of  $\mathcal{G}[N_f]$  and of  $\mathcal{G}[N_f, 1]$  as well. More in detail, one can impose
the following restriction on the spectrum of massless bound states (i.e. on the set of possible
representations): only states (representations) that can be interpolated by a class-A tensor are
assumed to have non-vanishing index. Given a spectrum  $\{\ell(r)\}$  fullfilling this hypothesis, with  $r \in \mathcal{R}[N_f]$ , it is possibile to define an uplifted spectrum of representations  $r' \in \mathcal{R}[N'_f]$  as follows:

$$\{\ell(r')\} = U^{(N'_f - N_f)}[\{\ell(r)\}]: \qquad \ell(r') = \begin{cases} \ell(r) & \text{if } r' = U^{(N'_f - N_f)}[r] \\ 0 & \text{otherwise} \end{cases}, \qquad (IV.15)$$

where  $U^{(N'_f - N_f)}[r]$  denotes the irrep of  $\mathcal{R}[N'_f]$  obtained by uplifting  $(N'_f - N_f)$  times the irrep r of  $\mathcal{R}[N_f]$ . One thus arrives at the following

**Lemma 2.** Let  $\{\ell(r)\}$  be a real solution of  $PMC[N_f, 1]$  such that the index of any state that cannot be interpolated by a class-A tensor vanishes. Then  $U^{(N'_f - N_f)}[\{\ell(r)\}]$  is a real solution of

<sup>&</sup>lt;sup>13</sup> Here and in the following, by class-A irrep we mean an irrep that is interpolated by a class-A tensor.

The proof follows easily from the definition of uplifting and the fact that it is an injective map.

### 3. Condition for $N_f$ -independence of $AMC[N_f]$

We now turn to the  $AMC[N_f]$ . These equations have an explicit dependence on  $N_f$  through the anomaly coefficients as well as an implicit one, since increasing the number of flavors gives rise to new tensor structures, hence new irreps. As for the PMC, one can consider Eq. (III.7) as a rule to generate a set of  $AMC[N_f]$  for any  $N_f$ , independently of the IR behavior of the corresponding theory. Then the following result holds true:

**Theorem 1** ( $N_f$  independence). Let  $\{\ell(r)\}$  be a real solution of  $AMC[N_f] \cup PMC[N_f, 1]$ such that the index of any state that cannot be interpolated by a class-A tensor vanishes. Then  $U^{(N'_f - N_f)}[\{\ell(r)\}]$  is a real solution of  $AMC[N'_f] \cup PMC[N'_f, 1]$  for any  $N'_f \ge N_f$ .

The proof is most easily obtained by induction. Let us start by considering the AMC for  $N_f + 1$  flavors,

$$N_c = \sum_{r' \in \mathcal{R}[N_f+1]} \ell(r') d(r'_R) \bar{D}_i(r'_L) , \qquad (IV.16)$$

which we want to prove to hold for  $\ell(r')$  defined in terms of  $\ell(r)$  as specified by the definition of uplifted solution of Eq. (IV.15):  $\{\ell(r')\} = U[\{\ell(r)\}]$ . In particular, the sum in Eq. (IV.16) runs over the representations  $r' = (r'_L, r'_R, b)$  of  $\mathcal{G}[N_f + 1]$  that are the uplift of class-A representations  $r = (r_L, r_R, b)$  of  $\mathcal{G}[N_f]$ . This means that any r' itself is interpolated by a class-A tensor of  $\mathcal{T}[N_f + 1]$ . It is possible to decompose the dimension d and the anomaly coefficient  $\overline{D}_i$  in terms of those of the CYT of  $SU(N_f)$ ; one has:

$$d(r'_R) = \sum_{r'_{1R}} k(r'_R \to r'_{1R}) \, d(r'_{1R}) \,, \tag{IV.17}$$

$$\bar{D}_i(r'_L) = \sum_{r'_{1L}} k(r'_L \to r'_{1L}) \,\bar{D}_i(r'_{1L}) \,\,, \qquad (\text{IV.18})$$

where the sums in Eqs. (IV.17) and (IV.18) run over all the irreps after decomposition. By inserting Eqs. (IV.17), (IV.18) into (IV.16) one obtains

$$N_{c} = \sum_{r_{1R}'} \sum_{r_{1L}'} d(r_{1R}') \,\bar{D}_{i}(r_{1L}') \sum_{r' \in \mathcal{R}[N_{f}+1]} \ell(r') \,k(r' \to r_{1}') \,, \qquad (\text{IV.19})$$

where  $r'_{1} = (r'_{1L}, r'_{1R}, H_{1}, b)$  is a generic irrep of  $\mathcal{G}[N_{f} + 1, 1]$ . By setting  $r'_{1L} = \{m_{L}; \bar{m}_{L}\}$  and  $r'_{1R} = \{m_{R}; \bar{m}_{R}\}$  one has  $H_{1} = bN_{c} - (m_{L} + m_{R}) + (\bar{m}_{L} + \bar{m}_{R})$ , and

$$k(r' \to r_1') = k(\{n_L; \bar{n}_L\} \to \{m_L; \bar{m}_L\}) \times k(\{n_R; \bar{n}_R\} \to \{m_R; \bar{m}_R\}).$$
(IV.20)

By Lemma 2,  $\{\ell(r')\}$  is a solution of PMC[ $N_f + 1, 1$ ], hence the last sum of Eq. (IV.19) vanishes unless  $H_1 = 0$ . One can thus rewrite Eq. (IV.19) as

$$N_c = \sum_r d(r_R) \,\bar{D}_i(r_L) \sum_{r' \in \mathcal{R}[N_f+1]} \ell(r') \,k(r' \to r) \,, \qquad (\text{IV.21})$$

where  $r = (r_L, r_R, b)$  is now a representation of  $\mathcal{R}[N_f]$  (since it belongs to the subset of representations  $r'_1$  with  $H_1 = 0$ , and  $\mathcal{R}_0[N_f + 1, 1] = \mathcal{R}[N_f]$ ).

The meaning of Eq. (IV.21) can be better understood by introducing the following definition of downlifting: given a spectrum  $\{\ell(r')\}$  of irreps  $r' \in \mathcal{R}[N_f + 1]$ , we define a downlifted spectrum of irreps  $r \in \mathcal{R}[N_f]$  as

$$\{\ell(r)\} = D\big[\{\ell(r')\}\big]: \qquad \ell(r) = \sum_{r' \in \mathcal{R}[N_f+1]} \ell\big(r'\big) \, k\big(r' \to r\big) \ . \tag{IV.22}$$

By  $k(r' \to r)$  here we mean the multiplicity for the decomposition of r' into the representation  $r'_1$ with  $H_1 = 0$  of  $\mathcal{R}_0[N_f + 1, 1]$  which corresponds to r by virtue of the identification  $\mathcal{R}_0[N_f + 1, 1] = \mathcal{R}[N_f]$ . It is possible to show that, for any class-A  $r \in \mathcal{R}[N_f]$ ,

$$\sum_{r' \in \mathcal{R}[N_f+1]} \ell(r') k(r' \to r) = \ell(U[r]) \qquad \text{that is:} \quad D[U[\{\ell(r)\}]] = \{\ell(r)\}. \tag{IV.23}$$

In other words: *downlifting is the inverse of uplifting*. Before providing the proof of this identity, we first discuss its implications.

The last sum in Eq. (IV.21) gives the downlift of the indices  $\ell(r')$ ; since we defined the latter to be the uplift of the  $\ell(r)$ , from the result stated in (IV.23) it follows that Eq. (IV.21) coincides with AMC[N<sub>f</sub>], which is identically satisfied. This proves that  $\{\ell(r')\}$ , defined as specified in Theorem 1, is a solution of AMC[N<sub>f</sub> + 1]. Similarly, one can uplift the solution from N<sub>f</sub> + 1 to N<sub>f</sub> + 2 and so on. We thus obtain that AMC[N<sub>f</sub>]  $\cup$  PMC[N<sub>f</sub>, 1] implies AMC[N'<sub>f</sub>]  $\cup$  PMC[N'<sub>f</sub>, 1] for any N'<sub>f</sub> > N<sub>f</sub>. This proves Theorem 1.

From Theorem 1, chiral symmetry breaking is proven as follows. Let us consider a confining theory with  $N_f$  flavors and assume that integer solutions of  $AMC[N_f] \cup PMC[N_f, 1]$  exist which satisfy the following property: Only massless bound states interpolated by class-A tensors have non-vanishing indices. Then Theorem 1 is valid and  $N_f$  independence of  $AMC[N_f]$  follows; by the argument presented at the beginning of this section, this in turn implies that no integer solution of  $AMC[N_f]$  can exist, contradicting the initial assumption. This proves that either chiral symmetry is spontaneously broken, or the assumed property of the spectrum is violated.

The  $N_f$ -independence of solutions of  $AMC[N_f] \cup PMC[N_f, 1]$  is sufficient to prove chiral symmetry breaking. Under the same assumptions of Theorem 1, on the other hand, it is possible to show that solutions of  $AMC[N_f] \cup PMC[N_f]$  are also  $N_f$ -independent. One can thus state the following

**Theorem 2** ( $N_f$  independence - stronger version). Let  $\{\ell(r)\}$  be a real solution of  $AMC[N_f] \cup PMC[N_f]$  such that the index of any state that cannot be interpolated by a class-A tensor vanishes. Then  $U^{(N'_f - N_f)}[\{\ell(r)\}]$  is a real solution of  $AMC[N'_f] \cup PMC[N'_f]$  for any  $N'_f \ge N_f$ .

For the proof, we need to show that  $\{\ell(r')\} = U^{(N'_f - N_f)}[\{\ell(r)\}]$  is also a solution of  $PMC[N'_f, i > 1]$ . Let us start by considering  $PMC[N_f + 1, 2]$ ; these equations are of the form:

$$0 = \ell(\hat{r}'_2) = \sum_{r'_1 \in \mathcal{R}_0[N_f + 1, 1]} k(r'_1 \to \hat{r}'_2) \ell(r'_1), \qquad \forall \ \hat{r}'_2 \in \hat{\mathcal{R}}[N_f + 1, 2], \qquad (IV.24)$$

where  $\ell(r'_1)$  is the multiplicity of the irrep  $r'_1$  with  $H_1 = 0$  which appears in the decomposition of r':

$$\ell(r'_1) = \sum_{r' \in \mathcal{R}[N_f + 1]} k(r' \to r'_1) \ell(r') \,. \tag{IV.25}$$

Since each  $r'_1$  in  $\mathcal{R}_0[N_f + 1, 1]$  can be identified with a corresponding irrep in  $\mathcal{R}[N_f]$ , it is clear from Eq. (IV.25) that the  $\ell(r'_1)$  are just the downlift of the  $\ell(r')$ . From Eq. (IV.23) and the definition of  $\ell(r')$ , it follows that  $\{\ell(r'_1)\} = D[\{\ell(r')\}] = D[U[\{\ell(r)\}]] = \{\ell(r)\}$ . We know from Section III B that  $PMC[N_f + 1, 2]$  are identical to  $PMC[N_f, 1]$  after setting  $\ell(r'_1) = \ell(r)$ ; therefore, this proves that  $\{\ell(r')\}$  is a solution of  $PMC[N_f + 1, 2]$ . It is straightforward to show that  $\{\ell(r')\}$  is also a solution of  $PMC[N_f + 1, i]$  for any i > 1, since the latter equations coincide with  $PMC[N_f, i - 1]$ . Similarly, by uplifting to higher values of  $N_f$  one can easily show that  $\{\ell(r')\}$  is a solution of  $PMC[N'_f, i > 1]$  for any  $N'_f > N_f$ . This proves Theorem 2.

Finally, we now provide the proof of Eq. (IV.23).

### 4. Proof of Eq. (IV.23): downlifting is the inverse of uplifting

Let us start by noticing that if one assumes a purely baryonic spectrum, then Eq. (IV.23) follows easily. Indeed, in that case  $\bar{n}_{L,R}$  and  $\bar{m}_{L,R}$  vanish in Eq. (IV.20), and  $H_1 = 0$  implies  $m_L = n_L$ ,  $m_R = n_R$  (because  $H_1 = n_L + n_R - m_L - m_R$  and  $m_L \le n_L$ ,  $m_R \le n_R$ ). Therefore, r' and r are interpolated by the same tensor, that is: the only irrep r' contributing in the sum of Eq. (IV.23) is r' = U[r]. Since  $k(U[r] \rightarrow r) = 1$ , then Eq. (IV.23) follows. This justifies the results of the early works [2, 10, 11].

For a spectrum with exotics the proof is less straightforward and, to the best of our knowledge, cannot be found in previous literature. Here we rely on the assumption that only bound states interpolated by class-A tensors have non-vanishing index. The spectrum of corresponding representations is finite, since the number of upper and lower indices of class-A tensors must satisfy  $n + \bar{n} < N_f$ . In particular, class-A tensors can have a maximum number of lower indices  $\bar{n}_{max} < (N_f - bN_c)/2$ . In this section we will assume  $\bar{n}_{max} = 1$  for simplicity, the generalization to  $\bar{n}_{max} > 1$  is discussed in Appendix D. Since baryon number is unchanged when decomposing a tensor, i.e. bound states of different baryon numbers do not mix in PMC, we work at fixed b in the following.

For  $\bar{n}_{\text{max}} = 1$ , there are only two kinds of traceless tensors of  $\mathcal{T}[N_f + 1]$  to be considered:

$$T^{\{bN_c+1\}}_{\{1\}}$$
 and  $T^{\{bN_c\}}_{\{0\}}$ . (IV.26)

Those interpolating the states r' in Eq. (IV.23) will have this form and, by the assumptions of Theorem 1, can be chosen to belong to class A. Tensors interpolating the irreps  $r'_1$  of  $\mathcal{G}[N_f + 1, 1]$ with  $H_1 = i \ge 0$  will be of the form  $T_{\{1\}}^{\{bN_c+1-i\}}$  and  $T_{\{0\}}^{\{bN_c-i\}}$ . In particular, those with  $H_1 = 0$  are of the same form as in Eq. (IV.26).

First, we consider the PMC relative to irreps  $\hat{r}'_1$  of  $\mathcal{G}[N_f + 1, 1]$  that have  $H_1 = i > 0$  and that are interpolated by tensors with  $(bN_c + 1 - i)$  upper indices and 1 lower index. For each such representation there is one PMC of the form

$$0 = \ell(\hat{r}'_{1})$$

$$= \sum_{r' \in \mathcal{R}[N_{f}+1]} \ell(r') k \left( T^{\{bN_{c}+1\}}_{\{1\}} \to T^{\{bN_{c}+1-i\}}_{\{1\}} \right)$$

$$= \sum_{\{bN_{c}+1\}} \ell(r') k (\{bN_{c}+1\} \to \{bN_{c}+1-i\}) ,$$
(IV.27)

where  $T_{\{1\}}^{\{bN_c+1\}}$  transforms as the representation r' of  $\mathcal{G}[N_f+1]$ . Next, we compute the sum of Eq. (IV.23) for representations r (with  $H_1 = 0$ ) that are interpolated by tensors  $T_{\{0\}}^{\{bN_c\}}$ . One has (using  $k(\{1\} \to \{0\}) = 1$ ):

$$\sum_{r'} \ell(r') k(r' \to r) = \ell(U[r]) + \sum_{r' \neq U[r]} \ell(r') k\left(T_{\{1\}}^{\{bN_c+1\}} \to T_{\{0\}}^{\{bN_c\}}\right)$$
  
=  $\ell(U[r]) + \sum_{\{bN_c+1\}} \ell(r') k(\{bN_c+1\} \to \{bN_c\})$ . (IV.28)

The sum in the second line coincides with that appearing in the last line of Eq. (IV.27) for i = 1; hence it vanishes and we obtain Eq. (IV.23). Finally, Eq. (IV.23) follows trivially if r is interpolated by a tensor  $T_{\{1\}}^{\{bN_c+1\}}$ , since in this case r' can only be interpolated by the same tensor, i.e. r' = U[r]. This concludes our proof for  $\bar{n}_{\text{max}} = 1$ .

### V. CONCLUSIONS

In this work we have critically reconsidered the argument based on 't Hooft anomaly matching that aims at proving chiral symmetry breaking in QCD-like theories. The main line of reasoning, first suggested by 't Hooft [1] and later developed by many other authors [2, 8–13], relies on  $N_f$ -independence. Such property was assumed to hold in previous works based on qualitative arguments, and it was never proven rigorously. In this paper we provided a proof of  $N_f$ -independence by establishing an injective map between massless bound states of theories with different number of flavors. We made use of tensor notation and found that such map can be established only on bound states that are interpolated from the vacuum by class-A tensors (see Sec. II B for a definition). We therefore conclude that  $N_f$ -independence holds, hence chiral symmetry breaking is proven, only by making a restriction on the possible massless bound states in the spectrum. Our analysis includes exotic bound states interpolated by tensor operators with lower (antiquark) indices, which were not considered in the past literature except for the work of Ref. [9].

Another interesting strategy to prove chiral symmetry breaking using 't Hooft anomaly matching and  $N_f$ -independence was proposed by Cohen and Frishman in Ref. [12]. These authors focused on the purely baryonic case and suggested that, although  $AMC[N_f]$  and  $PMC[N_f, 1]$  equations depend on  $N_f$  in general, there might be one linear combination of them which does not. Such ' $N_f$ -equation' can be derived from AMC at large  $N_f$  and, once evaluated at  $N_f = 0$  as justified by its  $N_f$ -independence, it has no integer solutions. Proving the existence of the  $N_f$ -equation is not straightforward, since the system of  $AMC[N_f]$  and  $PMC[N_f, 1]$  does not remain the same as  $N_f$  is varied, and the arguments provided in Ref. [12] are not conclusive. The best we can infer from our work is that the derivation of this equation is not in contradiction with the downlifting procedure used in Ref. [7].

The alternative approach proposed by Schwimmer in Ref. [8] makes use of the  $SU(N_f|N_f)$ superalgebra and does not rely on  $N_f$ -independence. As such, it is a promising candidate for a purely algebraic proof of chiral symmetry breaking. We have explicitly verified that, in the case of QCD-like theories with  $N_c = 3$  and 5, the superalgebra solution coincides with the generic solution of PMC equations if one assumes a purely baryonic spectrum of massless states. This result holds for any number of flavors  $N_f > 2$ , i.e. even in presence of equivalent tensors. It would be desirable to extend the work of Schwimmer to the case of a massless spectrum with exotics, and prove that a generic PMC solution can be always written in terms of irreps of  $SU(N_f|N_f)$ . In this way one would obtain an algebraic proof of chiral symmetry breaking valid for any  $N_f > 2$ . We leave such

The notion of global symmetries has been recently generalized [34], providing some new perspective on strongly-coupled gauge dynamics. Among the many papers on this subject that appeared in the literature, those which use 't Hooft anomaly matching of generalized symmetries include studies analyzing the vacuum of gauge theories at non-vanishing theta angle [35, 36], constraining the dynamics in the space of coupling constants [37, 38], constraining the phase diagram of QCD at finite temperature and/or finite density [39], excluding an exotic chiral symmetry breaking phase of QCD [40], constraining the infrared dynamics of QCD-like theories with quarks in higher-dimensional representations with non-trivial N-ality [41] and of chiral gauge theories (see e.g. Ref. [42] for a recent review and references therein). Such line of research is extremely interesting and might lead to additional insight on the occurrence of the ordinary chiral symmetry breaking phase in QCD-like theories where more traditional arguments like those used in this paper fail.

study to a future work.

In this work we focused on non-supersymmetric vector-like gauge theories with fermions in the fundamental representation of the color group. It would be interesting to extend our results to theories with fermions in different representations, or with different gauge groups. Studies of chiral symmetry breaking in some of these theories have been made in the literature, but we expect that our approach can be used to obtain further progress in a more systematic way.

### ACKNOWLEDGMENTS

We would like to thank Marco Bochicchio, Ethan Neil, Mauro Papinutto, Slava Rychkov, Yael Shadmi and Giovanni Villadoro for useful discussions and comments. This research was supported in part by the MIUR under contract 2017FMJFMW (PRIN2017) and performed in part at Aspen Center for Physics, which is supported by National Science Foundation grant PHY-2210452. The work of R.C. was partly supported by a grant from the Simons Foundation and by the Munich Institute for Astro- and Particle Physics (MIAPP), which is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany Excellence Strategy - EXC-2094 - 390783311. The work of A.L. was partially supported by the INFN special initiative grant "GAST" (Gauge and String Theories). L.X.X. would like to thank Scuola Normale Superiore in Pisa, where this project was initiated, for its warm hospitality. L.X.X. was supported in part by the National Science Foundation of China under Grants No. 11635001 and No. 11875072 while he was working on this project at Peking University.

#### **Appendix A: Technicalities on Anomalies**

This Appendix further clarifies some technicalities encountered in the previous sections.

For quarks in the fundamental representation, the anomaly traces are

$$D_{i=2}(\Box) \,\delta_{ab} = \frac{1}{N_c} \operatorname{tr}\left[\{\lambda_a^F, \lambda_b^F\}\right] \equiv \frac{1}{N_c} \delta_{ab} \,,$$
  

$$D_{i=3}(\Box) \,d_{abc} = \operatorname{tr}\left[\{\lambda_a^F, \lambda_b^F\}\lambda_c^F\right] \equiv d_{abc} \,.$$
(A.1)

Here  $\lambda_a^F$   $(a = 1, 2, \dots, N_f^2 - 1)$  are the generators of  $SU(N_f)$  in the fundamental representation, whose normalization satisfies  $tr[\lambda_a^F \lambda_b^F] = \delta_{ab}/2$ . The anomaly traces for bound states that appear in Eq. (III.4) are defined as

$$D_{i=2}(r) \,\delta_{ab} = b \,\operatorname{tr}[\{\lambda_a, \lambda_b\}] ,$$

$$D_{i=3}(r) \,d_{abc} = \operatorname{tr}[\{\lambda_a, \lambda_b\}\lambda_c] .$$
(A.2)

Here  $\lambda_a$   $(a = 1, 2, \dots, N_f^2 - 1)$  are the generators of  $SU(N_f)$  in the representation r. The constants  $\overline{D}_i(r)$  are conveniently defined as the ratios of anomaly traces, see Eq. (III.6). We notice that the values of  $\overline{D}_i(r)$  do not depend on which generators are chosen in calculating the traces; without loss of generality, one can choose generators belonging to the  $SU(N_f - 1)$  subalgebra.

It is worthwhile to comment more on Eq. (IV.17) and Eq. (IV.18), given their importance in the proof of  $N_f$ -independence. Equation (IV.17) just reflects the fact that the dimensionality of a representation is equal to the sum of the dimensionalities of the representations appearing in its decomposition. Equation (IV.18) is less trivial to justify, and we do so by considering the following facts:

1. The constants  $\overline{D}_i(r')$ , where r' is an irrep of  $SU(N_f + 1)$ , can be calculated by using the generators belonging to the  $SU(N_f)$  subalgebra of the original  $SU(N_f + 1)$ ; these act non-trivially on r' only when its indices are charged under  $SU(N_f)$ . This can be schematically indicated by

$$\bar{D}_i(r') = \bar{D}_i(r')|_{SU(N_f)}, \qquad (A.3)$$

where on the right-hand side we mean that  $D_i$  can be evaluated by using the  $SU(N_f)$  subalgebra.

2. The irrep r' is reducible under the  $SU(N_f)$  subalgebra, i.e. it can be decomposed into a series of irreducible representations (r, H) of  $SU(N_f) \times U(1)_H$ ,

$$r' = \bigoplus_{r} (r, H) , \qquad (A.4)$$

where each index of r' varies from 1 to  $N_f + 1$ , while each index of r goes from 1 to  $N_f$ . The  $U(1)_H$  charge of the decomposed representations is given by  $H = (n - m) - (\bar{n} - \bar{m})$ , where n and  $\bar{n}$  (m and  $\bar{m}$ ) are the numbers of upper and lower indices of r' (of r). The YTs characterizing r' and r can be identical, in which case H = 0. It follows that

$$\bar{D}_i(r')|_{SU(N_f)} = \bar{D}_i\left(\bigoplus_r r\right) . \tag{A.5}$$

3. Equation (IV.18) follows from Eqs. (A.3), (A.5) and the identity

$$\bar{D}_i \left( r_1 \bigoplus r_2 \right) = \bar{D}_i (r_1) + \bar{D}_i (r_2) \tag{A.6}$$

valid for any two irreps  $r_1$  and  $r_2$  of  $SU(N_f)$ .

### **Appendix B: Equivalent Tensors**

This Appendix contains the proof of Lemma 1 on the absence of equivalent tensors stated in Sec. II. Lemma 1 applies to class-A tensors, for which

$$n + \bar{n} < N_f \,. \tag{B.1}$$

In particular, we will prove that this is a necessary and sufficient condition for the absence of equivalent tensors, hence defining the largest set of non-equivalent tensors.

As explained in Sec. II, one way to construct equivalent traceless tensors of  $\mathcal{T}[N_f]$  is to let their YTs differ by the presence of one or more sets of  $N_f$  fully antisymmetrized indices (flavor singlets). In this case, the CYTs also differ by columns with  $N_f$  boxes. Clearly, for  $n + \bar{n} < N_f$  no such tensors can arise. This condition also ensures that no equivalent traceless tensors arise after decomposition which transform under the same representation of  $\mathcal{G}[N_f, 1]$  and whose YT differ only by flavor singlets. We thus consider the less trivial case where the equivalence arises by combining the YT of upper and lower indices. For simplicity, we will focus on tensors with only left or right indices, the generalization to tensors with both kinds of indices is straightforward. Let

$$\left\{T_{\{\bar{n}_1\}}^{\{n_1\}}, T_{\{\bar{n}_2\}}^{\{n_2\}}, \cdots, T_{\{\bar{n}_\kappa\}}^{\{n_\kappa\}}\right\}$$
(B.2)

be a class of  $\kappa$  equivalent traceless tensors of this kind. Since no flavor singlets can arise due to Eq. (B.1), all these tensors must have the same CYT. Let  $\mathcal{R} = \mathcal{P} + \mathcal{Q}$  be the number of columns of the CYT, where  $\mathcal{P}$  of them originate from the YT of upper (quark) indices, and the remaining  $\mathcal{Q}$ from the dual YT of lower (antiquark) indices. For a given tensor  $T_{\{\bar{n}_j\}}^{\{n_j\}}$ , we will indicate with  $\zeta_i^{(j)}$ the length of the *i*-th column of  $\{n_j\}$ , for  $i = 1, \ldots, \mathcal{P}$ , and with  $\zeta_i^{(j)}$  the length of the *i*-th column of  $\{\bar{n}_j\}$ , for  $i = 1, \ldots, \mathcal{Q}$ . The total number of boxes in the CYT then equals

$$n_{\text{box}} = \sum_{i=1}^{\mathcal{P}} \zeta_i^{(j)} + \sum_{i=1}^{\mathcal{Q}} \tilde{\zeta}_i^{(j)} = n_j + \mathcal{Q}N_f - \bar{n}_j = bN_c + \mathcal{Q}N_f \,, \tag{B.3}$$

where the last equality follows from the sum rule (II.1). Since  $n_{\text{box}}$  does not depend on j, all tensors in the equivalent class have the same  $\mathcal{Q}$ , and thus the same  $\mathcal{P}$  as well. In other words, the class is characterized by  $(\mathcal{P}, \mathcal{Q})$  and by the baryon number b. The number of equivalent tensors in a class  $(\mathcal{P}, \mathcal{Q})_b$  is given by the binomial coefficient:

$$\kappa = C_{\mathcal{R}}^{\mathcal{P}} = C_{\mathcal{R}}^{\mathcal{Q}} \equiv \frac{\mathcal{R}!}{\mathcal{P}!\mathcal{Q}!} \,. \tag{B.4}$$

Indeed, starting with a total of  $\mathcal{R}$  columns in the CYT, each tensor corresponds to a different way of assigning  $\mathcal{P}$  of them to the YT of upper indices (so that the remaining ones are columns of the dual YT of lower indices); there are  $C_{\mathcal{R}}^{\mathcal{P}}$  such possible assignments.

By defining  $n_{max} = \max_{j} \{n_j\}$  and  $\bar{n}_{max} = \max_{j} \{\bar{n}_j\}$  as respectively the maximum number of quarks and antiquarks of any tensor in the class, these two inequalities follow:

$$\sum_{j=1}^{n} \bar{n}_j \le \kappa \,\bar{n}_{max} = C_{\mathcal{R}}^{\mathcal{P}} \,\bar{n}_{max} \tag{B.5}$$

$$\sum_{j=1}^{\kappa} n_j \le \kappa \, n_{max} = C_{\mathcal{R}}^{\mathcal{P}} \left( \bar{n}_{max} + b N_c \right) \,. \tag{B.6}$$

Furthermore, the following identity holds:

$$\mathcal{P}\sum_{j=1}^{\kappa} \bar{n}_j + \mathcal{Q}\sum_{j=1}^{\kappa} n_j = C_{\mathcal{R}}^{\mathcal{P}} \mathcal{P} \mathcal{Q} N_f .$$
(B.7)

We postpone the proof to the end and focus first on its implications. Using (B.7) together with (B.5) and (B.6), one obtains

$$N_f \le \frac{\bar{n}_{max}}{\mathcal{Q}} + \frac{\bar{n}_{max} + bN_c}{\mathcal{P}} \,. \tag{B.8}$$

For fixed  $N_f$ , the weakest lower bound on  $\bar{n}_{max}$  and b comes from the classes with  $\mathcal{P} = \mathcal{Q} = 1$  and reads  $N_f \leq 2\bar{n}_{max} + bN_c$ . Any class of this kind with  $\mathcal{P} = \mathcal{Q} = 1$  contains a pair of equivalent tensors  $(\kappa = 2)$  with numbers of quarks and antiquarks related as follows:  $n_1 = N_f - \bar{n}_2$ ,  $n_2 = N_f - \bar{n}_1$ .<sup>14</sup>. The Eq. (B.8) implies that either  $2\bar{n}_1 + bN_c \leq N_f$  or  $2\bar{n}_2 + bN_c \leq N_f$ . Therefore, we see that Eq. (B.1) is a necessary requirement for avoiding pairs of equivalent tensors. We notice that in fact no equivalent traceless tensors with identical CYTs can arise even for  $2\bar{n} + bN_c = N_f$ , when  $\bar{n} \neq 0$ . Indeed, if the inequality of Eq. (B.8) is saturated, then the second tensor has (as a consequence of the relations written above) the same n and  $\bar{n}$ . In other words, in this case the two equivalent tensors are actually identical. The stronger condition of Eq. (B.1) is on the other hand required to exclude the existence of equivalent tensors whose YTs differ by flavor singlets.

The final step is to prove that Eq. (B.1) is a sufficient condition for the absence of equivalent tensors. Let us consider a generic pair of equivalent tensors  $T_{\{\bar{n}_1\}}^{\{n_1\}}, T_{\{\bar{n}_2\}}^{\{n_2\}}$  and suppose  $n_1 + \bar{n}_1 < N_f$ . Since they have the same  $\mathcal{P}$  and  $\mathcal{Q}$  they will be connected by exchanging the same number rof columns of quarks and antiquarks. If we call  $\delta$  and  $\bar{\delta}$  respectively the number of quarks and antiquarks of  $T_{\{\bar{n}_1\}}^{\{n_1\}}$  that were exchanged, we have that  $n_2 + \bar{n}_2 = n_1 + (rN_f - \bar{\delta}) - \delta + \bar{n}_1 + (rN_f - \delta) - \bar{\delta}$ . Since  $\delta + \bar{\delta} \leq n_1 + \bar{n}_1 < N_f$ , this imply that  $n_2 + \bar{n}_2 > (2r - 1)N_f \geq N_f$ . This means that there cannot be equivalent tensors both satisfying Eq. (B.1) and concludes our proof.

A few additional considerations show that when Eq. (B.1) is satified, no equivalent traceless tensors of  $\mathcal{G}[N_f, 1]$  can possibly arise as well. The argument leading to Eq. (B.8) can be repeated for traceless tensors of  $\mathcal{G}[N_f, 1]$  by replacing  $n_j$  and  $\bar{n}_j$  with the number of massless quarks  $n'_j$  and  $\bar{n}'_j$  characterizing a tensor of the class. The values of  $n'_j$  and  $\bar{n}'_j$  satisfy the sum rule

$$n'_{j} - \bar{n}'_{j} = bN_{c} - H$$
, (B.9)

where H is the number of massive quarks minus the number of massive antiquarks in the bound state. Since H corresponds to the charge under the  $U(1)_H$  factor inside  $\mathcal{G}[N_f, 1]$ , all tensors in the same class must have the same H. Since for every tensor  $n'_j + \bar{n}'_j \leq n_j + \bar{n}_j - 1$ , then Eq. (B.1) ensures that no singlets of  $SU(N_f - 1)$  can appear in either of the upper and lower YT of the decomposed tensor. The steps that led to Eq. (B.8) now imply

$$N_f - 1 \le \frac{\bar{n}'_{max}}{\mathcal{Q}} + \frac{n'_{max}}{\mathcal{P}}.$$
(B.10)

The strongest constraint is obtained by setting  $\mathcal{P} = \mathcal{Q} = 1$ .

<sup>&</sup>lt;sup>14</sup> These identities follow by requiring that the CYT be the same for the two tensors. Indeed, the CYT has two columns. In the case of the first tensor, the first column originates from its quark indices and the second column originates from its antiquark indices, while for the second tensor the role of quark and antiquark indices is exchanged.

- If H > 0, then there are more massive quarks than antiquarks, and  $n'_{max} \leq n_{max} 1$ ,  $\bar{n}'_{max} \leq \bar{n}_{max}$ .
- Similarly, if H < 0 there are more massive antiquarks than quarks, and  $n'_{max} \leq n_{max}$ ,  $\bar{n}'_{max} \leq \bar{n}_{max} - 1$ .
- Finally, if H = 0 there is the same number of massive quarks and antiquarks; hence  $n'_{max} \le n_{max} 1$  and  $\bar{n}'_{max} \le \bar{n}_{max} 1$ .

From these inequalities and Eq. (B.10) it follows  $N_f - 1 \leq 2\bar{n}_{max} + bN_c - 1$ , so that no equivalent traceless tensors of  $\mathcal{G}[N_f, 1]$  exist if Eq. (B.1) is satisfied. Considerations similar to those made after Eq. (B.8) show that no equivalent tensors can arise with identical CYT even when  $N_f = 2\bar{n}_{max} + bN_c$ ; this limit case must be however excluded to avoid the presence of equivalent tensors whose YT differ by flavor singlets.

To complete our proof we have to establish the validity of Eq. (B.7). We start by considering the following sums:

$$\sum_{j=1}^{\kappa} n_j = \sum_{j=1}^{\kappa} \sum_{i=1}^{\mathcal{P}} \zeta_i^{(j)}$$
(B.11)

$$\sum_{j=1}^{\kappa} \bar{n}_j = \sum_{j=1}^{\kappa} \sum_{i=1}^{\mathcal{Q}} (N_f - \tilde{\zeta}_i^{(j)}) = C_{\mathcal{R}}^{\mathcal{P}} \mathcal{Q} N_f - \sum_{j=1}^{\kappa} \sum_{i=1}^{\mathcal{Q}} \tilde{\zeta}_i^{(j)}.$$
 (B.12)

It is not difficult to perform the sums on the right-hand sides of Eqs.(B.11), (B.12). For example,  $\sum_{j=1}^{\kappa} \sum_{i=1}^{\mathcal{P}} \zeta_i^{(j)}$  can be computed by counting how many times a given column of length  $\zeta$  appears in all possible YTs of upper indices within the class. This number is equal to the binomial coefficient  $C_{\mathcal{R}-1}^{\mathcal{P}-1}$ , which corresponds to the number of YTs one can form with that column and other  $\mathcal{P}-1$ chosen from the  $\mathcal{R}-1$  available in the CYT. This multiplicity does not depend on which column is considered initially. Similarly, the sum  $\sum_{j=1}^{\kappa} \sum_{i=1}^{\mathcal{Q}} \tilde{\zeta}_i^{(j)}$  is computed by noticing that a given column of length  $\tilde{\zeta}$  appears  $C_{\mathcal{R}-1}^{\mathcal{Q}-1}$  times in all possible dual YTs of lower indices; again, this multiplicity does not depend on the column considered. One thus obtains

$$\sum_{j=1}^{\kappa} \sum_{i=1}^{\mathcal{P}} \zeta_i^{(j)} = C_{\mathcal{R}-1}^{\mathcal{P}-1} n_{\text{box}}$$
(B.13)

$$\sum_{j=1}^{\kappa} \sum_{i=1}^{Q} \tilde{\zeta}_{i}^{(j)} = C_{\mathcal{R}-1}^{Q-1} n_{\text{box}}, \qquad (B.14)$$

where  $n_{\text{box}}$  is the total number of boxes in the CYT, see Eq. (B.3). Since

$$\mathcal{P}C_{\mathcal{R}-1}^{\mathcal{Q}-1} = \frac{(\mathcal{R}-1)!}{(\mathcal{P}-1)!(\mathcal{Q}-1)!} = \mathcal{Q}C_{\mathcal{R}-1}^{\mathcal{P}-1},$$
(B.15)

from Eqs. (B.11)-(B.14) it follows Eq. (B.7).

#### Appendix C: PMC with equal masses

In this Appendix, we argue that the PMC that one obtains by considering k massive quarks with two or more equal masses are implied by  $PMC[N_f, 1], \ldots, PMC[N_f, k]$ . First, we prove the statement for two equal masses, and then we sketch how to extend the proof to the general case. Our proof relies on the very reasonable assumption that the spectrum of massless fermions with non-zero index is finite, i.e. there is an arbitrary though finite number of irreps  $r \in \mathcal{R}[N_f]$  such that  $\ell(r) \neq 0$ .<sup>15</sup>

If one gives two flavors the same mass,  $\mathcal{G}[N_f, 2]$  is enlarged to <sup>16</sup>

$$\mathcal{G}[N_f,\underline{2}] = SU(N_f - 2)_L \times SU(N_f - 2)_R \times U(1)_B \times SU(2)_H \times U(1)_H \quad , \tag{C.1}$$

where  $SU(2)_H$  acts on the doublet of massive quarks, and the  $U(1)_H$  symmetry counts their overall number. Here and in the following we underline quantities related to the two equal masses. Clearly,  $\mathcal{G}[N_f, 2]$  is a subgroup of  $\mathcal{G}[N_f, \underline{2}]$ .

It is useful to denote an irrep of  $\mathcal{G}[N_f, \underline{2}]$  with a tuple,  $\{A, h, \underline{j}\}$ , where A is an  $SU(N_f - 2)_L \times SU(N_f - 2)_R \times U(1)_B$  representation, h is the  $U(1)_H$  charge, and  $\underline{j}$  is an  $SU(2)_H$  irrep with spin  $\underline{j}$ . If h is even (odd), then  $\underline{j}$  must be integer (half-integer); to see this, just notice that an operator interpolating  $\{A, h, \underline{j}\}$  has a number of massive quarks equal to h, up to an arbitrary number of massive quark-antiquark pairs. <sup>17</sup> We further denote an irrep of  $\mathcal{G}[N_f, 2]$  with a tuple  $\{A, h, m_j\}$ . Here h and  $m_j$  codify the charges under  $U(1)_{H_1}$  and  $U(1)_{H_2}$  as  $h_1 = \frac{1}{2}h + m_j$  and  $h_2 = \frac{1}{2}h - m_j$ , respectively. Notice that  $h_1$  and  $h_2$  are always integers due to the connection between h and  $\underline{j}$ . By using the Vafa-Witten theorem (the same argument used in Section. III B), one can prove that

$$\sum_{\in \mathcal{R}[N_f]} \ell(r) \ k(r \to \{A, h, \underline{j}\}) = 0 \tag{C.2}$$

when h and j do not vanish simultaneously. Our goal is to show that these equations can be obtained from  $PMC[N_f, 1]$  and  $PMC[N_f, 2]$ .

The first step is to notice that  $PMC[N_f, 1]$  and  $PMC[N_f, 2]$  imply that

$$\sum_{r \in \mathcal{R}[N_f]} \ell(r) \ k(r \to \{A, h, m_j\}) = 0 \ . \tag{C.3}$$

<sup>&</sup>lt;sup>15</sup> It is believed that in a quantum field theory the number of massless degrees of freedom in the IR cannot be larger than the number of UV degrees of freedom, see for example [43, 44].

<sup>&</sup>lt;sup>16</sup> Here we are ignoring, as done before in this paper, any discrete identification. In particular, the actual symmetry acting on the massive flavors is  $U(2)_H = (SU(2)_H \times U(1)_H)/\mathbb{Z}_2$ .

<sup>&</sup>lt;sup>17</sup> From a more abstract viewpoint, the overlap between the  $SU(2)_H$  center and  $U(1)_H$  ensures that h + 2j = 0 mod 2.

for any irrep  $\{A, h, m_j\}$  of  $\mathcal{G}[N_f, 2]$  with  $m_j$  and h not simultaneously equal to zero. Let us call this set of equations  $\overline{PMC}[N_f, 2]$ . We assume that the spectrum of irreps r with non-vanishing index is finite, i.e. the sum in Eq. (C.3) is over a finite number of terms. Then it follows that the set of irreps  $\{A, h, \underline{j}\}$  of  $\mathcal{G}[N_f, \underline{2}]$  obtained from the decomposition of the original massless spectrum is also finite. Let us fix A and h and call  $\underline{j}_{max}$  the highest spin of  $SU(2)_H$  that appears in the decomposition. In general, we can decompose each  $\{A, h, \underline{j}\}$  further in irreps of  $\mathcal{G}[N_f, \underline{2}]$  as

$$\{A, h, \underline{j}\} = \{A, h, j\} \oplus \{A, h, j-1\} \oplus ... \oplus \{A, h, -j\}.$$
 (C.4)

Now we can proceed with a 'peeling off' argument. The decomposition (C.4) implies that, being  $j_{max}$  the maximal  $SU(2)_H$  spin in our spectrum,  $\{A, h, j_{max}\}$  appears only in the decomposition of  $\{A, h, \underline{j}_{max}\}$ , so

$$\sum_{e \in \mathcal{R}(N_f)} \ell(r) \, k(r \to \{A, h, \underline{j}_{max}\}) = \sum_{r \in \mathcal{R}(N_f)} \ell(r) \, k(r \to \{A, h, j_{max}\}) = 0 \,. \tag{C.5}$$

This is the  $PMC[N_f, \underline{2}]$  equation for  $\{A, h, \underline{j}_{max}\}$ , and we are done with the first layer.

Let us consider the second layer. The irrep  $\{A, h, j_{max} - 1\}$  appears in the decomposition of  $\{A, h, j_{max}\}$  and  $\{A, h, j_{max} - 1\}$ , in both cases only once. Hence,

$$\sum_{r \in \mathcal{R}(N_f)} \ell(r) \ k(r \to \{A, h, \underline{j_{max}}\}) + \sum_{r \in \mathcal{R}(N_f)} \ell(r) \ k(r \to \{A, h, \underline{j_{max} - 1}\})$$

$$= \sum_{r \in \mathcal{R}(N_f)} \ell(r) \ k(r \to \{A, h, j_{max} - 1\}) = 0 \ .$$
(C.6)

The first term in the first line of (C.6) vanishes because of Eq. (C.5), leaving the PMC[ $N_f, \underline{2}$ ] equation for  $\{A, h, \underline{j_{max} - 1}\}$ .

One can apply this same argument iteratively, obtaining the PMC equation for  $\{A, h, \underline{j_{max}} - i\}$ , with  $i < j_{max}$ , from the  $\overline{PMC}[N_f, 2]$  equation for  $\{A, h, j_{max} - i\}$  and the  $PMC[N_f, 2]$  equations for  $\{A, h, \underline{j_{max}} - k\}$  with k < i. If h is even and non-vanishing, then one can proceed further and prove also the PMC equation for  $\{A, h, \underline{0}\}$  by using the PMC equation for  $\{A, h, 0\}$ . We have thus proven that the  $PMC[N_f, 2]$  can be derived from  $PMC[N_f, 1]$  and  $PMC[N_f, 2]$  (more precisely, from  $\overline{PMC}[N_f, 2]$ ). Clearly, it is also true that  $\overline{PMC}[N_f, 2]$  can be derived from  $PMC[N_f, 2]$ . Indeed,

$$\sum_{\in \mathcal{R}(N_f)} \ell(r)k(r \to \{A, h, j\}) = \sum_{j' \ge j} \sum_{r \in \mathcal{R}(N_f)} \ell(r)k(r \to \{A, h, \underline{j'}\}) = 0.$$
(C.7)

The first straightforward generalization of this proof is to consider k equal masses. In this case, the symmetry group is <sup>18</sup>

$$G[N_f, \underline{k}] = SU(N_f - k) \times SU(N_f - k) \times U(1)_B \times SU(k)_H \times U(1)_H$$
(C.8)

 $<sup>^{18}</sup>$  Again, we ignore the discrete identification in the sketch of the proof.

 $G[N_f, k]$  is a subgroup of  $G[N_f, \underline{k}]$ . In particular one can identify  $U(1)_{H_1} \times U(1)_{H_2} \times ... \times U(1)_{H_k}$ with  $U(1)_H \times U(1)^{k-1}$ , where  $U(1)^{k-1}$  is the Cartan subgroup of  $SU(k)_H$ . We can generalize the notation  $\{A, h, \underline{j}\}$  used before with  $\{A, h, \underline{\mu}\}$ , where we label the  $SU(k)_H$  irrep with its highest weight  $\overline{\mu} = (\mu_1, ..., \mu_{k-1})$ . We remind that: *i*) the  $\mu_i$  are eigenvalues of the Cartan operators; *ii*) it is possible to define an ordering between weight vectors, e.g. the lexicographic one:  $\overline{\mu} > \overline{\mu}'$  if the first non-zero element of  $\overline{\mu} - \overline{\mu}'$  is positive. As in the k = 2 case, we further label the irreps  $\{A, h, \overline{\mu}\}$ of  $\mathcal{G}[N_f, k]$  with their  $U(1)_H \times U(1)^{k-1}$  charges, h and  $\overline{\mu}$ , instead of the  $U(1)_{H_1} \times ... \times U(1)_{H_k}$  ones. Upon decomposing an irrep of  $\mathcal{G}[N_f, \underline{k}]$  into irreps of  $\mathcal{G}[N_f, k]$ , one obtains

$$\{A, h, \underline{\vec{\mu}}\} = \{A, h, \vec{\mu}\} \bigoplus_{i} \{A, h, \vec{\mu} - \vec{\alpha}_i\} \oplus \dots$$
(C.9)

Here  $\vec{\alpha}_i$  are the positive roots of the su(k) algebra, and the dots denote terms obtained by further subtractions of the  $\vec{\alpha}_i$ .

We can repeat the 'peeling off' argument given before. Let us fix A and h and single out, in the (finite) space of all the irreps of  $\mathcal{G}[N_f, \underline{k}]$ , the one with the highest vector  $\underline{\mu}_{max}$ . Since the unique irrep  $\{A, h, \underline{\mu}_{max}\}$  in the decomposed spectrum comes from the decomposition of  $\{A, h, \underline{\mu}_{max}\}$ , <sup>19</sup> one has

$$\sum_{r \in \mathcal{R}[N_f]} \ell(r)k(r \to \{A, h, \underline{\vec{\mu}}_{max}\}) = \sum_{r \in \mathcal{R}[N_f]} \ell(r)k(r \to \{A, h, \vec{\mu}_{max}\}) = 0.$$
(C.10)

This is the PMC[ $N_f, \underline{k}$ ] for  $\{A, h, \underline{\vec{\mu}}_{max}\}$ .

We can thus proceed to the second layer, i.e. the second highest weight in the spectrum,  $\vec{\mu}'_{max}$ . In general, this state can appear only in the decomposition of  $\underline{\vec{\mu}}_{max}$  and of  $\underline{\vec{\mu}'}_{max}$ . Therefore, we have

$$0 = \sum_{r \in \mathcal{R}[N_f]} \ell(r) \, k(r \to \{A, h, \vec{\mu}'_{max}\})$$
  
= 
$$\sum_{r \in \mathcal{R}[N_f]} \ell(r) \, k(r \to \{A, h, \underline{\vec{\mu}}_{max}\}) \, k(\{A, h, \underline{\vec{\mu}}_{max}\} \to \{A, h, \vec{\mu}'_{max}\})$$
  
+ 
$$\sum_{r \in \mathcal{R}[N_f]} \ell(r) \, k(r \to \{A, h, \underline{\vec{\mu}}'_{max}\}) \,,$$
(C.11)

where we used that  $k(\{A, h, \underline{\vec{\mu}}_{max}\}) \to \{A, h, \underline{\vec{\mu}}_{max}\}) = 1$ . The sum in the second line vanishes because of Eq. (C.10) and we thus obtain the PMC[ $N_f, \underline{k}$ ] for  $\{A, h, \underline{\vec{\mu}}_{max}\}$ .

Similarly to the k = 2 case, one can proceed in this way layer by layer until  $\vec{\mu} = 0$ . Again, the case  $\vec{\mu} = 0$  can be included only if  $h \neq 0$ .

<sup>&</sup>lt;sup>19</sup> The maximal weight vector, i.e. the element of the irrep space with the maximal weight, is unique in any SU(k) irrep (see [45]).

In the most generic case, one has several blocks of flavors with equal masses. It is possible to generalize the proof in this case by proceeding block by block. One can start from the first block of degenerate quarks and prove that one obtains the same PMC equations if this degeneracy is broken. Then, one can proceed inductively with the remaining blocks.

### Appendix D: General Proof of Eq. (IV.23)

In this Appendix we generalize the proof of Eq. (IV.23) given in Section IV A 4 by considering a generic spectrum of massless bound states. The assumption is that each irrep with nonvanishing index can be interpolated by a class-A tensor, i.e. by a traceless tensor  $T_{\{\bar{n}\}}^{\{n\}}$  with  $n+\bar{n}=2\bar{n}+bN_c < N_f$ . This implies a finite maximal number of antiquarks  $\bar{n}_{max} < (N_f - bN_c)/2$ . The strategy is to decompose tensors starting from those with the highest rank.

We first consider representations  $r \in \mathcal{R}[N_f]$  interpolated by tensors with the highest rank, i.e. tensors  $T_{\{\bar{n}_{\max}\}}^{\{bN_c+\bar{n}_{\max}\}}$ . In this case, only the representation r' interpolated by the same tensor can appear in the sum of Eq. (IV.23), i.e. r' = U[r]. Hence, Eq. (IV.23) follows easily.

Next, we consider the PMC relative to irreps  $\hat{r}'_1$  of  $\mathcal{G}[N_f + 1, 1]$  interpolated by tensors with  $(bN_c + \bar{n}_{\max} - i)$  upper indices and  $\bar{n}_{\max}$  lower indices, i.e.  $T^{\{bN_c + \bar{n}_{\max} - i\}}_{\{\bar{n}_{\max}\}}$ . Here i > 0 and the  $U(1)_{H_1}$  charge of the irrep is  $H_1 = i > 0$ . For each such irrep of  $\mathcal{G}[N_f + 1, 1]$  one has a PMC of the form

$$0 = \ell(\hat{r}'_{1})$$

$$= \sum_{r' \in \mathcal{R}[N_{f}+1]} \ell(r') k \left( T^{\{bN_{c}+\bar{n}_{\max}\}}_{\{\bar{n}_{\max}\}} \to T^{\{bN_{c}+\bar{n}_{\max}-i\}}_{\{\bar{n}_{\max}\}} \right)$$

$$= \sum_{\{bN_{c}+\bar{n}_{\max}\}} \ell(r') k (\{bN_{c}+\bar{n}_{\max}\} \to \{bN_{c}+\bar{n}_{\max}-i\}) .$$
(D.1)

We thus compute the sum in Eq. (IV.23) for representations r (with  $H_1 = 0$ ) interpolated by

tensors  $T_{\{\bar{n}_{\max}-1\}}^{\{bN_c+\bar{n}_{\max}-1\}}$ . By making use of Eq. (D.1) with i = 1 we find

$$\sum_{r'} \ell(r') k(r' \to r) = \ell(U[r]) + \sum_{r' \neq U[r]} \ell(r') k\left(T_{\{\bar{n}_{\max}\}}^{\{bN_c + \bar{n}_{\max}\}} \to T_{\{\bar{n}_{\max} - 1\}}^{\{bN_c + \bar{n}_{\max} - 1\}}\right)$$

$$= \ell(U[r]) + \sum_{\{\bar{n}_{\max}\}} k(\{\bar{n}_{\max}\} \to \{\bar{n}_{\max} - 1\})$$

$$\times \underbrace{\left(\sum_{\{bN_c + \bar{n}_{\max}\}} \ell(r') k(\{bN_c + \bar{n}_{\max}\} \to \{bN_c + \bar{n}_{\max} - 1\})\right)}_{=0}$$

$$= \ell(U[r]) , \qquad (D.2)$$

which verifies Eq. (IV.23).

Third, let us consider  $PMC[N_f + 1, 1]$  relative to irreps  $\hat{r}'_1$  of  $\mathcal{G}[N_f + 1, 1]$  interpolated by tensors with  $bN_c + \bar{n}_{max} - 1 - i'$  upper indices and  $\bar{n}_{max} - 1$  lower indices, whose  $U(1)_{H_1}$  charge is  $H_1 = i' > 0$ . These irreps appear in the decomposition of tensors with the second highest rank  $T^{\{bN_c + \bar{n}_{max}-1\}}_{\{\bar{n}_{max}\}}$  and the highest rank  $T^{\{bN_c + \bar{n}_{max}\}}_{\{\bar{n}_{max}\}}$ . For each such irrep  $\hat{r}'_1$  one has a PMC

$$0 = \ell(\hat{r}'_{1})$$

$$= \sum_{\{bN_{c} + \bar{n}_{\max} - 1\}} \ell(r') k(\{bN_{c} + \bar{n}_{\max} - 1\} \rightarrow \{bN_{c} + \bar{n}_{\max} - 1 - i'\})$$

$$+ \sum_{\{\bar{n}_{\max}\}} k(\{\bar{n}_{\max}\} \rightarrow \{\bar{n}_{\max} - 1\})$$

$$\times \sum_{\{bN_{c} + \bar{n}_{\max}\}} \ell(r') k(\{bN_{c} + \bar{n}_{\max}\} \rightarrow \{bN_{c} + \bar{n}_{\max} - 1 - i'\}) .$$

$$= 0$$
(D.3)

The last line vanishes because of Eq. (D.1); hence we obtain

$$0 = \ell(\hat{r}'_1) = \sum_{\{bN_c + \bar{n}_{\max} - 1\}} \ell(r') k (\{bN_c + \bar{n}_{\max} - 1\} \to \{bN_c + \bar{n}_{\max} - 1 - i'\}),$$
(D.4)

which has the same form as Eq. (D.1), but the rank of the corresponding interpolating tensors is reduced by one. We thus compute the sum in Eq. (IV.23) for representations r (with  $H_1 = 0$ ) interpolated by tensors  $T_{\{\bar{n}_{\max}-2\}}^{\{bN_c+\bar{n}_{\max}-2\}}$ . By making use of Eq. (D.1) with i = 2 and Eq. (D.4) with i' = 1 we find

$$\sum_{r'} \ell(r')k(r' \to r)$$

$$= \ell(U[r]) + \sum_{\{\bar{n}_{\max}\}} k\left(\{\bar{n}_{\max}\} \to \{\bar{n}_{\max} - 2\}\right)$$

$$\times \underbrace{\sum_{\{bN_c + \bar{n}_{\max}\}} \ell(r') k\left(\{bN_c + \bar{n}_{\max}\} \to \{bN_c + \bar{n}_{\max} - 2\}\right)}_{=0}$$

$$+ \sum_{\{\bar{n}_{\max} - 1\}} k\left(\{\bar{n}_{\max} - 1\} \to \{\bar{n}_{\max} - 2\}\right)$$

$$\times \underbrace{\sum_{\{bN_c + \bar{n}_{\max} - 1\}} \ell(r') k\left(\{bN_c + \bar{n}_{\max} - 1\} \to \{bN_c + \bar{n}_{\max} - 2\}\right)}_{=0}$$

$$= \ell(U[r]) ,$$
(D.5)

which verifies Eq. (IV.23).

We have thus proven Eq. (IV.23) for irreps r interpolated by tensors with the highest rank  $T_{\{\bar{n}_{\max}\}}^{\{bN_c+\bar{n}_{\max}-1\}}$ , the second highest rank  $T_{\{\bar{n}_{\max}-1\}}^{\{bN_c+\bar{n}_{\max}-1\}}$ , and the third highest rank  $T_{\{\bar{n}_{\max}-2\}}^{\{bN_c+\bar{n}_{\max}-2\}}$ . By repeating the same steps, one can prove Eq. (IV.23) for irreps interpolated by the tensors of any lower rank.

- Gerard 't Hooft, "Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking," NATO Sci. Ser. B 59, 135–157 (1980).
- [2] Y. Frishman, A. Schwimmer, Tom Banks, and S. Yankielowicz, "The Axial Anomaly and the Bound State Spectrum in Confining Theories," Nucl. Phys. B 177, 157–171 (1981).
- [3] Sidney R. Coleman and Bernard Grossman, "'t Hooft's Consistency Condition as a Consequence of Analyticity and Unitarity," Nucl. Phys. B 203, 205–220 (1982).
- [4] Sidney R. Coleman and Edward Witten, "Chiral Symmetry Breakdown in Large N Chromodynamics," Phys. Rev. Lett. 45, 100 (1980).
- [5] John Preskill and Steven Weinberg, "Decoupling' Constraints On Massless Composite Particles," Phys. Rev. D 24, 1059 (1981).
- [6] C. Vafa and Edward Witten, "Restrictions on Symmetry Breaking in Vector-Like Gauge Theories," Nucl. Phys. B 234, 173–188 (1984).
- [7] Luca Ciambriello, Roberto Contino, Andrea Luzio, Marcello Romano, and Ling-Xiao Xu, "A Novel Strategy to Prove Chiral Symmetry Breaking Through in QCD-like Theories," (), work in progress.
- [8] A. Schwimmer, "Anomaly Equations and Superalgebras," Nucl. Phys. B 198, 269–279 (1982).

- [9] Glennys R. Farrar, "Anomaly Nonmatching in Composite Models," Phys. Lett. B 96, 273–275 (1980).
- [10] Seiichiro Takeshita, Hiromasa Komatsu, Akira Kakuto, and Kenzo Inoue, "Variety of Solutions of 't Hooft's Metacolor," Prog. Theor. Phys. 66, 2221 (1981).
- [11] Romesh K. Kaul and R. Rajaraman, "Derivation of the Decoupling Relations for Composite Models From Anomaly Constraints and N Independence," Phys. Lett. B 110, 385–390 (1982).
- [12] E. Cohen and Y. Frishman, "Anomaly Equations and the Persistent Mass Condition," Phys. Lett. B 109, 35–39 (1982).
- [13] Itzhak Bars, "Spontaneous Chiral Symmetry Breaking in Quantum Chromodynamics," Phys. Lett. B 109, 73–78 (1982).
- [14] Luca Ciambriello, Roberto Contino, and Ling-Xiao Xu, "On the Proof of Chiral Symmetry Breaking through Anomaly Matching in QCD-like Theories: An Exemplification," (), work in progress.
- [15] Don Weingarten, "Mass Inequalities for QCD," Phys. Rev. Lett. 51, 1830 (1983).
- [16] Shmuel Nussinov and Melissa A. Lampert, "QCD inequalities," Phys. Rept. 362, 193–301 (2002), arXiv:hep-ph/9911532.
- [17] Nathan Seiberg, "Exact results on the space of vacua of four-dimensional SUSY gauge theories," Phys. Rev. D 49, 6857–6863 (1994), arXiv:hep-th/9402044.
- [18] N. Seiberg, "Electric magnetic duality in supersymmetric nonAbelian gauge theories," Nucl. Phys. B 435, 129–146 (1995), arXiv:hep-th/9411149.
- [19] Ofer Aharony, Jacob Sonnenschein, Michael E. Peskin, and Shimon Yankielowicz, "Exotic nonsupersymmetric gauge dynamics from supersymmetric QCD," Phys. Rev. D 52, 6157–6174 (1995), arXiv:hepth/9507013.
- [20] Hsin-Chia Cheng and Yael Shadmi, "Duality in the presence of supersymmetry breaking," Nucl. Phys. B 531, 125–150 (1998), arXiv:hep-th/9801146.
- [21] Nima Arkani-Hamed and Riccardo Rattazzi, "Exact results for nonholomorphic masses in softly broken supersymmetric gauge theories," Phys. Lett. B 454, 290–296 (1999), arXiv:hep-th/9804068.
- [22] Markus A. Luty and Riccardo Rattazzi, "Soft supersymmetry breaking in deformed moduli spaces, conformal theories, and N=2 Yang-Mills theory," JHEP 11, 001 (1999), arXiv:hep-th/9908085.
- [23] Steven Abel, Matthew Buican, and Zohar Komargodski, "Mapping Anomalous Currents in Supersymmetric Dualities," Phys. Rev. D 84, 045005 (2011), arXiv:1105.2885 [hep-th].
- [24] Hitoshi Murayama, "Some Exact Results in QCD-like Theories," Phys. Rev. Lett. 126, 251601 (2021), arXiv:2104.01179 [hep-th].
- [25] Andrea Luzio and Ling-Xiao Xu, "On the derivation of chiral symmetry breaking in QCD-like theories and S-confining theories," JHEP 08, 016 (2022), arXiv:2202.01239 [hep-th].
- [26] Michael Dine and Yan Yu, "Challenges to Obtaining Results for Real QCD from SUSY QCD," (2022), arXiv:2205.00115 [hep-ph].
- [27] Steven Weinberg and Edward Witten, "Limits on Massless Particles," Phys. Lett. B 96, 59–62 (1980).
- [28] W. K. Tung, Group Theory in Physics (1985).

- [29] Steven Weinberg, The quantum theory of fields. Vol. 2: Modern applications (Cambridge University Press, 2013).
- [30] Cumrun Vafa and Edward Witten, "Parity Conservation in QCD," Phys. Rev. Lett. 53, 535 (1984).
- [31] R. Aloisio, V. Azcoiti, G. Di Carlo, A. Galante, and A. F. Grillo, "Probability distribution function of the diquark condensate in two colors QCD," Nucl. Phys. B 606, 322–336 (2001), arXiv:hep-lat/0011079.
- [32] V. Azcoiti, G. diCarlo, E. Follana, and A. Vaquero, "Parity Realization in Lattice QCD with Ginsparg-Wilson Fermions," JHEP 07, 047 (2010), arXiv:1004.3463 [hep-lat].
- [33] Matteo Giordano, "Impossibility of spontaneous vector flavor symmetry breaking on the lattice," Phys. Rev. D 107, 114509 (2023), arXiv:2303.03109 [hep-lat].
- [34] Davide Gaiotto, Anton Kapustin, Nathan Seiberg, and Brian Willett, "Generalized Global Symmetries," JHEP 02, 172 (2015), arXiv:1412.5148 [hep-th].
- [35] Davide Gaiotto, Anton Kapustin, Zohar Komargodski, and Nathan Seiberg, "Theta, Time Reversal, and Temperature," JHEP 05, 091 (2017), arXiv:1703.00501 [hep-th].
- [36] Davide Gaiotto, Zohar Komargodski, and Nathan Seiberg, "Time-reversal breaking in QCD<sub>4</sub>, walls, and dualities in 2 + 1 dimensions," JHEP 01, 110 (2018), arXiv:1708.06806 [hep-th].
- [37] Clay Córdova, Daniel S. Freed, Ho Tat Lam, and Nathan Seiberg, "Anomalies in the Space of Coupling Constants and Their Dynamical Applications I," SciPost Phys. 8, 001 (2020), arXiv:1905.09315 [hepth].
- [38] Clay Córdova, Daniel S. Freed, Ho Tat Lam, and Nathan Seiberg, "Anomalies in the Space of Coupling Constants and Their Dynamical Applications II," SciPost Phys. 8, 002 (2020), arXiv:1905.13361 [hepth].
- [39] Yuya Tanizaki, Yuta Kikuchi, Tatsuhiro Misumi, and Norisuke Sakai, "Anomaly matching for the phase diagram of massless  $\mathbb{Z}_N$ -QCD," Phys. Rev. D **97**, 054012 (2018), arXiv:1711.10487 [hep-th].
- [40] Yuya Tanizaki, "Anomaly constraint on massless QCD and the role of Skyrmions in chiral symmetry breaking," JHEP 08, 171 (2018), arXiv:1807.07666 [hep-th].
- [41] Mohamed M. Anber and Erich Poppitz, "On the baryon-color-flavor (BCF) anomaly in vector-like theories," JHEP 11, 063 (2019), arXiv:1909.09027 [hep-th].
- [42] Stefano Bolognesi, Kenichi Konishi, and Andrea Luzio, "Anomalies and phases of strongly-coupled chiral gauge theories: recent developments," (2021), arXiv:2110.02104 [hep-th].
- [43] Zohar Komargodski and Adam Schwimmer, "On Renormalization Group Flows in Four Dimensions," JHEP 12, 099 (2011), arXiv:1107.3987 [hep-th].
- [44] Thomas Appelquist, Andrew G. Cohen, and Martin Schmaltz, "A New constraint on strongly coupled gauge theories," Phys. Rev. D 60, 045003 (1999), arXiv:hep-th/9901109.
- [45] Howard Georgi, Lie algebras in particle physics, 2nd ed., Vol. 54 (Perseus Books, Reading, MA, 1999).