

Lattice QCD at non-zero temperature and density

F. Karsch

Fakultät für Physik, Universität Bielefeld, D-33615 Bielefeld, Germany

Abstract

Soon after the formulation of Quantum Chromodynamics in 1972 its regularization on Euclidean space-time lattices had been introduced by Kenneth Wilson [1]. This paved ground for numerical studies of non-perturbative aspects of QCD, first shown by Michael Creutz [2]. Ever since these first lattice QCD calculations the exploration of the QCD phase diagram and the thermodynamics of strong-interaction matter at non-zero temperature and density was pursued vigorously. In this brief review I try to highlight some of the results on QCD thermodynamics obtain during the last 42 years through lattice QCD calculations.

Article to appear in a special EPJC Volume in celebration of
'50 Years of Quantum Chromodynamics'
 edited by Franz Gross and Eberhard Klempt

1. QCD thermodynamics on Euclidean Lattices

The path integral formulation of QCD can easily be applied to cases of non-vanishing temperature (T) and other external control parameters, e.g. the chemical potentials (μ_f) that couple to the conserved currents of quark-flavor number.

Using the lattice regularization scheme of QCD, introduced by K. Wilson [1], QCD thermodynamics is formulated on Euclidean space-time lattices of size $N_\sigma^3 N_\tau$ where, for a given lattice spacing (a), the lattice extent in Euclidean time controls the temperature $T = 1/N_\tau a$ and the spatial extent is related to the volume of the thermodynamic system, $V = (N_\sigma a)^3$. The chemical potentials enter directly in the fermion matrices, M_f , which arise from the QCD Lagrangian after integrating out the fermion fields.

Bulk thermodynamics can then be derived from the lattice regularized partition function,

$$Z = \int \prod_{x_0=1}^{N_\tau} \prod_{x_i=1}^{N_\sigma} \prod_{\nu \hat{=} 0}^3 \mathcal{D}U_{x,\hat{\nu}} e^{-S_G} \prod_{f=u,d,s..} \det M_f(m_f, \mu_f) , \quad (1)$$

where $x = (x_0, \vec{x})$ labels the sites of the 4-dimensional lattice, S_G denotes the gluonic part of the Euclidean action, which is expressed in terms of $SU(3)$ matrices $U_{x,\hat{\nu}}$ and M_f is the fermion matrix for quark flavor f . It is a function of quark mass, m_f and flavor chemical potential $\hat{\mu}_f \equiv \mu_f/T$. Basic bulk thermodynamic observables (equation of state, susceptibilities, etc.) can then be obtained from the logarithm of the partition function, Z , which defines the pressure, P , as

$$P/T = \frac{1}{V} \ln Z(T, V, \vec{\mu}, \vec{m}) . \quad (2)$$

Applying standard thermodynamic relations one obtains other observables of interest; e.g. the energy density is related to the trace anomaly of the energy-momentum tensor, $\Theta^{\mu\mu}$,

$$\frac{\Theta^{\mu\mu}}{T^4} = \frac{\epsilon - 3P}{T^4} \equiv T \frac{\partial P/T^4}{\partial T} , \quad (3)$$

and the conserved charge densities are obtained as,

$$\frac{n_X}{T^3} = \frac{\partial P/T^4}{\partial \hat{\mu}_X} , \quad X = B, Q, S . \quad (4)$$

While the framework of lattice QCD provides easy access to QCD thermodynamics at vanishing values of the chemical potentials, major difficulties arise at $\mu_f \neq 0$. The fermion determinants, $\det M_f(m_f, \mu_f)$, are no longer positive definite when the real part of the chemical potential is non-zero, $\text{Re} \hat{\mu}_f \neq 0$. This includes the physically relevant case of strictly real chemical potentials. The presence of a complex valued integrand in the path integral makes the application of standard Monte Carlo techniques, which rely on a probabilistic interpretation of integration measures, impossible. The two most common approaches to circumvent this problem are to either (i) perform numerical calculations at imaginary values of the chemical potential, $\hat{\mu}_f^2 < 0$ [3, 4], or to (ii) perform Taylor series expansions around $\hat{\mu}_f = 0$ [5, 6].

In the former case numerical results need to be analytically continued to real values of μ_f . In the latter case the QCD partition function is written as,

$$P/T^4 = \frac{1}{VT^3} \ln Z(T, V, \vec{\mu}) = \sum_{i,j,k=0}^{\infty} \frac{\chi_{ijk}^{BQS}}{i!j!k!} \hat{\mu}_B^i \hat{\mu}_Q^j \hat{\mu}_S^k, \quad (5)$$

with $\chi_{000}^{BQS} \equiv P(T, V, \vec{0})/T^4$ and expansion coefficients,

$$\chi_{ijk}^{BQS}(T) = \left. \frac{\partial P/T^4}{\partial \hat{\mu}_B^i \partial \hat{\mu}_Q^j \partial \hat{\mu}_S^k} \right|_{\hat{\mu}=0}, \quad (6)$$

can be determined in Monte Carlo simulations performed at $\hat{\mu}_X = 0$.

The phase structure of QCD can be explored using suitable observables that are sensitive to the spontaneous breaking and the eventual restoration of global symmetries. They can act as order parameters in certain limits of the parameter space spanned by the quark masses. In QCD exact symmetries exist either in the chiral limit, *i.e.* at vanishing values of n_f quark masses, or for infinitely heavy quarks, *i.e.* in pure $SU(N_c)$ gauge theories, with N_c denoting the number of colors.

In order to probe the restoration of the global chiral symmetries one analyzes the chiral condensate and its susceptibilities,

$$\langle \bar{\chi} \chi \rangle_f = \frac{T}{V} \frac{\partial}{\partial m_f} \ln Z = \frac{T}{V} \langle \text{Tr} M_f^{-1} \rangle, \quad (7)$$

$$\chi_m^{fg} = \frac{\partial \langle \bar{\chi} \chi \rangle_f}{\partial m_g}, \quad \chi_t^f = T \frac{\partial \langle \bar{\chi} \chi \rangle_f}{\partial T}. \quad (8)$$

The former is an order parameter for the restoration of the $SU(n_f)_L \times SU(n_f)_R$ chiral flavor symmetry of QCD and distinguishes, in the limit of vanishing quark masses, a symmetry broken phase at low temperature from a chiral symmetry restored phase at high temperature,

$$\lim_{m_\ell \rightarrow 0} \langle \bar{\chi} \chi \rangle_\ell \begin{cases} > 0 & , \quad T < T_\chi \\ = 0 & , \quad T \geq T_\chi \end{cases}. \quad (9)$$

Similarly one considers the Polyakov loop $\langle L \rangle$ and its susceptibility χ_L ,

$$\begin{aligned}
\langle L \rangle &= \frac{1}{N_\sigma^3} \langle \sum_{\vec{x}} \text{Tr} L_{\vec{x}} \rangle, & L_{\vec{x}} &= \prod_{x_0=1}^{N_\tau} U_{(x_0, \vec{x}), \hat{0}}, \\
\chi_L &= N_\sigma^3 (\langle L^2 \rangle - \langle L \rangle^2), & &
\end{aligned} \tag{10}$$

to probe the breaking and restoration of the global $Z(N_c)$ center symmetry of pure $SU(N_c)$ gauge theories; *i.e.* $SU(N_c)$ gauge theories at finite temperature, formulated on Euclidean lattices, are invariant under global rotation of all temporal gauge field variables, $U_{\vec{x}, \hat{0}} \rightarrow z U_{\vec{x}, \hat{0}}$, with $z \in Z(N_c)$. The Polyakov loop expectation value vanishes as long as this center symmetry is not spontaneously broken.

The Polyakov loop expectation value also reflects the long distance behavior of Polyakov loop correlation functions,

$$|\langle L \rangle|^2 \equiv \lim_{|\vec{x}| \rightarrow \infty} G_L(\vec{x}) \begin{cases} = 0 \Leftrightarrow F_q = \infty, T \leq T_d \\ > 0 \Leftrightarrow F_q < \infty, T > T_d \end{cases} \tag{11}$$

where

$$G_L(\vec{x}) = e^{-F_{\bar{q}q}(\vec{x}, T)} = \langle \text{Tr} L_{\vec{0}} \text{Tr} L_{\vec{0}}^\dagger \rangle \tag{12}$$

is the correlation function of two Polyakov loops. It denotes the change in free energy (excess free energy, $F_{\bar{q}q}$), that is due to the presence of two static quark sources introduced in a thermal medium. At zero temperature this free energy reduces to the potential between static quark sources.

At least in the case of pure gauge theories this provides a connection between the confinement-deconfinement phase transition and the breaking of a global symmetry, the $Z(N_c)$ center symmetry of the $SU(N_c)$ gauge group. This symmetry, however, is explicitly broken in the presence of dynamical quarks with mass $m_f < \infty$. Unlike chiral symmetry restoration, deconfinement thus is not expected to be related to a phase transition in QCD with physical quark masses. Nonetheless, the consequences of deconfinement, related to the dissolution of hadronic bound states, becomes clearly visible in many thermodynamic observables.

2. Early lattice QCD calculations at non-zero temperature

Almost immediately after the formulation of QCD as the theory of strong interaction physics, its consequences for strong interaction matter at non-zero temperature were examined [10, 11]. It rapidly became obvious that




Figure 1: First evidence for the existence of a deconfinement phase transition in $SU(2)$ gauge theories using the Polyakov loop expectation value as an order parameter (left) [7] and a first extrapolation of the phase transition temperature to the continuum limit (middle) [8]. The right hand figure shows a first comparison of the temperature dependence of the Polyakov loop ($W \equiv \langle |L| \rangle$) and chiral condensate ($\langle \bar{\psi}\psi \rangle$) order parameters in a $SU(3)$ gauge theory [9].

fundamental properties of QCD, confinement and asymptotic freedom on the one hand [11, 12], and chiral symmetry breaking on the other hand [13], are likely to trigger a phase transition in strong interaction matter that separates a phase being dominated by hadrons as the relevant degrees of freedom from that of almost free quarks and gluons. The notion of a quark-gluon plasma was coined at that time [14].

Soon after these early, conceptually important developments it was realized that the formulation of QCD on discrete space-time lattices, which was introduced by K. Wilson as a regularization scheme in QCD [1], also provides a powerful framework for the analysis of non-perturbative properties of strong interaction matter through Monte-Carlo simulations [2]. This led to a first determination of a phase transition temperature in $SU(2)$ [7, 8] and $SU(3)$ [9, 15, 16] gauge theories, and a first determination of the equation of state of purely gluonic matter [17, 18]. The interplay between deconfinement on the one hand and chiral symmetry restoration on the other hand also was studied [9] early on and the question whether or not these two aspects of QCD may lead to two distinct phase transitions in QCD has been considered ever since. Some results from these first lattice QCD studies of the thermodynamics of strong interaction matter are shown in Fig. 1.

At physical values of the quark masses, neither deconfinement nor the effective restoration of chiral symmetry leads to a true phase transition. Still the transition from the low temperature hadronic to the high temperature




Figure 2: The so-called color averaged, heavy quark free energy ($F_{av} \equiv F_{\bar{q}q}$) in the vicinity of the pseudo-critical transition temperature (T_{pc}) in 2-flavor QCD [19]. Results shown cover a temperature range from $T/T_{pc} \simeq 0.75$ to $T/T_{pc} \simeq 2$.




Figure 3: Quark mass dependence of chiral order parameter, M , defined in Eq. 14 for QCD with two degenerate light quark masses and a strange quark mass tuned to its physical value. Shown are results from calculations on lattices with temporal extent $N_\tau = 8$ performed for several values of the light quark masses [20, 21]. The light quark masses, m_ℓ , are expressed in units of the strange quark mass, $H = m_\ell/m_s$. In the figure we give $1/H = m_s/m_\ell$ together with the corresponding values of the Goldstone pion mass.

partonic phase of QCD is clearly visible in the pseudo-critical behavior of the heavy quark free energy and the chiral condensate respectively. Some recent results on these observables, obtained in simulations of QCD with light, dynamical quark degrees of freedom, are shown in Figs. 2 and 3.

3. Global symmetries and the QCD phase diagram

The early studies of QCD thermodynamics made it clear that universality arguments and renormalization group techniques, successfully developed in condensed matter physics and applied in statistical physics to the analysis of phase transitions, also can be carried over to the analysis of the phase structure of quantum field theories [22, 23]. The renormalization group based arguments for the existence of a second order phase transition in the universality class of the 3-d Ising model in a $SU(2)$ gauge theory, and a first order transition for the $SU(3)$ color group of QCD [24] have been confirmed by detailed lattice QCD calculations [25, 26].

In the presence of n_f light, dynamical quarks, distinguished by a flavor quantum number, it is the chiral symmetry of QCD that triggers the occurrence of phase transitions [13]. In addition to a global $U(1)$ symmetry that reflects the conservation of baryon number and is unbroken at all temperatures and densities, the massless QCD Lagrangian is invariant under the symmetry group

$$U(1)_A \times SU(n_f)_L \times SU(n_f)_R . \quad (13)$$

The $SU(n_f)_L \times SU(n_f)_R$ symmetry corresponds to chiral rotations of n_f massless quark fields in flavor space. This symmetry is spontaneously broken at low temperatures, giving rise to $n_f^2 - 1$ massless Goldstone modes, which for $n_f = 2$ are the three light pions of QCD. They have a non-vanishing mass only because of the explicit breaking of chiral symmetry by a mass term in the QCD Lagrangian that couples to the chiral order parameter field $\bar{\chi}_f \chi_f$. The axial $U(1)_A$ group corresponds to global rotations of quark fields for a given flavor f . Although it is an exact symmetry of the classical Lagrangian, it is explicitly broken in the quantized theory. This explicit breaking of a global symmetry, arising from fluctuations on the quantum level, is known as the $U(1)_A$ anomaly.

The renormalization group based analysis of the chiral phase transition, performed by Pisarski and Wilczek [13], made it clear that the chiral phase transition is sensitive to the number of light quark flavors that become massless. Furthermore, it has been argued in [13] that the order of the transition may be sensitive to the magnitude of the axial anomaly at non-zero temperature, which is closely related to the temperature dependence of topological non-trivial field configurations.

Although it was generally expected that the chiral phase transition in 3-flavor QCD becomes a first order phase transition in the chiral limit [13], there is currently no direct evidence for this from lattice QCD calculations. In fact, the current understanding is that the chiral phase transition is second order for all $n_f \leq 6$ [27].

In Fig. 4 (left) we show the original version of the QCD phase diagram in the plane of two degenerate light (m_ℓ) and strange (m_s) quark masses, proposed in 1990 [28], together with an updated version from 2021 [27]. Here m_ℓ denotes the two degenerate up and down quark masses, $m_\ell \equiv m_u = m_d$. This sketch of our current understanding of the 3-flavor phase diagram also is supported by the increasing evidence for a non-singular crossover transition in QCD with physical light and strange quark masses and the absence of




Figure 4: Sketch of the phase diagram of QCD in the plane of degenerate, light up and down quark masses and a strange quark mass (Columbia plot). The figure shows the original version from 1990 [28] (left) and an updated version from 2021 [27] (right).

any evidence for a first order phase transition at lighter-than-physical values of the light and strange quark masses [27, 29]. In the chiral limit, *i.e.* for vanishing up and down quark masses¹, a second order phase transition will then occur.

4. The chiral phase transition at vanishing chemical potential

The occurrence of the chiral phase transition is signaled by the vanishing of the light quark chiral condensate. In order to remove multiplicative and additive divergences in $\langle \bar{\chi}\chi \rangle_\ell$ one considers instead the order parameter M which is a combination of light and strange quark condensates,

$$M = 2 (m_s \langle \bar{\psi}\psi \rangle_\ell - m_\ell \langle \bar{\psi}\psi \rangle_s) / f_K^4, \quad (14)$$

and its derivative with respect to the light quark masses, *i.e.* the chiral susceptibility χ_M

$$\chi_M = m_s \left(\frac{\partial M}{\partial m_u} + \frac{\partial M}{\partial m_d} \right)_{m_u=m_d \equiv m_\ell}. \quad (15)$$

¹Lattice QCD studies of the (2+1)-flavor phase diagram generally are performed with degenerate up and down quark masses.

Here the kaon decay constant $f_K = 156.1(9)/\sqrt{2}$ MeV, has been used to introduce a dimensionless order parameter. The scaling behavior of M and χ_M , have been used to characterize the chiral phase transition,

$$M \quad \widetilde{m}_\ell \rightarrow 0 \quad \begin{cases} A \left(\frac{T_c^0 - T}{T_c^0} \right)^\beta & , \quad T < T_c^0 \\ 0 & , \quad T \geq T_c^0 \end{cases} \quad (16)$$

$$\chi_M \quad \widetilde{m}_\ell \rightarrow 0 \quad \begin{cases} \infty & , \quad T \leq T_c^0 \\ C \left(\frac{T - T_c^0}{T_c^0} \right)^{-\gamma} & , \quad T > T_c^0 \end{cases} \quad (17)$$

where β , γ are critical exponents.

We note that the low temperature behavior of the order parameter susceptibility, χ_M , is quite different from that known, for instance, in the 3- d Ising model. The susceptibility diverges in the massless limit at all values of the temperature, $T \leq T_c^0$. This is a consequence of the breaking of a continuous rather than a discrete symmetry. The former gives rise to Goldstone modes, the pions in QCD, which contribute to the chiral condensate and as such to the order parameter M , *i.e.*,

$$M \sim a(T) \sqrt{m_\ell} \quad , \quad T < T_c^0 \quad . \quad (18)$$

As a consequence the chiral susceptibility diverges below T_c^0 , $\chi_M \sim 1/\sqrt{m_\ell}$, while at T_c^0 its divergence is controlled by the critical exponent $\delta = 1 + \gamma/\beta$,

$$\chi_M \sim \begin{cases} H^{-1/2} & , \quad T < T_\chi \\ H^{1/\delta-1} & , \quad T = T_\chi \end{cases} \quad (19)$$

with $H = m_\ell/m_s$. As $1 - 1/\delta > 1/2$ in all relevant universality classes χ_M develops a pronounced peak at small, but non-zero values of the quark masses,

$$\chi_M^{peak} \equiv \chi_M(T_{pc}(H)) \sim H^{1/\delta-1} \quad , \quad H = m_\ell/m_s \quad . \quad (20)$$

The location of such a peak in either χ_M or similarly in $T\partial M/\partial T$, defines pseudo-critical temperatures, $T_{pc}(H)$, which converge to the unique chiral phase transition, T_c^0 , at $H = 0$. Some results on the quark mass dependence of M and χ_M are shown in Figs. 3 and 5, respectively. A scaling analysis of




Figure 5: same as Fig. 3 but for the chiral susceptibility.

these observables, performed in [20], led to the determination of the chiral phase transition temperature [20],

$$T_c^0 = 132_{-6}^{+3} \text{ MeV} . \quad (21)$$

Similar results have also been obtained in [30] where a quite different discretization scheme for the fermion sector of QCD has been used.

For physical light and strange quark masses, corresponding to $H \simeq 1/27$, one finds as a pseudo-critical temperature [31],

$$T_{pc} = 156.5(1.5) \text{ MeV} , \quad (22)$$

which is in good agreement with other determinations of pseudo-critical temperatures in $(2+1)$ -flavor QCD [32, 33, 34].

The chiral symmetry group $SU(2)_L \times SU(2)_R$ is isomorphic to the rotation group $O(4)$. It thus is expected that the chiral phase transition for two vanishing light quark masses is in the same universality class as 3- d , $O(4)$ symmetric spin models. In fact, the rapid rise of χ_M , shown in Fig. 5, is consistent with a critical exponent in this universality class, $\delta = 4.824$ [35]. However, a precise determination of this exponent in 2-flavor QCD is not yet possible. This leaves open the possibility for other symmetry breaking patterns and other universality classes playing a role in the chiral limit of 2-flavor QCD [36]. In fact, the discussion of such possibilities is closely related to the yet unsettled question concerning the influence of the axial $U(1)_A$ symmetry on the chiral phase transition. For a recent review on this question see, for instance [37].




Figure 6: Temperature dependence of masses of parity partners in the baryon octet [38].

Thermal masses and screening masses: The restoration of symmetries is reflected also in the modification of the hadron spectrum at non-zero temperature. Interactions in a thermal medium lead to modifications of resonance peaks that can modify the location of maxima and the width of spectral functions that control properties of hadron correlation functions. This gives rise to so-called thermal masses as well as thermal screening masses that control the long-distance behavior of hadron correlation functions in Euclidean time and spatial directions, respectively.

A consequence of $U(1)_A$ breaking in the vacuum or at low temperature is that masses of hadronic states that are related to each other through a $U(1)_A$ transformation differ, while they become identical, or close to each other, when the $U(1)_A$ symmetry is effectively restored. This is easily seen to happen at high temperature. The crucial question, of relevance for the QCD phase transition, however, is to which extent $U(1)_A$ symmetry breaking is reduced, or already disappeared at the chiral phase transition temperature. Settling this question requires the analysis of observables sensitive to $U(1)_A$ breaking close to T_c^0 and for smaller-than-physical light quark masses.

The calculation of in-medium modifications of hadron masses is difficult, but has been attempted for quark masses close to their physical values [38]. Results for the temperature dependence of the mass-splitting of parity partners in the baryon octet [38] are shown in Fig. 6. These results suggest a strong temperature dependence of the negative parity states while the positive parity partners are not sensitive to temperature changes. At T_{pc} the masses of parity partners are almost degenerate.




Figure 7: Screening masses (left) and the related susceptibilities (right) of scalar and pseudo-scalar mesons [39, 40].

More easily accessible are so-called screening masses, which also are obtained from ordinary hadron correlation functions and can be analyzed close to the chiral limit. Rather than analyzing the long-distance behavior of hadron correlation functions in Euclidean time, one extracts a so-called screening mass from the long-distance behavior in one of the spatial directions [41, 42]. Finite temperature meson screening correlators, projected onto lowest Matsubara frequency of a bosonic state, $p_0 \equiv \omega_0 = 0$, and zero transverse momentum, $\mathbf{p}_\perp \equiv (p_x, p_y) = 0$, are defined by

$$G_\Gamma(z, T) = \int_0^\beta d\tau \int dxdy \left\langle \mathcal{M}_\Gamma(\vec{r}, \tau) \overline{\mathcal{M}_\Gamma}(\vec{0}, 0) \right\rangle$$

$$z \xrightarrow{\sim} \infty \quad e^{-m_\Gamma(T)z} \quad , \quad \vec{r} \equiv (x, y, z) \quad , \quad (23)$$

where $\mathcal{M}_\Gamma \equiv \bar{\psi}\Gamma\psi$ is a meson operator that projects onto a quantum number channel that is selected through an appropriate choice of Γ -matrices [39, 41]. At large distances this permits the extraction of the screening mass, m_Γ , in the quantum number channel selected by Γ from the exponential fall-off of these correlation functions. In Fig. 7 (left) we show results for the scalar and pseudo-scalar screening masses obtained in $(2+1)$ -flavor QCD calculations for different values of the light to strange quark mass ratio. The integrated correlation functions define susceptibilities in these quantum number channels, which also should be degenerate, if $U(1)_A$ is effectively restored. Both observables seem to suggest that there remains a significant remnant of $U(1)_A$ breaking at the chiral phase transition temperature, T_c^0 , which however reduces quickly above the chiral transition and gives rise to an effective restoration of $U(1)_A$ at $T \simeq 1.1T_c^0$.




Figure 8: Eigenvalue density of the overlap fermion matrix obtained in calculations with dynamical overlap fermions [43].

In the region $T > T_c^0$ the difference between pseudo-scalar and scalar susceptibilities is related to the so-called disconnected part, χ_{dis} , of the chiral susceptibility, $\chi_M = \chi_{dis} + \chi_{con}$, with

$$\chi_{dis} = \frac{1}{4N_\tau N_\sigma^3} (\langle (\text{Tr} M_\ell^{-1})^2 \rangle - \langle \text{Tr} M_\ell^{-1} \rangle^2) , \quad (24)$$

$$\chi_{con} = \frac{1}{2N_\tau N_\sigma^3} \langle \text{Tr} M_\ell^{-2} \rangle . \quad (25)$$

The disconnected chiral susceptibility can be expressed by an integral over the eigenvalue density, $\rho(\lambda)$, of the fermion matrix M_f ,

$$\chi_{dis} = \int_0^\infty d\lambda \rho(\lambda) \frac{2m_\ell^2}{(\lambda^2 + m_\ell^2)^2} . \quad (26)$$

In the chiral symmetric phase the density of vanishing eigenvalues, $\rho(0)$, vanishes. In order for χ_{dis} to be nonetheless non-zero in the chiral limit, the density of near-zero eigenvalues needs to converge to a non-vanishing value (δ -function) at $\lambda = 0$ in the limit $m_\ell \rightarrow 0$ and $V \rightarrow \infty$. Controlling the various limits involved and also taking into account that the pseudo-critical transition temperature, $T_{pc}(H)$, has a sizeable quark mass dependence is difficult. Nonetheless, studies of the temperature dependence of the eigenvalue

density of the Dirac matrix are crucial for a detailed understanding of the influence of the $U(1)_A$ anomaly on the QCD phase transition. Not surprisingly, it turns out that at non-zero values of the lattice spacing the spectrum of low lying eigenvalues is quite sensitive to the fermion discretization scheme. Using fermions with good chirality even at non-zero lattice spacing seems to be advantageous, although after having performed the extrapolation to the chiral limit, they should lead to results identical with those obtained, e.g. within the staggered discretization scheme. Current results are ambiguous. We show in Fig. 8 results from a calculation of eigenvalue distributions obtained from calculations with dynamical overlap fermions [43, 44]. These calculations provide evidence for a large density of near-zero eigenvalues and a non-zero eigenvalue density, possibly building up at $\lambda = 0$. This is in contrast to calculations performed with domain wall fermions [45] as well as so-called partially quenched calculations that use the overlap fermion operator to calculate eigenvalue distributions on gauge field configurations generated with dynamical staggered fermions [46]. Obviously this subtle aspect of the chiral phase transition is not yet resolved and the analysis of $U(1)_A$ restoration will remain to be a central topic in finite temperature QCD in the years to come.

5. The chiral phase transition at non-vanishing chemical potential

In the studies of QCD at non-vanishing baryon chemical potential the search for the existence of a second order phase transition at physical values of the quark masses, the critical end point (CEP), finds particular attention. It separates the crossover regime at small values of the chemical potential from a region of first order phase transitions, which is predicted in many phenomenological models to exist at high density. The CEP is searched for extensively in heavy ion experiments and, if confirmed, would provide a solid prediction for the existence of first order phase transitions in dense stellar matter, e.g. in neutron stars.

The dependence of the transition temperature on the chemical potentials, e.g. $T_{pc}(\mu_B)$, can be deduced from the μ_B -dependent shift of the peak in the chiral susceptibility. At non-vanishing values of the baryon chemical potential, μ_B , the QCD phase transition temperature in the chiral limit as well as the region of pseudo-critical behavior in QCD with its physical quark mass values shifts to smaller values of the temperature. This shift has been determined in calculations with imaginary values of the chemical potentials as well as from Taylor series expansions of the order parameter M and its




Figure 9: Sketch of a possible QCD phase diagram in the space of temperature (T), baryon chemical potential (μ_B) and light quark masses ($m_{u,d}$).

susceptibility χ_M . Using a Taylor series ansatz for $T_{pc}(\mu_B)$,

$$T_c(\mu_B) = T_c^0 \left(1 - \kappa_2^B \left(\frac{\mu_B}{T_c^0} \right)^2 - \kappa_4^B \left(\frac{\mu_B}{T_c^0} \right)^4 \right) \quad (27)$$

one finds for the curvature coefficients $\kappa_2^B \simeq 0.012$ while the next correction is consistent with zero in all current studies, e.g. $\kappa_4^B = 0.00032(67)$ [34]. The pseudo-critical temperature T_{pc} at physical values of the light and strange quark masses thus drops to about 150 MeV at $\mu_B \simeq 2T_{pc}$. This is still considerably larger than the chiral phase transition temperature, T_c^0 , determined at $\mu_B = 0$. As various model calculations [47, 48] suggest that the CEP at non-zero μ_B is located at a temperature below T_c^0 one thus needs to get access to thermodynamics at large chemical potentials. Assuming that the curvature of the pseudo-critical line does not change drastically at large values of the chemical potentials, our current understanding of the QCD phase diagram in the m_ℓ - T - μ_B space (see Fig. 9) suggests that a possible CEP in the phase diagram may exist only at a temperature,

$$T^{CEP}(\mu_B^{CEP}) < 130 \text{ MeV} , \mu_B^{CEP} > 400 \text{ MeV} . \quad (28)$$

Reaching the region $\mu_B/T > 3$ is a major challenge for any of the currently




Figure 10: *Left*: Pressure, energy and entropy densities in (2+1)-flavor QCD at vanishing chemical potential. The figure is taken from [49]. Also shown in the figure are results obtained with the stout discretization scheme for staggered fermions [50]. *Right*: The speed of sound as function of energy density.

used approaches in lattice QCD calculations as well as for collider based heavy ion experiments that search for the CEP.

6. Equation of state of strongly interacting matter

The equation of state (EoS) of strongly interacting matter, *i.e.* the pressure and its derivatives with respect to temperature and chemical potentials provides the basic information on the phase structure of QCD. It is of central importance not only for the analysis of critical behavior in QCD but also for the analysis of experimental results on strong interaction thermodynamics that are obtained in relativistic heavy ion collision experiments.

At vanishing values of the chemical potentials the QCD EoS is well controlled and consistent results for pressure, energy and entropy densities, as well as derived observables such as the speed of sound or specific heat, have been obtained by several groups [50, 49]. We show results for some of these observables in Fig. 10. The figure on the right shows the square of the speed of sound, c_s^2 , as function of the energy density. It can be seen that c_s^2 has a minimum in the transition region, sometimes called the softest point of the QCD EoS [51]. The energy density in the vicinity of the pseudo-critical temperature ($T_{pc} \simeq 155$ MeV) is found to be,

$$\epsilon_c \simeq (350 \pm 150) \text{ MeV}/\text{fm}^3, \quad (29)$$

which is compatible with the energy density of the nucleon, $m_N/(4\pi r_N^3/3)$ for nucleon radii in the range $r_N = (0.8 - 1)$ fm. Also shown in the top figure




Figure 11: μ_B -dependent contribution to the pressure (left) and net baryon number density (right) in (2+1)-flavor QCD at several values of the baryon chemical potential, $\mu/T_B = 1.0, 1.5, 2.0, 2.5$, (bottom to top) and for $\hat{\mu}_B = 2.0$. Shown are results from Taylor expansion up to eighth order in $\hat{\mu}_B$ in the pressure series for isospin symmetric ($\mu_Q = 0$) strangeness neutral ($n_S = 0$) matter and corresponding Padé approximants obtained from these Taylor expansion coefficients. The figures are taken from [52].

is the trace of the energy-momentum tensor, $(\epsilon - 3P)/T^4$. Its deviation from zero gives some hint to the relevance of interactions in the medium (for an ideal gas as well as to leading order in high temperature perturbation theory one has $\epsilon = 3P$). Not unexpected this is largest close to the transition region and decreases only slowly in the high temperature regime. This large deviations from ideal gas or perturbative behavior is seen in many observables at temperature $T_{pc} < T < 2T_{pc}$.

Calculations of the equation of state as a function of T and μ_B have been performed using direct simulations at imaginary chemical potentials, which then get analytically continued to real values of the chemical potentials [53], as well as calculations using up to eighth order Taylor expansions in μ_B [52]. Results of such calculations agree well for $\mu_B/T \leq (2 - 2.5)$. In Fig. 11 we show results for the μ_B -dependent contribution to the pressure and net baryon number density. Comparing Fig. 11 (left) with Fig. 10 (left) shows that at $\mu_B/T \simeq 2$ and $T \simeq T_{pc}$ the pressure increases by about 30%, which is due to the increase in number of baryons in the medium.

At larger values of the baryon chemical potential the Taylor series will not converge due to the presence of either poles in the complex μ_B -plane or a real pole, that may correspond to the searched for CEP. The occurrence of poles in the complex plane also generates problems for the analytic continuation of results obtained in simulations at imaginary values of μ_B as a suitable ansatz for the continuation needs to be found. Many approaches

to improve over straightforward Taylor series approaches or simulations at imaginary chemical potential are currently being discussed [54, 55, 56, 57]. In the context of Taylor expansions a natural way to proceed is to use Padé approximants, which provide a resummation of the Taylor series and reproduce this series, when expanded for small μ_B [58, 52]. Results from [4,4] and [3,4] Padé approximants for the pressure and number density series, respectively, are also shown in Fig. 11. The good agreement with the Taylor series for $\mu_B/T \leq 2.5$ gives confidence in the validity of the Taylor series results and once more seems to rule out the occurrence of a CEP in this parameter range.

7. Outlook

Achieving better control over the influence of the axial anomaly on the QCD phase transition in the chiral limit at vanishing chemical potentials and getting better control over the structure of the QCD phase diagram at large non-zero values of the chemical potentials certainly are the two largest challenges in studies of QCD thermodynamics for the next decade.

Acknowledgements

This work was supported by the DFG Collaborative Research Centre 315477589-TRR 211, "Strong interaction matter under extreme conditions".

References

- [1] K. G. Wilson, Confinement of Quarks, Phys. Rev. D 10 (1974) 2445–2459. doi:10.1103/PhysRevD.10.2445.
- [2] M. Creutz, Monte Carlo Study of Quantized SU(2) Gauge Theory, Phys. Rev. D 21 (1980) 2308–2315. doi:10.1103/PhysRevD.21.2308.
- [3] M. D’Elia, M.-P. Lombardo, Finite density QCD via imaginary chemical potential, Phys. Rev. D 67 (2003) 014505. arXiv:hep-lat/0209146, doi:10.1103/PhysRevD.67.014505.
- [4] P. de Forcrand, O. Philipsen, The QCD phase diagram for small densities from imaginary chemical potential, Nucl. Phys. B 642 (2002) 290–306. arXiv:hep-lat/0205016, doi:10.1016/S0550-3213(02)00626-0.

- [5] R. V. Gavai, S. Gupta, Quark number susceptibilities, strangeness and dynamical confinement, *Phys. Rev. D* 64 (2001) 074506. [arXiv:hep-lat/0103013](#), [doi:10.1103/PhysRevD.64.074506](#).
- [6] C. R. Allton, S. Ejiri, S. J. Hands, O. Kaczmarek, F. Karsch, E. Laermann, C. Schmidt, L. Scorzato, The QCD thermal phase transition in the presence of a small chemical potential, *Phys. Rev. D* 66 (2002) 074507. [arXiv:hep-lat/0204010](#), [doi:10.1103/PhysRevD.66.074507](#).
- [7] L. D. McLerran, B. Svetitsky, A Monte Carlo Study of SU(2) Yang-Mills Theory at Finite Temperature, *Phys. Lett. B* 98 (1981) 195–198. [doi:10.1016/0370-2693\(81\)90986-2](#).
- [8] J. Kuti, J. Polonyi, K. Szlachanyi, Monte Carlo Study of SU(2) Gauge Theory at Finite Temperature, *Phys. Lett. B* 98 (1981) 199–204. [doi:10.1016/0370-2693\(81\)90987-4](#).
- [9] J. B. Kogut, M. Stone, H. W. Wyld, W. R. Gibbs, J. Shigemitsu, S. H. Shenker, D. K. Sinclair, Deconfinement and Chiral Symmetry Restoration at Finite Temperatures in SU(2) and SU(3) Gauge Theories, *Phys. Rev. Lett.* 50 (1983) 393. [doi:10.1103/PhysRevLett.50.393](#).
- [10] J. C. Collins, M. J. Perry, Superdense Matter: Neutrons Or Asymptotically Free Quarks?, *Phys. Rev. Lett.* 34 (1975) 1353. [doi:10.1103/PhysRevLett.34.1353](#).
- [11] N. Cabibbo, G. Parisi, Exponential Hadronic Spectrum and Quark Liberation, *Phys. Lett. B* 59 (1975) 67–69. [doi:10.1016/0370-2693\(75\)90158-6](#).
- [12] G. Baym, Confinement of quarks in nuclear matter, *Physica A* 96 (1-2) (1979) 131–135. [doi:10.1016/0378-4371\(79\)90200-0](#).
- [13] R. D. Pisarski, F. Wilczek, Remarks on the Chiral Phase Transition in Chromodynamics, *Phys. Rev. D* 29 (1984) 338–341. [doi:10.1103/PhysRevD.29.338](#).
- [14] E. V. Shuryak, Quark-Gluon Plasma and Hadronic Production of Leptons, Photons and Psions, *Phys. Lett. B* 78 (1978) 150. [doi:10.1016/0370-2693\(78\)90370-2](#).

- [15] K. Kajantie, C. Montonen, E. Pietarinen, Phase Transition of SU(3) Gauge Theory at Finite Temperature, *Z. Phys. C* 9 (1981) 253. doi:10.1007/BF01410665.
- [16] L. Yaffe, B. Svetitsky, First Order Phase Transition in the SU(3) Gauge Theory at Finite Temperature, *Phys. Rev. D* 26 (1982) 963. doi:10.1103/PhysRevD.26.963.
- [17] J. Engels, F. Karsch, H. Satz, I. Montvay, High Temperature SU(2) Gluon Matter on the Lattice, *Phys. Lett. B* 101 (1981) 89. doi:10.1016/0370-2693(81)90497-4.
- [18] J. Engels, F. Karsch, H. Satz, I. Montvay, Gauge Field Thermodynamics for the SU(2) Yang-Mills System, *Nucl. Phys. B* 205 (1982) 545–577. doi:10.1016/0550-3213(82)90077-3.
- [19] O. Kaczmarek, F. Zantow, Static quark anti-quark interactions in zero and finite temperature QCD. I. Heavy quark free energies, running coupling and quarkonium binding, *Phys. Rev. D* 71 (2005) 114510. arXiv:hep-lat/0503017, doi:10.1103/PhysRevD.71.114510.
- [20] H. T. Ding, et al., Chiral Phase Transition Temperature in (2+1)-Flavor QCD, *Phys. Rev. Lett.* 123 (6) (2019) 062002. arXiv:1903.04801, doi:10.1103/PhysRevLett.123.062002.
- [21] O. Kaczmarek, F. Karsch, A. Lahiri, C. Schmidt, Universal scaling properties of QCD close to the chiral limit, *Acta Phys. Polon. Supp.* 14 (2021) 291. arXiv:2010.15593, doi:10.5506/APHYSPOLBSUPP.14.291.
- [22] K. Rajagopal, F. Wilczek, Static and dynamic critical phenomena at a second order QCD phase transition, *Nucl. Phys. B* 399 (1993) 395–425. arXiv:hep-ph/9210253, doi:10.1016/0550-3213(93)90502-G.
- [23] K. Rajagopal, F. Wilczek, The Condensed matter physics of QCD, published in: *At the frontier of particle physics. Handbook of QCD.* Vol. 1-3 (Edts. M.Shifman and B. Ioffe), 2000, pp. 2061–2151. arXiv:hep-ph/0011333, doi:10.1142/9789812810458_0043.
- [24] B. Svetitsky, L. G. Yaffe, Critical Behavior at Finite Temperature Confinement Transitions, *Nucl. Phys. B* 210 (1982) 423–447. doi:10.1016/0550-3213(82)90172-9.

- [25] F. R. Brown, N. H. Christ, Y. F. Deng, M. S. Gao, T. J. Woch, Nature of the Deconfining Phase Transition in SU(3) Lattice Gauge Theory, *Phys. Rev. Lett.* 61 (1988) 2058. doi:10.1103/PhysRevLett.61.2058.
- [26] J. Engels, J. Fingberg, M. Weber, Finite Size Scaling Analysis of SU(2) Lattice Gauge Theory in (3+1)-dimensions, *Nucl. Phys. B* 332 (1990) 737–759. doi:10.1016/0550-3213(90)90010-B.
- [27] F. Cuteri, O. Philipsen, A. Sciarra, On the order of the QCD chiral phase transition for different numbers of quark flavours, *JHEP* 11 (2021) 141. arXiv:2107.12739, doi:10.1007/JHEP11(2021)141.
- [28] F. R. Brown, F. P. Butler, H. Chen, N. H. Christ, Z.-h. Dong, W. Schaffer, L. I. Unger, A. Vaccarino, On the existence of a phase transition for QCD with three light quarks, *Phys. Rev. Lett.* 65 (1990) 2491–2494. doi:10.1103/PhysRevLett.65.2491.
- [29] L. Dini, P. Hegde, F. Karsch, A. Lahiri, C. Schmidt, S. Sharma, Chiral phase transition in three-flavor QCD from lattice QCD, *Phys. Rev. D* 105 (3) (2022) 034510. arXiv:2111.12599, doi:10.1103/PhysRevD.105.034510.
- [30] A. Y. Kotov, M. P. Lombardo, A. Trunin, QCD transition at the physical point, and its scaling window from twisted mass Wilson fermions, *Phys. Lett. B* 823 (2021) 136749. arXiv:2105.09842, doi:10.1016/j.physletb.2021.136749.
- [31] A. Bazavov, et al., Chiral crossover in QCD at zero and non-zero chemical potentials, *Phys. Lett. B* 795 (2019) 15–21. arXiv:1812.08235, doi:10.1016/j.physletb.2019.05.013.
- [32] C. Bonati, M. D’Elia, M. Mariti, M. Mesiti, F. Negro, F. Sanfilippo, Curvature of the chiral pseudocritical line in QCD: Continuum extrapolated results, *Phys. Rev. D* 92 (5) (2015) 054503. arXiv:1507.03571, doi:10.1103/PhysRevD.92.054503.
- [33] C. Bonati, M. D’Elia, F. Negro, F. Sanfilippo, K. Zambello, Curvature of the pseudocritical line in QCD: Taylor expansion matches analytic continuation, *Phys. Rev. D* 98 (5) (2018) 054510. arXiv:1805.02960, doi:10.1103/PhysRevD.98.054510.

- [34] S. Borsanyi, Z. Fodor, J. N. Guenther, R. Kara, S. D. Katz, P. Parotto, A. Pasztor, C. Ratti, K. K. Szabo, QCD Crossover at Finite Chemical Potential from Lattice Simulations, *Phys. Rev. Lett.* 125 (5) (2020) 052001. [arXiv:2002.02821](#), [doi:10.1103/PhysRevLett.125.052001](#).
- [35] R. Guida, J. Zinn-Justin, Critical exponents of the N vector model, *J. Phys. A* 31 (1998) 8103–8121. [arXiv:cond-mat/9803240](#), [doi:10.1088/0305-4470/31/40/006](#).
- [36] A. Pelissetto, E. Vicari, Relevance of the axial anomaly at the finite-temperature chiral transition in QCD, *Phys. Rev. D* 88 (10) (2013) 105018. [arXiv:1309.5446](#), [doi:10.1103/PhysRevD.88.105018](#).
- [37] A. Lahiri, Aspects of finite temperature QCD towards the chiral limit, *PoS LATTICE2021* (2022) 003. [arXiv:2112.08164](#), [doi:10.22323/1.396.0003](#).
- [38] G. Aarts, C. Allton, D. De Boni, B. Jäger, Hyperons in thermal QCD: A lattice view, *Phys. Rev. D* 99 (7) (2019) 074503. [arXiv:1812.07393](#), [doi:10.1103/PhysRevD.99.074503](#).
- [39] A. Bazavov, et al., Meson screening masses in (2+1)-flavor QCD, *Phys. Rev. D* 100 (9) (2019) 094510. [arXiv:1908.09552](#), [doi:10.1103/PhysRevD.100.094510](#).
- [40] S. Dentinger, O. Kaczmarek, A. Lahiri, Screening masses towards chiral limit, *Acta Phys. Polon. Supp.* 14 (2021) 321. [arXiv:2102.09916](#), [doi:10.5506/APHYSPOLBSUPP.14.321](#).
- [41] C. E. Detar, J. B. Kogut, The Hadronic Spectrum of the Quark Plasma, *Phys. Rev. Lett.* 59 (1987) 399. [doi:10.1103/PhysRevLett.59.399](#).
- [42] C. E. Detar, J. B. Kogut, Measuring the Hadronic Spectrum of the Quark Plasma, *Phys. Rev. D* 36 (1987) 2828. [doi:10.1103/PhysRevD.36.2828](#).
- [43] G. Cossu, S. Aoki, H. Fukaya, S. Hashimoto, T. Kaneko, H. Matsufuru, J.-I. Noaki, Finite temperature study of the axial U(1) symmetry on the lattice with overlap fermion formulation, *Phys. Rev. D* 87 (11) (2013) 114514, [Erratum: *Phys.Rev.D* 88, 019901 (2013)]. [arXiv:1304.6145](#), [doi:10.1103/PhysRevD.87.114514](#).

- [44] A. Tomiya, G. Cossu, S. Aoki, H. Fukaya, S. Hashimoto, T. Kaneko, J. Noaki, Evidence of effective axial $U(1)$ symmetry restoration at high temperature QCD, Phys. Rev. D 96 (3) (2017) 034509, [Addendum: Phys.Rev.D 96, 079902 (2017)]. [arXiv:1612.01908](#), doi: 10.1103/PhysRevD.96.034509.
- [45] M. I. Buchoff, et al., QCD chiral transition, $U(1)_A$ symmetry and the dirac spectrum using domain wall fermions, Phys. Rev. D 89 (5) (2014) 054514. [arXiv:1309.4149](#), doi:10.1103/PhysRevD.89.054514.
- [46] V. Dick, F. Karsch, E. Laermann, S. Mukherjee, S. Sharma, Microscopic origin of $U_A(1)$ symmetry violation in the high temperature phase of QCD, Phys. Rev. D 91 (9) (2015) 094504. [arXiv:1502.06190](#), doi: 10.1103/PhysRevD.91.094504.
- [47] A. M. Halasz, A. D. Jackson, R. E. Shrock, M. A. Stephanov, J. J. M. Verbaarschot, On the phase diagram of QCD, Phys. Rev. D 58 (1998) 096007. [arXiv:hep-ph/9804290](#), doi:10.1103/PhysRevD.58.096007.
- [48] M. Buballa, S. Carignano, Inhomogeneous chiral phases away from the chiral limit, Phys. Lett. B 791 (2019) 361–366. [arXiv:1809.10066](#), doi:10.1016/j.physletb.2019.02.045.
- [49] A. Bazavov, et al., Equation of state in (2+1)-flavor QCD, Phys. Rev. D 90 (2014) 094503. [arXiv:1407.6387](#), doi:10.1103/PhysRevD.90.094503.
- [50] S. Borsanyi, Z. Fodor, C. Hoelbling, S. D. Katz, S. Krieg, K. K. Szabo, Full result for the QCD equation of state with 2+1 flavors, Phys. Lett. B 730 (2014) 99–104. [arXiv:1309.5258](#), doi:10.1016/j.physletb.2014.01.007.
- [51] C. M. Hung, E. V. Shuryak, Hydrodynamics near the QCD phase transition: Looking for the longest lived fireball, Phys. Rev. Lett. 75 (1995) 4003–4006. [arXiv:hep-ph/9412360](#), doi:10.1103/PhysRevLett.75.4003.
- [52] D. Bollweg, J. Goswami, O. Kaczmarek, F. Karsch, S. Mukherjee, P. Petreczky, C. Schmidt, P. Scior, Taylor expansions and Padé approximants for cumulants of conserved charge fluctuations at nonvanishing chemical

- potentials, Phys. Rev. D 105 (7) (2022) 074511. [arXiv:2202.09184](#), [doi:10.1103/PhysRevD.105.074511](#).
- [53] S. Borsanyi, G. Endrodi, Z. Fodor, S. D. Katz, S. Krieg, C. Ratti, K. K. Szabo, QCD equation of state at nonzero chemical potential: continuum results with physical quark masses at order mu^2 , JHEP 08 (2012) 053. [arXiv:1204.6710](#), [doi:10.1007/JHEP08\(2012\)053](#).
 - [54] S. Mukherjee, V. Skokov, Universality driven analytic structure of the QCD crossover: radius of convergence in the baryon chemical potential, Phys. Rev. D 103 (7) (2021) L071501. [arXiv:1909.04639](#), [doi:10.1103/PhysRevD.103.L071501](#).
 - [55] S. Mondal, S. Mukherjee, P. Hegde, Lattice QCD Equation of State for Nonvanishing Chemical Potential by Resumming Taylor Expansions, Phys. Rev. Lett. 128 (2) (2022) 022001. [arXiv:2106.03165](#), [doi:10.1103/PhysRevLett.128.022001](#).
 - [56] P. Dimopoulos, L. Dini, F. Di Renzo, J. Goswami, G. Nicotra, C. Schmidt, S. Singh, K. Zambello, F. Ziesché, Contribution to understanding the phase structure of strong interaction matter: Lee-Yang edge singularities from lattice QCD, Phys. Rev. D 105 (3) (2022) 034513. [arXiv:2110.15933](#), [doi:10.1103/PhysRevD.105.034513](#).
 - [57] S. Borsanyi, Z. Fodor, J. N. Guenther, R. Kara, P. Parotto, A. Pasztor, C. Ratti, K. K. Szabo, Resummed lattice QCD equation of state at finite baryon density: Strangeness neutrality and beyond, Phys. Rev. D 105 (11) (2022) 114504. [arXiv:2202.05574](#), [doi:10.1103/PhysRevD.105.114504](#).
 - [58] S. Datta, R. V. Gavai, S. Gupta, Quark number susceptibilities and equation of state at finite chemical potential in staggered QCD with $N_t=8$, Phys. Rev. D 95 (5) (2017) 054512. [arXiv:1612.06673](#), [doi:10.1103/PhysRevD.95.054512](#).