

# Conductivity in flat bands from the Kubo-Greenwood formula

Kukka-Emilia Huhtinen<sup>1,\*</sup> and Päivi Törmä<sup>1,†</sup>

<sup>1</sup>*Department of Applied Physics, Aalto University School of Science, FI-00076 Aalto, Finland*  
(Dated: December 7, 2022)

Conductivity in a multiband system can be divided into intra- and interband contributions, and the latter further into symmetric and antisymmetric parts. In a flat band, intraband conductivity vanishes and the antisymmetric interband contribution, proportional to the Berry curvature, corresponds to the anomalous Hall effect. We investigate whether the symmetric interband conductivity, related to the quantum metric, can be finite in the zero frequency and flat band limit. Starting from the Kubo-Greenwood formula with a finite scattering rate  $\eta$ , we show that the DC conductivity is zero in a flat band when taking the clean limit ( $\eta \rightarrow 0$ ). If commonly used approximations involving derivatives of the Fermi distribution are used, finite conductivity appears at zero temperature  $T = 0$ , we show however that this is an artifact due to the lack of Fermi surfaces in a (partially) flat band. We then analyze the DC conductivity using the Kubo-Streda formula, and note similar problems at  $T = 0$ . The predictions of the Kubo-Greenwood formula (without the approximation) and the Kubo-Streda formula differ significantly at low temperatures. We illustrate the results within the Su-Schrieffer-Heeger model where one expects vanishing DC conductivity in the dimerized limit as the unit cells are disconnected. We discuss the implications of our results to previous work which has proposed the possibility of finite flat-band DC conductivity proportional to the quantum metric. Our results also highlight that care should be taken when applying established transport and linear response approaches in the flat band context, since many of them utilize the existence of a Fermi surface and assume scattering to be weak compared to kinetic energy.

## I. INTRODUCTION

Quantum geometry is key to understanding multiband systems. For instance, quantum geometry has been related to superconductivity in flat bands,<sup>1–11</sup> orbital magnetic susceptibility,<sup>12,13</sup> light-matter interactions,<sup>14,15</sup> the intrinsic anomalous Hall effect,<sup>16–20</sup> and other physical phenomena.<sup>21–27</sup> Quantum geometric quantities determine the phase and amplitude distances between quantum states, and are represented by the quantum geometric tensor<sup>28</sup> whose imaginary (antisymmetric) part is the Berry curvature and real (symmetric) part the quantum metric (Fubini-Study metric). The quantum metric and Berry curvature are particularly central in the properties of flat bands systems. Flat bands are interesting platforms for strongly correlated quantum phenomena, and have attracted increased interest due to their relevance in moiré materials.<sup>29–39</sup>

Non-interacting particles on flat bands are localized and have a diverging effective mass. However, recent results have predicted a nonzero DC conductivity weakly sensitive to the inelastic scattering rate,<sup>40,41</sup> also found in disorder-induced quasilocalized zero-energy modes in graphene.<sup>42</sup> This result is sensitive to the used approach, and for instance wave-packet propagation methods predict a vanishing conductivity of the zero-energy modes.<sup>43,44</sup> A nonzero or even diverging DC conductivity has also been predicted in disordered non-isolated flat bands.<sup>45,46</sup> In perfectly flat bands, the zero-temperature DC conductivity has been derived to be proportional to the quantum metric.<sup>47–49</sup>

The conductivity is often computed using the Kubo-Greenwood formula, which is the non-interacting version of the exact Kubo formula. The Kubo-Greenwood for-

mula can be sensitive to the order the relevant limits (zero temperature, zero scattering, zero frequency) are taken<sup>50</sup> or to approximations made for instance in the delta functions.<sup>51</sup> Here, we show that when applied to flat bands, the Kubo-Streda formula can give drastically different results than the Kubo-Greenwood formula obtained directly via an independent particle approximation of the Kubo formula. In particular, if the zero temperature limit is taken before taking the scattering rate to zero, it can yield a conductivity proportional to the quantum metric in the clean limit, which does not appear when applying the Kubo-Greenwood formula which predicts vanishing DC conductivity.

In Section II, we first derive the conductivity within the Kubo-Greenwood formula dividing it into intra- and interband contributions and their symmetric and antisymmetric parts. We show how in a flat band, where the intraband conductivity is zero and the antisymmetric interband contributions give the anomalous Hall effect, the symmetric intraband contribution vanishes at small temperatures in the clean limit. This means no DC conductivity. In Section III, we then compute the conductivity using the Kubo-Streda formula, and show, using also the sawtooth ladder and the dimerized SSH chain as examples, that the results differ dramatically from the Kubo-Greenwood formula at low temperatures. We explain the origin of the discrepancy. In Section IV, we summarize and discuss our results.

## II. KUBO-GREENWOOD FORMULA

We consider fermionic multiband models described by the tight-binding Hamiltonian  $H = \sum_{i\alpha,j\beta} t_{i\alpha,j\beta} c_{i\alpha}^\dagger c_{j\beta} -$




FIG. 1. (a) Sketch of the dimerized SSH model and the sawtooth ladder studied here. (b-c) DC conductivity in the SSH model at  $\mu = -t$ , when the chemical potential is in the lowest flat band (b), and at  $\mu = -0.9t$  (c). The conductivity obtained from the Kubo-Greenwood formula consists of solely interband contributions, and vanishes at all temperatures in the clean limit  $\eta \rightarrow 0^+$ . The DC conductivity obtained from the Kubo-Streda formula remains pinned at  $1/(8\pi)$  at  $T = 0$ , but vanishes for nonzero temperatures. When the chemical potential is tuned away from the flat band, the limit  $\eta \rightarrow 0^+$  from the Kubo-Greenwood and Kubo-Streda formulae is the same, although the behavior at nonzero  $\eta$  is drastically different. (d) Interband and intraband contributions obtained from the Kubo-Greenwood formula in the sawtooth ladder when the chemical potential is in the flat band. The intraband contribution from the dispersive band diverges as  $\eta \rightarrow 0^+$ , whereas the interband contribution vanishes. (e-f) DC conductivity from the Kubo-Greenwood and Kubo-Streda formula when the chemical potential is (e) in the flat band and (f) in the middle of the dispersive band. In a dispersive band, both methods give the same results, whereas in a flat band, they give drastically different results. In the flat band, the conductivity from the Kubo-Streda formula remains pinned to  $2/(3\sqrt{3}\pi)$  at  $T = 0$ .

$\mu \sum_{i\alpha} n_{i\alpha}$ , where  $t_{i\alpha,j\beta}$  is the hopping amplitude from site  $j\beta$  to  $i\alpha$ . The unit cells are labeled with  $i, j$ , while  $\alpha, \beta$  indicate the orbitals within a unit cell. By taking the Fourier transformation  $c_{i\alpha} = (1/\sqrt{N_c}) \sum_{\mathbf{k}} c_{\mathbf{k}\alpha} e^{i\mathbf{k} \cdot (\mathbf{R}_i + \boldsymbol{\delta}_\alpha)}$  the Hamiltonian becomes  $H = \sum_{\mathbf{k}} H_{\mathbf{k}}$ , where

$$H_{\mathbf{k}} = \sum_i \sum_{\alpha\beta} t_{i\alpha,0\beta} e^{-i\mathbf{k} \cdot (\mathbf{R}_i + \boldsymbol{\delta}_\alpha - \boldsymbol{\delta}_\beta)}. \quad (1)$$

Here,  $\mathbf{R}_i$  is the position of the  $i$ :th unit cell, and  $\boldsymbol{\delta}_\alpha = \mathbf{r}_{i\alpha} - \mathbf{R}_i$ , with  $\mathbf{r}_{i\alpha}$  the position of site  $i\alpha$ . The eigenvalues and eigenvectors give the band dispersion relations  $\epsilon_n(\mathbf{k})$  and the periodic parts of the Bloch functions  $|n_{\mathbf{k}}\rangle$ , respectively.

Note that in a multiband lattice taking the orbital positions  $\boldsymbol{\delta}_\alpha$  into account in the Fourier transformation is essential in order to obtain the correct conductivity.<sup>52–54</sup> This is in contrast to the superfluid weight, which is independent of the particular choice of  $\boldsymbol{\delta}_\alpha$  provided it is computed accurately.<sup>10</sup> In other words, the conductivity

is generally geometry-dependent, using the terminology introduced in Ref. 55.

The conductivity tensor  $\sigma_{ij}(\omega) \equiv \sigma_{ij}(\omega, \mathbf{q} = \mathbf{0})$  in a non-interacting system is given by the Kubo-Greenwood formula<sup>56</sup>

$$\sigma_{\mu\nu}(\omega) = \frac{e^2}{i\hbar V} \sum_{\mathbf{k}} \sum_{mn} \frac{n_F(\epsilon_n(\mathbf{k})) - n_F(\epsilon_m(\mathbf{k}))}{\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k})} \times \frac{[j_\mu(\mathbf{k})]_{nm} [j_\nu(\mathbf{k})]_{mn}}{\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}) + \hbar\omega + i\eta}, \quad (2)$$

which is obtained from the Kubo formula<sup>57</sup> by performing an independent electron approximation. The prefactor involving the Fermi-Dirac distribution  $n_F(\epsilon) = 1/(e^{\beta\epsilon} + 1)$ , with  $\beta = 1/(k_B T)$ , should be understood as  $\partial_\epsilon n_F(\epsilon)|_{\epsilon=\epsilon_n}$  when  $\epsilon_n(\mathbf{k}) = \epsilon_m(\mathbf{k})$ .<sup>51</sup> The infinitesimal imaginary shift  $\eta$  added to the frequency acts as a small inelastic scattering rate or relaxation rate. The current operators  $j_\mu$  are obtained from the momentum deriva-

tives of  $H_{\mathbf{k}}$ ,  $j_{\mu}(\mathbf{k}) = \partial_{k_{\mu}} H_{\mathbf{k}}$ , and

$$[j_{\mu}(\mathbf{k})]_{mn} = \partial_{k_{\mu}} \epsilon_m(\mathbf{k}) \delta_{mn} + (\epsilon_m(\mathbf{k}) - \epsilon_n(\mathbf{k})) \langle \partial_{k_{\mu}} m_{\mathbf{k}} | n_{\mathbf{k}} \rangle. \quad (3)$$

Another widely used form of the Kubo-Greenwood formula for the diagonal components of  $\sigma_{\mu\nu}$  is

$$\sigma_{\mu\mu} = -\frac{e^2}{\hbar\pi V} \sum_{\mathbf{k}} \sum_{mn} \int_{-\infty}^{\infty} d\epsilon \frac{\partial n_F(\epsilon)}{\partial \epsilon} \times \text{Tr}[\text{Im}[G_{\mathbf{k}}(\epsilon + i\eta)] j_{\mu}(\mathbf{k}) \text{Im}[G_{\mathbf{k}}(\epsilon + i\eta)] j_{\mu}(\mathbf{k})], \quad (4)$$

where  $G_{\mathbf{k}}(E) = (E - H_{\mathbf{k}})^{-1}$  is the Green's function. This is simply the diagonal components of the more general Bastin<sup>58</sup> and Streda<sup>59</sup> formulae. In this work, we show that when applied to flat bands, Eqs. (4) and (2) give drastically different results at  $T = 0$ . In order to avoid confusion between the two forms, we will refer to Eq. (2) as the Kubo-Greenwood formula, and to Eq. (4) as the Kubo-Streda formula.

Let us first derive  $\text{Re}[\sigma_{\mu\nu}]$  from Eq. (2). The real part of the conductivity can be decomposed in several ways, for instance into so-called Fermi surface and Fermi sea contributions.<sup>59-61</sup> In our case, we choose to split  $\sigma_{\mu\nu}$  to intraband, symmetric interband and antisymmetric interband contributions, similarly to the decomposition

used in Ref. 54. The advantage of this split when considering flat bands is apparent: the intraband contribution from a perfectly flat band vanishes exactly, and only the interband part remains. The antisymmetric part of the latter is related to the intrinsic anomalous Hall effect.

In the thermodynamic limit, the intraband contribution to the real part of the conductivity  $\sigma_{\mu\nu}^{\text{intra}}$  obtained from Eq. (2) is

$$\text{Re } \sigma_{\mu\nu}^{\text{intra}}(\omega) = -\frac{e^2}{\hbar} \sum_n \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} \frac{\partial n_F(E)}{\partial E} \Big|_{E=\epsilon_n(\mathbf{k})} \times [j_{\mu}(\mathbf{k})]_{nn} [j_{\nu}(\mathbf{k})]_{nn} \frac{\eta}{(\hbar\omega)^2 + \eta^2} \quad (5)$$

$$= -\frac{e^2}{\hbar} \sum_n \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} \frac{\partial n_F(E)}{\partial E} \Big|_{E=\epsilon_n(\mathbf{k})} \times \partial_{k_{\mu}} \epsilon_n(\mathbf{k}) \partial_{k_{\nu}} \epsilon_n(\mathbf{k}) \frac{\eta}{(\hbar\omega)^2 + \eta^2}. \quad (6)$$

We have replaced the momentum summation by an integral over the first Brillouin zone,  $(1/V) \sum_{\mathbf{k}} \rightarrow (1/2\pi)^D \int_{\text{B.z.}} d^D \mathbf{k}$ , where  $D$  is the dimension of the system. This contribution to the conductivity is the only component present in a single-band model, and gives the same result as the semiclassical Boltzmann theory of transport when taking  $\tau = 1/\eta$  as a momentum-independent relaxation time. The intraband contribution is clearly zero in a perfectly dispersionless band.

The total interband contribution is

$$\sigma_{\mu\nu}^{\text{inter}} = -i \frac{e^2}{\hbar} \sum_{n \neq m} \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} \frac{n_F(\epsilon_n(\mathbf{k}))}{\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k})} \left( \frac{[j_{\mu}(\mathbf{k})]_{nm} [j_{\nu}(\mathbf{k})]_{mn}}{\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}) + \hbar\omega + i\eta} + \frac{[j_{\nu}(\mathbf{k})]_{nm} [j_{\mu}(\mathbf{k})]_{mn}}{\epsilon_m(\mathbf{k}) - \epsilon_n(\mathbf{k}) + \hbar\omega + i\eta} \right). \quad (7)$$

Using Eq. (3), we can express the symmetric and antisymmetric parts of the real part of  $\sigma_{\mu\nu}^{\text{inter}}$  as

$$\sigma_{\mu\nu}^s(\omega) = -\frac{e^2}{\hbar} \sum_{n \neq m} \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} n_F(\epsilon_n(\mathbf{k})) \text{Re}[\langle \partial_{k_{\mu}} n_{\mathbf{k}} | m_{\mathbf{k}} \rangle \langle m_{\mathbf{k}} | \partial_{k_{\nu}} n_{\mathbf{k}} \rangle] \times \left( \frac{\eta(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}))}{(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}) + \hbar\omega)^2 + \eta^2} + \frac{\eta(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}))}{(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}) - \hbar\omega)^2 + \eta^2} \right), \quad (8)$$

$$\sigma_{\mu\nu}^a(\omega) = \frac{e^2}{\hbar} \sum_{\mathbf{k}} \sum_{n \neq m} \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} n_F(\epsilon_n(\mathbf{k})) \text{Im}[\langle \partial_{k_{\mu}} n_{\mathbf{k}} | m_{\mathbf{k}} \rangle \langle m_{\mathbf{k}} | \partial_{k_{\nu}} n_{\mathbf{k}} \rangle] \times \left( \frac{(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}))(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}) + \hbar\omega)}{(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}) + \hbar\omega)^2 + \eta^2} + \frac{(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}))(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}) - \hbar\omega)}{(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}) - \hbar\omega)^2 + \eta^2} \right), \quad (9)$$

The interband contribution is thus related to components of the quantum geometric tensor,<sup>62</sup> which for the  $n$ :th band is defined as

$$\mathcal{G}_{\mu\nu}^n(\mathbf{k}) = 2 \langle \partial_{k_{\mu}} n_{\mathbf{k}} | (1 - |n_{\mathbf{k}}\rangle \langle n_{\mathbf{k}}|) | \partial_{k_{\nu}} n_{\mathbf{k}} \rangle \quad (10)$$

$$= 2 \sum_{m:m \neq n} \langle \partial_{k_{\mu}} n_{\mathbf{k}} | m_{\mathbf{k}} \rangle \langle m_{\mathbf{k}} | \partial_{k_{\nu}} n_{\mathbf{k}} \rangle. \quad (11)$$

The real and imaginary parts of  $\mathcal{G}_{\mu\nu}^n = \mathcal{M}_{\mu\nu}^n + i\mathcal{B}_{\mu\nu}^n$  are the quantum metric and Berry curvature, respectively. Since the quantum geometric tensor is Hermitian, the quantum metric is symmetric while the Berry curvature is antisymmetric.

In the limit  $\omega \rightarrow 0$ ,  $\eta \rightarrow 0^+$ , we recover for  $\sigma_{\mu\nu}^a$  the well-known relationship between the intrinsic anomalous

Hall conductivity and the Berry curvature,<sup>16–20,63</sup>

$$\sigma_{\mu\nu}^a(\omega=0) = \frac{e^2}{\hbar} \sum_n \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} n_F(\epsilon_n(\mathbf{k})) \mathcal{B}_{\mu\nu}^n(\mathbf{k}). \quad (12)$$

Notably, the antisymmetric part of the conductivity can be nonzero even on a flat band in the clean limit  $\eta \rightarrow 0^+$ .

$$\begin{aligned} \sigma_{\mu\nu}^s(\omega) = & -\frac{e^2\pi}{\hbar} \sum_{n \neq m} \int_{\text{B.z.}} n_F(\epsilon_n(\mathbf{k})) \text{Re} [\langle \partial_{k_\mu} n_{\mathbf{k}} | m_{\mathbf{k}} \rangle \langle m_{\mathbf{k}} | \partial_{k_\nu} n_{\mathbf{k}} \rangle] (\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k})) \times \\ & \times (\delta(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}) + \hbar\omega) + \delta(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}) - \hbar\omega)). \end{aligned} \quad (13)$$

When all bands are isolated from each other,  $|\epsilon_n - \epsilon_m| \geq E_{\text{gap,min}}$  for  $m \neq n$ , where  $E_{\text{gap,min}}$  is the smallest interband gap. One can see from the delta-functions in Equation (13) that the interband conductivity then vanishes for any frequency  $\hbar\omega < E_{\text{gap,min}}$  in the clean limit. It follows that the symmetric part of the DC conductivity in an isolated flat band is zero in the limit  $\eta \rightarrow 0^+$ . At nonzero scattering rate, the DC conductivity can acquire a nonzero value because of the spread of the Lorentzian functions centered at frequencies  $\hbar\omega \geq E_{\text{gap,min}}$ ; this is just the finite linewidth of interband transitions resonant with an AC field. For  $\eta \ll E_{\text{gap,min}}$ , this nonzero interband contribution is approximately

$$\begin{aligned} \sigma_{\mu\nu}^s(\omega=0) \approx & -2\eta \frac{e^2}{\hbar} \sum_n \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} n_F(\epsilon_n(\mathbf{k})) \times \\ & \times \sum_{m:m \neq n} \frac{\text{Re}[\langle \partial_{k_\mu} n_{\mathbf{k}} | m_{\mathbf{k}} \rangle \langle m_{\mathbf{k}} | \partial_{k_\nu} n_{\mathbf{k}} \rangle]}{\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k})}, \end{aligned} \quad (14)$$

where we have approximated  $\eta(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k})) / [(\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}))^2 + \eta^2] \approx \eta / (\epsilon_n(\mathbf{k}) - \epsilon_m(\mathbf{k}))$ . The linear dependence on the scattering rate is consistent with the result obtained in Refs. 47 and 54 for small scattering rates in a dispersive band. Here, this holds also for perfectly flat bands. We note that an expression which relates the DC conductivity directly to the quantum metric can be obtained from Eq. (13) by using that

$$\begin{aligned} & \frac{n_F(\epsilon_n) - n_F(\epsilon_m)}{\epsilon_n - \epsilon_m} \delta(\epsilon_n - \epsilon_m + \hbar\omega) \\ & = \frac{n_F(\epsilon_n + \hbar\omega) - n_F(\epsilon_n)}{\hbar\omega} \delta(\epsilon_n - \epsilon_m + \hbar\omega). \end{aligned} \quad (15)$$

The prefactor on the second line should be, as mentioned earlier, understood as the derivative  $\partial n_F(E) / \partial E|_{E=\epsilon_n(\mathbf{k})}$  when  $\omega \rightarrow 0$ . If the delta functions are replaced by Lorentzian functions after this substitution, the DC con-

ductivity for  $\eta \ll E_{\text{gap,min}}$  becomes

$$\begin{aligned} \sigma_{\mu\nu}^s(\omega=0) = & -\eta \frac{e^2}{\hbar} \sum_n \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} \frac{\partial n_F(E)}{\partial E} \Big|_{E=\epsilon_n(\mathbf{k})} \times \\ & \times \sum_{m \neq n} \text{Re}[\langle \partial_{k_\mu} n | m \rangle \langle m | \partial_{k_\nu} n \rangle]. \end{aligned} \quad (16)$$

This form is more reminiscent of results obtained for instance by computing the conductivity in Matsubara space.<sup>47,54</sup> It should however be stressed that here this is an approximation, as Eq. (15) no longer holds if the delta-function is replaced by a Lorentzian function with a finite spread. In the limit  $\omega \rightarrow 0$ , the prefactor on the right-hand side of Equation (15) is replaced by a derivative of the Fermi-Dirac distribution at  $\epsilon_n$ , which approximates  $(n_F(\epsilon_n) - n_F(\epsilon_m)) / (\epsilon_n - \epsilon_m)$  on the left hand side only when  $\epsilon_n$  and  $\epsilon_m$  are close. If the bands are well isolated, this does not hold. Equation (16) is also problematic at  $T = 0$  if the Fermi energy is in a perfectly flat band, as the derivative of the Fermi-Dirac distribution would then become  $-\delta(E_{FB} - \mu) = -\delta(0)$ , where  $E_{FB}$  is the energy of the flat band, at all points of the Brillouin zone. Transforming the integral over the momentum to an integral over the energy does not help, as it would involve the density of states on the flat band, which diverges. However, at any nonzero temperature, equations (16) and (14) always give the same vanishing interband DC conductivity in the  $\eta \rightarrow 0^+$  limit.

A vanishing DC conductivity on a noninteracting isolated flat band is unsurprising, since single particles are localized. However, it contrasts with recent results at  $T = 0$ , which have found a nonzero DC conductivity proportional to the quantum metric even in the limit  $\eta \rightarrow 0^+$ .<sup>41,47–49</sup> In 47 and 54, it was shown that the interband contribution vanishes linearly with  $\eta$  on dispersive bands, and this finite conductivity in the clean limit thus appears only in perfectly flat bands. In the following, we show that such results can arise when applying the Kubo-Streda formula when the Fermi energy is in a (partially) flat band. This is related to contributions from states

at exactly the Fermi energy which do not vanish in the clean limit, present in the Streda formula but absent in the Kubo-Greenwood formula (Eq. (2)). These contributions only become meaningful in systems without a Fermi surface, such as a flat band.

### III. FLAT BAND CONDUCTIVITY FROM THE KUBO-STREDA FORMULA

The Kubo-Streda formula gives the symmetric part of the DC conductivity as<sup>59–61</sup>

$$\sigma_{\mu\nu}^{\text{sym}} = -\frac{e^2}{\hbar\pi} \int_{-\infty}^{\infty} d\epsilon \frac{\partial n_F(\epsilon)}{\partial \epsilon} \text{Tr}[\text{Im}[G_{\mathbf{k}}(\epsilon + i\eta)]j_{\mu}(\mathbf{k})\text{Im}[G_{\mathbf{k}}(\epsilon + i\eta)]j_{\nu}(\mathbf{k})]. \quad (17)$$

This equation can be derived from the Kubo-Greenwood formula directly, or can be obtained from the exact Kubo formula by computing the current-current response function in Matsubara space when vertex corrections are ignored. When applied to dispersive bands, Equations (2) and (17) usually give very close results, especially at low  $\eta$ , provided we take  $\eta_{K-G} = 2\eta_{\text{Streda}}$ . However, when applied to flat bands, the DC conductivities can differ drastically especially at low temperatures.

To illustrate this, we consider two one-dimensional flat band systems: the sawtooth ladder and the dimerized limit of the Su-Schrieffer-Heeger (SSH) model (see Fig. 1a). The sawtooth ladder features a perfectly flat band at energy  $E = 2t$ , isolated from a dispersive band  $\epsilon(\mathbf{k}) = -(2 + 2\cos(k))t$ . The dimerized SSH model has two exactly flat bands at energies  $E = \pm t$ . Importantly, the dimerized SSH model consists of two-site clusters that are completely disconnected from each other. It is thus reasonable to expect the DC conductivity to vanish.

As can be seen from Fig. 1, when the chemical potential is tuned into the flat band, the interband conductivity obtained from Eq. (2) vanishes in the clean limit  $\eta \rightarrow 0^+$  in all cases. In the SSH model, the intraband conductivity is exactly zero because the system contains only flat bands, and the Kubo-Greenwood formula predicts a vanishing  $\sigma(\omega = 0)$ , Fig. 1b,c. In the sawtooth ladder (Fig. 1d,e), the intraband contribution from the dispersive band is nonzero at any  $T > 0$ , and diverges in the limit  $\eta \rightarrow 0^+$ . However, it is highly suppressed at low temperatures even for small  $\eta$ . At the values of  $\eta$  used here, the interband conductivity is dominant up to a temperature  $k_B T \sim 0.3$ . This threshold temperature decreases with the scattering rate  $\eta$ .

The conductivity obtained from Eq. (17), the Kubo-Streda formula, is drastically different. At exactly  $T = 0$ , it retains a nonzero value proportional to the integrated quantum metric even when  $\eta \rightarrow 0^+$  (Fig. 1b,e). For  $T > 0$ , the DC conductivity vanishes in the clean limit in the dimerized SSH model (Fig. 1b). As a consequence, the limits  $\lim_{T \rightarrow 0^+}$  and  $\lim_{\eta \rightarrow 0^+}$  do not commute: taking  $\lim_{T \rightarrow 0^+}$  before  $\lim_{\eta \rightarrow 0^+}$  gives a nonzero conductivity,

whereas the inverted order gives  $\sigma(\omega = 0) = 0$ . This  $T = 0$  behavior only exists when the chemical potential is tuned exactly into the flat band: if it is slightly shifted away from the flat band, a bump in  $\sigma(\omega = 0)$  appears at a nonzero temperature, as shown in Fig. 1c, but the conductivity vanishes in the clean limit even at  $T = 0$ .

Even when the clean limit given by Eqs. (17) and (2) is the same, the behavior at finite scattering rates is very different whenever a flat band is close to the Fermi energy. In a dispersive band, however, both methods give close results (see Fig. 1f). The discrepancy between the Kubo-Greenwood and Kubo-Streda formulae is thus a flat-band effect.

#### A. Origin of the discrepancy

In order to understand the differences in the results obtained from Eq. (17) and Eq. (2), let us derive Eq. (17) from Eq. (2).

Starting from Eq. (2), we write

$$\sigma_{\mu\nu}(\omega = 0) = -i\frac{e^2}{\hbar} \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} \int_{-\infty}^{\infty} d\epsilon \sum_{\mathbf{k}} \sum_{m \neq n} [j_{\mu}]_{nm} [j_{\nu}]_{mn} \left( \frac{n_F(\epsilon)\delta(\epsilon - \epsilon_n)}{(\epsilon - \epsilon_m)(\epsilon - \epsilon_m + i\eta)} - \frac{n_F(\epsilon)\delta(\epsilon - \epsilon_m)}{(\epsilon_n - \epsilon)(\epsilon_n - \epsilon + i\eta)} \right), \quad (18)$$

Using that  $\lim_{\eta \rightarrow 0^+} 1/[(\epsilon - \epsilon_n)(\epsilon - \epsilon_n + i\eta)] = -\lim_{\eta \rightarrow 0^+} \partial_{\epsilon}(\epsilon - \epsilon_n + i\eta)^{-1}$ , we obtain the Kubo-Bastin formula

$$\sigma_{\mu\nu}(\omega = 0) = i\frac{e^2}{\hbar} \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} \int_{-\infty}^{\infty} d\epsilon n_F(\epsilon) \text{Tr} \left[ j_{\mu} \frac{\partial G_{\mathbf{k}}(\epsilon + i\eta)}{\partial \epsilon} j_{\nu} \delta(\epsilon - H) - j_{\mu} \delta(\epsilon - H) j_{\nu} \frac{\partial G_{\mathbf{k}}(\epsilon - i\eta)}{\partial \epsilon} \right], \quad (19)$$

A detailed derivation of the full Kubo-Streda formula from this form is given by Crépieux et al. in Ref. 60. Here, we will focus on the symmetric part of the conductivity, which reads

$$\sigma_{\mu\nu}^s(\omega = 0) = \frac{e^2}{\hbar} \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} \int_{-\infty}^{\infty} d\epsilon n_F(\epsilon) \times \text{Tr} \left[ j_{\mu} \frac{\partial}{\partial \epsilon} \text{Im}[G(\epsilon - i\eta)] j_{\nu} \delta(\epsilon - H) + j_{\mu} \delta(\epsilon - H) j_{\nu} \frac{\partial}{\partial \epsilon} \text{Im}[G(\epsilon - i\eta)] \right] \quad (20)$$

If we replace the delta functions by a Lorentzian, we can write  $\pi\delta(\epsilon - H) = \lim_{\eta' \rightarrow 0^+} \text{Im}G(\epsilon - i\eta')$ , where  $\eta'$  can generally be different from the scattering rate in  $G(\epsilon - i\eta)$ . With  $\eta' = \eta$ , we obtain precisely the formula (4) through integration by parts. Taking  $\eta' = \eta$  should not change the result in the clean limit as long as the limit  $\lim_{\eta' \rightarrow 0^+, \eta \rightarrow 0^+}$  does not depend on the direction it is taken in. However, this is not always the case



at  $T = 0$ . At exactly zero temperature, Eq. (20) becomes

$$\sigma_{\mu\nu}^s(\omega = 0) = \sigma_{\mu\nu}^I + \sigma_{\mu\nu}^{II}, \text{ where}$$

$$\sigma_{\mu\nu}^I = \frac{e^2}{2\hbar\pi} \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} \text{Tr} [j_\mu \text{Im}[G(\mu - i\eta)] j_\nu \text{Im}[G(\mu - i\eta')] + j_\mu \text{Im}[G(\mu - i\eta')] j_\nu \text{Im}[G(\mu - i\eta)]] \quad (21)$$

$$= \frac{e^2}{2\hbar\pi} \sum_{mn} \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} \left( \frac{\eta\eta'}{[(\epsilon_n - \mu)^2 + \eta^2][(\epsilon_m - \mu)^2 + \eta'^2]} + \frac{\eta\eta'}{[(\epsilon_n - \mu)^2 + \eta'^2][(\epsilon_m - \mu)^2 + \eta^2]} \right) [j_\mu]_{mn} [j_\nu]_{nm} \quad (22)$$

$$\sigma_{\mu\nu}^{II} = \frac{e^2}{2\hbar\pi} \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} \int_{-\infty}^{\mu} d\epsilon \text{Tr} [j_\mu \frac{\partial}{\partial \epsilon} \text{Im}[G(\epsilon - i\eta)] j_\nu \text{Im}[G(\epsilon - i\eta')] + j_\mu \text{Im}[G(\epsilon - i\eta')] j_\nu \frac{\partial}{\partial \epsilon} \text{Im}[G(\epsilon - i\eta)] \\ - j_\mu \text{Im}[G(\epsilon - i\eta)] j_\nu \frac{\partial}{\partial \epsilon} \text{Im}[G(\epsilon - i\eta')] - j_\mu \frac{\partial}{\partial \epsilon} \text{Im}[G(\epsilon - i\eta')] j_\nu \text{Im}[G(\epsilon - i\eta)]]]. \quad (23)$$

The contribution  $\sigma_{\mu\nu}^I$  causes the discrepancy between the Kubo-Greenwood and the Kubo-Streda formula in flat bands. Note that the double limit  $\lim_{\eta \rightarrow 0^+, \eta' \rightarrow 0^+}$  of  $\eta\eta'/[(\epsilon_n - \mu)^2 + \eta^2][(\epsilon_m - \mu)^2 + \eta'^2]$  depends on the direction of the limits whenever only one of  $\epsilon_n$  or  $\epsilon_m$  is equal to  $\mu$ . For instance, if  $\epsilon_n = \mu$  and  $\epsilon_m \neq \mu$ , the limit diverges if we first take  $\eta \rightarrow 0^+$ , vanishes if we take  $\eta' \rightarrow 0^+$  before  $\eta \rightarrow 0^+$ , and gives the nonzero finite result  $1/(\epsilon_m - \epsilon_n)^2$  if  $\eta = \eta'$ . The integrand in Eq. (22) is thus problematic whenever one of  $\epsilon_n$  or  $\epsilon_m$  is equal to the Fermi energy. In other words, the contribution coming from states at the Fermi energy to the interband part of the conductivity can be inaccurate.

In a dispersive band where the Fermi surface is  $D-1$  dimensional, the states at exactly the Fermi energy will not contribute in the thermodynamic limit, since their area in the Brillouin zone vanishes. However, if the Fermi energy is in a (partially) flat band, these contributions appear in the final result. In particular, if we set  $\eta = \eta'$  when taking the clean limit, the resulting DC conductivity when the chemical potential is tuned into a flat band is proportional to the integrated quantum metric of the flat band  $n$ ,

$$\sigma_{\mu\nu} = \frac{e^2}{\hbar\pi} \int_{\text{B.z.}} \frac{d^D \mathbf{k}}{(2\pi)^D} \sum_{m \neq n} \langle \partial_\mu n_{\mathbf{k}} | m_{\mathbf{k}} \rangle \langle m_{\mathbf{k}} | \partial_\nu n_{\mathbf{k}} \rangle. \quad (24)$$

However, this is only the case when taking the limit along  $\eta = \eta'$ , and the double limit is actually not well-defined.

The source of the problem is the introduction of the derivative of the Fermi distribution. Above, we already mentioned that integrating  $\partial n_F(\epsilon)/\partial \epsilon|_{\epsilon=\epsilon_n(\mathbf{k})}$  over the Brillouin zone causes problems when  $\epsilon_n(\mathbf{k})$  is constant. Here, it may appear that the same problem is no longer present, because we could integrate  $\partial n_F(\epsilon)/\partial \epsilon|_{\epsilon=\epsilon_n(\mathbf{k})}$  over the energy  $\epsilon$ . However, if we take  $\eta' \rightarrow 0^+$  first before  $\eta \rightarrow 0^+$ , we recover the delta-function first introduced in Eq. (18). Now that we already integrated over the energy to get rid of the delta function coming from  $\partial n_F(\epsilon)/\partial \epsilon|_{\epsilon=\epsilon_n(\mathbf{k})}$ , the remaining delta-function has transformed into  $\delta(\mu - \epsilon_n(\mathbf{k}))$ . Again, the integral

over the Brillouin zone is no longer well-defined when the chemical potential is in a dispersionless flat band.

#### IV. CONCLUSIONS AND DISCUSSION

We calculated the DC conductivity in a multiband system using the Kubo-Greenwood formula and the Kubo-Streda formula, and scrutinized various approximations used in the literature. Our focus was analyzing the DC conductivity in an isolated (gapped from other bands) flat band and its potential connection to quantum geometry. We summarize here our findings and discuss their implications.

The Kubo-Greenwood formalism, without approximate use of derivatives of the Fermi function, predicts vanishing DC conductivity in a flat band in the clean limit  $\eta \rightarrow 0$ . This is physically intuitive considering that single particles have infinite effective mass and DC conductivity is essentially single particle transport in a system with no correlations but only a (vanishingly small) scattering rate  $\eta$ . The Kubo-Greenwood formula gives a vanishing DC conductivity in the clean limit in the dimerized SSH chain, consistent with the fact that transport through the chain is impossible since it is disconnected. In our view, the Kubo-Greenwood result of zero DC conductivity is the physically correct description of non-interacting electron transport in a flat band.

At finite  $\eta$ , the Kubo-Greenwood formula gives a DC conductivity related to the quantum metric, but this is simply the DC tail of the AC conductivity resonance at the band gap frequency. We argue that this contribution, proportional to  $\eta$ , is not physically meaningful in a flat band where  $\eta$  cannot be claimed to be both finite and “small”, since there is no kinetic energy (band width) to compare it with. In other words, any small scattering, if considered to be truly finite, i.e.  $\eta \neq 0$ , is actually a strong interaction/perturbation in a flat band. Thus, various approximations done in the derivation of the Kubo-Greenwood and related formulas, such as inde-

pendent electrons, or neglecting vertex corrections, can be questioned.

The Kubo-Streda formula gives results very different from the Kubo-Greenwood one in a flat band, and we argue they do not describe the limit of a dispersionless band correctly. This becomes apparent when they predict finite DC conductivity at zero temperature even in the completely disconnected SSH model where a DC current through the system clearly cannot flow. We point out in detail where the problems with this formula arise: essentially, they boil down to the lack of a Fermi surface. In a flat band, the Fermi energy is massively degenerate and forms a volume (3D) or a surface (2D) in momentum space. Therefore any unphysical features at the Fermi energy arising from approximations become finite, while in a dispersive band they would vanish within an integral over the Brillouin zone since there the Fermi surface is an area (3D) or a line (2D) of zero measure. One might think this could be solved by changing the integration variable from momentum to energy, but that would involve introducing the density of states which diverges in a flat band; the same problem dressed in a different way. Thus, in general, in studies of linear response phenomena in the flat band limit, one needs to be cautious with commonly applied approximations and formulas, since many of them are valid and physically meaningful only in the presence of a Fermi surface. Several recent studies of conductivity in a flat band<sup>40–42,48,49</sup> should probably be revisited to understand the potential implications of our results there.

Our results relate also to the subtle connections between quantum geometry, physical observables, and the orbital positions in a lattice system. The symmetric and antisymmetric components of the quantum geomet-

ric tensor, namely the quantum metric and Berry curvature, depend on orbital positions (i.e. change of orbital coordinates while keeping the connectivity (hopping) between the orbitals the same). It is well known that many physical observables, such as the anomalous Hall conductivity, are determined by the Berry curvature and thus depend on orbital positions. Flat band superconductivity was predicted to be proportional to the quantum metric (which is orbital dependent) within a large number of studies (see<sup>1,39</sup> and references therein), while the definition of the superfluid weight is clearly independent of orbital positions. This discrepancy was solved only recently by showing that the superfluid weight is actually related to the minimal quantum metric, an orbital-independent quantity.<sup>10</sup> Here we showed that the flat band AC conductivity can have a relation to the quantum metric, while the DC conductivity obviously is not related to any quantum geometric quantity as it vanishes. We believe that it is probably possible to determine, at a general level, which physical observables are orbital-position dependent simply based on Maxwell's equations, general properties of the (superconducting or single electron) wavefunction, and gauge invariance. We leave this to a future work. Another important future research problem is to analyze transport in the case where other bands are touching the flat band.

## ACKNOWLEDGMENTS

We thank Johannes Mitscherling for useful discussions. We acknowledge support by the Academy of Finland under the project number 349313. K.-E.H. acknowledges financial support by the Magnus Ehrnrooth Foundation.

---

\* [kukka-emilia.huhtinen@aalto.fi](mailto:kukka-emilia.huhtinen@aalto.fi)

† [paivi.torma@aalto.fi](mailto:paivi.torma@aalto.fi)

<sup>1</sup> Sebastiano Peotta and Päivi Törmä, “Superfluidity in topologically nontrivial flat bands,” *Nature Communications* **6**, 8944 (2015).

<sup>2</sup> Long Liang, Tuomas I. Vanhala, Sebastiano Peotta, Topi Siro, Ari Harju, and Päivi Törmä, “Band geometry, Berry curvature, and superfluid weight,” *Physical Review B* **95**, 024515 (2017).

<sup>3</sup> Aleksi Julku, Sebastiano Peotta, Tuomas I. Vanhala, Dong-Hee Kim, and Päivi Törmä, “Geometric Origin of Superfluidity in the Lieb-Lattice Flat Band,” *Physical Review Letters* **117**, 045303 (2016).

<sup>4</sup> Enrico Rossi, “Quantum metric and correlated states in two-dimensional systems,” *Current Opinion in Solid State and Materials Science* **25**, 100952 (2021).

<sup>5</sup> P. Törmä, L. Liang, and S. Peotta, “Quantum metric and effective mass of a two-body bound state in a flat band,” *Physical Review B* **98**, 220511 (2018).

<sup>6</sup> A. Julku, T. J. Peltonen, L. Liang, T. T. Heikkilä, and P. Törmä, “Superfluid weight and Berezinskii-Kosterlitz-Thouless transition temperature of twisted bi-

layer graphene,” *Physical Review B* **101**, 060505 (2020).

<sup>7</sup> Xiang Hu, Timo Hyart, Dmitry I. Pikulin, and Enrico Rossi, “Geometric and Conventional Contribution to the Superfluid Weight in Twisted Bilayer Graphene,” *Physical Review Letters* **123**, 237002 (2019).

<sup>8</sup> Fang Xie, Zhida Song, Biao Lian, and B. Andrei Bernevig, “Topology-Bounded Superfluid Weight in Twisted Bilayer Graphene,” *Physical Review Letters* **124**, 167002 (2020).

<sup>9</sup> M. Iskin, “Two-body problem in a multiband lattice and the role of quantum geometry,” *Phys. Rev. A* **103**, 053311 (2021).

<sup>10</sup> Kukka-Emilia Huhtinen, Jonah Herzog-Arbeitman, Aaron Chew, Bogdan A. Bernevig, and Päivi Törmä, “Revisiting flat band superconductivity: Dependence on minimal quantum metric and band touchings,” *Phys. Rev. B* **106**, 014518 (2022).

<sup>11</sup> M. Iskin, “Effective-mass tensor of the two-body bound states and the quantum-metric tensor of the underlying Bloch states in multiband lattices,” *Phys. Rev. A* **105**, 023312 (2022).

<sup>12</sup> Frédéric Piéchon, Arnaud Raoux, Jean-Noël Fuchs, and Gilles Montambaux, “Geometric orbital susceptibility:

- Quantum metric without berry curvature,” *Phys. Rev. B* **94**, 134423 (2016).
- <sup>13</sup> Yang Gao, Shengyuan A. Yang, and Qian Niu, “Geometrical effects in orbital magnetic susceptibility,” *Phys. Rev. B* **91**, 214405 (2015).
  - <sup>14</sup> Tobias Holder, Daniel Kaplan, and Binghai Yan, “Consequences of time-reversal-symmetry breaking in the light-matter interaction: Berry curvature, quantum metric, and diabatic motion,” *Phys. Rev. Research* **2**, 033100 (2020).
  - <sup>15</sup> Gabriel E. Topp, Christian J. Eckhardt, Dante M. Kennes, Michael A. Sentef, and Päivi Törmä, “Light-matter coupling and quantum geometry in moiré materials,” *Phys. Rev. B* **104**, 064306 (2021).
  - <sup>16</sup> D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, “Quantized hall conductance in a two-dimensional periodic potential,” *Phys. Rev. Lett.* **49**, 405–408 (1982).
  - <sup>17</sup> Qian Niu, D. J. Thouless, and Yong-Shi Wu, “Quantized hall conductance as a topological invariant,” *Phys. Rev. B* **31**, 3372–3377 (1985).
  - <sup>18</sup> Mahito Kohmoto, “Topological invariant and the quantization of the hall conductance,” *Annals of Physics* **160**, 343–354 (1985).
  - <sup>19</sup> Masaru Onoda and Naoto Nagaosa, “Topological nature of anomalous hall effect in ferromagnets,” *Journal of the Physical Society of Japan* **71**, 19–22 (2002), <https://doi.org/10.1143/JPSJ.71.19>.
  - <sup>20</sup> T. Jungwirth, Qian Niu, and A. H. MacDonald, “Anomalous hall effect in ferromagnetic semiconductors,” *Phys. Rev. Lett.* **88**, 207208 (2002).
  - <sup>21</sup> Jun-Won Rhim, Kyoo Kim, and Bohm-Jung Yang, “Quantum distance and anomalous landau levels of flat bands,” *Nature* **584**, 59–63 (2020).
  - <sup>22</sup> Ahmed Abouelkomsan, Kang Yang, and Emil J. Bergholtz, “Quantum metric induced phases in moiré materials,” (2022), [arXiv:2202.10467](https://arxiv.org/abs/2202.10467).
  - <sup>23</sup> Yang Gao and Di Xiao, “Nonreciprocal directional dichroism induced by the quantum metric dipole,” *Phys. Rev. Lett.* **122**, 227402 (2019).
  - <sup>24</sup> Junyeong Ahn, Guang-Yu Guo, Naoto Nagaosa, and Ashvin Vishwanath, “Riemannian geometry of resonant optical responses,” *Nature Physics* (2021), [10.1038/s41567-021-01465-z](https://doi.org/10.1038/s41567-021-01465-z).
  - <sup>25</sup> M. Iskin, “Geometric contribution to the goldstone mode in spin-orbit coupled fermi superfluids,” *Physica B: Condensed Matter* **592**, 412260 (2020).
  - <sup>26</sup> Aleks Juku, Georg M. Bruun, and Päivi Törmä, “Quantum geometry and flat band Bose-Einstein condensation,” *Phys. Rev. Lett.* **127**, 170404 (2021).
  - <sup>27</sup> A. Gianfrate, O. Bleu, L. Dominici, V. Ardizzone, M. De Giorgi, D. Ballarini, G. Lerario, K. West, L. N. Pfeiffer, D. D. Solnyshkov, D. Sanvitto, and G. Malpuech, “Measurement of the Quantum Geometric Tensor and of the Anomalous Hall Drift,” *Nature* **578**, 381–385 (2020).
  - <sup>28</sup> J. P. Provost and G. Vallee, “Riemannian structure on manifolds of quantum states,” *Communications in Mathematical Physics* **76**, 289 – 301 (1980).
  - <sup>29</sup> Yuan Cao, Valla Fatemi, Shiang Fang, Kenji Watanabe, Takashi Taniguchi, Efthimios Kaxiras, and Pablo Jarillo-Herrero, “Unconventional superconductivity in magic-angle graphene superlattices,” *Nature* **556**, 43–50 (2018).
  - <sup>30</sup> Yuan Cao, Valla Fatemi, Ahmet Demir, Shiang Fang, Spencer L. Tomarken, Jason Y. Luo, Javier D. Sanchez-Yamagishi, Kenji Watanabe, Takashi Taniguchi, Efthimios Kaxiras, Ray C. Ashoori, and Pablo Jarillo-Herrero, “Correlated insulator behaviour at half-filling in magic-angle graphene superlattices,” *Nature* **556** (2018).
  - <sup>31</sup> Matthew Yankowitz, Shaowen Chen, Hryhorii Polshyn, Yuxuan Zhang, K. Watanabe, T. Taniguchi, David Graf, Andrea F. Young, and Cory R. Dean, “Tuning superconductivity in twisted bilayer graphene,” *Science* **363**, 1059–1064 (2019).
  - <sup>32</sup> Jeong Min Park, Yuan Cao, Kenji Watanabe, Takashi Taniguchi, and Pablo Jarillo-Herrero, “Tunable strongly coupled superconductivity in magic-angle twisted trilayer graphene,” *Nature* **590**, 249–255 (2021).
  - <sup>33</sup> Cheng Shen, Yanbang Chu, QuanSheng Wu, Na Li, Shuopei Wang, Yanchong Zhao, Jian Tang, Jieying Liu, Jinpeng Tian, Kenji Watanabe, Takashi Taniguchi, Rong Yang, Zi Yang Meng, Dongxia Shi, Oleg V. Yazyev, and Guangyu Zhang, “Correlated states in twisted double bilayer graphene,” *Nature Physics* **16**, 520–525 (2020).
  - <sup>34</sup> Yuan Cao, Daniel Rodan-Legrain, Oriol Rubies-Bigorda, Jeong Min Park, Kenji Watanabe, Takashi Taniguchi, and Pablo Jarillo-Herrero, “Tunable correlated states and spin-polarized phases in twisted bilayer-bilayer graphene,” *Nature* **583** (2020).
  - <sup>35</sup> Xiaobo Lu, Petr Stepanov, Wei Yang, Ming Xie, Mohammed Ali Aamir, Ipsita Das, Carles Urgell, Kenji Watanabe, Takashi Taniguchi, Guangyu Zhang, Adrian Bachtold, Allan H. MacDonald, and Dmitri K. Efetov, “Superconductors, orbital magnets and correlated states in magic-angle bilayer graphene,” *Nature* **574**, 653–657 (2019).
  - <sup>36</sup> Haidong Tian, Shi Che, Tianyi Xu, Patrick Cheung, Kenji Watanabe, Takashi Taniguchi, Mohit Randeria, Fan Zhang, Chun Ning Lau, and Marc W. Bockrath, “Evidence for flat band Dirac superconductor originating from quantum geometry,” (2021), [arXiv:2112.13401 \[cond-mat.supr-con\]](https://arxiv.org/abs/2112.13401).
  - <sup>37</sup> Leon Balents, Cory R. Dean, Dmitri K. Efetov, and Andrea F. Young, “Superconductivity and strong correlations in moiré flat bands,” *Nature Physics* **16**, 725–733 (2020).
  - <sup>38</sup> Eva Y. Andrei, Dmitri K. Efetov, Pablo Jarillo-Herrero, Allan H. MacDonald, Kin Fai Mak, T. Senthil, Emanuel Tutuc, Ali Yazdani, and Andrea F. Young, “The marvels of moiré materials,” *Nature Reviews Materials* **6**, 201–206 (2021).
  - <sup>39</sup> Päivi Törmä, Sebastiano Peotta, and Bogdan A. Bernevig, “Superconductivity, superfluidity and quantum geometry in twisted multilayer systems,” *Nature Reviews Physics* (2022), [10.1038/s42254-022-00466-y](https://doi.org/10.1038/s42254-022-00466-y).
  - <sup>40</sup> G. Bouzerar and D. Mayou, “Quantum transport in self-similar graphene carpets,” *Phys. Rev. Research* **2**, 033063 (2020).
  - <sup>41</sup> G. Bouzerar and D. Mayou, “Quantum transport in flat bands and supermetallicity,” *Phys. Rev. B* **103**, 075415 (2021).
  - <sup>42</sup> Aires Ferreira and Eduardo R. Mucciolo, “Critical delocalization of chiral zero energy modes in graphene,” *Phys. Rev. Lett.* **115**, 106601 (2015).
  - <sup>43</sup> Zheyong Fan, Andreas Uppstu, and Ari Harju, “Anderson localization in two-dimensional graphene with short-range disorder: One-parameter scaling and finite-size effects,” *Phys. Rev. B* **89**, 245422 (2014).
  - <sup>44</sup> Alessandro Cresti, Frank Ortmann, Thibaud Louvet, Dinh Van Tuan, and Stephan Roche, “Broken symmetries, zero-energy modes, and quantum transport in disordered



- graphene: From supermetallic to insulating regimes,” *Phys. Rev. Lett.* **110**, 196601 (2013).
- <sup>45</sup> Máté Vigh, László Oroszlány, Szabolcs Vajna, Pablo San-Jose, Gyula Dávid, József Cserti, and Balázs Dóra, “Diverging dc conductivity due to a flat band in a disordered system of pseudospin-1 dirac-weyl fermions,” *Phys. Rev. B* **88**, 161413 (2013).
- <sup>46</sup> J. Wang, J. F. Liu, and C. S. Ting, “Recovered minimal conductivity in the  $\alpha$ - $T_3$  model,” *Phys. Rev. B* **101**, 205420 (2020).
- <sup>47</sup> Johannes Mitscherling and Tobias Holder, “Bound on resistivity in flat-band materials due to the quantum metric,” *Phys. Rev. B* **105**, 085154 (2022).
- <sup>48</sup> Bruno Mera and Johannes Mitscherling, “Nontrivial quantum geometry of degenerate flat bands,” *Phys. Rev. B* **106**, 165133 (2022).
- <sup>49</sup> G. Bouzerar, “Giant boost of the quantum metric in disordered one-dimensional flat-band systems,” *Phys. Rev. B* **106**, 125125 (2022).
- <sup>50</sup> K. Ziegler, “Minimal conductivity of graphene: Nonuniversal values from the kubo formula,” *Phys. Rev. B* **75**, 233407 (2007).
- <sup>51</sup> L. Calderín, V.V. Karasiev, and S.B. Trickey, “Kubo-greenwood electrical conductivity formulation and implementation for projector augmented wave datasets,” *Computer Physics Communications* **221**, 118–142 (2017).
- <sup>52</sup> Jan M. Tomczak and Silke Biermann, “Optical properties of correlated materials: Generalized peierls approach and its application to  $\text{VO}_2$ ,” *Phys. Rev. B* **80**, 085117 (2009).
- <sup>53</sup> R. Nourafkan and A.-M. S. Tremblay, “Hall and faraday effects in interacting multiband systems with arbitrary band topology and spin-orbit coupling,” *Phys. Rev. B* **98**, 165130 (2018).
- <sup>54</sup> Johannes Mitscherling, “Longitudinal and anomalous hall conductivity of a general two-band model,” *Phys. Rev. B* **102**, 165151 (2020).
- <sup>55</sup> Steven H. Simon and Mark S. Rudner, “Contrasting lattice geometry dependent versus independent quantities: Ramifications for berry curvature, energy gaps, and dynamics,” *Phys. Rev. B* **102**, 165148 (2020).
- <sup>56</sup> D A Greenwood, “The boltzmann equation in the theory of electrical conduction in metals,” *Proceedings of the Physical Society* **71**, 585 (1958).
- <sup>57</sup> Ryogo Kubo, “Statistical-mechanical theory of irreversible processes. i. general theory and simple applications to magnetic and conduction problems,” *Journal of the Physical Society of Japan* **12**, 570–586 (1957), <https://doi.org/10.1143/JPSJ.12.570>.
- <sup>58</sup> A. Bastin, C. Lewiner, O. Betbedermatibet, and P. Nozieres, “Quantum oscillations of the hall effect of a fermion gas with random impurity scattering,” *Journal of Physics and Chemistry of Solids* **32**, 1811–1824 (1971).
- <sup>59</sup> P Streda, “Theory of quantised hall conductivity in two dimensions,” *Journal of Physics C: Solid State Physics* **15**, L717 (1982).
- <sup>60</sup> A. Crépieux and P. Bruno, “Theory of the anomalous hall effect from the kubo formula and the dirac equation,” *Phys. Rev. B* **64**, 014416 (2001).
- <sup>61</sup> Varga Bonbien and Aurélien Manchon, “Symmetrized decomposition of the kubo-bastin formula,” *Phys. Rev. B* **102**, 085113 (2020).
- <sup>62</sup> R. Resta, “The insulating state of matter: a geometrical theory,” *The European Physical Journal B* **79**, 121 (2011).
- <sup>63</sup> Di Xiao, Ming-Che Chang, and Qian Niu, “Berry phase effects on electronic properties,” *Rev. Mod. Phys.* **82**, 1959–2007 (2010).