Symmetries in one loop solutions: The AV, AVV, and AVVV diagrams, from 2D, 4D, and 6D dimensions and the role of breaking integration linearity.

L. Ebani^{1*}, T. J. Girardi^{1†}, J. F. Thuorst^{1‡}

¹CBPF-Centro Brasileiro de Pesquisas Físicas, 22290-180, Rio de Janeiro, RJ, Brazil

We investigated relations among green functions defined in the context of an alternative strategy for coping with the divergences, also called Implicit Regularization. Our targets are fermionic amplitudes in even space-time dimensions, where anomalous tensors connect to finite amplitudes. Those tensors depend on surface terms, whose non-zero values arise from finite amplitudes as requirements of consistency with the linearity of integration and uniqueness. Maintaining these terms implies breaking momentum-space homogeneity and in a later step the Ward identities. Meanwhile, eliminating them allows more than one mathematical expression for the same amplitude. That is a consequence of choices related to the involved Dirac traces. Independently of divergences, it is impossible to satisfy all symmetry implications that require the vanishing of surface terms and linearity simultaneously. Nonetheless, the symmetry violations are globally independent of divergences and can be allocated appropriately. From this perspective, we cast all the choices involved and the different meanings, whose implications go beyond the scenario described.

Keywords: Perturbative calculations, anomalies, linearity and uniqueness

1. INTRODUCTION

Since their inception, anomalies have been part of the culture of practitioners of quantum field theory (QFT). The issue surges in multiple avatars and is intricately linked to the conservation or not of the classical currents in the operator framework of quantum theory. Met in the end forties and, at the outset of the fifties by the authors [1], [2], [3] and [4]. Rediscovered in two dimensions (2D) [5] and in four dimensions (4D) by [6] [7] [8], the ABBJ anomaly of the triangle's graph. Later, many studies considered perturbative and non-perturbative approaches to investigate these phenomena, mainly the Fujikawa interpretation [9], [10] of the path-integral's measure, and heat kernel expansions, see [11] for a review, as well as cohomological methods [9]. One of the reasons for its importance comes from the fact that anomalies prevent the quantum expectation value of Noether currents from satisfying their classical symmetries. Hence, the anomaly manifestation breaks Ward-Takahashi identities (WI) or Slavnov-Taylor identities to non-abelian gauge symmetry, which are needed for the perturbative renormalizability of the gauge models, even when spontaneous symmetry breaking is present as in the case of the physical theory Standard Model. Another one the role in the phenomenological description of particle decays, the neutral pion the most emblematic one.

Their general and most notable manifestation is in the perturbative scenario where correlators of the axial and vector currents that are odd, linear divergent tensors corresponding to the AV^n amplitudes, in d = 2n dimensions, can not have all their WIs satisfied. All of them have three structural properties.

First, in the last instance, they are (n + 1)-rank tensor of odd parity, functions of n momenta variables, that for such features posses, through contractions with their momenta variables, a set of low energy theorems, such theorems come as a consequence of their WIs. One of these identities,

^{*} luci.ebani@gmail.com

[†] thalisjg@gmail.com

[‡] jfernando.th@gmail.com, corresponding author

axial-WI, relates the same tensor, in some way, to the amplitude PV^n , whose perturbative expression is a finite integral, and this expression behaves in the point where the low energy theorem is stated in such a way that either the PV^n have a value that contradicts its explicit form or some Ward identity is unavoidably violated. The anomaly.

Second, a property that is shared by those tensors, they have minimal dimensionality such that they rise from Dirac traces that cast two more gamma matrices than the dimension they are defined. This trace of 2n + 2 gamma matrices that are combinations of metric and Levi-Civita tensors are known to exhibit a multitude of identical expressions, where they display explicitly, different sets of indexes, signs, and number of monomials. All of them differ by sums of totally antisymmetric tensors in 2n + 1 indexes, therefore they are zero at 2n dimensions. The choice of one or another form is arbitrary nonetheless.

Finally, their power counting leaves the possible presence of non-identically vanishing surface terms opened, making these structures depending on how the external momenta percolate the graphs. That means, the final results generally show up many possibilities, since perturbative solutions may unavoidably depend on arbitrary choices for routings and Dirac traces.

This last proposition is inseparable from the fact that in perturbation theory, divergences are the rule, in order to get some model predictions of QFT, which means to acquire some information of the kinematical dependence, symmetries, and so on, regularization techniques are adopted in an attempt to circumvent these problems, as some examples, cut-off, Pauli-Villars, analytic regularization, dimensional regularization (DR) [12] [13], high covariant regularization, differential renormalization, and new methods have, until recently, been proposed to deal with multi-loop calculation and aiming to an algorithmic implementation for high precision numerical predictions [14] [15]. Modifying, invariably, the amplitudes by making divergent amplitudes finite, manipulations forbidden, or not guaranteed to the original expressions become valid, such as shifts in the integration variable. And determining how and which properties and parts of the algebras present must be dealt with [16][17][18][19][20][21], things that are not inherent to the Feynman rules defining the objects under investigation. As in the case of DR, this process is guided by symmetry preservation and implies the elimination of surface terms. Then in a later step, combining regularization and renormalization, predictions can be established and compared to the experimental data resulting in the success known, for example, in the Quantum Electrodynamics/Standard Model.

On the other hand, Feynman integrals of power counting linear or higher and logarithmic, but tensors ones, own surface terms. For the linear ones, a shift in the integration variable is acknowledged to require compensation through non-zero surface terms [22] [23] [9], hence they can not be free-shifted, arbitrary routings are the rule. It is a manifestation of the internal momenta arbitrariness, notwithstanding they satisfy the energy-momentum conservation, as the differences in the routings are functions of the physical momenta, by themselves and the sums of these routings are arbitrary, and they may even assume non-covariant expressions [24]. Given the fact that nonzero surface terms imply the breaking of translational symmetry in the momentum space, and this operation is thoroughly needed to prove WIs, then it is reasonable that other violations of symmetries occur in such calculations, and in fact, they do.

The question that is proposed and answered in this contribution is: What is the extension of the consequences resulting from the three properties of the tensors mentioned, over symmetries of the amplitudes and the mathematical content of the diagrams, that means, the integration linearity, translational symmetry, and uniqueness¹? That is independent of a particular set of rules, even if some rule should be elected in some very final level of analysis.

For such purpose, we use a general model where these aspects can be generated: Spin-1/2

 $^{^{1}}$ To uniqueness, which needs a particular definition to work its consequences, we provide it along the paper.

fermions coupled with boson fields of spin-zero and one, of even and odd parity. The *n*-vertices polygon graphs of spin 1/2 internal propagators are the center of the analysis, specifically the 2D-AV and VA bubbles, 4D-AVV, VAV, and VVA triangles, and 6D-AVVV box. The results are obtained within the context of an alternative procedure to handle divergent and finite integrals also, that are independent of a particular regularization, introduced in the Ph.D. thesis of O.A. Battistel [25]. This strategy has been applied in a large number of investigations, for example in two dimensions [26] [27] [28], in four dimensions [29] [30], in six dimensions [31], in odd five-dimensions (5D) [32], and in even and odd, from two to six, dimensions [33]. An incarnation of this strategy has been known in the literature as Implicit Regularization (IREG), and it has been applied in similar investigations [34] [35] [36].

The idea is very simple, the divergences are isolated by means of an identity of universal applicability that does not interfere with the Feynman rules used. Insofar as diverging integrals are not evaluated explicitly and the amplitudes are not modified at early steps of the calculations. Arbitrary routings are adopted for the momenta of internal lines to preserve their intrinsic arbitrariness which is a feature of such perturbative calculations. Furthermore, in the strategy, we devise a notational scheme to systematize the finite integrals and their divergent parts that appear in this work, based on previous works about the subject, ref. [37] for single-mass integrals, and ref.[38][39] for multiples masses case. There are three ingredients for such: irreducible divergent objects, tensor surface terms, and finite functions. It is precisely this organization that allows a clear view of the relevant points of our discussion.

The unique assumption made is that the linearity applies to Feynman integrals, typical quantities of the perturbative calculations. At the amplitude's level, the linearity of integration arises in the relations among green functions (RAGFs) derived for the well-defined integrands, that when integrated, the critical step of computations, embodies the referred property. This aspect is one of the main points of the investigation since if it was present automatically, it would be possible to prove all of Ward's identities by methods blessed by translational symmetry as the DR.

Our stance on the undetermined character of perturbation theory quantities enables a clear view of the active elements that produce the variety of results, notably of how and where the presence of surface terms in amplitudes is connected with ambiguities and symmetry violations. As a result, in the first place, if one asks: Is there a unique tensor of the external momenta? The answer will be, to even amplitudes, yes, any interpretation of the divergences that make zero the surface terms render the results unique and, as a by-product, symmetric. But if one asks soon after: Does such action make all the amplitudes, including the odd ones, unique functions of the external momenta? The answer is no, in other words, there is more than one answer to be obtained, even if the divergences are not touched at all and are taken consistently with the even amplitudes, such attitude wreck the integration linearity and arbitrary combinations of equal integrands give rise to arbitrary distinct polynomials in the integrals, that means, once you lose uniqueness an uncountable (literally) number of tensors can be reached from the same expression.

The other side is deep as well, once the value, the unique one, that saves linearity and uniqueness is adopted, no matter what manipulation is used in the traces, even if bilinears are not reduced in the splitting of divergent parts, it provides one and only one tensor, here is the catch, of the routing variables, implying that physical interpretation asks for arbitrary parameters to account for the realization of the symmetries. In the odd amplitudes, such freedom will, as in any other arbitrariness situation, enable one to fix the known and desired content of the results. And the striking consequence is that if universality is asked to play a role, even amplitudes will always violate their WI.

The lack of momentum ambiguities leads to a lack of uniqueness in a sub-class of pseudo-tensor amplitudes, where belongs precisely the anomalous ones in even space-time dimensions, among them the AV^n amplitudes. Moreover, in dimensions equal to four or higher, more amplitudes show

4

the behavior described here, AV^r , for $n \leq r < 2n$. As an example, the 4D-AVVV box, will show a dependence on surface terms, but not on internal momenta, and a triangle topology with a tensor vertex, TAV, suffers from the same properties we present in this work.

To appreciate these statements we organized the work as follows. In section (2), we have the general model, definitions, and a preliminary discussion. Section (3) deals with the alternative to regularizations strategy in handling the divergences, where we give the general definition of the irreducible objects and tensor surface terms that appear everywhere in the work. A detailed compilation of the effects of traces and surface terms in two dimensions appears in section (4), there, for the first time, and in a simplified situation the linearity of integration and uniqueness are fully analyzed through complete and independent computation of all quantities involved in the relation among green functions, the consequences of adopting results saving linearity and of saving translational symmetry are presented, and interpreted in light of low-energy theorems. The most lengthy section (5), handles with all odd triangles, their RAGFs, and the nature of uniqueness that is more convoluted in that case, sub-sections (5.2) and (5.3) deal with the general properties of lowenergy theorems and offer a theorem connecting linearity, low-energy behavior of finite amplitudes in general, and surface terms. The last section (6), extends seamlessly the propositions put in the realm of six dimensions, [31] has already worked with one of the possibilities, and among all the other possibilities we chose one more to illustrate the behavior we have presented in two and four dimensions.

Finally in section (7), Final Remarks and Perspectives, we present comments on some fine points of the work and a timeline of the arguments leading to our main results and the character of some of our conceptual tools. Just as importantly, and integral part is the appendices, in (A) we show how the attitude present in the main body of the work, in four dimensions, is enough to give account for any non-trivial proposition. Appendices (B, C, and D) contain the divergent and finite parts, obtained through the strategy delineated in section (3) for any structure used in the paper, in addition to all the reductions and identities needed. The last appendix (E) is a compilation of results required in section (5) not present in the text due to size reasons.

2. NOTATION, DEFINITIONS, MODEL AND PRELIMINARIES

The Feynman rules, vertexes, and propagators, employed in this investigation come from a model where fermionic currents coupled to bosonic fields of even and odd parity $\{\Phi(x), V_{\mu}(x), \Pi(x), A_{\mu}(x)\}$ through the general interacting action

$$S_{I} = \int d^{2n}x \left[e_{S}S(x) \Phi(x) + e_{\Pi}P(x) \Pi(x) + e_{V}J^{\mu}(x) V_{\mu}(x) + e_{A}J^{\mu}_{*}(x) A_{\mu}(x) \right].$$
(2.1)

The currents $\{S, P, J_{\mu}, J_{*\mu}\}$ are bilinears in the fermionic fields, $J_i = \bar{\psi}(x) \Gamma_i \psi(x)$, and they deliver the vertexes proportional to

$$\Gamma_i \in (S, P, V, A) = (1, \gamma_*, \gamma_\mu, \gamma_* \gamma_\mu), \qquad (2.2)$$

the proportionality comes from the coupling constants $\{e_S, e_{\Pi}, e, e_A\}$ that are taken as the unit for our purposes, as they can be easily recovered if needed. The elements γ_{μ} are the generators of the Clifford algebra of Dirac matrices satisfying $\{\gamma^{\mu_1}, \gamma^{\mu_2}\} = 2g^{\mu_1\mu_2}$. The highest-weight element of the algebra, in d = 2n, is the chiral matrix of that dimension that satisfies $\{\gamma_*, \gamma^{\mu_k}\} = 0$, explicitly

$$\gamma_* = i^{n-1} \gamma_0 \gamma_1 \cdots \gamma_{2n-1} = i^{n-1} \prod_{s=0}^{2n-1} \gamma_s = \frac{i^{n-1}}{(2n)!} \varepsilon_{\nu_1 \cdots \nu_{2n}} \gamma^{\nu_1 \cdots \nu_{2n}}.$$
 (2.3)

We often will adopt a notation of merging to the product of matrices $\gamma^{\nu_1\cdots\nu_{2n}} = \gamma^{\nu_1}\gamma^{\nu_2}\cdots\gamma^{\nu_{2n}}$ and will adapt to the Lorentz indexes, $\mu_1\mu_2\cdots\mu_s = \mu_{12\cdots s}$ when convenient and clear by context. The behavior under the permutation of the indexes is determined by the objects: $g_{\mu_1\mu_2} = g_{\mu_{12}} = g_{\mu_{21}}$ or $\varepsilon_{\mu_1\mu_2\cdots\mu_{2n}} = \varepsilon_{\mu_{12}\cdots\mu_{2n}} = -\varepsilon_{\mu_{21}\cdots\mu_{2n}}$. For the 2*n*-dimensional, follow the normalization $\varepsilon^{0123\cdots 2n-1} = 1$.

The elements of the algebra are the antisymmetrized products of gamma matrices

$$\gamma_{[\mu_1 \cdots \mu_r]} = \frac{1}{r!} \sum_{\pi \in S_r} \operatorname{sign}(\pi) \gamma_{\mu_{\pi(1)} \cdots \mu_{\pi(r)}}, \qquad (2.4)$$

that satisfies the general identities, see by example the appendix of the ref. [40].

$$\gamma_* \gamma_{[\mu_1 \cdots \mu_r]} = \frac{i^{n-1+r(r+1)}}{(2n-s)!} \varepsilon_{\mu_1 \cdots \mu_r} {}^{\nu_{r+1} \cdots \nu_{2n}} \gamma_{[\nu_{r+1} \cdots \nu_{2n}]}.$$
(2.5)

These identities are needed when taking traces with the chiral matrix.

The spinorial Feynman propagators come naturally from the standard kinetic term of Dirac fermions

$$S_F(K_i) = \frac{1}{(k_i - m + i0^+)} = \frac{(k_i + m)}{D_i}$$
(2.6)

where $D_i = K_i^2 - m^2$ and the momentum flowing through it

$$K_i = k + k_i, \tag{2.7}$$

where the k is the unrestricted loop momentum, and k_i are the routings that keep tracking of the flux of external momenta through the graph, see ref.[24]². They can not be written as a function of the kinematical data in the divergent integrals. In our approach, they codify the conditions of the satisfaction of symmetries or lack thereof. Nonetheless, their differences are related to the external momenta through the definition

$$p_{ij} = k_i - k_j, \tag{2.8}$$

using the momenta conservation in the vertexes of the diagram in the fig. (1)

From the diagram and Feynman rules follow the integrand of the amplitude

$$t^{\Gamma_{1}\Gamma_{2}\cdots\Gamma_{n_{1}}}\left(k_{1},\cdots,k_{n_{1}}\right) = \operatorname{tr}\left[\Gamma_{1}S_{F}\left(K_{1}\right)\Gamma_{2}S_{F}\left(K_{2}\right)\cdots\Gamma_{n_{1}}S_{F}\left(K_{n_{1}}\right)\right]$$

that is a well-defined function of the external momenta as the undetermined (by momentum conservation) sums

$$P_{ij} = k_i + k_j. \tag{2.9}$$

Often we are going to adopt the simplification $S(i) \equiv S_F(K_i)$, where numerical index *i* represents all the parameters of the corresponding line. The total amplitude comes from integration in the loop momenta

$$T^{\Gamma_1\Gamma_2\cdots\Gamma_s}(1,\cdots,s) = \int \frac{\mathrm{d}^{2n}k}{(2\pi)^{2n}} t^{\Gamma_1\Gamma_2\cdots\Gamma_s}(1,\cdots,s).$$
(2.10)

 $^{^{2}}$ Specifically in the section (4.1) for a rare comment on the level of arbitrariness of these routings.




FIG. 1: General diagram for the one-loop amplitudes of this work.

The vertexes Γ_i , when replaced by specific ones (2.2), the notation accompanies the Lorentz indexes.

It is possible to establish identities among the Green functions exhibiting Lorentz indexes from the vector and axial currents. They are commonly called relations among green functions (RAGF), see some references about [29][26][33]. The application of the relations has been used in a wide range of perturbative investigations in the scenario of IREG. Without claiming as such, they are often used in WI investigations. However, they can be considered as conditions on the linearity of integration before any symmetry problems arise. They function as constraints or guides even before symmetry-specific WIs are asked to play a consistent judgment role in Feynman perturbation diagrams.

As a working example let us take the amplitude AV^{r-1} ,

$$t^{AV\cdots V}_{\mu_1\mu_2\cdots\mu_r} = \operatorname{tr}[\gamma_*\gamma_{\mu_1}S(1)\gamma_{\mu_2}S(2)\cdots\gamma_{\mu_r}S(r)]$$
(2.11)

when contracted with $p_{21}^{\mu_2}$ in the vectorial vertex γ_{μ_2} it can have one of its propagators removed using (2.7) and (2.6) in the standard manipulation $p_{21} = k_2 - k_1 = S^{-1}(2) - S^{-1}(1)$. The result is direct the VRAGF

$$p_{21}^{\mu_2} t_{\mu_1 \mu_2 \cdots \mu_r}^{AV \cdots V} = \operatorname{tr}[\gamma_* \gamma_{\mu_1} S(1) \gamma_{\mu_3} S(3) \cdots \gamma_{\mu_r} S(r)] - \operatorname{tr}[\gamma_* \gamma_{\mu_1} S(2) \cdots \gamma_{\mu_r} S(r)].$$

The result is again an amplitude built out of the same rules. In this case, a difference between two such amplitudes

$$p_{21}^{\mu_2} t_{\mu_1 \mu_2 \cdots \mu_r}^{AV \cdots V} = t_{\mu_1 \hat{\mu}_2 \cdots \mu_r}^{AV \cdots V} (1, \hat{2}, \cdots, r) - t_{\mu_1 \hat{\mu}_2 \cdots \mu_r}^{AV \cdots V} (\hat{1}, 2, \cdots, r).$$
(2.12)

The hats mean the omission of the propagator corresponding to that routing and vertexes corresponding to the Lorentz index as well. In other words, the RHS contains lower point functions that in general (but not always) are singular under integration.

For the axial vertex

$$p_{r1}^{\mu_1} t_{\mu_{12}\cdots\mu_r}^{AV\cdots V} = \operatorname{tr}[S(r)\gamma_*S^{-1}(r)S(1)\gamma_{\mu_2}S(2)\cdots\gamma_{\mu_{r-1}}S(r-1)\gamma_{\mu_r}] \\ -\operatorname{tr}[\gamma_*\gamma_{\mu_2}S(2)\cdots\gamma_{\mu_r}S(r)].$$

We must use $S(r) \gamma_* S^{-1}(r) = (-\gamma_* - 2mS(r) \gamma_*)$, and the commutation product of the chiral and Dirac matrices. Thus follow the ARAGF

$$p_{r1}^{\mu_1} t_{\mu_{12}\cdots\mu_r}^{AV\cdots V} = t_{\mu_r\hat{\mu}_1\mu_2\cdots\mu_{r-1}}^{AVV\cdots V} (1, 2, \cdots, \hat{r}) - t_{\hat{\mu}_1\mu_2\cdots\mu_r}^{AVV\cdots V} (\hat{1}, 2, \cdots, r) - 2m t_{\mu_2\cdots\mu_r}^{PV\cdots V}.$$
(2.13)

After integration they become

$$p_{r1}^{\mu_1} T_{\mu_{12}\cdots\mu_r}^{AV\cdots V} = T_{\mu_r\hat{\mu}_1\cdots\mu_{r-1}}^{AVV\cdots V} (1, 2, \cdots, \hat{r}) - T_{\hat{\mu}_1\cdots\mu_r}^{AVV\cdots V} (\hat{1}, 2, \cdots, r) - 2m T_{\mu_2\cdots\mu_r}^{PV\cdots V}, \text{ and } p_{21}^{\mu_2} T_{\mu_{12}\cdots\mu_r}^{AV\cdots V} = T_{\mu_1\hat{\mu}_2\cdots\mu_r}^{AV\cdots V} (1, \hat{2}, \cdots, r) - T_{\mu_1\hat{\mu}_2\cdots\mu_r}^{AV\cdots V} (\hat{1}, 2, \cdots, r).$$

From these equations, it is clear that they embody the assumptions of linearity of integration in perturbative computations. This characteristic is not guaranteed for divergent amplitudes. We will expose this scenario through the full calculations of the amplitudes and their relations. Although these equations can be an important structural property, they are not a priori linked to the particularities of the model and its symmetries. However, after summing up all the contributions from the cross diagram (if applicable), the contraction with momenta, assuming some properties for the lower-point green functions, must correspond to the WIs.

The WIs are equations satisfied by the Green functions as a consequence of the continuous symmetries of the action. They are valid in the perturbative approximations built on Feynman's rules unless they are inevitably anomalous. They arise from the joint application of the algebra of the quantized currents and the motion's equations to the currents: $\partial_{\mu}J^{\mu} = 0$ and $\partial_{\mu}J^{\mu}_{*} = -2miP$, remember the currents are bilinears in the fermion. Their expression in the position space of the AWI

$$\frac{\partial}{\partial x_1^{\mu_1}} \left\langle J_*^{\mu_1}\left(x_1\right) J_{\mu_2}\left(x_2\right) \cdots J_{\mu_n}\left(x_n\right) \right\rangle = -2mi \left\langle P\left(x_1\right) J_{\mu_2}\left(x_2\right) \cdots J_{\mu_n}\left(x_n\right) \right\rangle, \qquad (2.14)$$

where $\langle \cdots \rangle = \langle 0 | T [\cdots] | 0 \rangle$, and of the VWI to the same correlator

$$\frac{\partial}{\partial x_2^{\mu_1}} \left\langle J_{*\mu_1}\left(x_1\right) J^{\mu_2}\left(x_2\right) \cdots J_{\mu_n}\left(x_n\right) \right\rangle = 0.$$
(2.15)

In our notation for the perturbative amplitudes, we must have an analogous equation

$$q_1^{\mu_1} T_{\mu_{12}\cdots\mu_r}^{A\to V\cdots V} = -2m T_{\mu_{2}\cdots\mu_r}^{PV\cdots V}; \quad q_2^{\mu_2} T_{\mu_{12}\cdots\mu_r}^{A\to V\cdots V} = 0; \cdots \quad q_r^{\mu_r} T_{\mu_{12}\cdots\mu_r}^{A\to V\cdots V} = 0.$$

The connection between the RAGF and the WIs is direct. The violation of the RAGF implies violations in the WIs. In this way, the satisfaction of the total set of WI will be conditional on satisfying all RAGFs plus translational invariant amplitudes in momentum space, a requirement that we will show to be impossible in general.

The last point in this section is related to the calculations themselves. To compute the amplitudes, we have got to take the Dirac traces using commutation relations of the algebra, after that, any amplitude is always expressed as linear combinations of bare Feynman integrals to which we adopt a definition

$$\bar{J}_{n_2}^{\mu_1\mu_2\cdots\mu_{n_1}}(1,2,\cdots,n_2) = \int \frac{\mathrm{d}^{2n}k}{(2\pi)^{2n}} \frac{K_i^{\mu_1}\cdots K_i^{\mu_{n_1}}}{D_1 D_2\cdots D_{n_2}},\tag{2.16}$$

simplifying the dependence of the functions on their arguments $f(k_1, k_2, \dots) = f(1, 2, \dots)$ and when convenient omitting them at all. The combination $K_i = k + k_i$ in the definition is only a convenience. To change from a reference routing k_j to k_i it is just a matter of recognizing the definition of p_{ij} in (2.8) and writing $K_i = K_j + p_{ij}$. These integrals have power-counting given by

$$\omega = 2n + n_1 - 2n_2, \tag{2.17}$$

being n_1 the tensor rank and n_2 the number of denominators present. Just a set of five types of integral will arise in each amplitude that we will investigate in this work, they are the theme of the section (3.2).

The amplitudes whose properties will be detailed are:

- The 2D bubbles: $T^{AV}_{\mu_1\mu_2}$; $T^{VA}_{\mu_1\mu_2}$
- The 4D triangles: $T^{AVV}_{\mu_1\mu_2\mu_3}$; $T^{VAV}_{\mu_1\mu_2\mu_3}$; $T^{VVA}_{\mu_1\mu_2\mu_3}$; $T^{AAA}_{\mu_1\mu_2\mu_3}$
- The 6D box: $T^{AVVV}_{\mu_1\mu_2\mu_3\mu_4}$

All of them are divergent odd tensors of rank at least two, in two dimensions a logarithmic power counting and in four and six dimensions a linear power counting after taking the Dirac traces. Because of their power counting, to determine the validity of linearity and the symmetries required for such tensor we will devise the procedure to handle the divergent integrals in the next section.

3. PROCEDURE TO HANDLE THE DIVERGENCES AND THE FINITE INTEGRALS

Before presenting the strategy used to solve the divergent amplitudes, let us make a digression about the divergent-integrals issue present in QFT.

It is known that the products of propagators, that are not regular distribution, are in general ill-defined, as a good example is the equation

$$I = \int \frac{d^4k}{(2\pi)^4} \operatorname{tr} \left[S_F(k-p) \, S_F(k) \right] = \int d^4x \operatorname{tr} \left[S_F^2(x) \right] e^{-ip \cdot x},\tag{3.1}$$

where in the LHS lays a divergent convolution integral of two Fourier transformed Feynman propagators in momentum space, as the RHS is the Fourier transform of the square of the propagator in position space. Both sides are not defined as distributions for the fact when the point-wise product of distributions does not exist the convolution product of their Fourier transform doesn't as well and vice versa. These are the short-distance UV singularities that are manifested in the divergences of the loop momentum integrals.

Their origins can be rigorously traced back to multiplications of singular distributions by discontinuous step function in the chronological ordering of operators in the interaction picture that leads, through the Wick theorem, to the Feynman rules, e.g., G.Scharf [41, 42], originally in Epstein and Glaser [43]. Although, the undefined Feynman diagrams can be averted by carefully studying the splitting of distributions with causal support in the setting of causal perturbation theory [44][45], where no divergent integral appears at all, we are going to keep working with the Feynman rules in the context of regularizations.

However, the systematic procedure to handle the divergences we will employ is slightly different from the usual regularizations. It is the framework known as Implicit Regularization (IREG) a method that was introduced and developed at the end of 1990's in the Ph.D. thesis of O.A. Battistel [25], whose the first references about the subject are [46, 47].

It has the objective to keep the connection at all moments with the expression of the bare Feynman rules removing all the parameters such as routings and masses from divergent integrals and putting them in, strictly, finite integrals that are integrated without restriction, while the divergent ones do not suffer any modification besides an organization in surface terms and irreducible scalar integrals.

This objective is realized by noticing that all Feynman integrals possess propagators-like functions, remember that $D_i = (k + k_i)^2 - m^2$ in eq. (2.6), that can be rewritten as

$$\frac{1}{D_i} = \frac{1}{D_\lambda + A_i} = \frac{1}{D_\lambda} \frac{1}{[1 - (-A_i/D_\lambda)]},$$
(3.2)

where $D_{\lambda} = k^2 - \lambda^2$ and $A_i = 2k \cdot k_i + (k_i^2 + \lambda^2 - m^2)$. It is worth noting that in this decomposition, the dependence of the parameters that are not integrated is contained only at A_i .

To motivate the first element used in this strategy, which means a general identity capable of realizing the aforementioned objectives, we shall digress shortly about the sum of the geometric progression of order N and ratio x and its relation with the behavior of the propagator, namely the sum $s = 1 + x + \cdots + x^N$, that is easily expressed in closed form, and then we can write succinctly

$$(1-x)^{-1} = \sum_{r=0}^{N} x^r + x^{N+1} (1-x)^{-1}, \qquad (3.3)$$

where immediately it is possible to determine the asymptotic behavior of the terms in this progression when is identified x in our organization of the propagator

$$x = -\frac{A_i}{D_\lambda} = \frac{2k \cdot k_i + (k_i^2 + \lambda^2 - m^2)}{(k^2 - \lambda^2)},$$
(3.4)

we see that the asymptotic behavior at infinity is $||k||^{-1}$. Thus follow that the sequence $\sum_{r=0}^{N} x^r$ depends on the routings only in the numerator, and as a polynomial, decays stronger as bigger it is N.

Being valid for arbitrary N and an identity, to any power counting, in a product of propagators, is always possible to obtain the external momenta, through the definition (2.8), in finite integrals because with the help of eq.'s (3.3) and (3.2), we get

$$\frac{1}{D_i} = \sum_{r=0}^{N} (-1)^r \frac{A_i^r}{D_\lambda^{r+1}} + (-1)^{N+1} \frac{A_i^{N+1}}{D_\lambda^{N+1} D_i}.$$
(3.5)

Making N equals the power counting ω , the last integral is finite and regularization independent. After this first step, we must mention that for any N the identity is independent of λ^2 , as can be verified by taking the derivative in relation to this parameter, in the end, this generates a connection among the divergent and finite parts of the integrals and/or amplitudes. It implies specific behavior to the basic divergent scalar integrals and it the straightforwardly satisfied, thus we will adopt the mass of the propagator $\lambda^2 = m^2$ as such a scale in this work.

Here again, to modularize the analysis and make clear the effects of divergent and finite parts over the definition of the amplitudes, we adopted this systematic: the finite integrals will be solved without restrictions, and the divergences will be kept without any further modification. They will not be resolved just standardized and basic properties for them will be established, as we will see next.

3.1. Divergent Terms

After applying conveniently the identity (3.5), the content of the Feynman integrals is going to be specified through, surface terms, irreducible divergent objects, and finite functions. To clarify, the divergent terms will appear as a set of pure integration-momentum integrals

$$\int \frac{\mathrm{d}^{2n}k}{(2\pi)^{2n}} \frac{1}{D_{\lambda}^{a}}, \quad \int \frac{\mathrm{d}^{2n}k}{(2\pi)^{2n}} \frac{k_{\mu_{1}}k_{\mu_{2}}}{D_{\lambda}^{a+1}}, \quad \int \frac{\mathrm{d}^{2n}k}{(2\pi)^{2n}} \frac{k_{\mu_{1}}k_{\mu_{2}}\cdots k_{\mu_{2b-1}}k_{\mu_{2b}}}{D_{\lambda}^{a+b}}, \tag{3.6}$$

in which $n \ge a$. A convenient systematization emerges naturally. Since they have the same power counting, it is always possible to combine the integrals above as surface terms noticing that

$$-\frac{\partial}{\partial k^{\mu_1}}\frac{k_{\mu_2}\cdots k_{\mu_{2n}}}{D^a_\lambda} = 2a\frac{k_{\mu_1}k_{\mu_2}\cdots k_{\mu_{2n}}}{D^{a+1}_\lambda} - g_{\mu_1\mu_2}\frac{k^{\mu_3}\cdots k^{\mu_{2n}}}{D^a_\lambda} - \text{permutations.}$$
(3.7)

As can be seen, all the terms of the RHS have equal power counting and the iterative use of this observation allow to recombine all of the integrals above as surface terms.

In short, the highest-rank surface term generates a tower through linear combinations of lowerrank surface terms up to scalar integrals that encode exactly the divergent content of the original expression. They will keep the possibility or not of shifting the integration variable, which means we are trading the freedom of the operation of translation in the momentum-space for the arbitrary choice of the routings in these perturbative corrections.

For linear and higher divergent or logarithmic-divergent tensor integrals, these surface terms are always present, nonetheless, to the former, they have some of its coefficients the ambiguous momenta defined in eq. (2.9) as for logarithmic power counting, the coefficients are the external momenta defined (2.8).

For our purposes, in this work we need to define the following combinations

$$\Delta_{(n+1);\mu_1\mu_2}^{2n}(\lambda^2) = \int \frac{\mathrm{d}^{2n}k}{(2\pi)^{2n}} \left(\frac{2nk_{\mu_1}k_{\mu_2}}{D_{\lambda}^{n+1}} - g_{\mu_1\mu_2}\frac{1}{D_{\lambda}^n}\right) = -\int \frac{\mathrm{d}^{2n}k}{(2\pi)^{2n}}\frac{\partial}{\partial k^{\mu_1}}\frac{k_{\mu_2}}{D_{\lambda}^n},\tag{3.8}$$

in two, four, and six dimensions, n = 1, n = 2, and n = 3 respectively indicated by the superscript.

The irreducible scalar is defined also by dimension and will be

$$I_{\log}^{(2n)}(\lambda^2) = \int \frac{\mathrm{d}^{2n}k}{(2\pi)^{2n}} \frac{1}{D_{\lambda}^n}.$$
(3.9)

As we have seen, the separation highlights diverging structures that organize them without performing any analytic operation. Moreover, it makes clear that the divergent content is a local polynomial in the ambiguous and physical momenta which is obtained without expansions or limits. The finite parts are regularization free and obtained by integrating just finite integrals that will be explained in the section (3.2).

3.2. Finite Functions

After separating the finite part using the identity (3.5), we solve the integrals using the usual techniques of perturbative calculations. It is possible to project the results into a family of functions. For the two-point basic functions they are given by

$$Z_{n_1}^{(-1)} = \int_0^1 \mathrm{d}x \frac{x^{n_1}}{Q}; \quad Z_{n_1}^{(0)} = \int_0^1 \mathrm{d}x x^{n_1} \log \frac{Q}{-m^2}$$
(3.10)

where Q is a polynomial given by

$$Q = p^2 x \left(1 - x\right) - m^2 \tag{3.11}$$

For the three-point basic we have³

$$Q = p^{2}x_{1}(1 - x_{1}) + q^{2}x_{2}(1 - x_{2}) - 2(p \cdot q)x_{1}x_{2} - m^{2}$$
(3.12)

and the basic functions

$$Z_{n_1 n_2}^{(-1)} = \int_0^1 \mathrm{d}x_1 \int_0^{1-x_1} \mathrm{d}x_2 \frac{x_1^{n_1} x_2^{n_2}}{Q}$$
(3.13)

$$Z_{n_1n_2}^{(0)} = \int_0^1 \mathrm{d}x_1 \int_0^{1-x_1} \mathrm{d}x_2 x_1^{n_1} x_2^{n_2} \log \frac{Q}{-m^2}$$
(3.14)

³ Thes polynomials can be written as a quadric form with the Symanzik polynomials constructed using the spanning trees and two-forests of the graph.

And finally, for the four-point is

$$Q = p^{2}x_{1}(1 - x_{1}) + q^{2}x_{2}(1 - x_{2}) + r^{2}x_{3}(1 - x_{3}) -2(p \cdot q)x_{1}x_{2} - 2(p \cdot r)x_{1}x_{3} - 2(q \cdot r)x_{2}x_{3} - m^{2}$$
(3.15)

and the basic functions

$$Z_{n_1n_2n_3}^{(-1)} = \int_0^1 \mathrm{d}x_1 \int_0^{1-x_1} \mathrm{d}x_2 \int_0^{1-x_1-x_2} \mathrm{d}x_3 \frac{x_1^{n_1} x_2^{n_2} x_3^{n_3}}{Q}$$
(3.16)

$$Z_{n_1n_2n_3}^{(0)} = \int_0^1 \mathrm{d}x_1 \int_0^{1-x_1} \mathrm{d}x_2 \int_0^{1-x_1-x_2} \mathrm{d}x_3 x_1^{n_1} x_2^{n_2} x_3^{n_3} \log \frac{Q}{-m^2}$$
(3.17)

It is possible to write the parameters in terms of derivatives of the polynomials and, then, use partial integration to obtain relations among these functions. More precisely, they are reductions of the parameter powers $n_1 + n_2$, for the equation (3.13) and $n_1 + n_2 + n_3$, for the equation (3.16) (see appendices C, and D), they were approched in the papers [37], [38], and [39]. This resource is necessary to perform the operations displayed in this paper.

3.3. Basis of Feynman Integrals

From the general definition presented in eq. (2.16) for the integrals that appear soon after taking Dirac traces, we will describe in a nutshell the ones that arise in the AV^n amplitudes in d = 2n, more generally, any amplitude of n + 1 points of odd parity. They are

$$\left(\bar{J}_n; \bar{J}_n^{\mu_1}\right) = \int \frac{\mathrm{d}^{2n}k}{(2\pi)^{2n}} \frac{(1; K_1^{\mu_1})}{D_{12\cdots n}}$$
(3.18)

and

$$\left(\bar{J}_{n+1}; \bar{J}_{n+1}^{\mu_1}; \bar{J}_{n+1}^{\mu_{12}}\right) = \int \frac{\mathrm{d}^{2n}k}{(2\pi)^{2n}} \frac{(1; K_1^{\mu_1}; K_1^{\mu_{12}})}{D_{12\cdots n+1}},\tag{3.19}$$

the power counting of *n*-point integrals are $\omega(J_n^{\mu_1}) = 1$, $\omega(J_n) = 0$ and of the (n + 1)-point integrals are $\omega(J_{n+1}^{\mu_{12}}) = 0$, $\omega(J_{n+1}^{\mu_1}) = -1$, and $\omega(J_{n+1}) = -2$. Observe our notational conventions $K_{12\cdots i}^{\nu_{12}\cdots\nu_{i}} = K_1^{\nu_1}K_2^{\nu_2}\cdots K_i^{\nu_i}$ and $D_{12\cdots i} = D_1D_2\cdots D_i$.

Therefore, as anticipated by the power counting, some of these integrals contain finite and divergent parts, as is the case with $\bar{J}_{n+1}^{\mu_{12}}$, $\bar{J}_n^{\mu_1}$, and \bar{J}_n . The integrals $J_{n+1}^{\mu_1}$ and J_{n+1} are finite and then will not get an overbar. All the time, we are working with the strictly finite part of the divergent integrals, they will come free of the overbar.

As a quite important example, we chose to compute the highest power-counting integral in d = 4 to illustrate some of the features of our treatment, the vector two-point integral

$$\bar{J}_2^{\mu_1} = \int \frac{\mathrm{d}^4 k}{(2\pi)^4} \frac{K_1^{\mu_1}}{D_{12}} \tag{3.20}$$

using the identity (3.5) with N = 1

$$\frac{1}{D_i} = \frac{1}{D_\lambda} - \frac{A_i}{D_\lambda^2} + \frac{A_i^2}{D_\lambda^2 D_i}$$
(3.21)

corresponding to its linear divergence, we get

$$\frac{K_1^{\mu_1}}{D_{12}} = \frac{K_1^{\mu_1}}{D_{\lambda}^2} - \frac{(A_1 + A_2)}{D_{\lambda}^3} K_1^{\mu_1} + \frac{A_1 A_2}{D_{\lambda}^4} K_1^{\mu_1} \\
+ \left[\frac{A_1^2}{D_{\lambda}^3 D_1} + \frac{A_2^2}{D_{\lambda}^3 D_2} - \frac{A_1 A_2^2}{D_{\lambda}^4 D_2} - \frac{A_2 A_1^2}{D_{\lambda}^4 D_1} + \frac{A_1^2 A_2^2}{D_{\lambda}^4 D_{12}} \right] K_1^{\mu_1},$$
(3.22)

collecting the purely divergent integrals and integrating the remaining finite integrals comes

$$\bar{J}_{2}^{\mu_{1}} = J_{2}^{\mu_{1}}(p_{21}) - \frac{1}{2} \left[P_{21}^{\nu_{1}} \Delta_{3\nu_{1}}^{(4)\mu_{1}} + p_{21}^{\mu_{1}} I_{\log}^{(4)} \right]$$
(3.23)

where

$$J_2^{\mu_1}(p_{21}) = \frac{i}{(4\pi)^2} p_{21}^{\mu_1} Z_1^{(0)}(p_{21})$$
(3.24)

showing all the elements we have presented before, the local divergent terms organized, and the finite part integrated without restrictions.

The same type of steps leads, in d = 6 dimensions, to

$$\bar{J}_{3}^{\mu_{1}} = \int \frac{\mathrm{d}^{6}k}{(2\pi)^{6}} \frac{K_{1}^{\mu_{1}}}{D_{123}}$$
(3.25)

$$\bar{J}_{3}^{\mu_{1}} = -\frac{1}{3} \left(k_{1}^{\nu_{1}} + k_{2}^{\nu_{1}} + k_{3}^{\nu_{1}} \right) \Delta_{4\nu_{1}}^{(6)\mu_{1}} - \frac{1}{3} \left(p_{21}^{\mu_{1}} + p_{31}^{\mu_{1}} \right) I_{\log}^{(6)} + J_{3}^{\mu_{1}}, \tag{3.26}$$

where the finite part is simply

$$J_{3}^{\mu_{1}}(p_{21}, p_{31}) = \frac{i}{(4\pi)^{3}} \left[p_{21}^{\mu_{1}} Z_{10}^{(0)}(p_{21}, p_{31}) + p_{31}^{\mu_{1}} Z_{01}^{(0)}(p_{21}, p_{31}) \right], \qquad (3.27)$$

as in two dimensions, we have

$$\bar{J}_{1\mu_1} = \int \frac{\mathrm{d}^2 k}{(2\pi)^2} \frac{K_{1\mu_1}}{D_1} = -k_1^{\nu_1} \Delta_{2\nu_1\mu_1}^{(2)}, \qquad (3.28)$$

that is a pure surface term, and this is the reason to illustrate for d = 4 dimensions first.

For all explicit results used see the appendices (C) and (D).

4. TWO DIMENSIONAL AV AND VA TWO-POINT FUNCTIONS

To establish the connection among linearity, uniqueness, and WIs we study the two Lorentz indexes amplitudes AV, and VA. Amid this process, the relation among all two-point functions will emerge. As the power counting is zero, it is not expected they depend on the routings through their sums besides they are a function of only two routings, hence we will adopt the simplification $q = p_{21} = k_2 - k_1$, when seen as suit.

Therefore, to start with, we have the RAGFs coming from the vector vertex

$$q^{\mu_2} t^{AV}_{\mu_{12}} = t^A_{\mu_1} \left(1\right) - t^A_{\mu_1} \left(2\right) \tag{4.1}$$

$$q^{\mu_1} t^{VA}_{\mu_{12}} = t^A_{\mu_2} \left(1\right) - t^A_{\mu_2} \left(2\right), \qquad (4.2)$$

and from the axial vertex

$$q^{\mu_1} t^{AV}_{\mu_{12}} = t^A_{\mu_2} (1) - t^A_{\mu_2} (2) - 2m t^{PV}_{\mu_2}$$

$$\tag{4.3}$$

$$q^{\mu_2} t^{VA}_{\mu_{12}} = t^A_{\mu_1} (1) - t^A_{\mu_1} (2) + 2m t^{VP}_{\mu_1}, \qquad (4.4)$$

obtained using the procedure delineated in the section (2). Taking their integrals, we should have

$$q^{\mu_2}T^{AV}_{\mu_{12}} = T^A_{\mu_1}(1) - T^A_{\mu_1}(2)$$
(4.5)

$$q^{\mu_1}T^{AV}_{\mu_{12}} = T^A_{\mu_2}(1) - T^A_{\mu_2}(2) - 2mT^{PV}_{\mu_2}, \qquad (4.6)$$

only by the linearity of integration. Similarly to VA amplitudes.

On the other hand, the WIs will require

$$q^{\mu_2}T^{AV}_{\mu_{12}} = 0; \quad q^{\mu_1}T^{AV}_{\mu_{12}} = -2mT^{PV}_{\mu_2}, \tag{4.7}$$

that in turn, through the general tensor structure of these amplitudes, imply kinematic properties to the scalar invariants of these tensors with the same status as the symmetry properties.

As an example, to the AV amplitude, we can write

$$T_{\mu_{12}}^{AV} = \varepsilon_{\mu_{1}\mu_{2}}F_{1} + \varepsilon_{\mu_{1}\nu}q^{\nu}q_{\mu_{2}}F_{2} + \varepsilon_{\mu_{2}\nu}q^{\nu}q_{\mu_{1}}F_{3}, \qquad (4.8)$$

being the F_i the scalar invariants, then by contracting with the external momenta in the respective indexes, we get

$$q^{\mu_2} T^{AV}_{\mu_{12}} = \varepsilon_{\mu_1 \nu} q^{\nu} \left(q^2 F_2 + F_1 \right)$$
 and (4.9)

$$q^{\mu_1}T^{AV}_{\mu_{12}} = \varepsilon_{\mu_2\nu}q^{\nu} \left(q^2 F_3 - F_1\right), \qquad (4.10)$$

by vector conservation, in the first equation, we trade $F_1 = -q^2 F_2$ in the second equation to obtain

$$q^{\mu_1}T^{AV}_{\mu_{12}} = \varepsilon_{\mu_2\nu}q^{\nu}q^2 \left(F_3 + F_2\right), \qquad (4.11)$$

and finally, under the hypothesis of regularity, we have the low energy theorem for the contraction with the index of the axial current $q^{\mu_1}T^{AV}_{\mu_{12}}|_{q^2=0} = 0$. However, if the WI that relates this contraction to the PV function is satisfied, comes the consequence

$$q^{\mu_1} T^{AV}_{\mu_{12}} \Big|_0 = -2m T^{PV}_{\mu_2} \Big|_0 = \varepsilon_{\mu_2\nu} q^{\nu} \Omega^{PV} \left(q^2 = 0 \right) = 0.$$
(4.12)

Therefore, if the symmetries were respected and the hypothesis of regular form factors is met, such behavior must be attained and in this sense, we have said that it has the status of symmetry.

All these constraints must be seen in the light of explicit computations that will be unfolded and analyzed in the sequel. Following the definitions of the previous section, after integration the amplitude becomes

$$T^{\Gamma_{1}\Gamma_{2}}(1,2) = \int \frac{\mathrm{d}^{2}k}{(2\pi)^{2}} t^{\Gamma_{1}\Gamma_{2}}(1,2)$$

expanding the terms of mass and momentum,

$$t^{\Gamma_{1}\Gamma_{2}} = +K_{12}^{\nu_{12}} \operatorname{tr} \left[\Gamma_{1}\gamma_{\nu_{1}}\Gamma_{2}\gamma_{\nu_{2}}\right] \frac{1}{D_{12}}$$

$$+mK_{1}^{\nu_{1}} \operatorname{tr} \left[\Gamma_{1}\gamma_{\nu_{1}}\Gamma_{2}\right] \frac{1}{D_{12}}$$

$$+mK_{2}^{\nu_{1}} \operatorname{tr} \left[\Gamma_{1}\Gamma_{2}\gamma_{\nu_{1}}\right] \frac{1}{D_{12}}$$

$$+m^{2} \operatorname{tr} \left[\Gamma_{1}\Gamma_{2}\right] \frac{1}{D_{12}} ,$$

$$(4.13)$$

from the formula above, choosing appropriately the vertexes and keeping the non-zero traces, we have

$$\begin{split} t^{AV}_{\mu_{12}} &= K^{\nu_{12}}_{12} \mathrm{tr}(\gamma_* \gamma_{\mu_1 \nu_1 \mu_2 \nu_2}) \frac{1}{D_{12}} + m^2 \mathrm{tr}(\gamma_* \gamma_{\mu_1 \mu_2}) \frac{1}{D_{12}}, \\ t^{VA}_{\mu_{12}} &= K^{\nu_{12}}_{12} \mathrm{tr}(\gamma_* \gamma_{\mu_1 \nu_1 \mu_2 \nu_2}) \frac{1}{D_{12}} - m^2 \mathrm{tr}(\gamma_* \gamma_{\mu_1 \mu_2}) \frac{1}{D_{12}}. \end{split}$$

The main point is that the trace of four gamma matrices, which is a linear combination of the metric and epsilon tensor, can be expressed in a variety of forms, obtained using the substitutions in the identity (2.5), that means

$$\gamma_* = \varepsilon_{\nu_{12}} \gamma^{\nu_{12}}/2; \quad \gamma_* \gamma_{\mu_1} = -\varepsilon_{\mu_1 \nu_1} \gamma^{\nu_1}; \quad \gamma_* \gamma_{[\mu_1 \mu_2]} = -\varepsilon_{\mu_1 \mu_2}$$

They lead to expressions that are not automatically equal after integration. Effectively they turn the RAGFs into equations among functions. To unfold the rationale, let us apply the definition of the chiral matrix in the form

$$\operatorname{tr} \left(\gamma_* \gamma_{abcd} \right) = \varepsilon_{\alpha_{12}} \operatorname{tr} \left(\gamma^{\alpha_{12}} \gamma_{abcd} \right) / 2 = 2 \left[-g_{ab} \varepsilon_{cd} + g_{ac} \varepsilon_{bd} - g_{ad} \varepsilon_{bc} - g_{bc} \varepsilon_{ad} + g_{bd} \varepsilon_{ac} - g_{cd} \varepsilon_{ab} \right],$$

where the Latin indexes make it simple to perform substitutions to obtain the trace when the definition of the chiral matrix is deployed adjacent to the first or the second vertex, by example $(a, b, c, d) = (\mu_1, \nu_1, \mu_2, \nu_2)$ and $(a, b, c, d) = (\mu_2, \nu_2, \mu_1, \nu_1)$. These traces will differ by the signs of the terms only and have all the indexes of the trace explicitly present although distinctly displayed, these seemingly innocuous observations have far-reaching implications. With these aspects in mind, let us call these two expressions version one and two of the traces and carry over to the amplitudes the same nomenclature, as we will demonstrate they are enough to reach any other expression.

First Version:

$$\begin{split} K_{12}^{\nu_{12}} \mathrm{tr} \left(\gamma_* \gamma_{\mu_1} \gamma_{\nu_1} \gamma_{\mu_2} \gamma_{\nu_2} \right) \; = \; -2\varepsilon_{\mu_1\nu_1} \left(K_{1\mu_2} K_2^{\nu_1} + K_{2\mu_2} K_1^{\nu_1} \right) - 2\varepsilon_{\mu_2\nu_1} \left(K_{1\mu_1} K_2^{\nu_1} - K_{2\mu_1} K_1^{\nu_1} \right) \\ & + 2\varepsilon_{\mu_1\mu_2} \left(K_1 \cdot K_2 \right) + 2g_{\mu_1\mu_2} \varepsilon_{\nu_1\nu_2} K_{12}^{\nu_{12}} \end{split}$$

Second Version:

$$K_{12}^{\nu_{12}} \operatorname{tr}(\gamma_* \gamma_{\mu_2} \gamma_{\nu_2} \gamma_{\mu_1} \gamma_{\nu_1}) = +2\varepsilon_{\mu_1\nu_1} \left(K_{1\mu_2} K_2^{\nu_1} - K_{2\mu_2} K_1^{\nu_1} \right) - 2\varepsilon_{\mu_2\nu_1} \left(K_{1\mu_1} K_2^{\nu_1} + K_{2\mu_1} K_1^{\nu_1} \right) \\ -2\varepsilon_{\mu_1\mu_2} \left(K_1 \cdot K_2 \right) - 2g_{\mu_1\mu_2} \varepsilon_{\nu_1\nu_2} K_{12}^{\nu_12}.$$

Here, we have already contracted with $K_{12}^{\nu_{12}}$. Now we note that in the first row of each version, it is possible to identify, as will be done in four and six dimensions, a common tensor

$$t_{\mu_2}^{(s)\nu_1} = \left(K_{1\mu_2}K_2^{\nu_1} + sK_{2\mu_2}K_1^{\nu_1}\right)\frac{1}{D_{12}},\tag{4.14}$$

where $s = \pm 1$. And in the other rows appear the amplitude SP obtained when substituting the respective vertexes in (4.13), $t^{SP} = 2\varepsilon_{\nu_{12}}K_{12}^{\nu_{12}}/D_{12} = -t^{PS}$. Completing the two amplitudes summing the mass terms using $\operatorname{tr}(\gamma_*\gamma_{\mu_{12}}) = -2\varepsilon_{\mu_{12}}$, we get the

Completing the two amplitudes summing the mass terms using $\operatorname{tr}(\gamma_*\gamma_{\mu_{12}}) = -2\varepsilon_{\mu_{12}}$, we get the first and the second versions of the AV

similar expressions can be obtained to the VA amplitude.

In the above relations, we have identified the scalar two-point amplitudes

$$t^{PP} = q^2 \frac{1}{D_{12}} - \frac{1}{D_1} - \frac{1}{D_2}$$
(4.15)

$$t^{SS} = (4m^2 - q^2)\frac{1}{D_{12}} + \frac{1}{D_1} + \frac{1}{D_2}, \qquad (4.16)$$

to obtain these amplitudes we have used

$$S_{ij} = K_i \cdot K_j - m^2 = \frac{1}{2} (D_i + D_j - p_{ij}^2), \qquad (4.17)$$

to reduce the bilinears that appear in their definitions in the eq.(4.13). So it is straightforward to identify in the middle of our expressions,

$$t_{\mu_1\mu_2}^{VV} = 2t_{\mu_1\mu_2}^{(+)} + g_{\mu_1\mu_2}t^{PP}; \text{ and } t_{\mu_1\mu_2}^{AA} = 2t_{\mu_1\mu_2}^{(+)} - g_{\mu_1\mu_2}t^{SS},$$

they follow immediately after the traces are taken in the respective definitions and clearly will relate the odd amplitudes to the even ones.

When integrated, it is always possible to see that some terms are finite and vanishing. Namely $\varepsilon_{\mu_i\nu}T_{\mu_j}^{(-)\nu} = 0$ and $T^{SP} = 0$, due of the properties of the finite vector integral that is proportional to momentum $q = p_{21}$ and the scalar integral J_2 see in (B1). Therefore, the integrals will provide the basic relations

However, if one applies the second relation $\gamma_*\gamma_{\mu_1} = -\varepsilon_{\mu_1\nu_1}\gamma^{\nu_1}$ around the same vertexes, we get directly

$$(t^{AV}_{\mu_{12}})_1 \,\tilde{=} \, - \, \varepsilon_{\mu_1}^{\ \nu_1} t^{VV}_{\nu_1 \mu_2}; \qquad (t^{AV}_{\mu_{12}})_2 \,\tilde{=} \, - \, \varepsilon_{\mu_2}^{\ \nu_1} t^{AA}_{\mu_1 \nu_1}.$$

The sign \approx means they are equal up to terms that are finite and vanish under integration. The other independent version is obtained using the third relation $\gamma_*\gamma_{[\mu_1\mu_2]} = -\varepsilon_{\mu_1\mu_2}$, in the form $\gamma_*\gamma_{\mu_1}\gamma_{\nu_1} = -\varepsilon_{\mu_1\nu_1} + g_{\mu_1\nu_1}\gamma_*$, given in the trace the expression $-2(\varepsilon_{\mu_1\nu_1}g_{\mu_2\nu_2} + \varepsilon_{\mu_2\nu_2}g_{\mu_1\nu_1})$, that in the amplitudes enable us to arrange the result

thereby their integrals provide us with the following

$$(T_{\mu_{12}}^{AV})_3 = \frac{1}{2} \left[(T_{\mu_{12}}^{AV})_1 + (T_{\mu_{12}}^{AV})_2 \right]; \quad (T_{\mu_{12}}^{VA})_3 = \frac{1}{2} \left[(T_{\mu_{12}}^{VA})_1 + (T_{\mu_{12}}^{VA})_2 \right].$$
 (4.18)

This form is present in the equation (85) of the paper [48], for example. This last form, obtained a linear combination of the other two is a particular aspect that is made clear in the section (5), here is present because it comes from the identities we introduced and we wish to make a pedestrian approach in two dimensions.

To put the consequences of versions for amplitudes into perspective, we need integrated expressions. It will be possible to see that the sampling of the indices that appear between the finite and divergent parts makes the expressions not automatically equal when integrated. However, the versions are related through linearity violations reflected in the RAGFs. Using the explicit results found in the appendix (B), we will have, from the expressions (4.15) and (4.16),

$$T^{PP} = q^2 J_2 - 2I_{\log} \tag{4.19}$$

$$T^{SS} = (4m^2 - p^2) J_2 + 2I_{\log}, \qquad (4.20)$$

and for the sign tensor (4.14)

$$T_{\mu_1\mu_2}^{(+)} = 2\left(\bar{J}_{2\mu_1\mu_2} + q_{\mu_1}J_{2\mu_2}\right) = 2\theta_{\mu_1\mu_2}\left(q\right)\left(i/4\pi + m^2J_2\right)$$
(4.21)

$$-\frac{1}{2}g^{\mu_1\mu_2}q^2J_2 + \left(\Delta_{2\mu_1\mu_2} + g_{\mu_1\mu_2}I_{\log}\right), \qquad (4.22)$$

to the two-rank, two-point, even amplitudes

$$T_{\mu_1\mu_2}^{VV} = 2\Delta_{2\mu_1\mu_2} + 4\theta_{\mu_1\mu_2} \left(m^2 J_2 + i/4\pi\right), \qquad (4.23)$$

$$T_{\mu_1\mu_2}^{AA} = 2\Delta_{2\mu_1\mu_2} + 4\theta_{\mu_1\mu_2} \left(m^2 J_2 + i/4\pi\right) - g_{\mu_1\mu_2} \left(4m^2 J_2\right), \qquad (4.24)$$

where $\theta_{\alpha\lambda}(q) = (g_{\alpha\lambda}q^2 - q_{\alpha}q_{\lambda})/q^2$ is the transversal projector. And in this way, the odd amplitude AV will turn up with the explicitly expressions

$$\left(T_{\mu_{12}}^{AV}\right)_{1} = -2\varepsilon_{\mu_{1}}{}^{\nu}\Delta_{2\mu_{2}\nu} - 4\varepsilon_{\mu_{1}\nu}\theta_{\mu_{2}}^{\nu}\left(m^{2}J_{2} + i/4\pi\right)$$
(4.25)

$$\left(T^{AV}_{\mu_{12}}\right)_{2} = -2\varepsilon_{\mu_{2}}{}^{\nu}\Delta_{2\mu_{1}\nu} - 4\varepsilon_{\mu_{2}\nu}\theta^{\nu}_{\mu_{1}}\left(m^{2}J_{2} + i/4\pi\right) - \varepsilon_{\mu_{1}\mu_{2}}\left(4m^{2}J_{2}\right).$$
(4.26)

The expression for (4.18) is obtained as a combination of the expressions above.

The two-point functions, that are finite, and appear in the RAGFs

$$T_{\mu}^{PV} = -T_{\mu}^{VP} = \varepsilon_{\mu\nu}q^{\nu} \left[-2mJ_{2}\left(q\right)\right].$$

$$T_{\mu}^{PA} = -T_{\mu}^{AP} = q_{\mu} \left[+2mJ_{2}\left(q\right)\right].$$
(4.27)

And the one-point ones arising in that relations as well

$$T^V_{\mu}(i) = -2k^{\nu}_i \Delta_{2\mu\nu}$$

$$T^A_{\mu}(i) = -2\varepsilon_{\mu\nu_1}k^{\nu_2}_i \Delta^{\nu_1}_{2\nu_2}.$$

Now it is possible to state that same if all integrands are the same, the same is not true for the integrals. In the case of even amplitudes VV and AA, the expressions depend in a unique way on their divergences once a method is chosen. In the case of odd amplitudes, the expressions depend on the version of the trace used. We will show that this dependence manifests itself in a predictable way in all dimensions in the corresponding amplitudes.

Where is the reason for such discrepancies? Even if it has been applied to just identities, deploying the definition of the chiral matrix around the first or the second vertexes has sampled that the indexes among the finite and divergent parts are not automatically equal after integration. This statement becomes clear when we subtract one expression from the other

$$(T^{AV}_{\mu_{12}})_1 - (T^{AV}_{\mu_{12}})_2 = -2 \left[\varepsilon_{\mu_1\nu} \Delta^{\nu}_{2\mu_2} - \varepsilon_{\mu_2\nu} \Delta^{\nu}_{2\mu_1} \right] - 4 \left[\varepsilon_{\mu_1\nu_1} \theta^{\nu}_{\mu_2} - \varepsilon_{\mu_2\nu_1} \theta^{\nu}_{\mu_1} \right] (m^2 J_2 + i/4\pi)$$

$$+ 4\varepsilon_{\mu_1\mu_2} m^2 J_2.$$

For the rearranging of the indexes in the finite part and in the object $\Delta_{2\mu\nu}$ we use the Schouten identity⁴ in two dimensions $\varepsilon_{[\mu_1\nu}\Delta^{\nu}_{2\mu_2]} = 0$ and $\varepsilon_{[\mu_1\nu}\theta^{\mu}_{\mu_2]} = 0$, the difference between the versions is reduced to

$$\left(T^{AV}_{\mu_{12}}\right)_1 - \left(T^{AV}_{\mu_{12}}\right)_2 = -\varepsilon_{\mu_1\mu_2} \left(2\Delta^{\alpha}_{2\alpha} + i/\pi\right).$$
(4.30)

The linearity can be translated as, if over the true equation $(t_{\mu_{12}}^{AV})_1 - (t_{\mu_{12}}^{AV})_2 = 0$ we apply an operation to replace the integral, that is defined to be a linear operation, then one should have the equation above identically vanishing, meantime, for this to be satisfied we would have a condition about the value of the object $\Delta_{2\alpha}^{\alpha}$, determined by the unique relation (4.30). Before delving into this issue, let us analyze how this condition is manifested in the RAGFs.

4.1. Verification and Consequences of the RAGFs

The RAGFs are given by the procedure illustrated in the section (2). The even ones readily comply with their relations determined to their integrands, which means

$$q^{\mu_1}T^{VV}_{\mu_{12}} = T^V_{\mu_2}(1) - T^V_{\mu_2}(2) = 2q^{\nu_1}\Delta_{2\mu_2\nu_1}$$
$$q^{\mu_1}T^{AA}_{\mu_{12}} + 2mT^{PA}_{\mu_2} = T^V_{\mu_2}(1) - T^V_{\mu_2}(2) = 2q^{\nu_1}\Delta_{2\mu_2\nu_1},$$

where we have used the following result $T^V_{\mu}(i) = -2k^{\nu}_i \Delta_{2\mu\nu}$ and the presence of the projector to eliminate some finite parts by $q^{\mu_2} \theta^{\nu}_{\mu_2} = 0$. Exactly the same results with the RAGF to the index μ_2 .

On the other hand, the odd amplitudes have a substantially more subtle behavior. Contracting the first version of AV, eq.(4.25), in the vector vertex we get

$$q^{\mu_2} \left(T^{AV}_{\mu_{12}} \right)_1 = -2\varepsilon_{\mu_1\nu_1} q^{\nu_2} \Delta^{\nu_1}_{2\nu_2} = \left[T^A_{\mu_1} \left(1 \right) - T^A_{\mu_1} \left(2 \right) \right],$$

where $T^{A}_{\mu}(i) = -2\varepsilon_{\mu\nu_{1}}k_{i}^{\nu_{2}}\Delta_{2\nu_{2}}^{\nu_{1}}$. Note this happen without any restriction.

Quite a different situation occurs when contracting with the axial vertex, there we have

$$q^{\mu_1} \left(T^{AV}_{\mu_{12}} \right)_1 = -2q^{\mu_1} \varepsilon_{\mu_1 \nu} \Delta^{\nu}_{2\mu_2} - 4q^{\mu_1} \varepsilon_{\mu_1 \nu} \theta^{\nu}_{\mu_2} \left(m^2 J_2 + i/4\pi \right) + \frac{1}{2} \left(m^2 J_2 + i/4\pi \right) +$$

Using the Schouten identity as (4.28) and (4.29), and projector properties, $\theta_{\nu}^{\nu} = 1$ and transversality, it is then obtained

$$q^{\mu_1} \left(T^{AV}_{\mu_{12}} \right)_1 = \left[T^A_{\mu_2} \left(1 \right) - T^A_{\mu_2} \left(2 \right) \right] - 2m T^{PV}_{\mu_2} + \varepsilon_{\mu_2 \nu_1} q^{\nu_1} \left[2\Delta^{\nu_2}_{2\nu_2} + \left(i/\pi \right) \right], \tag{4.31}$$

where the PV amplitude is given by (4.27). Note the last term in (4.31) spoils the automatic satisfaction of this RAGF.

$$A_{[\alpha_1 \cdots \alpha_r} B_{\alpha_{r+1} \cdots \alpha_s]} = \frac{1}{s!} \sum_{\pi \in S_s} \operatorname{sign}(\pi) A_{\alpha_{\pi(1)} \cdots \alpha_{\pi(r)}} B_{\alpha_{\pi(r+1)} \cdots \alpha_{\pi(s)}}$$

the normalizing factor is irrelevant to all the identities used. Through the antisymmetry of the Levi-Civita tensor, follows

$$\varepsilon_{\mu_1\nu}\Delta^{\nu}_{2\mu_2} + \varepsilon_{\mu_2\mu_1}\Delta^{\nu}_{2\nu} + \varepsilon_{\nu\mu_2}\Delta^{\nu}_{2\mu_1} = 0$$

$$(4.28)$$

$$\varepsilon_{\mu_1\nu}\theta^{\nu}_{\mu_2} + \varepsilon_{\mu_2\mu_1}\theta^{\nu}_{\nu} + \varepsilon_{\nu\mu_2}\theta^{\nu}_{\mu_1} = 0.$$
(4.29)

⁴ The notation to antisymmetrization of indexes adopted by us is

Following the same reasoning for the expression (4.26), we see the opposite behavior,

$$q^{\mu_2} \left(T^{AV}_{\mu_{12}} \right)_2 = \left[T^A_{\mu_1} \left(1 \right) - T^A_{\mu_1} \left(2 \right) \right] + \varepsilon_{\mu_1 \nu} q^{\nu} \left[2\Delta^{\alpha}_{2\alpha} + (i/\pi) \right]$$
(4.32)

$$q^{\mu_1} \left(T^{AV}_{\mu_{12}} \right)_2 = \left[T^A_{\mu_2} \left(1 \right) - T^A_{\mu_2} \left(2 \right) \right] - 2m T^{PV}_{\mu_2}, \tag{4.33}$$

namely, the ARAGF coming from the contraction in the axial vertex is satisfied without restrictions, however, the VRAGF arises conditioned by the value of the object $\Delta_{2\mu\nu}$.

From the relations ARAGF in (4.31) and VRAGF in (4.32), it is straightforward to see that for the expression (4.18), of the third version, both vertexes have potential violating terms, because it is given by a combination of the other two. For that version, none of the RAGFs is automatically satisfied.

The same happens to VA amplitude. The vertex having the offending term corresponds to the version in question. For the first version, the relation (4.2) is satisfied, while for the second version, the relation (4.4) is satisfied, and the possibly violating terms occur in

$$q^{\mu_1} \left(T^{VA}_{\mu_{12}} \right)_1 = \varepsilon_{\mu_2\nu_1} q^{\nu_1} \left[2\Delta^{\alpha}_{2\alpha} + (i/\pi) \right] + T^A_{\mu_2} \left(1 \right) - T^A_{\mu_2} \left(2 \right)$$
$$q^{\mu_2} \left(T^{VA}_{\mu_{12}} \right)_2 = \varepsilon_{\mu_1\nu_1} q^{\nu_1} \left[2\Delta^{\alpha}_{2\alpha} + (i/\pi) \right] + T^A_{\mu_1} \left(1 \right) - T^A_{\mu_1} \left(2 \right) + 2m T^{VP}_{\mu_1}.$$

From the RAGFs verified above, we can conclude that even amplitudes have their relations satisfied for any values of the surface term and therefore do not violate linearity. On the other hand, odd amplitudes require the condition

$$\Delta^{\alpha}_{2\alpha} = -\frac{i}{2\pi}.\tag{4.34}$$

The location of this term is in the vertex where the version is defined. Choosing a finite value for the surface term also requires that the expression (4.30) is null, ensuring that the two versions have the same content. However, this requirement implies that $[T^V_{\mu}(1) - T^V_{\mu}(2)] = -(i/2\pi) q_{\mu}$ and for the axial function is the same because the amplitudes are related through $T^A_{\mu}(k_i) = -\varepsilon_{\mu} \nu [T^V_{\nu}(k_i)]$. The implications are obvious but we need to analyze the consequences for the IWs.

4.2. Ward Identities

When a WI is unavoidably violated, we have an anomaly. With this simple assertion in mind, we will establish from now on all possible scenarios for the calculations outlined above and how the satisfaction or not of the RAGF affects the WIs. The VWI, in turn, asks for the identical vanishing of the one-point functions, for example, for the vector current conservation

$$q^{\mu_1}T^{VV}_{\mu_{12}} = \left[T^V_{\mu_2}\left(1\right) - T^V_{\mu_2}\left(2\right)\right] = 2q^{\nu_1}\Delta_{2\mu_2\nu_1} = 0$$

As for the axial current, we have the partial conservation of the axial current

$$q^{\mu_1}T^{AV}_{\mu_{12}} = \left[T^A_{\mu_2}\left(1\right) - T^A_{\mu_2}\left(2\right)\right] - 2mT^{PV}_{\mu_2} = -2mT^{PV}_{\mu_2}.$$

If this result was straightforward, it would be enough to enforce the vanishing of $\Delta_{2\mu\nu} = 0$. For the logarithmic divergences, it is possible to see which coefficients are not ambiguous (as will see for linear divergences as happens in 4D-AVV) and depends only on the external momentum (2.8). Nonetheless in this case linearity is violated in one or the other version and the violating term ends up offending the WIs. The condition is reflected in the WIs of the double vector function, as well, see table (I) for global results.

	-	
	$PV_{\mu_2} + (i/\pi) \varepsilon_{\mu_2 \nu_1} q^{\nu_1}$	$q^{\mu_2} \left(T^{AV}_{\mu_{12}} \right)_1 = 0$
$q^{\mu_1} \left(T^{AV}_{\mu_{12}}\right)_2 = -2mT_{\mu_{12}}$	$\frac{PV}{\mu_2}$	$q^{\mu_2} \left(T^{AV}_{\mu_{12}} \right)_2 = (i/\pi) \varepsilon_{\mu_1 \nu_1} q^{\nu_1}$
$q^{\mu_1} \left(T^{AV}_{\mu_{12}} \right)_3 = -2mT_{\mu_{12}}$	$PV_{\mu_2} + (i/2\pi) \varepsilon_{\mu_2\nu_1} q^{\nu_1}$	$q^{\mu_2} \left(T^{AV}_{\mu_{12}} \right)_3 = (i/2\pi) \varepsilon_{\mu_1 \nu_1} q^{\nu_1}$
$q^{\mu_1} T^{VV}_{\mu_{12}} = 0$		$q^{\mu_2}T^{VV}_{\mu_{12}} = 0$
$q^{\mu_1}T^{AA}_{\mu_{12}} = -2mT^{PA}_{\mu_2}$		$q^{\mu_2}T^{AA}_{\mu_{12}} = 2mT^{AP}_{\mu_2}$

TABLE I: Violations for vanishing surface term in each version.

TABLE II: Violations for unique amplitudes

$q^{\mu_1}T^{AV}_{\mu_{12}} = -2mT^{PV}_{\mu_2} + (i/2\pi)\varepsilon_{\mu_2\nu}q$	$^{\nu} \left q^{\mu_2} T^{AV}_{\mu_{12}} = (i/2\pi) \varepsilon_{\mu_1 \nu} q^{\nu} \right $
$q^{\mu_1} T^{VV}_{\mu_{12}} = -\left(i/2\pi\right) q_{\mu_2}$	$q^{\mu_2} T^{VV}_{\mu_{12}} = -\left(i/2\pi\right) q_{\mu_2}$
$q^{\mu_1}T^{AA}_{\mu_{12}} = -2mT^{PA}_{\mu_2} - (i/2\pi) q_{\mu_2}$	$q^{\mu_2}T^{AA}_{\mu_{12}} = 2mT^{AP}_{\mu_2} - (i/2\pi) q_{\mu_2}$

But if linearity is preserved and we assume the result (4.34) as valid, the amplitudes are unique but violate all WIs, see the table (II), notice we do not need to index the version. Nonetheless, in this scenario where all possible manipulations led to only one answer, one consequence over the VV, as we saw through $\left[T^V_{\mu}(1) - T^V_{\mu}(2)\right] = -(i/2\pi) q_{\mu}$, is that its WIs are violated.

The scenario is typical of anomalies where it is inevitable some sort of violation in the WIs. The singularities of perturbation theory are the main reason presented for such a state of affairs. However, we will also establish a finite reason for such observations, through a low-energy property of a finite function. At the beginning of this section we establish that if the two WIs to AV are satisfied by hypothesis, therefore we should have

$$q^{\mu_1} T^{AV}_{\mu_{12}} \big|_{q^2 = 0} = \varepsilon_{\mu_2 \nu} q^{\nu} \Omega^{PV} \left(0 \right) = 0,$$

but if we look at the PV amplitude, eq. (4.27), it is immediate to note that it is finite with the form factor given by $\Omega^{PV} = (i/\pi) m^2 Z_0^{(-1)}(q)^5$, and in the point $q^2 = 0$, happens that $Q(q^2 = 0) = -m^2$, thereby we have

$$\Omega^{PV}(q^2 = 0) = -i/\pi, \tag{4.35}$$

that it is the opposite of the result deduced in the equation above, to the satisfaction of both WIs, hence there is no possibility of satisfaction of both, even if all the elements involved were finite, as long as they are connected to the finite PV amplitude. The fact that remains is that if the VWI is observed, the violation of the AWI is the amount corresponding to the negative of the $\Omega^{PV}(0)$, only due to reasons of tensor structure.

An analogous conclusion follows if in the contractions of the general tensor representing the AV structure, eq.(4.8), we adopt the axial WI as the hypothesis, that means, in the eq.(4.10) we make $F_1 = q^2 F_3 - \Omega^{PV}$, and by substitution in the eq. (4.9), for the contraction in the vector vertex, one would get

$$q^{\mu_2} T^{AV}_{\mu_{12}} = \varepsilon_{\mu_1 \nu} q^{\nu} \left[q^2 \left(F_2 + F_3 \right) - \Omega^{PV} \right], \qquad (4.36)$$

and by the regularity of the form factors, we reach the conclusion

$$q^{\mu_2} T^{AV}_{\mu_{12}} \big|_{q^2 = 0} = -\varepsilon_{\mu_1 \nu} q^{\nu} \Omega^{PV} \left(0 \right) = \left(\frac{i}{\pi} \right) \varepsilon_{\mu_1 \nu} p^{\nu}, \tag{4.37}$$

$$m^{2} Z_{0}^{(-1)} \left(q^{2}\right) = m^{2} \int_{0}^{1} \mathrm{d}x \frac{1}{\left[q^{2} x \left(1-x\right)-m^{2}\right]} \bigg|_{q^{2}=0} = -1$$

5

hence it is clear that the kinematical behavior of the PV finite amplitude is responsible for the series of violations observed in the odd correlators. In four dimensions, we will show that, even if it is obtained violations in all WIs, a specific combination is always dictated by the kinematical properties of finite functions, and such result applies here, as well.

The conclusions drawn are immune to the divergent character of the AV or VA amplitude, and its eventual role in answering the question of inevitable symmetry breaking is, in the limit we are treating, obfuscated by the finite structure mentioned, notwithstanding it is possible to show, in general, that if linearity is preserved as condition, we have a connection $\Omega^{PV}(0) = 2\Delta_{2\alpha}^{\alpha}$, see the analogous result fully demonstrated in 4D eq. (5.76).

From now on, we will enter into a more elaborated scenario, and we will show that in the physical dimension the same conclusions can be drawn for the case of odd triangles. The presence of the anomaly can be anticipated through conclusions similar to those drawn for the two-dimensional case.

5. FOUR DIMENSIONS THREE POINT FUNCTIONS

In this dimension, the amplitudes that exhibit the claimed behavior are the rank three triangles, AVV its permutations VAV, VVA, and AAA. The way they displace the violating terms, their uniqueness properties, and violation of RAGF equations are only consequences of the ambiguity of the integrated expression provoked by the traces of six gamma matrices and an odd number of chiral matrices.

Their computations then, boil down to twenty-four triangles of rank one, twelve parity-even triangles, VPP, ASP, and all their permutations, and twelve parity-odd tensors ASS, APP, VPS, and all its permutations, in addition to three tensors as in two dimensions that can't be written down as other amplitudes and depend only on the leading odd trace of six gamma matrices.

Given the multitude and ambiguities in the path to express all the results, we will get excessively pedantic and detailed in some steps.

A general three-point function is obtained through the appropriate choice of vertexes factors Γ_i in

$$t^{\Gamma_{1}\Gamma_{2}\Gamma_{3}} = \operatorname{tr}\left[\Gamma_{1}S\left(1\right)\Gamma_{2}S\left(2\right)\Gamma_{3}S\left(3\right)\right],\tag{5.1}$$

and as usual in this text, the integrated form gets a capital letter

$$T^{\Gamma_1\Gamma_2\Gamma_3} = \int \frac{\mathrm{d}^4k}{(2\pi)^4} t^{\Gamma_1\Gamma_2\Gamma_3},$$

in the expansion of the terms of mass and momenta, it is possible to see that in the rank zero and rank two triangles the non-zero terms are the odd powers of the mass, and to the rank one and rank three the even power of mass have non-zero traces.

The backbone of amplitudes we are interested in are the odd tensors

$$\begin{aligned} t_{\mu_{123}}^{AVV} &= \mathrm{tr} \left[\gamma_* \gamma_{\mu_1} S \left(1 \right) \gamma_{\mu_2} S \left(2 \right) \gamma_{\mu_3} S \left(3 \right) \right] \\ t_{\mu_{123}}^{VAV} &= \mathrm{tr} \left[\gamma_{\mu_1} S \left(1 \right) \gamma_* \gamma_{\mu_2} S \left(2 \right) \gamma_{\mu_3} S \left(3 \right) \right] \\ t_{\mu_{123}}^{VVA} &= \mathrm{tr} \left[\gamma_{\mu_1} S \left(1 \right) \gamma_{\mu_2} S \left(2 \right) \gamma_* \gamma_{\mu_3} S \left(3 \right) \right] \\ t_{\mu_{123}}^{AAA} &= \mathrm{tr} \left[\gamma_* \gamma_{\mu_1} S \left(1 \right) \gamma_* \gamma_{\mu_2} S \left(2 \right) \gamma_* \gamma_{\mu_3} S \left(3 \right) \right] \end{aligned}$$

$$t_{\mu_{123}}^{AVV} = K_{123}^{\nu_{123}} \operatorname{tr}(\gamma_{*\mu_{1}\nu_{1}\mu_{2}\nu_{2}\mu_{3}\nu_{3}}) \frac{1}{D_{123}} + m^{2} \varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{1}} \left(K_{1}^{\nu_{1}} - K_{2}^{\nu_{1}} + K_{3}^{\nu_{1}}\right) \frac{1}{D_{123}}$$
(5.2)

$$t_{\mu_{123}}^{VAV} = K_{123}^{\nu_{123}} \operatorname{tr}(\gamma_{*\mu_{1}\nu_{1}\mu_{2}\nu_{2}\mu_{3}\nu_{3}}) \frac{1}{D_{123}} + m^{2} \varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{1}} \left(K_{1}^{\nu_{1}} + K_{2}^{\nu_{1}} - K_{3}^{\nu_{1}}\right) \frac{1}{D_{123}}$$
(5.3)

$$t_{\mu_{123}}^{VVA} = K_{123}^{\nu_{123}} \operatorname{tr}(\gamma_{*\mu_{1}\nu_{1}\mu_{2}\nu_{2}\mu_{3}\nu_{3}}) \frac{1}{D_{123}} - m^{2} \varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{1}} \left(K_{1}^{\nu_{1}} - K_{2}^{\nu_{1}} - K_{3}^{\nu_{1}}\right) \frac{1}{D_{123}}$$
(5.4)

$$t_{\mu_{123}}^{AAA} = K_{123}^{\nu_{123}} \operatorname{tr}(\gamma_{*\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3}) \frac{1}{D_{123}} - m^2 \varepsilon_{\mu_1\mu_2\mu_3\nu_1} \left(K_1^{\nu_1} + K_2^{\nu_1} + K_3^{\nu_1}\right) \frac{1}{D_{123}}.$$
 (5.5)

Apart from the other traces, the leading trace in all these pre-diagrams containing six matrices can be computed in lots of ways. We will show after how the protocol we are presenting is enough to achieve any possible result of any identity in the appendix (A).

Starting with the definition

$$\gamma_* = \frac{i}{4!} \varepsilon_{\nu_{1234}} \gamma^{\nu_{1234}} \tag{5.6}$$

the formula below gives all results to the trace of six matrices using the definition anywhere in the string

$$(4i)^{-1} \operatorname{tr} (\gamma_* \gamma_{abcdef}) = +g_{ab} \varepsilon_{cdef} + g_{ad} \varepsilon_{bcef} + g_{af} \varepsilon_{bcde}$$

$$+g_{bc} \varepsilon_{adef} + g_{cd} \varepsilon_{abef} + g_{cf} \varepsilon_{abde}$$

$$+g_{be} \varepsilon_{acdf} + g_{de} \varepsilon_{abcf} + g_{ef} \varepsilon_{abcd}$$

$$-g_{bd} \varepsilon_{acef} - g_{df} \varepsilon_{abce} - g_{bf} \varepsilon_{acde}$$

$$-g_{ac} \varepsilon_{bdef} - g_{ce} \varepsilon_{abdf} - g_{ae} \varepsilon_{bcdf}.$$

$$(5.7)$$

Although valid the identities

$$\operatorname{tr}(\gamma_*\gamma_{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3}) = \operatorname{tr}(\gamma_*\gamma_{\mu_2\nu_2\mu_3\nu_3\mu_1\nu_1}) = \operatorname{tr}(\gamma_*\gamma_{\mu_3\nu_3\mu_1\nu_1\mu_2\nu_2})$$
(5.8)

by using the eq. (5.7) above and adopting $(a, b, c, d, e, f) = (\mu_1, \nu_1, \mu_2, \nu_2, \mu_3, \nu_3)$, $(a, b, c, d, e, f) = (\mu_2, \nu_2, \mu_3, \nu_3, \mu_1, \nu_1)$ and $(a, b, c, d, e, f) = (\mu_3, \nu_3, \mu_1, \nu_1, \mu_2, \nu_2)$ the resulting expressions differ only in a couple of signs leading, when integrated, to three not automatically equivalent expressions to the odd rank-three amplitudes.

To these results for the traces and after contracting with factor $K_{123}^{\nu_{123}}$ we display in the equations below, because the place we deploy the definition of the chiral matrix is related to the position of the vertex we adopt a label of the version to these expressions, namely

First Version

$$\begin{aligned} K_{123}^{\nu_{123}} \mathrm{tr} \left(\gamma_* \gamma_{\mu_1 \nu_1 \mu_2 \nu_2 \mu_3 \nu_3} \right) &= -\varepsilon_{\mu_2 \mu_3 \nu_1 \nu_2} \left[K_{1\mu_1} K_{23}^{\nu_{12}} - K_{2\mu_1} K_{13}^{\nu_{12}} + K_{3\mu_1} K_{12}^{\nu_{12}} \right] \\ &- \varepsilon_{\mu_1 \mu_3 \nu_1 \nu_2} \left[K_{1\mu_2} K_{23}^{\nu_{12}} + K_{2\mu_2} K_{13}^{\nu_{12}} - K_{3\mu_3} K_{12}^{\nu_{12}} \right] \\ &+ \varepsilon_{\mu_1 \mu_2 \nu_1 \nu_2} \left[K_{1\mu_3} K_{23}^{\nu_{12}} - K_{2\mu_3} K_{13}^{\nu_{12}} - K_{3\mu_3} K_{12}^{\nu_{12}} \right] \\ &+ \varepsilon_{\mu_1 \mu_2 \mu_3 \nu_1} \left[-K_1^{\nu_1} \left(K_2 \cdot K_3 \right) + K_2^{\nu_1} \left(K_1 \cdot K_3 \right) - K_3^{\nu_1} \left(K_1 \cdot K_2 \right) \right] \\ &+ \left[-g_{\mu_1 \mu_2} \varepsilon_{\mu_3 \nu_1 \nu_2 \nu_3} - g_{\mu_2 \mu_3} \varepsilon_{\mu_1 \nu_1 \nu_2 \nu_3} + g_{\mu_1 \mu_3} \varepsilon_{\mu_2 \nu_1 \nu_2 \nu_3} \right] K_{123}^{\nu_{123}} \end{aligned}$$
(5.9)

Second Version

$$K_{123}^{\nu_{123}} \operatorname{tr} \left(\gamma_* \gamma_{\mu_2 \nu_2 \mu_3 \nu_3 \mu_1 \nu_1} \right) = + \varepsilon_{\mu_1 \mu_3 \nu_1 \nu_2} \left[K_{1\mu_2} K_{23}^{\nu_{12}} - K_{2\mu_2} K_{13}^{\nu_{12}} + K_{3\mu_2} K_{12}^{\nu_{12}} \right]$$

$$- \varepsilon_{\mu_1 \mu_2 \nu_1 \nu_2} \left[K_{1\mu_3} K_{23}^{\nu_{23}} + K_{2\mu_3} K_{13}^{\nu_{13}} + K_{3\mu_3} K_{12}^{\nu_{12}} \right]$$

$$- \varepsilon_{\mu_2 \mu_3 \nu_1 \nu_2} \left[K_{1\mu_1} K_{23}^{\nu_{23}} + K_{2\mu_1} K_{13}^{\nu_{13}} - K_{3\mu_1} K_{12}^{\nu_{12}} \right]$$

$$+ \varepsilon_{\mu_1 \mu_2 \mu_3 \nu_1} \left[-K_1^{\nu_1} \left(K_2 \cdot K_3 \right) - K_2^{\nu_1} \left(K_1 \cdot K_3 \right) + K_3^{\nu_1} \left(K_1 \cdot K_2 \right) \right]$$

$$+ \left[g_{\mu_1 \mu_2} \varepsilon_{\mu_3 \nu_1 \nu_2 \nu_3} - g_{\mu_1 \mu_3} \varepsilon_{\mu_2 \nu_1 \nu_2 \nu_3} - g_{\mu_2 \mu_3} \varepsilon_{\mu_1 \nu_1 \nu_2 \nu_3} \right] K_{123}^{\nu_{123}}$$
(5.10)

$$K_{123}^{\nu_{123}} \operatorname{tr} \left(\gamma_* \gamma_{\mu_3 \nu_3 \mu_1 \nu_1 \mu_2 \nu_2} \right) = -\varepsilon_{\mu_1 \mu_2 \nu_1 \nu_2} \left[K_{1\mu_3} K_{23}^{\nu_{12}} - K_{2\mu_3} K_{13}^{\nu_{12}} + K_{3\mu_3} K_{12}^{\nu_{12}} \right]$$

$$-\varepsilon_{\mu_2 \mu_3 \nu_1 \nu_2} \left[K_{1\mu_1} K_{23}^{\nu_{12}} - K_{2\mu_1} K_{13}^{\nu_{12}} - K_{3\mu_1} K_{12}^{\nu_{12}} \right]$$

$$-\varepsilon_{\mu_1 \mu_3 \nu_1 \nu_2} \left[K_{1\mu_2} K_{23}^{\nu_{12}} + K_{2\mu_2} K_{13}^{\nu_{12}} + K_{3\mu_2} K_{12}^{\nu_{12}} \right]$$

$$+\varepsilon_{\mu_1 \mu_2 \mu_3 \nu_1} \left[K_1^{\nu_1} \left(K_2 \cdot K_3 \right) - K_2^{\nu_1} \left(K_1 \cdot K_3 \right) - K_3^{\nu_1} \left(K_1 \cdot K_2 \right) \right]$$

$$+ \left[-g_{\mu_1 \mu_2} \varepsilon_{\mu_3 \nu_1 \nu_2 \nu_3} - g_{\mu_1 \mu_3} \varepsilon_{\mu_2 \nu_1 \nu_2 \nu_3} + g_{\mu_2 \mu_3} \varepsilon_{\mu_1 \nu_1 \nu_2 \nu_3} \right] K_{123}^{\nu_{123}}$$
(5.11)

Preceding the discussion of any particular third rank amplitude, we shall introduce here some definitions and general results which will be useful in what follows. If one looks at the three first rows of (5.9), (5.10) and (5.11), they all can be written in terms of tensors, which we call sign tensors (equivalently to 2D..)

$$\varepsilon_{\mu_{ab}\nu_{12}}t_{\mu_c}^{\nu_{12}(s_1s_2)} = \varepsilon_{\mu_{ab}\nu_{12}} \left(K_{1\mu_c}K_{23}^{\nu_{12}} + s_1K_{2\mu_c}K_{13}^{\nu_{12}} + s_2K_{3\mu_c}K_{12}^{\nu_{12}} \right) \frac{1}{D_{123}}$$
(5.12)

where $s_i = \pm 1$, thus using $K_i = K_j + p_{ij}$ and $\varepsilon_{\mu_{ab}\nu_{12}}K_{ij}^{\nu_{12}} = \varepsilon_{\mu_{ab}\nu_{12}}p_{ji}^{\nu_2}K_i^{\nu_1}$ and writing $p_{32} = p_{31} - p_{21}$ when necessary, we have

$$\varepsilon_{\mu_{ab}\nu_{12}}t_{\mu_{c}}^{\nu_{12}(s_{1}s_{2})} = \varepsilon_{\mu_{ab}\nu_{12}}\left[\left(1+s_{1}\right)p_{31}^{\nu_{2}}-\left(1-s_{2}\right)p_{21}^{\nu_{2}}\right]\frac{K_{1}^{\nu_{1}}K_{1\mu_{c}}}{D_{123}} + \varepsilon_{\mu_{ab}\nu_{12}}\left[p_{21}^{\nu_{1}}p_{32}^{\nu_{2}}K_{1\mu_{c}}+\left(s_{1}p_{21\mu_{c}}p_{31}^{\nu_{2}}+s_{2}p_{31\mu_{c}}p_{21}^{\nu_{2}}\right)K_{1}^{\nu_{1}}\right]\frac{1}{D_{123}}$$
(5.13)

and will be clear that the integral of one of these tensors is finite, namely

$$\varepsilon_{\mu_{ab}\nu_{12}}t_{\mu_c}^{\nu_{12}(-,+)} = \varepsilon_{\mu_{ab}\nu_{12}} \left[p_{21}^{\nu_1} p_{32}^{\nu_2} K_{1\mu_c} + \left(-p_{21\mu_c} p_{31}^{\nu_2} + p_{31\mu_c} p_{21}^{\nu_2} \right) K_1^{\nu_1} \right] \frac{1}{D_{123}},\tag{5.14}$$

because the presence of only one factor of the vector K_i or in other words only vector three-point integrals of power counting $\omega = -1$ appear here. After integration, and with the help of the result

$$J_3^{\mu} = -p_{21}^{\mu} Z_{10}^{(-1)} - p_{31}^{\mu} Z_{01}^{(-1)}, \qquad (5.15)$$

and the anti-symmetry of ε -tensor, we get the first relevant result

$$\varepsilon_{\mu_{ab}\nu_{12}}T^{\nu_{12}(-+)}_{\mu_c} = \varepsilon_{\mu_{ab}\nu_{12}} \left[p^{\nu_1}_{21} p^{\nu_2}_{32} J_{3\mu_c} + \left(-p_{21\mu_c} p^{\nu_2}_{31} + p_{31\mu_c} p^{\nu_2}_{21} \right) J^{\nu_1}_3 \right] = 0.$$
(5.16)

The other three non-zero ones, containing the log-diverging integral $\bar{J}_{3\mu_c}^{\nu_1}$ responsible for the divergent content, we present below

$$\varepsilon_{\mu_{ab}\nu_{12}}T^{\nu_{12}(+-)}_{\mu_{c}} = 2\varepsilon_{\mu_{ab}\nu_{12}} \left[p^{\nu_{2}}_{31} \left(J^{\nu_{1}}_{3\mu_{c}} + p_{21\mu_{c}} J^{\nu_{1}}_{3} \right) - p^{\nu_{2}}_{21} \left(J^{\nu_{1}}_{3\mu_{c}} + p_{31\mu_{c}} J^{\nu_{1}}_{3} \right) \right] + \frac{1}{2} \left(\varepsilon_{\mu_{ab}\nu_{12}} p^{\nu_{2}}_{32} \Delta^{\nu_{1}}_{3\mu_{c}} + \varepsilon_{\mu_{abc}\nu_{1}} p^{\nu_{1}}_{32} I_{\log} \right)$$
(5.17)

$$\varepsilon_{\mu_{ab}\nu_{12}}T^{\nu_{12}(--)}_{\mu_c} = -2\varepsilon_{\mu_{ab}\nu_{12}}p^{\nu_2}_{21}\left(J^{\nu_1}_{3\mu_c} + p_{31\mu_c}J^{\nu_1}_3\right) - \frac{1}{2}\left(\varepsilon_{\mu_{ab}\nu_{12}}p^{\nu_2}_{21}\Delta^{\nu_1}_{3\mu_c} + \varepsilon_{\mu_{abc}\nu_1}p^{\nu_1}_{21}I_{\log}\right)$$
(5.18)

$$\varepsilon_{\mu_{ab}\nu_{12}}T^{\nu_{12}(++)}_{\mu_c} = +2\varepsilon_{\mu_{ab}\nu_{12}}p^{\nu_2}_{31}\left(J^{\nu_1}_{3\mu_c} + p_{21\mu_c}J^{\nu_1}_3\right) + \frac{1}{2}\left(\varepsilon_{\mu_{ab}\nu_{12}}p^{\nu_2}_{31}\Delta^{\nu_1}_{3\mu_c} + \varepsilon_{\mu_{abc}\nu_1}p^{\nu_1}_{31}I_{\log}\right)$$
(5.19)

Hence the three versions will turn up each with a tensor that is shared with all rank-three pseudo triangles using that version of the trace we define them as

$$c_{1\mu_{123}} = -\varepsilon_{\mu_{23}\nu_{12}}t_{\mu_1}^{\nu_{12}(-+)} - \varepsilon_{\mu_{13}\nu_{12}}t_{\mu_2}^{\nu_{12}(+-)} + \varepsilon_{\mu_{12}\nu_{12}}t_{\mu_3}^{\nu_{12}(--)}$$
(5.20)

$$c_{2\mu_{123}} = +\varepsilon_{\mu_{13}\nu_{12}}t_{\mu_{2}}^{\nu_{12}(-+)} - \varepsilon_{\mu_{12}\nu_{12}}t_{\mu_{3}}^{\nu_{12}(++)} - \varepsilon_{\mu_{23}\nu_{12}}t_{\mu_{1}}^{\nu_{12}(+-)}$$
(5.21)

$$c_{3\mu_{123}} = -\varepsilon_{\mu_{12}\nu_{12}}t_{\mu_3}^{\nu_{12}(-+)} - \varepsilon_{\mu_{23}\nu_{12}}t_{\mu_1}^{\nu_{12}(--)} - \varepsilon_{\mu_{13}\nu_{12}}t_{\mu_2}^{\nu_{12}(++)}$$
(5.22)

whose integrals in terms of the previous results eq.(5.16) (5.17) (5.18) (5.19) are

$$C_{1\mu_{123}} = -\varepsilon_{\mu_{13}\nu_{12}}T^{\nu_{12}(+-)}_{\mu_2} + \varepsilon_{\mu_{12}\nu_{12}}T^{\nu_{12}(--)}_{\mu_3}$$
(5.23)

$$C_{2\mu_{123}} = -\varepsilon_{\mu_{12}\nu_{12}}T^{\nu_{12}(++)}_{\mu_3} - \varepsilon_{\mu_{23}\nu_{12}}T^{\nu_{12}(+-)}_{\mu_1}$$
(5.24)

$$C_{3\mu_{123}} = -\varepsilon_{\mu_{23}\nu_{12}}T^{\nu_{12}(--)}_{\mu_1} - \varepsilon_{\mu_{13}\nu_{12}}T^{\nu_{12}(++)}_{\mu_2}$$
(5.25)

Here another important point to notice is that the sampling of indexes mentioned reflects in the absence of the index μ_i of the vertex Γ_i in the T's of the C_i because it is present through the tensor $\varepsilon_{\mu_{ab}\nu_{12}}T_{\mu_c}^{\nu_{12}(-+)}$ in the eq.(5.16) that is finite and identically zero. This will enable us to anticipate violations of either WIs or RAGFs.

The next step is the vanishing of rank-one odd triangles, noticed that in the last rows of eq.'s (5.9), (5.10) and (5.11) they appear as components. It happens they all differ only by a sign and turn out to be finite and null, as an example

$$t_{\mu_i}^{ASS} = 4i\varepsilon_{\mu_i\nu_{123}}K_{123}^{\nu_{123}}\frac{1}{D_{123}} = 4i\varepsilon_{\mu_i\nu_{123}}p_{21}^{\nu_2}p_{31}^{\nu_3}\frac{K_1^{\nu_1}}{D_{123}}$$
(5.26)

due to the triple contraction, therefore using (5.15)

$$T^{ASS}_{\mu_i} = 4i\varepsilon_{\mu_i\nu_{123}}p^{\nu_2}_{21}p^{\nu_3}_{31}J^{\nu_1}_3 = 0$$
(5.27)

and the same for

$$T_{\mu_i}^{ASS} = T_{\mu_i}^{SAS} = T_{\mu_i}^{SSA} = T_{\mu_i}^{APP} = T_{\mu_i}^{PAP} = T_{\mu_i}^{PPA} = 0$$
(5.28)

$$T_{\mu_i}^{VSP} = T_{\mu_i}^{PVS} = T_{\mu_i}^{SPV} = T_{\mu_i}^{VPS} = T_{\mu_i}^{SVP} = T_{\mu_i}^{PSV} = 0$$
(5.29)

disappearing then from the integrated expressions.

The last feature on this web of relations is the distinguishing rank-one even triangles. Here we need a definite example, as the VPP which is defined by

$$\left(t^{VPP}\right)^{\nu_{1}} = \operatorname{tr}\left[\gamma^{\nu_{1}}S\left(1\right)\gamma_{*}S\left(2\right)\gamma_{*}S\left(3\right)\right] = 4\left(-K_{1}^{\nu_{1}}S_{23} + K_{2}^{\nu_{1}}S_{13} - K_{3}^{\nu_{1}}S_{12}\right)\frac{1}{D_{123}},\tag{5.30}$$

as always, before the integration during the trace operation, remember the combination $S_{ij} = K_i \cdot K_j - m^2$ was defined on eq. (4.17). Now is a direct task to see that combining the mass terms of the eq. (5.2) to the AVV integrand with the eq. (5.9) to the first version of the trace to get as a component sub-amplitude

$$\operatorname{sub}\left(t_{\mu_{123}}^{AVV}\right)_{1} = 4i\varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{1}}\left(-K_{1}^{\nu_{1}}S_{23} + K_{2}^{\nu_{1}}S_{13} - K_{3}^{\nu_{1}}S_{12}\right)\frac{1}{D_{123}} = i\varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{1}}\left(t^{VPP}\right)^{\nu_{1}}$$
(5.31)

where the pre-amplitude got the trace-version label. In the same vein then, it is easy to verify the table (III) accounts for all other possible combinations appearing in the explicit computation.

Version/Type	AVV	VAV	VVA	AAA
1	+VPP	+ASP	-APS	-VSS
2	-SAP	+PVP	+PAS	-SVS
3	+SPA	-PSA	+PPV	-SSV

TABLE III: Even sub-amplitudes corresponding to the vertex configuration of the rank-three odd ones and the version of the trace applied.

In other words, alongside the C_i tensor of each of the versions to each type of third-rank amplitude, there is one even and rank-one sub-amplitude. Back to the VPP example, its integrated form becomes

$$(T^{VPP})^{\nu_1} = -4 (p_{21} \cdot p_{32}) J_3^{\nu_1}$$

$$+ 2 [(p_{31}^{\nu_1} p_{21}^2 - p_{21}^{\nu_1} p_{31}^2) J_3 + p_{21}^{\nu_1} J_2 (p_{21}) - p_{32}^{\nu_1} J_2 (p_{32})]$$

$$+ 2 [P_{31}^{\nu_2} \Delta_{\nu_2}^{\nu_1} + (p_{21}^{\nu_1} - p_{32}^{\nu_1}) I_{\log}]$$

$$(5.32)$$

all the others are contemplated in the useful appendix (E), which is essential to verify all the expressions presented in the paper.

For completeness, in the first version of AVV, we note that the divergent part of the tensor $C_{1\mu_1\mu_2\mu_3}$, defined in eq. (5.23), can be read off from eq.'s (5.17) and (5.18), for $\varepsilon_{\mu_{13}\nu_{12}}T_{\mu_2}^{\nu_{12}(+-)}$ and $\varepsilon_{\mu_{12}\nu_{12}}T_{\mu_3}^{\nu_{12}(--)}$, as

$$4iC_{1\mu_{1}\mu_{2}\mu_{3}} = -2i\left(\varepsilon_{\mu_{13}\nu_{12}}p_{32}^{\nu_{2}}\Delta_{3\mu_{2}}^{\nu_{1}} + \varepsilon_{\mu_{12}\nu_{12}}p_{21}^{\nu_{2}}\Delta_{3\mu_{3}}^{\nu_{1}} + \varepsilon_{\mu_{123}\nu_{1}}\left(p_{21}^{\nu_{1}} - p_{32}^{\nu_{1}}\right)I_{\log}\right),$$

in such a way when combined with the divergent part of the sub-amplitude VPP in the eq. (5.32) above, it exactly cancels the object I_{log} . Then the total divergent content will be expressed exactly and solely in terms of surface terms

$$S_{1\mu_{1}\mu_{2}\mu_{3}} = -2i\left(\varepsilon_{\mu_{13}\nu_{12}}p_{32}^{\nu_{2}}\Delta_{3\mu_{2}}^{\nu_{1}} + \varepsilon_{\mu_{12}\nu_{12}}p_{21}^{\nu_{2}}\Delta_{3\mu_{3}}^{\nu_{1}}\right) + 2i\varepsilon_{\mu_{123}\nu_{1}}P_{31}^{\nu_{2}}\Delta_{3\nu_{2}}^{\nu_{1}}$$
(5.33)

hence, together with finite parts of the same eq.'s (5.17) and (5.18), the complete result assumes the form

$$(T^{AVV}_{\mu_{123}})_{1} = S_{1\mu_{1}\mu_{2}\mu_{3}}$$

$$-8i\varepsilon_{\mu_{13}\nu_{12}} \left[p_{31}^{\nu_{2}} \left(J^{\nu_{1}}_{3\mu_{2}} + p_{21\mu_{2}} J^{\nu_{1}}_{3} \right) - p_{21}^{\nu_{2}} \left(J^{\nu_{1}}_{3\mu_{2}} + p_{31\mu_{2}} J^{\nu_{1}}_{3} \right) \right]$$

$$-8i\varepsilon_{\mu_{12}\nu_{12}} p_{21}^{\nu_{2}} \left(J^{\nu_{1}}_{3\mu_{3}} + p_{31\mu_{3}} J^{\nu_{1}}_{3} \right)$$

$$-4i\varepsilon_{\mu_{123}\nu_{1}} \left(p_{21} \cdot p_{32} \right) J^{\nu_{1}}_{3} + 2i\varepsilon_{\mu_{123}\nu_{1}} \left[\left(p_{31}^{\nu_{1}} p_{21}^{2} - p_{21}^{\nu_{1}} p_{31}^{2} \right) J_{3} \right]$$

$$+2i\varepsilon_{\mu_{123}\nu_{1}} \left[p_{21}^{\nu_{1}} J_{2} \left(p_{21} \right) - p_{32}^{\nu_{1}} J_{2} \left(p_{32} \right) \right].$$

$$(5.34)$$

Some statements are in order here, the exact cancellation among the divergent objects I_{\log} of the common tensors and sub-amplitudes, separated from the finite integrals using the identity (3.5), happens for all versions and all amplitudes (AVV, VAV, VVA, AAA). And the surface terms content, $S_{1\mu_1\mu_2\mu_3}$, of the combined result is the same for all amplitudes, and only depends on the version in question, this being one of the reasons we did not attach a super index to these structures.

For later use, we must define the other two sets of surface terms appearing in versions two and three of any amplitude

$$S_{2\mu_{123}} = -2i\left(\varepsilon_{\mu_{12}\nu_{12}}p_{31}^{\nu_2}\Delta_{3\mu_3}^{\nu_1} + \varepsilon_{\mu_{23}\nu_{12}}p_{32}^{\nu_2}\Delta_{3\mu_1}^{\nu_1}\right) + 2i\varepsilon_{\mu_{123}\nu_1}P_{21}^{\nu_2}\Delta_{3\nu_2}^{\nu_1}$$
(5.36)

$$S_{3\mu_{123}} = -2i\left(\varepsilon_{\mu_{13}\nu_{12}}p_{31}^{\nu_2}\Delta_{3\mu_2}^{\nu_1} - \varepsilon_{\mu_{23}\nu_{12}}p_{21}^{\nu_2}\Delta_{3\mu_1}^{\nu_1}\right) + 2i\varepsilon_{\mu_{123}\nu_1}P_{32}^{\nu_2}\Delta_{3\nu_2}^{\nu_1}$$
(5.37)

$$\left(T_{\mu_{123}}^{AVV}\right)_{1} = 4iC_{1\mu_{123}} + i\varepsilon_{\mu_{123}\nu_{1}} \left(T^{VPP}\right)^{\nu_{1}}$$
(5.38)

$$\left(T^{AVV}_{\mu_{123}}\right)_2 = 4iC_{2\mu_{123}} - i\varepsilon_{\mu_{123}\nu_1} \left(T^{SAP}\right)^{\nu_1}$$
(5.39)

$$\left(T_{\mu_{123}}^{AVV}\right)_{3} = 4iC_{3\mu_{123}} + i\varepsilon_{\mu_{123}\nu_{1}} \left(T^{SPA}\right)^{\nu_{1}}$$
(5.40)

as for VAV, VVA and AAA they have the same structure

$$(T^{\Gamma_1\Gamma_2\Gamma_3}_{\mu_{123}})_i = 4iC_{i\mu_{123}} \pm i\varepsilon_{\mu_{123}\nu_1} (\text{Corresponding sub-amplitude})^{\nu_1}$$

where the sub-amplitude is given by the table (III) above.

The set of surface terms $S_{i\mu_1\mu_2\mu_3}$ defined in eq.'s (5.33), (5.36) and (5.37), pointing out again, are functions only of the version and not of the vertex content of the diagram. Their full verification is easily accomplished using the appendix (E) for the sub-amplitudes, where both divergent and finite parts are expressed explicitly. This is important because the mechanism of violation of RAGFs is thoroughly dependent on the topology and the traces of six matrices in the diagram.

The last element, that appear when investigating RAGFs, are the three-point and rank-two amplitudes, they are finite after integrated and given by

$$-2mT_{\mu_{23}}^{PVV} = \varepsilon_{\mu_{23}\nu_{12}} p_{21}^{\nu_1} p_{31}^{\nu_2} \left(8im^2 J_3\right)$$
(5.41)

$$2mT_{\mu_{13}}^{VPV} = \varepsilon_{\mu_{13}\nu_{12}} p_{21}^{\nu_1} p_{31}^{\nu_2} \left(8im^2 J_3\right)$$
(5.42)

$$2mT_{\mu_{12}}^{VVP} = \varepsilon_{\mu_{12}\nu_{12}} p_{21}^{\nu_1} p_{32}^{\nu_2} \left(-8im^2 J_3\right), \qquad (5.43)$$

the signs and the mass factor are only a matter of convenience for they appear in this form in the contractions with external momenta in the single axial triangles. To the triple axial triangle, in its contractions with the momenta, will appear the finite triangles below

$$-2mT_{\mu_{23}}^{PAA} = \varepsilon_{\mu_{23}\nu_{12}}p_{31}^{\nu_2} \left[8im^2 \left(2J_3^{\nu_1} + p_{21}^{\nu_1}J_3\right)\right]$$
(5.44)

$$2mT_{\mu_{13}}^{APA} = \varepsilon_{\mu_{13}\nu_{12}}p_{21}^{\nu_2} \left[-8im^2 \left(2J_3^{\nu_1} + p_{31}^{\nu_1}J_3\right)\right]$$
(5.45)

$$2mT_{\mu_{12}}^{AAP} = \varepsilon_{\mu_{12}\nu_{12}} p_{32}^{\nu_2} \left[8im^2 \left(2J_3^{\nu_1} + p_{21}^{\nu_1} J_3 \right) \right], \qquad (5.46)$$

the form factors of all these tensors can be obtaining retrieving the definitions of the scalar and vector 3pt-integrals

$$J_3 = \frac{i}{(4\pi)^2} Z_{00}^{(-1)}; \quad J_3^{\nu} = -\frac{i}{(4\pi)^2} \left[p_{21}^{\nu} Z_{10}^{(-1)} + p_{31}^{\nu} Z_{01}^{(-1)} \right]$$

and are being explicited for future use, resorting to their value in the point $q_i \cdot q_j = 0$

$$Z_{rs}^{(-1)}\Big|_0 = -\frac{r!s!}{m^2 \left[(r+s+2)!\right]},$$

where these momenta represent all the possible difference of routings, we can display a precise low energy behavior to these tensors

$$-2mT_{\mu_{23}}^{PVV}\big|_{0} = \frac{\varepsilon_{\mu_{23}\nu_{12}}p_{21}^{\nu_{1}}p_{32}^{\nu_{2}}}{(2\pi)^{2}}; \quad 2mT_{\mu_{13}}^{VPV}\big|_{0} = \frac{\varepsilon_{\mu_{13}\nu_{12}}p_{21}^{\nu_{1}}p_{32}^{\nu_{2}}}{(2\pi)^{2}}; \quad 2mT_{\mu_{12}}^{VVP}\big|_{0} = \frac{\varepsilon_{\mu_{12}\nu_{12}}p_{21}^{\nu_{1}}p_{32}^{\nu_{2}}}{(2\pi)^{2}}$$
(5.47)

and

$$-2mT_{\mu_{23}}^{PAA}\big|_{0} = \frac{\varepsilon_{\mu_{23}\nu_{12}}p_{21}^{\nu_{1}}p_{32}^{\nu_{2}}}{3\left(2\pi\right)^{2}}; \quad 2mT_{\mu_{13}}^{APA}\big|_{0} = \frac{\varepsilon_{\mu_{13}\nu_{12}}p_{21}^{\nu_{1}}p_{32}^{\nu_{2}}}{3\left(2\pi\right)^{2}}; \quad 2mT_{\mu_{12}}^{AAP}\big|_{0} = \frac{\varepsilon_{\mu_{12}\nu_{12}}p_{21}^{\nu_{1}}p_{32}^{\nu_{2}}}{3\left(2\pi\right)^{2}}$$
(5.48)

the consequences of these values will be determined and examined in section (5.2).

5.1. Relations Among Green Functions and Uniqueness

Recalling the chapter two the standard procedure to obtain the RAGFs, that allows to state to $t_{\mu_{123}}^{AVV}$

$$p_{31}^{\mu_1} t_{\mu_{123}}^{AVV} = t_{\mu_{32}}^{AV} (1,2) - t_{\mu_{23}}^{AV} (2,3) - 2m t_{\mu_{23}}^{PVV} p_{21}^{\mu_2} t_{\mu_{123}}^{AVV} = t_{\mu_{13}}^{AV} (1,3) - t_{\mu_{13}}^{AV} (2,3) p_{32}^{\mu_3} t_{\mu_{123}}^{AVV} = t_{\mu_{12}}^{AV} (1,2) - t_{\mu_{12}}^{AV} (1,3)$$

to $t_{\mu_{123}}^{VAV}$

$$p_{31}^{\mu_1} t_{\mu_{123}}^{VAV} = t_{\mu_{23}}^{AV} (2,1) - t_{\mu_{23}}^{AV} (2,3)$$

$$p_{21}^{\mu_2} t_{\mu_{123}}^{VAV} = t_{\mu_{31}}^{AV} (3,1) - t_{\mu_{13}}^{AV} (2,3) + 2m t_{\mu_{13}}^{VPV}$$

$$p_{32}^{\mu_3} t_{\mu_{123}}^{VAV} = t_{\mu_{21}}^{AV} (2,1) - t_{\mu_{21}}^{AV} (3,1)$$

to $t_{\mu_{123}}^{VVA}$

$$p_{31}^{\mu_1} t_{\mu_{123}}^{VVA} = t_{\mu_{32}}^{AV} (1,2) - t_{\mu_{32}}^{AV} (3,2) p_{21}^{\mu_2} t_{\mu_{123}}^{VVA} = t_{\mu_{31}}^{AV} (3,1) - t_{\mu_{31}}^{AV} (3,2) p_{32}^{\mu_3} t_{\mu_{123}}^{VVA} = t_{\mu_{12}}^{AV} (1,2) - t_{\mu_{21}}^{AV} (3,1) + 2m t_{\mu_{12}}^{VVP}$$

to $t_{\mu_{123}}^{AAA}$

$$\begin{array}{ll} p_{31}^{\mu_1}t_{\mu_{123}}^{AAA} &= t_{\mu_{23}}^{AV}\left(2,1\right) - t_{\mu_{32}}^{AV}\left(3,2\right) - 2mt_{\mu_{23}}^{PAA} \\ p_{21}^{\mu_2}t_{\mu_{123}}^{AAA} &= t_{\mu_{13}}^{AV}\left(1,3\right) - t_{\mu_{31}}^{AV}\left(3,2\right) + 2mt_{\mu_{13}}^{APA} \\ p_{32}^{\mu_3}t_{\mu_{123}}^{AAA} &= t_{\mu_{21}}^{AV}\left(2,1\right) - t_{\mu_{12}}^{AV}\left(1,3\right) + 2mt_{\mu_{12}}^{AAP}. \end{array}$$

The RHS of these identities show two and three-point functions, the two-point ones

$$t_{\mu_{ij}}^{AV}(a,b) = \operatorname{tr}[\gamma_*\gamma_{\mu_i}S(a)\gamma_{\mu_j}S(b)] = -4i\varepsilon_{\mu_i\mu_j\nu_1\nu_2}p_{ba}^{\nu_2}\frac{K_a^{\nu_1}}{D_{ab}}$$
(5.49)

are odd tensor functions of two variables, the external and ambiguous momenta (p_{ij}, P_{ij}) . After integration the result is proportional to the vector two-point function that has a naturally linear dependence on the ambiguous momenta, looking at the computed expression as an example in the section (3.3), eq. (3.23) it is straightforward to see that any of these amplitudes is a pure surface

$$T_{\mu_{ij}}^{AV}(a,b) = -4i\varepsilon_{\mu_i\mu_j\nu_1\nu_2} p_{ba}^{\nu_2} \bar{J}_2^{\nu_1}(a,b) = 2i\varepsilon_{\mu_i\mu_j\nu_1\nu_2} p_{ba}^{\nu_2} P_{ab}^{\nu_3} \Delta_{3\nu_3}^{\nu_1}$$
(5.50)

with the help of the relations among the (p_{ij}, P_{ij}) to AVV relations we obtain

$$T_{\mu_{32}}^{AV}(1,2) - T_{\mu_{23}}^{AV}(2,3) = -2i\varepsilon_{\mu_{23}\nu_{12}} \left(p_{21}^{\nu_2} P_{12}^{\nu_3} + p_{32}^{\nu_2} P_{32}^{\nu_3} \right) \Delta_{3\nu_3}^{\nu_1}$$
(5.51)

$$T_{\mu_{13}}^{AV}(1,3) - T_{\mu_{13}}^{AV}(2,3) = -2i\varepsilon_{\mu_{13}\nu_{12}} \left(p_{32}^{\nu_2} P_{32}^{\nu_3} - p_{31}^{\nu_2} P_{31}^{\nu_3} \right) \Delta_{3\nu_3}^{\nu_1}$$
(5.52)

$$T_{\mu_{12}}^{AV}(1,2) - T_{\mu_{12}}^{AV}(1,3) = -2i\varepsilon_{\mu_{12}\nu_{12}}\left(p_{31}^{\nu_2}P_{31}^{\nu_3} - p_{21}^{\nu_2}P_{21}^{\nu_3}\right)\Delta_{3\nu_3}^{\nu_1}$$
(5.53)

Such results could be different for the relations to the VAV, VVA, and AAA diagrams, but turn out they are not, and in truth depend only on the vertex contraction that the reader can be verify using the symmetry properties in the eq. (5.50), and thus will not be exhibited here.

Before going to the calculations we must remind the reader that if the RAGF were automatically valid, as one would be led to think because they are linear relations coming from integration, then the choice of to get rid of the surface terms could easily determine valid all the WI, the why of such a thing do not happen is all the specialty of these amplitudes.

As the contractions with the external momenta in the explicitly computed three rank amplitudes are concerned, it is necessary to state that the three non-vanishing sign tensors have a set of contraction properties, first to the momenta of each vertex one of them is null under contraction.

The finite part of the basic sign tensors on the eq.'s (5.17) (5.18) and (5.19) have a set of contractions with the external momenta that can are obtained using the appendix (C). The finite part of the tensors C_i can be shown to obey

$$p_{21}^{\mu_2}C_{1\mu_{123}} = +\varepsilon_{\mu_{13}\nu_1\nu_2}p_{32}^{\nu_2}p_{21}^2 \left(J_3^{\nu_1} + p_{21}^{\nu_1}J_3\right) - \frac{1}{2}\varepsilon_{\mu_{13}\nu_1\nu_2}p_{21}^{\nu_1}p_{31}^{\nu_2}J_2\left(p_{31}\right)$$

$$p_{32}^{\mu_3}C_{1\mu_{123}} = -\varepsilon_{\mu_{12}\nu_1\nu_2}p_{21}^{\nu_2}p_{32}^2J_3^{\nu_1} + \frac{1}{2}\varepsilon_{\mu_{12}\nu_1\nu_2}p_{21}^{\nu_1}p_{31}^{\nu_2}J_2\left(p_{31}\right)$$

$$p_{31}^{\mu_1}C_{2\mu_{123}} = \varepsilon_{\mu_{23}\nu_1\nu_2}p_{32}^{\nu_2}p_{31}^2 \left(J_3^{\nu_1} + p_{21}^{\nu_1}J_3\right) - \frac{1}{2}\varepsilon_{\mu_{23}\nu_1\nu_2}p_{21}^{\nu_1}p_{31}^{\nu_2}J_2\left(p_{21}\right)$$

$$p_{32}^{\mu_3}C_{2\mu_{123}} = \varepsilon_{\mu_{12}\nu_1\nu_2}p_{31}^{\nu_2}p_{32}^2J_3^{\nu_1} + \frac{1}{2}\varepsilon_{\mu_{12}\nu_1\nu_2}p_{21}^{\nu_1}p_{31}^{\nu_2}J_2\left(p_{21}\right)$$

$$p_{31}^{\mu_1}C_{3\mu_{123}} = +\varepsilon_{\mu_{23}\nu_1\nu_2}p_{21}^{\nu_2}p_{31}^2J_3^{\nu_1} - \frac{1}{2}\varepsilon_{\mu_{23}\nu_1\nu_2}p_{21}^{\nu_1}p_{31}^{\nu_2}J_2(p_{32})$$

$$p_{21}^{\mu_2}C_{3\mu_{123}} = -\varepsilon_{\mu_{13}\nu_1\nu_2}p_{31}^{\nu_2}p_{21}^2J_3^{\nu_1} - \frac{1}{2}\varepsilon_{\mu_{13}\nu_1\nu_2}p_{21}^{\nu_1}p_{31}^{\nu_2}J_2(p_{32})$$

and the contraction when the momenta being contracted matches the version

$$p_{31}^{\mu_1} C_{1\mu_{123}} = \varepsilon_{\mu_2\mu_3\nu_1\nu_2} \left[\left(p_{31}^{\nu_2} p_{21}^2 - p_{21}^{\nu_2} p_{31}^2 \right) J_3^{\nu_1} + p_{21}^{\nu_1} p_{32}^{\nu_2} J_2 \left(p_{32} \right) \right] \\ + \varepsilon_{\mu_2\mu_3\nu_1\nu_2} p_{21}^{\nu_1} p_{31}^{\nu_2} \left[2m^2 J_3 + i \left(4\pi\right)^{-2} \right]$$

$$p_{21}^{\mu_2} C_{2\mu_{123}} = \varepsilon_{\mu_1 \mu_3 \nu_1 \nu_2} \left[\left(p_{31}^{\nu_2} p_{21}^2 - p_{21}^{\nu_2} p_{31}^2 \right) J_3^{\nu_1} + p_{21}^{\nu_1} p_{32}^{\nu_2} J_2 \left(p_{32} \right) \right] \\ + \varepsilon_{\mu_1 \mu_3 \nu_1 \nu_2} p_{21}^{\nu_1} p_{31}^{\nu_2} \left[2m^2 J_3 + i \left(4\pi \right)^{-2} \right]$$

$$p_{32}^{\mu_3} C_{3\mu_{123}} = -\varepsilon_{\mu_1\mu_2\nu_1\nu_2} \left[\left(p_{31}^{\nu_2} p_{21}^2 - p_{21}^{\nu_2} p_{31}^2 \right) J_3^{\nu_1} + p_{21}^{\nu_1} p_{32}^{\nu_2} J_2 \left(p_{32} \right) \right] \\ -\varepsilon_{\mu_1\mu_2\nu_1\nu_2} p_{21}^{\nu_1} p_{31}^{\nu_2} \left[2m^2 J_3 + i \left(4\pi\right)^{-2} \right]$$

The equations were grouped in this way to emphasize that in each $C_{i\mu_1\mu_2\mu_3}$ the contraction with the momentum corresponding to μ_i needs the trace of $J_{3\mu\nu}$ see eq. (C2), or equivalently the reduction of $Z_{00}^{(0)}$ in terms of $Z_{rs}^{(-1)}$ and $Z_r^{(0)}$, eq. (C1), where appear a momentum-independent constant. It is essential to realize that all the previous results refer to definite relations among finite tensors and functions obtained free from any regularization procedure.

It was basically used

$$\varepsilon_{[\mu_1\mu_3\nu_1\nu_2}p_{ij\nu_3]}J_3^{\nu_1} = 0; \quad \varepsilon_{[\mu_a\nu_1\nu_2\nu_3}J_{3\mu_c]}^{\nu_1} = 0, \tag{5.54}$$

in order to search contractions of the J integrals with the momenta when they are not directly available, which enable us to reduce the functions by the formulas in the appendix. As a byproduct, we get the trace of the tensor integrals, and when there are three contractions involving the momenta and the vector integrals with the ε , such terms vanish⁶.

For the contractions with the set of surface terms $S_{i\mu_{123}}$, see eq.'s (5.33), (5.36) and (5.37), must be noted that the index μ_i does not appear in the tensor $\Delta_{3\mu\nu}$, only in the ε -tensor, and thus the contraction with the other indexes can be done directly, giving the term corresponding to the correct difference of two point AV functions, by example

$$p_{21}^{\mu_2} S_{1\mu_1\mu_2\mu_3} = -2i p_{21}^{\mu_2} \left(\varepsilon_{\mu_{13}\nu_{12}} p_{32}^{\nu_2} \Delta_{3\mu_2}^{\nu_1} + \varepsilon_{\mu_{12}\nu_{12}} p_{21}^{\nu_2} \Delta_{3\mu_3}^{\nu_1} \right) + 2i \varepsilon_{\mu_{123}\nu_1} p_{21}^{\mu_2} P_{31}^{\nu_2} \Delta_{3\nu_2}^{\nu_1} \\ = 2i \varepsilon_{\mu_{13}\nu_{12}} \left(-p_{21}^{\nu_3} p_{32}^{\nu_2} + p_{21}^{\nu_2} P_{31}^{\nu_3} \right) \Delta_{3\nu_3}^{\nu_1},$$
(5.55)

organizing the momenta (p_{ij}, P_{ij}) by $p_{ij} = P_{ir} - P_{jr}$, see the defining eq.'s (2.8) and (2.9), follows

$$p_{21}^{\mu_2} S_{1\mu_{123}} = -2i\varepsilon_{\mu_{13}\nu_{12}} \left(p_{32}^{\nu_2} P_{32}^{\nu_3} - p_{31}^{\nu_2} P_{31}^{\nu_3} \right) \Delta_{3\nu_3}^{\nu_1}.$$
(5.56)

The important feature here is that the contraction with $p_{31}^{\mu_1}$ only hits the index μ_1 in the ε -tensor letting the free indexes inside the tensors $\Delta_{3\mu\nu}$, however the AV functions have the indexes of its surface terms both contracted with the momenta, thereby to switch the indexes is just a matter of contract with that momentum and of using

$$\varepsilon_{\mu_1\mu_3\nu_1\nu_2}\Delta_{3\mu_2}^{\nu_1} - \varepsilon_{\mu_1\mu_2\nu_1\nu_2}\Delta_{3\mu_3}^{\nu_1} - \varepsilon_{\mu_2\mu_3\nu_1\nu_2}\Delta_{3\mu_1}^{\nu_1} - \varepsilon_{\mu_1\mu_2\mu_3\nu_1}\Delta_{3\nu_2}^{\nu_1} = -\varepsilon_{\mu_1\mu_2\mu_3\nu_2}\Delta_{3\nu_1}^{\nu_1}, \qquad (5.57)$$

plus a simple reorganization of the momenta, to reach at

$$p_{31}^{\mu_1} S_{1\mu_{123}} = -2i\varepsilon_{\mu_{23}\nu_{12}} \left(p_{21}^{\nu_2} P_{12}^{\nu_3} + p_{32}^{\nu_2} P_{32}^{\nu_3} \right) \Delta_{\nu_3}^{\nu_1} + 2i\varepsilon_{\mu_2\mu_3\nu_2\nu_3} p_{21}^{\nu_2} p_{31}^{\nu_3} \Delta_{3\nu_1}^{\nu_1}, \tag{5.58}$$

the remaining contraction with $p_{32}^{\mu_3}$ follows the example of the contraction with $p_{21}^{\mu_2}$, and is given by

$$p_{32}^{\mu_3} S_{1\mu_{123}} = -2i\varepsilon_{\mu_{12}\nu_{12}} \left(p_{31}^{\nu_2} P_{31}^{\nu_3} - p_{21}^{\nu_2} P_{21}^{\nu_3} \right) \Delta_{3\nu_3}^{\nu_1}.$$
(5.59)

The same phenomenon happens to the other two sets for which we quote the results

$$p_{31}^{\mu_1} S_{2\mu_{123}} = -2i\varepsilon_{\mu_{23}\nu_{12}} \left(p_{21}^{\nu_2} P_{12}^{\nu_3} + p_{32}^{\nu_2} P_{32}^{\nu_3} \right) \Delta_{3\nu_3}^{\nu_1} p_{21}^{\mu_2} S_{2\mu_{123}} = -2i\varepsilon_{\mu_{13}\nu_{12}} \left(p_{32}^{\nu_2} P_{32}^{\nu_3} - p_{31}^{\nu_2} P_{31}^{\nu_3} \right) \Delta_{3\nu_3}^{\nu_1} + 2i\varepsilon_{\mu_1\mu_3\nu_2\nu_3} p_{21}^{\nu_2} p_{31}^{\nu_3} \Delta_{3\nu_1}^{\nu_1}$$
(5.60)
$$p_{32}^{\mu_3} S_{2\mu_{123}} = -2i\varepsilon_{\mu_{12}\nu_{12}} \left(p_{31}^{\nu_2} P_{31}^{\nu_3} - p_{21}^{\nu_2} P_{21}^{\nu_3} \right) \Delta_{3\nu_3}^{\nu_1}$$

and for $S_{3\mu_1\mu_2\mu_3}$

$$p_{31}^{\mu_1} S_{3\mu_{123}} = -2i\varepsilon_{\mu_{23}\nu_{12}} \left(p_{21}^{\nu_2} P_{12}^{\nu_3} + p_{32}^{\nu_2} P_{32}^{\nu_3} \right) \Delta_{3\nu_3}^{\nu_1} p_{21}^{\mu_2} S_{3\mu_{123}} = -2i\varepsilon_{\mu_{13}\nu_{12}} \left(p_{32}^{\nu_2} P_{32}^{\nu_3} - p_{31}^{\nu_2} P_{31}^{\nu_3} \right) \Delta_{3\nu_3}^{\nu_1} p_{32}^{\mu_3} S_{3\mu_{123}} = -2i\varepsilon_{\mu_{12}\nu_{12}} \left(p_{31}^{\nu_2} P_{31}^{\nu_3} - p_{21}^{\nu_2} P_{21}^{\nu_3} \right) \Delta_{3\nu_3}^{\nu_1} - 2i\varepsilon_{\mu_1\mu_2\nu_2\nu_3} p_{21}^{\nu_2} p_{31}^{\nu_3} \Delta_{3\nu_3}^{\nu_1}$$

⁶ A common situation is when appearing a term like, $\varepsilon_{\mu_1\mu_3\nu_1\nu_2} \left[(p_{21} \cdot p_{31}) p_{21}^{\nu_2} - p_{31}^{\nu_2} p_{21}^{2} \right] J_3^{\nu_1}$, to find contraction with the momentum one just use $\varepsilon_{[\mu_1\mu_3\nu_1\nu_2}p_{21\nu_3]} = 0$, then multiplying by $p_{31}^{\nu_2} p_{21}^{\nu_3} J_3^{\nu_1}$, and get

$$\left[\varepsilon_{\mu_{1}\mu_{3}\nu_{1}\nu_{2}}p_{21}^{2}p_{31}^{\nu_{2}}J_{3}^{\nu_{1}}+\varepsilon_{\nu_{3}\mu_{1}\mu_{3}\nu_{1}}\left(p_{21}\cdot p_{31}\right)p_{21}^{\nu_{3}}J_{3}^{\nu_{1}}\right]=-\varepsilon_{\nu_{2}\nu_{3}\mu_{1}\mu_{3}}p_{31}^{\nu_{2}}p_{21}^{\nu_{3}}\left(p_{21\nu_{1}}J_{3}^{\nu_{1}}\right)$$

the RHS is the desired contraction, in this derivation terms like $\varepsilon_{\nu_1\nu_2\nu_3\mu_1}p_{31}^{\nu_2}p_{21}^{\nu_3}J_3^{\nu_1}$ vanish because the number of contraction and that the vector integral is proportional to the external momenta. Adjusting signs and indexes, the equation is exactly the one sought.

Now that we have laid down the tools to analyze the integrated expression, it is an issue of applying the devices presented until this moment to establish the contraction with the first version of the calculated three-point diagrams, starting with the AVV

$$p_{31}^{\mu_1} \left(T_{\mu_{123}}^{AVV} \right)_1 = T_{\mu_{32}}^{AV} \left(1, 2 \right) - T_{\mu_{23}}^{AV} \left(2, 3 \right) - 2m T_{\mu_{23}}^{PVV} + \underbrace{2i\varepsilon_{\mu_2\mu_3\nu_1\nu_2} p_{21}^{\nu_1} p_{31}^{\nu_2} \left[\Delta_{3\alpha}^{\alpha} + 2i \left(4\pi \right)^{-2} \right]}_{p_{21}^{\mu_2}} p_{21}^{AVV} \left(T_{\mu_{123}}^{AVV} \right)_1 = T_{\mu_{13}}^{AV} \left(1, 3 \right) - T_{\mu_{13}}^{AV} \left(2, 3 \right)$$

$$p_{32}^{\mu_3} \left(T_{\mu_{123}}^{AVV} \right)_1 = T_{\mu_{12}}^{AV} \left(1, 2 \right) - T_{\mu_{12}}^{AV} \left(1, 3 \right)$$

and noticing the same additional term will appear for the corresponding version of VAV in the contraction with the momentum $p_{31}^{\mu_1}$

as the other RAGFs will be satisfied without conditions and exactly the same thing for VVA and AAA.

It can be of no surprise that the second and third versions of all configurations of vertexes will show up the possibly violating factor $2i\varepsilon_{\mu_i\mu_j\nu_1\nu_2}p_{21}^{\nu_1}p_{31}^{\nu_2}\left[\Delta_{3\alpha}^{\alpha}+2i(4\pi)^{-2}\right]$ in the contraction with the momenta $p_{21}^{\mu_2}$ and $p_{32}^{\mu_3}$ entering in the second and third vertexes respectively.

The explicitly computed equations below subsume all the results to for the verification of the RAGFs

$$q_{1}^{\mu_{1}} \left[T_{\mu_{123}}^{\Gamma_{123}} \right]_{1}^{\text{viol}} = +2i\varepsilon_{\mu_{2}\mu_{3}\nu_{1}\nu_{2}} p_{21}^{\nu_{1}} p_{32}^{\nu_{2}} \left[\Delta_{3\alpha}^{\alpha} + 2i\left(4\pi\right)^{-2} \right]$$
(5.61)

$$q_{2}^{\mu_{2}} \left[T_{\mu_{123}}^{\Gamma_{123}} \right]_{2}^{\text{viol}} = +2i\varepsilon_{\mu_{1}\mu_{3}\nu_{1}\nu_{2}} p_{21}^{\nu_{1}} p_{32}^{\nu_{2}} \left[\Delta_{3\alpha}^{\alpha} + 2i\left(4\pi\right)^{-2} \right]$$
(5.62)

$$q_{3}^{\mu_{3}}\left[T_{\mu_{123}}^{\Gamma_{123}}\right]_{3}^{\text{viol}} = -2i\varepsilon_{\mu_{1}\mu_{2}\nu_{1}\nu_{2}}p_{21}^{\nu_{1}}p_{32}^{\nu_{2}}\left[\Delta_{3\alpha}^{\alpha} + 2i\left(4\pi\right)^{-2}\right],\tag{5.63}$$

where we adopt the notation to the routing differences $q_1 = p_{31}$, $q_2 = p_{21}$, and $q_3 = p_{32}$ to mark a convention for first, second, and third vertexes respectively, this has already appeared in the Fig.(1) for the general diagram. Whereas the symbol $\Gamma_{123} \equiv \Gamma_1 \Gamma_2 \Gamma_3$ encode all the possibilities of combination vertexes $\Gamma_i \in \{A, V\}$ for which the number of axial vertexes is odd. To visualize this violation pattern, we offer the schematic graph

$$q_{i}^{\mu_{i}} \left(\Gamma_{1} \underbrace{\left(\begin{array}{c} 1 \\ \Gamma_{1} \\ 3 \end{array}\right)}_{3} \\ \Gamma_{3} \end{array}\right)_{j}^{\nu_{1}} = 2i\delta_{ij}\varepsilon_{\mu_{i}\mu_{j}\nu_{1}\nu_{2}}p_{31}^{\nu_{1}}\left[\Delta_{3\alpha}^{\alpha} + 2i\left(4\pi\right)^{-2}\right]_{3}^{\nu_{1}}$$

FIG. 2: The violation factor of the RAGF established for the contraction with momenta $q_i^{\mu_1}$.

Uniqueness: The other subtle point is uniqueness. Any precise and unambiguous discussion about this point needs a concrete definition of uniqueness. Anyway, with the help of the explicit expressions we have determined, it is established the equations

$$\begin{bmatrix} T_{\mu_{123}}^{\Gamma_{123}} \end{bmatrix}_{1}^{2} - \begin{bmatrix} T_{\mu_{123}}^{\Gamma_{123}} \end{bmatrix}_{2}^{2} = +2i\varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{1}}p_{32}^{\nu_{1}} \left(\Delta_{3\alpha}^{\alpha} + 2i\left(4\pi\right)^{-2}\right) \\ \begin{bmatrix} T_{\mu_{123}}^{\Gamma_{123}} \end{bmatrix}_{1}^{2} - \begin{bmatrix} T_{\mu_{123}}^{\Gamma_{123}} \end{bmatrix}_{3}^{2} = -2i\varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{1}}p_{21}^{\nu_{1}} \left(\Delta_{3\alpha}^{\alpha} + 2i\left(4\pi\right)^{-2}\right) \\ \begin{bmatrix} T_{\mu_{123}}^{\Gamma_{123}} \end{bmatrix}_{2}^{2} - \begin{bmatrix} T_{\mu_{123}}^{\Gamma_{123}} \end{bmatrix}_{3}^{2} = -2i\varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{1}}p_{31}^{\nu_{1}} \left(\Delta_{3\alpha}^{\alpha} + 2i\left(4\pi\right)^{-2}\right),$$

obtained by the same means as for the momenta contractions. Uniqueness then, here, means to us, that any possible way to compute the same expression returns the same result, in other words, the RHS is conditioned by this definition to be zero.

Anyway, the integrands we have been treating so far are well-defined tensors and they obey

$$\left(t_{\mu_{123}}^{\Gamma_{123}} \right)_1 = \left(t_{\mu_{123}}^{\Gamma_{123}} \right)_2 = \left(t_{\mu_{123}}^{\Gamma_{123}} \right)_3,$$

well, after separating two sectors, one in which physical parameters are factored out of the integrals and another sector where it is not possible, but every integral is finite. It could seem that the above equation should be true and independent of interpretation given to the divergent aspects, insofar as the same values are given to the undetermined parts. Nonetheless, the sorting of indexes makes the results to the finite parts unequal at the same time making the surface terms carry configuration of indexes that condition the satisfaction of distinct sets of RAGF. What these features end up showing is that if the computations furnish unique answers as we have defined, then all RAGFs are satisfied and vice-versa.

Other identities of the Clifford algebra can lead to traces different from the one we have started this argumentation, but it is provable they end up in linear combinations of the ones we presented. Furthermore, in the section (5.3), it is going to become clear why the ones we chose are enough to disentangle any feature of the anomalous odd tensor in even space-time dimensions. In particular as the traces we employed can be linearly combined it is possible to reach the following expression as a result to the amplitude

$$\left[T_{\mu_{123}}^{\Gamma_{123}}\right]_{\{12\}} = \frac{1}{2} \left[\left(T_{\mu_{123}}^{\Gamma_{123}}\right)_1 + \left(T_{\mu_{123}}^{\Gamma_{123}}\right)_2 \right],$$

and the combination 13 and 23, which violates the RAGF to the vertexes 12, 13, and 23 respectively if the surface term is any other than the one producing unique amplitudes. These three combinations in reality are enough to reproduce the computations using any substitution coded in the identities among the antisymmetrized products, see eq.(2.5), by example, using a common substitution like

$$\gamma_* \gamma_{\mu_i \nu_i \mu_{i+1}} = i \varepsilon_{\mu_i \nu_i \mu_{i+1} \nu} \gamma^{\nu} + \gamma_* \left(g_{\nu_i \mu_{i+1}} \gamma_{\mu_i} - g_{\mu_i \mu_{i+1}} \gamma_{\nu_i} + g_{\mu_i \nu_i} \gamma_{\mu_{i+1}} \right),$$

the difference between the integrated expression and the combination $[T_{\mu_{123}}^{\Gamma_{123}}]_{\{i,i+1\}}$ are finite and identically vanishing integrals, in other words, the application of the identity above or the linear combination have equal results when integrated, without touching their surface terms. This happens to any manipulations anywhere in the string of six gamma matrices and γ_* . This proposition is outlined in the appendix (A).

Another example, that illustrates the use of the definition of the chiral matrix as nothing special, is the use of

$$\gamma_*\gamma_{\mu_i} = \frac{1}{3!} \varepsilon_{\mu_i\nu_{123}} \gamma^{\nu_{123}}$$

in the level of the trace, of six matrices, it produces ten terms, and the five terms different from the use of the version *i* are finite integrals that happens to vanish, precisely the $T_{\mu_i}^{(-+)}$ in eq. (5.16), and T_{μ}^{ASS} amplitude, eq. (5.27). The importance of defining the version of some amplitude in the way we did, stems from the fact they satisfy automatically the maximum possible number of RAGFs, and we can build arbitrary linear combinations of the basic building-block versions, that reproduce all the ones deduced in the appendix (A), exploiting the substitutions encompassed in the formulas (2.5), and any application of Schouten identities, precisely

$$\left[T_{\mu_{123}}^{\Gamma_{123}}\right]_{\{r_1r_2r_3\}} = \frac{1}{r_1 + r_2 + r_3} \sum_{i=1}^3 r_i \left(T_{\mu_{123}}^{\Gamma_{123}}\right)_i, \tag{5.64}$$

with their $r_1 + r_2 + r_3 \neq 0$, its significance is they correspond to identical integrands, but if the surface term is forced to be zero, then all of them become distinct and violate the RAGFs by arbitrary amounts, and as a function of the external momenta same for the WIs. It helps to explain certain violation amounts found in the literature as in [49].

5.2. A Low Energy Theorem to the $T^{\Gamma_1\Gamma_2\Gamma_3}$ -Triangles and its relation with the Ward Identities

Hitherto, we have shown how the dynamic of the traces and the surface terms interfere with the linearity of the contraction of the amplitudes with the external momenta and the uniqueness of the perturbative expressions of the nontrivial odd tensors in this dimension. Nonetheless, a question remains to be answered. Could one anticipate such verified properties? And are they unavoidable? And they are, then why? To spot the two main reasons why this happens, we present a two-part analysis of the most general tensor that can represent these amplitudes.

The cause of these phenomena start with a relation that a general tensor of rank three, that is function of two variables, and odd parity is required to have. Adopting the momenta such as $q_1 = q_2 + q_3$, where the q_2 and q_3 are incoming in the vertexes Γ_2 and Γ_3 a q_1 outgoing the vertex Γ_1 , the general tensor will be expressed as

$$F_{\mu_{1}\mu_{2}\mu_{3}} = \varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu} \left(q_{2}^{\nu}F_{1} + q_{3}^{\nu}F_{2}\right) + \varepsilon_{\mu_{1}\mu_{2}\nu_{1}\nu_{2}}q_{2}^{\nu_{1}}q_{3}^{\nu_{2}} \left(q_{2\mu_{3}}G_{1} + q_{3\mu_{3}}G_{2}\right)$$

$$+ \varepsilon_{\mu_{1}\mu_{3}\nu_{1}\nu_{2}}q_{2}^{\nu_{1}}q_{3}^{\nu_{2}} \left(q_{2\mu_{2}}G_{3} + q_{3\mu_{2}}G_{4}\right) + \varepsilon_{\mu_{2}\mu_{3}\nu_{1}\nu_{2}}q_{2}^{\nu_{1}}q_{3}^{\nu_{2}} \left(q_{2\mu_{1}}G_{5} + q_{3\mu_{1}}G_{6}\right),$$

$$(5.65)$$

that in the three contractions

$$q_1^{\mu_1} F_{\mu_1 \mu_2 \mu_3} = \varepsilon_{\mu_2 \mu_3 \nu_1 \nu_2} q_2^{\nu_1} q_3^{\nu_2} V_1$$

$$q_2^{\mu_2} F_{\mu_1 \mu_2 \mu_3} = \varepsilon_{\mu_1 \mu_3 \nu_1 \nu_2} q_2^{\nu_1} q_3^{\nu_2} V_2$$

$$q_3^{\mu_3} F_{\mu_1 \mu_2 \mu_3} = \varepsilon_{\mu_1 \mu_2 \nu_1 \nu_2} q_2^{\nu_1} q_3^{\nu_2} V_3$$

can be identified three functions written exclusively as functions of the ingredients if the general tensor, they are

$$V_1 = -F_1 + F_2 + (q_1 \cdot q_2) G_5 + (q_1 \cdot q_3) G_6$$

$$V_2 = -F_2 + q_2^2 G_3 + (q_2 \cdot q_3) G_4$$

$$V_3 = -F_1 + (q_2 \cdot q_3) G_1 + q_3^2 G_2,$$

and thus, at the point $q_i \cdot q_j = 0$, if the form factors are regular at most discontinuous, we obtain

$$V_1(0) = F_2 - F_1;$$
 $V_2(0) = -F_2;$ $V_3(0) = -F_1,$

where the zero in our notation will always mean the point where all the bilinears are zero $q_i \cdot q_j = 0$, therefore from the equations above follows

$$V_1(0) = V_3(0) - V_2(0). (5.66)$$

These relations among the form factors have the information about the symmetry or violation thereof, at one kinematical point, even if no particular symmetry was need to such relation. Its satisfaction does not guarantee symmetries, that this tensor may have, in all the points, but its violation in this point, imply the violation of symmetry anyway. The crucial feature is that if the divergence of the axial current is connected to the pseudo-scalar density in a correlator with two other vector currents, as by example in AVV, we have

$$\varepsilon_{\mu_2\mu_3\nu_1\nu_2} q_2^{\nu_1} q_3^{\nu_2} V_1(0) = -2m T_{\mu_2\mu_3}^{PVV}(0) = \varepsilon_{\mu_2\mu_3\nu_1\nu_2} q_2^{\nu_1} q_3^{\nu_2} \Omega_1^{PVV}(0)$$

simply reworded as $V_1(0) = 1/(2\pi)^2$ the result presented in eq. (5.47), thereby at least one of the vector currents will be violated as the eq.(5.66) above does not allow for $V_2 = 0$ and $V_3 = 0$ simultaneously in this case. Nonetheless, if the vector currents are divergenceless $V_2 = V_3 = 0$, then the parameters defining the tensor under consideration, that means, rank, parity, number of variables, and regularity, will imply the existence of an additional term in $V_1 = 1/(2\pi)^2 + A$, the anomaly, present simply by the reason that the eq.(5.66) relating the V's requires it. In which case $\mathcal{A} = -\Omega_1^{PVV}(0)$, furnishing a connection between an unavoidable property of a finite amplitude and the symmetry content of another tensor, emerges.

This is the starting point of the violation dynamics of anomalous amplitudes. If the AWI is preserved, that means, $V_1(0) = \Omega_1^{PVV}(0) = (2\pi)^{-2}$, and if at the same time VWIs are preserved $V_2(0) = V_3(0) = 0$, this implies an obvious violation the linear-algebra type solution (5.66), and the immediate conclusion is that no tensor, whatever its origin, can be at the same time connected to the finite PVV (take this as a boundary condition) and have vanishing contraction with the momenta $q_2^{\mu_2}$ and $q_3^{\mu_3}$.

On the other hand, constructing AVV such that $V_1 = \Omega_1^{PVV}$ + Anomaly, for the function V_1 to has the necessary zero to be compatible with the VWIs. This shows that whenever the function of the axial-vertex contraction of AVV is connected, anomalously or not, to the PVV there will be an anomaly in at least one of the currents.

The particular ingredients of its perturbative expression fully corroborate with these assertions, the computations cast in this contribution is one more expression of these facts. However, we will show more, the RAGFs embodying linearity of integration furnish an exact connection among the ultraviolet and infrared features of the amplitudes, namely

$$\Omega_1^{PVV}(0) = 2i\Delta_{3\alpha}^{\alpha} \tag{5.67}$$

it is the condition required by linearity and equivalently by a unique expression as functions of the internal momenta, a condition that will be detailed in the next section.

As will be demonstrated later in section (5.3) assuming the form $V_i = \Omega_i + \mathcal{A}_i$, where the V's are the result of the contraction with the momenta corresponding to the *i*-th vertex and the Ω 's are the odd rank-two finite amplitudes form factors. Even to classically non-conserved vector currents and for an amplitude with three arbitrary masses running in the loop, the \mathcal{A} 's obey

$$\mathcal{A}_3 - \mathcal{A}_2 - \mathcal{A}_1 = (2\pi)^{-2} \tag{5.68}$$

due to

$$\Omega_3(0) - \Omega_2(0) - \Omega_1(0) = -(2\pi)^{-2}, \qquad (5.69)$$

always and independent of the number of axial vertexes (odd ones for sure), permutations thereof, and the masses and if the vector currents are conserved or not. Independent of the masses is not the claim the anomaly is independent of masses because that is known, for us, it means that even when the finite amplitudes exhibit intricate dependency on the masses the relation they have in the point zero is all the time incompatible with the ones required to a tensor with the characteristics that the rank-three triangles must have, eq. (5.66), even if no divergence arises in perturbation theory.

The last claim, implicit in the equation (5.68), only explicit the common result that when the vector currents are preserved the value of the anomaly is unique, but even more, whatever explicit tensor obtained via regularization, or not, violating by any quantity all the currents, it will always obey the equation to the \mathcal{A} 's above, that in the very end are determined by the combinations at zero of finite functions representing the rank-two odd amplitudes.

Another striking feature of the way we pose the matter of divergences is the fact that some restriction over the undetermined surface terms can be anticipated only based on the characteristics of the general form this tensor can have and its connection to the AV two-point functions via linearity of integration. They are the theme of the next section where, in the end, the reasoning leading the the equation (5.67) is unfolded. After that we end up showing the simple, but non-obvious path leading to the general equation (5.69)

5.3. The RAGFs and the Kinematical Behavior of Arbitrary Amplitudes

In the section where the explicit result to the versions of the amplitudes was shown to not automatically satisfy all their RAGFs, that means, a condition connecting the surface terms and the finite part must be stated at least in one of the contractions with the external momenta. This result can be established without resorting to any explicit computation, only assuming the most general form that the undetermined part of the amplitude can assume, and studying the constraints given by the contractions relating it to the differences of two-point AV functions.

To demonstrate such a proposition, let us lay down the most general tensor of mass dimension one, built out of the kinematical data, the arbitrary momenta, and surface terms. First of all the q_i vectors—the kinematical data—are written as differences of the routings k_i , **but not the opposite**. Thus we replace the former by the latter and use, in equivalent form, the combinations we have defined and used in this paper, $P_{ij} = k_i + k_j$ in place of k_i , due to the size of the expressions. Hence, the most general set of surface terms for the odd amplitudes is

$$\begin{aligned} F^{\Delta}_{\mu_{123}} &= +\varepsilon_{\mu_{2}\mu_{3}\nu_{1}\nu_{2}} \left(A_{11}P_{21}^{\nu_{2}} + A_{12}P_{31}^{\nu_{2}} + A_{13}P_{32}^{\nu_{2}}\right)\Delta^{\nu_{1}}_{3\mu_{1}} \\ &+\varepsilon_{\mu_{1}\mu_{3}\nu_{1}\nu_{2}} \left(A_{21}P_{21}^{\nu_{2}} + A_{22}P_{31}^{\nu_{2}} + A_{23}P_{32}^{\nu_{2}}\right)\Delta^{\nu_{1}}_{3\mu_{2}} \\ &+\varepsilon_{\mu_{1}\mu_{2}\nu_{1}\nu_{2}} \left(A_{31}P_{21}^{\nu_{2}} + A_{32}P_{31}^{\nu_{2}} + A_{33}P_{32}^{\nu_{2}}\right)\Delta^{\nu_{1}}_{3\mu_{3}} \\ &+\varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{1}} \left(B_{1}P_{21}^{\nu_{2}} + B_{2}P_{31}^{\nu_{2}} + B_{3}P_{32}^{\nu_{2}}\right)\Delta^{\nu_{1}}_{3\nu_{2}} \\ &+\varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{1}} \left(C_{1}P_{21}^{\nu_{1}} + C_{2}P_{31}^{\nu_{1}} + C_{3}P_{32}^{\nu_{2}}\right)\Delta^{\nu_{2}}_{3\nu_{2}}. \end{aligned}$$

After contraction, we expect this tensor to be related to the two-point AV tensors, by example, $q_2^{\mu_2}T_{\mu_{123}}^{AVV} = T_{\mu_{13}}^{AV}(1,3) - T_{\mu_{13}}^{AV}(2,3)$. Those amplitudes have their indexes in the surface term both in contracted mode and without a term like the last one, a trace of the surface term. Thus one more property of this general tensor will be used, it has an identity that reduce the number of linearly independent constants

$$\varepsilon_{\mu_1\mu_2\mu_3\nu_1}\Delta_{3\nu_2}^{\nu_2} = \varepsilon_{\mu_1\mu_2\mu_3\nu_2}\Delta_{3\nu_1}^{\nu_2} - \varepsilon_{\mu_1\mu_2\nu_1\nu_2}\Delta_{3\mu_3}^{\nu_2} - \varepsilon_{\mu_1\mu_3\nu_1\nu_2}\Delta_{3\mu_2}^{\nu_2} - \varepsilon_{\mu_2\mu_3\nu_1\nu_2}\Delta_{3\mu_1}^{\nu_2}$$

using it, the most general tensor of these variables under the conditions stated become

$$F^{\Delta}_{\mu_{123}} = +\varepsilon_{\mu_{2}\mu_{3}\nu_{1}\nu_{2}} (a_{11}P_{21} + a_{12}P_{31} + a_{13}P_{32})^{\nu_{2}} \Delta^{\nu_{1}}_{3\mu_{2}} +\varepsilon_{\mu_{1}\mu_{3}\nu_{1}\nu_{2}} (a_{21}P_{21} + a_{22}P_{31} + a_{23}P_{32})^{\nu_{2}} \Delta^{\nu_{1}}_{3\mu_{2}} +\varepsilon_{\mu_{1}\mu_{2}\nu_{1}\nu_{2}} (a_{31}P_{21} + a_{32}P_{31} + a_{33}P_{32})^{\nu_{2}} \Delta^{\nu_{1}}_{3\mu_{3}} +\varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{1}} (b_{1}P_{21} + b_{2}P_{31} + b_{3}P_{32})^{\nu_{2}} \Delta^{\nu_{1}}_{3\mu_{3}}$$

where the new constants are now given by $a_{ij} = A_{ij} - C_j$ and $b_i = B_i + C_i$ and the range of $i, j \in \{1, 2, 3\}$.

The a_{ij} and b_j are twelve arbitrary constants that embody all the freedom present by such tensor: Function of three variables, essentially the routings of the diagram, the rank, and parity of the tensor, the power counting and mass dimension equal one. The *j* captures the *P* momenta in

the order (P_{21}, P_{31}, P_{32}) , and the index *i* is clearly linked to the index μ_i that is turn is associated with the vertex that will appear in the amplitudes $T^{\Gamma_{123}}_{\mu_{123}}$.

Now expressing for convenience the three independent differences of two-point AV functions (5.51), (5.52) and (5.53) as

$$T_{1(-)\mu_{23}}^{AV} = -2i\varepsilon_{\mu_{23}\nu_{12}} \left[-P_{21}^{\nu_2} P_{32}^{\nu_3} + P_{31}^{\nu_2} \left(P_{32}^{\nu_3} - P_{21}^{\nu_3} \right) + P_{32}^{\nu_2} P_{21}^{\nu_3} \right] \Delta_{3\nu_3}^{\nu_1}$$
(5.70)

$$T_{2(-)\mu_{13}}^{AV} = -2i\varepsilon_{\mu_{13}\nu_{12}} \left[+P_{21}^{\nu_2} \left(P_{31}^{\nu_3} - P_{32}^{\nu_3} \right) + P_{31}^{\nu_2} P_{32}^{\nu_3} - P_{32}^{\nu_2} P_{31}^{\nu_3} \right] \Delta_{3\nu_3}^{\nu_1}$$
(5.71)

$$T_{3(-)\mu_{12}}^{AV} = -2i\varepsilon_{\mu_{12}\nu_{12}} \left[-P_{21}^{\nu_2}P_{31}^{\nu_3} + P_{31}^{\nu_2}P_{21}^{\nu_3} + P_{32}^{\nu_2} \left(P_{31}^{\nu_3} - P_{21}^{\nu_3} \right) \right] \Delta_{3\nu_3}^{\nu_1}, \tag{5.72}$$

where the notation $T_{i(-)}^{AV}$ means they came from the μ_i -th contraction with the corresponding momenta.

Let us lay down the impossibility of satisfying all these relations without further conditions, beginning with the most general tensor of surface terms $F^{\Delta}_{\mu_1\mu_2\mu_3}$. Contracting $F^{\Delta}_{\mu_1\mu_2\mu_3}$ with $q_1^{\mu_1} = p_{31}^{\mu_1} = P_{32}^{\mu_1} - P_{21}^{\mu_1}$

$$\begin{split} p_{31}^{\mu_1} F_{\mu_{123}}^{\Delta} &= + \varepsilon_{\mu_3 \nu_1 \nu_2 \nu_3} \left[- \left(a_{21} + a_{23} \right) P_{21}^{\nu_2} P_{32}^{\nu_3} + a_{22} P_{31}^{\nu_2} \left(P_{21}^{\nu_3} - P_{32}^{\nu_3} \right) \right] \Delta_{3\mu_3}^{\nu_1} \\ &+ \varepsilon_{\mu_2 \nu_1 \nu_2 \nu_3} \left[- \left(a_{31} + a_{33} \right) P_{21}^{\nu_2} P_{32}^{\nu_3} + a_{32} P_{31}^{\nu_1} \left(P_{21}^{\nu_3} - P_{32}^{\nu_3} \right) \right] \Delta_{3\mu_3}^{\nu_1} \\ &+ \varepsilon_{\mu_2 \mu_3 \nu_1 \nu_2} \left[- \left(a_{11} - b_1 \right) P_{21}^{\nu_2} P_{21}^{\nu_3} + \left(a_{13} - b_3 \right) P_{32}^{\nu_2} P_{32}^{\nu_3} \right] \Delta_{3\nu_3}^{\nu_1} \\ &+ \varepsilon_{\mu_2 \mu_3 \nu_1 \nu_2} \left[+ \left(a_{11} + b_3 \right) P_{21}^{\nu_2} P_{32}^{\nu_3} - a_{12} P_{31}^{\nu_2} \left(P_{21}^{\nu_3} - P_{32}^{\nu_3} \right) \right] \Delta_{3\nu_3}^{\nu_1} \\ &+ \varepsilon_{\mu_2 \mu_3 \nu_1 \nu_2} \left[- \left(a_{13} + b_1 \right) P_{32}^{\nu_2} P_{21}^{\nu_3} + b_2 \left(P_{21}^{\nu_2} - P_{32}^{\nu_2} \right) P_{31}^{\nu_3} \right] \Delta_{3\nu_3}^{\nu_1} \end{split}$$

and from the first two rows we obtain $a_2 = (-a_{23}, 0, a_{23})$ and $a_3 = (-a_{33}, 0, a_{33})$, the remaining must be compared with

$$T_{1(-)\mu_{32}}^{AV} = 2i\varepsilon_{\mu_{23}\nu_{12}} \left(P_{21}^{\nu_2} P_{32}^{\nu_3} + P_{31}^{\nu_2} \left(P_{21}^{\nu_3} - P_{32}^{\nu_3} \right) - P_{32}^{\nu_2} P_{21}^{\nu_3} \right) \Delta_{3\nu_3}^{\nu_1}$$

giving

$$a_{11} + b_3 = 2i$$
, $a_{12} = -2i$, $a_{13} + b_1 = 2i$, $b_2 = 0$ $b_3 = 2i - b_1$

which rephrased in vector notation the full solution is

$$\begin{pmatrix} \mathbf{b} \\ \mathbf{a}_1 \\ \mathbf{a}_2 \\ \mathbf{a}_3 \end{pmatrix} = \begin{pmatrix} b_1 & 0 & 2i - b_1 \\ b_1 & -2i & 2i - b_1 \\ -a_{23} & 0 & a_{23} \\ -a_{33} & 0 & a_{33} \end{pmatrix}.$$
 (5.73)

First thing to note is the reduction from twelve parameters to just three, here the constants $\{a_{23}, a_{33}, b_1\}$, by requiring just one of the relations to be satisfied.

If it is asked for any other relation to be satisfied, the solution will be unique and it will have consequences over the last one. Repeating the analysis to the contraction of $F^{\Delta}_{\mu_1\mu_2\mu_3}$ with $q_2^{\mu_2} = p_{21}^{\mu_2} = P_{32}^{\mu_2} - P_{31}^{\mu_2}$, forming the system of linear equation by comparing $q_2^{\mu_2}F^{\Delta}_{\mu_{123}}$ with $T^{AV}_{2(-)\mu_{13}}$ follows the complete solution for the automatic satisfaction of the RAGF born out of the contraction with $q_2^{\mu_2}$

$$\begin{pmatrix} \mathbf{b} \\ \mathbf{a}_1 \\ \mathbf{a}_2 \\ \mathbf{a}_3 \end{pmatrix} = \begin{pmatrix} 0 & b_2 & 2i - b_2 \\ 0 & -a_{13} & a_{13} \\ 2i & -b_2 & b_2 - 2i \\ 0 & -a_{33} & a_{33} \end{pmatrix}.$$
 (5.74)

The contraction with $q_3^{\mu_3} = p_{32}^{\mu_3} = P_{31}^{\mu_3} - P_{21}^{\mu_3}$, allows us to determine the conditions for $q_3^{\mu_3}F_{\mu_{123}}^{\Delta} = T_{3(-)\mu_{12}}^{AV}$, and then emerges the solution to the automatic satisfaction of the RAGF, product of the contraction with the index and momenta of the third vertex

$$\begin{pmatrix} \mathbf{b} \\ \mathbf{a}_1 \\ \mathbf{a}_2 \\ \mathbf{a}_3 \end{pmatrix} = \begin{pmatrix} b_1 & 2i - b_1 & 0 \\ a_{11} & -a_{11} & 0 \\ a_{21} & -a_{21} & 0 \\ b_1 & 2i - b_1 & -2i \end{pmatrix}.$$
 (5.75)

The intersection of the solution (5.73) and (5.74), that means, the ones that satisfies automatically their RAGFs coming from the contraction with $q_1^{\mu_1}$ and $q_2^{\mu_2}$, leads to a unique solution with $b_1 = 0, b_2 = 0, b_3 = 2i$ and all the other coefficients determined. Putting all that values in the tensor, we get

$$\left(F_{\mu_{123}}^{\Delta}\right)_{12} = -2i \left[\varepsilon_{\mu_{2}\mu_{3}\nu_{1}\nu_{2}} \left(P_{32}^{\nu_{2}} - P_{31}^{\nu_{2}}\right) \Delta_{3\mu_{1}}^{\nu_{1}} + \varepsilon_{\mu_{1}\mu_{3}\nu_{1}\nu_{2}} \left(P_{21}^{\nu_{2}} - P_{32}^{\nu_{2}}\right) \Delta_{3\mu_{2}}^{\nu_{1}} + \varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{1}} P_{32}^{\nu_{2}} \Delta_{3\nu_{2}}^{\nu_{1}}\right],$$

where sub-index ij in $(F^{\Delta}_{\mu_{123}})_{ij}$ stands for the vertexes where the RAGFs are satisfied without further assumptions.

Each of the solutions presented depends on three parameters and are compatible with one another. However, once the coefficients are determined to the unique and unrestricted satisfaction of two RAGFs, the third solution will always exhibit an additional term, in other words

$$(F^{\Delta}_{\mu_{123}})_{23} \neq (F^{\Delta}_{\mu_{123}})_{13} \neq (F^{\Delta}_{\mu_{123}})_{12}$$

It is an easy task to verify that the solution obtained requiring at most two RAGF satisfied corresponds to versions of the traces computed explicitly, see (5.37),

$$(F^{\Delta}_{\mu_{123}})_{23} = S_{1\mu_{123}}; \quad (F^{\Delta}_{\mu_{123}})_{13} = S_{2\mu_{123}}; \quad (F^{\Delta}_{\mu_{123}})_{12} = S_{3\mu_{123}};$$

Owing to this behavior, whose hypotheses can be verified in the explicit computation, if desired, it can be drawn a parallel conclusion to the one stated in the section (5.2) where the value at zero of PVV has consequences over the symmetries, by example. Here this finite amplitude will establish a connection between the linearity embodied in the RAGFs and the low energy behavior of the same finite PVV, amplitude taken as example.

For such end, we have got to read this result in the light of the general form factors in the eq. (5.65). At this point, we will take the form factors as representing only the strictly finite part of any tensor under consideration.

Choosing the solution satisfying the RAGFs in the vertexes two and three

$$T_{\mu_{123}}^{\Gamma_{123}} = F_{\mu_{123}} + S_{1\mu_{123}}$$

to any combination that the vertexes might assume, once again due to the fact the two-point functions depend only on the vertex where we effect the contraction. Contracting with the momenta of the respective vertexes, we obtain

$$\begin{split} q_1^{\mu_1} T_{\mu_{123}}^{\Gamma_{123}} &= T_{1(-)\mu_{23}}^{AV} + \varepsilon_{\mu_2\mu_3\nu_1\nu_2} q_2^{\nu_1} q_3^{\nu_2} \left(V_1 + 2i\Delta_{3\alpha}^{\alpha}\right) \\ q_2^{\mu_2} T_{\mu_{123}}^{\Gamma_{123}} &= T_{2(-)\mu_{13}}^{AV} + \varepsilon_{\mu_1\mu_3\nu_1\nu_2} q_2^{\nu_1} q_3^{\nu_2} V_2 \\ q_3^{\mu_3} T_{\mu_{123}}^{\Gamma_{123}} &= T_{3(-)\mu_{12}}^{AV} + \varepsilon_{\mu_1\mu_2\nu_1\nu_2} q_2^{\nu_1} q_3^{\nu_2} V_3, \end{split}$$

remember here the V_i contains only dependence on the finite parts. The trace of the surface term must be put together with the finite part of the first contraction due to the equation obtained in (5.58)

$$q_1^{\mu_1} S_{1\mu_{123}} = T_{1(-)\mu_{23}}^{AV} + \varepsilon_{\mu_2\mu_3\nu_2\nu_3} q_2^{\nu_2} q_3^{\nu_3} \left(2i\Delta_{3\nu_1}^{\nu_1}\right).$$

Take an arbitrary sorting of vertexes that yields an odd tensor and let the Ω_i represent the scalar invariant that corresponds to the rank-two tensor obtained in the RAGF, which are finite functions, and write the equations that signify the hypothesis of satisfaction of the RAGF

$$\begin{split} q_1^{\mu_1} T_{\mu_{123}}^{\Gamma_{123}} &= T_{1(-)\mu_{23}}^{AV} + \varepsilon_{\mu_2\mu_3\nu_1\nu_2} q_2^{\nu_1} q_3^{\nu_2} \Omega_1 \\ q_2^{\mu_2} T_{\mu_{123}}^{\Gamma_{123}} &= T_{2(-)\mu_{13}}^{AV} + \varepsilon_{\mu_1\mu_3\nu_1\nu_2} q_2^{\nu_1} q_3^{\nu_2} \Omega_2 \\ q_3^{\mu_3} T_{\mu_{123}}^{\Gamma_{123}} &= T_{3(-)\mu_{12}}^{AV} + \varepsilon_{\mu_1\mu_2\nu_1\nu_2} q_2^{\nu_1} q_3^{\nu_2} \Omega_3, \end{split}$$

then look at the deduced behavior above where it is admitted that relation two and three are unrestricted satisfied, from there we get the final condition to our relations, namely

$$V_1 + 2i\Delta_{3\alpha}^{\alpha} = \Omega_1; \quad V_2 = \Omega_2; \quad V_3 = \Omega_3,$$

remain to observe the formulas

$$V_{1} = -F_{1} + F_{2} + q_{2}^{2}G_{5} + q_{3}^{2}G_{6} + (q_{2} \cdot q_{3})(G_{5} + G_{6})$$

$$V_{2} = -F_{2} + q_{2}^{2}G_{3} + (q_{2} \cdot q_{3})G_{4}$$

$$V_{3} = -F_{1} + (q_{2} \cdot q_{3})G_{1} + q_{3}^{2}G_{2},$$

and eliminate the F_i form factor to reach at

$$-q_2^2 (G_3 + G_5) + q_3^2 (G_2 - G_6) + (q_2 \cdot q_3) (G_1 - G_4 - G_5 - G_6)$$

= $2i\Delta_{3\alpha}^{\alpha} + \Omega_3 - \Omega_2 - \Omega_1$

now under the condition, that the G_i functions are regular at zero⁷, follows the master equation

$$-2i\Delta_{3\alpha}^{\alpha} = \Omega_3(0) - \Omega_2(0) - \Omega_1(0), \qquad (5.76)$$

and this trivially will happen irrespective of what set of RAGF is chosen to be satisfied without restriction, suppose one starts with a version with $S_{2\mu_{123}}$ that satisfies the RAGFs in the first and third vertex, to this tensor the term $\Delta_{3\alpha}^{\alpha}$ will appear in

$$q_2^{\mu_2} S_{2\mu_{123}} = T_{2(-)\mu_{13}}^{AV} + \varepsilon_{\mu_1\mu_3\nu_1\nu_2} q_2^{\nu_1} q_3^{\nu_2} \left(2i\Delta_{3\alpha}^{\alpha}\right),$$

a relation established in eq. (5.60), then from $V_1 = \Omega_1$ and $V_3 = \Omega_3$ because the RAGFs are unrestricted by hypothesis, we exchange the F_1 and F_2 by G_i multiplied by bilinears plus finite functions and then again in the point $q_i \cdot q_j = 0$ we retrieve the previous result.

Here is important to be aware of the fact that only hypotheses were employed, a tensor that has two RAGFs satisfied without restriction, that means connected to known differences of AV functions and PVV/PAA-like amplitudes without further ado, and from that point on the third

⁷ The functions $Z_{nm}^{(0)}, Z_{nm}^{(-1)}, Z_n^{(0)}$ that comprise the finite part of any of these amplitudes do not have kinematical singularities at the point $q_i \cdot q_j = 0$.

is necessarily bounded by the zero value of the finite rank-two amplitude. In any case, this requirements are always possible to attain in the explicit computations we have presented and shown how.

When assessing the values of $\Omega_i(0)$, see the results (5.47), $\Omega^{PVV} = \Omega^{VPV} = -\Omega^{VVP} = (2\pi)^{-2}$, we find out

$$\Omega_3(0) - \Omega_2(0) - \Omega_1(0) = -(2\pi)^{-2},$$

notice that for the AVV, VAV, and VVA where only one axial current appears, two of the Ω_i are zero to each amplitude or more precisely the result above represents in reality three situations. The same happens to the AAA triangle, on the other hand in the case the three contractions of the same amplitude are related to $T_{\mu_{23}}^{PAA}$, $T_{\mu_{13}}^{APA}$, $T_{\mu_{12}}^{AAP}$, thereby it only a matter of combining the constants cast in eq. (5.48), $\Omega^{PAA}(0) = \Omega^{APA}(0) = -\Omega^{AAP}(0) = \frac{1}{3(2\pi)^2}$, in other words

$$\Omega^{AAP}(0) - \Omega^{APA}(0) - \Omega^{PAA}(0) = -(2\pi)^{-2}$$

it shows that because the differences of AV structures depend only on the contraction with the momenta, eq.'s(5.51, 5.52 and 5.53), they are common to all cases analyzed, but as the correlators with the P density are finite but distinct, it could be (very unlikely) that distinct diagrams would require different numerical values to the surface term, notwithstanding what we get is

$$\text{RAGF} \Leftrightarrow \Delta_{3\alpha}^{\alpha} = -\frac{i}{2\left(2\pi\right)^2}$$

always, and to amplitudes where three distinct masses run in the internal lines, in that case, the vector currents are not classically conserved, the constrain remains the same.

To see this, let us consider, as an example, that three arbitrary masses run in the inner lines of the amplitude AVV, where the indexes of the propagators now account for the masses too, $S(a) = (\not{k}_a - m_a)^{-1}$, and then the standard identity $\not{p}_{ij} = S^{-1}(i) - S^{-1}(j) + (m_i - m_j)$ is applied to derive the relations

$$p_{31}^{\mu_1} t_{\mu_{123}}^{AVV} = t_{\mu_{32}}^{AV} (1,2) - t_{\mu_{23}}^{AV} (2,3) - (m_1 + m_3) t_{\mu_{23}}^{PVV} p_{21}^{\mu_2} t_{\mu_{123}}^{AVV} = t_{\mu_{13}}^{AV} (1,3) - t_{\mu_{13}}^{AV} (2,3) + (m_2 - m_1) t_{\mu_{13}}^{ASV} p_{32}^{\mu_3} t_{\mu_{123}}^{AVV} = t_{\mu_{12}}^{AV} (1,2) - t_{\mu_{12}}^{AV} (1,3) + (m_3 - m_2) t_{\mu_{12}}^{AVS},$$

now the vector currents aren't conserved, however the ASV, AVS, and PVV of arbitrary masses will not comply with the eq. (5.66), repeated here for clarity of the arguments, $V_3(0) - V_2(0) - V_1(0) = 0$.

Taking the integrals of the three-point rank-two finite amplitudes, we get

$$\begin{split} T^{PVV}_{\mu_{23}} &= \varepsilon_{\mu_{23}\nu_{12}} p^{\nu_1}_{21} p^{\nu_2}_{32} \left[(m_1 - m_2) \, Z^{(-1)}_{10} + (m_1 - m_3) \, Z^{(-1)}_{01} - m_1 Z^{(-1)}_{00} \right] \\ T^{ASV}_{\mu_{13}} &= \varepsilon_{\mu_{13}\nu_{12}} p^{\nu_1}_{21} p^{\nu_2}_{32} \left[(m_1 + m_2) \, Z^{(-1)}_{10} + (m_1 + m_3) \, Z^{(-1)}_{01} - m_1 Z^{(-1)}_{00} \right] \\ T^{AVS}_{\mu_{12}} &= \varepsilon_{\mu_{12}\nu_{12}} p^{\nu_1}_{21} p^{\nu_2}_{32} \left[(m_2 - m_1) \, Z^{(-1)}_{10} - (m_3 + m_1) \, Z^{(-1)}_{01} + m_1 Z^{(-1)}_{00} \right], \end{split}$$

identifying the form factor through

$$\begin{split} \varepsilon_{\mu_{23}\nu_{12}} p_{21}^{\nu_1} p_{32}^{\nu_2} \Omega_1^{PVV} &= -(m_1 + m_3) \, T_{\mu_{23}}^{PVV} \\ \varepsilon_{\mu_{13}\nu_{12}} p_{21}^{\nu_1} p_{32}^{\nu_2} \Omega_2^{ASV} &= +(m_2 - m_1) \, T_{\mu_{13}}^{ASV} \\ \varepsilon_{\mu_{12}\nu_{12}} p_{21}^{\nu_1} p_{32}^{\nu_2} \Omega_3^{AVS} &= +(m_3 - m_2) \, T_{\mu_{12}}^{AVS}, \end{split}$$

and combining them as dictated by eq. (5.66), we have

$$-\left(\Omega_3^{AVS} - \Omega_2^{ASV} - \Omega_1^{PVV}\right) = \frac{1}{2\pi^2} \left[\left(m_1^2 - m_2^2\right) Z_{10}^{(-1)} + \left(m_1^2 - m_3^2\right) Z_{01}^{(-1)} - m_1^2 Z_{00}^{(-1)} \right]$$
(5.77)

in the kinematical limit $q_i \cdot q_j = 0$, follows

$$\left(\Omega_3^{AVS} - \Omega_2^{ASV} - \Omega_1^{PVV}\right)(0) = -(2\pi)^{-2}, \qquad (5.78)$$

since in the definition for distinct masses we have⁸ $Q(0) = (m_1^2 - m_2^2) x_1 + (m_1^2 - m_3^2) x_2 - m_1^2$, hence

$$\left[\left(m_1^2 - m_2^2 \right) Z_{10}^{(-1)} + \left(m_1^2 - m_3^2 \right) Z_{01}^{(-1)} - m_1^2 Z_{00}^{(-1)} \right]_{q_i \cdot q_j = 0} = 1/2.$$

the integrals with distinct masses are more laborious, but this result can be obtained by integrating until the end all these functions in the limit under consideration.

What this says is that the kinematical limits of all rank-two odd and finite amplitudes are not compatible with the satisfaction of all Ward identities as already noted in the 2D-AV amplitude. This is independent of divergences or the particular aspects of perturbative solutions⁹, even if a solution coming from a divergent-free formalism were provided, but their contractions would have to be related to the finite tensors we have described, there should be an anomaly.

5.4. Parameter Landscape of the Violations¹⁰ and Commentary

From the consequences derived in the last section, we see that if the value attributed to our tensors is done in a form that obeys the linearity of integration, then the surface term present in their perturbative expressions must be non-zero. Since its coefficients are not combinations of only physical momenta, therefore choices must be made. As the choice dictates the symmetry and vice-versa, we could be guided by them.

Talking about choices, by example, the eventual role of a regularization argument in our strategy could be summarized schematically as: Consider a regularization obeying $\lim_{\Lambda\to\infty} G(\Lambda_i, k^2) = 1$, thus it yields tensors that are immune to the Dirac traces,

$$\lim_{\Lambda \to \infty} \int \frac{\mathrm{d}^4 k}{\left(2\pi\right)^4} G\left(\Lambda_i, k^2\right) \left(t_{\mu_{123}}^{\Gamma_{123}}\right)_r = \lim_{\Lambda \to \infty} \int \frac{\mathrm{d}^4 k}{\left(2\pi\right)^4} G\left(\Lambda_i, k^2\right) \left(t_{\mu_{123}}^{\Gamma_{123}}\right)_s,$$

if and only if $\Delta_{3\mu\nu} = -ig_{\mu\nu}/2 (4\pi)^2$.

But, if in the process we get $\Delta_{3\mu\nu} = 0$, then the linearity is violated by $q_i^{\mu_i} \left[T_{\mu_{123}}^{\Gamma_{123}} \right]_i^{\text{viol}} = \pm (2\pi)^{-2} \varepsilon_{\mu_k \mu_l \nu_1 \nu_2} q_2^{\nu_1} q_3^{\nu_2}$, see these results in the eq.'s (5.61), the sign is negative for i = 1, 2 and positive for i = 3, plus we arrange the terms with k < l. What comes next is independent of this type of arguments, one example of rules to achieve this is saw in [50].

$$Q = q_1^2 x_1 \left(1 - x_1\right) + q_2^2 x_2 \left(1 - x_2\right) - 2q_1 \cdot q_2 x_1 x_2 + \left(m_1^2 - m_2^2\right) x_1 + \left(m_1^2 - m_3^2\right) x_2 - m_1^2$$

and

$$Z_{rs}^{(-1)} = \int_0^1 \mathrm{d}x_1 \int_0^{1-x_1} \mathrm{d}x_2 \frac{x_1' x_2'}{Q(q_i^2, m_1^2, m_2^2, m_3^2)}$$

⁸ To arbitrary masses, the Feynman polynomial for the function involved in this derivation, reads

⁹ With the exception these finite functions are perturbative solutions

¹⁰ Throughout this section we factored out the three-point rank-two finite amplitudes from the discussion.

Therefore, combining arbitrarily, and before integration, the versions that saves the maximum number of RAGFs, observe that is clear in this point this claim can be stated independent of Dirac traces computation, we have

$$\begin{bmatrix} T_{\mu_{123}}^{\Gamma_{123}} \end{bmatrix}_{\{r_1 r_2 r_3\}} = \frac{1}{R} \left\{ r_1 \begin{bmatrix} T_{\mu_{123}}^{\Gamma_{123}} \end{bmatrix}_1 + r_2 \begin{bmatrix} T_{\mu_{123}}^{\Gamma_{123}} \end{bmatrix}_2 + r_3 \begin{bmatrix} T_{\mu_{123}}^{\Gamma_{123}} \end{bmatrix}_3 \right\}$$

$$R = r_1 + r_2 + r_3 \neq 0,$$

$$(5.79)$$

and as, the derivations about uniqueness in the end of section (5.1) showed, they are identically equal before integration, but when $\Delta_{3\mu\nu} = 0$, they become an infinity set of different tensors, obtained by the same rules from the same integrand. For zero surface term, their symmetry violations are

$$q_1^{\mu_1} \left[T_{\mu_{123}}^{\Gamma_{123}} \right]_{\{r_1 r_2 r_3\}} \sim -\frac{r_1}{R} \frac{1}{(2\pi)^2}; \quad q_2^{\mu_2} \left[T_{\mu_{123}}^{\Gamma_{123}} \right]_{\{r_1 r_2 r_3\}} \sim -\frac{r_2}{R} \frac{1}{(2\pi)^2}; \quad q_3^{\mu_3} \left[T_{\mu_{123}}^{\Gamma_{123}} \right]_{\{r_1 r_2 r_3\}} \sim \frac{r_3}{R} \frac{1}{(2\pi)^2};$$

and satisfies the equation determined to its anomalies, $\mathcal{A}_3 - \mathcal{A}_2 - \mathcal{A}_1 = (2\pi)^{-2}$, since the basic versions satisfies it.

If the surface term is considered an arbitrary parameter, any result and why they are such ones derives systematically from our strategy. Take the surface term determined by an arbitrary constant c_1 , see below, that is equal one for the satisfaction of RAGFs, or zero for the momentumspace translational invariance. Parametrize the internal lines chosing any of the sums $P_{ij} = k_i + k_j$, since any other is determined by the differences equal to the external momenta,

$$P_{31} = c_2 q_2 + c_3 q_3 \to P_{21} = c_2 q_2 + (c_3 - 1) q_3; \quad P_{32} = (c_2 + 1) q_2 + c_3 q_3,$$

and $\Delta_{3\mu_1\mu_2} = c_1 \left[-i/2 \left(4\pi\right)^2 \right] g_{\mu_1\mu_2}.$ (5.80)

the AV structures, see the derivations of section (5.1) and eq'.s (5.51, 5.52, and 5.53), assume the value

$$T_{1(-)\mu_{23}}^{AV} = \frac{c_1 \left(c_3 - c_2 - 2\right)}{4 \left(2\pi\right)^2}; \quad T_{2(-)\mu_{13}}^{AV} = -\frac{c_1 \left(c_3 + 1\right)}{4 \left(2\pi\right)^2}; \quad T_{3(-)\mu_{12}}^{AV} = -\frac{c_1 \left(c_2 - 1\right)}{4 \left(2\pi\right)^2}, \tag{5.81}$$

and the basic violations of the RAGFs, eq'.s (5.61), turn into

$$q_{1}^{\mu_{1}} \left[T_{\mu_{123}}^{\Gamma_{123}} \right]_{1}^{\text{ragf}} = \frac{(c_{1}-1)}{(2\pi)^{2}}; \quad q_{2}^{\mu_{2}} \left[T_{\mu_{123}}^{\Gamma_{123}} \right]_{2}^{\text{ragf}} = \frac{(c_{1}-1)}{(2\pi)^{2}}; \quad q_{3}^{\mu_{3}} \left[T_{\mu_{123}}^{\Gamma_{123}} \right]_{3}^{\text{ragf}} = -\frac{(c_{1}-1)}{(2\pi)^{2}}, \quad (5.82)$$

all the terms are multiplied by the adequate tensor, $\varepsilon_{\mu_i \mu_j \nu_1 \nu_2} q_2^{\nu_1} q_3^{\nu_2}$ with i < j.

Therefore, this is enough to get the total set of possible values to the contractions of the expression of the basic versions, with the caveat that only the contraction where both the AV's and the term that break of linearity appear are the *i*-th version with $q_i^{\mu_i}$, to the *i*-th version a contraction with q_j , $j \neq i$, only the AV's contributes. The arbitrary linear combination of the versions will assume the form

$$q_{1}^{\mu_{1}} \left[T_{\mu_{123}}^{\Gamma_{123}} \right]_{\{r_{1}r_{2}r_{3}\}} = \frac{1}{(2\pi)^{2}} \left\{ \frac{r_{1} \left(c_{1} - 1 \right)}{R} + \frac{1}{4} c_{1} \left(c_{3} - c_{2} - 2 \right) \right\}$$

$$q_{2}^{\mu_{2}} \left[T_{\mu_{123}}^{\Gamma_{123}} \right]_{\{r_{1}r_{2}r_{3}\}} = \frac{1}{(2\pi)^{2}} \left\{ \frac{r_{2} \left(c_{1} - 1 \right)}{R} - \frac{1}{4} c_{1} \left(c_{3} + 1 \right) \right\}$$

$$q_{3}^{\mu_{3}} \left[T_{\mu_{123}}^{\Gamma_{123}} \right]_{\{r_{1}r_{2}r_{3}\}} = \frac{1}{(2\pi)^{2}} \left\{ -\frac{r_{3} \left(c_{1} - 1 \right)}{R} - \frac{1}{4} c_{1} \left(c_{2} - 1 \right) \right\},$$
(5.83)

where in the first terms we have the contributions associated to the unique perspective, as we posed, as the second ones representing routing choice. Showing that the number of independent parameter is two

$$A = -\frac{1}{4}c_1(c_3+1) + \frac{r_2}{R}(c_1-1)$$
(5.84)

$$B = -\frac{1}{4}c_1(c_2 - 1) - \frac{r_3}{R}(c_1 - 1)$$
(5.85)

because the first line is simply

$$B - A - 4 = \frac{1}{4}c_1\left(c_3 - c_2 - 2\right) + \frac{r_1}{R}\left(c_1 - 1\right), \qquad (5.86)$$

since, as we deduced in the former sections, when two violations are given, no matter the path leading to them, a third is determined without ambiguity of any nature.

If $c_1 = 1$, there is no dependence in r_i , we have the unique solution that satisfies linearity but is momenta ambiguous. However, if $c_1 = 0$ will be no dependence in c_2 and c_3 , that account for the internal momenta ambiguities. This is the full range of possibilities and they comply with the constraint determined only by utilizing finite integrals, eq. (5.68) for the eventual arbitrary anomalies. The crossed diagrams work in the same form, and just add more parameters to the discussion, the break of linearity, ambiguities, and unavoidable violation of symmetries in the way we have described is independent of this feature.

The divergent character of the perturbative amplitudes is not the determining factor of the anomaly phenomena, but yes, it plays a role. The crucial factor is the kinematical behavior of finite, and as such independent of interpretation, functions that code the amplitudes for the pseudo-scalar density. Not demonstrated here is that this aspect gets mutated, in the massless limit, in the values to the residue of poles of the form factors, form factors that are regular in the massive case.

Breaking linearity has a function in the divergent amplitudes that corroborates, and should be, with the finite PV^n amplitude in dimension d = 2n. As implicitly said, it rises from the situation where integrating zero we obtain a non-zero result. To begin with, the unique surface-term value will guarantee the following identity

$$(\varepsilon_{\mu_{5123}}K_1^{\mu_5}K_{1\mu_4} + \varepsilon_{\mu_{4512}}K_1^{\mu_5}K_{1\mu_3} + \varepsilon_{\mu_{3451}}K_1^{\mu_5}K_{1\mu_2} + \varepsilon_{\mu_{2345}}K_1^{\mu_5}K_{1\mu_1})\frac{1}{D_{123}} + \varepsilon_{\mu_{1234}}\frac{1}{D_{23}} + \varepsilon_{\mu_{1234}}\frac{m^2}{D_{123}} = 0,$$
(5.87)

in four dimensions. That formula comes from the fact that $\varepsilon_{[\mu_{1234}}K_{1\mu_5]} = 0$, that was multiplied by $K_1^{\mu_5}/D_{123}$, and used $K_1^2 = D_1 + m^2$, it is a Schouten identity to the integrand of bare Feynman integral $\bar{J}_{3\mu\nu}$. The critical step arises when we separate, without commitment with a particular interpretation to the divergences, the finite and divergent parts,

$$\begin{split} \bar{J}_2 \left(2,3 \right) \;&=\; J_2 \left(p_{32} \right) + I_{\log} \\ J_3 \left(1,2,3 \right) \;&=\; J_3 \left(p_{21},p_{31} \right) \\ \bar{J}_{3\mu\nu} \left(1,2,3 \right) \;&=\; J_{3\mu\nu} \left(p_{21},p_{31} \right) + \left(\Delta_{3\mu\nu} + g_{\mu\nu} I_{\log} \right) /4 \end{split}$$

— ,

and recombine them by means of $\varepsilon_{[\mu_{1235}}\Delta^{\mu_5}_{3\mu_5]} = 0$ and $\varepsilon_{[\mu_{1235}}J^{\mu_5}_{3\mu_5]} = 0$. Thus, although the identity for the surface term is consistent to any value, constrained only by $\Delta_{3\mu\nu} = \left[g^{\alpha\beta}\Delta_{3\alpha\beta}\right]/4$, the same is not true to the bare integral $J_{3\mu\nu}$, after followed the steps indicated, the linearity-breaking phenomena arrive,

Eq. for
$$\bar{J}_{3\mu\nu}$$
 above $\rightarrow -\frac{1}{4}\varepsilon_{\mu_{1234}} \left[\Delta_{3\mu_5}^{\mu_5} + 2i/(4\pi)^2 \right] = 0,$ (5.88)

in other words, as a part of the Feynman integrals the mere satisfaction of the Schouten identity, which implies $\Delta_{3\mu\nu} = \left[g^{\alpha\beta}\Delta_{3\alpha\beta}\right]/4$, is not enough to make it valid when the full integrals are brought into light. We have used, see eq. (C2) in the appendix easily obtained with the formulas present there, the result

$$g^{\alpha\beta}J_{3\alpha\beta} = m^2 J_3 + J_2(p_{32}) + \frac{i}{2(4\pi)^2},$$
(5.89)

hence, the identity is respected only and only if

$$\Delta_{3\mu_5}^{\mu_5} = -2i/\left(4\pi\right)^2,\tag{5.90}$$

derived without ever manipulating a divergent integral.

We want to mention here the mere violation by an evanescent term as occur in dimensional methods¹¹, does not spoils the paradigm of linearity breaking, the finite value we are demonstrating to be necessary is not a function of dimension and in truth it corresponds to the low-energy limit of the integral J_3 , it can not be nullifyed in any limit and if not adopted violates the linearity and uniqueness of the perturbative amplitudes.

There is no more space to show the clear effect of what we claimed here, but as a title of comment, when one tries to establish the RAGF in the explicit traces, take by example the first version, before integration, for the contractions p_{21} and p_{32} no question arises, it is as if no trace need to be explicited at all, the startling thing is that in the contraction with p_{31} it is possible to organize the integrand in such a way the RAGF follows from the identity that we started this final discussion. Thereby, the condition to the validity of RAGFs is the validity, in general, of the vanishing of a total antisymmetric tensor of rank five or bigger, independent of its nature.

6. SIX DIMENSIONAL BOX AMPLITUDE AVVV

To complete our line of reasoning, let us make an exercise of our notation in two versions of the AVVV box. The sub-amplitudes for this tensor in six dimensions require vertexes not present in the model, the tensor and pseudo-tensor vertexes

$$T = \gamma_{[\alpha_{12}]} = \frac{1}{2} [\gamma_{\alpha_1}, \gamma_{\alpha_2}]; \text{ and } \tilde{T} = \gamma_* \gamma_{[\alpha_{12}]}.$$

Making the choices $\Gamma_1 = \gamma_* \gamma_{\mu_1}$ and $\Gamma_i = \gamma_{\mu_i}$ for i = 2, 3, 4, and keeping the non-zero terms we have

$$t_{\mu_{1234}}^{AVVV} = K_{1234}^{\nu_{1234}} \operatorname{tr}(\gamma_{*\mu_{1}\nu_{1}\mu_{2}\nu_{2}\mu_{3}\nu_{3}\mu_{4}\nu_{4}}) \frac{1}{D_{1234}} + m^{2} \operatorname{tr}(\gamma_{*\mu_{1}\mu_{2}\mu_{3}\mu_{4}\nu_{1}\nu_{2}}) \left(-K_{12}^{\nu_{12}} + K_{13}^{\nu_{12}} - K_{14}^{\nu_{12}} - K_{23}^{\nu_{12}} + K_{24}^{\nu_{12}} - K_{34}^{\nu_{12}}\right) \frac{1}{D_{1234}}.$$

¹¹ See [51][52] for this type of view in traditional approaches.

This tensor satisfies a class of relations, its RAGF, through the standard procedure outline in section (2)

$$\begin{split} p_{21}^{\mu_2} t_{\mu_{1234}}^{AVVV} &= t_{\mu_{134}}^{AVV} \left(1,3,4\right) - t_{\mu_{134}}^{AVV} \left(2,3,4\right) \\ p_{32}^{\mu_3} t_{\mu_{1234}}^{AVVV} &= t_{\mu_{124}}^{AVV} \left(1,2,4\right) - t_{\mu_{124}}^{AVV} \left(1,3,4\right) \\ p_{43}^{\mu_4} t_{\mu_{1234}}^{AVVV} &= t_{\mu_{123}}^{AVV} \left(1,2,3\right) - t_{\mu_{123}}^{AVV} \left(1,2,4\right) \\ p_{41}^{\mu_1} t_{\mu_{1234}}^{AVVV} &= t_{\mu_{423}}^{AVV} \left(1,2,3\right) - t_{\mu_{234}}^{AVV} \left(2,3,4\right) - 2m t_{\mu_{234}}^{PVVV} \end{split}$$

The integrated three point functions assume the setting

$$T_{\mu_1\mu_2\mu_3}^{AVV}(i,j,l) = \frac{4}{3} \varepsilon_{\mu_{123}\nu_{123}} p_{ji}^{\nu_2} p_{li}^{\nu_3} \left(P_{ij} + P_{il} + P_{jl} \right)^{\nu_4} \Delta_{\nu_4}^{\nu_1}$$
(6.1)

and the finite box arising from the contraction with axial-vertex momentum is

$$T_{\mu_{234}}^{PVVV} = -8m\varepsilon_{\mu_{234}\nu_{123}}p_{21}^{\nu_1}p_{32}^{\nu_2}p_{42}^{\nu_3}J_4, \tag{6.2}$$

where J_4 has the property

$$J_4\left(p_{ij}^2=0\right) = -\frac{i}{3!\left(4\pi\right)^3 m^2}.$$

To start with, one can use the general formula in the dimension d = 2n, to the string of 2n + 2 gamma matrices plus γ_* using the definition, as follows

$$\operatorname{tr}(\gamma_*\gamma_{a_1a_2\cdots a_{2n+1}a_{2n+2}}) = \operatorname{tr}(\mathbf{1}_{2^n\times 2^n}) i^{3n-1} \sum_{k=1}^{2n+1} \sum_{j=k+1}^{2n+2} (-1)^{j+k+1} g_{a_ka_j} \varepsilon_{a_1\cdots \hat{a}_k\cdots \hat{a}_j\cdots (2n+2)},$$

where the hat means omission of that indexes, the formula was indexed in such a way to facilitate its use by means of substitutions, effectively encompassing all the results present in this paper. The first version, that comes from the substitution adjacent to the matrix γ_{μ_1} , is obtained through the choices of n = 3 and the sequence $a_1 a_2 \cdots a_8$ as $\mu_1 \nu_1 \mu_2 \nu_2 \mu_3 \nu_3 \mu_4 \nu_4$.

First Version: The sign tensors came from the first four rows of the trace of eight matrices below

$$\begin{split} &\frac{1}{8}K_{1234}^{\nu_{1234}}\operatorname{tr}(\gamma_*\gamma_{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3\mu_4\nu_4}) \\ &= -\varepsilon_{\mu_2\mu_3\mu_4\nu_{123}}\left[K_{234}^{\nu_{123}}K_{1\mu_1} - K_{134}^{\nu_{123}}K_{2\mu_1} + K_{124}^{\nu_{123}}K_{3\mu_1} - K_{123}^{\nu_{123}}K_{4\mu_1}\right] \\ &-\varepsilon_{\mu_1\mu_3\mu_4\nu_{123}}\left[K_{234}^{\nu_{123}}K_{1\mu_2} + K_{134}^{\nu_{123}}K_{2\mu_2} - K_{124}^{\nu_{123}}K_{3\mu_2} + K_{123}^{\nu_{123}}K_{4\mu_2}\right] \\ &+\varepsilon_{\mu_1\mu_2\mu_4\nu_{123}}\left[K_{234}^{\nu_{123}}K_{1\mu_3} - K_{134}^{\nu_{123}}K_{2\mu_3} - K_{124}^{\nu_{123}}K_{3\mu_4} + K_{123}^{\nu_{123}}K_{4\mu_3}\right] \\ &-\varepsilon_{\mu_1\mu_2\mu_3\nu_{123}}\left[K_{234}^{\nu_{123}}K_{1\mu_4} - K_{134}^{\nu_{123}}K_{2\mu_4} + K_{124}^{\nu_{123}}K_{3\mu_4} + K_{123}^{\nu_{123}}K_{4\mu_4}\right] \\ &+\varepsilon_{\mu_1\mu_2\mu_3\mu_4\nu_1\nu_2}\left[K_{34}^{\nu_{12}}\left(K_1 \cdot K_2\right) - K_{24}^{\nu_{12}}\left(K_1 \cdot K_3\right) + K_{23}^{\nu_{12}}\left(K_1 \cdot K_4\right)\right] \\ &+\varepsilon_{\mu_1\mu_2\mu_3\mu_4\nu_1\nu_2}\left[K_{14}^{\nu_{12}}\left(K_2 \cdot K_3\right) - K_{13}^{\nu_{12}}\left(K_2 \cdot K_4\right) + K_{12}^{\nu_{12}}\left(K_3 \cdot K_4\right)\right] \\ &+\left[g_{\mu_1\mu_2}\varepsilon_{\mu_3\mu_4\nu_1\nu_2\nu_3\nu_4} - g_{\mu_1\mu_3}\varepsilon_{\mu_2\mu_4\nu_1\nu_2\nu_3\nu_4} + g_{\mu_3\mu_4}\varepsilon_{\mu_1\mu_2\nu_1\nu_2\nu_3\nu_4}\right]K_{1234}^{\nu_{1234}}, \end{split}$$

the general form of that tensors is given by

$$\varepsilon_{\mu_{abc}\nu_{123}}t_{\mu_d}^{(s_1s_2s_3)\nu_{123}} = \varepsilon_{\mu_{abc}\nu_{123}}\left[K_{1\mu_d}K_{234}^{\nu_{123}} + s_1K_{2\mu_d}K_{134}^{\nu_{123}} + s_2K_{3\mu_d}K_{124}^{\nu_{123}} + s_3K_{4\mu_d}K_{123}^{\nu_{123}}\right]\frac{1}{D_{1234}},$$

and their integrals, following our procedure, are

$$\begin{split} \varepsilon_{\mu_{abc}\nu_{123}}T^{(s_{1}s_{2}s_{3})\nu_{123}}_{\mu_{d}} &= (1+s_{1})\,\varepsilon_{\mu_{abc}\nu_{123}}p_{31}^{\nu_{2}}p_{41}^{\nu_{3}}\left(J^{\nu_{1}}_{4\mu_{d}}+p_{21\mu_{d}}J^{\nu_{1}}_{4}\right) \\ &\quad -(1-s_{2})\,\varepsilon_{\mu_{abc}\nu_{123}}p_{21}^{\nu_{2}}p_{41}^{\nu_{3}}\left(J^{\nu_{1}}_{4\mu_{d}}+p_{31\mu_{d}}J^{\nu_{1}}_{4}\right) \\ &\quad +(1+s_{3})\,\varepsilon_{\mu_{abc}\nu_{123}}p_{21}^{\nu_{2}}p_{31}^{\nu_{3}}\left(J^{\nu_{1}}_{4\mu_{d}}+p_{41\mu_{d}}J^{\nu_{1}}_{4}\right) \\ &\quad +\frac{1}{6}\varepsilon_{\mu_{abc}\nu_{123}}\left[(1+s_{1})\,p_{31}^{\nu_{2}}p_{41}^{\nu_{3}}-(1-s_{2})\,p_{21}^{\nu_{2}}p_{41}^{\nu_{3}}+(1+s_{3})\,p_{21}^{\nu_{2}}p_{31}^{\nu_{3}}\right]\Delta_{4\mu_{d}}^{\nu_{1}} \\ &\quad +\frac{1}{6}\varepsilon_{\mu_{abc}\nu_{12}}\left[(1+s_{1})\,p_{31}^{\nu_{1}}p_{41}^{\nu_{2}}-(1-s_{2})\,p_{21}^{\nu_{1}}p_{41}^{\nu_{2}}+(1+s_{3})\,p_{21}^{\nu_{1}}p_{31}^{\nu_{2}}\right]I_{\mathrm{log}},\end{split}$$

a general property to note is that one of the tensors is finite and zero

$$\varepsilon_{\mu_{abc}\nu_{123}}T^{(-+-)\nu_{123}}_{\mu_d} = 0$$

and as always some odd and rank-two sub-amplitudes are finite and vanishing, all equal to $\pm T^{AVPP}_{\mu\nu}$, as in 4D is the case of the T^{ASS}_{μ} like terms. To the first version, we obtain the result

$$\left(T_{\mu_{1234}}^{AVVV}\right)_1 = -8\varepsilon_{\mu_{134}\nu_{123}}T_{\mu_2}^{(+-+)\nu_{123}} + 8\varepsilon_{\mu_{124}\nu_{123}}T_{\mu_3}^{(--+)\nu_{123}} - 8\varepsilon_{\mu_{123}\nu_{123}}T_{\mu_4}^{(-++)\nu_{123}} - \frac{1}{2}\varepsilon_{\mu_{1234}}\nu_{\mu_2}^{TPPP},$$

where the sub-amplitude, resulting of sum of the bilinear terms above with the mass terms, coming from the trace $\operatorname{tr}(\gamma_*\gamma_{\mu_{1234}\nu_{12}}) = 8\varepsilon_{\mu_{1234}\nu_{12}}$, has the integrand

$$\begin{aligned} \frac{1}{2} \varepsilon_{\mu_{1234}} {}^{\nu_{12}} t_{\nu_{12}}^{\tilde{T}PPP} &= +8\varepsilon_{\mu_{1234}\nu_{12}} \left[-K_{12}^{\nu_{12}}S_{34} + K_{13}^{\nu_{12}}S_{24} - K_{14}^{\nu_{12}}S_{23} \right] \frac{1}{D_{1234}} \\ &+8\varepsilon_{\mu_{1234}\nu_{12}} \left[-K_{23}^{\nu_{12}}S_{14} + K_{24}^{\nu_{12}}S_{13} - K_{34}^{\nu_{12}}S_{12} \right] \frac{1}{D_{1234}}, \end{aligned}$$

its finite part is given by

$$\begin{aligned} -\frac{1}{2}\varepsilon_{\mu_{1234}}^{\nu_{12}}T_{\nu_{12}}^{\tilde{T}PPP} &= 8\varepsilon_{\mu_{1234}\nu_{12}}\left[\left(p_{31}\cdot p_{43}\right)p_{21}^{\nu_{2}} - \left(p_{21}\cdot p_{42}\right)p_{31}^{\nu_{2}} + \left(p_{21}\cdot p_{32}\right)p_{41}^{\nu_{2}}\right]J_{4}^{\nu_{1}} \\ &+ 4\varepsilon_{\mu_{1234}\nu_{12}}\left(p_{21}^{\nu_{1}}p_{41}^{\nu_{2}}p_{31}^{2} - p_{31}^{\nu_{1}}p_{41}^{\nu_{2}}p_{21}^{2} - p_{21}^{\nu_{1}}p_{31}^{\nu_{2}}p_{41}^{2}\right)J_{4} \\ &+ 4\varepsilon_{\mu_{1234}\nu_{12}}\left[2p_{43}^{\nu_{2}}J_{3}^{\nu_{1}}\left(p_{31}, p_{41}\right) + p_{31}^{\nu_{1}}p_{41}^{\nu_{2}}J_{3}\left(p_{31}, p_{41}\right)\right] \\ &+ 4\varepsilon_{\mu_{1234}\nu_{12}}\left[2p_{21}^{\nu_{2}}J_{3}^{\nu_{1}}\left(p_{21}, p_{41}\right) - p_{21}^{\nu_{1}}p_{41}^{\nu_{2}}J_{3}\left(p_{21}, p_{41}\right)\right] \\ &+ 4\varepsilon_{\mu_{1234}\nu_{12}}\left[p_{32}^{\nu_{1}}p_{43}^{\nu_{2}}J_{3}\left(p_{32}, p_{42}\right) + p_{21}^{\nu_{1}}p_{31}^{\nu_{2}}J_{3}\left(p_{21}, p_{31}\right)\right],\end{aligned}$$

note the fact that this vertex is a pseudo-tensor one, as we anticipated. Gathering all the divergent objects that appear in the sub-structure and the combination of sign tensors, we get the set of surface terms

$$S_{1\mu_{1234}} = -\frac{8}{3} \left[\varepsilon_{\mu_{134}\nu_{123}} p_{32}^{\nu_2} p_{42}^{\nu_3} \Delta_{4\mu_2}^{\nu_1} + \varepsilon_{\mu_{124}\nu_{123}} p_{21}^{\nu_2} p_{43}^{\nu_3} \Delta_{4\mu_3}^{\nu_1} + \varepsilon_{\mu_{123}\nu_{123}} p_{21}^{\nu_2} p_{31}^{\nu_3} \Delta_{4\mu_4}^{\nu_1} \right] \\ -\frac{8}{6} \varepsilon_{\mu_{1234}\nu_{12}} \left[p_{43}^{\nu_2} \left(P_{13} + P_{34} + P_{14} \right)^{\nu_4} + p_{21}^{\nu_2} \left(P_{12} + P_{24} + P_{14} \right)^{\nu_4} \right] \Delta_{4\nu_4}^{\nu_1} \right]$$

meanwhile the irreducible object cancels exactly. With the tools, we have been using up to now, these results are a quite direct, but a long, task.

Second Version: Without any new protocol, only computing the trace with the definition of γ_* in the right or left of γ_{μ_2} , and following exactly the same steps for two, four, and that first case in six dimensions, we have the formula

$$(T_{\mu_{1234}}^{AVVV})_2 = -8\varepsilon_{\mu_{234}\nu_{123}}T_{\mu_1}^{(+-+)\nu_{123}} - 8\varepsilon_{\mu_{124}\nu_{123}}T_{\mu_3}^{(++-)\nu_{123}} + 8\varepsilon_{\mu_{123}\nu_{123}}T_{\mu_4}^{(+--)\nu_{123}} - \frac{1}{2}\varepsilon_{\mu_{1234}}\nu_{\mu_2}T_{\nu_1\nu_2}^{STPP},$$

here the sub-amplitude is STPP, where we see another vertex a two-rank tensor one, its finite part is

$$\begin{split} &-\frac{1}{2}\varepsilon_{\mu_{1234}} ^{\nu_{12}}T_{\nu_{1}\nu_{2}}^{STPP} \\ &= 8\varepsilon_{\mu_{1234}\nu_{12}}\left[-\left(p_{41}\cdot p_{43}\right)p_{21}^{\nu_{2}}+\left(p_{41}\cdot p_{42}\right)p_{31}^{\nu_{2}}-\left(p_{31}\cdot p_{32}\right)p_{41}^{\nu_{2}}\right]J_{4}^{\nu_{1}} \\ &+4\varepsilon_{\mu_{1234}\nu_{12}}\left[p_{31}^{\nu_{1}}p_{43}^{\nu_{2}}p_{21}^{2}-p_{21}^{\nu_{1}}p_{42}^{\nu_{2}}p_{31}^{2}+p_{21}^{\nu_{1}}p_{32}^{\nu_{2}}p_{41}^{2}-4m^{2}p_{32}^{\nu_{1}}p_{42}^{\nu_{2}}\right]J_{4} \\ &+8\varepsilon_{\mu_{1234}\nu_{12}}\left[p_{41}^{\nu_{2}}J_{3}^{\nu_{1}}\left(p_{21},p_{41}\right)-p_{32}^{\nu_{2}}J_{3}^{\nu_{1}}\left(p_{21},p_{31}\right)\right]-4\varepsilon_{\mu_{1234}\nu_{12}}p_{32}^{\nu_{1}}p_{43}^{\nu_{2}}J_{3}\left(p_{42},p_{43}\right) \\ &+4\varepsilon_{\mu_{1234}\nu_{12}}\left[-p_{31}^{\nu_{1}}p_{43}^{\nu_{2}}J_{3}\left(p_{31},p_{41}\right)+p_{21}^{\nu_{1}}p_{42}^{\nu_{2}}J_{3}\left(p_{21},p_{41}\right)-p_{21}^{\nu_{1}}p_{32}^{\nu_{2}}J_{3}\left(p_{21},p_{31}\right)\right]. \end{split}$$

The divergent terms are summed between the two sectors, and provide us with

$$S_{2\mu_{1234}} = +\frac{8}{3} \left[-\varepsilon_{\mu_{234}\nu_{123}} p_{32}^{\nu_2} p_{43}^{\nu_3} \Delta_{4\mu_1}^{\nu_1} - \varepsilon_{\mu_{124}\nu_{123}} p_{31}^{\nu_2} p_{41}^{\nu_3} \Delta_{4\mu_3}^{\nu_1} + \varepsilon_{\mu_{123}\nu_{123}} p_{32}^{\nu_2} p_{41}^{\nu_3} \Delta_{4\mu_4}^{\nu_1} \right] \\ +\frac{8}{3} \varepsilon_{\mu_{1234}\nu_{12}} \left[(k_1^{\nu_3} + k_2^{\nu_3} + k_3^{\nu_3}) p_{32}^{\nu_2} - (k_1^{\nu_3} + k_2^{\nu_3} + k_4^{\nu_3}) p_{41}^{\nu_2} \right] \Delta_{4\nu_3}^{\nu_1},$$

owing this elements the computations follow in the same vein as the first version and all the other scenarios we presented.

6.1. RAGF, Linearity and Uniqueness

RAGF: First Version. For the contraction associated to the first vector vertex, the divergent terms obey

$$p_{21}^{\mu_2} \left(T_{\mu_{1234}}^{AVVV} \right)_1 = T_{\mu_1\mu_3\mu_4}^{AVV} \left(1, 3, 4 \right) - T_{\mu_1\mu_3\mu_4}^{AVV} \left(2, 3, 4 \right),$$

and to the finite part, it is necessary to use identities¹² to exchange the position of indexes as done in 4D and the results in the appendix (D), to determine that it obeys $p_{21}^{\mu_2} \left(T_{\mu_{1234}}^{AVVV}\right)_1^{\text{finite}} = 0$, and it is quite direct to verify that the part of the surface terms comply exactly with

$$\begin{split} p^{\mu_3}_{32} \left(T^{AVVV}_{\mu_{1234}}\right)_1 \ &= \ T^{AVV}_{\mu_1\mu_2\mu_4} \left(1,2,4\right) - T^{AVV}_{\mu_1\mu_2\mu_4} \left(1,3,4\right) \\ p^{\mu_4}_{43} \left(T^{AVVV}_{\mu_{1234}}\right)_1 \ &= \ T^{AVV}_{\mu_1\mu_2\mu_3} \left(1,2,3\right) - T^{AVV}_{\mu_1\mu_2\mu_3} \left(1,2,4\right). \end{split}$$

For the contraction with the momentum corresponding to the axial vertex, we must find among other things the expression

$$T_{\mu_4\mu_2\mu_3}^{AVV}(1,2,3) - T_{\mu_2\mu_3\mu_4}^{AVV}(2,3,4) = \frac{8}{6} \varepsilon_{\mu_{234}\nu_{123}} p_{32}^{\nu_2} \left[2p_{21}^{\nu_3} p_{41}^{\nu_4} - p_{41}^{\nu_3} \left(P_{23} + P_{24} + P_{43} \right)^{\nu_4} \right] \Delta_{4\nu_4}^{\nu_1}, \quad (6.3)$$

but in the contraction of the computed four-point function, as expected, we must use

$$p_{21}^{\nu_2} p_{31}^{\nu_3} p_{41}^{\nu_4} \left[-\varepsilon_{\mu_{34}\nu_{1234}} \Delta_{4\mu_2}^{\nu_1} + \varepsilon_{\mu_{24}\nu_{1234}} \Delta_{4\mu_3}^{\nu_1} - \varepsilon_{\mu_{23}\nu_{1234}} \Delta_{4\mu_4}^{\nu_1} \right] \\ = -\varepsilon_{\mu_{234}\nu_{234}} p_{41}^{\nu_4} p_{32}^{\nu_2} p_{42}^{\nu_3} \Delta_{4\rho}^{\rho} + \varepsilon_{\mu_{234}\nu_{123}} \left[-p_{41}^{\nu_3} p_{32}^{\nu_2} p_{42}^{\nu_4} + p_{41}^{\nu_3} p_{42}^{\nu_2} p_{32}^{\nu_4} + p_{32}^{\nu_2} p_{42}^{\nu_4} + p_{42}^{\nu_3} p_{42}^{\nu_4} + p_{32}^{\nu_4} p_{42}^{\nu_3} p_{42}^{\nu_4} \right] \Delta_{4\nu_4}^{\nu_1}$$

$$\varepsilon_{\mu_{134}\nu_{123}} \left[p_{31}^{\nu_2} p_{41}^{\nu_3} p_{21}^2 - (p_{21} \cdot p_{41}) p_{21}^{\nu_3} p_{31}^{\nu_2} + (p_{21} \cdot p_{31}) p_{21}^{\nu_3} p_{41}^{\nu_2} \right] J_4^{\nu_1} = \varepsilon_{\mu_{134}\nu_{234}} p_{31}^{\nu_2} p_{41}^{\nu_3} p_{21}^{\nu_4} (p_{21\nu_1} J_4^{\nu_1})$$

 $^{^{12}}$ One exemple is

that after added to the other terms of the expression, with some trivial algebraic manipulations to find the configuration of P_{ij} in the expression of the AVV parts, we find out

$$p_{41}^{\mu_1} \left(T_{\mu_{1234}}^{AVVV} \right)_1 = T_{\mu_4\mu_2\mu_3}^{AVV} \left(1, 2, 3 \right) - T_{\mu_2\mu_3\mu_4}^{AVV} \left(2, 3, 4 \right) + \frac{8}{3} \varepsilon_{\mu_{234}\nu_{234}} p_{41}^{\nu_4} p_{32}^{\nu_2} p_{42}^{\nu_3} \Delta_{4\rho}^{\rho},$$

where we see an additional term depending on the external momenta and the surface's term trace. On the other hand, to the finite part we use using the identity¹³ $\varepsilon_{[\mu_{34}\nu_{1234}}J^{\nu_1}_{4\mu_2]} = 0$ to exchange indexes in search of contractions with the momenta, in addition to¹⁴ $\varepsilon_{[\mu_{234}\nu_{123}}p_{41\nu_4]}J^{\nu_1}_4$, and the reductions of the six dimensional basic finite functions, we reach at the finite part of the contraction. Summing up both parts the total amplitude obeys

$$p_{41}^{\mu_1} \left(T_{\mu_{1234}}^{AVVV} \right)_1 = -2m T_{\mu_{234}}^{PVVV} + T_{\mu_4\mu_2\mu_3}^{AVV} \left(1, 2, 3 \right) - T_{\mu_2\mu_3\mu_4}^{AVV} \left(2, 3, 4 \right) \\ + \frac{8}{3} \varepsilon_{\mu_{234}\nu_{123}} p_{21}^{\nu_1} p_{31}^{\nu_2} p_{41}^{\nu_3} \left[\Delta_{4\rho}^{\rho} + i \left(4\pi \right)^{-3} \right].$$

RAGF: Second Version

The pattern is the same, the conditioning term for the satisfaction of the relations appears in the second vertex, and the three other are found without restriction, for this reason we only list the conditioned one

$$p_{21}^{\mu_2} \left(T_{\mu_{1234}}^{AVVV} \right)_2 = T_{\mu_{134}}^{AVV} \left(1, 3, 4 \right) - T_{\mu_{134}}^{AVV} \left(2, 3, 4 \right) + \frac{8}{3} \varepsilon_{\mu_{134}\nu_{123}} p_{21}^{\nu_1} p_{31}^{\nu_2} p_{41}^{\nu_3} \left[\Delta_{4\rho}^{\rho} + i \left(4\pi \right)^{-3} \right] + \frac{1}{3} \varepsilon_{\mu_{134}\nu_{123}} p_{21}^{\nu_1} p_{31}^{\nu_2} p_{41}^{\nu_3} \left[\Delta_{4\rho}^{\rho} + i \left(4\pi \right)^{-3} \right] + \frac{1}{3} \varepsilon_{\mu_{134}\nu_{134}} p_{21}^{\nu_1} p_{31}^{\nu_2} p_{41}^{\nu_3} \left[\Delta_{4\rho}^{\rho} + i \left(4\pi \right)^{-3} \right] + \frac{1}{3} \varepsilon_{\mu_{134}\nu_{134}} p_{41}^{\nu_2} p_{41}^{\nu_3} \left[\Delta_{4\rho}^{\rho} + i \left(4\pi \right)^{-3} \right] + \frac{1}{3} \varepsilon_{\mu_{134}\nu_{134}} p_{41}^{\nu_2} p_{41}^{\nu_3} \left[\Delta_{4\rho}^{\rho} + i \left(4\pi \right)^{-3} \right] + \frac{1}{3} \varepsilon_{\mu_{134}\nu_{134}} p_{41}^{\nu_2} p_{41}^{\nu_3} \left[\Delta_{4\rho}^{\rho} + i \left(4\pi \right)^{-3} \right] + \frac{1}{3} \varepsilon_{\mu_{134}\nu_{134}} p_{41}^{\nu_3} p_{41}^{\nu_4} \left[\Delta_{4\rho}^{\rho} + i \left(4\pi \right)^{-3} \right] + \frac{1}{3} \varepsilon_{\mu_{134}\nu_{134}} p_{41}^{\nu_4} p_$$

For the uniqueness, we subtract the expressions and using the same manipulations for the RAGFs and get

$$\left(T_{\mu_{1234}}^{AVVV}\right)_1 - \left(T_{\mu_{1234}}^{AVVV}\right)_2 = -\frac{8}{3}\varepsilon_{\mu_{1234}\nu_{12}}p_{32}^{\nu_1}p_{43}^{\nu_2}\left[\Delta_{4\rho}^{\rho} + \frac{i}{(4\pi)^3}\right].$$

Showing once more that connection among linearity and uniqueness in the sense we have posed it, different formulae to the traces do not deliver identical tensor and their equivalence depends on the value of surface term. These results are interrelated to a low energy theorem, or the behavior of the $T^{PVVV}_{\mu_{234}}$ amplitude for all bilinears set to zero.

$$2mT^{PVVV}_{\mu_{234}} = \frac{8i}{3(4\pi)^3} \varepsilon_{\mu_{234}\nu_{123}} p^{\nu_1}_{21} p^{\nu_2}_{32} p^{\nu_3}_{42} \neq 0$$

Adopting $q_1 = q_2 + q_3 + q_4$, where q_2 , q_3 , q_4 the incoming momenta and q_1 is outgoing, as in 4D. The tensor casting the properties of the AVVV box will be given by

$$\begin{split} F_{\mu_{1234}} &= \varepsilon_{\mu_{1234}\nu_{12}} q_2^{\nu_1} q_3^{\nu_2} F_1 + \varepsilon_{\mu_{1234}\nu_{12}} q_2^{\nu_1} q_4^{\nu_2} F_2 + \varepsilon_{\mu_{1234}\nu_{12}} q_3^{\nu_1} q_4^{\nu_2} F_3 \\ &+ \varepsilon_{\mu_{234}\nu_{123}} q_2^{\nu_1} q_3^{\nu_2} q_4^{\nu_3} \left[q_{2\mu_1} G_1 + q_{3\mu_1} G_2 + q_{4\mu_1} G_3 \right] \\ &+ \varepsilon_{\mu_{134}\nu_{123}} q_2^{\nu_1} q_3^{\nu_2} q_4^{\nu_3} \left[q_{2\mu_2} G_4 + q_{3\mu_2} G_5 + q_{4\mu_2} G_6 \right] \\ &+ \varepsilon_{\mu_{124}\nu_{123}} q_2^{\nu_1} q_3^{\nu_2} q_4^{\nu_3} \left[q_{2\mu_3} G_7 + q_{3\mu_3} G_8 + q_{4\mu_3} G_9 \right] \\ &+ \varepsilon_{\mu_{123}\nu_{123}} q_2^{\nu_1} q_3^{\nu_2} q_4^{\nu_3} \left[q_{2\mu_4} G_{10} + q_{3\mu_4} G_{11} + q_{4\mu_4} G_{12} \right], \end{split}$$

¹³ The specific substitution necessary is

$$\varepsilon_{\mu_{34}\nu_{1234}}J_{4\mu_{2}}^{\nu_{1}} - \varepsilon_{\mu_{24}\nu_{1234}}J_{4\mu_{3}}^{\nu_{1}} + \varepsilon_{\mu_{23}\nu_{1234}}J_{4\mu_{4}}^{\nu_{1}} = -\varepsilon_{\mu_{234}\nu_{123}}J_{4\nu_{4}}^{\nu_{1}} + \varepsilon_{\mu_{234}\nu_{124}}J_{4\nu_{3}}^{\nu_{1}} - \varepsilon_{\mu_{234}\nu_{134}}J_{4\nu_{2}}^{\nu_{1}} + \varepsilon_{\mu_{234}\nu_{234}}J_{4\nu_{1}}^{\nu_{1}} + \varepsilon_{\mu_{234}\nu_{1234}}J_{4\nu_{2}}^{\nu_{1}} + \varepsilon_{\mu_{234}\nu_{1234}}J_{4\nu_{2}}^{\nu_{1}} + \varepsilon_{\mu_{234}\nu_{234}}J_{4\nu_{2}}^{\nu_{1}} + \varepsilon_{\mu_{234}\nu_{234}}J_{4\nu_{2}}^{\nu_{2}} + \varepsilon_{\mu_{234}\nu_{2}}^{\nu_{2}} + \varepsilon$$

 14 When multiplied by $p_{21}^{\nu_2}p_{31}^{\nu_3}p_{41}^{\nu_4}J_4^{\nu_1}$ the desired result takes the form of

 $\varepsilon_{\mu_{234}\nu_{123}} \left[p_{21}^{\nu_2} p_{31}^{\nu_3} p_{41}^2 - \left(p_{41} \cdot p_{31} \right) p_{21}^{\nu_2} p_{41}^{\nu_3} + \left(p_{41} \cdot p_{21} \right) p_{31}^{\nu_2} p_{41}^{\nu_3} \right] J_4^{\nu_1} \\ = \varepsilon_{\mu_{234}\nu_{234}} p_{21}^{\nu_2} p_{31}^{\nu_3} p_{41}^{\nu_4} \left[p_{41\nu_1} J_4^{\nu_1} \right] .$

and then the four contractions $q_i^{\mu_i} F_{\mu_{1234}} = \varepsilon_{\mu_1...i\dots 4\nu_{123}} q_2^{\nu_1} q_3^{\nu_2} q_4^{\nu_3} V_i$, the hat means to suppress that index, they allow to trace the following fact, in the point $q_i \cdot q_j = 0$, it is straightforward to obtain $V_1(0) + V_2(0) - V_3(0) + V_4(0) = 0$. Then, what we have independent of the consistency of the method or manipulations is that if the WI built out of divergence of the axial current is satisfied or not, we have

$$V_1(0) = \Omega^{PVVV}(0) \neq 0,$$

hence there are vector vertexes where the WIs are not satisfied, and vice-versa, if the vector currents are conserved the axial must not be due to

$$V_1(0) + \Omega^{PVVV}(0) + \mathcal{A} = 0,$$

and the series of propositions derived for 4D, in sections (5.2 and 5.3), follow here by analogous constructions, that we see as unnecessary, it would connect the consequences of kinematical properties of finite functions to the obstructions, inconsistencies, linearity violation, lack of uniqueness, and all that. The value of the investigation proposed here is that this results are the same in all dimensions.

7. FINAL REMARKS AND PERSPECTIVES

In this study, a detailed probe of a significant number of pseudo-tensor diagrams that correspond to anomalous amplitudes in two, four, and six dimensions is performed, following a strategy to cope with the divergences introduced in the thesis of O.A. Battistel [25]. Using the idea that is possible to separate in two sets the integrand of an arbitrary perturbative amplitude using systematically an identity that localizes the divergent parts in a set where the physical parameters, masses, and momenta, can be factored out of the integrals and other where only finite integrals reside and are freely integrated.

This procedure is applied to the bubbles, triangles, and box, enabling us to write down any expression as a sum of standard tensors, that is comprised of what we have called sign tensors, and another amplitude, of parity even, formed by vertexes of the same nature the main amplitude is composed. In four dimensions, for example, we get schematically

 $\left(T_{\mu_{123}}^{\Gamma_1\Gamma_2\Gamma_3}\right)_i = 4iC_{i\mu_{123}} \pm i\varepsilon_{\mu_{123}\nu_1} \left(\text{Corresponding sub-amplitude}\right)^{\nu_1}.$

Thus, after splitting off and organizing the divergent parts, without further action, the finite ones are integrated. In this point, summing up these two parts, the scalar objects $I_{\log}^{(2n)}$ exactly cancel in all cases, letting the final result as a sum of finite tensors and surface terms, $\Delta_{n+1;\mu_{12}}^{2n}$, as defined along this work. Such recipe crucially relies on the principle that it is possible to write the integral of a sum as the sum of their integrals the linearity of integration.

The role of this aspect as a fundamental element in this discussion then emerges, writing in 4D three equations $q_i^{\mu_i}T_{\mu_{123}}^{\Gamma_{123}} = T_{(-)i}^{AV} + \varepsilon_{\mu_{ab}\nu_{12}}q_2^{\nu_1}q_3^{\nu_2}\Omega_i$ for the RAGFs, it follows that, if the vanishing of the AV or their difference, in that equations were a possibility, then that would allow the vector and partial axial symmetry to be true. Such hope could be based on the fact that if these structures were only functions of routing differences, then using the charge conjugation matrix, $C\gamma_{\mu}C^{-1} = -\gamma_{\mu}^{T}$, the properties of the spinor-propagators, and traces, it would be possible to prove that $T_{\mu\nu}^{AV}(p) = -T_{\mu\nu}^{AV}(p)$, or that they are vanishing. To have this property means to have translational invariance or translational invariance in momentum-space, but computations reveal that such structures, in principle, depend on the unphysical and arbitrary sum of routings and are proportional to a surface term, $T^{AV} \sim \varepsilon_{\alpha\mu\nu_1\nu_2} p_{ij}^{\nu_1} P_{ji}^{\nu_3} \Delta_{\nu_3}^{\nu_2}$, that violates the mentioned symmetry

Nevertheless, as demonstrated in section (5.2), regarding low-energy theorems, a tensor with the characteristics of AVV, for example, a function of the differences among the routings related by contraction to the well-defined tensor PVV, must satisfy, in this case, $p_{31}^{\mu_1}T_{\mu_{123}}^{AVV}|_0 = 0 \neq -2mT_{\mu_{23}}^{PVV}|_0$, what is impossible. At this point, anyone should notice that the satisfaction of the RAGF, or linearity, can not be satisfied for any value of the surface term, in particular, not for vanishing value.

Therefore, the next step is to consider that the undetermined content of all the tensors we investigated are always combinations of routings, surface terms, and the ε -tensor. With these assumptions in mind, in the form of hypotheses, plus the known RHS of the relations, the AVpart, we lay down, in definitive: it is impossible without additional conditions to satisfy all the RAGF, in other words, they are not valid for any value of the surface term, a result developed in the section (5.3). Beyond this, the obligation to satisfy all of them, the RAGFs, makes the kinematical property, at zero, of the PVV, the value and the reason why the surface term can not be vanishing, see eq. (5.76).For this reason, translational invariance in momentum space, and linearity are incompatible properties for these perturbative amplitudes. Moreover, if one could rule out any role for the internal momenta by adopting $\Delta_{3\mu\nu} = 0$ and have linearity, one would be wrestling with the fact that three-point amplitudes related by contraction do not vanish in zero,

$$2\Delta_{3\alpha}^{\alpha} = i \left[\Omega_3(0) - \Omega_2(0) - \Omega_1(0)\right] = -i/(2\pi)^2,$$

and, by necessity, when choosing the linear scenario, we can transform by use of linear combinations the internal momenta in terms of external ones. Because there are two independent external variables, we have two parameters available, that is not enough to keep all WIs. As expected, due non-ambiguous kinematical reasons.

The explicitly finite values of Ω_i always satisfies the last equality above, for arbitrary masses as well. Nevertheless, such a result is irreconcilable with that coming from a tensor with the properties of $T_{\mu_{123}}^{\Gamma_{123}}$, in which case the result should be equal to zero, but then, there must be some way to understand why this does not happen in any, minimally consistent, manipulation. After writing the internal momenta in terms of the external ones, assuming all arbitrary violations of the RAGF taking the surface term as an arbitrary quantity, we reach a tensor under the hypotheses stated in section (5.2). Follows that, all violations are encompassed by $V_i = \Omega_i + A_i$ and obey $A_3 - A_2 - A_1 = (2\pi)^{-2}$, demonstrated through the eq.'s (5.83). In it reside the straightforward fact that when two WI are satisfied, the third is violated by a unique amount independent of any consistent computational philosophy because there is no ambiguity in the values of finite amplitudes.

In what concerns the consequence of the Dirac traces, surface terms, and Schouten identities, in all these amplitudes arises the trace of 2n + 2 Dirac matrices and an odd number of the chiral matrices, schematically

$$\operatorname{tr}(\gamma_*\gamma_{\mu_1}\cdots\gamma_{\mu_{2n+2}})\sim\sum\pm g_{\mu_i\mu_j}\varepsilon_{\mu_1\cdots\hat{\mu}_i\cdots\hat{\mu}_j\cdots\mu_{2n}}.$$

Applying the definition of $\gamma_* = \varepsilon_{\nu_{1234}} \gamma^{\nu_{1234}}/24$ or using the identity $\gamma_* \gamma_{\mu_i} = \varepsilon_{\mu_i \nu_{123}} \gamma^{\nu_{123}}/6$ in the adjacent position of the matrix γ_{μ_i} , we have shown the tensors calculated to correspond to the versions defined as the main ingredients of the investigation, that a priori are not equivalent for $i \neq j$. If the surface term is adopted zero, the obtained set of tensors have the property they violate the RAGF around that vertex, that means, to the vertex corresponding to γ_{μ_i} , at the level of the diagram, the WI gets violated in the same vertex. These specific types of substitutions deliver different expressions in the number of monomials, but their difference, after integration,

is a combination of finite and null integrals. Their linear combination, of the versions, forms the building block of any other identity, and all the common substitutions are a subset of these possibilities. Arbitrary combinations of these building blocks can be used to explain any result obtained in scenarios without internal momentum ambiguities.

Adopting $\Delta_{3\mu\nu} = 0$ makes the amplitudes to depend on the traces used, the Schouten identity inside the integral that connects the integrands ceases to make it in the final integrated results. At the end of the day this is what breaks the linearity of integration and violates the RAGFs. Different formulae for the traces do not deliver identical tensors, and the equivalence depends on the value of the surface term.

For the term uniqueness that we have employed, some definition is necessary for it to work as a concrete criterion. A criterion that makes the amplitudes unique in a universal sense is impossible since they are divergent quantities. After renormalized, they become dependent on an arbitrary mass scale, and this is beginning of renormalization group equations.

However once an expression is attributed by a regularization, there is no other way to get another result of the same procedure in even tensors. On the other hand, to our amplitudes, adopting the same interpretation to the surface terms led to various different tensors. In this narrow sense we have defined uniqueness: if the stance on the divergent quantities is the same, uniqueness implies only one answer. Through this definition, all mathematical manipulations leads to one result. Apart the manipulations used in this work, when bilinears present in the expressions are not reduced or taking traces after the integrals manipulated, the result is the same. Here comes the point of having a narrow definition the unique answer is a function of the routings $k_1, k_2, \dots, k_i \cdots$ taken as independent variables. The consequence is that choices that break momentum-space homogeneity must be adopted and establishes that one does not have a unique function of the external momenta. The mathematically unique answer, that does not depend on any sequence of algebraic operations, is not unique in another form, in a way that preserves homogeneity, that is, freedom of the origin of integration.

As of rule, there is the attractive option of making the surface term zero as done in even amplitudes and by a convenient choice of trace to obtain the symmetry content with the condition that not all symmetries can be present and not all distribution of anomalies is possible. In this scenario, there is a myriad of tensors to represent some amplitude, but it is a choice that can be made. Notwithstanding, there is one keeping linearity of integration, turning amplitudes unique functions of their routings, violating momentum-space homogeneity, and then to make physical interpretation is necessary to write the routings as combinations of the physical momenta. The conclusions about the symmetries are the same, but now in a different context: The integral of the sum is the sum of the integrals.

Appendix A: Traces of a String of Six Gamma and the Chiral Matrix

The a way to insert a Levi-Civita tensor in the traces with the chiral matrix come from the use of

$$\gamma_*\gamma_{[\mu_1\cdots\mu_r]} = \frac{i^{n-1+r(r+1)}}{(2n-r)!}\varepsilon_{\mu_1\cdots\mu_r\nu_{r+1}\cdots\nu_{2n}}\gamma^{[\nu_{r+1}\cdots\nu_{2n}]}$$

in 2n = 4 dimensions they are the identities with 0, 1, 2, 3, 4 antisymmetrized products, giving rise in traces of a string of six gamma matrices to (15, 10, 7, 6, 7) monomials respectively.

Trace Using $\gamma_* = i \varepsilon_{\nu_1 \nu_2 \nu_3 \nu_4} \gamma^{\nu_1 \nu_2 \nu_3 \nu_4} / 4!$ (Definition)

The three main positions to deploy the definition of the chiral matrix is around the gamma matrices present in the vertexes Γ_1, Γ_2 , and Γ_3 , in the left or the right they return the same integrated results.

Distinct positions of the chiral matrix in the trace. First one

$$t_{1} = \operatorname{tr} \left(\gamma_{*} \gamma_{\mu_{1}\nu_{1}\mu_{2}\nu_{2}\mu_{3}\nu_{3}} \right) = i\varepsilon^{\alpha_{1}\alpha_{2}\alpha_{3}\alpha_{4}} \operatorname{tr} \left(\gamma_{\alpha_{1}\alpha_{2}\alpha_{3}\alpha_{4}} \gamma_{\mu_{1}\nu_{1}\mu_{2}\nu_{2}\mu_{3}\nu_{3}} \right) / 4!$$

$$= +g_{\mu_{1}\nu_{1}}\varepsilon_{\mu_{2}\nu_{2}\mu_{3}\nu_{3}} - g_{\mu_{1}\mu_{2}}\varepsilon_{\nu_{1}\nu_{2}\mu_{3}\nu_{3}} + g_{\mu_{1}\nu_{2}}\varepsilon_{\nu_{1}\mu_{2}\mu_{3}\nu_{3}} - g_{\mu_{1}\mu_{3}}\varepsilon_{\nu_{1}\mu_{2}\nu_{2}\nu_{3}} + g_{\mu_{1}\nu_{3}}\varepsilon_{\nu_{1}\mu_{2}\nu_{2}\mu_{3}} + g_{\nu_{1}\mu_{2}}\varepsilon_{\mu_{1}\nu_{2}\mu_{3}\nu_{3}} - g_{\nu_{1}\nu_{2}}\varepsilon_{\mu_{1}\nu_{2}\mu_{3}\nu_{3}} + g_{\nu_{1}\mu_{3}}\varepsilon_{\mu_{1}\mu_{2}\nu_{2}\nu_{3}} - g_{\nu_{1}\nu_{3}}\varepsilon_{\mu_{1}\mu_{2}\nu_{2}\mu_{3}} + g_{\mu_{2}\nu_{2}}\varepsilon_{\mu_{1}\nu_{1}\mu_{3}\nu_{3}} - g_{\mu_{2}\mu_{3}}\varepsilon_{\mu_{1}\nu_{1}\nu_{2}\nu_{3}} + g_{\mu_{2}\nu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\mu_{3}} + g_{\nu_{2}\mu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\nu_{3}} - g_{\nu_{2}\nu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\mu_{3}} + g_{\mu_{3}\nu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\nu_{2}}$$

Second one

$$t_{2} = \operatorname{tr} \left(\gamma_{\mu_{1}\nu_{1}}\gamma_{*}\gamma_{\mu_{2}\nu_{2}\mu_{3}\nu_{3}} \right) = i\varepsilon^{\alpha_{1}\alpha_{2}\alpha_{3}\alpha_{4}} \operatorname{tr} \left(\gamma_{\mu_{1}\nu_{1}}\gamma_{\alpha_{1}\alpha_{2}\alpha_{3}\alpha_{4}}\gamma_{\mu_{2}\nu_{2}\mu_{3}\nu_{3}} \right) / 4!$$

$$= +g_{\mu_{1}\nu_{1}}\varepsilon_{\mu_{2}\nu_{2}\mu_{3}\nu_{3}} + g_{\mu_{1}\mu_{2}}\varepsilon_{\nu_{1}\nu_{2}\mu_{3}\nu_{3}} - g_{\mu_{1}\nu_{2}}\varepsilon_{\mu_{2}\nu_{2}\mu_{3}\nu_{3}} + g_{\mu_{1}\mu_{3}}\varepsilon_{\nu_{1}\mu_{2}\nu_{2}\nu_{3}} - g_{\mu_{1}\nu_{3}}\varepsilon_{\mu_{1}\mu_{2}\mu_{2}\nu_{2}\mu_{3}} + g_{\mu_{2}\nu_{2}}\varepsilon_{\mu_{1}\nu_{1}\mu_{3}\nu_{3}} - g_{\mu_{2}\mu_{3}}\varepsilon_{\mu_{1}\nu_{1}\nu_{2}\nu_{3}} + g_{\mu_{2}\nu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\nu_{3}} + g_{\mu_{2}\nu_{2}}\varepsilon_{\mu_{1}\nu_{1}\mu_{3}\nu_{3}} - g_{\mu_{2}\mu_{3}}\varepsilon_{\mu_{1}\nu_{1}\nu_{2}\nu_{3}} + g_{\mu_{2}\nu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\mu_{3}} + g_{\mu_{2}\nu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\nu_{3}} - g_{\mu_{2}\mu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\mu_{3}} + g_{\mu_{3}\nu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\mu_{3}} + g_{\mu_{3}\nu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\nu_{2}}$$

Third one

$$t_{3} = \operatorname{tr}(\gamma_{\mu_{1}\nu_{1}\mu_{2}\nu_{2}}\gamma_{*}\gamma_{\mu_{3}\nu_{3}}) = i\varepsilon^{\alpha_{1}\alpha_{2}\alpha_{3}\alpha_{4}}\operatorname{tr}(\gamma_{\mu_{1}\nu_{1}\mu_{2}\nu_{2}}\gamma_{\alpha_{1}\alpha_{2}\alpha_{3}\alpha_{4}}\gamma_{\mu_{3}\nu_{3}})/4!$$

$$= +g_{\mu_{1}\nu_{1}}\varepsilon_{\mu_{2}\nu_{2}\mu_{3}\nu_{3}} - g_{\mu_{1}\mu_{2}}\varepsilon_{\nu_{1}\nu_{2}\mu_{3}\nu_{3}} + g_{\mu_{1}\nu_{2}}\varepsilon_{\mu_{2}\nu_{2}\mu_{3}\nu_{3}} + g_{\mu_{1}\mu_{3}}\varepsilon_{\nu_{1}\mu_{2}\nu_{2}\nu_{3}} - g_{\mu_{1}\nu_{3}}\varepsilon_{\nu_{1}\mu_{2}\nu_{2}\mu_{3}}$$

$$+g_{\nu_{1}\mu_{2}}\varepsilon_{\mu_{1}\nu_{2}\mu_{3}\nu_{3}} - g_{\nu_{1}\nu_{2}}\varepsilon_{\mu_{1}\mu_{2}\mu_{3}\nu_{3}} - g_{\nu_{1}\mu_{3}}\varepsilon_{\mu_{1}\mu_{2}\nu_{2}\nu_{3}} + g_{\mu_{1}\nu_{3}}\varepsilon_{\mu_{1}\mu_{2}\nu_{2}\mu_{3}} + g_{\mu_{2}\nu_{2}}\varepsilon_{\mu_{1}\nu_{1}\mu_{3}\nu_{3}}$$

$$+g_{\mu_{2}\mu_{3}}\varepsilon_{\mu_{1}\nu_{1}\nu_{2}\nu_{3}} - g_{\mu_{2}\nu_{3}}\varepsilon_{\mu_{1}\nu_{1}\nu_{2}\mu_{3}} - g_{\nu_{2}\mu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\nu_{3}} + g_{\mu_{2}\nu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\mu_{3}} + g_{\mu_{3}\nu_{3}}\varepsilon_{\mu_{1}\nu_{1}\mu_{2}\nu_{2}}$$

we omit the imaginary unit. Now, these three expressions cast all the indexes of the tensor and they have fifteen terms each, in a narrow sense they could be called symmetric and considered to be as respecting all the symmetry among the indexes, see, by example, the appendix of the ref. [53] or the refs. [49][34], we do not focus on such adjectives, but on the fact they are enough to obtain any other result by a careful analysis, and encompassing any possible manipulations with these structures.

First things first, the sign differences are the unique distinguishing factor in that traces, they effectively sample the indexes among finite and surface terms in the real calculations. The aim is to demonstrate that any expression to the triangles investigated are just linear combinations of the ones we have detailed in the main body of this work.

Making the combinations, only using sums and not Schouten identities

$$t_{ij} = \frac{1}{2} \left(t_i + t_j \right),$$

we will have

$$t_{12} = -g_{\mu_1\nu_1}\varepsilon_{\mu_2\mu_3\nu_2\nu_3} - g_{\mu_2\nu_2}\varepsilon_{\mu_1\mu_3\nu_1\nu_3} + g_{\mu_2\nu_3}\varepsilon_{\mu_1\mu_3\nu_1\nu_2} - g_{\nu_2\mu_3}\varepsilon_{\mu_1\mu_2\nu_1\nu_3} - g_{\mu_3\nu_3}\varepsilon_{\mu_1\mu_2\nu_1\nu_2} - g_{\mu_2\mu_3}\varepsilon_{\mu_1\mu_1\nu_2\nu_3} - g_{\nu_2\nu_3}\varepsilon_{\mu_1\mu_2\mu_3\nu_1},$$

$$t_{13} = -g_{\mu_3\nu_3}\varepsilon_{\mu_1\mu_2\nu_1\nu_2} - g_{\mu_1\nu_1}\varepsilon_{\mu_2\mu_3\nu_2\nu_3} + g_{\mu_1\nu_2}\varepsilon_{\mu_2\mu_3\nu_1\nu_3} -g_{\nu_1\mu_2}\varepsilon_{\mu_1\mu_3\nu_2\nu_3} - g_{\mu_2\nu_2}\varepsilon_{\mu_1\mu_3\nu_1\nu_3} - g_{\mu_1\mu_2}\varepsilon_{\mu_3\nu_1\nu_2\nu_3} - g_{\nu_1\nu_2}\varepsilon_{\mu_1\mu_2\mu_3\nu_3},$$

$$t_{23} = -g_{\mu_2\nu_2}\varepsilon_{\mu_1\mu_3\nu_1\nu_3} - g_{\mu_1\nu_1}\varepsilon_{\mu_2\mu_3\nu_2\nu_3} + g_{\mu_1\nu_3}\varepsilon_{\mu_2\mu_3\nu_1\nu_2} - g_{\nu_1\mu_3}\varepsilon_{\mu_1\mu_2\nu_2\nu_3} - g_{\mu_3\nu_3}\varepsilon_{\mu_1\mu_2\nu_1\nu_2} - g_{\mu_1\mu_3}\varepsilon_{\mu_2\nu_1\nu_2\nu_3} - g_{\nu_1\nu_3}\varepsilon_{\mu_1\mu_2\mu_3\nu_2},$$

and we can employ the identities involving the antisymmetrized products to compute the same trace as well obtaining other formulas.

Trace Using $\gamma_* \gamma_{\mu_1} = -i\varepsilon_{\mu_1\nu_1\nu_2\nu_3} \gamma^{\nu_1\nu_2\nu_3}/3!$ The straightforward application

$$\eta_1(a) = \operatorname{tr}(\gamma_*\gamma_{abcdef}) = g_{bc}\varepsilon_{adef} - g_{bd}\varepsilon_{acef} + g_{be}\varepsilon_{acdf} - g_{bf}\varepsilon_{acde} + g_{cd}\varepsilon_{abef} - g_{ce}\varepsilon_{abdf} + g_{cf}\varepsilon_{abde} + g_{de}\varepsilon_{abcf} + g_{ef}\varepsilon_{abcd} - g_{df}\varepsilon_{abce}$$

the notation means that it uses a product with one Dirac matrix with index a in the substitution of $\gamma_*\gamma_a$

$$\eta_1(a) = -i\varepsilon_a \,^{\nu_1\nu_2\nu_3} \mathrm{tr}\left(\gamma_{\nu_1\nu_2\nu_3}\gamma_{bcdef}\right)/6.$$

The Trace Using $\gamma_*\gamma_{[ab]} = -i\varepsilon_{ab\nu_1\nu_2}\gamma^{\nu_1\nu_2}/2!$

The application of this one requires to express the ordinary product in terms of the antisymmetrized one

$$\gamma_*\gamma_{ab} = -\frac{i}{2}\varepsilon_{ab\nu_1\nu_2}\gamma^{\nu_1\nu_1} + g_{ab}\gamma_*,$$

thereby follows

$$\eta_2(ab) = \operatorname{tr}(\gamma_*\gamma_{abcdef}) = g_{ab}\varepsilon_{cdef} + g_{cd}\varepsilon_{abef} - g_{ce}\varepsilon_{abdf} + g_{cf}\varepsilon_{abde} + g_{de}\varepsilon_{abcf} - g_{df}\varepsilon_{abce} + g_{ef}\varepsilon_{abcd}$$

The Trace Using $\gamma_*\gamma_{[abc]} = i\varepsilon_{abc\nu}\gamma^{\nu}$

Expressing the antisymmetric product as common products we get

$$\gamma_*\gamma_{abc} = i\varepsilon_{abc\nu}\gamma^\nu + \gamma_*\left(g_{bc}\gamma_a - g_{ac}\gamma_b + g_{ab}\gamma_c\right)$$

with arbitrary indexes we get

$$\eta_3 (abc) = \operatorname{tr} \left(\gamma_* \gamma_{abcdef} \right) = g_{ab} \varepsilon_{cdef} - g_{ac} \varepsilon_{bdef} + g_{bc} \varepsilon_{adef} + g_{de} \varepsilon_{abcf} - g_{df} \varepsilon_{abce} + g_{ef} \varepsilon_{abcd}$$

where the notation means that we absorb the indexes a, b and c, with the identity and compute the resulting trace, it can be used to apply the substitution in any place desired. The use of this identity is a common choice on computation of this type of diagrams. In them, and all other possible results, after integration, we get some of the results obtained through the linear combinations t_{12} , t_{13} , and t_{23} .

The Trace Using $\gamma_*\gamma_{[abcd]} = i\varepsilon_{abcd}$ With the help of

$$\gamma_* \gamma_{abcd} = i \varepsilon_{abcd} \mathbf{1} + g_{ab} \gamma_* \gamma_{[cd]} - g_{ac} \gamma_* \gamma_{[bd]} + g_{ad} \gamma_* \gamma_{[bc]} + g_{bc} \gamma_* \gamma_{[ad]} - g_{bd} \gamma_* \gamma_{[ac]} + g_{cd} \gamma_* \gamma_{[ab]} + (g_{ab} g_{cd} - g_{ac} g_{bd} + g_{ad} g_{bc}) \gamma_*$$

under the trace and with arbitrary indexes, we get

$$\eta_4 (abcd) = \operatorname{tr} (\gamma_* \gamma_{abcdef}) = +g_{ab} \varepsilon_{cdef} - g_{ac} \varepsilon_{bdef} + g_{ad} \varepsilon_{bcef} +g_{bc} \varepsilon_{adef} - g_{bd} \varepsilon_{acef} + g_{cd} \varepsilon_{abef} + g_{ef} \varepsilon_{abcd}.$$

The Interconnection Among the Formulas: The difference on the integrated amplitudes either will identically vanishing as the integrand are exactly equal as in

$$[t_{12} - \eta_2(\mu_1\nu_1)] = 0; \qquad [t_{13} - \eta_4(\mu_1\nu_1\mu_2\nu_2)] = 0,$$

that is why is needless to say anything more, or will vanish because the difference inside a explicit computation always corresponds, when integrated, to finite null integrals, see

$$\begin{split} \left[t_{1}-\eta_{1}\left(\mu_{1}\right)\right]\frac{K_{123}^{\nu_{123}}}{D_{123}} &= \varepsilon_{\mu_{2}\mu_{3}\nu_{1}\nu_{2}}t_{\mu_{1}}^{(-+)\nu_{12}} + g_{\mu_{1}\mu_{2}}t_{\mu_{3}}^{ASS} - g_{\mu_{1}\mu_{3}}t_{\mu_{2}}^{ASS} \\ \left[t_{12}+\eta_{2}\left(\nu_{1}\mu_{2}\right)\right]\frac{K_{123}^{\nu_{123}}}{D_{123}} &= -\varepsilon_{\mu_{2}\mu_{3}\nu_{1}\nu_{2}}t_{\mu_{1}}^{(-+)\nu_{12}} + \varepsilon_{\mu_{1}\mu_{3}\nu_{1}\nu_{2}}t_{\mu_{2}}^{(-+)\nu_{12}} - g_{\mu_{2}\mu_{3}}t_{\mu_{1}}^{ASS} + g_{\mu_{1}\mu_{3}}t_{\mu_{2}}^{ASS} \\ \left[t_{12}-\eta_{3}\left(\mu_{1}\nu_{1}\mu_{2}\right)\right]\frac{K_{123}^{\nu_{123}}}{D_{123}} &= +\varepsilon_{\mu_{1}\mu_{3}\nu_{1}\nu_{2}}t_{\mu_{2}}^{(-+)\nu_{12}} + g_{\mu_{1}\mu_{2}}t_{\mu_{3}}^{ASS} - g_{\mu_{2}\mu_{3}}t_{\mu_{1}}^{ASS} \\ \left[t_{13}+\eta_{4}\left(\nu_{1}\mu_{2}\nu_{2}\mu_{3}\right)\right]\frac{K_{123}^{\nu_{123}}}{D_{123}} &= -\varepsilon_{\mu_{1}\mu_{2}\nu_{1}\nu_{2}}t_{\mu_{3}}^{(-+)\nu_{12}} - \varepsilon_{\mu_{2}\mu_{3}\nu_{1}\nu_{2}}t_{\mu_{1}}^{(-+)\nu_{12}} + g_{\mu_{2}\mu_{3}}t_{\mu_{1}}^{ASS} - g_{\mu_{1}\mu_{2}}t_{\mu_{3}}^{ASS} \end{split}$$

and as was showed in the text the well defined integrals corresponding to $\varepsilon_{\alpha\beta\nu_1\nu_2}t_{\rho}^{(-+)\nu_{12}}$ eq. (5.16) and t_{μ}^{ASS} in eq. (5.27), are null

$$\varepsilon_{\alpha\beta\nu_1\nu_2}T_{\rho}^{(-+)\nu_{12}} = T_{\mu}^{ASS} = 0$$

delivering the conclusion that any form of substitution or manipulation is accounted by the linear combination of the version one, two or three replacing the definition of γ_* left or right of the matrices γ_{μ_1} , γ_{μ_2} , and γ_{μ_3} . Whose consequence is that it is enough to unfold any feature of such calculations with the basic versions we described. What we showed here is the forms that identically correspond, not that all differences are finite and vanishing. The form obtained from t_{12} is not identical without conditions to any t_i , for example.

Appendix B: The Integrals in Two Dimensions

$$\bar{J}_{1}(k_{i}) = I_{\log}^{(2)}; \quad \bar{J}_{1}^{\mu}(k_{i}) = -k_{i}^{\nu} \Delta_{2\nu}^{(2)\mu}
J_{2} = i (4\pi)^{-1} \left[Z_{0}^{(-1)} \left(p^{2}, m^{2} \right) \right]$$

$$J_{2}^{\mu_{1}} = i (4\pi)^{-1} \left[-p^{\mu_{1}} Z_{1}^{(-1)} \right]$$

$$J_{2}^{\mu_{1}\mu_{2}} = i (4\pi)^{-1} \left[-\frac{1}{2} g^{\mu_{1}\mu_{2}} Z_{0}^{(0)} + p^{\mu_{1}} p^{\mu_{2}} Z_{2}^{(-1)} \right]$$

$$\bar{J}_{2}^{\mu_{1}\mu_{2}} = J_{2}^{\mu_{1}\mu_{2}} + \frac{1}{2} (\Delta_{2}^{(2)\mu_{1}\mu_{2}} + g^{\mu_{1}\mu_{2}} I_{\log}^{(2)})$$
(B1)
(B2)

Reductions

$$(n+1) p^{2} Z_{n+2}^{(-1)} = (n+1) p^{2} Z_{n+1}^{(-1)} - (n+1) m^{2} Z_{n}^{(-1)} - 1$$

$$(n+2) p^{2} Z_{n+1}^{(0)} = (n+1) p^{2} Z_{n}^{(0)} - m^{2} n Z_{n-1}^{(0)} - \frac{n}{(n+1)(n+2)} p^{2}; \quad n = 0, 1, 3, \cdots$$

$$Z_{0}^{(0)} = 2 p^{2} Z_{2}^{(-1)} - p^{2} Z_{1}^{(-1)}$$

They imply

$$J_2^{\mu_1} = -\frac{1}{2}p^{\mu_1}J_2; \quad p_{\mu_1}J_2^{\mu_1} = -\frac{1}{2}p^2J_2$$
$$p_{\mu_1}J_2^{\mu_1\mu_2} = -\frac{1}{2}p^2J_2^{\mu_2}; \quad g_{\mu_1\mu_2}J_2^{\mu_1\mu_2} = m^2J_2 + \frac{i}{4\pi}$$

Appendix C: The Integrals in Four Dimensions

Two Point:

$$J_{2}(p_{ij}) = -\frac{i}{(4\pi)^{2}} Z_{0}^{(0)} \left(p_{ij}^{2}, m^{2}\right); \text{ and } \qquad J_{2\mu}(p_{ij}) = \frac{i}{(4\pi)^{2}} p_{ij\mu} Z_{1}^{(0)} \left(p_{ij}^{2}, m^{2}\right)$$
$$\bar{J}_{2} = I_{\log}^{(4)} + J_{2}$$
$$\bar{J}_{2\mu} = \int \frac{d^{4}k}{(2\pi)^{4}} \frac{K_{i\mu}}{D_{ij}} = -\frac{1}{2} (P_{ij}^{\nu} \Delta_{3\mu\nu}^{(4)} + p_{ji\mu} I_{\log}^{(4)}) + J_{2\mu}$$

Three Point:

$$J_{3} = i (4\pi)^{-2} \left[Z_{00}^{(-1)} (p,q) \right]$$

$$J_{3\mu} = i (4\pi)^{-2} \left[-p_{\mu} Z_{10}^{(-1)} - q_{\mu} Z_{01}^{(-1)} \right]$$

$$J_{3\mu_{1}\mu_{2}} = i (4\pi)^{-2} \left[p_{\mu_{1}} p_{\mu_{2}} Z_{20}^{(-1)} + q_{\mu_{1}} q_{\mu_{2}} Z_{02}^{(-1)} + (p_{\mu_{1}} q_{\mu_{2}} + p_{\mu_{2}} q_{\mu_{1}}) Z_{11}^{(-1)} - \frac{1}{2} g_{\mu_{1}\mu_{2}} Z_{00}^{(0)} \right]$$

$$\bar{J}_{3\mu_{1}\mu_{2}} = J_{3\mu_{1}\mu_{2}} + \frac{1}{4} \left(\Delta_{3\mu_{1}\mu_{2}}^{(4)} + g_{\mu_{1}\mu_{2}} I_{\log}^{(4)} \right),$$

it is worth mention that the arguments p and q are only general variables that tag the entries of the functions, they must be carefully substitute for the ones that appear in a particular part of the investigation. **Reductions** of the basic functions. The two point basic function that appear satisfy a simple relation $2Z_1^{(0)} = Z_0^{(0)}$, as the three point obey

$$2\left[p^{2}Z_{10}^{(-1)} + (p \cdot q) Z_{01}^{(-1)}\right] = p^{2}Z_{00}^{(-1)} + \left[Z_{0}^{(0)}(q) - Z_{0}^{(0)}(q - p)\right]$$

$$2\left[q^{2}Z_{01}^{(-1)} + (p \cdot q) Z_{10}^{(-1)}\right] = q^{2}Z_{00}^{(-1)} + \left[Z_{0}^{(0)}(p) - Z_{0}^{(0)}(q - p)\right]$$

$$2\left[p^{2}Z_{20}^{(-1)} + (p \cdot q) Z_{11}^{(-1)}\right] = p^{2}Z_{10}^{(-1)} + Z_{00}^{(0)} - Z_{1}^{(0)}(q - p)$$

$$2\left[q^{2}Z_{02}^{(-1)} + (p \cdot q) Z_{11}^{(-1)}\right] = q^{2}Z_{01}^{(-1)} + Z_{00}^{(0)} - Z_{1}^{(0)}(q - p)$$

$$2\left[p^{2}Z_{11}^{(-1)} + (p \cdot q) Z_{02}^{(-1)}\right] = p^{2}Z_{01}^{(-1)} + \left[Z_{1}^{(0)}(q) - Z_{1}^{(0)}(q - p)\right]$$

$$2\left[q^{2}Z_{11}^{(-1)} + (p \cdot q) Z_{20}^{(-1)}\right] = q^{2}Z_{10}^{(-1)} + \left[Z_{1}^{(0)}(p) - Z_{1}^{(0)}(q - p)\right]$$

$$2Z_{00}^{(0)} = \left[p^{2}Z_{10}^{(-1)} + q^{2}Z_{01}^{(-1)}\right] - 2m^{2}Z_{00}^{(-1)} - 1 + 2Z_{1}^{(0)}(q - p)$$
(C1)

therefore it is possible to show that the tensors J satisfy

$$p^{\mu_{1}}J_{3\mu_{1}} = -\frac{1}{2}p^{2}J_{3} + \frac{1}{2}\left[J_{2}\left(q\right) - J_{2}\left(q - p\right)\right]$$

$$q^{\mu_{1}}J_{3\mu_{1}} = -\frac{1}{2}q^{2}J_{3} + \frac{1}{2}\left[J_{2}\left(p\right) - J_{2}\left(q - p\right)\right]$$

$$p^{\mu_{1}}J_{3\mu_{1}\mu_{2}} = -\frac{1}{2}p^{2}J_{3\mu_{2}} + \frac{1}{2}\left[J_{2\mu_{2}}\left(q\right) + J_{2\mu_{2}}\left(q - p\right) + q_{\mu_{2}}J_{2}\left(q - p\right)\right]$$

$$q^{\mu_{1}}J_{3\mu_{1}\mu_{2}} = -\frac{1}{2}q^{2}J_{3\mu_{2}} + \frac{1}{2}\left[J_{2\mu_{2}}\left(p\right) + J_{2\mu_{2}}\left(q - p\right) + q_{\mu_{2}}J_{2}\left(q - p\right)\right]$$

$$g^{\mu_{1}\mu_{2}}J_{3\mu_{1}\mu_{2}} = m^{2}J_{3} + \frac{i}{2\left(4\pi\right)^{2}} + J_{2}\left(q - p\right)$$
(C2)

Appendix D: The Integrals in Six Dimensions

Three Point Functions

$$J_3(p,q) = i (4\pi)^{-3} \left[-Z_{00}^{(0)}(p,q) \right]; \qquad \bar{J}_3 = I_{\log}^{(6)} + J_3$$

$$J_{3\mu_{1}}(k_{1},k_{2},k_{3}) = i (4\pi)^{-3} \left[p_{21\mu_{1}} Z_{10}^{(0)} + p_{31\mu_{1}} Z_{01}^{(0)} \right]$$

$$\bar{J}_{3}^{\mu_{1}}(k_{1},k_{2},k_{3}) = -\frac{1}{3} \left(k_{1}^{\nu_{1}} + k_{2}^{\nu_{1}} + k_{3}^{\nu_{1}} \right) \Delta_{4\nu_{1}}^{(6)\mu_{1}} - \frac{1}{3} \left(p_{21}^{\mu_{1}} + p_{31}^{\mu_{1}} \right) I_{\log}^{(6)} + J_{3}^{\mu_{1}},$$

Four Point Functions

$$J_{4} = i (4\pi)^{-3} \left[Z_{000}^{(-1)} (p,q,r) \right]$$

$$J_{4\mu_{1}} = i (4\pi)^{-3} \left[-p_{\mu_{1}} Z_{100}^{(-1)} - q_{\mu_{1}} Z_{010}^{(-1)} - r_{\mu_{1}} Z_{001}^{(-1)} \right]$$

$$J_{4\mu_{1}\mu_{2}} = i (4\pi)^{-3} \left[-\frac{1}{2} g_{\mu_{1}\mu_{2}} Z_{000}^{(0)} + p_{\mu_{1}} p_{\mu_{2}} Z_{200}^{(-1)} + q_{\mu_{1}} q_{\mu_{2}} Z_{020}^{(-1)} + r_{\mu_{1}} r_{\mu_{2}} Z_{002}^{(-1)} \right] + i (4\pi)^{-3} \left[(p_{\mu_{1}} q_{\mu_{2}} + p_{\mu_{2}} q_{\mu_{1}}) Z_{110}^{(-1)} + (p_{\mu_{1}} r_{\mu_{2}} + p_{\mu_{2}} r_{\mu_{1}}) Z_{101}^{(-1)} + (q_{\mu_{1}} r_{\mu_{2}} + r_{\mu_{1}} q_{\mu_{2}}) Z_{011}^{(-1)} \right] \bar{J}_{4\mu_{1}\mu_{2}} = J_{4\mu_{1}\mu_{2}} + \frac{1}{6} (\Delta_{4\mu_{1}\mu_{2}}^{(6)} + g_{\mu_{1}\mu_{2}} I_{\log}^{(6)})$$

Reductions

$$2\left[p^{2}Z_{100}^{(-1)} + (p \cdot q) Z_{010}^{(-1)} + (p \cdot r) Z_{001}^{(-1)}\right] = p^{2}Z_{000}^{(-1)} + Z_{00}^{(0)}(q,r) - Z_{00}^{(0)}(p_{42},p_{43})$$

$$2\left[q^{2}Z_{010}^{(-1)} + (p \cdot q) Z_{100}^{(-1)} + (q \cdot r) Z_{001}^{(-1)}\right] = q^{2}Z_{000}^{(-1)} + Z_{00}^{(0)}(p,r) - Z_{00}^{(0)}(p_{42},p_{43})$$

$$2\left[r^{2}Z_{001}^{(-1)} + (p \cdot r) Z_{100}^{(-1)} + (q \cdot r) Z_{010}^{(-1)}\right] = r^{2}Z_{000}^{(-1)} + Z_{00}^{(0)}(p,q) - Z_{00}^{(0)}(p_{42},p_{43})$$

$$2\left[p^{2}Z_{200}^{(-1)} + (p \cdot q)Z_{110}^{(-1)} + (p \cdot r)Z_{101}^{(-1)}\right] = p^{2}Z_{100}^{(-1)} + Z_{000}^{(0)} - Z_{10}^{(0)}(p_{42}, p_{43})$$

$$2\left[p^{2}Z_{110}^{(-1)} + (p \cdot q)Z_{020}^{(-1)} + (p \cdot r)Z_{011}^{(-1)}\right] = p^{2}Z_{010}^{(-1)} + Z_{10}^{(0)}(p_{31}, p_{41}) - Z_{01}^{(0)}(p_{42}, p_{43})$$

$$2\left[p^{2}Z_{101}^{(-1)} + (p \cdot q)Z_{011}^{(-1)} + (p \cdot r)Z_{002}^{(-1)}\right] = p^{2}Z_{001}^{(-1)} + Z_{01}^{(0)}(q, r)$$

$$- \left[Z_{00}^{(0)} - Z_{10}^{(0)} - Z_{01}^{(0)}\right](p_{42}, p_{43})$$

$$2\left[q^{2}Z_{020}^{(-1)} + (p \cdot q)Z_{110}^{(-1)} + (q \cdot r)Z_{011}^{(-1)}\right] = q^{2}Z_{010}^{(-1)} + Z_{000}^{(0)} - Z_{01}^{(0)}(p_{42}, p_{43})$$

$$2\left[q^{2}Z_{110}^{(-1)} + (p \cdot q)Z_{200}^{(-1)} + (q \cdot r)Z_{101}^{(-1)}\right] = q^{2}Z_{100}^{(-1)} + Z_{10}^{(0)}(p,r) - Z_{10}^{(0)}(p_{42}, p_{43})$$

$$2\left[q^{2}Z_{011}^{(-1)} + (p \cdot q)Z_{101}^{(-1)} + (q \cdot r)Z_{002}^{(-1)}\right] = q^{2}Z_{001}^{(-1)} + Z_{01}^{(0)}(p,r) - Z_{10}^{(0)}(p_{42}, p_{43})$$

$$- \left[Z_{00}^{(0)} - Z_{10}^{(0)} - Z_{01}^{(0)}\right](p_{42}, p_{43})$$

$$2\left[r^{2}Z_{002}^{(-1)} + (q \cdot r) Z_{011}^{(-1)} + (p \cdot r) Z_{101}^{(-1)}\right] = r^{2}Z_{001}^{(-1)} + Z_{000}^{(0)} - \left[Z_{00}^{(0)} - Z_{10}^{(0)} - Z_{01}^{(0)}\right] (p_{42}, p_{43}) 2\left[r^{2}Z_{011}^{(-1)} + (p \cdot r) Z_{110}^{(-1)} + (q \cdot r) Z_{020}^{(-1)}\right] = r^{2}Z_{010}^{(-1)} + Z_{01}^{(0)} (p, q) - Z_{01}^{(0)} (p_{42}, p_{43}) 2\left[r^{2}Z_{101}^{(-1)} + (p \cdot r) Z_{200}^{(-1)} + (q \cdot r) Z_{110}^{(-1)}\right] = r^{2}Z_{100}^{(-1)} + Z_{10}^{(0)} (p, q) - Z_{10}^{(0)} (p_{42}, p_{43}) -3Z_{000}^{(0)} = +\frac{1}{3} + 2m^{2}Z_{000}^{(-1)} - \left[p^{2}Z_{100}^{(-1)} + q^{2}Z_{010}^{(-1)} + r^{2}Z_{001}^{(-1)}\right] - Z_{00}^{(0)} (p_{42}, p_{43})$$

they imply the relations

$$2p^{\mu_1}J_{4\mu_1} = -p^2J_4 + J_3(q,r) - J_3(r-p,r-q)$$

$$2q^{\mu_1}J_{4\mu_1} = -q^2J_4 + J_3(p,r) - J_3(r-p,r-q)$$

$$2r^{\mu_1}J_{4\mu_1} = -r^2J_4 + J_3(p,q) - J_3(r-p,r-q)$$

 $2p^{\mu_1}J_{4\mu_1\mu_2} = -p^2 J_{4\mu_2} + J_{3\mu_2} (p_{42}, p_{43}) + J_{3\mu_2} (p_{31}, p_{41}) + p_{41\mu_2} J_3 (p_{42}, p_{43})$ $2q^{\mu_1}J_{4\mu_1\mu_2} = -q^2 J_{4\mu_2} + J_{3\mu_2} (p_{42}, p_{43}) + J_{3\mu_2} (p_{21}, p_{41}) + p_{41\mu_2} J_3 (p_{42}, p_{43})$ $2r^{\mu_1}J_{4\mu_1\mu_2} = -r^2 J_{4\mu_2} + J_{3\mu_2} (p_{42}, p_{43}) + J_{3\mu_2} (p_{21}, p_{31}) + p_{41\mu_2} J_3 (p_{42}, p_{43})$

$$2g^{\mu_{12}}J_{4\mu_{1}\mu_{2}} = \frac{i}{3(4\pi)^{3}} + 2m^{2}J_{4} + 2J_{3}(p_{42}, p_{43})$$

$$p = p_{21}; \quad q = p_{31}; \quad r = p_{41}, \text{ in the computations}$$

Appendix E: Subamplitudes

The SAP subamplitude Integrand

$$S_{ij} = \left(K_i \cdot K_j - m^2\right)$$

$$\left(t^{SAP}\right)^{\nu_1} = 4 \left[K_1^{\nu_1} S_{23} + K_2^{\nu_1} \left(S_{13} + 2m^2\right) - K_3^{\nu_1} \left(S_{12} + 2m^2\right)\right] \frac{1}{D_{123}}$$

$$(T^{SAP})^{\nu_1} = -4 (p_{32} \cdot p_{31}) J_3^{\nu_1} + 2 [(p_{31}^{\nu_1} p_{21}^2 - p_{21}^{\nu_1} p_{31}^2 - 4m^2 p_{32}^{\nu_1}) J_3 - p_{32}^{\nu_1} J_2 (p_{32}) - p_{31}^{\nu_1} J_2 (p_{31})] - 2 [P_{21}^{\nu_2} \Delta_{3\nu_2}^{\nu_1} + (p_{32}^{\nu_1} + p_{31}^{\nu_1}) I_{\log}]$$

The SPA subamplitude

$$(T^{SPA})^{\nu_1} = +4 (p_{21} \cdot p_{31}) J_3^{\nu_1} + 2 [(p_{31}^{\nu_1} p_{21}^2 + p_{21}^{\nu_1} p_{31}^2 - 4m^2 p_{21}^{\nu_1}) J_3 - p_{21}^{\nu_1} J_2 (p_{21}) - p_{31}^{\nu_1} J_2 (p_{31})] + 2 [P_{32}^{\nu_2} \Delta_{\nu_2}^{\nu_1} - (p_{21}^{\nu_1} + p_{31}^{\nu_1}) I_{\log}]$$

The ASP subamplitude

$$(T^{ASP})^{\nu_1} = -4 (p_{21} \cdot p_{32}) J_3^{\nu_1} + 2 [(p_{31}^{\nu_1} p_{21}^2 - p_{21}^{\nu_1} p_{31}^2 - 4m^2 p_{32}^{\nu_1}) J_3 + p_{21}^{\nu_1} J_2 (p_{21}) - p_{32}^{\nu_1} J_2 (p_{32})] + 2 [P_{31}^{\nu_2} \Delta_{\nu_2}^{\nu_1} + (p_{21}^{\nu_1} - p_{32}^{\nu_1}) I_{\log}]$$

The PVP subamplitude

$$(T^{PVP})^{\nu_1} = +4 (p_{32} \cdot p_{31}) J_3^{\nu_1} + 2 [(p_{21}^{\nu_1} p_{31}^2 - p_{31}^{\nu_1} p_{21}^2) J_3 + p_{32}^{\nu_1} J_2 (p_{32}) + p_{31}^{\nu_1} J_2 (p_{31})] + 2 [P_{21}^{\nu_2} \Delta_{3\nu_2}^{\nu_1} + (p_{32}^{\nu_1} + p_{31}^{\nu_1}) I_{\log}]$$

The *PSA* subamplitude

$$(T^{PSA})^{\nu_1} = -4 (p_{21} \cdot p_{31}) J_3^{\nu_1} - 2 \left[\left(p_{21}^{\nu_1} p_{31}^2 + p_{31}^{\nu_1} p_{21}^2 - 4m^2 p_{31}^{\nu_1} \right) J_3 - p_{21}^{\nu_1} J_2 (p_{21}) - p_{31}^{\nu_1} J_2 (p_{31}) \right] + 2 \left[-P_{32}^{\nu_2} \Delta_{3\nu_2}^{\nu_1} + \left(p_{21}^{\nu_1} + p_{31}^{\nu_1} \right) I_{\log} \right]$$

The APS subamplitude

$$(T^{APS})^{\nu_1} = +4 (p_{21} \cdot p_{32}) J_3^{\nu_1} + 2 [(p_{21}^{\nu_1} p_{31}^2 - p_{31}^{\nu_1} p_{21}^2 - 4m^2 p_{21}^{\nu_1}) J_3 + p_{32}^{\nu_1} J_2 (p_{32}) - p_{21}^{\nu_1} J_2 (p_{21})] - 2 [P_{31}^{\nu_2} \Delta_{3\nu_2}^{\nu_1} - (p_{32}^{\nu_1} - p_{21}^{\nu_1}) I_{\log}]$$

The PAS subamplitude

$$(T^{PAS})^{\nu_1} = +4 (p_{32} \cdot p_{31}) J_3^{\nu_1} + 2 [(p_{21}^{\nu_1} p_{31}^2 - p_{31}^{\nu_1} p_{21}^2 + 4m^2 p_{31}^{\nu_1}) J_3 + p_{32}^{\nu_1} J_2 (p_{32}) + p_{31}^{\nu_1} J_2 (p_{31})] + 2 [P_{21}^{\nu_2} \Delta_{3\nu_2}^{\nu_1} + (p_{32}^{\nu_1} + p_{31}^{\nu_1}) I_{\log}]$$

The PPV subamplitude

$$(T^{PPV})^{\nu_1} = +4 (p_{21} \cdot p_{31}) J_3^{\nu_1} + 2 [(p_{31}^{\nu_1} p_{21}^2 + p_{21}^{\nu_1} p_{31}^2) J_3 - [p_{21}^{\nu_1} J_2 (p_{21}) + p_{31}^{\nu_1} J_2 (p_{31})]] + 2 [P_{32}^{\nu_2} \Delta_{\nu_2}^{\nu_1} - (p_{21}^{\nu_1} + p_{31}^{\nu_1}) I_{\log}]$$

The VSS subamplitude

$$(T^{VSS})^{\nu_1} = 4 \left[(p_{21} \cdot p_{32}) + 4m^2 \right] J_3^{\nu_1} + 2 \left(p_{21}^{\nu_1} p_{31}^2 - p_{31}^{\nu_1} p_{21}^2 + 4m^2 p_{31}^{\nu_1} \right) J_3 + 2 \left[p_{32}^{\nu_1} J_2 \left(p_{32} \right) - p_{21}^{\nu_1} J_2 \left(p_{21} \right) \right] - 2 \left[P_{31}^{\nu_2} \Delta_{\nu_2}^{\nu_1} - \left(p_{32}^{\nu_1} - p_{21}^{\nu_1} \right) I_{\log} \right]$$

The SVS subamplitude

$$(T^{SVS})^{\nu_1} = 4 \left[-(p_{32} \cdot p_{31}) + 4m^2 \right] J_3^{\nu_1} + 2 \left(p_{31}^{\nu_1} p_{21}^2 - p_{21}^{\nu_1} p_{31}^2 + 4m^2 p_{21}^{\nu_1} \right) J_3 - 2 \left[p_{32}^{\nu_1} J_2 \left(p_{32} \right) + p_{31}^{\nu_1} J_2 \left(p_{31} \right) \right] - 2 \left[P_{21}^{\nu_2} \Delta_{\nu_2}^{\nu_1} + \left(p_{32}^{\nu_1} + p_{31}^{\nu_1} \right) I_{\log} \right]$$

The SSV subamplitude

$$(T^{SSV})^{\nu_1} = +4 \left[-(p_{21} \cdot p_{31}) + 4m^2 \right] J_3^{\nu_1} + 2 \left[\left(-p_{31}^{\nu_1} p_{21}^2 - p_{21}^{\nu_1} p_{31}^2 + 4m^2 \left(p_{21}^{\nu_1} + p_{31}^{\nu_1} \right) \right) J_3 \right] + 2 \left[p_{21}^{\nu_1} J_2 \left(p_{21} \right) + p_{31}^{\nu_1} J_2 \left(p_{31} \right) \right] - 2 \left[P_{32}^{\nu_2} \Delta_{\nu_2}^{\nu_1} - \left(p_{21}^{\nu_1} + p_{31}^{\nu_1} \right) I_{\log} \right]$$

Acknowledgments

The authors would like to thank Sebastião A. Dias for helpful suggestions and support in this investigation. L. Ebani, T. J. Girardi and J.F. Thuorst, acknowledge the financial support of CAPES and CNPQ.

[1] Fukuda, H.; Miyamoto, Y. On the γ-decay of neutral meson. Prog. Theor. Phys. 1949, 4, 347-357.

- [2] Steinberger, J. On the Use of Subtraction Fields and the Lifetimes of Some Types of Meson Decay. *Phys. Rev.* 1949, 77, 1180-1186.
- [3] Schwinger, J. On Gauge Invariance and Vacuum Polarization. *Phys. Rev.* **1951**, 82, 664-679.
- [4] Rosenberg, L. Electromagnetic Interactions of Neutrinos. Phys. Rev. 1963, 129, 2786.
- [5] Johnson, K. γ_5 Invariance. *Phys. Lett.* **1963**, 5, 253.
- [6] Adler, S. L. Axial-Vector vertex in spinor electrodynamics. Phys. Rev. 1969, 177, 2426-2438.
- [7] Bardeen, W. A. Anomalous Ward identities in spinor field theories. Phys. Rev. 1969, 184, 1848-1857.
- [8] Bell, J. S.; Jackiw, R. A PCAC puzzle: $\pi^0 \to \gamma\gamma$ in the σ -model. Nuovo Cim. A **1969**, 60, 47-61.
- [9] Bertlmann, R. A. Anomalies in Quantum Field Theory, 1st ed.; Oxford University Press Inc: New York, United States, 1996.
- [10] Bastianelli, F.; Van Nieuwenhuizen, P. Path Integrals and Anomalies in Curved Space, 1st ed.; Cambridge University Press: Cambridge: Great Britain, 2006. doi:10.1017/CBO9780511535031
- [11] Vassilevich, D.V. Heat kernel expansion: user's manual. *Physics Reports* **2003**, 388, 279-360.
- [12] Bollini, C. G.; Giambiagi, J. J. Dimensional renormalization: The number of dimensions as a regularizing parameter. *Phys. Lett. B* 1972, 40, 566.
- [13] 't Hooft, G.; Veltman, M. Regularization and renormalization of gauge fields. Nucl. Phys. B 1972, 44, 189.
- [14] Pittau, R. A four-dimensional approach to quantum field theories. JHEP 2012, 11, 024.
- [15] Gnendiger, C.; Signer, A.; Stöckinger, D.; Broggio, A.; Cherchiglia, A. L.; Driencourt-Mangin, F.; Fazio, A. R.; Hiller, B.; Mastrolia, P.; Peraro, T.; Pittau, R.; Pruna, G. M.; Rodrigo, G.; Sampaio, M.; Sborlini, G.; Bobadilla, W. J. T.; Tramontano, F.; Ulrich, Y.; Visconti, A. To d, or not to d: recent developments and comparisons of regularization schemes. *Eur. Phys. J. C* **2017**, 77, 471.
- [16] Breitenlohner, P.; Maison D. Dimensional renormalization and the action Principle. Commun. math. Phys. 1977, 52, 11–38.
- [17] Jegerlehner, F. Facts of life with γ_5 . Eur. Phys. J. C 2001, 18, 673-679.
- [18] Tsai, Er-C. Gauge invariant treatment of γ_5 in the scheme of 't Hooft and Veltman. *Phys. Rev. D* **2011**, 83, 025020.
- [19] Tsai, Er-C. Maintaining gauge symmetry in renormalizing chiral gauge theories. *Phys. Rev. D* 2011, 83, 065011.
- [20] Ferrari, R. Managing γ_5 in Dimensional Regularization II: the Trace with more γ_5 's. Int. J. Theor. Phys. 2017, 56, 691-705.
- [21] Bruque, A. M.; Cherchiglia, A. L.; Pérez-Victoria, M. Dimensional regularization vs methods in fixed dimension with and without γ_5 . *JHEP* **2018**, 08 109.
- [22] Treiman, S. B.; Jackiw, R.; Zumino, B.; Witten, E. Current algebra and anomalies. Princeton University Press: Princeton, New Jersey, 1985.
- [23] Cheng, T.P.; Li, L.F. Gauge theory of elementary particle physics, 1st ed.; Oxford University Press: Oxford, UK, 1984.
- [24] Sterman, G. An Introduction to Quantum Field Theory, 1st ed.; Cambridge University Press: Cambridge, Great Britain, 1993; pp. 94–98.
- [25] Battistel, O. A. A new strategy for manipulations e calculations involving divergences in QFT. Ph.D. Thesis, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil, 1999.
- [26] Battistel, O. A.; Fonseca, M. V. S.; Dallabona, G. Anomalies in finite amplitudes: Two-dimensional single axial-vector triangle. *Phys. Rev. D* 2012, 85, 085007.
- [27] Battistel, O. A.; Traboussy, F.; Dallabona, G. Anomalies in finite amplitudes: Two-dimensional single and triple axial-vector triangles. Int. J. Mod. Phys. A 2018 33, 1850136.
- [28] Battistel, O.A.; Dallabona, G.; Fonseca, M.V.; Ebani, L. Can Really Regularized Amplitudes Be Obtained as Consistent with Their Expected Symmetry Properties? 2018, *Journal of Modern Physics*, 9, 1153-1178.
- [29] Battistel, O. A.; Dallabona, G. From arbitrariness to ambiguities in the evaluation of perturbative physical amplitudes and their symmetry relations. *Phys. Rev. D* **2002**, 65, 125017.
- [30] Battistel, O. A.; Dallabona, G. Anomalies dismissed of ambiguities and the neutral pion decay. J. Phys. G: Nucl. Part. Phys. 2002, 28, 2539.
- [31] Fonseca, M. V. S.; Dallabona, G.; Battistel, O. A. Perturbative calculations in space time having extra dimensions: The 6D single axial box anomaly. Int. J. Mod. Phys. A 2014, 29, 1450168.
- [32] Fonseca, M. V. S.; Girardi, T. J.; Dallabona, G.; Battistel, O. A. Ambiguities and symmetry relations

in five-dimensional perturbative calculations: The explicit evaluation of the QED₅ vacuum polarization tensor. Int. J. Mod. Phys. A **2013**, 28, 1350135-1350160.

- [33] Battistel, O. A.; Dallabona, G. Consistency and universality in odd and even dimensional space time QFT perturbative calculations. Int. J. Mod. Phys. A 2014, 29, 1450068.
- [34] Viglioni, A. C. D.; Cherchiglia, A. L.; Vieira, A. R.; Hiller, B.; Sampaio, M. γ_5 algebra ambiguities in Feynman amplitudes: Momentum routing invariance and anomalies in D = 4 and D = 2. *Phys. Rev.* D **2016**, 94, 065023.
- [35] Vieira, A. R.; Cherchiglia, A. L.; Sampaio, M. Momentum routing invariance in extended QED: Assuring gauge invariance beyond tree level. *Phys. Rev. D* **2016**, 93, 025029.
- [36] Ferreira, L. C.; Cherchiglia, A. L.; Hiller, B.; Sampaio, M.; Nemes, M. C. Momentum routing invariance in Feynman diagrams and quantum symmetry breakings, *Phys. Rev. D* 2012, 86, 025016.
- [37] Battistel, O. A.; Dallabona, G. A systematization for one-loop 4D Feynman integrals. Eur. Phys. J. C 2006, 45, 721.
- [38] Battistel, O. A.; Dallabona, G. A Systematization for One-Loop 4D Feynman Integrals-Different Species of Massive Fields. *Journal of Modern Physics*, **2012**, 3, 1408-1449.
- [39] Sun, Y.; Chang H-R. One loop integrals reduction. *Chinese Physics C* 2012, 36, 1055-1064.
- [40] Wit, B. de.; Smith, J. Field theory in particle physics. Volume 1. 1st ed.; North-Holland Physics Publishing: Amsterdam: The Netherlands, 1986; pp. 457–466.
- [41] Scharf, G. Finite Quantum Electrodynamics: The Causal Approach, 3rd ed.; Dover Publications Inc: Mineloa, New York, 2014.
- [42] Aste, A.; Arx, C. von; Scharf, G. Regularization in quantum field theory from the causal point of view. Progress in Particle and Nuclear Physics. 2010, 64, 61–119.
- [43] Epstein, H.; Glaser, V. The role of locality in perturbation theory. Ann. IHP, Phys. théor. 1973, 19, 211-295.
- [44] Aste, A. Two-Loop Diagrams in Causal Perturbation Theory. Ann. Phys. (N. Y.) 1997, 257, 158–204.
- [45] Aste, A.; Trautmann, D. Finite calculation of divergent self-energy diagrams. Can. J. Phys. 2003, 8, 1433–1445.
- [46] Battistel, O. A.; Mota, A. L.; Nemes, M. C. Consistency Conditions for 4-D Regularizations. Mod. Phys.Lett. A 1998, 13, 1597-1610.
- [47] Battistel, O. A.; Nemes, M. C. Consistency in regularizations of the gauged NJL model at the one loop level. *Phys. Rev. D* 1999, 59, 055010.
- [48] Battistel, O. A. From arbitrariness to anomalies in two-dimensional perturbative calculations. J. Phys. G: Nucl. Part. Phys. 2004,30, 543–564
- [49] Ma, Y-L.; Wu, Y-L.; Anomaly and Anomaly-Free Treatment of QFT's Based on Symmetry-Preserving Loop Regularization. Int. J. Mod. Phys. A 2006, 21, 6383-6456.
- [50] Cynolter, G.; Lendvai, E. Note on triangle anomaly with improved momentum cutoff. Mod. Phys. Lett. A 2011, 26,1537-1545
- [51] Elias, V.; McKeon, G.; Mann, R. B. VVA Triangle graph ambiguities in four and N dimensions. Nucl. Phys. B 1983, 229, 487-498.
- [52] Chowdhury, A. M.; McKeon, D. G. C.; Mann, R. B. Ambiguities of the chiral-anomaly graph in higher dimensions. *Phys. Rev. D* 1986, 33, 3090.
- [53] Aguila, F. del.; Pérez-Victoria, M. Differential Renormalization of Gauge Theories. Acta Physica Polonica B. 1998, 28, 2857-2863.