Yuki Sato

Abstract Two-dimensional causal dynamical triangulations (2d CDT) is a lattice model of quantum geometry. In 2d CDT, one can deal with the quantum effects analytically and explore the physics through the continuum limit. The continuum theory is known to be two-dimensional projectable Hořava-Lifshitz quantum gravity (2d projectable HL QG). In this chapter, we wish to review the very relation between 2d CDT and 2d projectable HL QG in detail.

Keywords

Lattice quantum gravity; Causal dynamical triangulations (CDT); Hořava-Lifshitz quantum gravity; Two-dimensional models; Wormholes

Note This is a preprint of the chapter in the Handbook of Quantum Gravity. It belongs to the section entitled Causal Dynamical Triangulations.

Yuki Sato

National Institute of Technology, Tokuyama College, Gakuendai, Shunan, Yamaguchi 745-8585, Japan, e-mail: sato@tokuyama.ac.jp

Nagoya University, Chikusaku, Nagoya 464-8602, Japan, e-mail: ysato.phys@gmail.com

1 Introduction

Two-dimensional toy models of quantum gravity are a very useful playground for understanding the quantum nature of geometry quantitatively. This is because many of them are simple enough to be dealt with analytically and complex enough to observe nontrivial quantum effects. Lattice regularizations in particular are known to be quite powerful tools for investigating non-perturbative quantum effects analytically. Among these, two-dimensional Euclidean dynamical triangulations (2d EDT) [1–6] (see a pedagogical textbook [7]) and two-dimensional causal dynamical triangulations (2d CDT) [9] (see a detailed review [8]) are good practical examples. The former and the latter, respectively, are Euclidean and Lorentzian lattice models based on Regge's discretization of geometries [10].

2d EDT discretizes Euclidean geometries by equilateral triangles and defines a regularized quantum amplitude as a sum over distinct triangulated geometries. Matrix models and combinatorics can be used for calculating such a statistical sum analytically (whenever possible). By virtue of analytic treatments, one can explicitly remove the regularization through the continuum limit to calculate physical observables. What is remarkable is that exactly the same value of observables can be reproduced from a genuine continuum field theory called the Liouville quantum gravity [11–14]. This means that 2d EDT serves as a well-defined regularization of the Liouville quantum gravity.

2d CDT is a Lorentzian lattice model of quantum geometry which respects a global time foliation and prohibits the creation of so-called baby universes. One can calculate the sum over such Lorentzian triangulated geometries using simple combinatorics, and take the continuum limit to remove the regularization. All these processes can be performed analytically at least for the plain model without coupling to a matter. It has been shown in Ref. [15] that the resulting continuum theory is known to be in the same universality class of projectable Hořava-Lifshitz quantum gravity (projectable HL QG) [16] in two dimensions, which is different from the Liouville quantum gravity ¹.

2d HL QG is a quantum field theory in two dimensions, which has a preferred foliation structure. This model is invariant only under the subclass of diffeomorphisms that respects the foliation, known as the foliation-preserving diffeomorphisms². 2d projectable HL QG is a certain version of HL QG where a part of the time-time component of the metric called the lapse function is projectable, i.e. a function only

¹ In fact, it has been shown that a direct lattice discretization of 2d projectable HL gravity, which has a lattice action different from that of 2d CDT, reproduces the same large-scale physics in the continuum limit [18].

² At the cost of full diffeomorphism invariance, HL QG has been designed originally as a model of quantum gravity in higher dimensions such that it has a good convergence at UV in keeping with unitarity, and would approximately recover the diffeomorphism invariance at IR [16]

of time. In this chapter, we wish to explain in detail the relation between projectable HL QG and CDT in two dimensions³.

In fact one can generalize 2d CDT in such a way that the creation and annihilation of (a finite number of) baby universes, and the formation of wormholes (handles) are allowed to occur in keeping with the foliation structure. This model is called the generalized CDT (GCDT) introduced first as a continuum theory [21,22] and later defined at the discrete level [23,24]. The full continuum description of GCDT is given by the so-called string field theory for CDT [22] in which the string means the one-dimensional closed spatial universe, and the baby universes and wormholes can be realized in terms of the splitting and joining interactions of strings.

One of remarkable facts is that focusing on a certain amplitude, i.e. loop-to-loop amplitude, one can read off a one-dimensional effective theory that takes in all possible baby universe and wormhole contributions in an effective manner [25, 26]: The 1d effective theory is a one-body quantum theory even though GCDT is a many-body theory that allows both creation and annihilation of strings. It is known that one can correctly reproduce the 1d effective theory if quantizing the projectable HL gravity with a certain bi-local interaction term [28, 29]. This topic will be treated in this chapter.

Furthermore, the 1d effective theory that includes all contributions of baby universes and wormholes is known to be reproduced if one assumes that the cosmological constant of the continuum limit of 2d CDT is not really a constant but fluctuates in time [30]. This idea leads to a certain realization of Coleman's mechanism about the cosmological constant [27] in the context of CDT [30], which will be also explained in this chapter.

The rest of this chapter is organized as follows. In Sec. 2, a self-contained introduction to 2d CDT is presented. We show that taking the continuum limit the physics of 2d CDT can be described as a one-dimensional quantum system with a Hamiltonian. 2d projectable HL QG is explained in Sec. 3. Through the pathintegral quantization we read off the quantum Hamiltonian that is equivalent to the one obtained in the continuum limit of 2d CDT. Thereby one can confirm that the continuum limit of 2d CDT is 2d projectable HL QG. In Sec. 4, we introduce GCDT that is a generalization of 2d CDT such that baby universes and wormholes are introduced so as to be compatible with the foliation, and determine the 1d effective theory obtained through the sum over all genera. In particular, we explain in detail that quantizing 2d projectable HL gravity with a bi-local interaction yields the 1d effective theory, and discuss Coleman's mechanism in 2d CDT. Sec. 5 is devoted to summary.

³ The relation between HL QG and CDT in four dimension has been pointed out first in Ref. [19] by looking at an observable called the spectral dimension, and the resemblance of the CDT phase diagram to a Lifshitz phase diagram has been shown in Ref. [20].

2 2d causal dynamical triangulations

Two-dimensional causal dynamical triangulations (2d CDT) [9] is a lattice model of quantum geometries based on Regge's discretization [10]. In this section, we give an overview of 2d CDT, and in particular explain how to obtain the quantum Hamiltonian through the continuum limit.

We start with a two-dimensional globally hyperbolic manifold equipped with a global time foliation:

$$\mathcal{M} = \bigcup_{t \in \mathbb{R}} \Sigma_t , \qquad (1)$$

where each leaf Σ_t is a one-dimensional Cauchy "surface" (line). One approximates the manifold with a foliation in such a way that the continuous label *t* is discretized by integers, i.e. $t \in \mathbb{Z}$; each leaf (line) is partitioned by vertices connected by isometric edges; vertices among neighboring time steps are connected by isometric edges to form a triangulation of strip (see Fig. 1). The edges at a given time step and those



Fig. 1 A triangulation of a strip: Thick and thin lines are space-like and time-like edges, respectively.

connecting vertices in different time steps are, respectively, space-like and time-like edges since the squared edge lengths of the space-like edge a_s^2 and the time-like edge a_t^2 are given by

$$a_s^2 = \varepsilon^2 , \quad a_t^2 = -\alpha \varepsilon^2 ,$$
 (2)

where α is a positive number and ε is a lattice spacing that serves as a UV cutoff.

2d CDT deals with a set of restricted class of Lorentzian triangulations as discussed above. In particular, we consider that the topology of the one-dimensional universe (a graph consisting of vertices and edges at a given time) is either S^1 or [0, 1], and the topology will not change during (discrete) time propagation. Since the topology is fixed the curvature term plays no role in two dimensions and only the discrete analogue of the cosmological constant term, $\Lambda_0 \int d^2x \sqrt{-g}$ is used as the lattice action of 2d CDT:

$$S_T[\lambda, \alpha] = -\frac{\lambda}{\varepsilon^2} \left(\frac{\sqrt{4\alpha + 1}}{4} \varepsilon^2 n(T) \right), \qquad (3)$$

where λ/ϵ^2 is the bare cosmological constant with the dimensionless number λ , n(T) the number of triangles in a triangulation T, and the term inside the parentheses denotes the total area of the triangulation. It is useful to rotate to the Euclidean signature which can be performed by changing $\alpha \rightarrow -\alpha - i0$. Accordingly the lattice action (3) changes as follows:

$$iS_T[\lambda,\alpha] \to iS_T[\lambda,-\alpha-i0] = -\lambda \frac{\sqrt{4\alpha-1}}{4} n(T) \equiv -\lambda n(T) ,$$
 (4)

where α is chosen to be greater than 1/4; otherwise, the triangle inequalities will not be satisfied after the rotation. In any case we have absorbed the parameter α by the redefinition of the dimensionless cosmological constant λ .

The amplitude of the one-dimensional universe that starts with ℓ_1 edges and ends up with ℓ_2 edges after the discrete time step t is given by the sum over all allowed triangulations:

$$G_{\lambda}^{(a)}(\ell_1, \ell_2; t) = \sum_{T \in \mathcal{T}^{(a)}(\ell_1, \ell_2, t)} e^{-\lambda n(T)} = \sum_n e^{-\lambda n} \mathcal{N}^{(a)}(\ell_1, \ell_2, n) , \qquad (5)$$

where $\mathcal{T}^{(a)}$ is a set of triangulations whose topology is $[0, 1] \times [0, 1]$ for a = 0 and $S^1 \times [0, 1]$ for a = 1, and

$$\mathcal{N}^{(a)}(\ell_1, \ell_2, n) = \# \left\{ T \in \mathcal{T}^{(a)} \mid n(T) = n \right\} .$$
(6)

When defining the amplitude (5), we do not allow the one-dimensional universe to vanish during the discrete time propagation. For later convenience, we introduce a marked amplitude:

$$G_{\lambda}^{(-1)}(\ell_1, \ell_2; t) = \ell_1 G_{\lambda}^{(1)}(\ell_1, \ell_2; t) , \qquad (7)$$

where one of the edges in the initial one-dimensional universe is marked. This is because there exist ℓ_1 possible ways of marking the edges. The three kinds of amplitude should satisfy the composition law:

$$G_{\lambda}^{(1)}(\ell_1, \ell_2; t_1 + t_2) = \sum_{\ell=1}^{\infty} G_{\lambda}^{(1)}(\ell_1, \ell; t_1) \ \ell \ G_{\lambda}^{(1)}(\ell, \ell_2; t_2) , \qquad (8)$$

$$G_{\lambda}^{(0)}(\ell_1, \ell_2; t_1 + t_2) = \sum_{\ell=1}^{\infty} G_{\lambda}^{(0)}(\ell_1, \ell; t_1) \ G_{\lambda}^{(0)}(\ell, \ell_2; t_2) , \qquad (9)$$

$$G_{\lambda}^{(-1)}(\ell_1, \ell_2; t_1 + t_2) = \sum_{\ell=1}^{\infty} G_{\lambda}^{(-1)}(\ell_1, \ell; t_1) \ G_{\lambda}^{(-1)}(\ell, \ell_2; t_2) , \qquad (10)$$

where for a = 1 one needs to multiply the amputated amplitudes by ℓ since there exist ℓ possible ways of gluing to recover the whole amplitude.

It is convenient to introduce the generating function of the number of triangulations. Using the notation

$$g = e^{-\lambda} , \qquad (11)$$

we define the generating function:

$$\widetilde{G}^{(a)}(g, x, y; t) = \sum_{\ell_1=1}^{\infty} \sum_{\ell_2=1}^{\infty} x^{\ell_1} y^{\ell_2} G_{\lambda}^{(a)}(\ell_1, \ell_2; t)$$
$$= \sum_{\ell_1=1}^{\infty} \sum_{\ell_2=1}^{\infty} \sum_{n=1}^{\infty} x^{\ell_1} y^{\ell_2} g^n \mathcal{N}^{(a)}(\ell_1, \ell_2, n) , \qquad (12)$$

where in the context of quantum gravity, x and y are related to the boundary cosmological constants, λ_1 and λ_2 , that control the size of the boundaries:

$$x = e^{-\lambda_1}, \quad y = e^{-\lambda_2}. \tag{13}$$

One can reconstruct the amplitude from the generating function through the following relation:

$$G_{\lambda}^{(a)}(\ell_1, \ell_2; t) = \oint_{C_1} \frac{dx}{2\pi i x^{\ell_1 + 1}} \oint_{C_2} \frac{dy}{2\pi i y^{\ell_2 + 1}} \widetilde{G}^{(a)}(g, x, y; t) , \qquad (14)$$

where the contour C_1 (C_2) is chosen to enclose x = 0 (y = 0) and to ensure the convergence of $\tilde{G}^{(a)}(g, x, y; t)$. One can derive the relation (14) using the identity:

$$\oint_C \frac{dz}{2\pi i z^{n+1}} = \delta_{n,0} , \quad (n \in \mathbb{Z}) ,$$
(15)

where the contour C encloses z = 0. We provide the composition law for the generating function when a = 0, -1 in preparation for later calculations:

$$\widetilde{G}^{(a)}(g,x,y;t_1+t_2) = \oint_C \frac{dz}{2\pi i z} \widetilde{G}^{(a)}(g,x,z^{-1};t_1) \ \widetilde{G}^{(a)}(g,z,y;t_2) \ , \ (a=0,-1) \ ,$$
(16)

where the contour encloses z = 0, and for fixed g, x and y lies inside the radius of convergence for $\tilde{G}^{(a)}(g, x, z^{-1}; t_1)$ as the series in 1/z and for $\tilde{G}^{(a)}(g, z, y; t_2)$ as the series in z, which is possible as we will see.

In what follows, we will discuss the one-step amplitude $G_{\lambda}^{(a)}(\ell_1, \ell_2; 1)$. This is because it becomes an important object when computing the whole amplitude.

2.1 Counting triangulations

In this section we focus on the one-step amplitude $G_{\lambda}^{(a)}(\ell_1, \ell_2; 1)$ which is the sum over triangulations of a strip as shown in Fig. 1:

$$G_{\lambda}^{(a)}(\ell_1, \ell_2; 1) = e^{-\lambda(\ell_1 + \ell_2)} \mathcal{N}^{(a)}(\ell_1, \ell_2, n = \ell_1 + \ell_2), \qquad (17)$$

and count the number of triangulations $\mathcal{N}^{(a)}(\ell_1, \ell_2, n = \ell_1 + \ell_2)$. Based on simple combinatorics, one can calculate the case of a = 1 which has the $S^1 \times [0, 1]$ topology:

$$\mathcal{N}^{(1)}(\ell_1, \ell_2, n = \ell_1 + \ell_2) = \frac{1}{\ell_1 + \ell_2} \begin{pmatrix} \ell_1 + \ell_2 \\ \ell_1 \end{pmatrix} = \frac{(\ell_1 + \ell_2 - 1)!}{\ell_1! \ell_2!} .$$
(18)

Because of the property (7), one can easily compute the case of a = -1 that the topology is $S^1 \times [0, 1]$ and one of the edges in the initial one-dimensional universe is marked:

$$\mathcal{N}^{(-1)}(\ell_1, \ell_2, n = \ell_1 + \ell_2) = \ell_1 \mathcal{N}^{(1)}(\ell_1, \ell_2, n = \ell_1 + \ell_2) = \frac{\ell_1(\ell_1 + \ell_2 - 1)!}{\ell_1!\ell_2!} .$$
(19)

Concerning the case of a = 0 whose topology is $[0, 1] \times [0, 1]$, there exist several possibilities depending on the restriction on the leftmost and rightmost triangles. If the rightmost triangle is the upward triangle (downward triangle) and the leftmost triangles is the downward triangle (upward triangle), then the counting of triangulations yields

$$\mathcal{N}^{(0)}(\ell_1, \ell_2, n = \ell_1 + \ell_2) = \begin{pmatrix} \ell_1 + \ell_2 - 2\\ \ell_1 - 1 \end{pmatrix} = \frac{(\ell_1 + \ell_2 - 2)!}{(\ell_1 - 1)!(\ell_2 - 1)!} .$$
(20)

In the following, we will use eq. (20) in the case of a = 0 for computational simplicity⁴.

The one-step generating functions can be derived inserting eqs. (18), (19) and (20) into eq. (12):

⁴ If we do not impose any restriction on the leftmost and rightmost triangles, the number of triangulations becomes $\mathcal{N}^{(0)}(\ell_1, \ell_2, n = \ell_1 + \ell_2) = (\ell_1 + \ell_2)!/(\ell_1!\ell_2!)$.

$$\widetilde{G}^{(1)}(g, x, y; 1) = \sum_{\ell_1=1}^{\infty} \sum_{\ell_2=1}^{\infty} x^{\ell_1} y^{\ell_2} g^{\ell_1+\ell_2} \mathcal{N}^{(1)}(\ell_1, \ell_2, n = \ell_1 + \ell_2)$$
$$= -\ln\left(\frac{1-gx-gy}{(1-gx)(1-gy)}\right);$$
(21)

$$\widetilde{G}^{(-1)}(g, x, y; 1) = \sum_{\ell_1=1}^{\infty} \sum_{\ell_2=1}^{\infty} x^{\ell_1} y^{\ell_2} g^{\ell_1+\ell_2} \mathcal{N}^{(-1)}(\ell_1, \ell_2, n = \ell_1 + \ell_2)$$
$$= \frac{g^2 x y}{(1 - gx)(1 - gx - gy)};$$
(22)

$$\widetilde{G}^{(0)}(g, x, y; 1) = \sum_{\ell_1=1}^{\infty} \sum_{\ell_2=1}^{\infty} x^{\ell_1} y^{\ell_2} g^{\ell_1+\ell_2} \mathcal{N}^{(0)}(\ell_1, \ell_2, n = \ell_1 + \ell_2)$$
$$= \frac{g^2 x y}{1 - g x - g y} .$$
(23)

In fact, one can also obtain eq. (22) through $\widetilde{G}^{(-1)}(g, x, y; 1) = x \frac{\partial}{\partial x} \widetilde{G}^{(1)}(g, x, y; 1)$. Alternatively, it is possible to compute the one-step generating functions directly by simple combinatorics:

$$\widetilde{G}^{(1)}(g,x,y;1) = \sum_{s=1}^{\infty} \frac{1}{s} \left(\sum_{k=1}^{\infty} (gx)^k \sum_{l=1}^{\infty} (gy)^l \right)^s = -\ln\left(\frac{1-gx-gy}{(1-gx)(1-gy)}\right); \quad (24)$$
$$\widetilde{G}^{(-1)}(g,x,y;1) = \sum_{k=0}^{\infty} \left(gx \sum_{l=0}^{\infty} (gy)^l \right)^k - \sum_{k=0}^{\infty} (gx)^k = \frac{g^2xy}{(1-gx)(1-gx-gy)}; \quad (25)$$

$$\widetilde{G}^{(0)}(g,x,y;1) = \sum_{s=1}^{\infty} \frac{1}{s} \left(\sum_{k=1}^{\infty} (gx)^k \sum_{l=1}^{\infty} (gy)^l \right)^s = \frac{g^2 x y}{1 - gx - gy} .$$
(26)

2.2 Continuum limit

All is now set for computing the amplitude in the continuum limit. In this section, however, instead of directly computing the amplitude in the continuum limit, we will derive the differential equation that the continuum amplitude satisfies.

Before going into details any further, let us explain some basic facts of the continuum limit. In order to remove the cutoff ε through the continuum limit, one has to tune the bare coupling constants (g, x, y) to their critical values (g_c, x_c, y_c) . At the critical values, the generating function hits the radii of convergence and therefore becomes non-analytic. Approaching such a critical point, infinitely many triangles and boundary edges become important in the summation of the generating function, i.e., essentially, the average number of triangles and boundary edges become infinity at the critical point. Having this in mind, one may intuitively understand that the

continuous surface would be obtained if $(g, x, y) \rightarrow (g_c, x_c, y_c)$ and $\varepsilon \rightarrow 0$ in a correlated manner.

Introducing $\lambda_c = -\ln[g_c]$, $\lambda_{1c} = -\ln[x_c]$ and $\lambda_{2c} = -\ln[y_c]$, one can transmute the dimension of the lattice spacing ε into the dimension of the renormalized bulk and boundary cosmological constants through the continuum limit:

$$\Lambda = \lim_{\substack{\lambda \to \lambda_c \\ \varepsilon \to 0}} \frac{\lambda - \lambda_c}{\varepsilon^2} , \quad X = \lim_{\substack{\lambda_1 \to \lambda_{1c} \\ \varepsilon \to 0}} \frac{\lambda_1 - \lambda_{1c}}{\varepsilon} , \quad Y = \lim_{\substack{\lambda_2 \to \lambda_{2c} \\ \varepsilon \to 0}} \frac{\lambda_2 - \lambda_{2c}}{\varepsilon} , \quad (27)$$

where Λ is the renormalized bulk cosmological constant, and *X* and *Y* are the renormalized boundary cosmological constants. Therefore, the divergent bare cosmological constants get additive renormalizations so as to obtain the finite renormalized cosmological constants that set the scale at IR.

In the following, we discuss the continuum limit in detail with respect to each topology of spacetime.

2.2.1 $S^1 \times [0, 1]$ topology

We consider the case of a = -1, i.e. $S^1 \times [0, 1]$ topology with a marked boundary. Using the composition law (16) and the one-step generating function (25), one obtains

$$\begin{split} \widetilde{G}^{(-1)}(g,x,y;t+1) &= \oint_C \frac{dz}{2\pi i z} \widetilde{G}^{(-1)}(g,x,z^{-1};1) \ \widetilde{G}^{(-1)}(g,z,y;t) \\ &= \oint_C \frac{dz}{2\pi i} \frac{g^2 x}{(1-gx)^2 (z-g/(1-gx))} \frac{\widetilde{G}^{(-1)}(g,z,y;t)}{z} \\ &= \frac{gx}{1-gx} \ \widetilde{G}^{(-1)}\left(g,\frac{g}{1-gx},y;t\right) \ . \end{split}$$
(28)

In the last equality, we have picked up a pole at z = g/(1 - gx), and there exists no pole at z = 0 since $\frac{\tilde{G}^{(-1)}(g,z,y;t)}{z}$ is regular. Through iterative use of eq. (28), one can analytically compute the generating function, and extract the information of the critical point [9]. However, we do not compute the generating function directly to obtain the critical point. Instead, we follow the procedure shown in [8]: One assumes the existence of the critical point, and determines the value of the critical coupling constants from the consistency.

We assume the critical point characterized by the critical coupling constants (g_c, x_c, y_c) and use the following parametrization:

$$g = g_c e^{-\varepsilon^2 \Lambda}, \quad x = x_c e^{-\varepsilon X}, \quad y = y_c e^{-\varepsilon Y}.$$
 (29)

Assuming the scalings

$$T = \varepsilon t$$
, $L_1 = \varepsilon \ell_1$, $L_2 = \varepsilon \ell_2$, (30)

we introduce the renormalized amplitude and the renormalized generating function at the critical point by the multiplicative renormalizations:

$$G_{\Lambda}^{(-1)}(L_1, L_2; T) = \lim_{\varepsilon \to 0} C_{\varepsilon} \ G_{\lambda}^{(-1)}(\ell_1, \ell_2; t) , \qquad (31)$$

$$\widetilde{G}_{\Lambda}^{(-1)}(X,Y;T) = \lim_{\varepsilon \to 0} \widetilde{C}_{\varepsilon} \ \widetilde{G}^{(-1)}(g,x,y;t) , \qquad (32)$$

where C_{ε} and \tilde{C}_{ε} are real functions of ε that will be fixed later. One can determine the critical coupling constants for the consistency. Using the scaling behavior (32), eq. (28) can yield the sensible continuum limit if the critical coupling constants satisfy

$$\frac{g_c x_c}{1 - g_c x_c} = 1 , \quad \frac{g_c}{1 - g_c x_c} = x_c \quad \Rightarrow \quad g_c = \frac{1}{2} , \quad x_c = 1 .$$
(33)

Let us determine the function C_{ε} in such a way that the composition law (10) holds in the continuum limit as

$$G_{\Lambda}^{(-1)}(L_1, L_2; T_1 + T_2) = \int_0^\infty dL \ G_{\Lambda}^{(-1)}(L_1, L; T_1) G_{\Lambda}^{(-1)}(L, L_2; T_2) , \qquad (34)$$

which is possible if $C_{\varepsilon} = \varepsilon^{-1}$. The function $\widetilde{C}_{\varepsilon}$ can be determined in such way that eq. (12) makes sense in the continuum limit, i.e.

$$\widetilde{G}_{\Lambda}^{(-1)}(X,Y;T) = \int_0^\infty dL_1 \int_0^\infty dL_2 \ e^{-XL_1} e^{-YL_2} G_{\Lambda}^{(-1)}(L_1,L_2;T) , \qquad (35)$$

which is possible if $\tilde{C}_{\varepsilon} = \varepsilon$.

Now we wish to take the continuum limit of eq. (28). For notational convenience, we redefine the renormalized coupling constants as follows:

$$g = \frac{1}{2}e^{-\varepsilon^2 \Lambda} \equiv \frac{1}{2}\left(1 - \frac{1}{2}\varepsilon^2 \Lambda\right), \quad x = e^{-\varepsilon X} \equiv 1 - \varepsilon X, \quad y = e^{-\varepsilon Y} \equiv 1 - \varepsilon Y.$$
(36)

Plugging eqs. (30), (36) into eq. (28), one obtains the differential equation:

$$\frac{\partial}{\partial T}\widetilde{G}_{\Lambda}^{(-1)}(X,Y;T) = -\frac{\partial}{\partial X}\left[(X^2 - \Lambda)\widetilde{G}_{\Lambda}^{(-1)}(X,Y;T) \right] .$$
(37)

Doing a little math, one can also derive the continuum description of eq. (14):

$$G_{\Lambda}^{(-1)}(L_1, L_2; T) = \int_{c-i\infty}^{c+i\infty} \frac{dX}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{dY}{2\pi i} e^{L_1 X} e^{L_2 Y} \tilde{G}_{\Lambda}^{(-1)}(X, Y; T) , \quad (38)$$

where c is a suitable real number. Using the inverse Laplace transform (38) and eq. (37), one obtains the differential equation that the continuum amplitude satisfies:

$$\frac{\partial}{\partial T}G_{\Lambda}^{(-1)}(L_1, L_2; T) = -\hat{H}^{(-1)}(L_1) G_{\Lambda}^{(-1)}(L_1, L_2; T) , \qquad (39)$$

where

$$\hat{H}^{(-1)}(L) = -L\frac{\partial^2}{\partial L^2} + \Lambda L .$$
(40)

As a result, one can interpret the continuum limit of 2d CDT as a quantum system of the one-dimensional universe with length L that propagates in time T following the quantum Hamiltonian (40). The quantum Hamiltonian (40) is Hermitian with respect to the inner product:

$$\int_0^\infty \frac{dL}{L} \phi^*(L) \, (\hat{H}^{(-1)}\psi)(L) = \int_0^\infty \frac{dL}{L} (\hat{H}^{(-1)}\phi)^*(L) \, \psi(L) \,. \tag{41}$$

The differential equation for the un-marked amplitude can be easily read off inserting the continuum limit of eq. (7)

$$G_{\Lambda}^{(-1)}(L_1, L_2; T) = L_1 G_{\Lambda}^{(1)}(L_1, L_2; T) , \qquad (42)$$

into eq. (39):

$$\frac{\partial}{\partial T} G_{\Lambda}^{(1)}(L_1, L_2; T) = -\hat{H}^{(1)}(L_1) \ G_{\Lambda}^{(1)}(L_1, L_2; T) , \qquad (43)$$

where

$$\hat{H}^{(1)}(L) = -\frac{\partial^2}{\partial L^2}L + \Lambda L .$$
(44)

The quantum Hamiltonian (44) is Hermitian with respect to the inner product:

$$\int_0^\infty L dL \phi^*(L) \ (\hat{H}^{(1)}\psi)(L) = \int_0^\infty L dL (\hat{H}^{(1)}\phi)^*(L) \ \psi(L) \ . \tag{45}$$

2.2.2 [0,1] × [0,1] topology

Let us consider the case of a = 0, i.e. $[0, 1] \times [0, 1]$ topology. We basically follow the procedure shown in Sect. 2.2.1. Using the composition law (16) and the one-step generating function (26), one obtains

$$\begin{split} \widetilde{G}^{(0)}(g,x,y;t+1) &= \oint_C \frac{dz}{2\pi i z} \widetilde{G}^{(0)}(g,x,z^{-1};1) \ \widetilde{G}^{(0)}(g,z,y;t) \\ &= \oint_C \frac{dz}{2\pi i} \frac{g^2 x}{(1-gx)(z-g/(1-gx))} \frac{\widetilde{G}^{(0)}(g,z,y;t)}{z} \\ &= gx \ \widetilde{G}^{(0)}\left(g,\frac{g}{1-gx},y;t\right) \ . \end{split}$$
(46)

From eq. (46), one may obtain a sensible continuum limit if the critical coupling constants are the same as before, i.e. $(g_c, x_c, y_c) = (1/2, 1, 1)$, and if the multiplicative renormalization is treated carefully:

$$\widetilde{G}_{\Lambda}^{(0)}(X,Y;T) = \lim_{\varepsilon \to 0} \frac{\varepsilon}{2^t} \widetilde{G}^{(0)}(g,x,y;t) .$$
(47)

In fact, this assumption yields the correct continuum limit. Plugging eqs. (30), (36) into eq. (46) and using eq. (47), one obtains the differential equation:

$$\frac{\partial}{\partial T}\widetilde{G}^{(0)}_{\Lambda}(X,Y;T) = -\left(X + (X^2 - \Lambda)\frac{\partial}{\partial X}\right)\widetilde{G}^{(0)}_{\Lambda}(X,Y;T) .$$
(48)

Defining the continuum amplitude in such a way that the inverse Laplace transform (38) holds in the case of a = 0 as well, i.e.

$$G_{\Lambda}^{(0)}(L_1, L_2; T) = \int_{c-i\infty}^{c+i\infty} \frac{dX}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{dY}{2\pi i} e^{L_1 X} e^{L_2 Y} \tilde{G}_{\Lambda}^{(0)}(X, Y; T) , \qquad (49)$$

and using eq. (49), the differential equation (48) becomes

$$\frac{\partial}{\partial T}G_{\Lambda}^{(0)}(L_1, L_2; T) = -\hat{H}^{(0)}(L_1) G_{\Lambda}^{(0)}(L_1, L_2; T) , \qquad (50)$$

where $\hat{H}^{(0)}$ is the quantum Hamiltonian obtained in Refs. [31, 32]:

$$\hat{H}^{(0)}(L) = -\frac{\partial}{\partial L}L\frac{\partial}{\partial L} + \Lambda L .$$
(51)

The quantum Hamiltonian (51) is hermitian with respect to the inner product:

$$\int_0^\infty dL\phi^*(L) \; (\hat{H}^{(0)}\psi)(L) = \int_0^\infty dL(\hat{H}^{(0)}\phi)^*(L) \; \psi(L) \; . \tag{52}$$

2.3 Short summary of 2d CDT

As discussed in Sect. 2.2, the continuum limit of 2d CDT is described by the quantum mechanics of a one-dimensional universe with length *L* that propagates in time *T* based on the Hamiltonian $\hat{H}^{(a)}$:

$$\hat{H}^{(-1)} = -L\frac{\partial^2}{\partial L^2} + \Lambda L , \quad \hat{H}^{(1)} = -\frac{\partial^2}{\partial L^2}L + \Lambda L , \quad \hat{H}^{(0)} = -\frac{\partial}{\partial L}L\frac{\partial}{\partial L} + \Lambda L ,$$
(53)

where the label *a* classifies the topology of the one-dimensional universe: S^1 and [0, 1] for a = 1 and a = 0, respectively. When a = -1, the closed one-dimensional universe is marked. Let us define the eigenstates of *L* as $|L\rangle_a$ that satisfy the completeness relation:

$$\hat{1} = \int_0^\infty L^a dL \, |L\rangle_{aa} \langle L| \quad \Leftrightarrow \quad {}_a \langle L'|L\rangle_a = \frac{1}{L^a} \delta(L - L') \,. \tag{54}$$

Note that $|L\rangle_{-1} = L|L\rangle_{1}$. One can then express the amplitudes as matrix elements:

$$G_{\Lambda}^{(1)}(L_1, L_2; T) = {}_1 \langle L_2 | e^{-T\hat{H}^{(1)}} | L_1 \rangle_1 , \qquad (55)$$

$$G_{\Lambda}^{(-1)}(L_1, L_2; T) = {}_1 \langle L_2 | e^{-T\hat{H}^{(-1)}} | L_1 \rangle_{-1} , \qquad (56)$$

$$G_{\Lambda}^{(0)}(L_1, L_2; T) = {}_0 \langle L_2 | e^{-T\hat{H}^{(0)}} | L_1 \rangle_0 , \qquad (57)$$

Using eq. (54), one can show that the composition laws hold: For a = -1, 0,

$$G_{\Lambda}^{(a)}(L_1, L_2; T_1 + T_2) = \int_0^\infty dL \ G_{\Lambda}^{(a)}(L_1, L; T_1) G_{\Lambda}^{(a)}(L, L_2; T_2) , \qquad (58)$$

and for a = 1,

$$G_{\Lambda}^{(1)}(L_1, L_2; T_1 + T_2) = \int_0^\infty dL \ G_{\Lambda}^{(1)}(L_1, L; T_1) L G_{\Lambda}^{(1)}(L, L_2; T_2) \ . \tag{59}$$

3 2d projectable Hořava-Lifshitz quantum gravity

We wish to introduce the classical field theory that reproduces the continuum limit of 2d CDT once it is quantized. The field theory is a certain version of the twodimensional Hořava-Lifshitz gravity (2d HL gravity).

The starting point is the same class of manifold with a foliation (1) where Σ_t is a one-dimensional space labelled by *t*:

$$\Sigma_t = \{ x^{\mu} \in \mathcal{M} \mid f(x^{\mu}) = t \}, \text{ with } \mu = 0, 1.$$
(60)

Choosing that $f(x^{\mu}) = x^0$, the time direction can be decomposed into the two directions, i.e. the normal and the tangential to Σ_t :

$$(\partial_t)^{\mu} = \frac{\partial x^{\mu}}{\partial t} = Nn^{\mu} + N^1 E_1^{\mu} , \qquad (61)$$

where n^{μ} and E_1^{μ} are, respectively, the unit normal vector and the tangent vector defined as

$$n^{\mu} = \left(\frac{1}{N}, -\frac{N^{1}}{N}\right), \quad E_{1}^{\mu} = \delta_{1}^{\mu}.$$
 (62)

Here N and N¹ are the Lapse function and the shift vector. Through the use of eq. (61), one can parametrize the metric $g_{\mu\nu}$ on \mathcal{M} as follows:

$$ds^{2} = g_{\mu\nu}dx^{\mu}dx^{\nu} = -N^{2}dt^{2} + h_{11}\left(dx + N^{1}dt\right)\left(dx + N^{1}dt\right), \quad (63)$$

where $t = x^0$ and $x = x^1$; h_{11} is the spatial metric on Σ_t defined as $h_{11} = E_1^{\mu} E_1^{\nu} g_{\mu\nu}$.

2d HL gravity is a field theory that preserves the structure of the time foliation, or in other words, it is invariant under the foliation-preserving diffeomorphisms (FPD):

$$t \to t + \xi^0(t) , \quad x \to x + \xi^1(t, x) .$$
 (64)

The fields transform under FPD as follows:

$$\delta_{\xi} h_{11} = \xi^0 \partial_0 h_{11} + \xi^1 \partial_1 h_{11} + 2h_{11} \partial_1 \xi^1 , \qquad (65)$$

$$\delta_{\xi} N_1 = \xi^{\mu} \partial_{\mu} N_1 + N_1 \partial_{\mu} \xi^{\mu} + h_{11} \partial_0 \xi^1 , \qquad (66)$$

$$\delta_{\xi} N = \xi^{\mu} \partial_{\mu} N + N \partial_0 \xi^0 . \tag{67}$$

where $N_1 = h_{11}N^1$. Here if a function is a constant on each foliation Σ_t , such a function is called projectable. In fact, implementing FPD the projectable Lapse function, i.e. N = N(t), stays as a function only of time. The HL gravity with the projectable Lapse function is dubbed the projectable HL gravity. Since it is 2d projectable HL gravity that reproduces the continuum limit of 2d CDT once it is quantized, from now we focus on this special version of HL gravity.

The action of 2d projectable HL gravity is given by

$$I = \int dt \,\mathcal{L} = \frac{1}{\kappa} \int dt dx \, N(t) \sqrt{h(t,x)} \left((1-\eta) K^2(t,x) - 2\widetilde{\Lambda} \right) \,, \tag{68}$$

where \mathcal{L} is the Lagrangian; η , $\tilde{\Lambda}$ and κ are a dimensionless parameter, the cosmological constant and the (dimensionless) gravitational coupling constant, respectively; h is the determinant of the spatial metric h_{11} , i.e. $h = h_{11}$; K is the trace of the extrinsic curvature K_{11} given by

$$K_{11} = \frac{1}{2N} \left(\partial_0 - 2\nabla_1 N_1 \right) , \quad \text{with} \quad \nabla_1 N_1 = \partial_1 N_1 - \Gamma_{11}^1 N_1 . \tag{69}$$

Here Γ_{11}^1 is the spatial Christoffel symbol:

$$\Gamma_{11}^1 = \frac{1}{2} h^{11} \partial_1 h_{11} . \tag{70}$$

One can in principle add higher spatial derivative terms to the action (68). However, such terms are not necessary because the model is renormalizable in two dimensions without introducing them, and therefore we omit such terms.

The continuum limit of 2d CDT can be precisely obtained if quantizing 2d projectable HL gravity with the following identification of parameters:

$$\Lambda = \frac{\widetilde{\Lambda}}{2(1-\eta)}, \quad \eta < 1, \quad \kappa = 4(1-\eta), \quad (71)$$

where Λ is the renormalized cosmological constant of CDT defined by eq. (36).

3.1 Quantization

Let us overview the quantization of 2d projectable HL gravity shown in Ref. [15] (see also Ref. [17] for another article examining this issue).

We introduce the conjugate momentum of \sqrt{h} as Π , which satisfy the Poisson bracket:

$$\left\{\sqrt{h(t,x)}, \Pi(t,x')\right\} = \delta(x-x') .$$
(72)

Through the Legendre transformation of the Lagrangian (68), one obtains the Hamiltonian of 2d projectable HL gravity:

$$H = \int dx \left(\Pi(t, x) \partial_t \sqrt{h(t, x)} \right) - \mathcal{L} = N(t)C(t) + \int dx \, N_1(t, x)C^1(t, x) \,. \tag{73}$$

Since 2d projectable HL gravity is a singular system due to the invariance under FPD, there exist two kinds of constraint:

$$C^{1}(t,x) = -\frac{\partial_{1}\Pi(t,x)}{\sqrt{h(t,x)}} \approx 0, \qquad (74)$$

$$C(t) = \int dx \, \left(\frac{\kappa}{4(1-\eta)} \Pi^2(t,x) \sqrt{h(t,x)} + \frac{2}{\kappa} \widetilde{\Lambda} \sqrt{h(t,x)}\right) \approx 0 \,, \qquad (75)$$

where $C^1(t, x) \approx 0$ is the momentum constraint, and $C(t) \approx 0$ is the Hamiltonian constraint which is global because of the projectable Lapse function⁵.

The strategy is to solve the momentum constraint (74) at the level of classical theory, i.e.

$$C^{1}(t,x) = 0 \quad \Rightarrow \quad \Pi(t,x) = \Pi(t) , \tag{76}$$

⁵ The Hamiltonian and the momentum constraints come from the consistency conditions that the primary constraints, $\Pi_N \approx 0$ and $\Pi_{N_1} \approx 0$, should be preserved under the time flow where Π_N and Π_{N_1} are the conjugate momenta of N and N₁, respectively.

meaning that the conjugate momentum becomes a function only of time. Applying eq. (76), the Hamiltonian (73) reduces to the one for the one-dimensional system:

$$H = N(t) \left(\frac{\kappa}{4(1-\eta)} \Pi^2(t) L(t) + \frac{2}{\kappa} \widetilde{\Lambda} L(t) \right), \quad \text{with} \quad L(t) = \int dx \sqrt{h(t,x)}, \quad (77)$$

where L(t) is the invariant length of the one-dimensional universe. Let us discuss solutions to the Hamiltonian constraint. If $(\eta - 1)\widetilde{\Lambda} > 0$, one has a solution:

$$\Pi^2 = \frac{8(\eta - 1)}{\kappa^2} \tilde{\Lambda} , \qquad (78)$$

which means that the extrinsic curvature is a constant. On the other hand, if $(\eta - 1)\widetilde{\Lambda} < 0$, the only solution is

$$L = 0$$
. (79)

Hereafter we apply the parametrization (71): We choose $(\eta - 1)\widetilde{\Lambda} < 0$, set the unimportant dimensionless gravitational constant as $\kappa = 4(1 - \eta)$, and redefine the cosmological constant as $\Lambda = \frac{\widetilde{\Lambda}}{2(1-\eta)}$. Since $\kappa > 0$, this means that $\eta < 1$ which selects the correct sign of the kinetic term, and the positive cosmological constant $\widetilde{\Lambda} > 0$. The dynamics of the classical 1d system with the Hamiltonian (77) can be alternatively described by the following action:

$$S = \int_0^1 dt \left(\frac{\dot{L}^2(t)}{4N(t)L(t)} - \Lambda N(t)L(t) \right) \,, \tag{80}$$

where $\dot{L}(t) := \frac{d}{dt}L(t)$. We then introduce the proper time:

$$\tau(s) = \int_0^s dt \, N(t) \, , \quad s \in [0, 1] \, . \tag{81}$$

Since the proper time (81) is invariant under the reparametrization of time, $t \rightarrow t + \xi^0(t)$, if one fixes the Lapse function as $N(\tau) = 1$, the length of the onedimensional universe $L(\tau)$ is also invariant under the time redefinition. Therefore, it makes sense to discuss the amplitude such that the one-dimensional universe with the length $L_1 := L(\tau = 0)$ propagates in the proper time τ , and ends up with the universe whose length is given by $L_2 := L_2(\tau = T)$.

With this understanding, we consider such an amplitude based on the path integral. For convenience, we rotate $t \rightarrow it$, which is possible thanks to the foliation and introduce the Euclidean action:

$$S_E = \int_0^1 dt \left(\frac{\dot{L}^2(t)}{4N(t)L(t)} + \Lambda N(t)L(t) \right) \,. \tag{82}$$

Using the Euclidean action (82), the amplitude becomes

$$\mathcal{G}_{\Lambda}(L_1, L_2; T) = \int \frac{\mathcal{D}N(t)}{\text{Diff}[0, 1]} \int_{L(0)=L_1}^{L(1)=L_2} \mathcal{D}L(t) \ e^{-S_E[N(t), L(t)]} , \qquad (83)$$

where

$$T := \int_0^1 dt \, N(t) \,. \tag{84}$$

We fix the Lapse function as $N(\tau) = 1$ introducing the corresponding Faddeev-Popov (FP) determinant. Since the FP determinant only gives an overall constant, we will omit it in the following. After the gauge fixing, the amplitude (83) becomes

$$\mathcal{G}_{\Lambda}(L_1, L_2; T) = \int_{L(0)=L_1}^{L(T)=L_2} \mathcal{D}L(\tau) \exp\left[-\int_0^T d\tau \left(\frac{\dot{L}^2(\tau)}{4L(\tau)} + \Lambda L(\tau)\right)\right], \quad (85)$$

where $\dot{L}(\tau) := \frac{d}{d\tau}L(\tau)$.

So far, we have not specified the integral measure $\mathcal{D}L(\tau)$. We apply the three kinds of measure given by

$$\mathcal{D}^{(a)}L(\tau) = \prod_{\tau=0}^{\tau=T} L^a(\tau) \ dL(\tau) \ , \quad (a=0,\pm 1) \ . \tag{86}$$

Accordingly, we consider the three kinds of amplitude, i.e. $\mathcal{G}_{\Lambda}^{(a)}(L_1, L_2; T)$, and rewrite them introducing the quantum Hamiltonian $\hat{H}^{(a)}$:

$$\mathcal{G}_{\Lambda}^{(a)}(L_1, L_2; T) = {}_a \langle L_2 | e^{-T\hat{H}^{(a)}} | L_1 \rangle_a , \qquad (87)$$

where the eigenstates of L satisfy the completeness relation:

$$1 = \int_0^\infty L^a dL \, |L\rangle_{aa} \langle L| \quad \Leftrightarrow \quad {}_a \langle L'|L\rangle_a = \frac{1}{L^a} \delta(L - L') \,. \tag{88}$$

In order to read off the quantum Hamiltonian $\hat{H}^{(a)}$, we discretize the proper time interval in steps of ε , and calculate the one-step matrix element $\mathcal{G}_{\Lambda}^{(a)}(L, L'; \varepsilon)$. The normalization can be fixed so as to satisfy the following equation:

$$\lim_{\varepsilon \to 0} \int_0^\infty L^a dL \, \mathcal{G}_{\Lambda}^{(a)}(L,L';\varepsilon) = 1 \,, \tag{89}$$

which comes from the completeness relation (88). The result is

$$\mathcal{G}_{\Lambda}^{(a)}(L,L';\varepsilon) = \frac{(LL')^{(1-a)/2}}{L'\sqrt{4\pi\varepsilon L'}} e^{-\frac{(L-L')^2}{4\varepsilon L'} - \Lambda\varepsilon L'} .$$
(90)

Integrating the one-step amplitude together with a function, $\psi_a(L) = {}_a\langle L|\psi\rangle$, for $\varepsilon \ll 1$, one can read off the quantum Hamiltonian:

$$\psi_{a}(L';\varepsilon) = {}_{a}\langle L'|e^{-\varepsilon H^{(a)}}|\psi\rangle$$

=
$$\int_{0}^{\infty} L^{a} dL {}_{a}\langle L'|e^{-\varepsilon \hat{H}^{(a)}}|L\rangle_{aa}\langle L|\psi\rangle$$

$$\cong \psi_{a}(L') - \varepsilon \hat{H}^{(a)}\psi_{a}(L') + O(\varepsilon^{3/2}) .$$
(91)

Using eq. (90), one obtains

$$\hat{H}^{(-1)}(L) = -L\frac{d^2}{dL^2} + \Lambda L , \quad \hat{H}^{(0)}(L) = -\frac{d}{dL}L\frac{d}{dL} + \Lambda L , \quad \hat{H}^{(1)}(L) = -\frac{d^2}{dL^2}L + \Lambda L$$
(92)

 $\hat{\Delta}(a)$

The quantum Hamiltonians (92) obtained by quantizing 2d projectable HL gravity are precisely equivalent to those obtained by the continuum limit of 2d CDT (see eqs. (40), (44) and (51)). The amplitudes are related as follows:

$$\mathcal{G}_{\Lambda}^{(-1)}(L,L';T) = L'G_{\Lambda}^{(-1)}(L,L';T) , \quad \mathcal{G}_{\Lambda}^{(a)}(L,L';T) = G_{\Lambda}^{(a)}(L,L';T) , \quad (a = 0, 1) .$$
(93)

Thereby, we understand that the classical field theory that reproduces the continuum limit of 2d CDT once it is quantized is indeed 2d projectable HL gravity. The projectable Lapse function allows us to introduce the reparametrization-invariant proper time, and to reduce the 2d field theory to the 1d system.

4 Sum over all wormholes and baby universes

In the CDT model, the spatial topology change is not allowed to occur by definition. One can generalize the 2d CDT model in such a way that spatial topology changes do occur in keeping with the foliation structure, and the universality class is the same as that of 2d CDT. Such a model is called generalized CDT (GCDT). GCDT can be constructed as both discretized and continuum models. Here of course the continuum model can be obtained by the continuum limit of the discretized model, but one can directly construct the continuum GCDT model promoting the onedimensional quantum-mechanical system discussed in Sec. 2 to a 2d field theory that includes the splitting and joining interactions of the one-dimensional spatial universe. Such a field theory is dubbed the string field theory for CDT, in which the string means the one-dimensional universe [22]. In this section, we introduce the string field theory for CDT, and briefly explain the fact that one can take the sum over all wormholes (i.e. handles) and baby universes [25,26]. Here the baby universe is a portion of geometry that is pinched off from the "parent universe" and vanishes into the vacuum. We also introduce an effective one-body theory that reproduces the many-body effects coming from the splitting and joining interactions. We then discuss those effects in the context of HL gravity [28]. In the end, we show that a

sort of Coleman's mechanism works when taking into account all contributions of wormholes and baby universes non-perturbatively [30].

We introduce an operator that creates a marked closed string (i.e. a marked closed one-dimensional universe) with length L, $\Psi^{\dagger}(L)$, and an operator that annihilates a length-L marked closed string, $\Psi(L)$. These operators satisfy the following commutators:

$$[\Psi(L), \Psi^{\dagger}(L')] = L\delta(L - L'), \quad [\Psi(L), \Psi(L')] = [\Psi^{\dagger}(L), \Psi^{\dagger}(L')] = 0.$$
(94)

The vacuum state $|vac\rangle$ is defined by the equation: $\Psi(L)|vac\rangle = 0$. The CDT amplitude (57) can be expressed by sandwiching the one-body Hamiltonian:

$$\mathcal{G}_{\Lambda}^{(-1)}(L_1, L_2; T) = L_2 G_{\Lambda}^{(-1)}(L_1, L_2; T) = \langle \operatorname{vac} | \Psi(L_2) \ e^{-T \mathcal{H}^{(-1)}_{\text{free}}} \ \Psi^{\dagger}(L_1) | \operatorname{vac} \rangle ,$$
(95)

where

$$\mathcal{H}_{\text{free}}^{(-1)} = \int_0^\infty \frac{dL}{L} \Psi^{\dagger}(L) \left(-L \frac{\partial^2}{\partial L^2} + \Lambda L \right) \Psi(L) . \tag{96}$$

Hereafter we omit the superscript (-1) for avoiding notational complexity. Adding splitting and joining interactions into the free Hamiltonian (96), one obtains the full Hamiltonian of the string field theory for CDT:

$$\mathcal{H} = \mathcal{H}_{\text{free}} - g_s \int_0^\infty dL_1 \int_0^\infty dL_2 \Psi^{\dagger}(L_1) \Psi^{\dagger}(L_2) \Psi(L_1 + L_2)$$
$$- \alpha g_s \int_0^\infty dL_1 \int_0^\infty dL_2 \Psi^{\dagger}(L_1 + L_2) \Psi(L_1) \Psi(L_2)$$
$$- \int_0^\infty \frac{dL}{L} \delta(L) \Psi(L) , \qquad (97)$$

where the second, third and fourth terms respectively mean the splitting interaction with the string coupling constant g_s , the joining interaction with the coupling constant αg_s , and the term associated with a string vanishing into the vacuum. Here the parameter α is introduced for counting the number of handles (i.e. wormholes). One can in principle calculate the amplitude for the process such that *m* closed strings propagate in time and end up with *n* closed strings:

$$A(L_1, \cdots, L_m; L'_1, \cdots, L'_n; T) = \langle \operatorname{vac} | \Psi(L'_1) \cdots \Psi(L'_n) e^{-T\mathcal{H}} \Psi^{\dagger}(L_1) \cdots \Psi^{\dagger}(L_m) | \operatorname{vac} \rangle,$$
(98)

4.1 Effective theory

Let us consider the full propagator $A(L_1; L_2; T)$ that includes the sum over all genera and baby universes. We can set $\alpha = 1$ without loss of generality since the parameter α plays a supplementary role and we are interested in taking the sum over all genus contributions. Somewhat miraculously, the full propagator defined in the many-body system with the Hamiltonian (97) can be effectively described by the one-body system [25, 26]:

$$A(L_1; L_2; T) = \langle L_2 | e^{-TH_{\text{eff}}(L_1)} | L_1 \rangle , \qquad (99)$$

where \hat{H} is the effective Hamiltonian given by

$$\hat{H}_{\text{eff}}(L) = -L \frac{d^2}{dL^2} + \Lambda L - g_s L^2 .$$
 (100)

This is possible because there exists a bijection called Ambjørn-Budd bijection [24] such that one can map each geometry generated in GCDT to a branched polymer with loops at the discrete level. The last term in the Hamiltonian (100), $-g_s L^2$, expresses all the effects originated with the baby universes and the wormholes. Note that the Hamiltonian (100) is not bounded from below because of the last term, but in fact this system is known to be "classical incomplete," which means that the Hamiltonian has discrete energy spectra, and a set of square integrable eigenfunctions (see e.g. Ref. [33] for a pedagogical explanation about the classical incomplete systems). A similar deformation has been observed in the c = 1 non-critical string theory [34,35].

The full propagator (100) can be also described in terms of the path-integral:

$$A(L_1; L_2; T) = \int_{L(0)=L_1}^{L(T)=L_2} \mathcal{D}L(\tau) \exp\left[-\int_0^T d\tau \left(\frac{\dot{L}^2(\tau)}{4L(\tau)} + \Lambda L(\tau) - g_s L^2(\tau)\right)\right],$$
(101)

where the integral measure is given by

$$\mathcal{D}L(\tau) = \prod_{\tau=0}^{\tau=T} L^{-1}(\tau) dL(\tau) .$$
 (102)

In order for the functional integral (101) to be well defined, one need to choose the boundary conditions on $L(\tau)$ at infinity such that the kinetic term counteracts the unboundedness of the potential. If one generalizes the integral measure (102) as

$$\mathcal{D}^{(a)}L(\tau) = \prod_{\tau=0}^{\tau=T} L^a(\tau) dL(\tau) , \quad (a = 0, \pm 1) , \qquad (103)$$

one can recover all possible orderings of the effective Hamiltonian (100) following the procedure explained in Sec. 3.1.

Interestingly, one can reproduce the full propagator (101) if one considers that the cosmological constant Λ in eq. (85) is not a constant but fluctuates independently in time around Λ , following the Gaussian distributions with a standard deviation $\sigma = 2\sqrt{g_s}$:

$$A(L_1; L_2; T) = \int \mathcal{D}\nu(\tau) \ e^{-\frac{1}{4g_s} \int_0^T d\tau \ \nu^2(\tau)} \mathcal{G}_{\Lambda+\nu}(L_1, L_2; T) \ , \qquad (104)$$

where

$$\mathcal{G}_{\Lambda+\nu}(L_1, L_2; T) := \int_{L(0)=L_1}^{L(T)=L_2} \mathcal{D}L(\tau) \exp\left[-\int_0^T d\tau \left(\frac{\dot{L}^2(\tau)}{4L(\tau)} + (\Lambda+\nu(\tau))L(\tau)\right)\right]$$
(105)

Therefore, all the contributions coming from the sum over all wormholes and baby universes can be fully taken in if the cosmological "constant" in (the continuum limit of) 2d CDT or projectable HL quantum gravity where no wormholes and baby universes exist is not really a constant but fluctuates in time. This would lead to a realization of Coleman's mechanism that will be discussed in Sec. 4.3.

In the next section, we will show that the full propagator can be also obtained if quantizing 2d projectable HL gravity with an effective wormhole interaction term.

4.2 Wormhole interaction in 2d projectable HL gravity

Let us consider the 2d projectable HL gravity with a space-like wormhole interaction given by the following action:

$$I_{\rm w} = \frac{1}{\kappa} \int dt dx \ N(t) \sqrt{h(t,x)} \left((1-\eta) K^2(t,x) - 2\widetilde{\Lambda} \right) + \beta \int dt N(t) \int dx_1 dx_2 \sqrt{h(t,x_1)} \sqrt{h(t,x_2)} , \qquad (106)$$

where β is a dimension-full coupling constant. The last bi-local term can be interpreted as an effective interaction term for a space-like wormhole connecting two distant regions at a given *t*. This bi-local term is allowed to be included since it is invariant under FPD.

Following essentially the same procedure explained in Sec. 3.1, let us quantize the system defined by the action (106). Introducing the conjugate momentum of the density \sqrt{h} as Π , we introduce the Poisson bracket (72). Implementing the Legendre transform, one obtains the corresponding Hamiltonian:

$$H_{\rm w} = N(t) C_{\rm w}(t) + \int dx N_1(t,x) C_{\rm w}^1(t,x) , \qquad (107)$$

where

$$C_{\rm w}^1(t,x) = -\frac{\partial_1 \Pi(t,x)}{\sqrt{h(t,x)}} \approx 0, \qquad (108)$$

$$C_{\rm w}(t) = \int dx \left(\frac{\kappa}{4(1-\eta)} \Pi^2(t,x) \sqrt{h(t,x)} + \frac{2}{\kappa} \widetilde{\Lambda} \sqrt{h(t,x)} - \beta \sqrt{h(t,x)} \int dx_2 \sqrt{h(t,x_2)} \right) \approx 0.$$
(109)

The constraints (108) and (109) are the momentum constraint and the Hamiltonian constraint, respectively. Solving the momentum constraint (108) at the classical level as before, the Hamiltonian (107) reduces to the following one-dimensional one:

$$H_{\rm w} = N(t) \left(\frac{\kappa}{4(1-\eta)} \Pi^2(t) L(t) + \frac{2}{\kappa} \widetilde{\Lambda} L(t) - \beta L^2(t) \right) , \qquad (110)$$

where $L(t) := \int dx \sqrt{h(t,x)}$. The Hamiltonian (110) is subject to the Hamiltonian constraint:

$$L(t)\left(\frac{\kappa}{4(1-\eta)}\Pi^2(t) + \frac{2}{\kappa}\widetilde{\Lambda} - \beta L(t)\right) \approx 0.$$
(111)

Here we choose the CDT parametrization (71). A solution to the Hamiltonian constraint (111) is

$$\Pi^2 = -\Lambda + \beta L \ge 0, \quad \text{for} \quad \sqrt{\Lambda}L \ge 1/\xi, \qquad (112)$$

where ξ is a dimensionless parameter given by $\xi = \beta / \Lambda^{3/2}$. For $\sqrt{\Lambda}L < 1/\xi$, the only allowed solution is L = 0.

When quantizing the system, if we follow the same procedure described in Sec. 3.1, and set $\beta = g_s$, one can reproduce the path-integral of the full propagator (101). Remember the boundary condition for the path-integral (101), i.e. the kinetic term should counteract the unboundedness of the potential term at $L = \infty$. This balance between the kinetic and potential terms is precisely what is reflected in the classical Hamiltonian constraint (112).

4.3 Coleman's mechanism

In this section, we discuss a sort of Coleman's mechanism in the context of twodimensional gravity based on CDT briefly.

Let us define the two kinds of Wheeler-deWitt equation:

$$\hat{H}W_0(L) = 0$$
, $\hat{H}_{\text{eff}}W(L) = 0$, (113)

where $\hat{H} := \hat{H}^{(-1)}$ introduced in eq. (53). The solutions to the Wheeler-deWitt equations are the Hartle-Hawking wave functions given by

$$W_0(L) = e^{-\sqrt{\Lambda}L} , \quad W(L) = \frac{\operatorname{Bi}(\xi^{-2/3} - \xi^{1/3}\sqrt{\Lambda}L)}{\operatorname{Bi}(\xi^{-2/3})} + c \operatorname{Ai}(\xi^{-2/3} - \xi^{1/3}\sqrt{\Lambda}L) ,$$
(114)

where Ai and Bi are the standard Airy functions, ξ is a dimensionless string coupling constant measured by the cosmological constant, i.e., $\xi := g_s/\Lambda^{3/2}$; and *c* is an undetermined dimensionless constant. The Hartle-Hawking wave function $W_0(L)$ is the one for (the continuum theory of) 2d CDT, i.e. neither baby universe nor wormhole contributions are included. On the other hand, W(L) is the Hartle-Hawking wave function including all possible contributions of baby universes and wormholes non-perturbatively.

We wish to explore the behavior of the non-perturbative Hartle-Hawking wave function W(L) (see Fig. 2). For $\sqrt{\Lambda}L \ll 1/\xi$, one obtains the asymptotic expansion:



Fig. 2 A plot of the Hartle-Hawking wave functions, W_0 (dashed line) and W (solid line), for $\xi = 1/3$ and c = 0: The horizontal axis is $\sqrt{\Lambda}L$ and the vertical axis is either W_0 or W.

$$W(L) \sim e^{-\sqrt{\Lambda L}} = W_0(L) . \tag{115}$$

Therefore, when the size of the one-dimensional universe is small enough, the physics is very close to the one without baby universes and wormholes, and it is essentially governed by the cosmological constant. The wave function in this region decreases exponentially, and this behavior does not change at any finite order of perturbation.

However, once the size of the universe is large enough, i.e. $\sqrt{\Lambda}L > 1/\xi$, the wave function starts oscillating, and the behavior is governed by the string coupling constant instead of the cosmological constant:

$$W(L) \sim 1/(g_s^{1/3}L)^{1/4}$$
 (116)

This drastic change happens due to the infinitely many wormholes and baby universes. The similar behavior has been observed in the context of non-critical string theory [34, 35].

From the discussion above, we observe that a sort of Coleman's mechanism works: For a large universe, the cosmological constant is not important enough to govern the physics⁶.

5 Summary

We have reviewed the relation between two-dimensional causal dynamical triangulations (2d CDT) and two-dimensional projectable Hořava-Lifshitz quantum gravity (2d projectable HL QG).

In the first part, it has been shown that the physics described by the continuum limit of 2d CDT coincides with the one obtained quantizing 2d projectable HL gravity. This is confirmed because the quantum Hamiltonians of both models are exactly the same. The system is expressed in terms of quantum mechanics of a 1d extended object, i.e. a 1d universe.

It would be too hasty to consider that this scenario also holds for the higher dimensional cases. In fact, it has been shown that in 2 + 1 dimensions numerical studies of the so-called locally causal dynamical triangulations that relax the proper time foliation of CDT and require the local causality reproduce an intriguing specialty of CDT, an emergence of the de-Sitter-like geometry [36] (see e.g. Ref. [37] for the higher-dimensional CDT). On the other hand, a Landau theory approach suggests a relation between CDT and theories invariant under the foliation-preserving diffeomorphisms in 2 + 1 dimensions [38, 39] (see also the dedicated chapter of the Handbook of Quantum Gravity [40]). This issue should be investigated further.

In the second part, we have introduced the generalized CDT (GCDT) that permits baby universes and wormholes to form in keeping with the foliation, and in particular the construction based on the string field theory for CDT has been explained. Here the string means the 1d universe, and the string field theory is constructed in such a way that the free part reproduces the CDT amplitudes, and the splitting and joining interactions of string are introduced to create baby universes and wormholes.

Focusing on the loop-to-loop amplitude, we have introduced an effective 1d theory that includes all the contributions coming from the sum over all possible baby universes and wormholes. From the point of view of HL gravity, the effective theory can be precisely reproduced if introducing a bi-local interaction term into the action of 2d projectable HL gravity and if quantizing the system. In addition, the effective theory can be also obtained considering that the cosmological constant of 2d CDT is not a constant but it fluctuates in time. This leads to Coleman's mechanism in

⁶ The Hartle-Hawking wave function W(L) is not normalizable, but similar arguments are valid on the normalizable energy eigenstates [30].

2d CDT such that for a large universe the cosmological constant is not important enough to govern the physics.

Although we have not discussed issues of the coupling to matter, 2d CDT coupled to Yang-Mills theory has been solved analytically in Ref. [41], and it has been shown that the quantum Hamiltonian obtained in Ref. [41] can be reproduced quantizing 2d projectable HL gravity coupled to Yang-Mills theory [42]. In fact, we know very little about the analytical treatment of the coupling to matter compared to the situation of 2d dynamical triangulations and the Liouville quantum gravity. This direction needs to be explored in the future.

What is remarkable is that following the standard Wilsonian renormalization group, one can take the continuum limit of the lattice model, 2d CDT, and find the continuum quantum field theory, 2d projectable HL QG, which is in the same universality class of 2d CDT. A missing piece is the continuum quantum field theory of GCDT that is described by metric components and allows us to compute all the amplitudes defined by the string field theory for CDT, although we have the effective field theory, the 2d projectable HL gravity with a bi-local interaction, that reproduces the restricted class of GCDT amplitudes once it is quantized. We wish to unveil the underlying continuum quantum field theory of GCDT, through which we can understand something inherently interesting about quantum geometries for sure.

Acknowledgements YS would like to thank Jan Ambjørn, Lisa Glaser, Yuki Hiraga, Yoshiyasu Ito and Yoshiyuki Watabiki, for wonderful collaborations on the topics related to this chapter. Great ideas appeared in this chapter attribute to them. However, if conceptual mistakes exist, YS is responsible for them.

References

- J. Ambjørn, B. Durhuus and J. Frohlich, "Diseases of Triangulated Random Surface Models, and Possible Cures," Nucl. Phys. B 257 (1985) 433. doi:10.1016/0550-3213(85)90356-6
- J. Ambjørn, B. Durhuus, J. Frohlich and P. Orland, "The Appearance of Critical Dimensions in Regulated String Theories," Nucl. Phys. B 270 (1986) 457. doi:10.1016/0550-3213(86)90563-8
- F. David, "Planar Diagrams, Two-Dimensional Lattice Gravity and Surface Models," Nucl. Phys. B 257 (1985) 45. doi:10.1016/0550-3213(85)90335-9
- A. Billoire and F. David, "Microcanonical Simulations of Randomly Triangulated Planar Random Surfaces," Phys. Lett. 168B (1986) 279. doi:10.1016/0370-2693(86)90979-2
- V. A. Kazakov, A. A. Migdal and I. K. Kostov, "Critical Properties of Randomly Triangulated Planar Random Surfaces," Phys. Lett. 157B (1985) 295. doi:10.1016/0370-2693(85)90669-0
- D. V. Boulatov, V. A. Kazakov, I. K. Kostov and A. A. Migdal, "Analytical and Numerical Study of the Model of Dynamically Triangulated Random Surfaces," Nucl. Phys. B 275 (1986) 641. doi:10.1016/0550-3213(86)90578-X
- J. Ambjørn, B. Durhuus and T. Jonsson, "Quantum Geometry : A Statistical Field Theory Approach," Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 1997.
- 8. J. Ambjørn, "Elementary Quantum Geometry," [arXiv:2204.00859 [hep-th]].

- J. Ambjørn and R. Loll, "Nonperturbative Lorentzian quantum gravity, causality and topology change," Nucl. Phys. B 536 (1998), 407-434 doi:10.1016/S0550-3213(98)00692-0 [arXiv:hepth/9805108 [hep-th]].
- T. Regge, "GENERAL RELATIVITY WITHOUT COORDINATES," Nuovo Cim. 19 (1961), 558-571 doi:10.1007/BF02733251
- A. M. Polyakov, "Quantum Geometry of Bosonic Strings," Phys. Lett. B 103 (1981), 207-210 doi:10.1016/0370-2693(81)90743-7
- V. G. Knizhnik, A. M. Polyakov and A. B. Zamolodchikov, "Fractal Structure of 2D Quantum Gravity," Mod. Phys. Lett. A 3 (1988), 819 doi:10.1142/S0217732388000982
- F. David, "Conformal Field Theories Coupled to 2D Gravity in the Conformal Gauge," Mod. Phys. Lett. A 3 (1988), 1651 doi:10.1142/S0217732388001975
- J. Distler and H. Kawai, "Conformal Field Theory and 2D Quantum Gravity," Nucl. Phys. B 321 (1989), 509-527 doi:10.1016/0550-3213(89)90354-4
- J. Ambjørn, L. Glaser, Y. Sato and Y. Watabiki, "2d CDT is 2d Hořava–Lifshitz quantum gravity," Phys. Lett. B **722** (2013), 172-175 doi:10.1016/j.physletb.2013.04.006 [arXiv:1302.6359 [hep-th]].
- P. Hořava, "Quantum gravity at a Lifshitz point," Phys. Rev. D 79 (2009) 084008, [arXiv:0901.3775, hep-th].
- B. F. Li, A. Wang, Y. Wu and Z. C. Wu, "Quantization of (1+1)-dimensional Hořava-Lifshitz theory of gravity," Phys. Rev. D 90 (2014) no.12, 124076 doi:10.1103/PhysRevD.90.124076 [arXiv:1408.2345 [hep-th]].
- L. Glaser, T. P. Sotiriou and S. Weinfurtner, "Extrinsic curvature in twodimensional causal dynamical triangulation," Phys. Rev. D 94 (2016) no.6, 064014 doi:10.1103/PhysRevD.94.064014 [arXiv:1605.09618 [hep-th]].
- P. Hořava, "Spectral Dimension of the Universe in Quantum Gravity at a Lifshitz Point," Phys. Rev. Lett. **102** (2009), 161301 doi:10.1103/PhysRevLett.102.161301 [arXiv:0902.3657 [hep-th]].
- J. Ambjørn, A. Gorlich, S. Jordan, J. Jurkiewicz and R. Loll, "CDT meets Horava-Lifshitz gravity," Phys. Lett. B 690 (2010), 413-419 doi:10.1016/j.physletb.2010.05.054 [arXiv:1002.3298 [hep-th]].
- J. Ambjørn, R. Loll, W. Westra and S. Zohren, "Putting a cap on causality violations in CDT," JHEP 12 (2007), 017 doi:10.1088/1126-6708/2007/12/017 [arXiv:0709.2784 [gr-qc]].
- J. Ambjørn, R. Loll, Y. Watabiki, W. Westra and S. Zohren, "A String Field Theory based on Causal Dynamical Triangulations," JHEP 05 (2008), 032 doi:10.1088/1126-6708/2008/05/032 [arXiv:0802.0719 [hep-th]].
- J. Ambjørn, R. Loll, Y. Watabiki, W. Westra and S. Zohren, "A New continuum limit of matrix models," Phys. Lett. B 670 (2008), 224-230 doi:10.1016/j.physletb.2008.11.003 [arXiv:0810.2408 [hep-th]].
- J. Ambjørn and T. G. Budd, "Trees and spatial topology change in CDT," J. Phys. A: Math. Theor. 46 (2013), 315201 doi:10.1088/1751-8113/46/31/315201 [arXiv:1302.1763 [hep-th]].
- J. Ambjørn, R. Loll, W. Westra and S. Zohren, "Stochastic quantization and the role of time in quantum gravity," Phys. Lett. B 680 (2009), 359-364 doi:10.1016/j.physletb.2009.09.016 [arXiv:0908.4224 [hep-th]].
- J. Ambjørn, R. Loll, W. Westra and S. Zohren, "Summing over all Topologies in CDT String Field Theory," Phys. Lett. B 678 (2009), 227-232 doi:10.1016/j.physletb.2009.06.031 [arXiv:0905.2108 [hep-th]].
- S. R. Coleman, "Why There Is Nothing Rather Than Something: A Theory of the Cosmological Constant," Nucl. Phys. B 310 (1988), 643-668 doi:10.1016/0550-3213(88)90097-1
- J. Ambjørn, Y. Hiraga, Y. Ito and Y. Sato, "Wormholes in 2d Hořava-Lifshitz quantum gravity," Phys. Lett. B 816 (2021), 136205 doi:10.1016/j.physletb.2021.136205 [arXiv:2101.07401 [hep-th]].
- J. Ambjørn, Y. Hiraga, Y. Ito and Y. Sato, "Wormhole interaction in 2d Horava-Lifshitz quantum gravity," [arXiv:2110.08479 [hep-th]].

- J. Ambjørn, Y. Sato and Y. Watabiki, "Wormholes, a fluctuating cosmological constant and the Coleman mechanism," Phys. Lett. B 815 (2021), 136152 doi:10.1016/j.physletb.2021.136152 [arXiv:2101.00478 [hep-th]].
- P. Di Francesco, E. Guitter and C. Kristjansen, "Generalized Lorentzian triangulations and the Calogero Hamiltonian," Nucl. Phys. B 608 (2001), 485-526 doi:10.1016/S0550-3213(01)00239-5 [arXiv:hep-th/0010259 [hep-th]].
- B. Durhuus and C. W. H. Lee, "A String bit Hamiltonian approach to two-dimensional quantum gravity," Nucl. Phys. B 623 (2002), 201-219 doi:10.1016/S0550-3213(01)00628-9 [arXiv:hep-th/0108149 [hep-th]].
- J. Ambjørn and C. F. Kristjansen, "Nonperturbative 2-d quantum gravity and Hamiltonians unbounded from below," Int. J. Mod. Phys. A 8 (1993), 1259-1282, [arXiv:hep-th/9205073 [hep-th]].
- 34. G. W. Moore, "Double scaled field theory at c = 1," Nucl. Phys. B 368 (1992), 557-590 doi:10.1016/0550-3213(92)90214-V
- P. Betzios and O. Papadoulaki, "Liouville theory and Matrix models: A Wheeler DeWitt perspective," JHEP 09 (2020), 125 doi:10.1007/JHEP09(2020)125 [arXiv:2004.00002 [hepth]].
- S. Jordan and R. Loll, "Causal Dynamical Triangulations without Preferred Foliation," Phys. Lett. B 724 (2013), 155-159 doi:10.1016/j.physletb.2013.06.007 [arXiv:1305.4582 [hep-th]].
- R. Loll, "Quantum Gravity from Causal Dynamical Triangulations: A Review," Class. Quant. Grav. 37 (2020) no.1, 013002 doi:10.1088/1361-6382/ab57c7 [arXiv:1905.08669 [hep-th]].
- D. Benedetti and J. Henson, "Spacetime condensation in (2+1)-dimensional CDT from a Hořava–Lifshitz minisuperspace model," Class. Quant. Grav. 32 (2015) no.21, 215007 doi:10.1088/0264-9381/32/21/215007 [arXiv:1410.0845 [gr-qc]].
- D. Benedetti and J. P. Ryan, "Capturing the phase diagram of (2 + 1)-dimensional CDT using a balls-in-boxes model," Class. Quant. Grav. 34 (2017) no.10, 105012 doi:10.1088/1361-6382/aa6b5d [arXiv:1612.09533 [hep-th]].
- 40. D. Benedetti, "Landau Theory of Causal Dynamical Triangulations," Handbook of Quantum Gravity, Springer Singapore (2023) [arXiv:2212.11043 [hep-th]].
- J. Ambjørn and A. Ipsen, "Two-dimensional causal dynamical triangulations with gauge fields," Phys. Rev. D 88 (2013) no.6, 067502 doi:10.1103/PhysRevD.88.067502 [arXiv:1305.3148 [hep-th]].
- 42. A. C. Ipsen, "Quantum gravity in two dimensions," Ph.D. Thesis.