arXiv:2212.03561v1 [hep-ph] 7 Dec 2022

Searching for $a_0(980)$-meson parton distribution function

Zai-Hui Wu, Hai-Bing Fu,* Tao Zhong,† Yu Chen, and Ya-Hong Dai
Department of Physics, Guizhou Minzu University, Guiyang 550025, P.R. China
(Dated: December 8, 2022)

In this paper, we calculate the scalar $a_0(980)$-meson leading-twist wave function by using light-cone harmonic oscillator model (LCHO). In which the model parameters are determined by fitting the ξ-moments $\langle\xi^n_{a_0}\rangle_\zeta$ of its light-cone distribution amplitudes. Then, the $a_0(980)$-meson leading-twist light-cone distribution amplitudes with three different scales $\zeta = (1.0, 2.0, 5.2)$ GeV are given. After constructing the relationship between $a_0(980)$-meson leading-twist parton distribution functions/valence quark distribution function and its LCHO wave function, we exhibit the $q^{a_0}(x,\zeta)$ and $xq^{a_0}(x,\zeta)$ with different scales. Furthermore, we also calculate the Mellin moments of the $a_0(980)$-meson's valence quark distribution function $\langle x^n q^{a_0}\rangle_\zeta$ with $n = (1, 2, 3)$, i.e. $\langle xq^{a_0}\rangle_{\zeta_5} = 0.026$, $\langle x^2 q^{a_0}\rangle_{\zeta_5} = 0.017$ and $\langle x^3 q^{a_0}\rangle_{\zeta_5} = 0.012$. Finally, the scale evolution for the ratio of the Mellin moments $\chi^n_{a_0}(\zeta,\zeta_k)$ are presented.

PACS numbers: 12.38.-t, 12.38.Bx, 14.40.Aq

1. Introduction — The exploration of quark-gluon structure in hadrons is a cross-cutting edge issue between particle physics and medium-high energy nuclear physics in recent years. In which quarks and gluons, called partons, are the fundamental degrees of freedom of quantum chromodynamics (QCD). Although the parton can not be directly observed, the QCD factorization theorem allows one to express the information of the parton inside the nucleon in terms of nonperturbative functions [1]. At the same time, the parton distribution function (PDF) is considered to be the most important nonperturbative function, which plays an important role in describing the nonperturbative QCD for the internal structure of hadronic bound states [2]. In addition, it also gives the probability of finding quarks and gluons inside a hadron. In the infinite momentum coordinate system [3–6], the PDFs are used to describe the one-dimensional momentum distributions of quarks and gluons. Therefore, the internal structure of hadrons can be studied by calculating the meson's PDF.

The PDF constitute the basic limit of Higgs boson representation in terms of coupling, which is the main system for standard model (SM) measurement such as W-boson mass. And it is also still the largest uncertainty outside the production of SM heavy particles, so it has important phenomenological value. The MMHT[7], CT [8], NNPDF [9], HERAPDF [10], and JAM [11] have made substantial efforts to determine PDFs and their uncertainties. The pion deemed to the lightest bound state of QCD and kaon have been predicted by many theoretical calculations, chiral-quark model [12–14], Nambu-Jona-Lasinio model [15], light-front holographic QCD (LHFQCD) [16–19], light front quantization [20–22], maximum entropy method [23, 24], Dyson-Schwinger equations (DSEs) [25–35] and lattice QCD [36–45] for the valence quark PDF. In most cases, the light front wave function (LFWF) is adopted to predict the PDFs. The LFWF is obtained by calculating the Bethe-Salpeter wave functions by the covariant DSEs and projecting the quantity in the Bethe-Salpeter wave functions to the light front in Ref. [34]. At the same time, other models have also analyzed LFWF, such as LHFQCD [16–19] and BLFQ [20–22]. Besides, there are few papers on the PDFs of the scalar meson $a_0(980)$ with the same quark $q\bar{q}$ state, which motivates us a research motivation.

One of the earliest predictions for $J = 0$ meson valence-quark distribution function for large-x behaviour within the QCD improved parton model [46–48], has the following expression:

$$q^M(x;\zeta) \overset{x\simeq 1}{=} c(\zeta)(1-x)^{\beta_\zeta},\ \beta_\zeta = 2, \tag{1}$$

where $c(\zeta)$ is independent of x and the ζ stands for the resolving scale. Since the meson's PDF can be obtained directly from its wave function (WF), a more accurate $a_0(980)$-meson WF is crucial for us to determine its PDF. The light-cone harmonic oscillator model (LCHO) for the light or heavy meson WF is mainly based on the Brodsky-Huang-Lepage (BHL) prescription [49], which have been used in many cases [50–58]. For this model, the total WF can be separated into spin-space WF $\chi^{\lambda_1\lambda_2}_M(x,\mathbf{k}_\perp)$ and spatial wave function $\psi^R_M(x,\mathbf{k}_\perp)$. The spatial WF is divided into x-dependence part and $\mathbf{k}_\perp$-dependence part for calculation. The $\mathbf{k}_\perp$-dependence part derives from the approximate bound-state solution and the x-dependence part $\varphi_M(x)$ can be expanded in Gegenbauer polynomials.

Furthermore, the meson light-cone distribution amplitudes (LCDAs) can also been related to its WF, which leads to the indirect relationship between meson LCDAs and PDF. In many applications of LCDAs, one usually takes a truncated form to determine DAs, involving only the first few terms of the Gegenbauer expansion series. With the increase of n, there will exist dimensional anomalies which lead to spurious oscillations. In addition, one of the most important fac-

*Electronic address: fuhb@cqu.edu.cn (Corresponding author)
†Electronic address: zhongtao1219@sina.com

tors is the unreliability of the higher-order Gegenbauer moments. In order to improve this phenomenon, one can adopt LCHO model to deal with the meson LCDAs. In our previous works [51, 59], meson's the leading-twist LCDAs are studied by using the LCHO model, and then the model parameters are determined by fitting moments with the least squares method. Therefore, the $a_0(980)$-meson PDFs will be studied based on BHL prescription in this paper.

2. Theoretical Framework — If one wants to use the typical probability expression of quantum mechanics to describe the measurable properties of a given hadron, the first thing one needs to do is finding the WF. Each element of the WF fock-space decomposition represents the probability amplitude of finding n components in the hadron. However, the PDFs describe the longitudinal momentum distribution parton of hadron. To derive the $a_0(980)$-meson leading-twist PDF, the following expression can be used [35]

$$q^{a_0}(x,\zeta) = \int_{|\mathbf{k}_\perp|^2\leq\zeta^2} d^2\mathbf{k}_\perp |\psi_{a_0}(x,\mathbf{k}_\perp)|^2. \tag{2}$$

where $\psi_{a_0}(x,\mathbf{k}_\perp)$ is WF and ζ stands for the scale. In order to calculate the PDF, the connection between distribution amplitudes and distribution function can be established to achieve the purpose. Exploiting this relationship, one can predict $a_0(980)$-meson leading-twist PDFs with the LCHO model based on BHL description [49]. The LCHO model of the $a_0(980)$-meson leading-twist WF is denoted by

$$\psi_{a_0}(x,\mathbf{k}_\perp) = \sum_{\lambda_1\lambda_2} \chi_{a_0}^{\lambda_1\lambda_2}(x,\mathbf{k}_\perp)\psi_{a_0}^R(x,\mathbf{k}_\perp), \tag{3}$$

with $\mathbf{k}_\perp$ is transverse momentum. What's more, λ_1 and λ_2 are the helicities of the two constituent quark. The spin-space WF $\chi_{a_0}^{\lambda_1\lambda_2}(x,\mathbf{k}_\perp)$ comes from the Wigner-Melosh rotation. The different forms for $\lambda_1\lambda_2$ can also be found in Ref. [56]. Thus the sum of the spin-space WF have the following form

$$\sum_{\lambda_1\lambda_2} \chi_{a_0}^{\lambda_1\lambda_2}(x,\mathbf{k}_\perp) = \frac{m_q^2}{\sqrt{\mathbf{k}_\perp^2+m_q^2}}, \tag{4}$$

with $m_q = m_u = m_d$. On the other hand, the BHL description [56, 60] proposed the assumption that the valence Fock WF depends only on the energy variable ϵ outside the shell. At the same time, the connection between the equal-time WF in the rest frame and the light-cone wave function in the infinite frame by equating the energy propagator $\epsilon = M^2 - (\sum_{i=1}^n k_i)^2$ is proposed. The propagators in different frames are as follws

$$\epsilon_1 = M^2 - \left(\sum_{i=1}^n q_i^0\right)^2, \qquad \sum_{i=1}^n \mathbf{q}^i = 0$$

$$\epsilon_2 = M^2 - \left(\sum_{i=1}^n \frac{\mathbf{k}_{\perp i}^2+m_i^2}{x_i}\right), \qquad \sum_{i=1}^n \mathbf{k}_{\perp i}^2 = 0, \quad \sum_{i=1}^n x^i = 1. \tag{5}$$

For the two-particle system, one can get

$$\mathbf{q} \leftrightarrow \frac{\mathbf{k}_{\perp i}^2+m_q^2}{4x(1-x)} - m_q^2 \tag{6}$$

where $q_1^0 = q_2^0$. Besides, there is approximate connection between $\psi_{\rm CM}(\mathbf{q})$ and $\psi_{\rm LC}(x,\mathbf{k})$,

$$\psi_{\rm CM}(\mathbf{q}^2) \leftrightarrow \psi_{\rm LC}\left(\frac{\mathbf{k}_\perp^2+m_q^2}{4x(1-x)} - m_q^2\right) \tag{7}$$

Based on the approximation for bound state solution of the meson quark model, the WF of the harmonic oscillator model in the rest frame is expressed as

$$\psi_{\rm CM}(\mathbf{q}^2) = A\exp\left(-\frac{\mathbf{q}^2}{2\beta^2}\right) \tag{8}$$

Combining Eqs. (7) and (8), the spatial WF is

$$\psi_{a_0}^R(x,\mathbf{k}_\perp) = A_{a_0}\varphi_{a_0}(x)\exp\left[-\frac{\mathbf{k}_\perp^2+m_q^2}{8\beta_{a_0}^2 x\bar{x}}\right]. \tag{9}$$

where A_{a_0} is the normalization constant and $\varphi_{a_0}(x) = (x\bar{x})^{\alpha_{a_0}} C_1^{3/2}(2x-1)$. Thus, one can get the $a_0(980)$-meson leading-twist WF

$$\psi_{a_0}(x,\mathbf{k}_\perp) = \frac{m_q^2 A_{a_0}\varphi_{a_0}(x)}{\sqrt{\mathbf{k}_\perp^2+m_q^2}}\exp\left[-\frac{\mathbf{k}_\perp^2+m_q^2}{8\beta_{a_0}^2 x\bar{x}}\right]. \tag{10}$$

Furthermore, the $a_0(980)$-meson leading-twist valence-quark distribution function can be obtained by integrating over the squared transverse momentum, i.e. Eq. (2), which leads to the following formula

$$q^{a_0}(x,\zeta) = \frac{A_{a_0}^2 m_q^2 (1-2x)^2 (x\bar{x})^{2\alpha_{a_0}}}{512\pi^5} \times\left\{\mathrm{Ei}\left[\sqrt{\frac{m_q^2+\zeta^2}{4\beta_{a_0}^2 x\bar{x}}}\right] - \mathrm{Ei}\left[\sqrt{\frac{m_q^2}{4\beta_{a_0}^2 x\bar{x}}}\right]\right\}, \tag{11}$$

with $\mathrm{Ei}(x) = -\int_{-z}^{\infty}\frac{e^{-t}}{t}dt$. It guarantees the relationship

$$\int_0^1 dx q^{a_0}(x,\zeta) = 1 \tag{12}$$

which is in agreement with the cases of pion and kaon. For further research, the process-independent effective charge is used to redesign the process-dependent-charge alternative and implement evolution to integrate the one-loop Dokshitzer-Gribov-Lipatov-Altarelli-Parisi

(DGLAP) equations with describing the evolution of quarks or gluons fragmenting into hadrons and identifying only a hadron at a time. The relevant explanation can be found in Ref. [35]. Using this process, one can get the Mellin moments of $a_0(980)$-meson's valence-quark distribution function:

$$\langle x^n q^{a_0}\rangle_\zeta = \int_0^1 x^n q^{a_0}(x,\zeta)dx \tag{13}$$

Then, the ratio of Mellin moments is also a point of interest, and its expression is as follows

$$\varkappa^n_{a_0}(\zeta,\zeta_k) = \frac{\langle x^n q^{a_0}\rangle_\zeta}{\langle x^n q^{a_0}\rangle_{\zeta_k}} \tag{14}$$

Another significant physical quantity associated with $a_0(980)$-meson PDF is its LCDA. The relationship between the $a_0(980)$-meson leading-twist LCDA and the WF is

$$\phi_{a_0}(x,\zeta) = \int_{|\mathbf{k}_\perp|^2\le\zeta^2} \frac{d^2\mathbf{k}_\perp}{16\pi^3}\psi_{a_0}(x,\mathbf{k}_\perp). \tag{15}$$

After integrating over the squared transverse momentum, one can get the LCDA formula

$$\begin{aligned}\phi_{a_0}(x,\zeta) &= \frac{A_{a_0} m_q \beta_{a_0}}{4\sqrt{2}\pi^{3/2}}\sqrt{x\bar{x}}\varphi_{a_0}(x)\\ &\times\left\{\mathrm{Erf}\left[\sqrt{\frac{m_q^2+\zeta^2}{8\beta_{a_0}^2 x\bar{x}}}\right]-\mathrm{Erf}\left[\sqrt{\frac{m_q^2}{8\beta_{a_0}^2 x\bar{x}}}\right]\right\},\end{aligned} \tag{16}$$

where $\mathrm{Erf}(x) = 2\int_0^x e^{-t^2}dx/\sqrt{\pi}$ is the error function. In order to determine the free model parameters A_{a_0}, β_{a_0} and α_{a_0}, we should use the ξ-moments of the $a_0(980)$-meson leading-twist LCDA, which has the following definition

$$\langle \xi^n_{a_0}\rangle_\zeta = \int_0^1 \xi^n \phi_{a_0}(x,\zeta)dx \tag{17}$$

On the other hand, the ξ-moments can be calculated by QCD sum rule approach within background field theory (BFTSR). The two-point correlation function is taken as

$$\Pi^{(n,0)}_{a_0} = i\int d^4x e^{iq\cdot x}\langle 0|T\{J^V_n(x), J^{S,\dagger}_0(0)\}|0\rangle, \tag{18}$$

with n will take the odd numbers, while the even order will vanish due to the G-parity. The currents are $J^V_n(x) = \bar{q}_1(x)\not{z}(iz\cdot\overleftrightarrow{D})^n q_2(x)$ and $J^S_0(0) = \bar{q}_1(0)q_2(0)$. The detail calculation process for the ξ-moments are given in our recent paper [61].

Then, one can adopt the least squares method to fit ξ-moments $\langle\xi^n_{a_0}\rangle_\zeta$ in determining the free model parameters. The purpose of the least squares method is to obtain the optimal value of the fitting parameter θ through minimizing the likelihood function

$$\chi^2(\theta) = \sum_{i=1}^N \frac{y_i-\mu(x_i,\theta)}{\sigma_i^2} \tag{19}$$

$\mu(x_i,\theta)$ is the $a_0(980)$-meson ξ-moments $\langle\xi^n_{a_0}\rangle_\zeta$ of combining Eqs. (16) and (17). The value of y_i and its variance σ_i are defined the value of ξ-moments calculated by QCD sum rule. And beyond that, making use of the probability density function $f(y,n_d) = (1/\Gamma(\frac{n_d}{2})2^{n_d/2})y^{n_d/2-1}e^{-\frac{y}{2}}$ of χ^2, one can get the goodness of fit with the following probability P_{χ^2}

$$P_{\chi^2} = \int_0^1 (f_y; n_d)dy \tag{20}$$

with $P_{\chi^2}\in(0,1)$. The closer the goodness of fit is to 1, the better the parameters are obtained. Incorporating the effect of scale ζ, according to the renormalization group equations (RGE) of Gegenbauer moments of the $a_0(980)$-meson leading-twist LCDA,

$$a^{a_0}_n(\zeta) = a^{a_0}_n(\zeta_0)E_n(\zeta,\zeta_0), \tag{21}$$

where $E_n(\zeta,\zeta_0) = [\alpha_s(\zeta)/\alpha_s(\zeta_0)]^{-(\gamma^{(0)}_n+4)/b}$ and the coefficient $b = (33-2n_f)/3$ [62]. n_f is the number of active quark flavors. The ζ_0 and ζ are considered as the initial scale and the running scale. Here we make a notation that the ξ-moments can translate into a_n-moment directly. The one-loop anomalous dimensions is

$$\gamma^{(0)}_n = C_F\left[1-\frac{2}{(n+1)(n+2)}+4\sum_{j=2}^{n+1}\frac{1}{j}\right] \tag{22}$$

with $C_F = 4/3$. Then, one can gain the ξ-moments at the arbitrary scales ζ. Finally, after fitting the moments $\langle\xi^n_{a_0}\rangle_\zeta$ with the least squares method, we can predict the $a_0(980)$-meson valence quark distribution function, Mellin moments $\langle x^n q^{a_0}\rangle_\zeta$ and ratio $\varkappa^n_{a_0}(\zeta,\zeta_k)$.

3. Numerical analysis — To do the numerical analysis, the following input parameters are used. The mass of $a_0(980)$-meson is taken as $m_{a_0} = 0.980\pm0.020$ GeV. The current light quark-mass, charm quark mass, the values of the non-perturbative vacuum condensates, the contin-

TABLE I: The $a_0(980)$-meson leading-twist LCDA moments $\langle\xi^n_{a_0}\rangle_\zeta$ at scale $\zeta = (1.0, 2.0, 5.2)$ GeV.

	ζ_0	ζ_2	ζ_5
$\langle\xi^1_{a_0}\rangle_\zeta$	-0.307 ± 0.043	-0.212 ± 0.030	-0.158 ± 0.022
$\langle\xi^3_{a_0}\rangle_\zeta$	-0.181 ± 0.034	-0.091 ± 0.017	-0.048 ± 0.009
$\langle\xi^5_{a_0}\rangle_\zeta$	-0.078 ± 0.028	-0.055 ± 0.017	-0.039 ± 0.011
$\langle\xi^7_{a_0}\rangle_\zeta$	-0.049 ± 0.026	-0.035 ± 0.015	-0.025 ± 0.010
$\langle\xi^9_{a_0}\rangle_\zeta$	-0.036 ± 0.024	-0.011 ± 0.012	0.000 ± 0.006




FIG. 1: The $a_0(980)$-meson leading-twist LCDA $\phi_{a_0}(x,\zeta)$ changed with three different scales $\zeta = (1.0, 2.0, 5.2)$ GeV.

TABLE II: The LCHO model parameters A_{a_0} (in unit: GeV^{-1}), β_{a_0} (in unit: GeV), α_{a_0} and goodness of fit $P_{\chi^2_{\rm min}}$ changed with the factorization scale $\zeta = (1.0, 2.0, 5.2)$ GeV.

ζ	A_{a_0}	β_{a_0}	α_{a_0}	$P_{\chi^2_{\rm min}}$
1.0	-260	0.5	-0.43	0.773
2.0	-380	0.5	-0.05	0.833
5.2	-1580	0.5	0.85	0.129

uum threshold s_0 and the corresponding Borel windows used in ξ-moments BFTSR are consistent with our previous work [61]. The light quark constitute mass is taken as $m_q = 250$ MeV. In this paper, we take three typical scales, the initial scales $\zeta_0 = 1.0$ GeV, the processes scale $\zeta_2 = 2.0$ GeV and the scale for π-nucleon Drell-Yan experiment [63] or the E615 experiment $\zeta_5 = 5.2$ GeV, which agree with the pion cases [35].

Firstly, we list the $a_0(980)$-meson leading-twist LCDA ξ-moments $\langle\xi^n_{a_0}\rangle_\zeta$ with three different scales $\zeta = (1.0, 2.0, 5.2)$ GeV in Table III. In which, the accurate of our calculation is up to 9th-order. It can be seen that the absolute value of ξ-moments decreases as the scale ζ increases. Secondly, the absolute value of ξ-moments decreases with the n increases, which shows that our calculation has good convergence. Then, we adopt the least squares method to fit the ξ-moments $\langle\xi^n_{a_0}\rangle_\zeta$. Next, the fitting model parameters with different scale ζ are given in Table II. Based on the experience of other mesons, we take the WF model parameter $\beta_{a_0} = 0.5$. Obviously, A_{a_0} gradually decreases with the increment of the scale ζ. However, the goodness of fit $P_{\chi^2_{\rm min}}$ increases firstly and then decreases.

With the resultant LCHO model parameters, the curves of $a_0(980)$-meson leading-twist LCDA with three scales ζ are shown in Fig. 1. The figure shows that




FIG. 2: (Color online) The $a_0(980)$-meson valence quark distribution function $q^{a_0}(x,\zeta)$ and $xq^{a_0}(x,\zeta)$ with different scales.

- The behaivor of the three curves are tend to antisymmetric, which will equal to zero when the LCDA integrate with respect to x, e.g.

$$\int_0^1 dx\phi_{a_0}(x,\zeta) = 0. \tag{23}$$

  Meanwhile, the three curves go through the zero at the location $x = 0.5$.

- The absolute value of the peaks are decreased with the increase of ζ and the x-location of the peaks are tend to 0.5 with the ζ increase. When the scale tend to infinity i.e. $\zeta \to \infty$, the curve of $a_0(980)$-meson LCDA will tend to asymptotic form $\phi_{a_0}(x,\infty) = 0$

Secondly, after taking the LCHO parameters into the $a_0(980)$-meson valence-quark distribution function, e.g. Eq. (11), the predictions of $q^{a_0}(x,\zeta)$ can be obtained. The curves of $a_0(980)$-meson valence-quark distribution

TABLE III: The Mellin moments for $a_0(980)$-meson leading-twist distribution function $\langle x^n q^{a_0}\rangle_\zeta$ with different scales $\zeta = (1.0, 2.0, 5.2)$ GeV.

ζ	$\langle xq^{a_0}\rangle_\zeta$	$\langle x^2q^{a_0}\rangle_\zeta$	$\langle x^3q^{a_0}\rangle_\zeta$
1.0	0.100	0.074	0.062
2.0	0.044	0.031	0.024
5.2	0.026	0.017	0.012




FIG. 3: (Color online) The predicted ratio of Mellin moments of the $a_0(980)$-meson valence-quark distribution function $\chi^n_{a_0}(\zeta, \zeta_k)$ with three fixed scales $\zeta_k = (1.0, 2.0, 5.2)$ GeV changed with arbitrary scales ζ in the range $\zeta = [1, 6]$ GeV. In which, the n is taken as $n = (1, 2, 3)$ respectively.

function $q^{a_0}(x,\zeta)$ and $xq^{a_0}(x,\zeta)$ with different scales ζ are shown in Fig. 2, which shows that the value of peaks decrease with the increase of scale ζ. Since the $a_0(980)$-meson leading-twist LCDA is antisymmetric behavior under $u \to (1-u)$ interchange in the $\mathrm{SU}_f(3)$ limit, its valence-quark distribution function $xq^{a_0}(x,\zeta)$ is tend to zero at $x = 0.5$. Additionally, valence-quark distribution function tends to bimodal behavior. In general, the valence-quark distribution functions of pion and kaon tend to a unimodal behavior [35].

Using the meson's valence quark distribution function, we can get the Mellin moments $\langle x^n q^{a_0}\rangle_\zeta$ of the $a_0(980)$-meson valence-quark distribution function, which are presented in Table III. From the table, we can see that the Mellin moments are convergence with the order n increased. Meanwhile, the Mellin moments are also convergence with the scale ζ increased. This agrees with the Mellin moments of pion's valence-quark distribution function decreases with the increase of scale. The greater ζ, the smaller the value of the moments in Refs. [35]. It proves our prediction of the Mellin moments $\langle x^n q^{a_0}\rangle_\zeta$ are reasonable.

Finally, we also calculate the ratio $\chi^n_{a_0}(\zeta, \zeta_k)$ of Mellin moments changed with the scale ζ. The predictions of the ratio of Mellin moments with three fixed scales $\zeta_k = (1.0, 2.0, 5.2)$ GeV and different order $n = (1, 2, 3)$ are depicted in Fig. 3. From which the $\chi^n_{a_0}(\zeta, \zeta_k)$ are increased with index n before the point of ζ_k, and decreased with n after ζ_k. The curves of $\chi^n_{a_0}(\zeta, \zeta_k)$ are decreasing with the ζ increases. The curves will coincide with each other when the scale ζ and ζ_k tend to infinity.

Summary — In this paper, we fit moments $\langle x^n_{a_0}\rangle_\zeta$ with the least squares method to obtain the free model parameters A_{a_0}, β_{a_0} and α_{a_0} at the scales $\zeta = (1.0, 2.0, 5.2)$ GeV. Meanwhile, the goodness of fit $P_{\chi^2_{\min}}$ is also given. Then, we present the curves of $a_0(980)$-meson leading-twist LCDA shown in Fig. 1. After constructing the relationship between $a_0(980)$-meson leading-twist WF and PDFs, the $a_0(980)$-meson valence quark distribution function $q^{a_0}(x,\zeta)$ and $xq^{a_0}(x,\zeta)$ with different scales are shown in Fig. 2, which tends to bimodal behavior. The LCDA and PDFs are tend to zero at the location $x = 0.5$ due to the antisymmetry of the WF. Based on the $a_0(980)$-meson valence quark distribution function, we can get the first three order Mellin moments $\langle xq^{a_0}\rangle_\zeta$ of the $a_0(980)$-meson valence quark DF shown in Table III. Referring to the predicted pion's Mellin moments, our predicted result is quite reasonable. At the same time, we also give the ratio $\chi^n_{a_0}(\zeta, \zeta_k)$ of Mellin moments with $\zeta = (1.0, 2.0, 5.2)$ GeV shown in Fig. 3. The ratio $\chi^n_{a_0}(\zeta, \zeta_k)$ shows a downward tendency with the increase of ζ.

Acknowledgments — This work was supported in part by the National Natural Science Foundation of China under Grant No.12265010, No.12265009, the Project of Guizhou Provincial Department of Science and Technology under Grant No.ZK[2021]024, the Project of Guizhou Provincial Department of Education under Grant No.KY[2021]030.

[1] J. C. Collins, D. E. Soper and G. F. Sterman, *Factorization of hard processes in QCD*, Adv. Ser. Direct. High Energy Phys. **5**, 1-91 (1989). [hep-ph/0409313]

[2] E. L. Berger and S. J. Brodsky, *Quark structure functions of mesons and the drell-yan process*, Phys. Rev. Lett. **42**, 940-944 (1979).

[3] J. D. Bjorken, *Asymptotic sum rules at infinite momentum*, Phys. Rev. **179**, 1547-1553 (1969).

[4] R. P. Feynman, *Very high-energy collisions of hadrons*,

Phys. Rev. Lett. **23**, 1415-1417 (1969).
[5] J. D. Bjorken and E. A. Paschos, *Inelastic electron proton and gamma proton scattering, and the structure of the nucleon*, Phys. Rev. **185**, 1975-1982 (1969).
[6] V. N. Gribov, *Space-time description of hadron interactions at high-energies*, [hep-ph/0006158].
[7] L. A. Harland-Lang, A. D. Martin, P. Motylinski and R. S. Thorne, *Parton distributions in the LHC era: MMHT 2014 PDFs*, Eur. Phys. J. C **75**, 204 (2015). [arXiv:1412.3989]
[8] S. Dulat, T. J. Hou, J. Gao, M. Guzzi, J. Huston, P. Nadolsky, J. Pumplin, C. Schmidt, D. Stump and C. P. Yuan, *New parton distribution functions from a global analysis of quantum chromodynamics*, Phys. Rev. D **93**, 033006 (2016). [arXiv:1506.07443]
[9] R. D. Ball *et al.* [NNPDF], *Parton distributions from high-precision collider data*, Eur. Phys. J. C **77**, 663 (2017). [arXiv:1706.00428]
[10] S. Alekhin, J. Blümlein, S. Moch and R. Placakyte, *Parton distribution functions, α_s, and heavy-quark masses for LHC run II*, Phys. Rev. D **96**, 014011 (2017). [arXiv:1701.05838]
[11] J. J. Ethier, N. Sato and W. Melnitchouk, *First simultaneous extraction of spin-dependent parton distributions and fragmentation functions from a global QCD analysis*, Phys. Rev. Lett. **119**, 132001 (2017). [arXiv:1705.05889]
[12] S. I. Nam, *Parton-distribution functions for the pion and kaon in the gauge-invariant nonlocal chiral-quark model*, Phys. Rev. D **86**, 074005 (2012). [arXiv:1205.4156]
[13] A. Watanabe, C. W. Kao and K. Suzuki, *Meson cloud effects on the pion quark distribution function in the chiral constituent quark model*, Phys. Rev. D **94**, 114008 (2016). [arXiv:1610.08817]
[14] A. Watanabe, T. Sawada and C. W. Kao, *Kaon quark distribution functions in the chiral constituent quark model*, Phys. Rev. D **97**, 074015 (2018). [arXiv:1710.09529]
[15] P. T. P. Hutauruk, I. C. Cloet and A. W. Thomas, *Flavor dependence of the pion and kaon form factors and parton distribution functions*, Phys. Rev. C **94**, 035201 (2016). [arXiv:1604.02853]
[16] G. F. de Teramond *et al.* [HLFHS], *Universality of generalized parton distributions in light-front holographic QCD*, Phys. Rev. Lett. **120**, 182001 (2018). [arXiv:1801.09154]
[17] L. Chang, K. Raya and X. Wang, *Pion parton distribution function in light-front holographic QCD*, Chin. Phys. C **44**, 114105 (2020). [arXiv:2001.07352]
[18] A. Watanabe, T. Sawada and M. Huang, *Extraction of gluon distributions from structure functions at small x in holographic QCD*, Phys. Lett. B **805**, 135470 (2020). [arXiv:1910.10008]
[19] J. Lan and C. Mondal, *Pion-nucleus induced drell-yan cross section in models inspired by light-front holography*, Phys. Lett. B **807**, 135613 (2020). [arXiv:2007.05858]
[20] J. Lan, C. Mondal, S. Jia, X. Zhao and J. P. Vary, *Parton distribution functions from a light front hamiltonian and QCD evolution for light mesons*, Phys. Rev. Lett. **122**, 172001 (2019). [arXiv:1901.11430]
[21] J. Lan, C. Mondal, S. Jia, X. Zhao and J. P. Vary, *Pion and kaon parton distribution functions from basis light front quantization and QCD evolution*, Phys. Rev. D **101**, 034024 (2020). [arXiv:1907.01509]
[22] J. Lan *et al.* [BLFQ], *Light meson parton distribution functions from basis light-front quantization and QCD evolution*, arXiv:2003.12249.
[23] C. Han, H. Xing, X. Wang, Q. Fu, R. Wang and X. Chen, *Pion valence quark distributions from maximum entropy method*, Phys. Lett. B **800**, 135066 (2020). [arXiv:1809.01549]
[24] C. Han, G. Xie, R. Wang and X. Chen, *An analysis of parton distribution functions of the pion and the kaon with the maximum entropy input*, Eur. Phys. J. C **81**, 302 (2021). [arXiv:2010.14284]
[25] L. Chang, C. Mezrag, H. Moutarde, C. D. Roberts, J. Rodríguez-Quintero and P. C. Tandy, *Basic features of the pion valence-quark distribution function*, Phys. Lett. B **737**, 23-29 (2014). [arXiv:1406.5450]
[26] L. Chang and A. W. Thomas, *Pion valence-quark parton distribution function*, Phys. Lett. B **749**, 547-550 (2015). [arXiv:1410.8250]
[27] L. Chang and C. D. Roberts, *Regarding the distribution of glue in the pion*, Chin. Phys. Lett. **38**, 081101 (2021). [arXiv:2106.08451]
[28] C. Chen, L. Chang, C. D. Roberts, S. Wan and H. S. Zong, *Valence-quark distribution functions in the kaon and pion*, Phys. Rev. D **93**, 074021 (2016). [arXiv:1602.01502]
[29] C. Shi, C. Mezrag and H. S. Zong, *Pion and kaon valence quark distribution functions from dyson-schwinger equations*, Phys. Rev. D **98**, 054029 (2018). [arXiv:1806.10232]
[30] K. D. Bednar, I. C. Cloët and P. C. Tandy, *Distinguishing quarks and gluons in pion and kaon parton distribution functions*, Phys. Rev. Lett. **124**, 042002 (2020). [arXiv:1811.12310]
[31] M. Ding, K. Raya, D. Binosi, L. Chang, C. D. Roberts and S. M. Schmidt, *Symmetry, symmetry breaking, and pion parton distributions*, Phys. Rev. D **101**, 054014 (2020). [arXiv:1905.05208]
[32] A. Freese, I. C. Cloët and P. C. Tandy, *Gluon PDF from quark dressing in the nucleon and pion*, Phys. Lett. B **823**, 136719 (2021). [arXiv:2103.05839]
[33] Z. F. Cui, M. Ding, J. M. Morgado, K. Raya, D. Binosi, L. Chang, J. Papavassiliou, C. D. Roberts, J. Rodríguez-Quintero and S. M. Schmidt, *Concerning pion parton distributions*, Eur. Phys. J. A **58**, 10 (2022). [arXiv:2112.09210]
[34] Z. F. Cui, M. Ding, J. M. Morgado, K. Raya, D. Binosi, L. Chang, F. De Soto, C. D. Roberts, J. Rodríguez-Quintero and S. M. Schmidt, *Emergence of pion parton distributions*, Phys. Rev. D **105**, L091502 (2022). [arXiv:2201.00884]
[35] Z. F. Cui, M. Ding, F. Gao, K. Raya, D. Binosi, L. Chang, C. D. Roberts, J. Rodríguez-Quintero and S. M. Schmidt, *Kaon and pion parton distributions*, Eur. Phys. J. C **80**, 1064 (2020).
[36] X. Gao, A. D. Hanlon, N. Karthik, S. Mukherjee, P. Petreczky, P. Scior, S. Shi, S. Syritsyn, Y. Zhao and K. Zhou, *Continuum-extrapolated NNLO valence PDF of pion at the physical point*, [arXiv:2208.02297].
[37] X. Gao, L. Jin, C. Kallidonis, N. Karthik, S. Mukherjee, P. Petreczky, C. Shugert, S. Syritsyn and Y. Zhao, *Valence parton distribution of the pion from lattice QCD: approaching the continuum limit*, Phys. Rev. D **102**, 094513 (2020). [arXiv:2007.06590]
[38] X. Gao, A. D. Hanlon, S. Mukherjee, P. Petreczky, P. Scior, S. Syritsyn and Y. Zhao, *Lattice QCD*

determination of the bjorken-x dependence of parton distribution functions at next-to-next-to-leading order, Phys. Rev. Lett. **128**, 142003 (2022). [arXiv:2112.02208]

[39] R. S. Sufian, C. Egerer, J. Karpie, R. G. Edwards, B. Joó, Y. Q. Ma, K. Orginos, J. W. Qiu and D. G. Richards, *Pion valence quark distribution from current-current correlation in lattice QCD*, Phys. Rev. D **102**, 054508 (2020). [arXiv:2001.04960]

[40] H. W. Lin, J. W. Chen, Z. Fan, J. H. Zhang and R. Zhang, *Valence-quark distribution of the kaon and pion from lattice QCD*, Phys. Rev. D **103**, 014516 (2021). [arXiv:2003.14128]

[41] B. Joó, J. Karpie, K. Orginos, A. V. Radyushkin, D. G. Richards, R. S. Sufian and S. Zafeiropoulos, *Pion valence structure from Ioffe-time parton pseudodistribution functions*, Phys. Rev. D **100**, 114512 (2019). [arXiv:1909.08517]

[42] C. Shugert, X. Gao, T. Izubichi, L. Jin, C. Kallidonis, N. Karthik, S. Mukherjee, P. Petreczky, S. Syritsyn and Y. Zhao, *Pion valence quark PDF from lattice QCD*, [arXiv:2001.11650].

[43] T. Izubuchi, L. Jin, C. Kallidonis, N. Karthik, S. Mukherjee, P. Petreczky, C. Shugert and S. Syritsyn, *Valence parton distribution function of pion from fine lattice*, Phys. Rev. D **100**, 034516 (2019). [arXiv:1905.06349]

[44] R. S. Sufian, J. Karpie, C. Egerer, K. Orginos, J. W. Qiu and D. G. Richards, *Pion valence quark distribution from matrix element calculated in lattice QCD*, Phys. Rev. D **99**, 074507 (2019). [arXiv:1901.03921]

[45] J. H. Zhang, J. W. Chen, L. Jin, H. W. Lin, A. Schäfer and Y. Zhao, *First direct lattice-QCD calculation of the x-dependence of the pion parton distribution function*, Phys. Rev. D **100**, 034505 (2019). [arXiv:1804.01483]

[46] Z. F. Ezawa, *Wide-Angle Scattering in Softened Field Theory*, Nuovo Cim. A **23**, 271-290 (1974).

[47] G. R. Farrar and D. R. Jackson, *Pion and Nucleon Structure Functions Near* $x = 1$, Phys. Rev. Lett. **35**, 1416 (1975).

[48] R. J. Holt and C. D. Roberts, *Distribution Functions of the Nucleon and Pion in the Valence Region*, Rev. Mod. Phys. **82**, 2991-3044 (2010). [arXiv:1002.4666]

[49] S. J. Brodsky, T. Huang, and G. P. Lepage, in Particles and Fields-2, Proceedings of the Banff Summer Institute, Ban8; Alberta, 1981, edited by A. Z. Capri and A. N. Kamal (Plenum, New York, 1983), p. 143; G. P. Lepage, S. J. Brodsky, T. Huang, and P. B.Mackenize, ibid., p. 83; T. Huang, in Proceedings of XXth International Conference on High Energy Physics, Madison, Wisconsin, 1980, edited by L. Durand and L. G Pondrom, AIP Conf. Proc. No. 69 (AIP, New York, 1981), p. 1000.

[50] T. Zhong, X. G. Wu, Z. G. Wang, T. Huang, H. B. Fu and H. Y. Han, *Revisiting the pion leading-twist distribution amplitude within the QCD background field theory*, Phys. Rev. D **90**, 016004 (2014). [arXiv:1405.0774]

[51] T. Zhong, Z. H. Zhu, H. B. Fu, X. G. Wu and T. Huang, *Improved light-cone harmonic oscillator model for the pionic leading-twist distribution amplitude*, Phys. Rev. D **104**, 016021 (2021). [arXiv:2102.03989]

[52] X. G. Wu, T. Huang and T. Zhong, *Information on the Pion Distribution Amplitude from the Pion-Photon Transition Form Factor with the Belle and BaBar Data*, Chin. Phys. C **37**, 063105 (2013). [arXiv:1206.0466]

[53] T. Huang, X. G. Wu and T. Zhong, *Finding a way to determine the pion distribution amplitude from the experimental data*, Chin. Phys. Lett. **30**, 041201 (2013). [arXiv:1303.2301]

[54] T. Huang, T. Zhong and X. G. Wu, *Determination of the pion distribution amplitude*, Phys. Rev. D **88**, 034013 (2013). [arXiv:1305.7391]

[55] T. Zhong, X. G. Wu and T. Huang, *The longitudinal and transverse distributions of the pion wave function from the present experimental data on the pion–photon transition form factor*, Eur. Phys. J. C **76**, 390 (2016). [arXiv:1510.06924]

[56] T. Huang, B. Q. Ma and Q. X. Shen, *Analysis of the pion wave function in light cone formalism*, Phys. Rev. D **49**, 1490-1499 (1994). [arXiv:9402285]

[57] T. Huang, X. G. Wu and X. H. Wu, *Pion form-factor in the* k_T *factorization formalism*, Phys. Rev. D **70**, 053007 (2004). [hep-ph/0404163]

[58] X. G. Wu and T. Huang, *Constraints on the light pseudoscalar meson distribution amplitudes from their meson-photon transition form factors*, Phys. Rev. D **84**, 074011 (2011). [arXiv:1106.4365]

[59] D. D. Hu, H. B. Fu, T. Zhong, Z. H. Wu and X. G. Wu, $a_1(1260)$*-meson longitudinal twist-2 distribution amplitude and the* $D \to a_1(1260)\ell^+\nu_\ell$ *decay processes*, Eur. Phys. J. C **82**, 603 (2022). [arXiv:2107.02758]

[60] X. H. Guo and T. Huang, *Hadronic wave functions in D and B decays*, Phys. Rev. D **43**, 2931-2938 (1991).

[61] Z. H. Wu, H. B. Fu, T. Zhong, D. Huang, D. D. Hu and X. G. Wu, $a_0(980)$*-meson twist-2 distribution amplitude within the QCD sum rules and investigation of* $D \to a_0(980)(\to \eta\pi)e^+\nu_e$, arXiv:2211.05390.

[62] H. Y. Cheng, C. K. Chua and K. C. Yang, *Charmless hadronic B decays involving scalar mesons: Implications to the nature of light scalar mesons*, Phys. Rev. D **73**, 014017 (2006). [hep-ph/0508104]

[63] J. S. Conway, C. E. Adolphsen, J. P. Alexander, K. J. Anderson, J. G. Heinrich, J. E. Pilcher, A. Possoz, E. I. Rosenberg, C. Biino and J. F. Greenhalgh, *et al. Experimental Study of Muon Pairs Produced by 252-GeV Pions on Tungsten*, Phys. Rev. D **39**, 92-122 (1989).