

Set-valued tableaux for Macdonald polynomials

Zajj Daugherty email: zdaugherty@gmail.com
 Arun Ram email: aram@unimelb.edu.au

In memory of Georgia Benkart

Abstract

Set-valued tableaux formulas play an important role in Schubert calculus. Using the box greedy reduced word for the construction of the Macdonald polynomials, we convert the alcove walk formula for Macdonald polynomials to a set-valued tableaux formula for Macdonald polynomials. Our hope is that providing set-valued tableaux formulas for Macdonald polynomials will help to strengthen the analogies and possible connections between the calculus of Macdonald polynomials and Schubert calculus.

Key words— Macdonald polynomials, tableaux formulas, Schubert calculus

1 Introduction

The goal of this paper is to give set-valued tableaux formulas for Macdonald polynomials. As evidenced by the work of [Bu02], [KMY09], [MS19], [PS19], [Wei20], [Yu21] and others, set-valued tableaux formulas play an important role in Schubert calculus and our hope is that providing set-valued tableaux formulas for Macdonald polynomials will help to strengthen the analogies and possible connections between the calculus of Macdonald polynomials and Schubert calculus.

Our set-valued tableaux formulas for Macdonald polynomials are derived by making a bijection between set-valued tableaux with alcove walks. Using this bijection and converting the alcove walks formula from [RY08, Theorem 2.2] produces a set-valued tableaux formula for Macdonald polynomials. We follow the framework of [GR21, §1 and §2], which gives an exposition of the alcove walks formula in the GL_n context and analyzes the favourite reduced for the n -periodic permutation u_μ that is used to construct the relative Macdonald polynomial E_μ^z .

Providing a set-valued tableaux formula for E_μ^z also provides a set-valued tableau formula for the symmetric Macdonald polynomials P_λ since if λ is a partition then (see [GR21b, (1.4),(1.5) and (1.10)])

$$P_\lambda = (\text{const}) \sum_{z \in S_n \lambda} E_\lambda^z.$$

The set-valued tableaux formulas given in Theorem 2.1 (respectively, Theorem 2.2) are specializable at $q = 0$ and $t = 0$ (respectively, $q = \infty$ and $t = \infty$). The specializations of E_μ^z at $t = 0$ and $t = \infty$ have interpretations in terms of crystals for level 1 and level 0 affine Demazure characters (see, for example, [Kt16, Theorem C and Corollary D] or [MRY19, (2.12), Theorem 1.1 and §2.4]). It would be interesting to write the root operators for these crystals explicitly on the set-valued tableaux (it is likely that the crystals of [Yu21, §4] cover some cases of these crystals).

2 Set-valued tableaux for Macdonald polynomials

We begin with combinatorial definitions necessary for stating the set-valued tableaux formula for Macdonald polynomials. To aid in processing these definitions, we encourage the reader to follow Example 2.1 in tandem.

Fix $n \in \mathbb{Z}_{>0}$ and let S_n denote the symmetric group on n letters. Identify $\mu = (\mu_1, \dots, \mu_n) \in \mathbb{Z}_{\geq 0}^n$ with *the set of boxes in μ* ,

$$\mu = \{(r, c) \mid r \in \{1, \dots, n\} \text{ and } c \in \{1, \dots, \mu_r\}\}.$$

2.1 Definition of the shift and height statistics

The minimal length permutation $v_\mu \in S_n$ such that $v_\mu \mu$ is weakly increasing is given by

$$v_\mu(r) = 1 + \#\{r' \in \{1, \dots, r-1\} \mid \mu_{r'} \leq \mu_r\} + \#\{r' \in \{r+1, \dots, n\} \mid \mu_{r'} < \mu_r\},$$

for $r \in \{1, \dots, n\}$. For a box $(r, c) \in \mu$ and $i \in \{1, \dots, u_\mu(r, c)\}$ define

$$\text{sh}(r, c) = \mu_r - c + 1 \quad \text{and} \quad \text{ht}(i, r, c) = v_\mu(r) - i. \quad (2.1)$$

2.2 Definition of set-valued tableaux

For $(r, c) \in \mu$ define

$$u_\mu(r, c) = \#\{r' \in \{1, \dots, r-1\} \mid \mu_{r'} < c \leq \mu_r\} + \#\{r' \in \{r+1, \dots, n\} \mid \mu_{r'} < c-1 < \mu_r\}. \quad (2.2)$$

The values $u_\mu(r, c)$ play a similar role to the flagging in the use of set-valued tableaux to give formulas for Grothendieck polynomials (see [Wei20, §2.6 and Theorem 7.1]). A *set-valued tableau T of shape μ* is a choice of subset $T(r, c) \subseteq \{1, \dots, u_\mu(r, c)\}$ for each box $(r, c) \in \mu$. More formally, a set-valued tableau T of shape μ is a function

$$T: \mu \rightarrow \{\text{subsets of } \{1, \dots, n\}\} \quad \text{such that} \quad T(r, c) \subseteq \{1, \dots, u_\mu(r, c)\}.$$

Let $|T|$ denote the total number of entries in T .

2.3 Definition of $z_{(r,c)}^T$ and x^T

Using one-line notation, let γ_n be the n -cycle in the symmetric group S_n given by

$$\gamma_n = (n \ 1 \ 2 \ \cdots \ n-1).$$

For positive integers $k_1, \dots, k_\ell$ such that $k_1 + \dots + k_\ell = n$, let

$$\gamma_{k_1} \times \cdots \times \gamma_{k_\ell} \quad \text{be the disjoint product of cycles in} \quad S_{k_1} \times \cdots \times S_{k_\ell} \subseteq S_n.$$

Fix a permutation $z \in S_n$. Order the boxes of μ down columns and then left to right (i.e., in increasing order of the values $r + nc$). Starting from the permutation z , associate a permutation $z_{(r,c)}^T$ to each box (r, c) of μ as follows:

If $(r, c) \in \mu$ and $T(r, c) = \{m_1, \dots, m_p\}$ with $1 \leq m_1 < \dots < m_p \leq u_\mu(r, c)$ define

$$z_{(r,c)}^T = z_{(r',c')}^T (\gamma_{m_1-0} \times \gamma_{m_2-m_1} \cdots \times \gamma_{m_p-m_{p-1}} \times \gamma_{u_\mu(r,c)+1-m_p} \times (\gamma_1)^{\times(n-u_\mu(r,c)-1)}) \gamma_n^{-1}, \quad (2.3)$$

where $(r', c') \in \mu$ denotes the box before (r, c) in μ . In the case that (r, c) is the first box in μ we let $z_{(r', c')}^T = z$. Then define

$$x^T = \prod_{(r, c) \in \mu} x_{z_{(r, c)}^T(n)}, \quad (2.4)$$

where $z_{(r, c)}^T(n)$ is the value of the permutation $z_{(r, c)}^T$ at n .

The permutation $z_{\text{init}}^T = z$ is the *left key* (or *initial direction*) and the permutation z_{fin}^T corresponding to the last box in μ is the *right key* (or *final direction*). These left and right keys play important roles in Schubert calculus: for example in [Yu21, §3] and in the statement of Pieri-Chevalley formulas (compare [FL94, Theorem 1] and [PR98, Corollary]).

2.4 Definition of the cov and maj statistics.

Keep the notation $T(r, c) = \{m_1, \dots, m_p\}$ with $m_1 < \dots < m_p$. Define

$$\begin{aligned} T_{<}^z(r, c) &= \{m_k \in T(r, c) \mid z_{(r, c)}^T(m_{k-1}) < z_{(r, c)}^T(m_k)\} \quad \text{and} \\ T_{>}^z(r, c) &= \{m_k \in T(r, c) \mid z_{(r, c)}^T(m_{k-1}) > z_{(r, c)}^T(m_k)\}, \end{aligned} \quad (2.5)$$

where we make the convention that $m_0 = n$. Then define

$$\begin{aligned} \text{maj}_{>}^z(T) &= \sum_{(r, c) \in \mu} \text{sh}(r, c) \cdot |T_{>}^z(r, c)|, \\ \text{cov}_{>}^z(T) &= \left(\sum_{(r, c) \in \mu} \sum_{m \in T_{>}^z(r, c)} \text{ht}(m, r, c) \right) + \frac{1}{2} \left(\ell(z_{\text{fin}}^T) - \ell(z_{\text{init}}^T v_{\mu}^{-1}) - |T| \right), \end{aligned}$$

where $\ell(w)$ denotes the length of the permutation w in S_n .

2.5 Set-valued tableaux formulas for E_{μ}^z .

The relative Macdonald polynomial $E_{\mu}^z = E_{\mu}^z(x_1, \dots, x_n; q, t)$ is what is termed the “permuted basement nonsymmetric Macdonald polynomial” in [Al16] (see [GR21, (3.7)] for further references and a definition in terms of Cherednik-Dunkl operators and the double affine Hecke algebra action on polynomials).

Theorem 2.1. *Let $z \in S_n$ and $\mu \in \mathbb{Z}_{\geq 0}^n$. Then the relative Macdonald polynomial E_{μ}^z is*

$$E_{\mu}^z = \sum_T q^{\text{maj}_{>}^z(T)} t^{\text{cov}_{>}^z(T)} \left(\prod_{(r, c) \in \mu} \prod_{m \in T(r, c)} \frac{1 - t}{1 - q^{\text{sh}(r, c)} t^{\text{ht}(m, r, c)}} \right) x^T,$$

where the sum is over set-valued tableaux T of shape μ .

Because the powers of q and t in formula in Theorem 2.1 are nonnegative integers, this formula is well suited to the specializations $q = 0$ and/or $t = 0$. The following Theorem provides an alternate formula which is better for identifying the specializations of E_{μ}^z at $q = \infty$ and/or $t = \infty$ (i.e., specializations at $q^{-1} = 0$ and $t^{-1} = 0$).

$$\begin{aligned} \text{maj}_{<}^z(T) &= \sum_{(r, c) \in \mu} \text{sh}(r, c) \cdot |T_{<}^z(r, c)|, \\ \text{cov}_{<}^z(T) &= \left(\sum_{(r, c) \in \mu} \sum_{m \in T_{<}^z(r, c)} \text{ht}(m, r, c) \right) - \frac{1}{2} \left(\ell(z_{\text{fin}}^T) - \ell(z_{\text{init}}^T v_{\mu}^{-1}) - |T| \right). \end{aligned}$$

Theorem 2.2. *Let $z \in S_n$ and $\mu \in \mathbb{Z}_{\geq 0}^n$. Then the relative Macdonald polynomial E_μ^z is*

$$E_\mu^z = \sum_T q^{-\text{maj}_z^<(T)} t^{-\text{cov}_z^<(T)} \left(\prod_{(r,c) \in \mu} \prod_{m \in T(r,c)} \frac{1 - t^{-1}}{1 - q^{-\text{sh}(r,c)} t^{-\text{ht}(m,r,c)}} \right) x^T,$$

where the sum is over set-valued tableaux T of shape μ .

The proof of Theorem 2.2 is obtained from Theorem 2.1 by multiplying numerator and denominator of each coefficient by

$$\prod_{(r,c) \in \mu} \prod_{m \in T(r,c)} q^{-\text{sh}(r,c)} t^{-\text{ht}(m,r,c)}.$$

The proof of Theorem 2.1 is obtained by considering the reduced word for the n -periodic permutation

$$u_\mu = \prod_{(r,c) \in \mu} s_{u_\mu(r,c)-1} \cdots s_2 s_1 \pi \quad (\text{product taken in increasing order of boxes})$$

studied in [GR21, Propositions 2.1 and 2.2] (see also [Mac03, (2.4.3)]) and using an easy bijection (see §3.3)

$$\{\text{alcove walks of type } \vec{u}_\mu\} \longleftrightarrow \{\text{set-valued tableaux of shape } \mu\}$$

This bijection gives a conversion between the alcove walks formula for E_μ^z in [GR21, Theorem 1.1(a)] (see also [RY08, Theorem 2.2] for this alcove walks formula in a root system language) and the set-valued tableaux formula in Theorem 2.1.

Example 2.1. Let $n = 5$ and $\mu = (0, 4, 5, 1, 4)$. Then the box arrangement associated to μ is as follows, with boxes placed on a labeled grid, filled with values $u_\mu(r, c)$, and marked in the lower corner by their cylindrical numbers $r + nc$.

		column c					
		1	2	3	4	5	
1							
2		1 7	1 12	2 17	2 22		
3		1 8	1 13	2 18	2 23	3 28	
4		1 9					
5		1 10	2 15	2 20	2 25		

(2.6)

Box by box, the corresponding factors $\frac{(1-t)}{1-q^{\text{sh}(r,c)}t^{\text{ht}(i,r,c)}}$ are

$$\frac{(1-t)}{1-q^{\text{sh}}t^{\text{ht}}} =$$

$\frac{(1-t)}{(1-q^4t^2)}$	$\frac{(1-t)}{(1-q^3t^2)}$	$\frac{(1-t)}{(1-q^2t^2)}$	$\frac{(1-t)}{(1-qt^2)}$	
$\frac{(1-t)}{(1-q^5t^4)}$	$\frac{(1-t)}{(1-q^4t^4)}$	$\frac{(1-t)}{(1-q^3t^4)}$	$\frac{(1-t)}{(1-q^2t^4)}$	$\frac{(1-t)}{(1-qt^4)}$
		$\frac{(1-t)}{(1-q^3t^3)}$	$\frac{(1-t)}{(1-q^2t^3)}$	$\frac{(1-t)}{(1-qt^3)}$
$\frac{(1-t)}{(1-qt)}$				
$\frac{(1-t)}{(1-q^4t^3)}$	$\frac{(1-t)}{(1-q^3t^3)}$	$\frac{(1-t)}{(1-q^2t^3)}$	$\frac{(1-t)}{(1-qt^3)}$	

One set-valued tableau of shape λ is

$$T =$$

$\{1\}$	$\emptyset$	$\{1\}$	$\{2\}$	
$\{1\}$	$\emptyset$	$\{1,2\}$	$\emptyset$	$\{1,3\}$
$\emptyset$				
$\emptyset$	$\emptyset$	$\{1,2\}$	$\emptyset$	

$$=$$

1		1	2	
1		12		13
		12		

(2.7)

which has size $|T| = 1 + 1 + 1 + 2 + 2 + 1 + 2 = 10$ (we shall often omit the set notation as in the right-most picture above). The product in Theorem 2.1 corresponding to this set-valued tableau T is

$$\prod_{(r,c) \in \mu} \prod_{m \in T(r,c)} \frac{(1-t)}{1-q^{\text{sh}(r,c)}t^{\text{ht}(m,r,c)}} =$$

$\frac{(1-t)}{(1-q^4t^2)}$	1	$\frac{(1-t)}{(1-q^2t^2)}$	$\frac{(1-t)}{(1-qt)}$	
$\frac{(1-t)}{(1-q^5t^4)}$	1	$\frac{(1-t)}{(1-q^3t^4)}$	$\frac{(1-t)}{(1-q^3t^3)}$	$\frac{(1-t)}{(1-qt^4)}$
1				
1	1	$\frac{(1-t)}{(1-q^2t^3)}$	$\frac{(1-t)}{(1-q^2t^2)}$	1

Let $z = \text{id}$ be the identity. The *box-by-box permutation sequence* of T with initial direction z (written in 1-line notation $w = (w(1) \cdots w(n))$) is

$$z^T =$$

<u>(23451)</u>	<u>(32451)</u>	<u>(14253)</u>	<u>(23145)</u>	
<u>(34512)</u>	<u>(34512)</u>	<u>(42531)</u>	<u>(24351)</u>	<u>(53142)</u>
<u>(35124)</u>				
<u>(31245)</u>	<u>(34125)</u>	<u>(25314)</u>	<u>(23514)</u>	

(2.8)

In each box we have underlined the positions specified by the entries of T ; these positions are the $m_1, \dots, m_p$ in Section 2.5. The red highlighted entry indicates $z_{(r,c)}^T(n)$ which is used in the formula

for x^T in Section 2.3, and also as $z_{(r,c)}^T(m_0)$ in the definition of $T_{<}^z(r, c)$ and $T_{>}^z(r, c)$ in Section 2.5. So we have

$$x^T = \begin{array}{|c|} \hline \\ \hline x_1 & x_1 & x_3 & x_5 \\ \hline x_2 & x_2 & x_1 & x_1 & x_2 \\ \hline x_4 & & & & \\ \hline x_5 & x_5 & x_4 & x_4 \\ \hline \end{array} = x_1^4 x_2^3 x_3 x_4^3 x_5^3 ;$$

and, following the definition in Section 2.5 and splitting T into *ascents* and *descents* in the sequence of underlined numbers (preceded by the red number $z_{(r,c)}^T(n)$):

$$T_{<}^z = \begin{array}{|c|} \hline \\ \hline 1 & & & \\ \hline 1 & & & 1 \\ \hline & & & \\ \hline & & 2 & \\ \hline \end{array} \quad \text{and} \quad T_{>}^z = \begin{array}{|c|} \hline \\ \hline & & 1 & 2 \\ \hline & & 2 & 3 \\ \hline & & & \\ \hline & & 1 & \\ \hline \end{array} .$$

The initial and final directions of T are

$$z_{\text{init}}^T = \text{id} = (12345) \quad \text{and} \quad z_{\text{fin}}^T = z_{(3,5)}^T = (53142),$$

where $z_{(3,5)}^T$ indicates the permutation in box $(3, 5)$ of z^T . Since $v_\mu = (13524)$ then

$$\ell(z_{\text{fin}}^T) = 4 + 2 + 0 + 1 = 7 \quad \text{and} \quad \ell(z_{\text{init}}^T v_\mu^{-1}) = 3,$$

so that

$$\frac{1}{2}(\ell(z_{\text{fin}}^T) - \ell(z_{\text{init}}^T v_\mu^{-1}) - |T|) = \frac{1}{2}(7 - 3 - 10) = -3.$$

Then

$$q^{\text{maj}_{>}^z(T)} t^{\text{cov}_{>}^z(T)} = t^{\frac{1}{2}(7-3-10)} \cdot \begin{array}{|c|} \hline \\ \hline 1 & 1 & q^2 t^2 & qt \\ \hline 1 & 1 & q^3 t^3 & 1 & qt^2 \\ \hline 1 & & & & \\ \hline 1 & 1 & q^2 t^3 & 1 \\ \hline \end{array} = t^{-3} q^9 t^{11} = q^9 t^8.$$

3 Proof of Theorem 2.1

In this section we describe the conversion from alcove walks to set-valued tableaux. We follow the framework of [GR21, §1], which gives an exposition of the alcove walks formula in the GL_n context and analyzes the favourite reduced for the n -periodic permutation u_μ that is used to construct the relative Macdonald polynomial E_μ^z .

3.1 Inversions and the box-greedy reduced word

An n -periodic permutation is a bijection $w: \mathbb{Z} \rightarrow \mathbb{Z}$ such that $w(i+n) = w(i) + n$. Since an n -periodic permutation w is determined by the values $w(1), \dots, w(n)$ then a permutation $w \in S_n$ extends uniquely to an n -periodic permutation. For $i \in \{1, \dots, n-1\}$, let s_i be the transposition in S_n switching i and $i+1$. Define $\pi: \mathbb{Z} \rightarrow \mathbb{Z}$ by $\pi(i) = i+1$.

An *inversion* of an n -periodic permutation w is an element of the set

$$\text{Inv}(w) = \{(i, k) \mid i \in \{1, \dots, n\}, k \in \mathbb{Z}, i < k \text{ and } w(i) > w(k)\}.$$

If $i, j \in \{1, \dots, n\}$ with $i < j$ and $\ell \in \mathbb{Z}$ then the *shift* and *height* of $(i, j + \ell n)$ are defined to be

$$\text{sh}(i, j + \ell n) = \ell \quad \text{and} \quad \text{ht}(i, j + \ell n) = j - i.$$

Let $\mu = (\mu_1, \dots, \mu_n) \in \mathbb{Z}_{\geq 0}^n$ and let u_μ be the n -periodic permutation defined by

$$u_\mu(i) = v_\mu^{-1}(i) + n\mu_i, \quad \text{for } i \in \{1, \dots, n\}. \quad (3.1)$$

Let $u_\mu(r, c)$ be as defined in (2.2). Following [GR21, Prop. 2.2(1)], the *box-greedy reduced word* for u_μ (an abstract word in symbols $s_1, \dots, s_{n-1}$ and π) is

$$u_\mu^\square = \prod_{\text{boxes } (r, c) \text{ in } dg(\mu)} (s_{u_\mu(r, c)} \cdots s_2 s_1 \pi). \quad (3.2)$$

and the product is taken in increasing order of the values $r + nc$. As in [GR21, (2.12)], this reduced word induces a partition of the elements of $\text{Inv}(u_\mu)$ according to the boxes (r, c) of μ , and an ordering of the inversions in each box. For $(r, c) \in \mu$ and $i \in \{1, \dots, u_\mu(r, c)\}$,

$$\text{the } i\text{th inversion in the box } (r, c) \text{ is } \beta(i, r, c) = (v_\mu(r), i + n \cdot (\mu_r - c + 1)).$$

The inversion $\beta(i, r, c)$ has shift and height

$$\text{sh}(r, c) = \text{sh}(\beta(i, r, c)) = \mu_r - c + 1 \quad \text{and} \quad \text{ht}(i, r, c) = \text{ht}(\beta(i, r, c)) = v_\mu(r) - i. \quad (3.3)$$

Example 3.1. If $\mu = (0, 4, 5, 1, 4)$ then $n = 5$, the box-greedy reduced word for u_μ is

$$u_\mu^\square = (s_1\pi)^6 (s_2s_1\pi)^7 (s_3s_2s_1\pi) =$$

$s_1\pi$	$s_1\pi$	$s_2s_1\pi$	$s_2s_1\pi$	
$s_1\pi$	$s_1\pi$	$s_2s_1\pi$	$s_2s_1\pi$	$s_3s_2s_1\pi$
$s_1\pi$				
$s_1\pi$	$s_2s_1\pi$	$s_2s_1\pi$	$s_2s_1\pi$	

(3.4)

and the inversion set of u_μ is

$$\text{Inv}(u_\mu) =$$

$(3, 1 + 4 \cdot 5)$	$(3, 1 + 3 \cdot 5)$	$(3, 1 + 2 \cdot 5)$ $(3, 2 + 2 \cdot 5)$	$(3, 1 + 1 \cdot 5)$ $(3, 2 + 1 \cdot 5)$	
$(5, 1 + 5 \cdot 5)$	$(5, 1 + 4 \cdot 5)$	$(5, 1 + 3 \cdot 5)$ $(5, 2 + 3 \cdot 5)$	$(5, 1 + 2 \cdot 5)$ $(5, 2 + 2 \cdot 5)$	$(5, 1 + 1 \cdot 5)$ $(5, 2 + 1 \cdot 5)$ $(5, 3 + 1 \cdot 5)$
$(2, 1 + 1 \cdot 5)$				
$(4, 1 + 4 \cdot 5)$	$(4, 1 + 3 \cdot 5)$ $(4, 2 + 3 \cdot 5)$	$(4, 1 + 2 \cdot 5)$ $(4, 2 + 2 \cdot 5)$	$(4, 1 + 1 \cdot 5)$ $(4, 2 + 1 \cdot 5)$	

3.2 Alcove walks and permutation sequences

Let $z \in S_n$ and choose a subset F of the s_i factors in $u_\mu^\square = w_1 w_2 \cdots w_r$ to cross out. The corresponding *alcove walk of type* $(z, u_\mu^\square)$ is the sequence $p(F) = (p_0, p_1, \dots, p_r)$ of n -periodic permutations given by

$$p_0 = z \quad \text{and} \quad p_k = \begin{cases} p_{k-1}\pi, & \text{if } w_k = \pi, \\ p_{k-1}, & \text{if } w_k \in F \text{ (so that the factor } w_k \text{ is crossed out),} \\ p_{k-1}w_k, & \text{if } w_k \notin F \text{ (so that the factor } w_k \text{ is not crossed out).} \end{cases}$$

Following [GR21, (1.14)], the *permutation sequence of* $p(F)$ is the sequence of permutations in S_n given by

$$\bar{p}(F) = (\bar{p}_0, \bar{p}_1, \dots, \bar{p}_r), \quad \text{where } \bar{w}(i) = w(i) \bmod n.$$

Note that $\overline{w_1 w_2} = \bar{w}_1 \bar{w}_2$ and $\bar{\pi} = c_n = s_1 \cdots s_{n-1}$ and $\bar{s}_j = s_j$ for $j \in \{1, \dots, n-1\}$.

3.3 Converting to set-valued tableaux

The bijection between alcove walks and set-valued tableaux is

$$\begin{array}{ccc} \{\text{alcove walks of type } u_\mu^\square\} & \longleftrightarrow & \{\text{set-valued tableaux of shape } \mu\} \\ p(F) & \longmapsto & T \end{array}$$

where, in the set-valued tableau T corresponding to the alcove walk $p(F)$, the set $T(r, c)$ in box (r, c) specifies which factors are crossed out in that box: if a box in T contains i then delete s_i in the corresponding word.

Example 3.2. In the case $\mu = (0, 4, 5, 1, 4)$, where the box-greedy reduced word for u_μ is as given in (3.4), there are $5 + 2 \cdot 8 + 3 = 24$ factors of the form s_i in $u_\mu^\square$ and so there are a total of 2^{24} alcove walks of type $(z, u_\mu^\square)$ (for any fixed permutation $z \in S_n$). Each alcove walk corresponds to a choice of s_i factors in $u_\mu^\square$ to cross out.

The tableau T in Example 2.1 corresponds to the subset

$$F = \begin{array}{|c|c|c|c|c|} \hline & & & & \\ \hline \cancel{s_1}\pi & s_1\pi & s_2\cancel{s_1}\pi & \cancel{s_1}s_1\pi & \\ \hline \cancel{s_1}\pi & s_1\pi & \cancel{s_1}\cancel{s_2}\pi & s_2s_1\pi & \cancel{s_1}s_2\cancel{s_1}\pi \\ \hline s_1\pi & & & & \\ \hline s_1\pi & s_2s_1\pi & \cancel{s_1}\cancel{s_2}\pi & s_2s_1\pi & \\ \hline \end{array}, \quad (3.5)$$

which has alcove walk

$$p(F) = (p_0, p_1, \dots, p_{37}) = (z, z, z\pi, z\pi, z\pi^2, z\pi^2s_1, z\pi^2s_1\pi, z\pi^2s_1\pi s_1, z\pi^2s_1\pi s_1\pi, \dots)$$

(there is a repeat entry in $p(F)$ each time there is an s_i crossed out in F). Using one line notation $w = (w(1) \ w(2) \ \cdots \ w(n))$ for n -periodic permutations (underlining 2-digit terms for emphasis), a

box-by-box formulation in the case that $z = \text{id} = (12345)$ is

$$p(F) = \begin{array}{|c|c|c|c|c|} \hline & & & & \\ \hline (1\ 2\ 3\ 4\ 5) & (6\ 3\ 7\ 9\ \underline{10}) & (3\ \underline{11}\ 9\ \underline{12}\ \underline{15}) & (\underline{12}\ \underline{15}\ 8\ \underline{16}\ \underline{14}) & \\ (2\ 3\ 4\ 5\ 6) & (3\ 7\ 9\ \underline{10}\ \underline{11}) & (3\ \underline{11}\ 9\ \underline{12}\ \underline{15}) & (\underline{15}\ \underline{12}\ 8\ \underline{16}\ \underline{14}) & \\ & & (\underline{11}\ 9\ \underline{12}\ \underline{15}\ 8) & (\underline{12}\ 8\ \underline{16}\ \underline{14}\ \underline{20}) & \\ \hline (2\ 3\ 4\ 5\ 6) & (7\ 3\ 9\ \underline{10}\ \underline{11}) & (\underline{11}\ 9\ \underline{12}\ \underline{15}\ 8) & (\underline{12}\ \underline{16}\ 8\ \underline{14}\ \underline{20}) & (\underline{12}\ 8\ \underline{20}\ \underline{21}\ \underline{19}) \\ (3\ 4\ 5\ 6\ 7) & (3\ 9\ \underline{10}\ \underline{11}\ \underline{12}) & (\underline{11}\ 9\ \underline{12}\ \underline{15}\ 8) & (\underline{16}\ \underline{12}\ 8\ \underline{14}\ \underline{20}) & (\underline{12}\ \underline{20}\ 8\ \underline{21}\ \underline{19}) \\ & & (9\ \underline{12}\ \underline{15}\ 8\ \underline{16}) & (\underline{12814}\ \underline{20}\ \underline{21}) & (\underline{12}\ \underline{20}\ 8\ \underline{21}\ \underline{19}) \\ & & & & (\underline{20}\ 8\ \underline{21}\ \underline{19}\ \underline{17}) \\ \hline (4\ 3\ 5\ 6\ 7) & & & & \\ (3\ 5\ 6\ 7\ 9) & & & & \\ \hline (5\ 3\ 6\ 7\ 9) & (3\ \underline{10}\ 9\ \underline{11}\ \underline{12}) & (9\ \underline{12}\ \underline{15}\ 8\ \underline{16}) & (\underline{12}\ \underline{14}\ 8\ \underline{20}\ \underline{21}) & \\ (3\ 6\ 7\ 9\ \underline{10}) & (\underline{10}\ 3\ 9\ \underline{11}\ \underline{12}) & (9\ \underline{12}\ \underline{15}\ 8\ \underline{16}) & (\underline{14}\ \underline{12}\ 8\ \underline{20}\ \underline{21}) & \\ & (3\ 9\ \underline{11}\ \underline{12}\ \underline{15}) & (\underline{12}\ \underline{15}\ 8\ \underline{16}\ \underline{14}) & (\underline{12}\ 8\ \underline{20}\ \underline{21}\ \underline{19}) & \\ \hline \end{array}$$

The permutation sequence is

$$\bar{p}(F) = (\bar{p}_0, \bar{p}_1, \dots, \bar{p}_{37}) = (z, z, zc_n, zc_n, zc_n^2, zc_n^2s_1, zc_n^2s_1c_n, zc_n^2s_1c_ns_1, zc_n^2s_1c_ns_1c_n, \dots)$$

(obtained by replacing each π by the n -cycle $c_n = s_1 \cdots s_{n-1} = \gamma_n^{-1}$). Namely, $\bar{p}(F)$ is obtained by taking all the values in $p(F) \bmod n$ (here $n = 5$ for $\mu = (0, 4, 1, 5, 4)$).

$$\bar{p}(F) = \begin{array}{|c|c|c|c|c|} \hline & & & & \\ \hline (\underline{1}2345) & (13245) & (31425) & (2\underline{5}314) & \\ (23451) & (32451) & (\underline{3}1425) & (52314) & \\ & & (14253) & (23145) & \\ \hline (\underline{2}3451) & (23451) & (1\underline{4}253) & (21345) & (23\underline{5}14) \\ (34512) & (34512) & (\underline{1}4253) & (12345) & (25314) \\ & & (42531) & (24351) & (\underline{2}5314) \\ & & & & (53142) \\ \hline (43512) & & & & \\ (35124) & & & & \\ \hline (53124) & (35412) & (4\underline{2}531) & (24351) & \\ (31245) & (53412) & (\underline{4}2531) & (42351) & \\ & (34125) & (25314) & (23514) & \\ \hline \end{array} \tag{3.6}$$

The *box-by-box permutation sequence* of F is obtained by recording the last permutation in each box, which is z^T in (2.8). The elements $z_{(r,c)}^T$ in each box of z^T are also obtained from $z = z_{\text{init}}^T$ and T by the formula in Section 2.3.

We have used the blue highlighted numbers in (3.6) to record the positions $i, i+1$ when there is a repeat entry in the sequence $\bar{p}(F)$ coming from a crossed out s_i . These record the information of F . This same information is translated into the underlines in z^T in (2.8), and the underlines exactly specify the entries of T . The set-valued tableau indicating the positions of the underlined numbers in each box is the tableau in (2.7).

3.4 Conversion of the statistics

The proof of Theorem 2.1 is completed by matching up the statistics in Theorem 2.1 with the statistics that appear in [RY08, Theorem 3.1]. In Example 3.2, we have used the blue highlighted numbers

in (3.6) and the underlines in (2.8) to illustrate how the information of the ‘folds’ in [RY08, Theorem 3.1] is equivalent to the information of the set valued tableau T . In the context of [RY08, (2.36)], the set $T(r, c)$ exactly records the folds coming from the box (r, c) . By (3.3), the factor $\prod_{(r,c) \in \mu} \prod_{m \in T(r,c)} \frac{(1-t)}{1-q^{\text{sh}(r,c)} t^{\text{ht}(m,r,c)}}$ which appears in the product in Theorem 2.1 corresponds to

$$\text{the factors} \quad \left(\prod_{k \in f^+(p)} \frac{(1-t_{\beta_k^\vee})}{1-q^{\langle -\beta_k^\vee, \rho_c \rangle}} \right) \left(\prod_{k \in f^-(p)} \frac{(1-t_{\beta_k^\vee})}{1-q^{\langle -\beta_k^\vee, \rho_c \rangle}} \right) \quad \text{in [RY08, Theorem 3.1].}$$

The sets $T_{\leq}^z$ and $T_{\geq}^z$ correspond to the sets $f^+(p)$ and $f^-(p)$ of positive and negative folds in [RY08, Theorem 3.1]. The permutation z_{fin}^T is the permutation denoted $\varphi(p)$ in [RY08, Theorem 3.1] (denoted z_r in [GR21, (1.14)]), and the permutation $z_{\text{init}}^T v_\mu^{-1}$ is the permutation m of [RY08, Theorem 3.1]. With these conversions, the last term $\frac{1}{2}(\ell(z_{\text{fin}}^T) - \ell(z_{\text{init}}^T v_\mu^{-1}) - |T|)$ in the definition of $\text{cov}_{\varphi(p)}^z(T)$ in Section 2.5 corresponds to the combination of the factors $t_{\varphi(p)}^{\frac{1}{2}}$ and $t_{\beta_k^\vee}^{-\frac{1}{2}}$ in [RY08, Theorem 3.1] and the factor $t_m^{\frac{1}{2}}$ in [RY08, Remark 3.2].

Example 3.3. Suppose that $n = 14$, $\mu = (\mu_1, \dots, \mu_{14}) \in (\mathbb{Z}_{\geq 0})^{14}$, (r, c) is a box in μ and (r', c') is the box before (r, c) in μ . Say

$$u_{(r,c)} = 11, \quad T(r, c) = \{3, 4, 8, 10\} \quad \text{and} \quad z_{(r',c')}^T = (u_1 u_2 u_3 u_4 u_5 u_6 u_7 u_8 u_9 u_{10} u_{11} u_{12} u_{13} u_{14}),$$

so that $u_i = z_{(r',c')}^T(i)$ for $i \in \{1, \dots, 14\}$. Then the box (r, c) of F contains

$$\begin{aligned} s_{11} \cancel{s_{10}} s_9 \cancel{s_8} s_7 s_6 s_5 \cancel{s_4} \cancel{s_3} s_2 s_1 \pi &= (\gamma_3 \times \gamma_1 \times \gamma_4 \times \gamma_2 \times \gamma_2 \times \gamma_1 \times \gamma_1) \pi \\ &= (\gamma_{3-0} \times \gamma_{4-3} \times \gamma_{8-4} \times \gamma_{10-8} \times \gamma_{12-10} \times (\gamma_1)^{\times(14-12)}) \pi \end{aligned}$$

and the entries in box (r, c) of the permutation sequence $\bar{p}(F)$ are

$$\begin{aligned} z_{(r',c')}^T s_{11} &= z_{(r',c')}^T s_{11} \cancel{s_{10}} = (u_1 u_2 u_3 u_4 u_5 u_6 u_7 u_8 u_9 u_{10} u_{11} u_{13} u_{14}) \\ z_{(r',c')}^T s_{11} \cancel{s_{10}} s_9 &= z_{(r',c')}^T s_{11} \cancel{s_{10}} s_9 \cancel{s_8} = (u_1 u_2 u_3 u_4 u_5 u_6 u_7 u_8 u_{10} u_9 u_{12} u_{11} u_{13} u_{14}) \\ z_{(r',c')}^T s_{11} \cancel{s_{10}} s_9 \cancel{s_8} s_7 &= (u_1 u_2 u_3 u_4 u_5 u_6 u_8 u_7 u_{10} u_9 u_{12} u_{11} u_{13} u_{14}) \\ z_{(r',c')}^T s_{11} \cancel{s_{10}} s_9 \cancel{s_8} s_7 s_6 &= (u_1 u_2 u_3 u_4 u_5 u_8 u_6 u_7 u_{10} u_9 u_{12} u_{11} u_{13} u_{14}) \\ z_{(r',c')}^T s_{11} \cancel{s_{10}} s_9 \cancel{s_8} s_7 s_6 s_5 &= (u_1 u_2 u_3 u_4 u_8 u_5 u_6 u_7 u_{10} u_9 u_{12} u_{11} u_{13} u_{14}) \\ z_{(r',c')}^T s_{11} \cancel{s_{10}} s_9 \cancel{s_8} s_7 s_6 s_5 \cancel{s_4} \cancel{s_3} s_2 &= (u_1 u_3 u_2 u_4 u_8 u_5 u_6 u_7 u_{10} u_9 u_{12} u_{11} u_{13} u_{14}) \\ z_{(r',c')}^T s_{11} \cancel{s_{10}} s_9 \cancel{s_8} s_7 s_6 s_5 \cancel{s_4} \cancel{s_3} s_2 s_1 &= (u_3 u_1 u_2 u_4 u_8 u_5 u_6 u_7 u_{10} u_9 u_{12} u_{11} u_{13} u_{14}) \\ z_{(r,c)}^T &= (u_1 u_2 \underline{u_4 u_8} u_5 u_6 u_7 \underline{u_{10}} u_9 \underline{u_{12}} u_{11} u_{13} u_{14} \mathbf{u_3}) \end{aligned}$$

and

$$\begin{aligned} z_{(r,c)}^T &= z_{(r',c')}^T s_{11} \cancel{s_{10}} s_9 \cancel{s_8} s_7 s_6 s_5 \cancel{s_4} \cancel{s_3} s_2 s_1 \gamma_{14}^{-1} \\ &= z_{(r',c')}^T (\gamma_3 \times \gamma_1 \times \gamma_4 \times \gamma_2 \times \gamma_2 \times \gamma_1 \times \gamma_1) \gamma_{14}^{-1} \\ &= z_{(r',c')}^T (\gamma_{3-0} \times \gamma_{4-3} \times \gamma_{8-4} \times \gamma_{10-8} \times \gamma_{12-10} \times (\gamma_1)^{\times(14-12)}) \gamma_{14}^{-1} \end{aligned}$$

so that $z_{(r,c)}^T(n) = u_3$, $z_{(r,c)}^T(3) = u_4$, $z_{(r,c)}^T(4) = u_8$, $z_{(r,c)}^T(8) = u_{10}$, $z_{(r,c)}^T(10) = u_{12}$. Then $T_{<}^z(r, c)$ and $T_{>}^z(r, c)$ are determined by

$$\begin{array}{llll} \text{if } u_3 < u_4 & \text{then} & 3 \in T_{<}^z(r, c) & \text{and otherwise } 3 \in T_{>}^z(r, c); \\ \text{if } u_4 < u_8 & \text{then} & 4 \in T_{<}^z(r, c) & \text{and otherwise } 4 \in T_{>}^z(r, c); \\ \text{if } u_8 < u_{10} & \text{then} & 8 \in T_{<}^z(r, c) & \text{and otherwise } 8 \in T_{>}^z(r, c); \\ \text{if } u_{10} < u_{12} & \text{then} & 10 \in T_{<}^z(r, c) & \text{and otherwise } 10 \in T_{>}^z(r, c). \end{array}$$

4 Example: Computing $E_{(2,2,1,1,0,0)}$

Here we compute $E_{(2,2,1,1,0,0)} = E_{(2,2,1,1,0,0)}^{\text{id}}$ using the set-valued tableaux formula of Theorem ?? . For comparison, this same Macdonald polynomial is computed in terms of nonattacking fillings in [GR21, example following Theorem 1.1] and in terms of multiline queues in [CMW18, Figure 4].

Let

$$z = \text{id} = (123456), \quad \text{and} \quad \mu = (2, 2, 1, 1, 0, 0) \quad \text{with} \quad v_\mu = (563412) \quad \text{and} \quad \ell(v_\mu) = 12.$$

Then

$$u_\mu = \pi^4(s_2 s_1 \pi)^2 = \begin{array}{|c|c|} \hline \pi & s_2 s_1 \pi \\ \hline \pi & s_2 s_1 \pi \\ \hline \pi & \\ \hline \pi & \\ \hline \end{array} \quad \text{and} \quad \text{Inv}(u_\mu) = \begin{array}{|c|c|} \hline \emptyset & (5, 1 + 1 \cdot 6) \\ \hline \emptyset & (5, 2 + 1 \cdot 6) \\ \hline \emptyset & (6, 1 + 1 \cdot 6) \\ \hline \emptyset & (6, 2 + 1 \cdot 6) \\ \hline \emptyset & \\ \hline \emptyset & \\ \hline \end{array}$$

Box by box, the corresponding factors $\frac{(1-t)}{1 - q^{\text{sh}(r,c)} t^{\text{ht}(i,r,c)}}$ are

$$\frac{(1-t)}{1 - q^{\text{sh}} t^{\text{ht}}} = \begin{array}{|c|c|} \hline & \frac{(1-t)}{(1-qt^4)} \\ \hline & \frac{(1-t)}{(1-qt^3)} \\ \hline & \frac{(1-t)}{(1-qt^4)} \\ \hline & \frac{(1-t)}{(1-qt^3)} \\ \hline & \\ \hline & \\ \hline \end{array}$$

For a set valued tableau T let

$$\text{wt}(T) = q^{\text{maj}_{>}^z(T)} t^{\text{cov}_{>}^z(T)} \left(\prod_{(r,c) \in \mu} \prod_{m \in T(r,c)} \frac{1-t}{1 - q^{\text{sh}(r,c)} t^{\text{ht}(m,r,c)}} \right)$$

where

$$\text{cov}_{>}^z(T) = \left(\sum_{(r,c) \in \mu} \sum_{m \in T_{>}^z(r,c)} \text{ht}(m, r, c) \right) + \frac{1}{2} \left(\ell(z_{\text{fin}}^T) - \ell(z_{\text{init}}^T v_\mu^{-1}) - |T| \right).$$

For this example, $\text{sh}(1, 2) = \text{sh}(2, 2) = 1$ so that $\text{maj}_{>}^z(T) = |T_{>}^z(r, c)|$.

There are 16 terms in the expansion of $E_{(2,2,1,1,0,0)}$ corresponding to the following permutation sequences z^T .

$$z^{(1)} = \begin{array}{|c|c|} \hline (23456\mathbf{1}) & (\underline{6}1234\mathbf{5}) \\ \hline (34561\mathbf{2}) & (\underline{1}2345\mathbf{6}) \\ \hline (45612\mathbf{3}) & \\ \hline (56123\mathbf{4}) & \\ \hline \end{array} \quad \text{with} \quad \begin{aligned} x^T &= x_1x_2x_3x_4x_5x_6 \\ \text{wt}(T) &= q^2 \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^5)} \\ \text{cov}_{\geq}^z(T) &= 3 + 5 + \frac{1}{2}(0 - 12 - 4) = 0. \end{aligned}$$

$$z^{(2)} = \begin{array}{|c|c|} \hline (23456\mathbf{1}) & (\underline{6}1234\mathbf{5}) \\ \hline (34561\mathbf{2}) & (\underline{2}1345\mathbf{6}) \\ \hline (45612\mathbf{3}) & \\ \hline (56123\mathbf{4}) & \\ \hline \end{array} \quad \text{with} \quad \begin{aligned} x^T &= x_1x_2x_3x_4x_5x_6 \\ \text{wt}(T) &= q^2t \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^5)} \\ \text{cov}_{\geq}^z(T) &= 3 + 5 + \frac{1}{2}(1 - 12 - 3) = 1. \end{aligned}$$

$$z^{(3)} = \begin{array}{|c|c|} \hline (23456\mathbf{1}) & (\underline{6}1234\mathbf{5}) \\ \hline (34561\mathbf{2}) & (\underline{6}2345\mathbf{1}) \\ \hline (45612\mathbf{3}) & \\ \hline (56123\mathbf{4}) & \\ \hline \end{array} \quad \text{with} \quad \begin{aligned} x^T &= x_1x_2x_3x_4x_5x_1 \\ \text{wt}(T) &= q \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^4)} \\ \text{cov}_{\geq}^z(T) &= 3 + \frac{1}{2}(9 - 12 - 3) = 0. \end{aligned}$$

$$z^{(4)} = \begin{array}{|c|c|} \hline (23456\mathbf{1}) & (\underline{6}1234\mathbf{5}) \\ \hline (34561\mathbf{2}) & (61345\mathbf{2}) \\ \hline (45612\mathbf{3}) & \\ \hline (56123\mathbf{4}) & \\ \hline \end{array} \quad \text{with} \quad \begin{aligned} x^T &= x_1x_2x_3x_4x_5x_2 \\ \text{wt}(T) &= q \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^4)} \\ \text{cov}_{\geq}^z(T) &= 3 + \frac{1}{2}(8 - 12 - 2) = 0. \end{aligned}$$

$$z^{(5)} = \begin{array}{|c|c|} \hline (23456\mathbf{1}) & (\underline{1}6234\mathbf{5}) \\ \hline (34561\mathbf{2}) & (\underline{6}2345\mathbf{1}) \\ \hline (45612\mathbf{3}) & \\ \hline (56123\mathbf{4}) & \\ \hline \end{array} \quad \text{with} \quad \begin{aligned} x^T &= x_1x_2x_3x_4x_5x_1 \\ \text{wt}(T) &= q^2t^5 \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^5)} \\ \text{cov}_{\geq}^z(T) &= 4 + 4 + \frac{1}{2}(9 - 12 - 3) = 5. \end{aligned}$$

$z^{(6)} =$ <table style="border-collapse: collapse; margin-left: 10px;"> <tr><td style="border: 1px solid black; padding: 2px;">(234561)</td><td style="border: 1px solid black; padding: 2px;">(<u>1</u>62345)</td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(345612)</td><td style="border: 1px solid black; padding: 2px;">(2<u>6</u>3451)</td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(456123)</td><td></td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(561234)</td><td></td></tr> <tr><td style="border: 1px solid black; padding: 2px;"> </td><td></td></tr> </table>	(23456 1)	(<u>1</u> 6234 5)	(34561 2)	(2 <u>6</u> 345 1)	(45612 3)		(56123 4)				with	$x^T = x_1 x_2 x_3 x_4 x_5 x_1$ $\text{wt}(T) = qt \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^5)}$ $\text{cov}_{>}^z(T) = 4 + \frac{1}{2}(8 - 12 - 2) = 1.$
(23456 1)	(<u>1</u> 6234 5)											
(34561 2)	(2 <u>6</u> 345 1)											
(45612 3)												
(56123 4)												
$z^{(7)} =$ <table style="border-collapse: collapse; margin-left: 10px;"> <tr><td style="border: 1px solid black; padding: 2px;">(234561)</td><td style="border: 1px solid black; padding: 2px;">(<u>1</u>62345)</td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(345612)</td><td style="border: 1px solid black; padding: 2px;">(1<u>2</u>3456)</td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(456123)</td><td></td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(561234)</td><td></td></tr> <tr><td style="border: 1px solid black; padding: 2px;"> </td><td></td></tr> </table>	(23456 1)	(<u>1</u> 6234 5)	(34561 2)	(1 <u>2</u> 345 6)	(45612 3)		(56123 4)				with	$x^T = x_1 x_2 x_3 x_4 x_5 x_6$ $\text{wt}(T) = q^2 t \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^4)}$ $\text{cov}_{>}^z(T) = 4 + 4 + \frac{1}{2}(0 - 12 - 2) = 1.$
(23456 1)	(<u>1</u> 6234 5)											
(34561 2)	(1 <u>2</u> 345 6)											
(45612 3)												
(56123 4)												
$z^{(8)} =$ <table style="border-collapse: collapse; margin-left: 10px;"> <tr><td style="border: 1px solid black; padding: 2px;">(234561)</td><td style="border: 1px solid black; padding: 2px;">(<u>1</u>62345)</td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(345612)</td><td style="border: 1px solid black; padding: 2px;">(513462)</td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(456123)</td><td></td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(561234)</td><td></td></tr> <tr><td style="border: 1px solid black; padding: 2px;"> </td><td></td></tr> </table>	(23456 1)	(<u>1</u> 6234 5)	(34561 2)	(51346 2)	(45612 3)		(56123 4)				with	$x^T = x_1 x_2 x_3 x_4 x_5 x_2$ $\text{wt}(T) = qt \frac{(1-t)}{(1-qt^4)}$ $\text{cov}_{>}^z(T) = 4 + \frac{1}{2}(7 - 12 - 1) = 1.$
(23456 1)	(<u>1</u> 6234 5)											
(34561 2)	(51346 2)											
(45612 3)												
(56123 4)												
$z^{(9)} =$ <table style="border-collapse: collapse; margin-left: 10px;"> <tr><td style="border: 1px solid black; padding: 2px;">(234561)</td><td style="border: 1px solid black; padding: 2px;">(5<u>1</u>2346)</td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(345612)</td><td style="border: 1px solid black; padding: 2px;">(12<u>3</u>465)</td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(456123)</td><td></td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(561234)</td><td></td></tr> <tr><td style="border: 1px solid black; padding: 2px;"> </td><td></td></tr> </table>	(23456 1)	(5 <u>1</u> 234 6)	(34561 2)	(12 <u>3</u> 46 5)	(45612 3)		(56123 4)				with	$x^T = x_1 x_2 x_3 x_4 x_6 x_5$ $\text{wt}(T) = q^2 t \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^5)}$ $\text{cov}_{>}^z(T) = 3 + 5 + \frac{1}{2}(1 - 12 - 3) = 1.$
(23456 1)	(5 <u>1</u> 234 6)											
(34561 2)	(12 <u>3</u> 46 5)											
(45612 3)												
(56123 4)												
$z^{(10)} =$ <table style="border-collapse: collapse; margin-left: 10px;"> <tr><td style="border: 1px solid black; padding: 2px;">(234561)</td><td style="border: 1px solid black; padding: 2px;">(5<u>1</u>2346)</td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(345612)</td><td style="border: 1px solid black; padding: 2px;">(<u>2</u>13465)</td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(456123)</td><td></td></tr> <tr><td style="border: 1px solid black; padding: 2px;">(561234)</td><td></td></tr> <tr><td style="border: 1px solid black; padding: 2px;"> </td><td></td></tr> </table>	(23456 1)	(5 <u>1</u> 234 6)	(34561 2)	(<u>2</u> 1346 5)	(45612 3)		(56123 4)				with	$x^T = x_1 x_2 x_3 x_4 x_6 x_5$ $\text{wt}(T) = q^2 t^2 \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^5)}$ $\text{cov}_{>}^z(T) = 3 + 5 + \frac{1}{2}(2 - 12 - 2) = 2.$
(23456 1)	(5 <u>1</u> 234 6)											
(34561 2)	(<u>2</u> 1346 5)											
(45612 3)												
(56123 4)												

$$z^{(11)} = \begin{array}{|c|c|} \hline (23456\mathbf{1}) & (\mathbf{5}1234\mathbf{6}) \\ \hline (34561\mathbf{2}) & (\mathbf{5}2346\mathbf{1}) \\ \hline (45612\mathbf{3}) & \\ \hline (56123\mathbf{4}) & \\ \hline \end{array} \quad \text{with} \quad \begin{aligned} x^T &= x_1x_2x_3x_4x_6x_1 \\ \text{wt}(T) &= q \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^4)} \\ \text{cov}_{>}^z(T) &= 3 + \frac{1}{2}(8 - 12 - 2) = 0. \end{aligned}$$

$$z^{(12)} = \begin{array}{|c|c|} \hline (23456\mathbf{1}) & (\mathbf{5}1234\mathbf{6}) \\ \hline (34561\mathbf{2}) & (\mathbf{5}1346\mathbf{2}) \\ \hline (45612\mathbf{3}) & \\ \hline (56123\mathbf{4}) & \\ \hline \end{array} \quad \text{with} \quad \begin{aligned} x^T &= x_1x_2x_3x_4x_6x_2 \\ \text{wt}(T) &= q \frac{(1-t)}{(1-qt^3)} \\ \text{cov}_{>}^z(T) &= 3 + \frac{1}{2}(7 - 12 - 1) = 0. \end{aligned}$$

$$z^{(13)} = \begin{array}{|c|c|} \hline (23456\mathbf{1}) & (\mathbf{5}6234\mathbf{1}) \\ \hline (34561\mathbf{2}) & (\mathbf{6}2341\mathbf{5}) \\ \hline (45612\mathbf{3}) & \\ \hline (56123\mathbf{4}) & \\ \hline \end{array} \quad \text{with} \quad \begin{aligned} x^T &= x_1x_2x_3x_4x_1x_5 \\ \text{wt}(T) &= qt \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^5)} \\ \text{cov}_{>}^z(T) &= 4 + \frac{1}{2}(8 - 12 - 2) = 1. \end{aligned}$$

$$z^{(14)} = \begin{array}{|c|c|} \hline (23456\mathbf{1}) & (\mathbf{5}6234\mathbf{1}) \\ \hline (34561\mathbf{2}) & (\mathbf{2}6341\mathbf{5}) \\ \hline (45612\mathbf{3}) & \\ \hline (56123\mathbf{4}) & \\ \hline \end{array} \quad \text{with} \quad \begin{aligned} x^T &= x_1x_2x_3x_4x_6x_5 \\ \text{wt}(T) &= qt^2 \frac{(1-t)}{(1-qt^5)} \\ \text{cov}_{>}^z(T) &= 5 + \frac{1}{2}(7 - 12 - 1) = 2. \end{aligned}$$

$$z^{(15)} = \begin{array}{|c|c|} \hline (23456\mathbf{1}) & (\mathbf{5}6234\mathbf{1}) \\ \hline (34561\mathbf{2}) & (\mathbf{5}2341\mathbf{6}) \\ \hline (45612\mathbf{3}) & \\ \hline (56123\mathbf{4}) & \\ \hline \end{array} \quad \text{with} \quad \begin{aligned} x^T &= x_1x_2x_3x_4x_6x_1 \\ \text{wt}(T) &= qt \frac{(1-t)}{(1-qt^4)} \\ \text{cov}_{>}^z(T) &= 4 + \frac{1}{2}(7 - 12 - 1) = 1. \end{aligned}$$

$$z^{(16)} = \begin{array}{|c|c|} \hline (23456\mathbf{1}) & (56234\mathbf{1}) \\ \hline (34561\mathbf{2}) & (56341\mathbf{2}) \\ \hline (45612\mathbf{3}) & \\ \hline (56123\mathbf{4}) & \\ \hline \end{array} \quad \text{with} \quad \begin{array}{l} x^T = x_1x_2x_3x_4x_1x_2 \\ \text{wt}(T) = 1 \\ \text{cov}_{>}^z(T) = 0 + \frac{1}{2}(12 - 12 - 0) = 0. \end{array}$$

In summary,

$$\begin{aligned} & (x_1x_2x_3x_4)^{-1}E_{(2,2,1,1,0,0)} \\ &= q^3 \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^5)} x_5x_6 + q^2t \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^5)} x_5x_6 \\ &+ q \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^4)} x_5x_1 + q \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^4)} x_5x_2 \\ &+ q^2t^5 \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^5)} x_5x_1 + qt \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^5)} x_5x_1 \\ &+ q^2t \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^4)} x_5x_6 + qt \frac{(1-t)}{(1-qt^4)} x_5x_2 \\ &+ q^2t \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^5)} x_6x_5 + q^2t^2 \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^5)} x_6x_5 \\ &+ q \frac{(1-t)}{(1-qt^3)} \frac{(1-t)}{(1-qt^4)} x_6x_1 + q \frac{(1-t)}{(1-qt^3)} x_6x_2 \\ &+ qt \frac{(1-t)}{(1-qt^4)} \frac{(1-t)}{(1-qt^5)} x_1x_5 + qt^2 \frac{(1-t)}{(1-qt^5)} x_6x_5 + qt \frac{(1-t)}{(1-qt^4)} x_6x_1 + x_1x_2. \end{aligned}$$

Acknowledgements. We thank A. Garbali and Tianyi Yu for very helpful proofreading and corrections.

References

- [Al16] P. Alexandersson, *Non-symmetric Macdonald polynomials and Demazure-Lusztig operators*, Sémin. Lothar. Combin. B76d (2019), MR4063906, arXiv:1602.05153. 2.5
- [Bu02] A.S. Buch, *A Littlewood-Richardson rule for the K-theory of Grassmannians*, Acta Math. **189** (2002) 37-78, MR1946917, arXiv:math/0004137. 1
- [CMW18] S. Corteel, O. Mandelshtam and L. Williams, *From multiline queues to Macdonald polynomials via the exclusion process*, Amer. J. Math. **144** (2022) 395-436, MR4401508, arXiv:1811.01024. 4
- [FL94] W. Fulton and A. Lascoux, *A Pieri formula in the Grothendieck ring of a flag bundle*, Duke Math. J. **76** (1994) 711-729, MR1309327. 2.3
- [GR21] W. Guo and A. Ram, *Comparing formulas for type GL_n Macdonald polynomials*, to appear in Algebraic Combinatorics, arXiv:2104.02942. 1, 2.5, 2.5, 3, 3.1, 3.1, 3.2, 3.4, 4

- [GR21b] W. Guo and A. Ram, *Comparing formulas for type GL_n Macdonald polynomials, Supplement*, to appear in Algebraic Combinatorics, arXiv:2104.04578. 1
- [Kt16] S. Kato, *Demazure character formula for semi-infinite flag varieties*, Math. Ann. **371** (2018) 1769-1801, MR3831285, arXiv:1605.04953. 1
- [KMY09] A. Knutson, E. Miller and A. Yong, *Gröbner geometry of vertex decompositions and of flagged tableaux*, J. Reine Angew. Math. **630** (2009) 1-31, MR2526784, arXiv:math/0502144. 1
- [Mac] I.G. Macdonald, *Symmetric functions and Hall polynomials*, Second edition, Oxford Mathematical Monographs, Oxford University Press, New York, 1995. ISBN: 0-19-853489-2, MR1354144.
- [Mac95] I.G. Macdonald, *Affine Hecke algebras and orthogonal polynomials*, Séminaire Bourbaki 1994/95, Astérisque **237** (1996) Exp. No. 797, 4, 189-207. MR1423624.
- [Mac03] I.G. Macdonald, *Affine Hecke Algebras and Orthogonal Polynomials*, Cambridge Tracts in Mathematics, vol. **157**, Cambridge University Press, Cambridge, 2003. MR1976581. 2.5
- [MS19] T. Matsumura and S. Sugimoto, *Factorial flagged Grothendieck polynomials*, in *Schubert calculus and its applications in combinatorics and representation theory*, Springer Proc. Math. Stat. **332**, Springer, Singapore (2020) 1-15, MR4167510, arXiv:1903.02169. 1
- [MRY19] F. McGlade, A. Ram and Y. Yang, *Positive level, negative level and level 0, Schubert calculus and its applications in combinatorics and representation theory*, Springer Proc. Math. Stat. **332**, Springer, Singapore (2020) 153-194, MR4167516, arXiv:1907.11796. 1
- [PS19] O. Pechenik and T. Scrimshaw, *K-theoretic crystals for set-valued tableaux of rectangular shapes*, Algebraic Combin. **5** (2022) 515-536, arXiv:1904.09674. 1
- [PR98] H. Pittie and A. Ram, *A Pieri-Chevalley formula in the K-theory of a G/B -bundle*, Electron. Res. Announc. Amer. Math. Soc. **5** (1999) 102-107, MR1701888, arXiv:math/0401331. 2.3
- [RY08] A. Ram and M. Yip, *A combinatorial formula for Macdonald polynomials*, Adv. Math. **226** (2011) 309-331, arXiv:0803.1146, MR2735761. 1, 2.5, 3.4
- [Wei20] A. Weigandt, *Bumpless pipe dreams and alternating sign matrices*, J. Combin. Theory Ser. A **182** (2021) Paper No. 105470, MR4258766, arXiv:2003.07342. 1, 2.2
- [Yu21] T. Yu, *Set-valued tableaux rule for Lascoux polynomials*, arXiv:2110.00164. 1, 2.3