# PAWŁUCKI-PLEŚNIAK EXTENSION OPERATOR FOR NON-MARKOV SETS

#### ALEXANDER GONCHAROV AND YAMAN PAKSOY

ABSTRACT. We show that Pawłucki-Pleśniak's operator is bounded for some non-Markov sets.

# 1. Introduction

By Whitney's extension theorem [18], for each compact set  $K \subset \mathbb{R}^d$ , there is a continuous linear operator extending jets of finite order from  $\mathscr{E}^p(K)$  to functions defined on the whole space, preserving the order of differentiability. However, for  $p = \infty$ , such an operator does not exist in the general case. Let us say that K has the *extension property* (EP) if there exists a linear continuous extension operator  $W : \mathscr{E}(K) \longrightarrow C^{\infty}(\mathbb{R}^d)$ . By Tidten ([17]), a set K has EP if and only if the space  $\mathscr{E}(K)$  possesses the dominating norm. By [8], there is no complete description of EP in terms of densities of measures, Hausdorff contents or related characteristics. For a short review of known extension operators we refer the reader to Section 2 in [8].

One of the approaches is due to W. Pawłucki and W. Pleśniak. In [13], they present an extension operator W in the form of a telescoping series containing Lagrange interpolating polynomials with Fekete nodes. The operator initially was considered for uniformly polynomially cuspidal compact sets. Later, in [14], the result was extended to any Markov set. By T.3.3 in [14], for each  $C^{\infty}$  determining compact set K, the suggested operator is continuous in the so-called Jackson topology  $\tau_J$  (see [14] and Section 2 in [1] for the definition and some properties of  $\tau_J$ ) if and only if  $\tau_J$  coincides with the natural topology  $\tau$  of the space  $\mathscr{E}(K)$  and this happens if and only if the set K is Markov. Since  $\tau_J$  is not stronger then  $\tau$ , each Markov set has EP. However, by [4] and [5], the inverse implication is not valid. Thus, in the case of non-Markov compact set with EP, the Pawłucki-Pleśniak extension operator is not continuous in  $\tau_J$ , yet this does not exclude the possibility for it to be bounded in  $\tau$ . Our aim is to check continuity of the operator in the natural topology of the space.

In the construction of the extension operator in [14], Pleśniak used four main components:

- 1) Markov property of the set *K*, which implies
- 2) the possibility of a suitable individual extension of polynomials from the set K to some neighborhood of the set, while preserving the norm of polynomials
- 3) a moderate growth of the Lebesgue constants corresponding to interpolation at Fekete points
- 4) the classical Jackson theorem on the polynomial approximation of  $C^{\infty}$  functions on the interval.

Date: Received: date / Accepted: date.

<sup>2010</sup> Mathematics Subject Classification. Primary 46E10; Secondary 41A05, 41A10.

Key words and phrases. Whitney functions, extension operators, Cantor-type sets, Markov's Property.

We analyze these components for non-Markov sets with EP. The paper is organized as follows. In Section 2, we repeat the argument from [14] for a Markov set K on the line such that K contains an array of interpolating nodes with the polynomial growth of the corresponding Lebesgue constants. Section 3 considers the family  $K^{\alpha}$ ,  $\alpha > 1$ , of non-Markov Cantor-type sets. They have EP provided  $\alpha < 2$ . Endpoints of intervals in Cantor procedure are enumerated in a special way. Some properties of the resulting sequence  $(x_k)_{k=1}^{\infty}$ are considered, which makes it possible to estimate the corresponding Lebesgue constants. In Section 4, we introduce the Markov  $M_N^{(p)}$  factor as the norm of the operator of *p*-fold differentiation in the space of polynomials of degree at most N. In contrast to the Markov case, for small sets, the value of  $M_N^{(p)}$  is substantially less than the p-th power of the usual Markov factor  $M_N$ . The fast growth of  $M_N^{(p)}$  can be neutralized (Section 5) by an ultra-fast rate of polynomial approximation of functions from  $\mathscr{E}(K^{\alpha})$ . In Section 6 we show that the boundedness of the operator depends not on the rate of growth of  $M_N^{(p)}$  factors but rather on the possibility of suitable individual extensions of polynomials. In the considered case, the polynomials involved in the interpolation process form a topological basis in the space  $\mathscr{E}(K)$ . This creates a bridge to Mityagin's method of extension [11]. Section 7 contains the main result: at least in the considered case, Pawłucki-Pleśniak's operator is continuous in the natural topology of the space. Moreover, Pleśniak's argument can be used for the given non-Markov set as well.

# 2. PAWŁUCKI-PLEŚNIAK'S OPERATOR FOR MARKOV SETS

For the convenience of the reader, we repeat the main points of [14]. We shall restrict the discussion to a perfect compact subset K of the line with a closed interval I containing K.

Let  $\mathscr{P}_n$  denote the set of all polynomials of degree at most n and  $|\cdot|_K$  (later  $|\cdot|_{0,K}$ ) stand for the uniform norm on K. We say that K is Markov (or K has Markov's Property P) if there are constants  $C_1$  and P with

$$(2.1) |Q^{(p)}|_K \le C_1 n^{rp} |Q|_K for all Q \in \mathscr{P}_n, p \in \mathbb{N}.$$

Given  $\delta > 0$ , let  $K_{\delta}$  denote  $\{x \in \mathbb{R} : \operatorname{dist}(x,K) \leq \delta\}$  and  $u_{\delta}$  be a  $C^{\infty}$  function with the properties:  $0 \leq u_{\delta} \leq 1, u_{\delta} = 1$  on  $K, u_{\delta} = 0$  on  $\mathbb{R} \setminus K_{\delta}$ , and  $|u_{\delta}^{(j)}(x)| \leq c_{j}\delta^{-j}$  for each  $j \in \mathbb{N}, x \in \mathbb{R}$ . The constants  $c_{j}$  do not depend on K. Although the existence of such a function is well known, a specific version of  $u_{\delta}$  is considered in Section 6. Suppose K satisfies (2.1). For each polynomial Q of degree n let us take  $\delta = n^{-r}$ . By means of Taylor's expansion, it's easy to show that

$$(2.2) |Q(x)| \le C_1 e |Q|_K for Q \in \mathscr{P}_n, x \in K_{n^{-r}}.$$

In addition, if  $p \le n$  then for each  $x \in \mathbb{R}$  we have

$$(2.3) |(Q \cdot u_{\delta})^{(p)}(x)| \le C_2 n^{rp} |Q|_K,$$

where  $C_2$  depends only on p. Indeed, it is evident for  $x \in K$  or  $x \notin K_{\delta}$ . Otherwise, by Leibnitz's rule,

$$|(Q \cdot u_{\delta})^{(p)}(x)| \le \sum_{j=0}^{p} {p \choose j} c_{j} n^{rj} C_{1} e |Q^{(p-j)}|_{K} \le C_{2} n^{rp} |Q|_{K}$$

with 
$$C_2 := C_1^2 e \sum_{j=0}^p {p \choose j} c_j$$
.

Let  $\mathscr{X}=(x_{k,n})_{k=1,n=1}^{n,\infty}$  be an infinite triangular matrix of points from K such that each row  $\mathscr{X}_n$  consists of distinct elements. For a fixed n, the points of  $\mathscr{X}_n$  determine the polynomial  $\omega_n(x)=\prod_{k=1}^n(x-x_{k,n})$ , the fundamental Lagrange polynomials  $l_{k,n}(x)=\frac{\omega_n(x)}{(x-x_{k,n})\omega_n'(x_{k,n})}$  with  $1\leq k\leq n$ , the Lebesgue function  $\lambda_n(x)=\sum_{k=1}^n|l_{k,n}(x)|$ , and the Lebesgue constant  $\Lambda_n(K)=\sup_{x\in K}\lambda_n(x)$ . Given function f defined on f, by f by f we denote the corresponding Lagrange interpolating polynomial, by f by f the interpolating projection, so f is the sup-norm of f defined on f by f by the interpolating projection, so f is the sup-norm of f defined on f by f by the interpolating projection, so f is chosen in a such way that the sequence f in f has at most polynomial growth: there are constants f and f such that

(2.4) 
$$\Lambda_{n+1}(K) \le C_3 n^R \text{ for all } n \in \mathbb{N}.$$

Of course, Fekete points provide this condition with R = 1, see for instance Section 2 in [14]. Our main object is the operator  $W : \mathcal{E}(K) \longrightarrow C^{\infty}(\mathbb{R})$  which is defined as follows

$$(2.5) W(f,x) = L_1(f,x;\mathscr{X}_1) \cdot u_{\delta_1}(x) + \sum_{n=1}^{\infty} [L_{n+1}(f,x;\mathscr{X}_{n+1}) - L_n(f,x;\mathscr{X}_n)] \cdot u_{\delta_n}(x).$$

Recall that, by [18], the Whitney space  $\mathcal{E}(K)$  consists of traces on K of functions from  $C^{\infty}(I)$  and the Whitney topology  $\tau$  is given by the seminorms

$$||f||_q = |f|_{q,K} + \sup \left\{ |(R_y^q f)^{(k)}(x)| \cdot |x - y|^{k-q} : x, y \in K, x \neq y, k = 0, 1, ...q \right\}$$

for  $q \in \mathbb{Z}_+$ , where  $|f|_{q,K} = \sup\{|f^{(k)}(x)| : x \in K, k \le q\}$  and  $R^q_y f(x) = f(x) - T^q_y f(x)$  is the Taylor remainder. By the open mapping theorem, for any q there exists C such that

(2.6) 
$$\inf |F|_{q,I} \le C ||f||_q,$$

where the infimum is taken over all extensions of f to  $F \in C^{\infty}(I)$ , see, e.g., (2.3) in [10].

Following Zerner [19], let us consider the given below seminorms in  $\mathscr{E}(K)$ 

$$d_{-1}(f) = |f|_K$$
,  $d_0(f) = E_0(f, K)$ ,  $d_k(f) = \sup_{n \ge 1} n^k E_n(f, K)$  for  $k \ge 1$ ,

where  $E_n(f,K) := \min_{P \in \mathscr{P}_n} |f-P|_K$  is the best approximation of f by n-th degree polynomials. Since K is perfect, the *Jackson topology*  $\tau_J$ , given by  $(d_k)$ , is Hausdorff. By Jackson's theorem,  $\tau_J$  is well-defined and is not stronger than  $\tau$  (see [1] for more details). By Pleśniak (T.3.3 in [14]), the operator W is continuous in  $\tau_J \Leftrightarrow \tau_J = \tau \Leftrightarrow (\mathscr{E}(K), \tau_J)$  is complete  $\Leftrightarrow K$  is Markov. We now extract a part of this theorem in the following form.

**Theorem 2.1.** ([14]) Let  $K \subset \mathbb{R}$  be Markov and the operator W be given by an array  $\mathscr{X}$  satisfying (2.4). Then W is bounded in  $\tau$ .

*Proof.* Let  $G_n$  denote the n-th term of the series (2.5) with  $\delta_n = n^{-r}$ , where r is taken from (2.1). Fix  $p \in \mathbb{N}$  and  $x \in \mathbb{R}$ . The expression in square brackets is a polynomial of degree n, so, by (2.3),  $|G_n^{(p)}(x)| \leq C_2 n^{rp} |L_{n+1}(f) - L_n(f)|_K$ . Lebesgue's Lemma (see, e.g., [3], Chapter 2, Prop. 4.1) now yields  $|L_{n+1}(f) - L_n(f)|_K \leq |L_{n+1}(f) - f|_K + |L_n(f) - f|_K \leq (\Lambda_{n+1}(K) + 1)E_n(f,K) + (\Lambda_n(K) + 1)E_{n-1}(f,K)$ . By (2.4),  $|G_n^{(p)}(x)| \leq C_4 n^{rp+R} E_{n-1}(f,K)$ , where  $C_4$  does not depend on n and f. Clearly,  $E_{n-1}(f,K) \leq E_{n-1}(F,I)$ , where F is any extensions of f to  $F \in C^{\infty}(I)$ . Jackson's Theorem (see, e.g., [3], Chapter 7, Cor. 6.5) and

(2.6) show that  $|G_n^{(p)}(x)|$  is a term of a uniformly convergent series, which completes the proof.

We aim to show that a modified version of this theorem can also be applied to some non-Markov sets.

#### 3. Non-Markov sets with the extension property

We consider the family  $K^{\alpha}$  of sets proposed in [5]. A geometrically symmetric Cantor set K is the intersection  $\bigcap_{s=0}^{\infty} E_s$ , where  $E_0 = [0,1]$  and for each  $s \in \mathbb{N}$ , the set  $E_s$  is a union of  $2^s$  closed *basic* intervals  $I_{j,s}$ ,  $j=1,2,...,2^s$  of length  $\ell_s$ . Recursively,  $E_{s+1}$  is obtained by replacing each interval  $I_{j,s}$  by two *adjacent* subintervals  $I_{2j-1,s+1}$  and  $I_{2j,s+1}$ , where the distance between them is  $h_s = \ell_s - 2\ell_{s+1}$ .

distance between them is  $h_s = \ell_s - 2\ell_{s+1}$ . For  $\alpha > 1$  and  $\ell_1 < 1/2$  with  $2\ell_1^{\alpha - 1} < 1$ , the set  $K^{\alpha}$  is the Cantor set associated to the lengths of intervals satisfying  $\ell_s = \ell_{s-1}^{\alpha} = \ell_1^{\alpha^{s-1}}$  for  $s \ge 2$ . By [15],  $K^{\alpha}$  are not Markov. On the other hand, by [5] and [6], the set  $K^{\alpha}$  has the extension property if and only if  $\alpha \le 2$ .

We follow the notation of [9]. Let  $X_0 := \{0,1\}$  and, for  $k \in \mathbb{N}$ , let  $X_k$  be the set of endpoints of intervals from  $E_k$  that are not endpoints of intervals from  $E_{k-1}$ . Thus,  $X_1 := \{\ell_1, 1 - \ell_1\}, X_2 := \{\ell_2, \ell_1 - \ell_2, 1 - \ell_1 + \ell_2, 1 - \ell_2\}$ , etc. We refer s-th type points to the elements of  $X_s$ . Set  $Y_s = \bigcup_{k=0}^s X_k$ . Clearly,  $\#(X_s) = 2^s$  for  $s \in \mathbb{N}$  and  $\#(Y_s) = 2^{s+1}$  for  $s \in \mathbb{Z}_+$ . Here and below, #(Z) denotes the cardinality of a finite set Z.

Let  $Z = (z_k)_{k=1}^M \subset \mathbb{R}$  and  $\omega_M(x) = \prod_1^M (x - z_k)$ . For a fixed  $x \in \mathbb{R}$ , by  $d_k(x, Z)$  we denote the distances  $|x - z_{j_k}|$  from x to points of Z, where these distances are arranged in the nondecreasing order, so  $d_k(x, Z) \leq d_{k+1}(x, Z)$  for  $k = 1, 2, \cdots, M-1$ . Then  $|\omega_M(x)| = \prod_{k=1}^M d_k(x, Z)$  and, given p < M, the p-th derivative of  $\omega_M$  at the point x is the sum of  $\frac{M!}{(M-p)!}$  products, where each product contains M - p terms of the type  $(x - z_k)$ . Hence

(3.1) 
$$|\omega_{M}^{(p)}(x)| \le M^{p} \prod_{k=p+1}^{M} d_{k}(x, Z).$$

Suppose we are given a finite set  $Z = (z_k)_{k=1}^M \subset K^{\alpha}$ . Let  $m_{j,s}(Z) := \#(Z \cap I_{j,s})$ . We say that points of Z are *uniformly distributed* on  $K^{\alpha}$  if for each  $k \in \mathbb{N}$  and  $i, j \in \{1, 2, ..., 2^k\}$  we have

$$|m_{i,k}(Z) - m_{i,k}(Z)| \le 1.$$

As in [8], see also [9], we put all points from  $\bigcup_{k=0}^{\infty} X_k$  in order by the rule of increase of type. First, we enumerate points from  $Y_0 = X_0 : x_1 = 0, x_2 = 1$ . After this we include points from  $X_1$  by increase the index of each point by 2:  $x_3 = \ell_1, x_4 = 1 - \ell_1$ . Thus,  $Y_1$  in ascending order is  $\{x_1, x_3, x_4, x_2\}$ . Increasing the index of each point by 4 gives the order  $X_2 = \{x_5, x_7, x_8, x_6\}$  with  $Y_2 = \{x_1, x_5, x_7, x_3, x_4, x_8, x_6, x_2\}$ . Continuing in this fashion, we use  $Y_{k-1} = \{x_{i_1}, x_{i_2}, \cdots, x_{i_{2k}}\}$  to define the ordering  $X_k = \{x_{i_1+2^k}, x_{i_2+2^k}, \cdots, x_{i_{2k}+2^k}\}$ . We see that the points with odd indices are on the left part of  $K^{\alpha}$ , whereas  $x_{2n} \in I_{2,1}$ . For each M, the first M points chosen by the above rule are uniformly distributed on  $K^{\alpha}$ .

Let us consider the location of these points in more detail. Suppose  $k \in \mathbb{N}$  and  $1 \le j \le 2^k$ . Then  $(x_i)_{i=1}^{2^k}$  is  $Y_{k-1}$  that is the set of all endpoints of types  $\le k-1$ , whereas  $x_{2^k+j}=$ 

 $x_j \pm \ell_k$ , where the sign is determined as follows: if  $2^k + j = 2^k + 2^m + \cdots + 2^r + 1$  with  $\kappa := \#\{k, m, \dots, r\}$  then the sign is  $(-1)^{\kappa - 1}$ .

In what follows, we will interpolate polynomials at points of  $Z = (x_k)_{k=1}^{N+1}$ : if  $Q \in \mathcal{P}_N$  then

(3.3) 
$$Q(x) = \sum_{k=1}^{N+1} Q(x_k) \frac{a_k(x)}{a_k(x_k)},$$

where  $a_k(x) = \prod_{i=1, i \neq k}^{N+1} (x - x_i)$ . Let us fix N with the binary decomposition for N+1

$$(3.4) N+1 = 2^s + 2^{s_1} + 2^{s_2} + \dots + 2^{s_m} \text{ with } 0 \le s_m < \dots < s_1 < s_0 := s.$$

By the above,

$$(3.5) x_{N+2} = \ell_{s_m} - \ell_{s_{m-1}} + \ell_{s_{m-2}} - \dots + (-1)^m \ell_{s_0}.$$

The representation (3.4) gives the decomposition

$$(3.6) Z = A_s \cup A_{s_1} \cup \cdots \cup A_{s_m}$$

with  $\#(A_{s_j}) = 2^{s_j}$  for  $0 \le j \le m$ . Here,  $A_s = (x_k)_{k=1}^{2^s} = (x_{k,s})_{k=1}^{2^s}$ , with  $x_{1,s} = 0 < x_{2,s} = \ell_{s-1} < \cdots < x_{2^s,s} = 1$ . Similarly,  $A_{s_1} = (x_k)_{k=2^s+1}^{2^s+2^{s_1}} = (x_{k,s_1})_{k=1}^{2^{s_1}}, \cdots, A_{s_m} = (x_k)_{k=N-2^{s_m}+2}^{N+1} = (x_{k,s_m})_{k=1}^{2^{s_m}}$  with  $x_{k,s_j} \nearrow$  as k increases and j is fixed.

Our next objective is to determine the location of points from  $A_{s_j}$ . Let  $I_{i,k} = [a_{i,k}, b_{i,k}]$  for  $1 \le i \le 2^k$ . Then  $A_s = Y_{s-1} = \bigcup_{i=1}^{2^{s-1}} \{a_{i,s-1}, b_{i,s-1}\}$ . Each  $I_{i,s_1-1}$  contains two points of  $A_{s_1}$  that are symmetric with respect to the endpoints of

Each  $I_{i,s_1-1}$  contains two points of  $A_{s_1}$  that are symmetric with respect to the endpoints of  $I_{i,s_1-1}$ . For example,  $x_{1,s_1} = x_{2^s+1} = \ell_s$  is symmetric to  $x_{2,s_1} = \ell_{s_1-1} - \ell_s$ , which is  $x_{2^s+2^{s_1-1}+1}$ . The largest point of  $A_{s_1}$ , namely  $x_{2^{s_1},s_1} = 1 - \ell_s$  is  $x_k$  with  $k = 2^s + 2 = 2^s + 2^0 + 1$ . Thus,  $A_{s_1} = \bigcup_{i=1}^{2^{s_1}-1} \{a_{i,s_1-1} + \ell_s, b_{i,s_1-1} - \ell_s\}$ . Likewise,  $x_{1,s_2} = x_{2^s+2^{s_1}+1} = \ell_{s_1} - \ell_s$  is symmetric in regard to  $I_{1,s_2-1}$  with  $x_{2,s_2} = \ell_{s_2-1} - \ell_{s_1} + \ell_s = x_{2^s+2^{s_1}+2^{s_2-1}+1}$ , so  $A_{s_2} = \bigcup_{i=1}^{2^{s_2-1}} \{a_{i,s_2-1} + \ell_{s_1} - \ell_s, b_{i,s_2-1} - \ell_{s_1} + \ell_s\}$ . Continuing in this fashion, we get  $A_{s_m} = \bigcup_{i=1}^{2^{s_m}-1} \{a_{i,s_m-1} + \ell_{s_{m-1}} - \ell_{s_{m-2}} + \cdots - (-1)^m \ell_s, b_{i,s_m-1} - \ell_{s_{m-1}} + \ell_{s_{m-2}} + \cdots + (-1)^m \ell_s\}$ .

Each  $x \in K^{\alpha}$  determines the increasing chain of basic intervals:

$$(3.7) x \in I_{j,s} \subset I_{j_1,s-1} \subset I_{j_2,s-2} \subset \cdots \subset I_{j_s,0} = [0,1].$$

Let  $J_s$  and  $I_{j,s}$  be the adjacent subintervals of  $I_{j_1,s-1}$  and, more generally, for  $1 \le n \le s-1$ , let  $J_n := (I_{j_{s-n+1},n-1} \setminus I_{j_{s-n},n}) \cap E_n$ . Obviously,  $\#(A_s \cap I_{j,s}) = \#(A_s \cap J_s) = 1, \dots, \#(A_s \cap J_n) = 2^{s-n}$ , so  $\prod_{x_k \in A_s} |x - x_k| \le \pi_0 := \ell_s \ell_{s-1} \ell_{s-2}^2 \cdots \ell_0^{2^{s-1}}$ .

The same reasoning applies to  $1 \le j \le m$ . Let  $\pi_j := \ell_{s_j} \ell_{s_j-1} \ell_{s_j-2}^2 \cdots \ell_0^{2^{s_j-1}}$ . Then  $|\omega_{N+1}(x)| = \prod_{0 \le j \le m} \prod_{x_k \in A_{s_j}} |x - x_k| \le \prod_{j=0}^m \pi_j$ .

We shall use two more representations of the product above

(3.8) 
$$|\omega_{N+1}(x)| \le \prod_{i=0}^{m} \pi_i = \prod_{i=0}^{s} \ell_i^{\lambda_i} = \prod_{k=1}^{N+1} \rho_k,$$

where  $(\rho_k)_{k=1}^{N+1}$  are all terms  $\ell_i$  of the product arranged in nondecreasing order. We see that the real multiplication in  $\prod_{i=0}^{s} \ell_i^{\lambda_i}$  starts at i=1 because  $\ell_0=1$ , so at least half of the terms  $\rho_k$  are 1.

The inequality (3.8) is exact in powers in the following sense

$$|\omega_{N+1}(x_{N+2})| \geq \prod_{i=0}^{s} h_i^{\lambda_i}.$$

Indeed,  $\prod_{x_k \in A_s} |x_{N+2} - x_k| \ge h_s h_{s-1} h_{s-2}^2 \cdots h_0^{2^{s-1}}$ . Taking into account (3.5) and the location of points in  $A_{s_j}$ , we have a similar estimate of  $\prod_{x_k \in A_{s_j}} |x_{N+2} - x_k|$  for  $1 \le j \le m$ . By Lemma 2.1 in [9], the ratio  $h_n/\ell_n$  increases with n. Hence,  $h_n \ge \ell_n \cdot h_0$ . Therefore,

(3.9) 
$$h_0^{N+1} \prod_{i=0}^s \ell_i^{\lambda_i} \le |\omega_{N+1}(x_{N+2})| \le \prod_{i=0}^s \ell_i^{\lambda_i}.$$

As above, let  $Z = (x_k)_{k=1}^{N+1}$ . Fix k with  $1 \le k \le N+1$  and  $I_{j,s}$  containing  $x_k$ . Our next goal is to analyze the function  $a_k$ . The chain (3.7) for  $x_k$  determines intervals  $(J_n)_{n=1}^s$  and degrees  $\mu_n := \#(J_{n+1} \cap Z \setminus \{x_k\})$  for  $0 \le n \le s-1$ ,  $\mu_s := \#(I_{j,s} \cap Z \setminus \{x_k\})$ . Clearly,  $\mu_s = m_{j,s} - 1$  and  $\sum_{n=0}^s \mu_n = N$ . Let us consider the interval  $I_{j,s}$ . If  $m_{j,s} = 1$ , then  $\mu_s = 0$ . Otherwise,  $m_{j,s} = 2$ . Here, by the choice procedure,  $d_1(x_k, Z \setminus \{x_k\}) = \ell_s$ . In both cases this gives the factor  $\ell_s^{\mu_s}$  in  $|a_k(x_k)|$  corresponding to  $I_{j,s}$ . If  $x_i \in J_{n+1}$  then  $h_n \le |x_k - x_i| \le \ell_n$ . It follows that

(3.10) 
$$h_0^N \prod_{i=0}^s \ell_i^{\mu_i} \le |a_k(x_k)| \le \prod_{i=0}^s \ell_i^{\mu_i}.$$

Suppose  $x \in I_{j,s}$ , so  $x_k$  and x have the same degrees  $(\mu_i)_{i=0}^s$ . As above,  $|a_k(x)| \le \prod_{i=0}^s \ell_i^{\mu_i}$ . Thus, if x and  $x_k$  are on the same interval of the s-th level, then  $|a_k(x)| \le h_0^{-N} |a_k(x_k)|$ . Let us show that a similar upper bound holds for each  $x \in K^{\alpha}$ .

**Lemma 3.1.** Given  $N \in \mathbb{N}$ , let  $Z = (x_k)_{k=1}^{N+1}$  be chosen in  $K^{\alpha}$  as above,  $1 \le k \le N+1$ . Then  $\max_{x \in K^{\alpha}} |a_k(x)| \le h_0^{-N} |a_k(x_k)|$ .

*Proof.* Fix k and  $\tilde{x}$  such that  $\max_{x \in K^{\alpha}} |a_k(x)| = |a_k(\tilde{x})|$ . For  $x_k$  we take the chain (3.7) and degrees  $(\mu_i)_{i=0}^s$  as above. On the other hand, the point  $\tilde{x}$  determines the chain  $\tilde{x} \in \tilde{I}_{j,s} \subset \tilde{I}_{j_1,s-1} \subset \cdots \subset \tilde{I}_{j_s,0} = [0,1]$  with the corresponding  $(\tilde{J}_n)_{n=1}^s$  and  $v_n := \#(\tilde{J}_{n+1} \cap Z \setminus \{x_k\})$  for  $0 \le n \le s-1, v_s(x) := \#(\tilde{I}_{j,s} \cap Z \setminus \{x_k\})$ . We aim to show that for  $1 \le i \le s$ 

$$(3.11) \mu_s + \mu_{s-1} + \dots + \mu_i \le \nu_s + \nu_{s-1} + \dots + \nu_i.$$

Let n be the largest level for each  $I_{j_{s-n},n} = \tilde{I}_{j_{s-n},n}$ . Then the intervals  $J_i$  and  $\tilde{J}_i$  coincide for  $0 \le i \le n$  and  $\mu_i = \nu_i$  for  $0 \le i \le n-1$ . The interval  $I_{j_{s-n},n}$  contains  $N - \sum_{q=0}^{n-1} \mu_q$  which is  $\sum_{q=n}^s \mu_q$ . The same is valid for  $\nu_q$ . Hence, (3.11) is valid for i=n and we need only prove it for  $i \ge n+1$ . We note that  $x_k$  and  $\tilde{x}$  are on different subintervals of the n+1-st level of  $I_{j_{s-n},n}$ , namely,  $x_k \in \tilde{J}_{n+1}, \tilde{x} \in J_{n+1}$ .

Let us show (3.11) for i = s. Since the set Z is distributed uniformly, each interval of s-th level may contain not less than one and not more that two points of Z. Thus,  $1 \le v_s$  and  $\mu_s \le 1$ , as the point  $x_k$  is excluded. Similarly, by (3.2),

$$\mu_s + \mu_{s-1} = m_{j_1,s-1} - 1 \le \#(\tilde{I}_{j_1,s-1} \cap Z) = \#(\tilde{I}_{j_1,s-1} \cap Z \setminus \{x_k\}) = v_s + v_{s-1},$$

since  $x_k \notin \tilde{I}_{j_1,s-1}$ . We can repeat the argument for other  $i \ge n+1$ , since for such values i the interval  $\tilde{I}_{j_s-i,i}$  does not contain  $x_k$ . Thus, (3.11) is valid for all i.

We proceed to show that the desired assertion follows from (3.11). On the one hand,  $\max_{x \in K^{\alpha}} |a_k(x)| \le \prod_{j=0}^s \ell_j^{\nu_j} = \ell_1^{\nu_1 + \alpha \nu_2 + \dots + \alpha^{s-1} \nu_s}$ . On the other hand,  $|a_k(x_k)| \ge \prod_{j=0}^s h_j^{\mu_j} = \prod_{j=0}^s \ell_j^{\mu_j} \cdot \prod_{j=0}^s (h_j/\ell_j)^{\mu_j}$ , where, as above,  $h_j/\ell_j \ge h_0$ . Hence,  $|a_k(x_k)| \ge h_0^N \prod_{j=0}^s \ell_j^{\mu_j}$  and it remains to prove that

(3.12) 
$$\prod_{j=0}^{s} \ell_{j}^{v_{j}} \leq \prod_{j=0}^{s} \ell_{j}^{\mu_{j}},$$

or, what is equivalent, that

$$\mu_1 + \alpha \mu_2 + \dots + \alpha^{s-1} \mu_s \le v_1 + \alpha v_2 + \dots + \alpha^{s-1} v_s$$

The left side can be written as  $(\mu_s + \mu_{s-1} + \cdots + \mu_1) + (\alpha - 1)(\mu_s + \mu_{s-1} + \cdots + \mu_2) + \cdots + (\alpha^{s-1} - \alpha^{s-2})\mu_s$ . Similar representation for the right side and (3.11) completes the proof.

One may conjecture that the coefficient  $h_0^{-N}$  in the above lemma can be reduced. However, it cannot be replaced by a factor that increases polynomially with N. Let's show this.

**Example 3.2.** Let  $N = 2^s + 2$ ,  $\alpha = 2$ ,  $\ell_1 \le 1/4$  and a constant r be fixed. Then for  $Z = (x_k)_{k=1}^{N+1}$ , k = N+1,  $y = \ell_2 - \ell_s$  we have  $N^r |a_k(x_k)| < |a_k(y)|$  for large s.

Here,  $x_k = \ell_1 - \ell_s$ ,  $a_k(x) = \prod_{j=1}^N (x - x_j)$ . We proceed to show that  $\frac{|a_k(x_k)|}{|a_k(y)|}$  is exponentially small (with respect to N) for large enough s. If the set  $(x_j)_{j=1}^{2^s+2}$  is decomposed in the form  $Y_{s-1} \cup A$  with  $A = (x_j)_{j=2^s+1}^{2^s+2} = \{\ell_s, 1 - \ell_s\}$ , then  $|a_k(x)| = \prod_{x_j \in Y_{s-1}} |x - x_j| \prod_{x_j \in A} |x - x_j|$ . For the second part, we have  $\prod_{x_j \in A} \frac{|x_k - x_j|}{|y - x_j|} = \frac{(\ell_1 - 2\ell_s)(1 - \ell_1)}{(\ell_2 - 2\ell_s)(1 - \ell_2)} < M := \frac{2}{\ell_1(1 - \ell_2)}$  that does not depend on s. It remains to estimate  $\prod_{x_j \in Y_{s-1}} \frac{|x_k - x_j|}{|y - x_j|} = \prod_{j=1}^{2^s} \frac{d_j(x_k)}{d_j(y)}$ , where for brevity, we drop the argument  $Y_{s-1}$  in  $d_j(x, Y_{s-1})$ . By symmetry,  $d_j(x_k) = d_j(y)$  for  $1 \le j \le 2^{s-2}$ , that is for the points  $x_i$  on the nearest (to the argument of  $d_j$ ) interval of the second level. Let us take  $\tilde{y} := \ell_1 - \ell_2 + \ell_s$ , which is symmetric to y with respect to  $I_{1,1}$ . Then  $d_j(\tilde{y}) = d_j(y)$  for  $2^{s-2} + 1 \le j \le 2^{s-1}$ . Since  $x_k - \tilde{y} = \ell_2 - 2\ell_s$ , we get  $\prod_{j=2^{s-1}-2+1}^{2^{s-1}} \frac{d_j(x_k)}{d_j(y)} = \prod_{j=2^{s-2}-2+1}^{2^{s-1}} (1 + \frac{\ell_2 - 2\ell_s}{D_i})$ . Here and for the remaining two intervals of the 2-nd level we express the corresponding products in terms of  $D_i := d_{2^{s-2}+i}$  with  $1 \le i \le 2^{s-2}$ . For these intervals  $d_j(x_k) = d_j(y) - \ell_1 + \ell_2$ .

If 
$$x_i \in I_{3,2}$$
 then  $d_{2^{s-1}+i} = D_i + 1 - 2\ell_1 + \ell_s$ , so  $\prod_{j=2^{s-1}+1}^{3 \cdot 2^{s-2}} \frac{d_j(x_k)}{d_j(y)} = \prod_{i=1}^{2^{s-2}} \left(1 - \frac{\ell_1 - \ell_2}{D_i + 1 - 2\ell_1 + \ell_2}\right)$ .  
If  $x_i \in I_{4,2}$  then  $d_{3 \cdot 2^{s-2} + i} = D_i + 1 - \ell_1$  and  $\prod_{j=3 \cdot 2^{s-2} + 1}^{2^s} \frac{d_j(x_k)}{d_j(y)} = \prod_{i=1}^{2^{s-2}} \left(1 - \frac{\ell_1 - \ell_2}{D_i + 1 - \ell_1}\right)$ .

Therefore,  $\prod_{j=1}^{2^s} \frac{d_j(x_k)}{d_j(y)} = \prod_{i=1}^{2^{s-2}} (1 + \frac{\ell_2 - 2\ell_s}{D_i}) (1 - \frac{\ell_1 - \ell_2}{D_{i+1} - 2\ell_1 + \ell_2}) (1 - \frac{\ell_1 - \ell_2}{D_{i+1} - \ell_1})$  with admissible values  $\ell_1 - 2\ell_2 + \ell_s \leq D_i \leq \ell_1 - \ell_2 + \ell_s$ . The general term of the product (we denote it briefly by  $b_i$ ) consists of three parts. It can be increased only if  $D_i$  is replaced by  $\ell_1 - 2\ell_2$  in the first part and by the maximum  $D_i$  in the 2nd and 3rd parts. Hence,  $b_i \leq \frac{\ell_1 - \ell_2}{\ell_1 - 2\ell_2} \frac{1 - 2\ell_1 + \ell_2 + \ell_s}{1 - \ell_1 + \ell_s} \frac{1 - \ell_2 + \ell_s}{1 + \ell_1 - 2\ell_2 + \ell_s}$ . For large s, the right side of the expression is as close to  $\sigma := \frac{1 - \ell_1}{1 - 2\ell_1} \frac{1 - 2\ell_1 + \ell_2}{1 - \ell_1} \frac{1 - \ell_2}{1 + \ell_1 - 2\ell_2}$  as we want it to be. A straightforward computation shows that  $\sigma < 1$ . Let  $\sigma < \sigma_1 < 1$ . Then  $\frac{|a_k(x_k)|}{|a_k(y)|} < M \sigma_1^{2^{s-2}}$ , which is the desired conclusion.

**Remark 3.3.** Here,  $\ell_1$  may be arbitrary small.

**Remark 3.4.** The example shows that the sequence  $(x_n)_{n=1}^{\infty}$  is not Leja. Recall that a sequence  $(y_n)_{n=1}^{\infty} \subset K$  has Leja's property if  $|y_1 - y_2| = \operatorname{diam}(K)$  and, once  $y_1, y_2, \dots, y_n$  have been determined,  $y_{n+1}$  is chosen so that it provides the maximum modulus of the polynomial  $(x - y_1) \cdots (x - y_n)$  on K. In our case,  $|\omega_{2^s + 2}(x_{2^s + 3})| < |\omega_{2^s + 2}(y)|$ . However, using the example technique, it can be shown that  $(x_n)_{n=1}^{\infty}$  is a Leja sequence for  $K^{\alpha}$  if  $\alpha > 2$ .

**Remark 3.5.** Similarly, the set  $(x_n)_{n=1}^M$  is not a Fekete M-tuple for  $M=2^s+3$ . Indeed, let  $V(t_1, \dots, t_M)$  be the Vandermonde determinant. In our case,  $\frac{|V(x_1, \dots, x_M)|}{|V(x_1, \dots, x_{M-1}, y)|} = \frac{|\omega_{2^s+2}(x_M)|}{|\omega_{2^s+2}(y)|} < 1$ , whereas a Fekete M-tuple must realize the maximum modulus of the Vandermonde determinant. minant.

4. Markov 
$$M_N^{(p)}$$
 factors

As a first application of Lemma 3.1, we can estimate the Lebesgue constants for Newton's interpolation at points  $(x_n)_{n=1}^{\infty}$ .

**Proposition 4.1.** Given  $N \in \mathbb{N}$ , let  $(x_k)_{k=1}^{N+1}$  be chosen by the rule of increase of type and  $\Lambda_{N+1}(K^{\alpha})$  be the corresponding Lebesgue constant. Then

$$\Lambda_{N+1}(K^{\alpha}) \leq h_0^{-N} \cdot (N+1).$$

*Proof.* It is evident in view of (3.3) and Lemma 3.1.

From now on, for any product  $\prod_{k=1}^{M} t_k$  with  $t_k \ge 0$  and p < M, we will use the symbol  $p(\prod_{k=1}^M t_k)$  to denote the product of p smallest terms  $t_k$ . Also, let  $(\prod_{k=1}^M t_k)_p$  be the original product without p smallest terms, so  $\prod_{k=1}^M t_k =_p (\prod_{k=1}^M t_k) \cdot (\prod_{k=1}^M t_k)_p$ . Of course, in general,  $\prod_{k=1}^M \tau_k \leq \prod_{k=1}^M t_k$  does not imply

$$\left(\prod_{k=1}^{M} \tau_{k}\right)_{p} \leq \left(\prod_{k=1}^{M} t_{k}\right)_{p}.$$

But (4.1) is trivially valid provided additional condition:  $0 \le \tau_k \le t_k$  for all k.

Let us show that in (3.11) and (3.12)  $v_i$  can be replaced by  $\lambda_i$ . The argument of Lemma 3.1 can be applied to each point  $x \in K^{\alpha}$  instead of  $\tilde{x}$ . Let us apply it to  $x_{N+2}$ , for which we have some degrees  $\overline{v}_j$  and (3.11) with  $\overline{v}_j$  instead of  $v_j$ . If we add the point  $x_k$  to  $Z \setminus \{x_k\}$  then one of  $\overline{V}_i$  will increase by one and new powers become  $\lambda_i$ . Thus,

for  $1 \le i \le s$ . By this, as above, we have  $\prod_{j=0}^{s} \ell_j^{\lambda_j} \le \prod_{j=0}^{s} \ell_j^{\mu_j}$ . Note that the values of  $\mu_j, \overline{\nu}_j$ depend on k, while  $\lambda_i$  is only defined by N. The left product has N+1 terms, and the right product has N. Of course, we can start multiplication starting from j = 1. By (4.2), it can be started from each i. In fact, we can make a more general estimate. Let  $p \le N$ . Then

$$(4.3) p\left(\prod_{j=0}^{s} \ell_{j}^{\lambda_{j}}\right) \leq p\left(\prod_{j=0}^{s} \ell_{j}^{\mu_{j}}\right).$$

The proof is by induction on p. Let p=1. The smallest term of the  $\lambda$ -product is  $\ell_s$  as for each N with (3.4), the value of  $\lambda_s$  is equal to one. On the other hand,  $\mu_s=m_{j,s}-1$  with  $m_{j,s}\in\{1,2\}$  as was discussed before Lemma 3.1. Hence, the smallest term of the  $\mu$ -product is  $\ell_s$  or  $\ell_{s-1}$  and (4.3) is valid. Suppose it is true for some  $p\geq 1$ . Let  $p=\lambda_s+\cdots+\lambda_i+\tau$  with  $0\leq \tau<\lambda_{i-1}$ . Then  $(p+1)\left(\prod_{j=0}^s\ell_j^{\lambda_j}\right)=p\left(\prod_{j=0}^s\ell_j^{\lambda_j}\right)\cdot\ell_{i-1}$ . In its turn,  $(p+1)\left(\prod_{j=0}^s\ell_j^{\mu_j}\right)=p\left(\prod_{j=0}^s\ell_j^{\lambda_j}\right)\cdot t$ . By (4.2),  $\mu_s+\mu_{s-1}+\cdots+\mu_i\leq p$ . Consequently, the product  $p\left(\prod_{j=0}^s\ell_j^{\mu_j}\right)$  consists of the corresponding powers of  $\ell_s,\ldots,\ell_i$ , possibly with some number of larger terms. Therefore, for p+1-st, the term t cannot be less than  $\ell_{i-1}$  and (4.3) is valid for each p.

The  $(\cdot)_p$ -version of (4.3)

$$\left(\prod_{j=0}^{s} \ell_{j}^{\lambda_{j}}\right)_{p} \leq \left(\prod_{j=0}^{s} \ell_{j}^{\mu_{j}}\right)_{p}$$

is also correct. Let us first show a stronger result.

**Lemma 4.2.** Given  $N \in \mathbb{N}$ , let  $Z = (x_k)_{k=1}^{N+1}$  be chosen in  $K^{\alpha}$  by the rule of increase of type. Suppose  $1 \le k \le N+1$  and  $(\mu_j)_{j=0}^s$  are degrees corresponding to  $x_k$ . For a fixed  $x \in K^{\alpha}$ , let  $(v_j)_{j=0}^s$  be defined as in Lemma 3.1 with x instead of  $\tilde{x}$ . Then for  $1 \le p \le N$  we have

$$\left(\prod_{j=0}^{s} \ell_{j}^{\nu_{j}}\right)_{p} \leq \left(\prod_{j=0}^{s} \ell_{j}^{\mu_{j}}\right)_{p}.$$

*Proof.* Let  $\sigma_q(\mu) = \mu_s + \mu_{s-1} + \cdots + \mu_q$  for  $0 \le q \le s$  with a similar definition of  $\sigma_q(v)$ . As in Lemma 3.1,  $I_{j_{s-n},n} = \tilde{I}_{j_{s-n},n}$  is the smallest basic interval containing both points  $x_k$  and x. Then  $\sigma_n(\mu) = \sigma_n(v)$  as  $\mu_i = v_i$  for  $0 \le i \le n-1$ . It follows that (4.5) is the equality for  $\sigma_n(\mu) \le p \le N$ . In particular,  $\left(\prod_{j=0}^s \ell_j^{\mu_j}\right)_{\sigma_n(\mu)} = \ell_{n-1}^{\mu_{n-1}} \cdots \ell_0^{\mu_0} = \ell_{n-1}^{v_{n-1}} \cdots \ell_0^{v_0}$ . For brevity, we denote this product by A.

By decreasing induction on p, suppose that  $\sigma_{n+1}(\mu) \leq p < \sigma_n(\mu)$  so  $p = \sigma_{n+1}(\mu) + \tau$  with  $0 \leq \tau < \mu_n$ . Removing the smallest p terms from  $\prod_{j=0}^s \ell_j^{\mu_j}$  gives  $\left(\prod_{j=0}^s \ell_j^{\mu_j}\right)_p = \ell_n^{\mu_n - \tau} \cdot A$ . It follows that to get  $\left(\prod_{j=0}^s \ell_j^{\nu_j}\right)_p$  we must multiply the product A by the  $\mu_n - \tau$  largest terms from  $\ell_s^{\nu_s} \cdots \ell_n^{\nu_n}$ . Since these terms do not exceed  $\ell_n$ , we have (4.5) for a given p.

Suppose, (4.5) is valid with the subscript  $\sigma_q(\mu)$  for n+1 < q < s. Let  $p = \sigma_{q+1}(\mu) + \tau$  with  $0 \le \tau < \mu_q$ . Then  $(\prod_{j=0}^s \ell_j^{\mu_j})_p = \ell_q^{\mu_q - \tau} \cdot (\prod_{j=0}^s \ell_j^{\mu_j})_{\sigma_q(\mu)}$ . On the other hand,  $(\prod_{j=0}^s \ell_j^{\nu_j})_p = t_1 \cdots t_{\mu_q - \tau} \cdot (\prod_{j=0}^s \ell_j^{\nu_j})_{\sigma_q(\mu)}$ , where  $(\prod_{j=0}^s \ell_j^{\nu_j})_{\sigma_q(\mu)} \le (\prod_{j=0}^s \ell_j^{\mu_j})_{\sigma_q(\mu)}$  by the induction hypothesis and  $(t_i)_{i=1}^{\mu_q - \tau}$  are the next descending members of  $\ell_s^{\nu_s} \cdots \ell_0^{\nu_0}$  after the largest  $\sigma_q(\mu)$  terms have been removed. By (3.11), these  $t_i$  are among  $\ell_s^{\nu_s} \cdots \ell_q^{\nu_q}$ , which completes the proof.

**Remark**. Multiplying the left side of (4.5) by some additional term  $\ell_j$  can only reduce it. This implies the inequality (4.4).

Now the task is to find an analog of the Markov property for the sets under consideration.

Let  $K \subset \mathbb{R}$  be a compact set of infinite cardinality. A sequence of *Markov's factors* for K is defined as  $M_N(K) = \inf\{M : |Q'|_K \le M |Q|_K, Q \in \mathscr{P}_N\}$  for  $N \in \mathbb{N}$ . Thus, there are constants C and r with  $M_N(K) \leq CN^r$  for all N under the Markov property of K. We see that  $M_N(K)$  is the norm of the differentiation operator D in the space  $(\mathscr{P}_N, |\cdot|_K)$ .

Given  $p, N \in \mathbb{N}$ , we define Markov's N-th factor of p-th order as the norm of  $D^p$ :

$$M_N^{(p)}(K) = \inf\{M : |Q^{(p)}|_K \le M |Q|_K, Q \in \mathscr{P}_N\}.$$

Clearly,  $M_N^{(p)}(K) \leq M_N(K)^p$ . This estimate is not rough for Markov sets. For example, if K = [-1,1] then, see, e.g., [3], p. 132,  $M_N^{(p)}(K) = \frac{N^2 \cdot (N^2 - 1) \cdots (N^2 - (p-1)^2)}{1 \cdot 3 \cdots (2p-1)}$  with  $M_N(K) = N^2$ . For the Cantor sets under consideration, the difference between  $M_N^{(p)}(K)$  and  $M_N(K)^p$  is

essential.

In the next lemma, given  $N \in \mathbb{N}$ , we use (3.4) and (3.8) for N+1,  $\rho_1 \cdots \rho_p =_p \left(\prod_{j=0}^s \ell_j^{\lambda_j}\right)$ .

**Theorem 4.3.** Given N and  $1 \le p < N$ , we have  $M_N^{(p)}(K^{\alpha}) \le h_0^{-N} \frac{(N+1)N^p}{\rho_1 \cdots \rho_n}$ .

*Proof.* Fix  $Q \in \mathscr{P}_N$ . There is no loss of generality in assuming  $|Q|_{K^{\alpha}} = 1$ . In view of (3.3), it suffices to show that

(4.6) 
$$\frac{|a_k^{(p)}(x)|}{|a_k(x_k)|} \le N^p h_0^{-N} (\rho_1 \cdots \rho_p)^{-1} \text{ for } 1 \le k \le N+1, x \in K^{\alpha}.$$

Fix k and x. Let degrees  $(v_j)_{j=0}^s$  correspond to x, as it was in Lemma 3.1 for  $\tilde{x}$ . Then  $|a_k(x)| = \prod_{i=1}^N d_i \le \prod_{j=0}^s \ell_j^{v_j}$  with  $d_i := d_i(x, Z \setminus \{x_k\})$ . Suppose  $v_s \ge 1$ . Then  $0 \le d_1 \le \ell_s$ . Due to the choice of the degrees, for every  $2 \le i \le N$  there exists n(i) with  $h_{n(i)} \le d_i \le \ell_{n(i)}$ . If  $v_s = 0$  this is also true for i = 1 with n(1) = s - 1. Here x and  $x_k$  belong to the same  $I_{j,s}$  with  $m_{j,s} = 1$ . Since  $p \ge 1$ , by (4.1),  $(\prod_{j=0}^{s} h_j^{v_j})_p \le (\prod_{i=1}^{N} d_i)_p \le (\prod_{j=0}^{s} \ell_j^{v_j})_p$ . By (3.1),  $|a_k^{(p)}(x)| \le N^p(\prod_{j=0}^s \ell_j^{v_j})_p$ . Applying Lemma 4.2 and (3.10) yields

$$\frac{|a_k^{(p)}(x)|}{|a_k(x_k)|} \le \frac{N^p(\prod_{j=0}^s \ell_j^{\mu_j})_p}{h_0^N \prod_{j=0}^s \ell_j^{\mu_j}} = \frac{N^p}{h_0^N p(\prod_{j=0}^s \ell_j^{\mu_j})},$$

which gives (4.6), by (4.3).

The inequality in the previous proposition is exact with respect to the terms  $\rho_i$ .

**Example.** Let  $N=2^s$ . Consider a polynomial  $\omega_N(x)=\prod_{k=1}^N(x-x_k)$  that has zeros at all points from  $Y_{s-1}$ . Then  $|\omega_N|_{K^{\alpha}}=|\omega_N(\ell_s)|\leq \ell_s\ell_{s-1}\ell_{s-2}^2\cdots\ell_0^{2^{s-1}}=\prod_{i=1}^N\rho_i$ . The exact value of  $|\omega_N|_0$  is  $\ell_s \prod_{i=2}^N d_i(x)$  with  $d_i(x) = d_i(x, (x_k)_{k=1}^N)$ . As above,  $h_0 \rho_i \le d_i(x) \le \rho_i$  for  $i \ge 2$ . Then  $|\omega_N^{(p)}(0)| \ge \prod_{i=p+1}^N d_i(0)$ , because  $\omega_N^{(p)}(0)$  is a sum of products of the same sign and one of them is  $\prod_{i=p+1}^N d_i(0)$ . Consequently,  $M_N^{(p)}(K^{\alpha}) \geq \frac{|\omega_N^{(p)}(0)|}{|\omega_N|_{\kappa^{\alpha}}} \geq h_0^{N-p}(\rho_1 \cdots \rho_p)^{-1}$ .

# 5. Jackson's type inequality by means of Faber bases

Suppose that X(K) is a space of functions on K, containing polynomials. A polynomial topological basis  $(Q_n)_{n=0}^{\infty}$  in X is called a Faber (or strict polynomial) basis if  $\deg Q_n = n$ for all n. Thus, for each  $f \in X$  there is a unique number sequence  $(\xi_n(f))_{n=1}^{\infty}$  such that the series  $\sum_{n=0}^{\infty} \xi_n(f) Q_n$  converges to f in the topology of X. This gives an estimate of the best uniform approximation of f by polynomials:

(5.1) 
$$E_N(f,K) \le \sup_{x \in K} |\sum_{n=N+1}^{\infty} \xi_n(f) Q_n(x)|.$$

**Example.** By Lemma 25 in [11], the Chebyshev polynomials  $(T_n)_{n=0}^{\infty}$  form a basis in the space  $C^{\infty}[-1,1]$ . The corresponding biorthogonal functionals are given as follows  $\xi_0(f) = \frac{1}{\pi} \int_0^{\pi} f(\cos t) dt$ ,  $\xi_n(f) = \frac{2}{\pi} \int_0^{\pi} f(\cos t) \cos nt dt$ ,  $n \in \mathbb{N}$ . By (44) in [11] (Jackson's theorem is not used!),  $|\xi_n(f)| \leq \frac{C_k |f|_k}{n^k}$  for each k. Since the basis is absolute and  $|T_n|_0 = 1$ , we get  $E_N(f, [-1, 1]) \leq \sum_{n=N+1}^{\infty} |\xi_n(f)|$ . This gives  $E_N(f, [-1, 1]) \leq \frac{C_q |f|_{q+1}}{N^q}$  for  $q \in \mathbb{N}$ .

In the case of small sets K, the phenomenon of ultra-fast convergence of polynomials to functions from  $\mathscr{E}(K)$  is observed. Let  $\omega_0 = 1$  and  $\omega_n(x) = \prod_{k=1}^n (x - x_k)$  for  $n \in \mathbb{N}$ , where the points  $(x_k)_{k=1}^n$  are chosen in  $K^{\alpha}$  by the rule of increase of type. Given  $f \in X(K)$  and  $n \in \mathbb{Z}_+$ , by  $\xi_n(f)$  we denote the divided difference  $[x_1, x_2, \cdots, x_{n+1}]f$ . The functionals  $(\xi_n)_{n=0}^{\infty}$  are biorthogonal to  $(\omega_n)_{n=0}^{\infty}$ . If  $\alpha \geq 2$  then, by Theorem 1 in [7], the sequence  $(\omega_n)_{n=0}^{\infty}$  is a basis in the space  $\mathscr{E}(K^{\alpha})$ . This allows us to evaluate  $E_N(f, K^{\alpha})$  in terms of the values  $(\rho_i)_{i=1}^{N+1}$  determined in (3.8). In order to do this, we define analogous  $\rho_k(n)$  for another n.

Let  $2^r \le n < 2^{r+1}$ . Then  $n = 2^r + 2^{r_1} + \dots + 2^{r_m}$  with  $0 \le r_m < \dots < r_1 < r_0 := r$ . In the same way as  $\pi_j$  for N+1, we define  $\pi_j(n)$  and  $(\lambda_j(n))_{j=0}^r$  so that  $\prod_{j=0}^m \pi_j(n) = \prod_{i=0}^r \ell_i^{\lambda_i(n)} = \prod_{k=1}^n \rho_k(n)$  with nondecreasing  $\rho_k(n)$ . Let us point out some obvious properties of the degrees  $\lambda_j(n)$ . First,  $\lambda_j(n) \le \lambda_j(n+1)$  with  $\lambda_j(n) = \lambda_j(n+1)$  for all j except some  $j_0$  for which  $\lambda_{j_0}(n+1) = \lambda_{j_0}(n) + 1$ . Secondly,  $\prod_{i=0}^r \ell_i^{\lambda_i(2^r)} = \pi_0(2^r) = \ell_r \ell_{r-1} \ell_{r-2}^2 \cdots \ell_0^{2^{r-1}}$ , whereas  $\prod_{i=0}^r \ell_i^{\lambda_i(2^{r+1}-1)} = \ell_r \ell_{r-1}^2 \ell_{r-2}^4 \cdots \ell_0^{2^r}$ .

Therefore, if n is as above, then  $\lambda_j(2^r) \leq \lambda_j(n) \leq \lambda_j(2^{r+1}-1)$  with  $\lambda_r(n)=1$  and  $2^{r-j-1} \leq \lambda_j(n) \leq 2^{r-j}$  for  $0 \leq j \leq r-1$ . If  $q=2^w < n$  then

(5.2) 
$$q\left(\prod_{j=0}^{r} \ell_{j}^{\lambda_{j}(n)}\right) \leq q\left(\prod_{j=0}^{r} \ell_{j}^{\lambda_{j}(2^{r})}\right) = \ell_{r}\ell_{r-1}\cdots\ell_{r-w}^{2^{w-1}}.$$

As in Proposition 4.3, for a given N with  $2^s \le N+1 < 2^{s+1}$ , we use (3.4) and (3.8) for N+1 and the corresponding  $(\rho_i)_{i=1}^{N+1}$  with  $\rho_i = \rho_i(N+1)$ .

**Theorem 5.1.** Suppose  $\alpha \ge 2$  and  $\ell_1 \le 1/4$ . Let N be as above. Then for each  $f \in \mathcal{E}(K^{\alpha})$  and  $q = 2^w$  with w < s - 8 we have  $E_N(f, K^{\alpha}) \le C_q \rho_1 \cdots \rho_q ||f||_{q_1}$ , where  $C_q$  does not depend on f and  $N, q_1 = 2^{w+8} + 1$ .

*Proof.* Fix N,q and f as above. By (5.1),  $E_N(f,K^\alpha) \leq \sum_{n=N+1}^\infty |\xi_n(f)| \cdot |\omega_n|_0$  with the decomposition  $\sum_{n=N+1}^\infty = \sum_{n=N+1}^{2^{s+1}-1} + \sum_{r=s+1}^\infty \sum_{n=2^r}^{2^{r+1}-1} =: \Sigma_1 + \Sigma_2$ . Let  $2^r \leq n < 2^{r+1}$  with  $r \geq s+1$ . To estimate  $|\xi_n(f)| \cdot |\omega_n|_0$  from above, we use the arguments of Theorem 1 in [7] with minor modifications. For each  $x \in K^\alpha$ , the estimate

Let  $2^r \le n < 2^{r+1}$  with  $r \ge s+1$ . To estimate  $|\xi_n(f)| \cdot |\omega_n|_0$  from above, we use the arguments of Theorem 1 in [7] with minor modifications. For each  $x \in K^{\alpha}$ , the estimate  $|\omega_n(x)| \le \prod_{j=0}^r \ell_j^{\lambda_j(n)}$  holds true. We apply (2) in [7] with  $q_1$  instead of q using the following two improvements. The Open Mapping Theorem can be applied to the space  $\mathcal{E}^{q_1}(K^{\alpha})$ , giving the  $q_1$  version of (2.6) instead of (3) in [7]. And three lines above (3) in [7] we do not

replace  $1 - 2\ell_1 = h_0$  by  $\ell_1$ . This gives  $|\xi_n(f)| \le C||f||_{q_1} 2^n \cdot h_0^{-n} (\prod_{j=0}^r \ell_j^{\lambda_j(n)})_{q_1}^{-1}$ . Hence,

$$|\xi_n(f)| \cdot |\omega_n|_0 \le C ||f||_{q_1} \left(\frac{2}{h_0}\right)^n q_1 \left(\prod_{j=0}^r \ell_j^{\lambda_j(n)}\right).$$

Let's show that  $\sum_{n=2^r}^{2^{r+1}-1} \left(\frac{2}{h_0}\right)^n q_1 \left(\prod_{j=0}^r \ell_j^{\lambda_j(n)}\right) \leq q \left(\prod_{j=0}^r \ell_j^{\lambda_j(2^r)}\right)$ . By condition,  $\frac{2}{h_0} \leq \frac{1}{\ell_1}$ , so  $\left(\frac{2}{h_0}\right)^n < \ell_1^{-2^{r+1}}$ . On the other hand, the second term of the product in the sum above takes maximum value if  $n=2^r$ . Therefore the whole sum does not exceed  $2^r \ell_1^{-2^{r+1}} q_1 \left(\prod_{j=0}^r \ell_j^{\lambda_j(2^r)}\right)$ . Here the last term is  $q \left(\prod_{j=0}^r \ell_j^{\lambda_j(2^r)}\right) \cdot \{\ell_{r-w-1}^{2^w} \cdots \ell_{r-w-8}^{2^{w+7}} \cdot \ell_{r-w-8}\}$ . Of course,  $2^r \ell_{r-w-8} < 1$  for sufficiently large r. Also we have in braces 8 terms of the type  $\ell_{r-k-1}^2$ . Since  $\alpha \geq 2$ , for each of them we have  $\ell_{r-k-1}^{2^k} = \ell_1^{\alpha^{r-k-2}2^k} \leq \ell_1^{2^{r-2}}$ , so their product neutralizes  $\ell_1^{-2^{r+1}}$ . From this

$$\Sigma_2 \leq C ||f||_{q_1} \sum_{r=s+1}^{\infty} q \left( \prod_{j=0}^{r} \ell_j^{\lambda_j(2^r)} \right).$$

It is easy to check that the first term in the above sum dominates, so the whole sum does not exceed twice the first term. By monotonicity,  $\lambda_j(2^{s+1}) \ge \lambda_j(N+1)$  for all j. Therefore,

$$\Sigma_2 \leq 2C ||f||_{q_1} \cdot_q \left( \prod_{j=0}^s \ell_j^{\lambda_j(N+1)} \right) = 2C ||f||_{q_1} \rho_1 \cdots \rho_q.$$

Similar arguments apply to  $\Sigma_1$  with  $N+1 \leq n < 2^{s+1}$ , but now we estimate  $|\xi_n(f)| \cdot |\omega_n|_0$  directly using  $q\left(\prod_{j=0}^r \ell_j^{\lambda_j(N+1)}\right)$ . This gives the desired result.

**Remark.** The condition  $\ell_1 \leq 1/4$  is not particularly restrictive. Enlarging  $q_1$  allows us to neutralize  $\left(\frac{2}{h_0}\right)^n$  for larger values of  $\ell_1$  as well. However, the condition  $\alpha \geq 2$  is important here since the sequence  $(\omega_n)_{n=0}^{\infty}$  is not a basis in  $\mathscr{E}(K^{\alpha})$  for  $\alpha < 2$ . We believe that these spaces also have Faber interpolation bases with a different, more complex choice of interpolation nodes, but we cannot present them. For this reason, our main result is given only for  $\mathscr{E}(K^2)$ .

Comparison of Theorems 4.3 and 5.1 shows that  $E_N(f, K^{\alpha})$  successfully neutralizes the fast growth of factors  $M_N^{(p)}(K^{\alpha})$  for all  $\alpha \geq 2$ . Let us show that an even stronger fact holds for  $\alpha = 2$ .

**Proposition 5.2.** For each p and r > p there exists  $r_1$  such that

$$M_N^{(p)}(K^2) \cdot E_{N-1}(f, K^2) \le \rho_{p+1}(N+1) \cdots \rho_r(N+1) ||f||_{r_1}$$

for sufficiently large N. Here we assume  $\ell_1 \leq 1/3$ .

*Proof.* As above,  $2^s \le N+1 < 2^{s+1}$ . Recall that  $M_N^{(p)}(K^2)$  is determined by  $\rho_k(N+1)$ , whereas  $E_{N-1}(f,K^2)$  is given in terms of  $\rho_k(N)$ . Fix p and r > p. There is no loss of generality in assuming  $r = 2^w$ . Set  $r_1 = 2^{w+10}$ . We apply Theorem 5.1 with r instead of q. We

need to show

$$C_r(N+1)N^ph_0^{-N}\frac{\rho_1(N)\cdots\rho_{r_1}(N)}{\rho_1(N+1)\cdots\rho_p(N+1)} \le \rho_{p+1}(N+1)\cdots\rho_r(N+1)$$

or equivalently

$$C_r(N+1)N^ph_0^{-N}rac{
ho_1(N)\cdots
ho_{r-1}(N)}{
ho_1(N+1)\cdots
ho_r(N+1)}
ho_r(N)\cdots
ho_{r_1}(N)\leq 1.$$

The fraction in the middle is  $\frac{r-1\left(\prod_{j=0}^{s}\ell_{j}^{\lambda_{j}(N)}\right)}{r\left(\prod_{j=0}^{s}\ell_{j}^{\lambda_{j}(N+1)}\right)}$ , where the denominator contains all the factors

of the numerator, so the fraction is  $\frac{1}{\ell_j}$  with some  $0 \le j \le s$ . In the worst case it is  $\frac{1}{\ell_s}$ . In addition,  $h_0 \ge \ell_1$  and  $N < 2^{s+1}$  imply  $h_0^{-N} < \ell_1^{-2^{s+1}} = \ell_s^{-4}$ . It suffices to prove that

$$(5.3) C_r(N+1)N^p \rho_r(N) \cdots \rho_{r_1}(N) \le \ell_s^5.$$

Since  $N \geq 2^s-1$ , we have  $\prod_{j=0}^N \rho_j(N) \leq \prod_{j=0}^N \rho_j(2^s-1) = (\ell_s \ell_{s-1}^2 \cdots \ell_{s-w}^{2^{w-1}}) \cdot \ell_{s-w-1}^{2^w} \cdot \ell_0^{2^s}$ , where the product in parentheses contains r-1 terms. We can only enlarge the left side of (5.3) by replacing  $\rho_r(N) \cdots \rho_{r_1}(N)$  with  $\ell_{s-w-1}^{2^w} \cdots \ell_{s-w-10}^{2^{w+9}} \cdot \ell_{s-w-11}$ , containing exactly  $r_1-r+1$  terms. Now the product  $C_r(N+1)N^p \cdot \ell_{s-w-11}$  does not exceed 1 for sufficiently large N. Also,  $\ell_{s-k-1}^{2^k} = \ell_1^{2^{s-2}}$ , and we have 10 such terms, so their product is  $\ell_1^{10 \cdot 2^{s-2}}$ , which is equal to  $\ell_s^5$ . This is the desired conclusion.

#### 6. SIMULTANEOUS EXTENSIONS OF BASIC POLYNOMIALS

How does  $K^2$  with the extension property differ from  $K^{\alpha}$ ,  $\alpha > 2$ , without it? Let us show that the difference depends mainly on the possibility of suitable individual extensions of the basic interpolation polynomials.

Let K and  $Q \in \mathscr{P}_N$  be such as in Section 2. We fix  $\delta > 0$  and a segment I containing  $K_{\delta}$ . Write  $\tilde{Q} = Q \cdot u_{\delta}$ . Clearly,  $|Q|_{0,K} \leq |\tilde{Q}|_{0,I}$ . We will use the notation  $\tilde{Q} \sim Q$  if  $|\tilde{Q}|_{0,I} \leq C|Q|_{0,K}$  with some C independent of Q and N. If K is Markov, then by (2.2) the choice  $\delta = N^{-r}$  provides  $|\tilde{Q}|_{0,I} \leq C|Q|_{0,K}$ . In addition, suppose that Q realizes  $M_N^{(p)}(K)$  for p < N. This means that the converse of (2.1) is true with some constant  $c_1$  instead of  $C_1$ , similar to how Chebyshev polynomials implement  $M_N^{(p)}([-1,1])$ . Then by (2.3),  $\tilde{Q}^{(p)} \sim Q^{(p)}$ , so there exists  $\delta = \delta(N)$  which can be applied to extend both Q and its derivatives.

In general,  $\delta$  depends on p. Let us illustrate this with an apparent example. From now on,  $u_{\delta}$  is determined by the function  $\varphi$  with  $\varphi(x) = 1$  for  $x \le 0$ ,  $\varphi(x) = 0$  for  $x \ge 1$  and

(6.1) 
$$\varphi(x) = \exp\left[\frac{1}{x-1}\exp\left(-\frac{1}{x}\right)\right]$$

for 0 < x < 1. We see that  $\varphi \in C^{\infty}(\mathbb{R})$  with  $\varphi(\frac{1}{2}) > \frac{1}{2}$ . Given  $\delta > 0$ , let  $\varphi_{\delta}(x) = \varphi(x/\delta)$ . Given K with a complementary interval (a,b) with  $b-a \geq 2\delta$ , we define  $u_{\delta} = 1$  on K,  $u_{\delta}(x) = \varphi_{\delta}(x-a)$  for  $a < x < a+\delta, u_{\delta}(x) = b-x$  for  $b-\delta < x < b$ , and  $u_{\delta}(x) = 0$  for  $a+\delta \leq x \leq b-\delta$ . If  $b-a < 2\delta$  then  $u_{\delta} = 1$  on (a,b). Then  $u_{\delta}$  has the desired properties indicated in Section 2.

**Example.** Let  $K = [-\varepsilon, \varepsilon]$  for a small  $\varepsilon$ . Of course, K is Markov, but with the constant  $C_1$  in (2.1) depending essentially on  $\varepsilon$ . Indeed, fix  $\delta > 0$ . As above,  $\tilde{Q} = Q \cdot u_{\delta}$ . Suppose that  $|\tilde{Q}|_{0,I} \leq C|Q|_{0,K}$  and  $|\tilde{Q}''|_{0,I} \leq Cn^2|Q|_{2,K}$  hold for  $Q \in \mathscr{P}_n$ , where I = [-1,1]. Then for Q(x) = x we have  $|\tilde{Q}|_{0,I} \geq \tilde{Q}(\varepsilon + \frac{\delta}{2}) > (\varepsilon + \frac{\delta}{2})\frac{1}{2} > \frac{\delta}{4}$  as  $\varphi_{\delta}(\frac{\delta}{2}) > \frac{1}{2}$ . On the other hand,  $Q'(\varepsilon) = 1, Q'(\varepsilon + \delta) = 0$ , so, by the mean value theorem,  $|\tilde{Q}''|_{0,I} \geq \frac{1}{\delta}$ . Here,  $|Q|_{0,K} = \varepsilon, |Q|_{2,K} = 1$ . Hence, if the inequalities above hold then  $\frac{\delta}{4} \leq C\varepsilon$  and  $\frac{1}{\delta} \leq C$ , which means that  $C \geq \frac{1}{2\sqrt{\varepsilon}}$ .

An extension of polynomials that provides the above equivalence can be called *strong* simultaneous extensions. The term simultaneous extensions was used in [12] for the existence of a linear extension operator. In our case, instead of (2.2), we consider the following condition for the sequence  $(\omega_n)_{n=0}^{\infty}$  defined in Section 5

$$(6.2) \exists (\delta_n)_{n=1}^{\infty} : \forall p \exists q, C : |\tilde{\omega}_n|_p \leq C|\omega_n|_q, \ n \in \mathbb{N}.$$

**Proposition 6.1.** Choosing  $\delta_n = \ell_s$  for  $2^s \le n < 2^{s+1}$  provides (6.2) on  $K^2$ . Here we assume  $\ell_1 \le 1/3$ .

*Proof.* The previous formulas are given mainly for N+1, so we prove (6.2) for n=N+1. Both parts of (6.2) will be expressed in terms of  $\rho_k(N+1)$  with N as in (3.4). Thus,  $\delta_{N+1} = \ell_s$ . Since  $\delta_{N+1}$  is not included in the right-hand side of (6.2), we first estimate  $||\omega_{N+1}||_q$  for a given  $1 \le q < N+1$ . Henceforth  $q=2^w+1$ , where w will be defined later depending on p. Our claim is that

(6.3) 
$$|\omega_{N+1}^{(q)}| \ge h_0^{N+1-q} \cdot \rho_q \cdots \rho_{N+1}.$$

Recall that  $\omega_{N+1}(x) = \prod_{i=1}^{N+1}(x-x_i)$ , where  $Z=(x_k)_{k=1}^{N+1}$  are chosen by the rule of increase of type, so  $x_1=0$ . Then  $a_1(x)=\prod_{i=2}^{N+1}(x-x_i)$ . Let  $d_k:=d_k(0,Z)$ . Then, of course,  $d_1=0$  and  $d_{k+1}=d_k(0,Z\setminus\{0\})$  for  $1\leq k< N$ . The function  $a_1$  determines the powers  $(\mu_j)_{j=0}^s$  such that (3.10) holds for  $|a_1(0)|=\prod_{k=1}^N d_k(0,Z\setminus\{0\})$ . As in Theorem 4.3, for each  $1\leq k< N$  there is an index j with  $h_j\leq d_k(0,Z\setminus\{0\})\leq \ell_j$ . Removing the q-1 smallest terms from the above product gives  $\prod_{k=q}^N d_k(0,Z\setminus\{0\})\geq \left(\prod_{j=0}^s h_j^{\mu_j}\right)_{q-1}\geq h_0^{N+1-q}\left(\prod_{j=0}^s \ell_j^{\mu_j}\right)_{q-1}$ .

We now turn to  $\omega_{N+1}^{(q)}$ . It is a sum of  $\frac{(N+1)!}{(N+1-q)!}$  products, each containing N+1-q terms, so they all have the same sign, and one of them is  $d_{q+1}\cdots d_{N+1}$ . Therefore,

$$|\omega_{N+1}^{(q)}| \ge \prod_{k=q+1}^{N+1} d_k = \prod_{k=q}^{N} d_k(0, Z \setminus \{0\}) \ge h_0^{N+1-q} \left(\prod_{j=0}^{s} \ell_j^{\mu_j}\right)_{q-1}.$$

Applying (4.4) yields (6.3).

We proceed to estimate from above  $|\tilde{\omega}_{N+1}^{(p)}(x)|$  for fixed p and x. It is clear that only x outside the set should be considered. Fix x with  $0 < \operatorname{dist}(x, K^2) = |x - y| < \ell_s$ . By (3.8),  $|\omega_{N+1}(y)| \le \prod_{k=1}^{N+1} \rho_k$ , so  $|\omega_{N+1}(x)| \le \prod_{k=1}^{N+1} (\rho_k + \ell_s)$ . Here,  $\rho_1 = \ell_s$ ,  $\rho_2 = \ell_{s-1}$  and  $\rho_3 = \ell_{s-1}$  or  $\ell_{s-2}$ , depending on the value of N+1. In a fairly straightforward way, one can show that  $\prod_{k=2}^{N+1} (\rho_k + \ell_s) \le 2 \prod_{k=2}^{N+1} \rho_k$ . This and (3.1) give  $|\omega_{N+1}^{(j)}(x)| \le 4(N+1)^j \prod_{k=j+1}^{N+1} \rho_k$ . Also,

 $|u_{\delta_{N+1}}^{(i)}(x)| \le c_i \ell_s^{-i}$ . Without loss of generality, we can assume that  $c_i$  increases with i. Then, by Leibnitz's rule,

$$|(\omega_{N+1} \cdot u_{\delta})^{(p)}(x)| \le 4c_p \sum_{j=0}^p {p \choose j} (N+1)^j t_j,$$

where  $t_j := \ell_s^{-p+j} \prod_{k=j+1}^{N+1} \rho_k$ . We see that  $t_0 = t_1 = \ell_s^{-p+1} \cdot \rho_2 \cdots \rho_{N+1}$ , while the following terms decrease very rapidly. Hence,  $|(\omega_{N+1} \cdot u_\delta)^{(p)}(x)| \le 4c_p t_0 [1+p(N+1)+o(1)] \le c_p' N t_0$ , where  $c_p'$  does not depend on N. Combining this with (6.3) we reduce the desired inequality to  $N \cdot \rho_2 \cdots \rho_{q-1} \le C \ell_1^{N+1-q} \cdot \ell_s^{p-1}$ , because  $\ell_1 \le h_0$  by condition. Let us replace  $\rho_k$  with  $\rho_k'$  determined by  $2^s$ . Then  $\prod_{k=2}^{q-1} \rho_k' = \ell_{s-1} \ell_{s-2}^2 \cdots \ell_{s-w}^{2^{w-1}} \ge \prod_{k=2}^{q-1} \rho_k$ . It remains to prove that

(6.4) 
$$N \cdot \ell_{s-1} \ell_{s-2}^2 \cdots \ell_{s-w}^{2^{w-1}} \le C \ell_1^{N+1-q} \cdot \ell_s^{p-1}.$$

In the case under consideration, we have  $\ell_k = \ell_1^{2^{k-1}}$ . Therefore, the left side of (6.4) is  $N \ell_1^{w2^{s-2}}$  and  $C \ell_1^{N+1-q+(p-1)2^{s-1}}$  is on the right. Since  $N+1 < 2^{s+1}$ , we reduce (6.4) to  $2^{s+1} \ell_1^{2^{s-2}(w-2p-6+q)} \le C$ , which holds for  $s \ge 3$  provided w = 2p+6.

Small values of s do not cause problems, since  $|\omega_{N+1}^{(q)}| = (N+1)!$  for such s and given q and  $|\tilde{\omega}_n|_p \le c_p' N t_0$  with  $N \le 6$  and  $t_0 \le \ell_2^{-p}$ .

**Proposition 6.2.** If  $\alpha > 2$  then (6.2) is not valid on  $K^{\alpha}$ .

*Proof.* By [7], the polynomials  $(\omega_n)_{n=0}^{\infty}$  form an absolute topological basis in the space  $\mathscr{E}(K^{\alpha})$ . If (6.2) holds then

(6.5) 
$$W(f) = \sum_{n=0}^{\infty} \xi_n(f)\tilde{\omega}_n$$

is a linear continuous extension operator. However, by [5],  $\mathscr{E}(K^{\alpha})$  does not have EP for  $\alpha > 2$ .

The following example illustrates the above proposition. We will directly show the absence of (6.2) for  $\alpha > 2$ . We restrict ourselves to the same value of  $\delta$  as in Proposition 6.1. Let us consider the function  $\varphi$  in more detail. Let  $\tau(x) := \exp\left(-\frac{1}{x}\right)$ ,  $Q_0(x) := x - 1 - x^2$ . Then  $\varphi'(x) = \varphi(x) \tau(x) (x - x^2)^{-2} Q_0(x)$  and by induction

(6.6) 
$$\varphi^{(k)}(x) = \varphi(x) \tau(x) (x - x^2)^{-2k} Q_{k-1}(\tau(x))$$

for  $k \geq 2$ . Here,  $Q_k(\tau(x)) = Q_0(x)Q_{k-1}(\tau(x))\tau(x) + Q_{k-1}(\tau(x))r_k(x) + (1-x)^2\frac{dQ_{k-1}}{d\tau}\tau(x)$  with  $r_k(x) := (1-x)^2 + 2kx(1-x)(2x-1)$  for  $k \in \mathbb{N}$ . From this it follows that the signs of  $\varphi^{(k)}(x)$  and  $Q_{k-1}(\tau(x))$  coincide and  $Q_k(\tau(1)) = (-1)^{k+1}e^{-k}$ . Assume that  $k \in \mathbb{N}$  is even. Then  $\varphi^{(k)}(x) > 0$  and  $\varphi^{(k-1)}(x) < 0$  near point 1.

We will denote by  $\eta_k$  the number  $\min\{\eta: |Q_k(\tau(x)) - Q_k(\tau(1))| \leq \frac{1}{2e^k} \text{ for } \eta \leq x \leq 1\}$ . Let  $\theta_k := \max\{\eta_k, \eta_{k-1}, 1 - \frac{1}{4\sqrt{ke}}\}$ . Then

(6.7) 
$$\theta_k Q_{k-1}(\tau(\theta_k)) - 2k(\theta_k - \theta_k^2)^2 |Q_{k-2}(\tau(\theta_k))| > \frac{1}{2e^k}.$$

Indeed,  $\theta_k > \frac{7}{8}$  for  $k \geq 2$ ,  $Q_{k-1}(\tau(\theta_k)) \geq \frac{1}{2e^{k-1}}$  and  $|Q_{k-2}(\tau(\theta_k))| \leq \frac{3}{2e^{k-2}}$ . Also,  $\theta_k^2 < 1$  and  $(1-\theta_k)^2 \leq \frac{1}{16ke}$ . Therefore, the left side of (6.7) exceeds  $\frac{7}{16e^{k-1}} - \frac{3}{16e^{k-1}}$ , which gives the desired inequality. Let  $A_k := \varphi(\theta_k)\tau(\theta_k)(\theta_k - \theta_k^2)^{-2k}\frac{1}{2e^k}$ .

**Example.** Assume that  $\alpha > 2$ . Let  $N = 2^s$ ,  $\delta = \ell_s$ . Then there is p and z such that for each q and C we have for sufficiently large s

$$|\tilde{\boldsymbol{\omega}}_{N}^{(p)}(z)| > C|\boldsymbol{\omega}_{N}|_{q}.$$

Let, as above,  $Z = (x_k)_{k=1}^N, d_k(x) := d_k(x, Z)$ . In view of the structure of the set, we have  $d_k(1) \ge d_k(x)$  for each  $x \in K^{\alpha}$  if  $k \ge 2$ , so, by (3.1),  $|\omega_N|_q \le N^q \prod_{k=q+1}^N d_k(1)$  for  $q \ge 1$ .

Let us fix an even  $p > \frac{2\alpha - 3}{\alpha - 2}$  and  $z = 1 + \theta_p \delta$ . We aim to estimate  $|\tilde{\omega}_N^{(p)}(z)|$  from below. Since  $u_{\delta}(x) = \varphi(\frac{x-1}{\delta})$  for  $1 < x < 1 + \delta$ , we have  $u_{\delta}^{(j)}(z) = \ell_s^{-j} \varphi^{(j)}(\theta_p)$ . It follows that

$$\tilde{\omega}_{N}^{(p)}(z) = \omega_{N}(z) \, \ell_{s}^{-p} \varphi^{(p)}(\theta_{p}) + p \omega_{N}'(z) \, \ell_{s}^{-p+1} \varphi^{(p-1)}(\theta_{p}) + \sum_{i=2}^{p} \binom{p}{j} \omega_{N}^{(j)}(z) \ell_{s}^{-p+j} \varphi^{(p-j)}(\theta_{p}).$$

Since p is even, the first term on the right-hand side is positive and the second is negative. In our case, all values  $\omega_N^{(j)}(z)$  are positive. In particular,  $\omega_N(z) = \prod_{k=1}^N d_k(z) = \theta_p \ell_s \cdot \prod_{k=2}^N d_k(z)$  and  $\omega_N'(z) = \sum_{i=1}^N \prod_{k=1, k \neq i}^N d_k(z) = \prod_{k=2}^N d_k(z) [1 + d_1(z) \sum_{i=2}^N d_k^{-1}(z)]$ , where the expression in square brackets is smaller than 2, as is easy to check. By (3.1),  $\omega_N^{(j)}(z) \leq N^j \prod_{k=j+1}^N d_k(z)$ , so the modulus of the last sum in the above representation of  $\tilde{\omega}_N^{(p)}(z)$  can be estimated from above by  $C_1 \sum_{j=2}^p t_j$ , where  $C_1 = \max_{0 \leq j \leq p-2} |\varphi|_{j,[0,1]}$  and  $t_j = \binom{p}{j} N^j \ell_s^{-p+j} \prod_{k=j+1}^N d_k(z)$ . It is a simple matter to show  $2t_{j+1} < t_j$ . Hence,  $C_1 \sum_{j=2}^p t_j \leq 2C_1 t_2$ . Therefore,

$$\tilde{\omega}_{N}^{(p)}(z) \geq \ell_{s}^{-p+1} \prod_{k=2}^{N} d_{k}(z) [\theta_{p} \varphi^{(p)}(\theta_{p}) - 2p \varphi^{(p-1)}(\theta_{p}) - 2C_{1} p^{2} N^{2} \ell_{s} d_{3}^{-1}(z)].$$

By (6.6),  $\theta_p \varphi^{(p)}(\theta_p) - 2p \varphi^{(p-1)}(\theta_p) = A_p$ . The last term in brackets is arbitrarily small for sufficiently large s. It remains to prove that for each C and q there is  $s_0$  such that  $\ell_s^{-p+1} \prod_{k=2}^N d_k(z) > CN^q \prod_{k=q+1}^N d_k(1)$  for  $s > s_0$ . Of course,  $d_k(z) > d_k(1)$  for each k, so we reduce the desired inequality to  $CN^q \ell_s^{p-1} < \prod_{k=2}^q d_k(1)$ . Let  $2^{w-1} < q \le 2^w$  for some w. Then  $\prod_{k=2}^q d_k(1) \ge \prod_{k=2}^{2^w} d_k(1)$ . Here,  $d_2(1) = \ell_{s-1}$  In its turn,  $\ell_{s-2} - \ell_{s-1} \le d_3, d_4 \le \ell_{s-2}, \dots, \ell_{s-w} - \ell_{s-w+1} \le d_{2^{w-1}+1}, \dots, d_{2^w} \le \ell_{s-w}$ . Thus  $\prod_{k=2}^{2^w} d_k(1) \ge \pi_w \cdot \ell_{s-1} \ell_{s-2}^2 \cdots \ell_{s-w}^{2^{w-1}}$ , where  $\pi_w = \prod_{k=1}^{w-1} \left(1 - \frac{\ell_{s-k}}{\ell_{s-k-1}}\right)^{2^k}$ . An easy computation shows that  $\pi_w > \frac{1}{2}$ , so (6.8) holds if

$$2C2^{sq}\ell_s^{p-1} < \ell_{s-1}\ell_{s-2}^2 \cdots \ell_{s-w}^{2^w}.$$

Here, the right side is  $\ell_1^{\kappa}$  with  $\kappa = \alpha^{s-2} + 2\alpha^{s-3} + \dots + 2^{w-1}\alpha^{s-w-1} = \alpha^{s-2}[1 + \frac{2}{\alpha} + \dots + (\frac{2}{\alpha})^{w-1}] < \frac{\alpha^{s-1}}{\alpha-2}$ . On the other hand,  $\ell_s^{p-1} = \ell_1^{(p-1)\alpha^{s-1}}$  with  $(p-1)\alpha^{s-1} - \kappa > \alpha^{s-1}$ , due to the choice of p. Clearly,  $2C2^{sq}\ell_s < 1$  for sufficiently large s.

7. CONTINUITY OF PAWŁUCKI-PLEŚNIAK'S OPERATOR FOR A NON-MARKOV SET Let points  $Z_N = (x_k)_{k=1}^N$  be chosen in  $K^2$  by the rule of increase of type and

$$(7.1) W(f,x) = L_1(f,x;Z_1) \cdot u_{\delta_1}(x) + \sum_{N=1}^{\infty} [L_{N+1}(f,x;Z_{N+1}) - L_N(f,x;Z_N)] \cdot u_{\delta_N}(x),$$

where  $\delta_N = \ell_s$  for  $2^s \le N < 2^{s+1}$ .

Compare the proofs of the following theorem and T.2.1.

**Theorem 7.1.** The operator  $W : \mathcal{E}(K^2) \longrightarrow C^{\infty}([-2,2])$  is bounded.

*Proof.* Let  $G_N$  denote the N-th term of the series (7.1). Fix  $f \in \mathscr{E}(K^2), p \in \mathbb{N}$  and  $x \in \mathbb{R}$ . By Newton's form of the interpolation operator,  $G_N(f,x) = \xi_N(f)\tilde{\omega}_N$ . We aim to show that  $|G_N^{(p)}(f,x)|$  is a term of a series that converges uniformly with respect to x. Proposition 6.1 gives q and C with  $|G_N^{(p)}(f,x)| \leq C|\xi_N(f)| \cdot |\omega_N^{(q)}|_0 \leq CM_N^{(q)} \cdot |\xi_N(f)| \cdot |\omega_N|_0$ . Here and below, to simplify the writing, we omit the argument  $K^2$  of  $M_N^{(q)}(\cdot), \Lambda_j(\cdot)$ , and  $E_j(f,\cdot)$ .

Next,  $|\xi_N(f)| \cdot |\omega_N|_0 = |L_{N+1}(f) - L_N(f)| \le (\Lambda_{N+1} + 1) E_N(f) + (\Lambda_N + 1) E_{N-1}(f)$  by the argument from Lebesgue's Lemma, so  $|\xi_N(f)| \cdot |\omega_N|_0 \le (\Lambda_{N+1} + \Lambda_N + 2) E_{N-1}(f)$ . Applying Proposition 4.1 yields  $\Lambda_{N+1} + \Lambda_N + 2 \le 2h_0^{-N}N$ , as is easy to check. Therefore,

$$|G_N^{(p)}(f,x)| \le 2CNh_0^{-N}M_N^{(q)} \cdot E_{N-1}(f).$$

By Proposition 5.2, for each r > q there exists  $r_1$  such that

$$M_N^{(q)} \cdot E_{N-1}(f) \le \rho_{q+1}(N+1) \cdots \rho_r(N+1) ||f||_{r_1}.$$

Arguing in the same way as in the proof of Proposition 5.2, we can take a sufficiently large number of terms  $\rho_j(N+1)$  in such a way that  $\rho_{q+1}(N+1)\cdots\rho_m(N+1)\leq \ell_s$  and  $2CNh_0^{-N}\rho_{m+1}(N+1)\cdots\rho_r(N+1)\leq 1$ . This gives  $|G_N^{(p)}(f,x)|\leq \ell_s||f||_{r_1}$ . Of course, the series  $\sum_{s=1}^{\infty}2^s\ell_s$  converges.

**Conclusions.** At least for the considered case,

- 1. The operator W is continuous not only in  $\tau_I$ , but also in the stronger Whitney topology.
- 2. This can be shown by a modification of Pleśniak's argument.
- 3. The difference between sets  $K^{\alpha}$  with and without the extension property does not depend on the growth rate of Markov's factors but rather on the existence of suitable individual extensions of  $\omega_N$ .
- 4. Since these polynomials form a topological basis in the corresponding Whitney space, the operator (7.1) given by the interpolation method coincides with the operator (6.5) obtained by extensions of the basis elements. This method goes back to Mityagin [11].

This coincidence can be clearly observed in the following model case.

**Example.** Fekete points  $\mathscr{X}_n$  are known for K = [-1,1], see, e.g., [16], p.382. They are zeros of  $\omega_n(x) := (1-x^2)P_{n-2}^{1,1}(x)$ , where  $P_{n-2}^{1,1}$  is the Jacobi polynomial with the parameters  $\alpha = \beta = 1$ . By [2], for any admissible parameters, the Jacobi polynomials form a basis in  $C^{\infty}[-1,1]$ . The value  $\delta_n = n^{-2}$  defines  $\tilde{\omega}_n$  and the extension operator (7.1) that can be written as  $W(f) = \sum_{n=1}^{\infty} \xi_N(f) \tilde{\omega}_N$ .

# REFERENCES

- [1] M. Altun, A. Goncharov, A local version of the Pawlucki-Pleśniak extension operator, J. Approx. Th. 132 (2004), 34-41.
- [2] G.I. Balikov, *Polynomial bases in the space of infinitely differentiable functions on an interval*, C. R. Acad. Bulgare Sci. **34** (1981), 919–921.
- [3] R. A. DeVore, G. G. Lorentz, Constructive Approximation, Springer-Verlag, 1993.
- [4] A. Goncharov, A compact set without Markov's property but with an extension operator for  $C^{\infty}$  functions, Studia Math. 119 (1996), 27-35.
- [5] A. Goncharov, Perfect sets of finite class without the extension property, Studia Math. 126 (1997), 161-170.
- [6] A. Goncharov, On extension property of Cantor-type sets. Function spaces, 129–137, World Sci. Publ., River Edge, NJ, 2003.
- [7] A. Goncharov, Bases in the spaces of  $C^{\infty}$ -functions on Cantor-type sets, Constr. Approx. 23 (2006), no. 3, 351-360.
- [8] A. Goncharov, Z. Ural, *Mityagin's extension problem. Progress report*, J. Math. Anal. Appl. **448** (2017), no. 1, 357-375.
- [9] A. Goncharov, Y. Şengül, *Quasi-equivalence of bases in some Whitney spaces*, Canad. Math. Bull. 65 (2022), no. 1, 106–115.
- [10] A. Goncharov, *Bases in the spaces of Whitney jets*, Banach J. Math. Anal. **16** (2022), no. 1, Paper No. 16, 17 pp.
- [11] B. S. Mitjagin, *Approximate dimension and bases in nuclear spaces*, Uspehi Mat. Nauk **16**, 63-132 (1961), (Russian).
- [12] Z. Ogrodzka, On simultaneous extension of infinitely differentiable functions, Studia Math. 28 (1966/67), 193–207.
- [13] W. Pawłucki and W. Pleśniak, *Extension of C*<sup>∞</sup> functions from sets with polynomial cusps, Stud. Math. **88** (1988), 279-287.
- [14] W. Pleśniak, Markov's Inequality and the Existence of an Extension Operator for  $C^{\infty}$  functions, J. Approx. Th. **61** (1990), 106-117.
- [15] W. Pleśniak, A Cantor regular set which does not have Markov's property, Ann. Polon. Math. **51**, (1990), 269-274.
- [16] G. Szegő, *Orthogonal polynomials*, 4th ed., American Mathematical Society, Providence, R.I., 1975. American Mathematical Society, Colloquium Publications, Vol. XXIII.
- [17] M. Tidten, Fortsetzungen von  $C^{\infty}$  Funktionen, welche auf einer abgeschlossenen Menge in  $\mathbb{R}^n$  definiert sind, Manuscripta Math. 27 (1979), 291-312.
- [18] H. Whitney, *Analytic extensions of differentiable functions defined in closed sets*, Trans. Amer. Math. Soc. **36** (1934), 63-89.
- [19] M. Zerner, Développement en séries de polynômes orthonormaux des fonctions indéfiniment différentiables, C.R.Acad.Sci.Paris Sér.I, **268** (1969), 218-220.

DEPARTMENT OF MATHEMATICS, BILKENT UNIVERSITY, 06800 ANKARA, TURKEY *Email address*: goncha@fen.bilkent.edu.tr

DEPARTMENT OF MATHEMATICS, BILKENT UNIVERSITY, 06800 ANKARA, TURKEY *Email address*: yamanpaks@hotmail.com