

COMBINATORIAL IDENTITIES USING BERNOULLI GRAPHS

• Jacques Bourg

December 9, 2022

ABSTRACT

In here, I present a series of combinatorial equalities derived using a graph based approach. Different nodes in the graphs are visited following probabilistic dynamics of a moving dot. The results are presented in such a way that the generalisation (to more nodes, or dimensions) is straightforward. At an instant m , we 'take a picture' of the system and we compute the probabilities of being at particular positions in space. The sum of all these probabilities is equal to one.


Keywords Combinatorial identities

1 Introduction

There are hundreds of combinatorial identities [Gould, 1972a], [Gould, 1972b], the most known one being the binomial identity. In here we present some identities obtained using the following method: consider a particle displacing itself given simple probabilistic dynamics. On certain node configurations, there are multiple ways to attain a point at a given time point. The sum of the probabilities over space at a given time point must be equal to one.

2 Chain

Lets consider a point that advances from the state c_i to the state c_{i+1} with probability p , and with a waiting time of one unit. This point reaches also the state c_{i+1} with probability $1 - p$, but instantaneously. After N states, the point attains a terminal state.



A trajectory is described as an ordered tuple of probabilities, for instance the tuple $(p, 1 - p, p)$ describes a trajectory which passes through the first node, avoids the second node (2), and attains the third module, all in two time units. A similar trajectory which attains the third node in two time units is $(1 - p, p, p)$. The probability of attaining the third node in two time units is therefore $p \cdot (1 - p) \cdot p + (1 - p) \cdot p \cdot p = p(1 - p) \cdot p = p \binom{2}{1} p(1 - p)$. The probability of having attained a node at position k at time point m is given by:

$$p(x = k, t = m\tau^+) = p \binom{k-1}{m} p^m (1-p)^{k-1-m} \quad m \leq k-1$$

The probability of attaining the terminal node before the time m is:

$$p(x = N, t \leq m\tau^+) = \sum_{w=0}^m \binom{N-1}{w} p^w (1-p)^{N-1-w}$$

The sum of the last two probabilities is equal to one:

$$\sum_{k=m+1}^{N-1} p(x = k, t = m\tau^+) + p(x = N, t \leq m\tau^+) = 1$$

We then deduce the formula:

$$\sum_{k=m+1}^{N-1} \binom{k-1}{m} p^{m+1} (1-p)^{k-1-m} + \sum_{w=0}^m \binom{N-1}{w} p^w (1-p)^{N-1-w} = 1 \quad (1)$$





Figure 1: Probability decomposed as sum of A (first summand) and B (second), as function of time m , for $p = 0.1$ and $p = 0.9$. $N = 28$.

3 Two nodes



$$p(x = k, t = m\tau^+) = \sum_{\substack{2a+b+j=m \\ a,b \in \llbracket 0, m \rrbracket \\ j \in \{0,1\}}} \binom{k-1}{a} \binom{k-1-a}{b} (p_1 p_2^j) (p_1 p_2)^a (p_1 (1-p_2))^b (1-p_1)^{k-a-b-1}$$


$x = N$: terminal node.

$$p(x = N, t \leq m\tau^+) = \sum_{w=0}^m \sum_{\substack{2a+b=w \\ a,b \in \llbracket 0, m \rrbracket}} \binom{N-1}{a} \binom{N-1-a}{b} (p_1 p_2)^a (p_1 (1-p_2))^b (1-p_1)^{N-a-b-1}$$

Therefore:

$$\begin{aligned} & \sum_{k=1+\lfloor \frac{m}{2} \rfloor}^{N-1} p(x = k, t = m\tau^+) + p(x = N, t \leq m\tau^+) = 1 \\ 1 &= \sum_{k=1+\lfloor \frac{m}{2} \rfloor}^{N-1} \sum_{\substack{2a+b+j=m \\ a,b \in \llbracket 0, m \rrbracket \\ j \in \{0,1\}}} \binom{k-1}{a} \binom{k-1-a}{b} (p_1 p_2^j) (p_1 p_2)^a (p_1 (1-p_2))^b (1-p_1)^{k-a-b-1} \\ &+ \sum_{w=0}^m \sum_{\substack{2a+b=w \\ a,b \in \llbracket 0, m \rrbracket}} \binom{N-1}{a} \binom{N-1-a}{b} (p_1 p_2)^a (p_1 (1-p_2))^b (1-p_1)^{N-a-b-1} \end{aligned} \quad (2)$$

4 Three nodes



We give the rule for 3 nodes, which suggests the form of the general formula, for arbitrary number of nodes.

$$\begin{aligned}
 p(x = k, t = m\tau^+) &= \sum_{\substack{3a+2b+c+j+jl=m \\ a,b,c \in \llbracket 0,m \rrbracket \\ j,l \in \{0,1\}}} \binom{k-1}{a} \binom{k-1-a}{b} \binom{k-1-a-b}{c} (p_1 p_2^j p_3^{jl}) (p_1 p_2 p_3)^a (p_1 p_2 (1-p_3))^b \dots \\
 &\quad \dots (p_1 (1-p_2))^c (1-p_1)^{k-a-b-c-1} \\
 p(x = k, t \leq m\tau^+) &= \sum_{w=0}^m \sum_{\substack{3a+2b+c=m \\ a,b,c \in \llbracket 0,m \rrbracket}} \binom{k-1}{a} \binom{k-1-a}{b} \binom{k-1-a-b}{c} (p_1 p_2 p_3)^a (p_1 p_2 (1-p_3))^b \dots \\
 &\quad \dots (p_1 (1-p_2))^c (1-p_1)^{k-a-b-c-1}
 \end{aligned}$$

5 Random walk in two dimensions

Consider a point initially at the origin, doing random jumps on \mathbb{Z}^2 , on the possible directions, with probabilities $p_{1,2,3,4} : p_1 + p_2 + p_3 + p_4 = 1$. The jumps can be of arbitrary size and direction. Call T_i the travelling time ($T_i \in \mathbb{N}^*$). Considering a certain time point m ($m \in \mathbb{N}^*$), and a pair of coordinates $(h, v) \in \mathbb{Z}^2$. Let's study the case: $p_1 : (\rightarrow, J_1, T_1)$ $p_2 : (\leftarrow, J_2, T_2)$ $p_3 : (\uparrow, J_3, T_3)$ $p_4 : (\downarrow, J_4, T_4)$.

Then:

$$\begin{aligned}
 1 &= \sum_{\substack{h \in \llbracket J_2.m, J_1.m \rrbracket \\ v \in \llbracket J_4.m, J_3.m \rrbracket}} p\left(\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} h \\ v \end{pmatrix}, t = m\right) \\
 1 &= \sum_{\substack{h \in \llbracket J_2.m, J_1.m \rrbracket \\ v \in \llbracket J_4.m, J_3.m \rrbracket}} \sum_{\substack{J_1.a + J_2.b = h \\ J_3.c + J_4.d = v \\ a.T_1 + b.T_2 + c.T_3 + d.T_4 = m}} \binom{a+b}{a} \binom{c+d}{c} p_1^a \cdot p_2^b \cdot p_3^c \cdot p_4^d \quad (3)
 \end{aligned}$$

In the second summand, we iterate over all pair of values (a, b, c, d) which are solutions in \mathbb{N}^4 , of this equations system (two spatial equations and one temporal equation).

5.1 Simple random walk in 1D

Starting from the origin, consider a point which moves $+1$ space unit with probability p and -1 with probability $1-p$. The jumps last both one time unit. After m time steps, the point is situated in the interval $[-m, m]$. We will only observe the point position at "even" moments in time ($m = 2n, n \in \mathbb{N}$ even), and therefore we only consider even positions ($k = 2l, l \in \mathbb{Z}$ even). The probability of being in odd positions is nil at even times. We find the combinatorial sum:

$$1 = \sum_{\substack{k \in \llbracket -m, m \rrbracket \\ k \text{ even}}} p(x = k, t = m) \iff 1 = \sum_{\substack{k \in \llbracket -m, m \rrbracket \\ k \text{ even}}} \binom{m}{\frac{m+k}{2}} p^{\frac{m+k}{2}} (1-p)^{\frac{m-k}{2}} \quad (4)$$

5.2 Simple chain with delay and a terminal node

We can consider in 1D a chain in which elements go wright by one spatial unit with probability p (with a waiting time of 1 time unit), and go wright, with jumps of one spatial unit, with a probability of $1-p$, instantaneously. We find back the formula of the chain presented in the second paragraph.

5.3 Simple 2D chain with delay and a terminal barrier

We consider that a point has the following dynamics:

$$p_1(\rightarrow, J_1 = 1, T_1 = 1), p_2(\rightarrow, J_2 = 1, T_2 = 0), p_3(\uparrow, J_3 = 1, T_3 = 1), p_4(\uparrow, J_4 = 1, T_4 = 0).$$


$$p\left(\binom{x}{y} = \binom{h}{v}, t = m\right) = \sum_{\substack{a+c=m \\ a \in \llbracket 0, h \rrbracket \\ c \in \llbracket 0, v \rrbracket}} \binom{h}{a} \binom{v}{c} p_1^a \cdot p_2^{h-a} \cdot p_3^c \cdot p_4^{v-c}$$

Lets consider that there is a barrier \mathcal{B} delimited by the coordinates:

$$\mathcal{B} = \left\{ \binom{N}{v}, v \in \llbracket 0, N-1 \rrbracket \text{ or } \binom{h}{N}, h \in \llbracket 0, N-1 \rrbracket \right\}$$

Therefore, at time m , the probability of being in a certain position (that has not reached the barrier), plus the probability of having reached the barrier is equal to one.

$$\sum_{\substack{\binom{h}{v} \notin \mathcal{B} \\ m \leq h+v}} \sum_{\substack{a+c=m \\ a \in \llbracket 0, h \rrbracket \\ c \in \llbracket 0, v \rrbracket}} \binom{h}{a} \binom{v}{c} p_1^a \cdot p_2^{h-a} \cdot p_3^c \cdot p_4^{v-c} + \sum_{\substack{\binom{h}{v} \in \mathcal{B} \\ m \leq h+v}} \sum_{\substack{a+c=0 \\ a \in \llbracket 0, h \rrbracket \\ c \in \llbracket 0, v \rrbracket}} \binom{h}{a} \binom{v}{c} p_1^a \cdot p_2^{h-a} \cdot p_3^c \cdot p_4^{v-c} = 1 \quad (5)$$



6 Conclusion

The formula deduced in equation (1) has some ressemblance with Gosper's formula [Beeler et al., 1972].

$$1 = \sum_{k=0}^{N-1} \binom{N+k-1}{k} \left(p^N (1-p)^k + (1-p)^N p^k \right)$$

Consider a coin flip: it has a probability p of falling heads, and a probability $1-p$ of being tails. This formula reflects that if the flip the coin until one of its sides comes up N times, the sum of the probability all the probable states \mathcal{S} is equal to one.

$$\mathcal{S} = \{(N, 0), (N, 1), \dots, (N, N-1)\} \cup \{(0, N), (1, N), \dots, (N-1, N)\}$$

In equation (1), however, we "take the picture", not after N heads or tails like in the Gosper's formula, but at an arbitraty time m , ($m \leq N$). After m flips we compute the probabilities of having reached different states. In fact the sum of the exponents in Gopers Formula is equal to $N+k$, $k \in \llbracket 0, N-1 \rrbracket$, whereas in (1) it is equal to k ($k \in \llbracket m+1, N-1 \rrbracket$).

References

Henry Wadsworth Gould. *Combinatorial Identities: A standardized set of tables listing 500 binomial coefficient summations*. Gould, 1972a.

Henry Wadsworth Gould. Gould tables of combinatorial identities, edited by jocelyn quaintance. <https://web.archive.org/web/20190629193344/http://www.math.wvu.edu/~gould/>, 1972b.

Michael Beeler, R William Gosper, and Richard Schroepel. Hakmem. *MIT AI Memo 239*, 1972.