

# Quantum Magnetic Properties, Entanglement for Antiferromagnetic Spin 1 and 3/2 Cluster Models

N. Ananikian\*

*AANL, Alikhanian Br. 2, 0036 Yerevan, Armenia and  
CANDLE, Acharyan 31, 0040 Yerevan, Armenia*

Vl. V. Papoyan†

*BLTP, JINR, 141980 Dubna, Russian Federation and  
Dubna State University, Dubna, Russian Federation*

# Abstract

Entanglement, magnetization and magnetic susceptibility for 1D antiferromagnetic spin 1 and spin  $\frac{3}{2}$  Heisenberg  $XXX$  model with Dzyaloshinskii-Moriya interaction, single-ion anisotropy and external magnetic field on the finite chain are obtained.

## I. INTRODUCTION

Ferromagnetic and antiferromagnetic quantum magnetism has become an extremely active area of research [1, 2]. Experimental data on the supramolecular architecture, synthesis, magnetic susceptibility with dinuclear, tetranuclear, heptanuclear, decanuclear, and polynuclear complexes of  $\text{Cu}_2$  ions in the antiferromagnetic case are found [3]. The preparation of an antiferromagnetic trinuclear  $\text{Ni}_2$  ionic complex with spin 1 and magnetic susceptibility via temperature have been obtained [4].

X-ray crystallographic structures, magnetic susceptibility from 2 to 300K and the theoretical analysis of magnetism for a triangular and a tetranuclear molecule consisting of linked high-spin cobalt (II) centers were described by the inclusion of an antisymmetric exchange term (Dzyaloshinsky-Moriya interaction) [5, 6].

Thermal non-classical correlations quantified by concurrence entanglement, local quantum the uncertainty and quantum coherence in a four-qubit square chain (tetranuclear copper (II) complex) of mixed ferromagnetic-antiferromagnetic interactions between nearest neighboring spins with a magnetic field were examined exactly in [7]. Magnetic susceptibility can reveal spin entanglement between individual constituents of a solid, and magnetization describes their local properties [8].

In this paper, we consider analytically the thermal entanglement, magnetization and magnetic susceptibility of antiferromagnetic high-spin dinuclear and trinuclear  $\text{Ni}_2$  (spin 1) and  $\text{Co}_2$  (spin  $\frac{3}{2}$ ) metal ion complexes as a function of an external magnetic field and single-ion anisotropy at low temperatures.

---

\* ananik@yerphi.am

† vpap@theor.jinr.ru

## II. MODEL AND DEFINITIONS

The Hamiltonian for 1D antiferromagnetic Heisenberg XXX model with Dzyaloshinskii-Moriya (DM) interaction and single-ion anisotropy in external magnetic field given as

$$H = J \sum_{i=1}^N \vec{S}_i \vec{S}_{i+1} + G \sum_{i=1}^N (S_i^x S_{i+1}^y - S_i^y S_{i+1}^x) + D \sum_{i=1}^N (S_i^z)^2 - B \sum_{i=1}^N S_i^z. \quad (1)$$

where  $J = 1$  to fix the energy scale,  $G$  is DM interaction parameter,  $D$  represents uniaxial single-ion anisotropy,  $B$  is the controllable parameter. The local spin vector  $S_i$  for each site has the components of the spin-1 operators:

$$S_x = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad S_y = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -i & 0 \\ i & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad S_z = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

The components of the spin  $\frac{3}{2}$  operators are expressed as follows:

$$S_x = \frac{1}{2} \begin{pmatrix} 0 & \sqrt{3} & 0 & 0 \\ \sqrt{3} & 0 & 2 & 0 \\ 0 & 2 & 0 & \sqrt{3} \\ 0 & 0 & \sqrt{3} & 0 \end{pmatrix}, \quad S_y = \frac{1}{2} \begin{pmatrix} 0 & \sqrt{3} & 0 & 0 \\ \sqrt{3} & 0 & 2 & 0 \\ 0 & 2 & 0 & \sqrt{3} \\ 0 & 0 & \sqrt{3} & 0 \end{pmatrix}, \quad S_z = \frac{1}{2} \begin{pmatrix} 3 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -3 \end{pmatrix}.$$

Logarithmic negativity [9] is an entanglement measure defined as

$$LN = \log_2 (\| \rho_{ab}^{T_a} \|_1) = \log_2 \left( \sum_{k=1} |\lambda_k| \right) \quad (2)$$

where  $\lambda_k$  are the eigenvalues of the matrix  $\rho_{ab}^{T_a}$ .  $T_a$  denotes the partial pairwise transpose with respect to the first system and  $\rho_{ab}$  is partial trace of the density matrix of the whole system between quantum states  $a$  and  $b$ .

## III. CALCULATED RESULTS

The entanglement of the 1D antiferromagnetic spin 1 and spin  $\frac{3}{2}$  Heisenberg XXX model with Dzyaloshinskii-Moriya interaction, single-ion anisotropy and an external magnetic field on finite chain with sizes  $N = 2$ ,  $N = 3$  and  $N = 4$  is studied. The behavior of logarithmic negativity  $LN$  as a function of the external magnetic field parameter  $B$  for spin 1 and spin

$\frac{3}{2}$  is displayed in Fig. 1, for fixed  $J = 1$ ,  $D = 1$  and  $G = 1$ . The critical point  $B_c$  at  $B \geq B_c$  of the ground state, at which entanglement disappears, is shown if  $B > 0$ .

In Fig. 2 we present the behavior of logarithmic negativity  $LN$  via DM interaction parameter  $G$  at fixed  $J = 1$ ,  $D = 1$  for spin 1 at  $B = 5$  and for spin  $\frac{3}{2}$  at  $B = 9$ . We fixed these values of  $B$  near the  $B_c$  for spin 1 and spin  $\frac{3}{2}$  respectively.

In Fig. 1 and Fig. 2 (a, b, c) the upper curves show the logarithmic negativity of  $LN$  for spin  $\frac{3}{2}$ , and the lower curves for spin 1 ( $N = 2, 3, 4$ ). The same in plots (d, e, f) corresponds to a fixed temperature  $k_B T = 0.1$ .




FIG. 1. The logarithmic negativity  $LN$  versus of external magnetic field parameter  $B$

We found out also the phase diagrams of ground-state in the  $D - B$  and the  $G - B$  planes at fixed value  $J = 1$  for  $N = 2$ , see Fig. 3. The entanglement region is located below the linear line and the adjacent curve for the corresponding values of the parameter  $G$  (Fig. 3 (a) and (b)). For example for spin 1 and  $N = 2$  at  $G = 1$  equations of the linear line and the adjacent curve are obtained as follows:

$$B = 2 + 2\sqrt{2} + D \quad \left( \text{for } D > -\frac{2}{7} \left( 2 + 3\sqrt{2} \right) \right), \quad (3)$$

$$B = \frac{1}{2} \left( \sqrt{D^2 - 2D + 17} + D + 3 \right) \quad \left( \text{for } D < -\frac{2}{7} \left( 2 + 3\sqrt{2} \right) \right). \quad (4)$$

The curve connecting the corner points shows the possible values  $B$  and  $D$  of the triple




FIG. 2. The logarithmic negativity  $LN$  versus of DM interaction parameter  $G$

points for various quantities of  $G$  and in the case of spin 1 and  $N = 2$  is defined by:

$$B = 2\sqrt{\frac{1}{2} \left( D^2 + \sqrt{(D+1)^2 (D^2 + 1)} + D - 1 \right)} + 1 + D + 2. \quad (5)$$




FIG. 3. Ground-state phase diagrams in the  $D - B$  plane at fixed value  $J = 1$  and various values of  $G$  (from top to bottom  $G = 1, G = \frac{1}{2}, G = 0$ ), plot (a) corresponds the spin 1, plot (b) corresponds the spin  $\frac{3}{2}$ . Plot (c) corresponds the ground-state phase diagrams in the  $G - B$  plane at fixed values  $J = 1$  and  $D = 1$ . The entanglement area is between the upper and lower curves for the spin 1. The region between middle curves corresponds the entanglement states, for the spin  $\frac{3}{2}$ .

The ground state of the system in the non entanglement region ( $LN = 0$ ) has a wave function corresponding to the state with all spins down ( $\Psi_{GS} = |\downarrow\downarrow \dots \downarrow\rangle$ ). The first nonzero plateau ground state wave function for the spin 1 for chain sizes  $N = 2$  and  $N = 3$

is defined follow:

$$\Psi_{GS}(N = 2) = \frac{1}{\sqrt{2}}(|\downarrow\downarrow 0\rangle + (\cos(\frac{\pi}{4}) + i \cos(\frac{\pi}{4}))|0\downarrow\rangle) \quad (6)$$

$$\Psi_{GS}(N = 3) = \frac{1}{\sqrt{3}}(|0\downarrow\downarrow\rangle + i|\downarrow 0\downarrow\rangle - |\downarrow\downarrow 0\rangle) \quad (7)$$

The coefficients for the spin  $\frac{3}{2}$  are the same with replacements in the labels of vector of state  $\downarrow$  by  $-\frac{3}{2}$  and 0 by  $-\frac{1}{2}$ .

The entanglement values  $LN$  as a function of temperature  $T$  (from now on  $k_B = 1$ ) at fixed  $J = 1$ , DM interaction parameter  $G = 1$  and single-ion anisotropy  $D = 1$  are shown in Fig. 4




FIG. 4. The upper curves represent the behaviour of entanglement amount  $LN$  for the spin  $\frac{3}{2}$ , and lowest curves show the same for the spin 1. Plots (a), (b), (c) correspond  $N = 2$ ,  $N = 3$ ,  $N = 4$ , respectively.

The behavior of magnetization and magnetic susceptibility of the chain size  $N = 2$  are presented in Fig. 5. As it was found, the plateau of magnetization and the plateau of logarithmic negativity coincide. For chains of size  $N = 3$  and  $N = 4$ , we get the results that are qualitatively similar.

N. A. acknowledges the receipt of the grant No. SCS 21AG-1C006.

- 
- [1] T. Nagamiya, K. Yosida, R. Kubo, Antiferromagnetism, *Advances in Physics* 4 (13) (1955) 1-112. DOI.org/10.1080/00018735500101154.
  - [2] I. M. Georgescu, S. Ashhab, and F. Nori, Quantum Simulation, *Rev. Mod. Phys.* 86, 153 (2014). DOI: 10.1103/RevModPhys.86.153.




FIG. 5. The logarithmic negativity (red curve), magnetization (orange curve) and magnetic susceptibility (blue curve) for  $N = 2$  as a function of  $B$  at  $J = 1$ ,  $D = 1$ ,  $G = 1$  and  $k_B T = 0.05$ . Plots (a) and (b) corresponds spin 1 and  $\frac{3}{2}$ , respectively.

- [3] S. S. Tandon, S. D. Bunge, N. Patel, E. C. Wang and L. K. Thompson, Self-Assembly of Antiferromagnetically-Coupled Copper(II) Supramolecular Architectures with Diverse Structural Complexities, *Molecules* 25 (2020) 5549(pp 31). DOI:10.3390/molecules25235549.
- [4] N. M. Bonanno, A. J. Lough, M. T. Lemaire, A trinuclear nickel(II) cluster containing a ditopic redox active ligand: Structural and magnetic properties, *Polyhedron* 183 (2020) 114536 (pp 6). DOI:10.1016/j.poly.2020.114536.
- [5] J. F. Berry, F. A. Cotton, C. Y. Liu, T. Lu, C. A. Murillo, B. S. Tsukerblat, D. Villagran, and X. Wang, Modeling Spin Interactions in a Cyclic Trimer and a Cuboidal Co<sub>4</sub>O<sub>4</sub> Core with Co(II) in Tetrahedral and Octahedral Environments, *J. Am. Chem. Soc.* 127 (13) (2005) 4895–4902. DOI: 10.1021/ja044185b.
- [6] H. Arian Zad, R. Kenna and N. Ananikian, Magnetic and thermodynamic properties of the octanuclear nickel phosphonate-based cage, *Physica A* 538 (2020) 122841. DOI: 10.1016/j.physa.2019.122841.
- [7] F. Benabdallah, S. Haddadi, H. A. Zad, M. R. Pourkarimi, M. Daoud, N. Ananikian Pairwise quantum criteria and teleportation in a spin square complex, *Scientific Reports* 12(1) (2022) 1-12, DOI: 10.1038/s41598-022-10248-2.
- [8] M. Wieniak, V. Vedral and C. Brukner, Magnetic susceptibility as a macroscopic entanglement witness, *New J. Phys.* 7 (2005) 258 (pp 8). DOI: 10.1088/1367-2630/7/1/258.
- [9] G. Vidal, R. F. Werner, A computable measure of entanglement. *Phys. Rev. A.* 65 (3) (2002) 032314 (pp11). DOI: 10.1103/PhysRevA.65.032314.