Nonradial stability of expanding Goldreich-Weber stars

Mahir Hadžić, Juhi Jang, King Ming Lam [‡]

Abstract

Goldreich-Weber solutions constitute a finite-parameter of expanding and collapsing solutions to the mass-critical Euler-Poisson system. Two subclasses of this family correspond to compactly supported density profiles suitably modulated by the dynamic radius of the star that expands at the self-similar rate $\lambda(t)_{t\to\infty} \sim t^{\frac{2}{3}}$ and linear rate $\lambda(t)_{t\to\infty} \sim t$ respectively. We prove two results: any linearly expanding Goldreich-Weber star is nonlinearly stable, while any given self-similarly expanding Goldreich-Weber star is codimension-4 nonlinearly stable against irrotational perturbations.

The codimension-4 condition in the latter result is optimal and reflects the presence of 4 unstable directions in the linearised dynamics in self-similar coordinates, which are induced by the conservation of the energy and the momentum. This result can be viewed as a codimension-1 nonlinear stability of the moduli space of self-similarly expanding Goldreich-Weber stars against irrotational perturbations.

Contents

1	Intr	duction	2
	1.1	The Euler-Poisson system	2
	1.2	Goldreich-Weber stars	4
		1.2.1 Self-similarly expanding Goldreich-Weber stars	5
		1.2.2 Linearly expanding Goldreich-Weber stars	5
	1.3	Main results and review	6
2	Nota	ion and preliminary lemmas	7
	2.1	Essential lemmas and definitions	9
		2.1.1 Basic bounds on the gravity term \mathbf{G}	9
		2.1.2 Basic bounds on the pressure term P	12
3	Non	adial stability of self-similarly expanding Goldreich-Weber stars	15
	3.1	Formulation and statement of the result	15
		3.1.1 Equation in self-similar coordinates	15
		3.1.2 Total energy and momentum	16
		3.1.3 High-order energies and the main theorem	17
		3.1.4 Proof strategy	20
	3.2	Linearisation and coercivity	22
		3.2.1 The linear and non-linear part of Euler-Poisson system	22
		3.2.2 Coercivity of \mathbf{L}	25
	3.3	Momentum and energy	34
	3.4	Coercivity via irrotationality	37
		3.4.1 Lagrangian description of irrotationality	37
		3.4.2 Coercivity of \mathbf{L}	40
	3.5	Reduction to linear problem	42
		3.5.1 Estimating the non-linear part of the pressure term	42

^{*}Department of Mathematics, University College London, London WC1E 6XA, UK. Email: m.hadzic@ucl.ac.uk.

[†]Department of Mathematics, University of Southern California, Los Angeles, CA 90089, USA, and Korea Institute for Advanced Study, Seoul, Korea. Email: juhijang@usc.edu.

[‡]Department of Mathematics, University College London, London WC1E 6XA, UK. Email: king.lam.19@ucl.ac.uk. Delft Institute of Applied Mathematics, Delft University of Technology, 2628 CD Delft, Netherlands. Email: K.M.Lam@tudelft.nl.

		3.5.2 Estimating the linear and non-linear part of the gravity term	5
		3.5.3 Reduction to linear problem	9
	3.6	Energy estimates and proof of the main theorem	1
		3.6.1 Near boundary energy estimate	1
		3.6.2 Near origin energy estimate	2
		3.6.3 Bootstrapping scheme and final theorem	7
4	Non	radial stability of linearly expanding Goldreich-Weber stars 5	8
	4.1	Formulation and statement of the result	8
		4.1.1 Equation in linearly-expanding coordinates	8
		4.1.2 High-order energies and the main theorem	0
		4.1.3 Proof strategy	1
	4.2	Pressure estimates	1
	4.3	Gravity estimates	2
	4.4	Vorticity estimates	5
	4.5	Energy estimates and proof of the main theorem	7
A	Арр	endix 6	9
	A.1	Differentiation and commutation properties	9
	A.2	Spherical harmonics	1
	A.3	Hardy-Poincaré inequality and embeddings	3
		A.3.1 Embedding theorems for self-similarly expanding GW stars 73	5
		A.3.2 Embedding theorems for linearly expanding GW stars	5

1 Introduction

1.1 The Euler-Poisson system

We consider a fundamental model of a self-gravitating compressible fluid, given by the Euler-Poisson system. The unknowns are the fluid density $\rho \ge 0$, the velocity vector **u**, the fluid pressure $p \ge 0$, and the gravitational potential ϕ . They solve the system

$$\partial_t \rho + \nabla \cdot (\rho \mathbf{u}) = 0, \qquad \text{in } \Omega(t), \qquad (1.1)$$

$$\rho \left(\partial_t + \mathbf{u} \cdot \nabla\right) \mathbf{u} + \nabla p + \rho \nabla \phi = \mathbf{0}, \qquad \text{in } \Omega(t), \qquad (1.2)$$

$$\Delta \phi = 4\pi\rho, \qquad \text{in } \mathbb{R}^3. \tag{1.3}$$

Here the pressure p satisfies the mass-critical polytropic equation of state

$$p = \rho^{\frac{4}{3}},\tag{1.4}$$

and the star is isolated, which translates into the asymptotic boundary condition for the gravitational potential:

$$\lim_{|\mathbf{x}| \to \infty} \phi(t, \mathbf{x}) = 0.$$
(1.5)

We refer to the system (1.1)–(1.5) as the (EP)_{$\frac{4}{3}$}-system. Moreover $\Omega(t) := \{\mathbf{x} \mid \rho(t, \mathbf{x}) > 0\}$ is the interior of the support of star density. Note that in (1.3) the density is trivially extended by 0 to the complement of $\Omega(t)$. Since we allow the boundary to move, we must complement equations (1.1)–(1.5) with suitable boundary conditions at the vacuum free boundary $\partial \Omega(t)$. We assume the classical kinematic boundary condition

$$\mathcal{V}_{\partial\Omega(t)} = \mathbf{u} \cdot \mathbf{n} \qquad \text{on } \partial\Omega(t), \tag{1.6}$$

which states that the normal velocity of the boundary $\mathcal{V}_{\partial\Omega(t)}$ equals the normal component of the velocity vector field; here **n** is the outward pointing unit normal to $\partial\Omega(t)$. It is well-known that the presence of the

vacuum boundary complicates the local-in-time well-posedness problem, as the acoustic cones degenerate as the speed of sound c_s defined through

$$c_s^2 := \frac{dp}{d\rho} = \frac{4}{3}\rho^{\frac{1}{3}},$$

becomes 0 at the vacuum boundary. The resolution comes by imposing a condition on initial data, that specifies the rate of decay of the initial density to 0 as we approach the vacuum boundary. This condition is known as the physical vacuum condition and reads

$$\nabla(c_s^2) \cdot \mathbf{n} \Big|_{\partial\Omega_0} < 0. \tag{1.7}$$

The Euler-Poisson system (1.1)-(1.5) possesses the following important conserved quantities – mass, momentum and energy, given respectively by:

$$M[\rho] := \int_{\mathbb{R}^3} \rho \, \mathrm{d}\mathbf{x},\tag{1.8}$$

$$\mathbf{W}[\rho, \mathbf{u}] := \int_{\mathbb{R}^3} \rho \mathbf{u} \, \mathrm{d}\mathbf{x},\tag{1.9}$$

$$E[\rho, \mathbf{u}] := \int_{\mathbb{R}^3} \left(\frac{1}{2} \rho |\mathbf{u}|^2 + 3\rho^{\frac{4}{3}} + \frac{1}{2} \rho \phi \right) d\mathbf{x}.$$
 (1.10)

The mass-criticality associated with the polytropic index $\frac{4}{3}$ in (1.4) is simply a statement that the natural self-similar rescaling of the problem also preserves the total mass $M[\rho]$. Namely, for any $\lambda > 0$ and $\mathbf{x}_0 \in \mathbb{R}^3$ one can check that if (ρ, \mathbf{u}) is a classical solution of the (EP) $\frac{4}{2}$ -system, then $(\tilde{\rho}, \tilde{\mathbf{u}})$ defined by

$$\rho(t, \mathbf{x}) = \lambda^{-3} \tilde{\rho}\left(\frac{t}{\lambda^{\frac{3}{2}}}, \frac{\mathbf{x} - \mathbf{x}_0}{\lambda}\right)$$
(1.11)

$$\mathbf{u}(t,\mathbf{x}) = \lambda^{\frac{1}{2}} \tilde{\mathbf{u}}\left(\frac{t}{\lambda^{\frac{3}{2}}}, \frac{\mathbf{x} - \mathbf{x}_0}{\lambda}\right)$$
(1.12)

is also a solution to the $(EP)_{\frac{4}{2}}$ -system as functions of the rescaled variables (s, \mathbf{y}) :

$$s = \frac{t}{\lambda^{\frac{3}{2}}}, \qquad \mathbf{y} = \frac{\mathbf{x} - \mathbf{x}_0}{\lambda}$$

Relation (1.11) readily implies that the total mass is conserved under this change of variables.

We shall mostly work in the Lagrangian coordinates in this article, as they are particularly well suited to the analysis of fluids featuring a vacuum boundary. Let $\eta(t, \mathbf{x})$ be the fluid flow map, defined through

$$\partial_t \boldsymbol{\eta} = \mathbf{u} \circ \boldsymbol{\eta}$$
 with $\boldsymbol{\eta}(0, \mathbf{x}) = \boldsymbol{\eta}_0(\mathbf{x}),$

where $\mathbf{u} \circ \boldsymbol{\eta}(t, \mathbf{x}) = \mathbf{u}(t, \boldsymbol{\eta}(t, \mathbf{x}))$. The spatial domain is then fixed for all time as $\Omega_0 := \boldsymbol{\eta}_0^{-1}(\Omega(0))$. To reformulate the (EP) $\frac{4}{3}$ -system in the new variables, we introduce

$\mathbf{v} = \mathbf{u} \circ \boldsymbol{\eta}$	(Lagrangian velocity)
$f = ho \circ \boldsymbol{\eta}$	(Lagrangian density)
$\psi = \phi \circ oldsymbol{\eta}$	(Lagrangian potential)
$A = (\nabla \eta)^{-1}$	(inverse of the deformation tensor)
$J = \det(\nabla \boldsymbol{\eta})$	(Jacobian determinant)
a = JA	(cofactor matrix of the deformation tensor)

Under this change of coordinates, the continuity equation becomes $fJ = f_0 J_0$ and the momentum equation (1.2) in the domain Ω_0 reads

$$\partial_t \mathbf{v} + \frac{1}{w^3} \partial_k (A^k_{\bullet} (w^4 J^{-\frac{1}{3}}) + A \nabla \psi = \mathbf{0}, \qquad (1.13)$$

where $A\nabla := A^k \partial_k$ and we have introduced the enthalpy w

$$w := (f_0 J_0)^{\frac{1}{3}}.$$
(1.14)

Moreover, ψ solves the Poisson equation

$$(A\nabla) \cdot (A\nabla)\psi = 4\pi f_0 J_0 J^{-1}. \tag{1.15}$$

For details of the Lagrangian description of the Euler-Poisson system, we refer to [26].

1.2 Goldreich-Weber stars

The mass-criticality of the problem allows for the existence of a special class of expanding solutions, known as the Goldreich-Weber stars [16]. The reason such solutions exist is, roughly speaking, because the scaling properties of the Euler-Poisson system in the mass critical case allows us to scale solutions while maintain the overall mass. This suggests that natural solutions that evolve in time under this scaling exist (note that solutions must conserve overall mass in time). For reader's convenience we provide a brief summary of this special class of solutions of (1.1)–(1.5) which has been analysed in [16, 40, 15, 11]. A comprehensive overview can be found in [24].

We let

$$\mathscr{K} := 4\pi \Delta^{-1} \tag{1.16}$$

i.e. $\mathcal{K}f(\mathbf{x}) = -\int_{\mathbb{R}^3} \frac{f(\mathbf{y})}{|\mathbf{x}-\mathbf{y}|} d\mathbf{y}$ for any $f \in L^2(\mathbb{R}^3)$. Observe that

$$\begin{split} \psi(\mathbf{x}) &= (\mathscr{K}\rho)(\boldsymbol{\eta}(\mathbf{x})) = -\int \frac{\rho(\mathbf{y})}{|\boldsymbol{\eta}(\mathbf{x}) - \mathbf{y}|} \mathrm{d}\mathbf{y} = -\int \frac{f(\mathbf{z})J(\mathbf{z})}{|\boldsymbol{\eta}(\mathbf{x}) - \boldsymbol{\eta}(\mathbf{z})|} \mathrm{d}\mathbf{z} = -\int \frac{f_0(\mathbf{z})J_0(\mathbf{z})}{|\boldsymbol{\eta}(\mathbf{x}) - \boldsymbol{\eta}(\mathbf{z})|} \mathrm{d}\mathbf{z} \\ &= -\frac{1}{\lambda}\int \frac{f_0(\mathbf{z})}{|\mathbf{x} - \mathbf{z}|} \mathrm{d}\mathbf{z} = \frac{1}{\lambda} \mathscr{K}(f_0)(\mathbf{x}). \end{split}$$

We look for spherically symmetric solutions to the Euler-Poisson system of the form $\eta(t, \mathbf{x}) = \lambda(t)\mathbf{x}$, and assume without loss of generality that $\lambda(0) = 1$. Under this affine ansatz, the momentum equation (1.13) reduces to

$$\ddot{\lambda}\lambda^2 \mathbf{x} + \frac{1}{f_0}\nabla(f_0^{\frac{4}{3}}) + \nabla \mathscr{K}f_0 = \mathbf{0}$$

Assuming spherically symmetry we get

$$\ddot{\lambda}\lambda^2 + \frac{1}{rf_0}\partial_r(f_0^{\frac{4}{3}}) + \frac{1}{r}\partial_r\mathcal{K}f_0 = 0$$

Since we can separate variables above, we look for a $\delta \in \mathbb{R}$ and a δ -dependent solution $(\lambda, f_0) = (\lambda_{\delta}, f_0^{\delta})$ so that

$$\ddot{\lambda}(t)\lambda(t)^2 = \delta, \tag{1.17}$$

$$\frac{4}{r}\partial_r\bar{w}_\delta + \frac{1}{r}\partial_r\mathcal{K}(\bar{w}^3_\delta) = -\delta, \qquad (1.18)$$

where \bar{w}_{δ} is the enthalpy associated with f_0^{δ} satisfying

$$(\bar{w}_{\delta})^3 := f_0^{\delta}.$$
 (1.19)

We also equip (1.17) with initial data

$$\lambda(0) = 1, \qquad \dot{\lambda}(0) = \lambda_1 \in \mathbb{R}.$$
(1.20)

It can be shown that there exists a negative constant $\tilde{\delta} < 0$ such that the solution $(\lambda_{\delta}(t), \bar{w}_{\delta})$ to (1.17)–(1.20) exists for all $\delta \geq \tilde{\delta}$, see [15, 24], whereby $\lambda_{\delta}(\cdot)$ either blows up in finite positive time, or exists globally for all $t \geq 0$. Moreover, for any such $\delta \geq \tilde{\delta}$, the enthalpy profile \bar{w}_{δ} is compactly supported, has finite total mass, and by adapting the value $\bar{w}_{\delta}(0)$ it can be normalised to be supported on the interval $r \in [0, R]$ for a fixed R > 0. At the vacuum boundary, by analogy to the classical Lane-Emden stars [24], the Goldreich-Weber star satisfies the so-called physical vacuum condition, which in this context reads

$$\left. \bar{w}_{\delta}'(r) \right|_{r=R} < 0. \tag{1.21}$$

1.2.1 Self-similarly expanding Goldreich-Weber stars

The self-similarly expanding Goldreich-Weber stars are the subclass of solutions to (1.17)–(1.20) of total energy 0, for which $\lambda_{\delta}(\cdot)$ exists for all $t \ge 0$. Since the total conserved energy of the above affine motion is easily seen to be

$$E_{\delta}(t) = \left(\lambda_1^2 + 2\delta\right) \int 2\pi f_0^{\delta} z^4 \,\mathrm{d}z,\tag{1.22}$$

solutions with vanishing energy necessitate $\delta < 0$. For any such $\tilde{\delta} \leq \delta < 0$, equation (1.17) with (1.20) is explicitly solvable with

$$\lambda_{\delta}(t) = \left(1 + \frac{3}{2}\lambda_1 t\right)^{2/3}, \qquad \lambda_1^2 = -2\delta.$$
(1.23)

In particular, for any $\lambda_1 > 0$ we obtain an expanding solution with the explicit rate of expansion $\lambda_{\delta}(t) \sim_{t \to \infty} t^{\frac{2}{3}}$. This is the self-similarly expanding Goldreich-Weber solution.

Definition 1.1 (Self-similarly expanding Goldreich-Weber solutions). To any $\delta \in [\overline{\delta}, 0)$ we associate the Goldreich-Weber (GW) star which constitutes a solution of the mass-critical free-boundary Euler-Poisson system (1.1)–(1.5):

$$\bar{\rho}(t,\mathbf{x}) = \lambda_{\delta}(t)^{-3}\bar{w}_{\delta}^{3}\left(\frac{|\mathbf{x}|}{\lambda_{\delta}(t)}\right), \qquad \bar{\mathbf{u}}(t,\mathbf{x}) = \frac{\lambda_{\delta}(t)}{\lambda_{\delta}(t)}\mathbf{x}, \qquad \bar{\Omega}(t) = B_{\lambda_{\delta}(t)}(\mathbf{0}), \qquad (1.24)$$

with $\lambda_{\delta}(t)$ given by (1.23) with $\lambda_1 > 0$ and \bar{w}_{δ} the normalised solution to (1.18) as above.

These solutions are spherical symmetric about the origin, have zero momentum $\mathbf{W}[\bar{\rho}, \bar{\mathbf{u}}] = 0$ and zero energy $E[\bar{\rho}, \bar{\mathbf{u}}] = 0$. Without loss of generality, this can be assumed by setting our frame of reference.

Remark 1.2. The Galilean invariance of the Euler-Poisson system (1.1)–(1.5) implies the conservation of momentum. If we change our frame of reference, we can obtain an enlarged family of the GW-solutions with arbitrary momentum $\bar{\mathbf{W}} \in \mathbb{R}^3$. More precisely, for any motion $\mathbf{p}(t) = \mathbf{p}_0 + t\mathbf{p}_1$ we can obtain a new solution via

$$\bar{\rho}_{\mathbf{p}}(t, \mathbf{x}) = \bar{\rho}(t, \mathbf{x} - \mathbf{p}(t)),$$
$$\bar{\mathbf{u}}_{\mathbf{p}}(t, \mathbf{x}) = \bar{\mathbf{u}}(t, \mathbf{x} - \mathbf{p}(t)) + \mathbf{p}_{1},$$

or equivalently $\eta_{\mathbf{p}}(t, \mathbf{x}) = \eta(t, \mathbf{x}) + \mathbf{p}(t)$ in Lagrangian coordinates. It is easy to verify that $(\bar{\rho}_{\mathbf{p}}, \bar{\mathbf{u}}_{\mathbf{p}})$ so obtained solves the Euler-Poisson system with the total momentum $\mathbf{W}[\bar{\rho}_{\mathbf{p}}, \bar{\mathbf{u}}_{\mathbf{p}}] = M[\bar{\rho}, \bar{\mathbf{u}}]\mathbf{p}_1$ and energy $E[\bar{\rho}_{\mathbf{p}}, \bar{\mathbf{u}}_{\mathbf{p}}] = \frac{1}{2}M[\bar{\rho}, \bar{\mathbf{u}}]|\mathbf{p}_1|^2$. The freedom to choose $\mathbf{p}_1 \in \mathbb{R}^3$ thus parametrises the three degrees of freedom associated with the total linear momentum, and this will play a role in our analysis.

1.2.2 Linearly expanding Goldreich-Weber stars

In the case

$$\delta > 0$$
 or $\delta = 0$ with $\lambda_1 > 0$ or $\delta \in (\delta, 0)$ with $\lambda_1 > \sqrt{2|\delta|}$, (1.25)

the solution $\lambda_{\delta}(\cdot)$ exists for all $t \ge 0$ and expands indefinitely at a linear rate, i.e. there exists a constant c > 0 such that

$$\lim_{t \to \infty} \dot{\lambda}(t) = c$$

These solutions have strictly positive energy

$$E_{\delta,\lambda_1}(t) = \left(\lambda_1^2 + 2\delta\right) \int 2\pi f_0^{\delta} z^4 \, \mathrm{d}z > 0.$$
 (1.26)

We refer to such solutions of (1.17)–(1.20) as the *linearly expanding Goldreich-Weber* solutions.

Definition 1.3 (Linearly expanding Goldreich-Weber solutions). To any δ , λ_1 satisfying (1.25) we associate the Goldreich-Weber (GW) star which constitutes a solution of the mass-critical free-boundary Euler-Poisson system (1.1)–(1.5):

$$\bar{\rho}(t,\mathbf{x}) = \lambda_{\delta,\lambda_1}(t)^{-3} \bar{w}_{\delta}^3 \left(\frac{|\mathbf{x}|}{\lambda_{\delta,\lambda_1}(t)}\right), \quad \bar{\mathbf{u}}(t,\mathbf{x}) = \frac{\bar{\lambda}_{\delta,\lambda_1}(t)}{\lambda_{\delta,\lambda_1}(t)} \mathbf{x}, \quad \bar{\Omega}(t) = B_{\lambda_{\delta,\lambda_1}(t)}(\mathbf{0}), \quad (1.27)$$

with $\lambda_{\delta}(t)$ the solution to (1.17), (1.20) and \bar{w}_{δ} the normalised solution to (1.18) as above.

Unless stated otherwise, we shall drop the subscript δ in the definition of the GW-solution, as this will create no confusion in the analysis.

1.3 Main results and review

The two results to be presented in this paper are a generalisation of nonlinear stability of GW-stars against radial perturbations shown in [24] by the first two authors. We prove nonlinear stability against non-radial perturbations. In Section 3 and 4 respectively, we will prove the following two theorems.

Theorem 1.4 (Informal statement). *The class of self-similarly expanding Goldreich-Weber stars is codimension 1 non-linearly stable under irrotational perturbations.*

Theorem 1.5 (Informal statement). *The linearly expanding Goldreich-Weber stars are non-linearly stable (against general perturbations).*

The precise statements will be provided in Section 3 and 4 respectively. More precisely, Theorem 1.4 corresponds to Theorem 3.7 and Corollary 3.8, while Theorem 1.5 corresponds to Theorem 4.2.

First discovered class of nontrivial global solutions to the Euler-Poisson system are the classical Lane-Emden (LE) stars [5]. Their linear stability is well-known to depend on the size of the polytropic exponent in the general pressure law $p = \rho^{\gamma}$, $\frac{6}{5} \leq \gamma < 2$. Very few rigorous nonlinear results are available on the dynamics in the vicinity of LE-steady states, we refer to [44] for some rigorous statements about the stability in the subcritical range $\frac{4}{3} < \gamma < 2$ and to [29, 30] for rigorous nonlinear instability analysis in the supercritical range $\frac{6}{5} \leq \gamma < \frac{4}{3}$. In the context of nonradial stability, recent works [32, 35, 34] treat this question from the Lagrangian and the Eulerian perspective respectively. In the critical case $\gamma = \frac{4}{3}$, the very existence of the GW-stars shows that the LE-steady states are embedded in a larger family of collapsing/expanding solutions, and are therefore unstable. Our main result can be viewed as a definitive nonradial instability statement about the mass-critical LE-solutions, improving upon the radial nonlinear stability shown by the first two authors [24]. We emphasise that in the presence of viscosity, the parabolic effect takes over and various asymptotic stability results are available [37, 38]. We also mention recent works [7, 6] on global existence result with radial symmetry in the class of weak solutions and conditional behaviour of strong solutions.

The driving stabilisation mechanism that allows for the global existence in Theorem 3.7 is the expansion of the support of the background GW-star. Intuitively expansion translates into dispersion, since the total mass is preserved. When there is no vacuum boundary present, the dispersion induced by the expansion was used by Grassin [17], Serre [50], and Rozanova [48] to give examples of global-in-time solutions to the compressible Euler flows. We also mention here that there has been a recent surge of activity on the problem of existence of collapsing self-gravitating flows, which are characterised by the finite-time implosion of the fluid density. We refer the reader to [19, 20, 21, 22, 1, 49] and for a discussion of the various features of the collapsing and expanding stellar dynamics, we refer to the overview paper [23].

The GW-stars belong to a class of so-called affine motions. In the context of compressible flows the notion of an affine motion goes back to the works of Ovsiannikov [41] and Dyson [13]. In the presence of vacuum, Sideris [52] showed the existence of a finite-parameter family of compactly supported expanding affine flows, whose nonlinear stability was shown by Hadžić and Jang [25] and Shkoller and Sideris [51] for the pure Euler flows. For expanding profiles with small initial densities, but not necessarily close to the Sideris solutions, see [43]. Further results in this direction, in the nonisentropic setting and in the presence of heat convection can be found in [45, 46, 47]. A similar method works for the Euler-Poisson system and global-in-time flows were shown to exist in both the gravitational and electrostatic case [26], where the Euler part of the flow entirely dominates the gravitational/electrostatic response of the model.

Another application of an expansion-induced stabilisation is the work of Parmeshwar [42] where an *N*-body configuration of expanding stars is shown to exist globally in-time. If damping is present in Euler flows it can drive sublinear expansion of Barenblatt-like solutions, see [39, 53, 54].

Our result concerning the self-similarly expanding GW stars in Theorem 1.4 has one notable difference to the above results. The stability of linearly expanding GW stars in Theorem 1.5 is in fact easier than that for self-similarly expanding GW stars in Theorem 1.4. The reason is that the linearly expanding GW stars expand faster than the self-similarly expanding ones and thus the effects of dispersion-via-expansion are stronger in the proof of Theorem 1.5. One important consequence is that the gravitational forces in the linearly expanding case are of subleading order and from the analysis it is apparent that the results do not depend on the attractive/repulsive nature of the force field. By contrast, our result on the self-similarly expanding GW stars in Theorem 1.4 are profoundly sensitive to the attractive nature of the gravitational force and require more sophisticated estimates. This is particularly felt in the linearised stability analysis in Section 3.2.

The plan of the paper is as follows. In Section 2 we introduce important notational conventions and some key objects that will play a role throughout the paper. We introduce the gravity and the pressure operators **G** and **P** expressed in Lagrangian coordinates and include several preparatory lemmas. In Section 3 we provide a precise formulation of Theorem 1.4 and provide its proof. One of the main difficulties is to obtain coercivity of the associated linearised operator, see Section 3.2. In Section 4 we provide a rigorous formulation and proof of Theorem 1.5. Finally, in Appendix A we provide an overview of several technical tools used throughout the paper, including various properties of the spherical harmonics, as well as some weighted Poincaré inequalities.

Acknowledgments. M. Hadžić's research is supported by the EPSRC Early Career Fellowship EP/S02218X/1. J. Jang's research is supported in part by the NSF grants DMS-2009458 and DMS-2306910. K.-M. Lam was supported by the EPSRC studentship grant EP/R513143/1 when undertaking this research; now supported by the NWO grant OCENW.M20.194.

2 Notation and preliminary lemmas

As Section 3 and 4 are devoted to the self-similarly expanding GW stars and linearly expanding GW stars respectively, we will write \bar{w} to denote the the self-similarly expanding GW stars and linearly expanding GW stars enthalpy profile respectively (see Definition 1.1 and 1.3) in these sections.

Since the gaseous Euler-Poisson system is degenerate near the vacuum boundary, we will need to make use of weighted Sobolev spaces. Let $L^2(B_R, w)$ denote the L^2 space on B_R weighted by a non-negative weight w. Of crucial importance in this paper are the weighted inner products

$$\langle g,h\rangle_k := \int_{B_R} gh\bar{w}^k \mathrm{d}\mathbf{x},$$
(2.28)

$$\langle \mathbf{g}, \mathbf{h} \rangle_k := \int_{B_R} \mathbf{g} \cdot \mathbf{h} \bar{w}^k \mathrm{d} \mathbf{x},$$
 (2.29)

defined for any scalar fields $g, h \in L^2(B_R, \bar{w}^k)$ and vector fields $\mathbf{g}, \mathbf{h} \in L^2(B_R, \bar{w}^k)^3$. The weighted inner product for tensor fields are defined in the same way. The associated norm is then given by

$$||f||_{k}^{2} = \int_{B_{R}} |f(\mathbf{x})|^{2} \bar{w}(\mathbf{x})^{k} \mathrm{d}\mathbf{x}.$$
(2.30)

To capture the structure of the roughly spherical stars, we will need to use the following specially defined radial and tangential derivatives in our analysis. We define

$$X_r := x^i \partial_i = r \partial_r \tag{2.31}$$

$$\partial_i := \epsilon_{ijk} x^j \partial_k \tag{2.32}$$

$$\partial_{ij} := x^i \partial_j - x^j \partial_i \tag{2.33}$$

where ϵ_{ijk} is the alternating symbol (see Definition 2.1). Note that $\partial_{ij} = \epsilon_{ijk} \partial_k$. We denote

$$\begin{aligned} \operatorname{div} \boldsymbol{\theta} &:= \nabla \cdot \boldsymbol{\theta} \\ [\operatorname{curl} \boldsymbol{\theta}]_l^k &:= \partial_l \theta^k - \partial_k \theta^l \\ \operatorname{div}_{\mathscr{A}} \boldsymbol{\theta} &:= (\mathscr{A} \nabla) \cdot \boldsymbol{\theta} \\ [\operatorname{curl}_{\mathscr{A}} \boldsymbol{\theta}]_l^k &:= \mathscr{A} \partial_l \theta^k - \mathscr{A} \partial_k \theta^l \end{aligned}$$

where $\mathscr{A}\nabla := \mathscr{A}^k \partial_k$ and $\mathscr{A}\partial_i := \mathscr{A}^k_i \partial_k$.

Let $(\bar{\rho}, \bar{\mathbf{u}})$ be a given self-similarly or linearly expanding GW-flow from Definition 1.1 and 1.3 with the corresponding radius $R\lambda(t)$ and the associated enthalpy $\bar{w} : [0, R] \to \mathbb{R}_+$. In order to study the stability of the flow, we will follow the strategy introduced in [24, 25] and renormalise the equation by introducing a new unknown

$$\boldsymbol{\xi}(t, \mathbf{x}) = \frac{\boldsymbol{\eta}(t, \mathbf{x})}{\lambda(t)}.$$
(2.34)

We suitably renormalise the inverse of the Jacobian gradient and the Jacobian determinant, so that

$$\begin{split} \mathcal{A} &:= (\nabla \boldsymbol{\xi})^{-1} &= \lambda A \\ \mathcal{J} &:= \det(\nabla \boldsymbol{\xi}) &= \lambda^{-3} J \\ \boldsymbol{a} &:= \mathcal{J} \mathcal{A} &= \lambda^{-2} a \\ \Phi &:= -\int \frac{f_0(\mathbf{z}) \mathcal{J}_0(\mathbf{z})}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} d\mathbf{z} = \lambda \psi \end{split}$$

We will work mainly with the perturbation variable defined by

$$\boldsymbol{\theta}(\mathbf{x}) := \boldsymbol{\xi}(\mathbf{x}) - \mathbf{x},\tag{2.35}$$

which measures the deviation of the nonlinear flow to the background GW profile.

As will see later, in these new variable, the pressure and gravity term in the Euler-Poisson system take the following form

$$\mathbf{P} := \bar{w}^{-3} \partial_k (\bar{w}^4 (\mathscr{A}^k_{\bullet} \mathcal{J}^{-1/3} - I^k_{\bullet})), \qquad (2.36)$$

$$\mathbf{G} := \mathscr{A} \nabla \Phi - \mathscr{K} \nabla \bar{w}^3 \tag{2.37}$$

In the rest of the paper we will use some fairly standard notations which we collect here for reader's convenience.

Definition 2.1 (Standard notations).

- *i.* Greek letter superscript on derivatives are multi-index notation for derivatives. For example, $\partial^{\alpha} = \partial_1^{\alpha_1} \partial_2^{\alpha_2} \partial_3^{\alpha_3}$ where $\alpha = (\alpha_1, \alpha_2, \alpha_3)$. And $|\alpha| = \alpha_1 + \alpha_2 + \alpha_3$.
- ii. Roman letter indices such as i, j, k, l, m on derivatives and vector or tensor fields are assumed to range over $\{1, 2, 3\}$. However, this does not apply to s which we reserved to denote the rescaled time variable. Also, it does not apply when they are indices of non-vector or non-tensor objects, for example Ψ_{lm} and Λ_{lm} in Section 3.
- iii. The Einstein summation convention will be used, i.e. repeated indices on derivatives and vector or tensor fields are summed over. For example, $\partial_i \theta^i = \sum_{i=1}^3 \partial_i \theta^i$. However, this does not apply to non-vector or non-tensor objects, for example Ψ_{lm} and Λ_{lm} in Section 3.
- iv. I denotes the identity matrix, δ_{ij} or δ_j^i the Kronecker delta, ϵ_{ijk} the alternating symbol (Levi-Civita symbol).
- v. C will denote generic "analyst's constant", whose exact value can change from line to line and term to term. When appearing in equalities, it can potentially denote any real constant, but when appearing in inequalities, it is generally assumed to be non-negative. We will use subscript to emphasise its dependence on certain variables, for example C_{δ} is a constant that depends on δ .

vi. \mathbf{e}_i (i = 1, 2, 3) denotes the standard basis of \mathbb{R}^3 , while \mathbf{e}_r denotes the radial unit vector $\mathbf{x}/|\mathbf{x}|$.

Now we will define some important special new notations that the reader probably will not have seen before.

Definition 2.2 (Special notations).

- *i.* We will denote ∂_{\bullet} as a generic derivative, so it can be any of ∂_s , ∂_i , ∂ or X_r .
- ii. We will use to denote an unspecified index, or to emphasise the vectorial/tensorial nature of nonscalar quantities. For example if A is a matrix, we can write A^k_{\bullet} .
- iii. When the exact value/ordering of the indices is not important, we shall often write $\langle \star \rangle$ for a generic term that looks like \star to avoid invoking indices. For example, $\langle C \mathfrak{A} \nabla \theta \rangle$ could represent a term like $C\mathcal{A}_{i}^{i}\partial_{k}\theta^{l}$ for some i, j, k, l and constant $C \in \mathbb{R}$.
- iv. We will write $\Re[\star]$ to denote terms that can be bounded by \star , e.g. $|\Re[S_n E_n]| \leq S_n E_n$.
- *v.* We will write $\mathbf{1}_{\star}$ to denote the usual indicator function, and write $\mathbf{1}[\star]$ to denote the Iverson bracket. For example, $\mathbf{1}_A(\mathbf{x}) = \mathbf{1}[\mathbf{x} \in A]$.

2.1 Essential lemmas and definitions

We next state and prove some essential preparatory lemmas and definitions that will be used throughout the paper. The following Hodge-type estimate allows us to estimate the norm of the gradient of a quantity in terms of the div and curl of the quantity term and a lower order term.

Lemma 2.3 (Hodge-type bound). Let $k \ge 0$. For any θ we have

$$\|\nabla \boldsymbol{\theta}\|_{k+2}^2 \lesssim \|\nabla \cdot \boldsymbol{\theta}\|_{k+2}^2 + \|\nabla \times \boldsymbol{\theta}\|_{k+2}^2 + \|\boldsymbol{\theta}\|_k^2$$

Proof. We have

$$\begin{split} \int |\nabla \boldsymbol{\theta}|^2 \bar{w}^{k+2} \mathrm{d}\mathbf{x} &= \int (\partial_j \theta^i) (\partial_j \theta^i) \bar{w}^{k+2} \mathrm{d}\mathbf{x} = \int (\partial_j \theta^i) (\partial_i \theta^j + [\operatorname{curl} \boldsymbol{\theta}]_j^i) \bar{w}^{k+2} \mathrm{d}\mathbf{x} \\ &= \int (\partial_j \theta^i) (\partial_i \theta^j) \bar{w}^{k+2} \mathrm{d}\mathbf{x} + \int (\partial_j \theta^i) [\operatorname{curl} \boldsymbol{\theta}]_j^i \bar{w}^{k+2} \mathrm{d}\mathbf{x} \\ &= -\int (\partial_i \partial_j \theta^i) (\theta^j) \bar{w}^{k+2} \mathrm{d}\mathbf{x} + \int (\partial_j \theta^i) [\operatorname{curl} \boldsymbol{\theta}]_j^i \bar{w}^{k+2} \mathrm{d}\mathbf{x} \\ &- (k+2) \int (\partial_j \theta^j) (\theta^j) \bar{w}^{k+1} \partial_i \bar{w} \mathrm{d}\mathbf{x} \\ &= \int (\partial_i \theta^i) (\partial_j \theta^j) \bar{w}^{k+2} \mathrm{d}\mathbf{x} + \int (\partial_j \theta^i) [\operatorname{curl} \boldsymbol{\theta}]_j^i \bar{w}^{k+2} \mathrm{d}\mathbf{x} \\ &+ (k+2) \left(\int (\partial_i \theta^i) (\theta^j) \bar{w}^{k+1} \partial_j \bar{w} \mathrm{d}\mathbf{x} - \int (\partial_j \theta^i) (\theta^j) \bar{w}^{k+1} \partial_i \bar{w} \mathrm{d}\mathbf{x} \right) \\ &\lesssim \delta' \int |\nabla \boldsymbol{\theta}|^2 \bar{w}^{k+2} \mathrm{d}\mathbf{x} + \frac{1}{\delta'} \left(\int (|\nabla \cdot \boldsymbol{\theta}|^2 + |\nabla \times \boldsymbol{\theta}|^2) \bar{w}^{k+2} \mathrm{d}\mathbf{x} + \int |\boldsymbol{\theta}|^2 \bar{w}^k \mathrm{d}\mathbf{x} \right). \end{split}$$

Picking δ' small enough, we are done.

2.1.1 Basic bounds on the gravity term G

We recall here the definition (2.37) of G. The following lemma is a structural identity that allows us to estimate the gravitational term G more conveniently.

Lemma 2.4. Recall the definition (2.37) of the gravitational term **G**. We then have the identity:

$$\mathbf{G} = \mathcal{K}_{\boldsymbol{\xi}} \nabla \cdot (\mathcal{A}_{\bullet} \bar{w}^3) - \mathcal{K} \nabla \bar{w}^3 \tag{2.38}$$

$$= \mathcal{K}_{\boldsymbol{\xi}}((\mathcal{A}-I)\nabla\bar{w}^3 - \bar{w}^3\mathcal{A}_m^i\mathcal{A}_{\boldsymbol{\bullet}}^l\partial_l\partial_l\theta^m) + (\mathcal{K}_{\boldsymbol{\xi}} - \mathcal{K})\nabla\bar{w}^3,$$
(2.39)

where

$$(\mathscr{K}_{\boldsymbol{\xi}}g)(\mathbf{x}) := -\int \frac{g(\mathbf{z})}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} d\mathbf{z}$$
(2.40)

Proof. Note that formally

$$(\nabla \mathcal{K} \rho)(\mathbf{x}) = -\int \nabla_{\mathbf{x}} \frac{\rho(\mathbf{z})}{|\mathbf{x} - \mathbf{z}|} d\mathbf{z} = \int \nabla_{\mathbf{z}} \frac{\rho(\mathbf{z})}{|\mathbf{x} - \mathbf{z}|} d\mathbf{z} = -\int \frac{\nabla \rho(\mathbf{z})}{|\mathbf{x} - \mathbf{z}|} d\mathbf{z} = (\mathcal{K} \nabla \rho)(\mathbf{x})$$

and so

$$\begin{split} A\nabla\psi(\mathbf{x}) &= (\nabla\phi)(\boldsymbol{\eta}(\mathbf{x})) = (\nabla\mathcal{H}\rho)(\boldsymbol{\eta}(\mathbf{x})) = (\mathcal{H}\nabla\rho)(\boldsymbol{\eta}(\mathbf{x})) = -\int \frac{\nabla\rho(\mathbf{y})}{|\boldsymbol{\eta}(\mathbf{x}) - \mathbf{y}|} d\mathbf{x} \\ &= -\int \frac{A\nabla f(\mathbf{z})J(\mathbf{z})}{|\boldsymbol{\eta}(\mathbf{x}) - \boldsymbol{\eta}(\mathbf{z})|} d\mathbf{z} = -\int \frac{a\nabla(fJJ^{-1})(\mathbf{z})}{|\boldsymbol{\eta}(\mathbf{x}) - \boldsymbol{\eta}(\mathbf{z})|} d\mathbf{z} = -\int \frac{a\nabla(\bar{w}^3J^{-1})(\mathbf{z})}{|\boldsymbol{\eta}(\mathbf{x}) - \boldsymbol{\eta}(\mathbf{z})|} d\mathbf{z} \\ &= -\int \frac{\nabla \cdot (a\bar{w}^3J^{-1})(\mathbf{z})}{|\boldsymbol{\eta}(\mathbf{x}) - \boldsymbol{\eta}(\mathbf{z})|} d\mathbf{z} = -\int \frac{\nabla \cdot (A\bar{w}^3)(\mathbf{z})}{|\boldsymbol{\eta}(\mathbf{x}) - \boldsymbol{\eta}(\mathbf{z})|} d\mathbf{z} = \frac{1}{\lambda^2} (\mathcal{H}_{\boldsymbol{\xi}} \nabla \cdot (\mathcal{A}\bar{w}^3))(\mathbf{x}) d\mathbf{z}$$

where we denote $\nabla \cdot M = \partial_i M^i$ for a matrix M and recall (2.40). We then have

$$\mathscr{A}\nabla\Phi = \lambda^2 A \nabla\psi(\mathbf{x}) = \mathscr{K}_{\boldsymbol{\xi}}\nabla\cdot(\mathscr{A}\bar{w}^3).$$

Now we have

$$\mathbf{G} = \mathscr{K}_{\boldsymbol{\xi}} \nabla \cdot (\mathscr{A}_{\bullet} \bar{w}^3) - \mathscr{K} \nabla \bar{w}^3 = \mathscr{K}_{\boldsymbol{\xi}} (\nabla \cdot (\mathscr{A} \bar{w}^3) - \nabla \bar{w}^3) + (\mathscr{K}_{\boldsymbol{\xi}} - \mathscr{K}) \nabla \bar{w}^3$$
$$= \mathscr{K}_{\boldsymbol{\xi}} ((\mathscr{A} - I) \nabla \bar{w}^3 - \bar{w}^3 \mathscr{A}^i_m \mathscr{A}^l_{\bullet} \partial_i \partial_l \theta^m) + (\mathscr{K}_{\boldsymbol{\xi}} - \mathscr{K}) \nabla \bar{w}^3.$$

Since the gravity term is a non-local term, we need to estimate a convolution-like operator. However, rather than the convolution kernel $|\mathbf{x} - \mathbf{z}|^{-1}$ we actually need to estimate $|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|^{-1}$. The next lemma tells us how to reduce the latter to the former, which will allow us to estimate using the Young convolution inequality.

Lemma 2.5. Let $\boldsymbol{\xi}$ be as in (2.35). For any $\mathbf{x}, \mathbf{z} \in B_R$ we have

$$|\mathbf{x} - \mathbf{z}| \le \|\mathscr{A}\|_{L^{\infty}(B_R)} |\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|$$

 $|\partial_s^a \partial_{\mathbf{x}}^{\beta} \boldsymbol{\xi}(\mathbf{x}) - \partial_s^a \partial_{\mathbf{z}}^{\beta} \boldsymbol{\xi}(\mathbf{z})| \le \|\nabla \partial_s^a \partial_{\mathbf{x}}^{\beta} \boldsymbol{\xi}\|_{L^{\infty}(B_R)} |\mathbf{x} - \mathbf{z}|$

Proof. Using the mean value inequality we have

$$\begin{aligned} |\mathbf{x} - \mathbf{z}| &= |\boldsymbol{\xi}^{-1} \boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}^{-1} \boldsymbol{\xi}(\mathbf{z})| \\ &\leq \|\nabla \boldsymbol{\xi}^{-1}\|_{L^{\infty}(B_R)} |\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})| = \|\mathscr{A}\|_{L^{\infty}(B_R)} |\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})| \end{aligned}$$

and

$$|\partial_s^{a_i}\partial_{\mathbf{x}}^{\beta_i}\boldsymbol{\xi}(\mathbf{x}) - \partial_s^{a_i}\partial_{\mathbf{z}}^{\beta_i}\boldsymbol{\xi}(\mathbf{z})| \le \|\nabla\partial_s^{a_i}\partial^{\beta_i}\boldsymbol{\xi}\|_{L^{\infty}(B_R)}|\mathbf{x} - \mathbf{z}|.$$

Since we cannot commute extra weights into the non-local gravity term, the radial derivatives, which affect the powers in the weight, need to be estimated differently to avoid possible loss of regularity via unfavourable weights. Using methods from [26], the following two lemmas provide the way to do this. More precisely, the radial derivative can be estimated with curl, divergence and tangential derivatives. And this is useful because the curl and divergence of the gravity term consist only of local or non-linear terms, which we can estimate.

Lemma 2.6. For any vector field $\tilde{\mathbf{G}} \in H^1_{loc}$

$$|X_r \tilde{\mathbf{G}}|^2 \lesssim |r \nabla \cdot \tilde{\mathbf{G}}|^2 + |r \nabla \times \tilde{\mathbf{G}}|^2 + \sum_{k=1}^3 |\partial_k \tilde{\mathbf{G}}|^2,$$

where we recall the notation (2.31)-(2.33).

Proof. Note that

$$\begin{aligned} |\mathbf{x} \cdot \tilde{\mathbf{G}}|^2 &= (x^i \tilde{G}^i)(x^j \tilde{G}^j) = (x^j \tilde{G}^i)(x^i \tilde{G}^j) = (x^j \tilde{G}^i)(x^j \tilde{G}^i) + (x^j \tilde{G}^i)(x^i \tilde{G}^j - x^j \tilde{G}^i) \\ &= |\mathbf{x}|^2 |\tilde{\mathbf{G}}|^2 - \frac{1}{2} (x^i \tilde{G}^j - x^j \tilde{G}^i)(x^i \tilde{G}^j - x^j \tilde{G}^i) = r^2 |\tilde{\mathbf{G}}|^2 - |\mathbf{x} \times \tilde{\mathbf{G}}|^2 \end{aligned}$$

We have by definition

$$\partial_i = \frac{x^j}{r^2} \partial_{ji} + \frac{x^i}{r^2} X_r, \ i = 1, 2, 3.$$

The divergence and the curl of $\tilde{\mathbf{G}}$ can be written as

$$r^2 \nabla \cdot \tilde{\mathbf{G}} = x^j \partial_{ji} \tilde{\mathbf{G}}^i + \mathbf{x} \cdot X_r \tilde{\mathbf{G}}$$
 and $r^2 \nabla \times \tilde{\mathbf{G}} = x^j \partial_{j\bullet} \times \tilde{\mathbf{G}} + \mathbf{x} \times X_r \tilde{\mathbf{G}}$

We then obtain

$$r^{2}|X_{r}\tilde{\mathbf{G}}|^{2} = |r^{2}\nabla \cdot \tilde{\mathbf{G}} - x^{j}\partial_{ji}G^{i}|^{2} + |r^{2}\nabla \times \tilde{\mathbf{G}} - x^{j}\partial_{j\bullet} \times \tilde{\mathbf{G}}|^{2}$$

from which we deduce the result.

Lemma 2.7 (The div-curl structure of the gravitational term). Let G be as in (2.38). We have

$$\nabla \cdot \mathbf{G} = (I - \mathcal{A}) \nabla \cdot \mathbf{G} + (I - \mathcal{A}) \nabla \cdot \nabla \mathcal{K} \bar{w}^3 + 4\pi \bar{w}^3 (\mathcal{J}^{-1} - 1)$$
$$\nabla \times \mathbf{G} = (I - \mathcal{A}) \nabla \times \mathbf{G} + (I - \mathcal{A}) \nabla \times \nabla \mathcal{K} \bar{w}^3.$$

Proof. By definition $\mathbf{G} = \mathscr{A} \nabla \Phi - \nabla \mathscr{K} \bar{w}^3$, so

$$\mathscr{A}\nabla\cdot\mathbf{G} = (\mathscr{A}\nabla)\cdot(\mathscr{A}\nabla)\Phi - \mathscr{A}\nabla\cdot\nabla\mathscr{K}\bar{w}^3 = (\mathscr{A}\nabla)\cdot(\mathscr{A}\nabla)\Phi + (I-\mathscr{A})\nabla\cdot\nabla\mathscr{K}\bar{w}^3 - 4\pi\bar{w}^3$$

And we have

$$(\mathscr{A}\nabla) \cdot (\mathscr{A}\nabla)\Phi(\mathbf{x}) = \lambda^3 (A\nabla) \cdot (A\nabla)\psi(\mathbf{x}) = \lambda^3 (\nabla \cdot \nabla \mathscr{K}\rho)(\boldsymbol{\eta}(\mathbf{x}))$$
$$= \lambda^3 4\pi\rho(\boldsymbol{\eta}(\mathbf{x})) = \lambda^3 4\pi f(\mathbf{x}) = \lambda^3 4\pi \bar{w}^3 J^{-1} = 4\pi \bar{w}^3 \mathscr{J}^{-1}.$$
(2.41)

So we get the first formula. Proof for the second formula is similar but we use $(\mathscr{A}\nabla) \times (\mathscr{A}\nabla) = 0$.

The next lemma lets us deal with time and tangential derivatives on non-local terms and its kernel, as we will need when dealing with the gravity term.

Lemma 2.8. (i) For any $K: B_R \times B_R \to \mathbb{R}$ sufficiently smooth and $g \in H^1_0(B_R)$ we have

$$\partial_{i,\mathbf{x}} \int_{B_R} K(\mathbf{x}, \mathbf{z}) g(\mathbf{z}) d\mathbf{z} = \int_{B_R} \left(g(\mathbf{z}) (\partial_{i,\mathbf{x}} + \partial_{i,\mathbf{z}}) K(\mathbf{x}, \mathbf{z}) + K(\mathbf{x}, \mathbf{z}) \partial_{i,\mathbf{z}} g(\mathbf{z}) \right) d\mathbf{z},$$

where we recall the notation (2.31)-(2.33).

(ii) For any $\boldsymbol{\theta}: B_R \to \mathbb{R}^3$ sufficiently smooth and $\mathbf{x}, \mathbf{y} \in B_R$ we have

$$|\partial_s^a \partial^\beta \theta(\mathbf{x}) - \partial_s^a \partial^\beta \theta(\mathbf{z})| \le \|\nabla \partial_s^a \partial^\beta \theta\|_{L^\infty(B_R)} |\mathbf{x} - \mathbf{z}|$$
(2.42)

$$|\partial^{\beta} \mathbf{x} - \partial^{\beta} \mathbf{z}| \le |\mathbf{x} - \mathbf{z}| \tag{2.43}$$

Proof. For part (i), integrate by parts to get

$$\int_{B_R} K(\mathbf{x}, \mathbf{z}) \partial_{i, \mathbf{z}} g(\mathbf{z}) d\mathbf{z} = - \int_{B_R} g(\mathbf{z}) \partial_{i, \mathbf{z}} K(\mathbf{x}, \mathbf{z}) d\mathbf{z}.$$

For part (ii), use the mean value inequality to get (2.42). Bound (2.43) follows from $\partial_{ij}x^k = x^i \delta_j^k - x^j \delta_i^k$.

2.1.2 Basic bounds on the pressure term P

In order to apply the high order energy method, will need to estimate derivatives of the pressure term $\partial_s^a X_r^b \partial^\beta \mathbf{P}$ and $\partial^\gamma \mathbf{P}$. Recall that $\mathbf{P} := \bar{w}^{-3} \partial_k (\bar{w}^4 (\mathscr{A}^k \mathcal{F}^{-1/3} - I^k))$ by (2.36) and therefore we will need to compute the commutators between the operator $\partial_s^a X_r^b \partial^\beta$ and ∂^γ and the weighted derivative $\bar{w}^{-3} \partial_k (\bar{w}^4 \cdot)$. Lemma 2.9 deals with the case when no radial derivatives are present, while Lemma 2.10 includes the radial derivatives. Lemmas 2.9 and 2.10 are necessary to control all the non-"top-order" contributions coming from $\partial_s^a X_r^b \partial^\beta \mathbf{P}$ and $\partial^\gamma \mathbf{P}$ by our energy norms.

Lemma 2.9. For any tensor field T_i^k sufficiently smooth, we have

$$\partial^{\beta} \left(\bar{w}^{-3} \partial_k (\bar{w}^4 T_i^k) \right) = \bar{w}^{-3} \partial_k (\bar{w}^4 \partial^{\beta} T_i^k) + \sum_{|\beta'| \le |\beta| - 1} \langle C \bar{w}^{-3} \nabla (\bar{w}^4 \partial^{\beta'} T) \rangle$$
(2.44)

for i = 1, 2, 3, where we recall notations defined in Definition 2.2.

Proof. We will prove this by induction. Assume this is true for β , then we have

$$\begin{split} \partial_{j}\partial^{\beta}\left(\bar{w}^{-3}\partial_{k}(\bar{w}^{4}T_{i}^{k})\right) &= \bar{w}^{-3}\partial_{k}(\bar{w}^{4}\partial_{j}\partial^{\beta}T_{i}^{k}) - \bar{w}^{-3}\epsilon_{jkl}\partial_{l}(\bar{w}^{4}\partial^{\beta}T_{i}^{k}) + \partial_{j}\sum_{|\beta'| \leq |\beta|-1} \langle C\bar{w}^{-3}\nabla(\bar{w}^{4}\partial^{\beta'}T)\rangle \\ &= \bar{w}^{-3}\partial_{k}(\bar{w}^{4}\partial_{j}\partial^{\beta}T_{i}^{k}) + \sum_{|\beta'| \leq |\beta|} \langle C\bar{w}^{-3}\nabla(\bar{w}^{4}\partial^{\beta'}T)\rangle. \end{split}$$

where we used the commutation relation for $[\partial_j, \partial_k]$ from Lemma A.2.

The use of radial derivatives naturally changes the weighting structure, which is one of the key observations that makes the high-order energy argument possible and goes back to [31].

Lemma 2.10. For any tensor field T_i^k sufficiently smooth, we have

$$\begin{split} X_r \left(\bar{w}^{-c} \partial_k (\bar{w}^{1+c} T_i^k) \right) &= \bar{w}^{-(1+c)} \partial_k (\bar{w}^{2+c} X_r T_i^k) \\ &+ (1+c) (T_i^k X_r \partial_k \bar{w}) + (\partial_k \bar{w}) \partial_{kj} T_i^j - \bar{w} \partial_k T_i^k \\ \partial_j \left(\bar{w}^{-c} \partial_k (\bar{w}^{1+c} T_i^k) \right) &= \bar{w}^{-c} \partial_k (\bar{w}^{1+c} \partial_j T_i^k) \\ &- \epsilon_{jkl} ((1+c) (\partial_l \bar{w}) T_i^k + \bar{w} \partial_l T_i^k)) \\ \partial_j \left(\bar{w}^{-c} \partial_k (\bar{w}^{1+c} T_i^k) \right) &= \bar{w}^{-(2+c)} \partial_k (\bar{w}^{3+c} \partial_j T_i^k) \\ &+ (1+c) (T_i^k \partial_j \partial_k \bar{w}) + (\partial_j \bar{w}) \partial_k T_i^k - 2(\partial_j T_i^k) \partial_k \bar{w} \\ X_r^d \partial^\beta \left(\bar{w}^{-3} \partial_k (\bar{w}^4 T_i^k) \right) &= \bar{w}^{-(3+d)} \partial_k (\bar{w}^{4+d} X_r^d \partial^\beta T_i^k) \\ &+ \left(\sum_{\substack{d' \leq d \\ |\beta'| \leq |\beta| - 1}} + \sum_{\substack{d' \leq d-1 \\ |\beta'| \leq |\beta| + 1}} \right) \langle C \omega X_r^d \partial^{\beta'} \nabla T \rangle \\ &+ \left(\sum_{\substack{d' \leq d - 1 \\ |\beta'| \leq |\beta| - 1}} + \sum_{\substack{d' \leq d - 1 \\ |\beta'| \leq |\beta|} \right) \langle C \bar{w} X_r^d \partial^{\beta'} \nabla T \rangle \\ &+ \left(\sum_{\substack{d' \leq d - 1 \\ |\beta'| \leq |\beta| - 1}} + \sum_{\substack{d' \leq d - 1 \\ |\beta'| \leq |\beta|} \right) \langle C \bar{w} X_r^d \partial^{\beta'} \nabla T \rangle \\ &+ \left(\sum_{\substack{d' \leq d - 1 \\ |\beta'| \leq |\beta| - 1}} + \sum_{\substack{d' \leq d - 1 \\ |\beta'| \leq |\beta|} \right) \langle C \bar{w} X_r^d \partial^{\beta'} \nabla T \rangle \\ &+ \left(\sum_{\substack{d' \leq d - 1 \\ |\beta'| \leq |\beta| - 1}} + \sum_{\substack{d' \leq d - 1 \\ |\beta'| \leq |\beta|} \right) \langle C \bar{w} X_r^d \partial^{\beta'} \nabla T \rangle \\ &+ \sum_{|\gamma'| \leq |\gamma| - 1} \left(\langle C \omega \partial^{\gamma'} T \rangle + \langle C \omega \partial^{\gamma'} \nabla T \rangle \right) \end{split}$$

for any $c \ge 0$ and i = 1, 2, 3, where ω denotes some derivatives of \overline{w} . Here we used notations defined in Definition 2.2.

Proof. First note that

$$x^{i}X_{r}T^{i} = x^{i}x^{j}\partial_{j}T^{i} = r^{2}\partial_{i}T^{i} + x^{j}(x^{i}\partial_{j} - x^{j}\partial_{i})T^{i} = r^{2}\partial_{i}T^{i} - x^{i}\partial_{ij}T^{j}.$$

Using this we have

$$\begin{split} X_r\left(\bar{w}^{-c}\partial_k(\bar{w}^{1+c}T_i^k)\right) &= X_r\left((1+c)T_i^k\partial_k\bar{w} + \bar{w}\partial_kT_i^k\right) \\ &= (1+c)\left((X_rT_i^k)\partial_k\bar{w} + T_i^kX_r\partial_k\bar{w}\right) + (\mathbf{x}\cdot\nabla\bar{w})\partial_kT_i^k + \bar{w}X_r\partial_kT_i^k \\ &= (1+c)\left((X_rT_i^k)\partial_k\bar{w} + T_i^kX_r\partial_k\bar{w}\right) + (\mathbf{x}\cdot\nabla\bar{w})r^{-2}(x^kX_rT_i^k + x^k\partial_{kj}T_i^j) \\ &\quad + \bar{w}\partial_kX_rT_i^k - \bar{w}\partial_kT_i^k \\ &= (1+c)\left((X_rT_i^k)\partial_k\bar{w} + T_i^kX_r\partial_k\bar{w}\right) + (\partial_k\bar{w})(X_rT_i^k + \partial_{kj}T_i^j) \\ &\quad + \bar{w}\partial_kX_rT_i^k - \bar{w}\partial_kT_i^k \\ &= \bar{w}^{-(1+c)}\partial_k(\bar{w}^{2+c}X_rT_i^k) + (1+c)(T_i^kX_r\partial_k\bar{w}) + (\partial_k\bar{w})\partial_{kj}T_i^j - \bar{w}\partial_kT_i^k \\ \partial_j\left(\bar{w}^{-c}\partial_k(\bar{w}^{1+c}T_i^k)\right) &= \bar{w}^{-c}\partial_k(\bar{w}^{1+c}\partial_jT_i^k) - \epsilon_{jkl}\bar{w}^{-c}\partial_l(\bar{w}^{1+c}T_i^k) \\ \partial_j\left(\bar{w}^{-c}\partial_k(\bar{w}^{1+c}T_i^k)\right) &= \partial_j\left((1+c)T_i^k\partial_k\bar{w} + \bar{w}\partial_kT_i^k\right) \\ &= (1+c)\left((\partial_jT_i^k)\partial_k\bar{w} + T_i^k\partial_j\partial_k\bar{w}\right) + (\partial_j\bar{w})\partial_kT_i^k + \bar{w}\partial_j\partial_kT_i^k \\ &= \bar{w}^{-(2+c)}\partial_k(\bar{w}^{3+c}\partial_jT_i^k) + (1+c)(T_i^k\partial_j\partial_k\bar{w}) + (\partial_j\bar{w})\partial_kT_i^k - 2(\partial_jT_i^k)\partial_k\bar{w} \end{split}$$

where we used commutation relations from Lemma A.2. The final two formulas can be proven by induction. $\hfill \Box$

The next lemma deals with the terms we get when we apply $\partial_s^a X_r^b \partial^\beta$ or ∂^γ to $\mathscr{A} \mathcal{J}^{-1/3} - I$. Lemma 2.11. Let

$$T := \mathscr{A}\mathcal{J}^{-1/3} - I. \tag{2.46}$$

Recall notations defined in Definition 2.2. For $a \leq 0$ *and* $|\gamma| > 0$ *, we have*

$$\partial_{\bullet}T = T_T[\partial_{\bullet}\nabla\theta],\tag{2.47}$$

$$\partial_s^a X_r^d \partial^\beta T = T_T [\partial_s^a X_r^b \partial^\beta \nabla \theta] + T_{R:a,\beta,d}$$
(2.48)

$$\partial^{\gamma}T = T_T[\partial^{\gamma}\nabla\theta] + T_{R:\gamma}.$$
(2.49)

where

$$T_T[M]^k := -\mathcal{J}^{-1/3} \left(\mathscr{A}_m^k \mathscr{A}^l + \frac{1}{3} \mathscr{A}^k \mathscr{A}_m^l \right) M_l^m, \qquad k = 1, 2, 3$$
(2.50)

$$T_{R:\gamma} := \mathcal{J}^{-1/3} \sum_{\substack{c=2\\ \sum_{i=1}^{c} \gamma_i = \gamma\\ |\gamma_i| > 0}}^{|\gamma|} \langle C \rangle \langle \mathcal{A} \rangle^{1+c} \prod_{i=1}^{c} \langle \partial^{\gamma_i} \nabla \boldsymbol{\theta} \rangle.$$
(2.52)

We write $T_{R:\beta,d} := T_{R:0,\beta,d}$.

Proof. Applying Lemma A.1 we get that

$$\begin{split} \partial_{\bullet}(\mathscr{A}^{k}\mathscr{F}^{-1/3}-I^{k}) &= -\mathscr{F}^{-1/3}\mathscr{A}_{m}^{k}\mathscr{A}^{l}\partial\partial_{l}\theta^{m} - \frac{1}{3}\mathscr{F}^{-1/3}\mathscr{A}^{k}\mathscr{A}_{m}^{l}\partial_{\bullet}\partial_{l}\theta^{m} \\ &= -\mathscr{F}^{-1/3}\left(\mathscr{A}_{m}^{k}\mathscr{A}^{l} + \frac{1}{3}\mathscr{A}^{k}\mathscr{A}_{m}^{l}\right)\partial_{\bullet}\partial_{l}\theta^{m}. \end{split}$$

Hence $\partial_{\bullet} T^k = T_T [\partial_{\bullet} \nabla \theta]^k$. By repeated application of this we get the next two formulas.

We have from (3.62)

$$\partial_s^a X_r^b \partial^\beta \mathbf{P} = \partial_s^a X_r^b \partial^\beta \left(\bar{w}^{-3} \partial_k (\bar{w}^4 T^k) \right)$$

One can see from the last few lemmas (we will prove this properly in Section 3.5.1 and 4.2) that the leading order term is in fact $\mathbf{P}_d \partial_s^a X_r^b \partial^\beta \boldsymbol{\theta}$, where

$$\mathbf{P}_{d}\boldsymbol{\theta} := \bar{w}^{-3-d}\partial_{k}(\bar{w}^{4+d}T_{T}[\nabla\boldsymbol{\theta}]^{k})$$
(2.53)

$$= -\bar{w}^{-(3+d)}\partial_k \left(\bar{w}^{4+d} \left(\mathscr{A}_m^k \mathscr{A}^l + \frac{1}{3} \mathscr{A}^k \mathscr{A}_m^l \right) \partial_l \theta^m \right)$$
(2.54)

Let $\mathbf{P}_{d,L}$ be the linear part of \mathbf{P}_d , i.e.

$$\mathbf{P}_{d,L}\boldsymbol{\theta} := -\bar{w}^{-(3+d)}\partial_k \left(\bar{w}^{4+d} \left(I_m^k I^l + \frac{1}{3} I^k I_m^l \right) \partial_l \boldsymbol{\theta}^m \right) \\ = -\frac{1}{3\bar{w}^{3+d}} \nabla (\bar{w}^{4+d} \nabla \cdot \boldsymbol{\theta}) - \frac{1}{\bar{w}^{3+d}} \partial_k (\bar{w}^{4+d} \nabla \boldsymbol{\theta}^k)$$
(2.55)

In doing energy estimates, the term $\langle \partial_s^a X_r^b \partial^\beta \mathbf{P}, \partial_s^a X_r^b \partial^\beta \theta \rangle$ and $\langle \partial_s^a X_r^b \partial^\beta \mathbf{P}, \partial_s^{a+1} X_r^b \partial^\beta \theta \rangle$ will arise. Using the lemmas in this subsection, we will show in Section 3.5.1 and 4.2 that \mathbf{P} here can be reduced to $\mathbf{P}_{d,L}$ modulo remainder terms that can be estimated. The following identity is needed for that purpose.

Lemma 2.12. For any vector field θ_1, θ_2 sufficiently smooth we have

$$\langle \mathbf{P}_{d}\boldsymbol{\theta}_{1},\boldsymbol{\theta}_{2}\rangle_{3+d}$$

$$= \int \left((\mathscr{A}\partial_{m}\boldsymbol{\theta}_{1}) \cdot (\mathscr{A}\partial_{m}\boldsymbol{\theta}_{2}) + \frac{1}{3} (\operatorname{div}_{\mathscr{A}}\boldsymbol{\theta}_{1}) (\operatorname{div}_{\mathscr{A}}\boldsymbol{\theta}_{2}) - \frac{1}{2} [\operatorname{curl}_{\mathscr{A}}\boldsymbol{\theta}_{1}]_{j}^{m} [\operatorname{curl}_{\mathscr{A}}\boldsymbol{\theta}_{2}]_{j}^{m} \right) \mathcal{F}^{-1/3} \bar{w}^{4+d} \mathbf{dx} \quad (2.56)$$

$$\langle \mathbf{P}_{d,L}\boldsymbol{\theta}_{1},\boldsymbol{\theta}_{2}\rangle_{3+d}$$

$$= \int \left((\partial_{m}\boldsymbol{\theta}_{1}) \cdot (\partial_{m}\boldsymbol{\theta}_{2}) + \frac{1}{3} (\operatorname{div}\boldsymbol{\theta}_{1}) (\operatorname{div}\boldsymbol{\theta}_{2}) - \frac{1}{2} [\operatorname{curl}\boldsymbol{\theta}_{1}]_{j}^{m} [\operatorname{curl}\boldsymbol{\theta}_{2}]_{j}^{m} \right) \bar{w}^{4+d} \mathbf{dx} \quad (2.57)$$

Proof. We have

$$\langle \mathbf{P}_{d}\boldsymbol{\theta}_{1},\boldsymbol{\theta}_{2}\rangle_{3+d} = \int \left((\mathcal{A}\partial_{l}\theta_{1}^{m}) \cdot (\mathcal{A}\partial_{m}\theta_{2}^{l}) + \frac{1}{3} (\operatorname{div}_{\mathscr{A}}\boldsymbol{\theta}_{1}) (\operatorname{div}_{\mathscr{A}}\boldsymbol{\theta}_{2}) \right) \mathcal{J}^{-1/3} \bar{w}^{4+d} \mathrm{d}\mathbf{x}.$$

We are done for \mathbf{P}_d noting that

$$[\operatorname{curl}_{\mathscr{A}} \boldsymbol{\theta}_{1}]_{j}^{m} [\operatorname{curl}_{\mathscr{A}} \boldsymbol{\theta}_{2}]_{j}^{m} = (\mathscr{A}\partial_{j}\theta_{1}^{m} - \mathscr{A}\partial_{m}\theta_{1}^{j})(\mathscr{A}\partial_{j}\theta_{2}^{m} - \mathscr{A}\partial_{m}\theta_{2}^{j})$$
$$= 2(\mathscr{A}\partial_{j}\theta_{1}^{m})(\mathscr{A}\partial_{j}\theta_{2}^{m}) - 2(\mathscr{A}\partial_{j}\theta_{1}^{m})(\mathscr{A}\partial_{m}\theta_{2}^{j}).$$

Similarly for $\mathbf{P}_{d,L}$.

3 Nonradial stability of self-similarly expanding Goldreich-Weber stars

3.1 Formulation and statement of the result

3.1.1 Equation in self-similar coordinates

In this section we will take the enthalpy \bar{w} to be the profile associated with the self-similarly expanding GW star from Definition 1.1. To study the stability of self-similarly expanding GW stars, we want to write our variables as a perturbation from the model GW star. To that end we will use the rescaled variable ξ (equation (2.34)) introduced in Section 2 adapted to the expanding background profile and also write the problem in self-similar time variables. We introduce the self-similar time coordinate *s* adapted to the expanding profile via

$$\frac{ds}{dt} = \lambda(t)^{-\frac{3}{2}}$$

We then have the following change of coordinate formula $\partial_t = \lambda^{-3/2} \partial_s$. The condition $\ddot{\lambda} \lambda^2 = \delta$ (1.17) becomes

$$\delta = \lambda^{1/2} \partial_s (\lambda^{-3/2} \partial_s \lambda) = \frac{\partial_s^2 \lambda}{\lambda} - \frac{3}{2} \frac{(\partial_s \lambda)^2}{\lambda^2} = \partial_s \left(\frac{\partial_s \lambda}{\lambda}\right) - \frac{1}{2} \frac{(\partial_s \lambda)^2}{\lambda^2} = -\frac{1}{2} \theta^2$$
(3.58)

where

Then the Euler-Poisson equations (1.13) becomes

$$\mathbf{0} = \partial_t \mathbf{v} + (f_0 J_0)^{-1} \partial_k (A^k (f_0 J_0)^{4/3} J^{-1/3}) + A \nabla \psi$$

= $\lambda^{-3/2} \partial_s (\lambda^{-3/2} \partial_s (\lambda \boldsymbol{\xi})) + \lambda^{-2} (f_0 J_0)^{-1} \partial_k (\mathcal{A}^k (f_0 J_0)^{4/3} \mathcal{J}^{-1/3}) + \lambda^{-2} \mathcal{A} \nabla \Phi$

Times the equation by λ^2 we get

$$\begin{aligned} \mathbf{0} &= \lambda^{1/2} \partial_s (\lambda^{-3/2} \partial_s (\lambda \boldsymbol{\xi})) + (f_0 J_0)^{-1} \partial_k (\mathscr{A}^k (f_0 J_0)^{4/3} \mathcal{F}^{-1/3}) + \mathscr{A} \nabla \Phi \\ &= \left(\partial_s^2 \boldsymbol{\xi} + \frac{1}{2} \frac{\partial_s \lambda}{\lambda} \partial_s \boldsymbol{\xi} + \left(\frac{\partial_s^2 \lambda}{\lambda} - \frac{3}{2} \frac{(\partial_s \lambda)^2}{\lambda^2} \right) \boldsymbol{\xi} \right) + (f_0 J_0)^{-1} \partial_k (\mathscr{A}^k (f_0 J_0)^{4/3} \mathcal{F}^{-1/3}) + \mathscr{A} \nabla \Phi \\ &= \left(\partial_s^2 \boldsymbol{\xi} - \frac{1}{2} \mathscr{C} \partial_s \boldsymbol{\xi} + \delta \boldsymbol{\xi} \right) + (f_0 J_0)^{-1} \partial_k (\mathscr{A}^k (f_0 J_0)^{4/3} \mathcal{F}^{-1/3}) + \mathscr{A} \nabla \Phi \end{aligned}$$

So the Euler-Poisson equations in terms of ξ (2.34) is:

$$\partial_s^2 \boldsymbol{\xi} - \frac{1}{2} \boldsymbol{\ell} \partial_s \boldsymbol{\xi} + \delta \boldsymbol{\xi} + \frac{1}{f_0 J_0} \partial_k (\mathcal{A}^k (f_0 J_0)^{4/3} \boldsymbol{\mathcal{J}}^{-1/3}) + \mathcal{A} \nabla \Phi = \boldsymbol{0}.$$
(3.60)

The self-similarly expanding GW-star is a particular *s*-independent solution of (3.60) of the form $\boldsymbol{\xi}(\mathbf{x}) \equiv \mathbf{x}$ and $f_0 = \bar{w}^3$. Before formulating the stability problem, we must first make the use of the labelling gauge freedom and fix the choice of the initial enthalpy $(f_0 J_0)^{1/3}$ for the general perturbation to be exactly identical to the background enthalpy \bar{w} , i.e. we set

$$(f_0 J_0)^{1/3} = \bar{w}$$
 on $B_R(\mathbf{0})$. (3.61)

Equation (3.61) can be re-written in the form $\rho_0 \circ \eta_0 \det[\nabla \eta_0] = \bar{w}^3$ on the initial domain $B_R(\mathbf{0})$. By a result of Dacorogna-Moser [10] and similarly to [24, 25] there exists a choice of an initial bijective map $\eta_0 : B_R(\mathbf{0}) \to \Omega(\mathbf{0})$ so that (3.61) holds true. The gauge fixing condition (3.61) is necessary as it constrains the freedom to arbitrary relabel the particles at the initial time.

Lemma 3.1 (Euler-Poisson in self-similar coordinate). With respect to the self-similarly expanding profile (λ, \bar{w}) from Definition 1.1, the perturbation θ defined in (2.35) formally solves

$$\partial_s^2 \boldsymbol{\theta} - \frac{1}{2} \boldsymbol{\theta} \partial_s \boldsymbol{\theta} + \delta \boldsymbol{\theta} + \mathbf{P} + \mathbf{G} = \mathbf{0}, \qquad (3.62)$$

where the nonlinear pressure operator \mathbf{P} and the nonlinear gravity operator \mathbf{G} are defined in (2.36) and (2.38).

Proof. Recall that the GW-enthalpy satisfies

$$\mathbf{0} = \delta \mathbf{x} + 4\nabla \bar{w} + \nabla \mathcal{K} \bar{w}^3 \tag{3.63}$$

Using the gauge condition (3.61), the momentum equation (3.60) becomes

$$\bar{w}^3 \left(\partial_s^2 \boldsymbol{\theta} - \frac{1}{2} \boldsymbol{\theta} \partial_s \boldsymbol{\theta} + \delta \boldsymbol{\theta} \right) + \partial_k (\bar{w}^4 (\mathcal{A}^k \mathcal{J}^{-1/3} - I^k)) + \bar{w}^3 (\mathcal{A} \nabla \Phi - \nabla \mathcal{K} \bar{w}^3) = \mathbf{0}.$$

Hence, we can write the momentum equation as

$$\mathbf{0} = \partial_s^2 \boldsymbol{\theta} - \frac{1}{2} \boldsymbol{\theta} \partial_s \boldsymbol{\theta} + \delta \boldsymbol{\theta} + \underbrace{\bar{w}^{-3} \partial_k (\bar{w}^4 (\mathcal{A}^k \mathcal{F}^{-1/3} - I^k))}_{\mathbf{P}} + \underbrace{\mathcal{A} \nabla \Phi - \nabla \mathcal{K} \bar{w}^3}_{\mathbf{G}}.$$

3.1.2 Total energy and momentum

Next we will give expressions for the total momentum and energy in terms of θ . We will write the expressions in a way that separates the linear and non-linear terms of θ clearly. To that end, we first derive the following identity.

Lemma 3.2. For any θ sufficiently smooth we have the identity

$$\int_{B_R} \left(\bar{w}^4 \nabla \cdot \boldsymbol{\theta} + \frac{1}{2} \delta^2 \bar{w}^3 \boldsymbol{\theta} \cdot \mathbf{x} - \frac{1}{2} \bar{w}^3 (\mathcal{K}_{\boldsymbol{\xi}}^{(1)} \bar{w}^3) \right) d\mathbf{x} = 0,$$

where

$$(\mathscr{K}_{\boldsymbol{\xi}}^{(1)}g)(\mathbf{x}) := \int \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^3} g(\mathbf{z}) d\mathbf{z},$$
(3.64)

and we recall (3.59).

Proof. We have

$$\begin{split} \int \bar{w}(\mathbf{x})^3 (\mathscr{K}_{\boldsymbol{\xi}}^{(1)} \bar{w}^3)(\mathbf{x}) \mathrm{d}\mathbf{x} &= \int \int \bar{w}(\mathbf{x})^3 \bar{w}(\mathbf{z})^3 \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^3} \mathrm{d}\mathbf{z} \mathrm{d}\mathbf{x} \\ &= 2 \int \int \bar{w}(\mathbf{x})^3 \bar{w}(\mathbf{z})^3 \frac{(\mathbf{x} - \mathbf{z}) \cdot \boldsymbol{\theta}(\mathbf{x})}{|\mathbf{x} - \mathbf{z}|^3} \mathrm{d}\mathbf{z} \mathrm{d}\mathbf{x} \\ &= 2 \int \bar{w}(\mathbf{x})^3 \boldsymbol{\theta}(\mathbf{x}) \cdot \int \bar{w}(\mathbf{z})^3 \frac{\mathbf{x} - \mathbf{z}}{|\mathbf{x} - \mathbf{z}|^3} \mathrm{d}\mathbf{z} \mathrm{d}\mathbf{x} = 2 \int \bar{w}^3 \boldsymbol{\theta} \cdot \nabla \mathcal{K} \bar{w}^3 \mathrm{d}\mathbf{x} \end{split}$$

Also, using (3.63), we have

$$\int \left(\bar{w}^4 \nabla \cdot \boldsymbol{\theta} - w^3 \boldsymbol{\theta} \cdot \nabla \mathcal{K} \bar{w}^3\right) d\mathbf{x} = \int \left(-\boldsymbol{\theta} \cdot \nabla \bar{w}^4 - w^3 \boldsymbol{\theta} \cdot \nabla \mathcal{K} \bar{w}^3\right) d\mathbf{x}$$
$$= \int \delta \bar{w}^3 \boldsymbol{\theta} \cdot \mathbf{x} d\mathbf{x} = -\frac{1}{2} \theta^2 \int \bar{w}^3 \boldsymbol{\theta} \cdot \mathbf{x} d\mathbf{x}.$$

With this identity we can now derive the expression for the momentum and the energy in terms of θ .

Lemma 3.3 (Momentum and energy in self-similar coordinate). Fix $a \delta \in [\tilde{\delta}, 0)$. In self-similar Lagrangian coordinates introduced above, the total momentum (1.9) and energy (1.10) are respectively denoted by

$$\begin{split} \mathbf{W}_{\delta}[\boldsymbol{\theta}](s) &:= \mathbf{W}_{\delta}(s, \boldsymbol{\theta}(s), \partial_{s}\boldsymbol{\theta}(s)), \\ E_{\delta}[\boldsymbol{\theta}](s) &:= E_{\delta}(s, \boldsymbol{\theta}(s), \partial_{s}\boldsymbol{\theta}(s)), \end{split}$$

where

$$\begin{split} \mathbf{W}_{\delta}(s,\boldsymbol{\theta}(s),\partial_{s}\boldsymbol{\theta}(s)) &= \bar{\mathbf{W}} + \frac{1}{\lambda(s)^{1/2}} \int (\partial_{s}\boldsymbol{\theta}(s) - \boldsymbol{\vartheta}\boldsymbol{\theta}(s))\bar{w}^{3} \mathrm{d}\mathbf{x}, \\ E_{\delta}(s,\boldsymbol{\theta}(s),\partial_{s}\boldsymbol{\theta}(s)) &= \bar{E} + \frac{1}{\lambda(s)} \int \left(\frac{1}{2}\bar{w}^{3} \left(|\partial_{s}\boldsymbol{\theta}(s) - \boldsymbol{\vartheta}\boldsymbol{\theta}(s)|^{2} - \boldsymbol{\vartheta}\mathbf{x} \cdot \left(2\partial_{s}\boldsymbol{\theta}(s) - \frac{5}{2}\boldsymbol{\vartheta}\boldsymbol{\theta}(s)\right)\right)\right) \mathrm{d}\mathbf{x} \\ &+ \frac{1}{\lambda(s)} \int \left(3\bar{w}^{4} \left(\boldsymbol{\mathscr{J}}(s)^{-\frac{1}{3}} - 1 + \frac{1}{3}\nabla\cdot\boldsymbol{\theta}(s)\right) + \frac{1}{2}\bar{w}^{3}(\boldsymbol{\mathscr{K}}_{\boldsymbol{\xi}} - \boldsymbol{\mathscr{K}} - \boldsymbol{\mathscr{K}}_{\boldsymbol{\xi}}^{(1)})\bar{w}^{3}(s)\right) \mathrm{d}\mathbf{x}, \end{split}$$

and $\bar{\mathbf{W}} := \mathbf{W}_{\delta}[\mathbf{0}] = \mathbf{0}$ and $\bar{E} := E_{\delta}[\mathbf{0}] = 0$ are respectively the momentum and energy of the GW star given by Definition 1.1.

Proof. We clearly have

$$\begin{split} \mathbf{W}[\rho,\mathbf{u}] &= \int f J \partial_t \boldsymbol{\eta} \, \mathrm{d}\mathbf{x} = \int f_0 J_0 \partial_t \boldsymbol{\eta} \, \mathrm{d}\mathbf{x} = \int \bar{w}^3 \lambda^{-3/2} \partial_s (\lambda(\mathbf{x}+\boldsymbol{\theta})) \mathrm{d}\mathbf{x} \\ &= \bar{\mathbf{W}} + \frac{1}{\lambda^{1/2}} \int (\partial_s \boldsymbol{\theta} - \boldsymbol{\theta} \boldsymbol{\theta}) \bar{w}^3 \mathrm{d}\mathbf{x}, \\ E[\rho,\mathbf{u}] &= \int \left(\frac{1}{2} f |\partial_t \boldsymbol{\eta}|^2 + 3f^{\frac{4}{3}} + \frac{1}{2} f \boldsymbol{\psi}\right) J \mathrm{d}\mathbf{x} = \int \left(\frac{1}{2} f_0 J_0 |\partial_t \boldsymbol{\eta}|^2 + 3J^{-\frac{1}{3}} (f_0 J_0)^{\frac{4}{3}} + \frac{1}{2} f_0 J_0 \boldsymbol{\psi}\right) \mathrm{d}\mathbf{x} \\ &= \int \left(\frac{1}{2} \bar{w}^3 |\lambda^{-3/2} \partial_s (\lambda(\mathbf{x}+\boldsymbol{\theta}))|^2 + \frac{3}{\lambda} \mathcal{J}^{-\frac{1}{3}} \bar{w}^4 + \frac{1}{2\lambda} \bar{w}^3 \Phi\right) \mathrm{d}\mathbf{x} \\ &= \bar{E} + \frac{1}{\lambda} \int \left(\frac{1}{2} \bar{w}^3 (|\partial_s \boldsymbol{\theta} - \boldsymbol{\theta} \boldsymbol{\theta}|^2 - 2\boldsymbol{\theta} \mathbf{x} \cdot (\partial_s \boldsymbol{\theta} - \boldsymbol{\theta} \boldsymbol{\theta})) + 3(\mathcal{J}^{-\frac{1}{3}} - 1) \bar{w}^4 + \frac{1}{2} \bar{w}^3 (\mathcal{H}_{\boldsymbol{\xi}} - \mathcal{H}) \bar{w}^3 \right) \mathrm{d}\mathbf{x} \\ &= \bar{E} + \frac{1}{\lambda} \int \left(\frac{1}{2} \bar{w}^3 \left(|\partial_s \boldsymbol{\theta} - \boldsymbol{\theta} \boldsymbol{\theta}|^2 - \boldsymbol{\theta} \mathbf{x} \cdot \left(2\partial_s \boldsymbol{\theta} - \frac{5}{2} \boldsymbol{\theta} \boldsymbol{\theta}\right)\right)\right) \mathrm{d}\mathbf{x} \\ &+ \frac{1}{\lambda} \int \left(3 \bar{w}^4 \left(\mathcal{J}^{-\frac{1}{3}} - 1 + \frac{1}{3} \nabla \cdot \boldsymbol{\theta}\right) + \frac{1}{2} \bar{w}^3 (\mathcal{H}_{\boldsymbol{\xi}} - \mathcal{H} - \mathcal{H}_{\boldsymbol{\xi}}^{(1)}) \bar{w}^3\right) \mathrm{d}\mathbf{x}. \end{split}$$

When re-writing $E[\rho, \mathbf{u}]$ above, we have used Lemma 3.2 and (3.63).

Remark 3.4. If instead we consider the GW solutions $\eta_{\mathbf{p}}$ translated at constant velocity \mathbf{p}_1 as in Remark 1.2, we will get $\mathbf{\bar{W}} = M\mathbf{p}_1$ and $\bar{E} = \frac{1}{2}M|\mathbf{p}_1|^2$ instead of $\mathbf{\bar{W}} = \mathbf{0}$ and $\bar{E} = 0$.

3.1.3 High-order energies and the main theorem

We now introduce high-order weighted Sobolev norm that measures the size of the deviation θ without time derivatives. Recall the notation in Section 2. Assuming that $(s, \mathbf{y}) \mapsto \theta(s, \mathbf{y})$ is a sufficiently smooth field, for any $n \in \mathbb{N}_0$ and $s \ge 0$ we let

$$Z_n(s) := \sum_{|\beta|+b \le n} \|X_r^b \partial^\beta \theta\|_{3+b} + \sum_{c \le n} \|\nabla^c \theta\|_{3+2c}$$
(3.65)

$$\mathfrak{X}_n(s) := \sup_{\tau \in [0,s]} Z_n(\tau). \tag{3.66}$$

Next we define energy norms *with* time-derivatives - they will be a basis of our high-order energy method explained in Section 3.5.

$$S_{n}(s) := \sum_{\substack{a+|\beta|+b \le n \\ a>0}} \left(\|\partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\theta\|_{3+b}^{2} + \|\partial_{s}^{a}X_{r}^{b}\partial^{\beta}\theta\|_{3+b}^{2} + \|\partial_{s}^{a}\nabla X_{r}^{b}\partial^{\beta}\theta\|_{4+b}^{2} \right)$$

$$S_{n,c}(s) := \sum_{\substack{a+|\beta|+b \le n \\ a>0 \\ |\beta|+b \le c}} \left(\|\partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\theta\|_{3+b}^{2} + \|\partial_{s}^{a}X_{r}^{b}\partial^{\beta}\theta\|_{3+b}^{2} + \|\partial_{s}^{a}\nabla X_{r}^{b}\partial^{\beta}\theta\|_{4+b}^{2} \right)$$

$$S_{n,c,d}(s) := \sum_{\substack{a+|\beta|+b \le n \\ a>0 \\ |\beta|+b \le c \\ b \le d}} \left(\|\partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\theta\|_{3+b}^{2} + \|\partial_{s}^{a}X_{r}^{b}\partial^{\beta}\theta\|_{3+b}^{2} + \|\partial_{s}^{a}\nabla X_{r}^{b}\partial^{\beta}\theta\|_{4+b}^{2} \right)$$

$$Q_{n}(s) := \sum_{\substack{a+c \le n+1 \\ a>0 \\ c \le d+1}} \|\partial_{s}^{a}\nabla^{c}\theta\|_{3+2c}^{2}$$

Note that $S_{n,n,n} = S_n$. We will also use the convention that $S_{n,-1} = 0$ etc.

Remark 3.5. The indexing above is needed to describe qualitative differences between taking the time, the angular, and the radial derivatives. We shall need this distinction to later close our estimates via a delicate induction argument in high-order spaces, which not only depends on the number of derivatives, but also on the order in which the derivatives are taken.

We define the total instant energy via

$$E_n := S_n + Q_n. \tag{3.67}$$

We shall run the energy identity using E_n ; energies S_n and Q_n will be used for high-order estimates near the vacuum boundary and near the origin respectively. In particular, the control afforded by Q_n is stronger near the origin, while S_n is stronger near the boundary. Finally we define

$$\mathcal{S}_{\bullet}(s) := \sup_{\tau \in [0,s]} S_{\bullet}(\tau) + \int_0^s S_{\bullet}(\tau) \mathrm{d}\tau, \qquad (3.68)$$

$$\mathbb{Q}_{\bullet}(s) := \sup_{\tau \in [0,s]} Q_{\bullet}(\tau) + \int_0^s Q_{\bullet}(\tau) \mathrm{d}\tau, \qquad (3.69)$$

$$\mathscr{C}_{n}(s) := \sup_{\tau \in [0,s]} E_{n}(\tau) + \int_{0}^{s} E_{n}(\tau) \mathrm{d}\tau, \qquad (3.70)$$

where • stands for indices of the form n; n, c; n, c, d in (3.68), and of the form n; n, c in (3.69). The norms (3.68)–(3.70) will play the role of the "left hand side" in the high-order energy identities.

Remark 3.6. We emphasise that the higher order energies E_n we defined (always with a subscript $n \in \mathbb{N}_0$) are different from the total conserved energy E (and E_{δ}) defined in (1.10). Where no confusion arises, we will refer to both as "energy".

In this section, we make the following a priori assumption:

A priori assumption:
$$E_{\bullet}, Z_{\bullet} \leq \epsilon$$
 where $\epsilon > 0$ is some small constant. (3.71)

We now state our main theorem.

Theorem 3.7 (Nonlinear stability of GW stars). Let $n \ge 21$. There exists $\tilde{\delta} \le \delta^* < 0$ such that for any $\delta \in (\delta^*, 0)$ the associated GW expanding star from Definition 1.1 is codimension-4 nonlinearly stable in

the class of irrotational perturbations. More precisely, there exists an $\epsilon_0 > 0$ such that for any initial data $(\boldsymbol{\theta}(0), \partial_s \boldsymbol{\theta}(0))$ satisfying

$$E_n(0) + Z_n(0)^2 \le \epsilon_0 \tag{3.72}$$

$$\mathbf{W}_{\delta}(0,\boldsymbol{\theta}(0),\partial_{s}\boldsymbol{\theta}(0)) = \mathbf{W}_{\delta}[\mathbf{0}] =: \bar{\mathbf{W}} = \mathbf{0}$$
(3.73)

$$\mathbf{W}_{\delta}(0,\boldsymbol{\theta}(0),\partial_{s}\boldsymbol{\theta}(0)) = \mathbf{W}_{\delta}[\mathbf{0}] =: \mathbf{W} = \mathbf{0}$$

$$E_{\delta}(0,\boldsymbol{\theta}(0),\partial_{s}\boldsymbol{\theta}(0)) = E_{\delta}[\mathbf{0}] =: \bar{E} = 0$$
(3.74)

$$\operatorname{curl}_A \partial_t \boldsymbol{\eta}(0) = \mathbf{0},\tag{3.75}$$

the associated solution $s \mapsto (\theta(s, \cdot), \partial_s \theta(s, \cdot))$ to (3.62) exists for all $s \ge 0$ and is unique in the class of all data with finite norm $E_n + Z_n^2$. Moreover, there exists a constant C > 0 such that

$$E_n(s) + Z_n(s)^2 \le C\epsilon_0$$
 for all $s \ge 0$,

and $E_n(s)$ decays exponentially fast in s.

Note that condition (3.75) is the Lagrangian statement that the fluid velocity is irrotational. The momentum and energy constraints (3.73)-(3.74) define the codimension-4 "manifold" of initial data.

Heuristically speaking, since momentum and energy are conserved quantities for the Euler-Poisson system, it is necessary that our perturbation does not alter the momentum and energy if our background solution were to be the right asymptotic-in-time "limit". Indeed, the momentum and energy constraints (3.73)-(3.74) are necessary in our stability analysis due to the presence of growing modes of the linearised operator in self-similar coordinates induced by the conservation of the energy and momentum. However if our perturbation does alter the momentum W and energy E away from 0 and 0 such that $E = \frac{1}{2} |\mathbf{W}|^2 / M$, our proof can be easily adapted to show that it still leads to global existence with the solution staying close to a GW star for all time, but one translated at constant velocity \mathbf{p}_1 with $\mathbf{W} = M\mathbf{p}_1$ and $E = \frac{1}{2}M|\mathbf{p}_1|^2$ as described in Remarks 1.2 and 3.4. In this sense the "manifold" of GW-solutions is codimension-1 nonlinearly stable in the class of irrotational perturbations, even though each individual GW-star is only codimension-4 stable. In particular, given any initial data (ρ_0, \mathbf{u}_0) such that $E[\rho_0, \mathbf{u}_0] = \frac{1}{2} |\mathbf{W}[\rho_0, \mathbf{u}_0]|^2 / M[\rho_0, \mathbf{u}_0]$, we can change our frame of reference and subtract a constant velocity of $\mathbf{p}_1 = \mathbf{\bar{W}}[\rho_0, \mathbf{u}_0] |/M[\rho_0, \mathbf{u}_0]$ from \mathbf{u}_0 to obtain

$$\begin{split} \mathbf{W}[\rho_0, \mathbf{u}_0 - \mathbf{p}_1] &= \mathbf{W}[\rho_0, \mathbf{u}_0] - M[\rho_0, \mathbf{u}_0] \mathbf{p}_1 = \mathbf{0} \\ E[\rho_0, \mathbf{u}_0 - \mathbf{p}_1] &= E[\rho_0, \mathbf{u}_0] - \int_{\mathbb{R}^3} \rho_0 \mathbf{u}_0 \cdot \mathbf{p}_1 d\mathbf{x} + \frac{1}{2} \int_{\mathbb{R}^3} \rho_0 |\mathbf{p}_1|^2 d\mathbf{x} \\ &= E[\rho_0, \mathbf{u}_0] - \mathbf{p}_1 \cdot \mathbf{W}[\rho_0, \mathbf{u}_0] + \frac{1}{2} |\mathbf{p}_1|^2 M[\rho_0, \mathbf{u}_0] = 0 \end{split}$$

So in this new frame of reference, the constraints (3.73) and (3.74) are satisfied.

To formalise this, note that for any $\mathbf{p}_1 \in \mathbb{R}^3$, $\boldsymbol{\theta} = \mathbf{p}_1 t \lambda(t)^{-1}$ is a global-in-time solution to (3.62) which corresponds to a Lagrangian description of a GW-star translated by a constant velocity. Then, as a corollary of Theorem 3.7 we have the following result.

Corollary 3.8. Let $n \ge 21$. There exists $\tilde{\delta} \le \delta^* < 0$ such that for any $\delta \in (\delta^*, 0)$, the "manifold" of *GW*-stars $(\bar{\rho}_{\mathbf{p}}, \bar{\mathbf{u}}_{\mathbf{p}})$, $\mathbf{p} = \mathbf{p}_1 t$, $\mathbf{p}_1 \in \mathbb{R}^3$ (from Remark 1.2) is codimension-1 nonlinearly stable in the class of irrotational perturbations. More precisely, for given any initial data $(\tilde{\theta}(0), \partial_s \tilde{\theta}(0))$ define

$$\boldsymbol{\theta}_0 = \boldsymbol{\theta}(0)$$
$$(\partial_s \boldsymbol{\theta})_0 = \partial_s \tilde{\boldsymbol{\theta}}(0) - \mathbf{p}_1,$$

where

$$\mathbf{p}_1 = \frac{W_{\delta}(0, \tilde{\boldsymbol{\theta}}(0), \partial_s \tilde{\boldsymbol{\theta}}(0))}{M[\bar{w}^3]}.$$

Then, there exists an $\epsilon_0 > 0$ such that for any initial data $(\tilde{\theta}(0), \partial_s \tilde{\theta}(0))$ such that

$$(E_n + Z_n^2)[\boldsymbol{\theta}_0, (\partial_s \boldsymbol{\theta})_0] \le \epsilon_0$$

$$E_{\delta}(0, \tilde{\boldsymbol{\theta}}(0), \partial_{s} \tilde{\boldsymbol{\theta}}(0)) = \frac{1}{2} |\mathbf{W}_{\delta}(0, \tilde{\boldsymbol{\theta}}(0), \partial_{s} \tilde{\boldsymbol{\theta}}(0))|^{2} / M[\bar{w}^{3}]$$

curl_A $\partial_{t} \boldsymbol{\eta}(0) = \mathbf{0}$

the associated solution $s \mapsto (\tilde{\theta}(s, \cdot), \partial_s \tilde{\theta}(s, \cdot))$ to (3.62) exists for all $s \ge 0$ and is unique in the class of all data with finite norm $(E_n + Z_n^2)[\tilde{\theta}]$. Moreover, there exists a constant C > 0 such that

$$(E_n + Z_n^2)[\boldsymbol{\theta}](s) \le C\epsilon_0 \quad \text{for all} \quad s \ge 0,$$

where $\theta = \tilde{\theta} - \mathbf{p}_1 t \lambda(t)^{-1}$, and $E_n[\theta](s)$ decays exponentially fast in s.

Remark 3.9. Our goal is not to optimise the number n of derivatives in our spaces. As usual, the size of n is conditioned by the Hardy-Sobolev type embeddings, which allow us to bound the L^{∞} -norms of contributions with less than $|\frac{n}{2}|$ derivatives by \overline{w}^k -weighted Sobolev norms.

Remark 3.10. The subclass of expanding GW-stars with non-zero total energy (in the frame of reference of **0** momentum) consists of stars that expand at a linear rate in time, i.e. not at the self-similar rate considered above. This problem leads to a stronger damping effect which allows the "Euler part" of the flow to dominate the dynamics. The stability of such GW-stars is the content of Section 4.

Local-in-time well-posedness. The presence of vacuum is known to pose challenges in the well-posedness theory for compressible fluid flows. To develop a satisfactory local existence and uniqueness theory, one needs to impose an additional assumption on the initial data - the so-called physical vacuum condition (1.7). In the works of Jang and Masmoudi [31] and independently Coutand and Shkoller [8] the local well-posedness for the compressible Euler equations was shown in the Lagrangian coordinates (for a more recent treatment in Eulerian coordinates see [27]). From the point of view of regularity theory, gravity represents a lower order term, so the techniques from [31, 8] can be adapted to obtain a local-in-time well-posedness result for the free boundary EP-system [30, 18, 36, 26]. In particular, a simple adaptation of the methods in [31, 26] yields the following local well-posedness result in the weighted high-order energy space $E_n + Z_n^2$.

Theorem 3.11 (Local well-posedness). Let $n \ge 21$. Then for any given initial data $(\theta(0), \partial_s \theta(0))$ such that $E_n(0) + Z_n(0)^2 < \infty$, there exist some T > 0 and a unique solution $(\theta, \partial_s \theta) : [0, T] \times B_R \to \mathbb{R}^3 \times \mathbb{R}^3$ to (3.62) such that $E_n(s) + Z_n(s)^2 \le 2(E_n(0) + Z_n(0)^2)$ for all $s \in [0, T]$.

Theorem 3.11 is a starting point for the continuity argument that will culminate in the proof of Theorem 3.7.

3.1.4 Proof strategy

The basic idea behind the global existence in Theorem 3.7 is the presence of the damping term $-\frac{1}{2}\theta \partial_s \theta$ in (3.62), which clearly suggests a stabilising mechanism in the problem. Such a term appears as a direct consequence of the expanding character of the underlying GW-motion (and it would be of the opposite sign if we were linearising about a collapsing GW-star). This stabilisation effect was first exhibited in [24] where the purely radial version of Theorem 3.7 was established.

Since the problem features the vacuum free boundary satisfying the physical vacuum condition (1.7), we use weighted high-order energy spaces introduced by Jang and Masmoudi [31]. The key idea to overcome a possible loss of derivatives is to introduce increasing powers of \bar{w} into the function spaces, as we increase the number of radial derivatives, but not the tangential ones. In particular, the proof of the main result is based on a high-order energy method which necessitates commuting the equation (3.62) with operators of the form $\partial_s^a X_r^n \partial^\beta$. To understand the energy contribution from the combined pressure and gravity term $\mathbf{P} + \mathbf{G}$ (see (2.36)–(2.38)), we must linearise (3.62). As shown in Lemma 3.12, this linearisation reads

$$\partial_s^2 \boldsymbol{\theta} - \frac{1}{2} \boldsymbol{\theta} \partial_s \boldsymbol{\theta} + \mathbf{L} \boldsymbol{\theta} = \mathbf{0}, \qquad (3.76)$$

where the linearised operator \mathbf{L} takes the form

$$\mathbf{L}\boldsymbol{\theta} := -\frac{4}{3}\nabla(\bar{w}^{-2}\nabla\cdot(\bar{w}^{3}\boldsymbol{\theta})) - \nabla\mathcal{K}\nabla\cdot(\bar{w}^{3}\boldsymbol{\theta}), \qquad (3.77)$$

where we recall (3.59) and (1.16). The fundamental challenge with respect to the radial result [24] is to show the coercivity of the operator L in suitably weighted spaces, dictated by our local-in-time well-posedness theory.

The difficulty in proving a useful coercivity bound for the operator L lies in the antagonism between the nonlocal nature of the gravitational interaction described by G in (2.38), and the Lagrangian perspective, which is naturally imposed on us by the problem. The operator L has a nontrivial unstable space, spanned by the eigenvectors x and the standard basis e_i , i = 1, 2, 3. The 4-dimensional nature of the unstable space is a reflection of the energy and momentum conservation laws, which in self-similar variables induce formally unstable modes.

Nonradial linearised analysis around the Lane-Emden stars ($\delta = \lambda_1 = 0$) is given in [32] where the non-negativity of the associated quadratic form is shown using the expansion in spherical harmonics. In this work we work in a similar spirit, but our linear analysis around the GW-stars improves upon [32] considerably, as we show strict quantitative coercivity bound

$$\langle \mathbf{L}\boldsymbol{\theta}, \boldsymbol{\theta} \rangle_3 \gtrsim \int_{B_R} \bar{w}^{-2} |\Delta \Psi|^2 d\mathbf{x} + \int_{\mathbb{R}^3} |\nabla \Psi|^2 d\mathbf{x},$$
 (3.78)

under the crucial orthogonality conditions

$$\langle \boldsymbol{\theta}, \mathbf{x} \rangle_3 = 0 = \langle \boldsymbol{\theta}, \mathbf{e}_i \rangle_3, \qquad i = 1, 2, 3, \qquad (3.79)$$

where $\Delta \Psi = \operatorname{div}(\bar{w}^3\theta)$. This is the central estimate of Section 3.2 (see Theorem 3.16) and it relies on a careful decomposition in spherical harmonics. It is non-trivial as it requires a careful use of the above orthogonality conditions to obtain quantitative lower bounds for the 0-th and the 1-st order spherical harmonics. In the former case, the problem essentially reduces to the radial coercivity bound from [24], while the analysis of the projection of L onto 1-st order spherical harmonics requires a careful use of Sturm-Liouville theory, see Lemma 3.21, a related argument was used in [32].

One of the main challenges is that the quantity $\int_{B_R} \bar{w}^{-2} |\Delta \Psi|^2 d\mathbf{x} + \int_{\mathbb{R}^3} |\nabla \Psi|^2 d\mathbf{x}$ on the right-hand side of (3.78) a priori does not appear useful for the energy estimates as we need to control the norms $\|\theta\|_3^2 + \|\nabla \theta\|_4^2$, which are localised to the set B_R by definition, see (2.30). An intermediate step towards a resolution of this issue is to relate the general estimate (3.78) (which holds for any sufficiently smooth map θ), to the nonlinear dynamics. In Section 3.3, by linearising the nonlinear energy-momentum constraints

$$E[\rho, \mathbf{u}] = \overline{E}, \qquad \mathbf{W}[\rho, \mathbf{u}] = \overline{\mathbf{W}}, \qquad (3.80)$$

we obtain effective ODEs (modulo lower order nonlinear terms) that allow to dynamically control the inner products $\langle \theta, \mathbf{x} \rangle_3$ and $\langle \theta, \mathbf{e}_i \rangle_3$. With this in hand we prove in Proposition 3.27 a high-order differentiated version of the bound (3.78) for the solutions of (3.76) satisfying the constraints (3.80):

$$\int_{B_R} \bar{w}^{-2} |\nabla \cdot (\bar{w}^3 \partial_s^a \partial_s^\beta \theta)|^2 \mathrm{d}\mathbf{x} + \|\partial_s^{a+1} \partial_s^\beta \theta\|_3^2 \lesssim \langle \mathbf{L} \partial_s^a \partial_s^\beta \theta, \partial_s^a \partial_s^\beta \theta \rangle_3 + \|\partial_s^{a+1} \partial_s^\beta \theta\|_3^2 + \text{l.o.t.}$$
(3.81)

The final and crucial step toward useful lower bounds is to exploit the irrotationality assumption $\nabla \times \mathbf{u} = 0$ to obtain a dynamic control over $\|\boldsymbol{\theta}\|_3^2 + \|\nabla\boldsymbol{\theta}\|_4^2$. Looking at (3.81), this necessitates a careful examination of the \bar{w} -weighted divergence appearing on the left-hand side. It is clear that any vectorfield such that $\nabla \cdot (\bar{w}^3 \boldsymbol{\theta}) = 0$ formally belongs to the kernel of \mathbf{L} and therefore, to obtain strict coercivity, we must mod out this infinite-dimensional kernel. The orthogonal complement with respect to the $\langle \cdot, \cdot \rangle_3$ -inner product consists precisely of the gradients, so the first key observation is the content of Lemma 3.28, which roughly states that $\partial_s^a \boldsymbol{\theta}$ is a gradient modulo "good" terms for $a \ge 1$, assuming $\nabla \times \mathbf{u} = 0$. Here, in simplest possible terms, the issue is that the irrotationality in Lagrangian variables creates error terms that a priori seem problematic, but luckily all such terms can be absorbed into a pure gradient. The second key ingredient is Lemma 3.33, which is an exact identity relating the norm of the weighted divergence of $\boldsymbol{\theta}$ to the weighted norms of the derivative of $\boldsymbol{\theta}$. This can be viewed as a form of "elliptic regularity". Finally, we use these ingredients in the central statement of Section 3.4 - Proposition 3.34 - to show that natural energy norms obtained via integration-by-parts from (3.76) control the weighted norms of the pure time derivatives

of θ . In Proposition 3.36 we treat also the angular derivatives in our operators, and the same statement as in the previous proposition holds, modulo the presence of a linear (and therefore not small) contribution, which fortunately involves one angular derivative less. This decoupling structure enables us to use a careful inductive procedure to eventually close the nonlinear estimates.

The nonlinear arguments are presented in Sections 3.5 and 3.6. The global nonlinear stability will follow from the bound

$$\mathscr{E}_n \lesssim \mathscr{E}_n(0) + (\mathscr{E}_n + \mathscr{Z}_n^2)^{1/2} \mathscr{E}_n \tag{3.82}$$

in the regime where $E_n + Z_n^2$ is sufficiently small. To prove such a bound, we commute (3.62) with highorder derivatives and while the discussion above refers to the extraction of coercive bounds for the linear part of the operator, we are still left with the nonlinear estimates. Propositions 3.38, 3.39, and 3.42 show that the deviation of the pressure term **P** and the gravity term **G** from its linearisation, can be controlled by the good trilinear error $(\mathcal{E}_n + \mathcal{I}_n^2)^{1/2} \mathcal{E}_n$ modulo some terms that scale like the linear norms, but always decouple at the top order of differentiation, so that they involve, for example, "one spatial derivative less and one time derivative more". This decoupling is crucial for the closure of the estimates, and the key effective reduction to the linear problem is formulated in Theorem 3.44. This feature of the problem suggests that we can show (3.82) inductively by taking derivatives in the right order. Key energy bounds for the nonlinear contributions from the pressure and the gravity are presented in Sections 3.6.1 and 3.6.2 respectively. The final continuity argument and the exponential decay based on (3.82) is presented in Section 3.6.3.

3.2 Linearisation and coercivity

3.2.1 The linear and non-linear part of Euler-Poisson system

The proof of Theorem 3.7 crucially relies on good coercive properties of the linearisation around the background GW-star. In the next lemma we formally derive the linearised Euler-Poisson system.

Lemma 3.12 (Linearised Euler-Poisson). The formal linearisation of (3.62) reads

$$\partial_s^2 \boldsymbol{\theta} - \frac{1}{2} \boldsymbol{\theta} \partial_s \boldsymbol{\theta} + \mathbf{L} \boldsymbol{\theta} = \mathbf{0}$$
(3.83)

where

$$\mathbf{L}\boldsymbol{\theta} := -\frac{4}{3}\nabla(\bar{w}^{-2}\nabla\cdot(\bar{w}^{3}\boldsymbol{\theta})) - \nabla\mathcal{K}\nabla\cdot(\bar{w}^{3}\boldsymbol{\theta})$$
(3.84)

and we recall (3.59). Moreover, the formal linearisation of the gravitational contribution **G** (2.38) is given by the operator

$$\mathbf{G}_{L}\boldsymbol{\theta} := \boldsymbol{\theta} \cdot \nabla \nabla \mathcal{K} \bar{w}^{3} - \nabla \mathcal{K} \nabla \cdot (\bar{w}^{3}\boldsymbol{\theta}) = \mathcal{K}_{\boldsymbol{\xi}}^{(1)} \nabla \bar{w}^{3} - \mathcal{K} \partial_{i} (\bar{w}^{3} \nabla \theta^{i}), \qquad (3.85)$$

where we recall (2.40) and (3.64).

Proof. Since $\nabla \boldsymbol{\xi} = I + \nabla \boldsymbol{\theta}$, to first order (in $\boldsymbol{\theta}$) we have $\mathcal{A} = I - \nabla \boldsymbol{\theta}$ and $\mathcal{J} = 1 + \nabla \cdot \boldsymbol{\theta}$. So to first order we have

$$\mathscr{A}\mathcal{J}^{-1/3} = (I - \nabla\theta)(1 + \nabla \cdot \theta)^{-1/3} = (I - \nabla\theta)\left(1 - \frac{1}{3}\nabla \cdot \theta\right) = \left(1 - \frac{1}{3}\nabla \cdot \theta\right)I - \nabla\theta$$

and

$$\begin{aligned} \frac{1}{\bar{w}^3} \partial_k (\bar{w}^4 (\mathscr{A}^k \mathscr{F}^{-1/3} - I^k)) &= -\frac{1}{3\bar{w}^3} \nabla (\bar{w}^4 \nabla \cdot \boldsymbol{\theta}) - \frac{1}{\bar{w}^3} \partial_k (\bar{w}^4 \nabla \theta^k) \\ &= -\frac{4}{3} \nabla (\bar{w} \nabla \cdot \boldsymbol{\theta}) - 4 \nabla (\boldsymbol{\theta} \cdot \nabla \bar{w}) + 4 \boldsymbol{\theta} \cdot \nabla \nabla \bar{w} \\ &= -\frac{4}{3} \nabla (\bar{w}^{-2} \nabla \cdot (\bar{w}^3 \boldsymbol{\theta})) + 4 \boldsymbol{\theta} \cdot \nabla \nabla \bar{w} \\ &= -\frac{4}{3} \nabla (\bar{w}^{-2} \nabla \cdot (\bar{w}^3 \boldsymbol{\theta})) - \boldsymbol{\theta} \cdot \nabla (\delta \mathbf{x} + \nabla \mathscr{K} \bar{w}^3) \end{aligned}$$

$$= -\frac{4}{3}\nabla(\bar{w}^{-2}\nabla\cdot(\bar{w}^{3}\boldsymbol{\theta})) - \delta\boldsymbol{\theta} - \boldsymbol{\theta}\cdot\nabla\nabla\mathcal{K}\bar{w}^{3}$$

Since

$$\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|^2 = |\mathbf{x} - \mathbf{z} + \boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^2 = |\mathbf{x} - \mathbf{z}|^2 + 2(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})) + |\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^2,$$

to first order we have

$$\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} = \frac{1}{|\mathbf{x} - \mathbf{z}|} \left(1 - \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^2} \right).$$

So to first order we have

$$\begin{split} \Phi(\mathbf{x}) &= -\int \frac{\bar{w}(\mathbf{z})^3}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} d\mathbf{z} = -\int \frac{\bar{w}(\mathbf{z})^3}{|\mathbf{x} - \mathbf{z}|} d\mathbf{z} + \int \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^3} \bar{w}(\mathbf{z})^3 d\mathbf{z} \\ &= (\mathcal{K}\bar{w}^3)(\mathbf{x}) + \int \left(-\boldsymbol{\theta}(\mathbf{x}) \cdot \nabla_{\mathbf{x}} \frac{1}{|\mathbf{x} - \mathbf{z}|} - \boldsymbol{\theta}(\mathbf{z}) \cdot \nabla_{\mathbf{z}} \frac{1}{|\mathbf{x} - \mathbf{z}|}\right) \bar{w}(\mathbf{z})^3 d\mathbf{z} \\ &= (\mathcal{K}\bar{w}^3)(\mathbf{x}) + \boldsymbol{\theta} \cdot \nabla(\mathcal{K}\bar{w}^3)(\mathbf{x}) - (\mathcal{K}\nabla \cdot (\bar{w}^3\boldsymbol{\theta}))(\mathbf{x}) \end{split}$$

and

$$\begin{aligned} \mathcal{A}\nabla\Phi &= (I^{i} - \nabla\theta^{i})\partial_{i}(\mathcal{K}\bar{w}^{3} + \boldsymbol{\theta}\cdot\nabla\mathcal{K}\bar{w}^{3} - \mathcal{K}\nabla\cdot(\bar{w}^{3}\boldsymbol{\theta})) \\ &= \nabla\mathcal{K}\bar{w}^{3} - (\nabla\theta^{i})\partial_{i}\mathcal{K}\bar{w}^{3} + \nabla(\boldsymbol{\theta}\cdot\nabla\mathcal{K}\bar{w}^{3}) - \nabla\mathcal{K}\nabla\cdot(\bar{w}^{3}\boldsymbol{\theta}) \\ &= \nabla\mathcal{K}\bar{w}^{3} + \boldsymbol{\theta}\cdot\nabla\nabla\mathcal{K}\bar{w}^{3} - \nabla\mathcal{K}\nabla\cdot(\bar{w}^{3}\boldsymbol{\theta}) = \nabla\mathcal{K}\bar{w}^{3} + \mathbf{G}_{L}\boldsymbol{\theta}, \end{aligned}$$

where we have used (3.85) in the last line. Therefore the linearisation of the momentum equation (3.62) takes the form (3.83). Note that

$$\begin{split} \mathbf{G}_{L}\boldsymbol{\theta} &= \boldsymbol{\theta} \cdot \nabla \mathcal{K} \nabla \bar{w}^{3} - \mathcal{K} \partial_{i} (\nabla \bar{w}^{3} \theta^{i}) - \mathcal{K} \partial_{i} (\bar{w}^{3} \nabla \theta^{i}) \\ &= \int \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^{3}} \nabla \bar{w}(\mathbf{z})^{3} \mathrm{d}\mathbf{z} - \mathcal{K} \partial_{i} (\bar{w}^{3} \nabla \theta^{i}) = \mathcal{K}_{\boldsymbol{\xi}}^{(1)} \nabla \bar{w}^{3} - \mathcal{K} \partial_{i} (\bar{w}^{3} \nabla \theta^{i}), \end{split}$$

which completes the proof of the lemma.

Finally, it will be important to keep track of the precise structure of the nonlinear correction $\mathbf{G} - \mathbf{G}_L \boldsymbol{\theta}$, which is given in the next lemma.

Lemma 3.13 (Non-linear part of gravity term). We have

$$\mathbf{G} - \mathbf{G}_{L}\boldsymbol{\theta} = \mathscr{K}_{\boldsymbol{\xi}}(\mathscr{A}_{l}^{i}(\partial_{k}\theta^{l})(\nabla\theta^{k})\partial_{i}\bar{w}^{3} - \bar{w}^{3}(\mathscr{A}_{m}^{i}\mathscr{A}_{\bullet}^{l} - I_{m}^{i}I_{\bullet}^{l})\partial_{i}\partial_{l}\theta^{m}) - (\mathscr{K}_{\boldsymbol{\xi}} - \mathscr{K})\partial_{i}(\bar{w}^{3}\nabla\theta^{i}) + (\mathscr{K}_{\boldsymbol{\xi}} - \mathscr{K} - \mathscr{K}_{\boldsymbol{\xi}}^{(1)})\nabla\bar{w}^{3}$$
(3.86)

Proof. Since $\mathcal{A} = (\nabla \boldsymbol{\xi})^{-1}$, we have

$$I_j^i = \mathscr{A}_k^i \partial_j \xi^k = \mathscr{A}_k^i (I_j^k + \partial_j \theta^k)$$

Therefore $\mathcal{A}^i_j - I^i_j = -\mathcal{A}^i_k \partial_j \theta^k$. We have

$$\begin{aligned} \mathbf{G} - \mathbf{G}_{L}\boldsymbol{\theta} &= \mathfrak{K}_{\boldsymbol{\xi}}((\mathfrak{A} - I)\nabla\bar{w}^{3} - \bar{w}^{3}\mathcal{A}_{m}^{i}\mathcal{A}_{\bullet}^{l}\partial_{i}\partial_{l}\theta^{m}) + (\mathfrak{K}_{\boldsymbol{\xi}} - \mathfrak{K} - \mathfrak{K}_{\boldsymbol{\xi}}^{(1)})\nabla\bar{w}^{3} + \mathfrak{K}\partial_{i}(\bar{w}^{3}\nabla\theta^{i}) \\ &= \mathfrak{K}_{\boldsymbol{\xi}}((\nabla\theta^{i} - \mathcal{A}_{k}^{i}\nabla\theta^{k})\partial_{i}\bar{w}^{3} - \bar{w}^{3}(\mathcal{A}_{m}^{i}\mathcal{A}_{\bullet}^{l} - I_{m}^{i}I_{\bullet}^{l})\partial_{i}\partial_{l}\theta^{m}) \\ &+ (\mathfrak{K} - \mathfrak{K}_{\boldsymbol{\xi}})\partial_{i}(\bar{w}^{3}\nabla\theta^{i}) + (\mathfrak{K}_{\boldsymbol{\xi}} - \mathfrak{K} - \mathfrak{K}_{\boldsymbol{\xi}}^{(1)})\nabla\bar{w}^{3} \\ &= \mathfrak{K}_{\boldsymbol{\xi}}(\mathcal{A}_{l}^{i}(\partial_{k}\theta^{l})(\nabla\theta^{k})\partial_{i}\bar{w}^{3} - \bar{w}^{3}(\mathcal{A}_{m}^{i}\mathcal{A}_{\bullet}^{l} - I_{m}^{i}I_{\bullet}^{l})\partial_{i}\partial_{l}\theta^{m}) \\ &- (\mathfrak{K}_{\boldsymbol{\xi}} - \mathfrak{K})\partial_{i}(\bar{w}^{3}\nabla\theta^{i}) + (\mathfrak{K}_{\boldsymbol{\xi}} - \mathfrak{K} - \mathfrak{K}_{\boldsymbol{\xi}}^{(1)})\nabla\bar{w}^{3}. \end{aligned}$$

We next derive helpful identities for the operators $\mathcal{K}_{\xi} - \mathcal{K}$ and $\mathcal{K}_{\xi} - \mathcal{K} - \mathcal{K}_{\xi}^{(1)}$ appearing on the right-hand side of (3.86). We first note that

$$(\mathscr{K}_{\boldsymbol{\xi}} - \mathscr{K})g(\mathbf{x}) = -\int_{\mathbb{R}^3} K_1(\mathbf{x}, \mathbf{z})g(\mathbf{z})d\mathbf{z},$$
(3.87)

$$(\mathscr{K}_{\boldsymbol{\xi}} - \mathscr{K} - \mathscr{K}_{\boldsymbol{\xi}}^{(1)})g(\mathbf{x}) = -\int_{\mathbb{R}^3} K_2(\mathbf{x}, \mathbf{z})g(\mathbf{z})d\mathbf{z},$$
(3.88)

where

$$K_1(\mathbf{x}, \mathbf{z}) := \frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} - \frac{1}{|\mathbf{x} - \mathbf{z}|},$$
(3.89)

$$K_2(\mathbf{x}, \mathbf{z}) := \frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} - \frac{1}{|\mathbf{x} - \mathbf{z}|} + \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^3}.$$
(3.90)

In the following lemma, we write K_1 and K_2 explicitly in terms of θ , which will play a role in our energy estimates. In particular, we see that θ appears at least linearly in K_1 , and at least quadratically in K_2 . Lemma 3.14. We have

$$K_{2}(\mathbf{x}, \mathbf{z}) = K_{1}(\mathbf{x}, \mathbf{z}) + \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^{3}}$$

$$= -\frac{1}{2} \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^{2}}{|\mathbf{x} - \mathbf{z}|^{3}} + \frac{3}{4|\mathbf{x} - \mathbf{z}|} \left(2 \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^{2}} + \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^{2}}{|\mathbf{x} - \mathbf{z}|^{2}} \right)^{2}$$

$$\varpi_{\frac{1}{2}} \left(2 \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^{2}} + \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^{2}}{|\mathbf{x} - \mathbf{z}|^{2}} \right), \quad (3.91)$$

where

$$\varpi_q(y) := \int_0^1 \frac{1-z}{(1+yz)^{q+2}} \mathrm{d}z, \qquad y > -1, \quad q \in \mathbb{R}.$$
(3.92)

Proof. Let $q \in \mathbb{R} \setminus \{-1, 0\}$. Then for y > -1 and $y \neq 0$

$$\int_0^1 \frac{1-z}{(1+yz)^{q+2}} dz = -\frac{1}{(q+1)y} \left[\frac{1-z}{(1+yz)^{q+1}} \right]_0^1 - \frac{1}{(q+1)y} \int_0^1 \frac{1}{(1+yz)^{q+1}} dz$$
$$= \frac{1}{(q+1)y} \left(1 + \frac{1}{qy} \left[\frac{1}{(1+yz)^q} \right]_0^1 \right) = \frac{1}{q(q+1)y^2} \left(-1 + qy + \frac{1}{(1+y)^q} \right)$$

and thus

$$\frac{1}{(1+y)^q} = 1 - qy + q(q+1)y^2 \varpi_q(y), \qquad y > -1$$
(3.93)

where we note that (3.93) trivially holds for y = 0 and q = -1, 0. Since

$$|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|^2 = |\mathbf{x} - \mathbf{z} + \boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^2 = |\mathbf{x} - \mathbf{z}|^2 + 2(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})) + |\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^2,$$

we have

$$\frac{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|^2}{|\mathbf{x} - \mathbf{z}|^2} = 1 + 2\frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^2} + \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^2}{|\mathbf{x} - \mathbf{z}|^2}.$$

Hence by applying (3.93) with $y = 2 \frac{(\mathbf{x}-\mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x}-\mathbf{z}|^2} + \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^2}{|\mathbf{x}-\mathbf{z}|^2}$ and $q = \frac{1}{2}$, we see that

$$\frac{|\mathbf{x} - \mathbf{z}|}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} = 1 - \left(\frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^2} + \frac{1}{2} \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^2}{|\mathbf{x} - \mathbf{z}|^2}\right) \\ + \frac{3}{4} \left(2 \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^2} + \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^2}{|\mathbf{x} - \mathbf{z}|^2}\right)^2$$

$$\varpi_{\frac{1}{2}}\left(2\frac{(\mathbf{x}-\mathbf{z})\cdot(\boldsymbol{\theta}(\mathbf{x})-\boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x}-\mathbf{z}|^2}+\frac{|\boldsymbol{\theta}(\mathbf{x})-\boldsymbol{\theta}(\mathbf{z})|^2}{|\mathbf{x}-\mathbf{z}|^2}\right).$$

Therefore, we obtain

$$K_{2}(\mathbf{x}, \mathbf{z}) = \underbrace{\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|}_{=K_{1}(\mathbf{x}, \mathbf{z})} - \frac{1}{|\mathbf{x} - \mathbf{z}|}_{=K_{1}(\mathbf{x}, \mathbf{z})} + \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^{3}}$$

$$= -\frac{1}{2} \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^{2}}{|\mathbf{x} - \mathbf{z}|^{3}} + \frac{3}{4|\mathbf{x} - \mathbf{z}|} \left(2\frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^{2}} + \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^{2}}{|\mathbf{x} - \mathbf{z}|^{2}}\right)^{2}$$

$$\varpi_{\frac{1}{2}} \left(2\frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^{2}} + \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^{2}}{|\mathbf{x} - \mathbf{z}|^{2}}\right).$$

3.2.2 Coercivity of L

A fundamental prerequisite for the understanding of the nonlinear stability is a good linear stability theory. This entails a precise understanding of the coercivity properties of the operator \mathbf{L} and this is the subject of this section.

For sufficiently smooth θ , we have

$$\langle \mathbf{L}\boldsymbol{\theta}_1,\boldsymbol{\theta}_2\rangle_3 = \int_{B_R} \left(\frac{4}{3}\bar{w}^{-2}\nabla\cdot(\bar{w}^3\boldsymbol{\theta}_2)\nabla\cdot(\bar{w}^3\boldsymbol{\theta}_1) + \nabla\cdot(\bar{w}^3\boldsymbol{\theta}_2)\mathscr{K}\nabla\cdot(\bar{w}^3\boldsymbol{\theta}_1)\right) \mathrm{d}\mathbf{x}$$

Note this is defined in a weak sense for θ_i (i = 1, 2) such that $\nabla \cdot (\bar{w}^3 \theta_i) \in L^2(B_R, \bar{w}^{-2})$. We see that L is symmetric under $\langle \cdot, \cdot \rangle_3$ since

$$\int \nabla \cdot (\bar{w}^3 \boldsymbol{\theta}_2) \mathscr{K} \nabla \cdot (\bar{w}^3 \boldsymbol{\theta}_1) d\mathbf{x} = - \int \int \frac{\nabla \cdot (\bar{w}^3 \boldsymbol{\theta}_2)(\mathbf{x}) \nabla \cdot (\bar{w}^3 \boldsymbol{\theta}_1)(\mathbf{y})}{|\mathbf{x} - \mathbf{y}|} d\mathbf{x} d\mathbf{y}.$$

Before stating the main theorem, we first characterise the growing modes for the linearised dynamics.

Proposition 3.15 (Growing modes). Let \mathbf{e}_i (i = 1, 2, 3) be the standard basis of \mathbb{R}^3 . Then \mathbf{e}_i and \mathbf{x} are eigenfunctions for \mathbf{L} with eigenvalue δ and 3δ respectively.

Proof. Let $\mathbf{f} \in \mathbb{R}^3$ be a constant vector. Since $\mathbf{0} = \delta \mathbf{x} + 4\nabla \bar{w} + \nabla \mathcal{K} \bar{w}^3$, we have

$$\mathbf{L}\mathbf{f} = -\nabla\left(\frac{4}{3}\bar{w}^{-2}\nabla\cdot(\bar{w}^{3}\mathbf{f}) + \mathcal{K}\nabla\cdot(\bar{w}^{3}\mathbf{f})\right) = -\nabla\left(\frac{4}{3}\bar{w}^{-2}\mathbf{f}\cdot\nabla\bar{w}^{3} + \mathcal{K}\mathbf{f}\cdot\nabla\bar{w}^{3}\right)$$
$$= -\nabla\left(4\mathbf{f}\cdot\nabla\bar{w} + \mathbf{f}\cdot\nabla\mathcal{K}\bar{w}^{3}\right) = \nabla\left(\delta\mathbf{f}\cdot\mathbf{x}\right) = \delta\mathbf{f}$$

And

$$\mathbf{L}\mathbf{x} = -\nabla\left(\frac{4}{3}\bar{w}^{-2}\nabla\cdot(\bar{w}^{3}\mathbf{x}) + \mathcal{K}\nabla\cdot(\bar{w}^{3}\mathbf{x})\right) = -\nabla\left(4\mathbf{x}\cdot\nabla\bar{w} + \mathcal{K}\mathbf{x}\cdot\nabla\bar{w}^{3} + 4\bar{w} + 3\mathcal{K}\bar{w}^{3}\right)$$
$$= -\nabla\left(4\mathbf{x}\cdot\nabla\bar{w} + \mathbf{x}\cdot\nabla\mathcal{K}\bar{w}^{3} + 4\bar{w} + \mathcal{K}\bar{w}^{3}\right) = \nabla\left(\delta\mathbf{x}\cdot\mathbf{x} - 4\bar{w} - \mathcal{K}\bar{w}^{3}\right) = 2\delta\mathbf{x} + \delta\mathbf{x} = 3\delta\mathbf{x}$$

where we have used

$$\begin{split} \mathscr{K}(\mathbf{x} \cdot \nabla \bar{w}^3)(\mathbf{y}) &= -\int \frac{\mathbf{x} \cdot \nabla \bar{w}(\mathbf{x})^3}{|\mathbf{y} - \mathbf{x}|} d\mathbf{x} = \int \left(\frac{\bar{w}(\mathbf{x})^3 \nabla \cdot \mathbf{x}}{|\mathbf{y} - \mathbf{x}|} + \bar{w}(\mathbf{x})^3 \mathbf{x} \cdot \nabla_{\mathbf{x}} \frac{1}{|\mathbf{y} - \mathbf{x}|} \right) d\mathbf{x} \\ &= \int \left(\frac{3 \bar{w}(\mathbf{x})^3}{|\mathbf{y} - \mathbf{x}|} + \bar{w}(\mathbf{x})^3 \mathbf{x} \cdot \frac{\mathbf{y} - \mathbf{x}}{|\mathbf{y} - \mathbf{x}|^3} \right) d\mathbf{x} = \int \left(\frac{2 \bar{w}(\mathbf{x})^3}{|\mathbf{y} - \mathbf{x}|} + \bar{w}(\mathbf{x})^3 \mathbf{y} \cdot \frac{\mathbf{y} - \mathbf{x}}{|\mathbf{y} - \mathbf{x}|^3} \right) d\mathbf{x} \\ &= -2 \mathscr{K} \bar{w}^3 + \mathbf{y} \cdot \nabla \mathscr{K} \bar{w}^3. \end{split}$$

The main result of this section states that if the perturbation θ is orthogonal to the four eigenvectors from Proposition 3.15, then the operator L is non-negative and we provide a quantitative lower bound.

Theorem 3.16 (Non-negativity of L). Recall that $\bar{w} = \bar{w}_{\delta}$ and L (3.84) depends on δ . There exists $\epsilon > 0$ such that for any $\delta \in (-\epsilon, 0)$ the following holds. If θ is such that $\|\theta\|_3 + \|\nabla\theta\|_4 < \infty$ and

$$\langle \boldsymbol{\theta}, \mathbf{x} \rangle_3 = 0 = \langle \boldsymbol{\theta}, \mathbf{e}_i \rangle_3, \qquad i = 1, 2, 3$$
(3.94)

then we have

$$\langle \mathbf{L}\boldsymbol{\theta}, \boldsymbol{\theta} \rangle_3 \gtrsim \int_{B_R} \bar{w}^{-2} |\Delta \Psi|^2 d\mathbf{x} + \int_{\mathbb{R}^3} |\nabla \Psi|^2 d\mathbf{x}$$
 (3.95)

where the constants do not depend on δ , and Ψ is the gravitational potential induced by the flow disturbance $\bar{w}^3 \theta$:

$$\Psi := \frac{1}{4\pi} \mathscr{K} \nabla \cdot (\bar{w}^3 \theta) \in H^1(\mathbb{R}^3) \cap C^1(\mathbb{R}^3)$$
$$\Delta \Psi = \nabla \cdot (\bar{w}^3 \theta) \in L^2(B_R, \bar{w}^{-2})$$

The proof of Theorem 3.16 is a simple consequence of Lemmas 3.19–3.22. Our strategy is to use spherical harmonics to break down the problem into a sequence of scalar problems for each individual mode, by analogy to [32]. The modes l = 0, 1 correspond to radial and translational motion, and therefore, although formally unstable, can be factored out from the dynamics through suitable orthogonality conditions.

Lemma 3.17 (Spherical harmonics decomposition). Suppose θ is such that $\|\theta\|_3 + \|\nabla\theta\|_4 < \infty$. Then

$$g := \nabla \cdot (\bar{w}^3 \boldsymbol{\theta}) \in L^2(B_R, \bar{w}^{-2})$$
(3.96)

$$\Psi(\mathbf{x}) := \frac{1}{4\pi} \mathscr{K}g(\mathbf{x}) \in H^1(\mathbb{R}^3) \cap C^1(\mathbb{R}^3),$$
(3.97)

and they can be expanded in spherical harmonics

$$g(\mathbf{x}) = \sum_{l=0}^{\infty} \sum_{m=-l}^{l} g_{lm}(r) Y_{lm}(\mathbf{x}) \qquad on \ B_R,$$
(3.98)

$$\Psi(\mathbf{x}) = \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \Psi_{lm}(r) Y_{lm}(\mathbf{x}) \qquad on \ \mathbb{R}^3, \tag{3.99}$$

that converge in $L^2(B_R, \bar{w}^{-2})$ and $L^2(\mathbb{R}^3)$ respectively, where the spherical harmonics Y_{lm} are introduced in Appendix A.2. Moreover, Ψ_{lm} are related to g_{lm} by

$$\Psi_{lm}(r) = \frac{-1}{2l+1} \left(\int_0^r \frac{y^{l+2}}{r^{l+1}} g_{lm}(y) \mathrm{d}y + \int_r^R \frac{r^l}{y^{l-1}} g_{lm}(y) \mathrm{d}y \right)$$
(3.100)

$$g_{lm} = \Delta^{\langle l \rangle} \Psi_{lm} := \left(\frac{1}{r^2} \left(r^2 \Psi'_{lm} \right)' - \frac{l(l+1)}{r^2} \Psi_{lm} \right).$$
(3.101)

With this, the following identity holds:

$$\langle \mathbf{L}\boldsymbol{\theta}, \boldsymbol{\theta} \rangle_3 = \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \Lambda_{lm},$$
(3.102)

where

$$\Lambda_{lm} := \int_0^R \left(\frac{4}{3} \bar{w}^{-2} g_{lm}^2 + 4\pi g_{lm} \Psi_{lm} \right) r^2 \mathrm{d}r, \qquad l \ge 0, \ m \in \{-l, \dots, l\}.$$
(3.103)

Proof. From $\|\boldsymbol{\theta}\|_3 + \|\nabla\boldsymbol{\theta}\|_4 < \infty$, Corollary A.5 of Hardy-Poincaré inequality means that we have $\|\boldsymbol{\theta}\|_2 + \|\nabla\boldsymbol{\theta}\|_4 < \infty$. This immediately gives that $g \in L^2(B_R, \bar{w}^{-2})$. Since Ψ is a convolution of g with the kernel $\|\cdot\|^{-1}$, where g is trivially extended by 0 on $\mathbb{R}^3 \setminus B_R$, standard computation shows $\Psi \in C^1(\mathbb{R}^3) \cap H^1(\mathbb{R}^3)$. Since spherical harmonics form an L^2 basis (see [2, 28, 9] and Appendix A.2), we have the spherical harmonics expansion (3.98)-(3.99) for g and Ψ in L^2 .

By Lemma A.3 we have

$$\frac{1}{|\mathbf{x} - \mathbf{y}|} = 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \frac{1}{2l+1} \frac{\min\{|\mathbf{x}|, |\mathbf{y}|\}^{l}}{\max\{|\mathbf{x}|, |\mathbf{y}|\}^{l+1}} Y_{lm}(\mathbf{y}) Y_{lm}(\mathbf{x})$$

which converge uniformly on all compact set in $\{(\mathbf{x}, \mathbf{y}) : |\mathbf{x}| \neq |\mathbf{y}|\}$. So we have

$$\begin{aligned} \mathscr{K}g(\mathbf{x}) &= -4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \frac{1}{2l+1} Y_{lm}(\mathbf{x}) \left(\int_{B_{|\mathbf{x}|}(\mathbf{0})} \frac{|\mathbf{y}|^{l}}{|\mathbf{x}|^{l+1}} g Y_{lm} d\mathbf{y} + \int_{B_{|\mathbf{x}|}(\mathbf{0})^{c}} \frac{|\mathbf{x}|^{l}}{|\mathbf{y}|^{l+1}} g Y_{lm} d\mathbf{y} \right) \\ &= -4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \frac{1}{2l+1} Y_{lm}(\mathbf{x}) \left(\int_{0}^{|\mathbf{x}|} \frac{y^{l+2}}{|\mathbf{x}|^{l+1}} g_{lm} dy + \int_{|\mathbf{x}|}^{R} \frac{|\mathbf{x}|^{l}}{y^{l-1}} g_{lm} dy \right). \end{aligned}$$

We therefore conclude that

$$\Psi_{lm}(r) = \frac{-1}{2l+1} \left(\int_0^r \frac{y^{l+2}}{r^{l+1}} g_{lm}(y) \mathrm{d}y + \int_r^R \frac{r^l}{y^{l-1}} g_{lm}(y) \mathrm{d}y \right)$$

since spherical harmonics expansion is unique (using standard Hilbert space theory and the fact that spherical harmonics forms a L^2 basis for L^2 functions on the sphere). Inverting this expression, we get (3.101).

Now using the spherical harmonics expansion for g and Ψ , we get

$$egin{aligned} \langle \mathbf{L}m{ heta},m{ heta}
angle_3 &= \int \left(rac{4}{3} ar{w}^{-2} |
abla \cdot (ar{w}^3m{ heta})|^2 +
abla \cdot (ar{w}^3m{ heta}) \mathcal{K}
abla \cdot (ar{w}^3m{ heta})
ight) \mathrm{d} \mathbf{x} \ &= \int \left(rac{4}{3} ar{w}^{-2} |g|^2 + 4\pi g \Psi
ight) \mathrm{d} \mathbf{x} = \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \Lambda_{lm}, \end{aligned}$$

with Λ_{lm} as in (3.103).

From [24] we have the following lemma.

Lemma 3.18. There exists $\epsilon > 0$ such that for any $\delta \in (-\epsilon, 0)$ and the associated $\bar{w} = \bar{w}_{\delta}$, we have

 $\langle \mathscr{L}\varphi,\varphi\rangle_{\bar{w}^3r^4}\gtrsim \|\varphi'\|_{\bar{w}^4r^4}^2+\|\varphi\|_{\bar{w}^3r^4}^2\qquad \text{whenever}\qquad \langle\varphi,1\rangle_{\bar{w}^3r^4}=0$

where the constants do not depend on δ , and

$$\begin{aligned} \mathscr{L}\varphi &:= -\frac{4}{3\bar{w}^3r^4}\partial_r\left(\bar{w}^4r^4\partial_r\varphi\right) + 3\delta\varphi\\ \langle f,g\rangle_{\bar{w}^kr^4} &:= \int_0^R f(r)g(r)\bar{w}(r)^kr^4\mathrm{d}r. \end{aligned}$$

We shall use Lemma 3.18 to obtain coercivity for the quadratic form Λ_{00} under the orthogonality assumption $\langle \boldsymbol{\theta}, \mathbf{x} \rangle_3 = 0$.

Lemma 3.19 (l = 0 mode bound). Suppose θ is as in Lemma 3.17 and $\langle \theta, \mathbf{x} \rangle_3 = 0$. Then we have $\Psi'_{00}(r) = 0$ for $r \ge R$ and

$$\Lambda_{00} \gtrsim \int_0^R \left(\bar{w}^{-2} g_{00}^2 + (\Psi_{00}')^2 \right) r^2 \mathrm{d}r, \tag{3.104}$$

where we recall (3.103).

Proof. From $\|\boldsymbol{\theta}\|_3 + \|\nabla\boldsymbol{\theta}\|_4 < \infty$, Corollary A.5 of Hardy-Poincaré inequality means that we have $\|\boldsymbol{\theta}\|_2 + \|\nabla\boldsymbol{\theta}\|_4 < \infty$. It follows that $\bar{w}^3\boldsymbol{\theta}$ is well defined on ∂B_R (trace theorem) and must vanish there. Since $\bar{w}^3\boldsymbol{\theta} = \nabla\Psi + \mathbf{C}$ where \mathbf{C} is divergence-free, we have

$$\int_{\partial B_R} \partial_r \Psi \, \mathrm{d}S = \int_{\partial B_R} \nabla \Psi \cdot \mathrm{d}\mathbf{S} = \int_{\partial B_R} \bar{w}^3 \boldsymbol{\theta} \cdot \mathrm{d}\mathbf{S} = 0.$$

It follows that $\Psi'_{00}(R) = 0$. Now taking the derivative of (3.100) and using $g_{00}(r) = 0$ for r > R, we see that in fact we must have

$$\Psi'_{00}(r) = 0$$
 for $r \ge R.$ (3.105)

From the orthogonality condition $\langle \boldsymbol{\theta}, \mathbf{x} \rangle_3 = 0$ we infer that

$$0 = \langle \boldsymbol{\theta}, \mathbf{x} \rangle_3 = \frac{1}{2} \int \bar{w}^3 \boldsymbol{\theta} \cdot \nabla |\mathbf{x}|^2 \, \mathrm{d}\mathbf{x} = \frac{1}{2} \int g |\mathbf{x}|^2 \, \mathrm{d}\mathbf{x}.$$

This means

$$\int_0^R g_{00}(r) r^4 \mathrm{d}r = 0. \tag{3.106}$$

and therefore by (3.101) and (3.105) in terms of Ψ_{00} ,

$$\int_{0}^{R} \Psi_{00}'(r) r^{3} \mathrm{d}r = 0.$$
(3.107)

Since $\Psi \in H^1(\mathbb{R}^3) \cap C^1(\mathbb{R}^3)$, we have that $\partial_r \Psi \in L^2(\mathbb{R}^3) \cap C(\mathbb{R}^3 \setminus \{\mathbf{0}\})$. So $\partial_r \Psi$ has spherical harmonics expansion $\partial_r \Psi = \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \Psi_{r,lm} Y_{lm}$ in $L^2(\mathbb{R}^3)$ with

$$\Psi_{r,lm}(r) = \frac{1}{4\pi r^2} \int_{\partial B_r} (\partial_r \Psi) Y_{lm} \mathrm{d}S = \frac{1}{4\pi r^2} \partial_r \int_{B_r} \Psi Y_{lm} \mathrm{d}S = \partial_r \Psi_{lm}(r) = \Psi'_{lm}(r). \tag{3.108}$$

If we denote

$$\varphi := \Psi_{00}'/(r\bar{w}^3), \tag{3.109}$$

then by (3.107) we have

$$0 = \int_0^R \Psi'_{00}(r) r^3 \mathrm{d}r = \int_0^R \varphi(r) \bar{w}^3 r^4 \mathrm{d}r$$

and thus $\langle \varphi, 1 \rangle_{\bar{w}^3 r^4} = 0$. Using (3.101) and (3.105), we get

$$\begin{split} \Lambda_{00} &= \int_0^R \left(\frac{4}{3} \bar{w}^{-2} g_{00}^2 + 4\pi g_{00} \Psi_{00} \right) r^2 \mathrm{d}r = \int_0^R \left(\frac{4}{3r^2} \bar{w}^{-2} \left(\left(r^2 \Psi_{00}' \right)' \right)^2 + 4\pi \left(r^2 \Psi_{00}' \right)' \Psi_{00} \right) \mathrm{d}r \\ &= \int_0^R \left(\frac{4}{3r^2} \bar{w}^{-2} \left(\left(r^2 \Psi_{00}' \right)' \right)^2 - 4\pi r^2 (\Psi_{00}')^2 \right) \mathrm{d}r + 4\pi R^2 \Psi_{00}' (R) \Psi_{00}(R) \\ &= \int_0^R \left(\frac{4}{3r^2} \bar{w}^{-2} \left(\left(r^3 \bar{w}^3 \varphi \right)' \right)^2 - 4\pi \varphi^2 \bar{w}^6 r^4 \right) \mathrm{d}r \end{split}$$

Now since $0 = 3\delta + 4\Delta \bar{w} + 4\pi \bar{w}^3$ as in (3.63), we see that

$$\begin{split} \Lambda_{00} &= \int_{0}^{R} \left(\frac{4}{3r^{2}} \bar{w}^{-2} \left(\left(r^{3} \bar{w}^{3} \varphi \right)' \right)^{2} + (3\delta + 4\Delta \bar{w}) \varphi^{2} \bar{w}^{3} r^{4} \right) \mathrm{d}r \\ &= \int_{0}^{R} \left(\frac{4}{3r^{2}} \bar{w}^{-2} \left(3r^{2} \bar{w}^{3} \varphi + 3r^{3} \bar{w}^{2} \bar{w}' \varphi + r^{3} \bar{w}^{3} \varphi' \right)^{2} + 4(r^{2} \bar{w}')' \varphi^{2} \bar{w}^{3} r^{2} + 3\delta \varphi^{2} \bar{w}^{3} r^{4} \right) \mathrm{d}r \end{split}$$

$$\begin{split} &= \int_0^R \left(\frac{4}{3} \left(3r\bar{w}^2 \varphi + 3r^2 \bar{w}\bar{w}'\varphi + r^2 \bar{w}^2 \varphi' \right)^2 - 4r^2 \bar{w}' (\varphi^2 \bar{w}^3 r^2)' + 3\delta \varphi^2 \bar{w}^3 r^4 \right) \mathrm{d}r \\ &= \int_0^R \left(\frac{4}{3} \left(9r^2 \bar{w}^4 \varphi^2 + 9r^4 \bar{w}^2 (\bar{w}')^2 \varphi^2 + r^4 \bar{w}^4 (\varphi')^2 + 18r^3 \bar{w}^3 \bar{w}' \varphi^2 + 6r^4 \bar{w}^3 \bar{w}' \varphi \varphi' + 6r^3 \bar{w}^4 \varphi \varphi' \right) \\ &\quad - 4(2\varphi\varphi' \bar{w}^3 \bar{w}' r^4 + 3\varphi^2 \bar{w}^2 (\bar{w}')^2 r^4 + 2\varphi^2 \bar{w}^3 \bar{w}' r^3) + 3\delta \varphi^2 \bar{w}^3 r^4 \right) \mathrm{d}r \\ &= \int_0^R \left(\frac{4}{3} \left(9r^2 \bar{w}^4 \varphi^2 + r^4 \bar{w}^4 (\varphi')^2 + 12r^3 \bar{w}^3 \bar{w}' \varphi^2 + 6r^3 \bar{w}^4 \varphi \varphi' \right) + 3\delta \varphi^2 \bar{w}^3 r^4 \right) \mathrm{d}r \\ &= \int_0^R \left(\frac{4}{3} r^4 \bar{w}^4 (\varphi')^2 + 3\delta \varphi^2 \bar{w}^3 r^4 \right) \mathrm{d}r = \langle \mathscr{L}\varphi, \varphi \rangle_{\bar{w}^3 r^4} \\ &\gtrsim \|\varphi'\|_{\bar{w}^4 r^4}^2 + \|\varphi\|_{\bar{w}^3 r^4}^2 = \int_0^R ((\varphi')^2 \bar{w}^4 + \varphi^2 \bar{w}^3) r^4 \mathrm{d}r \end{split}$$

Then for ϵ small enough, we get

$$\begin{split} \Lambda_{00} \gtrsim & \int_{0}^{R} \left(\epsilon \left(\frac{4}{3} r^{4} \bar{w}^{4} (\varphi')^{2} + 3\delta \varphi^{2} \bar{w}^{3} r^{4} \right) + \varphi^{2} \bar{w}^{3} r^{4} \right) \mathrm{d}r \\ & = \int_{0}^{R} \left(\epsilon \left(\frac{4}{3} \bar{w}^{-2} g_{00}^{2} - 4\pi (\Psi_{00}')^{2} \right) + \bar{w}^{-3} (\Psi_{00}')^{2} \right) r^{2} \mathrm{d}r. \end{split}$$

Finally by choosing ϵ small enough we get (3.104).

In order to prove positivity of the higher modes, we will need the following lemma which provides an estimate from below for Λ_{lm} by an elliptic operator; a related bound was also used in [32].

Lemma 3.20. Suppose θ is as in Lemma 3.17. Then for any $l \ge 0, m \in \{-l, \ldots, l\}$, we have

$$\Lambda_{lm} \ge 4\pi \int_0^R \left(-\Delta^{\langle l \rangle} - 3\pi \bar{w}^2 \right) (\Psi_{lm}) \Psi_{lm} r^2 \mathrm{d}r.$$

Proof. We have

$$\begin{split} \Lambda_{lm} &= \int_0^R \left(\frac{4}{3} \bar{w}^{-2} g_{lm}^2 + 4\pi g_{lm} \Psi_{lm} \right) r^2 \mathrm{d}r \\ &= \int_0^R \left| \frac{2}{\sqrt{3}\bar{w}} g_{lm} + 2\pi \sqrt{3} \bar{w} \Psi_{lm} \right|^2 r^2 \mathrm{d}r - 4\pi \int_0^R \left(g_{lm} \Psi_{lm} + 3\pi \bar{w}^2 \Psi_{lm}^2 \right) r^2 \mathrm{d}r \\ &\geq 4\pi \int_0^R \left(-\Delta^{\langle l \rangle} - 3\pi \bar{w}^2 \right) (\Psi_{lm}) \Psi_{lm} r^2 \mathrm{d}r. \end{split}$$

With this bound from below by an elliptic operator, we can prove the positivity of Λ_{lm} using elliptic ODE theory.

Lemma 3.21 (l = 1 modes bound). Suppose θ is as in Lemma 3.17 and $\langle \theta, \mathbf{e}_i \rangle_3 = 0$ for i = 1, 2, 3. Then we have $\Psi_{1m}(r) = 0$ for $r \ge R$ and

$$\Lambda_{1m} \gtrsim \int_0^R \left(\bar{w}^{-2} g_{1m}^2 r^2 + \Psi_{1m}'^2 r^2 + \Psi_{1m}^2 \right) \mathrm{d}r, \qquad m = -1, 0, 1, \tag{3.110}$$

where we recall (3.103).

Proof. For these modes, we adapt the method of proof as found in [32] that makes use of the Sturm-Liouville theory. We have by Lemma 3.20

$$\Lambda_{1m} \ge 4\pi \langle A_1 \Psi_{1m}, \Psi_{1m} \rangle_{r^2}$$

where $\langle y_1, y_2 \rangle_{r^2} := \int_0^R y_1 y_2 r^2 \mathrm{d}r$ and

$$A_1 := -\Delta^{\langle 1 \rangle} - 3\pi \bar{w}^2. \tag{3.111}$$

As this operator A_1 resembles the operator A analyzed in [32] (cf. (7.15) of [32]), by arguing analogously, we deduce that the operator A_1 has the Friedrichs extension in the Hilbert space induced by the inner product $\langle y_1, y_2 \rangle_{r^2}$, denoted by the same A_1 . Moreover it is of Sturm-Liouville type and the eigenvalues are simple under the Dirichlet boundary condition on r = R, i.e. y(R) = 0 (cf. Section VII of [32]).

We next claim the least eigenvalue μ_1 of A_1 is strictly positive. Let ϕ_1 be an associated eigenfunction such that $A_1\phi_1 = \mu_1\phi_1$. Since ϕ_1 must have no zeros on (0, R) by Sturm-Liouville theory, we may assume that $\phi_1(r) > 0$ for $r \in (0, R)$ so that $\phi'_1(R) \leq 0$ and $\phi_1(R) = 0$. In fact we must have $\phi'_1(R) < 0$, for if $\phi'_1(R) = 0$, then ϕ_1 must be the zero function, which is a contradiction. To see the latter assertion, note that A_1 is a second order ODE operator with C^1 coefficients away from the origin. Picard-Lindelöf existence theorem implies that for any $\epsilon > 0$ the solution u on $(\epsilon, R]$ satisfying u'(R) = u(R) = 0 must be unique. Since u = 0 is such a solution, we must have $\phi'_1 = u = 0$. On the other hand, recalling $\Delta(4\bar{w}) = -3\delta - 4\pi\bar{w}^3$, we see that $A_1\bar{w}' = 0$. Note that $\bar{w}'(R) \neq 0$, so $\bar{w}' \notin \text{Dom } A_1$ where $\text{Dom } A_1$ denotes the domain of A_1 under the Sturm-Liouville theory framework. By using $A_1\bar{w}' = 0$, the properties of ϕ_1 and integration by parts, we have

$$0 = \langle A_1 \bar{w}', \phi_1 \rangle_{r^2} = \langle \bar{w}', A_1 \phi_1 \rangle_{r^2} + R^2 \bar{w}'(R) \phi_1'(R) = \mu_1 \langle \bar{w}', \phi_1 \rangle_{r^2} + R^2 \bar{w}'(R) \phi_1'(R).$$

Since $\bar{w}'(r) < 0$ for $r \in (0, R]$, we see that $\langle \bar{w}', \phi_1 \rangle_{r^2} < 0$. Also $R^2 \bar{w}'(R) \phi_1'(R) > 0$. Therefore we must have $\mu_1 > 0$.

By the orthogonality condition

$$0 = \langle \boldsymbol{\theta}, \mathbf{e}_i \rangle_3 = \int_{B_R} \bar{w}^3 \boldsymbol{\theta} \cdot \nabla x^i \, \mathrm{d}\mathbf{x} = \int_{B_R} g x^i \, \mathrm{d}\mathbf{x},$$

we conclude that $0=\int_0^R g_{1m}r^3\mathrm{d}r$ and therefore

$$\Psi_{1m}(r) = 0$$
 for $r \ge R.$ (3.112)

This (3.112) means that $\Psi_{1m} \in \text{Dom } A_1$, it follows that

$$\Lambda_{1m} \ge 4\pi \langle A_1 \Psi_{1m}, \Psi_{1m} \rangle_{r^2} \ge 4\pi \mu_1 \langle \Psi_{1m}, \Psi_{1m} \rangle_{r^2} \ge 0.$$
(3.113)

The second inequality of (3.113) implies

$$\int_0^R \left(\Psi_{1m}^{\prime 2} + \frac{2}{r^2} \Psi_{1m}^2 - 3\pi \bar{w}^2 \Psi_{1m}^2 \right) r^2 \mathrm{d}r \ge \mu_1 \int_0^R \Psi_{1m}^2 r^2 \mathrm{d}r$$

which we can rewrite as

$$(1+\epsilon) \int_0^R \left(\Psi_{1m}^{\prime 2} + \frac{2}{r^2} \Psi_{1m}^2 - 3\pi \bar{w}^2 \Psi_{1m}^2 \right) r^2 \mathrm{d}r$$

$$\geq \epsilon \int_0^R \left(\Psi_{1m}^{\prime 2} + \frac{2}{r^2} \Psi_{1m}^2 - 3\pi \bar{w}^2 \Psi_{1m}^2 \right) r^2 \mathrm{d}r + \mu_1 \int_0^R \Psi_{1m}^2 r^2 \mathrm{d}r$$

$$\geq \epsilon \int_0^R \left(\Psi_{1m}^{\prime 2} r^2 + 2\Psi_{1m}^2 \right) \mathrm{d}r + (\mu_1 - 3\epsilon\pi \bar{w}(0)^2) \int_0^R \Psi_{1m}^2 r^2 \mathrm{d}r$$

Chose ϵ small enough so that the last term is non-negative. Hence we see that

$$\int_0^R \left(\Psi_{1m}^{\prime 2} + \frac{2}{r^2} \Psi_{1m}^2 - 3\pi \bar{w}^2 \Psi_{1m}^2 \right) r^2 \mathrm{d}r \gtrsim \int_0^R \left(\Psi_{1m}^{\prime 2} r^2 + \Psi_{1m}^2 \right) \mathrm{d}r$$

Together with (3.113) we deduce that

$$\Lambda_{1m} = \int_0^R \left(\frac{4}{3}\bar{w}^{-2}g_{1m}^2 + 4\pi g_{1m}\Psi_{1m}\right)r^2 \mathrm{d}r \gtrsim \int_0^R \left(\Psi_{1m}'^2 r^2 + \Psi_{1m}^2\right) \mathrm{d}r$$

We can rewrite this as, for some C > 0,

$$\begin{split} (1+\epsilon)\Lambda_{1m} &\geq \epsilon \int_0^R \left(\frac{4}{3}\bar{w}^{-2}g_{1m}^2 + 4\pi g_{1m}\Psi_{1m}\right)r^2\mathrm{d}r + C\int_0^R \left(\Psi_{1m}'^2r^2 + \Psi_{1m}^2\right)\mathrm{d}r \\ &= \epsilon \int_0^R \left(\frac{4}{3}\bar{w}^{-2}g_{1m}^2r^2 + 4\pi \left((r^2\Psi_{1m}')' - 2\Psi_{1m}\right)\Psi_{1m}\right)\mathrm{d}r + C\int_0^R \left(\Psi_{1m}'^2r^2 + \Psi_{1m}^2\right)\mathrm{d}r \\ &= \epsilon \int_0^R \left(\frac{4}{3}\bar{w}^{-2}g_{1m}^2r^2 - 4\pi \left(\Psi_{1m}'^2r^2 + 2\Psi_{1m}^2\right)\right)\mathrm{d}r + C\int_0^R \left(\Psi_{1m}'^2r^2 + \Psi_{1m}^2\right)\mathrm{d}r \end{split}$$

Choosing ϵ small enough we obtain (3.110).

Lemma 3.22 $(l \ge 2 \text{ modes bound})$. Suppose θ is as in Lemma 3.17. Then for $l \ge 2$,

$$\Lambda_{lm} \gtrsim \int_0^R \bar{w}^{-2} g_{lm}^2 r^2 \mathrm{d}r + \int_0^\infty \left(\Psi_{lm}'^2 r^2 + l(l+1) \Psi_{lm}^2 \right) \mathrm{d}r, \qquad m \in \{-l, \dots, l\}.$$
(3.114)

Proof. For these higher modes, we use a continuity argument. We have by Lemma 3.20

$$\begin{split} \Lambda_{lm} &\geq 4\pi \int_{0}^{R} \left(-\Delta^{\langle l \rangle} - 3\pi \bar{w}^{2} \right) (\Psi_{lm}) \Psi_{lm} r^{2} \mathrm{d}r \\ &= 4\pi \int_{0}^{R} \left(\Psi_{lm}^{\prime 2} + \frac{l(l+1)}{r^{2}} \Psi_{lm}^{2} - 3\pi \bar{w}^{2} \Psi_{lm}^{2} \right) r^{2} \mathrm{d}r - 4\pi R^{2} \Psi_{lm} (R) \Psi_{lm}^{\prime} (R) \\ &= 4\pi \int_{0}^{R} \left(\Psi_{lm}^{\prime 2} + \frac{l(l+1)}{r^{2}} \Psi_{lm}^{2} - 3\pi \bar{w}^{2} \Psi_{lm}^{2} \right) r^{2} \mathrm{d}r + 4\pi \left(\int_{R}^{\infty} \left(\Psi_{lm}^{\prime 2} r^{2} + (r^{2} \Psi_{lm}^{\prime})^{\prime} \Psi_{lm} \right) \mathrm{d}r \right) \\ &= 4\pi \int_{0}^{\infty} \left(\Psi_{lm}^{\prime 2} + \frac{l(l+1)}{r^{2}} \Psi_{lm}^{2} - 3\pi \bar{w}^{2} \Psi_{lm}^{2} \right) r^{2} \mathrm{d}r \end{split}$$
(3.115)

$$=4\pi \int_{0}^{\infty} \left(-\Delta^{\langle 1\rangle} - 3\pi \bar{w}^{2}\right) (\Psi_{lm}) \Psi_{lm} r^{2} \mathrm{d}r + 4\pi \int_{0}^{\infty} (l(l+1)-2) \Psi_{lm}^{2} \mathrm{d}r$$
(3.116)

where we used

$$g_{lm}(r) = \Delta^{\langle l \rangle} \Psi_{lm}(r) = \frac{1}{r^2} (r^2 \Psi'_{lm})' - \frac{l(l+1)}{r^2} \Psi_{lm} = 0 \quad \text{for} \quad r > R.$$

Recall that $\bar{w} = \bar{w}_{\delta}$ depends on δ . In the proof of Lemma 3.21 we have shown that

$$\int_{0}^{R} \left(-\Delta^{\langle 1 \rangle} - 3\pi \bar{w}_{\delta}^{2} \right) (y) y r^{2} \mathrm{d}r \geq 0$$

for all $y \in H^2([0,R],r^2)$ such that y(R) = 0. In fact when $\bar{w}_{\delta} = \bar{w}_0$ (the Lane-Emden star), the same analysis can be extended to any $R' \ge R$ to give rise to

$$\int_{0}^{R'} \left(-\Delta^{\langle 1 \rangle} - 3\pi \bar{w}_0^2 \right) (y) y r^2 \mathrm{d}r \ge 0 \tag{3.117}$$

for all $y \in H^2([0, R'], r^2)$ such that y(R') = 0. To do so, we replace \bar{w}'_0 (used to argue the non-negativity of the least eigenvalue) with \tilde{w}' , where $\tilde{w} := -\frac{1}{4} \mathcal{K} \bar{w}^3_0$ (recall $\bar{w}_0 = 0$ for r > R). Note that \tilde{w} is $C^3(\mathbb{R}^3)$, or $C^3([0, \infty))$ as a function of the radial variable. By (1.18), we see that $\tilde{w}' = \bar{w}'_0$ on [0, R]. Moreover, $\tilde{w}' < 0$ on $(0, \infty)$. Since $\Delta \tilde{w} = -\pi \bar{w}^3_0$, taking ∂_r we get $\Delta^{\langle 1 \rangle} \tilde{w} = -3\pi \bar{w}^2_0 \bar{w}'_0 = -3\pi \bar{w}^2_0 \tilde{w}'$. So we have $(-\Delta^{\langle 1 \rangle} - 3\pi \bar{w}^2_0) \tilde{w}' = 0$ on $[0, \infty)$ which allows us to apply the same proof in Lemma 3.21.

Let

$$y_{R'}(r) = \Psi_{lm}(r) - \Psi_{lm}(R) \left(\frac{R}{R'}\right)^{l+1} \frac{r}{R'}$$

From (3.100) we see that $y_{R'}(R') = 0$. By using $\Delta^{\langle 1 \rangle} r = 0$ and applying (3.117) with $y = y_{R'}$, we obtain

$$\begin{split} &\int_{0}^{R'} \left(-\Delta^{\langle 1 \rangle} - 3\pi \bar{w}_{0}^{2} \right) (\Psi_{lm}) \Psi_{lm} r^{2} \mathrm{d}r \\ &= \int_{0}^{R'} \left(-\Delta^{\langle 1 \rangle} - 3\pi \bar{w}_{0}^{2} \right) (y_{R'}(r)) \left(y_{R'}(r) + \Psi_{lm}(R) \left(\frac{R}{R'} \right)^{l+1} \frac{r}{R'} \right) r^{2} \mathrm{d}r \\ &- \int_{0}^{R} 3\pi \bar{w}_{0}^{2} \Psi_{lm}(r) \Psi_{lm}(R) \left(\frac{R}{R'} \right)^{l+1} \frac{r}{R'} r^{2} \mathrm{d}r \\ &\geq \int_{0}^{R'} \left(-\Delta^{\langle 1 \rangle} - 3\pi \bar{w}_{0}^{2} \right) (y_{R'}(r)) \Psi_{lm}(R) \left(\frac{R}{R'} \right)^{l+1} \frac{r^{3}}{R'} \mathrm{d}r - \int_{0}^{R} 3\pi \bar{w}_{0}^{2} \Psi_{lm}(r) \Psi_{lm}(R) \left(\frac{R}{R'} \right)^{l+1} \frac{r^{3}}{R'} \mathrm{d}r \\ &\geq \int_{0}^{R'} \left(-\Delta^{\langle 1 \rangle} - 3\pi \bar{w}_{0}^{2} \right) (y_{R'}(r)) \Psi_{lm}(R) \left(\frac{R}{R'} \right)^{l+1} \frac{r^{3}}{R'} \mathrm{d}r - \int_{0}^{R} 3\pi \bar{w}_{0}^{2} \Psi_{lm}(r) \Psi_{lm}(R) \left(\frac{R}{R'} \right)^{l+1} \frac{r^{3}}{R'} \mathrm{d}r \end{split}$$

Denote the last two integral terms by K. By integrating by parts and using the boundary condition $y_{R'}(R') = 0$,

$$\begin{split} K &= -R'^2 y'_{R'}(R') \Psi_{lm}(R) \left(\frac{R}{R'}\right)^{l+1} - \int_0^R 3\pi \bar{w}_0^2 y_{R'}(r) \Psi_{lm}(R) \left(\frac{R}{R'}\right)^{l+1} \frac{r^3}{R'} \mathrm{d}r \\ &- \int_0^R 3\pi \bar{w}_0^2 \Psi_{lm}(r) \Psi_{lm}(R) \left(\frac{R}{R'}\right)^{l+1} \frac{r^3}{R'} \mathrm{d}r \\ &= -R'^2 \Psi'_{lm}(R') \Psi_{lm}(R) \left(\frac{R}{R'}\right)^{l+1} + R' (\Psi_{lm}(R))^2 \left(\frac{R}{R'}\right)^{2l+2} \\ &+ 3\pi \int_0^R \bar{w}_0^2 (\Psi_{lm}(R))^2 \left(\frac{R}{R'}\right)^{2l+2} \frac{r^4}{R'^2} \mathrm{d}r - 6\pi \int_0^R \bar{w}_0^2 \Psi_{lm}(r) \Psi_{lm}(R) \left(\frac{R}{R'}\right)^{l+1} \frac{r^3}{R'} \mathrm{d}r \\ &\to 0 \qquad \text{as} \qquad R' \to \infty \end{split}$$

when $l \ge 1$, where we used (3.100) to see for example that $\Psi'_{lm}(R') \to 0$ as $R' \to \infty$.

Therefore we have proven¹ that for any $l \ge 1$,

$$\int_{0}^{\infty} \left(\Psi_{lm}^{\prime 2} + \frac{l(l+1)}{r^{2}} \Psi_{lm}^{2} - 3\pi \bar{w}_{0}^{2} \Psi_{lm}^{2} \right) r^{2} \mathrm{d}r \geq \int_{0}^{\infty} (l(l+1) - 2) \Psi_{lm}^{2} \mathrm{d}r \quad \text{for all} \quad \Psi_{lm}. \quad (3.118)$$

So we have

$$\begin{split} \int_{0}^{\infty} \left(\Psi_{lm}^{\prime 2} + \frac{l(l+1)}{r^{2}} \Psi_{lm}^{2} - 3\pi \bar{w}_{\delta}^{2} \Psi_{lm}^{2} \right) r^{2} \mathrm{d}r \\ & \geq \underbrace{\int_{0}^{\infty} (l(l+1) - 2) \Psi_{lm}^{2} \mathrm{d}r - 3\pi \left\| (\bar{w}_{\delta}^{2} - \bar{w}_{0}^{2}) r^{2} \right\|_{L^{\infty}} \int_{0}^{\infty} \Psi_{lm}^{2} \mathrm{d}r}_{=M} \\ & = M \end{split}$$

For sufficiently small δ we have

$$M \ge (l(l+1)-3) \int_0^\infty \Psi_{lm}^2 \mathrm{d}r$$

which leads to

$$\Lambda_{lm} \geq 4\pi (l(l+1)-3) \int_0^\infty \Psi_{lm}^2 \mathrm{d} r \geq 0$$

¹The proof of (3.118) can be easily adapted to correct an inconsistency appearing in [32] and establish the non-negativity of the quadratic form $\langle \mathbf{L}\boldsymbol{\theta}, \boldsymbol{\theta} \rangle_3$ around the Lane-Emden stars.

Observe that

$$(1+\epsilon) \int_0^\infty \left(\Psi_{lm}^{\prime 2} + \frac{l(l+1)}{r^2} \Psi_{lm}^2 - 3\pi \bar{w}_{\delta}^2 \Psi_{lm}^2 \right) r^2 \mathrm{d}r \\ \ge \epsilon \int_0^\infty \left(\Psi_{lm}^{\prime 2} + \frac{l(l+1)}{r^2} \Psi_{lm}^2 - 3\pi \bar{w}_{\delta}^2 \Psi_{lm}^2 \right) r^2 \mathrm{d}r + (l(l+1)-3) \int_0^\infty \Psi_{lm}^2 \mathrm{d}r.$$

Choosing $\epsilon>0$ small enough we see that

$$\begin{split} \int_0^\infty \left(\Psi_{lm}'^2 + \frac{l(l+1)}{r^2} \Psi_{lm}^2 - 3\pi \bar{w}_{\delta}^2 \Psi_{lm}^2 \right) r^2 \mathrm{d}r \gtrsim \int_0^\infty \left(\Psi_{lm}'^2 r^2 + (l(l+1) - 4) \Psi_{lm}^2 \right) \mathrm{d}r \\ \gtrsim \int_0^\infty \left(\Psi_{lm}'^2 r^2 + l(l+1) \Psi_{lm}^2 \right) \mathrm{d}r. \end{split}$$

We have

$$\Lambda_{lm} = \int_0^R \left(\frac{4}{3}\bar{w}^{-2}g_{lm}^2 + 4\pi g_{lm}\Psi_{lm}\right)r^2 \mathrm{d}r \gtrsim \int_0^\infty \left(\Psi_{lm}'^2 r^2 + l(l+1)\Psi_{lm}^2\right)\mathrm{d}r.$$

We can rewrite this as, for some C > 0,

$$\begin{split} (1+\epsilon)\Lambda_{lm} &\geq \epsilon \int_0^R \left(\frac{4}{3}\bar{w}^{-2}g_{lm}^2 + 4\pi g_{lm}\Psi_{lm}\right)r^2\mathrm{d}r + C\int_0^\infty \left(\Psi_{lm}'^2r^2 + l(l+1)\Psi_{lm}^2\right)\mathrm{d}r\\ &= \epsilon \int_0^R \left(\frac{4}{3}\bar{w}^{-2}g_{lm}^2r^2 + 4\pi \left((r^2\Psi_{lm}')' - l(l+1)\Psi_{lm}\right)\Psi_{lm}\right)\mathrm{d}r\\ &+ C\int_0^\infty \left(\Psi_{lm}'^2r^2 + l(l+1)\Psi_{lm}^2\right)\mathrm{d}r\\ &= \frac{4}{3}\epsilon \int_0^R \bar{w}^{-2}g_{lm}^2r^2\mathrm{d}r + 4\pi\epsilon \int_0^\infty \left(\Psi_{lm}'^2r^2 - l(l+1)\Psi_{lm}^2\right)\mathrm{d}r\\ &+ C\int_0^\infty \left(\Psi_{lm}'^2r^2 + l(l+1)\Psi_{lm}^2\right)\mathrm{d}r \end{split}$$

where we used the fact that

$$4\pi \int_0^R \left(\Psi_{lm}^{\prime 2} + \frac{l(l+1)}{r^2} \Psi_{lm}^2 \right) r^2 dr - 4\pi R^2 \Psi_{lm}(R) \Psi_{lm}^{\prime}(R) = 4\pi \int_0^\infty \left(\Psi_{lm}^{\prime 2} + \frac{l(l+1)}{r^2} \Psi_{lm}^2 \right) r^2 dr$$

oved in (3.115). Choosing $\epsilon > 0$ small enough we obtain the desired (3.114).

proved in (3.115). Choosing $\epsilon > 0$ small enough we obtain the desired (3.114).

Proof of Theorem 3.16. Combining all the bounds we have for each l, m from Lemmas 3.19–3.22, we have

$$\langle \mathbf{L}\boldsymbol{\theta}, \boldsymbol{\theta} \rangle_{3} = \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \Lambda_{lm} \gtrsim \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \int_{0}^{R} \bar{w}^{-2} g_{lm}^{2} r^{2} dr + \int_{0}^{\infty} \left(\Psi_{lm}^{\prime 2} r^{2} + l(l+1) \Psi_{lm}^{2} \right) dr.$$

We know

$$\int_{B_R} \bar{w}^{-2} |\Delta \Psi|^2 \mathrm{d}\mathbf{x} = 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^l \int_0^R \bar{w}^{-2} g_{lm}^2 r^2 \mathrm{d}r.$$

It remains to show that

$$\int_{\mathbb{R}^3} |\nabla \Psi|^2 d\mathbf{x} = 4\pi \sum_{l=0}^\infty \sum_{m=-l}^l \int_0^\infty \left(\Psi_{lm}'^2 r^2 + l(l+1)\Psi_{lm}^2 \right) dr$$
(3.119)

Since $\nabla \Psi \in L^2(\mathbb{R}^3)^3$, it has a vector spherical harmonics expansion in $L^2(\mathbb{R}^3)^3$ [3, 14],

$$\nabla \Psi = \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \left(\Psi_{lm}^{[0]} \mathbf{Y}_{lm}^{[0]} + \Psi_{lm}^{[1]} \mathbf{Y}_{lm}^{[1]} + \Psi_{lm}^{[2]} \mathbf{Y}_{lm}^{[2]} \right).$$
(3.120)

where

$$\mathbf{Y}_{lm}^{[0]} = Y_{lm}\hat{\mathbf{r}}, \qquad \mathbf{Y}_{lm}^{[1]} = r\nabla Y_{lm}, \qquad \mathbf{Y}_{lm}^{[2]} = \mathbf{r} \times \nabla Y_{lm}$$

are the vector spherical harmonics [3, 14]. We have

$$\Psi_{lm}^{[0]}(r) = \frac{1}{r^2} \int_{\partial B_r} \nabla \Psi \cdot \mathbf{Y}_{lm}^{[0]} \mathrm{d}S = \frac{1}{r^2} \int_{\partial B_r} (\partial_r \Psi) Y_{lm} \mathrm{d}S = \Psi_{lm}'(r)$$

using (3.108). And

$$\begin{split} \Psi_{lm}^{[1]}(r) &= \frac{1}{l(l+1)r^2} \int_{\partial B_r} \nabla \Psi \cdot \mathbf{Y}_{lm}^{[1]} \mathrm{d}S = -\frac{1}{l(l+1)r^2} \int_{\partial B_r} (\Psi r \Delta Y_{lm} + \Psi \hat{\mathbf{r}} \cdot \nabla Y_{lm}) \mathrm{d}S \\ &= \frac{1}{r^3} \int_{\partial B_r} \Psi Y_{lm} \mathrm{d}S = \frac{1}{r} \Psi_{lm}(r) \end{split}$$

where we used the fact that $\Delta Y_{lm} = -l(l+1)r^{-2}Y_{lm}$. Also,

$$\Psi_{lm}^{[2]}(r) = \frac{1}{l(l+1)r^2} \int_{\partial B_r} \nabla \Psi \cdot \mathbf{Y}_{lm}^{[2]} \mathrm{d}S = -\frac{1}{l(l+1)r^2} \int_{\partial B_r} \Psi \nabla \cdot (\mathbf{r} \times \nabla Y_{lm}) \mathrm{d}S = 0.$$

Evaluating $\int_{\mathbb{R}^3} |\nabla \Psi|^2 d\mathbf{x}$ using (3.120) we get (3.119). This completes the proof of (3.119).

3.3 Momentum and energy

The energy and momentum conservation account for a four-dimensional freedom in the parameter space of the self-similar Goldreich-Weber solutions, see Definition 1.1. We shall require that the initial perturbation belongs to a codimension 4 "manifold" of initial data so that they have the same total momentum and total energy as the background GW star, i.e. (3.73) and (3.74). We will show that the linearisation of this requirement allows us to dynamically control the inner products

$$\langle \boldsymbol{\theta}, \mathbf{x} \rangle_3, \quad \langle \boldsymbol{\theta}, \mathbf{e}_i \rangle_3, \qquad i = 1, 2, 3,$$

modulo nonlinear terms, which is necessary for the proof of linear coercivity in Theorem 3.16. Hence, by fixing the total momentum and energy, we will be able to apply the non-negativity results we have for the linear operator **L** to control $\int_{B_R} \bar{w}^{-2} |\nabla \cdot (\bar{w}^3 \partial_s^a \partial^\beta \theta)|^2 dx$ with $\langle \mathbf{L} \partial_s^a \partial^\beta \theta, \partial_s^a \partial^\beta \theta \rangle_3 + ||\partial_s^{a+1} \partial^\beta \theta||_3^2$ modulo a correction involving non-linear terms. This is the main result of this section, stated and proved in Proposition 3.27.

Firstly, the momentum condition (3.73) gives us the following.

Lemma 3.23. Let θ be a solution of (3.62) in the sense of Theorem 3.11, and such that $\mathbf{W} = \overline{\mathbf{W}}$ (3.73). *Then*

$$-\frac{1}{2}\langle \partial_s^{a+1}\partial\!\!\!/^\beta \boldsymbol{\theta}, \mathbf{e}_i\rangle_3^2 = \delta \langle \partial_s^a \partial\!\!\!/^\beta \boldsymbol{\theta}, \mathbf{e}_i\rangle_3^2, \qquad a \ge 0, \, |\beta| \ge 0.$$

Proof. From Lemma 3.3, we see that when $\mathbf{W}_{\delta}[\boldsymbol{\theta}] = \bar{\mathbf{W}}$ we have

$$\langle \partial_s \boldsymbol{ heta}, \mathbf{e}_i
angle_3 = \mathfrak{G} \langle \boldsymbol{ heta}, \mathbf{e}_i
angle_3 \qquad ext{for} \qquad i=1,2,3.$$

and hence for any a with θ sufficiently smooth,

$$\langle \partial_s^{a+1} \boldsymbol{\theta}, \mathbf{e}_i \rangle_3 = \ell \langle \partial_s^a \boldsymbol{\theta}, \mathbf{e}_i \rangle_3 \quad \text{for} \quad i = 1, 2, 3.$$
 (3.121)

Now note that, using integration by parts,

$$\langle \partial_s^{a+1} \partial_j \partial^{\beta'} \boldsymbol{\theta}, \mathbf{e}_i \rangle_3 = -\langle \partial_s^{a+1} \partial^{\beta'} \boldsymbol{\theta}, \partial_j \mathbf{e}_i \rangle_3 \tag{3.122}$$

$$= 0 = \ell \langle \partial_s^a \partial_j \partial^{\beta'} \boldsymbol{\theta}, \mathbf{e}_i \rangle_3 \quad \text{for} \quad i = 1, 2, 3.$$
 (3.123)

We are done noting $\delta = -\frac{1}{2}\delta^2$ (3.58).

We now turn our attention to the energy condition (3.74).

Lemma 3.24. Let θ be a solution of (3.62) in the sense of Theorem 3.11, and such that $E = \overline{E}$ (3.74). Then

$$\frac{5}{2} \delta^2 \langle \partial_s^a \boldsymbol{\theta}, \mathbf{x} \rangle_3 = 2 \delta \langle \partial_s^{a+1} \boldsymbol{\theta}, \mathbf{x} \rangle_3 - \int \left(\bar{w}^3 \partial_s^a | \partial_s \boldsymbol{\theta} - \delta \boldsymbol{\theta} |^2 + 6 \bar{w}^4 \partial_s^a \left(\mathcal{F}^{-\frac{1}{3}} - 1 + \frac{1}{3} \nabla \cdot \boldsymbol{\theta} \right) \right) d\mathbf{x} \\
- \int \bar{w}^3 \partial_s^a (\mathcal{H}_{\boldsymbol{\xi}} - \mathcal{H} - \mathcal{H}_{\boldsymbol{\xi}}^{(1)}) \bar{w}^3 d\mathbf{x}.$$
(3.124)

Proof. From Lemma 3.3, we see that when $E_{\delta}[\theta] = \overline{E}$ we have

$$\begin{split} \frac{5}{2} \delta^2 \langle \boldsymbol{\theta}, \mathbf{x} \rangle_3 &= 2 \delta \langle \partial_s \boldsymbol{\theta}, \mathbf{x} \rangle_3 \\ &- \int \left(\bar{w}^3 |\partial_s \boldsymbol{\theta} - \delta \boldsymbol{\theta}|^2 + 6 \bar{w}^4 \left(\mathcal{J}^{-\frac{1}{3}} - 1 + \frac{1}{3} \nabla \cdot \boldsymbol{\theta} \right) + \bar{w}^3 (\mathcal{K}_{\boldsymbol{\xi}} - \mathcal{K} - \mathcal{K}_{\boldsymbol{\xi}}^{(1)}) \bar{w}^3 \right) \mathrm{d}\mathbf{x}. \end{split}$$

And hence for any $a \ge 0$ the identity (3.124) easily follows.

The next lemma is needed to estimate the term with $\Re_{\xi} - \Re - \Re_{\xi}^{(1)}$. Lemma 3.25. Let $n \ge 20$ and $a + |\beta| \le n$ with a > 0. We have

$$\begin{aligned} |\partial_s^a (\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^\beta K_2(\mathbf{x}, \mathbf{z})| &\lesssim \frac{(E_n + Z_n^2)^{1/2}}{|\mathbf{x} - \mathbf{z}|^2} \sum_{\substack{0 < a' \leq a \\ \beta' \leq \beta}} |\partial_s^{a'} \partial_{\beta'} \theta(\mathbf{x}) - \partial_s^{a'} \partial_{\beta'} \theta(\mathbf{z})| \\ &+ \frac{E_n^{1/2}}{|\mathbf{x} - \mathbf{z}|^2} \sum_{\beta' \leq \beta} |\partial_{\beta'} \theta(\mathbf{x}) - \partial_{\beta'} \theta(\mathbf{z})|, \end{aligned}$$

where we recall K_2 (3.90).

Proof. From Lemma 3.14,

$$K_{2}(\mathbf{x}, \mathbf{z}) = -\frac{1}{2} \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^{2}}{|\mathbf{x} - \mathbf{z}|^{3}} + \frac{3}{4|\mathbf{x} - \mathbf{z}|} \left(2 \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^{2}} + \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^{2}}{|\mathbf{x} - \mathbf{z}|^{2}} \right)^{2}$$
$$\varpi_{\frac{1}{2}} \underbrace{\left(2 \frac{(\mathbf{x} - \mathbf{z}) \cdot (\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z}))}{|\mathbf{x} - \mathbf{z}|^{2}} + \frac{|\boldsymbol{\theta}(\mathbf{x}) - \boldsymbol{\theta}(\mathbf{z})|^{2}}{|\mathbf{x} - \mathbf{z}|^{2}} \right)}_{:=y(\mathbf{x},\mathbf{z})}$$

Note that $|y(\mathbf{x}, \mathbf{z})| \leq \|\nabla \theta\|_{L^{\infty}}$. Our a priori assumption (3.71) together with the embedding theorems A.8 and A.9 mean that $\|\nabla \theta\|_{L^{\infty}}$ is bounded by a small constant. So we can assume $|y(\mathbf{x}, \mathbf{z})| \leq 1/2$. Then from the definition of ϖ_q (3.92) we can see that

$$\varpi_{\frac{1}{2}}^{(k)}(y(\mathbf{x}, \mathbf{z})) \lesssim 1 \quad \text{for any} \quad k \ge 0.$$

Now using part (ii) of Lemma 2.8, chain and product rule for derivatives and the embedding theorems A.8 and A.9, we can see that $\partial_s^a (\partial_x + \partial_z)^\beta K_2(\mathbf{x}, \mathbf{z})$ satisfies the stated bounds.

So the energy condition (3.74) gives us the following.

Lemma 3.26. Let $n \ge 20$ and $a + |\beta| \le n$ with $a > 0, |\beta| \ge 0$. Let θ be a solution of (3.62) in the sense of Theorem 3.11, and such that $E = \overline{E}$ (3.74). Then

$$\left| 3\delta \langle \partial_s^a \partial^\beta \boldsymbol{\theta}, \mathbf{x} \rangle_3^2 + \frac{24}{25} \langle \partial_s^{a+1} \partial^\beta \boldsymbol{\theta}, \mathbf{x} \rangle_3^2 \right| \lesssim S_{n,|\beta|-1,0} + C_{\delta} (E_n + Z_n^2)^{1/2} E_n$$

Proof. First we deal with the case $|\beta| = 0$. From (3.124) we get

$$2\langle \partial_s^{a+1}\boldsymbol{\theta}, \mathbf{x} \rangle_3 = \frac{5}{2} \mathscr{C} \langle \partial_s^a \boldsymbol{\theta}, \mathbf{x} \rangle_3 + \mathscr{C}^{-1} \int \left(\bar{w}^3 \partial_s^a |\partial_s \boldsymbol{\theta} - \mathscr{C} \boldsymbol{\theta}|^2 + 6 \bar{w}^4 \partial_s^a \left(\mathscr{J}^{-\frac{1}{3}} - 1 + \frac{1}{3} \nabla \cdot \boldsymbol{\theta} \right) \right) d\mathbf{x} \\ + \mathscr{C}^{-1} \int \bar{w}^3 \partial_s^a (\mathscr{K}_{\boldsymbol{\xi}} - \mathscr{K} - \mathscr{K}_{\boldsymbol{\xi}}^{(1)}) \bar{w}^3 d\mathbf{x}.$$

$$(3.125)$$

With the embedding theorems A.8 and A.9, it is easy to see that

$$\left|\int \left(\bar{w}^3 \partial_s^a \left|\partial_s \boldsymbol{\theta} - \boldsymbol{\vartheta} \boldsymbol{\theta}\right|^2 + 6\bar{w}^4 \partial_s^a \left(\boldsymbol{\mathcal{J}}^{-\frac{1}{3}} - 1 + \frac{1}{3} \nabla \cdot \boldsymbol{\theta}\right)\right) \mathrm{d}\mathbf{x}\right| \lesssim_{\delta} (E_n + Z_n^2)^{1/2} E_n^{1/2}.$$

Now using Lemmas 2.8-3.25 and Young's convolution inequality we have

$$\left| \int \bar{w}^3 \partial_s^a (\mathcal{K}_{\boldsymbol{\xi}} - \mathcal{K} - \mathcal{K}_{\boldsymbol{\xi}}^{(1)}) \bar{w}^3 \mathrm{d}\mathbf{x} \right| = \left| \int \int \bar{w}^3 (\mathbf{x}) \bar{w}^3 (\mathbf{z}) \partial_s^a K_2(\mathbf{x}, \mathbf{z}) \mathrm{d}\mathbf{x} \mathrm{d}\mathbf{z} \right| \lesssim (E_n + Z_n^2)^{1/2} E_n^{1/2}.$$
(3.126)

Therefore, upon taking the square of (3.125) and using the simple bound $|\langle \partial_s^a \theta, \mathbf{x} \rangle_3| \lesssim E_n^{\frac{1}{2}}$, we obtain

$$\left|4\langle\partial_s^{a+1}\boldsymbol{\theta},\mathbf{x}\rangle_3^2 - \frac{25}{4}\boldsymbol{\vartheta}^2\langle\partial_s^a\boldsymbol{\theta},\mathbf{x}\rangle_3^2\right| \lesssim_{\delta} (E_n + Z_n^2)^{1/2}E_n,$$

which concludes the proof for when $|\beta| = 0$ since $\delta = -\frac{1}{2}\delta^2$ (recall (3.58)).

Now note that, using integration by parts,

$$|\langle \partial_s^{a+1} \partial_j \partial^{\beta'} \boldsymbol{\theta}, \mathbf{x} \rangle_3| = |\langle \partial_s^{a+1} \partial^{\beta'} \boldsymbol{\theta}, \partial_j \mathbf{x} \rangle_3| \lesssim S_{n,|\beta'|,0}^{1/2}$$

Similarly we have $|\langle \partial_s^a \partial_j \partial^{\beta'} \theta, \mathbf{x} \rangle_3| \lesssim S_{n,|\beta'|,0}^{1/2}$. Noting that $|\delta| \lesssim 1$ and we are done.

Finally, the momentum condition (3.73) and the energy condition (3.74) together gives us the following proposition.

Proposition 3.27. Let $n \ge 20$ and $a + |\beta| \le n$ with a > 0. Let θ be a solution of (3.62) in the sense of Theorem 3.11 such that $\mathbf{W} = \overline{\mathbf{W}}$ (3.73) and $E = \overline{E}$ (3.74). Then

$$\begin{split} |\mathfrak{C}|^{2} \int_{B_{R}} \bar{w}^{-2} |\nabla \cdot (\bar{w}^{3} \partial_{s}^{a} \partial_{s}^{\beta} \theta)|^{2} \mathrm{d}\mathbf{x} &\lesssim \langle \mathbf{L} \partial_{s}^{a} \partial_{s}^{\beta} \theta, \partial_{s}^{a} \partial_{s}^{\beta} \theta \rangle_{3} + \frac{49}{50} \|\partial_{s}^{a+1} \partial_{s}^{\beta} \theta\|_{3}^{2} \\ &+ S_{n,|\beta|-1,0} + C_{\delta} (E_{n} + Z_{n}^{2})^{1/2} E_{n}. \end{split}$$
(3.127)

Proof. Let

$$\tilde{\boldsymbol{\theta}} = \partial_s^a \partial^\beta \boldsymbol{\theta} - \frac{\langle \partial_s^a \partial^\beta \boldsymbol{\theta}, \mathbf{x} \rangle_3}{\|\mathbf{x}\|_3^2} \mathbf{x} - \sum_{i=1}^3 \frac{\langle \partial_s^a \partial^\beta \boldsymbol{\theta}, \mathbf{e}_i \rangle_3}{\|\mathbf{e}_i\|_3^2} \mathbf{e}_i$$
(3.128)

$$\boldsymbol{\theta}' = \partial_s^{a+1} \partial^\beta \boldsymbol{\theta} - \frac{\langle \partial_s^{a+1} \partial^\beta \boldsymbol{\theta}, \mathbf{x} \rangle_3}{\|\mathbf{x}\|_3^2} \mathbf{x} - \sum_{i=1}^3 \frac{\langle \partial_s^{a+1} \partial^\beta \boldsymbol{\theta}, \mathbf{e}_i \rangle_3}{\|\mathbf{e}_i\|_3^2} \mathbf{e}_i$$
(3.129)

Then $\langle \tilde{\theta}, \mathbf{x} \rangle_3 = 0 = \langle \tilde{\theta}, \mathbf{e}_i \rangle_3$ for i = 1, 2, 3, and

$$\|\partial_s^{a+1}\partial^{\beta}\boldsymbol{\theta}\|_3^2 = \|\boldsymbol{\theta}'\|_3^2 + \frac{\langle\partial_s^{a+1}\partial^{\beta}\boldsymbol{\theta}, \mathbf{x}\rangle_3^2}{\|\mathbf{x}\|_3^2} + \sum_{i=1}^3 \frac{\langle\partial_s^{a+1}\partial^{\beta}\boldsymbol{\theta}, \mathbf{e}_i\rangle_3^2}{\|\mathbf{e}_i\|_3^2}.$$
(3.130)

Since x and \mathbf{e}_i are eigenfunctions of L with eigenvalues 3δ and δ respectively, we have

$$\langle \mathbf{L} \partial_s^a \partial^\beta \boldsymbol{\theta}, \partial_s^a \partial^\beta \boldsymbol{\theta} \rangle_3$$
$$= \left\langle \mathbf{L}\tilde{\boldsymbol{\theta}} + 3\delta \frac{\langle \partial_s^a \partial^\beta \boldsymbol{\theta}, \mathbf{x} \rangle_3}{\|\mathbf{x}\|_3^2} \mathbf{x} + \sum_{i=1}^3 \delta \frac{\langle \partial_s^a \partial^\beta \boldsymbol{\theta}, \mathbf{e}_i \rangle_3}{\|\mathbf{e}_i\|_3^2} \mathbf{e}_i, \tilde{\boldsymbol{\theta}} + \frac{\langle \partial_s^a \partial^\beta \boldsymbol{\theta}, \mathbf{x} \rangle_3}{\|\mathbf{x}\|_3^2} \mathbf{x} + \sum_{i=1}^3 \frac{\langle \partial_s^a \partial^\beta \boldsymbol{\theta}, \mathbf{e}_i \rangle_3}{\|\mathbf{e}_i\|_3^2} \mathbf{e}_i \right\rangle_3$$
$$= \langle \mathbf{L}\tilde{\boldsymbol{\theta}}, \tilde{\boldsymbol{\theta}} \rangle_3 + 3\delta \frac{\langle \partial_s^a \partial^\beta \boldsymbol{\theta}, \mathbf{x} \rangle_3^2}{\|\mathbf{x}\|_3^2} + \sum_{i=1}^3 \delta \frac{\langle \partial_s^a \partial^\beta \boldsymbol{\theta}, \mathbf{e}_i \rangle_3^2}{\|\mathbf{e}_i\|_3^2}.$$
(3.131)

We use Lemmas 3.23 and 3.26 to control the last two terms on the right-most side of (3.131) to get

$$\langle \mathbf{L}\tilde{\boldsymbol{\theta}}, \tilde{\boldsymbol{\theta}} \rangle_{3} \leq \langle \mathbf{L}\partial_{s}^{a} \boldsymbol{\vartheta}^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} \boldsymbol{\vartheta}^{\beta} \boldsymbol{\theta} \rangle_{3} + \frac{24}{25} \frac{\langle \partial_{s}^{a+1} \boldsymbol{\vartheta}^{\beta} \boldsymbol{\theta}, \mathbf{x} \rangle_{3}^{2}}{\|\mathbf{x}\|_{3}^{2}} + \frac{1}{2} \sum_{i=1}^{3} \frac{\langle \partial_{s}^{a+1} \boldsymbol{\vartheta}^{\beta} \boldsymbol{\theta}, \mathbf{e}_{i} \rangle_{3}^{2}}{\|\mathbf{e}_{i}\|_{3}^{2}} + CS_{n,|\beta|-1,0} + C_{\delta} (E_{n} + Z_{n}^{2})^{1/2} E_{n}$$

$$\leq \langle \mathbf{L} \partial_{s}^{a} \boldsymbol{\vartheta}^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} \boldsymbol{\vartheta}^{\beta} \boldsymbol{\theta} \rangle_{s} + \frac{24}{24} \|\partial_{s}^{a+1} \boldsymbol{\vartheta}^{\beta} \boldsymbol{\theta}\|^{2} + CS_{n} + CS_{n} + C_{\delta} (E_{n} + Z_{n}^{2})^{1/2} E_{n}$$

$$\leq \langle \mathbf{L} \partial_{s}^{a} \boldsymbol{\vartheta}^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} \boldsymbol{\vartheta}^{\beta} \boldsymbol{\theta} \rangle_{s} + \frac{24}{24} \|\partial_{s}^{a+1} \boldsymbol{\vartheta}^{\beta} \boldsymbol{\theta}\|^{2} + CS_{n} + CS$$

$$\leq \langle \mathbf{L}\partial_s^a \partial^\beta \boldsymbol{\theta}, \partial_s^a \partial^\beta \boldsymbol{\theta} \rangle_3 + \frac{24}{25} \|\partial_s^{a+1} \partial^\beta \boldsymbol{\theta}\|_3^2 + CS_{n,|\beta|-1,0} + C_\delta (E_n + Z_n^2)^{1/2} E_n, \qquad (3.133)$$

where we have used (3.130) in the last line. We now use the decomposition (3.128) and then apply Theorem 3.16 (with $\theta = \tilde{\theta}$) to obtain

$$\begin{aligned} \epsilon \int_{B_{R}} \bar{w}^{-2} |\nabla \cdot (\bar{w}^{3} \partial_{s}^{a} \partial^{\beta} \theta)|^{2} \mathrm{d}\mathbf{x} \\ &\leq C \epsilon \int_{B_{R}} \bar{w}^{-2} |\nabla \cdot (\bar{w}^{3} \tilde{\theta})|^{2} \mathrm{d}\mathbf{x} + C \epsilon \frac{\langle \partial_{s}^{a} \partial^{\beta} \theta, \mathbf{x} \rangle_{3}^{2}}{\|\mathbf{x}\|_{3}^{2}} + C \epsilon \sum_{i=1}^{3} \frac{\langle \partial_{s}^{a} \partial^{\beta} \theta, \mathbf{e}_{i} \rangle_{3}^{2}}{\|\mathbf{e}_{i}\|_{3}^{2}} \\ &\leq \langle \mathbf{L} \tilde{\theta}, \tilde{\theta} \rangle_{3} + \frac{C \epsilon}{|\delta|} \|\partial_{s}^{a+1} \partial^{\beta} \theta\|_{3}^{2} + C S_{n,|\beta|-1,0} + C_{\delta} (E_{n} + Z_{n}^{2})^{1/2} E_{n} \\ &\leq \langle \mathbf{L} \partial_{s}^{a} \partial^{\beta} \theta, \partial_{s}^{a} \partial^{\beta} \theta \rangle_{3} + \frac{24}{25} \|\partial_{s}^{a+1} \partial^{\beta} \theta\|_{3}^{2} + \frac{C \epsilon}{|\delta|} \|\partial_{s}^{a+1} \partial^{\beta} \theta\|_{3}^{2} + C S_{n,|\beta|-1,0} + C_{\delta} (E_{n} + Z_{n}^{2})^{1/2} E_{n} \\ &\leq \langle \mathbf{L} \partial_{s}^{a} \partial^{\beta} \theta, \partial_{s}^{a} \partial^{\beta} \theta \rangle_{3} + \frac{49}{50} \|\partial_{s}^{a+1} \partial^{\beta} \theta\|_{3}^{2} + C S_{n,|\beta|-1,0} + C_{\delta} (E_{n} + Z_{n}^{2})^{1/2} E_{n}, \end{aligned}$$
(3.134)

where we have chosen ϵ small enough so that $C\epsilon \lesssim 1$ in the second line and then further shrink ϵ so that $\frac{C\epsilon}{|\delta|} < \frac{1}{50}$ in the fourth line. Note that since $\delta = -\frac{1}{2}t^2$ (recall (3.58)), the dependence of ϵ on t is $\epsilon \sim |t|^2$. We have used Lemmas 3.23 and 3.26 in the second bound, and (3.133) in the third bound.

3.4 Coercivity via irrotationality

Note that Proposition 3.27 only controls the weighted divergence $g = \nabla \cdot (\bar{w}^3 \theta)$ and not the norms of θ in our energy spaces. It is therefore still not strong enough for our energy estimates in Sections 3.5–3.6. To derive the coercivity we seek, we must mod out the kernel of L, i.e. the subspace of θ with weighted divergence g = 0. This is naturally linked to the assumption of irrotationality (3.75) which guarantees, we show this in the key result of this section – Proposition 3.36, that we can in fact dynamically control $\|\partial_s^a \partial^\beta \theta\|_3^2 + \|\partial_s^a \nabla \partial^\beta \theta\|_4^2$ modulo lower order nonlinear terms.

3.4.1 Lagrangian description of irrotationality

From (3.84) it is clear that any H^2 vectorfield $\boldsymbol{\theta}$ such that $g = \nabla \cdot (\bar{w}^3 \boldsymbol{\theta}) = 0$ is in the kernel of the operator **L**. In particular, to obtain strict coercivity of **L** we restrict ourselves to $\langle \cdot, \cdot \rangle_3$ -orthogonal complement of $K = \{\boldsymbol{\theta} : \nabla \cdot (\bar{w}^3 \boldsymbol{\theta}) = 0\}$. Note that $\{\boldsymbol{\theta} = \nabla \vartheta\} \subseteq K^{\perp}$ since for any $\boldsymbol{\theta}_0 \in K$ we have

$$\langle \nabla \vartheta, \boldsymbol{\theta}_0 \rangle_3 = \int \nabla \vartheta \cdot \boldsymbol{\theta}_0 \bar{w}^3 d\mathbf{x} = \int \vartheta \nabla \cdot (\boldsymbol{\theta}_0 \bar{w}^3) d\mathbf{x} = 0.$$

Therefore, the natural assumption to hope for the strict coercivity of the term on the left-hand side of (3.127) is that θ is in fact a gradient. In this section we show that this is true to the top order if we assume that the fluid is irrotational. The challenge is that the irrotationality condition in the Lagrangian variables (3.75) is expressed at the level of the *s*-derivative of the flow map, and a careful analysis is necessary to obtain satisfactory lower bounds.

Lemma 3.28. Let θ be a solution of (3.62) in the sense of Theorem 3.11, given on its maximal interval of existence. Assume further that the fluid is irrotational, i.e. initially (3.75) holds. Then for a > 0 we have

$$\partial_s \boldsymbol{\theta} = \nabla \left(\tilde{H} + \frac{1}{2} \boldsymbol{\theta} |\boldsymbol{\theta} + \mathbf{x}|^2 \right) - (\partial_s \theta^k) \nabla \theta^k$$
(3.135)

$$\partial_s^a \boldsymbol{\theta} = \nabla H_a - \sum_{j=0}^{\lfloor \frac{a-j}{2} \rfloor} C_{a,j} (\partial_s^{a-j} \boldsymbol{\theta}^k) \nabla \partial_s^j \boldsymbol{\theta}^k$$
(3.136)

for some real constants $C_{a,j}$, $j \in \{1, \ldots, \lfloor \frac{a-1}{2} \rfloor\}$ and H^1 -functions H_a and \tilde{H} .

Proof. Since the Euler-Poisson equation preserves the fluid irrotational condition, (3.75) implies that $\operatorname{curl}_A \partial_t \eta = \mathbf{0}$ for t, or equivalently in Eulerian coordinates $\nabla \times \mathbf{u} = 0$. Since any curl-free vector field can be written as a gradient, we have $\mathbf{u} = \nabla \hat{H}$ for some \hat{H} , or equivalently $\partial_t \eta = A \nabla H$ for some H in Lagrangian coordinates. Since

$$\partial_t \boldsymbol{\eta} = \lambda^{-3/2} \partial_s (\lambda(\boldsymbol{\theta} + \mathbf{x})) = \lambda^{-3/2} ((\boldsymbol{\theta} + \mathbf{x}) \partial_s \lambda + \lambda \partial_s \boldsymbol{\theta}) = \lambda^{-1/2} (\partial_s \boldsymbol{\theta} - \boldsymbol{\vartheta}(\boldsymbol{\theta} + \mathbf{x})),$$

this means on the level of θ we have

$$\partial_s \boldsymbol{\theta} - \boldsymbol{\vartheta} (\boldsymbol{\theta} + \mathbf{x}) = \mathscr{A} \nabla \tilde{H}.$$

Hence we have

$$\partial_{s}\boldsymbol{\theta} = \nabla\tilde{H} + (\boldsymbol{\mathcal{A}} - I)\nabla\tilde{H} + \boldsymbol{\mathcal{C}}(\boldsymbol{\theta} + \mathbf{x}) = \nabla\tilde{H} + (I - \boldsymbol{\mathcal{A}}^{-1})\boldsymbol{\mathcal{A}}\nabla\tilde{H} + \boldsymbol{\mathcal{C}}(\boldsymbol{\theta} + \mathbf{x})$$
$$= \nabla\tilde{H} - (\partial_{s}\theta^{k} - \boldsymbol{\mathcal{C}}(\theta^{k} + x^{k}))\nabla\theta^{k} + \boldsymbol{\mathcal{C}}(\boldsymbol{\theta} + \mathbf{x}) = \nabla\left(\tilde{H} + \frac{1}{2}\boldsymbol{\mathcal{C}}|\boldsymbol{\theta} + \mathbf{x}|^{2}\right) - (\partial_{s}\theta^{k})\nabla\theta^{k}.$$

This proves (3.135). To prove (3.136), we will use induction. We have shown that it is true for a = 1. Suppose it is true for some $a \ge 1$. Then

$$\begin{split} \partial_s^{a+1} \boldsymbol{\theta} &= \nabla \partial_s H_a - \partial_s \sum_{j=0}^{\lfloor \frac{a-1}{2} \rfloor} C_{a,j} (\partial_s^{a-j} \theta^k) \nabla \partial_s^j \theta^k \\ &= \nabla \partial_s H_a - \sum_{j=0}^{\lfloor \frac{a-1}{2} \rfloor} C_{a,j} (\partial_s^{a+1-j} \theta^k) \nabla \partial_s^j \theta^k - \sum_{j=0}^{\lfloor \frac{a-1}{2} \rfloor} C_{a,j} (\partial_s^{a-j} \theta^k) \nabla \partial_s^{j+1} \theta^k \\ &= \nabla \partial_s H_a - \sum_{j=0}^{\lfloor \frac{a-1}{2} \rfloor} C_{a,j} (\partial_s^{a+1-j} \theta^k) \nabla \partial_s^j \theta^k - \sum_{j=1}^{\lfloor \frac{a-1}{2} \rfloor + 1} C_{a,j-1} (\partial_s^{a+1-j} \theta^k) \nabla \partial_s^j \theta^k. \end{split}$$

Note that $\lfloor \frac{a-1}{2} \rfloor + 1 > \lfloor \frac{a}{2} \rfloor$ if and only if a is odd. Assume therefore that a = 2a' + 1 for some $a' \in \mathbb{N} \cup \{0\}$. Then $\lfloor \frac{a-1}{2} \rfloor + 1 = a' + 1$ and $\lfloor \frac{a}{2} \rfloor = a'$. When j = a' + 1 in the last sum, we have

$$C_{2a'+1,a'}(\partial_s^{2a'+2-(a'+1)}\theta^k)\nabla\partial_s^{a'+1}\theta^k = C_{2a'+1,a'}(\partial_s^{a'+1}\theta^k)\nabla\partial_s^{a'+1}\theta^k = \frac{1}{2}C_{2a'+1,a'}\nabla(\partial_s^{a'+1}\theta^k)^2$$

which can be absorbed into H_{a+1} . Therefore

$$\begin{split} \partial_s^{a+1} \boldsymbol{\theta} &= \nabla \left(\partial_s H_a + \frac{1}{2} \mathbf{1} [a \text{ odd}] C_{a,a'} \nabla (\partial_s^{a'+1} \theta^k)^2 \right) \\ &- \sum_{j=0}^{\lfloor \frac{a-1}{2} \rfloor} C_{a,j} (\partial_s^{a+1-j} \theta^k) \nabla \partial_s^j \theta^k - \sum_{j=1}^{\lfloor \frac{a}{2} \rfloor} C_{a,j-1} (\partial_s^{a+1-j} \theta^k) \nabla \partial_s^j \theta^k \\ &= \nabla H_{a+1} - \sum_{j=0}^{\lfloor \frac{a}{2} \rfloor} C_{a+1,j} (\partial_s^{a-j} \theta^k) \nabla \partial_s^j \theta^k, \end{split}$$

where $\mathbf{1}[\star]$ denotes the Iverson bracket (see Definition 2.2). This completes the induction argument.

Remark 3.29. The above lemma is a purely structural statement about (suitably smooth) irrotational fields. Strictly speaking we do not need θ to be a solution of the Euler-Poisson system (3.62).

With this we can now show that the curl of $\partial_s^a X_r^b \partial^\beta \theta$ equals lower order terms and non-linear terms.

Lemma 3.30. Let θ be a solution of (3.62) in the sense of Theorem 3.11, given on its maximal interval of existence. Assume further that the fluid is irrotational, i.e. initially (3.75) holds. Then for a > 0 we have

$$\nabla \times \partial_s^a \boldsymbol{\theta} = -\partial_s^{a-1}((\nabla \partial_s \theta^k) \times \nabla \theta^k).$$
(3.137)

Moreover, for some constants $C_{\gamma,\beta} > 0$ we have

$$\nabla \times \partial_s^a \partial^\beta \boldsymbol{\theta} = -\partial_s^{a-1} \partial^\beta ((\nabla \partial_s \theta^k) \times \nabla \theta^k) + \sum_{|\gamma| < |\beta|} C_{\gamma,\beta} \langle \nabla \partial_s^a \partial^\gamma \boldsymbol{\theta} \rangle, \tag{3.138}$$

$$\nabla \times \partial_s^a X_r^b \partial^\beta \boldsymbol{\theta} = -\partial_s^{a-1} X_r^b \partial^\beta ((\nabla \partial_s \theta^k) \times \nabla \theta^k) + \sum_{|\gamma|+d < |\beta|+b} C_{\gamma,\beta} \langle \nabla \partial_s^a X_r^d \partial^\gamma \boldsymbol{\theta} \rangle,$$
(3.139)

where we recall notations defined in Definition 2.2.

Proof. Apply $\nabla \times \partial_s^{a-1}$ to (3.135) to get (3.137). Formulas (3.138)–(3.139) follow trivially when $|\beta| = 0 = b$. Now assume formula (3.138) is true for a multi-index β , $|\beta| \ge 0$. Then

$$\begin{split} \nabla \times \partial_s^a \partial_j \partial^\beta \theta &= \langle \nabla \partial_s^a \partial^\beta \theta \rangle + \partial_j \nabla \times \partial_s^a \partial^\beta \theta \\ &= \langle \nabla \partial_s^a \partial^\beta \theta \rangle - \partial_s^{a-1} \partial_j \partial^\beta ((\nabla \partial_s \theta^k) \times \nabla \theta^k) + \sum_{|\gamma| < |\beta|} C_{\gamma,\beta} \langle \partial_j \nabla \partial_s^a \partial^\gamma \theta \rangle \\ &= -\partial_s^{a-1} \partial_j \partial^\beta ((\nabla \partial_s \theta^k) \times \nabla \theta^k) + \sum_{|\gamma| < |\beta| + 1} C'_{\gamma,\beta} \langle \nabla \partial_s^a \partial^\gamma \theta \rangle, \end{split}$$

where we recall the notation from Definition 2.2 and the commutation relation $[\partial_j, \nabla] = \langle \nabla \rangle$ from Lemma A.2. The proof then follows by induction. The proof of (3.139) is similar, using the commutation relation $[X_r, \nabla] = \langle \nabla \rangle$ from Lemma A.2.

Corollary 3.31. Let θ be a solution of (3.62) in the sense of Theorem 3.11, given on its maximal interval of existence. Assume further that the fluid is irrotational, i.e. initially (3.75) holds. Let $n \ge 20$.

i. For $a + |\beta| \le n$ with a > 0 we have

$$\|\nabla \times \partial_s^a \partial^\beta \theta\|_4^2 \lesssim S_{n,|\beta|-1,0} + (E_n + Z_n^2) E_n$$

ii. For $a + |\beta| + b \le n$ with a > 0 we have

$$\|\nabla \times \partial_s^a X_r^b \partial^\beta \theta\|_{4+b}^2 \lesssim S_{n,|\beta|+b-1} + (E_n + Z_n^2) E_n.$$

Proof. Use Lemma 3.30 and note that

$$\begin{split} \left\| \partial_s^{a-1} \partial^{\beta} ((\nabla \partial_s \theta^k) \times \nabla \theta^k) \right\|_4^2 &\lesssim (E_n + Z_n^2) E_n \\ \left\| \sum_{|\gamma| < |\beta|} C_{\gamma,\beta} \langle \nabla \partial_s^a \partial^{\gamma} \theta \rangle \right\|_4^2 &\lesssim S_{n,|\beta|-1,0}, \end{split}$$

which yields the first claim. The second claim follows similarly.

3.4.2 Coercivity of L

The lemmas in the last subsection showed that $\partial_s^a \theta$ is a gradient on the linear level, which will ultimately help us show that $\|\partial_s^a \partial^\beta \theta\|_3^2 + \|\partial_s^a \nabla \partial^\beta \theta\|_4^2$ can be "controlled" by the linearised dynamics. We start by showing we can control $\|\partial_s^a \theta\|_3^2$ in the following lemma.

Lemma 3.32. Let θ be a solution of (3.62) in the sense of Theorem 3.11, given on its maximal interval of existence. Assume further that the fluid is irrotational, i.e. initially (3.75) holds. Let $n \ge 20$. Then we have the bound

$$\|\partial_s^a \theta\|_3^2 \lesssim \int_{B_R} \bar{w}^{-2} |\nabla \cdot (\bar{w}^3 \partial_s^a \theta)|^2 d\mathbf{x} + (E_n + Z_n^2)^{1/2} E_n \quad \text{for} \quad 0 < a \le n.$$
(3.140)

Proof. Let $g = \nabla \cdot (\bar{w}^3 \partial_s^a \theta)$. Multiply both sides of this equation by H_a and integrate over B_R to get

$$\begin{split} \int_{B_R} \bar{w}^3 (\nabla H_a) \cdot \partial_s^a \boldsymbol{\theta} \, \mathrm{d}\mathbf{x} &= -\int_{B_R} g H_a \mathrm{d}\mathbf{x} = -\int_{B_R} g (H_a - (H_a)_{B_{2R/3}}) \mathrm{d}\mathbf{x} \\ &\leq \epsilon^{-1} \int_{B_R} \bar{w}^{-2} g^2 \mathrm{d}\mathbf{x} + \epsilon \int_{B_R} (H_a - (H_a)_{B_{2R/3}})^2 \bar{w}^2 \mathrm{d}\mathbf{x} \\ &\leq \epsilon^{-1} \int_{B_R} \bar{w}^{-2} g^2 \mathrm{d}\mathbf{x} + \epsilon C' \int_{B_R} |\nabla H_a|^2 \bar{w}^4 \mathrm{d}\mathbf{x} \end{split}$$

where we have used the Hardy-Poincaré inequality in the last line, see Theorem A.4. From this and Lemma 3.28 we get

$$\int_{B_R} \bar{w}^3 |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} \le \epsilon^{-1} \int_{B_R} \bar{w}^{-2} g^2 \mathrm{d}\mathbf{x} + \epsilon C' \int_{B_R} |\partial_s^a \boldsymbol{\theta}|^2 \bar{w}^4 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' (E_n + Z_n^2)^{1/2} E_n |\partial_s^a \boldsymbol{\theta}|^2 \mathrm{d}\mathbf{x} + (1+\epsilon) C'' |\partial_s^a \boldsymbol{\theta}$$

where we bound for example

$$\begin{split} \left| \int_{B_R} \bar{w}^3 \left(\sum_{j=0}^{\lfloor \frac{a-1}{2} \rfloor} C_{a,j}(\partial_s^{a-j} \theta^k) \nabla \partial_s^j \theta^k \right) \cdot \partial_s^a \boldsymbol{\theta} \, \mathrm{d} \mathbf{x} \right| \\ \lesssim (E_n + Z_n^2)^{1/2} \left| \int_{B_R} \bar{w}^3 \left(\sum_{j=0}^{\lfloor \frac{a-1}{2} \rfloor} C_{a,j}(\partial_s^{a-j} \theta^k) \right) \cdot \partial_s^a \boldsymbol{\theta} \, \mathrm{d} \mathbf{x} \right| \lesssim (E_n + Z_n^2)^{1/2} S_n \end{split}$$

Choosing ϵ small enough, we get (3.140).

Before proving the key result of this section, we have the following structural decomposition, which holds for any sufficiently smooth vectorfield θ .

Lemma 3.33. For any θ such that $\|\theta\|_3 + \|\nabla\theta\|_4 < \infty$ we have

$$\begin{split} &\int_{B_R} \frac{4}{3} \bar{w}^{-2} |\nabla \cdot (\bar{w}^3 \boldsymbol{\theta})|^2 \mathrm{d}\mathbf{x} \\ &= \int_{B_R} \left(\bar{w}^4 \left(\frac{1}{3} |\nabla \cdot \boldsymbol{\theta}|^2 + |\nabla \boldsymbol{\theta}|^2 + [\operatorname{curl} \boldsymbol{\theta}]_l^k \partial_k \theta^l \right) - 4 \bar{w}^3 \theta^k \theta^l \partial_k \partial_l \bar{w} \right) \mathrm{d}\mathbf{x} \\ &= \int_{B_R} \left(\bar{w}^4 \left(\frac{1}{3} |\nabla \cdot \boldsymbol{\theta}|^2 + |\nabla \boldsymbol{\theta}|^2 - \frac{1}{2} |\operatorname{curl} \boldsymbol{\theta}|^2 \right) - \bar{w}^3 \left(\bar{w}'' |\boldsymbol{\theta} \cdot \mathbf{e}_r|^2 + \frac{\bar{w}'}{r} (|\boldsymbol{\theta}|^2 - |\boldsymbol{\theta} \cdot \mathbf{e}_r|^2) \right) \right) \mathrm{d}\mathbf{x}. \end{split}$$

where \mathbf{e}_r denotes the radial unit vector $\mathbf{x}/|\mathbf{x}|$.

Proof. The first line follows from the following identity:

$$-\frac{4}{3}\nabla(\bar{w}^{-2}\nabla\cdot(\bar{w}^{3}\boldsymbol{\theta})) = -\frac{1}{3\bar{w}^{3}}\nabla(\bar{w}^{4}\nabla\cdot\boldsymbol{\theta}) - \frac{1}{\bar{w}^{3}}\partial_{k}(\bar{w}^{4}\nabla\theta^{k}) - 4\boldsymbol{\theta}\cdot\nabla\nabla\bar{w}$$
$$= -\frac{1}{3\bar{w}^{3}}\nabla(\bar{w}^{4}\nabla\cdot\boldsymbol{\theta}) - \frac{1}{\bar{w}^{3}}\partial_{k}(\bar{w}^{4}(\partial_{k}\boldsymbol{\theta} + [\operatorname{curl}\boldsymbol{\theta}]_{\bullet}^{k}) - 4\boldsymbol{\theta}\cdot\nabla\nabla\bar{w}.$$

And then the second line follows from

$$[\operatorname{curl} \boldsymbol{\theta}]_{l}^{k} [\operatorname{curl} \boldsymbol{\theta}]_{l}^{k} = (\partial_{l} \theta^{k} - \partial_{k} \theta^{l}) (\partial_{l} \theta^{k} - \partial_{k} \theta^{l}) = (\partial_{k} \theta^{l} - \partial_{l} \theta^{k}) \partial_{k} \theta^{l} - (\partial_{l} \theta^{k} - \partial_{k} \theta^{l}) \partial_{k} \theta^{l} = -2(\partial_{l} \theta^{k} - \partial_{k} \theta^{l}) \partial_{k} \theta^{l} = -2[\operatorname{curl} \boldsymbol{\theta}]_{l}^{k} \partial_{k} \theta^{l}$$

and

$$\theta^k \theta^l \partial_k \partial_l \bar{w} = \theta^k \theta^l \partial_k \left(\bar{w}' \frac{x^l}{r} \right) = \theta^k \theta^l \left(\bar{w}'' \frac{x^l x^k}{r^2} + \bar{w}' \frac{\delta_k^l}{r} - \bar{w}' \frac{x^l x^k}{r^3} \right) = \bar{w}'' |\boldsymbol{\theta} \cdot \mathbf{e}_r|^2 + \frac{\bar{w}'}{r} (|\boldsymbol{\theta}|^2 - |\boldsymbol{\theta} \cdot \mathbf{e}_r|^2).$$

Using this, we can now prove that we can control $\|\partial_s^a \theta\|_3^2 + \|\partial_s^a \nabla \theta\|_4^2$.

Proposition 3.34. Let $n \ge 20$. Let θ be a solution of (3.62) in the sense of Theorem 3.11, given on its maximal interval of existence. Assume further that the energy, momentum, and irrotationality constraints (3.73), (3.74), and (3.75) hold respectively. Then for any $0 < a \le n$ we have

$$\|\partial_s^a \boldsymbol{\theta}\|_3^2 + \|\partial_s^a \nabla \boldsymbol{\theta}\|_4^2 \lesssim |\boldsymbol{b}|^{-2} \left(\frac{49}{50} \left\|\partial_s^{a+1} \boldsymbol{\theta}\right\|_3^2 + \langle \mathbf{L} \partial_s^a \boldsymbol{\theta}, \partial_s^a \boldsymbol{\theta} \rangle\right) + C_{\delta} (E_n + Z_n^2)^{1/2} E_n \tag{3.141}$$

Proof. Combining Proposition 3.27 and Lemma 3.32 we have, for small ϵ ,

$$\underbrace{\epsilon \int_{B_R} \bar{w}^{-2} |\nabla \cdot (\bar{w}^3 \partial_s^a \boldsymbol{\theta})|^2 \mathrm{d}\mathbf{x} + \|\partial_s^a \boldsymbol{\theta}\|_3^2}_{:=M} \lesssim |\boldsymbol{b}|^{-2} \left(\langle \mathbf{L} \partial_s^a \boldsymbol{\theta}, \partial_s^a \boldsymbol{\theta} \rangle_3 + \frac{49}{50} \|\partial_s^{a+1} \boldsymbol{\theta}\|_3^2 \right) + C_{\delta} (E_n + Z_n^2)^{1/2} E_n.$$

Note that by Corollary 3.31 $\|\operatorname{curl} \partial_s^a \theta\|_4^2 \leq (E_n + Z_n^2) E_n$. Using Lemma 3.33 we have

$$\begin{split} M &= \int_{\mathbb{R}^3} \left(\epsilon \bar{w}^{-2} |\nabla \cdot (\bar{w}^3 \partial_s^a \boldsymbol{\theta})|^2 + \bar{w}^3 |\partial_s^a \boldsymbol{\theta}|^2 \right) \mathrm{d}\mathbf{x} \\ &\geq \int_{\mathbb{R}^3} \left(\epsilon \frac{3}{4} \bar{w}^4 \left(|\nabla \partial_s^a \boldsymbol{\theta}|^2 - \frac{1}{2} |\operatorname{curl} \partial_s^a \boldsymbol{\theta}|^2 \right) - \epsilon \frac{3}{4} \bar{w}^3 \bar{w}'' |\partial_s^a \boldsymbol{\theta} \cdot \mathbf{e}_r|^2 + \bar{w}^3 |\partial_s^a \boldsymbol{\theta}|^2 \right) \mathrm{d}\mathbf{x} \\ &\geq \int_{\mathbb{R}^3} \left(\epsilon \frac{3}{4} \bar{w}^4 |\nabla \partial_s^a \boldsymbol{\theta}|^2 - \epsilon \frac{3}{4} \bar{w}^3 \bar{w}'' |\partial_s^a \boldsymbol{\theta} \cdot \mathbf{e}_r|^2 + \bar{w}^3 |\partial_s^a \boldsymbol{\theta}|^2 \right) \mathrm{d}\mathbf{x} - \epsilon C (E_n + Z_n^2) E_n \end{split}$$

Choosing ϵ small enough, we then have

$$M + (E_n + Z_n^2) S_n \gtrsim \int_{\mathbb{R}^3} \left(\bar{w}^4 |\nabla \partial_s^a \boldsymbol{\theta}|^2 + \bar{w}^3 |\partial_s^a \boldsymbol{\theta}|^2 \right) \mathrm{d}\mathbf{x}.$$

Next we will upgrade our estimate to control $\|\partial_s^a \partial^\beta \theta\|_3^2 + \|\partial_s^a \nabla \partial^\beta \theta\|_4^2$ for $|\beta| > 0$. First we will need the following lemma.

Lemma 3.35. For any vector field θ and $\epsilon > 0$,

$$\|\boldsymbol{\theta}\|_3^2 \lesssim \epsilon \|X_r \boldsymbol{\theta}\|_4^2 + (1+\epsilon^{-1}) \|\boldsymbol{\theta}\|_4^2.$$

Proof. We have

$$\begin{split} -\int_{B_R} |\boldsymbol{\theta}|^2 r \bar{w}' \bar{w}^3 \mathrm{d}\mathbf{x} &= -\frac{1}{4} \int_{B_R} |\boldsymbol{\theta}|^2 X_r \bar{w}^4 \mathrm{d}\mathbf{x} = \frac{1}{2} \int_{B_R} \bar{w}^4 \boldsymbol{\theta} \cdot X_r \boldsymbol{\theta} \, \mathrm{d}\mathbf{x} + \frac{3}{4} \int_{B_R} |\boldsymbol{\theta}|^2 \bar{w}^4 \mathrm{d}\mathbf{x} \\ &\leq \epsilon \|X_r \boldsymbol{\theta}\|_4^2 + \frac{1}{4} (3 + \epsilon^{-1}) \|\boldsymbol{\theta}\|_4^2. \end{split}$$

Now

$$\|\boldsymbol{\theta}\|_3^2 \lesssim \|\boldsymbol{\theta}\|_4^2 - \int_{B_R} |\boldsymbol{\theta}|^2 r \bar{w}' \bar{w}^3 \mathrm{d}\mathbf{x} \le \epsilon \|X_r \boldsymbol{\theta}\|_4^2 + \frac{1}{4} (7 + \epsilon^{-1}) \|\boldsymbol{\theta}\|_4^2.$$

Proposition 3.36. Let $n \ge 20$. Let θ be a solution of (3.62) in the sense of Theorem 3.11, given on its maximal interval of existence. Assume further that the energy, momentum, and irrotationality constraints (3.73), (3.74), and (3.75) hold respectively. Then for any $a + |\beta| \le n$ with $a, |\beta| > 0$ we have

$$\left\| \partial_s^a \partial^\beta \boldsymbol{\theta} \right\|_3^2 + \left\| \partial_s^a \nabla \partial^\beta \boldsymbol{\theta} \right\|_4^2$$

$$\lesssim |\boldsymbol{\varepsilon}|^{-2} \left(\frac{49}{50} \left\| \partial_s^{a+1} \partial^\beta \boldsymbol{\theta} \right\|_3^2 + \langle \mathbf{L} \partial_s^a \partial^\beta \boldsymbol{\theta}, \partial_s^a \partial^\beta \boldsymbol{\theta} \rangle \right) + CS_{n,|\beta|-1,0} + C_{\delta} (E_n + Z_n^2)^{1/2} E_n \quad (3.142)$$

Proof. By Proposition 3.27 and Lemma 3.33 we have

$$\begin{split} \|\nabla\partial_s^a \partial^\beta \boldsymbol{\theta}\|_4^2 &- \frac{1}{2} \|\operatorname{curl} \partial_s^a \partial^\beta \boldsymbol{\theta}\|_4^2 - \int_{B_R} \bar{w}'' \bar{w}^3 |\partial_s^a \partial^\beta \boldsymbol{\theta} \cdot \mathbf{e}_r|^2 \mathrm{d}\mathbf{x} \\ &\leq \int_{B_R} \frac{4}{3} \bar{w}^{-2} |\nabla \cdot (\bar{w}^3 \partial_s^a \partial^\beta \boldsymbol{\theta})|^2 \mathrm{d}\mathbf{x} \\ &\lesssim |\boldsymbol{\ell}|^{-2} \left(\langle \mathbf{L} \partial_s^a \partial^\beta \boldsymbol{\theta}, \partial_s^a \partial^\beta \boldsymbol{\theta} \rangle_3 + \frac{49}{50} \|\partial_s^{a+1} \partial^\beta \boldsymbol{\theta}\|_3^2 \right) + C_{\delta} (E_n + Z_n^2) E_n \end{split}$$

By Corollary 3.31 we have

$$\begin{split} \|\nabla\partial_s^a \partial^\beta \boldsymbol{\theta}\|_4^2 &- \int_{B_R} \bar{w}'' \bar{w}^3 |\partial_s^a \partial^\beta \boldsymbol{\theta} \cdot \mathbf{e}_r|^2 \mathrm{d} \mathbf{x} \\ &\lesssim |\boldsymbol{\delta}|^{-2} \left(\langle \mathbf{L} \partial_s^a \partial^\beta \boldsymbol{\theta}, \partial_s^a \partial^\beta \boldsymbol{\theta} \rangle_3 + \frac{49}{50} \|\partial_s^{a+1} \partial^\beta \boldsymbol{\theta}\|_3^2 \right) + CS_{n,|\beta|-1,0} + C_{\delta}(E_n + Z_n^2)E_n. \end{split}$$

Now by Lemma 3.35, we have $\frac{1}{2\epsilon} \|\partial_s^a \partial^\beta \theta\|_3^2 - \frac{1}{2} \|\nabla \partial_s^a \partial^\beta \theta\|_4^2 \lesssim_{\epsilon} \|\partial_s^a \partial^\beta \theta\|_4^2 \leq S_{n,|\beta|-1,0}$. Adding this and the above equation, and chosing ϵ small enough we get

$$\begin{aligned} \|\partial_s^a \partial^\beta \boldsymbol{\theta}\|_3^2 + \|\nabla \partial_s^a \partial^\beta \boldsymbol{\theta}\|_4^2 \\ \lesssim |\boldsymbol{\ell}|^{-2} \left(\langle \mathbf{L} \partial_s^a \partial^\beta \boldsymbol{\theta}, \partial_s^a \partial^\beta \boldsymbol{\theta} \rangle_3 + \frac{49}{50} \|\partial_s^{a+1} \partial^\beta \boldsymbol{\theta}\|_3^2 \right) + CS_{n,|\beta|-1,0} + C_{\delta}(E_n + Z_n^2)E_n. \end{aligned}$$

Remark 3.37. Estimate (3.142) features an order 1 term $CS_{n,|\beta|-1,0}$ on the right-hand side. This could be problematic for the closure of the estimates, but the key point is that this term is effectively decoupled, as it features one tangential derivative less. This will allows us later to close the estimates via induction on the order of derivatives in the problem.

3.5 Reduction to linear problem

In order to prove the bound (3.82), we will need to apply the coercivity estimates from Section 3.4. In particular, we must control the non-linear terms in order to effectively reduce the problem to a linear one. In Sections 3.5.1 and 3.5.2 we will prove high-order energy bounds for the nonlinear contributions from the pressure and the gravity term respectively. We will also prove high-order energy bounds for the full gravity term (including the linear part) in Section 3.5.2 that we will need for induction on radial derivatives. Then using these, we will reduce the full non-linear problem to the linear one in Section 3.5.3. This will then allow us to prove energy estimates and our main theorem in Section 3.6.

3.5.1 Estimating the non-linear part of the pressure term

In this subsection we will estimate the non-linear part of the pressure term $\partial_s^a X_r^b \partial^\beta \mathbf{P}$ (3.62), and show that it can be bounded by $(\mathcal{E}_n + \mathcal{Z}_n^2)^{1/2} \mathcal{E}_n$. More precisely, when doing energy estimates, the term $\langle \partial_s^a X_r^b \partial^\beta \mathbf{P}, \partial_s^a X_r^b \partial^\beta \theta \rangle$ and $\langle \partial_s^a X_r^b \partial^\beta \mathbf{P}, \partial_s^{a+1} X_r^b \partial^\beta \theta \rangle$ will arise, we will show that \mathbf{P} here can be reduced to $\mathbf{P}_{d,L}$ modulo remainder terms that can be estimated. We will use results from section 2.1.2.

Using Lemma 2.12, we will now estimate the difference between " \mathbf{P}_b " and " $\mathbf{P}_{b,L}$ ".

Proposition 3.38. Let $n \ge 20$ and $a + |\beta| + b \le n$ with a > 0. For any θ that satisfies our a priori assumption (3.71) we have

$$\begin{aligned} \left| \int_{0}^{s} \langle \mathbf{P}_{b} \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta}, \partial_{s}^{a+1} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta} \rangle_{3+b} \mathrm{d}\tau - \frac{1}{2} \langle \mathbf{P}_{b,L} \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta} \rangle_{3+b} \right|_{0}^{s} \right| \lesssim (\mathcal{E}_{n} + \mathcal{I}_{n}^{2})^{1/2} \mathcal{E}_{n} \\ \left| \int_{0}^{s} \langle \mathbf{P}_{b} \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta} \rangle_{3+b} \mathrm{d}\tau - \int_{0}^{s} \langle \mathbf{P}_{b,L} \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta} \rangle_{3+b} \mathrm{d}\tau \right| \lesssim (\mathcal{E}_{n} + \mathcal{I}_{n}^{2})^{1/2} \mathcal{E}_{n} \end{aligned}$$

Proof. By Lemma 2.12

$$\begin{split} \langle \mathbf{P}_{b} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}, \partial_{s}^{a+1} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta} \rangle_{3+b} \\ &= \int \left((\mathscr{A} \partial_{m} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}) \cdot (\mathscr{A} \partial_{m} \partial_{s}^{a+1} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}) + \frac{1}{3} (\operatorname{div}_{\mathscr{A}} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}) (\operatorname{div}_{\mathscr{A}} \partial_{s}^{a+1} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}) \\ &\quad - \frac{1}{2} [\operatorname{curl}_{\mathscr{A}} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}]_{j}^{m} [\operatorname{curl}_{\mathscr{A}} \partial_{s}^{a+1} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}]_{j}^{m} \Big) \mathcal{F}^{-1/3} \bar{w}^{4+b} \mathrm{d} \mathbf{x} \\ &= \int \left((\mathscr{A} \partial_{m} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}) \cdot \partial_{s} (\mathscr{A} \partial_{m} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}) + \frac{1}{3} (\operatorname{div}_{\mathscr{A}} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}) \partial_{s} (\operatorname{div}_{\mathscr{A}} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}) \\ &\quad - \frac{1}{2} [\operatorname{curl}_{\mathscr{A}} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}]_{j}^{m} \partial_{s} [\operatorname{curl}_{\mathscr{A}} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}]_{j}^{m} \Big) \mathcal{F}^{-1/3} \bar{w}^{4+b} \mathrm{d} \mathbf{x} \\ &\quad - \frac{1}{2} [\operatorname{curl}_{\mathscr{A}} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}]_{j}^{m} \partial_{s} [\operatorname{curl}_{\mathscr{A}} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}]_{j}^{m} \Big) \mathcal{F}^{-1/3} \bar{w}^{4+b} \mathrm{d} \mathbf{x} \\ &\quad - \frac{1}{2} [\operatorname{curl}_{\mathscr{A}} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}]_{j}^{m} \partial_{s} [\operatorname{curl}_{\mathscr{A}} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}]_{j}^{m} \Big) \mathcal{F}^{-1/3} \bar{w}^{4+b} \mathrm{d} \mathbf{x} \\ &\quad + \mathcal{R} [(E_{n} + Z_{n}^{2})^{1/2} E_{n}] \\ &= \frac{1}{2} \partial_{s} \int \left(|\mathscr{A} \partial_{m} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}|^{2} + \frac{1}{3} |\operatorname{div}_{\mathscr{A}} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}|^{2} - \frac{1}{2} |[\operatorname{curl}_{\mathscr{A}} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}]|^{2} \right) \bar{w}^{4+b} \mathrm{d} \mathbf{x} \\ &\quad + \partial_{s} \mathcal{R} [(E_{n} + Z_{n}^{2})^{1/2} E_{n}] + \mathcal{R} [(E_{n} + Z_{n}^{2})^{1/2} E_{n}] \\ &= \frac{1}{2} \partial_{s} \langle \mathbf{P}_{b,L} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta} \rangle_{3+b} + \partial_{s} \mathcal{R} [(E_{n} + Z_{n}^{2})^{1/2} E_{n}] + \mathcal{R} [(E_{n} + Z_{n}^{2})^{1/2} E_{n}] \\ &= \frac{1}{2} \partial_{s} \langle \mathbf{P}_{b,L} \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} X_{r}^{b} \partial_{\beta}^{\beta} \boldsymbol{\theta} \rangle_{3+b} + \partial_{s} \mathcal{R} [(E_{n} + Z_{n}^{2})^{1/2} E_{n}] + \mathcal{R} [(E_{n} + Z_{n}^{2})^{1/2} E_{n}] \\ &= \frac{1}{2} \partial_{s} \langle$$

where we recall notation $\Re[\star]$ introduced in Definition 2.2. Integrating in time we get the first equation. For the second equation, note that

$$\langle \mathbf{P}_b \partial_s^a X_r^b \partial^\beta \boldsymbol{\theta}, \partial_s^a X_r^b \partial^\beta \boldsymbol{\theta} \rangle_{3+b} = \langle \mathbf{P}_{b,L} \partial_s^a X_r^b \partial^\beta \boldsymbol{\theta}, \partial_s^a X_r^b \partial^\beta \boldsymbol{\theta} \rangle_3 + \Re[(E_n + Z_n^2)^{1/2} E_n].$$

Integrating in time we get the second equation.

And now we will estimate the difference between "**P**" and "**P**_b". **Proposition 3.39.** Let $n \ge 20$. For any θ that satisfies our a priori assumption (3.71) we have

i. For $a + |\beta| \le n$ with a > 0 we have

$$\left| \int_{0}^{s} \langle \partial_{s}^{a} \partial^{\beta} \mathbf{P} - \mathbf{P}_{0} \partial_{s}^{a} \partial^{\beta} \theta, \partial_{s}^{a+1} \partial^{\beta} \theta \rangle_{3} \mathrm{d}\tau \right| \lesssim \delta_{n,|\beta|-1,0}^{1/2} \delta_{n,|\beta|,0}^{1/2} + (\mathcal{E}_{n} + \mathcal{Z}_{n}^{2})^{1/2} \mathcal{E}_{n} \\ \left| \int_{0}^{s} \langle \partial_{s}^{a} \partial^{\beta} \mathbf{P} - \mathbf{P}_{0} \partial_{s}^{a} \partial^{\beta} \theta, \partial_{s}^{a} \partial^{\beta} \theta \rangle_{3} \mathrm{d}\tau \right| \lesssim \delta_{n,|\beta|-1,0}^{1/2} \delta_{n,|\beta|,0}^{1/2} + (\mathcal{E}_{n} + \mathcal{Z}_{n}^{2})^{1/2} \mathcal{E}_{n}$$

ii. For $a + |\beta| + b \le n$ with a > 0 we have

$$\begin{split} \left| \int_{0}^{s} \langle \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \mathbf{P} - \mathbf{P}_{b} \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta}, \partial_{s}^{a+1} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta} \rangle_{3+b} \mathrm{d}\tau \right| \\ \lesssim (S_{n,|\beta|+b-1}^{1/2} + S_{n,|\beta|+b,b-1}^{1/2}) S_{n,|\beta|+b}^{1/2} + (\mathcal{E}_{n} + \mathcal{Z}_{n}^{2})^{1/2} \mathcal{E}_{n} \end{split}$$

$$\begin{split} \left| \int_{0}^{s} \langle \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \mathbf{P} - \mathbf{P}_{b} \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \theta, \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \theta \rangle_{3+b} \mathrm{d}\tau \right| \\ \lesssim (\mathcal{S}_{n,|\beta|+b-1}^{1/2} + \mathcal{S}_{n,|\beta|+b,b-1}^{1/2}) \mathcal{S}_{n,|\beta|+b}^{1/2} + (\mathcal{C}_{n} + \mathcal{Z}_{n}^{2})^{1/2} \mathcal{C}_{n} \\ \end{split}$$

Proof. i. Using Lemma 2.9 we have

$$\begin{split} & \left| \int_{0}^{s} \langle \partial_{s}^{a} \partial^{\beta} \mathbf{P} - \mathbf{P}_{L} \partial_{s}^{a} \partial^{\beta} \theta, \partial_{s}^{a+1} \partial^{\beta} \theta \rangle_{3} \mathrm{d}\tau \right| \\ & \leq \left| \int_{0}^{s} \int_{B_{R}} \partial_{k} (\bar{w}^{4}(T_{R:a,\beta})_{i}^{k}) \partial_{s}^{a+1} \partial^{\beta} \theta^{i} \, \mathrm{d}x \mathrm{d}\tau \right| \\ & + \left| \int_{0}^{s} \int_{B_{R}} \sum_{|\beta'| \leq |\beta| - 1} \langle C \nabla(\bar{w}^{4} \partial_{s}^{a} \partial^{\beta'} T) \rangle \langle \partial_{s}^{a+1} \partial^{\beta} \theta \rangle \, \mathrm{d}x \mathrm{d}\tau \right| \\ & \leq \left| \int_{0}^{s} \int_{B_{R}} \bar{w}^{4} \partial_{s}(T_{R:a,\beta})_{i}^{k} \partial_{s}^{a} \partial_{k} \partial^{\beta} \theta^{i} \, \mathrm{d}x \mathrm{d}\tau \right| + \left| \int_{0}^{s} \int_{B_{R}} \sum_{|\beta'| \leq |\beta| - 1} \bar{w}^{4} \langle C \partial_{s}^{a+1} \partial^{\beta'} T \rangle \langle \partial_{s}^{a} \nabla \partial^{\beta} \theta \rangle \, \mathrm{d}x \mathrm{d}\tau \right| \\ & + (E_{n} + Z_{n}^{2})(0)^{1/2} E_{n}(0) + (E_{n} + Z_{n}^{2})(s)^{1/2} E_{n}(s) \\ & + S_{n,|\beta| - 1,0}(0)^{1/2} S_{n,|\beta|,0}(0)^{1/2} + S_{n,|\beta| - 1,0}(s)^{1/2} S_{n,|\beta|,0}(s)^{1/2} \\ & \lesssim \delta_{n,|\beta| - 1,0}^{1/2} \delta_{n,|\beta|,0}^{1/2} + (\mathfrak{C}_{n} + \mathfrak{T}_{n}^{2})^{1/2} \mathfrak{C}_{n}. \end{split}$$

Proof of the second formula is similar and easier.

ii. By Lemma 2.10 we need to estimate the following.

$$\begin{split} \left| \int_{0}^{s} \int_{B_{R}} \left(\sum_{\substack{b' \leq b \\ |\beta'| \leq |\beta| - 1}} + \sum_{\substack{b' \leq b - 1 \\ |\beta'| \leq |\beta| + 1}} \right) \langle C\omega \partial_{s}^{a} X_{r}^{b'} \partial^{\beta'} T \rangle \langle \partial_{s}^{a+1} X_{r}^{b} \partial^{\beta} \theta \rangle \bar{w}^{3+b} \mathrm{dx} \mathrm{d\tau} \right| \\ \lesssim \left| \int_{0}^{s} \int_{B_{R}} \sum_{\substack{b' \leq b \\ |\beta'| \leq |\beta| - 1}} \langle C\omega T_{T} [\partial_{s}^{a+1} X_{r}^{b'} \partial^{\beta'} \nabla \theta] \rangle \langle \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \theta \rangle \bar{w}^{3+b} \mathrm{dx} \mathrm{d\tau} \right| \\ + \left| \int_{0}^{s} \int_{B_{R}} \sum_{\substack{b' \leq b - 1 \\ |\beta'| \leq |\beta| + 1}} \langle C\omega T_{T} [\partial_{s}^{a} X_{r}^{b'} \partial^{\beta'} \nabla \theta] \rangle \langle \partial_{s}^{a+1} X_{r}^{b} \partial^{\beta} \theta \rangle \bar{w}^{3+b} \mathrm{dx} \mathrm{d\tau} \right| \\ + \delta_{n,|\beta|+b-1}^{1/2} \delta_{n,|\beta|+b}^{1/2} + (\mathcal{E}_{n} + \mathcal{E}_{n}^{2})^{1/2} \mathcal{E}_{n} \\ \lesssim (\delta_{n,|\beta|+b-1}^{1/2} + \delta_{n,|\beta|+b,b-1}^{1/2}) \delta_{n,|\beta|+b}^{1/2} + (\mathcal{E}_{n} + \mathcal{E}_{n}^{2})^{1/2} \mathcal{E}_{n} \\ \left| \int_{0}^{s} \int_{$$

$$\begin{split} \left| \int_{0}^{s} \int_{B_{R}} \left(\sum_{\substack{b' \leq b-1 \\ |\beta'| \leq |\beta|-1}} + \sum_{\substack{b' \leq b-2 \\ |\beta'| \leq |\beta|}} \right) \langle C \partial_{s}^{a} X_{r}^{b'} \partial^{\beta'} \nabla T \rangle \langle \partial_{s}^{a+1} X_{r}^{b} \partial^{\beta} \theta \rangle \bar{w}^{3+b} \mathrm{dx} \mathrm{d\tau} \right| \\ \lesssim \left| \int_{0}^{s} \int_{B_{R}} \left(\sum_{\substack{b' \leq b-1 \\ |\beta'| \leq |\beta|-1}} + \sum_{\substack{b' \leq b-2 \\ |\beta'| \leq |\beta|}} \right) \langle C \partial_{s}^{a+1} X_{r}^{b'} \partial^{\beta'} T \rangle \langle \partial_{s}^{a} \nabla X_{r}^{b} \partial^{\beta} \theta \rangle \bar{w}^{3+b} \mathrm{dx} \mathrm{d\tau} \right| \\ + \left| \int_{0}^{s} \int_{B_{R}} \left(\sum_{\substack{b' \leq b-1 \\ |\beta'| \leq |\beta|-1}} + \sum_{\substack{b' \leq b-2 \\ |\beta'| \leq |\beta|}} \right) \langle C \partial_{s}^{a+1} X_{r}^{b'} \partial^{\beta'} T \rangle \langle \omega \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \theta \rangle \bar{w}^{2+b} \mathrm{dx} \mathrm{d\tau} \right| \\ + \mathcal{S}_{n,|\beta|+b-1}^{1/2} \mathcal{S}_{n,|\beta|+n}^{1/2} + (\mathcal{C}_{n} + \mathcal{T}_{n}^{2})^{1/2} \mathcal{C}_{n} \\ \lesssim \mathcal{S}_{n,|\beta|+b-1}^{1/2} \mathcal{S}_{n,|\beta|+n}^{1/2} + (\mathcal{C}_{n} + \mathcal{T}_{n}^{2})^{1/2} \mathcal{C}_{n} \end{split}$$

$$\begin{split} \left| \int_{0}^{s} \int_{B_{R}} \left(\sum_{\substack{b' \leq b \\ |\beta'| \leq |\beta| - 1}} + \sum_{\substack{b' \leq b - 1 \\ |\beta'| \leq |\beta|}} \right) \langle C\bar{w}X_{r}^{b'}\partial^{\beta'}\nabla T \rangle \langle \partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\theta \rangle \bar{w}^{3+b} \mathrm{dxd\tau} \right| \\ \lesssim \left| \int_{0}^{s} \int_{B_{R}} \left(\sum_{\substack{b' \leq b \\ |\beta'| \leq |\beta| - 1}} + \sum_{\substack{b' \leq b - 1 \\ |\beta'| \leq |\beta|}} \right) \langle C\partial_{s}^{a+1}X_{r}^{b'}\partial^{\beta'}T \rangle \langle \partial_{s}^{a}\nabla X_{r}^{b}\partial^{\beta}\theta \rangle \bar{w}^{4+b} \mathrm{dxd\tau} \right| \\ + \left| \int_{0}^{s} \int_{B_{R}} \left(\sum_{\substack{b' \leq b \\ |\beta'| \leq |\beta| - 1}} + \sum_{\substack{b' \leq b - 1 \\ |\beta'| \leq |\beta|}} \right) \langle C\partial_{s}^{a+1}X_{r}^{b'}\partial^{\beta'}T \rangle \langle \omega\partial_{s}^{a}X_{r}^{b}\partial^{\beta}\theta \rangle \bar{w}^{3+b} \mathrm{dxd\tau} \right| \\ + \mathcal{S}_{n,|\beta|+b-1}^{1/2} \mathcal{S}_{n,|\beta|+n}^{1/2} + (\mathcal{C}_{n} + \mathcal{Z}_{n}^{2})^{1/2} \mathcal{C}_{n} \\ \lesssim \mathcal{S}_{n,|\beta|+b-1}^{1/2} \mathcal{S}_{n,|\beta|+n}^{1/2} + (\mathcal{C}_{n} + \mathcal{Z}_{n}^{2})^{1/2} \mathcal{C}_{n} \end{split}$$

This proves the first formula. Proof of the second formula is similar and easier.

3.5.2 Estimating the linear and non-linear part of the gravity term

In this subsection we will estimate the gravity term $\partial_s^a X_r^b \partial^\beta \mathbf{G}$ (3.62) and show that it can be bounded by E_n . We will also estimate the non-linear part of $\partial_s^a \partial^\beta \mathbf{G}$, and show that it can be bounded by $(\mathcal{E}_n + \mathcal{E}_n^2)^{1/2} \mathcal{E}_n$. We will use results from Section 2.1.1.

Since the gravity term is a non-local term, we need to estimate convolution-like operator. However, rather than the convolution kernel $|\mathbf{x} - \mathbf{z}|^{-1}$ we actually need to estimate $|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|^{-1}$. Lemma 2.5 and the following lemma tell us how to reduce the latter to the former, which will allows us to estimate using the Young's convolution inequality.

Lemma 3.40. Let $\boldsymbol{\xi}$ and $\boldsymbol{\theta}$ be as in (2.35), and $\boldsymbol{\theta}$ satisfies our a priori assumption (3.71). Let $n \geq 21$ and $a + |\beta| \leq n$ with a > 0.

i. When $a + |\beta| > n/2$ we have

$$\begin{split} \left| \partial_s^a (\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^\beta \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} \right) \right| \lesssim \frac{1}{|\mathbf{x} - \mathbf{z}|^2} \sum_{\substack{n/2 < a' + |\gamma| \le n \\ a' > 0}} |\partial_s^{a'} \partial_{\mathbf{x}}^{\gamma} \boldsymbol{\theta}(\mathbf{x}) - \partial_s^{a'} \partial_{\mathbf{z}}^{\gamma} \boldsymbol{\theta}(\mathbf{z}) \\ + \frac{E_n^{1/2}}{|\mathbf{x} - \mathbf{z}|^2} \sum_{n/2 < |\gamma| \le n} |\partial_{\mathbf{x}}^{\gamma} \boldsymbol{\xi}(\mathbf{x}) - \partial_{\mathbf{z}}^{\gamma} \boldsymbol{\xi}(\mathbf{z})| \end{split}$$

ii. When $|\beta| > n/2$ we have

$$\left| (\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^{\beta} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} \right) \right| \lesssim \frac{1}{|\mathbf{x} - \mathbf{z}|^2} \sum_{n/2 < |\gamma| \le n} |\partial_{\mathbf{x}}^{\gamma} \boldsymbol{\xi}(\mathbf{x}) - \partial_{\mathbf{z}}^{\gamma} \boldsymbol{\xi}(\mathbf{z})|$$

iii. When $a + |\beta| \le n/2$ we have

$$\left|\partial_{i,\mathbf{z}}\partial_s^a(\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^\beta \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|}\right)\right| \lesssim \frac{E_n^{1/2}}{|\mathbf{x} - \mathbf{z}|^2}$$

iv. When $|\beta| \leq n/2$ we have

$$\left|\partial_{i,\mathbf{z}}(\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^{\beta}\left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|}\right)\right| \lesssim \frac{1}{|\mathbf{x} - \mathbf{z}|^2}$$

Proof. These follows from Lemma 2.5, the embedding theorems A.8 and A.9, the a priori bounds $E_n, Z_n \lesssim 1$ (3.71), and the following.

$$\partial_{s}^{a}(\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^{\beta} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} \right) = \sum_{m=1}^{a+|\beta|} \sum_{\substack{\sum_{i=1}^{m} (a_{i}+a_{i}')=a \\ \sum_{i=1}^{m} (\beta_{i}+\beta_{i}')=\beta \\ |a_{i}|+|\beta_{i}|>0}} \frac{(-1)^{m}(2m)!}{m!2^{m}} \frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|^{1+2m}} \prod_{i=1}^{m} (\partial_{s}^{a_{i}}\partial_{\mathbf{x}}^{\beta_{i}}\boldsymbol{\xi}(\mathbf{x}) - \partial_{s}^{a_{i}}\partial_{\mathbf{z}}^{\beta_{i}}\boldsymbol{\xi}(\mathbf{z})) \cdot (\partial_{s}^{a_{i}'}\partial_{\mathbf{x}}^{\beta_{i}'}\boldsymbol{\xi}(\mathbf{x}) - \partial_{s}^{a_{i}'}\partial_{\mathbf{x}}^{\beta_{i}'}\boldsymbol{\xi}(\mathbf{z})).$$

We will now prove the main results of this subsection.

Proposition 3.41. Let $n \ge 21$ and suppose θ satisfies our a priori assumption (3.71). For $a + |\beta| + b \le n$ with a > 0 we have

$$\|\partial_s^a X_r^b \partial^\beta \mathbf{G}\|_{3+b}^2 \lesssim E_n. \tag{3.143}$$

Proof. By definition

$$\mathbf{G} = \mathscr{K}_{\boldsymbol{\xi}} \nabla \cdot (\mathscr{A} \bar{w}^3) - \mathscr{K} \nabla \bar{w}^3 = -\int_{\mathbb{R}^3} \frac{\partial_k (\mathscr{A}^k \bar{w}^3)}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} d\mathbf{z} + \int_{\mathbb{R}^3} \frac{\nabla \bar{w}^3}{|\mathbf{x} - \mathbf{z}|} d\mathbf{z}$$

Consider first when b = 0. Since a > 0, by Lemma 2.8 we have

$$\begin{split} \partial_s^a \partial^\beta \mathbf{G}(\mathbf{x}) &= \partial_s^a \partial^\beta \mathcal{H}_{\boldsymbol{\xi}} \nabla \cdot (\mathscr{A} \bar{w}^3)(\mathbf{x}) = -\partial_s^a \partial^\beta \int_{\mathbb{R}^3} \frac{\partial_k (\mathscr{A}^k \bar{w}^3)}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} \mathrm{d}\mathbf{z} \\ &= -\int_{\mathbb{R}^3} \sum_{\substack{a_1 + a_2 = a \\ \beta_1 + \beta_2 = \beta}} \partial_s^{a_1} (\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^{\beta_1} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|}\right) \partial_s^{a_2} \partial_{\mathbf{z}}^{\beta_2} \partial_k (\mathscr{A}^k \bar{w}^3)(\mathbf{z}) \mathrm{d}\mathbf{z} \\ &= -\int_{\mathbb{R}^3} \sum_{\substack{a_1 + a_2 = a \\ \beta_1 + \beta_2 = \beta}} \partial_s^{a_1} (\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^{\beta_1} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|}\right) \partial_s^{a_2} \partial_{\mathbf{z}}^{\beta_2} \partial_k (\mathscr{A}^k \bar{w}^3)(\mathbf{z}) \mathrm{d}\mathbf{z} \\ &- \int_{\mathbb{R}^3} \sum_{\substack{a_1 + a_2 = a \\ \beta_1 + \beta_2 = \beta}} \partial_s^{a_1} (\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^{\beta_1} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|}\right) \partial_s^{a_2} \partial_{\mathbf{z}}^{\beta_2} \partial_k (\mathscr{A}^k \bar{w}^3)(\mathbf{z}) \mathrm{d}\mathbf{z} \\ &= -\int_{\mathbb{R}^3} \sum_{\substack{a_1 + a_2 = a \\ \beta_1 + \beta_2 = \beta}} \partial_s^{a_1} (\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^{\beta_1} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|}\right) \partial_s^{a_2} \partial_{\mathbf{z}}^{\beta_2} \partial_k (\mathscr{A}^k \bar{w}^3)(\mathbf{z}) \mathrm{d}\mathbf{z} \\ &+ \int_{\mathbb{R}^3} \sum_{\substack{a_1 + a_2 = a \\ \beta_1 + \beta_2 = \beta}} \partial_s^{a_1} (\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^{\beta_1} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|}\right) (\langle \partial_s^{a_2} \partial_{\mathbf{z}}^{\beta_2} \partial_k (\mathscr{A}^k \bar{w}^3)(\mathbf{z}) \mathrm{d}\mathbf{z} \\ &+ \int_{\mathbb{R}^3} \sum_{\substack{a_1 + a_2 = a \\ \beta_1 + \beta_2 \leq \beta}} \langle \nabla_{\mathbf{z}} \rangle \partial_s^{a_1} (\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^{\beta_1} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|}\right) (\langle \partial_s^{a_2} \partial_{\mathbf{z}}^{\beta_2} \mathcal{A} \rangle \bar{w}^3)(\mathbf{z}) \mathrm{d}\mathbf{z} \end{split}$$

Now using Lemma 3.40, we get

$$egin{aligned} |\partial_s^a \partial^eta \mathbf{G}(\mathbf{x})| \lesssim & \int_{\mathbb{R}^3} rac{1}{|\mathbf{x} - \mathbf{z}|^2} \sum_{n/2 < a' + |\gamma| \le n} |\partial_s^{a'} \partial_{\mathbf{x}}^\gamma m{ heta}(\mathbf{x}) - \partial_s^{a'} \partial_{\mathbf{z}}^\gamma m{ heta}(\mathbf{z})| \mathrm{d}\mathbf{z} \ & + \int_{\mathbb{R}^3} rac{E_n^{1/2}}{|\mathbf{x} - \mathbf{z}|^2} \sum_{n/2 < |\gamma| \le n} |\partial_{\mathbf{x}}^\gamma m{\xi}(\mathbf{x}) - \partial_{\mathbf{z}}^\gamma m{\xi}(\mathbf{z})| \mathrm{d}\mathbf{z} \end{aligned}$$

$$+\int_{\mathbb{R}^3}\sum_{\substack{0< a_2\leq a\\\beta_2\leq\beta}}\frac{1}{|\mathbf{x}-\mathbf{z}|^2}(\langle\partial_s^{a_2}\partial\!\!\!/ \mathbf{z}^{\beta_2}\mathcal{A}\rangle\bar{w}^3)(\mathbf{z})\mathrm{d}\mathbf{z}+\int_{\mathbb{R}^3}\sum_{\beta_2\leq\beta}\frac{E_n^{1/2}}{|\mathbf{x}-\mathbf{z}|^2}(\langle\partial\!\!\!/ \mathbf{z}^{\beta_2}\mathcal{A}\rangle\bar{w}^3)(\mathbf{z})\mathrm{d}\mathbf{z}$$

Now using Young's convolution inequality we get

$$\|\partial_s^a \partial^\beta \mathbf{G}(\mathbf{x})\|_{L^2(\mathbb{R}^3)} \lesssim E_n^{1/2}$$

Hence $\|\partial_s^a \partial^\beta \mathbf{G}\|_3^2 \lesssim E_n$. From the above proof, with small modification, we can further see that

$$\begin{aligned} \|\partial_s^a \partial^\beta \mathbf{G}(\mathbf{x})\|_{L^\infty(\mathbb{R}^3)} &\lesssim E_n^{1/2} \quad \text{when} \quad a+|\beta| \le n/2\\ \|\partial^\beta \mathbf{G}(\mathbf{x})\|_{L^\infty(\mathbb{R}^3)} &\lesssim 1 \quad \text{when} \quad |\beta| \le n/2. \end{aligned}$$

Now we deal with the case b > 0. Let

$$W_{n,c} = \sum_{\substack{a+|\beta|+b\leq n\\a>0\\|\beta|+b\leq c}} \|\partial_s^a X_r^b \partial^\beta \mathbf{G}\|_{3+b}^2$$
$$W_{n,c,d} = \sum_{\substack{a+|\beta|+b\leq n\\\beta|+b\leq c\\b\leq d}} \|\partial_s^a X_r^b \partial^\beta \mathbf{G}\|_{3+b}^2$$
$$V_{n,c} = \sum_{\substack{a+|\beta|+b\leq c\\\beta|+b\leq c\\|\beta|+b\leq c}} \sup_{\mathbb{R}^3} \left(\bar{w}^b |\partial_s^a X_r^b \partial^\beta \mathbf{G}|^2\right)$$
$$V_{n,c,d} = \sum_{\substack{a+|\beta|+b\leq n\\\beta|+b\leq c\\\beta|+b\leq c\\\beta|+b\leq c\\\beta|+b\leq c\\\beta|+b\leq c}} \sup_{\mathbb{R}^3} \left(\bar{w}^b |\partial_s^a X_r^b \partial^\beta \mathbf{G}|^2\right).$$

For $a + |\beta| + b \le n/2$, using the above lemmas 2.6 and 2.7 we have

$$\begin{split} \bar{w}^{b} |\partial_{s}^{a} X_{r}^{b} \partial^{\beta} \mathbf{G}|^{2} &\lesssim \bar{w}^{b} |r \partial_{s}^{a} \nabla \cdot X_{r}^{b-1} \partial^{\beta} \mathbf{G}|^{2} + \bar{w}^{b} |r \partial_{s}^{a} \nabla \times X_{r}^{b-1} \partial^{\beta} \mathbf{G}|^{2} + \sum_{k=1}^{3} \bar{w}^{b} |\partial_{s}^{a} X_{r}^{b-1} \partial_{k} \partial^{\beta} \mathbf{G}|^{2} \\ &\lesssim \bar{w}^{b} |r \partial_{s}^{a} X_{r}^{b-1} \partial^{\beta} \nabla \cdot \mathbf{G}|^{2} + \bar{w}^{b} |r \partial_{s}^{a} X_{r}^{b-1} \partial^{\beta} \nabla \times \mathbf{G}|^{2} + V_{n,b+|\beta|-1} + V_{n,b+|\beta|,b-1} \\ &\lesssim \bar{w}^{b} |r \partial_{s}^{a} X_{r}^{b-1} \partial^{\beta} (I - \mathcal{A}) \nabla \cdot \mathbf{G}|^{2} + \bar{w}^{b} |r \partial_{s}^{a} X_{r}^{b-1} \partial^{\beta} (I - \mathcal{A}) \nabla \times \mathbf{G}|^{2} \\ &\quad + E_{n} + V_{n,b+|\beta|-1} + V_{n,b+|\beta|,b-1} \\ &\lesssim \bar{w}^{b} |r (I - \mathcal{A}) \partial_{s}^{a} X_{r}^{b-1} \partial^{\beta} \nabla \cdot \mathbf{G}|^{2} + \bar{w}^{b} |r (I - \mathcal{A}) \partial_{s}^{a} X_{r}^{b-1} \partial^{\beta} \nabla \times \mathbf{G}|^{2} \\ &\quad + E_{n} + V_{n,b+|\beta|-1} + V_{n,b+|\beta|,b-1} \\ &\lesssim \bar{w}^{b} (E_{n} + Z_{n}^{2}) |r \partial_{s}^{a} X_{r}^{b-1} \partial^{\beta} \nabla \mathbf{G}|^{2} + E_{n} + V_{n,b+|\beta|-1} + V_{n,b+|\beta|,b-1} \end{split}$$

So

$$V_{n,b+|\beta|,b} \lesssim (E_n + Z_n^2) V_{n,b+|\beta|,b} + E_n + V_{n,b+|\beta|-1} + V_{n,b+|\beta|,b-1}$$

By a priori assumption (3.71), we have $E_n + Z_n^2 \ll 1$, so

$$V_{n,b+|\beta|,b} \lesssim E_n + V_{n,b+|\beta|-1} + V_{n,b+|\beta|,b-1}.$$

We know $V_{n',c,0} \lesssim E_n$ for all $c \leq n' \leq n/2$, so by induction we get $V_{n',c,d} \lesssim E_n$ for all $d \leq c \leq n' \leq n/2$. Now for $a + |\beta| + b \leq n$, using the above lemmas 2.6 and 2.7 and results for V we have

$$\|\partial_s^a X_r^b \partial^\beta \mathbf{G}\|_{3+b}^2 \lesssim \|r\partial_s^a \nabla \cdot X_r^{b-1} \partial^\beta \mathbf{G}\|_{3+b}^2 + \|r\partial_s^a \nabla \times X_r^{b-1} \partial^\beta \mathbf{G}\|_{3+b}^2 + \sum_{k=1}^3 \|\partial_s^a X_r^{b-1} \partial_k \partial^\beta \mathbf{G}\|_{3+b}^2$$

$$\lesssim \|r\partial_s^a X_r^{b-1} \partial^\beta \nabla \cdot \mathbf{G}\|_{3+b}^2 + \|r\partial_s^a X_r^{b-1} \partial^\beta \nabla \times \mathbf{G}\|_{3+b}^2 + W_{n,b+|\beta|-1} + W_{n,b+|\beta|,b-1}$$

$$\lesssim \|r\partial_s^a X_r^{b-1} \partial^\beta (1-\mathcal{A}) \nabla \cdot \mathbf{G}\|_{3+b}^2 + \|r\partial_s^a X_r^{b-1} \partial^\beta (1-\mathcal{A}) \nabla \times \mathbf{G}\|_{3+b}^2$$

$$+ E_n + W_{n,b+|\beta|-1} + W_{n,b+|\beta|,b-1}$$

$$\lesssim \|r(1-\mathcal{A}) \partial_s^a X_r^{b-1} \partial^\beta \nabla \cdot \mathbf{G}\|_{3+b}^2 + \|r(1-\mathcal{A}) \partial_s^a X_r^{b-1} \partial^\beta \nabla \times \mathbf{G}\|_{3+b}^2$$

$$+ E_n + W_{n,b+|\beta|-1} + W_{n,b+|\beta|,b-1}$$

$$\lesssim (E_n + Z_n^2) \|r\partial_s^a X_r^{b-1} \partial^\beta \nabla \mathbf{G}\|_{3+b}^2 + E_n + W_{n,b+|\beta|-1} + W_{n,b+|\beta|,b-1}$$

So

$$W_{n,b+|\beta|,b} \lesssim (E_n + Z_n^2) W_{n,b+|\beta|,b} + E_n + W_{n,b+|\beta|-1} + W_{n,b+|\beta|,b-1}$$

By a priori assumption (3.71), we have $E_n + Z_n^2 \ll 1$, so

$$W_{n,b+|\beta|,b} \lesssim E_n + W_{n,b+|\beta|-1} + W_{n,b+|\beta|,b-1}.$$

We know $W_{n,c,0} \lesssim E_n$ for all c, so by induction we get $W_{n,c,d} \lesssim E_n$ for all $d \leq c \leq n$.

We are now in the position to estimate the difference between the high order derivatives of nonlinear gravity term G (2.38) and its linearised part G_L (3.85).

Proposition 3.42. Let $n \ge 21$ and suppose θ satisfies our a priori assumption (3.71). For $a + |\beta| \le n$ with a > 0 we have

$$\left| \int_0^s \langle \partial_s^a \partial^\beta \mathbf{G} - \mathbf{G}_L \partial_s^a \partial^\beta \boldsymbol{\theta}, \partial_s^{a+1} \partial^\beta \boldsymbol{\theta} \rangle_3 \mathrm{d}\tau \right| \lesssim (\mathscr{C}_n + \mathfrak{X}_n^2)^{1/2} \mathscr{C}_n \\ \left| \int_0^s \langle \partial_s^a \partial^\beta \mathbf{G} - \mathbf{G}_L \partial_s^a \partial^\beta \boldsymbol{\theta}, \partial_s^a \partial^\beta \boldsymbol{\theta} \rangle_3 \mathrm{d}\tau \right| \lesssim (\mathscr{C}_n + \mathfrak{X}_n^2)^{1/2} \mathscr{C}_n$$

Proof. Since $\|\partial_s^{a+1}\partial^\beta \theta\|_3 + \|\partial_s^a \partial^\beta \theta\|_3 \lesssim E_n^{1/2}$, it suffice to prove that

$$\|\partial_s^a \partial^\beta \mathbf{G} - \mathbf{G}_L \partial_s^a \partial^\beta \boldsymbol{\theta}\|_3 \lesssim (E_n + Z_n^2)^{1/2} E_n^{1/2}.$$

Recall from Lemma 3.13 that

$$\mathbf{G} - \mathbf{G}_{L}\boldsymbol{\theta} = \underbrace{\mathcal{K}_{\boldsymbol{\xi}}(\mathcal{A}_{l}^{i}(\partial_{k}\theta^{l})(\nabla\theta^{k})\partial_{i}\bar{w}^{3} - \bar{w}^{3}(\mathcal{A}_{m}^{i}\mathcal{A}_{\bullet}^{l} - I_{m}^{i}I_{\bullet}^{l})\partial_{i}\partial_{l}\theta^{m})}_{:=M_{1}}_{:=M_{1}} \underbrace{-(\mathcal{K}_{\boldsymbol{\xi}} - \mathcal{K})\partial_{i}(\bar{w}^{3}\nabla\theta^{i})}_{:=M_{2}} + \underbrace{(\mathcal{K}_{\boldsymbol{\xi}} - \mathcal{K} - \mathcal{K}_{\boldsymbol{\xi}}^{(1)})\nabla\bar{w}^{3}}_{:=M_{3}}.$$

Now $\|\partial_s^a \partial^\beta M_1\|_3$ can be estimated in a similar way as the previous Proposition 3.41, and $\|\partial_s^a \partial^\beta M_3\|_3$ can be estimated in the same way as in Lemma 3.26 in equation (3.126). Now in the same way as in Lemma 3.25 and recalling K_1 (3.89) we can show that

$$\begin{aligned} |\partial_s^a (\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^\beta K_1(\mathbf{x}, \mathbf{z})| &\lesssim \frac{1}{|\mathbf{x} - \mathbf{z}|^2} \sum_{\substack{0 < a' \leq a \\ \beta' \leq \beta}} |\partial_s^{a'} \partial_{\beta'} \theta(\mathbf{x}) - \partial_s^{a'} \partial_{\beta'} \theta(\mathbf{z})| + \frac{E_n^{1/2}}{|\mathbf{x} - \mathbf{z}|^2} \sum_{\beta' \leq \beta} |\partial_{\beta'} \theta(\mathbf{x}) - \partial_{\beta'} \theta(\mathbf{z})| \\ |(\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^\beta K_1(\mathbf{x}, \mathbf{z})| &\lesssim \frac{1}{|\mathbf{x} - \mathbf{z}|^2} \sum_{\beta' \leq \beta} |\partial_{\beta'} \theta(\mathbf{x}) - \partial_{\beta'} \theta(\mathbf{z})| \end{aligned}$$

And when $a + |\beta| \le n/2$,

$$|\partial_{i,\mathbf{z}}\partial_s^a(\mathbf{\partial}_{\mathbf{x}} + \mathbf{\partial}_{\mathbf{z}})^\beta K_1(\mathbf{x}, \mathbf{z})| \lesssim \frac{E_n^{1/2}}{|\mathbf{x} - \mathbf{z}|^2}$$

and when $|\beta| \leq n/2$,

$$|\partial_{i,\mathbf{z}}(\boldsymbol{\partial}_{\mathbf{x}} + \boldsymbol{\partial}_{\mathbf{z}})^{\beta} K_1(\mathbf{x}, \mathbf{z})| \lesssim \frac{(E_n + Z_n^2)^{1/2}}{|\mathbf{x} - \mathbf{z}|^2}$$

Using these bounds (in the same way we use Lemma 3.40 in the proof of the previous Proposition 3.41), we can estimate $\|\partial_s^a \partial^\beta M_2\|_3$.

3.5.3 Reduction to linear problem

Having estimated the non-linear parts of the equation in the last two subsections, in this section we will use them to reduce our problem to the linear problem for which we have the coercivity result that we can apply. We only need to do this for the case with no radial derivatives, the case with radial derivatives can be obtained by induction.

Lemma 3.43. For any θ that satisfies our a priori assumption (3.71) we have

$$\begin{split} \int_{0}^{s} \langle \mathbf{G}_{L} \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta}, \partial_{s}^{a+1} \partial^{\beta} \boldsymbol{\theta} \rangle_{3} \mathrm{d}\tau &= \frac{1}{2} \left\langle \mathbf{G}_{L} \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta} \rangle_{3} \right|_{0}^{s} \\ \langle \mathbf{L} \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta} \rangle_{3} &= \delta \| \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta} \|_{3}^{2} + \langle \mathbf{P}_{0,L} \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta} \rangle_{3} + \langle \mathbf{G}_{L} \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta} \rangle_{3}, \end{split}$$

where we recall (3.84), (3.85) and (2.55).

Proof. We have from (3.85)

$$\begin{split} \langle \mathbf{G}_L \partial_s^a \partial \!\!\!\!\!\partial^\beta \boldsymbol{\theta}, \partial_s^{a+1} \partial \!\!\!\!\partial^\beta \boldsymbol{\theta} \rangle_3 &= \int \left((\partial_s^a \partial \!\!\!\!\!\partial^\beta \theta^i) (\partial_s^{a+1} \partial \!\!\!\!\partial^\beta \theta^j) \bar{w}^3 \partial_i \partial_j \mathcal{K} \bar{w}^3 \\ &- (4\pi)^{-1} (\nabla \mathcal{K} \nabla \cdot (\bar{w}^3 \partial_s^a \partial \!\!\!\!\partial^\beta \theta)) \cdot (\nabla \mathcal{K} \nabla \cdot (\bar{w}^3 \partial_s^{a+1} \partial \!\!\!\!\partial^\beta \theta)) \right) \mathrm{d} \mathbf{x} \\ &= \frac{1}{2} \partial_s \int \left((\partial_s^a \partial \!\!\!\!\partial^\beta \theta^i) (\partial_s^a \theta^j) \bar{w}^3 \partial_i \partial_j \mathcal{K} \bar{w}^3 - (4\pi)^{-1} |\nabla \mathcal{K} \nabla \cdot (\bar{w}^3 \partial_s^a \partial \!\!\!\!\partial^\beta \theta)|^2 \right) \mathrm{d} \mathbf{x} \\ &= \frac{1}{2} \partial_s \langle \mathbf{G}_L \partial_s^a \partial \!\!\!\!\partial^\beta \theta, \partial_s^a \partial \!\!\!\!\partial^\beta \theta \rangle_3. \end{split}$$

The second formula follows from the definition of \mathbf{L} , $\mathbf{P}_{0,L}$ and \mathbf{G}_L .

The following theorem reduces the full non-linear problem to the linear one.

Theorem 3.44. Let $n \ge 20$ and suppose θ satisfies our a priori assumption (3.71). For $a + |\beta| \le n$ with a > 0 we have

$$\left| \int_{0}^{s} \langle \partial_{s}^{a} \partial^{\beta} (\delta \boldsymbol{\theta} + \mathbf{P} + \mathbf{G}), \partial_{s}^{a+1} \partial^{\beta} \boldsymbol{\theta} \rangle_{3} \mathrm{d}\tau - \frac{1}{2} \left\langle \mathbf{L} \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta} \rangle_{3} \right|_{0}^{s} \right|$$

$$\lesssim S_{n,|\beta|-1,0}^{1/2} S_{n,|\beta|,0}^{1/2} + (\mathcal{C}_{n} + \mathcal{Z}_{n}^{2})^{1/2} \mathcal{C}_{n} \quad (3.144)$$

$$\left| \int_{0}^{s} \langle \partial_{s}^{a} \partial^{\beta} (\delta \boldsymbol{\theta} + \mathbf{P} + \mathbf{G}), \partial_{s}^{a} \boldsymbol{\theta} \rangle_{3} \mathrm{d}\tau - \int_{0}^{s} \langle \mathbf{L} \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta}, \partial_{s}^{a} \partial^{\beta} \boldsymbol{\theta} \rangle_{3} \mathrm{d}\tau \right|$$

$$\lesssim \delta_{n,|\beta|-1,0}^{1/2} \delta_{n,|\beta|,0}^{1/2} + (\mathcal{C}_{n} + \mathcal{Z}_{n}^{2})^{1/2} \mathcal{C}_{n} \quad (3.145)$$

Proof. Using Lemma 3.43 and Propositions 3.38, 3.39, and 3.42, we conclude the proof.

This theorem above reduces the non-linear problem for time and tangential derivatives to the linear problem. Now applying our linear coercivity results from before, we get the following coercivity result for our non-linear problem, allowing us to control $\|\partial_s^{a+1}\partial^\beta\theta\|_{3+b}^2 + \|\partial_s^a\partial^\beta\theta\|_{3+b}^2 + \|\partial_s^a\nabla\partial^\beta\theta\|_{4+b}^2$.

Corollary 3.45. Let $n \ge 20$. Let θ be a solution of (3.62) in the sense of Theorem 3.11, given on its maximal interval of existence. Assume further that the energy, momentum, and irrotationality constraints (3.73), (3.74), and (3.75) hold respectively. Then for $a + |\beta| \le n$ with a > 0 we have

$$\begin{aligned} \left\|\partial_{s}^{a+1}\partial^{\beta}\boldsymbol{\theta}\right\|_{3}^{2} + \left\|\partial_{s}^{a}\partial^{\beta}\boldsymbol{\theta}\right\|_{3}^{2} + \left\|\partial_{s}^{a}\nabla\partial^{\beta}\boldsymbol{\theta}\right\|_{4}^{2} \\ \lesssim \left|\boldsymbol{\delta}\right|^{-2} \left(CS_{n,|\beta|,0}(0) + \frac{1}{2}\left\|\partial_{s}^{a+1}\partial^{\beta}\boldsymbol{\theta}\right\|_{3}^{2}\right|_{0}^{s} + \int_{0}^{s} \langle\partial_{s}^{a}\partial^{\beta}(\delta\boldsymbol{\theta} + \mathbf{P} + \mathbf{G}), \partial_{s}^{a+1}\partial^{\beta}\boldsymbol{\theta}\rangle_{3} \mathrm{d}\tau\right) \\ + C\left(\delta_{n,|\beta|-1,0} + \left|\boldsymbol{\delta}\right|^{-2}\delta_{n,|\beta|-1,0}^{1/2}\delta_{n,|\beta|,0}^{1/2}\right) + C_{\delta}(\mathcal{C}_{n} + \mathcal{Z}_{n}^{2})^{1/2}\mathcal{C}_{n} \end{aligned}$$
(3.146)

$$\int_{0}^{s} \left(\left\| \partial_{s}^{a+1} \partial^{\beta} \theta \right\|_{3}^{2} + \left\| \partial_{s}^{a} \partial^{\beta} \theta \right\|_{3}^{2} + \left\| \partial_{s}^{a} \nabla \partial^{\beta} \theta \right\|_{4}^{2} \right) d\tau$$

$$\lesssim |t|^{-2} \int_{0}^{s} \left(\left\| \partial_{s}^{a+1} \partial^{\beta} \theta \right\|_{3}^{2} + \langle \partial_{s}^{a} \partial^{\beta} (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_{s}^{a} \partial^{\beta} \theta \rangle_{3} \right) d\tau$$

$$+ C \left(\delta_{n,|\beta|-1,0} + |t|^{-2} \delta_{n,|\beta|-1,0}^{1/2} \delta_{n,|\beta|,0}^{1/2} \right) + C_{\delta} (\mathcal{E}_{n} + \mathcal{E}_{n}^{2})^{1/2} \mathcal{E}_{n}$$
(3.147)

Proof. Combining Theorem 3.44 and Propositions 3.34 and 3.36 we conclude the proof.

To control the version with radial derivative $\|\partial_s^{a+1}X_r^b\partial^\beta\theta\|_{3+b}^2 + \|\partial_s^aX_r^b\partial^\beta\theta\|_{3+b}^2 + \|\partial_s^a\nabla X_r^b\partial^\beta\theta\|_{4+b}^2$, we do not need to apply the linear coercivity result like Theorem 3.44 above. This is because we get control of $\|\partial_s^a\nabla X_r^b\partial^\beta\theta\|_{4+b}^2$ directly from the pressure term, while the control of $\|\partial_s^{a+1}X_r^b\partial^\beta\theta\|_{3+b}^2 + \|\partial_s^aX_r^b\partial^\beta\theta\|_{3+b}^2$ and the gravity term we get automatically from induction from the step with one less space derivative, as follows.

Corollary 3.46. Let $n \ge 21$ and suppose θ satisfies our a priori assumption (3.71). For $a + |\beta| + b \le n$ with a, b > 0 we have

$$\begin{split} \left\|\partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{3+b}^{2} + \left\|\partial_{s}^{a}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{3+b}^{2} + \left\|\partial_{s}^{a}\nabla X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{4+b}^{2} \\ \lesssim CS_{n,|\beta|+b,b}(0) + \frac{1}{2}\left\|\partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{3+b}^{2} + \int_{0}^{s}\langle\partial_{s}^{a}X_{r}^{b}\partial^{\beta}(\delta\boldsymbol{\theta} + \mathbf{P} + \mathbf{G}), \partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\rangle_{3+b}d\tau \\ + C\left(\delta_{n,|\beta|+b-1} + (\delta_{n,|\beta|+b-1}^{1/2} + \delta_{n,|\beta|+b,b-1}^{1/2})\mathfrak{E}_{n}^{1/2} + (\mathfrak{E}_{n} + \mathfrak{E}_{n}^{2})^{1/2}\mathfrak{E}_{n}\right)$$
(3.148)

$$\int_{0}^{s} \left(\left\| \partial_{s}^{a+1} X_{r}^{b} \partial^{\beta} \theta \right\|_{3+b}^{2} + \left\| \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \theta \right\|_{3+b}^{2} + \left\| \partial_{s}^{a} \nabla X_{r}^{b} \partial^{\beta} \theta \right\|_{4+b}^{2} \right) \mathrm{d}\tau$$

$$\lesssim \int_{0}^{s} \left(\left\| \partial_{s}^{a+1} X_{r}^{b} \partial^{\beta} \theta \right\|_{3+b}^{2} + \langle \partial_{s}^{a} X_{r}^{b} \partial^{\beta} (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_{s}^{a} X_{r}^{b} \partial^{\beta} \theta \rangle_{3+b} \right) \mathrm{d}\tau$$

$$+ C \left(\delta_{n,|\beta|+b-1} + (\delta_{n,|\beta|+b-1}^{1/2} + \delta_{n,|\beta|+b,b-1}^{1/2}) \mathcal{E}_{n}^{1/2} + (\mathcal{E}_{n} + \mathcal{E}_{n}^{2})^{1/2} \mathcal{E}_{n} \right)$$
(3.149)

Proof. By Propositions 3.39 and 3.38, we can replace $\partial_s^a X_r^b \partial^\beta \mathbf{P}$ by $\mathbf{P}_{b,L} \partial_s^a X_r^b \partial^\beta \boldsymbol{\theta}$. Now by Lemma 2.12 we have

$$\begin{split} &\frac{1}{2} \left[\|\partial_s^a \nabla X_r^b \partial^\beta \theta\|_{4+b}^2 + \frac{1}{3} \|\partial_s^a \nabla \cdot (X_r^b \partial^\beta \theta)\|_{4+b}^2 - \frac{1}{2} \|[\partial_s^a \operatorname{curl} X_r^b \partial^\beta \theta]\|_{4+b}^2 \right]_0^s \\ &= \int_0^s \langle \mathbf{P}_{b,L} \partial_s^a X_r^b \partial^\beta \theta, \partial_s^{a+1} X_r^b \partial^\beta \theta \rangle_{3+b} \mathrm{d}\tau \end{split}$$

Now using Corollary 3.31 we get

$$\begin{aligned} \|\partial_s^a \nabla X_r^b \partial^\beta \boldsymbol{\theta}\|_{4+b}^2 &\lesssim CS_{n,|\beta|+b,b}(0) + \int_0^s \langle \mathbf{P}_{b,L} \partial_s^a X_r^b \partial^\beta \boldsymbol{\theta}, \partial_s^{a+1} X_r^b \partial^\beta \boldsymbol{\theta} \rangle_{3+b} \mathrm{d}\tau \\ &+ C \left(S_{n,|\beta|+b-1} + (E_n + Z_n^2) E_n \right). \end{aligned}$$

Furthermore, note that

$$\begin{split} \left\|\partial_s^{a+1} X_r^b \partial^\beta \boldsymbol{\theta}\right\|_{3+b}^2 + \left\|\partial_s^a X_r^b \partial^\beta \boldsymbol{\theta}\right\|_{3+b}^2 \lesssim \left\|\partial_s^{a+1} \nabla X_r^{b-1} \partial^\beta \boldsymbol{\theta}\right\|_{4+(b-1)}^2 + \left\|\partial_s^a \nabla X_r^{b-1} \partial^\beta \boldsymbol{\theta}\right\|_{4+(b-1)}^2 \\ \lesssim S_{n,|\beta|+b-1}. \end{split}$$

Now note that, using Proposition 3.41,

$$\left| \int_{0}^{s} \langle \delta \partial_{s}^{a} X_{r}^{b} \boldsymbol{\partial}^{\beta} \boldsymbol{\theta}, \partial_{s}^{a+1} X_{r}^{b} \boldsymbol{\partial}^{\beta} \boldsymbol{\theta} \rangle_{3+b} \mathrm{d}\tau \right| \lesssim \mathcal{S}_{n,|\beta|+b-1} \\ \left| \int_{0}^{s} \langle \partial_{s}^{a} X_{r}^{b} \boldsymbol{\partial}^{\beta} \mathbf{G}, \partial_{s}^{a+1} X_{r}^{b} \boldsymbol{\partial}^{\beta} \boldsymbol{\theta} \rangle_{3+b} \mathrm{d}\tau \right| \lesssim \mathcal{S}_{n,|\beta|+b-1}^{1/2} \mathcal{S}_{n}^{1/2}$$

then we are done for the first formula. Proof for the second formula is similar.

3.6 Energy estimates and proof of the main theorem

In this section we finally commute the momentum equation (3.62) and then derive the high-order energy estimates. Since the bounds near the vacuum boundary are more delicate as they are sensitive to the weights, we present them in Section 3.6.1 and the estimates away from the vacuum boundary in Section 3.6.2. Then finally we will prove our main theorem in section 3.6.3 using the energy estimates.

3.6.1 Near boundary energy estimate

In this subsection we will prove the energy estimate for S_n (recall (3.68)).

Theorem 3.47 (Near boundary energy estimate). Let $n \ge 21$, and assume that $\epsilon > 0$ and $|\delta|$ are sufficiently small. Let θ be a solution of (3.62) in the sense of Theorem 3.11, given on its maximal interval of existence. Assume further that the energy, momentum, and irrotationality constraints (3.73), (3.74), and (3.75) hold respectively. Then there exist m > 0 such that

$$\mathcal{S}_n - C\epsilon \mathcal{E}_n \lesssim_{\epsilon} |\mathfrak{G}|^{-m} \mathcal{S}_n(0) + C_{\delta} (\mathcal{E}_n + \mathfrak{X}_n^2)^{1/2} \mathcal{E}_n \tag{3.150}$$

whenever our a priori assumption (3.71) is satisfied. Here we recall Definition (3.70) of the total norm \mathcal{E}_n .

Proof. Let $a + |\beta| + b \le n$. Apply $\partial_s^a X_r^b \partial^\beta$ to the momentum equation (3.62) to get

$$\partial_s^{a+2} X_r^b \partial^\beta \theta - \frac{\theta}{2} \partial_s^{a+1} X_r^b \partial^\beta \theta + \partial_s^a X_r^b \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}) = 0$$

Taking the $\langle \cdot, \cdot \rangle_{3+b}$ -inner with $\partial_s^{a+1} X_r^b \partial^\beta \theta$ we get

$$0 = \frac{1}{2}\partial_s \|\partial_s^{a+1} X_r^b \partial^\beta \theta\|_{3+b}^2 + \langle \partial_s^a X_r^b \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_s^{a+1} X_r^b \partial^\beta \theta \rangle_{3+b} - \frac{\theta}{2} \|\partial_s^{a+1} X_r^b \partial^\beta \theta\|_{3+b}^2$$

On the other hand, taking inner product of the equation with $\partial_s^a X_r^b \partial^\beta \theta$ we get

$$0 = \partial_s \langle \partial_s^{a+1} X_r^b \partial^\beta \theta, \partial_s^a X_r^b \partial^\beta \theta \rangle_{3+b} - \|\partial_s^{a+1} X_r^b \partial^\beta \theta\|_{3+b}^2 - \frac{b}{4} \partial_{3+b} \|\partial_s^a X_r^b \partial^\beta \theta\|_{3+b}^2 + \langle \partial_s^a X_r^b \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_s^a X_r^b \partial^\beta \theta \rangle_{3+b}$$

where we used the identiity $\langle \partial_s^{a+2} X_r^b \partial^\beta \theta, \partial_s^a X_r^b \partial^\beta \theta \rangle = \partial_s \langle \partial_s^{a+1} X_r^b \partial^\beta \theta, \partial_s^a X_r^b \partial^\beta \theta \rangle - \|\partial_s^{a+1} X_r^b \partial^\beta \theta\|^2$. Multiply the latter equation by c, add to it two times the equation before, and then integrate w.r.t. s to obtain

$$\begin{split} 0 &= \left(\frac{1}{2} \|\partial_s^{a+1} X_r^b \partial^\beta \boldsymbol{\theta}\|_{3+b}^2 + c \langle \partial_s^{a+1} X_r^b \partial^\beta \boldsymbol{\theta}, \partial_s^a X_r^b \partial^\beta \boldsymbol{\theta} \rangle_{3+b} - \frac{c \delta}{4} \|\partial_s^a X_r^b \partial^\beta \boldsymbol{\theta}\|_{3+b}^2 \right) \Big|_0^s \\ &+ \int_0^s \left(\langle \partial_s^a X_r^b \partial^\beta (\delta \boldsymbol{\theta} + \mathbf{P} + \mathbf{G}), \partial_s^{a+1} X_r^b \partial^\beta \boldsymbol{\theta} \rangle_{3+b} + c \langle \partial_s^a X_r^b \partial^\beta (\delta \boldsymbol{\theta} + \mathbf{P} + \mathbf{G}), \partial_s^a X_r^b \partial^\beta \boldsymbol{\theta} \rangle_{3+b} \\ &- \left(c + \frac{\delta}{2} \right) \|\partial_s^{a+1} X_r^b \partial^\beta \boldsymbol{\theta}\|_{3+b}^2 \right) \mathrm{d}\tau. \end{split}$$

i. When b = 0, using Corollary 3.45 we get

$$\begin{split} \left\|\partial_{s}^{a+1}\partial^{\beta}\boldsymbol{\theta}\right\|_{3}^{2} + \left\|\partial_{s}^{a}\partial^{\beta}\boldsymbol{\theta}\right\|_{3}^{2} + \left\|\partial_{s}^{a}\nabla\partial^{\beta}\boldsymbol{\theta}\right\|_{4}^{2} + c\int_{0}^{s}\left(\left\|\partial_{s}^{a+1}\partial^{\beta}\boldsymbol{\theta}\right\|_{3}^{2} + \left\|\partial_{s}^{a}\partial^{\beta}\boldsymbol{\theta}\right\|_{3}^{2} + \left\|\partial_{s}^{a}\nabla\partial^{\beta}\boldsymbol{\theta}\right\|_{4}^{2}\right)\mathrm{d}\tau \\ + |\boldsymbol{\ell}|^{-2}\left(c\langle\partial_{s}^{a+1}\partial^{\beta}\boldsymbol{\theta},\partial_{s}^{a}\partial^{\beta}\boldsymbol{\theta}\rangle_{3} - \frac{c\boldsymbol{\ell}}{4}\|\partial_{s}^{a}\partial^{\beta}\boldsymbol{\theta}\|_{3}^{2}\right)\Big|_{0}^{s} - |\boldsymbol{\ell}|^{-2}\int_{0}^{s}\left(2c + \frac{\boldsymbol{\ell}}{2}\right)\|\partial_{s}^{a+1}\partial^{\beta}\boldsymbol{\theta}\|_{3}^{2}\mathrm{d}\tau \\ \lesssim |\boldsymbol{\ell}|^{-2}S_{n,|\beta|,0}(0) + S_{n,|\beta|-1,0} + |\boldsymbol{\ell}|^{-2}S_{n,|\beta|-1,0}^{1/2}S_{n,|\beta|,0}^{1/2} + C_{\delta}(\boldsymbol{\mathcal{E}}_{n} + \boldsymbol{\mathcal{X}}_{n}^{2})^{1/2}\boldsymbol{\mathcal{E}}_{n}. \end{split}$$

Choosing c small enough (e.g. $c = |b|^2/100$ when $b \ll 1$), we get

$$\left\|\partial_{s}^{a+1}\partial^{\beta}\boldsymbol{\theta}\right\|_{3}^{2}+\left\|\partial_{s}^{a}\partial^{\beta}\boldsymbol{\theta}\right\|_{3}^{2}+\left\|\partial_{s}^{a}\nabla\partial^{\beta}\boldsymbol{\theta}\right\|_{4}^{2}+\int_{0}^{s}\left(\left\|\partial_{s}^{a+1}\partial^{\beta}\boldsymbol{\theta}\right\|_{3}^{2}+\left\|\partial_{s}^{a}\partial^{\beta}\boldsymbol{\theta}\right\|_{3}^{2}+\left\|\partial_{s}^{a}\nabla\partial^{\beta}\boldsymbol{\theta}\right\|_{4}^{2}\right)\mathrm{d}\tau$$

$$\lesssim |\mathfrak{C}|^{-2} (|\mathfrak{C}|^{-2} S_{n,|\beta|,0}(0) + S_{n,|\beta|-1,0} + |\mathfrak{C}|^{-2} S_{n,|\beta|-1,0}^{1/2} S_{n,|\beta|,0}^{1/2} + C_{\delta} (\mathfrak{C}_{n} + \mathfrak{X}_{n}^{2})^{1/2} \mathfrak{C}_{n}),$$

and so (noting that the constant implicit in the notation \leq do not depend on s)

$$\mathcal{S}_{n,|\beta|,0} \lesssim |\mathfrak{C}|^{-4} \mathcal{S}_{n,|\beta|,0}(0) + |\mathfrak{C}|^{-2} \mathcal{S}_{n,|\beta|-1,0} + |\mathfrak{C}|^{-4} \mathcal{S}_{n,|\beta|-1,0}^{1/2} \mathcal{S}_{n,|\beta|,0}^{1/2} + C_{\delta} (\mathfrak{C}_{n} + \mathfrak{X}_{n}^{2})^{1/2} \mathfrak{C}_{n}.$$

In particular when $|\beta| = 0$ we have $S_{n,0,0} \leq |\delta|^{-4} S_{n,0,0}(0) + C_{\delta} (\mathcal{E}_n + \mathcal{Z}_n^2)^{1/2} \mathcal{E}_n$. And so using Young's inequality and by induction on $|\beta|$ we have

$$\mathcal{S}_{n,|\beta|,0} \lesssim |\mathfrak{C}|^{-4-8|\beta|} \mathcal{S}_{n,|\beta|,0}(0) + C_{\delta}(\mathscr{C}_n + \mathscr{X}_n^2)^{1/2} \mathscr{C}_n \tag{3.151}$$

for all $|\beta| \leq n$.

ii. When b > 0, using Corollary 3.46 we get

$$\begin{split} \left\|\partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{3+b}^{2} + \left\|\partial_{s}^{a}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{3+b}^{2} + \left\|\partial_{s}^{a}\nabla X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{4+b}^{2} \\ + c\int_{0}^{s}\left(\left\|\partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{3+b}^{2} + \left\|\partial_{s}^{a}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{3+b}^{2} + \left\|\partial_{s}^{a}\nabla X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{4+b}^{2}\right)\mathrm{d}\tau \\ + \left(c\langle\partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta},\partial_{s}^{a}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\rangle_{3} - \frac{c\delta}{4}\|\partial_{s}^{a}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\|_{3}^{2}\right)\Big|_{0}^{s} - \int_{0}^{s}\left(2c + \frac{\delta}{2}\right)\|\partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\|_{3}^{2}\mathrm{d}\tau \\ \lesssim S_{n,|\beta|+b,b}(0) + \delta_{n,|\beta|+b-1} + (\delta_{n,|\beta|+b-1}^{1/2} + \delta_{n,|\beta|+b,b-1}^{1/2})\mathcal{E}_{n}^{1/2} + C_{\delta}(\mathcal{E}_{n} + \mathcal{E}_{n}^{2})^{1/2}\mathcal{E}_{n} \end{split}$$

Choosing c small enough we get

$$\begin{split} \left\|\partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{3+b}^{2} + \left\|\partial_{s}^{a}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{3+b}^{2} + \left\|\partial_{s}^{a}\nabla X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{4+b}^{2} \\ + \int_{0}^{s} \left(\left\|\partial_{s}^{a+1}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{3+b}^{2} + \left\|\partial_{s}^{a}X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{3+b}^{2} + \left\|\partial_{s}^{a}\nabla X_{r}^{b}\partial^{\beta}\boldsymbol{\theta}\right\|_{4+b}^{2}\right) \mathrm{d}\tau \\ \lesssim |\mathfrak{E}|^{-1}(S_{n,|\beta|+b,b}(0) + \mathcal{S}_{n,|\beta|+b-1} + (\mathcal{S}_{n,|\beta|+b-1}^{1/2} + \mathcal{S}_{n,|\beta|+b,b-1}^{1/2})\mathcal{E}_{n}^{1/2} + C_{\delta}(\mathcal{E}_{n} + \mathcal{E}_{n}^{2})^{1/2}\mathcal{E}_{n}) \end{split}$$

and so

$$\begin{split} & \mathcal{S}_{n,|\beta|+b,b} \lesssim |\mathfrak{C}|^{-1} (\mathcal{S}_{n,|\beta|+b,b}(0) + \mathcal{S}_{n,|\beta|+b-1} + \mathcal{S}_{n,|\beta|+b,b-1} \\ & + (\mathcal{S}_{n,|\beta|+b-1}^{1/2} + \mathcal{S}_{n,|\beta|+b,b-1}^{1/2}) \mathcal{C}_n^{1/2} + C_{\delta} (\mathcal{C}_n + \mathcal{Z}_n^2)^{1/2} \mathcal{C}_n). \end{split}$$

or equivalently

$$S_{n,c,d} \lesssim |\mathfrak{b}|^{-1} (S_{n,c,d}(0) + S_{n,c-1} + S_{n,c,d-1} + (S_{n,c-1}^{1/2} + S_{n,c,d-1}^{1/2}) \mathcal{E}_n^{1/2} + C_{\delta} (\mathcal{E}_n + \mathcal{X}_n^2)^{1/2} \mathcal{E}_n).$$

We already know $\mathscr{S}_{n,c,0} \lesssim |\mathscr{C}|^{-4-8c} \mathscr{S}_{n,c,0}(0) + C_{\delta}(\mathscr{C}_n + \mathscr{Z}_n^2)^{1/2} \mathscr{C}_n$. And so using Young's inequality and by induction on c and d we have

$$\mathcal{S}_{n,c,d} - C\epsilon \mathfrak{E}_n \lesssim_{\epsilon} |\mathfrak{E}|^{-m} \mathcal{S}_{n,c,d}(0) + C_{\delta} (\mathfrak{E}_n + \mathfrak{X}_n^2)^{1/2} \mathfrak{E}_n.$$

for all $d \leq c \leq n$. This means we have $\mathscr{S}_n - C \epsilon \mathscr{C}_n \lesssim_{\epsilon} |\mathfrak{G}|^{-m} \mathscr{S}_n(0) + C_{\delta}(\mathscr{C}_n + \mathfrak{L}_n^2)^{1/2} \mathscr{C}_n$.

3.6.2 Near origin energy estimate

In this subsection we will prove the energy estimate for Q_n , see (3.69).

Using Lemma 2.3, the following lemma shows that in fact we only need to control the divergence $\partial_s^a \nabla^c \nabla \cdot \boldsymbol{\theta}$ in order to control the near origin energy Q_n .

Lemma 3.48. Let $n \ge 20$ and $c \le n$. Let θ be a solution of (3.62) in the sense of Theorem 3.11, given on its maximal interval of existence. Assume further that the irrotationality constraint (3.75). Then

$$Q_{n,c} \lesssim \|\partial_s^a \nabla^c \nabla \cdot \boldsymbol{\theta}\|_{3+2(c+1)}^2 + Q_{n,c-1} + (E_n + Z_n^2) E_n.$$

Proof. Let $a + c \le n$ with a > 0. Using the previous Lemma 2.3, we have

$$\begin{aligned} \|\partial_{s}^{a}\nabla^{c+1}\boldsymbol{\theta}\|_{3+2(c+1)}^{2} &\lesssim \|\partial_{s}^{a}\nabla^{c}\nabla\cdot\boldsymbol{\theta}\|_{3+2(c+1)}^{2} + \|\partial_{s}^{a}\nabla^{c}\nabla\times\boldsymbol{\theta}\|_{3+2(c+1)}^{2} + \|\partial_{s}^{a}\nabla^{c}\boldsymbol{\theta}\|_{3+2(c+1)}^{2} \\ &\leq \|\partial_{s}^{a}\nabla^{c}\nabla\cdot\boldsymbol{\theta}\|_{3+2(c+1)}^{2} + \|\partial_{s}^{a}\nabla^{c}\nabla\times\boldsymbol{\theta}\|_{3+2(c+1)}^{2} + Q_{n,c-1}. \end{aligned}$$

Recalling (3.135) we have $\partial_s^a \nabla^c \nabla \times \boldsymbol{\theta} = -\partial_s^{a-1} \nabla^c ((\partial_s \nabla \theta^k) \times \nabla \theta^k)$ and therefore

$$\|\partial_s^a \nabla^c \nabla \times \boldsymbol{\theta}\|_{3+2(c+1)}^2 \lesssim (E_n + Z_n^2) E_n.$$

		I
		I
		I

Lemma 3.49. For any tensor field T smooth enough we have

$$\partial^{\gamma} \left(\bar{w}^{-3} \partial_k (\bar{w}^4 T^k) \right) = \bar{w} \partial^{\gamma} \partial_k T^k + \sum_{|\gamma'| \le |\gamma|} \bar{w}^{|\gamma| - |\gamma'|} \langle C \partial^{\gamma'} T^k \rangle$$

where we recall notations introduced in Definition 2.2.

Proof. The statement follows easily by induction.

Theorem 3.50 (Near origin energy estimate). Let $n \ge 21$ and δ small. Let θ be a solution of (3.62) in the sense of Theorem 3.11, given on its maximal interval of existence. Assume further that the energy, momentum, and irrotationality constraints (3.73), (3.74), and (3.75) hold respectively. Then we have

$$\mathbb{Q}_n \lesssim |\mathfrak{C}|^{-4} \mathscr{E}_n(0) + C_\delta (\mathscr{E}_n + \mathscr{Z}_n^2)^{1/2} \mathscr{E}_n \tag{3.152}$$

whenever our a priori assumption (3.71) is satisfied.

Proof. Recall the momentum equation (3.62) is

$$\mathbf{0} = \partial_s^2 \boldsymbol{\theta} - \frac{1}{2} \boldsymbol{\vartheta} \partial_s \boldsymbol{\theta} + \delta \boldsymbol{\theta} + \bar{w}^{-3} \partial_k (\bar{w}^4 (\underbrace{\mathscr{A}^k \mathcal{J}^{-1/3} - I^k}_{=T})) + \mathscr{A} \nabla \Phi - \nabla \mathcal{K} \bar{w}^3.$$

where we recall T in (2.46). Also recall from (2.41) $(\mathscr{A}\nabla) \cdot (\mathscr{A}\nabla)\Phi(\mathbf{x}) = 4\pi \bar{w}^3 \mathscr{J}^{-1}$. So taking the divergence of the gravity term makes it easy to estimate. From Lemma 3.48 we also know that to control Q_n it suffices to estimate the divergence. Let $a + |\gamma| + 1 \le n$. Evaluating the dot product of (3.62) with $\partial_s^a \partial^{\gamma} \mathscr{A} \nabla$ we get

$$\begin{split} \mathbf{0} &= \partial_s^a \partial^\gamma \mathcal{A} \nabla \cdot \partial_s^2 \boldsymbol{\theta} - \frac{1}{2} \mathcal{C} \partial_s^a \partial^\gamma \mathcal{A} \nabla \cdot \partial_s \boldsymbol{\theta} + \delta \partial_s^a \partial^\gamma \mathcal{A} \nabla \cdot \boldsymbol{\theta} \\ &+ \partial_s^a \partial^\gamma \mathcal{A} \nabla \cdot \left(\bar{w}^{-3} \partial_k (\bar{w}^4 T^k) \right) + 4\pi \partial_s^a \partial^\gamma (\bar{w}^3 \mathcal{F}^{-1}) - (\partial_s^a \partial^\gamma \mathcal{A} \nabla) \cdot \nabla \mathcal{K} \bar{w}^3 \\ &= \partial_s^a \partial^\gamma \mathcal{A} \nabla \cdot \partial_s^2 \boldsymbol{\theta} - \frac{1}{2} \mathcal{C} \partial_s^a \partial^\gamma \mathcal{A} \nabla \cdot \partial_s \boldsymbol{\theta} + \delta \partial_s^a \partial^\gamma \mathcal{A} \nabla \cdot \boldsymbol{\theta} \\ &+ \partial_s^a \partial^\gamma \mathcal{A} \nabla \cdot \left(\bar{w}^{-3} \partial_k (\bar{w}^4 T^k) \right) + 4\pi \partial_s^a \partial^\gamma (\bar{w}^3 (\mathcal{F}^{-1} - 1)) - (\partial_s^a \partial^\gamma (\mathcal{A} - I) \nabla) \cdot \nabla \mathcal{K} \bar{w}^3 \end{split}$$

From here we will the do two things (i) and (ii) as follows

(i) Times the equation with $\bar{w}^{6+2|\gamma|}\partial_s^{a+1}\partial^{\gamma}\mathscr{A}\nabla\cdot\boldsymbol{\theta}$ and integrate in time and space we get

$$\begin{split} 0 &= \int_0^s \left(\left\langle \partial_s^a \partial^\gamma \mathscr{A} \nabla \cdot \partial_s^2 \theta - \frac{1}{2} \mathscr{b} \partial_s^a \partial^\gamma \mathscr{A} \nabla \cdot \partial_s \theta + \delta \partial_s^a \partial^\gamma \mathscr{A} \nabla \cdot \theta, \partial_s^{a+1} \partial^\gamma \mathscr{A} \nabla \cdot \theta \right\rangle_{6+2|\gamma|} \\ &+ \left\langle \partial_s^a \partial^\gamma \mathscr{A} \nabla \cdot \left(\bar{w}^{-3} \partial_k (\bar{w}^4 T^k) \right), \partial_s^{a+1} \partial^\gamma \mathscr{A} \nabla \cdot \theta \right\rangle_{6+2|\gamma|} \\ &+ \left\langle 4\pi \partial_s^a \partial^\gamma (\bar{w}^3 (\mathcal{J}^{-1} - 1)) - (\partial_s^a \partial^\gamma (\mathscr{A} - I) \nabla) \cdot \nabla \mathcal{K} \bar{w}^3, \partial_s^{a+1} \partial^\gamma \mathscr{A} \nabla \cdot \theta \right\rangle_{6+2|\gamma|} \right) \mathrm{d} \pi \mathcal{K}$$

Now commuting \mathscr{A} with space and time derivatives, we get a non-linear remainder $\mathscr{R}[(\mathscr{C}_n + \mathscr{Z}_n^2)^{1/2} \mathscr{C}_n]$ (recall notation $\mathscr{R}[\star]$ defined in Definition 2.2),

$$\begin{split} 0 &= \int_{0}^{s} \left(\left\langle \partial_{s} \mathcal{A} \nabla \cdot \partial_{s}^{a+1} \partial^{\gamma} \boldsymbol{\theta} - \frac{1}{2} \mathcal{E} \mathcal{A} \nabla \cdot \partial_{s}^{a+1} \partial^{\gamma} \boldsymbol{\theta}, \mathcal{A} \nabla \partial_{s}^{a+1} \partial^{\gamma} \cdot \boldsymbol{\theta} \right\rangle_{6+2|\gamma|} \\ &+ \left\langle \delta \mathcal{A} \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \boldsymbol{\theta}, \partial_{s} \mathcal{A} \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \boldsymbol{\theta} \right\rangle_{6+2|\gamma|} \\ &+ \left\langle \mathcal{A} \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \left(\bar{w}^{-3} \partial_{k} (\bar{w}^{4} T^{k}) \right), \partial_{s} \mathcal{A} \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \boldsymbol{\theta} \right\rangle_{6+2|\gamma|} \\ &+ \left\langle 4\pi (\mathcal{F}^{-1} - 1) \partial_{s}^{a} \partial^{\gamma} (\bar{w}^{3}) - (\mathcal{A} - I) \nabla \cdot \nabla \mathcal{K} \partial_{s}^{a} \partial^{\gamma} \bar{w}^{3}, \partial_{s}^{a+1} \partial^{\gamma} \mathcal{A} \nabla \cdot \boldsymbol{\theta} \right\rangle_{6+2|\gamma|} \right) \mathrm{d} \tau \\ &+ \mathcal{R} [(\mathcal{E}_{n} + \mathcal{X}_{n}^{2})^{1/2} \mathcal{E}_{n}] \end{split}$$

Now terms in the first two line we factorised, and terms in the last line in the integral we can estimate by $\mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}$ and $\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}$,

$$\begin{split} 0 &= \int_0^s \left(\frac{1}{2} \partial_s \left\| \mathscr{A} \nabla \cdot \partial_s^{a+1} \partial^{\gamma} \boldsymbol{\theta} \right\|_{6+2|\gamma|}^2 - \frac{1}{2} \mathscr{E} \left\| \mathscr{A} \nabla \cdot \partial_s^{a+1} \partial^{\gamma} \boldsymbol{\theta} \right\|_{6+2|\gamma|}^2 + \frac{1}{2} \delta \partial_s \left\| \mathscr{A} \nabla \cdot \partial_s^a \partial^{\gamma} \boldsymbol{\theta} \right\|_{6+2|\gamma|}^2 \\ &+ \left\langle \mathscr{A} \nabla \cdot \left(\bar{w} \partial_s^a \partial^{\gamma} \partial_k T^k \right), \partial_s \mathscr{A} \nabla \cdot \partial_s^a \partial^{\gamma} \boldsymbol{\theta} \right\rangle_{6+2|\gamma|} \right) \mathrm{d} \tau \\ &+ \mathscr{R} [(\mathscr{E}_n + \mathscr{E}_n^2)^{1/2} \mathscr{E}_n] + \mathscr{R} [\mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathscr{R} [\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] \end{split}$$

Now terms that are full time derivatives can be evaluated, and $\partial_s^a \partial^\gamma \partial_k T^k$ can be converted to $T_T[\partial_s^a \partial^\gamma \partial_k \nabla \theta]^k$ (recall Lemma 2.11) leaving a reminder that we can estimate with $(\mathcal{C}_n + \mathcal{Z}_n^2)^{1/2} \mathcal{C}_n$.

$$\begin{split} 0 &= \frac{1}{2} \left(\left\| \mathscr{A} \nabla \cdot \partial_s^{a+1} \partial^{\gamma} \boldsymbol{\theta} \right\|_{6+2|\gamma|}^2 + \delta \left\| \mathscr{A} \nabla \cdot \partial_s^a \partial^{\gamma} \boldsymbol{\theta} \right\|_{6+2|\gamma|}^2 \right) \Big|_0^s \\ &+ \int_0^s \left(-\frac{1}{2} \delta \left\| \mathscr{A} \nabla \cdot \partial_s^{a+1} \partial^{\gamma} \boldsymbol{\theta} \right\|_{6+2|\gamma|}^2 + \left\langle \mathscr{A} \nabla \cdot \left(\bar{w} T_T [\partial_s^a \partial^{\gamma} \partial_k \nabla \boldsymbol{\theta}]^k \right), \partial_s \mathscr{A} \nabla \cdot \partial_s^a \partial^{\gamma} \boldsymbol{\theta} \right\rangle_{6+2|\gamma|} \right) \mathsf{d}\tau \\ &+ \mathscr{R} [(\mathscr{E}_n + \mathscr{E}_n^2)^{1/2} \mathscr{E}_n] + \mathscr{R} [\mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathscr{R} [\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] \end{split}$$

Now all the term before the term with T_T can be bounded by $\mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}$ and $\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}$, and we integrate by parts on the term with T_T ,

$$0 = -\int_{0}^{s} \left\langle \bar{w}T_{T}[\partial_{s}^{a}\partial^{\gamma}\partial_{k}\nabla\boldsymbol{\theta}]^{k}, \partial_{s}\mathcal{A}\nabla(\mathcal{A}\nabla\cdot\partial_{s}^{a}\partial^{\gamma}\boldsymbol{\theta})\right\rangle_{6+2|\gamma|} \mathrm{d}\tau \\ + \mathcal{R}[(\mathcal{C}_{n} + \mathcal{Z}_{n}^{2})^{1/2}\mathcal{C}_{n}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}^{1/2}\mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathcal{R}[\mathbb{Q}_{n-1}^{1/2}\mathbb{Q}_{n,|\gamma|+1}^{1/2}]$$

Now we expend the terms by definition and simplify,

$$\begin{split} 0 &= \int_0^s \left\langle \mathcal{F}^{-1/3} \left(\mathcal{A}_m^k \mathcal{A}^l + \frac{1}{3} \mathcal{A}^k \mathcal{A}_m^l \right) \partial_s^a \partial^\gamma \partial_k \partial_l \theta^m, \partial_s (\mathcal{A}^j \mathcal{A}_i^\ell \partial_j \partial_\ell \partial_s^a \partial^\gamma \theta^i) \right\rangle_{7+2|\gamma|} \mathrm{d}\tau \\ &+ \mathcal{R}[(\mathcal{E}_n + \mathcal{X}_n^2)^{1/2} \mathcal{E}_n] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathcal{R}[\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] \\ &= \int_0^s \left\langle \mathcal{F}^{-1/3} \left(\mathcal{A}_m^k \mathcal{A}_o^l + \frac{1}{3} \mathcal{A}_o^k \mathcal{A}_m^l \right) \partial_k \partial_l \partial_s^a \partial^\gamma \theta^m, \partial_s (\mathcal{A}_o^j \mathcal{A}_i^\ell \partial_j \partial_\ell \partial_s^a \partial^\gamma \theta^i) \right\rangle_{7+2|\gamma|} \mathrm{d}\tau \\ &+ \mathcal{R}[(\mathcal{E}_n + \mathcal{X}_n^2)^{1/2} \mathcal{E}_n] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathcal{R}[\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] \\ &= \int_0^s \frac{4}{3} \left\langle \mathcal{F}^{-1/3} \mathcal{A}_o^k \mathcal{A}_m^l \partial_k \partial_l \partial_s^a \partial^\gamma \theta^m, \partial_s (\mathcal{A}_o^j \mathcal{A}_i^\ell \partial_j \partial_\ell \partial_s^a \partial^\gamma \theta^i) \right\rangle_{7+2|\gamma|} \mathrm{d}\tau \\ &+ \mathcal{R}[(\mathcal{E}_n + \mathcal{X}_n^2)^{1/2} \mathcal{E}_n] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathcal{R}[\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] \end{split}$$

Now the term in the integral can be factorised into a time derivative,

$$0 = \frac{2}{3} \int_0^s \int_{B_R} \mathcal{F}^{-1/3} \partial_s \left\| \mathcal{A}^k \mathcal{A}_m^l \partial_k \partial_l \partial_s^a \partial^\gamma \theta^m \right\|^2 \bar{w}^{7+2|\gamma|} \mathrm{d}\mathbf{x} \mathrm{d}\tau$$

$$+ \mathscr{R}[(\mathscr{C}_n + \mathscr{Z}_n^2)^{1/2} \mathscr{C}_n] + \mathscr{R}[\mathbb{Q}_{n,|\gamma|}] + \mathscr{R}[\mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathscr{R}[\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}]$$

Now we can evaluate the time integral using integration by parts, leaving a remainder term that can be estimated with $(\mathcal{C}_n + \mathcal{X}_n^2)^{1/2} \mathcal{C}_n$ when the time derivative falls on $\mathcal{J}^{-1/3}$,

$$\begin{split} 0 &= \frac{2}{3} \int_{B_R} \mathcal{J}^{-1/3} \left\| \mathcal{A}^k \mathcal{A}_m^l \partial_k \partial_l \partial_s^a \partial^\gamma \theta^m \right\|^2 \bar{w}^{7+2|\gamma|} \mathbf{dx} \bigg|_0^s \\ &+ \mathfrak{R}[(\mathcal{E}_n + \mathcal{X}_n^2)^{1/2} \mathcal{E}_n] + \mathfrak{R}[\mathfrak{Q}_{n,|\gamma|}] + \mathfrak{R}[\mathfrak{Q}_{n,|\gamma|}^{1/2} \mathfrak{Q}_{n,|\gamma|+1}^{1/2}] + \mathfrak{R}[\mathfrak{Q}_{n-1}^{1/2} \mathfrak{Q}_{n,|\gamma|+1}^{1/2}] \\ &= \frac{2}{3} \int_{B_R} \left\| \nabla \nabla \cdot \partial_s^a \partial^\gamma \theta \right\|^2 \bar{w}^{7+2|\gamma|} \mathbf{dx} \bigg|_0^s \\ &+ \mathfrak{R}[(\mathcal{E}_n + \mathcal{X}_n^2)^{1/2} \mathcal{E}_n] + \mathfrak{R}[\mathfrak{Q}_{n,|\gamma|}] + \mathfrak{R}[\mathfrak{Q}_{n,|\gamma|}^{1/2} \mathfrak{Q}_{n,|\gamma|+1}^{1/2}] + \mathfrak{R}[\mathfrak{Q}_{n-1}^{1/2} \mathfrak{Q}_{n,|\gamma|+1}^{1/2}] \end{split}$$

It follows that

$$\begin{split} \|\nabla \nabla \cdot \partial_s^a \partial^{\gamma} \boldsymbol{\theta}\|_{3+2(2+|\gamma|)}^2 \\ \lesssim \mathbb{Q}_{n,|\gamma|+1}(0) + \mathbb{Q}_{n,|\gamma|} + \mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2} + \mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2} + (\mathscr{E}_n + \mathscr{Z}_n^2)^{1/2} \mathscr{E}_n \end{split}$$

Using Lemma 3.48 we get

$$Q_{n,|\gamma|+1} \lesssim \mathbb{Q}_{n,|\gamma|+1}(0) + \mathbb{Q}_{n,|\gamma|} + \mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2} + \mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2} + (\mathcal{E}_n + \mathcal{Z}_n^2)^{1/2} \mathcal{E}_n$$

(ii) Times the equation with $\bar{w}^{6+2|\gamma|}\partial_s^a\partial^\gamma \mathscr{A}\nabla\cdot\boldsymbol{\theta}$ and integrate in time and space we get

$$\begin{split} 0 &= \int_{0}^{s} \left(\left\langle \partial_{s}^{a} \partial^{\gamma} \mathscr{A} \nabla \cdot \partial_{s}^{2} \theta - \frac{1}{2} \mathscr{b} \partial_{s}^{a} \partial^{\gamma} \mathscr{A} \nabla \cdot \partial_{s} \theta + \delta \partial_{s}^{a} \partial^{\gamma} \mathscr{A} \nabla \cdot \theta, \partial_{s}^{a} \partial^{\gamma} \mathscr{A} \nabla \cdot \theta \right\rangle_{6+2|\gamma|} \\ &+ \left\langle \partial_{s}^{a} \partial^{\gamma} \mathscr{A} \nabla \cdot \left(\bar{w}^{-3} \partial_{k} (\bar{w}^{4} T^{k}) \right), \partial_{s}^{a} \partial^{\gamma} \mathscr{A} \nabla \cdot \theta \right\rangle_{6+2|\gamma|} \\ &+ \left\langle 4 \pi \partial_{s}^{a} \partial^{\gamma} (\bar{w}^{3} (\mathcal{F}^{-1} - 1)) - (\partial_{s}^{a} \partial^{\gamma} (\mathscr{A} - I) \nabla) \cdot \nabla \mathcal{K} \bar{w}^{3}, \partial_{s}^{a} \partial^{\gamma} \mathscr{A} \nabla \cdot \theta \right\rangle_{6+2|\gamma|} \right) \mathrm{d}\tau \end{split}$$

Now commuting \mathscr{A} with space and time derivatives, we get a non-linear remainder $\mathscr{R}[(\mathscr{C}_n + \mathscr{L}_n^2)^{1/2} \mathscr{C}_n]$,

$$\begin{split} 0 &= \int_{0}^{s} \left(\left\langle \partial_{s} \mathscr{A} \nabla \cdot \partial_{s}^{a+1} \partial^{\gamma} \theta - \frac{1}{2} \mathscr{E} \mathscr{A} \nabla \cdot \partial_{s}^{a+1} \partial^{\gamma} \theta, \mathscr{A} \nabla \partial_{s}^{a} \partial^{\gamma} \cdot \theta \right\rangle_{6+2|\gamma|} \\ &+ \left\langle \delta \mathscr{A} \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \theta, \mathscr{A} \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \theta \right\rangle_{6+2|\gamma|} \\ &+ \left\langle \mathscr{A} \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \left(\bar{w}^{-3} \partial_{k} (\bar{w}^{4} T^{k}) \right), \mathscr{A} \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \theta \right\rangle_{6+2|\gamma|} \\ &+ \left\langle 4\pi (\mathscr{F}^{-1} - 1) \partial_{s}^{a} \partial^{\gamma} (\bar{w}^{3}) - (\mathscr{A} - I) \nabla \cdot \nabla \mathscr{K} \partial_{s}^{a} \partial^{\gamma} \bar{w}^{3}, \partial_{s}^{a} \partial^{\gamma} \mathscr{A} \nabla \cdot \theta \right\rangle_{6+2|\gamma|} \right) \mathrm{d}\tau \\ &+ \mathscr{R} [(\mathscr{E}_{n} + \mathscr{X}_{n}^{2})^{1/2} \mathscr{E}_{n}] \end{split}$$

Now all the terms, apart from the top order term involving T_T from the pressure, can be bounded by $(\mathscr{C}_n + \mathscr{Z}_n^2)^{1/2} \mathscr{C}_n] + \mathscr{R}[\mathbb{Q}_{n,|\gamma|} + \mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2} + \mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2},$

$$\begin{split} 0 &= -\int_{0}^{s} \left(\left\langle \mathscr{A} \nabla \cdot \partial_{s}^{a+1} \partial^{\gamma} \boldsymbol{\theta}, \partial_{s} \mathscr{A} \nabla \partial_{s}^{a} \partial^{\gamma} \cdot \boldsymbol{\theta} \right\rangle_{6+2|\gamma|} + \left\langle \mathscr{A} \nabla \cdot \left(\bar{w} \partial_{s}^{a} \partial^{\gamma} \partial_{k} T^{k} \right), \mathscr{A} \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \boldsymbol{\theta} \right\rangle_{6+2|\gamma|} \right) \mathrm{d}\tau \\ &+ \mathfrak{R}[(\mathfrak{C}_{n} + \mathfrak{X}_{n}^{2})^{1/2} \mathfrak{C}_{n}] + \mathfrak{R}[\mathfrak{Q}_{n,|\gamma|}^{1/2}] + \mathfrak{R}[\mathfrak{Q}_{n,|\gamma|+1}^{1/2}] + \mathfrak{R}[\mathfrak{Q}_{n-1}^{1/2} \mathfrak{Q}_{n,|\gamma|+1}^{1/2}] \\ &= \int_{0}^{s} \left\langle \mathscr{A} \nabla \cdot \left(\bar{w} T_{T}[\partial_{s}^{a} \partial^{\gamma} \partial_{k} \nabla \boldsymbol{\theta}]^{k} \right), \mathscr{A} \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \boldsymbol{\theta} \right\rangle_{6+2|\gamma|} \mathrm{d}\tau \\ &+ \mathfrak{R}[(\mathfrak{C}_{n} + \mathfrak{X}_{n}^{2})^{1/2} \mathfrak{C}_{n}] + \mathfrak{R}[\mathfrak{Q}_{n,|\gamma|}] + \mathfrak{R}[\mathfrak{Q}_{n,|\gamma|+1}^{1/2}] + \mathfrak{R}[\mathfrak{Q}_{n-1}^{1/2} \mathfrak{Q}_{n,|\gamma|+1}^{1/2}] \end{split}$$

Now we integrate by parts,

$$0 = -\int_{0}^{s} \left\langle \bar{w} T_{T} [\partial_{s}^{a} \partial^{\gamma} \partial_{k} \nabla \boldsymbol{\theta}]^{k}, \mathscr{A} \nabla (\mathscr{A} \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \boldsymbol{\theta}) \right\rangle_{6+2|\gamma|} \mathrm{d}\tau \\ + \mathscr{R} [(\mathscr{C}_{n} + \mathscr{Z}_{n}^{2})^{1/2} \mathscr{C}_{n}] + \mathscr{R} [\mathfrak{Q}_{n,|\gamma|}] + \mathscr{R} [\mathfrak{Q}_{n,|\gamma|}^{1/2} \mathfrak{Q}_{n,|\gamma|+1}^{1/2}] + \mathscr{R} [\mathfrak{Q}_{n-1}^{1/2} \mathfrak{Q}_{n,|\gamma|+1}^{1/2}]$$

Now we expend the terms by defintion and simplify,

$$\begin{split} 0 &= \int_{0}^{s} \left\langle \mathcal{F}^{-1/3} \left(\mathcal{A}_{m}^{k} \mathcal{A}^{l} + \frac{1}{3} \mathcal{A}^{k} \mathcal{A}_{m}^{l} \right) \partial_{s}^{a} \partial^{\gamma} \partial_{k} \partial_{l} \theta^{m}, \mathcal{A}^{j} \mathcal{A}_{i}^{\ell} \partial_{j} \partial_{\ell} \partial_{s}^{a} \partial^{\gamma} \theta^{i} \right\rangle_{7+2|\gamma|} \mathrm{d}\tau \\ &+ \mathcal{R}[(\mathcal{C}_{n} + \mathcal{X}_{n}^{2})^{1/2} \mathcal{C}_{n}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathcal{R}[\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] \\ &= \int_{0}^{s} \left\langle \mathcal{F}^{-1/3} \left(\mathcal{A}_{m}^{k} \mathcal{A}_{o}^{l} + \frac{1}{3} \mathcal{A}_{o}^{k} \mathcal{A}_{m}^{l} \right) \partial_{k} \partial_{l} \partial_{s}^{a} \partial^{\gamma} \theta^{m}, \mathcal{A}_{o}^{j} \mathcal{A}_{i}^{\ell} \partial_{j} \partial_{\ell} \partial_{s}^{a} \partial^{\gamma} \theta^{i} \right\rangle_{7+2|\gamma|} \mathrm{d}\tau \\ &+ \mathcal{R}[(\mathcal{C}_{n} + \mathcal{X}_{n}^{2})^{1/2} \mathcal{C}_{n}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathcal{R}[\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] \\ &= \int_{0}^{s} \frac{4}{3} \left\langle \mathcal{F}^{-1/3} \mathcal{A}_{o}^{k} \mathcal{A}_{m}^{l} \partial_{k} \partial_{l} \partial_{s}^{a} \partial^{\gamma} \theta^{m}, \mathcal{A}_{o}^{j} \mathcal{A}_{i}^{\ell} \partial_{j} \partial_{\ell} \partial_{s}^{a} \partial^{\gamma} \theta^{i} \right\rangle_{7+2|\gamma|} \mathrm{d}\tau \\ &+ \mathcal{R}[(\mathcal{C}_{n} + \mathcal{X}_{n}^{2})^{1/2} \mathcal{C}_{n}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathcal{R}[\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] \\ &= \frac{4}{3} \int_{0}^{s} \int_{B_{R}} \mathcal{F}^{-1/3} \left\| \mathcal{A}^{k} \mathcal{A}_{m}^{l} \partial_{k} \partial_{l} \partial_{s}^{a} \partial^{\gamma} \theta^{m} \right\|^{2} \bar{w}^{7+2|\gamma|} \mathrm{d}x \mathrm{d}\tau \\ &+ \mathcal{R}[(\mathcal{C}_{n} + \mathcal{X}_{n}^{2})^{1/2} \mathcal{C}_{n}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathcal{R}[\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] \\ &= \frac{4}{3} \int_{0}^{s} \int_{B_{R}} \| \nabla \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \theta \|^{2} \bar{w}^{7+2|\gamma|} \mathrm{d}x \mathrm{d}\tau \\ &+ \mathcal{R}[(\mathcal{C}_{n} + \mathcal{X}_{n}^{2})^{1/2} \mathcal{C}_{n}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathcal{R}[\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] \\ &= \frac{4}{3} \int_{0}^{s} \int_{B_{R}} \| \nabla \nabla \cdot \partial_{s}^{a} \partial^{\gamma} \theta \|^{2} \bar{w}^{7+2|\gamma|} \mathrm{d}x \mathrm{d}\tau \\ &+ \mathcal{R}[(\mathcal{C}_{n} + \mathcal{X}_{n}^{2})^{1/2} \mathcal{C}_{n}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|}] + \mathcal{R}[\mathbb{Q}_{n,|\gamma|+1}^{1/2}] + \mathcal{R}[\mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2}] \\ &= \frac{4}{3} \int_{0}^{s} \mathcal{L}_{0}^{1/2} \mathcal{L$$

It follows that

Using Lemma 3.48 we get

$$\int_0^s Q_{n,|\gamma|+1} \mathrm{d}\tau \lesssim \mathbb{Q}_{n,|\gamma|+1}(0) + \mathbb{Q}_{n,|\gamma|} + \mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2} + \mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2} + (\mathscr{C}_n + \mathcal{Z}_n^2)^{1/2} \mathscr{C}_n$$

Combining the results of (i) and (ii) and noting that \lesssim does not depend on *s*, we get that

$$\begin{split} \mathbb{Q}_{n,|\gamma|+1} &\lesssim \mathbb{Q}_{n,|\gamma|+1}(0) + \mathbb{Q}_{n,|\gamma|} + \mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2} + \mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2} + (\mathcal{E}_n + \mathcal{X}_n^2)^{1/2} \mathcal{E}_n \\ &\lesssim \mathcal{E}_n(0) + \mathbb{Q}_{n,|\gamma|} + \mathbb{Q}_{n,|\gamma|}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2} + \mathbb{Q}_{n-1}^{1/2} \mathbb{Q}_{n,|\gamma|+1}^{1/2} + (\mathcal{E}_n + \mathcal{X}_n^2)^{1/2} \mathcal{E}_n \end{split}$$

We have by definition and equation (3.151) in the previous theorem

$$\mathfrak{Q}_{n,0} \lesssim \mathfrak{S}_{n,0} \lesssim |\mathfrak{E}|^{-4} \mathfrak{S}_n(0) + C_{\delta}(\mathfrak{C}_n + \mathfrak{T}_n^2)^{1/2} \mathfrak{C}_n \leq |\mathfrak{E}|^{-4} \mathfrak{C}_n(0) + C_{\delta}(\mathfrak{C}_n + \mathfrak{T}_n^2)^{1/2} \mathfrak{C}_n$$

And so using Young's inequality and by induction we have

$$\mathbb{Q}_{n,d} \lesssim |\mathcal{E}|^{-4} \mathcal{E}_n(0) + C_{\delta} (\mathcal{E}_n + \mathcal{Z}_n^2)^{1/2} \mathcal{E}_n$$

for all $d \leq n$. Therefore we have $\mathbb{Q}_n \lesssim |\mathfrak{E}|^{-4} \mathfrak{C}_n(0) + C_{\delta} (\mathfrak{C}_n + \mathfrak{X}_n^2)^{1/2} \mathfrak{C}_n$.

3.6.3 Bootstrapping scheme and final theorem

In this subsection we will prove our main theorem that the energy E_n decays exponentially while Z_n remains bounded. To do so we will use the bootstrapping scheme in the following lemma and proposition.

Lemma 3.51. Suppose $E: [0,T] \rightarrow [0,\infty]$ is continuous and

$$E(t) \le C_1 E(0) + C_2 E(t)^{3/2}$$
 whenever $\sup_{\tau \in [0,t]} E(\tau) \le C_3.$

where $C_1 \ge 1$. Then $E \le 2C_1 E(0)$ whenever $E(0) \le \min\{(2^5 C_1 C_2^2)^{-1}, C_3/2C_1\}$.

Proof. We will prove this by a standard bootstrap argument. Let

$$I = \left\{ t \in [0,T] : \sup_{\tau \in [0,t]} E(\tau) \le \min\{2C_1 E(0), C_3\} \right\}.$$

Then I is non-empty (since $0 \in I$) and closed (since E is continuous). If I = [0, T], then we are done. Otherwise, let $t_0 = \inf\{t \in [0, T] : t \notin I\}$. We must have $t_0 \in I$ since $0 \in I$ and I is closed. Then we have

$$E(t_0) \le C_1 E(0) + C_2 (2C_1 E(0))^{3/2} \le \frac{3}{2} C_1 E(0) \le \frac{3}{4} C_3.$$

So by continuity of E, a neighbourhood of t_0 must lie in I. But this contradicts the definition of t_0 . So we must have I = [0, T].

Proposition 3.52. Suppose $E, Z : [0, \infty) \to [0, \infty]$ are continuous and for all $t \ge t_0 \ge 0$ we have

$$\mathscr{E}_{t_0}(t) \le C_0 \mathscr{E}_{t_0}(t_0) + C_1 Z(t_0) \mathscr{E}_{t_0}(t) + C_2 (1 + (t - t_0)^k) \mathscr{E}_{t_0}(t)^{3/2}$$

$$Z(t) \le Z(t_0) + C_3 (t - t_0)^l \mathscr{E}_{t_0}^{1/2}(t)$$

whenever $\sup_{\tau \in [t_0,t]} (E(\tau) + Z(\tau)) \le C_4$, where $k, l \ge 0$ and

$$\mathscr{E}_{t_0}(t) = \sup_{\tau \in [t_0,t]} E(\tau) + \int_{t_0}^t E(\tau) \mathrm{d}\tau.$$

Then there exist $\epsilon > 0$ such that $\mathfrak{C}_0 \leq 6C_0\mathfrak{C}_0(0)$ whenever $\mathfrak{C}_0(0), Z(0) \leq \epsilon$. Moreover, $E(t) \leq 16(4^{-t/32C_0})C_0E(0)$.

Proof. Let $T = 32C_0$ and $C_1Z(0) < \min\{1/4, C_1C_4/4\} \le 1/2$. Then by the above Lemma 3.51, for small enough ϵ , we have $\mathscr{C}_0 \le 4C_0\mathscr{C}_0(0)$ on [0, T] whenever $\mathscr{C}_0(0) \le \epsilon$. On [T/2, T], there must exist a point T_1 such that $E(T_1) \le \frac{1}{4}\mathscr{C}_0(0)$, otherwise $\mathscr{C}_0(T) > 4C_0\mathscr{C}_0(0)$.

By having a small enough ϵ , we can assume $2C_0^{1/2}C_1C_3T^{l}\mathcal{E}_0(0)^{1/2} < \min\{1/8, C_1C_4/8\}$. Now

$$C_1 Z(T_1) \le C_1 Z(0) + 2C_0^{1/2} C_1 C_3 T^l \mathscr{E}_0(0)^{1/2}$$

$$\le \min\left\{\frac{1}{4}, \frac{C_1 C_4}{4}\right\} + \min\left\{\frac{1}{8}, \frac{C_1 C_4}{8}\right\} \le \min\left\{\frac{1}{2}, \frac{C_1 C_4}{2}\right\}.$$

Then by the above Lemma 3.51, we get that $\mathscr{C}_{T_1} \leq 4C_0 \mathscr{C}_{T_1}(T_1) = 4C_0 E(T_1) \leq C_0 \mathscr{C}_0(0)$ on $[T_1, T_1 + T]$. On $[T_1 + T/2, T_1 + T]$, there must exist a point T_2 such that $E(T_2) \leq \frac{1}{4} \mathscr{C}_{T_1}(T_1) = \frac{1}{4} E(T_1)$, otherwise $\mathscr{C}_{T_1}(T) > 4C_0 \mathscr{C}_{T_1}(T_1)$.

Repeating inductively, we can get $T_n \in [T_{n-1} + T/2, T_{n-1} + T]$ such that

$$C_1 Z(T_n) \le C_1 Z(T_{n-1}) + C_1 C_3 T^l \mathscr{C}_{T_{n-1}}(T_n)^{1/2}$$
$$\le \left(\frac{1}{4} \sum_{m=0}^{n-1} \frac{1}{2^m} + \frac{1}{4} \frac{1}{2^n}\right) \min\{1, C_1 C_4\} \le \frac{1}{2} \min\{1, C_1 C_4\}$$

$$\mathscr{C}_{T_n} \le 4^{1-n} C_0 \mathscr{C}_0(0) \qquad \text{on} \qquad [T_n, T_n + T].$$

Now

$$\mathscr{E}_0 \leq \mathscr{E}_0(\infty) \leq \mathscr{E}_0(T_1) + \mathscr{E}_{T_1}(T_2) + \mathscr{E}_{T_2}(T_3) + \dots \leq 6C_0 \mathscr{E}_0(0).$$

Π

Finally, before proving the main theorem, we provide a simple lemma based on the fundamental theorem of calculus, which relates the \mathcal{Z}_n -norm (3.66) to the total energy norm \mathcal{C}_n (3.70). Lemma 3.53. We have

$$\mathfrak{X}_{1}(s) \leq \mathfrak{X}_{1}(0) + Cs^{1/2}\mathfrak{S}_{1}^{1/2}(s)$$

 $\mathfrak{X}_{n}(s) \leq \mathfrak{X}_{n}(0) + Cs^{1/2}\mathfrak{S}_{n}^{1/2}(s)$

Proof. We have $h(s) - h(0) = \int_0^s \partial_s h(\tau) d\tau$ and therefore

$$(h(s) - h(0))^2 = \left(\int_0^s \partial_s h(\tau) \mathrm{d}\tau\right)^2 \le s \int_0^s (\partial_s h(\tau))^2 \mathrm{d}\tau.$$

This easily gives $\|h(s) - h(0)\|_k^2 \leq s \int_0^s \|\partial_s h(\tau)\|_k^2 d\tau$ and thus $\|h(s)\|_k \leq \|h(0)\|_k + s^{1/2} \left(\int_0^s \|\partial_s h(\tau)\|_k^2 d\tau\right)^{1/2}$, which concludes the proof.

Theorem 3.54. Let $n \ge 21$ and δ small. Let $(\theta, \partial_s \theta)$ be a solution of (3.62) in the sense of Theorem 3.11 and that satisfies (3.73), (3.74) and (3.75) (i.e. the perturbation does not change the momentum or energy of the star, and correspond to an irrotational flow). Then there is some m > 0 such that we have

$$\mathscr{E}_n(s) \lesssim |\mathscr{E}|^{-m} \mathscr{E}_n(0) + C_\delta \left(\mathscr{Z}_n(0) \mathscr{E}_n(s) + (1 + s^{1/2}) \mathscr{E}_n(s)^{3/2} \right)$$
(3.153)

whenever our a priori assumption (3.71) is satisfied. Moreover, there exists $\epsilon_0 > 0$ such that if $E_n(0) + Z_n(0)^2 \leq \epsilon_0$, then we have $\mathcal{C}_n \lesssim \mathcal{C}_n(0)$ with $E_n(s) \lesssim e^{-C|\beta|^m s}$ (decaying exponentially on $[0,\infty)$) and Z_n bounded on $[0,\infty)$.

Proof. By the energy estimates in Theorem 3.47 and 3.50 (for \mathscr{S}_n and \mathfrak{Q}_n) in the last two subsections and Lemma 3.53 we have (choosing ϵ small enough)

$$\begin{split} & \mathcal{E}_n \lesssim \mathcal{S}_n + \mathcal{Q}_n - C\epsilon \mathcal{E}_n \lesssim_{\epsilon} |\mathfrak{b}|^{-m} \mathcal{E}_n(0) + C_{\delta} (\mathcal{E}_n + \mathcal{I}_n^2)^{1/2} \mathcal{E}_n \\ & \lesssim |\mathfrak{b}|^{-m} \mathcal{E}_n(0) + C_{\delta} \left(\mathcal{I}_n(0) \mathcal{E}_n(s) + (1 + s^{1/2}) \mathcal{E}_n(s)^{3/2} \right) \end{split}$$

Using Proposition 3.52 above, we get $\mathscr{C}_n \lesssim \mathscr{C}_n(0)$ with $E_n(s) \lesssim e^{-C|\mathscr{E}|^m s}$ and Z_n bounded on $[0,\infty)$. \Box

4 Nonradial stability of linearly expanding Goldreich-Weber stars

4.1 Formulation and statement of the result

4.1.1 Equation in linearly-expanding coordinates

In this section we will take the enthalpy \bar{w} to be the profile associated with the linearly expanding GW star from Definition 1.3. To study the linearly expanding GW stars, we want to write our variables as a perturbation from the model GW star. To that end we will use the rescaled variable ξ (equation (2.34)) introduced in Section 2 adapted to the expanding background profile and also write the problem in "linear" time variables. We introduce the "linear" time coordinate *s* adapted to the expanding profile via

$$\frac{ds}{dt} = \lambda(t)^{-1}.$$

In this new coordinate, $\lambda(s)$ is an increasing function such that

$$\lambda(s) \sim e^{s\sqrt{\lambda_1^2 + 2\delta}}$$
 as $s \to \infty$. (4.154)

We have the following change of coordinate formula $\partial_t = \lambda^{-1} \partial_s$. The condition $\ddot{\lambda} \lambda^2 = \delta$ (1.17) becomes

$$\delta = \lambda \partial_s (\lambda^{-1} \partial_s \lambda) = \partial_s^2 \lambda - \frac{(\partial_s \lambda)^2}{\lambda}$$
(4.155)

Then the Euler-Poisson equations (1.13) becomes

$$\mathbf{0} = \partial_t \mathbf{v} + (f_0 J_0)^{-1} \partial_k (A^k (f_0 J_0)^{4/3} J^{-1/3}) + A \nabla \psi$$

= $\lambda^{-1} \partial_s (\lambda^{-1} \partial_s (\lambda \boldsymbol{\xi})) + \lambda^{-2} (f_0 J_0)^{-1} \partial_k (\mathcal{A}^k (f_0 J_0)^{4/3} \boldsymbol{\xi}^{-1/3}) + \lambda^{-2} \mathcal{A} \nabla \Phi$

Times the equation by λ^2 we get

$$\begin{aligned} \mathbf{0} &= \lambda \partial_s (\lambda^{-1} \partial_s (\lambda \boldsymbol{\xi})) + (f_0 J_0)^{-1} \partial_k (\mathcal{A}^k (f_0 J_0)^{4/3} \mathcal{F}^{-1/3}) + \mathcal{A} \nabla \Phi \\ &= \lambda \left(\partial_s^2 \boldsymbol{\xi} + \frac{\partial_s \lambda}{\lambda} \partial_s \boldsymbol{\xi} + \left(\frac{\partial_s^2 \lambda}{\lambda} - \frac{(\partial_s \lambda)^2}{\lambda^2} \right) \boldsymbol{\xi} \right) + (f_0 J_0)^{-1} \partial_k (\mathcal{A}^k (f_0 J_0)^{4/3} \mathcal{F}^{-1/3}) + \mathcal{A} \nabla \Phi \\ &= \left(\lambda \partial_s^2 \boldsymbol{\xi} + (\partial_s \lambda) \partial_s \boldsymbol{\xi} + \delta \boldsymbol{\xi} \right) + (f_0 J_0)^{-1} \partial_k (\mathcal{A}^k (f_0 J_0)^{4/3} \mathcal{F}^{-1/3}) + \mathcal{A} \nabla \Phi \end{aligned}$$

So the Euler-Poisson equations in terms of $\boldsymbol{\xi}$ (2.34) is:

$$\lambda \partial_s^2 \boldsymbol{\xi} + \lambda' \partial_s \boldsymbol{\xi} + \delta \boldsymbol{\xi} + \frac{1}{f_0 J_0} \partial_k (\mathscr{A}^k (f_0 J_0)^{4/3} \boldsymbol{\mathcal{J}}^{-1/3}) + \mathscr{A} \nabla \Phi = \mathbf{0},$$
(4.156)

where $\lambda' := \partial_s \lambda$.

The GW-star is a particular s-independent solution of (4.156) of the form $\boldsymbol{\xi}(\mathbf{x}) \equiv \mathbf{x}$ and $f_0 = \bar{w}^3$. Before formulating the stability problem, we must first make the use of the labelling gauge freedom and fix the choice of the initial enthalpy $(f_0 J_0)^{1/3}$ for the general perturbation to be exactly identical to the background enthalpy \bar{w} , i.e. we set

$$(f_0 J_0)^{1/3} = \bar{w}$$
 on $B_R(\mathbf{0})$. (4.157)

Equation (4.157) can be re-written in the form $\rho_0 \circ \eta_0 \det[\nabla \eta_0] = \bar{w}^3$ on the initial domain $B_R(\mathbf{0})$. By a result of Dacorogna-Moser [10] and similarly to [24, 25] there exists a choice of an initial bijective map $\eta_0 : B_R(\mathbf{0}) \to \Omega(\mathbf{0})$ so that (4.157) holds true. The gauge fixing condition (4.157) is necessary as it constraints the freedom to arbitrary relabel the particles at the initial time.

Lemma 4.1 (Euler-Poisson in linearly-expanding coordinate). With respect to the linearly expanding profile (λ, \bar{w}) from Definition 1.3, the perturbation θ defined in (2.35) formally solves

$$\lambda \partial_s^2 \theta + \lambda' \partial_s \theta + \delta \theta + \mathbf{P} + \mathbf{G} = \mathbf{0}, \tag{4.158}$$

where the nonlinear pressure operator \mathbf{P} and the nonlinear gravity operator \mathbf{G} are defined in (2.36) and (2.37).

Proof. Recall that the GW-enthalpy satisfies

$$\mathbf{0} = \delta \mathbf{x} + 4\nabla \bar{w} + \nabla \mathcal{K} \bar{w}^3 \tag{4.159}$$

Using the gauge condition (4.157), the momentum equation (4.156) becomes

$$\bar{w}^3 \left(\lambda \partial_s^2 \boldsymbol{\theta} + \lambda' \partial_s \boldsymbol{\theta} + \delta \boldsymbol{\theta} \right) + \partial_k (\bar{w}^4 (\mathcal{A}^k \mathcal{J}^{-1/3} - I^k)) + \bar{w}^3 (\mathcal{A} \nabla \Phi - \nabla \mathcal{K} \bar{w}^3) = \mathbf{0}.$$

Hence, we can write the momentum equation as

$$\mathbf{0} = \lambda \partial_s^2 \boldsymbol{\theta} + \lambda' \partial_s \boldsymbol{\theta} + \delta \boldsymbol{\theta} + \underbrace{\bar{w}^{-3} \partial_k (\bar{w}^4 (\mathcal{A}^k \mathcal{F}^{-1/3} - I^k))}_{\mathbf{P}} + \underbrace{\mathcal{A} \nabla \Phi - \mathcal{K} \nabla \bar{w}^3}_{\mathbf{G}}.$$

4.1.2 High-order energies and the main theorem

We now introduce high-order weighted Sobolev norm that we will use for our high-order energy method explained in Section 4.5. Recall the notation in Section 2. Assuming that $(s, \mathbf{y}) \mapsto \boldsymbol{\theta}(s, \mathbf{y})$ is a sufficiently smooth field, for any $n \in \mathbb{N}_0$ we let

$$S_{n}(s) := \sum_{|\beta|+b \le n} \left(\lambda \|X_{r}^{b} \partial^{\beta} \partial_{s} \theta\|_{3+b}^{2} + \|X_{r}^{b} \partial^{\beta} \theta\|_{3+b}^{2} + \|\nabla X_{r}^{b} \partial^{\beta} \theta\|_{4+b}^{2} \right)$$
$$Q_{n}(s) := \sum_{c \le n} \left(\lambda \|\nabla^{c} \partial_{s} \theta\|_{3+2c}^{2} + \|\nabla^{c} \theta\|_{3+2c}^{2} + \|\nabla^{c+1} \theta\|_{4+2c}^{2} \right)$$
$$Z_{n}(s) := \sum_{|\beta|+b=n} \lambda \|X_{r}^{b} \partial^{\beta} (\mathscr{A} \nabla \times \partial_{s} \theta)\|_{4+b}^{2} + \lambda \|\nabla^{n} (\mathscr{A} \nabla \times \partial_{s} \theta)\|_{4+2n}^{2}$$

We define the total instant energy via

$$E_n := S_n + Q_n + Z_n. (4.160)$$

We shall run the energy identity using E_n ; Z_n controls the curl of the velocity, while the energies S_n and Q_n will be used for high-order estimates near the vacuum boundary and near the origin respectively. In particular, the control afforded by Q_n is stronger near the origin, while S_n is stronger near the boundary. Finally we define

$$\mathcal{S}_n(s) := \sup_{\tau \in [0,s]} S_n(\tau),$$
(4.161)

$$\mathfrak{Q}_n(s) := \sup_{\tau \in [0,s]} Q_n(\tau), \tag{4.162}$$

$$\mathscr{E}_n(s) := \sup_{\tau \in [0,s]} E_n(\tau), \tag{4.163}$$

The norms (4.161)–(4.163) will play the role of the "left hand side" in the high-order energy identities.

In this section, we make the following a priori assumption:

A priori assumption: $E_n \le \epsilon$ where $\epsilon > 0$ is some small constant. (4.164)

We now state our main theorem.

Theorem 4.2 (Nonlinear stability of GW stars). Let $n \ge 21$. The linearly expanding GW star from Definition 1.3 nonlinearly stable. More precisely, there exists an $\epsilon^* > 0$ such that for any initial data $(\theta(0), \partial_s \theta(0))$ satisfying

$$E_n(0) \le \epsilon^*, \tag{4.165}$$

the associated solution $s \mapsto (\theta(s, \cdot), \partial_s \theta(s, \cdot))$ to (4.158) exists for all $s \ge 0$ and is unique in the class of all data with finite norm E_n . Moreover, there exists a constant C > 0 such that

 $E_n(s) \le C\epsilon^*$ for all $s \ge 0$.

Remark 4.3. Like in the last section (cf. Remark 3.9), it is not our goal to optimise the number n of derivatives in our spaces.

Local-in-time well-posedness. The same process as described in section 3.1.3 for the self-similarly expanding GW star can be use to obtain the equivalent well-posedness result in the weighted high-order energy space E_n defined in the current section for the linearly expanding GW star.

Theorem 4.4 (Local well-posedness). Let $n \ge 21$. Then for any given initial data $(\theta(0), \partial_s \theta(0))$ such that $E_n(0) < \infty$, there exist some T > 0 and a unique solution $(\theta, \partial_s \theta) : [0, T] \times B_R \to \mathbb{R}^3 \times \mathbb{R}^3$ to (4.158) such that $E_n(s) \le 2E_n(0)$ for all $s \in [0, T]$.

Theorem 4.4 is a starting point for the continuity argument that will culminate in the proof of Theorem 4.2.

4.1.3 **Proof strategy**

The basic idea behind the global existence in Theorem 4.2 is similar to that of the self-similarly expanding GW case in the last section. In fact, it is more straightforward here than in the last section owing to the fact the linearly expanding GW star expands at a faster rate that the self-similarly expanding GW star, and hence there is a stronger dispersion effect.

In particular, we have the exponentially increasing $\lambda(s)$ factor in the first term in (4.158). This leads to a λ -factor in front of the "velocity" terms in the higher order energy in (4.160). This gives terms on the velocity level (terms with at least one time derivatives) an extra decay that effectively make it subleading order on par with the non-linear term and hence negligible in the dynamics.

This in particular renders the effect of gravity in the dynamics to be secondary:

$$\int_{s_0}^s \langle X_r^b \partial^\beta \mathbf{G}, \partial_s X_r^b \partial^\beta \theta \rangle_{3+b} \mathrm{d}\tau \lesssim \mathfrak{E}_n(s) \int_{s_0}^s \lambda^{-1/2} \mathrm{d}\tau.$$

Here $\mathscr{C}_n(s) \int_{s_0}^s \lambda^{-1/2} d\tau$ has and effect similar to the non-linear term $\mathscr{C}^{3/2}$, see Proposition 4.15 and Theorem 4.16.

This leads to a key simplification - we do not need a precise coercivity result like in the self-similarly expanding GW case for the operator L in Section 3. In particular we do not need to make the assumption that the fluid is irrotational in this case – we allow nontrivial vorticity initially and control its time evolution by the curl estimates (Section 4.4), similar to [25]. Note also that linear motion is secondary (bounded) in a linearly expanding coordinate. So a non-zero momentum in the initial data, which in theory should make the overall GW star to travel at constant speed in the direction of the momentum, is automatically encapsulated by the linear expanding coordinate about a linearly expanding GW star centred at the origin.

Many terms that appeared on the primary "linear level" in self-similarly expanding case of the last section are now not at leading order any more. As a result, higher time derivatives can be avoided in our higher order energy in (4.160), and we do not need the sophisticated triple induction scheme on the higher order energies that we had to carry out in the proof of Theorem 3.7.

4.2 Pressure estimates

In this section we will estimate the non-linear part of the pressure term $X_r^b \partial^\beta \mathbf{P}$ and $\partial^\gamma \mathbf{P}$ (4.158). More precisely, when doing energy estimates, terms like $\langle X_r^b \partial^\beta \mathbf{P}, \partial_s X_r^b \partial^\beta \theta \rangle$ will arise, we will show that \mathbf{P} here can be reduced to $\mathbf{P}_{d,L}$ modulo remainder terms that can be estimated. We will use results from section 2.1.2.

Using Lemma 2.12, we will now estimate the difference between " \mathbf{P}_b " and " $\mathbf{P}_{b,L}$ ".

Proposition 4.5. Let $n \ge 20$. Let $|\beta| + b \le n$ and $|\gamma| \le n$. Suppose θ satisfies our a priori assumption (4.164). Then, for any $0 \le s_0 \le s$, we have

$$\left| \int_{s_0}^s \langle \mathbf{P}_b X_r^b \partial^\beta \boldsymbol{\theta}, \partial_s X_r^b \partial^\beta \boldsymbol{\theta} \rangle_{3+b} \mathrm{d}\tau - \frac{1}{2} \left\langle \mathbf{P}_{b,L} X_r^b \partial^\beta \boldsymbol{\theta}, X_r^b \partial^\beta \boldsymbol{\theta} \rangle_{3+b} \right|_{s_0}^s \right| \lesssim \mathfrak{E}_n(s)^{3/2} \\ \left| \int_{s_0}^s \langle \mathbf{P}_{2|\gamma|} \partial^\gamma \boldsymbol{\theta}, \partial_s \partial^\gamma \boldsymbol{\theta} \rangle_{3+2|\gamma|} \mathrm{d}\tau - \frac{1}{2} \left\langle \mathbf{P}_{2|\gamma|,L} \partial^\gamma \boldsymbol{\theta}, \partial^\gamma \boldsymbol{\theta} \rangle_{3+2|\gamma|} \right|_{s_0}^s \right| \lesssim \mathfrak{E}_n(s)^{3/2}$$

Proof. The proof is similar to Proposition 3.38 except the reminder term here will be $\partial_s \Re[E_n^{3/2}] + \Re[\lambda^{-1/2}E_n^{3/2}]$. Since $\int_0^\infty \lambda^{-1/2} ds < \infty$, integrating in time we get the first equation. Proof for the second formula is similar.

And now we will estimate the difference between " \mathbf{P} " and " \mathbf{P}_b ".

Proposition 4.6. Let $n \ge 20$ and $|\beta| + b \le n$. Suppose θ satisfies our a priori assumption (4.164). Then, for any $0 \le s_0 \le s$, we have

$$\int_{s_0}^s \langle X_r^b \partial^\beta \mathbf{P} - \mathbf{P}_b X_r^b \partial^\beta \boldsymbol{\theta}, \partial_s X_r^b \partial^\beta \boldsymbol{\theta} \rangle_{3+b} \mathrm{d}\tau \bigg| \lesssim \int_{s_0}^s \lambda^{-\frac{1}{2}} E_n \mathrm{d}\tau$$

$$\left|\int_{s_0}^s \langle \partial^{\gamma} \mathbf{P} - \mathbf{P}_{2|\gamma|} \partial^{\gamma} \boldsymbol{\theta}, \partial_s \partial^{\gamma} \boldsymbol{\theta} \rangle_{3+2|\gamma|} \mathrm{d}\tau\right| \lesssim \int_{s_0}^s \lambda^{-\frac{1}{2}} E_n \mathrm{d}\tau.$$

Proof. By Lemma 2.10 we need to estimate the following.

$$\begin{split} & \left| \int_{s_0}^s \int_{B_R} \left(\sum_{\substack{b' \leq b \\ |\beta'| \leq |\beta| - 1}} + \sum_{\substack{b' \leq b - 1 \\ |\beta'| \leq |\beta| + 1}} \right) \langle C\omega X_r^{b'} \partial^{\beta'} T \rangle \langle \partial_s X_r^{b} \partial^{\beta} \theta \rangle \bar{w}^{3+b} \mathrm{dxd}\tau \right| \\ & + \left| \int_{s_0}^s \int_{B_R} \left(\sum_{\substack{b' \leq b - 1 \\ |\beta'| \leq |\beta| - 1}} + \sum_{\substack{b' \leq b - 2 \\ |\beta'| \leq |\beta| - 1}} \right) \langle CX_r^{b'} \partial^{\beta'} \nabla T \rangle \langle \partial_s X_r^{b} \partial^{\beta} \theta \rangle \bar{w}^{3+b} \mathrm{dxd}\tau \right| \\ & + \left| \int_{s_0}^s \int_{B_R} \left(\sum_{\substack{b' \leq b \\ |\beta'| \leq |\beta| - 1}} + \sum_{\substack{b' \leq b - 1 \\ |\beta'| \leq |\beta|}} \right) \langle C\bar{w} X_r^{b'} \partial^{\beta'} \nabla T \rangle \langle \partial_s X_r^{b} \partial^{\beta} \theta \rangle \bar{w}^{3+b} \mathrm{dxd}\tau \right| \\ & \lesssim \int_{s_0}^s \lambda^{-\frac{1}{2}} E_n \mathrm{d}\tau \end{split}$$

where the $\lambda^{-1/2}$ factor comes from estimating $\|\partial_s X_r^b \partial^\beta \theta\|_{3+b} \leq \lambda^{-1/2} E_n^{1/2}$. The terms with T can be estimated noting the structure given in Lemma 2.11. This proves the first formula. The proof for the second formula is similar.

4.3 Gravity estimates

In this subsection we will estimate the gravity term $X_r^b \partial^\beta \mathbf{G}$ and $\partial^\gamma \mathbf{G}$ (4.158) and show that it can be bounded by E_n . We will use results from Section 2.1.1.

Since the gravity term is a non-local term, we need to estimate convolution-like operator. However, rather than the convolution kernel $|\mathbf{x} - \mathbf{z}|^{-1}$ we actually need to estimate $|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|^{-1}$. Lemma 2.5 and the following lemma tell us how to reduce the latter to the former, which will allows us to estimate using the Young's convolution inequality.

Lemma 4.7. Let $\boldsymbol{\xi}$ and $\boldsymbol{\theta}$ be as in (2.35), and $\boldsymbol{\theta}$ satisfies our a priori assumption (4.164). Let $n \geq 21$ and $|\beta| \leq n$.

i. When $|\beta| > n/2$ we have

$$\left| (\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^{\beta} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} \right) \right| \lesssim \frac{1}{|\mathbf{x} - \mathbf{z}|^2} \sum_{n/2 < |\gamma| \le n} |\partial_{\mathbf{x}}^{\gamma} \boldsymbol{\xi}(\mathbf{x}) - \partial_{\mathbf{z}}^{\gamma} \boldsymbol{\xi}(\mathbf{z})|$$

ii. When $|\beta| \leq n/2$ we have

$$\left|\partial_{i,\mathbf{z}}(\mathbf{\partial}_{\mathbf{x}} + \mathbf{\partial}_{\mathbf{z}})^{\beta} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|}\right)\right| \lesssim \frac{1}{|\mathbf{x} - \mathbf{z}|^2}$$

Proof. The proof proceed like Lemma 3.40 but using instead the embedding theorems A.10 and A.11 and the a priori bounds $E_n \leq 1$ (4.164).

We next derive an helpful lemma for derivatives on $\mathcal{K}_{\xi} - \mathcal{K}$. Recall K_1 from (3.89) with

$$(\mathcal{H}_{\boldsymbol{\xi}} - \mathcal{H})g(\mathbf{x}) = -\int_{\mathbb{R}^3} K_1(\mathbf{x}, \mathbf{z})g(\mathbf{z})d\mathbf{z},$$
(4.166)

Lemma 4.8. Let $n \ge 20$ and $|\beta| \le n$. We have

$$|(\partial_{\mathbf{x}} + \partial_{\mathbf{z}})^{eta} K_1(\mathbf{x}, \mathbf{z})| \lesssim rac{1}{|\mathbf{x} - \mathbf{z}|^2} \sum_{eta' \leq eta} |\partial^{eta'} m{ heta}(\mathbf{x}) - \partial^{eta'} m{ heta}(\mathbf{z})|$$

Proof. The proof proceed like Lemma 3.25 but using instead the embedding theorems A.10 and A.11 and the a priori bounds $E_n \leq 1$ (4.164).

Finally we can prove the main results of this subsection.

Proposition 4.9 (Gravity estimates). Let $n \ge 21$ and suppose θ satisfies our a priori assumption (4.164). Then we have

$$\begin{split} \|X_r^b \partial^{\beta} \mathbf{G}\|_{3+b}^2 \lesssim E_n & \text{when} & |\beta| + b \le n, \\ \|\bar{w}^{b/2} X_r^b \partial^{\beta} \mathbf{G}\|_{L^{\infty}(\mathbb{R}^3)}^2 \lesssim E_n & \text{when} & |\beta| + b \le n/2, \\ \|\nabla^c \mathbf{G}\|_{3+2c}^2 \lesssim E_n & \text{when} & c \le n, \\ \|\bar{w}^{c/2} \nabla^c \mathbf{G}\|_{L^{\infty}(\mathbb{R}^3)}^2 \lesssim E_n & \text{when} & c \le n/2. \end{split}$$

Proof. By definition

$$\begin{split} \mathbf{G} &= \mathscr{K}_{\boldsymbol{\xi}} \nabla \cdot (\mathscr{A} \bar{w}^3) - \mathscr{K} \nabla \bar{w}^3 = -\int_{\mathbb{R}^3} \frac{\partial_k (\mathscr{A}^k \bar{w}^3)}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} d\mathbf{z} + \int_{\mathbb{R}^3} \frac{\nabla \bar{w}^3}{|\mathbf{x} - \mathbf{z}|} d\mathbf{z} \\ &= -\int_{\mathbb{R}^3} \frac{\partial_k ((\mathscr{A}^k - I^k) \bar{w}^3)}{|\boldsymbol{\xi}(\mathbf{x}) - \boldsymbol{\xi}(\mathbf{z})|} d\mathbf{z} - \int_{\mathbb{R}^3} K_1(\mathbf{x}, \mathbf{z}) \nabla \bar{w}^3 d\mathbf{z} \end{split}$$

By Lemma 2.8 we have

$$\begin{split} \vartheta^{\beta}\mathbf{G}(\mathbf{x}) &= -\vartheta^{\beta}\left(\int_{\mathbb{R}^{3}} \frac{\partial_{k}((\mathscr{A}^{k}-I^{k})\bar{w}^{3})}{|\boldsymbol{\xi}(\mathbf{x})-\boldsymbol{\xi}(\mathbf{z})|} \mathrm{d}\mathbf{z} + \int_{\mathbb{R}^{3}} K_{1}(\mathbf{x},\mathbf{z})\nabla\bar{w}^{3}\mathrm{d}\mathbf{z}\right) \\ &= -\int_{\mathbb{R}^{3}} \sum_{\beta_{1}+\beta_{2}=\beta} (\vartheta_{\mathbf{x}}+\vartheta_{\mathbf{z}})^{\beta_{1}} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x})-\boldsymbol{\xi}(\mathbf{z})|}\right) \vartheta_{\mathbf{z}}^{\beta_{2}}\partial_{k}((\mathscr{A}^{k}-I^{k})\bar{w}^{3})(\mathbf{z})\mathrm{d}\mathbf{z} \\ &- \int_{\mathbb{R}^{3}} \sum_{\beta_{1}+\beta_{2}=\beta} (\vartheta_{\mathbf{x}}+\vartheta_{\mathbf{z}})^{\beta_{1}} K_{1}(\mathbf{x},\mathbf{z})(\vartheta_{\mathbf{z}}^{\beta_{2}}\nabla\bar{w}^{3})(\mathbf{z})\mathrm{d}\mathbf{z} \\ &= -\int_{\mathbb{R}^{3}} \sum_{\substack{\beta_{1}+\beta_{2}=\beta\\|\beta_{1}|>n/2}} (\vartheta_{\mathbf{x}}+\vartheta_{\mathbf{z}})^{\beta_{1}} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x})-\boldsymbol{\xi}(\mathbf{z})|}\right) \vartheta_{\mathbf{z}}^{\beta_{2}}\partial_{k}((\mathscr{A}^{k}-I^{k})\bar{w}^{3})(\mathbf{z})\mathrm{d}\mathbf{z} \\ &- \int_{\mathbb{R}^{3}} \sum_{\substack{\beta_{1}+\beta_{2}=\beta\\|\beta_{1}|\leq n/2}} (\vartheta_{\mathbf{x}}+\vartheta_{\mathbf{z}})^{\beta_{1}} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x})-\boldsymbol{\xi}(\mathbf{z})|}\right) \vartheta_{\mathbf{z}}^{\beta_{2}}\partial_{k}((\mathscr{A}^{k}-I^{k})\bar{w}^{3})(\mathbf{z})\mathrm{d}\mathbf{z} \\ &- \int_{\mathbb{R}^{3}} \sum_{\substack{\beta_{1}+\beta_{2}=\beta\\|\beta_{1}|\leq n/2}} (\vartheta_{\mathbf{x}}+\vartheta_{\mathbf{z}})^{\beta_{1}} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x})-\boldsymbol{\xi}(\mathbf{z})|}\right) \vartheta_{\mathbf{z}}^{\beta_{2}}\partial_{k}((\mathscr{A}^{k}-I^{k})\bar{w}^{3})(\mathbf{z})\mathrm{d}\mathbf{z} \\ &+ \int_{\mathbb{R}^{3}} \sum_{\substack{\beta_{1}+\beta_{2}=\beta\\|\beta_{1}|\leq n/2}} (\vartheta_{\mathbf{x}}+\vartheta_{\mathbf{z}})^{\beta_{1}} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x})-\boldsymbol{\xi}(\mathbf{z})|}\right) (\langle\vartheta_{\mathbf{z}}^{\beta_{2}}(\mathscr{A}-I^{k})\bar{w}^{3})(\mathbf{z})\mathrm{d}\mathbf{z} \\ &+ \int_{\mathbb{R}^{3}} \sum_{\substack{\beta_{1}+\beta_{2}=\beta\\|\beta_{1}|\leq n/2}} (\vartheta_{\mathbf{x}}+\vartheta_{\mathbf{z}})^{\beta_{1}} \left(\frac{1}{|\boldsymbol{\xi}(\mathbf{x})-\boldsymbol{\xi}(\mathbf{z})|}\right) (\langle\vartheta_{\mathbf{z}}^{\beta_{2}}(\mathscr{A}-I)\rangle\bar{w}^{3})(\mathbf{z})\mathrm{d}\mathbf{z} \\ &- \int_{\mathbb{R}^{3}} \sum_{\substack{\beta_{1}+\beta_{2}=\beta\\|\beta_{1}+\beta_{2}=\beta}} (\vartheta_{\mathbf{x}}+\vartheta_{\mathbf{z}})^{\beta_{1}} K_{1}(\mathbf{x},\mathbf{z})(\vartheta_{\mathbf{z}}^{\beta_{2}}\nabla\bar{w}^{3})(\mathbf{z})\mathrm{d}\mathbf{z} \end{split}$$

Now using Lemma 4.7 and 4.8, we get

$$|\partial_s^a \partial^eta^eta \mathbf{G}(\mathbf{x})| \lesssim \int_{\mathbb{R}^3} rac{E_n^{1/2}}{|\mathbf{x} - \mathbf{z}|^2} \sum_{n/2 < |\gamma| \leq n} |\partial_{\mathbf{x}}^\gamma \boldsymbol{\xi}(\mathbf{x}) - \partial_{\mathbf{z}}^\gamma \boldsymbol{\xi}(\mathbf{z})| ar{w}^2 \mathrm{d}\mathbf{z}$$

$$\begin{split} &+ \int_{\mathbb{R}^3} \sum_{\beta_2 \leq \beta} \frac{1}{|\mathbf{x} - \mathbf{z}|^2} (\langle \partial_{\mathbf{z}}^{\beta_2} (\mathcal{A} - I) \rangle \bar{w}^3)(\mathbf{z}) d\mathbf{z} \\ &+ \int_{\mathbb{R}^3} \sum_{\beta_1 + \beta_2 = \beta} \sum_{\beta' \leq \beta_1} \frac{|\partial^{\beta'} \boldsymbol{\theta}(\mathbf{x}) - \partial^{\beta'} \boldsymbol{\theta}(\mathbf{z})|}{|\mathbf{x} - \mathbf{z}|^2} (\partial_{\mathbf{z}}^{\beta_2} \nabla \bar{w}^3)(\mathbf{z}) d\mathbf{z} \end{split}$$

Now using Young's convolution inequality we get

$$\|\partial^{\beta}\mathbf{G}(\mathbf{x})\|_{L^{2}(\mathbb{R}^{3})} \lesssim E_{n}^{1/2}$$

Hence $\|\partial^{\beta} \mathbf{G}\|_{3}^{2} \lesssim E_{n}$. From the above proof, with small modification, we can further see that

$$\|\partial^{\beta} \mathbf{G}(\mathbf{x})\|_{L^{\infty}(\mathbb{R}^3)} \lesssim E_n^{1/2} \quad \text{when} \quad |\beta| \le n/2$$

Now we deal with the case b > 0. Let

$$W_{n} = \sum_{\substack{|\beta|+b \leq n \\ b \leq d}} \|X_{r}^{b} \partial^{\beta} \mathbf{G}\|_{3+b}^{2}$$
$$W_{n,d} = \sum_{\substack{|\beta|+b \leq n \\ b \leq d}} \|X_{r}^{b} \partial^{\beta} \mathbf{G}\|_{3+b}^{2}$$
$$V_{n} = \sum_{\substack{|\beta|+b \leq n \\ b \leq d}} \sup_{\mathbb{R}^{3}} \left(\bar{w}^{b} |X_{r}^{b} \partial^{\beta} \mathbf{G}|^{2}\right)$$
$$V_{n,d} = \sum_{\substack{|\beta|+b \leq n \\ b \leq d}} \sup_{\mathbb{R}^{3}} \left(\bar{w}^{b} |X_{r}^{b} \partial^{\beta} \mathbf{G}|^{2}\right).$$

For $|\beta| + b \le n/2$, using the above lemmas 2.6 and 2.7 we have

$$\begin{split} \bar{w}^{b} |X_{r}^{b} \partial^{\beta} \mathbf{G}|^{2} &\lesssim \bar{w}^{b} |r \nabla \cdot X_{r}^{b-1} \partial^{\beta} \mathbf{G}|^{2} + \bar{w}^{b} |r \nabla \times X_{r}^{b-1} \partial^{\beta} \mathbf{G}|^{2} + \sum_{k=1}^{3} \bar{w}^{b} |X_{r}^{b-1} \partial_{k} \partial^{\beta} \mathbf{G}|^{2} \\ &\lesssim \bar{w}^{b} |r X_{r}^{b-1} \partial^{\beta} \nabla \cdot \mathbf{G}|^{2} + \bar{w}^{b} |r X_{r}^{b-1} \partial^{\beta} \nabla \times \mathbf{G}|^{2} + V_{b+|\beta|-1} + V_{b+|\beta|,b-1} \\ &\lesssim \bar{w}^{b} |r X_{r}^{b-1} \partial^{\beta} ((I - \mathcal{A}) \nabla \cdot \mathbf{G})|^{2} + \bar{w}^{b} |r X_{r}^{b-1} \partial^{\beta} ((I - \mathcal{A}) \nabla \times \mathbf{G})|^{2} \\ &+ E_{n} + V_{b+|\beta|-1} + V_{b+|\beta|,b-1} \\ &\lesssim \bar{w}^{b} |r (I - \mathcal{A}) X_{r}^{b-1} \partial^{\beta} \nabla \cdot \mathbf{G}|^{2} + \bar{w}^{b} |r (I - \mathcal{A}) X_{r}^{b-1} \partial^{\beta} \nabla \times \mathbf{G}|^{2} \\ &+ E_{n} + V_{b+|\beta|-1} + V_{b+|\beta|,b-1} \\ &\lesssim \bar{w}^{b} E_{n} |r X_{r}^{b-1} \partial^{\beta} \nabla \mathbf{G}|^{2} + E_{n} + V_{b+|\beta|-1} + V_{b+|\beta|,b-1} \end{split}$$

So

$$V_{b+|\beta|,b} \lesssim E_n V_{b+|\beta|,b} + E_n + V_{b+|\beta|-1} + V_{b+|\beta|,b-1}$$

By a priori assumption (4.164), we have $E_n \ll 1$, so

$$V_{b+|\beta|,b} \lesssim E_n + V_{b+|\beta|-1} + V_{b+|\beta|,b-1}.$$

We know $V_{n',0} \lesssim E_n$ for all $n' \leq n/2$, so by induction we get $V_{n',d} \lesssim E_n$ for all $d \leq n' \leq n/2$.

Now for $|\beta| + b \le n$, using the above lemmas 2.6 and 2.7 and results for V we have

$$\begin{split} \|X_r^b \partial^{\beta} \mathbf{G}\|_{3+b}^2 &\lesssim \|r \nabla \cdot X_r^{b-1} \partial^{\beta} \mathbf{G}\|_{3+b}^2 + \|r \nabla \times X_r^{b-1} \partial^{\beta} \mathbf{G}\|_{3+b}^2 + \sum_{k=1}^3 \|X_r^{b-1} \partial_k \partial^{\beta} \mathbf{G}\|_{3+b}^2 \\ &\lesssim \|r X_r^{b-1} \partial^{\beta} \nabla \cdot \mathbf{G}\|_{3+b}^2 + \|r X_r^{b-1} \partial^{\beta} \nabla \times \mathbf{G}\|_{3+b}^2 + W_{b+|\beta|-1} + W_{b+|\beta|,b-1} \\ &\lesssim \|r X_r^{b-1} \partial^{\beta} ((1-\mathcal{A}) \nabla \cdot \mathbf{G})\|_{3+b}^2 + \|r X_r^{b-1} \partial^{\beta} ((1-\mathcal{A}) \nabla \times \mathbf{G})\|_{3+b}^2 \end{split}$$

$$+ E_n + W_{b+|\beta|-1} + W_{b+|\beta|,b-1} \lesssim \|r(1 - \mathscr{A})X_r^{b-1}\partial^{\beta}\nabla \cdot \mathbf{G}\|_{3+b}^2 + \|r(1 - \mathscr{A})X_r^{b-1}\partial^{\beta}\nabla \times \mathbf{G}\|_{3+b}^2 + E_n + W_{b+|\beta|-1} + W_{b+|\beta|,b-1} \lesssim E_n \|rX_r^{b-1}\partial^{\beta}\nabla \mathbf{G}\|_{3+b}^2 + E_n + W_{b+|\beta|-1} + W_{b+|\beta|,b-1}$$

So

$$W_{b+|\beta|,b} \lesssim E_n W_{b+|\beta|,b} + E_n + W_{b+|\beta|-1} + W_{b+|\beta|,b-1}$$

By a priori assumption (4.164), we have $E_n \ll 1$, so

$$W_{b+|\beta|,b} \lesssim E_n + W_{b+|\beta|-1} + W_{b+|\beta|,b-1}.$$

We know $W_{n,0} \lesssim E_n$, so by induction we get $W_{n,d} \lesssim E_n$ for all $d \le n$. Let

$$Y_n = \sum_{c \le n} \|\nabla^c \mathbf{G}\|_{3+2c}^2$$

By Sobolev embeddings like those used to prove the embedding theorems A.10 and A.11, we have that

$$\|\bar{w}^{c/2}\nabla^{c}\mathbf{G}\|_{L^{\infty}(\mathbb{R}^{3})}^{2} \lesssim E_{n} + Y_{n}$$
 when $c \leq n/2$.

Now for $c \leq n$, using the above lemmas 2.3 and 2.7 we have

$$\begin{aligned} \|\nabla^{c}\mathbf{G}\|_{3+2c}^{2} &\lesssim \|\nabla^{c-1}\nabla\cdot\mathbf{G}\|_{3+2c}^{2} + \|\nabla^{c-1}\nabla\times\mathbf{G}\|_{3+2c}^{2} + \|\nabla^{c-1}\mathbf{G}\|_{3+2(c-1)}^{2} \\ &\lesssim \|\nabla^{c-1}((1-\mathscr{A})\nabla\cdot\mathbf{G})\|_{3+2c}^{2} + \|\nabla^{c-1}((1-\mathscr{A})\nabla\times\mathbf{G})\|_{3+2c}^{2} + E_{n} + Y_{c-1} \\ &\lesssim \|(1-\mathscr{A})\nabla^{c-1}\nabla\cdot\mathbf{G}\|_{3+2c}^{2} + \|(1-\mathscr{A})\nabla^{c-1}\nabla\times\mathbf{G}\|_{3+2c}^{2} + E_{n}(1+Y_{n}) + Y_{c-1} \\ &\lesssim E_{n}\|\nabla^{c-1}\nabla\mathbf{G}\|_{3+2c}^{2} + E_{n}(1+Y_{n}) + Y_{c-1} \end{aligned}$$

So

$$Y_c \lesssim E_n Y_n + E_n + Y_{c-1}$$

By induction on $c \leq n$ we have

 $Y_n \lesssim E_n Y_n + E_n$

By a priori assumption (4.164), we have $E_n \ll 1$, so we get $Y_n \lesssim E_n$.

4.4 Vorticity estimates

In this section we will estimate the vorticity Z_n , $\|\nabla \times X_r^b \partial^\beta \theta\|_{4+b}$ and $\|\nabla \times \partial^\gamma \theta\|_{4+2|\gamma|}$ which will be needed to control the "curl" part of the pressure term as seem in Lemma 2.12.

By taking curl to the Euler-Poisson equation (4.158) we can essentially get rid of the pressure and gravity terms, which allows us to estimate the curl separately.

Lemma 4.10. Let θ be a solution of (4.158) in the sense of Theorem 4.4. Then for any $s \ge s_0 \ge 0$ we have

$$\begin{split} \lambda(s)^{\frac{1}{2}}(\mathscr{A}\nabla\times\partial_{s}\boldsymbol{\theta})(s) &= \frac{\lambda(s_{0})}{\lambda(s)^{1/2}}(\mathscr{A}\nabla\times\partial_{s}\boldsymbol{\theta})(s_{0}) + \lambda(s)^{-\frac{1}{2}}\int_{s_{0}}^{s}(\partial_{s}\mathscr{A})\nabla\times\partial_{s}\boldsymbol{\theta}\mathrm{d}s'\\ (\mathscr{A}\nabla\times\boldsymbol{\theta})(s) &= (\mathscr{A}\nabla\times\boldsymbol{\theta})(s_{0}) + \int_{s_{0}}^{s}(\partial_{s}\mathscr{A})\nabla\times\boldsymbol{\theta}\mathrm{d}s' + \int_{s_{0}}^{s}\frac{\lambda(s_{0})}{\lambda(s)}\mathrm{d}s'(\mathscr{A}\nabla\times\partial_{s}\boldsymbol{\theta})(s_{0})\\ &+ \int_{s_{0}}^{s}\lambda(s')^{-1}\int_{s_{0}}^{s'}(\partial_{s}\mathscr{A})\nabla\times\partial_{s}\boldsymbol{\theta}\mathrm{d}s''\mathrm{d}s'. \end{split}$$

Proof. Recall (4.156) is

$$\mathbf{0} = \lambda \partial_s^2 \boldsymbol{\xi} + \lambda' \partial_s \boldsymbol{\xi} + \delta \boldsymbol{\xi} + \frac{1}{w^3} \partial_k (\mathcal{A}^k w^4 \mathcal{J}^{-1/3}) + \mathcal{A} \nabla \Phi$$
(4.167)

Now note that

$$\frac{1}{w^3}\partial_k(\mathscr{A}^k w^4 \mathscr{F}^{-1/3}) = \bar{w}^{-3}\partial_k(\bar{w}^4 a^k \mathscr{F}^{-4/3}) = \bar{w}^{-3} a^k \partial_k(\bar{w}^4 \mathscr{F}^{-4/3}) = (\bar{w}\mathscr{F}^{-1/3})^{-1} \mathscr{A}^k \partial_k(\bar{w}^4 \mathscr{F}^{-4/3}) = \frac{4}{3} \mathscr{A} \nabla(\bar{w}^3 \mathscr{F}^{-1})$$

and $\mathscr{A}\nabla \times \boldsymbol{\xi} = \epsilon_{\bullet jk} \mathscr{A}_{j}^{l} \partial_{l} \xi_{k} = \epsilon_{\bullet jk} \delta_{jk} = \mathbf{0}$. So taking $\mathscr{A}\nabla \times$ to (4.156) we get

$$\begin{aligned} \mathbf{0} &= \lambda \mathscr{A} \nabla \times \partial_s^2 \boldsymbol{\theta} + \lambda' \mathscr{A} \nabla \times \partial_s \boldsymbol{\theta} = \lambda \left(\partial_s (\mathscr{A} \nabla \times \partial_s \boldsymbol{\theta}) - (\partial_s \mathscr{A}) \nabla \times \partial_s \boldsymbol{\theta} \right) + \lambda' \mathscr{A} \nabla \times \partial_s \boldsymbol{\theta} \\ &= \partial_s (\lambda \mathscr{A} \nabla \times \partial_s \boldsymbol{\theta}) - (\partial_s \mathscr{A}) \nabla \times \partial_s \boldsymbol{\theta} \end{aligned}$$

So

$$\lambda(s)(\mathscr{A}\nabla\times\partial_s\boldsymbol{\theta})(s) = \lambda(s_0)(\mathscr{A}\nabla\times\partial_s\boldsymbol{\theta})(s_0) + \int_{s_0}^s(\partial_s\mathscr{A})\nabla\times\partial_s\boldsymbol{\theta}\mathrm{d}s'$$

So

$$\lambda \partial_s (\mathscr{A} \nabla \times \boldsymbol{\theta})(s) = \lambda (\partial_s \mathscr{A}) \nabla \times \boldsymbol{\theta}(s) + \lambda (s_0) (\mathscr{A} \nabla \times \partial_s \boldsymbol{\theta})(s_0) + \int_{s_0}^s (\partial_s \mathscr{A}) \nabla \times \partial_s \boldsymbol{\theta} \mathrm{d}s'$$

So

$$\begin{split} (\mathscr{A}\nabla\times\boldsymbol{\theta})(s) &= (\mathscr{A}\nabla\times\boldsymbol{\theta})(s_0) + \int_{s_0}^s (\partial_s\mathscr{A})\nabla\times\boldsymbol{\theta} \, \mathrm{d}s' + \int_{s_0}^s \frac{\lambda(s_0)}{\lambda(s)} \mathrm{d}s'(\mathscr{A}\nabla\times\partial_s\boldsymbol{\theta})(s_0) \\ &+ \int_{s_0}^s \lambda(s')^{-1} \int_{s_0}^{s'} (\partial_s\mathscr{A})\nabla\times\partial_s\boldsymbol{\theta} \mathrm{d}s'' \mathrm{d}s'. \end{split}$$

Proposition 4.11. Let θ be a solution of (4.158) in the sense of Theorem 4.4. Let $n \ge 21$. Then for any $s \ge s_0 \ge 0$, we have

$$Z_n(s) \lesssim Z_n(s_0) + \frac{(s-s_0)^2}{\lambda(s)} \mathscr{C}_n(s) + \lambda(s)^{-1} \mathscr{C}_n(s)^2.$$

Proof. Take $X_r^b \partial^{\beta} (b + \beta = n)$ to the first equation in Lemma 4.10 we get

$$\lambda(s)^{\frac{1}{2}}X_{r}^{b}\partial^{\beta}(\mathscr{A}\nabla\times\partial_{s}\boldsymbol{\theta})(s) = \frac{\lambda(s_{0})}{\lambda(s)^{1/2}}X_{r}^{b}\partial^{\beta}(\mathscr{A}\nabla\times\partial_{s}\boldsymbol{\theta})(s_{0}) + \lambda(s)^{-\frac{1}{2}}X_{r}^{b}\partial^{\beta}\int_{s_{0}}^{s}(\partial_{s}\mathscr{A})\nabla\times\partial_{s}\boldsymbol{\theta}\mathrm{d}s'$$

Since λ is an increasing function (4.154), we have $\lambda(s_0)^{1/2}/\lambda(s)^{1/2} \leq q$ which we can used to bound the first term on the RHS. And for the other terms we can estimate for example

$$\begin{split} &\int_{B_R} \left| \lambda(s)^{-\frac{1}{2}} \int_{s_0}^s (X_r^b \partial^\beta \partial_s \mathscr{A}) \nabla \times \partial_s \boldsymbol{\theta} \, \mathrm{d}s' \right|^2 \bar{w}^{4+b} \mathrm{d}\mathbf{x} \\ &\leq \lambda(s)^{-1} \int_{B_R} \left| \int_{s_0}^s (X_r^b \partial^\beta \partial_s \mathscr{A}) \nabla \times \partial_s \boldsymbol{\theta} \, \mathrm{d}s' \right|^2 \bar{w}^{4+b} \mathrm{d}\mathbf{x} \\ &\lesssim \lambda(s)^{-1} \int_{B_R} \left| \int_{s_0}^s (X_r^b \partial^\beta \mathscr{A}) \nabla \times \partial_s^2 \boldsymbol{\theta} \, \mathrm{d}s' \right|^2 \bar{w}^{4+b} \mathrm{d}\mathbf{x} + \lambda(s)^{-1} \mathscr{C}_n(s)^2 \\ &\lesssim \frac{(s-s_0)^2}{\lambda(s)} \mathscr{C}_n(s) + \lambda(s)^{-1} \mathscr{C}_n(s)^2 \end{split}$$

where we used $\lambda \partial_s^2 \theta = -\lambda' \partial_s \theta - \delta \theta - \mathbf{P} - \mathbf{G}$ (4.158).

We can then repeat this process for $\partial^{\gamma} (|\gamma| = n)$ in place of $X_r^b \partial^{\beta}$ to complete the proof.

Proposition 4.12. Let θ be a solution of (4.158) in the sense of Theorem 4.4. Let $n \ge 21$. Then for any $s \ge s_0 \ge 0$ and $n' \le n$, we have

$$\sum_{b+\beta \le n'} \|\nabla \times X_r^b \partial^\beta \boldsymbol{\theta}(s)\|_{4+b}^2 \lesssim E_n(s_0) + S_{n'-1}(s) + \mathcal{C}_n(s)^2 + \int_{s_0}^s \frac{1 + (s' - s_0)^2}{\lambda(s')} \mathrm{d}s' \, \mathcal{C}_n(s)$$
$$\sum_{|\gamma| \le n} \|\nabla \times \partial^\gamma \boldsymbol{\theta}(s)\|_{4+2|\gamma|}^2 \lesssim E_n(s_0) + \mathcal{C}_n(s)^2 + \int_{s_0}^s \frac{1 + (s' - s_0)^2}{\lambda(s')} \mathrm{d}s' \, \mathcal{C}_n(s).$$

Proof. Take $X_r^b \partial^\beta (b + \beta = n)$ to the second equation in Lemma 4.10 and then estimate in a similar way to Lemma 4.11. We estimate

$$\begin{split} \left\| \int_{s_0}^s \lambda(s')^{-1} \int_{s_0}^{s'} \star \mathrm{d}s'' \mathrm{d}s' \right\|_{4+b}^2 &\leq \int_{B_R} \left(\int_{s_0}^s \lambda(s')^{-1} \mathrm{d}s' \right) \left(\int_{s_0}^s \lambda(s')^{-1} \left| \int_{s_0}^{s'} \star \mathrm{d}s'' \right|^2 \mathrm{d}s' \right) \bar{w}^{4+b} \mathrm{d}\mathbf{x} \\ &\lesssim \int_{s_0}^s \lambda(s')^{-1} \int_{B_R} \left| \int_{s_0}^{s'} \star \mathrm{d}s'' \right|^2 \bar{w}^{4+b} \mathrm{d}\mathbf{x} \mathrm{d}s' \\ &\lesssim \int_{s_0}^s \frac{1 + (s' - s_0)^2}{\lambda(s')} \mathrm{d}s' \, \mathscr{C}_n(s) \end{split}$$

where we used the Cauchy–Schwarz inequality to get the first inequality, and the fact that $\int_0^\infty \lambda^{-1} ds' < \infty$ for the second inequality (see (4.154)). And

$$\sum_{\substack{b+\beta \leq n \\ b+\beta \leq n}} \| [\nabla \times, X_r^b \partial^\beta] \boldsymbol{\theta}(s) \|_{4+b}^2 \lesssim S_{n'-1}(s)$$
$$\sum_{\substack{b+\beta \leq n \\ b+\beta \leq n}} \| X_r^b \partial^\beta ((\mathcal{A} - I) \nabla \times \boldsymbol{\theta})(s) \|_{4+b}^2 \lesssim \mathscr{E}_n(s)^2.$$

Then we get the first formula. Proof for the second formula is similar.

4.5 Energy estimates and proof of the main theorem

In this section we finally commute the momentum equation (4.158) and then derive the high-order energy estimates. Then finally we will prove our main theorem using the energy estimates.

Theorem 4.13 (Energy estimates). Let $n \ge 21$. Let θ be a solution of (4.158) in the sense of Theorem 4.4, given on its maximal interval of existence. Then

$$\mathscr{C}_n(s) \lesssim \mathscr{C}_n(s_0) + \mathscr{C}_n(s)^{3/2} + \mathscr{C}_n(s) \int_{s_0}^s \lambda^{-1/2} \mathrm{d}\tau.$$
(4.168)

for any $s \ge s' \ge 0$ whenever our a priori assumption (4.164) is satisfied. Here we recall Definition (4.163) of the total norm \mathcal{C}_n .

Proof. Since $\mathscr{C}_n = \mathscr{S}_n + \mathscr{Q}_n + \mathscr{X}_n$, we need to prove the formula with LHS each of these three component terms.

We first deal with the S_n part. Let $|\beta| + b \le n$. Apply $X_r^b \partial^\beta$ to the momentum equation (4.158) to get

$$\lambda \partial_s^2 X_r^b \partial^\beta \theta + \lambda' \partial_s X_r^b \partial^\beta \theta + X_r^b \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}) = 0$$

Taking the $\langle \cdot, \cdot \rangle_{3+b}$ -inner with $\partial_s X_r^b \partial^\beta \theta$ we get

$$0 = \frac{1}{2}\lambda\partial_{s}\|\partial_{s}X_{r}^{b}\partial^{\beta}\theta\|_{3+b}^{2} + \lambda'\|\partial_{s}X_{r}^{b}\partial^{\beta}\theta\|_{3+b}^{2} + \langle X_{r}^{b}\partial^{\beta}(\delta\theta + \mathbf{P} + \mathbf{G}), \partial_{s}X_{r}^{b}\partial^{\beta}\theta\rangle_{3+b}$$
$$= \frac{1}{2}\partial_{s}\left(\lambda\|\partial_{s}X_{r}^{b}\partial^{\beta}\theta\|_{3+b}^{2}\right) + \frac{1}{2}\lambda'\|\partial_{s}X_{r}^{b}\partial^{\beta}\theta\|_{3+b}^{2} + \langle X_{r}^{b}\partial^{\beta}(\delta\theta + \mathbf{P} + \mathbf{G}), \partial_{s}X_{r}^{b}\partial^{\beta}\theta\rangle_{3+b}$$

Integrate in time we get

$$0 = \frac{1}{2} \left(\lambda \|\partial_s X^b_r \partial^\beta \theta\|^2_{3+b} \right) \Big|_{s_0}^s + \frac{1}{2} \int_{s_0}^s \lambda' \|\partial_s X^b_r \partial^\beta \theta\|^2_{3+b} \mathrm{d}\tau + \int_{s_0}^s \langle X^b_r \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_s X^b_r \partial^\beta \theta \rangle_{3+b} \mathrm{d}\tau + \int_{s_0}^s \langle X^b_r \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_s X^b_r \partial^\beta \theta \rangle_{3+b} \mathrm{d}\tau + \int_{s_0}^s \langle X^b_r \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_s X^b_r \partial^\beta \theta \rangle_{3+b} \mathrm{d}\tau + \int_{s_0}^s \langle X^b_r \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_s X^b_r \partial^\beta \theta \rangle_{3+b} \mathrm{d}\tau + \int_{s_0}^s \langle X^b_r \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_s X^b_r \partial^\beta \theta \rangle_{3+b} \mathrm{d}\tau + \int_{s_0}^s \langle X^b_r \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_s X^b_r \partial^\beta \theta \rangle_{3+b} \mathrm{d}\tau + \int_{s_0}^s \langle X^b_r \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_s X^b_r \partial^\beta \theta \rangle_{3+b} \mathrm{d}\tau + \int_{s_0}^s \langle X^b_r \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_s X^b_r \partial^\beta \theta \rangle_{3+b} \mathrm{d}\tau + \int_{s_0}^s \langle X^b_r \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_s X^b_r \partial^\beta \theta \rangle_{3+b} \mathrm{d}\tau + \int_{s_0}^s \langle X^b_r \partial^\beta (\delta \theta + \mathbf{P} + \mathbf{G}), \partial_s X^b_r \partial^\beta \theta \rangle_{3+b} \mathrm{d}\tau$$

By Proposition 4.6, 4.5, 4.9 and Lemma 2.12 we get

$$\begin{split} &\frac{1}{2} \bigg(\lambda \|\partial_s X_r^b \partial^\beta \boldsymbol{\theta}\|_{3+b}^2 + \|\nabla X_r^b \partial^\beta \boldsymbol{\theta}\|_{4+b}^2 + \frac{1}{3} \|\operatorname{div} X_r^b \partial^\beta \boldsymbol{\theta}\|_{4+b}^2 - \frac{1}{2} \|\operatorname{curl} X_r^b \partial^\beta \boldsymbol{\theta}\|_{4+b}^2 \bigg) \Big|_{s_0}^s \\ &+ \frac{1}{2} \int_{s_0}^s \lambda' \|\partial_s X_r^b \partial^\beta \boldsymbol{\theta}\|_3^2 \mathrm{d}\tau \\ &\lesssim \mathfrak{E}_n(s)^{3/2} + \int_{s_0}^s \lambda^{-1/2} E_n \mathrm{d}\tau \end{split}$$

Using Proposition 4.12 we get

$$\begin{split} \lambda(s) \|\partial_s X_r^b \partial^\beta \theta(s)\|_{3+b}^2 + \|\nabla X_r^b \partial^\beta \theta(s)\|_{4+b}^2 \\ \lesssim E_n(s_0) + E_{n-1}(s) + \mathfrak{E}_n(s)^2 + \int_{s_0}^s \frac{1 + (s' - s_0)^2}{\lambda(s')} \mathrm{d}\tau \ \mathfrak{E}_n(s) + \mathfrak{E}_n(s)^{3/2} + \int_{s_0}^s \lambda^{-1/2} E_n \mathrm{d}\tau \\ \lesssim E_n(s_0) + E_{n-1}(s) + \mathfrak{E}_n(s)^{3/2} + \mathfrak{E}_n(s) \int_{s_0}^s \lambda^{-1/2} \mathrm{d}\tau \end{split}$$

where we used (4.154) and (4.164). Add to it

$$\begin{split} \|X_r^b \partial^{\beta} \boldsymbol{\theta}(s)\|_{3+b}^2 &= \|X_r^b \partial^{\beta} \boldsymbol{\theta}(s_0)\|_{3+b}^2 + 2\int_{s_0}^s \langle X_r^b \partial^{\beta} \boldsymbol{\theta}(s), \partial_s X_r^b \partial^{\beta} \boldsymbol{\theta}(s) \rangle_{3+b} \mathrm{d}\tau \\ &\lesssim E_n(s_0) + \int_{s_0}^s \lambda^{-1/2} E_n \mathrm{d}\tau, \end{split}$$

sum over $|\beta| + b \le n' \le n$ and we get

$$S_{n'}(s) \lesssim E_n(s_0) + S_{n'-1}(s) + \mathcal{C}_n(s)^{3/2} + \mathcal{C}_n(s) \int_{s_0}^s \lambda^{-1/2} \mathrm{d}\tau.$$

Induct on n' we get

$$S_n(s) \lesssim E_n(s_0) + \mathscr{C}_n(s)^{3/2} + \mathscr{C}_n(s) \int_{s_0}^s \lambda^{-1/2} \mathrm{d}\tau$$

To prove the \mathbb{Q}_n part, we repeat the above with ∂^{γ} in place of $X_r^b \partial^{\beta}$, and weight $3 + 2|\gamma|$ instead of weight 3 + b.

Finally the \mathfrak{X}_n part is given by Proposition 4.11 noting that

$$\lambda(s)^{-1} \mathfrak{C}_n(s)^2 + \frac{(s-s_0)^2}{\lambda(s)} \mathfrak{C}_n(s) \lesssim \mathfrak{C}_n(s)^{3/2} + \mathfrak{C}_n(s) \int_{s_0}^s \lambda^{-1/2} \mathrm{d}\tau.$$

where we used (4.154) and (4.164).

To proof our main theorem that the energy E_n remains bounded, we will use the bootstrapping scheme in the following lemma and proposition.

Lemma 4.14. Suppose $E : [0,T] \rightarrow [0,\infty]$ is continuous and

$$E(t) \le C_1 E(0) + C_2 E(t)^{3/2}$$
 whenever $\sup_{\tau \in [0,t]} E(\tau) \le C_3$

where $C_1 \ge 1$. Then $E \le 2C_1 E(0)$ whenever $E(0) \le \min\{(2^5 C_1 C_2^2)^{-1}, C_3/2C_1\}$.

Proof. Same as Lemma 3.51.

Proposition 4.15. Suppose $E: [0, \infty) \to [0, \infty]$ are continuous and for all $s \ge s_0 \ge 0$ we have

$$E(s) \le C_1 E(s_0) + C_2 E(s)^{3/2} + C_3 F(s_0, s) E(s) \quad \text{whenever} \quad \sup_{\tau \in [s_0, s]} E(\tau) \le C_4$$

where $F : \{(s_0, s) \in [0, \infty) \times [0, \infty) : s \ge s_0\} \to [0, \infty)$ is a function such that

i. $\lim_{s_0\to\infty}\sup_{s\ge s_0}F(s_0,s)=0;$

ii. $\lim_{\delta' \to 0} \sup_{|s-s_0| \le \delta'} F(s_0, s) = 0.$

Then there exist $\epsilon^* > 0$ such that $E \leq_{C_1} E(0)$ whenever $E(0) \leq \epsilon^*$.

Proof. Pick s_{∞} large enough so that

$$C_3 \sup_{s \ge s_\infty} F(s_\infty, s) < \frac{1}{2}.$$

Then by Lemma 4.14 there exist $\epsilon_{\infty} > 0$ such that $\sup_{s \in [s_{\infty},\infty)} \leq 4C_1 E(s_{\infty})$ whenever $E(s_{\infty}) \leq \epsilon_{\infty}$.

Now pick δ' small enough so that

$$C_3 \sup_{|s-s_0| \le \delta'} F(s_0, s) < \frac{1}{2}.$$

Then by Lemma 4.14 there exist $\epsilon_0 > 0$ such that $\sup_{s \in [m\delta', (m+1)\delta']} \leq 4C_1 E(m\delta')$ whenever $E(m\delta') \leq \epsilon_0$.

Let $\epsilon^* \leq \min\{\epsilon_0, \epsilon_\infty\}/(4C_1)^{\lceil s_\infty/\delta'\rceil}$. Then $E(s) \leq (4C_1)^{\lceil s_\infty/\delta'\rceil+1}E(0)$ for all $s \geq 0$ whenever $E(0) \leq \epsilon^*$.

Theorem 4.16. Let $n \ge 21$. Let $(\theta, \partial_s \theta)$ be a solution of (4.158) in the sense of Theorem 4.4. Then there exists $\epsilon^* > 0$ such that if $E_n(0) \le \epsilon^*$, then we have $\mathfrak{C}_n \lesssim \mathfrak{C}_n(0)$.

Proof. By the energy estimates in Theorem 4.13 we have

$$\mathscr{C}_n(s) \lesssim \mathscr{C}_n(s_0) + \mathscr{C}_n(s)^{3/2} + \mathscr{C}_n(s) \int_{s_0}^s \lambda^{-1/2} \mathrm{d}\tau.$$
(4.169)

Applying Proposition 4.15 above with $E = \mathscr{C}_n$ and $F(s_0, s) = \int_{s_0}^s \lambda^{-1/2} d\tau$ (which satisfies the properties required for the proposition because of (4.154)) we get the desired result.

A Appendix

A.1 Differentiation and commutation properties

Here we first collect some standard results on how derivatives interact with \mathcal{J} and \mathcal{A} , which can be found for example in [31]. After that we state various derivative commutators frequently used in the article. **Lemma A.1.** *Recall notations defined in Definition 2.2. We have*

$$\begin{split} \mathcal{A}_{j}^{i} - I_{j}^{i} &= -\mathcal{A}_{k}^{i}\partial_{j}\theta^{k} \\ \partial_{\bullet}\mathcal{J} &= \mathcal{J}\mathcal{A}_{m}^{l}\partial_{\bullet}\partial_{l}\theta^{m} \\ \partial_{\bullet}\mathcal{A}_{j}^{i} &= -\mathcal{A}_{m}^{i}\mathcal{A}_{j}^{l}\partial_{\bullet}\partial_{l}\theta^{m} \end{split}$$

Proof. Since $\mathcal{A} = (\nabla \boldsymbol{\xi})^{-1}$, we have

$$I_j^i = \mathscr{A}_k^i \partial_j \xi^k = \mathscr{A}_k^i (I_j^k + \partial_j \theta^k)$$

It can be proven that if $U: t \mapsto U(t)$ is a differentiable map of invertible square matrices, then

i.
$$\frac{\mathrm{d}\det U}{\mathrm{d}t} = \det(U)\tilde{\rho}\left(U^{-1}\frac{\mathrm{d}U}{\mathrm{d}t}\right);$$

ii.
$$\frac{\mathrm{d}U^{-1}}{\mathrm{d}t} = -U^{-1}\frac{\mathrm{d}A}{\mathrm{d}t}U^{-1}.$$

Using i. we get $\partial_{\bullet}J = JA_i^k \partial_{\bullet}\partial_k \eta^i$, and using ii. we get $\partial_{\bullet}A_j^i = -A_k^i (\partial_{\bullet}\partial_l \eta^k)A_j^l$. Converting to \mathcal{A} and \mathcal{F} by tracing the definition and keeping track of the factors of λ , we get the stated formulas.

We commonly use various commutation properties between the Cartesian, radial, angular derivatives, and their Lagrangian counterparts.

Lemma A.2 (Commutation relations). We have the following commutation relations

$$\begin{split} [X_r, \nabla] &= -\nabla \\ [X_r, \mathbf{x}] &= \mathbf{x} \\ [\partial_i, \partial_j] &= -\epsilon_{ijk}\partial_k \\ [\partial_i, x^j] &= -\epsilon_{ijk}x^k \\ [X_r, \mathcal{K}] &= 2\mathcal{K} \\ [\partial_i, \mathcal{K}] &= 0 \\ [\partial_s, \mathcal{A}\partial_j] &= -(\mathcal{A}\partial_j\partial_s\theta^m)\mathcal{A}\partial_m \\ [\nabla, \mathcal{A}\partial_j] &= -(\mathcal{A}\partial_j\nabla\theta^m)\mathcal{A}\partial_m \\ [X_r, \mathcal{A}\partial_j] &= -(\mathcal{A}_j^l X_r \partial_l \theta^m)\mathcal{A}\partial_m - \mathcal{A}\partial_j \\ [\partial_i, \mathcal{A}\partial_j] &= -(\mathcal{A}_j^l \partial_i \partial_l \theta^m)\mathcal{A}\partial_m - \epsilon_{ikl}\mathcal{A}_j^k \partial_l \mathcal{A}_k \\ [X_r, \partial_i] &= 0 \\ [\mathcal{M}_i, \partial_i'] &= -\partial_{ii'}. \end{split}$$

Proof. We have

$$\begin{split} & [X_r, \partial_j] = x^i \partial_i \partial_j - \partial_j x^i \partial_i = -\partial_j \\ & [X_r, x^j] = x^i \partial_i x^j - x^j x^i \partial_i = x^j \\ & [\vartheta_i, \partial_j] = \epsilon_{ilk} x^l \partial_k \partial_j - \epsilon_{ilk} \partial_j x^l \partial_k = -\epsilon_{ijk} \partial_k \\ & [\vartheta_i, x^j] = \epsilon_{ilk} x^l \partial_k x^j - \epsilon_{ilk} x^j x^l \partial_k = \epsilon_{ilj} x^l \\ & [\partial_s, \mathcal{A}\partial_j] = \partial_s \mathcal{A}_j^l \partial_i - \mathcal{A}_j^j \partial_i \partial_s = -\mathcal{A}_m^l \mathcal{A}_j^l (\partial_l \partial_s \theta^m) \partial_i \\ & [\partial_i, \mathcal{A}\partial_j] = \partial_i \mathcal{A}_j^k \partial_k - \mathcal{A}_j^k \partial_k \partial_i = -\mathcal{A}_m^k \mathcal{A}_j^l (\partial_l \partial_i \theta^m) \partial_k \\ & [X_r, \mathcal{A}\partial_j] = X_r (\mathcal{A}_j^k \partial_k) - \mathcal{A}_j^k \partial_k X_r = -(\mathcal{A}_m^k \mathcal{A}_j^l X_r \partial_l \theta^m) \partial_k + \mathcal{A}_j^k X_r \partial_k - \mathcal{A}_j^k \partial_k X_r \\ & = -(\mathcal{A}_m^k \mathcal{A}_j^l X_r \partial_l \theta^m) \partial_k - \mathcal{A}_j^k \partial_k \\ & [\vartheta_i, \mathcal{A}\partial_j] = \vartheta_i (\mathcal{A}_j^k \partial_k) - \mathcal{A}_j^k \partial_k \vartheta_i = -(\mathcal{A}_m^k \mathcal{A}_j^l \partial_i \partial_l \theta^m) \partial_k + \mathcal{A}_j^k \partial_i \partial_k - \mathcal{A}_j^k \partial_k \partial_i \\ & = -(\mathcal{A}_j^l \partial_i \partial_l \theta^m) \mathcal{A}_m - \epsilon_{ikl} \mathcal{A}_j^k \partial_l \\ & = -(\mathcal{A}_j^l \partial_i \partial_l \theta^m) \mathcal{A}_m - \epsilon_{ikl} \mathcal{A}_j^k \partial_l \\ & = -(\mathcal{A}_j^l \partial_i \partial_k - \mathcal{A}_j^l \partial_k) - \mathcal{A}_j^i \partial_k (x^l \partial_l) = \epsilon_{ijk} \left(\delta_l^j x^l \partial_k + x^l x^j \partial_l \partial_k - \delta_k^l x^j \partial_l - x^j x^l \partial_k \partial_l \right) \\ & = \epsilon_{ijk} \left(x^j \partial_k - x^j \partial_k \right) = 0 \\ & [\vartheta_i, \vartheta_{i'}] = \epsilon_{ijk} \epsilon_{i'j'k'} \left(x^j \partial_k (x^{j'} \partial_{k'}) - x^{j'} \partial_{k'} (x^j \partial_k) \right) = \epsilon_{ijk} \epsilon_{i'j'k'} \left(\delta_k^{j'} x^j \partial_{k'} - \delta_{k'}^j x^{j'} \partial_k \right) \\ & = \epsilon_{ijk} \epsilon_{i'k'k'} x^j \partial_{k'} - \epsilon_{ik'k} \epsilon_{i'j'k'} x^{j'} \partial_k = \epsilon_{kij} \epsilon_{kk'i'} x^{j} \partial_{k'} - \epsilon_{k'k} \epsilon_{k'i'j'} x^{j'} \partial_k \\ & = \left(\delta_{ik'} \delta_{ji'} - \delta_{ii'} \delta_{jk'} \right) \mathcal{A}_k - \left(\frac{g(\mathbf{x}) \nabla \cdot \mathbf{x}}{|\mathbf{y} - \mathbf{x}|} + g(\mathbf{x}) \mathbf{x} \cdot \nabla_{\mathbf{x}} \frac{1}{|\mathbf{y} - \mathbf{x}|} \right) \mathbf{dx} \\ & = \int \left(\frac{3g(\mathbf{x})}{|\mathbf{y} - \mathbf{x}|} + g(\mathbf{x}) \mathbf{x} \cdot \frac{\mathbf{y} - \mathbf{x}}{|\mathbf{y} - \mathbf{x}|^3} \right) \mathbf{dx} = \int \left(\frac{2g(\mathbf{x})}{|\mathbf{y} - \mathbf{x}|^3} \right) \mathbf{dx} \end{aligned}$$

$$\begin{split} &= -2\mathscr{K}g + \mathbf{y} \cdot \nabla \mathscr{K}g \\ \mathscr{K}(\mathbf{\partial}_{i}g)(\mathbf{y}) = -\int \frac{\epsilon_{ijk}x^{j}\partial_{k}g(\mathbf{x})}{|\mathbf{y} - \mathbf{x}|} \mathrm{d}\mathbf{x} = \int g(\mathbf{x})\epsilon_{ijk}x^{j}\frac{y^{k} - x^{k}}{|\mathbf{y} - \mathbf{x}|^{3}} \mathrm{d}\mathbf{x} \\ &= \int g(\mathbf{x})\epsilon_{ijk}x^{j}\frac{y^{k}}{|\mathbf{y} - \mathbf{x}|^{3}} \mathrm{d}\mathbf{x} = \int g(\mathbf{x})\epsilon_{ijk}y^{j}\frac{-x^{k}}{|\mathbf{y} - \mathbf{x}|^{3}} \mathrm{d}\mathbf{x} \\ &= \int g(\mathbf{x})\epsilon_{ijk}y^{j}\frac{y^{k} - x^{k}}{|\mathbf{y} - \mathbf{x}|^{3}} \mathrm{d}\mathbf{x} = (\mathbf{\partial}_{i}\mathscr{K}g)(\mathbf{y}). \end{split}$$

A.2 Spherical harmonics

Spherical harmonics has a real as well as complex version. For the definition and basic properties of the complex version, see [28]. The relation between complex spherical harmonics $Y_l^m : S^2 \to \mathbb{C}$ and real spherical harmonics $Y_{lm} : S^2 \to \mathbb{R}$ are

$$Y_l^m = \begin{cases} \frac{1}{\sqrt{2}}(Y_{l,-m} - iY_{lm}) & m < 0\\ Y_{l0} & m = 0\\ \frac{(-1)^m}{\sqrt{2}}(Y_{lm} + iY_{l,-m}) & m > 0 \end{cases}$$

We also have the relation $(Y_l^m)^* = (-1)^m Y_l^{-m}$. The zeroth and first order real spherical harmonics are given by

$$Y_{0,0}(\mathbf{x}) = \frac{1}{\sqrt{4\pi}}, \qquad Y_{1,-1}(\mathbf{x}) = \sqrt{\frac{3}{4\pi}} \frac{x^2}{|\mathbf{x}|},$$
$$Y_{1,0}(\mathbf{x}) = \sqrt{\frac{3}{4\pi}} \frac{x^3}{|\mathbf{x}|}, \qquad Y_{1,1}(\mathbf{x}) = \sqrt{\frac{3}{4\pi}} \frac{x^1}{|\mathbf{x}|}.$$

The spherical harmonics satisfy the following orthonormal conditions

$$\int_{S^2} Y_{lm} Y_{l'm'} \mathrm{d}S = \delta_{ll'} \delta_{mm'} = \int_{S^2} Y_m^m (Y_{l'}^{m'})^* \mathrm{d}S$$

and they form a basis for $L^2(S^2)$ [2] so that, in particular, any function $g \in L^2(S^2)$ has a spherical harmonics expansion

$$g = \sum_{l=0}^{\infty} \sum_{m=-l}^{l} g_{lm} Y_{lm}, \qquad g_{lm} \in \mathbb{R}$$

that converge in $L^2(S^2)$. More generally, a function $g \in L^2(B_R)$ has a spherical harmonics expansion in $L^2(B_R)$,

$$g = \sum_{l=0}^{\infty} \sum_{m=-l}^{l} g_{lm}(r) Y_{lm}, \qquad g_{lm} : [0, R] \to \mathbb{R}.$$
 (1.170)

Indeed, since $L^{2}(B_{R}) = L^{2}([0, R]; L^{2}(S^{2}), r^{2}) = L^{2}([0, R]; L^{2}(\partial B_{r}))$, or in other words

$$\int_{B_R} |\cdot| \, \mathrm{d}\mathbf{x} = \int_0^R \int_{\partial B_r} |\cdot| \, \mathrm{d}S \mathrm{d}r,$$

 $g|_{\partial B_r}$ must be in $L^2(\partial B_r)$ for almost every $r \in [0, R]$. So a spherical harmonics expansion exist for almost every r. Now

$$\left\|g - \sum_{l=0}^{N} \sum_{m=-l}^{l} g_{lm} Y_{lm}\right\|_{L^{2}(B_{R})}^{2} = \int_{0}^{R} \left\|g - \sum_{l=0}^{N} \sum_{m=-l}^{l} g_{lm} Y_{lm}\right\|_{L^{2}(\partial B_{r})}^{2} \mathrm{d}r$$

$$\rightarrow 0$$
 as $N \rightarrow \infty$

by dominated convergence theorem (where the dominating function is $4||g||^2_{L^2(\partial B_r)}$). Hence (1.170) converge in $L^2(B_R)$. Similarly, functions in $L^2(B_R, \bar{w}^{-2})$ and $L^2(\mathbb{R}^3)$ have a spherical harmonics expansion.

The following lemma allows us to expand gravitational potentials in spherical harmonics. Lemma A.3. For $\mathbf{x}, \mathbf{y} \in \mathbb{R}^3$ we have

$$\frac{1}{|\mathbf{x} - \mathbf{y}|} = 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \frac{1}{2l+1} \frac{\min\{|\mathbf{x}|, |\mathbf{y}|\}^{l}}{\max\{|\mathbf{x}|, |\mathbf{y}|\}^{l+1}} Y_{lm}(\mathbf{y}) Y_{lm}(\mathbf{x})$$

and this expression converge uniformly for (\mathbf{x}, \mathbf{y}) in any compact set in $\{(\mathbf{r}, \mathbf{r}') \in \mathbb{R}^6 : |\mathbf{r}| \neq |\mathbf{r}'|\}$.

Proof. From [28] we have

$$\frac{1}{|\mathbf{x} - \mathbf{y}|} = 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \frac{1}{2l+1} \frac{\min\{|\mathbf{x}|, |\mathbf{y}|\}^{l}}{\max\{|\mathbf{x}|, |\mathbf{y}|\}^{l+1}} Y_{l}^{m}(\mathbf{y})^{*} Y_{l}^{m}(\mathbf{x})$$

One derivation of this formula is as follows. Assume $r' = |\mathbf{r}'| < |\mathbf{r}| = r$, otherwise swap \mathbf{r}' and \mathbf{r} . By the law of cosines,

$$\frac{1}{|\mathbf{r} - \mathbf{r}'|} = \frac{1}{\sqrt{r^2 + (r')^2 - 2rr'\cos\gamma}} = \frac{1}{r\sqrt{1 + h^2 - 2h\cos\gamma}} \quad \text{with} \quad h := \frac{r'}{r}.$$

We find here the generating function of the Legendre polynomials $P_{\ell}(\cos \gamma)$:

$$\frac{1}{\sqrt{1+h^2 - 2h\cos\gamma}} = \sum_{\ell=0}^{\infty} h^{\ell} P_{\ell}(\cos\gamma).$$
(1.171)

Use of the spherical harmonic addition theorem

$$P_{\ell}(\cos\gamma) = \frac{4\pi}{2\ell+1} \sum_{m=-\ell}^{\ell} (-1)^m Y_{\ell}^{-m}(\theta,\varphi) Y_{\ell}^m(\theta',\varphi')$$

gives our first formula. Since $|P_{\ell}(\cos \gamma)| \leq 1$ for all ℓ , the power series in (1.171) has radius of convergence 1, and uniform convergence for any compact set in B_1 . By Identity theorem for analytic functions, the equality of (1.171) holds for h < 1. We thus conclude that the expansion for $|\mathbf{r} - \mathbf{r}'|^{-1}$ converge uniformly on any compact set in $\{(\mathbf{r}, \mathbf{r}') \in \mathbb{R}^6 : |\mathbf{r}| \neq |\mathbf{r}'|\}$. Moreover, in real spherical harmonics,

$$\begin{split} \frac{1}{|\mathbf{x} - \mathbf{y}|} &= 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \frac{1}{2l+1} \frac{\min\{|\mathbf{x}|, |\mathbf{y}|\}^{l}}{\max\{|\mathbf{x}|, |\mathbf{y}|\}^{l+1}} Y_{l}^{m}(\mathbf{y})^{*} Y_{l}^{m}(\mathbf{x}) \\ &= 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \frac{(-1)^{m}}{2l+1} \frac{\min\{|\mathbf{x}|, |\mathbf{y}|\}^{l}}{\max\{|\mathbf{x}|, |\mathbf{y}|\}^{l+1}} Y_{l}^{-m}(\mathbf{y}) Y_{l}^{m}(\mathbf{x}) \\ &= 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \frac{1}{2l+1} \frac{\min\{|\mathbf{x}|, |\mathbf{y}|\}^{l}}{\max\{|\mathbf{x}|, |\mathbf{y}|\}^{l+1}} (Y_{l,m}(\mathbf{y}) - iY_{l,-m}(\mathbf{y})) (Y_{lm}(\mathbf{x}) + iY_{l,-m}(\mathbf{x})) \\ &+ 4\pi \sum_{l=0}^{\infty} \frac{1}{2l+1} \frac{\min\{|\mathbf{x}|, |\mathbf{y}|\}^{l}}{\max\{|\mathbf{x}|, |\mathbf{y}|\}^{l+1}} Y_{l0}(\mathbf{y}) Y_{l0}(\mathbf{x}) \\ &+ 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^{-1} \frac{1}{2l+1} \frac{\min\{|\mathbf{x}|, |\mathbf{y}|\}^{l}}{\max\{|\mathbf{x}|, |\mathbf{y}|\}^{l+1}} (Y_{l,-m}(\mathbf{y}) + iY_{lm}(\mathbf{y})) (Y_{l,-m}(\mathbf{x}) - iY_{lm}(\mathbf{x})) \\ &= 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^{l} \frac{1}{2l+1} \frac{\min\{|\mathbf{x}|, |\mathbf{y}|\}^{l}}{\max\{|\mathbf{x}|, |\mathbf{y}|\}^{l+1}} Y_{lm}(\mathbf{y}) Y_{lm}(\mathbf{x}) \end{split}$$
There also exist a vectorial version of spherical harmonics which allow the expansion of L^2 vector fields, details can be found in [3, 14].

A.3 Hardy-Poincaré inequality and embeddings

Here we will state and prove a version of the Hardy-Poincaré inequality and the embedding theorems, which are important for our analysis.

Denote $B_R = B_R(\mathbb{R}^m)$ the ball of radius R in \mathbb{R}^m , and $L^2(B_R, w)$ the L^2 space on B_R weighted by w. Denote $d_{\partial B_R}$ the distance function to ∂B_R .

Theorem A.4 (Hardy-Poincaré inequality). Let R > 0 and $k \ge 0$. For any $\theta \in H^1_{loc}(B_R(\mathbb{R}^m))$ we have

$$\|\theta - \theta_{B_{(m-1)R/m}}\|_{L^{2}(B_{R}, d^{k}_{\partial B_{R}})} \lesssim \|\nabla\theta\|_{L^{2}(B_{R}, d^{k+2}_{\partial B_{R}})},$$
(1.172)

where θ_{B_r} denotes the average of θ on B_r .

Proof. Using only the standard Poincaré inequality and elementary methods, we will provide here a proof for the case k > 0. However, in this paper we also used the case k = 0. A slightly different version of the case k = 0 was first proven in [4] which makes use of the Hardy inequality. A proof of the k = 0 case for the version here (and a more general form) can be found in [12] by Drelichman and Durán, the proof of which (with very slight modification) will work for the k = 0 as well as the k > 0 case.

It suffice to show this for $\theta \in C^1(\overline{B}_R)$ since $C^1(\overline{B}_R)$ is dense in the type of Sobolev spaces we are considering [33]. In particular we can assume $\lim_{|\mathbf{x}|\to R} \theta(\mathbf{x}) d_{\partial B_R}(\mathbf{x})^{(k+1)/2} = 0$ for integration by parts later.

Let w be a smooth function on B_R such that $d_{\partial B_R} \leq w \leq d_{\partial B_R}$ and $w = d_{\partial B_R}$ on $B_R \setminus B_{(m-1)R/m}$. Let $G = \theta w^{k/2}$. Then

$$\begin{split} \int_{B_R} |\nabla \theta|^2 w^{k+2} \mathrm{d}\mathbf{x} &= \int_{B_R} |\nabla (w^{-k/2}G)|^2 w^{k+2} \mathrm{d}\mathbf{x} = \int_{B_R} |w^{-k/2} \nabla G - \frac{k}{2} G w^{-k/2-1} \nabla w|^2 w^{k+2} \mathrm{d}\mathbf{x} \\ &= \int_{B_R} |w \nabla G - \frac{k}{2} G \nabla w|^2 \mathrm{d}\mathbf{x} = \int_{B_R} \left(w^2 |\nabla G|^2 - k(G \nabla G)(w \nabla w) + \frac{k^2}{4} |\nabla w|^2 G^2 \right) \mathrm{d}\mathbf{x} \\ &= \int_{B_R} \left(w^2 |\nabla G|^2 - \frac{k}{4} (\nabla w^2)(\nabla G^2) + \frac{k^2}{4} |\nabla w|^2 G^2 \right) \mathrm{d}\mathbf{x} \\ &= \int_{B_R} \left(w^2 |\nabla G|^2 + \frac{k}{4} \left(\Delta w^2 + k |\nabla w|^2 \right) G^2 \right) \mathrm{d}\mathbf{x} \end{split}$$

On $B_R \setminus B_{(m-1)R/m}$ we have

$$k(\Delta w^{2} + k|\nabla w|^{2}) = \frac{k}{r^{n-1}} \frac{\mathrm{d}}{\mathrm{d}r} \left(r^{n-1} \frac{\mathrm{d}(R-r)^{2}}{\mathrm{d}r} \right) + k^{2} \left(\frac{\mathrm{d}(R-r)}{\mathrm{d}r} \right)^{2}$$
$$= \frac{k}{r^{n-1}} \frac{\mathrm{d}}{\mathrm{d}r} \left(r^{n-1} (-2R+2r) \right) + k^{2} = k \left(-2(n-1)\frac{R}{r} + 2n \right) + k^{2}$$
$$\geq \begin{cases} k^{2} & k \ge 0\\ k^{2} + 2k & k \le 0 \end{cases}$$

which is strictly positive when $k \in \mathbb{R} \setminus [-2, 0]$. So we have

$$\begin{split} \frac{k^2 + \min\{0, 2k\}}{4} \int_{B_R} \theta^2 w^k \mathrm{d}\mathbf{x} &= \frac{k^2 + \min\{0, 2k\}}{4} \int_{B_R} G^2 \mathrm{d}\mathbf{x} \\ &\leq \int_{B_R} |\nabla \theta|^2 w^{k+2} \mathrm{d}\mathbf{x} + \frac{k}{4} (k + \|\Delta w^2\|_{\infty} + k\|\nabla w\|_{\infty}) \int_{B_{(m-1)R/m}} G^2 \mathrm{d}\mathbf{x} \\ &\lesssim \int_{B_R} |\nabla \theta|^2 w^{k+2} \mathrm{d}\mathbf{x} + \int_{B_{(m-1)R/m}} \theta^2 \mathrm{d}\mathbf{x} \end{split}$$

$$egin{aligned} &\lesssim \int_{B_R} |
abla heta|^2 w^{k+2} \mathrm{d} \mathbf{x} + \left(\int_{B_{(m-1)R/m}} heta \mathrm{d} \mathbf{x}
ight)^2 + \int_{B_{(m-1)R/m}} |
abla heta|^2 \mathrm{d} \mathbf{x} \ &\lesssim \int_{B_R} |
abla heta|^2 w^{k+2} \mathrm{d} \mathbf{x} + \left(\int_{B_{(m-1)R/m}} heta \mathrm{d} \mathbf{x}
ight)^2 \end{aligned}$$

where we used the Poincaré inequality on $B_{(m-1)R/m}$. Replacing θ with $\theta - \theta_{B_{(m-1)R/m}}$ (valid when k > -1) we see that

$$\int_{B_R} (\theta - \theta_{B_{(m-1)R/m}})^2 w^k \mathrm{d}\mathbf{x} \lesssim \int_{B_R} |\nabla \theta|^2 w^{k+2} \mathrm{d}\mathbf{x}.$$

An immediate corollary of the Hardy-Poincaré inequality is the following. Corollary A.5. Let $k, l \ge 0$. We have

$$\|\theta\|_k \lesssim \|\theta\|_l + \|\nabla\theta\|_{k+2} \tag{1.173}$$

Proof. We have

$$\begin{aligned} \|\theta\|_{k} &\leq \|\theta - \theta_{B_{(m-1)R/m}}\|_{k} + \|\theta_{B_{(m-1)R/m}}\|_{k} \lesssim \theta_{B_{(m-1)R/m}}\|1\|_{k} + \|\nabla\theta\|_{k+2} \\ &\leq \|\theta\|_{L^{2}(B_{(m-1)R/m})}\|1\|_{L^{2}(B_{(m-1)R/m})}\|1\|_{k} + \|\nabla\theta\|_{k+2} \lesssim \|\theta\|_{l} + \|\nabla\theta\|_{k+2}. \end{aligned}$$

Using this corollary we will next derive the embedding theorems. We will show that terms with less than n/2 the derivatives can be estimated in the L^{∞} norm by E_n or $E_n + Z_n^2$, which is relevant for the energy estimates. For this we will need the following lemmas.

Lemma A.6. We have

$$\begin{split} \sum_{c=0}^{n} \|\nabla^{n-c} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta}\|_{\max\{3+b+n-2c,0\}}^{2} \lesssim \sum_{c=0}^{n} \|\nabla^{c} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta}\|_{3+b+c}^{2} \\ \|\bar{w}^{\lfloor b/2 \rfloor} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta}(s)\|_{H^{n}}^{2} \lesssim \sum_{c=0}^{4+2n} \|\nabla^{c} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta}\|_{3+b+c}^{2} \\ \|\bar{w}^{\lfloor b/2 \rfloor} \nabla X_{r}^{b} \partial^{\beta} \boldsymbol{\theta}(s)\|_{H^{n}}^{2} \lesssim \sum_{c=1}^{6+2n} \|\nabla^{c} X_{r}^{b} \partial^{\beta} \boldsymbol{\theta}\|_{3+b+c}^{2} \end{split}$$

Proof. The first formula follows from repeated application of the above Corollary A.5. The formulas after follows from the first with n replaced by 4 + 2n and 6 + 2n respectively.

Lemma A.7. We have

$$\begin{split} \|\bar{w}^{\lfloor b/2 \rfloor} X_r^b \partial\!\!\!\!\partial^{\beta} \boldsymbol{\theta}(s)\|_{L^{\infty}}^2 \lesssim \sum_{c=0}^8 \|\nabla^c X_r^b \partial\!\!\!\!\partial^{\beta} \boldsymbol{\theta}\|_{3+b+c}^2 \\ \lesssim \sum_{b'+|\beta'| \le 8+b+|\beta|} \|X_r^{b'} \partial\!\!\!\!\partial^{\beta'} \boldsymbol{\theta}\|_{3+b'}^2 + \sum_{c \le 8+b+|\beta|} \|\nabla^c \boldsymbol{\theta}\|_{3+2c}^2 \\ \|\bar{w}^{\lfloor b/2 \rfloor} \nabla X_r^b \partial\!\!\!\!\partial^{\beta} \boldsymbol{\theta}(s)\|_{L^{\infty}}^2 \lesssim \sum_{c=1}^{10} \|\nabla^c X_r^b \partial\!\!\!\!\partial^{\beta} \boldsymbol{\theta}\|_{3+b+c}^2 \\ \lesssim \sum_{b'+|\beta'| \le 10+b+|\beta|} \|X_r^{b'} \partial\!\!\!\!\partial^{\beta'} \boldsymbol{\theta}\|_{3+b'}^2 + \sum_{c \le 10+b+|\beta|} \|\nabla^c \boldsymbol{\theta}\|_{3+2c}^2 \end{split}$$

Proof. This follows from Lemma A.6 and the embedding $H^2 \hookrightarrow L^{\infty}$.

A.3.1 Embedding theorems for self-similarly expanding GW stars

Using Lemmas A.6 and A.7 we can derive the Embedding theorems for self-similarly expanding GW stars, relevant for Section 3.

Theorem A.8 (Near boundary embedding theorem). We have

$$\begin{split} \sum_{\substack{a+|\beta|+b\leq n \\ a>0}} \|\bar{w}^{\lfloor b/2 \rfloor} \partial_s^a X_r^b \partial^{\beta} \theta(s)\|_{L^{\infty}}^2 \lesssim \sum_{\substack{a+|\beta|+b\leq 8+n \\ s \in R+8 \\ s = R+8 \\ \sum_{\substack{a+|\beta|+b\leq n \\ a>0}} \|\bar{w}^{\lfloor b/2 \rfloor} \nabla \partial_s^a X_r^b \partial^{\beta} \theta(s)\|_{L^{\infty}}^2 \lesssim \sum_{\substack{a+|\beta|+b\leq 8+n \\ a>0}} \|\partial_s^a X_r^b \partial^{\beta} \theta\|_{3+b}^2 + \sum_{\substack{a+c\leq 8+n \\ a>0}} \|\partial_s^a \nabla^c \theta\|_{3+2c}^2 \\ \lesssim E_{n+8} \\ \sum_{\substack{a+|\beta|+b\leq n \\ s>0}} \|\bar{w}^{\lfloor b/2 \rfloor} \nabla \partial_s^a X_r^b \partial^{\beta} \theta(s)\|_{L^{\infty}}^2 \lesssim \sum_{\substack{a+|\beta|+b\leq 10+n \\ a>0}} \|\partial_s^a X_r^b \partial^{\beta} \theta\|_{3+b}^2 + \sum_{\substack{a+c\leq 10+n \\ a+c\leq 10+n \\ s>0}} \|\partial_s^a \nabla^c \theta\|_{3+2c}^2 \\ \lesssim E_{n+10} + Z_{n+10}^2 \\ \sum_{\substack{a+|\beta|+b\leq 10+n \\ a>0}} \|\bar{w}^{\lfloor b/2 \rfloor} \nabla \partial_s^a X_r^b \partial^{\beta} \theta(s)\|_{L^{\infty}}^2 \lesssim \sum_{\substack{a+|\beta|+b\leq 10+n \\ a>0}} \|\partial_s^a X_r^b \partial^{\beta} \theta\|_{3+b}^2 + \sum_{\substack{a+c\leq 10+n \\ a>0}} \|\partial_s^a \nabla^c \theta\|_{3+2c}^2 \\ \lesssim E_{n+10} \\ \lesssim E_{n+10} \end{split}$$

Proof. The proof is a direct consequence of Lemma A.7.

Theorem A.9 (Near origin embedding theorem). We have

$$\sum_{a+c \le n+1} \|\bar{w}^c \partial_s^a \nabla^c \boldsymbol{\theta}(s)\|_{L^{\infty}}^2 \lesssim E_{n+10} + Z_{n+10}^2$$

Proof. Similar to above A.8.

A.3.2 Embedding theorems for linearly expanding GW stars

Using Lemmas A.6 and A.7 we can derive the Embedding theorems for linearly expanding GW stars, relevant for Section 4.

Theorem A.10 (Near boundary embedding theorem). We have

$$\begin{split} \sum_{|\beta|+b\leq n} \|\bar{w}^{\lfloor b/2 \rfloor} X_r^b \partial^{\beta} \theta(s)\|_{L^{\infty}}^2 &\lesssim \sum_{|\beta|+b\leq 8+n} \|X_r^b \partial^{\beta} \theta\|_{3+b}^2 + \sum_{c\leq 8+n} \|\nabla^c \theta\|_{3+2c}^2 \\ &\lesssim E_{n+8} \\ \sum_{|\beta|+b\leq n} \|\bar{w}^{\lfloor b/2 \rfloor} \partial_s X_r^b \partial^{\beta} \theta(s)\|_{L^{\infty}}^2 &\lesssim \sum_{|\beta|+b\leq 8+n} \|\partial_s X_r^b \partial^{\beta} \theta\|_{3+b}^2 + \sum_{c\leq 8+n} \|\partial_s \nabla^c \theta\|_{3+2c}^2 \\ &\lesssim \lambda^{-1} E_{n+8} \\ \sum_{|\beta|+b\leq n} \|\bar{w}^{\lfloor b/2 \rfloor} \nabla X_r^b \partial^{\beta} \theta(s)\|_{L^{\infty}}^2 &\lesssim \sum_{|\beta|+b\leq 10+n} \|X_r^b \partial^{\beta} \theta\|_{3+b}^2 + \sum_{c\leq 10+n} \|\nabla^c \theta\|_{3+2c}^2 \\ &\lesssim E_{n+10} \end{split}$$

Proof. The proof is a direct consequence of Lemma A.7.

Theorem A.11 (Near origin embedding theorem). We have

$$\sum_{\substack{c \le n+1 \\ c \le n}} \|\bar{w}^c \nabla^c \boldsymbol{\theta}(s)\|_{L^{\infty}}^2 \lesssim E_{n+10}$$
$$\sum_{\substack{c \le n}} \|\bar{w}^c \partial_s \nabla^c \boldsymbol{\theta}(s)\|_{L^{\infty}}^2 \lesssim \lambda^{-1} E_{n+10}$$

Proof. Similar to above A.10.

References

- C. Alexander, M. Hadžić, M., Schrecker, Supersonic Gravitational Collapse for Non-Isentropic Gaseous Stars. *Preprint, arXiv:2311.18795* (2023)
- [2] Atkinson, K., Han, W., *Spherical Harmonics and Approximations on the Unit Sphere: An Introduction*. Lecture Notes in Mathematics, Springer, 2012.
- [3] Barrera, R. G., Estévez, G. A., Giraldo, J., Vector spherical harmonics and their applications to magnetostatics. *Eur. J. Phys.*, **6**, 287–294 (1985)
- [4] Boas, H. B., Straube, E. J., Integral inequalities of Hardy and Poincaré type. *Proc. Amer. Math. Soc.*, 103, 172–176 (1988)
- [5] Chandrasekhar, S., An introduction to the study of stellar structure. University of Chicago Press, Chicago, 1939.
- [6] Cheng, M., Cheng, X., Lin, Z., Expanding solutions near unstable Lane-Emden stars. *Preprint*, arXiv:2305.10218 (2023)
- [7] Chen G-Q. G., He, L., Wang, Y., Yuan D., Global Solutions of the Compressible Euler-Poisson Equations with Large Initial Data of Spherical Symmetry. *Comm. Pure Appl. Math.* (2023), online first
- [8] Coutand, D., Shkoller, S., Well-posedness in smooth function spaces for the moving boundary threedimensional compressible Euler equations in physical vacuum. *Arch. Ration. Mech. Anal.*, 206, no. 2, 515–616 (2012)
- [9] Courant, R., Hilbert, D., Methods of mathematical physics. Vol. 1. New York: Interscience 1953.
- [10] Dacorogna B., Moser, J., On a partial differential equation involving the Jacobian determinant. Ann. Inst. H. Poincaré Anal. Non Linéaire, 7, 1, 1–26 (1990)
- [11] Deng, Y., Xiang, J., Yang, T., Blowup phenomena of solutions to Euler-Poisson equations. J. Math. Anal. Appl., 286, 295–306 (2003)
- [12] Drelichman, I., Durán, R. G., Improved Poincaré inequalities with weights. *Journal of Mathematical Analysis and Applications*, **347**, 286–293 (2008)
- [13] Dyson F. J., Dynamics of a Spinning Gas Cloud. J. Math. Mech., 18, no. 1, (1968) 91–101
- [14] Freeden, W., Schreiner, M., *Spherical Functions of Mathematical Geosciences*, Geosystems Mathematics, Birkhäuser, 2.ed., Heidelberg, 2022.
- [15] Fu, C.-C., Lin, S.-S., On the critical mass of the collapse of a gaseous star in spherically symmetric and isentropic motion. *Japan J. Indust. Appl. Math.*, **15**, no. 3, 461–469 (1998)
- [16] Goldreich, P., Weber, S., Homologously collapsing stellar cores. Astrophys. J., 238, 991–997 (1980)
- [17] Grassin, M., Global smooth solutions to Euler equations for a perfect gas. *Indiana Univ. Math. J.*, 47, 1397-1432 (1998)
- [18] Gu, X., Lei, Z., Local Well-posedness of the three dimensional compressible Euler–Poisson equations with physical vacuum. *Journal de Mathématiques Pures et Appliquées*, **105**, 5, 662–723 (2016)
- [19] Guo, Y., Hadžić, M., Jang, J., Continued gravitational collapse for Newtonian stars. Arch. Rat. Mech. Anal. 239, 431–552 (2021)
- [20] Guo, Y., Hadžić, M., Jang, J., Larson-Penston Self-similar Gravitational Collapse. Comm. Math. Phys., 386, 1551–1601 (2021)
- [21] Guo, Y., Hadžić, M., Jang, J., Naked singularities in the Einstein-Euler system. Annals of PDE, 9, 4 (2023)
- [22] Guo, Y., Hadžić, M., Jang, J., Schrecker, M., Gravitational Collapse for Polytropic Gaseous Stars: Self-similar Solutions. Arch. Rat. Mech. Anal., 246, 957–1066 (2022)
- [23] Hadžić, M., Star dynamics: Collapse vs. expansion. Quarterly of Applied Math., 81, 329–365 (2023)
- [24] Hadžić, M., Jang, J., Nonlinear stability of expanding star solutions in the radially-symmetric masscritical Euler-Poisson system. *Comm. Pure Appl. Math.*, 71, no. 5, 827–891 (2018)

- [25] Hadžić, M., Jang, J., Expanding large global solutions of the equations of compressible fluid mechanics. *Invent. Math.*, 1–62 (2018)
- [26] Hadžić, M., Jang, J., A class of global solutions to the Euler-Poisson system. *Commun. Math. Phys.*, 370, no 2, 475–505 (2019)
- [27] Ifrim, M., Tataru, D., The compressible Euler equations in a physical vacuum: a comprehensive Eulerian approach. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2023), online first
- [28] Jackson, J. D., *Classical Electrodynamics*. Wiley, NY, 1962.
- [29] Jang, J., Nonlinear Instability in Gravitational Euler-Poisson system for $\gamma = 6/5$. Arch. Ration. Mech. Anal., **188**, 265–307 (2008)
- [30] Jang, J., Nonlinear Instability Theory of Lane-Emden stars. Comm. Pure Appl. Math., 67, no. 9, 1418– 1465 (2014)
- [31] Jang, J., Masmoudi, N., Well-posedness of compressible Euler equations in a physical vacuum. Comm. Pure Appl. Math., 68, no. 1, 61–111 (2015)
- [32] Jang, J., Makino, T., Linearized analysis of barotropic perturbations around spherically symmetric gaseous stars governed by the Euler–Poisson equations. *J. Math. Phys.*, **61**, 051508 (2020)
- [33] Kufner A., Weighted Sobolev Spaces. Teubner-Texte zur Mathematik, Volume 31, Teubner, 1980
- [34] Lin, Z., Wang, Y., Zhu, H., Nonlinear stability of non-rotating gaseous stars. *Preprint*, arXiv:2305.07088 (2023)
- [35] Lin, Z., Zeng, C., Separable Hamiltonian PDEs and Turning Point Principle for Stability of Gaseous Stars. *Comm. Pure Appl. Math.*, **75**, no. 11, 2511–2572 (2021)
- [36] Luo, T., Xin, Z., Zeng, H., Well-posedness for the motion of physical vacuum of the three-dimensional compressible Euler equations with or without self-gravitation. *Arch. Ration. Mech. Anal.*, 213, no. 3, 763–831 (2014)
- [37] Luo, T., Xin, Z., Zeng, H., On nonlinear asymptotic stability of the Lane-Emden solutions for the viscous gaseous star problem. *Adv. Math.*, **291**, 90–182 (2016)
- [38] Luo, T., Xin, Z., Zeng, H., Nonlinear Asymptotic Stability of the Lane-Emden Solutions for the Viscous Gaseous Star Problem with Degenerate Density Dependent Viscosities. *Comm. Math. Phys.*, 347, no. 3, 657–702 (2016)
- [39] Luo, T., Zeng, H., Global Existence of Smooth Solutions and Convergence to Barenblatt Solutions for the Physical Vacuum Free Boundary Problem of Compressible Euler Equations with Damping. *Comm. Pure Appl. Math.*, 69, 1354–1396 (2016)
- [40] Makino, T., Blowing up solutions of the Euler-Poisson equation for the evolution of gaseous stars. *Transport Theory Statist. Phys.*, 21, 615–624 (1992)
- [41] Ovsiannikov, L. V., New solution of hydrodynamic equations. Dokl. Akad. Nauk SSSR, Vol 111, N l, (1956) 47–49.
- [42] Parmeshwar, S., Global Existence for the N Body Euler-Poisson System. Arch. Ration. Mech. Anal., 244, no. 2, 157–208 (2022)
- [43] Parmeshwar, S., Hadžić, M., Jang, J., Global expanding solutions of compressible Euler equations with small initial densities. *Quart. Appl. Math.*, **79**, no. 2, 273–334 (2021)
- [44] Rein, G., Non-linear stability of gaseous stars. Arch. Ration. Mech. Anal., 168, no. 2, 115–130 (2003)
- [45] Rickard, C., Global solutions to the compressible Euler equations with heat transport by convection around Dyson's isothermal affine solutions. *Arch. Ration. Mech. Anal.*, **241**, no. 2, 947–1007 (2021)
- [46] Rickard, C., The vacuum boundary problem for the spherically symmetric compressible Euler equations with positive density and unbounded entropy. *J. Math. Phys.*, **62**(2), 021504 (2021)
- [47] Rickard, C., Hadžić, M., Jang, J., Global existence of the nonisentropic compressible Euler equations with vacuum boundary surrounding a variable entropy state. *Nonlinearity*, **34** no. 1, 33–91 (2021)
- [48] Rozanova, O., Solutions with linear profile of velocity to the Euler equations in several dimensions. *Hyperbolic problems: theory, numerics, applications*, 861-870, Springer, Berlin, 2003

- [49] Sandine, E., Hunter self-similar implosion profiles for the gravitational Euler-Poisson system, *Preprint, arXiv:2310.18870* (2023)
- [50] Serre, D., Solutions classiques globales des équations d'Euler pour un fluide parfait compressible. Annales de l'Institut Fourier, **47**, 139–153 (1997)
- [51] Shkoller, S., Sideris, T., Global existence of near-affine solutions to the compressible Euler equations. *Arch. Rational Mech. Anal.*, **234**, 115–180 (2019)
- [52] Sideris, T., C., Global existence and asymptotic behavior of affine motion of 3D ideal fluids surrounded by vacuum. *Arch. Ration. Mech. Anal.*, **225**, no. 1, 141–176 (2017)
- [53] Zeng, H., Global resolution of the physical vacuum singularity for three-dimensional isentropic inviscid flows with damping in spherically symmetric motions. *Arch. Ration. Mech. Anal.*, 226, 33–82 (2017)
- [54] Zeng, H., Almost global solutions to the three-dimensional isentropic inviscid flows with damping in physical vacuum around Barenblatt solutions. Arch. Ration. Mech. Anal., 239, 553–597 (2021)