

Matrix Product Renormalization Group: Potential Universal Quantum Many-Body Solver

Masahiko G. Yamada,^{1,2,*} Takumi Sanno,² Masahiro O. Takahashi,²
Yutaka Akagi,³ Hidemaro Suwa,³ Satoshi Fujimoto,^{2,4} and Masafumi Udagawa¹

¹*Department of Physics, Gakushuin University, Mejiro, Tokyo, 171-8588, Japan*

²*Department of Materials Engineering Science, Osaka University, Toyonaka 560-8531, Japan*

³*Department of Physics, The University of Tokyo, Hongo, Tokyo, 113-0033, Japan*

⁴*Center for Quantum Information and Quantum Biology, Osaka University, Toyonaka 560-8531, Japan*

(Dated: December 29, 2022)

The density matrix renormalization group (DMRG) is a celebrated tensor network algorithm, which computes the ground states of one-dimensional quantum many-body systems very efficiently. Here we propose an improved formulation of continuous tensor network algorithms, which we name a matrix product renormalization group (MPRG). MPRG is a universal quantum many-body solver, which potentially works at both zero and finite temperatures, in two and higher dimensions, and is even applicable to open quantum systems. Furthermore, MPRG does not rely on any variational principles and thus supports any kind of non-Hermitian systems in any dimension. As a demonstration, we present critical properties of the Yang-Lee edge singularity in one dimension as a representative non-Hermitian system.

Introduction. — Variational principles appear everywhere in physics. For example, classical mechanics can be formulated by the principle of least action. Even in quantum systems, the variational principle is the central assumption to solve various problems efficiently, which is especially useful to obtain the ground state of Hermitian systems.

One of the most successful applications of the quantum variational principle is the density matrix renormalization group (DMRG) algorithm for one-dimensional (1D) quantum many-body systems [1, 2]. DMRG is a tensor network algorithm which systematically computes the ground state of 1D quantum many-body systems based on a variational principle [3]. However, due to the variational principle, most of the methods, including DMRG, mainly deal with Hermitian quantum systems, and it still remains a challenging problem to develop a technique for open quantum systems like non-Hermitian systems [4].

Non-Hermitian systems attract attention these days due to the recent experimental progress [5–9]. While the development in the realization of open quantum systems or engineering non-Hermitian terms in cold atomic systems is remarkable, the dissipation effect is ubiquitous and inevitable in any kind of realistic quantum systems, and thus dealing with non-Hermitian dissipation terms should be at the heart of the application of quantum mechanics to the real world.

Variational principles are not directly applicable to non-Hermitian quantum systems, and thus solving interacting open quantum many-body systems has long been a very difficult task. While non-variational method like infinite time-evolving block decimation (iTEBD) [10, 11] has been used to simulate non-Hermitian 1D quantum many-body systems [12], it requires a large bond dimension to obtain a reliable result from iTEBD. We need a more efficient algorithm which can solve non-Hermitian

systems with a direct method.

However, obtaining the direct solution of the many-body Schrödinger equation without a variational principle is highly challenging. One of the main reasons is the numerical instability of solving the generalized eigenvalue problem. We find that this instability can be removed by using the so-called *continuous* tensor product state [13–16], which can be regarded as a hyperdual-number-coefficient tensor network state [17]. This formalism allows us to diagonalize general many-body Hamiltonians efficiently based on tensor network ansatz for both Hermitian and non-Hermitian quantum systems in a nonvariational way. Indeed, the use of hyperdual numbers drastically simplifies the algorithm, and facilitates the implementation with modern programming languages. We refer to a continuous tensor network algorithm with this hyperdual number eigensolver as a matrix product renormalization group (MPRG).

Compared with other methods, including a quantum Monte Carlo, MPRG has the following three advantages: (i) the absence of numerical sign problems, (ii) the absence of Trotter errors, and (iii) the absence of finite-size effects. MPRG is a potential universal quantum many-body solver, which can compute the correct thermodynamic quantities in the large-bond-dimension limit.

In this Letter, we propose MPRG as a universal quantum many-body solver. MPRG potentially works at both zero and finite temperature, in any dimensions, and is even applicable to open quantum systems without a variational principle. As a demonstration, we formulate MPRG in the 1D case, and show that MPRG can solve various types of many-body problems very efficiently in both Hermitian and non-Hermitian cases. Specifically, we present a benchmark result for the Yang-Lee edge singularity of the complex-valued-transverse-field Ising model [18–20]. Furthermore, the extension to two and

higher dimensions is straightforward. This will eventually give rise to an ultimate solution to notorious numerical sign problems.

Continuous matrix product state and continuous projected entangled-pair states. — For 1D chain systems, the tensor product state ansatz is usually called a matrix product state (MPS). MPS is used in many quantum many-body solvers, such as DMRG and iTEBD. Those algorithms use a discrete version of MPS, which is very efficient due to the existence of the so-called canonical form. The generalization to higher dimensions is called projected entangled-pair states (PEPS), and again it is useful to solve the ground states of various quantum many-body systems at zero temperature.

Recently, a continuous variant of tensor product states appeared, and it is found that the continuous version is useful, especially for finite-temperature systems. For example, by using a continuous MPS (cMPS), we can eliminate the error associated with the Trotter decomposition, leading to an accurate computation of the partition function at finite temperature. Later we will see that cMPS can be reformulated as a hyper-dual-number-coefficient MPS in cases we are interested in, and this view will simplify various technical difficulties in the calculation. Similarly, we define a continuous PEPS (cPEPS) as a hyper-dual-number-coefficient PEPS for higher dimensions [21].

Matrix product renormalization group. — A hyper-dual number is an extended version of a dual number. We have two elements ε_1 and ε_2 that satisfy $\varepsilon_1^2 = 0$, $\varepsilon_2^2 = 0$, and ε_1 commutes with ε_2 . These relations are enough to regard ε_1 and ε_2 as two independent infinitesimal values. Any hyper-dual number can be denoted by $a + b\varepsilon_1 + c\varepsilon_2 + d\varepsilon_1\varepsilon_2$, where a , b , c , and d are complex numbers, and using this quantity, we can compute the derivative of functions up to the second order.

Similarly, the Taylor expansion about $\sqrt{\Delta\tau}$ can be possible up to the second order, where $\Delta\tau$ is a Trotter error. From a simple fact $[(\varepsilon_1 + \varepsilon_2)/\sqrt{2}]^2 = \varepsilon_1\varepsilon_2$, we can easily identify $(\varepsilon_1 + \varepsilon_2)/\sqrt{2} = \sqrt{\Delta\tau}$ and $\varepsilon_1\varepsilon_2 = \Delta\tau$ to translate the continuous tensor network into the hyper-dual-number-coefficient tensor network.

In the appendix of Ref. [16], many continuous matrix product operators (cMPOs) for various quantum models are provided, and we can easily translate them into hyper-dual-number-coefficient MPOs. For example, cMPO for the transverse-field Ising model is

$$M = \begin{pmatrix} I + \Gamma S^x & \sqrt{J/2}(\varepsilon_1 + \varepsilon_2)S^z \\ \sqrt{J/2}(\varepsilon_1 + \varepsilon_2)S^z & 0 \end{pmatrix}, \quad (1)$$

where M is a rank-4 cMPO, $S^{x,y,z}$ is the spin-1/2 operator, and I and 0 are identity and zero matrices, respectively. The parameters J and Γ are model parameters for the transverse-field Ising model defined later. After this identification, we still need to make a contraction of these infinite tensor networks. However, we will see that

the contraction is much easier than conventional discrete tensor networks, thanks to the introduction of hyper-dual numbers.

To do this contraction with hyper-dual numbers, we propose another type of continuous tensor network method named MPRG. Let us discuss the ground state calculation of MPRG. The algorithm for the ground state is similar to infinite DMRG (iDMRG), or the variational uniform matrix product state (VUMPS) ansatz [22]. The difference lies in the fact that the transformation to a canonical form of MPS is no longer necessary, and cMPS environmental tensors can be computed directly in a translation-invariant form. While we need to solve the generalized eigenvalue problem in this case, for hyper-dual-number-coefficient cMPS, it is enough to compute the *derivative* of the generalized eigenvalue problem. The derivative of the generalized eigenvalue problem can be computed directly with a forward-mode automatic differentiation, or we can simply use the power method. We found that the power method has a better performance in both Hermitian and non-Hermitian cases to *avoid* solving the full generalized eigenvalue problem. This is the basic idea of MPRG.

The generalization to the finite temperature is straightforward. We can even use the *same* environmental tensor calculated for the ground state. This is a remarkable fact because it means that a single simulation for the $T \rightarrow 0$ limit is enough for all physical temperature T . This is due to the generalized Hellmann-Feynman theorem for continuous tensor networks [23]. Indeed, from this theorem, it is easy to show that the free energy does not depend on $\beta = 1/T$ *implicitly*, and it means that the β dependence of the environmental tensor does not affect any thermodynamic quantities computed from the free energy.

We note that all the calculations in this work have been done using the Julia language [24] with a forward-mode automatic differentiation library Handagote.jl [25].

Fixed point iteration. — Based on this observation, we can easily formulate the fixed point iteration for the cMPS environmental tensor. The fixed point iteration for MPRG is similar to that for VUMPS, which is the sequence of the iteration of the environment calculation, the power method for hyper-dual-number-valued matrices, and the cMPS tensor update.

First, the environment calculation is done simply by diagonalizing the $\varepsilon_1\varepsilon_2$ -part of the hyper-dual-number-coefficient transfer matrix. This is justified by the fact that after the convergence, the transfer matrix always becomes $I - \varepsilon_1\varepsilon_2 H_{\text{eff}}$, where I is an identity matrix and H_{eff} is an effective Hamiltonian. Thus, it is enough to diagonalize the $\varepsilon_1\varepsilon_2$ -part of the transfer matrix, and environments are always pure complex tensors; In the degenerate perturbation theory, the first-order perturbation affects only the zeroth-order part of the wavefunction. This fact is not altered for non-Hermitian systems, i.e., the same

method can be applied to non-Hermitian systems.

Next, the power method is used to solve the generalized eigenvalue problem in terms of hyper-dual numbers. In order to solve the problem of the form $A\psi = \lambda B\psi$, where (A, B) is a given matrix pencil and (λ, ψ) is a resulting eigenpair, we compute B^{-1} first using the derivative of the matrix inverse. The eigenvector ψ with the largest eigenvalue is computed by applying $B^{-1}A$ many times, or more simply by solving some consistency equation derived from the fixed point iteration. We found that the latter is better for our simulation. The details of this approach will be discussed in a future publication.

Finally, When updating the cMPS environmental tensor, it is important to stabilize the whole fixed point iteration by the following important modification. If the fixed point iteration from the old tensor E_i to the new tensor E_{i+1} has the form of $E_{i+1} = f(E_i)$, then the modified (stabilized) version of this fixed point iteration should be $E_{i+1} = (1 - \alpha)E_i + \alpha f(E_i)$. Here $\alpha \in (0, 1]$ is a small number chosen carefully to stabilize the fixed point iteration. The optimal α strongly depends on the model and the parameter, and we can only know its value empirically.

Comparison with an exact solution in Hermitian systems. — After the above modification, we succeed in stabilizing the calculation for various 1D Hermitian systems. As the simplest benchmark, we demonstrate a simulation for the 1D transverse-field Ising model, and compare the result with the exact one. The Hamiltonian is

$$H_{\text{Ising}} = - \sum_{j=1}^N (JS_j^z S_{j+1}^z + \Gamma S_j^x), \quad (2)$$

where $S_j^{x,y,z}$ is the spin-1/2 operator at site j and N is the system size. Here, we impose periodic boundary conditions, and set $\lambda \equiv J/2\Gamma$ with $\Gamma = 1$. The parameter λ is normalized so that $\lambda = 1$ corresponds to the critical point, and we use this normalization from now on.

While the calculation for gapped cases $\lambda \neq 1$ converges very fast and accurately as by DMRG and VUMPS, the most remarkable thing of MPRG is the accuracy at the critical point $\lambda = 1$. First, we show the ground state energy calculated for various bond dimensions χ by MPRG at $\lambda = 1$, and it converges very rapidly to the exact value calculated for the $N = 10^7$ system, as shown in Table I. The calculation with $\chi = 8$ is already very accurate, and

χ	E_G/N
2	-0.63627091
4	-0.63659594
8	-0.63660486
exact ($N = 10^7$)	-0.63661982

TABLE I. Ground state energy E_G/N for the transverse-field Ising model at $\lambda = 1$.




FIG. 1. Specific heat for the 1D transverse-field Ising model calculated by MPRG (solid lines) for different bond dimensions χ . The exact values ($N = 10^4$) are shown in purple dots.

this means that MPRG has the ability to represent a gapless state with a very small bond dimension.

The high representation ability of MPRG is more evident in the finite-temperature calculation. As shown in Fig. 1, the heat capacity calculated for $\chi = 8$ by MPRG, shown in the green line, is almost converged to the exact result ($N = 10^4$) shown in purple dots. While we see an oscillation at low temperature for small bond dimensions $\chi = 2, 4$, the accuracy of $\chi = 8$ is remarkable as we see the correct gapless behavior at the lowest temperature. Although MPS methods usually fail to capture the gapless behavior [26], it is shown that cMPS is more suitable for gapless (critical) systems.

Demonstration in Non-Hermitian systems. — Next, in order to demonstrate the potential of MPRG, we simulate a more exotic critical behavior, which has never been discussed in previous variational approaches. As a benchmark result for non-Hermitian systems, we reproduce the phase diagram of the Yang-Lee Ising spin model. The Yang-Lee Ising spin model is the 1D transverse-field Ising spin model with an additional pure imaginary magnetic field h :

$$H_{\text{YL}} = - \sum_{j=1}^N (JS_j^z S_{j+1}^z + \Gamma S_j^x + ih S_j^z). \quad (3)$$

We also impose periodic boundary conditions on the system. The magnetization of this model exhibits singular behavior with a negative exponent close to the critical magnetic field h_c . This *non-unitary* critical behavior is called Yang-Lee edge singularity [18–20]. This critical phenomenon can be described by the *non-unitary* conformal field theory (CFT) with a central charge $c = -22/5$ [27]. This model also possesses the *Parity-Time* (\mathcal{PT}) symmetry, which imposes a constraint on the spectrum of this model. From the eigenvalue spectrum theory, the spectrum is real when the ground state is invariant with the \mathcal{PT} symmetry, while after breaking the \mathcal{PT}




FIG. 2. Imaginary part of E_G/N . These values are numerically zero up to h_c .

symmetry, the spectrum changes to complex conjugate pairs [28, 29]. Therefore, the Yang-Lee model attracts much attention from viewpoints of the *non-unitary* CFT and as the \mathcal{PT} -symmetric system. Notably, elementary excitations close to h_c could be *non-unitary non-Abelian anyons* and satisfy the same fusion rule as that of Fibonacci anyons [30–32].

Breaking the \mathcal{PT} symmetry, the spectra of the Yang-Lee model drastically change from real to complex spectra. The imaginary part of the spectra directly corresponds to the phase diagram of the Yang-Lee model. Fig. 2 shows the imaginary part of the ground state energy per site E_G/N [33]. The line in Fig. 2 means the critical magnetic field h_c . Table II shows some results

λ	h_c	λ	h_c
0.10	0.636	0.50	0.154
0.20	0.457	0.60	0.095
0.30	0.328	0.70	0.052
0.40	0.230	0.80	0.021

TABLE II. Critical magnetic field h_c .

of the critical magnetic field h_c . This numerical result is consistent with the previous study based on the exact diagonalization method [34].

Moreover, we would like to discuss the magnetization of the Yang-Lee model as a characteristic feature of the *non-unitary* critical phenomena. From the renormalization group method and the *non-unitary* CFT, the exponent σ of the magnetization, i.e., $\langle S^z \rangle \sim (h - h_c)^\sigma$, is $-1/6$ in the thermodynamic limit. Fig. 3 shows the magnetization of the Yang-Lee model. The real part of the magnetization drastically changes from zero to a finite value by breaking the \mathcal{PT} symmetry. These numerical results would imply that the exponent of the magnetization is equal to a negative value as a function of the mag-




FIG. 3. The magnetization S^z as a function of h at $\lambda = 0.1$. The inset shows the log-log plot of the real part of S^z for $h > h_c$ region with a power function fitting.

netic field h . In fact, by breaking the \mathcal{PT} symmetry, we succeed in estimating the component as $\sigma = -0.165(2)$ as shown in the inset of Fig. 3 in the case $\lambda = 0.1$, which is consistent with the exact value. Note that, through our calculations for non-Hermitian systems, the bond dimension is set $\chi = 2$, which indicates that the computational cost is low enough.

Conclusion. — We have proposed a universal quantum many-body solver MPRG, and MPRG supports infinite systems, finite temperature, and non-Hermitian systems. We have studied the Yang-Lee edge singularity of the complex-valued-transverse-field Ising model to show the high potential of this algorithm. The representation ability is remarkable, as it can accurately represent the criticality described by the *non-unitary* CFT.

Finally, we should mention that the extension of MPRG to two or higher dimensions is straightforward by using cPEPS. We will present more systematic studies for various quantum systems in a future publication.

We thank T. Okubo, F. Pollmann, W. Tang, H.-H. Tu, and L. Wang for fruitful discussions. M.G.Y. is supported by JST PRESTO Grant No. JPMJPR225B. T.S. and M.O.T. are supported by a JSPS Fellowship for Young Scientists. Y.A. is supported by JST PRESTO Grant No. JPMJPR2251. S.F. is supported by JST CREST Grant No. JPMJCR19T5, Japan. This work was supported by JSPS KAKENHI Grant Nos. JP20H05655, JP20K14411, JP20J20468, JP21H01039, JP22K03508, JP22K14005, JP22H01147, and JP22J20066, and JSPS Grant-in-Aid for Scientific Research on Innovative Areas “Quantum Liquid Crystals” (KAKENHI Grant No. JP22H04469).

* masahiko.yamada@gakushuin.ac.jp

[1] S. R. White, Phys. Rev. Lett. **69**, 2863 (1992).
[2] S. R. White, Phys. Rev. B **48**, 10345 (1993).

- [3] See e.g. Ref. [35] for review.
- [4] K. Yamamoto, M. Nakagawa, M. Tezuka, M. Ueda, and N. Kawakami, Phys. Rev. B **105**, 205125 (2022).
- [5] M. Müller, S. Diehl, G. Pupillo, and P. Zoller, in *Advances in Atomic, Molecular, and Optical Physics*, Vol. 61 (Academic Press, 2012) pp. 1–80.
- [6] A. J. Daley, Adv. Phys. **63**, 77 (2014).
- [7] X. Peng, H. Zhou, B.-B. Wei, J. Cui, J. Du, and R.-B. Liu, Phys. Rev. Lett. **114**, 010601 (2015).
- [8] L. M. Sieberer, M. Buchhold, and S. Diehl, Rep. Prog. Phys. **79**, 096001 (2016).
- [9] Y. Ashida, Z. Gong, and M. Ueda, Adv. Phys. **69**, 249 (2020).
- [10] G. Vidal, Phys. Rev. Lett. **98**, 070201 (2007).
- [11] R. Orús and G. Vidal, Phys. Rev. B **78**, 155117 (2008).
- [12] Y. Ashida, S. Furukawa, and M. Ueda, Nat. Commun. **8**, 15791 (2017).
- [13] F. Verstraete and J. I. Cirac, Phys. Rev. Lett. **104**, 190405 (2010).
- [14] M. B. Hastings and R. Mahajan, Phys. Rev. A **91**, 032306 (2015).
- [15] E. Tirrito, N. J. Robinson, M. Lewenstein, S.-J. Ran, and L. Tagliacozzo, arXiv:1810.08050 (2018).
- [16] W. Tang, H.-H. Tu, and L. Wang, Phys. Rev. Lett. **125**, 170604 (2020).
- [17] In this Letter, we restrict the definition of continuous tensor networks to those where only the imaginary time direction is made continuous. Spatial directions are always discrete and should be treated differently.
- [18] C. N. Yang and T. D. Lee, Phys. Rev. **87**, 404 (1952).
- [19] T. D. Lee and C. N. Yang, Phys. Rev. **87**, 410 (1952).
- [20] M. E. Fisher, Phys. Rev. Lett. **40**, 1610 (1978).
- [21] The use of hyper-dual numbers is only sufficient for systems with only one- or two-body interactions. If there is a three-body interaction, a higher object, like hyper-hyper-dual numbers, will be required.
- [22] V. Zauner-Stauber, L. Vanderstraeten, M. T. Fishman, F. Verstraete, and J. Haegeman, Phys. Rev. B **97**, 045145 (2018).
- [23] This form of the Hellmann-Feynman theorem generalized to higher-order derivatives has been proven in Ref. [16] up to the second order. The extension of this proof to higher orders is straightforward and will be discussed in a future publication.
- [24] J. Bezanson, A. Edelman, S. Karpinski, and V. B. Shah, SIAM Review **59**, 65 (2017).
- [25] <https://github.com/MGYamada/Handagote.jl>.
- [26] Y. Huang, arXiv:1505.00772 (2015).
- [27] J. L. Cardy, Phys. Rev. Lett. **54**, 1354 (1985).
- [28] C. M. Bender and S. Boettcher, Phys. Rev. Lett. **80**, 5243 (1998).
- [29] C. M. Bender, V. Branchina, and E. Messina, Phys. Rev. D **87**, 085029 (2013).
- [30] E. Ardonne, J. Gukelberger, A. W. W. Ludwig, S. Trebst, and M. Troyer, New J. Phys. **13**, 045006 (2011).
- [31] M. H. Freedman, J. Gukelberger, M. B. Hastings, S. Trebst, M. Troyer, and Z. Wang, Phys. Rev. B **85**, 045414 (2012).
- [32] T. Sanno, M. G. Yamada, T. Mizushima, and S. Fujimoto, arXiv:2208.01803 (2022).
- [33] Here the ground state means that the real part of the energy is at its minimum.
- [34] G. von Gehlen, J. Phys. A: Math. Gen. **24**, 5371 (1991).
- [35] U. Schollwöck, Ann. Phys. **326**, 96 (2011), january 2011 Special Issue.