What is expected for China's SARS-CoV-2 epidemic?

Carlos Hernandez-Suarez¹, Efrén Murillo-Zamora^{2,*},

¹Instituto de Ciencias Tecnología e Innovación, Universidad Francisco Gavidia, El Progreso St., No. 2748, Colonia Flor Blanca, San Salvador, El Salvador.

²Departamento de Epidemiología, Unidad de Medicina Familiar No. 19, IMSS, Av. Javier Mina 301, 28000, Colima, Colima. MEXICO.

Correspondence*: Corresponding Author efren.murilloza@imss.gob.mx

ABSTRACT

Recently, China announced that its "zero-covid' policy would end, which will bring serious challenges to the country's health system. In here we provide simple calculations that allows us to provide an estimate of what is expected as an outcome in terms of fatalities, using the fact that it is a highly contagious disease that will expose most of a highly vaccinated population to the virus. We use recent findings regarding the amount of reduction in the risk of severe outcome achieved by vaccination and arrive to an estimate of 1.1 m deaths, 60% of these are males. In our model, 84% percent of deaths occur in individuals with age 55 years or older. In a scenario in which this protection is completely lost due to waning and the infection fatality rate of the prevalent strain reaches similar levels to the observed in the beginning of the epidemic, the death toll could reach 2.4 m, 93% in 55 years or older.

Keywords: China, COVID-19, SARS-CoV-2, Immunity, Vaccines, Epidemic size

1 INTRODUCTION

China was the first country to report SARS-CoV-2 cases in December 2019 (Pekar et al., 2021) and to implement a "zero-covid' policy, consisting in mass testing, centralized quarantine, and lockdowns. On December 7th, 2022, the government announced that this policy is ending (Wall Street Journal, 2022) which may bring great challenges to the country (Chen and Chen, 2022). Several studies have shown that the basic reproductive number R_0 , the potential of disease transmission in the absence of any control or mitigation measures, may be as high as 5 (Sanche et al., 2020; D'Arienzo and Coniglio, 2020; Liu et al., 2020; Alimohamadi et al., 2020), which suggests that more than 98% of the Chinese population will be exposed to the virus. Nevertheless, 92.61% of the population has received at least one dose of vaccine (Johns Hopkins University, 2022a), which will affect the number of expected fatalities. To date (2022/12/24), there has been reported 16,628 deaths (Johns Hopkins University, 2022a)

out of a population of about 1.4 billion, which is a reflect of the stringent "zero-covid" policy that has been implemented.

Early in the pandemics, we made forecasts of the death toll in a Za'atari refugee camp in Jordan (Hernandez-Suarez et al., 2020) by analyzing the fatality rate of SARS-CoV-2 at different ages in Mexico and projecting these in the refugee camp by using its demography. Nevertheless, this was before vaccines were available. Now, there is ample evidence of a waning vaccine effect (Goldberg et al. 2021; Housset et al. 2022; Levine-Tiefenbrun et al., 2022; Coppeta et al., 2022; Berar-Yanay et al., 2021; Kolaric et al., 2021; Hernandez-Suarez and Murillo-Zamora, 2022) and there is evidence that vaccines increase in general Age₅₀ (the age at which there is a 50% probability of severe disease if infected) in about 15 years on average for all vaccines (Hernandez-Suarez and Murillo-Zamora, 2022). In what follows we take all these factors into consideration to build a projection on the expected number of fatalities in China.

There are several factors that we cannot consider here that are indeed relevant, for instance, the availability of medical resources, including doctors, nurses, hospitals, ICU beds, new medications and the distribution and number of known comorbidities in the population. Nevertheless, even a large availability of these resources could be overwhelmed if the epidemic in China grows exponentially. Also, there is little information on the lethality of current and possible new strains, therefore, our analysis relies on the observed average lethality, the expected protection of vaccines and the available knowledge on demographics.

2 METHODOLOGY

In Hernandez-Suarez et al. (2020), we reported the observed lethality per group age, shown in Table 1 using data from a Mexican unvaccinated population with 10,330 SARS-CoV-2 infections. Recall the numbers in Table 1 are the proportion of individuals of each group age and gender that die from the disease, if infected. As previously mentioned, given the high R_0 - value we assume that most of the Chinese population will be infected.

Table 1. Lethality of SARS-CoV-2 in an unvaccinated Mexican population

Age	Males	Females
0-9	0.000028	0.000008
10-19	0.000017	0.000015
20-29	0.000155	0.000145
30-39	0.000811	0.000381
40-49	0.001998	0.001258
50-59	0.003543	0.002012
60 +	0.005871	0.004596

Table 2 contains the Chinese demographics by group age (National Bureau of Statistics of China, 2021). Since Sinovac, the prevalent vaccine was found to increase

 AGE_{50} in 16 years on average, what is needed is to subtract 16 years to each individual and rebuild the age categories. Since available intervals are 5 years wide, the closest multiple of 5 to 16 years is 15, and so we subtract 15 years to each boundary. All negative boundaries were converted to zero. Then, we concentrated the data in seven age groups to match Table 1. The result is the adjusted Chinese population shown in Table 3.

3 RESULTS

The element-by-element product of Tables 1 and 3 yields Table 4, containing expected deaths by group age and gender, for an expected total of 1.1 m deaths. About 60% of these are males and about the 84% of deaths would occur in 55 years or older.

4 DISCUSSION

Initially, this methodology was developed as a preparation for the expected outcome in refugee camps, whose crowded conditions would guarantee a fast and broad dissemination of the virus. Nevertheless, as mentioned before, the basic reproductive number of the disease is high enough to guarantee that eventually, a broad dissemination of SARS-CoV-2 would occur in any population. Our model does not allow to provide any time frame, only death toll. In October 2020 we applied this methodology to the Za'atari refugee camp in Jordan and projected 92 deaths out of 83,465 refugees, whereas for Kutupalong-Balukhali Site in Bangladesh, with 600 k refugees, we projected 565 deaths. This implied a 0.1% death toll. So far, the death toll in Za'atari camp is 48 (UNHCR 2022) and in Kutupalong-Balukhali camp is 42 (WHO 2022). The large differences are due to our model is very sensitive to the demographics of the population, for instance a 60+ years infected individual is 7 times more likely to die than 30 years old, and the demographics of a refugee population is difficult to assess, and thus a lot of assumptions had to be made regarding demographics of the camps. Nevertheless, our main finding was that our model predicted a fifth of deaths of other studies (Truelove, 2020), and contributed to eradicate the idea that poor nourishment, lack of medical resources, poverty, crowdedness and other diseases, characteristics of refugee camps, would rampage the population.

When we applied our methodology to Cuba, a country with similar characteristics than China in vaccine distribution (94.63% vaccinated) which also developed and deployed a national vaccine, and whose demography is available (United Nations, 2021), we calculated an expected death toll of 10,588 (see detailed calculations in Supplementary material). To date, the reported number of deaths in Cuba is 8,530, and no deaths have been reported in the last six months (Johns Hopkins University, 2022b). Our estimate is not far from the observed especially considering the possibility of under-reporting for many natural reasons.

For China, The Economist (2022) predicted around 1.5 million deaths using a modification of the SEIR model of Cai et al. (2022) and Airfinity, a data firm, estimated between 1.3 and 2.1 million deaths if the "zero-covid" policy were to be lifted (Airfinity, 2022).

Our calculations assume a protective effect of the vaccine according to what it has been reported in Hernandez-Suarez and Murillo-Zamora, (2022). Nevertheless, in a scenario in which this protection is completely lost due to waning, and the infection fatality rate of the prevalent strain reaches values around those observed in the beginning of the epidemic, the death toll could reach 2.4 m. 93% in 55 years or older.

Table 2.	Demographics of	Chinese population	per group age.
----------	-----------------	--------------------	----------------

Age	Males	Females	Total
0-4	40,969,331	36,914,557	77,883,888
5-9	48,017,458	42,226,598	90,244,056
10-14	45,606,790	39,649,204	85,255,994
15-19	39,053,343	33,630,797	72,684,140
20-24	39,675,995	35,265,680	74,941,675
25-29	48,162,270	43,685,062	91,847,332
30-34	63,871,808	60,273,382	124,145,190
35-39	50,932,037	48,080,895	99,012,932
40-44	47,632,694	45,322,636	92,955,330
45-49	58,191,686	56,033,201	114,224,887
50-54	61,105,470	60,058,826	121,164,296
55-59	50,816,026	50,584,760	101,400,786
60-64	36,871,125	36,511,813	73,382,938
65-69	36,337,923	37,667,637	74,005,560
70-74	24,162,733	25,427,303	49,590,036
75-79	14,752,433	16,486,416	31,238,849
80+	15,257,272	20,543,563	35,800,835
	721,416,394	688,362,330	1,409,778,724

Table 3. Adjusted Demographics of Chinese population per group age (after reducing 15 years to boundaries to reflect the vaccine effect).

Age	Males	Females	Total
0-9	213,322,917	187,686,836	401,009,753
10-19	112,034,078	103,958,444	215,992,522
20-29	98,564,731	93,403,531	191,968,262
30-39	119,297,156	116,092,027	235,389,183
40-49	87,687,151	87,096,573	174,783,724
50-59	60,500,656	63,094,940	123,595,596
60 +	30,009,705	37,029,979	67,039,684
Total	721,416,394	688,362,330	1,409,778,724

Age	Males	Females	Total
0-24	5,973	1,501	7,475
25-34	1,905	1,559	3,464
35-44	15,278	13,544	28,821
45-54	96,750	44,231	140,981
55-64	175,199	109,567	284,766
65-74	214,354	126,947	341,301
75 +	176,187	170,190	346,377
Total	685,645	467,540	1,153,185

Table 4. Expected deaths per group age, China (15 years have been added to boundaries)

REFERENCES

Airfinity (2022). China risks between 1.3 and 2.1 million deaths if it ends its zero-COVID strategy. Tech. Rep., <u>https://www.airfinity.com/articles/china-risks-between-1-3-and-2-1-million-deaths-if-it-ends-its-zero-covid</u> [Accessed December 22, 2022].

Alimohamadi, Y., Taghdir, M., and Sepandi, M. (2020). The estimate of the basic reproduction number for novel coronavirus disease (COVID-19): a systematic review and meta-analysis. J Prev Med Public Health. 53(3), 151.

Berar-Yanay, N., Freiman, S., Shapira, M., Saffoury, A., Elemy, A., Hamze, M., et al. (2021). Waning humoral response 3 to 6 months after vaccination with the sars-cov-2 bnt162b2 mrna vaccine in dialysis patients. J Clin Med 11, 64

Cai, J., Deng, X., Yang, J., Sun, K., Liu, H., Chen, Z., et al. (2022). Modeling transmission of SARS-CoV-2 Omicron in China. Nat Med 28, 1468–1475.

Chen, J.-M. and Chen, Y.-Q. (2022). China can prepare to end its Zero-COVID policy. Nat Med 28, 1104–1105

Coppeta, L., Ferrari, C., Somma, G., Mazza, A., D'Ancona, U., Marcuccilli, F., et al. (2022). Reduced titers of circulating anti-sars-cov-2 antibodies and risk of covid-19 infection in healthcare workers during the nine months after immunization with the bnt162b2 mrna vaccine. Vaccines 10

D'Arienzo, M. and Coniglio, A. (2020). Assessment of the sars-cov-2 basic reproduction number, R0, based on the early phase of covid-19 outbreak in Italy. Bios Health 2, 57 - 59.

Goldberg, Y., Mandel, M., Bar-On, Y. M., Bodenheimer, O., Freedman, L., Haas, E. J., et al. (2021). Waning immunity after the bnt162b2 vaccine in Israel. New Eng J Med 385, E85.

Hernandez-Suarez, C. and Murillo-Zamora, E. (2022). Waning immunity to SARS-CoV-2 following vaccination or infection. Front Med 9

Hernandez-Suarez, C., Verme, P., Radyakin, S., and Efren-Murillo (2020). Covid-19 outbreaks in refugee camps. medRxiv

Housset, P., Kubab, S., Pardon, A., Vittoz, N., Bozman, D.-F., Hanafi, L., et al. (2022). Waning but persistent humoral response 6 months after the third dose of the mrna bnt162b2 vaccine in hemodialysis and peritoneal dialysis patients. J Neph , 1-3

Johns Hopkins University (2022a). Coronavirus Resource Center. <u>https://coronavirus.jhu.edu/region/china</u> [Accessed December 25, 2022].

Johns Hopkins University (2022b). Coronavirus Resource Center. <u>https://coronavirus.jhu.edu/region/cuba</u> [Accessed December 25, 2022].

Kolaric, B., Ambriovic-Ristov, A., Tabain, I., and Vilibic-Cavlek, T. (2021). Waning immunity six months after biontech/pfizer covid-19 vaccination among nursing home residents in zagreb, croatia. *Croatian Medical Journal* 62, 630–633

Levine-Tiefenbrun, M., Yelin, I., Alapi, H., Herzel, E., Kuint, J., Chodick, G., et al. (2022). Waning of sars-cov-2 booster viral-load reduction effectiveness. *Nature Communications* 13.

Liu, Y., Gayle, A. A., Wilder-Smith, A., and Rocklo[°]v, J. (2020). The reproductive number of COVID-19 is higher compared to SARS coronavirus. *J Travel Med* 27.

National Bureau of Statistics of China (2021). *China Statistical Yearbook 2021* (National Bureau of Statistics of China)

Pekar, J., Worobey, M., Moshiri, N., Scheffler, K., and Wertheim, J. O. (2021). Timing the SARS-CoV-2 index case in Hubei province. *Science* 372, 412–417

Sanche, S., Lin, Y. T., Xu, C., Romero-Severson, E., Hengartner, N., and Ke, R. (2020). High contagiousness and rapid spread of severe acute respiratory syndrome coronavirus 2. *Emerg Infect Dis* 26, 1470

The Economist (2022). Our model shows that China's covid death toll could be massive. https://www.economist.com/china/2022/12/15/our-model-shows-that-chinas-covid-death-tollcould-be-massive [Accessed December 22, 2022].

Truelove, S., Abrahim, O., Altare, C., Lauer, S. Grantz, K., Azman, A., Spiegel, P. (2020) The potential impact of covid-19 in refugee camps in Bangladesh and beyond: A modeling study. PLOS Medicine, 17(6):1–15

United Nations (2021). Demographic Yearbook 2021 (Statistics Division).

UNHCR (2022) *COVID-19 situation in refugee camps in Jordan*. UNHCR Jordan (May 2021) <u>https://data.unhcr.org/es/documents/download/87236</u> [Accessed December 25, 2022].

Wall Street Journal (2022). China scraps most covid testing, quarantine requirements in policy pivot. <u>https://www.wsj.com/articles/china-scraps-most-testing-quarantine-requirements-in-covid-19-policy-pivot-11670398522?mod=article_inline</u> [Accessed December 25, 2022].

WHO (2022) Emergency: Rohingya Cris. Monthly Situation Report # 4. <u>https://cdn.who.int/media/docs/default-source/searo/bangladesh/bangladesh---rohingya-crisis---pdf-reports/sitreps/2022/who-cox-s-bazar-monthly-situation-report-4.pdf?sfvrsn=eec0f56a_1 [Accessed December 25, 2022].</u>

Supplementary Material

Table S1. Lethality of SARS-CoV-2 in an unvaccinated Mexican population.

Age	Males	Females
0-9	0.000028	0.000008
10-19	0.000017	0.000015
20-29	0.000155	0.000145
30-39	0.000811	0.000381
40-49	0.001998	0.001258
50-59	0.003543	0.002012
60 +	0.005871	0.004596

Table S2. Demographics of Cuban population per group age.

Age	Males	Females	Total
0-4	284,758	265,909	550,666
5-9	317,987	297,439	615,426
10-14	302,634	285,574	588,208
15-19	321,819	304,354	626,173
20-24	358,731	338,129	696,860
25-29	366,550	343,788	710,338
30-34	408,870	386,649	795,519
35-39	353,346	340,870	694,216
40-44	322,242	316,024	638,266
45-49	450,573	454,723	905,296
50-54	480,598	498,770	979,368
55-59	462,561	492,306	954,867
60-64	304,748	333,958	638,706
65-69	256,611	287,734	544,345
70-74	210,979	243,047	454,026
75-79	153,365	186,785	340,150
80-84	97,694	124,981	222,675
85-89	48,696	66,613	115,309
90-94	20,235	30,807	51,042
95-99	7,092	10,628	17,720
100 +	3,487	4,731	8,218
Total	5,533,576	5,613,819	11,147,394

Supplementary Material

Table S3. Adjusted Demographics of Cuban population per group age (after reducing 15 years to boundaries to reflect the vaccine effect).

Age	Males	Females	Total
0-9	1,585,929	1,491,405	3,077,333
10-19	775,420	730,437	1,505,857
20-29	675,588	656,894	1,332,482
30-39	931,171	953,493	1,884,664
40-49	767,309	826,264	1,593,573
50-59	467,590	530,781	998,371
60 +	330,569	424,545	755,114
Total	5,533,576	5,613,819	11,147,394

Table S4. Expected deaths per group age, Cuba (obtained multiplying element-byelement Tables S1 and S3 and adding 15 years to boundaries).

Age	Males	Females	Total
0-24	44	12	56
25-34	13	11	24
35-44	105	95	200
45-54	755	363	1,118
55-64	1,533	1,039	2,573
65-74	1,657	1,068	2,725
75 +	1,941	1,951	3,892
Total	6,048	4,540	10,588