A blow-up formula for stationary quaternionic maps $*^{\dagger}$

Jiayu Li[‡] Chaona Zhu[§]

Abstract

Let $(M, J^{\alpha}, \alpha = 1, 2, 3)$ and $(N, \mathcal{J}^{\alpha}, \alpha = 1, 2, 3)$ be Hyperkähler manifolds. Suppose that u_k is a sequence of stationary quaternionic maps and converges weakly to u in $H^{1,2}(M, N)$, we derive a blow-up formula for $\lim_{k\to\infty} d(u_k^*\mathcal{J}^{\alpha})$, for $\alpha = 1, 2, 3$, in the weak sense. As a corollary, we show that the maps constructed by Chen-Li [CL2] and by Foscolo [F] can not be tangent maps (c.f [LT], Theorem 3.1) of a stationary quaternionic map satisfing $d(u^*\mathcal{J}^{\alpha}) = 0$.

1 Introduction and the main result

A hyperkähler manifold is a Riemannian manifold (M, g) with three parallel complex structures $\{J^1, J^2, J^3\}$ compatible with the metric g such that $(J^1)^2 = (J^2)^2 = (J^3)^2 = J^1 J^2 J^3 = -id$. The simplest hyperkähler manifold is the Euclidean space \mathbb{R}^{4m} . It is well-known that the only compact hyperkähler manifolds of dimension 4 are K3 surfaces and complex tori. Let $(M, g, J^{\alpha}, \alpha = 1, 2, 3)$ and $(N, h, \mathcal{J}^{\alpha}, \alpha = 1, 2, 3)$ be hyperkähler manifolds. Let $\omega_{\alpha}(\cdot, \cdot) = g(\cdot, J^{\alpha} \cdot)$ and $\Omega_{\alpha}(\cdot, \cdot) = h(\cdot, \mathcal{J}^{\alpha} \cdot), (\alpha = 1, 2, 3)$ be the Kähler forms on M and N respectively. A smooth map $u: M \to N$ is called a quaternionic map (triholomorphic map) if

$$A_{\alpha\beta}\mathcal{J}^{\beta} \circ du \circ J^{\alpha} = du \tag{1}$$

where $A_{\alpha\beta}$ denote the entries of a matrix A in SO(3). For simplicity, we choose $A_{\alpha\beta} = \delta_{\alpha\beta}$.

The quaternionic maps (triholomorphic maps) between Hyperkähler manifolds has been studied by many aothors (cf. [BT], [Ch], [CL1, [CL2], [FKS], [W]). Quaternionic maps automatically minimize the energy functional in their homotopy classes (cf. [Ch], [CL1] and [FKS]) and hence they are harmonic. It can be verified that holomorphic and anti-holomorphic maps with respect to some complex structures on M and N are quaternionic maps. However, Chen-Li constructed quaternionic maps which are not holomorphic with respect to any complex structures on M and N (cf. [CL1]).

^{*}This work is supported by NSF grant 11721101.

[†]MSC (2000): 53C26, 53C43, 58E12, 58E20. Keywords: Stationary harmonic maps, quaternionic maps, blow-up formula.

[‡]jiayuli@ustc.edu.cn

[§]zcn1991@mail.ustc.edu.cn

Definition 1.1 A map u from M to N is called a stationary quaternionic map if it is a stationary harmonic map and it is a quaternionic map outside its singular set.

It is clear that (c.f. [BT]), if u satisfies (1) almost everywhere, and

$$d(u^*\mathcal{J}^{\alpha}) = 0, \text{ for } \alpha = 1, 2, 3,$$
 (2)

then u is a stationary quaternionic map.

Chen-Li ([CL2]) proved that, if there is a harmonic sphere $\phi:\mathbb{S}^2\to N$ which satisfies

$$d\phi J_{\mathbb{S}^2} = -\sum_{k=1}^3 a_k \mathcal{J}^k \, d\phi,\tag{3}$$

where $\vec{a} = (a_1, a_2, a_3) : \mathbb{S}^2 \to \mathbb{S}^2$, and

$$\int_{\mathbb{S}^2} x_i |\nabla \phi|^2 d\sigma = 0, \ i = 1, 2, 3, \ (x_1, x_2, x_3) \in \mathbb{S}^2,$$
(4)

then

$$u(x, x^4) = \phi(\frac{x}{|x|})$$
 for any $x \in \mathbb{R}^3 \setminus \{0\}$

is a stationary quaternionic map with the x^4 -axis as its singular set.

Chen-Li ([CL2]) showed that there does exist a complete noncompact hyperkähler manifold, into which there is a harmonic S^2 which satisfies (3) and (4). Recently, Foscolo [F] showed that there exists a compact K3 surface with the above property. However, the map u constructed by Chen-Li or by Foscolo does not satisfy (2). Now the question is whether the maps constructed by Chen-Li or by Foscolo could be a tangent map of a stationary quaternionic map with identity (2), if not the singular set of a stationary quaternionic map with identity (2) might be of codimensional 4 (Remark 1.2 in [BT]).

Suppose that u_k is a sequence of stationary quaternionic maps with bounded energies $E(u_k) \leq \Lambda$. The blow-up set of u_k can be defined as

$$\Sigma = \bigcap_{r>0} \{ x \in M | \liminf_{k \to \infty} r^{2-m} \int_{B_r(x)} | \nabla u_k |^2 dy \ge \epsilon_0 \}.$$

We can always assume that $u_k \rightharpoonup u$ weakly in $W^{1,2}(M,N)$ and that

$$|\bigtriangledown u_k|^2 dx \rightharpoonup |\bigtriangledown u|^2 dx + \nu$$

in the sense of measure as $k \to \infty$. Here ν is a nonnegative Radon measure on M with support in Σ . It is known that Σ is a \mathcal{H}^{m-2} -rectifiable set, and we may write $\nu = \theta(x)H^{m-2}[\Sigma]$. It is clear that strongly convergence in $H^{1,2}(M, N)$ preserves the identity (2). In this paper we mainly prove the following blow-up formula for weakly convergence sequence of stationary quaternionic maps.

Theorem 1.2 Let u_k be a sequence of stationary quaternionic map with $E(u_k) \leq \Lambda$. Assume that $u_k \to u$ weakly in $H^1(M, N)$. Then there exist $(a^1, a^2, a^3) \in \mathbb{R}^3$ with $\sum_{\alpha=1}^3 (a^{\alpha})^2 = 1$ such that, for any smooth (m-3)-form η with compact support in M,

$$\lim_{k \to \infty} \sum_{\alpha=1}^{3} a^{\alpha} \int_{M} d\eta \wedge u_{k}^{*} \mathcal{J}^{\alpha} = \sum_{\alpha=1}^{3} a^{\alpha} \int_{M} d\eta \wedge u^{*} \mathcal{J}^{\alpha} + \int_{\Sigma} \theta d\eta|_{\Sigma}$$
(5)

and for any $(b^1, b^2, b^3) \perp (a^1, a^2, a^3)$, there holds

$$\lim_{k \to \infty} \sum_{\alpha=1}^{3} b^{\alpha} \int_{M} d\eta \wedge u_{k}^{*} \mathcal{J}^{\alpha} = \sum_{\alpha=1}^{3} b^{\alpha} \int_{M} d\eta \wedge u^{*} \mathcal{J}^{\alpha}.$$

As a corollary of the theorem, the maps constructed by Chen-Li [CL2] and by Foscolo [F] can not be tangent maps (c.f [LT], Theorem 3.1) of a stationary quaternionic map satisfing $d(u^*\mathcal{J}^{\alpha}) = 0$.

2 The proof of the blow-up formula

If u is a strong limit of a sequence of stationary quaternionic maps in $H^{1,2}(M, N)$, then it's easy to see that u satisfies (2). If u is just a weak limit, i.e. there exists a sequence of stationary quaternionic maps u_k satisfying $u_k \to u$ weakly in $H^{1,2}(M, N)$ and $|\nabla u_k|^2 dV \to |\nabla u|^2 dV + \theta H^{m-2}|_{\Sigma}$ in the sense of measure, we prove in this section a formula for the blow-up set $\theta H^{m-2}|_{\Sigma}$ and the limiting map u.

Without loss of generality, we may assume that m = 4. Because Σ is a H^{m-2} -rectifiable set, so we may assume that $\Sigma = \bigcup_{i=0}^{\infty} \Sigma_i, \Sigma_i \cap \Sigma_{i'} = \phi$ if $i \neq i', H^{m-2}(\Sigma_0) = 0, \Sigma_i \subset N_i$ and N_i $(i = 1, 2, \cdots)$ is an (m-2)-dimensional embedded C^1 submanifold of M. It is important that (see p. 61 in [Si]) $T_x \Sigma = T_x N_i$ for H^{m-2} -a.e. $x \in \Sigma_i$.

It is known that $\nu = \theta(x)H^{m-2}\lfloor\Sigma$, where $\theta(x)$ is upper semi-continuous with $\epsilon_0 \leq \theta(x) \leq C(l1)$ for H^{m-2} -a.e. $x \in \Sigma$, C(l1) is a positive constant depending only on M and l1 (cf. [Lin], Lemma 1.6). Since $H^{m-2}(\Sigma) < +\infty$, for any 1 > 0, there exist $\Sigma_1 \subset \Sigma$ and i_0 such that $H^{m-2}(\Sigma_1) < 1$, $\Sigma_1^c = \Sigma \setminus \Sigma_1 = \bigcup_{i=1}^{i_0} \Sigma_i^1$ where $\Sigma_i^1 \subset \Sigma_i$ $(i = 1, \dots, i_0)$ is a bounded closed set. We choose a covering $\{B_{r_n}|n = 1, 2, \dots\}$ of Σ_1 such that $\sum_n r_n^{m-2} < C_1$. Here and in the sequel, C always denotes a uniform constant depending only on M and N.

Suppose that $(x^1, ..., x^4)$ is a local normal coordinate system in $B_{\epsilon}(\Sigma_i^{\delta})$, and that (x^3, x^4) is the corresponding coordinate system in Σ_i , and the matrix expressions of the complex structures are given by (6), (7) and (8).

$$J^{1} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \quad A_{1\beta}\mathcal{J}^{\beta} = \begin{pmatrix} J^{1} & & & \\ & \cdot & & \\ & & \cdot & \\ & & \cdot & \\ & & & J^{1} \end{pmatrix}$$
(6)

$$J^{2} = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}, \quad A_{2\beta}\mathcal{J}^{\beta} = \begin{pmatrix} J^{2} & & \\ & \cdot & & \\ & & \cdot & \\ & & \cdot & \\ & & & J^{2} \end{pmatrix}$$
(7)

$$J^{3} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}, \quad A_{3\beta}\mathcal{J}^{\beta} = \begin{pmatrix} J^{3} & & \\ & \cdot & \\ & & \cdot & \\ & & \cdot & \\ & & \cdot & J^{3} \end{pmatrix}$$
(8)

where $A_{\alpha\beta}\mathcal{J}^{\beta}$ are $4n \times 4n$ -matrices, $A_{\alpha\beta}$ are the entries of a matrix A in SO(3). Then the quaternionic equation is

$$\begin{cases} u_1^1 + u_2^2 + u_3^3 + u_4^4 &= 0\\ u_1^2 - u_2^1 + u_3^4 - u_4^3 &= 0\\ u_1^3 - u_3^1 - u_2^4 + u_4^2 &= 0\\ u_1^4 - u_4^1 - u_3^2 + u_2^3 &= 0\\ u_1^5 + u_2^6 + u_3^7 + u_4^8 &= 0\\ u_1^6 - u_2^5 + u_3^8 - u_4^7 &= 0\\ u_1^7 - u_3^5 - u_2^8 + u_4^6 &= 0\\ u_1^8 - u_5^4 - u_3^6 + u_2^7 &= 0\\ \dots \\ u_1^{4n-3} + u_2^{4n-2} + u_3^{4n-1} + u_4^{4n} &= 0\\ u_1^{4n-2} - u_2^{4n-3} + u_3^{4n} - u_4^{4n-1} &= 0\\ u_1^{4n-1} - u_3^{4n-3} - u_2^{4n} + u_4^{4n-2} &= 0\\ u_1^{4n} - u_4^{4n-3} - u_3^{4n-2} + u_4^{4n-1} &= 0. \end{cases}$$
(9)

Theorem 2.1 For any smooth (m-3)-form η with compact support in M, we have

$$\lim_{k \to \infty} \sum_{\alpha=1}^{3} A_{\alpha\beta} \int_{M} d\eta \wedge u_{k}^{*} \mathcal{J}^{\beta} = \sum_{\alpha=1}^{3} A_{\alpha\beta} \int_{M} d\eta \wedge u^{*} \mathcal{J}^{\beta} + \int_{\Sigma} \theta d\eta|_{\Sigma}$$

and

$$\lim_{k \to \infty} A_{1\beta} \int_M d\eta \wedge u_k^* \mathcal{J}^\beta = A_{1\beta} \int_M d\eta \wedge u^* \mathcal{J}^\beta,$$
$$\lim_{k \to \infty} A_{3\beta} \int_M d\eta \wedge u_k^* \mathcal{J}^\beta = A_{3\beta} \int_M d\eta \wedge u^* \mathcal{J}^\beta,$$

Proof. Assume that $\eta = \sum_{I} \eta_{I} dx^{I}$. We have

$$\lim_{k \to \infty} \int_{M} d\eta \wedge u_{k}^{*}(A_{\alpha\beta}\mathcal{J}^{\beta}) = \int_{M} d\eta \wedge u^{*}(A_{\alpha\beta}\mathcal{J}^{\beta}) + \lim_{\delta \to 0} \lim_{\epsilon \to 0} \lim_{k \to \infty} \int_{B_{\epsilon}(\cup_{i=1}^{i_{0}} \Sigma_{i}^{\delta})} d\eta \wedge u_{k}^{*}(A_{\alpha\beta}\mathcal{J}^{\beta}) + \lim_{\delta \to 0} \lim_{\epsilon \to 0} \lim_{k \to \infty} \int_{\cup_{n} B_{r_{n}} \setminus B_{\epsilon}(\cup_{i=1}^{i_{0}} \Sigma_{i}^{\delta})} d\eta \wedge u_{k}^{*}(A_{\alpha\beta}\mathcal{J}^{\beta}).$$
(10)

It's easy to see that

$$\lim_{\delta \to 0} \lim_{k \to \infty} \lim_{k \to \infty} \int_{\bigcup_n B_{r_n}} d\eta \wedge u_k^*(\mathcal{J}^\beta) = 0$$
(11)

By Lemma 2.2 in [LT], we get

$$\lim_{\delta \to 0} \lim_{\epsilon \to 0} \lim_{k \to \infty} \int_{B_{\epsilon}(\Sigma_{i}^{\delta})} d\eta \wedge u_{k}^{*}(A_{\alpha\beta}\mathcal{J}^{\beta})$$

$$= \lim_{\delta \to 0} \lim_{\epsilon \to 0} \lim_{k \to \infty} \int_{B_{\epsilon}(\Sigma_{i}^{\delta})} 2\frac{\partial \eta_{I}}{\partial x^{l}} \frac{\partial u_{k}^{\sigma}}{\partial x^{1}} (A_{\alpha\beta}\mathcal{J}^{\beta})_{\sigma\gamma} \frac{\partial u_{k}^{\gamma}}{\partial x^{2}} dx^{l} \wedge dx^{1} \wedge dx^{2} \quad (12)$$

Substituting (9) to (12) and applying Lemma 2.2 in [LT], we have

$$\lim_{\delta \to 0} \lim_{\epsilon \to 0} \lim_{k \to \infty} \int_{B_{\epsilon}(\Sigma_{i}^{\delta})} d\eta \wedge u_{k}^{*}(A_{1\beta}\mathcal{J}^{\beta}) = \lim_{\delta \to 0} \lim_{\epsilon \to 0} \lim_{k \to \infty} \int_{B_{\epsilon}(\Sigma_{i}^{\delta})} d\eta \wedge u_{k}^{*}(A_{3\beta}\mathcal{J}^{\beta}) = 0$$

and

$$\lim_{\delta \to 0} \lim_{\epsilon \to 0} \lim_{k \to \infty} \int_{B_{\epsilon}(\Sigma_{i}^{\delta})} d\eta \wedge u_{k}^{*}(A_{2\beta}\mathcal{J}^{\beta}) = \lim_{\delta \to 0} \lim_{\epsilon \to 0} \lim_{k \to \infty} \int_{B_{\epsilon}(\Sigma_{i}^{\delta})} |\nabla u_{k}|^{2} d\eta \wedge dx^{1} \wedge dx^{2}$$
$$= \lim_{\delta \to 0} \lim_{\epsilon \to 0} (\int_{B_{\epsilon}(\Sigma_{i}^{\delta})} |\nabla u|^{2} d\eta \wedge dx^{1} \wedge dx^{2} + \int_{B_{\epsilon}(\Sigma_{i}^{\delta}) \cap \Sigma} \theta d\eta|_{\Sigma}) = \int_{\Sigma_{i}} \theta d\eta|_{\Sigma} \quad (13)$$

Then the proof of the theorem is completed.

Remark 2.2 From this theorem, we see that if u_k satisfies (2), the weak limit u still satisfies (2) if and only if $\theta = \text{constant}$.

As a corollary, we can derive that $\theta(x)$ is locally constant. Precisely,

Corollary 2.3 Under the assumption of Theorem 1.2, and assume that there is an open ball $B^m \subset M \setminus \text{Sing}_u$ with $H^{m-2}(\Sigma \cap B^m) > 0$. We have $\theta(x)$ is constant on $\Sigma \cap B^m$.

Proof. In (5), we choose cutoff function η such that $\operatorname{supp} \eta \subset B^m$. Since $B^m \subset M \setminus \operatorname{Sing}_u$, we have u is smooth on B^m . Then $du^* \mathcal{J}^\beta = 0$ on B^m for $\beta = 1, 2, 3$. In view of (5), we conclude that θ is constant on $\Sigma \cap B^m$.

Q.E.D.

Q.E.D.

Let $\phi : \mathbb{S}^2 \to N$ be a nonconstant smooth map satisfying (3) and (4). Set

$$u(x, x^4) = \phi(\frac{x}{|x|}) \quad \text{for any} \quad x \in \mathbb{R}^3 \setminus \{0\} \quad x^4 \in \mathbb{R}^{m-3}$$
(14)

as Chen-Li ([CL2]) did. Then we have

Proposition 2.4 For any smooth (m-3)-form η with compact support in \mathbb{R}^m , we have

$$\int_{\mathbb{R}^m} d\eta \wedge u^* \mathcal{J}^\alpha = -E_T^\alpha(\phi) \int_{\mathbb{R}^{m-3}} \eta(0, x^4), \tag{15}$$

where

$$E_T(\phi) = \int_{\mathbb{S}^2} \langle J^{\alpha}_{\mathbb{S}^2}, u^* \mathcal{J}^{\alpha} \rangle d\sigma.$$

Proof. We choose a spherical coordinate system (r, φ, θ) in \mathbb{R}^3 , because u is smooth for any r > 0, we have

$$\int_{\mathbb{R}^m} d\eta \wedge u^* \mathcal{J}^{\alpha}$$

$$= \int_{\mathbb{R}^{m-3}} \int_0^\infty \frac{\partial \eta_I}{\partial r} dr \wedge dx^I \int_{\mathbb{S}^2} \phi^* \mathcal{J}^{\alpha}$$

$$= -\int_{\mathbb{R}^{m-3}} \eta(0, x^4) \int_{\mathbb{S}^2} \phi^* \mathcal{J}^{\alpha}$$

$$= -E_T^{\alpha}(\phi) \int_{\mathbb{R}^{m-3}} \eta(0, x^4)$$

Q.E.D.

By Theorem 2.1 and Proposition 2.4, we have the following corollary.

Corollary 2.5 The map u defined in (14) can not be a tangent map (c.f [LT], Theorem 3.1) of a stationary quaternionic map with the property (2) at a singular point.

Proof. Suppose that u is defined as in (14). If it is a tangent map, then we have by Theorem 2.1,

$$\sum_{\alpha=1}^{3} A_{\alpha\beta} \int_{M} d\eta \wedge u^{*} \mathcal{J}^{\beta} + \int_{\Sigma} \theta d\eta|_{\Sigma} = 0.$$

By Proposition 2.4, we obtain

$$\sum_{\alpha=1}^{3} A_{\alpha\beta} E_T^{\beta}(\phi) \int_{\mathbb{R}^{m-3}} \eta(0, x^4) = \int_{\Sigma} \theta d\eta|_{\Sigma}.$$

Since u is stationary, by the blow-up formula of Li-Tian [LT], we have Σ is stationary. Using the constancy theorem (Theorem 41.1 in [Si]), it follows that the density function θ is constant in every connected component of Σ , which implies that ϕ is homotopy to a constant map. We therefore get a contradiction.

Q.E.D.

REFERENCES

[BT] C. Bellettini and G. Tian, Compactness results for triholomorphic maps, J. Eur. Math. Soc., 2(2019), 1271-1317.

- [Ch] J. Chen, Complex anti-self-dual connections on product of Calabi-Yau surfaces and triholomorphic curves, Commun. Math. Phys. 201(1999), 201-247.
- [CL1] J. Chen and J. Li, Quaternionic maps between Hyperkähler manifolds, J. Diff. Geom. 55(2000), no. 2, 355-384.
- [CL2] J. Chen and J. Li, Quarternionic maps and minimal surfaces, Ann. Sc. Norm. Super. Pisa Cl. Sci. 4 (2005), no. 3, 375-388.
- [FKS] J.M. Figuroa-O'Farrill, C. Köhl and B. Spence, Supersymmetric Yang-Mills, octonionic instantons and triholomorphic curves, Nucl. Phys. B 521 (1998) no. 3, 419-443.
- [F] L. Foscolo, ALF gravitational instantons and collapsing Ricci-flat metrics on the K3 surface, J. Diff. Geom., 112(2019), 79-120.
- [LT] J. Li, and G. Tian, A blow-up formula for stationary harmonic maps, IMRN, 14(1998), 735-755.
- [Lin] F.-H. Lin, Gradient estimates and blow-up analysis for stationary harmonic maps I, Ann. of Math. 149(1999), 785-829.
- [Si] L. Simon, Lectures on Geometric Measure Theory, Proc. Center Math. Anal. 3(1983), Australian National Univ. Press.
- [W] C. Wang, Energy quantization for triholomorphic maps, Calc. Var. PDE 18(2003), 145-158.