CURVATURE BOUND OF DYSON BROWNIAN MOTION

KOHEI SUZUKI

ABSTRACT. We show that a differential structure associated with the infinite particle Dyson Brownian motion satisfies the Bakry-Émery nonnegative lower Ricci curvature bound $\mathsf{BE}(0,\infty)$. Various functional inequalities follow including a local spectral gap inequality, the Lipschitz Feller property, the dimension-free Harnack inequality and the evolutional variation inequality. As a consequence, the infinite Dyson Brownian motion is characterised as a gradient flow of the Boltzmann-Shannon entropy associated with sine_{β} point processes with respect to a certain Benamou-Brenier-like Wasserstein distance. At the end, we provide a sufficient condition for $\mathsf{BE}(K,\infty)$ beyond the Dyson model and apply it to the infinite particle diffusions corresponding to the β -Riesz gas.

CONTENTS

1.	Introduction	1
2.	Notation and Preliminaries	8
3.	Curvature bound for finite-particle systems	15
4.	Curvature bound for infinite-particle systems	18
5.	Dimension-free and log Harnack inequalities	29
6.	Gradient flow	31
7.	Generalisation	34
Appendix A.		38
References		40

1. INTRODUCTION

The objective of this article is to reveal a *curvature bound* of a differential structure behind infinite particle systems with long-range interactions.

Infinite Dyson Brownian motion. One of the interacting particle systems studied in this article can be formally described as the following stochastic differential equation of infinitely many particles:

(1.1)
$$\mathrm{d}X_t^k = \frac{\beta}{2} \lim_{r \to \infty} \sum_{i \neq k: |X_t^k - X_t^i| < r} \frac{1}{X_t^k - X_t^i} \mathrm{d}t + \mathrm{d}B_t^k, \quad k \in \mathbb{N} \; ,$$

 $^{2020\} Mathematics\ Subject\ Classification.\ {\rm Primary}\ 60{\rm K}35,\ {\rm Secondary}\ 31{\rm C}25.$

Key words and phrases. Dyson Brownian motion, log-gas, Ricci curvature bound.

Department of Mathematical Science, Durham University E-mail: kohei.suzuki@durham.ac.uk .

whereby $\{B_{\cdot}^{k}\}_{k\in\mathbb{N}}$ are infinitely many independent Brownian motions on \mathbb{R} . The solution to (1.1) is called *infinite Dyson Brownian motion*, which has a particular importance in relation to random matrix theory (see [Dys62, Spo87, NF98, KT10, Osa96, Osa12, Osa13, Tsa16]). The infinite interacting diffusions (1.1) can be thought of as *a single* diffusion process on *the configuration spcae* $\Upsilon = \Upsilon(\mathbb{R})$ over \mathbb{R} (i.e., the space of locally finite point measures on \mathbb{R}). This diffusion process on Υ has an invariant measure μ , called sine_{β} ensemble (see § 2.7), that is known as the universal limit of the eigenvalue distributions of Gaussian random matrices.

Differential structure of interacting particles. The infinite Dyson Brownian motion (1.1) induces a differential structure on the infinite-dimensional space Υ , called *Dirichlet form*, that is, a closed symmetric bilinear form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ on $L^2(\Upsilon,\mu)$ satisfying a maximum principle (called Markovian property). The Dirichlet form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ corresponding to (1.1) is described as follows (see Dfn. 4.15):

$$\mathcal{E}^{\Upsilon,\mu}(u) := \frac{1}{2} \int_{\Upsilon} \Gamma^{\Upsilon}(u) \,\mathrm{d}\mu , \quad u \in \mathcal{D}(\mathcal{E}^{\Upsilon,\mu})$$

where Γ^{Υ} corresponds to the (infinite-dimensional) squared gradient operator on Υ called square field. The transition semigroup of (1.1) is associated with the L^2 -semigroup $\{T_t^{\Upsilon,\mu}\}_{t\geq 0}$ induced by the Dirichlet form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$. In other words, $\{T_t^{\Upsilon,\mu}\}_{t\geq 0}$ is the L^2 -gradient flow of $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$.

Bakry-Émery curvature bound. In the seminal paper Bakry-Émery [BÉ84], they observed that a complete Riemannian manifold (M, g) has a Ricci curvature lower bound Ric $\geq K$ for some $K \in \mathbb{R}$ if and only if the gradient estimate

$$(\mathsf{BE}(K,\infty)) \qquad \qquad |\nabla T_t u|^2 \le e^{-2Kt} T_t |\nabla u|^2 \ , \quad u \in W^{1,2}(M)$$

holds in terms of the heat semigroup $\{T_t\}_{t\geq 0}$, the gradient operator ∇ and the (1,2)-Sobolev space $W^{1,2}(M)$ on M. This observation opened a way to generalise the concept of lower Ricci curvature bound to singular spaces beyond manifolds such as metric measure spaces and infinite-dimensional spaces since the latter formulation $\mathsf{BE}(K,\infty)$ requires only a weak (Sobolev) differentiable structure, which does not require *Ricci curvature tensors* nor a C^2 -structure. This generalised concept of the lower Ricci curvature bound turned out to be very powerful and many functional inequalities and quantitative controls of geometry and analysis have been revealed as a consequence of $\mathsf{BE}(0,\infty)$. We refer the readers to e.g., [BGL14] and [Vil09] for comprehensive references.

Main results. The main result of this article is to show that the differentiable structure $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ induced by the infinite Dyson SDE (1.1) has the non-negative Ricci curvature bound "Ric ≥ 0 " in the sense of Bakry–Émery.

Theorem 1.1 (Thm. 4.19). Let $\beta > 0$ and μ be the sine_{β} ensemble. The form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ satisfies the Bakry-Émery estimate $\mathsf{BE}(0,\infty)$. Namely,

$$\Gamma^{\Upsilon}(T_t^{\Upsilon,\mu}u) \leq T_t^{\Upsilon,\mu}\Gamma^{\Upsilon}(u) , \quad u \in \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}) \quad t > 0 .$$

Theorem 1.1 says that the configuration space Υ endowed with the differential structure $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ can be understood as a "non-negatively curved" space. We note that the

curvature bound "Ric ≥ 0 " does not depend on β . We believe that the choice of β cannot improve the bound and $\mathsf{BE}(0,\infty)$ gives the optimal bound for every $\beta > 0$ (see *discussion for the best K* at the end of this introduction for more details).

Applications to the Dyson SDE (1.1). The Bakry–Émery lower curvature bound $\mathsf{BE}(0,\infty)$ brings various functional inequalities as byproducts, which provide applications to the corresponding infinite Dyson SDE (1.1). We start with the local Poincaré inequality, which bring a *local* spectral gap estimate of the corresponding particle dynamics (1.1) (see Rem. 4.21 for more details).

Corollary 1.2 (Cor. 4.20). Let $\beta > 0$ and μ be the sine_{β} ensemble. Then, the Local Poincaré inequality holds: for $u \in \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}), t > 0$,

$$\begin{split} T_t^{\Upsilon,\mu} u^2 &- (T_t^{\Upsilon,\mu} u)^2 \leq 2t T_t^{\Upsilon,\mu} \Gamma^{\Upsilon}(u) \ , \\ T_t^{\Upsilon,\mu} u^2 &- (T_t^{\Upsilon,\mu} u)^2 \geq 2t \Gamma^{\Upsilon}(T_t^{\Upsilon,\mu} u) \ . \end{split}$$

Optimal transport and the Dyson Brownian motion. The configuration space Υ has a metric structure lifted from the base space \mathbb{R} , called the L^2 -transportation distance d_{Υ} (or L^2 -optimal matching distance), where the cost function is the squared Euclidean distance $\mathsf{d}_{\mathbb{R}}^2$ (see (2.18)). We use a variant $\bar{\mathsf{d}}_{\Upsilon}$ of d_{Υ} called the L^2 -transportation-like distance (see (2.19)). We prove that the metric $\bar{\mathsf{d}}_{\Upsilon}$ has a consistency with the differential structure ($\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu})$) induced by the infinite Dyson SDE (1.1).

Theorem 1.3 (Prop. 4.16). Let $\beta > 0$ and μ be the sine_{β} ensemble. Then the Rademachertype property holds:

$$\operatorname{Lip}_{b}(\bar{\mathsf{d}}_{\Upsilon},\mu) \subset \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}) , \quad \Gamma^{\Upsilon,\mu}(u) \leq \operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(u)^{2}$$

where $\operatorname{Lip}_b(\bar{\mathsf{d}}_{\mathbf{\Upsilon}},\mu)$ denotes the space of bounded μ -measurable $\bar{\mathsf{d}}_{\mathbf{\Upsilon}}$ -Lipschitz functions and $\operatorname{Lip}_{\bar{\mathsf{d}}_{\mathbf{\Upsilon}}}(u)$ denotes the $\bar{\mathsf{d}}_{\mathbf{\Upsilon}}$ -Lipschitz constant of u.

Combined with the local Poincaré inequality, we have the following exponential decay estimate of the heat kernel measure $P_t^{\Upsilon,\mu}(\gamma, d\eta)$ (i.e., the transition probability of (1.1)) in terms of \bar{d}_{Υ} .

Corollary 1.4 (Cor. 4.22). Let $\beta > 0$ and μ be the sine β ensemble. If u is a \bar{d}_{Υ} -Lipschitz μ -measurable function with $\operatorname{Lip}_{\bar{d}_{\Upsilon}}(u) \leq 1$ and $|u(\gamma)| < \infty \mu$ -a.e. γ , then for every $s < \sqrt{2/t}$

$$\int_{\Upsilon} e^{su(\eta)} P_t^{\Upsilon,\mu}(\gamma,\mathrm{d}\eta) < \infty$$

Curvature bound in terms of the metric $\bar{\mathsf{d}}_{\Upsilon}$. In the case of Riemannian manifolds (M, g), the Ricci curvature lower bound Ric $\geq K$ is known to be equivalent to the dimension-free Harnack inequality ([Wan14, Thm. 2.3.3]): for $\alpha > 1$ and every bounded Borel function $u \geq 0$ on M

$$(T_t u)^{\alpha}(x) \le T_t u^{\alpha}(y) \exp\left\{\frac{\alpha K}{2(\alpha - 1)(1 - e^{-2Kt})} \mathsf{d}_g(x, y)^2\right\},$$

where d_g is the geodesic distance induced by g. This provides a characterisation of Ric $\geq K$ in terms of the metric d_g . In the following theorem, we prove that the dimension-free

Harnack inequality with K = 0 holds true for the infinite Dyson SDE (1.1) with respect to the L^2 -optimal transport-type distance $\bar{\mathsf{d}}_{\Upsilon}$. We furthermore prove the log-Harnack inequality, and the Lipschitz contraction estimate by $T_t^{\Upsilon,\mu}$, the latter of which can be understood as a metric counter-part of $\mathsf{BE}(0,\infty)$. As a byproduct, we obtain the Lipschitz regularisation property (Lipschitz-Feller property) of the semigroup $T_t^{\Upsilon,\mu}$.

Theorem 1.5 (Thm. 5.1). Let $\beta > 0$ and μ be the sine_{β} ensemble. Then, the following hold:

 Dimension-free Harnack inequality: for every non-negative u ∈ L[∞](Υ, μ), t > 0 and α > 1 there exists Ω ⊂ Υ so that μ(Ω) = 1 and

$$(T_t^{\Upsilon,\mu}u)^{\alpha}(\gamma) \leq T_t^{\Upsilon,\mu}u^{\alpha}(\eta) \exp\left\{\frac{\alpha}{2(\alpha-1)}\bar{\mathsf{d}}_{\Upsilon}(\gamma,\eta)^2\right\}\,, \quad \forall \gamma, \eta \in \Omega\,;$$

Log-Harnack inequality: for any non-negative u ∈ L[∞](Υ, μ), ε ∈ (0,1], t > 0, there exists Ω ⊂ Υ so that μ(Ω) = 1 and

$$T_t^{\Upsilon,\mu}\log(u+\varepsilon)(\gamma) \leq \log(T_t^{\Upsilon,\mu}u(\eta)+\varepsilon) + \bar{\mathsf{d}}_{\Upsilon}(\gamma,\eta)^2 , \quad \forall \gamma,\eta \in \Omega ;$$

• Lipschitz contraction: for every $u \in \operatorname{Lip}(\bar{\mathsf{d}}_{\Upsilon}, \mu)$ and t > 0

 $T_t^{\Upsilon,\mu}u$ has a $\bar{\mathsf{d}}_{\Upsilon}$ -Lipschitz μ -modification $\tilde{T}_t^{\Upsilon,\mu}u$

and the following Lipschitz contraction holds:

$$\operatorname{Lip}_{\bar{\mathsf{d}}_{\boldsymbol{\Upsilon}}}(\tilde{T}_t^{\boldsymbol{\Upsilon},\mu}u) \leq \operatorname{Lip}_{\bar{\mathsf{d}}_{\boldsymbol{\Upsilon}}}(u) ;$$

• $L^{\infty}(\mu)$ -to-Lip $(\bar{d}_{\Upsilon}, \mu)$ Feller property: For $u \in L^{\infty}(\mu)$ and t > 0,

$$T_t^{\Upsilon,\mu}u$$
 has a $\bar{\mathsf{d}}_{\Upsilon}$ -Lipschitz μ -modification $\tilde{T}_t^{\Upsilon,\mu}u$

and the following estimate holds:

$$\operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(\tilde{T}_t^{\Upsilon,\mu}u) \leq \frac{1}{\sqrt{2t}} \|u\|_{L^{\infty}(\mu)} .$$

Dyson Brownian motions as a gradient flow. Jordan, Kinderlehrer and Otto [JKO98] discovered a class of partial differential equations that can be realised as gradient flows in the space (\mathcal{P}_2, W_2) of probability measures with finite second moment endowed with the L^2 -Monge-Kantrovich-Rubinstein-Wasserstein distance W_2 . In particular, the dual flow of the heat equation in the Euclidean space \mathbb{R}^n , where the corresponding diffusion process is the Brownian motion in \mathbb{R}^n , is characterised as the W_2 -gradient flow

$$\partial_t \nu = -\nabla_{\mathsf{W}_2} \mathsf{Ent}(\nu)$$

of the Boltzmann-Shannon entropy $\operatorname{Ent}(\nu) = \int_{\mathbb{R}^n} \rho \log \rho \, dx$ with $d\nu = \rho \, dx$. Here, the W₂-gradient flow is defined as the energy dissipation equality:

(1.2)
$$\frac{\mathrm{d}}{\mathrm{d}t}\mathsf{Ent}(\nu_t) = -|\dot{\nu}_t|^2 = -|\mathsf{D}_{\mathsf{W}_2}^-\mathsf{Ent}|^2(\nu_t) \quad \text{a.e. } t > 0 \; .$$

where $|\dot{\nu}_r|$ denotes the metric speed of the curve (ν_r) and $|\mathsf{D}_{\mathsf{W}_2}^-\mathsf{Ent}|$ is the descending slope of Ent with respect to W_2 , see §2.3. The result in [JKO98] brought a new perspective of the Brownian motion in \mathbb{R}^n as a *steepest descent* of the Boltzmann-Shannon entropy with respect to W_2 . Exploiting Thm. 1.1, we can extend this perspective to the case of infinite Dyson Brownian motions in terms of the Boltzmann-Shannon entropy $\operatorname{Ent}_{\mu}(\nu) = \int_{\Upsilon} \rho \log \rho \, d\mu$ associated with the reference measure $\mu = \operatorname{sine}_{\beta}$ and a Benamou–Brenier-like extended distance $W_{\mathcal{E}}$, see (6.3). Let $\mathcal{P}(\Upsilon)$ be the space of all Borel probability measures in Υ and $\mathcal{D}(\operatorname{Ent}_{\mu}) := \{\nu \in \mathcal{P}(\Upsilon) : \operatorname{Ent}_{\mu}(\nu) < \infty\}$ be the domain of Ent_{μ} . Let $t \mapsto \mathcal{T}_{t}^{\Upsilon,\mu}\nu$ be the dual flow of the transition semigroup $T_{t}^{\Upsilon,\mu}$ of the infinite Dyson Brownian motion:

$$\mathcal{T}_t^{\Upsilon,\mu}\nu := (T_t^{\Upsilon,\mu}\rho)\cdot\mu , \quad \nu = \rho\cdot\mu\in\mathcal{P}(\Upsilon) .$$

Corollary 1.6 (Cor. 6.4). Let μ be the sine_{β} ensemble with $\beta > 0$.

• Evolutional variation inequality: For every $\nu, \sigma \in \mathcal{D}(\mathsf{Ent}_{\mu})$ with $\mathsf{W}_{\mathcal{E}}(\nu, \sigma) < \infty$,

$$\frac{1}{2} \frac{\mathrm{d}^+}{\mathrm{d}t} \mathsf{W}_{\mathcal{E}} \big(\mathcal{T}_t^{\Upsilon,\mu} \sigma, \nu \big)^2 \leq \mathsf{Ent}_{\mu}(\nu) - \mathsf{Ent}_{\mu}(\mathcal{T}_t^{\Upsilon,\mu} \sigma) \;, \quad t > 0 \;.$$

Geodesical convexity: The space (D(Ent_μ), W_ε) is an extended geodesic metric space. Namely, for every pair ν, σ ∈ D(Ent_μ) with W_ε(ν, σ) < ∞, there exists W_ε-Lipschitz curve ν. : [0, 1] → (D(Ent_μ), W_ε) so that

$$\nu_0 = \nu , \quad \nu_1 = \sigma , \quad \mathsf{W}_{\mathcal{E}}(\nu_t, \nu_s) = |t - s| \mathsf{W}_{\mathcal{E}}(\nu, \sigma) , \quad s, t \in [0, 1]$$

• Gradient flow: The dual flow $(\mathcal{T}_t^{\Upsilon,\mu}\nu_0)_{t>0}$ is the unique solution to the $W_{\mathcal{E}}$ -gradient flow of Ent_{μ} starting at ν_0 . Namely, for any $\nu_0 \in \mathcal{D}(\mathsf{Ent}_{\mu})$, the curve $[0,\infty) \ni t \mapsto \nu_t = \mathcal{T}_t^{\Upsilon,\mu}\nu_0 \in \mathcal{D}(\mathsf{Ent}_{\mu})$ is the unique solution to the energy equality:

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathsf{Ent}_{\mu}(\nu_t) = -|\dot{\nu}_t|^2 = -|\mathsf{D}_{\mathsf{W}_{\mathcal{E}}}^-\mathsf{Ent}_{\mu}|^2(\nu_t) \quad a.e. \ t > 0 \ .$$

Generalisation beyond $\operatorname{sine}_{\beta}$. At the end of this article, our results will be extended to $\mathsf{BE}(K,\infty)$ with $K \in \mathbb{R}$ for general point processes satisfying conditional geodesical Kconvexity, see Thm. 7.2. We apply Thm. 7.2 to prove $\mathsf{BE}(0,\infty)$ of the Dirichlet form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ with β -Riesz emsemble $\mu = \mu_{\beta}$ for $\beta > 0$.

Corollary 1.7 (Cor. 7.8). The Dirichlet form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ with the β -Riesz ensemble μ satisfies $\mathsf{BE}(0,\infty)$ for $\beta > 0$. Furthermore, all the statements in Thms. 1.1, 1.3, 1.5, Cors. 1.2, 1.4, 1.6 hold true in this case.

Comparison with Literature. To the author's best knowledge, this is the first article addressing the lower Ricci curvature bound on Υ under the presence of interactions. Even with a simpler interaction potential like compactly supported smooth pair potential with Ruelle condition, no result regarding the curvature bound has been known so far. In the non-interacting case where the invariant measure is the Poisson measure, the synthetic lower Ricci curvature bound has been established in the pioneering work [EH15] in the case where the base space is Riemannian manifolds, and in [DS22] in the case where the base space is general diffusion spaces. In [EHJM23], a specific entropy associated with the Poisson point processes and an optimal transport distance have been introduced in the space of stationary point processes. They established the evolutional variation inequality, the gradient flow property, the displacement convexity and the HWI inequality for the flows induced by independent Brownian particles starting at stationary measures. In the case of

finite particle systems, a variable Ricci curvature bound has been addressed in [GV20] for Coulomb-type potentials.

Up until now, only little is understood about the transition probability of interacting *infinite* particle diffusions. In particular so far, almost nothing is known about *quantitative* estimates of the transition semigroup corresponding to the infinite particle Dyson models. The functional inequalities in Cor. 1.2, Thm. 1.5 and the exponential decay estimate of the transition semigroup in Cor. 1.4 contribute to the quantitative side of the Dyson Brownian motion.

Furthermore, the dimension-free Harnack inequality in Thm. 1.5 provides quantitative estimates of the transition semigroup of the Dyson SDE (1.1) in term of the metric structure $\bar{\mathbf{d}}_{\Upsilon}$, which could give a new approach to study the Dyson SDEs in a geometric manner. We note that [KS21] provided an equivalence between a synthetic lower Ricci curvature bound (what is called RCD condition) and the Wang's dimension Harnack inequality in a framework of metric measure spaces. We cannot however apply their result to our setting because (a) we do not know if $(\Upsilon, \bar{\mathbf{d}}_{\Upsilon}, \mu)$ is an RCD space; (b) $(\Upsilon, \bar{\mathbf{d}}_{\Upsilon}, \mu)$ is not a metric measure space due to the fact that $\bar{\mathbf{d}}_{\Upsilon}$ does not generate the given topology (the vague topology) on Υ and $\bar{\mathbf{d}}_{\Upsilon}$ takes $+\infty$ on sets of positive measure with respect to μ . We therefore prove the dimension-free Harnack inequality through a finite-particle approximation.

On the qualitative side, we reveal in Cor. 1.6 that the infinite Dyson Brownian motion is the unique $W_{\mathcal{E}}$ -gradient flow of Ent_{μ} associated with $\mu = \mathsf{sine}_{\beta}$, which provides a new perspective of the Dyson Brownian motion as a steepest descent of the Boltzmann–Shannon entropy associated with sine_{β} ensemble.

As a final remark, we provide a construction of Dirichlet forms with $\operatorname{sine}_{\beta}$ for every $\beta > 0$ (cf., for the case of $\beta = 1, 2, 4$ in [Osa96, Osa13]). It should be noted that the domain of the Dirichlet form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ in this paper is different from the one constructed in [Osa96, Osa13] even for $\beta = 1, 2, 4$, due to the difference of the domain of the approximating form $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}))$ (see Dfn. 4.3 and Rem. 4.4). The domain used in this paper is tailor made to show BE $(0, \infty)$. We do not know if we can show BE $(0, \infty)$ with the domains used in [Osa96, Osa13]. The identification of these domains remains open. We note that the uniqueness result in [KOT21] cannot be directly applied to the setting of this paper since our domain $\mathcal{D}(\mathcal{E}^{\Upsilon,\mu})$ is not contained in their domain (called lower Dirichlet form).

Discussion for the best possible K. The curvature lower bound K = 0 in Thm. 1.1 does not depend on the inverse temperature β . One might wonder if there is a positive constant $K_{\beta} > 0$ depending on β so that the sharper curvature bound $\mathsf{BE}(K_{\beta}, \infty)$ holds. However, we do not believe the existence of such a positive constant and we believe that K = 0 is the best constant for the lower bound. This is due to the fact that the logarithmic potential $-\beta \log |x - y|$ cannot be K-convex with positive K > 0 for any $\beta > 0$, which indicates that the choice of β could not improve the curvature bound in the infinite-volume case. Indeed, the Hessian matrix of the Hamiltonian in the DLR equation is computed in (3.7), which suggests that the Hessian cannot be bounded from below by a fixed positive constant as the radii r, R go to infinity.

We stress that even with the bound K = 0, various new quantitative results regarding the transition probability of the infinite Dyson Brownian motion (1.1) follow as explained above. Furthermore, the bound K = 0 is enough to identify the Dyson Brownian motion with the gradient flow of the Boltzmann–Shannon entropy $\mathsf{Ent}_{\mathsf{sine}_\beta}$ in the space of probability measures as in Cor. 1.6, which sheds light on a new perspective of the Dyson Brownian motion.

Outlook for further study. In Thm. 7.2, we provide a sufficient condition for the Bakry– Émery lower Ricci curvature bound $\mathsf{BE}(K,\infty)$ in the case of general point processes. The sine_{β} ensemble, the β -Riesz ensemble and the Poisson ensemble are currently only the examples for which one can verify this sufficient condition. It would be interesting as a further study to explore the curvature bound for other 1-dimensional point processes such as β -Airy or Bessel ensembles. In this direction, one of the important points is to know the exact information of the Gibbs specification (i.e., DLR equation).

Outline of the article. In Section 2, the notation and the preliminary materials are presented. In Section 3, we discuss the lower Ricci curvature bound of finite particle systems, i.e., we discuss the Dirichlet forms

(1.3)
$$(\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}, \mathcal{D}(\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}))$$

on the configuration space $\Upsilon(B_r)$ over the closed metric ball B_r with radius r > 0 centred at 0, whose invariant measure is the projected regular conditional probability μ_r^{η} on $\Upsilon(B_r)$ conditioned at η on the compliment $B_r^c \subset \mathbb{R}$. The key point for the lower Ricci curvature bound of (1.3) is the geodesical convexity of the corresponding Hamiltonian on $(\Upsilon(B_r), \bar{\mathsf{d}}_{\Upsilon})$, i.e., the logarithm of the Radon–Nikodým density $\Psi_r^{\eta} := -\log(\mathrm{d}\mu_r^{\eta}/\mathrm{d}\pi_{\mathsf{m}_r})$, where π_{m_r} denotes the Poisson measure on $\Upsilon(B_r)$ with the intensity measure m_r being the Lebesgue measure restricted on B_r . This convexity is due to the following DLR (Dobrushin–Lanford– Ruelle) equation proven in [DHLM20, Thm.1.1]: for μ -a.e. η , there exists a unique $k = k(\eta) \in \mathbb{N}_0$ so that

$$\begin{split} \mathrm{d}\mu_r^{\eta} &= \frac{1}{Z_r^{\eta}} e^{-\Psi_r^{k,\eta}} \,\mathrm{d}\mathsf{m}_r^{\odot k} \;, \quad \gamma = \sum_{i=1}^k \delta_{x_i} \in \Upsilon(B_r) \\ \Psi_r^{k,\eta}(\gamma) &:= -\log \left(\prod_{i < j}^k |x_i - x_j|^{\beta} \prod_{i=1}^k \lim_{R \to \infty} \prod_{y \in \eta_{B_r^c}, |y| \le R} \left| 1 - \frac{x_i}{y} \right|^{\beta} \right) \;, \end{split}$$

where $\mathbf{m}_r^{\odot k}$ is the k-symmetric product measure of the Lebesgue measure \mathbf{m}_r restricted on $B_r \subset \mathbb{R}$ and Z_r^{η} is the normalising constant (note that the roles of the notation γ and η in [DHLM20] are opposite to this article).

In Section 4, we prove $\mathsf{BE}(0,\infty)$ of $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ in the following steps: we first construct the truncated form $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}))$ on Υ whose gradient operator is truncated up to configurations on B_r (Prop. 4.8). We then identify it with the superposition Dirichlet form $(\bar{\mathcal{E}}_r^{\Upsilon,\mu}, \mathcal{D}(\bar{\mathcal{E}}_r^{\Upsilon,\mu}))$ lifted from $(\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}, \mathcal{D}(\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}))$ with respect to the conditioning η (Thm. 4.12). By this identification, we can lift $\mathsf{BE}(0,\infty)$ from $(\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}, \mathcal{D}(\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}))$ onto the truncated form $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}))$. By the monotonicity of the form $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}))$ with respect to r and passing to the limit $r \to \infty$, we prove $\mathsf{BE}(0,\infty)$ for the limit form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ (Thm. 4.19). As a consequence of $\mathsf{BE}(0,\infty)$, we obtain the inte-

$$\begin{split} (\bar{\mathcal{E}}_{r}^{\Upsilon,\mu},\mathcal{D}(\bar{\mathcal{E}}_{r}^{\Upsilon,\mu}))^{identification}(\mathcal{E}_{r}^{\Upsilon,\mu},\mathcal{D}(\mathcal{E}_{r}^{\Upsilon,\mu})) \xrightarrow{monotone} (\mathcal{E}^{\Upsilon,\mu},\mathcal{D}(\mathcal{E}^{\Upsilon,\mu})) & \mathsf{BE}(0,\infty) \\ & \uparrow^{superposition} \\ (\mathcal{E}^{\Upsilon(B_{r}),\mu_{r}^{\eta}},\mathcal{D}(\mathcal{E}^{\Upsilon(B_{r}),\mu_{r}^{\eta}})) & \mathsf{BE}(0,\infty) \end{split}$$

FIGURE 1. $\mathsf{BE}(0,\infty)$ is transferred to $\mathcal{E}^{\Upsilon,\mu}$ via the vertical arrow \uparrow , the equality =, and the right arrow \rightarrow .

gral Bochner inequality and the local Poincaré inequality (Cor. 4.20) and an exponential decay of the transition semigroup (Cor. 4.22).

In Section 5, we prove the dimension-free Harnack inequality, the log-Harnack inequality, the Lipschitz contraction and $L^{\infty}(\mu)$ -to-Lip $(\bar{\mathbf{d}}_{\Upsilon})$ Feller properties (Thm. 5.1). In section 6, we discuss the optimal transport theory on Υ and prove that the flow in $\mathcal{P}(\Upsilon)$ induced by the Dyson Brownian motion is the unique gradient flow of the Boltzmann–Shannon entropy Ent_{μ} . In Section 7, we extend Theorem to the case of general point processes beyond sine_{β} (Thm. 7.2) and discuss β -Riesz ensembles.

Acknowledgement. The author expresses his great appreciation to Professor Thomas Leblé for making him aware of the β -Riesz ensemble as an example of Assumption 7.1. A large part of the current work has been completed while he was at Bielefeld University. He gratefully acknowledges funding by the Alexander von Humboldt Stiftung to support his stay.

Data Availability Statement. No datasets were generated or analysed during the current study.

2. NOTATION AND PRELIMINARIES

2.1. Numbers, Tensors, Function Spaces. We write $\mathbb{N} := \{1, 2, 3, ...\}, \mathbb{N}_0 = \{0, 1, 2, ...\}, \overline{\mathbb{N}} := \mathbb{N} \cup \{+\infty\}$ and $\overline{\mathbb{N}}_0 := \mathbb{N}_0 \cup \{+\infty\}$. The uppercase letter N is used for $N \in \overline{\mathbb{N}}_0$, while the lowercase letter n is used for $n \in \mathbb{N}_0$. We shall adhere to the following conventions:

- the superscript $\Box^{\times N}$ (the subscript $\Box_{\times N}$) denotes (N-fold) product objects;
- the superscript $\Box^{\otimes N}$ (the subscript $\Box_{\otimes N}$) denotes (N-fold) tensor objects;
- the superscript $\Box^{\odot N}$ (the subscript $\Box_{\odot N}$) denotes (N-fold) symmetric tensor objects;

Let (X, τ) be a topological space with σ -finite Borel measure ν . We use the following symbols:

(a) $L^p(\nu)$ $(1 \le p \le \infty)$ for the space of ν -equivalence classes of functions u with $|u|^p$ ν -integrable when $1 \le p < \infty$, and with u ν -essentially bounded when $p = \infty$. The $L^p(\nu)$ -norm is denoted by $||u||_{L^p(\nu)}^p := ||u||_p^p := \int_X |u|^p \, d\nu$ for $1 \le p < \infty$, and $||u||_{L^\infty(\nu)} := ||u||_\infty = \text{esssup}_X u$. In the case of p = 2, the inner-product is denoted by $(u, v)_{L^2(\nu)} := \int_X uv \, d\nu$;

- (b) $L_s^p(\nu^{\otimes n}) := \{u \in L^p(\nu^{\otimes n}) : u \text{ is symmetric}\}$ where u is said to be *symmetric* if and only if $u(x_1, \ldots, x_k) = u(x_{\sigma(1)}, \ldots, x_{\sigma(k)})$ for any element $\sigma \in \mathfrak{S}(k)$ in the k-symmetric group.
- (c) $C_b(X)$ for the space of τ -continuous bounded functions on X; if X is locally compact, $C_0(X)$ denotes the space of τ -continuous and compactly supported functions on X; $C_0^{\infty}(\mathbb{R})$ for the space of compactly supported smooth functions on \mathbb{R} ;
- (d) We write $\mathbf{1}_A$ for the indicator function on A, i.e., $\mathbf{1}_A(x) = 1$ if and only if $x \in A$, and $\mathbf{1}_A(x) = 0$ otherwise; δ_x for the Dirac measure at x, i.e., $\delta_x(A) = 1$ if and only if $x \in A$, and $\delta_x(A) = 0$ otherwise;
- (e) For a space of real-valued functions, the subscript + is used for the subspace of non-negative functions. For instance, $C_{b,+}(X) := \{u \in C_b(X) : u \ge 0\}.$

2.2. Dirichlet forms. We refer the reader to [MR90, BH91] for this subsection. Throughout this paper, a Hilbert space always means a separable Hilbert space with inner product $(\cdot, \cdot)_H$ taking value in \mathbb{R} .

Dirichlet forms. Given a bilinear form $(Q, \mathcal{D}(Q))$ on a Hilbert space H, we write

$$Q(u) \coloneqq Q(u, u) , \qquad Q_{\alpha}(u, v) \coloneqq Q(u, v) + \alpha(u, v)_{H} , \ \alpha > 0$$

Let (X, Σ, ν) be a σ -finite measure space. A symmetric Dirichlet form on $L^2(\nu)$ is a non-negative definite densely defined closed symmetric bilinear form $(Q, \mathcal{D}(Q))$ on $L^2(\nu)$ satisfying the Markov property

$$u_0 \coloneqq 0 \lor u \land 1 \in \mathcal{D}(Q)$$
 and $Q(u_0) \le Q(u)$, $u \in \mathcal{D}(Q)$.

Throughout this article, *Dirichlet form* always means symmetric Dirichlet form. If not otherwise stated, $\mathcal{D}(Q)$ is always regarded as a Hilbert space with norm

$$\|\cdot\|_{\mathcal{D}(Q)} \coloneqq Q_1(\cdot)^{1/2} := \sqrt{Q(\cdot) + \|\cdot\|_{L^2(\nu)}^2}$$

In order to distinguish Dirichlet forms defined in different base spaces with different reference measures, we often use the notation $Q^{X,\nu}$ to specify the base space X and the reference measure ν .

Square field. A Dirichlet form $(Q, \mathcal{D}(Q))$ admits square field Γ if there exists a dense subspace $H \subset \mathcal{D}(Q) \cap L^{\infty}(\nu)$ having the following property: for any $u \in H$, there exists $v \in L^{1}(\nu)$ so that

$$2Q(uh, u) - Q(h, u^2) = \int_X hv \, \mathrm{d}\nu \quad \forall h \in \mathcal{D}(Q) \cap L^{\infty}(\nu) \; .$$

Such v is denoted by $\Gamma(u)$. The square field Γ can be uniquely extended as an operator on $\mathcal{D}(Q) \times \mathcal{D}(Q) \to L^1(\nu)$ ([BH91, Thm. I.4.1.3]).

Semigroups and generators. We refer the reader to [MR90, Chap. I, Sec. 2] for the following contents. Let $(Q, \mathcal{D}(Q))$ be a symmetric closed form on a Hilbert space H. The infinitesimal generator $(A, \mathcal{D}(A))$ corresponding to $(Q, \mathcal{D}(Q))$ is the unique densely defined closed operator on H satisfying the following integration-by-parts formula:

$$-(u, Av)_H = Q(u, v) \quad \forall u \in \mathcal{D}(Q), \ v \in \mathcal{D}(A)$$
.

The resolvent operator $\{G_{\alpha}\}_{\alpha\geq 0}$ is the unique bounded linear operator on H satisfying

$$Q_{\alpha}(G_{\alpha}u, v) = (u, v)_H \quad \forall u \in H \quad v \in \mathcal{D}(Q)$$

The semigroup $\{T_t\}_{t>0}$ is the unique bounded linear operator on H satisfying

$$G_{\alpha}u = \int_0^{\infty} e^{-\alpha t} T_t u \,\mathrm{d}t \quad u \in H \;.$$

Locality. Let $(Q, \mathcal{D}(Q))$ be a Dirichlet form on $L^2(\nu)$. It is called *local* ([BH91, Def. 5.1.2 in Chap. I]) if for any $F, G \in C_c^{\infty}(\mathbb{R})$ and any $u \in \mathcal{D}(Q)$,

$$\operatorname{supp}[F] \cap \operatorname{supp}[G] = \emptyset \implies Q(F_0 \circ u, G_0 \circ u) = 0 ,$$

where $F_0(x) := F(x) - F(0)$ and $G_0(x) := G(x) - G(0)$.

2.3. Extended metric space. Let X be any non-empty set. A function $d: X^{\times 2} \to [0, \infty]$ is an extended distance if it is symmetric and satisfying the triangle inequality, and it does not vanish outside the diagonal in $X^{\times 2}$, i.e. d(x, y) = 0 iff x = y; a distance if it is finite. Let $x_0 \in X$ and $r \in [0, \infty)$. We write $B_r(x_0) := \{ d_{x_0} \leq r \}$, where $d_{x_0} := d(x_0, \cdot)$. A space X equipped with an extended distance (resp. a distance) is called an extended metric space (resp. a metric space) (X, d).

Lipschitz algebras. A function $f: X \to \mathbb{R}$ is d-Lipschitz if there exists a constant L > 0 so that

(2.1)
$$|u(x) - u(y)| \le L \operatorname{d}(x, y) , \qquad x, y \in X$$

The smallest constant L so that (2.1) holds is the (global) Lipschitz constant of u, denoted by $\operatorname{Lip}_{\mathsf{d}}(u)$. For any non-empty $A \subset X$, we write $\operatorname{Lip}(A, \mathsf{d})$, resp. $\operatorname{Lip}_b(A, \mathsf{d})$ for the family of all d -Lipschitz functions, resp. bounded d -Lipschitz functions on A. For simplicity of notation, further let

$$\operatorname{Lip}(\mathsf{d}) \coloneqq \operatorname{Lip}(X, \mathsf{d}) , \qquad \operatorname{Lip}_b(\mathsf{d}) \coloneqq \operatorname{Lip}_b(X, \mathsf{d}) .$$

Set also $\operatorname{Lip}^1(\mathsf{d}) := \{ u \in \operatorname{Lip}(\mathsf{d}) : \operatorname{Lip}_{\mathsf{d}}(u) \leq 1 \}$ and $\operatorname{Lip}_b^1(\mathsf{d}) := \operatorname{Lip}^1(\mathsf{d}) \cap \operatorname{Lip}_b(\mathsf{d})$. For a given measure ν , we set

$$\operatorname{Lip}(\mathsf{d},\nu) := \{ u \in \operatorname{Lip}(\mathsf{d}) : u \text{ is } \nu \text{-measurable} \}$$

as well as $\operatorname{Lip}_b(\mathsf{d},\nu)$ and $\operatorname{Lip}_b^1(\mathsf{d},\nu)$ denoting the corresponding subspaces of ν -measurable functions respectively.

Absolutely continuous curve. Let (X, d) be an extended metric space and $J \subset \mathbb{R}$ be an open interval. A continuous map $\rho_{\cdot} : J \to X$ is *p*-absolutely continuous and denoted by $\rho_{\cdot} \in AC^p(J, (X, \mathsf{d}))$ if there exists $g \in L^p(J, \mathrm{d}x)$ so that

(2.2)
$$\mathsf{d}(\rho_s, \rho_t) \le \int_s^t g(r) \, \mathrm{d}r \;, \quad s, t \in J \quad s < t$$

If p = 1, we simply say that ρ is absolutely continuous and denoted by $\rho \in AC(J, (X, \mathsf{d}))$. The minimal g among those satisfying (2.2) exists and this is identical to

(2.3)
$$|\dot{\rho}_t| := \lim_{s \to t} \frac{\mathsf{d}(\rho_s, \rho_t)}{|s - t|} , \quad \text{the limit exists in a.e. } t \in J ,$$

which is called *metric speed*, or metric derivative of ρ . See [AGS08, Thm. 1.1.2] for a proof. We say that an absolutely continuous curve ρ is constant speed if $|\dot{\rho}_t|$ is a constant for a.e. $t \in J$.

Geodesic space. Let (X, d) be an extended metric space. We say that $(x_t)_{t \in [0,1]}$ is a constant speed geodesic connecting x_0 and x_1 with $\mathsf{d}(x_0, x_1) < \infty$ if

(2.4)
$$\mathsf{d}(x_t, x_s) = |t - s| \mathsf{d}(x_0, x_1) , \quad \forall s, t \in [0, 1]$$

We denote by $\text{Geod}(X, \mathsf{d})$ the space of all constant speed geodesics on (X, d) . We say that (X, d) is a *geodesic* extended metric space if for every pair $x_0, x_1 \in X$ with $\mathsf{d}(x_0, x_1) < \infty$, there exists at least one $(x_t)_{t \in [0,1]} \in \mathsf{Geod}(X, \mathsf{d})$ connecting x_0 and x_1 .

Geodesical convexity. Let (X, d) be a geodesic extended metric space. We say that $U: X \to \mathbb{R} \cup \{+\infty\}$ is K-geodesically convex for $K \in \mathbb{R}$ if for any $x_0, x_1 \in \mathcal{D}(U) := \{x \in X : U(x) \in \mathbb{R}\}$ there exists a constant speed geodesic $\omega : [0, 1] \to X$ with $\omega_0 = x_0$ and $\omega_1 = x_1$ and

$$U(\omega_t) \le (1-t)U(\omega_0) + tU(\omega_1) - \frac{K}{2}t(1-t)\mathsf{d}^2(\omega_0,\omega_1) \quad \forall t \in [0,1] \; .$$

When K = 0, we say that U is geodesically convex.

Slopes. Let (X, d) be an extended metric space and $u : X \to \mathbb{R} \cup \{\pm \infty\}$ be a function. For $x \in \mathcal{D}(u) = \{x \in X : u(x) \in \mathbb{R}\}$, the slope of u at x is defined as

(2.5)
$$|\mathsf{D}_{\mathsf{d}}u|(x) := \begin{cases} \limsup_{y \to x} \frac{|u(y) - u(x)|}{\mathsf{d}(y, x)} & \text{if } x \text{ is not isolated;} \\ 0 & \text{otherwise }. \end{cases}$$

The ascending slope and the descending slope are defined correspondingly as

(2.6)
$$|\mathsf{D}_{\mathsf{d}}^{+}u|(x) := \begin{cases} \limsup_{y \to x} \frac{(u(y) - u(x))^{+}}{\mathsf{d}(y, x)} & \text{if } x \text{ is not isolated;} \\ 0 & \text{otherwise }. \end{cases}$$

(2.7)
$$|\mathsf{D}_{\mathsf{d}}^{-}u|(x) := \begin{cases} \limsup_{y \to x} \frac{(u(y) - u(x))^{-}}{\mathsf{d}(y, x)} & \text{if } x \text{ is not isolated;} \\ 0 & \text{otherwise }. \end{cases}$$

2.4. Cheeger energies. A complete separable geodesic metric space (X, d) equipped with fully supported Radon measure ν with finite total mass $\nu(X) < \infty$ is called *a metric measure space* in this article. Let (X, d, ν) be a metric measure space. The Cheeger energy $\mathsf{Ch}_{\mathsf{d},\nu} : L^2(\nu) \to \mathbb{R} \cup \{+\infty\}$ is defined as the $L^2(\nu)$ -lower semi-continuous envelope of $\int_X |\mathsf{D}_{\mathsf{d}}u|^2 \, \mathrm{d}\nu$:

$$\mathsf{Ch}_{\mathsf{d},\nu}(u) := \frac{1}{2} \inf \left\{ \liminf_{n \to \infty} \int_X |\mathsf{D}_{\mathsf{d}} u_n|^2 \, \mathrm{d}\nu : \ u_n \in \operatorname{Lip}(\mathsf{d}) \cap L^2(\nu) \xrightarrow{L^2} u \right\}.$$

The domain is denoted by $W^{1,2}(X, \mathsf{d}, \nu) := \{u \in L^2(\nu) : \mathsf{Ch}_{\mathsf{d},\nu}(u) < \infty\}$. The Cheeger energy $\mathsf{Ch}_{\mathsf{d},\nu}$ can be expressed by the following integration, see [AGS14a, Thm. 4.5] : there exists a measurable function $|\nabla u|_* \in L^2(\nu)$ so that $|\nabla u|_* \leq |\mathsf{D}_{\mathsf{d}} u|$ ν -a.e. for every $u \in \operatorname{Lip}(\mathsf{d})$ and

$$\mathsf{Ch}_{\mathsf{d},\nu}(u) = \frac{1}{2} \int_X |\nabla u|^2_* \, \mathrm{d}\nu \quad \forall u \in W^{1,2}(X,\mathsf{d},\nu) \ ,$$

where $|\nabla u|_*$ is called *minimal relaxed slope*.

2.5. Riemannian Curvature-dimension condition. Let (X, d, ν) be a metric measure space. The following definition is an equivalent characterisation of $\mathsf{RCD}(K, \infty)$ by [AGS15, Cor. 4.18]. We say that (X, d, ν) satisfies the *Riemannian Curvature-Dimension Condition* $\mathsf{RCD}(K, \infty)$ for $K \in \mathbb{R}$ if

- (i) $\mathsf{Ch}_{\mathsf{d},\nu}$ is quadratic, i.e., $\mathsf{Ch}_{\mathsf{d},\nu}(u+v) + \mathsf{Ch}_{\mathsf{d},\nu}(u-v) = 2\mathsf{Ch}_{\mathsf{d},\nu}(u) + 2\mathsf{Ch}_{\mathsf{d},\nu}(v);$
- (ii) Sobolev-to-Lipschitz property holds, i.e., every $u \in W^{1,2}(X, \mathsf{d}, \nu)$ with $|\nabla u|_* \leq 1$ has a d-Lipschitz ν -representative \tilde{u} with $\operatorname{Lip}(\tilde{u}) \leq 1$;
- (iii) $\mathsf{Ch}_{\mathsf{d},\nu}$ satisfies $\mathsf{BE}_2(K,\infty)$, i.e., $|\nabla T_t u|_*^2 \leq e^{-2Kt}T_t |\nabla u|_*^2$ for every $u \in W^{1,2}(X,\mathsf{d},\nu)$ and t > 0.

In this case, the Cheeger energy $Ch_{d,\nu}$ is a local Dirichlet form ([AGS14b, §4.3]). We note that, while [AGS15, Cor. 4.18] is stated in terms of the minimal weak upper gradient denoted by $|\nabla \cdot|_w$, it is identical to the minimal relaxed slope $|\nabla \cdot|_*$ due to [AGS14a, Thm. 6.2].

2.6. Configuration spaces. A configuration on a locally compact Polish space X is any $\overline{\mathbb{N}}_0$ -valued Radon measure γ on X, which can be expressed by $\gamma = \sum_{i=1}^N \delta_{x_i}$ for $N \in \overline{\mathbb{N}}_0$, where $x_i \in X$ for every i and $\gamma \equiv 0$ if N = 0. The configuration space $\Upsilon = \Upsilon(X)$ is the space of all configurations over X. The space Υ is equipped with the vague topology, i.e., the topology generated by the duality of the space $C_0(X)$ of continuous functions with compact support. We write the restriction $\gamma_A \coloneqq \gamma \mid_A$ for a Polish subspace $A \subset X$ and the corresponding restriction map is denoted by

(2.8)
$$\operatorname{pr}_A \colon \Upsilon \longrightarrow \Upsilon(A) \colon \gamma \longmapsto \gamma_A$$

The N-particle configuration space is denoted by

$$\mathbf{\Upsilon}^N \coloneqq \{\gamma \in \mathbf{\Upsilon} : \gamma(X) = N\}$$
, $N \in \overline{\mathbb{N}}_0$,

Let \mathfrak{S}_k be the k-symmetric group. It can be readily seen that the k-particle configuration space Υ^k is isomorphic to the quotient space $X^{\times k}/\mathfrak{S}_k$:

(2.9)
$$\Upsilon^k \cong X^{\odot k} := X^{\times k} / \mathfrak{S}_k , \quad k \in \mathbb{N}_0$$

The associated projection map from $X^{\times k}$ to the quotient space $X^{\times k}/\mathfrak{S}_k$ is denoted by P_k . For $\eta \in \Upsilon$ and r > 0, we set

(2.10)
$$\boldsymbol{\Upsilon}_r^{\eta} := \{ \gamma \in \boldsymbol{\Upsilon} : \gamma_{B_r^c} = \eta_{B_r^c} \} .$$

Conditional probability. Let μ be a Borel probability measure on Υ . Let

$$\mu(\cdot \mid \operatorname{pr}_{B_r^c}(\cdot) = \eta_{B_r^c})$$

denote the regular conditional probability of μ conditioned at $\eta \in \Upsilon$ with respect to the σ -field generated by the projection map $\gamma \in \Upsilon \mapsto \operatorname{pr}_{B_r}(\gamma) := \operatorname{pr}_r(\gamma) := \gamma_{B_r} \in \Upsilon(B_r)$

(see e.g., [DS21a, Def. 3.32] for the precise definition). Let μ_r^{η} be the probability measure on $\Upsilon(B_r)$ defined as

(2.11)
$$\mu_r^{\eta} := (\mathrm{pr}_r)_{\#} \mu(\cdot \mid \mathrm{pr}_{B_r^c}(\cdot) = \eta_{B_r^c}) ,$$

and its restriction on $\Upsilon^k(B_r)$ is denoted by $\mu_r^{k,\eta} := \mu_r^{\eta}|_{\Upsilon^k(B_r)}$.

Note: The conditional probability $\mu(\cdot | \operatorname{pr}_{B_r^c}(\cdot) = \eta_{B_r^c})$ is a probability measure on the whole space Υ whose support is contained in $\Upsilon_r^{\eta} = \{\gamma \in \Upsilon : \gamma_{B_r^c} = \eta_{B_r^c}\}$. We may project the conditional probability to the probability measure μ_r^{η} on $\Upsilon(B_r)$ as in (2.11) without loss of information in the sense that

(2.12)
$$\operatorname{pr}_r: \Upsilon^{\eta}_r \to \Upsilon(B_r)$$
 is a bi-measure-preserving bijection

Namely, the projection map pr_r is bijective with the inverse map pr_r^{-1} defined as $\operatorname{pr}_r^{-1}(\gamma) := \gamma + \eta$, and both pr_r and pr_r^{-1} are measure-preserving between the two measures $\mu(\cdot | \operatorname{pr}_{B_r^c}(\cdot) = \eta_{B_r^c})$ and μ_r^{η} .

For a measurable function $u: \Upsilon \to \mathbb{R}, r > 0$ and for $\eta \in \Upsilon$, we set

(2.13)
$$u_r^{\eta}(\gamma) \coloneqq u(\gamma + \eta_{B_r^c}) \qquad \gamma \in \Upsilon(B_r)$$

By the property of the conditional probability, it is straightforward to see that for any $u \in L^{1}(\mu)$,

(2.14)
$$\int_{\Upsilon} u \, \mathrm{d}\mu = \int_{\Upsilon} \left[\int_{\Upsilon(B_r)} u_r^{\eta} \, \mathrm{d}\mu_r^{\eta} \right] \mathrm{d}\mu(\eta)$$

See, e.g., [DS21a, Prop. 3.44]. For a measurable set $\Omega \subset \Upsilon$, define a section $\Omega_r^{\eta} \subset \Upsilon(B_r)$ at $\eta \in \Upsilon$ on B_r^c by

(2.15)
$$\Omega_r^{\eta} := \{ \gamma \in \Upsilon(B_r) : \gamma + \eta_{B_r^c} \in \Omega \} .$$

By applying the disintegration formula (2.14) to $u = \mathbf{1}_{\Omega}$, we obtain

(2.16)
$$\mu(\Omega) = \int_{\Upsilon} \mu_r^{\eta}(\Omega_r^{\eta}) \,\mathrm{d}\mu(\eta) \;.$$

Poisson measure. Let (X, τ, ν) be a locally compact Polish space with Radon measure ν satisfying $\nu(X) < \infty$. The Poisson measure π_{ν} on $\Upsilon(X)$ with intensity ν is defined in terms of the symmetric tensor measure ν^{\odot} as follows:

(2.17)
$$\pi_{\nu}(\cdot) := e^{-\nu(X)} \sum_{k=1}^{\infty} \nu^{\odot k} \left(\cdot \cap \Upsilon^{k}(X) \right) = e^{-\nu(X)} \sum_{k=1}^{\infty} \frac{1}{k!} (\mathsf{P}_{k})_{\#} \nu^{\otimes k} \left(\cdot \cap \Upsilon^{k}(X) \right) .$$

 L^2 -transportation distance. Let (X, d) be a locally compact complete separable metric space. For i = 1, 2 let $\operatorname{proj}_i \colon X^{\times 2} \to X$ denote the projection to the i^{th} coordinate for i = 1, 2. For $\gamma, \eta \in \Upsilon$, let $\operatorname{Cpl}(\gamma, \eta)$ be the set of all couplings of γ and η , i.e.,

$$\operatorname{Cpl}(\gamma,\eta) \coloneqq \{ q \in \mathscr{M}(X^{\times 2}) \colon (\operatorname{proj}_1)_{\sharp} q = \gamma \ , \ (\operatorname{proj}_2)_{\sharp} q = \eta \} \ .$$

Here $\mathscr{M}(X^{\times 2})$ denotes the space of all Radon measures on $X^{\times 2}$. The L²-transportation extended distance on $\Upsilon(X)$ is

(2.18)
$$\mathsf{d}_{\Upsilon}(\gamma,\eta) \coloneqq \inf_{q \in \operatorname{Cpl}(\gamma,\eta)} \left(\int_{X^{\times 2}} \mathsf{d}^2(x,y) \, \mathrm{d}q(x,y) \right)^{1/2} , \qquad \inf \mathscr{D} = +\infty .$$

We refer the readers to e.g., [DS21a, Prop. 4.27, 4.29, Thm. 4.37, Prop. 5.12] and [RS99, Lem. 4.1, 4.2] for details regarding the L^2 -transportation extended distance d_{Υ} and examples of d_{Υ} -Lipschitz functions. It is important to note that d_{Υ} is an *extended* distance, attaining the value $+\infty$ and d_{Υ} is lower semi-continuous with respect to the product vague topology $\tau_v^{\times 2}$ but never $\tau_v^{\times 2}$ -continuous.

We introduce a variant of the L^2 -transportation extended distance, called L^2 -transportationtype extended distance \bar{d}_{Υ} defined as

(2.19)
$$\bar{\mathsf{d}}_{\Upsilon}(\gamma,\eta) := \begin{cases} \mathsf{d}_{\Upsilon}(\gamma,\eta) & \text{if } \gamma_{B_r^c} = \eta_{B_r^c} \text{ for some } r > 0 \\ +\infty & \text{otherwise } . \end{cases}$$

By definition, $d_{\Upsilon} \leq \bar{d}_{\Upsilon}$ on Υ , and $d_{\Upsilon} = \bar{d}_{\Upsilon}$ on $\Upsilon(B_r)$ for any r > 0. In particular, we have

(2.20)
$$\operatorname{Lip}(\Upsilon, \mathsf{d}_{\Upsilon}) \subset \operatorname{Lip}(\Upsilon, \overline{\mathsf{d}}_{\Upsilon}) , \quad \operatorname{Lip}_{\overline{\mathsf{d}}_{\Upsilon}}(u) \leq \operatorname{Lip}_{\mathsf{d}_{\Upsilon}}(u) , \quad u \in \operatorname{Lip}(\Upsilon, \mathsf{d}_{\Upsilon}) .$$

It can be readily seen that

(2.21)
$$\bar{\mathsf{d}}_{\Upsilon}(\gamma,\eta) < \infty \quad \iff \quad \gamma_{B_r^c} = \eta_{B_r^c} , \ \gamma(B_r) = \eta(B_r) \quad \text{for some } r > 0$$

When we work with the configuration space $\Upsilon(\mathbb{R}^n)$ over the *n*-dimensional Euclidean space \mathbb{R}^n or over any Polish subset in \mathbb{R}^n , we always choose the Euclidean distance $\mathsf{d}(x, y) = |x - y|$ and the L^2 -transportation distance d_{Υ} and $\bar{\mathsf{d}}_{\Upsilon}$ associated with d .

2.7. sine_{β} ensemble. Let $\beta > 0$ and $C\beta E_k$ be the circular β ensemble on the k-particle configuration space, i.e., it is the probability measure $\mathbb{P}_{k,\beta}$ on the space $\Upsilon^k(\mathbb{S}^1)$ over the unit circle $\mathbb{S}^1 \subset \mathbb{C}$ defined as

$$\mathrm{d}\mathbb{P}_{k,\beta} := \frac{1}{Z_{k,\beta}} \prod_{1 \le j < l \le k} \left| e^{i\theta_j} - e^{i\theta_l} \right|^{\beta} \frac{\mathrm{d}\theta_1}{2\pi} \cdots \frac{\mathrm{d}\theta_k}{2\pi} \,,$$

where the normalisation constant $Z_{k,\beta}$ is given in terms of Gamma function Γ :

$$Z_{k,\beta} := \frac{\Gamma(\frac{1}{2}\beta k + 1)}{\Gamma(\frac{1}{2}\beta k + 1)^k} \,.$$

According to [KS09, Def. 1.6], the *circular* β ensemble C β E is defined as the limit probability measure \mathbb{P}_{β} whose Laplace transform is determined as

$$\int \exp\left(-\sum_{x\in\gamma} f(x)\right) d\mathbb{P}_{\beta}(\gamma) = \lim_{k\to\infty} \int \exp\left(-\sum_{i=1}^{k} f(k\theta_i)\right) d\mathbb{P}_{k,\beta}(\theta_1,\ldots,\theta_k) ,$$

for all $f \in C_0(\mathbb{R})$. In [VV09], a Borel probability measure μ_β on $\Upsilon(\mathbb{R})$ called sine β ensemble has been constructed by a limit of Gaussian β -ensemble. These two measures \mathbb{P}_β and μ_β turned out to be identical each other by the work of [Nak14]. Throughout the rest of the article, we use the symbol $\mu = \mu_\beta$ to denote sine β ensemble (equivalently, circular β ensemble) and we do not specify the inverse temperature β as there is no particular role played by a special β .

Number-rigidity. A Borel probability μ on $\Upsilon = \Upsilon(\mathbb{R}^n)$ is said to be number rigid (in short: (R)) if for any bounded domain $E \subset \mathbb{R}^n$, there exists $\Omega \subset \Upsilon$ so that $\mu(\Omega) = 1$ and,

for any $\gamma, \eta \in \Omega$

(R)
$$\gamma_{E^c} = \eta_{E^c}$$
 implies $\gamma E = \eta E$

Namely, the configuration outside E determines the number of particle inside E. The number-rigidity has been proven in [Gho15] for the sine₂ ensemble and in [NR18], [DHLM20] for the sine_{β} ensemble for general $\beta > 0$.

3. Curvature bound for finite-particle systems

In this section, we study Dirichlet forms on the configuration space $\Upsilon(B_r)$ over metric balls $B_r \subset \mathbb{R}$. We denoted by **m** and \mathbf{m}_r the Lebesgue measure on \mathbb{R} and its restriction on the metric ball $B_r := [-r, r]$ respectively, and take the Euclidean distance $\mathsf{d}(x, y) := |x - y|$ for $x, y \in B_r$.

3.1. Construction of Dirichlet forms on $\Upsilon^k(B_r)$. Let $W_s^{1,2}(\mathsf{m}_r^{\otimes k})$ be the space of $\mathsf{m}_r^{\otimes k}$ classes of (1, 2)-Sobolev and *symmetric* functions on the product space $B_r^{\times k}$, i.e.,

$$W^{1,2}_s(\mathsf{m}_r^{\otimes k}) := \left\{ u \in L^2_s(\mathsf{m}_r^{\otimes k}) : \int_{B_r^{\times k}} |\nabla^{\otimes k} u|^2 \, \mathrm{d}\mathsf{m}_r^{\otimes k} < \infty \right\}$$

where $\nabla^{\otimes k}$ denotes the weak derivative on $\mathbb{R}^{\times k}$: $\nabla^{\otimes k} u := (\partial_1 u, \ldots, \partial_k u)$. The space $W_s^{1,2}(\mathsf{m}_r^{\otimes k})$ consisting of symmetric functions, the projection $\mathsf{P}_k : B_r^{\times k} \to \Upsilon^k(B_r) \cong B_r^{\times k}/\mathfrak{S}_k$ naturally acts on $W_s^{1,2}(\mathsf{m}_r^{\otimes k})$ and the resulting quotient space is denoted by $W^{1,2}(\mathsf{m}_r^{\otimes k})$, which is the (1,2)-Sobolev space on $\Upsilon^k(B_r)$:

$$W^{1,2}(\mathsf{m}_r^{\odot k}) := \left\{ u \in L^2(\mathsf{m}_r^{\odot k}) : \int_{\mathbf{\Upsilon}^k(B_r)} |\nabla^{\odot k} u|^2 \, \mathrm{d}\mathsf{m}_r^{\odot k} < \infty \right\} \,,$$

where $\nabla^{\odot k}$ is the quotient operator of the weak gradient operator $\nabla^{\otimes k}$ through the projection P_k and $\mathsf{m}_r^{\odot k}$ is the symmetric product measure defined as

$$\mathsf{m}_r^{\odot k} := \frac{1}{k!} (\mathsf{P}_k)_{\#} \mathsf{m}_r^{\otimes k}$$

For $0 < r < R < \infty$, $k \in \mathbb{N}_0$ and $\eta \in \Upsilon(B_r^c)$, we introduce the following finite Borel measure on $\Upsilon^k(B_r)$: for $\gamma = \sum_{i=1}^k \delta_{x_i}$

(3.1)
$$d\mu_{r,R}^{k,\eta}(\gamma) := e^{-\Psi_{r,R}^{k,\eta}(\gamma)} dm_r^{\odot k}(\gamma) ,$$
$$\Psi_{r,R}^{k,\eta}(\gamma) := -\log\left(\prod_{i< j}^k |x_i - x_j|^\beta \prod_{i=1}^k \prod_{y \in \eta_{B_r^c}, |y| \le R} \left|1 - \frac{x_i}{y}\right|^\beta\right) .$$

The corresponding weighted Sobolev norm is denoted by

(3.2)
$$\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}}(u) := \frac{1}{2} \int_{\Upsilon^k(B_r)} |\nabla^{\odot k} u|^2 \,\mathrm{d}\mu_{r,R}^{k,\eta} \,, \quad u \in \mathrm{Lip}_b(\Upsilon^k(B_r),\mathsf{d}_\Upsilon) \,,$$

where we note that as $\operatorname{Lip}(\Upsilon^k, \mathsf{d}_{\Upsilon}) \subset W^{1,2}(\mathsf{m}_r^{\odot k})$ and $|\nabla^{\odot k}u| \leq \operatorname{Lip}_{\mathsf{d}_{\Upsilon}}(u)$ due to the Rademacher theorem descendent from the one in the product Sobolev space $W^{1,2}(\mathsf{m}_r^{\otimes k})$ through the quotient, the expression $|\nabla^{\odot k}u|$ and its integral against the probability measure $\mu_{r,R}^{k,\eta}$ make sense for $u \in \operatorname{Lip}_b(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon})$.

Proposition 3.1. The form (3.2) is well-defined and closable. The closure is a local Dirichlet form on $L^2(\mu_{r,R}^{k,\eta})$ and its domain is denoted by $\mathcal{D}(\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}})$.

Proof. The well-definedness follows from the following inequality:

(3.3)
$$\int_{\mathbf{\Upsilon}^{k}(B_{r})} |\nabla^{\odot k} u|^{2} \,\mathrm{d}\mu_{r,R}^{k,\eta} \leq \left\| e^{-\Psi_{r,R}^{k,\eta}} \right\|_{L^{\infty}(\mathbf{\Upsilon}^{k}(B_{r}),\mu_{r,R}^{k,\eta})} \int_{\mathbf{\Upsilon}^{k}(B_{r})} |\nabla^{\odot k} u|^{2} \,\mathrm{d}\mathbf{m}^{\odot k} < \infty \;.$$

The closability of $\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}}$ descends from the closability of the corresponding Dirichlet form on the product space $B_r^{\times k}$ defined on the space of symmetric $\mathsf{d}^{\times k}$ -Lipschitz functions:

$$\mathcal{E}^{B^{\times k}_r,\mu^{k,\eta}_{r,R}} := \frac{1}{2} \int_{B^{\times k}_r} |\nabla^{\otimes k} u|^2 e^{-\Psi^{k,\eta}_{r,R}} \,\mathrm{d}\mathbf{m}_r^{\otimes k}$$

where the closability of $\mathcal{E}^{B_r^{\times k}, \mu_{r,R}^{k,\eta}}$ is a consequence of the continuity of the density $e^{-\Psi_{r,R}^{k,\eta}}$ on $B_r^{\times k}$ and the standard Hamza-type argument by [MR85, Fuk97], see for an accessible reference, e.g., [MR90, pp. 44-45]. The locality of the form is an immediate consequence of the locality of the gradient operator $\nabla^{\odot k}$.

Let μ be the sine_{β} ensemble. Due to [DHLM20, Thm. 1.1], the following limit exists for μ -a.e. η , all $x \in B_r$ and r > 0:

$$\lim_{R \to \infty} \prod_{y \in \eta_{B_r^c}, |y| \le R} \left| 1 - \frac{x}{y} \right|^{\beta}.$$

Recall that μ_r^{η} has been defined in (2.11). By [DHLM20, Thm. 1.1] and the numberrigidity (R) of μ , for μ -a.e. η there exists $k = k(\eta)$ so that

(3.4)
$$\mu_r^{\eta}(\Upsilon^l(B_r)) > 0 \text{ if and only if } l = k(\eta)$$

and for $\gamma = \sum_{i=1}^{k} \delta_{x_i}$,

(3.5)
$$d\mu_{r}^{\eta} = d\mu_{r}^{k,\eta} = \frac{e^{-\Psi_{r}^{k,\eta}}}{Z_{r}^{\eta}} d\mathsf{m}_{r}^{\odot k} ,$$
$$\Psi_{r}^{k,\eta}(\gamma) := -\log\left(\prod_{i< j}^{k} |x_{i} - x_{j}|^{\beta} \prod_{i=1}^{k} \lim_{R \to \infty} \prod_{y \in \eta_{B_{r}^{c}}, |y| \le R} \left|1 - \frac{x_{i}}{y}\right|^{\beta}\right)$$

where Z_r^{η} is the normalising constant. Note that the roles of the notation γ and η in [DHLM20] are opposite to this article. The corresponding weighted Sobolev norm is defined as

(3.6)
$$\mathcal{E}^{\Upsilon(B_r),\mu_r^{k,\eta}}(u) := \frac{1}{2} \int_{\Upsilon^k(B_r)} |\nabla^{\odot k} u|^2 \,\mathrm{d}\mu_r^{k,\eta} \,, \quad u \in \mathrm{Lip}_b(\Upsilon^k(B_r),\mathsf{d}_\Upsilon) \,.$$

Proposition 3.2. Let μ be the sine_{β} ensemble for $\beta > 0$. The form (3.6) is well-defined and closable for μ -a.e. η . The closure is a local Dirichlet form on $L^2(\mu_r^{k,\eta})$ and its domain is denoted by $\mathcal{D}(\mathcal{E}^{\Upsilon(B_r),\mu_r^{k,\eta}})$.

Proof. As $e^{-\Psi_{r,R}^{k,\eta}} \xrightarrow{R \to \infty} e^{-\Psi_r^{k,\eta}}$ uniformly on $\Upsilon^k(B_r)$ for μ -a.e. η by [DHLM20, Lem. 2.3 and Proof of Thm. 2.1 in p. 183], the density $e^{-\Psi_r^{k,\eta}}$ is continuous on $B_r^{\odot k}$, hence the same proof as Prop. 3.1 applies to conclude the statement.

3.2. Curvature bound for finite-particle systems. We show that the potential $\Psi_{r,R}^{k,\eta}$ defined in (3.1) is geodesically convex in $(\Upsilon(B_r), \mathsf{d}_{\Upsilon})$.

Proposition 3.3. $\Psi_{r,R}^{k,\eta}$ is geodesically convex in $(\Upsilon^k(B_r), \mathsf{d}_\Upsilon)$ for any $0 < r < R < \infty$, $k \in \mathbb{N}$ and $\eta \in \Upsilon(B_r^c)$,

Proof. Note that if u_1, \ldots, u_k are convex and $\alpha_1, \ldots, \alpha_k \geq 0$, then $\sum_{i=1}^k \alpha_i u_i$ is again convex. Let H_{ij}, H_i^y be the Hessian matrices of the functions $(x_1, \ldots, x_k) \mapsto -\log |x_i - x_j|$ and $(x_1, \ldots, x_k) \mapsto -\log |1 - \frac{x_i}{y}|$ respectively. For any vector $\mathbf{v} = (v_1, \ldots, v_k) \in \mathbb{R}^k$,

(3.7)
$$\mathbf{v}H_{ij}\mathbf{v}^{t} = \frac{(v_{i} - v_{j})^{2}}{|x_{i} - x_{j}|^{2}}, \quad \mathbf{v}H_{i}^{y}\mathbf{v}^{t} = \frac{v_{i}^{2}}{|y - x_{i}|^{2}}.$$

Both H_{ij} and H_i^y are, therefore, positive semi-definite. Thus, for any 0 < r < R, any $y \in [-R, -r] \cup [r, R]$ and any $i, j \in \{1, 2, ..., k\}$ with $i \neq j$, the functions $(x_1, ..., x_k) \mapsto -\log |x_i - x_j|$ and $(x_1, ..., x_k) \mapsto -\log |1 - \frac{x_i}{y}|$ are convex in the following areas for any $\sigma \in \mathfrak{S}_k$:

$$\{(x_1, \ldots, x_k) \in B_r^{\times k} : x_{\sigma(1)} < x_{\sigma(2)} < \cdots < x_{\sigma(k)}\}.$$

In view of (2.9), the following expression, therefore, concludes that $\Psi_{r,R}^{k,\eta}$ is geodesically convex as a function on $\Upsilon^k(B_r)$: for any $\gamma = \sum_{i=1}^k \delta_{x_i}$

(3.8)
$$\Psi_{r,R}^{k,\eta}(\gamma) = -\beta \sum_{i$$

The proof is complete.

Thanks to the geodesical convexity of the potential $\Psi_{r,R}^{k,\eta}$ shown in Prop. 3.3, the Dirichlet form $(\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}}, \mathcal{D}(\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}}))$ satisfies the Riemannian Curvature Dimension condition $\mathsf{RCD}(0,\infty)$.

Proposition 3.4. The space $(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon}, \mu_{r,R}^{k,\eta})$ satisfies $\mathsf{RCD}(0,\infty)$ for every $k \in \mathbb{N}_0$, $0 < r < R < \infty$ and $\eta \in \Upsilon$, and it holds that

$$\left(\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}},\mathcal{D}(\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}})\right) = \left(\mathsf{Ch}_{\mathsf{d}_{\Upsilon},\mu_{r,R}^{k,\eta}},W^{1,2}(\Upsilon^k(B_r),\mathsf{d}_{\Upsilon},\mu_{r,R}^{k,\eta})\right) \,.$$

Proof. Noting that $B_r^{\times k}$ is a convex subset in \mathbb{R}^k , the space $(B_r^{\times k}, \mathsf{d}^{\times k}, \mathsf{m}_r^{\otimes k})$ is a geodesic subspace of \mathbb{R}^k and, therefore, satisfies $\mathsf{RCD}(0,\infty)$ by the Global-to-Local property of $\mathsf{RCD}(0,\infty)$, see [AGS14b, Thm. 6.20]. Noting that the k-particle configuration space $(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon}, \mu_{r,R}^{k,\eta})$ is the quotient space of $(B_r^{\times k}, \mathsf{d}^{\times k}, \mathsf{m}^{\otimes k})$ with respect to the symmetric group \mathfrak{S}_k and that the property $\mathsf{RCD}(0,\infty)$ is preserved under the quotient operation with respect to \mathfrak{S}_k thanks to [GKMS18], we obtain that $(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon}, \mathsf{m}^{\odot k})$ satisfies $\mathsf{RCD}(0,\infty)$ as well. By the geodesical convexity of the potential $\Psi_{r,R}^{k,\eta}$ shown in Prop. 3.3 and the continuity of the density $e^{-\Psi_{r,R}^{k,\eta}}$, the weighted space $(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon}, \mu_{r,R}^{k,\eta})$ satisfies $\mathsf{RCD}(0,\infty)$ by [AGS14b, Prop. 6.21].

To conclude the statement, it suffices to check the identity

$$\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{\kappa,\eta}} = \mathsf{Ch}_{\mathsf{d}_\Upsilon,\mu_{r,R}^{k,\eta}}$$

By the Rademacher theorem on $\Upsilon^k(B_r)$ descendent from the Rademacher theorem on $B_r^{\times k}$, the slope $|\mathsf{D}_{\mathsf{d}_{\Upsilon}} u|$ coincides with the gradient $|\nabla^{\odot k} u|$ for any $u \in \operatorname{Lip}(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon})$. Thus,

(3.9)
$$\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}}(u) = \frac{1}{2} \int_{\Upsilon^k(B_r)} |\mathsf{D}_{\mathsf{d}_{\Upsilon}}u|^2 \,\mathrm{d}\mu_{r,R}^{k,\eta} \quad u \in \mathrm{Lip}_b(\Upsilon^k(B_r),\mathsf{d}_{\Upsilon}) \,.$$

Since $\mathsf{Ch}_{\mathsf{d}_{\Upsilon},\mu_{r,R}^{k,\eta}}$ is the L^2 -lower semi-continuous envelope by definition, the functional $\mathsf{Ch}_{\mathsf{d}_{\Upsilon},\mu_{r,R}^{k,\eta}}$ is the maximal L^2 -lower semi-continuous functional satisfying

$$\mathsf{Ch}_{\mathsf{d}_{\Upsilon},\mu_{r,R}^{k,\eta}}(u) \leq \frac{1}{2} \int_{\Upsilon^k(B_r)} |\mathsf{D}_{\mathsf{d}_{\Upsilon}}u|^2 \,\mathrm{d}\mu_{r,R}^{k,\eta} \,.$$

As $\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}}$ is closed by Prop. 3.1, in particular, $\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}}$ is L^2 -lower semi-continuous. Therefore, combining the maximality of $\mathsf{Ch}_{\mathsf{d}_{\Upsilon},\mu_{r,R}^{k,\eta}}$ with (3.9), it holds that

$$\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}} \leq \mathsf{Ch}_{\mathsf{d}_{\Upsilon},\mu_{r,R}^{k,\eta}} \quad \text{and} \quad W^{1,2}(\Upsilon^k(B_r),\mathsf{d}_{\Upsilon},\mu_{r,R}^{k,\eta}) \subset \mathcal{D}(\mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}})$$

and

$$\mathsf{Ch}_{\mathsf{d}_{\Upsilon},\mu_{r,R}^{k,\eta}}(u) = \mathcal{E}^{\Upsilon(B_r),\mu_{r,R}^{k,\eta}}(u) \quad u \in \mathrm{Lip}_b(\Upsilon^k(B_r),\mathsf{d}_{\Upsilon}) \ .$$

As $\operatorname{Lip}_b(\Upsilon^k(B_r), \mathsf{d}_\Upsilon)$ is dense both in $\mathcal{D}(\mathcal{E}^{\Upsilon(B_r), \mu_{r,R}^{k,\eta}})$ and $W^{1,2}(\Upsilon^k(B_r), \mathsf{d}_\Upsilon, \mu_{r,R}^{k,\eta})$ by construction, the proof is completed.

In view of Prop. 3.4 and the approximation $\Psi_{r,R}^{k,\eta}$ to $\Psi_{r,R}^{k,\eta}$ as $R \to \infty$, we prove that $(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon}, \mu_r^{k,\eta})$ satisfies $\mathsf{RCD}(0, \infty)$ as well.

Proposition 3.5. Let μ be the sine_{β} ensemble for $\beta > 0$. For any $0 < r < \infty$ and μ a.e. $\eta \in \Upsilon$, the space $(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon}, \mu_r^{k,\eta})$ satisfies $\mathsf{RCD}(0, \infty)$, where $k = k(\eta)$ as in (3.4). Furthermore,

$$\left(\mathcal{E}^{\Upsilon(B_r),\mu_r^{k,\eta}}, \mathcal{D}(\mathcal{E}^{\Upsilon(B_r),\mu_r^{k,\eta}})\right) = \left(\mathsf{Ch}_{\mathsf{d}_{\Upsilon},\mu_r^{k,\eta}}, W^{1,2}(\Upsilon^k(B_r),\mathsf{d}_{\Upsilon},\mu_r^{k,\eta})\right)$$

Proof. Since the potential $\Psi_{r,R}^{k,\eta}$ is geodesically convex for any R and it converges pointwise to $\Psi_r^{k,\eta}$ as $R \to \infty$ for μ -a.e. η by [DHLM20, Lem. 2.3 and Proof of Thm. 2.1 in p. 183], the potential $\Psi_r^{k,\eta}$ is again geodesically convex on $(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon})$. Furthermore, as the density $e^{-\Psi_{r,R}^{k,\eta}}$ converges uniformly to $e^{-\Psi_r^{k,\eta}}$ on $\Upsilon^k(B_r)$ as $R \to \infty$ for μ -a.e. η by [DHLM20, Lem. 2.3 and Proof of Thm. 2.1 in p. 183], the density $e^{-\Psi_r^{k,\eta}}$ is continuous on $\Upsilon(B_r)$. Noting the fact that the constant multiplication (by the normalisation constant Z_r^{η}) does not change the lower Ricci curvature bound (see e.g., [Stu06, Prop. 4.13]), the same proof as Prop. 3.4 applies to conclude the statement.

4. CURVATURE BOUND FOR INFINITE-PARTICLE SYSTEMS

In this section, we construct a local Dirichlet form on $\Upsilon = \Upsilon(\mathbb{R})$ associated with $\operatorname{sine}_{\beta}$ ensemble μ and show the $\mathsf{BE}(0,\infty)$ property by the following steps: we first construct truncated Dirichlet forms on Υ whose gradient operators are truncated up to configurations inside B_r . We then identify them with the superposition Dirichlet forms lifted from $\Upsilon(B_r)$, thanks to which we can show $\mathsf{BE}(0,\infty)$ for the truncated forms. We take the monotone limit of the truncated forms to construct a Dirichlet form with invariant measure $\operatorname{sine}_{\beta}$ ensembles μ and $\mathsf{BE}(0,\infty)$ extends to the limit form. In the end of this section, we discuss several applications of the $\mathsf{BE}(0,\infty)$ property.

4.1. Truncated Dirichlet forms. In this subsection, we construct the truncated Dirichlet forms on Υ . We first construct square field operators on Υ and $\Upsilon(B_r)$ respectively. For so doing, we introduce a map $\mathcal{U}_{\gamma,x}$ transferring functions on the configuration space Υ to functions on the base space \mathbb{R} . For $u : \Upsilon \to \mathbb{R}$, define $\mathcal{U}_{\gamma,x}(u) : \mathbb{R} \to \mathbb{R}$ by

(4.1)
$$\mathcal{U}_{\gamma,x}(u)(y) := u\big(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma + \delta_y\big) - u\big(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma\big) , \quad \gamma \in \Upsilon, \quad x \in \gamma.$$

In the context of configuration spaces, the operation $\mathcal{U}_{\gamma,x}$ has been firstly discussed in [MR00, Lem. 1.2], see also [DS21a, Lem. 2.16]. We introduce the localisation of the operator $\mathcal{U}_{\gamma,x}$ on B_r . Recall that for a measurable function $u: \Upsilon \to \mathbb{R}, r > 0$ and for $\eta \in \Upsilon$, we set in (2.13)

$$u_r^{\eta}(\gamma) \coloneqq u(\gamma + \eta_{B_r^c}) \text{ for } \gamma \in \Upsilon(B_r)$$

Lemma 4.1. For $u : \Upsilon(B_r) \to \mathbb{R}$, define $\mathcal{U}_{\gamma,x}^r(u) : B_r \to \mathbb{R}$ by

$$\mathcal{U}_{\gamma,x}^{r}(u)(y) := u(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma + \delta_{y}) - u(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma) \quad \gamma \in \Upsilon(B_{r}), \ x \in \gamma .$$

The operation $\mathcal{U}_{\gamma,x}^r$ maps from $\operatorname{Lip}(\Upsilon(B_r), \mathsf{d}_\Upsilon)$ to $\operatorname{Lip}(B_r)$ and Lipschitz constants are contracted by $\mathcal{U}_{\gamma,x}^r$ for any r > 0:

$$\operatorname{Lip}(\mathcal{U}_{\gamma,x}^{r}(u)) \leq \operatorname{Lip}_{\mathsf{d}_{\boldsymbol{\Upsilon}}}(u) \quad \forall \gamma \in \boldsymbol{\Upsilon}(B_{r}) \quad \forall x \in \gamma \ .$$

Furthermore, for any $u: \Upsilon \to \mathbb{R}$,

$$\mathcal{U}^r_{\gamma_{B_r},x}(u^{\gamma}_r)(y) = \mathcal{U}_{\gamma,x}(u)(y) \quad for \ every \ \gamma \in \Upsilon, \ x \in \gamma_{B_r} \ and \ y \in B_r \ .$$

Proof. Let $u \in \text{Lip}(\Upsilon(B_r), \mathsf{d}_{\Upsilon})$. Then

$$\begin{aligned} |\mathcal{U}_{\gamma,x}^{r}(u)(y) - \mathcal{U}_{\gamma,x}^{r}(u)(z)| &= |u(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma + \delta_{y}) - u(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma + \delta_{z})| \\ &\leq \operatorname{Lip}_{\mathsf{d}_{\Upsilon}}(u)\mathsf{d}_{\Upsilon}(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma + \delta_{y}, \mathbf{1}_{X \setminus \{x\}} \cdot \gamma + \delta_{z}) \\ &= \operatorname{Lip}_{\mathsf{d}_{\Upsilon}}(u)|y - z| , \end{aligned}$$

which concludes the first assertion.

We verify the second assertion. For every $x \in \gamma_{B_r}$ and $y \in B_r$,

$$\begin{aligned} \mathcal{U}_{\gamma,x}(u)(y) &= u(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma + \delta_y) - u(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma) \\ &= u(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma_{B_r} + \gamma_{B_r^c} + \delta_y) - u(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma_{B_r} + \gamma_{B_r^c}) \\ &= u_{r,\gamma}(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma_{B_r} + \delta_y) - u_{r,\gamma}(\mathbf{1}_{X \setminus \{x\}} \cdot \gamma_{B_r}) \\ &= \mathcal{U}_{\gamma_{B_r},x}^r(u_r^\gamma)(y) \;. \end{aligned}$$

The proof is complete.

We now define a square field operator on Υ truncated up to particles inside B_r .

Definition 4.2 (Truncated square field on Υ). Let $u : \Upsilon \to \mathbb{R}$ be a measurable function so that $\mathcal{U}_{\gamma,x}(u)|_{B_r} \in W^{1,2}(\mathfrak{m}_r)$ for μ -a.e. γ and every $x \in \gamma_{B_r}$. The following operator is called the truncated square field Γ_r^{Υ} ,

(4.2)
$$\Gamma_r^{\Upsilon}(u)(\gamma) := \sum_{x \in \gamma_{B_r}} |\nabla \mathcal{U}_{\gamma,x}(u)|^2(x) \; .$$

Thanks to Lem. A.1, Formula (4.2) is well-defined for μ -a.e. γ . Indeed, as $\mathcal{U}_{\gamma,x}(u)|_{B_r} \in W^{1,2}_{loc}(\mathfrak{m}_r)$, the weak gradient $\nabla \mathcal{U}_{\gamma,x}(u)$ is well-defined pointwise on a measurable set $\Sigma \subset B_r$ with $\mathfrak{m}_r(\Sigma^c) = 0$. By applying Lem. A.1, Formula (4.2) is well-defined on the set $\Omega(r)$ of μ -full measure.

Based on the truncated square field Γ_r^{Υ} , we introduce the truncated form on Υ defined on a certain core.

Definition 4.3 (Core). For r > 0, let C_r be defined as the space of μ -classes of measurable functions u so that

- (a) $u \in L^{\infty}(\mu);$
- (b) $u_r^{\eta} \in \operatorname{Lip}_b(\Upsilon(B_r), \mathsf{d}_\Upsilon)$ for μ -a.e. η ;
- (c) The following integral is finite:

(4.3)
$$\mathcal{E}_r^{\Upsilon,\mu}(u) := \frac{1}{2} \int_{\Upsilon} \Gamma_r^{\Upsilon}(u) \,\mathrm{d}\mu < \infty$$

Note that, thanks to Lem. 4.1, if a measurable function $u : \Upsilon \to \mathbb{R}$ satisfies (b), then $\mathcal{U}_{\gamma,x}(u)|_{B_r} \in \operatorname{Lip}(B_r, \mathsf{d}) \subset W^{1,2}(\mathsf{m}_r)$. Thus, the expression $\Gamma_r^{\Upsilon}(u)$ in (4.3) is well-posed. It will be proved in Prop. 4.8 that \mathcal{C}_r is non-trivial in the sense that every μ -measurable bounded $\bar{\mathsf{d}}_{\Upsilon}$ -Lipschitz functions on Υ belongs to \mathcal{C}_r .

Remark 4.4. Note that the core C_r is much larger than the Lipschitz algebras $\operatorname{Lip}_b(\Upsilon, \bar{\mathsf{d}}_{\Upsilon}, \mu)$ because the condition (b) allows the Lipschitz constant $\operatorname{Lip}_{\Upsilon(B_r), \mathsf{d}_{\Upsilon}}(u_r^{\eta})$ to depend both on η and r.

Definition 4.5 (Square field on $\Upsilon(B_r)$). Fix r > 0 and $\eta \in \Upsilon$. For a μ -measurable function $u: \Upsilon(B_r) \to \mathbb{R}$ satisfying $u|_{\Upsilon^k(B_r)} \in \mathcal{D}(\mathcal{E}^{\Upsilon(B_r),\mu_r^{k,\eta}})$ for any $k \in \mathbb{N}_0$, we define the following square field operator on $\Upsilon(B_r)$:

(4.4)
$$\Gamma^{\Upsilon(B_r)}(u) := \sum_{k=0}^{\infty} \left| \nabla^{\odot k} \left(u |_{\Upsilon^k(B_r)} \right) \right|^2 \,,$$

and define the following form:

$$\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}(u) := \frac{1}{2} \int_{\Upsilon(B_r)} \Gamma^{\Upsilon(B_r)}(u) \, \mathrm{d}\mu_r^{\eta} ,$$

$$\mathcal{D}(\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}) := \{ u : \Upsilon(B_r) \to \mathbb{R}, \ \mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}(u) < \infty \} .$$

Due to the number-rigidity (R), the Dirichlet form $\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}$ is equal to $\mathcal{E}^{\Upsilon(B_r),\mu_r^{k,\eta}}$ for some $k = k(\eta)$ up to the normalising multiplicative constant, therefore, it is a Dirichlet form as well. The corresponding semigroup operator is denoted by $\{T_t^{\Upsilon(B_r),\mu_r^{\eta}}\}_{t\geq 0}$.

Remark 4.6. The number-rigidity (\mathbb{R}) is not essential here for $\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}$ to be a Dirichlet form. Any countable sum of Dirichlet forms is a Dirichlet form (see e.g., [MR90, Exercise 3.9 in p.31]).

Before discussing properties of truncated forms, we prepare a lemma, which states that the operation $(\cdot)_r^{\eta}$ defined in (2.13) maps from $\operatorname{Lip}(\Upsilon, \bar{\mathsf{d}}_{\Upsilon})$ to $\operatorname{Lip}(\Upsilon(B_r), \mathsf{d}_{\Upsilon})$ and contracts Lipschitz constants.

20

Lemma 4.7. Let $u \in \operatorname{Lip}(\Upsilon, \overline{\mathsf{d}}_{\Upsilon})$. Then, $u_r^{\eta} \in \operatorname{Lip}(\Upsilon(B_r), \mathsf{d}_{\Upsilon})$ and

(4.5)
$$\operatorname{Lip}_{\mathsf{d}_{\Upsilon}}(u_r^{\eta}) \leq \operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(u) , \quad \eta \in \Upsilon , \quad r > 0$$

Proof. Let $\gamma, \zeta \in \Upsilon(B_r)$ and $\eta \in \Upsilon$. Then,

$$\begin{aligned} |u_r^{\eta}(\gamma) - u_r^{\eta}(\zeta)| &= |u(\gamma + \eta_{B_r^c}) - u(\zeta + \eta_{B_r^c})| \le \operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(u)\bar{\mathsf{d}}_{\Upsilon}(\gamma + \eta_{B_r^c}, \zeta + \eta_{B_r^c}) \\ &= \operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(u)\mathsf{d}_{\Upsilon}(\gamma, \zeta) \;. \end{aligned}$$

The proof is completed.

The following proposition relates the two square fields Γ_r^{Υ} and $\Gamma^{\Upsilon(B_r)}$.

Proposition 4.8 (Truncated form). The following relations hold on C_r :

(4.6)
$$\Gamma_{r}^{\Upsilon}(u)(\gamma + \eta_{B_{r}^{c}}) = \Gamma^{\Upsilon(B_{r})}(u_{r}^{\eta})(\gamma) , \quad \mu\text{-a.e. } \eta, \quad \mu_{r}^{\eta}\text{-a.e. } \gamma \in \Upsilon(B_{r}) ,$$
$$\mathcal{E}_{r}^{\Upsilon,\mu}(u) = \int_{\Upsilon} \mathcal{E}^{\Upsilon(B_{r}),\mu_{r}^{\eta}}(u_{r}^{\eta}) \,\mathrm{d}\mu(\eta) , \quad u \in \mathcal{C}_{r} .$$

Furthermore, the Rademacher-type property holds: $\operatorname{Lip}_{h}(\bar{\mathsf{d}}_{\Upsilon},\mu) \subset \mathcal{C}_{r}$ and

(4.7)
$$\Gamma_r^{\Upsilon}(u) \le \operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(u)^2 \qquad u \in \operatorname{Lip}_b(\bar{\mathsf{d}}_{\Upsilon},\mu)$$

As a consequence, the form $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{C}_r)$ in (4.3) is a densely defined closable Markovian form and the closure $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}))$ is a local Dirichlet form on $L^2(\mu)$. The L^2 -semigroups corresponding to $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}))$ is denoted by $\{T_{r,t}^{\Upsilon,\mu}\}_{t\geq 0}$.

Proof. We first prove (4.6). Let $u \in C_r$. Thanks to (b) in Def. 4.3 and Lem. 4.1,

 $\mathcal{U}_{\gamma,x}(u) \in \operatorname{Lip}(B_r, \mathsf{d})$, μ -a.e. γ r > 0.

Thus, noting $\operatorname{Lip}(B_r, \mathsf{d}) \subset W^{1,2}(\mathsf{m}_r)$, the LHS of (4.6) is well-defined. Thus, there exists $\Omega \subset \Upsilon$ with $\mu(\Omega) = 1$ so that the LHS of (4.6) is well-defined everywhere on Ω . The RHS of (4.6) is also well-defined by (b) in Def. 4.3 and by $\operatorname{Lip}_b(\Upsilon(B_r), \mathsf{d}_\Upsilon) \subset \mathcal{D}(\mathcal{E}^{\Upsilon(B_r), \mu_r^{\eta}})$ by construction. Let Ω_r^{η} be a section as defined in (2.15). As μ_r^{η} is absolutely continuous with respect to the Poisson measure π_{m_r} and the Poisson measure does not have multiple points almost everywhere, we may assume that every $\gamma \in \Omega_r^{\eta}$ does not have multiple points, i.e., $\gamma(\{x\}) \in \{0,1\}$ for every $x \in B_r$. Let $\gamma \in \Omega_r^{\eta} \cap \Upsilon^k(B_r)$. Then, according to (4.4),

$$\begin{split} &\Gamma_{r}^{\boldsymbol{\Upsilon}}(u)(\gamma+\eta_{B_{r}^{c}})\\ &=\sum_{x\in\gamma}\left|\nabla\Big(u\big(\mathbf{1}_{X\setminus\{x\}}\cdot(\gamma+\eta_{B_{r}^{c}})+\delta_{\bullet}\big)-u\big(\mathbf{1}_{X\setminus\{x\}}\cdot(\gamma+\eta_{B_{r}^{c}})\big)\Big)\right|^{2}(x)\\ &=\sum_{x\in\gamma}\left|\nabla\Big(u_{r}^{\eta}\big(\mathbf{1}_{X\setminus\{x\}}\cdot\gamma+\delta_{\bullet}\big)-u_{r}^{\eta}\big(\mathbf{1}_{X\setminus\{x\}}\cdot\gamma\big)\Big)\right|^{2}(x)\\ &=\sum_{x\in\gamma}\left|\nabla u_{r}^{\eta}\big(\mathbf{1}_{X\setminus\{x\}}\cdot\gamma+\delta_{\bullet}\big)\right|^{2}(x)\\ &=\left|\nabla^{\odot k}\big(u_{r}^{\eta}\big)\right|^{2}(\gamma)\\ &=\Gamma^{\boldsymbol{\Upsilon}(B_{r})}(u_{r}^{\eta})(\gamma) \end{split}$$

where the first equality is the definition of the square field Γ_r^{Υ} ; the third equality holds as $u_r^{\eta}(\mathbf{1}_{X\setminus\{x\}},\gamma)$ does not depend on the variable denoted as \bullet on which the weak gradient ∇

operates; the fourth equality followed from the definition of the symmetric gradient operator $\nabla^{\odot k}$, for which we used the fact that $\gamma \in \Omega_r^{\eta}$ does not have multiple points. As this argument holds for arbitrary $k \in \mathbb{N}_0$, (4.6) has been shown. The locality and the Markov property of $\mathcal{E}_r^{\Upsilon,\mu}$ follow from (4.6) and the fact that $\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}$ possesses the corresponding properties by construction.

We now show the Rademacher-type property: $\operatorname{Lip}_b(\bar{\mathsf{d}}_{\Upsilon}, \mu) \subset \mathcal{C}_r$ and

(4.8)
$$\Gamma_r^{\Upsilon}(u) \le \operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(u)^2 \quad \forall u \in \operatorname{Lip}_b(\bar{\mathsf{d}}_{\Upsilon}, \mu) \quad \forall r > 0$$

We first show $\operatorname{Lip}_b(\bar{\mathsf{d}}_{\mathbf{\Upsilon}},\mu) \subset \mathcal{C}_r$. The verification of (a) in Def. 4.3 is obvious. The verification of (b) in Def. 4.3 follows from the Lipschitz contraction (4.5) of the operator $(\cdot)_r^{\eta}$. The verification of (c) in Def. 4.3 follows by showing (4.8) as μ is a probability measure.

We now prove (4.8). As the Cheeger energy $\mathsf{Ch}_{\mathsf{d}_{\Upsilon},\mu_r^{k,\eta}}$ coincided with the form $\mathcal{E}^{\Upsilon(B_r),\mu_r^{k,\eta}}$ by Prop. 3.5, the Rademacher-type property for $\mathcal{E}^{\Upsilon(B_r),\mu_r^{k,\eta}}$ follows from that for $\mathsf{Ch}_{\mathsf{d}_{\Upsilon},\mu_r^{k,\eta}}$, the latter of which is an immediate consequence by the definition of the Cheeger energy. Therefore, we have that

(4.9)
$$\Gamma^{\Upsilon(B_r)}(u) \leq \operatorname{Lip}_{\mathsf{d}_{\Upsilon}}(u)^2 \quad \forall u \in \operatorname{Lip}(\Upsilon(B_r), \mathsf{d}_{\Upsilon}) \quad \forall r > 0 .$$

In view of the relation between Γ_r^{Υ} and $\Gamma^{\Upsilon(B_r)}$ in (4.6) and the Lipschitz contraction (4.5) of the operator $(\cdot)_r^{\eta}$, we concluded (4.8).

Noting that $\operatorname{Lip}_b(\mathsf{d}_{\Upsilon},\mu) \subset L^2(\mu)$ is dense (e.g., [AGS14a, Prop. 4.1]) and the fact that $\operatorname{Lip}_b(\mathsf{d}_{\Upsilon},\mu) \subset \operatorname{Lip}_b(\bar{\mathsf{d}}_{\Upsilon},\mu) \subset \mathcal{C}_r$ by (2.20) and (4.8), we obtain that the form $(\mathcal{E}_r^{\Upsilon,\mu},\mathcal{C}_r)$ is densely defined.

We now show the closability. Noting that $\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}$ is closable for μ -a.e. η by Prop. 3.5, the superposition form $(\bar{\mathcal{E}}_r^{\Upsilon,\mu}, \mathcal{D}(\bar{\mathcal{E}}_r^{\Upsilon,\mu}))$ (defined below in Def. 4.9) is closable (indeed it is closed) by [BH91, Prop. V.3.1.1]. As the two forms $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{C}_r)$ and $(\bar{\mathcal{E}}_r^{\Upsilon,\mu}, \mathcal{D}(\bar{\mathcal{E}}_r^{\Upsilon,\mu}))$ coincide on \mathcal{C}_r by definition and $\mathcal{C}_r \subset \mathcal{D}(\bar{\mathcal{E}}_r^{\Upsilon,\mu})$ by construction, the closability of $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{C}_r)$ is inherited from the closedness of the superposition form $(\bar{\mathcal{E}}_r^{\Upsilon,\mu}, \mathcal{D}(\bar{\mathcal{E}}_r^{\Upsilon,\mu}))$. The proof is complete.

4.2. Superposition form. The superposition of the Dirichlet form $\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}$ onto Υ is now defined below.

Definition 4.9 (Superposition Dirichlet form, e.g., [BH91, Prop. V.3.1.1]).

(4.10)
$$\mathcal{D}(\bar{\mathcal{E}}_{r}^{\Upsilon,\mu}) := \left\{ u \in L^{2}(\mu) : \int_{\Upsilon} \mathcal{E}^{\Upsilon(B_{r}),\mu_{r}^{\eta}}(u_{r}^{\eta}) \,\mathrm{d}\mu(\eta) < \infty \right\},\ \bar{\mathcal{E}}_{r}^{\Upsilon,\mu}(u) := \int_{\Upsilon} \mathcal{E}^{\Upsilon(B_{r}),\mu_{r}^{\eta}}(u_{r}^{\eta}) \,\mathrm{d}\mu(\eta) .$$

It is known that $(\bar{\mathcal{E}}_r^{\Upsilon,\mu}, \mathcal{D}(\bar{\mathcal{E}}_r^{\Upsilon,\mu}))$ is a Dirichlet form on $L^2(\mu)$ [BH91, Prop. V.3.1.1]. The L^2 -semigroup and the infinitesimal generator corresponding to $(\bar{\mathcal{E}}_r^{\Upsilon,\mu}, \mathcal{D}(\bar{\mathcal{E}}_r^{\Upsilon,\mu}))$ are denoted by $\{\bar{T}_{r,t}^{\Upsilon,\mu}\}_{t\geq 0}$ and $(\bar{A}_r^{\Upsilon,\mu}, \mathcal{D}(\bar{A}_r^{\Upsilon,\mu}))$ respectively.

The semigroup $\{\bar{T}_{r,t}^{\Upsilon,\mu}\}_{t\geq 0}$ corresponding to the superposition form $\bar{\mathcal{E}}_r^{\Upsilon,\mu}$ can be obtained as the superposition of the semigroup $\{T_t^{\Upsilon(B_r),\mu_r^{\eta}}\}_{t\geq 0}$ associated with the form $\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}$. For the following proposition, we refer the reader to [Del21, (iii) Prop. 2.13]. **Proposition 4.10** ([Del21, (iii) Prop. 2.13]). The following holds:

(4.11)
$$\bar{T}_{r,t}^{\Upsilon,\mu}u(\gamma) = T_t^{\Upsilon(B_r),\mu_r^{\gamma}}u_r^{\gamma}(\gamma_{B_r}),$$

for μ -a.e. $\gamma \in \Upsilon$, any t > 0.

Remark 4.11. The proof of [Del21, (iii) Prop. 2.13] has been given in terms of direct integral in a general setting. As the measure μ_r^{η} can be identified to the conditional probability $\mu(\cdot | \cdot_{B_r^c} = \eta_{B_r^c})$ by a bi-measure-preserving isomorphism as remarked in (2.12), our setting is a particular case of direct integrals discussed in [Del21].

We now discuss the relation between $\mathcal{E}_r^{\Upsilon,\mu}$ and $\overline{\mathcal{E}}_r^{\Upsilon,\mu}$. As the former form is constructed as the smallest closed extension of $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{C}_r)$, it is clear by definition that

$$\mathcal{E}_r^{\Upsilon,\mu} = \bar{\mathcal{E}}_r^{\Upsilon,\mu} \quad \text{on} \quad \mathcal{C}_r \;, \quad \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}) \subset \mathcal{D}(\bar{\mathcal{E}}_r^{\Upsilon,\mu}) \;.$$

The following theorem proves that the opposite inclusion holds as well.

Theorem 4.12. $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu})) = (\bar{\mathcal{E}}_r^{\Upsilon,\mu}, \mathcal{D}(\bar{\mathcal{E}}_r^{\Upsilon,\mu})).$

Proof. The inclusion $\mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}) \subset \mathcal{D}(\bar{\mathcal{E}}_r^{\Upsilon,\mu})$ with the inequality $\bar{\mathcal{E}}_r^{\Upsilon,\mu} \leq \mathcal{E}_r^{\Upsilon,\mu}$ is straightforward by definition. Noting $\bar{\mathcal{E}}_r^{\Upsilon,\mu} = \mathcal{E}_r^{\Upsilon,\mu}$ on \mathcal{C}_r and $\mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu})$ is the closure of \mathcal{C}_r , it suffices to show that $\mathcal{C}_r \subset \mathcal{D}(\bar{\mathcal{E}}_r^{\Upsilon,\mu})$ is dense. Thanks to Lem. A.4, we only need to show that $\bar{T}_{r,t}^{\Upsilon,\mu}\mathcal{C}_r \subset \mathcal{C}_r$.

As $\bar{T}_{r,t}^{\mathbf{\Upsilon},\mu}$ is an L^{∞} -contraction semigroup by the sub-Markovian property of the semigroup (see, e.g., [MR90, Def. I.4.1]), we obtain $\bar{T}_{r,t}^{\mathbf{\Upsilon},\mu}\mathcal{C}_r \subset L^{\infty}(\mu)$, which verifies (a) in Def. 4.3

Verification of (b) in Def. 4.3. Let $u \in C_r$ and we show that $\overline{T}_{r,t}^{\Upsilon,\mu}u$ satisfies (b) in Def. 4.3. By Prop. 4.10, we can identify the following two operators:

$$\bar{T}_{r,t}^{\Upsilon,\mu}u = T_t^{\Upsilon(B_r),\mu_r}u_r^{\cdot}(\cdot_{B_r}) \ .$$

This implies that

$$\left(\bar{T}_{r,t}^{\Upsilon,\mu}u\right)_r^{\eta}(\cdot) = \bar{T}_{r,t}^{\Upsilon,\mu}u(\cdot+\eta_{B_r^c}) = T_t^{\Upsilon(B_r),\mu_r^{\eta}}u_r^{\eta}(\cdot) \ .$$

Take $k = k(\eta)$ as in (3.4). As the conditional probability μ_r^{η} is supported only on $\Upsilon^k(B_r)$, we only need to show

(4.12)
$$T_t^{\Upsilon(B_r),\mu_r^{k,\eta}} u_r^{\eta} \in \operatorname{Lip}_b(\Upsilon^k(B_r),\mathsf{d}_\Upsilon) .$$

As $(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon}, \mu_r^{k,\gamma})$ is $\mathsf{RCD}(0, \infty)$ for $k = k(\eta)$ for μ -a.e. η by Prop. 3.5, the corresponding semigroup satisfies $L^{\infty}(\mu_r^{k,\eta})$ -to- $\mathrm{Lip}_b(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon})$ -regularisation property ([AGS14a, Thm. 6.5]), which shows that for μ -a.e. η

$$T_t^{\Upsilon(B_r),\mu_r^{k,\eta}} v \in \operatorname{Lip}_b(\Upsilon^k(B_r), \mathsf{d}_\Upsilon) \quad \forall v \in L^\infty(\mu_r^{k,\eta})$$

and its Lipchitz constant is bounded as

$$\operatorname{Lip}_{\mathsf{d}_{\Upsilon}}(T_t^{\Upsilon(B_r),\mu_r^{k,\eta}}v) \le c(t,K) \|v\|_{L^{\infty}(\mu_r^{k,\eta})},$$

with constant c(t, K) depending only on t and the curvature bound K = 0 (to be more precise, $c(t, 0) = \frac{1}{\sqrt{2t}}$). This proves (4.12), which completes the verification of (b).

Verification of (c) in Def. 4.3. Let $u \in C_r$. Thanks to the verification of (b), the square field $\Gamma_r^{\Upsilon}(\bar{T}_{r,t}^{\Upsilon,\mu}u)$ is well-defined, and by (4.6) it holds that for μ -a.e. η

(4.13)
$$\Gamma_r^{\boldsymbol{\Upsilon}}(\bar{T}_{r,t}^{\boldsymbol{\Upsilon},\mu}u)(\gamma+\eta_{B_r^c}) = \Gamma^{\boldsymbol{\Upsilon}(B_r)}\big((\bar{T}_{r,t}^{\boldsymbol{\Upsilon},\mu}u)_r^{\eta}\big)(\gamma) \quad \mu_r^{\eta}\text{-a.e. } \gamma \in \boldsymbol{\Upsilon}(B_r) \ .$$

In view of the contraction property of the semigroup with respect to the form by general theory (see, e.g., [FOT11, p.23, Lem. 1.3.3]), viz.

$$\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}(T_t^{\Upsilon(B_r),\mu_r^{\eta}}u_r^{\eta}) \le \mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}}(u_r^{\eta})$$

as well as Prop. 4.10 and (4.13), we obtain

$$\begin{split} \frac{1}{2} \int_{\boldsymbol{\Upsilon}} \Gamma_r^{\boldsymbol{\Upsilon}}(\bar{T}_{r,t}^{\boldsymbol{\Upsilon},\mu}u) \, \mathrm{d}\mu &= \int_{\boldsymbol{\Upsilon}} \mathcal{E}^{\boldsymbol{\Upsilon}(B_r),\mu_r^{\eta}} \left((\bar{T}_{r,t}^{\boldsymbol{\Upsilon},\mu}u)_r^{\eta} \right) \, \mathrm{d}\mu(\eta) \\ &= \int_{\boldsymbol{\Upsilon}} \mathcal{E}^{\boldsymbol{\Upsilon}(B_r),\mu_r^{\eta}} (T_t^{\boldsymbol{\Upsilon}(B_r),\mu_r^{\eta}}u_r^{\eta}) \, \mathrm{d}\mu(\eta) \\ &\leq \int_{\boldsymbol{\Upsilon}} \mathcal{E}^{\boldsymbol{\Upsilon}(B_r),\mu_r^{\eta}}(u_r^{\eta}) \, \mathrm{d}\mu(\eta) \\ &= \mathcal{E}_r^{\boldsymbol{\Upsilon},\mu}(u) < \infty \; . \end{split}$$

The verification of (c) is completed. Therefore, we confirmed $\overline{T}_{r,t}^{\Upsilon,\mu}\mathcal{C}_r \subset \mathcal{C}_r$, which concludes the statement.

As a consequence of Thm. 4.12 and Prop. 4.10, we obtain the superposition formula for the semigroup $\{T_{r,t}^{\mathbf{\hat{\gamma}},\mu}\}_{t\geq 0}$ in terms of the semigroup $\{T_t^{\mathbf{\hat{\gamma}}(B_r),\mu_r^{\eta}}\}_{t\geq 0}$.

Corollary 4.13 (Coincidence of semigroups). The following three operators coincide:

(4.14)
$$T_{r,t}^{\Upsilon,\mu}u(\gamma) = \bar{T}_{r,t}^{\Upsilon,\mu}u(\gamma) = T_t^{\Upsilon(B_r),\mu_r^{\gamma}}u_r^{\gamma}(\gamma_{B_r}),$$

for μ -a.e. $\gamma \in \Upsilon$, any t > 0.

4.3. Monotone limit form. We now construct a Dirichlet form on Υ with sine_{β}-invariant measure μ as the monotone limit of $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu})$ as $r \to \infty$. The following proposition follows immediately from the definitions of the square field Γ_r^{Υ} and the core \mathcal{C}_r .

Proposition 4.14 (Monotonicity). The form $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}))$ and the square field Γ_r^{Υ} are monotone increasing as $r \uparrow \infty$, viz.,

$$\Gamma_r^{\Upsilon}(u) \leq \Gamma_s^{\Upsilon}(u) , \quad \mathcal{E}_r^{\Upsilon,\mu}(u) \leq \mathcal{E}_s^{\Upsilon,\mu}(u) , \quad \mathcal{D}(\mathcal{E}_s^{\Upsilon,\mu}) \subset \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}) \quad r \leq s .$$

Proof. As C_r is a core of the form $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}))$, it suffices to check $\mathcal{C}_s \subset \mathcal{C}_r$ and $\Gamma_r^{\Upsilon}(u) \leq \Gamma_s^{\Upsilon}(u)$ on \mathcal{C}_s . Let $u \in \mathcal{C}_s$ and we show $u \in \mathcal{C}_r$. By a simple reasoning similar to the proof of Lem. 4.7, we can see

 $u_r^\eta \in \operatorname{Lip}_b(\Upsilon(B_r), \mathsf{d}_\Upsilon) \quad \mu\text{-a.e. } \eta \quad \text{if} \ \ u_s^\eta \in \operatorname{Lip}_b(\Upsilon(B_s), \mathsf{d}_\Upsilon) \quad \mu\text{-a.e. } \eta \ .$

By Def. 4.2, it is straightforward to see $\Gamma_r^{\Upsilon}(u) \leq \Gamma_s^{\Upsilon}(u)$. Thus,

$$\mathcal{E}_{r}^{\Upsilon,\mu}(u) = \frac{1}{2} \int_{\Upsilon} \Gamma_{r}^{\Upsilon}(u) \,\mathrm{d}\mu \leq \frac{1}{2} \int_{\Upsilon} \Gamma_{s}^{\Upsilon}(u) \,\mathrm{d}\mu = \frac{1}{2} \int_{\Upsilon} \Gamma_{s}^{\Upsilon}(u) \,\mathrm{d}\mu = \mathcal{E}_{s}^{\Upsilon,\mu}(u) < \infty$$

Therefore, we conclude $u \in C_r$. The proof is completed.

We now define a Dirichlet form on Υ whose invariant measure is the $\operatorname{sine}_{\beta}$ measure μ by the monotone limit of $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}))$.

Definition 4.15 (Monotone limit form). The form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ is defined as the monotone limit:

(4.15)
$$\mathcal{D}(\mathcal{E}^{\Upsilon,\mu}) := \{ u \in \bigcap_{r>0} \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}) : \mathcal{E}^{\Upsilon,\mu}(u) = \lim_{r \to \infty} \mathcal{E}_r^{\Upsilon,\mu}(u) < \infty \}$$
$$\mathcal{E}^{\Upsilon,\mu}(u) := \lim_{r \to \infty} \mathcal{E}_r^{\Upsilon,\mu}(u) .$$

The form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ is a Dirichlet form on $L^2(\mu)$ as it is the monotone limit of Dirichlet forms (e.g., by [MR90, Exercise 3.9]). The square field Γ^{Υ} is defined as the monotone limit of Γ_r^{Υ} as well:

(4.16)
$$\Gamma^{\Upsilon}(u) := \lim_{r \to \infty} \Gamma^{\Upsilon}_r(u) \quad u \in \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}) .$$

The corresponding $L^2(\mu)$ -semigroup is denoted by $\{T_t^{\Upsilon,\mu}\}_{t\geq 0}$.

We now show that the form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ is a local Dirichlet form on $L^2(\mu)$ and satisfies the Rademacher-type property with respect to the L^2 -transportation-type distance $\bar{\mathsf{d}}_{\Upsilon}$.

Proposition 4.16. The form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ is a local Dirichlet form on $L^2(\mu)$. Furthermore, $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ satisfies Rademacher-type property:

(4.17)
$$\operatorname{Lip}(\bar{\mathsf{d}}_{\Upsilon},\mu) \subset \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}), \quad \Gamma^{\Upsilon}(u) \leq \operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(u)^2 \quad \forall u \in \operatorname{Lip}(\bar{\mathsf{d}}_{\Upsilon},\mu).$$

Proof. The local property of $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ follows from (4.16) and the locality of Γ_r^{Υ} for every r > 0. We show the Rademacher-type property. Since Γ^{Υ} is the limit square field of Γ_r^{Υ} as in (4.16), it suffices to show

$$\operatorname{Lip}(\bar{\mathsf{d}}_{\Upsilon},\mu) \subset \mathcal{C}_r \quad \text{and} \quad \Gamma_r^{\Upsilon}(u) \leq \operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(u)^2 \quad \forall u \in \operatorname{Lip}(\bar{\mathsf{d}}_{\Upsilon},\mu) \quad \forall r > 0 \ ,$$

which has been already proven in Prop. 4.8. We verified (4.17). The proof is complete.

Proposition 4.17. The semigroup $\{T_t^{\Upsilon,\mu}\}_{t\geq 0}$ is the $L^2(\mu)$ -strong operator limit of the semigroups $\{T_{r,t}^{\Upsilon,\mu}\}_{t\geq 0}$, viz.,

$$L^{2}(\mu) - \lim_{t \to \infty} T^{\mathbf{\Upsilon}, \mu}_{r, t} u = T^{\mathbf{\Upsilon}, \mu}_{t} u \quad \forall u \in L^{2}(\mu) , \quad t > 0 .$$

Furthermore, $\{T_t^{\Upsilon,\mu}\}_{t\geq 0}$ is also the $L^1(\mu)$ -strong operator limit of the semigroups $\{T_{r,t}^{\Upsilon,\mu}\}_{t\geq 0}$, viz.,

$$L^1(\mu) - \lim_{r \to \infty} T^{\Upsilon,\mu}_{r,t} u = T^{\Upsilon,\mu}_t u \quad \forall u \in L^1(\mu) , \quad t > 0 .$$

Proof. The first statement follows from the monotonicity of $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}) \text{ as } r \uparrow \infty$ proven in Prop. 4.14 and [RS80, S.14, p.373]. The latter statement is a standard consequence of the first statement for the strongly continuous Markovian contraction semigroup. We give a proof for the sake of the completeness. We note that the L^2 -operators $T_{r,t}^{\Upsilon,\mu}$ and $T_t^{\Upsilon,\mu}$ are uniquely extended to L^1 -strongly continuous Markovian contraction semigroups (see, e.g, [FOT11, (1.5.2) in §1.5]). As $L^1(\mu) \cap L^2(\mu)$ is dense in $L^1(\mu)$, for any $u \in L^1(\mu)$ and $\varepsilon > 0$, there exists $u_{\varepsilon} \in L^1(\mu) \cap L^2(\mu)$ so that $||u - u_{\varepsilon}||_{L^1(\mu)} < \varepsilon$ and

$$\begin{aligned} \|T_{r,t}^{\Upsilon,\mu}u - T_{t}^{\Upsilon,\mu}u\|_{L^{1}(\mu)} \\ &\leq \|T_{r,t}^{\Upsilon,\mu}u - T_{r,t}^{\Upsilon,\mu}u_{\varepsilon}\|_{L^{1}(\mu)} + \|T_{r,t}^{\Upsilon,\mu}u_{\varepsilon} - T_{t}^{\Upsilon,\mu}u_{\varepsilon}\|_{L^{1}(\mu)} + \|T_{t}^{\Upsilon,\mu}u_{\varepsilon} - T_{t}^{\Upsilon,\mu}u\|_{L^{1}(\mu)} \\ &\leq \|u - u_{\varepsilon}\|_{L^{1}(\mu)} + \|T_{r,t}^{\Upsilon,\mu}u_{\varepsilon} - T_{t}^{\Upsilon,\mu}u_{\varepsilon}\|_{L^{2}(\mu)} + \|u_{\varepsilon} - u\|_{L^{1}(\mu)} \end{aligned}$$

$$\xrightarrow{r \to \infty} \varepsilon + 0 + \varepsilon \; .$$

As $\varepsilon > 0$ is arbitrarily small, the proof is completed.

Corollary 4.18. For any fixed r > 0 and $u \in \mathcal{D}(\mathcal{E}^{\Upsilon,\mu})$,

(4.18)
$$T_{r',t}^{\Upsilon,\mu} u \xrightarrow{r' \to \infty} T_t^{\Upsilon,\mu} u \quad weakly \text{ in } \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}) \ .$$

In particular,

(4.19)
$$\int_{\mathbf{\Upsilon}} \Gamma_r^{\mathbf{\Upsilon}}(T_t^{\mathbf{\Upsilon},\mu}u)h \,\mathrm{d}\mu \leq \liminf_{r' \to \infty} \int_{\mathbf{\Upsilon}} \Gamma_r^{\mathbf{\Upsilon}}(T_{r',t}^{\mathbf{\Upsilon},\mu}u)h \,\mathrm{d}\mu ,$$

for all non-negative $h \in \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}) \cap L^{\infty}(\mu)$.

Proof. First of all, (4.18) is well-posed as $T_{r',t}^{\Upsilon,\mu}u, T_t^{\Upsilon,\mu}u \in \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu})$ thanks to the inclusion $\mathcal{D}(\mathcal{E}^{\Upsilon,\mu}) \subset \bigcap_{r>0} \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu})$ and the monotonicity $\mathcal{D}(\mathcal{E}_{r'}^{\Upsilon,\mu}) \subset \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu})$ for $r \leq r'$. By the monotonicity proven in Prop. 4.14 and the general contraction property of the semigroup with respect to the Dirichlet form, it holds that for $r \leq r'$

$$\mathcal{E}_{r}^{\Upsilon,\mu}(T_{r',t}^{\Upsilon,\mu}u) \leq \mathcal{E}_{r'}^{\Upsilon,\mu}(T_{r',t}^{\Upsilon,\mu}u) \leq \mathcal{E}_{r'}^{\Upsilon,\mu}(u) \leq \mathcal{E}^{\Upsilon,\mu}(u) < \infty$$

Noting also that the semigroup $T_{r',t}^{\Upsilon,\mu}$ contracts the $L^2(\mu)$ -norm by a general property of semigroups, $\{T_{r',t}^{\Upsilon,\mu}u\}_{r'}$ is bounded in $\mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu})$. Thanks to (4.17), $\{T_{r',t}^{\Upsilon,\mu}u\}_{r'}$ converges to $T_t^{\Upsilon,\mu}u$ weakly in $\mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu})$. The latter statement is a consequence of the first statement, see, e.g., [HR03, Lem. 2.4].

4.4. Bakry–Émery Curvature bound for $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$. In this subsection, we prove the Bakry–Émery curvature bound for the local Dirichlet form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$.

Theorem 4.19. Let $\beta > 0$ and μ be the sine_{β} ensemble. The local Dirichlet form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ satisfies the Bakry-Émery curvature dimension condition $\mathsf{BE}(0,\infty)$:

$$(\mathsf{BE}(0,\infty)) \qquad \Gamma^{\Upsilon}(T_t^{\Upsilon,\mu}u) \le T_t^{\Upsilon,\mu}(\Gamma^{\Upsilon}(u)) \quad \forall u \in \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}) \quad \forall t > 0$$

Proof. We first prove $\mathsf{BE}(0,\infty)$ for the form $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}))$. Let $u \in \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu})$. By Prop. 3.5, by the expression (3.4) of μ_r^{η} in terms of $\mu_r^{k,\eta}$ and by the definition (4.4) of $\Gamma^{\Upsilon(B_r)}$, there exists $\Xi_r^1 \subset \Upsilon$ with $\mu(\Xi_r^1) = 1$ so that for every $\eta \in \Xi_r^1$ there exists a measurable set $\Omega_r^{1,\eta} \subset \Upsilon(B_r)$ with $\mu_r^{\eta}(\Omega_r^{1,\eta}) = 1$ satisfying that for every $\gamma \in \Omega_r^{1,\eta}$, the following Bakry-Émery gradient estimate holds:

(4.20)
$$\Gamma^{\boldsymbol{\Upsilon}(B_r)}(T_t^{\boldsymbol{\Upsilon}(B_r),\mu_r^{\eta}}u_r^{\eta})(\gamma) \le T_t^{\boldsymbol{\Upsilon}(B_r),\mu_r^{\eta}}(\Gamma^{\boldsymbol{\Upsilon}(B_r)}(u_r^{\eta}))(\gamma)$$

By Prop. 4.8, there exists $\Xi_r^2 \subset \Upsilon$ with $\mu(\Xi_r^2) = 1$ so that for every $\eta \in \Xi_r^2$ there exists a measurable set $\Omega_r^{2,\eta} \subset \Upsilon(B_r)$ with $\mu_r^{\eta}(\Omega_r^{2,\eta}) = 1$ satisfying that for every $\gamma \in \Omega_r^{2,\eta}$

(4.21)
$$\Gamma_r^{\Upsilon}(T_{r,t}^{\Upsilon,\mu}u)(\gamma+\eta_{B_r^c}) = \Gamma^{\Upsilon(B_r)}\Big(\Big(T_{r,t}^{\Upsilon,\mu}u\Big)_r^{\eta}\Big)(\gamma) + \Gamma_r^{\Upsilon}(u)(\gamma+\eta_{B_r^c}) = \Gamma^{\Upsilon(B_r)}(u_r^{\eta})(\gamma) .$$

By Cor. 4.13, there exists $\Lambda_r^3 \subset \Upsilon$ with $\mu(\Lambda_r^3) = 1$ so that for every $\gamma \in \Lambda_r^3$

(4.22)
$$T_{r,t}^{\Upsilon,\mu}u(\gamma) = T_t^{\Upsilon(B_r),\mu_r^{\gamma}}u_r^{\gamma}(\gamma) \; .$$

26

By the standard disintegration argument, we can write

$$\Lambda^3_r = igcup_{\eta\in \Xi^3_r} \mathrm{pr}_r^{-1}(\Omega^{3,\eta}_r) \cap \mathbf{\Upsilon}^\eta_r$$
 .

where $\Omega_r^{3,\eta} = (\Lambda_r^3)_r^{\eta} := \{\gamma \in \Upsilon(B_r) : \gamma + \eta_{B_r^c} \in \Lambda_r^3\}$ and $\Xi_r^3 = \operatorname{pr}_{B_r^c}(\Lambda_r^3)$, and Υ_r^{η} has been defined in (2.10). By the disintegration formula (2.16), $\mu(\Xi_r^3) = 1$ and $\mu_r^{\eta}(\Omega_r^{3,\eta}) = 1$ for every $\eta \in \Xi_r^3$.

Let $\Xi_r := \Xi_r^1 \cap \Xi_r^2 \cap \Xi_r^3$ and $\Omega_r^\eta := \Omega_r^{1,\eta} \cap \Omega_r^{2,\eta} \cap \Omega_r^{3,\eta}$ for $\eta \in \Xi_r$. Set $\mathbf{K}_r := \bigcup_{\eta \in \Xi_r} \mathrm{pr}_r^{-1}(\Omega_r^\eta) \cap \mathbf{\Upsilon}_r^\eta .$

By construction, $\mu(\Xi_r) = 1$ and $\mu_r^{\eta}(\Omega_r^{\eta}) = 1$ for every $\eta \in \Xi_r$. By (4.20), (4.21) and (4.22), the following inequalities hold for every $\gamma \in K_r$:

(4.23)
$$\Gamma_{r}^{\Upsilon}(T_{r,t}^{\Upsilon,\mu}u)(\gamma) = \Gamma_{r}^{\Upsilon}(T_{r,t}^{\Upsilon,\mu}u)(\gamma_{B_{r}} + \gamma_{B_{r}^{c}})$$
$$= \Gamma^{\Upsilon(B_{r})}((T_{r,t}^{\Upsilon,\mu}u)_{r}^{\gamma})(\gamma_{B_{r}})$$
$$\leq T_{t}^{\Upsilon(B_{r}),\mu_{r}^{\gamma}}\Gamma^{\Upsilon(B_{r})}(u_{r}^{\gamma})(\gamma_{B_{r}})$$
$$= T_{t}^{\Upsilon(B_{r}),\mu_{r}^{\gamma}}\left(\Gamma_{r}^{\Upsilon}(u)_{r}^{\gamma}\right)(\gamma_{B_{r}})$$
$$= T_{r,t}^{\Upsilon,\mu}\Gamma_{r}^{\Upsilon}(u)(\gamma) .$$

Let $\Theta := \{\gamma \in \Upsilon : \Gamma_r^{\Upsilon}(T_{r,t}^{\Upsilon,\mu}u)(\gamma) \leq T_{r,t}^{\Upsilon,\mu}\Gamma_r^{\Upsilon}(u)(\gamma)\}$. Then Θ is μ -measurable by construction, and thanks to (4.23), it holds that $K_r \subset \Theta$. By applying Lem. A.2, we obtain $\mu(\Theta) = 1$, which concludes $\mathsf{BE}(0,\infty)$ for the truncated form $(\mathcal{E}_r^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}))$ for any r > 0.

We now prove $\mathsf{BE}(0,\infty)$ of the form $(\mathcal{E}^{\Upsilon,\mu},\mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$. It suffices to prove

(4.24)
$$\int_{\mathbf{\Upsilon}} \Gamma^{\mathbf{\Upsilon}}(T_t^{\mathbf{\Upsilon},\mu}u)h \,\mathrm{d}\mu \leq \int_{\mathbf{\Upsilon}} T_t^{\mathbf{\Upsilon},\mu}\Gamma^{\mathbf{\Upsilon}}(u)h \,\mathrm{d}\mu \;,$$

for all non-negative $h \in \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}) \cap L^{\infty}(\mu)$. Indeed, thanks to the Rademacher-type property proven in Prop. 4.8, we have

$$\operatorname{Lip}_{b,+}(\mathsf{d}_{\Upsilon}) \subset \mathcal{D}(\mathcal{E}_r^{\Upsilon,\mu}) \cap L^{\infty}_+(\mu).$$

As $\operatorname{Lip}_{b,+}(\mathsf{d}_{\Upsilon}) \cap C(\tau_{v})$ is point separating (see [DS21a, (a) in Rem. 5.13]), it is separating (i.e., measure determining) by e.g., [EK86, p.113, (a) in Thm. 4.5 in Chap. 3]. Thus, the inequality (4.24) implies $\Gamma^{\Upsilon}(T_{t}^{\Upsilon,\mu}u) \leq T_{t}^{\Upsilon,\mu}\Gamma^{\Upsilon}(u)$. We now prove (4.24). Let $u \in \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}) \subset \bigcap_{r>0} \mathcal{D}(\mathcal{E}_{r}^{\Upsilon,\mu})$. By making use of the monotonicity $\Gamma_{r}^{\Upsilon} \leq \Gamma_{r'}^{\Upsilon}$ for $r \leq r'$ (we will use it in the following displayed formulas in the first equality and in the second inequality), Cor. 4.18 (used in the first inequality below), BE(0, ∞) for the truncated form $(\mathcal{E}_{r}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}_{r}^{\Upsilon,\mu}))$ for any r > 0 (used in the third inequality below), and the convergence of $T_{r',t}^{\Upsilon,\mu}$ to $T_{t}^{\Upsilon,\mu}$ as $r' \to \infty$ in the L^{1} -strong operator sense by Prop. 4.17 (used in the last equality), the following inequalities hold true:

$$\int_{\Upsilon} \Gamma^{\Upsilon}(T_t^{\Upsilon,\mu}u)h \, \mathrm{d}\mu = \int_{\Upsilon} \lim_{r \to \infty} \Gamma_r^{\Upsilon}(T_t^{\Upsilon,\mu}u)h \, \mathrm{d}\mu$$
$$= \lim_{r \to \infty} \int_{\Upsilon} \Gamma_r^{\Upsilon}(T_t^{\Upsilon,\mu}u)h \, \mathrm{d}\mu$$

$$\leq \limsup_{r \to \infty} \liminf_{r' \to \infty} \int_{\mathbf{\Upsilon}} \Gamma_r^{\mathbf{\Upsilon}} (T_{r',t}^{\mathbf{\Upsilon},\mu} u) h \, \mathrm{d}\mu$$

$$\leq \limsup_{r' \to \infty} \int_{\mathbf{\Upsilon}} \Gamma_{r'}^{\mathbf{\Upsilon}} (T_{r',t}^{\mathbf{\Upsilon},\mu} u) h \, \mathrm{d}\mu$$

$$\leq \limsup_{r' \to \infty} \int_{\mathbf{\Upsilon}} T_{r',t}^{\mathbf{\Upsilon},\mu} \Gamma_{r'}^{\mathbf{\Upsilon}} (u) h \, \mathrm{d}\mu$$

$$= \int_{\mathbf{\Upsilon}} T_t^{\mathbf{\Upsilon},\mu} \Gamma^{\mathbf{\Upsilon}} (u) h \, \mathrm{d}\mu .$$

The last equality in the above displayed formulas followed by the L^1 -contraction property of $T_{r',t}^{\Upsilon,\mu}$ and the second statement of Prop. 4.17:

$$\begin{split} & \left\| T_{r',t}^{\boldsymbol{\Upsilon},\mu} \Gamma_{r'}^{\boldsymbol{\Upsilon}}(u) - T_{t}^{\boldsymbol{\Upsilon},\mu} \Gamma^{\boldsymbol{\Upsilon}}(u) \right\|_{L^{1}(\mu)} \\ & = \left\| T_{r',t}^{\boldsymbol{\Upsilon},\mu} \Gamma_{r'}^{\boldsymbol{\Upsilon}}(u) - T_{r',t}^{\boldsymbol{\Upsilon},\mu} \Gamma^{\boldsymbol{\Upsilon}}(u) \right\|_{L^{1}(\mu)} + \left\| T_{r',t}^{\boldsymbol{\Upsilon},\mu} \Gamma^{\boldsymbol{\Upsilon}}(u) - T_{t}^{\boldsymbol{\Upsilon},\mu} \Gamma^{\boldsymbol{\Upsilon}}(u) \right\|_{L^{1}(\mu)} \\ & \leq \left\| \Gamma_{r'}^{\boldsymbol{\Upsilon}}(u) - \Gamma^{\boldsymbol{\Upsilon}}(u) \right\|_{L^{1}(\mu)} + \left\| T_{r',t}^{\boldsymbol{\Upsilon},\mu} \Gamma^{\boldsymbol{\Upsilon}}(u) - T_{t}^{\boldsymbol{\Upsilon},\mu} \Gamma^{\boldsymbol{\Upsilon}}(u) \right\|_{L^{1}(\mu)} \xrightarrow{r' \to \infty} 0 \; . \end{split}$$

We verified (4.24), the proof is completed.

4.5. Integral Bochner and local Poicaré inequalities. As an application of $\mathsf{BE}(0,\infty)$ proven in Thm. 4.19, we show several functional inequalities. We define the integral Γ_2 -operator as follows:

(4.25)
$$\Gamma_{2}^{\Upsilon,\mu}(u,\varphi) := \int_{\Upsilon} \left(\frac{1}{2} \Gamma^{\Upsilon}(u) A^{\Upsilon,\mu} \varphi - \Gamma^{\Upsilon}(u, A^{\Upsilon,\mu}u) \varphi \right) d\mu ,$$
$$\mathcal{D}(\Gamma_{2}^{\Upsilon,\mu}) := \left\{ (u,\varphi) : \mathcal{D}(A^{\Upsilon,\mu})^{\times 2} : A^{\Upsilon,\mu}u \in \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}), \ \varphi, A^{\Upsilon,\mu}u \in L^{\infty}(\mu) \right\} ,$$

where $A^{\Upsilon,\mu}$ denotes the $L^2(\mu)$ -infinitesimal generator associated with the form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$.

Corollary 4.20. Let μ be the sine_{β} ensemble with $\beta > 0$. The following hold:

(a) (Integral Bochner inequality) for every $(u, \varphi) \in \mathcal{D}(\Gamma_2^{\Upsilon, \mu})$

$$\Gamma_2^{\Upsilon,\mu}(u,\varphi) \ge 0 ;$$

(b) (local Poincaré inequality) for $u \in \mathcal{D}(\mathcal{E}^{\Upsilon,\mu})$ and t > 0,

$$\begin{split} T_t^{\Upsilon,\mu} u^2 &- (T_t^{\Upsilon,\mu} u)^2 \leq 2t T_t^{\Upsilon,\mu} \Gamma^{\Upsilon}(u) \ , \\ T_t^{\Upsilon,\mu} u^2 &- (T_t^{\Upsilon,\mu} u)^2 \geq 2t \Gamma^{\Upsilon}(T_t^{\Upsilon,\mu} u) \ ; \end{split}$$

Proof. The statement (a) follows from $\mathsf{BE}(0,\infty)$ proven in Thm. 4.19 and [AGS15, Cor. 2.3]. The statement (b) is a consequence of $\mathsf{BE}(0,\infty)$, see e.g., [BGL14, Thm. 4.7.2].

Remark 4.21 (Local spectral gap inequality). Let $P_t^{\Upsilon,\mu}(\gamma, d\eta)$ denote the heat kernel measure, which is defined as (see, e.g., [BGL14, (1.2.4) in p.12])

$$T_t^{\boldsymbol{\Upsilon},\mu}u(\gamma) = \int_{\boldsymbol{\Upsilon}} u(\eta) P_t^{\boldsymbol{\Upsilon},\mu}(\gamma,\mathrm{d}\eta) \qquad \mu\text{-a.e. } \gamma$$

The local Poincaré inequality in Cor. 4.20 is the spectral gap inequality with respect to the heat kernel measure $P_t^{\Upsilon,\mu}(\gamma, \mathrm{d}\eta)$:

$$\int_{\mathbf{\Upsilon}} \left| u(\eta) - \int_{\mathbf{\Upsilon}} u(\eta) P_t^{\mathbf{\Upsilon},\mu}(\gamma,\mathrm{d}\eta) \right|^2 P_t^{\mathbf{\Upsilon},\mu}(\gamma,\mathrm{d}\eta) \le 2t \int_{\mathbf{\Upsilon}} \Gamma^{\mathbf{\Upsilon},\mu}(u)(\eta) P_t^{\mathbf{\Upsilon},\mu}(\gamma,\mathrm{d}\eta) \ .$$

The name "local" stems from the observation that $P_t^{\Upsilon,\mu}(\gamma, d\eta)$ is expected to be *concentrated around* γ when t is small (see [BGL14, §4.7 in p. 206]).

The following corollary provides a non-trivial tail estimate of the heat kernel measure $P_t^{\Upsilon,\mu}(\gamma, \mathrm{d}\eta)$, which decays sufficiently fast at the tail to make every (not necessarily bounded) 1-Lipschitz function exponentially integrable.

Corollary 4.22 (Exponential integrability of 1-Lipschitz functions). Let μ be the sine_{β} ensemble with $\beta > 0$. If u is a \bar{d}_{Υ} -Lipschitz function with $\operatorname{Lip}_{\bar{d}_{\Upsilon}}(u) \leq 1$ and $|u(\gamma)| < \infty$ μ -a.e. γ , then for every $s < \sqrt{2/t}$

$$\int_{\Upsilon} e^{su(\eta)} P_t^{\Upsilon,\mu}(\gamma,\mathrm{d}\eta) < \infty \; .$$

Proof. By the Rademacher-type theorem with respect to \bar{d}_{Υ} proven in Prop. 4.16 and the local Poincaré inequality in Cor. 4.20, the same proof works as in [BGL14, Prop. 4.4.2].

5. DIMENSION-FREE AND LOG HARNACK INEQUALITIES

In this section, we prove functional inequalities involving the Bakry–Émery curvature bound $\mathsf{BE}(0,\infty)$ and the L^2 -transportation-type extended distance $\bar{\mathsf{d}}_{\Upsilon}$.

Theorem 5.1. Let μ be the sine_{β} ensemble with $\beta > 0$. Then the following inequalities hold:

(a) (log-Harnack inequality) for every non-negative $u \in L^{\infty}(\Upsilon, \mu) \varepsilon \in (0, 1], t > 0$, there exists $\Omega \subset \Upsilon$ so that $\mu(\Omega) = 1$ and

$$T_t^{\mathbf{T},\mu}\log(u+\varepsilon)(\gamma) \le \log(T_t^{\mathbf{T},\mu}u(\eta)+\varepsilon) + \bar{\mathsf{d}}_{\mathbf{\Upsilon}}(\gamma,\eta)^2 , \quad every \ \gamma,\eta\in\Omega ;$$

(b) (dimension-free Harnack inequality) for every non-negative $u \in L^{\infty}(\Upsilon, \mu)$, $t > 0 \text{ and } \alpha > 1 \text{ there exists } \Omega \subset \Upsilon \text{ so that } \mu(\Omega) = 1 \text{ and}$

$$(T_t^{\Upsilon,\mu}u)^{\alpha}(\gamma) \leq T_t^{\Upsilon,\mu}u^{\alpha}(\eta) \exp\Big\{\frac{\alpha}{2(\alpha-1)}\bar{\mathsf{d}}_{\Upsilon}(\gamma,\eta)^2\Big\} , \quad for \; every \; \gamma,\eta \in \Omega \; ;$$

(c) (Lipschitz contraction) For $u \in \text{Lip}_b(\bar{d}_{\Upsilon}, \mu)$ and t > 0,

$$T_t^{\Upsilon,\mu}u$$
 has a $\bar{\mathsf{d}}_{\Upsilon}$ -Lipschitz μ -modification $\tilde{T}_t^{\Upsilon,\mu}u$

and the following estimate holds:

$$\operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(\tilde{T}_t^{\Upsilon,\mu}u) \leq \operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(u) ;$$

(d) $(L^{\infty}$ -to-Lip regularisation) For $u \in L^{\infty}(\mu)$ and any t > 0,

 $T_t^{\mathbf{\Upsilon},\mu}u$ has a $\bar{\mathsf{d}}_{\mathbf{\Upsilon}}$ -Lipschitz μ -modification $\tilde{T}_t^{\mathbf{\Upsilon},\mu}u$

and the following estimate holds:

$$\operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(\tilde{T}_t^{\Upsilon,\mu}u) \leq \frac{1}{\sqrt{2t}} \|u\|_{L^{\infty}(\mu)}$$

Proof. We prove (a). By the relation between $T_{r,t}^{\Upsilon,\mu}$ and $T_t^{\Upsilon(B_r),\mu_r}(\cdot_{B_r})$ in Prop. 4.13, there exists a measurable set $\Omega_{\mathsf{sem}}^r \subset \Upsilon$ with $\mu(\Omega_{\mathsf{sem}}^r) = 1$ so that for every $\eta \in \Omega_{\mathsf{sem}}^r$

(5.1)
$$T_{r,t}^{\Upsilon,\mu}(\eta) = T_t^{\Upsilon(B_r),\mu_r^{\eta}}(\eta_{B_r}) \ .$$

Let $u \in L^{\infty}(\mu)$. Thanks to Lem. A.3, there exists $\Omega_{\infty}^r \subset \Upsilon$ so that $\mu(\Omega_{\infty}^r) = 1$ and

$$u_r^\eta \in L^\infty(\mu_r^\eta), \quad \forall \eta \in \Omega_\infty^r, \quad \forall r \in \mathbb{N}$$

By Prop. 3.5, there exists a measurable set $\Omega_{\mathsf{rcd}}^r \subset \Upsilon$ so that $\mu(\Omega_{\mathsf{rcd}}^r) = 1$ and $(\Upsilon^k, \mathsf{d}_\Upsilon, \mu_r^\eta)$ is $\mathsf{RCD}(0, \infty)$ with $k = k(\eta)$ as in (3.4) for every $\eta \in \Omega_{\mathsf{rcd}}^r$.

Let $\Omega^r := \Omega^r_{sem} \cap \Omega^r_{\infty} \cap \Omega^r_{rcd}$. As the log-Harnack inequality holds in RCD spaces (see, [AGS15, Lem. 4.6]), the following holds for every $\eta \in \Omega^r$ and $k = k(\eta)$ and $\varepsilon \in (0, 1]$

(5.2)
$$T_t^{\mathbf{\Upsilon}^k(B_r),\mu_r^{k,\eta}}\log(u_r^{\eta}+\varepsilon)(\gamma) \le \log\left(T_t^{\mathbf{\Upsilon}^k(B_r),\mu_r^{k,\eta}}u_r^{\eta}(\zeta)+\varepsilon\right) + \mathsf{d}_{\mathbf{\Upsilon}}(\gamma,\zeta)^2 ,$$

for every $\gamma, \zeta \in \Upsilon^k(B_r)$.

Noting the convergence of the semigroups $\{T_{r,t}^{\Upsilon,\mu}\}_{t\geq 0}$ to $\{T_t^{\Upsilon,\mu}\}_{t\geq 0}$ in the $L^2(\mu)$ -operator sense by Prop. 4.17, there exist $\Omega_{\mathsf{con}} \subset \Upsilon$ with $\mu(\Omega_{\mathsf{con}}) = 1$ and a (non-relabelled) subsequence of $\{r\}$ so that for every $\gamma \in \Omega_{\mathsf{con}}$

(5.3)
$$T_{r,t}^{\Upsilon,\mu} \log(u+\varepsilon)(\gamma) \xrightarrow{r \to \infty} T_t^{\Upsilon,\mu} \log(u+\varepsilon)(\gamma) ,$$
$$\log(T_{r,t}^{\Upsilon,\mu}u(\gamma)+\varepsilon) \xrightarrow{r \to \infty} \log(T_t^{\Upsilon,\mu}u(\gamma)+\varepsilon) .$$

Let $\Omega' = \Omega_{\text{con}} \cap_{r \in \mathbb{N}} \Omega^r$, which by construction satisfies $\mu(\Omega') = 1$. Our goal is now to prove that there exists $\Omega \subset \Omega'$ with $\mu(\Omega) = 1$ so that

(5.4)
$$T_t^{\Upsilon,\mu} \log(u+\varepsilon)(\gamma) \le \log(T_t^{\Upsilon,\mu}u(\eta)+\varepsilon) + \bar{\mathsf{d}}_{\Upsilon}(\gamma,\eta)^2$$
, every $\gamma,\eta\in\Omega$.

Thanks to (5.3), Formula (5.4) comes down to the corresponding inequality for the semigroup $\{T_{r,t}^{\Upsilon,\mu}\}_{t\geq 0}$ for any r > 0:

(5.5)
$$T_{r,t}^{\Upsilon,\mu}\log(u+\varepsilon)(\gamma) \le \log(T_{r,t}^{\Upsilon,\mu}u(\eta)+\varepsilon) + \bar{\mathsf{d}}_{\Upsilon}(\gamma,\eta)^2$$
, every $\gamma,\eta\in\Omega$.

We prove (5.5) by contradiction. Suppose that for any $\Omega \subset \Omega'$ with $\mu(\Omega) = 1$, there exists $\gamma, \eta \in \Omega$ so that

(5.6)
$$T_{r,t}^{\Upsilon,\mu}\log(u+\varepsilon)(\gamma) \ge \log(T_{r,t}^{\Upsilon,\mu}u(\eta)+\varepsilon) + \bar{\mathsf{d}}_{\Upsilon}(\gamma,\eta)^2$$

We may assume that $\bar{d}_{\Upsilon}(\gamma, \eta) < \infty$ without loss of generality. Thus, by (2.21), there exists r > 0 so that

(5.7)
$$\gamma_{B_r^c} = \eta_{B_r^c} , \quad \gamma(B_r) = \eta(B_r) .$$

By making use of (5.1), (5.2), (5.7), we obtain

(5.8)
$$T_{r,t}^{\Upsilon,\mu} \log(u+\varepsilon)(\gamma) = T_{r,t}^{\Upsilon,\mu} \log(u+\varepsilon)(\gamma_{B_r}+\gamma_{B_r^c}) = T_t^{\Upsilon(B_r),\mu_r^{\gamma}} \log(u_r^{\gamma}+\varepsilon)(\gamma_{B_r}) \leq \log(T_t^{\Upsilon(B_r),\mu_r^{\gamma}}u_r^{\gamma}(\eta_{B_r})+\varepsilon) + \mathsf{d}_{\Upsilon}(\gamma_{B_r},\eta_{B_r})^2 = \log(T_{r,t}^{\Upsilon,\mu}u(\eta)+\varepsilon) + \bar{\mathsf{d}}_{\Upsilon}(\gamma,\eta)^2 ,$$

which contradicts (5.6), therefore, the proof of (a) is completed.

The proof of (b) follows precisely in the same strategy as above by replacing $T_t^{\Upsilon,\mu} \log(u + \varepsilon)$, $\log(T_t^{\Upsilon,\mu}u + \varepsilon)$ and $\bar{d}_{\Upsilon}(\gamma,\eta)^2$ by $(T_t^{\Upsilon,\mu}u)^{\alpha}$, $T_t^{\Upsilon,\mu}u^{\alpha}$ and $\frac{\alpha}{2(\alpha-1)}\bar{d}_{\Upsilon}(\gamma,\eta)^2$ respectively, and noting that the dimension-free Harnack inequality holds on $\mathsf{RCD}(K,\infty)$ spaces ([Li15, Thm. 3.1]).

30

The proof of (c): Note that $u_r^{\eta} \in \operatorname{Lip}(\Upsilon(B_r), \mathsf{d}_{\Upsilon})$ whenever $u \in \operatorname{Lip}(\Upsilon, \overline{\mathsf{d}}_{\Upsilon})$ and $\operatorname{Lip}_{\mathsf{d}_{\Upsilon}}(u_r^{\eta}) \leq \operatorname{Lip}_{\overline{\mathsf{d}}_{\Upsilon}}(u)$ by Lem. 4.7. Note also that the sought conclusion of (c) can be rephrased as

$$\tilde{T}_t^{\Upsilon,\mu}u(\gamma) - \tilde{T}_t^{\Upsilon,\mu}u(\eta) \leq \operatorname{Lip}_{\bar{\mathsf{d}}_\Upsilon}(u)\bar{\mathsf{d}}_\Upsilon(\gamma,\eta) \quad \forall \gamma,\eta \in \Upsilon \ .$$

Thus, by the same proof strategy as in (a) replacing $T_t^{\Upsilon,\mu} \log(u+\varepsilon)(\gamma)$ and $\log(T_t^{\Upsilon,\mu}u(\eta)+\varepsilon)$ with $T_t^{\Upsilon,\mu}u(\gamma)$ and $T_t^{\Upsilon,\mu}u(\eta)$, and noting that the Lipschitz contraction property holds on RCD spaces ([AGS14b, (iv) in Thm. 6.1]), we conclude that there exists $\Omega \subset \Upsilon$ with $\mu(\Omega) = 1$ so that

$$T_t^{\Upsilon,\mu}(\gamma) - T_t^{\Upsilon,\mu}(\eta) \leq \operatorname{Lip}_{\bar{\mathsf{d}}_\Upsilon}(u) \bar{\mathsf{d}}_\Upsilon(\gamma,\eta) \quad \forall \gamma, \eta \in \Omega \ .$$

The conclusion now follows from the McShane extension Theorem (for extended metric spaces, see [DS21b, Lem. 2.1]).

The proof of (d) is the same as that of (c) but using the L^{∞} -to-Lip property ([AGS14b, Thm. 6.5]) in $\mathsf{RCD}(K, \infty)$ spaces instead of [AGS14b, (iv) in Thm. 6.1]). The proof is complete.

Corollary 5.2. Let μ be the sine_{β} ensemble with $\beta > 0$. Then

$$\operatorname{Lip}_b(\mathsf{d}_{\Upsilon},\mu)$$
 is dense in $\mathcal{D}(\mathcal{E}^{\Upsilon,\mu})$

Proof. $\operatorname{Lip}_b(\bar{\mathsf{d}}_{\Upsilon}, \mu)$ is dense in $L^2(\mu)$ as noted in the fourth paragraph of the proof of Prop. 4.8. Thus, the statement follows from (c) of Thm. 5.1 and Lem. A.4.

6. GRADIENT FLOW

In this section, as an application of Thm. 4.19, we discuss properties of the dual flow in the space $\mathcal{P}(\Upsilon)$ of Borel probability measures on Υ induced by the transition semigroup $T_t^{\Upsilon,\mu}$ of the infinite Dyson Brownian motions. In particular, the Dyson Brownian motion is identified to the unique $W_{\mathcal{E}}$ -gradient flow of the Boltzmann–Shannon entropy associated with $\mu = \operatorname{sine}_{\beta}$, where $W_{\mathcal{E}}$ is a certain Benamou–Brenier type extended distance on $\mathcal{P}(\Upsilon)$.

Boltzmann-Shannon entropy and Fisher information. Let $(\mathcal{P}(\Upsilon), \tau_w)$ be the space of all Borel probability measures on (Υ, τ_v) equipped with the weak topology τ_w , i.e., the topology induced by the duality of $C_b(\Upsilon, \tau_v)$. Let $\mathcal{P}_{\mu}(\Upsilon)$ be the subspace of $\mathcal{P}(\Upsilon)$ consisting of measures absolutely continuous with respect to μ . We write $\nu = \rho \cdot \mu$ if $\rho = \frac{d\nu}{d\mu}$.

• The Boltzmann-Shannon entropy $\mathsf{Ent}_{\mu}: \mathcal{P}_{\mu}(\Upsilon) \to \mathbb{R} \cup \{+\infty\}$ is defined as

$$\mathsf{Ent}_{\mu}(\nu) := \int_{\mathbf{\Upsilon}} \rho \log \rho \, \mathrm{d}\mu \;, \quad \nu = \rho \cdot \mu \;.$$

The domain of Ent_{μ} is denoted by $\mathcal{D}(\mathsf{Ent}_{\mu}) := \{\nu \in \mathcal{P}_{\mu}(\Upsilon) : \mathsf{Ent}_{\mu}(\nu) < \infty\}.$

• The Fisher information $\mathsf{F}_{\mu}:\mathcal{P}_{\mu}(\Upsilon)\to\mathbb{R}\cup\{+\infty\}$ is defined as

$$\mathsf{F}_{\mu}(\nu) := 8\mathcal{E}^{\Upsilon,\mu}(\sqrt{\rho}) \;, \quad \nu = \rho \cdot \mu \;.$$

The domain of F_{μ} is denoted by $\mathcal{D}(\mathsf{F}_{\mu}) := \{ \nu \in \mathcal{P}_{\mu}(\Upsilon) : \mathsf{F}_{\mu}(\nu) < \infty \}.$

The L²-Monge-Kantorovich-Rubinstein-Wasserstein distance. For $\nu, \sigma \in \mathcal{P}(\Upsilon)$, we define $W_{d_{\Upsilon}}$ as

(6.1)
$$\mathsf{W}^{2}_{\mathsf{d}_{\Upsilon}}(\nu,\sigma) := \inf_{\mathsf{c}\in\mathsf{Cpl}(\nu,\sigma)} \int_{\Upsilon^{\times 2}} \mathsf{d}^{2}_{\Upsilon}(\gamma,\eta) \, \mathrm{d}\mathsf{c}(\gamma,\eta) \, ,$$

where $\mathsf{Cpl}(\nu, \sigma)$ is the space of all Borel probability measures on $(\Upsilon^{\times 2}, \tau_v^{\times 2})$ satisfying $\mathsf{c}(\Xi \times \Upsilon) = \nu(\Xi)$ and $\mathsf{c}(\Upsilon \times \Lambda) = \sigma(\Lambda)$ for every $\Xi, \Lambda \in \mathscr{B}(\Upsilon, \tau_v)$.

Benamou-Brenier-like distance. We define a sub-algebra \mathcal{L} in $\mathcal{D}(\mathcal{E}^{\Upsilon,\mu})$:

(6.2)
$$\mathcal{L} = \{ u \in \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}) : u \in L^{\infty}(\mu) , \ \Gamma^{\Upsilon,\mu}(u) \in L^{\infty}(\mu) \} .$$

Definition 6.1 (Continuity equaiton [AES16, (10.6)]). Given a family of probability measures $(\rho_t)_{t\in[0,T]}$, we write $(\rho_t)_{t\in[0,T]} \in \mathsf{CE}^2(\mathcal{E}^{\Upsilon,\mu})$ if there exists $c \in L^2(0,T)$ so that

(6.3)
$$\left| \int_{\mathbf{\Upsilon}} u\rho_t \, \mathrm{d}\mu - \int_{\mathbf{\Upsilon}} u\rho_s \, \mathrm{d}\mu \right| \le \int_s^t c(r) \left(\int_{\mathbf{\Upsilon}} \Gamma^{\mathbf{\Upsilon},\mu}(u)\rho_r \, \mathrm{d}\mu \right)^{1/2} \mathrm{d}r \; ,$$

for every $u \in \mathcal{L}$ and $0 \leq s \leq t \leq T$. The least c in (6.3) is denoted by $\|\rho'_t\|$.

Definition 6.2 (Benamou–Brenier-like extended distance [AES16, Dfn. 10.4]). For $\nu, \sigma \in \mathcal{P}_{\mu}(\Upsilon)$,

(6.4)
$$\mathsf{W}_{\mathcal{E}}(\nu,\sigma)^2 := \inf\left\{\int_0^1 \|\rho_t'\|^2 \,\mathrm{d}t : (\rho_t) \in \mathsf{CE}^2(\mathcal{E}^{\Upsilon,\mu}) , \ \nu = \rho_0 \cdot \mu , \ \sigma = \rho_1 \cdot \mu\right\}.$$

If there is no $(\rho_t) \in \mathsf{CE}^2(\mathcal{E}^{\Upsilon,\mu})$ connecting ν and σ , we define $\mathsf{W}_{\mathcal{E}}(\nu,\sigma) = +\infty$. We will see in Cor. 6.4 that $\mathsf{W}_{\mathcal{E}}(\mathcal{T}_t^{\Upsilon,\mu}\nu,\nu) < \infty$ for every $\nu \in \mathcal{D}(\mathsf{Ent}_{\mu})$ and t > 0, so that $\mathsf{W}_{\mathcal{E}}$ is non-trivial.

Proposition 6.3 (Properties of $W_{\mathcal{E}}$). The following hold:

- (i) $\mathsf{W}_{\mathcal{E}}$ is a complete length extended distance on $\mathcal{P}_{\mu}(\Upsilon)$. Furthermore, $\mathsf{W}_{\mathcal{E}}^2$ is jointly convex in $\mathcal{P}_{\mu}(\Upsilon)^{\times 2}$.
- (ii) The following inequality holds:

$$W_{d_{\Upsilon}} \leq W_{\mathcal{E}}$$
 .

(iii) Let
$$\nu_t = \rho_t \cdot \mu$$
 with $\rho_t = T_t^{\Upsilon,\mu} \rho_0$ and $\rho_0 \in L^2(\mu)$. Then, $(\nu_t)_{t \in [0,1]} \in \mathsf{CE}^2(\mathcal{E}^{\Upsilon,\mu})$ and
(6.5) $\|\rho_t'\|^2 \leq \mathsf{F}(\nu_t)$, $t > 0$.

Proof. (i): The statement follows from [AES16, 5th paragraph on p.113]. We note that the completeness follows from the completeness of $W_{d\gamma}$ and the inequality $W_{d\gamma} \leq W_{\mathcal{E}}$, the latter of which will be proved in (ii).

(ii): Let $\mathsf{d}_{\mathcal{E}}(\gamma, \eta) := \sup\{u(\gamma) - u(\eta) : \Gamma^{\Upsilon}(u) \leq 1, u \in C_b(\Upsilon, \tau_v)\}$ be the intrinsic distance associated with $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$. By the Rademacher-type property in Prop. 4.16, we have $\mathsf{d}_{\Upsilon} \leq \mathsf{d}_{\mathcal{E}}$ (see [DS21a, the first half of the proof of Thm. 5.25]). In particular,

$$\mathsf{W}_{\mathsf{d}_{\Upsilon}} \leq \mathsf{W}_{\mathsf{d}_{\mathcal{E}}}$$

where $W_{d_{\mathcal{E}}}$ denotes the extended distance (6.1) induced by $d_{\mathcal{E}}$ in place of d_{Υ} . By [AES16, (a) Prop. 7.4],

$$W_{d_{\mathcal{E}}} \leq W_{Ch_{d_{\mathcal{E}}}}$$
,

where $\mathsf{Ch}_{\mathsf{d}_{\mathcal{E}}}$ is the Cheeger energy associated with $(\Upsilon, \mathsf{d}_{\mathcal{E}}, \mu)$ (see [AES16, Dfn. 6.1]). Furthermore, by [AES16, Thm. 12.5], we have $\mathsf{W}_{\mathsf{Ch}_{\mathsf{d}_{\mathcal{E}}}} \leq \mathsf{W}_{\mathcal{E}}$, which completes the proof.

(iii): The is a consequence of [AES16, (10.10)].

Evolutional Variation Inequality. Recall that $(T_t^{\Upsilon,\mu})_{t>0}$ is the L^2 -semigroup associated with $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$, see (i) Prop. 4.16. For $\nu = \rho \cdot \mu \in \mathcal{P}_{\mu}(\Upsilon)$, we define the dual flow $(\mathcal{T}_t^{\Upsilon,\mu})_{t>0}$ as

$$\mathcal{T}_t^{\Upsilon,\mu}\nu = (T_t^{\Upsilon,\mu}\rho)\cdot\mu \ , \quad t>0$$

The following is the main result in this section as a corollary of Thm. 4.19.

Corollary 6.4 (EVI). Suppose that μ is the sine_{β} ensemble with $\beta > 0$. For every $\nu, \sigma \in \mathcal{D}(\mathsf{Ent}_{\mu})$ with $\mathsf{W}_{\mathcal{E}}(\nu, \sigma) < \infty$, the curve $t \mapsto \mathcal{T}_{t}^{\Upsilon, \mu} \sigma \in (\mathcal{P}(\Upsilon), \mathsf{W}_{\mathcal{E}})$ is locally absolutely continuous, $\mathsf{Ent}_{\mu}(\mathcal{T}_{t}^{\Upsilon, \mu}\nu) < \infty$, $\mathsf{W}_{\mathcal{E}}(\mathcal{T}_{t}^{\Upsilon, \mu}\nu, \sigma) < \infty$ for every t > 0 and

$$(\mathsf{EVI}(0,\infty)) \qquad \qquad \frac{1}{2} \frac{\mathrm{d}^+}{\mathrm{d}t} \mathsf{W}_{\mathcal{E}} \big(\mathcal{T}_t^{\Upsilon,\mu} \sigma, \nu \big)^2 \leq \mathsf{Ent}_{\mu}(\nu) - \mathsf{Ent}_{\mu}(\mathcal{T}_t^{\Upsilon,\mu} \sigma) \ , \qquad t > 0 \ .$$

Proof. This is a direct consequence of Thm. 4.19 and [AES16, Cor. 11.3].

As a consequence of $\mathsf{EVI}(0,\infty)$, we have the following corollary, see [AES16, Thm. 10.14, Cor.s 11.2, 11.5, Thm. 11.4].

Corollary 6.5 (Geodesical convexity). Suppose that μ is the sine_{β} ensemble with $\beta > 0$. The following hold:

(a) The space $(\mathcal{D}(\mathsf{Ent}_{\mu}), \mathsf{W}_{\mathcal{E}})$ is an extended geodesic metric space: for every pair $\nu, \sigma \in \mathcal{D}(\mathsf{Ent}_{\mu})$ with $\mathsf{W}_{\mathcal{E}}(\nu, \sigma) < \infty$, there exists $\mathsf{W}_{\mathcal{E}}$ -Lipschitz curve ν . : $[0,1] \rightarrow (\mathcal{D}(\mathsf{Ent}_{\mu}), \mathsf{W}_{\mathcal{E}})$ so that

 $\nu_0 = \nu, \quad \nu_1 = \sigma, \quad \mathsf{W}_{\mathcal{E}}(\nu_t, \nu_s) = |t - s| \mathsf{W}_{\mathcal{E}}(\nu, \sigma), \quad s, t \in [0, 1].$

(b) Displacement convexity: The entropy Ent_μ is W_ε-convex along every W_ε-geodesic (ν_t)_{t∈[0,1]}:

$$\mathsf{Ent}_{\mu}(\nu_t) \le (1-t)\mathsf{Ent}_{\mu}(\nu_0) + t\mathsf{Ent}_{\mu}(\nu_1) , \quad t \in [0,1] .$$

(c) Wasserstein contraction:

$$\mathsf{W}_{\mathcal{E}}(\mathcal{T}_{t}^{\Upsilon,\mu}\nu,\mathcal{T}_{t}^{\Upsilon,\mu}\sigma) \leq \mathsf{W}_{\mathcal{E}}(\nu,\sigma) , \quad t > 0 , \quad \nu,\sigma \in \mathcal{P}_{\mu}(\Upsilon) .$$

(d) The descending $W_{\mathcal{E}}$ -slope of Ent_{μ} coincides with the Fisher information:

$$|\mathsf{D}^-_{\mathsf{W}_{\mathcal{E}}}\mathsf{Ent}_{\mu}(\nu)|^2 = \mathsf{F}(
ho) \ , \quad \nu =
ho \cdot \mu \in \mathcal{D}(\mathsf{Ent}_{\mu}) \ .$$

- (e) The set $A_c := \{ \nu = \rho \cdot \mu \in \mathcal{P}_{\mu}(\Upsilon) : \|\rho\|_{L^{\infty}(\mu)} \leq c \}$ is geodesically convex with respect to $W_{\mathcal{E}}$ for every c > 0.
- (f) $L \log L$ -regularisation of $\mathcal{T}_t^{\Upsilon,\mu}$: For every $\nu = \rho \cdot \mu \in \mathcal{P}_{\mu}(\Upsilon)$ (not necessarily in $\mathcal{D}(\mathsf{Ent}_{\mu})$) and $\sigma \in \mathcal{D}(\mathsf{Ent}_{\mu})$,

$$\operatorname{Ent}_{\mu}(\mathcal{T}_{t}^{\Upsilon,\mu}\nu) \leq \operatorname{Ent}_{\mu}(\sigma) + \frac{1}{2t}\operatorname{W}_{\mathcal{E}}(\nu,\sigma)^{2}, \quad t > 0.$$

Finally, we show that the dual flow $(\mathcal{T}_t^{\Upsilon,\mu})$ is the $\mathsf{W}_{\mathcal{E}}$ -gradient flow of Ent_{μ} .

Corollary 6.6 (Gradient flow). Suppose that μ is the sine_{β} ensemble with $\beta > 0$. The dual flow $(\mathcal{T}_t^{\Upsilon,\mu})_{t>0}$ is the unique solution to the $W_{\mathcal{E}}$ -gradient flow of Ent_{μ} . Namely, for any $\nu_0 \in \mathcal{D}(\mathsf{Ent}_{\mu})$, the curve $[0,\infty) \ni t \mapsto \nu_t = \mathcal{T}_t^{\Upsilon,\mu}\nu_0 \in \mathcal{D}(\mathsf{Ent}_{\mu})$ is the unique solution to the energy equality:

(6.6)
$$\frac{\mathrm{d}}{\mathrm{d}t}\mathsf{Ent}_{\mu}(\nu_t) = -|\dot{\nu}_t|^2 = -|\mathsf{D}_{\mathsf{W}_{\mathcal{E}}}^-\mathsf{Ent}_{\mu}|^2(\nu_t) \quad a.e. \ t > 0 \ .$$

Proof. This is a consequence of Cor. 6.4 and [MS20, Thm. 3.15]. The uniqueness follows from [MS20, Thm. 4.2]. Note that although [MS20] works only in metric spaces (as opposed to extended metric spaces), we can apply the result to our setting by the following argument: Noting that $W_{\mathcal{E}}(\nu_t, \nu_s) < \infty$ for every $s, t \in [0, \infty)$ by Cor. 6.4 with $\sigma = \nu_0$, we can think of (ν_t) as a curve in a (non-extended) metric space $\{\sigma \in \mathcal{P}(\Upsilon) : W_{\mathcal{E}}(\nu_0, \sigma) < \infty\}$, to which we can apply the results in [MS20].

7. Generalisation

We have been so far working in the case of $\operatorname{sine}_{\beta}$ ensemble. In this section, we seek to generalise the aforementioned statements to general probability measures on $\Upsilon = \Upsilon(\mathbb{R}^n)$ for $n \in \mathbb{N}$. As an application, we prove $\mathsf{BE}(0,\infty)$ in the case of β -Riesz ensemble.

In this section, we denote by \mathbf{m} and \mathbf{m}_r the Lebesgue measure on \mathbb{R}^n and its restriction on $B_r(0)$ respectively, and we take the Euclidean distance $\mathsf{d}(x,y) := |x-y|$ for $x, y \in \mathbb{R}^n$. Let μ be a Borel probability on Υ . Let $\mathcal{K}(\mu_r^\eta) \subset \mathbb{N}_0$ be defined as

$$\mathcal{K}(\mu^\eta_r) := \{k \in \mathbb{N}_0 : \mu^{k,\eta}_r(\mathbf{\Upsilon}^k(B_r)) > 0\}$$
.

Assumption 7.1. Let $K \in \mathbb{R}$ and μ be a Borel probability. Assume the following conditions:

- (a) the measure μ_r^{η} is absolutely continuous with respect to the Poisson measure π_{m_r} , and $\mu_r^{k,\eta}$ is equivalent to $\pi_{\mathsf{m}_r}|_{\mathbf{\Upsilon}^k(B_r)}$ for any $k \in \mathcal{K}(\mu_r^{\eta})$, μ -a.e. η and any r > 0;
- (b) (conditional geodesical K-convexity) the density

$$\frac{\mathrm{d}\mu_r^{k,\eta}}{\mathrm{d}\pi_{\mathsf{m}_r}|_{\mathbf{\Upsilon}^k(B_r)}}$$

is $\tau_{\rm v}$ -continuous on $\Upsilon^k(B_r)$, and the logarithmic density

$$\Psi_r^{k,\eta} = -\log\left(\frac{\mathrm{d}\mu_r^{k,\eta}}{\mathrm{d}\pi_{\mathsf{m}_r}|_{\mathbf{\Upsilon}^k(B_r)}}\right) \quad \text{is } K\text{-geodesically convex}$$

with respect to d_{Υ} on $\Upsilon^k(B_r)$ for every $k \in \mathcal{K}(\mu_r^{\eta})$, μ -a.e. η and every r > 0.

Under (a) in Assumption, the local Dirichlet form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ is constructed in the same proof as in the case of $\operatorname{sine}_{\beta}$ ensemble as we have not use any particular property of K = 0. We further show the synthetic curvature bound for the form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ and related functional inequalities.

Theorem 7.2. Suppose that μ satisfies Assumption 7.1. Then the local Dirichlet form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ satisfies

(a) (Bakry–Émery inequality $BE(K,\infty)$)

$$\Gamma^{\Upsilon}(T_t^{\Upsilon,\mu}u) \le e^{-2Kt}T_t^{\Upsilon,\mu}\Gamma^{\Upsilon}(u) \quad \forall u \in \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}) ;$$

(b) (Integral Bochner inequality) for every $(u, \varphi) \in \mathcal{D}(\Gamma_2^{\Upsilon, \mu})$

$$\Gamma_2^{\Upsilon,\mu}(u,\varphi) \ge 2K \int_{\Upsilon} \Gamma^{\Upsilon}(u) \varphi \,\mathrm{d}\mu$$

(c) (local Poincaré inequality) for $u \in \mathcal{D}(\mathcal{E}^{\Upsilon,\mu})$ and t > 0,

$$\begin{split} T_t^{\mathbf{\Upsilon},\mu} u^2 &- (T_t^{\mathbf{\Upsilon},\mu} u)^2 \leq \frac{1 - e^{-2Kt}}{K} T_t^{\mathbf{\Upsilon},\mu} \Gamma^{\mathbf{\Upsilon}}(u) \\ T_t^{\mathbf{\Upsilon},\mu} u^2 &- (T_t^{\mathbf{\Upsilon},\mu} u)^2 \geq \frac{e^{2Kt} - 1}{K} \Gamma^{\mathbf{\Upsilon}}(T_t^{\mathbf{\Upsilon},\mu} u) ; \end{split}$$

(d) (Exponential integrability of 1-Lipschitz functions) If u is a \bar{d}_{Υ} -Lipschitz function with $\operatorname{Lip}_{\bar{d}_{\Upsilon}}(u) \leq 1$ and $|u(\gamma)| < \infty \mu$ -a.e. γ , then for every $s < \sqrt{\frac{8K}{1-e^{-2Kt}}}$

$$\int_{\Upsilon} e^{su(\eta)} P_t^{\Upsilon,\mu}(\gamma,\mathrm{d}\eta) < \infty$$

(e) (log Harnack inequality) for every non-negative $u \in L^{\infty}(\Upsilon, \mu)$, $\varepsilon \in (0, 1]$, t > 0, there exists $\Omega \subset \Upsilon$ so that $\mu(\Omega) = 1$ and

$$T_t^{\Upsilon,\mu}\log(u+\varepsilon)(\gamma) \le \log(T_t^{\Upsilon,\mu}u(\eta)+\varepsilon) + \frac{K}{2(1-e^{-2Kt})}\bar{\mathsf{d}}_{\Upsilon}(\gamma,\eta)^2 , \quad \forall \gamma,\eta \in \Omega ;$$

(f) (dimension-free Harnack inequality) for every non-negative $u \in L^{\infty}(\Upsilon, \mu)$, $t > 0 \text{ and } \alpha > 1 \text{ there exists } \Omega \subset \Upsilon \text{ so that } \mu(\Omega) = 1 \text{ and}$

$$(T_t^{\mathbf{\Upsilon},\mu}u)^{\alpha}(\gamma) \leq T_t^{\mathbf{\Upsilon},\mu}u^{\alpha}(\eta) \exp\left\{\frac{\alpha K}{2(\alpha-1)(1-e^{-2Kt})}\bar{\mathsf{d}}_{\mathbf{\Upsilon}}(\gamma,\eta)^2\right\}, \quad \forall \gamma, \eta \in \Omega ;$$

(g) (Lipschitz contraction) For $u \in \operatorname{Lip}(\bar{\mathsf{d}}_{\Upsilon}, \mu)$ and t > 0,

 $T_t^{\mathbf{\Upsilon},\mu}u$ has a $\bar{\mathsf{d}}_{\mathbf{\Upsilon}}$ -Lipschitz μ -modification $\tilde{T}_t^{\mathbf{\Upsilon},\mu}u$

and the following estimate holds:

$$\operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(\tilde{T}_{t}^{\Upsilon,\mu}u) \leq e^{-Kt}\operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(u) ;$$

(h) (L^{∞} -to-Lip regularisation) For $u \in L^{\infty}(\mu)$ and t > 0,

$$T_t^{\Upsilon,\mu}u$$
 has a $\bar{\mathsf{d}}_{\Upsilon}$ -Lipschitz μ -modification $\tilde{T}_t^{\Upsilon,\mu}u$

and the following estimate holds:

$$\operatorname{Lip}_{\bar{\mathsf{d}}_{\Upsilon}}(\tilde{T}_t^{\Upsilon,\mu}u) \leq \frac{1}{\sqrt{2I_{2K}(t)}} \|u\|_{L^{\infty}(\mu)} \quad \forall t > 0 ,$$

where $I_K(t) := \int_0^t e^{Kr} \,\mathrm{d}r;$

(i) (The density of Lipschitz algebra)

 $\operatorname{Lip}_b(\bar{\mathsf{d}}_{\Upsilon},\mu)$ is dense in $\mathcal{D}(\mathcal{E}^{\Upsilon,\mu})$.

(j) (Evolutional Variation Inequality) For every $\nu, \sigma \in \mathcal{D}(\mathsf{Ent}_{\mu})$ with $\mathsf{W}_{\mathcal{E}}(\nu, \sigma) < \infty$, the curve $t \mapsto \mathcal{T}_{t}^{\Upsilon,\mu} \sigma \in (\mathcal{P}(\Upsilon), \mathsf{W}_{\mathcal{E}})$ is locally absolutely continuous and

$$\frac{1}{2}\frac{\mathrm{d}^+}{\mathrm{d}t}\mathsf{W}_{\mathcal{E}}\big(\mathcal{T}_t^{\Upsilon,\mu}\sigma,\nu\big)^2 + \frac{K}{2}\mathsf{W}_{\mathcal{E}}\big(\mathcal{T}_t^{\Upsilon,\mu}\sigma,\nu\big)^2 \leq \mathsf{Ent}_{\mu}(\nu) - \mathsf{Ent}_{\mu}(\mathcal{T}_t^{\Upsilon,\mu}\sigma) \;, \quad t>0 \;.$$

(k) (**Displacement** K-convexity) The entropy Ent_{μ} is $W_{\mathcal{E}}$ -convex along every $W_{\mathcal{E}}$ -geodesic $(\nu_t)_{t\in[0,1]}$:

$$\mathsf{Ent}_{\mu}(\nu_t) \le (1-t)\mathsf{Ent}_{\mu}(\nu_0) + t\mathsf{Ent}_{\mu}(\nu_1) - \frac{K}{2}t(1-t)\mathsf{W}_{\mathcal{E}}(\nu_0,\nu_1)^2 , \quad t \in [0,1]$$

(l) (Wasserstein contraction)

$$\mathsf{W}_{\mathcal{E}}(\mathcal{T}_{t}^{\Upsilon,\mu}\nu,\mathcal{T}_{t}^{\Upsilon,\mu}\sigma) \leq e^{-2Kt}\mathsf{W}_{\mathcal{E}}(\nu,\sigma) , \quad t > 0 , \quad \nu,\sigma \in \mathcal{P}_{\mu}(\Upsilon) .$$

(m) (L log L-regularisation) For every $\nu = \rho \cdot \mu \in \mathcal{P}_{\mu}(\Upsilon)$ (not necessarily in $\mathcal{D}(\mathsf{Ent}_{\mu})$) and $\sigma \in \mathcal{D}(\mathsf{Ent}_{\mu})$,

$$\mathsf{Ent}_{\mu}(\mathcal{T}_{t}^{\Upsilon,\mu}\nu) \leq \mathsf{Ent}_{\mu}(\sigma) + \frac{K}{e^{2Kt}-1}\mathsf{W}_{\mathcal{E}}(\nu,\sigma)^{2} , \quad t > 0 .$$

(n) (**Gradient flow**) The dual flow $(\mathcal{T}_t^{\Upsilon,\mu})_{t>0}$ is the unique solution to the $W_{\mathcal{E}}$ -gradient flow of Ent_{μ} . Namely, for any $\nu_0 \in \mathcal{D}(\mathsf{Ent}_{\mu})$, the curve $[0,\infty) \ni t \mapsto \nu_t = \mathcal{T}_t^{\Upsilon,\mu}\nu_0 \in \mathcal{D}(\mathsf{Ent}_{\mu})$ is the unique solution to the energy equality:

(7.1)
$$\frac{\mathrm{d}}{\mathrm{d}t} \mathsf{Ent}_{\mu}(\nu_t) = -|\dot{\nu}_t|^2 = -|\mathsf{D}_{\mathsf{W}_{\mathcal{E}}}^-\mathsf{Ent}_{\mu}|^2(\nu_t) \quad a.e. \ t > 0 \ .$$

Proof. Thanks to Assumption 7.1, the space $(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon}, \mu_r^{k,\eta})$ satisfies $\mathsf{RCD}(K, \infty)$ for every $k \in \mathcal{K}(\mu_r^\eta)$ as in the same proof of Prop. 3.4. The rest of the proofs in Sections 4, 5, 6 work exactly in the same way up to the multiplicative constants (e.g., e^{-2Kt} instead of 1 for the $\mathsf{BE}(K, \infty)$ inequality).

Remark 7.3 (Non-necessity of the number rigidity (R)). Under the number rigidity (R), we have $\#\mathcal{K}(\mu_r^{\eta}) = 1$. This however has not been essentially used for the proofs in the case of $\operatorname{sine}_{\beta}$. We therefore do not need (R) in Thm. 7.2. See Remark 4.6 for the construction of Dirichlet forms. For the arguments in Section 4 and 5 involving the semigroup $T_t^{\Upsilon(B_r),\mu_r^{\eta}}$, we just need to observe that k-particle space $\Upsilon^k(B_r)$ is an invariant set of the semigroup $T_t^{\Upsilon(B_r),\mu_r^{\eta}}$ for every k, i.e.,

$$T_t^{\boldsymbol{\Upsilon}(B_r),\mu_r^{\eta}} u \mathbf{1}_{\boldsymbol{\Upsilon}^k(B_r)} = \mathbf{1}_{\boldsymbol{\Upsilon}^k(B_r)} T_t^{\boldsymbol{\Upsilon}(B_r),\mu_r^{\eta}} u , \quad u \in L^2(\boldsymbol{\Upsilon}(B_r),\mu_r^{\eta})$$

From the probabilistic viewpoint, this invariance follows from the fact the semigroup $T_t^{\Upsilon(B_r),\mu_r^{\eta}}$ corresponds to finitely many interacting diffusions on B_r with reflecting boundary condition (due to the choice of the domain $\mathcal{D}(\mathcal{E}^{\Upsilon(B_r),\mu_r^{\eta}})$), so that the number of particles in B_r is preserved under evolution. Thus, we may think of $\Upsilon(B_r)$ as the disjoint union $\sqcup_{k \in \mathcal{K}(\mu_r^{\eta})} \Upsilon^k(B_r)$ for μ -a.e. η regarding the semigroup action. Hence by applying the same proofs as in Section 4 and 5 to each $k \in \mathcal{K}(\mu_r^{\eta})$ (instead of using $k = k(\eta)$), Thm. 7.2 can be proved without number rigidity (**R**).

7.1. β -Riesz ensemble. In this section, we apply Thm. 7.2 to prove $\mathsf{BE}(0,\infty)$ in the case of β -Riesz ensemble $\mu = \mu_{\beta}$ for every $\beta > 0$ on $\Upsilon(\mathbb{R})$. We drop the subscript β as it does not play any particular role in the following argument. Let $g(x) = |x|^{-s}$ with $s \in (0,1)$ for $x \in \mathbb{R}$. Define

$$H_r^k(\gamma) := \sum_{i < j}^k g(x_i - x_j) , \quad M_{r,R}^{k,\eta}(\gamma,\eta) := \sum_{i=1}^k \sum_{y \in \eta_{B_r^c}, \ |y| \le R} \left(g(x_i - y) - g(y) \right) ,$$

$$\Psi_{r,R}^{k,\eta}(\gamma) := \beta \Big(H_r^k(\gamma) + M_{r,R}^{k,\eta}(\gamma,\eta) \Big) \quad \text{for } \gamma = \sum_{i=1}^k \delta_{x_i} \in \Upsilon^k(B_r) \text{ and } \eta \in \Upsilon(\mathbb{R})$$

Proposition 7.4. $\Psi_{r,R}^{k,\eta}$ is geodesically convex in $(\Upsilon^k(B_r), \mathsf{d}_{\Upsilon})$ for any $0 < r < R < \infty$, $k \in \mathbb{N}, \eta \in \Upsilon(B_r^c)$ and $\beta > 0$.

Proof. Let H_{ij}, H_i^y be the Hessian matrices of the functions $(x_1, \ldots, x_k) \mapsto g(x_i - x_j)$ and $(x_1, \ldots, x_k) \mapsto g(x_i - y) - g(y)$ respectively. By observing

(7.2)
$$\mathbf{v}H_{ij}\mathbf{v}^{t} = \frac{s(s+1)(v_{i}-v_{j})^{2}}{|x_{i}-x_{j}|^{s+2}}, \quad \mathbf{v}H_{i}^{y}\mathbf{v}^{t} = \frac{s(s+1)v_{i}^{2}}{|y-x_{i}|^{s+2}},$$
$$\mathbf{v} = (v_{1},\dots,v_{k}) \in \mathbb{R}^{k},$$

the same proof works as in Prop. 3.3.

Theorem 7.5 ([DV21, Thm. 1.8]). There exists a Borel probability measure $\mu = \mu_{\beta}$ so that the pointwise limit $\Phi_r^{k,\eta} := \lim_{R\to\infty} \Phi_{r,R}^{k,\eta}$ exists μ -a.e. η and satisfying the following DLR equation:

(7.3)
$$\mathrm{d}\mu_r^{k,\eta} = \frac{e^{-\Psi_r^{k,\eta}}}{Z_r^{\eta}} \,\mathrm{d}\mathbf{m}_r^{\odot k} \;, \quad k \in \mathcal{K}(\mu_r^{\eta})$$

where $\mu_r^{k,\eta}$ was defined after (2.11) and Z_r^{η} is the normalisation constant.

Remark 7.6. The measure $\mu = \mu_{\beta}$ was constructed as a subsequencial limit of certain finitevolume Gibbs measures. The uniqueness of the limit points seems still open, and any limit point is currently called β -circular Riesz gas (or ensemble), see e.g., [DV21, Prop. 1.5] for more details.

Corollary 7.7. Any β -circular Riesz ensemble μ satisfies Assumption 7.1 for $\beta > 0$.

Proof. The condition (a) and the geodesical convexity in (b) of Assumption 7.1 follow by Thm. 7.5 and Prop. 7.4. We only need to verify the continuity of the map

(7.4)
$$\Upsilon^k(B_r) \ni \gamma \mapsto e^{-\Psi_r^{k,\eta}(\gamma)}$$

for every $k \in \mathcal{K}(\mu_r^{\eta})$ and μ -a.e. η . Thanks to [DV21, Lem. 1.7] (note that the roles of γ and η there are opposite to this article), the following pointwise limit exists for μ -a.e. η

$$M_r^{k,\eta}(\gamma,\eta) := \lim_{R \to \infty} M_{r,R}^{k,\eta}(\gamma,\eta) < \infty , \quad k \in \mathcal{K}(\mu_r^\eta) ,$$

and $\Psi_r^{k,\eta}(\gamma,\eta)$ can be written as

$$\Psi_r^{k,\eta}(\gamma,\eta) = \lim_{R \to \infty} \Psi_{r,R}^{k,\eta}(\gamma,\eta) = \beta \Big(H_r^k(\gamma) + M_r^{k,\eta}(\gamma,\eta) \Big) \ .$$

Thus, it suffices for the continuity of (7.4) to show the continuity of the map $\gamma \mapsto M_r^{k,\eta}(\gamma,\eta)$ on $\Upsilon^k(B_r)$ for μ -a.e. η and each $k \in \mathcal{K}(\mu_r^{\eta})$. Let $\gamma_n = \sum_{i=1}^k \delta_{x_i^{(n)}} \in \Upsilon^k(B_r)$ converge to γ vaguely in $\Upsilon^k(B_r)$. Then, the continuity follows by observing

$$|M_r^{k,\eta}(\gamma,\eta) - M_r^{k,\eta}(\gamma_n,\eta)| = \lim_{R \to \infty} \left| \sum_{i=1}^k \sum_{y \in \eta_{B_r^c}, \ |y| \le R} \left(g(x_i - y) - g(x_i^{(n)} - y) \right) \right|$$

$$= \lim_{R \to \infty} \left| \sum_{i=1}^{k} \sum_{y \in \eta_{B_{r}^{c}}, |y| \le R} g'(c_{y}^{(n)})(x_{i} - x_{i}^{(n)}) \right|$$
$$= \left| \sum_{i=1}^{k} (x_{i} - x_{i}^{(n)}) \lim_{R \to \infty} \sum_{y \in \eta_{B_{r}^{c}}, |y| \le R} g'(c_{y}^{(n)}) \right|,$$

for some $c_y^{(n)} \in [x_i - y, x_i^{(n)} - y]$ when $x_i - y < x_i^{(n)} - y$ and $c_y^{(n)} \in [x_i^{(n)} - y, x_i - y]$ when $x_i - y > x_i^{(n)} - y$ by Mean Value Theorem. Note that the case of $x_i - y = x_i^{(n)} - y$ trivialises the argument as $g(x_i - y) - g(x_i^{(n)} - y) = 0$, which can be therefore ignored. Passing to the limit $n \to \infty$, the proof is completed.

Corollary 7.8. The Dirichlet form $(\mathcal{E}^{\Upsilon,\mu}, \mathcal{D}(\mathcal{E}^{\Upsilon,\mu}))$ in (4.15) with the β -Riesz ensemble μ satisfies $\mathsf{BE}(0,\infty)$. Furthermore, all the statements in Thm. 7.2 hold true with K = 0.

APPENDIX A.

Let **m** and \mathbf{m}_r be the Lebesgue measure on \mathbb{R}^n and its restriction on B_r respectively. Set $\Upsilon = \Upsilon(\mathbb{R}^n)$.

Lemma A.1. Let μ be a Borel probability on Υ satisfying that μ_r^{η} is absolutely continuous with respect to the Poisson measure π_{m_r} for any r > 0 and μ -a.e. η . Let $\Sigma \subset B_r$ so that $\mathsf{m}_r(\Sigma^c) = 0$. Let $\Omega(r) := \{\gamma \in \Upsilon : \gamma_{\Sigma} = \gamma_{B_r}\}$. Then,

$$\mu(\Omega(r)) = 1 \qquad \forall r > 0 .$$

Proof. We fix r > 0 and write simply $\Omega = \Omega(r)$. By the disintegration formula (2.16),

$$\mu(\Omega) = \int_{\Upsilon} \mu_r^{\eta}(\Omega_r^{\eta}) \,\mathrm{d}\mu(\eta) \;.$$

Thus, it suffices to show $\mu_r^{\eta}(\Omega_r^{\eta}) = 1$ for μ -a.e. η . This is equivalent to show

(A.1)
$$\mu_r^{\eta}(\Omega_r^{\eta}) = \sum_{k \in \mathbb{N}_0} \mu_r^{k,\eta}(\Omega_r^{\eta}) = 1 .$$

As $\mu_r^{k,\eta}$ is absolutely continuous with respect to $\pi_{\mathbf{m}_r}|_{\mathbf{\Upsilon}^k(B_r)}$, it suffices to prove

$$\pi_{\mathsf{m}_r}|_{\mathbf{\Upsilon}^k(B_r)}((\Omega_r^\eta)^c) = 0$$

for every $k \in \mathbb{N}_0$ and $\eta \in \Upsilon$.

We show that (recall the definition of symmetric product $\Sigma^{\odot k}$ in (2.9))

(A.2)
$$\Sigma^{\odot k} \subset \Omega^{\eta}_r \cap \Upsilon^k(B_r) \qquad \forall \eta \in \Upsilon .$$

Let $\gamma \in \Sigma^{\odot k}$. Then by the definition of Ω , it holds that $\gamma + \eta_{B_r^c} \in \Omega$ for any $\eta \in \Upsilon$. Thus, by recalling the definition (2.15) of Ω_r^{η} , we obtain $\gamma \in \Omega_r^{\eta} \cap \Upsilon^k(B_r)$. Thus, (A.2) holds true.

By using (A.2), $\pi_{\mathsf{m}_r}|_{\mathbf{\Upsilon}^k(B_r)} = e^{-\mathsf{m}_r(B_r)}\mathsf{m}_r^{\odot k}$ by (2.17) and $\mathsf{m}_r^{\odot k}((\Sigma^{\odot k})^c) = 0$ by hypothesis, we conclude that for every $\eta \in \mathbf{\Upsilon}$

$$\pi_{\mathsf{m}_r}|_{\mathbf{\Upsilon}^k(B_r)}((\Omega_r^{\eta})^c) = e^{-\mathsf{m}_r(B_r)}\mathsf{m}_r^{\odot k}\Big(\big(\Omega_r^{\eta} \cap \mathbf{\Upsilon}^k(B_r)\big)^c\Big) \le e^{-\mathsf{m}_r(B_r)}\mathsf{m}_r^{\odot k}\big((\Sigma^{\odot k})^c\big) = 0 \ .$$

The proof is complete.

We recall that for $\eta \in \Upsilon$, we set $\Upsilon_r^{\eta} := \{\gamma \in \Upsilon : \gamma_{B_r^c} = \eta_{B_r^c} \}.$

Lemma A.2 (disintegration lemma). Assume that there exists a measurable set $\Xi \subset \Upsilon$ with $\mu(\Xi) = 1$ so that for every $\eta \in \Xi$, there exists a family of measurable sets $\Omega^{\eta} \subset \Upsilon(B_r)$ so that $\mu_r^{\eta}(\Omega^{\eta}) = 1$ for every $\eta \in \Xi$. Let $\Omega \subset \Upsilon$ be the (not necessarily measurable) subset defined by

$$\Omega := \bigcup_{\eta \in \Xi} \operatorname{pr}_r^{-1}(\Omega^\eta) \cap \Upsilon_r^\eta \ .$$

Assume further that there exists a measurable set $\Theta \subset \Upsilon$ so that $\Omega \subset \Theta$. Then, $\mu(\Theta) = 1$.

Caveat. As the set Ω is defined as uncountable union of measurable sets, the measurability of Ω is not necessarily true in general. The disintegration formula (2.16) is, therefore, not necessarily applicable directly to Ω , which motivates the aforementioned lemma.

Proof of Lem. A.2. Let $\Theta_r^{\eta} = \{\gamma \in \Upsilon(B_r) : \gamma + \eta_{B_r^c} \in \Theta\}$ be a section of Θ at $\eta_{B_r^c}$ as in (2.15). Then, $\Omega^{\eta} \subset \Theta_r^{\eta}$ by assumption. Thus, $\mu_r^{\eta}(\Theta_r^{\eta}) \ge \mu_r^{\eta}(\Omega^{\eta}) \ge 1$. By the disintegration formula in (2.16), we have that

$$\mu(\Theta) = \int_{\Upsilon} \mu_r^{\eta}(\Theta_r^{\eta}) \,\mathrm{d}\mu(\eta) \ge 1 \;.$$

The proof is completed.

Lemma A.3. Let μ be a Borel probability on Υ . Let $\Omega \subset \Upsilon$ satisfy $\mu(\Omega) = 1$. Then, there exists $\Omega' \subset \Omega$ with $\mu(\Omega') = 1$ and

(A.3)
$$\mu_r^\eta(\Omega_r^\eta) = 1$$
, $\forall \eta \in \Omega'$.

Proof. By the disintegration formula (2.16),

$$1 = \mu(\Omega) = \int_{\mathbf{\Upsilon}} \mu_r^{\eta}(\Omega_r^{\eta}) \, \mathrm{d}\mu(\eta) = \int_{\Omega} \mu_r^{\eta}(\Omega_r^{\eta}) \, \mathrm{d}\mu(\eta)$$

by which the statement is readily concluded.

Lemma A.4. Let $(Q, \mathcal{D}(Q))$ be a symmetric closed form on a separable Hilbert space H. Let $\{T_t\}$ and $(A, \mathcal{D}(A))$ be the corresponding semigroup and infinitesimal generator respectively. Suppose that there exists an algebra $\mathcal{C} \subset \mathcal{D}(Q)$ so that $\mathcal{C} \subset H$ is dense and $T_t\mathcal{C} \subset \mathcal{C}$ for any t > 0. Then \mathcal{C} is dense in $\mathcal{D}(Q)$.

Proof. It holds that $T_t \mathcal{D}(A) \subset \mathcal{D}(A)$ by the general property of semigroups associated with symmetric closed forms. Thus, combining it with the hypothesis $T_t \mathcal{C} \subset \mathcal{C}$,

$$T_t(\mathcal{C} \cap \mathcal{D}(A)) \subset \mathcal{C} \cap \mathcal{D}(A)$$
.

Thus, by [RS75, Thm. X.49], $\mathcal{C} \cap \mathcal{D}(A)$ is dense in the graph norm in the space $(A, \mathcal{D}(A))$. Namely, we obtained

 $(A, \mathcal{C} \cap \mathcal{D}(A))$ is essentially self-adjoint.

The density $\mathcal{C} \subset \mathcal{D}(Q)$ now follows by the density of $\mathcal{C} \cap \mathcal{D}(A)$ in the graph norm, by the density of $\mathcal{D}(A) \subset \mathcal{D}(Q)$ due to the general property of symmetric closed forms, by the density of $\mathcal{C} \subset H$ and by a simple integration-by-parts argument

$$Q(u, u) = (-Au, u)_H \le ||Au||_H ||u||_H$$
.

The proof is complete.

References

- [AES16] Ambrosio, L., Erbar, M., and Savaré, G. Optimal transport, Cheeger energies and contractivity of dynamic transport distances in extended spaces. *Nonlinear Anal.*, 137:77–134, 2016.
- [AGS08] Ambrosio, L., Gigli, N., and Savaré, G. Gradient Flows in Metric Spaces and in the Space of Probability Measures. Lectures in Mathematics - ETH Zürich. Birkhäuser, 2nd edition, 2008.
- [AGS14a] Ambrosio, L., Gigli, N., and Savaré, G. Calculus and heat flow in metric measure spaces and applications to spaces with Ricci bounds from below. *Invent. Math.*, 395:289–391, 2014.
- [AGS14b] Ambrosio, L., Gigli, N., and Savaré, G. Metric measure spaces with Riemannian Ricci curvature bounded from below. *Duke Math. J.*, 163(7):1405–1490, 2014.
- [AGS15] Ambrosio, L., Gigli, N., and Savaré, G. Bakry-Émery Curvature-Dimension Condition and Riemannian Ricci Curvature Bounds. Ann. Probab., 43(1):339–404, 2015.
- [BGL14] Bakry, D., Gentil, I., and Ledoux, M. Analysis and Geometry of Markov Diffusion Operators, volume 348 of Grundlehren der mathematischen Wissenschaften. Springer, 2014.
- [BH91] Bouleau, N. and Hirsch, F. Dirichlet forms and analysis on Wiener space. De Gruyter, 1991.
- [BÉ84] Bakry, D. and Émery, M. Hypercontractivité de semi-groupes de diffusion. C. R. Acad. Sci. ParisSér. I Math., 299:775–778, 1984.
- [Del21] Dello Schiavo, L. Ergodic Decomposition of Dirichlet Forms via Direct Integrals and Applications. Potential Anal., 2021.
- [DHLM20] Dereudre, D., Hardy, A., Leblé, T., and Maïda, M. DLR Equations and Rigidity for the Sine-Beta Process. Commun. Pure Appl. Math., pages 172–222, 2020.
- [DS21a] Dello Schiavo, L. and Suzuki, K. Configuration spaces over singular spaces –I. Dirichlet-Form and Metric Measure Geometry –. arXiv:2109.03192v2 (version 2), 2021.
- [DS21b] Dello Schiavo, L. and Suzuki, K. On the Rademacher and Sobolev-to-Lipschitz Properties for Strongly Local Dirichlet Spaces. J. Func. Anal., 281(11):Online first, 2021.
- [DS22] Dello Schiavo, L. and Suzuki, K. Configuration Spaces over Singular Spaces II Curvature. arXiv:2205.01379, 2022.
- [DV21] Dereudre, D and Vasseur, T. Number-rigidity and β-circular riesz gas. Arxiv:2104.09408, 2021.
- [Dys62] F. J. Dyson. A brownian-motion model for the eigenvalues of a random matrix. J. Math. Phys, 3:1191–1198, 1962.
- [EH15] Erbar, M. and Huesmann, M. Curvature bounds for configuration spaces. Calc. Var., 54:307–430, 2015.
- [EHJM23] Erbar, M, Huesmann, M., Jalowy, J., and Müller, B. Optimal transport of stationary point processes: Metric structure, gradient flow and convexity of the specificentropy. arXiv: 2304.11145, 2023.
- [EK86] Ethier, S.N. and Kurtz, T.G. Markov Processes Characterization and Convergence—. Wiley Inter-science, A JOHN WILEY & SONS, INC., PUBLICATION, 1986.
- [FOT11] Fukushima, M., Oshima, Y., and Takeda, M. Dirichlet forms and symmetric Markov processes, volume 19 of De Gruyter Studies in Mathematics. de Gruyter, extended edition, 2011.
- [Fuk97] Fukushima, M. Distorted Brownian motions and BV functions. Trends in Probability and Analysis, N. Kono, N-R. Shieh, eds, pages 143–150, 1997.
- [Gho15] Ghosh, S. Determinantal processes and completeness of random exponentials: the critical case. Probab. Theory Relat. Fields, 163(3):643–665, 2015.
- [GKMS18] Galaz-García, F., Kell, M., Mondino, A., and Sosa, G. On quotients of spaces with Ricci curvature bounded below. J. Funct. Anal., 275:1368–1446, 2018.
- [GV20] Gyünesu, B. and Von Renesse, M. Molecules as metric measure spaces with kato-bounded ricci curvature. Comptes Rendus. Mathématique, 358:595–602, 2020.
- [HR03] Hino, M. and Ramírez, J. A. Small-Time Gaussian Behavior of Symmetric Diffusion Semigroups. Ann. Probab., 31(3):1254–1295, 2003.
- [JKO98] Jordan, R, Kinderlehrer, D., and Otto F. The variational formulation of the fokker-planck equation. SIAM Journal on Mathematical Analysis, 29:1–17, 1998.

- [KOT21] Kawamoto Y., Osada H., and Tanemura H. Uniqueness of dirichlet forms related to infinite systems of interacting brownian motions. *Potential Anal.*, 55:639–676, 2021.
- [KS09] Killip, R. and Stoiciu, M. Eigenvalue statistics for cmv matrices: from poisson to clock via random matrix ensembles. *Duke Math. J.*, 146 (3)::361–399, 2009.
- [KS21] Kopfer, E. and Sturm, K-Th. Functional inequalities for the heat flow on time-dependent metric measure spaces. J. London Math. Soc., 104-2:926–955, 2021.
- [KT10] Katori, M. and Tanemura, H. Non-equilibrium dynamics of Dyson's model with an infinite number of particles. Comm. Math. Phys., 293(2):469–497, 2010.
- [Li15] H. Li. Dimension-Free Harnack Inequalities on $\mathsf{RCD}(K, \infty)$ Spaces. J. Theoret. Probab., 29:1280–1297, 2015.
- [MR85] Ma, Z.-M. and Röckner, M. Dirichlet forms-closability and change of speed measure, Infinite dimensional analysis and stochastic processes. *Research Notes in Math. S. Albeverio, ed., Pitman*, 124:119–144, 1985.
- [MR90] Ma, Z.-M. and Röckner, M. Introduction to the Theory of (Non-Symmetric) Dirichlet Forms. Springer, 1990.
- [MR00] Ma, Z.-M. and Röckner, M. Construction of Diffusions on Configuration Spaces. Osaka J. Math., 37:273–314, 2000.
- [MS20] Muratori, M. and Savaré, G. Uniqueness of dirichlet forms related to infinite systems of interacting brownian motions. J. Funct. Anal., 278, 2020.
- [Nak14] Nakano, F. Level statistics for one-dimensional schrödinger operators and gaussian beta ensemble. J. Stat. Phys., 156(1):66–93, 2014.
- [NF98] Nagao, T and Forrester, P. J. Multilevel dynamical correlation functions for dyson's brownian motion model of random matrices. *Physics Letters A*, 247:801–850, 1998.
- [NR18] Najnundel, J. and Reda, C. Rigidity of the $Sine_{\beta}$ process. *Electron. Commun. Probab.*, 23:1–8, 2018.
- [Osa96] Osada, H. Dirichlet Form Approach to Infinite-Dimensional Wiener Processes with Singular Interactions. Comm. Math. Phys., 176:117–131, 1996.
- [Osa12] Osada, H. Infinite-dimensional stochastic differential equations related to random matrices. Prob. Theory Relat. Fields, 153(1):471–509, 2012.
- [Osa13] Osada, H. Interacting Brownian Motions in Infinite Dimensions with Logarithmic Interaction Potentials. Ann. Probab., 41(1):1–49, 2013.
- [RS75] Reed, M. and Simon, B. Methods of Modern Mathematical Physics II Fourier Analysis, Self-Adjointness. Academic Press, New York, London, 1975.
- [RS80] Reed, M. and Simon, B. Methods of Modern Mathematical Physics I Functional Analysis. Academic Press, New York, London, 1980.
- [RS99] Röckner, M. and Schied, A. Rademacher's Theorem on Configuration Spaces and Applications. J. Funct. Anal., 169(2):325–356, 1999.
- [Spo87] Spohn, H. Interacting Brownian Particles: A Study of Dyson's Model. Hydrodynamic Behavior and Interacting Particle Systems, pages 151–179, 1987.
- [Stu06] Sturm, K.-T. On the geometry of metric measure spaces. I. Acta Math., 196:65–131, 2006.
- [Tsa16] Tsai, L.-C. Infinite dimensional stochastic differential equations for dyson's model. Probability Theory and Related Fields, 166:801–850, 2016.
- [Vil09] Villani, C. Optimal transport, old and new, volume 338 of Grundlehren der mathematischen Wissenschaften. Springer-Verlag, 2009.
- [VV09] Valkó, B. and Virág, B. Continuum limits of random matrices and the brownian carousel. Invent. math., 177(3):463–508, 2009.
- [Wan14] Wang. F-Y. Analysis for diffusion processes on Riemannian manifolds, volume 18. World Scientific,, 2014.

DEPARTMENT OF MATHEMATICAL SCIENCE, DURHAM UNIVERSITY, SCIENCE LABORATORIES, SOUTH ROAD, DH1 3LE, UNITED KINGDOM