

Hybrid Inflation, Reheating and Dark Radiation in a IIB perturbative moduli stabilization scenario

Waqas Ahmed^{a 1}, Athanasios Karozas^{b 2}, George K. Leontaris^{b 3}, Ilias Tavellaris^{b 4}

^a *Center for Fundamental Physics and School of Mathematics and Physics, Hubei Polytechnic University
Huangshi 435003, China*

^b *Physics Department, University of Ioannina
45110, Ioannina, Greece*

Abstract

We study the cosmological implications of an effective field theory model derived within a configuration of D7 brane stacks in the framework of type-IIB string theory. We consider a suitable geometric setup where the Kähler moduli fields are stabilized and the parametric space is constrained so that a de Sitter vacuum is ensured. In addition to the moduli fields we also take into account the usual Higgs and matter fields included in the effective field theory. In this background, we implement the standard hybrid inflation scenario with a singlet scalar field acting as the inflaton and the Higgs states serving as waterfall fields. Radiative corrections and soft supersymmetry breaking terms play an essential role in the realization of a successful inflationary scenario consistent with the present cosmological data. Small tensor-to-scalar ratio values are predicted, which can be probed in future planned experiments. Further constraints on the model's parameters are derived from bounds on dark radiation which is measured as a contribution to the effective number of neutrino species N_{eff} . In particular, we find an excess of $\Delta N_{\text{eff}} \leq 0.95$ at 2σ confidence level with natural values of the involved couplings.

¹E-mail: waqasmit@hbpu.edu.cn

²E-mail: athanasioskarozas@gmail.com

³E-mail: leonta@uoi.gr

⁴E-mail: i.tavellaris@uoi.gr

Contents

1	Introduction	1
2	Description of the model and its constituents	3
3	The effective potential	6
3.1	Inflationary phase	10
3.2	Numerical results	12
4	Reheating and dark radiation	13
5	Conclusions	16
A	Coleman-Weinberg Corrections	18

1 Introduction

Cosmological inflation is one of the most successful candidate scenarios in explaining the evolution of the Universe and its large-scale structure observed today. Meanwhile, numerous effective quantum field theory models have been built with the aim of conciliating cosmic inflation with particle physics models describing the low energy observables. A key criterion on the quest for appropriate models would be to have an ultra-violet (UV) completion in a quantum theory of gravity, valid up to the Planck scale M_P . In this context, at present, string theory appears to be the only promising candidate for a consistent quantum theory at such a high scale which incorporates the Standard Model (SM) and its minimal supersymmetric extension (MSSM). String theory, however, is formulated in a ten-dimensional spacetime framework and, therefore, compactification of the six extra dimensions is required to achieve a four-dimensional effective field theory compatible with the observed world. The reduction of the corresponding higher dimensional string action to four spacetime dimensions, however, entails an immense set of string vacua which is commonly known as the string landscape. Yet, starting from a successful effective field theory that adequately describes the known physics phenomena, we cannot always embed it in a string theory framework ⁵.

On the other hand, consistent effective field theory models emerging after compactification should possess a number of important features. Among them, they should predict a positive tiny cosmological constant Λ of the order $\Lambda \approx 10^{-122} M_P^4$ which could account for the dark energy, as suggested by cosmological observations. A simple way to realize such a scenario is within an effective model involving a scalar field ϕ with a potential $V(\phi)$ which displays a (possibly metastable) positive minimum equal to the cosmological constant Λ . In fact, effective field theory models from strings compactified on Calabi-

⁵See reviews [1, 2, 3] and references therein.

Yau (CY) manifolds contain a vast number of moduli fields and some of them could play the role of the inflaton ϕ . In this context, it is inferred that the cosmological issues are intertwined with the well-known problem of moduli stabilization. In fact, moduli stabilization and (metastable) de Sitter vacua, play a key role in the successful implementation of the cosmological inflationary scenario in effective field theory (EFT) models of string origin. Therefore, reconciling these two issues is essential in the quest for a suitable non-vanishing effective potential of some scalar field enacting as the inflaton ϕ which rolls down to the minimum of its (relatively shallow) potential. This enables the required exponential growth of the Universe, provided that the trajectory length $\sim \Delta\phi$ of the field ϕ to reach the minimum is sufficiently long to trigger inflation.

A particular class of models involves constructions with large volume compactification scenarios [4, 5, 6, 7, 8, 9] and inflatons associated with the Kähler moduli fields $T_k = \tau_k + ia_k$. In previous scenarios [10, 11, 12, 13] considered in the context of type-IIB theory, it was shown that the internal volume modulus \mathcal{V} expressed in terms of the real components of Kähler moduli, $\text{Re}T_k = \tau_k$, is a suitable candidate for this role, ($\phi \propto \log \mathcal{V}$). Radiative corrections associated with intersecting space-filling D7 branes on the other hand, provide a stabilization mechanism for Kähler moduli, and an uplift to their scalar potential through their universal abelian factors, this way leading to a positive cosmological constant ⁶. In particular, Kähler moduli stabilization is achieved through a non-zero potential generated by α' and radiative (logarithmic) corrections induced when closed string loops traverse their codimension-two bulk towards localized gravity sources. Furthermore, the dS vacuum is obtained due to the positive D-term contributions, (originally proposed in [16]) coming from the intersecting D7-branes.

In the large volume limit, the induced effective potential for the Kähler moduli acquires a simple structure possessing two local extrema (a minimum and a maximum) and approaches zero for $\phi \rightarrow \infty$. The separation distance of its two local extrema is of the order, $\Delta\phi = \phi_{\max} - \phi_{\min} \propto \log(\mathcal{V}_{\max}/\mathcal{V}_{\min})$. In the minimal effective model involving only moduli fields, it can be parametrized in terms of a single non-negative parameter while the largest possible separation $\Delta\phi$ occurs at a critical value of this parameter where beyond this point only AdS solutions appear. A non-zero value of the aforementioned parameter exists at which a new inflationary small-field scenario is successfully implemented. In this novel scenario, most of the required number N_0 of efolds ($N_0 \sim 60$) are collected in the vicinity of the minimum of the potential and the prediction for the tensor-to-scalar ratio density fluctuations in the early universe is $r \approx 4 \times 10^{-4}$. Despite these successes of the model, the picture is not complete and a waterfall mechanism has to be realized in order to end inflation and bring the dS vacuum down to a lower value compatible with the cosmological constant. It has been shown [17] that when open string states appearing in the D7 brane intersections are included in the massless spectrum, appropriate magnetic fluxes and specific brane separations can be chosen in a way that, for \mathcal{V} less than some critical value, a charged open string scalar becomes tachyonic. In [17, 18] it was shown that such a state can represent a waterfall field. In possible generalizations of this scenario, one may include several of such fields, which end inflation and provide a deeper vacuum in accordance with the present value of Λ .

In the present work, an alternative scenario is proposed where it is taken into account that, in ad-

⁶There are many other approaches on these issues and a vast literature exists. For an updated list, see e.g. the recent review articles [14, 15].

dition to the moduli fields, the EFT model accommodates the ordinary fermion matter and the Higgs fields in appropriate representations on the gauge group stemming from the specific details of the compactification procedure. Then, a possible and interesting variation of the scenario described previously would be that the Higgs rolls down a potential hill towards a new lower minimum, where its initial condition is defined around the metastable vacuum of the moduli potential. In the present construction, the metastable vacuum is determined by the Kähler moduli and the associated compactification volume \mathcal{V} . More precisely, we implement the standard hybrid inflation [19] with a singlet scalar field acting as inflaton and the Higgses operating as waterfall fields. In this scenario, the vacuum energy is determined by the scalar field and waterfall fields. We consider radiative corrections which are essential in shaping the slope along the inflationary track. Since supersymmetry (SUSY) is broken during inflation, we also include SUSY soft terms as well. The soft terms play an important role in order to achieve spectral index (n_s) values consistent with the current experimental bounds. We find small tensor-to-scalar values that can be probed in future designed experiments. We also discuss dark radiation and show that change in the effective degree of neutrinos satisfies the bound 0.95% confidence level with natural values of the involved couplings.

The paper is organized as follows. In Section 2 we present the basic features of the model including the Kähler potential and the superpotential. An analysis of the effective potential is given in Section 3 followed by the presentation of the inflation setup along with our inflationary numerical predictions. In Section 4 we discuss reheating and dark radiation predictions. Section 5 concludes the paper.

2 Description of the model and its constituents

In this work, we consider a type-IIB string framework in ten dimensions where six of them are compactified on a Calabi-Yau threefold \mathcal{X} . We restrict our attention mainly to the moduli spectrum and use the following notation: ϕ represents the dilaton field, while T_i , and z_a denote the Kähler and the complex structure (CS) moduli respectively. Furthermore, we introduce the usual axion-dilaton combination

$$\tau = C_0 + i e^{-\phi} \equiv C_0 + \frac{i}{g_s}, \quad (2.1)$$

where g_s is the string coupling and C_0 a 0-form potential (RR-scalar). We assume a perturbative -flux induced- superpotential W_0 of the form proposed in [20]. At the classical level, W_0 is a holomorphic function which depends on the axion-dilaton modulus τ , and the CS moduli z_a ⁷. τ and z_a are stabilized in the standard supersymmetric way, by solving $D_\tau W_0 = 0$, $D_{z_a} W_0 = 0$, where $D_I = \partial_I W + W \partial_I K$ are the covariant derivatives.

We consider a geometric configuration of three intersecting D7-brane stacks equipped with magnetic fluxes. Regarding the Kähler potential, we will take into account α' corrections as well as the effects of a novel four-dimensional Einstein-Hilbert (EH) term (localized in the internal space) which is generated from higher derivative terms in the ten-dimensional string effective action [11]. This setup induces

⁷We dispense with the use of non-perturbative corrections which would also introduce the Kähler moduli through terms of the form $\mathcal{W}_{NP} \propto e^{-aT_k}$. As explained in the subsequent analysis, T_k can be perturbatively stabilized through one-loop corrected Kähler potential.

logarithmic corrections to the scalar potential via loop effects. Taking into account these corrections the relevant part of the Kähler potential receives the form [11]

$$K = -2M_P^2 \log(\mathcal{V} + \xi_0 + \eta_0 \log \mathcal{V}) + \dots, \quad (2.2)$$

where the dots stand for terms depending on the complex structure, z_a and Kähler moduli $T_k = \tau_k + ia_k$. The general form of the volume is given by $\mathcal{V} = \frac{1}{6} \kappa_{ijk} t^i t^j t^k$ where t^i are the two-cycle Kähler moduli fields and κ_{ijk} are triple intersection numbers on \mathcal{X} . In the subsequent, we work within a concrete scenario where the three Kähler moduli appear in the volume on equal footing and stabilize them through perturbative logarithmic loop corrections [21]. Following [22], in particular, we consider a framework that is based on a CY threefold corresponding to the polytope Id: 249 of the Kreuzer-Skarke [23] CY database [24]. In this particular CY manifold there are three Kähler moduli fields satisfying the simple relation $\tau_i = \alpha t^j t^k$ between the two- and four-cycle moduli, and α is a positive constant associated with the intersection number. Then, the volume is simply given by [22]

$$\mathcal{V} = \alpha t^1 t^2 t^3 = \tau_1 t^1 = \tau_2 t^2 = \tau_3 t^3 = \frac{1}{\sqrt{\alpha}} \sqrt{\tau_1 \tau_2 \tau_3}. \quad (2.3)$$

After the dimensional reduction, the effective field theory model could be either some Grand Unified Theory (GUT) or directly the MSSM model where the ordinary low energy (super)-fields appear in appropriate representations of the EFT gauge group. Thus, in addition to the quantum corrections considered above, we also include matter fields in the Kähler potential. These contributions are essential in studying soft supersymmetry breaking effects and cosmological inflation. Within the present context, in particular, we focus on the Higgs sector which plays a vital role in implementing the scenario of hybrid inflation and investigating the possible production of dark radiation. Thus, we consider a generic set of Higgs pairs Φ_i, Φ_j which are assumed to break the gauge group at some GUT scale much lower than the Planck scale M_P . We have also introduced a field S , representing a gauge singlet superfield which realizes the trilinear superpotential couplings of the form $S\Phi_1\Phi_2$. Such singlet fields are ubiquitous in effective string theory models. In the setup described so far, the relevant terms of the superpotential have the following generic form

$$W = W_0 + \kappa S(\Phi_1\Phi_2 - M^2) + \dots, \quad (2.4)$$

where κ is a coupling constant coefficient, M is a high scale mass parameter whose values are below the string scale depending on which scale the Higgs field obtains VEV, W_0 is the flux induced part introduced previously, and dots stand for possible terms irrelevant to our discussion.

In our study, the procedure for working with the Kähler metric and incorporating matter fields can be compared to the study by [25]. This comparison extends to the examination of chiral matter, which is localized on magnetized D7-branes, and the more comprehensively understood fractional D3-branes found at singularities. When considering the second term within our Kähler metric, we explore soft terms linked to massless open strings concentrated at the intersections between D3 and D7 branes. However, it's important to note that the presence of such massless string states cannot be derived from the Dirac-Born-Infeld (DBI) or Chern-Simons (CS) actions alone.

For the case of hybrid inflation within the framework proposed in our work, it becomes imperative to account for the contributions of scalar fields along with their fermionic superpartners in the Kähler poten-

tial. These contributions typically exhibit a generic dependence, expected to be in the form of $\tilde{K}_{ij}\Phi_i\bar{\Phi}_j$. Here, \tilde{K}_{ij} represents a function of the moduli $\tau_k = (T_k + \bar{T}_k)/2$ and S . The precise characterization of \tilde{K}_{ij} becomes clear upon identifying the origin of zero modes and in the context of type IIB theory, various possibilities arise (see for example [26]).

In the case of realistic constructions, chiral matter emerges from either the world volume of D7 brane stacks or at the intersections with other D7 branes. This is where gauge and scalar fields, such as Φ and $\bar{\Phi}$, manifest on the world volume by configuring D7 branes to wrap suitable divisors. The supermultiplets involving these scalar fields, Φ and $\bar{\Phi}$, exhibit a scaling dependence, with the leading contribution taking the form [27]:

$$\frac{\Phi_i\bar{\Phi}_i}{T_k + \bar{T}_k} \rightarrow \frac{\Phi_i\bar{\Phi}_i}{\gamma^{2/3}}.$$

Furthermore, it's worth mentioning that there are next-to-leading order α' contributions [25] of the form $\text{Re}(S)/(T_k + \bar{T}_k)$, however, for our specific purposes, these contributions remain insignificant and, hence, are disregarded.

Including contributions of the above matter fields in the Kähler potential (2.2), we obtain the following generic form

$$\mathcal{K} = -2M_P^2 \log \left[\left(\prod_{k=1}^3 (T_k + \bar{T}_k) \right)^{\frac{1}{2}} + \mathcal{C} \right] + \sum_{k=1}^3 \frac{a_k}{T_k + \bar{T}_k} f_k(\Phi_i, \Phi_2, \dots, S, \bar{S}) \quad (2.5)$$

where \mathcal{C} is a function with a logarithmic dependence on the three moduli fields $T_{1,2,3}$ defined as

$$\mathcal{C} = \xi_0 + \eta_0 \log \left(\prod_{k=1}^3 (T_k + \bar{T}_k) \right). \quad (2.6)$$

In the above equation the parameter ξ_0 stands for α'^3 corrections [28] and is proportional to the Euler characteristic χ_{CY} of the Calabi-Yau manifold

$$\xi_0 = -\frac{\zeta(3)}{4} \chi_{CY}, \quad (2.7)$$

and η_0 is an order one coefficient [11]. The functions f_k in Eq. (2.5) describe the visible Higgs sector. For simplicity, we will assume that all f_k are the same and have the following form

$$f(\Phi, \bar{\Phi}, S) = \alpha \Phi_1 \Phi_1^\dagger + \beta \Phi_2 \Phi_2^\dagger + \gamma S S^\dagger + \lambda (\Phi_1 \Phi_2 + h.c.), \quad (2.8)$$

where α, β, γ and λ are dimensionless couplings. Furthermore, we will adopt the approach used in [25] and we will express the matter contribution term in (2.5) in terms of the compactification volume (note that $T \propto \gamma^{2/3}$). Regarding the origin of these fields, in the present work, we focus on matter fields residing on magnetized $D7$ -branes, as well as chiral fields generated at $D7$ -brane intersections. Notice, however, that the general analysis also applies to states associated with the excitations of open strings stretching between $D7$ - $D3$ branes or having both ends on the same $D3$ brane. The choice of the configuration dictates the modular weights in the Kähler potential. For our purposes, we consider the modular

weights to be 1 [26]. Hence, in large volume compactifications, the Kähler potential in Eq. (2.5) takes the form [26, 29, 30]⁸

$$\mathcal{K} = -2\log[\mathcal{V} + \xi_0 + \eta_0\log(\mathcal{V})] + \frac{3a}{\mathcal{V}^{2/3}} \left[\alpha\Phi_1\Phi_1^\dagger + \beta\Phi_2\Phi_2^\dagger + \gamma SS^\dagger + \lambda(\Phi_1\Phi_2 + h.c.) \right] \quad (2.9)$$

where a is a dimensionless constant and \mathcal{V} is defined in (2.3). Note that in Eq. (2.9) and from now on we adopt $M_P = 1$ units.

Up to this point, we have presented the minimum number of moduli and matter fields that are necessary for our subsequent analysis. Next, we proceed with the computation of the scalar potential which is essential to investigate the properties of the model and compute the various cosmological and other phenomenological observables.

3 The effective potential

The scalar potential of the effective field theory model receives various contributions. As we will see shortly, in the present construction there are F- and D-terms associated with the moduli sector, and contributions from the EFT matter fields, as well as supersymmetry breaking terms. We start with the F-term potential which is given by the generic formula

$$V_F = e^G \left(G_i G_{ij}^{-1} G_{j^*} - 3 \right), \quad (3.1)$$

where

$$G = \mathcal{K} + \log|W|^2 \equiv \mathcal{K} + \log W + \log W^*,$$

and the indices i, j in (3.1) denote the derivatives with respect to the various moduli and other fields.

Computing the derivatives, and substituting in Eq. (3.1) while keeping only the leading order terms, the F-term potential receives the following simplified form⁹

$$V_F \simeq \frac{\kappa^2 \alpha \beta (M^2 - \varphi_1 \varphi_2)^2 + \gamma \kappa^2 S^2 (\alpha \varphi_1^2 + \beta \varphi_2^2)}{3a\alpha\beta\gamma\mathcal{V}^{4/3}} + \frac{3W_0^2(2\eta_0\log\mathcal{V} - 8\eta_0 + \xi_0)}{2\mathcal{V}^3}, \quad (3.2)$$

where φ_1 and φ_2 are the bosonic components of the superfields Φ_1 and Φ_2 .

At the extrema of the F-term potential, the fields take the following values¹⁰

$$S_o = 0, \quad \varphi_{1,o}\varphi_{2,o} = M^2, \quad \mathcal{V}_o = e^{\frac{13}{3} - \frac{\xi_0}{2\eta_0}}. \quad (3.3)$$

⁸Notice that in our case this formula is identified with the expansion of the warped form of the Kähler potential

$$K = -2\log\left((T + \bar{T})^{3/2} + \xi + \eta\log\mathcal{V} - \frac{a}{2}(T + \bar{T})^{1/2}\varphi_i\bar{\varphi}_i\right) = -2\log\left(\mathcal{V} + \xi + \eta\log\mathcal{V} - \frac{a}{2}\mathcal{V}^{1/3}\varphi_i\bar{\varphi}_i\right)$$

⁹The complex field S has the general form $S = |S| e^{i\theta}$ and we choose $\theta = 0$ to align it with the real axis. So in Eq. (3.2) and the subsequent analysis, S denotes the real part of the field.

¹⁰In more general EFT backgrounds it is possible that minimization with respect to the fields S, Φ_i leads to a potential of the form $V \sim \frac{a}{5\mathcal{V}^{4/3}} + \frac{1}{3}\frac{b + \eta\log\mathcal{V}}{\mathcal{V}^3}$. In this case it is possible to have a dS minimum with the volume acquiring a value $\mathcal{V}_o^{5/3} = \left(\frac{9n}{4a}\right) \mathcal{W}\left(\frac{4a}{9n}\left(e^{\frac{1}{3} - \frac{b}{n}}\right)^{5/3}\right)$, where \mathcal{W} is the product-log (Lambert) function.




Figure 1: Plots of the potential along the volume direction. The left panel shows the F-term potential, while in the right panel, the D-term potential has also been included. We choose $\xi_0 = 10$, $\eta_0 = -0.92$, $S = 0$, $\varphi_{1,o} = \varphi_{2,o} = M$, $\kappa = 0.1$ and $\gamma = 1$. Here x represents the volume, $x \equiv \mathcal{V}$.

Substituting the solution (3.3) into (3.2) we obtain

$$V_F^{extr.} = \frac{3|W_0|^2 \eta_0 (2 \log \mathcal{V}_o - 8 + \frac{\xi_0}{\eta_0})}{2\mathcal{V}_o^3} \equiv \eta_0 \frac{|W_0|^2}{\mathcal{V}_o^3}. \quad (3.4)$$

A simple analysis shows that this is a minimum of the potential as long as $\eta_0 < 0$. However, since $\frac{|W_0|^2}{\mathcal{V}_o^3} > 0$, the potential (3.4) at the minimum acquires a negative value, it turns out that the F-term potential predicts an anti-de Sitter (AdS) vacuum. In Fig. 1, left panel, the F-term potential with AdS minimum is plotted for a choice of the parameters ξ_0, η_0, W_0 .

Despite the forgoing negative F-term contribution, the potential can be lifted to a de Sitter minimum, once D-term contributions [16, 31] associated with the universal $U(1)$ symmetries of the D7-branes are taken into account [10]. In this context, the D-term potential is generated in hidden sectors by D7-branes wrapping the 3 divisors whose volumes are τ_i . More precisely, in the present geometric setting, there are D-term contributions due to the universal $U(1)$ factors associated with the $D7$ brane stacks. These terms have the general form [10, 31]

$$V_D = \sum_{i=1}^3 \frac{g_{D7_i}^2}{2} \left(Q_i \partial_{T_i} K + \sum_j q_j |\Theta^j|^2 \right)^2, \quad (3.5)$$

where $g_{D7_i} = (\text{Re} T_i)^{-1}$ and Q_j, q_j are “charges”, while Θ^j represent matter fields charged under the $U(1)$ gauge factors. Some aspects regarding strings at $D7$ intersections in particular have been recently discussed in [17]. The fields Θ^j carry charges under the $U(1)$ factors associated with the D-branes. A compelling question then arises regarding the possible contributions of these fields in the D-term, and hence, in the scalar potential. One possible scenario is that non-zero field vevs are chosen so that the D-term vanishes, and in this case, the uplift should be realized by a suitable modification (see footnote 10), or the standard procedure of introducing $\overline{D3}$ branes (see [32]). Here we make a simplified assumption that these singlets have vanishing vevs. Whether the vevs of these fields are zero or not, depends on the specific details of the effective model. One possibility is to assume that in the effective field theory limit,

all these fields Θ^j are minimized at $\langle \Theta^j \rangle = 0$. Alternatively, even in the case of non-zero vevs, however, it might happen that there could be accidental cancellations that can reduce the significance of their contributions. Some related discussions regarding these issues can be found also in [33]. Here, since the D-term potential is only used to uplift the non-supersymmetric AdS vacuum of the F-term potential, following closely previous works (see for example [10], [16] and [31]) we make the reasonable assumption that moduli fields dominate over other fields vevs. Therefore, it is adequate for our purposes to assume that the flux induced D-term piece $\propto Q_i \partial_{T_i} K$ dominates, and therefore we minimize the potential by setting $\langle \Theta^j \rangle$ vevs zero. For $\langle \Theta^j \rangle = 0$ the second term in the parenthesis vanishes and each component of the D-term acquires a simple -model independent- form $V_{D_i} \approx Q_i^2 / \tau_i^3$. Then, the total D-term potential, being the sum of three components, is approximated by [10]

$$V_{\mathcal{D}} = \sum_{i=1}^3 \frac{d_i}{\tau_i} \left(\frac{\partial \mathcal{K}}{\partial \tau_i} \right)^2 \approx \sum_{i=1}^3 \frac{d_i}{\tau_i^3} \equiv \frac{d_1}{\tau_1^3} + \frac{d_3}{\tau_3^3} + \frac{d_2 \tau_1^3 \tau_3^3}{\mathcal{V}^6}, \quad (3.6)$$

Here, d_i are positive constants related to the charges $d_i \sim Q_i^2 > 0$. Furthermore, using the volume formula $\mathcal{V}^2 = \tau_1 \tau_2 \tau_3$, we have substituted the modulus τ_2 with its equivalent $\tau_2 = \mathcal{V}^2 / (\tau_1 \tau_3)$. Then, the total effective potential is the sum $V_{\text{eff}} = V_F + V_D$ which can be minimized with respect to \mathcal{V} and the two remaining Kähler moduli $\tau_{1,2}$. We have assumed that the F-part of the potential depends on the total volume, hence the explicit dependence of V_{eff} on τ_1, τ_3 only comes through the V_D part. It is found that the two minimization conditions with respect to $\tau_{1,3}$ determine the ratios between the moduli, i.e., $\left(\frac{\tau_i}{\tau_j} \right)^3 = \frac{d_i}{d_j}$ [34]. Expressed in terms of the stabilized total volume \mathcal{V} , the conditions for the two τ_i can be written as

$$\tau_i^3 = \left(\frac{d_i^2}{d_k d_j} \right)^{\frac{1}{3}} \mathcal{V}^2,$$

where $i = 1, 3$. In this case, the D-term potential receives the simple form

$$V_D \approx \frac{d}{\mathcal{V}^2}, \quad \text{with} \quad d = 3(d_1 d_2 d_3)^{\frac{1}{3}}. \quad (3.7)$$

By choosing $\phi_1 = \phi_2 = \phi$ and $\alpha = \beta$, the D-term contribution from the Higgs fields vanishes. Then the effective potential has the following form

$$V_{\text{eff}} \simeq \frac{\kappa^2 \alpha (M^2 - \phi^2)^2 + 2\gamma \kappa^2 S^2 \phi^2}{3\alpha \gamma \mathcal{V}^{4/3}} + \frac{3W_0^2(2\eta_0 \log(\mathcal{V}) - 8\eta_0 + \xi_0)}{2\mathcal{V}^3} + \frac{d}{\mathcal{V}^2}. \quad (3.8)$$

The shape of the potential along the volume modulus \mathcal{V} when both F- and D-terms are included is shown in the right panel of Fig. 1. As can be seen, a positive D-term is sufficient to uplift the potential along the volume direction so that we achieve a de Sitter minimum. In order to find the extrema of the potential along the ϕ and S directions, we require the vanishing of its corresponding derivatives. Thus, for ϕ we impose the condition

$$\frac{dV_{\text{eff}}}{d\phi} = 0 \Rightarrow \frac{\kappa^2 (4\alpha \phi^3 - 4\alpha M^2 \phi + 4\gamma S^2 \phi)}{3\alpha \gamma \mathcal{V}^{4/3}} = 0, \quad (3.9)$$

which is solved for the following three ϕ -values

$$\phi = 0, \quad \phi_{\pm} = \pm \sqrt{M^2 - \frac{\gamma}{\alpha} S^2}. \quad (3.10)$$




Figure 2: The shape of the effective potential in $\phi - S$ plane for the choice of parameters $\xi_0 = 10$, $\eta_0 = -0.92$, $\gamma_0 = 32000$, $\kappa = 0.1$, $\gamma = 1$. and $d = 10^{-4.52}$.

Similarly along the S direction

$$\frac{dV_{\text{eff}}}{dS} = 0 \Rightarrow \frac{4\kappa^2 S \phi^2}{3a\alpha \gamma^{4/3}} = 0, \quad (3.11)$$

which yields

$$S = 0. \quad (3.12)$$

Combining Equations (3.10) and (3.12), in the large volume limit we obtain the following solutions

$$(S = 0, \phi = 0), \quad (S = 0, \phi = \pm M). \quad (3.13)$$

We have already dealt with the minimization of V_F with respect to the volume modulus. However, in the presence of D-terms, the minima along the volume direction are shifted. Thus, requiring the vanishing of the derivative of (3.8) with respect to γ , we obtain the equation

$$\begin{aligned} \frac{dV_{\text{eff}}}{d\gamma} = 0 \Rightarrow & -\frac{4\kappa^2 (\alpha\phi^4 + \alpha M^4 - 2\alpha M^2 \phi^2 + 2\gamma S^2 \phi^2)}{9a\alpha \gamma^{7/3}} - \frac{2d}{\gamma^3} \\ & - \frac{9(2\eta_0 W_0^2 \log(\gamma) - 8\eta_0 W_0^2 + \xi_0 W_0^2)}{2\gamma^4} + \frac{3\eta_0 W_0^2}{\gamma^4} = 0. \end{aligned} \quad (3.14)$$

In the large volume limit, to a good approximation, the above equation gives the solution

$$\gamma_o \approx \frac{9\eta_0 W_0^2}{2d} \gamma \left(\frac{2de^{\frac{13}{3} - \frac{\xi_0}{2\eta_0}}}{9\eta_0 W_0^2} \right), \quad (3.15)$$

where \mathcal{W} represents the product-log (Lambert) function. The shape of the scalar potential V_{eff} in the φ - S plane is displayed in Fig. 2. As discussed earlier the D-term contribution in the effective potential ensures the existence of de Sitter vacua when S approaches zero.

3.1 Inflationary phase

Up to this point, we have analyzed in detail the scalar potential of the effective theory and the role of the various fields in its final shape. Therefore we are now fully equipped with all the tools and the necessary ingredients to examine whether cosmological inflation is realized in the present model. In a previous approach, within the same type-IIB framework and the geometric setup of intersecting D7 brane stacks, the inflaton field was associated with the logarithm of the compactification volume modulus. The accumulation of the required 60 efolds to realize slow-roll inflation was converted to a lower bound on the minimum vacuum energy [34], yet much bigger than the cosmological constant. Subsequently, a new “waterfall” field was introduced which adds a new direction of the potential. The waterfall field rolls down towards the new lower minimum while at the same time, it ends inflation. It was shown [17] that this role can be realized by open string states oscillating near the intersections of the D7 stacks.

In the present case where the physical states from the effective theory model have been taken into account, new possibilities have emerged. At the minimum along the compactification volume modulus \mathcal{V} , the Higgs field φ , and the singlet S add new directions (transverse to that of \mathcal{V}) providing new lower minima for the scalar potential. As such, they are potential candidates for waterfall fields. In the present scenario inflation proceeds along the local minimum $\varphi = 0$ (the inflationary track), beginning at large S values. An instability occurs at the waterfall point $S_c^2 = M^2$, which is the value of S , such that $S_c = \frac{\partial^2 V}{\partial S^2} \big|_{S=0} = 0$. At this point, the field falls naturally into one of the two minima at $\varphi = \pm M$. At large S , the scalar potential is approximately quadratic in φ , whereas at $S = 0$, equation (3.8) becomes a Higgs potential. Along the inflationary track the constant term

$$V_0^{\text{vol}} = \frac{\kappa^2 M^4}{3a\mathcal{V}_o^{4/3}} + \frac{3W_0^2(2\eta_0 \log(\mathcal{V}_o) - 8\eta_0 + \xi_0)}{2\mathcal{V}_o^3},$$

is present at tree level, thus SUSY is broken during inflation. When SUSY breaks, splitting between fermionic and bosonic mass multiplets is created and contributions to radiative corrections occur. Following [35, 36], the soft terms are¹¹

$$\Delta V_{\text{soft}} = \left[(m_{3/2}^2 + V_0) - \frac{2}{3\mathcal{V}_o^2} W_o^2 + \dots \right] M^2 y^2 \frac{\mathcal{V}_o^{2/3}}{3a\gamma} = M_{S_c}^2 y^2, \quad (3.16)$$

where in the above equation, y defines the ratio $y = S/M$, and

$$M_{S_c} = \left[(m_{3/2}^2 + V_0) - \frac{2}{3\mathcal{V}_o^2} W_o^2 + \dots \right] M^2 \frac{\mathcal{V}_o^{2/3}}{3a\gamma},$$

¹¹Note that we use S for both the canonically normalized field and the non-canonically one. The scaling factor required for the canonically normalized field is adjusted by multiplying $\mathcal{V}_o^{2/3}/(3a\gamma)$.

represents the soft mass parameter for canonical normalized field y , and V_0 is the minimum of the potential (3.8).¹² For appropriate set of parameters, V_0 takes the value of the cosmological constant. The first extremum ($S = 0, \varphi = 0$) is the maximum of the potential. The $\varphi = 0$ corresponds to the trajectory of the standard hybrid inflation for which $\{\varphi = 0, S > M\}$. When the inflaton reaches $S = M$, the waterfall field takes over and the inflaton moves towards the minimum at $\varphi = \pm M$. Moreover, SUSY is broken along the inflationary track, and the radiative corrections along with the soft SUSY-breaking potential V_{soft} can lift its flatness, while also providing the necessary slope for driving inflation. The effective contribution of the one-loop radiative corrections can be calculated using the Coleman-Weinberg formula [38] as

$$\Delta V_{1\text{-loop}} = \frac{\kappa^4 M^4 y^4}{144\pi^2 a^2 \gamma^2 \gamma_o^{4/3}} \left[F(y) - \left(\frac{1}{54a^2 \alpha^2 \gamma^2} - \frac{3}{2} \gamma_o^{8/3} \right) \right], \quad (3.17)$$

where

$$F(y) = \frac{1}{81a^2 \alpha^2 \gamma^2} \log \left(\frac{\kappa^2 M^2 y^2}{27a^2 \alpha \gamma^2 Q^2 \gamma_o^{2/3}} \right) - \gamma_o^{8/3} \log \left(\frac{\kappa^2 M^2 y^2 \gamma_o^{2/3}}{3a\gamma Q^2} \right). \quad (3.18)$$

It should be observed that the Coleman-Weinberg correction is computed along the inflationary trajectory, wherein the field S takes a non-zero value while the field φ remains fixed at zero. Furthermore, we set $V_0 = 3.2 \times 10^4$ for the volume field, and the mass spectrum solely depends on S . The detailed calculations are provided in Appendix A.

Including the various contributions computed above, we may write the scalar potential along the inflationary trajectory (i.e. $\varphi_1 = \varphi_2 = 0$) as follows,

$$\begin{aligned} V &\simeq V_F + V_D + \Delta V_{1\text{-loop}} + \Delta V_{\text{soft}}, \\ &\simeq \kappa^2 M^4 \left(\frac{V_0^{\text{vol}}}{\kappa^2 M^4} + \frac{\kappa^2 y^4 F(y)}{144\pi^2 a^2 \gamma^2 \gamma_o^{4/3}} - \frac{\kappa^2 y^4}{144\pi^2 a^2 \gamma^2 \gamma_o^{4/3}} \left(\frac{1}{54a^2 \alpha^2 \gamma^2} - \frac{3}{2} \gamma_o^{8/3} \right) + \frac{M_{S_c}^2 y^2}{\kappa^2 M^2} \right), \end{aligned} \quad (3.19)$$

M_{S_c} is the soft-mass parameter of the field S , and $V_0 \approx 0$ due to its extremely small magnitude compared to $m_{3/2}^2$.

The predictions for the various inflationary observables are estimated using the standard slow-roll parameters defined as,

$$\begin{aligned} \varepsilon &= \frac{1}{2} \left(\frac{1}{M} \right)^2 \left(\frac{V'}{V} \right)^2, \quad \eta = \frac{1}{M^2} \left(\frac{V''}{V} \right), \\ \xi^2 &= \frac{1}{M^4} \left(\frac{V' V'''}{V^2} \right), \end{aligned} \quad (3.20)$$

where prime denotes the derivative with respect to y . In the slow-roll approximation, the scalar spectral index n_s , the tensor-to-scalar ratio r and the running of the scalar spectral index $\alpha_s \equiv dn_s/d \ln k$ are given by,

$$\begin{aligned} n_s &\simeq 1 + 2\eta - 4\varepsilon, \quad r \simeq 16\varepsilon, \\ \alpha_s &\simeq 16\varepsilon\eta - 24\varepsilon^2 - 2\xi^2. \end{aligned} \quad (3.21)$$

¹²The soft mass parameter, M_{S_c} , depends on both the gravitino mass, $m_{3/2}$, and the constant parameter W_0 . By choosing W_0 to be very small, the dominant term of the soft masses becomes dependent on the gravitino mass [37].




Figure 3: Variations of r and M in $\kappa - m_{3/2}$ plane. The upper line has M fixed at a value equal to M_{string} . The other two lines correspond to lower M values, as indicated in the plot.

The value of the scalar spectral index n_s in the Λ CDM model is $n_s = 0.9665 \pm 0.0038$ [39].

The amplitude of the scalar power spectrum is given by,

$$A_s(k_0) = \frac{1}{24\pi^2} \left(\frac{V(y_0)}{\varepsilon(y_0)} \right), \quad (3.22)$$

where $A_s(k_0) = 2.137 \times 10^{-9}$ at the pivot scale $k_0 = 0.05 \text{ Mpc}^{-1}$ as measured by Planck 2018 [39]. The relevant number of e-folds, N_0 , before the end of inflation is,

$$N_0 = 2M^2 \int_{y_e}^{y_0} \left(\frac{V}{V'} \right) dy, \quad (3.23)$$

where $y_0 \equiv y(k_0)$ is the field value at the pivot scale k_0 , and y_e is the field value at the end of inflation. As the case may be, the value of y_e is fixed either by the breakdown of the slow-roll approximation ($\eta(y_e) = -1$) or by a ‘waterfall’ destabilization occurring at the value $y_e = 1$.

3.2 Numerical results

The results of our numerical calculations are displayed in Fig. 3, which show the ranges of r , M in the $\kappa - M_{sc}$ plane. We consider up to second-order approximation on the slow-roll parameters, and for simplicity, we set $\gamma = 1$, $\alpha = 10^{-7}$, $\mathcal{V}_0 = 32000$ and $y_e = 1$. Moreover, we have fixed the spectral index n_s to the central value ($n_s = 0.96655$) from Planck’s data.

We further require $\kappa \leq 0.1$, the Higgs mass parameter $10^{15} \leq M \leq M_{string} \sim 1/\mathcal{V}_0^{1/2} = 5.5 \times 10^{-3} M_p = 1.36 \times 10^{16} \text{ GeV}$, and FT, which is defined as the difference of field value at pivot scale at

the end of inflation $FT \equiv y_0 - y_e \lesssim 0.1$. These constraints appear in Fig. 3 as the boundaries of the allowed region in the $\kappa - M_{S_c}$ plane. In our analysis, the soft SUSY contributions, along with the radiative corrections, play the dominant role in obtaining a parametric space consistent with experimental bounds, parametrized by M_{S_c} and a . Additionally, in the entire parameter space, a remains in close proximity to the value of 1.

In our analysis, we carefully selected a parameter space where the contribution of the logarithmic terms of in (3.18) is suppressed. However, for $\kappa \geq 0.1$, the logarithmic terms gradually become more dominant compared to the other components of the potential in (3.19). This dominance results in a deviation of the spectral index (n_s) from the Planck bound.

For the scalar spectral index n_s fixed at Planck's central value, $n_s = 0.96655$, according to our numerical analysis, the exact range of parameters where acceptable solutions occur is presented below

$$\begin{aligned} 1.6 \times 10^{-5} &\lesssim \kappa \lesssim 0.1, \\ (1 \times 10^{15} &\lesssim M \lesssim 1.3 \times 10^{16}) \text{ GeV}, \\ (4.6 \times 10^5 &\lesssim M_{S_c} \lesssim 1 \times 10^{11}) \text{ GeV}, \\ 1.9 \times 10^{-14} &\lesssim r \lesssim 1.4 \times 10^{-4}, \\ 0.86 &\lesssim a \lesssim 1. \end{aligned}$$

Examining the exact ranges of solutions, our calculations predict a low tensor-to-scalar ratio (r) compared to current experimental bounds. However, ongoing and future gravity waves experiments are expected to reach much smaller ranges of tensor-to-scalar ratio, comparable to our numerical predictions. Note that Fig. 3 shows the parametric space where $0.86 \lesssim a \lesssim 1$, aligning with the conditions for dark radiation, as discussed in the following section.

4 Reheating and dark radiation

After the end of inflation, the lightest moduli will begin to oscillate around their minima, acquiring a large energy density in the process. The decay products of the modulus fall into two categories. The first are decays that go to the visible sector that is, particles of the SM, or its extensions such as the MSSM. The decays to visible matter induce reheating, after which the standard hot Big Bang cosmology follows. In addition, there may also be decays to hidden sector states. The hidden sector contains several candidates for dark radiation, such as massless axions or light hidden gauge bosons. Let us consider the three Kähler moduli case, $T_k = \tau_k + ia_k$, $\mathcal{V} = \sqrt{\tau_1 \tau_2 \tau_3}$ where a_k is the RR-axion. Decay to the light axion a_k takes place primarily through the supergravity kinetic terms for the Kähler moduli which read as,

$$\mathcal{L} \supset K_{i\bar{j}} \partial_\mu T^i \partial^\mu \bar{T}^{\bar{j}}. \quad (4.1)$$

The tree level Kähler potential is,

$$K = -2 \log \sqrt{(T_1 + \bar{T}_1)(T_2 + \bar{T}_2)(T_3 + \bar{T}_3)} = -\log(\tau_1 \tau_2 \tau_3) + \dots, \quad (4.2)$$

where the dots represent constant terms that are ignored. Then, the Kähler matrix is found to be

$$K_{i\bar{j}} = \frac{1}{4} \text{diag} \left(\frac{1}{\tau_1^2}, \frac{1}{\tau_2^2}, \frac{1}{\tau_3^2} \right). \quad (4.3)$$

Therefore, Eq (4.1) can be rewritten as,

$$\mathcal{L} \supset \frac{1}{4} \frac{1}{\tau_i^2} \partial_\mu \tau_i \partial^\mu \tau_i.$$

Note here that we have set the reduced Planck mass $M_p = 1$. To put this into canonical form we need to find the transformation $\tau_i(u_i)$ so that

$$\mathcal{L} \supset \frac{1}{2} \sum_i \partial_\mu u_i \partial^\mu u_i.$$

This implies

$$\tau_i = e^{\sqrt{2}u_i}.$$

The moduli fields for canonical kinetic terms take the form

$$u_k = \frac{1}{\sqrt{2}} \log \tau_k.$$

The corresponding volume modulus is

$$t = \frac{u_1 + u_2 + u_3}{\sqrt{3}} = \frac{1}{\sqrt{3}} \frac{1}{\sqrt{2}} \sum_k \log \tau_k = \sqrt{\frac{2}{3}} \log \mathcal{V}.$$

The transverse directions are

$$u = \frac{u_1 - u_2}{\sqrt{2}} = \frac{1}{2} \log \frac{\tau_1}{\tau_2}, \quad v = \frac{u_1 + u_2 - 2u_3}{\sqrt{6}} = \frac{1}{\sqrt{3}} \log \frac{\tau_1 \tau_2}{\tau_3^2}.$$

We reverse the relations

$$\begin{pmatrix} u_1 \\ u_2 \\ u_3 \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & 0 & -\sqrt{\frac{2}{3}} \end{pmatrix} \begin{pmatrix} t \\ u \\ v \end{pmatrix}$$

while substituting to the Lagrangian part for axions we have

$$\begin{aligned} \mathcal{L} \supset & \frac{1}{\sqrt{3}} t (\partial_\mu a_1 \partial^\mu a_1 + \partial_\mu a_2 \partial^\mu a_2 + \partial_\mu a_3 \partial^\mu a_3) \\ & + \frac{1}{\sqrt{2}} u (\partial_\mu a_1 \partial^\mu a_1 - \partial_\mu a_2 \partial^\mu a_2) \\ & + \frac{1}{\sqrt{6}} v (\partial_\mu a_1 \partial^\mu a_1 + \partial_\mu a_2 \partial^\mu a_2 - 2 \partial_\mu a_3 \partial^\mu a_3). \end{aligned} \quad (4.4)$$

The decay rate of the lightest modulus u to axions is given by

$$\Gamma(u \rightarrow a_1 a_1) = \frac{1}{64\pi} m_u^3, \quad (4.5)$$

where m_u is the modulus mass. In the large volume scenario a distinct hierarchy of mass scales is generated, (for details see Ref [40]). The mass eigenstates after diagonalization in units of $M_p = 1$ can be written as,

$$m_t = m_u = m_v \sim \frac{1}{\mathcal{V}_o^{3/2}}, \quad m_{a_i} \sim e^{-2\pi\mathcal{V}_o^{2/3}}, \quad m_{\text{soft}} \sim \frac{1}{\mathcal{V}_o^{2/3}}, \quad m_{3/2} \sim \frac{1}{\mathcal{V}_o}, \quad M_{\text{string}} \sim \frac{1}{\mathcal{V}_o^{1/2}}.$$

Similarly, the dominant visible-sector decay channel is the decay to Higgs bosons. Specializing to the MSSM case, we can make the identifications $\Phi_1 = H_u$ and $\Phi_2 = H_d$. Each of these is a two-component complex field, so there are eight degrees of freedom. Then the decay rate can be derived by including the matter contribution to the Kähler potential

$$\mathcal{L} \supset \frac{3a\lambda}{\sqrt{2}} H_u H_d \square u + h.c. + \dots \quad (4.6)$$

The dominant contribution to the decay of light moduli u comes from the Giudice-Masiero coupling [41] $3a\lambda H_u H_d \square u$ as all other couplings are mass-suppressed [42]. Each field is a complex doublet, hence we include the partial widths from each of the four decay channels. This yields

$$\Gamma(u \rightarrow H_u H_d) = \frac{9a^2\lambda^2}{8\pi} m_u^3. \quad (4.7)$$

The present-day radiation content of the Universe can be described in terms of the energy density associated with each relativistic particle species at present. This radiation consists of photons and neutrinos plus any additional hidden components, which we call dark radiation (DR):

$$\rho_{\text{radiation}} = \rho_{\text{photon}} + \rho_{\text{neutrino}} + \rho_{\text{DR}}. \quad (4.8)$$

We can express this in terms of an effective number of neutrino species, N_{eff} , as follows

$$\rho_{\text{radiation}} = \rho_{\text{photon}} \left(1 + \frac{7}{8} \left(\frac{4}{11} \right)^{4/3} N_{\text{eff}} \right). \quad (4.9)$$

Any excess can be accounted for by the presence of DR which is expressed as

$$\rho_{\text{DR}} = \rho_{\text{photon}} \left(\frac{7}{8} \left(\frac{4}{11} \right)^{4/3} \Delta N_{\text{eff}} \right), \quad (4.10)$$

where $\Delta N_{\text{eff}} = N_{\text{eff}} - 3.046$ is the change in the effective number of neutrino species. If there were no dark radiation we would expect to find $N_{\text{eff}} = 3.046$ which is slightly greater than 3 to account for partial reheating due to e^+e^- annihilation. ΔN_{eff} can also be written in terms of decay rate channels

$$\Delta N_{\text{eff}} = \frac{43}{7} \left(\frac{10.75}{g_*(T_r)} \right)^{\frac{1}{3}} \frac{\Gamma_{\tau \rightarrow \text{DR}}}{\Gamma_{\tau \rightarrow \text{SM}}} = \frac{43}{7} \left(\frac{10.75}{g_*(T_r)} \right)^{\frac{1}{3}} \frac{1}{72a^2\lambda^2}, \quad (4.11)$$

where $g_*(T_R h)$ is the effective degree of freedom at the time of reheating the Universe. The measured values of N_{eff} require $\Delta N_{\text{eff}} \lesssim 0.95$ at the 95% confidence level, which translates into a bound on the parameters of the model, $a \cdot \lambda \gtrsim 0.1688$. In Fig. 3, the parametric space predicted by the inflationary analysis has $0.86 \leq a \leq 1$. This range aligns with the values predicted by $\Delta N_{\text{eff}} \lesssim 0.95$ and is consistent with dark radiation. Within this parametric space, we obtain a tensor-to-scalar ratio values of $r \leq 1.4 \times 10^{-4}$, $1 \times 10^{15} \lesssim M \lesssim 1.3 \times 10^{16}$ GeV, and a soft mass parameter $4.6 \times 10^5 \lesssim M_{S_c} \lesssim 1 \times 10^{11}$ GeV.

The isotropy of the cosmic microwave background (CMB) over large scales can be explained by inflation, which is followed by a period of reheating. During this process, the expansion rate slows, leading to the transfer of energy to SM particles that subsequently enter local thermal equilibrium. We




Figure 4: Variations of the reheating temperature (T_r) with respect to coefficient a consistent with dark radiation constraint ($\Delta N_{\text{eff}} \lesssim 0.95$) at 95% confidence level.

consider the scenario where the Universe is reheated by a modulus u decaying into SM particles. In this case the reheating temperature T_r , using Eq.(4.5) and Eq.(4.11) is defined as

$$T_r = \sqrt{\Gamma_u} = \sqrt{\frac{63}{344\pi} \Delta N_{\text{eff}} \left(\frac{g_*(T_r)}{10.75} \right)^{1/2} a^2 \lambda^2 m_u^3}. \quad (4.12)$$

For a SUSY scale at the TeV regime, the constraint on reheating temperature is $T_r \lesssim 1$ GeV, which corresponds to $g_*(T_r) = 224/7$, see Ref [29, 43]. For the present model, we have $\mathcal{V}_o \sim 3.2 \times 10^4$ which corresponds to a SUSY scale: $m_{\text{soft}} > 10$ TeV. As a result, the constraint on the reheating temperature is relaxed in this scenario. For SUSY scale greater than 10 TeV we have $g_*(T_r) = 106.75$, so the reheating temperature is $T_r \sim 10^7$ GeV as shown in Fig. 4.

5 Conclusions

We investigated the cosmological and low energy supersymmetry implications of an effective model [10] stemming from a geometric configuration of intersecting three D7-branes stacks, within the framework of type-IIB string theory. In this model perturbative string loop corrections which depend logarithmically on the compactification volume \mathcal{V} , and D-terms associated with the universal $U(1)$ factors of D7-brane stacks generate an effective scalar potential with de Sitter vacuum, and stabilize all three Kähler moduli fields of the specific geometric setting. In the present work we took into account the effects of ordinary matter contributions in the Kähler potential of the effective model and, in particular, we focused on the role of a generic pair of Higgs fields Φ_1, Φ_2 (related to the gauge group of the effective theory) on low en-

ergy phenomenology predictions and various cosmological observables. We included matter field content and soft-term contributions as well as Coleman-Weinberg corrections to the previously derived potential and studied the implementation of the standard hybrid inflationary scenario where a singlet gauge field sharing common couplings with the Higgs fields in the superpotential plays the role of the inflaton whilst the Higgs states act as waterfall fields. Fixing the spectral index at central value, $n_s = 0.96655$, we provided predictions for the remaining cosmological observables in accordance with the latest Planck data. In particular, we predicted the value of the tensor-to-scalar ratio to be $r \sim 2 \times 10^{-4}$ which is much smaller than the current experimental bounds, however, within the reach of future designed experiments. Next, we discussed the decay of the lighter Kähler moduli after the end of inflation, which includes modes to visible as well as invisible particles. In particular, in the context of an MSSM effective theory and the presence of a Giudice-Masiero coupling, the dominant decay of the lightest modulus is to the Higgs fields, in accordance with previous computations [29]. Furthermore, we investigated predictions of the model to dark radiation production and we found $\Delta N_{\text{eff}} \leq 0.95$ at 2σ confidence level. This requires the model parameters a and λ (associated with the couplings $\propto a\lambda(\Phi_1\Phi_2 + h.c)$ in the Kähler potential) to satisfy the bound $a\lambda \gtrsim 0.1688$, which for $\lambda \approx 1$ translates into a bound for a in the perturbative regime. Regarding other vital low energy predictions for a volume fixed at $\mathcal{V}_o \sim 3.2 \times 10^4$ which ensures a dS minimum, we found that the SUSY scale is greater than 10 TeV. As a result, the constraint imposed on the reheating temperature is relaxed in this scenario. For SUSY scale (>10 TeV) we have $g_*(T_r) = 106.75$ [29], and the reheating temperature is $T_r \sim 10^7$ GeV .

A Coleman-Weinberg Corrections

In this Appendix we present analytically the computation of the CW corrections. Using the effective potential

$$V_{\text{eff}} = \frac{\kappa^2 \alpha | (M^2 - \Phi_1 \Phi_2) |^2 + 2\gamma \kappa^2 S^2 (\alpha \Phi_1 \Phi_1^\dagger + \beta \Phi_2 \Phi_2^\dagger)}{3\alpha\beta\gamma\gamma^{4/3}} + \frac{3W_0^2(2\eta_0 \log(\gamma) - 8\eta_0 + \xi_0)}{2\gamma^3} + \frac{d}{\gamma^2}, \quad (\text{A.1})$$

the scalar mass matrix along the inflationary track read as,

$$M_S^2 = \begin{pmatrix} \frac{\kappa^2 S^2}{6\alpha\beta\gamma^{4/3}} & 0 & -\frac{\kappa^2 M^2}{3a\gamma^{4/3}} & 0 & 0 & 0 & 0 \\ 0 & \frac{\kappa^2 S^2}{6\alpha\beta\gamma^{4/3}} & 0 & \frac{\kappa^2 M^2}{3a\gamma^{4/3}} & 0 & 0 & 0 \\ -\frac{\kappa^2 M^2}{3a\gamma^{4/3}} & 0 & \frac{\kappa^2 S^2}{6\alpha\gamma^{4/3}} & 0 & 0 & 0 & 0 \\ 0 & \frac{\kappa^2 M^2}{3a\gamma^{4/3}} & 0 & \frac{\kappa^2 S^2}{6\alpha\gamma^{4/3}} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{28\kappa^2 M^4}{27a\gamma^{10/3}} + \dots \end{pmatrix} \quad (\text{A.2})$$

After diagonalization and choosing $\alpha = \beta$ the scalar mass spectrum is

$$M_S^2 = \left[0, 0, \frac{\kappa^2 M^2 \left(y^2 - \frac{2\alpha}{\gamma} \right)}{6a\gamma^{4/3}\alpha}, \frac{\kappa^2 M^2 \left(y^2 - \frac{2\alpha}{\gamma} \right)}{6a\gamma^{4/3}\alpha}, \frac{\kappa^2 M^2 \left(y^2 + \frac{2\alpha}{\gamma} \right)}{6a\gamma^{4/3}\alpha}, \frac{\kappa^2 M^2 \left(y^2 + \frac{2\alpha}{\gamma} \right)}{6a\gamma^{4/3}\alpha}, \frac{28\kappa^2 M^4}{27a\gamma^{10/3}} + \dots \right] \quad (\text{A.3})$$

where $y^2 = S^2/M^2$. Similarly, the generic fermionic mass matrix is defined as

$$M_{Fij} = e^{\mathcal{K}/2} \left(W_{ij} + \mathcal{K}_{ij} W + \mathcal{K}_i W_j + \mathcal{K}_j W_i + \mathcal{K}_i \mathcal{K}_j W - \mathcal{K}^{k\bar{l}} \mathcal{K}_{ij\bar{l}} D_k W \right). \quad (\text{A.4})$$

Using the Superpotential (2.4) and the Kähler potential in (2.9) we obtain

$$M_{Fij} = \begin{pmatrix} -\frac{6a\gamma\kappa M^2 S}{\gamma^{2/3}} + \frac{9a^2\gamma^2 S^2}{\gamma^{4/3}} + \dots & 0 & 0 & \frac{2\kappa M^2}{\gamma} + \frac{2a\gamma S(\kappa M^2 S - 4)}{\gamma^{5/3}} + \dots \\ 0 & 0 & \kappa S & 0 \\ 0 & \kappa S & 0 & 0 \\ \frac{2\kappa M^2}{\gamma} + \frac{2a\gamma S(\kappa M^2 S - 4)}{\gamma^{5/3}} + \dots & 0 & 0 & \frac{6}{\gamma^2} + \frac{34}{3}a\gamma S^2 \left(\frac{1}{\gamma} \right)^{8/3} + \dots \end{pmatrix}. \quad (\text{A.5})$$

After diagonalization, the fermionic masses are

$$M_{Fij}^2 = \left[\kappa^2 M^2 y^2, \kappa^2 M^2 y^2, \frac{36a^2 \kappa^2 M^6 y^2 \gamma^2}{\gamma^{4/3}} + \dots, \frac{36a^2 \kappa^2 M^6 y^2 \gamma^2}{\gamma^{4/3}} + \dots \right]. \quad (\text{A.6})$$

The fermionic mass $\frac{36a^2 \kappa^2 M^6 y^2 \gamma^2}{\gamma^{4/3}}$ and the bosonic mass $\frac{28\kappa^2 M^4}{27a\gamma^{10/3}}$ are small compared to other masses, so their contribution to the Coleman-Weinberg potential is suppressed. Note that we use the symbol y for both the canonically, and the non-canonically normalized field. The scaling factor required for the canonically normalized field is adjusted by multiplying $\gamma_0^{2/3}/(3a\gamma)$. Additionally, we focus on the region where $(y^2 \gg 2\alpha/\gamma)$. The effective contribution of the one-loop radiative corrections can be calculated using the Coleman-Weinberg formula

$$\Delta V_{1\text{-loop}} = \frac{1}{64\pi^2} \left[M_S^4 \log \left(\frac{M_S^2}{Q^2} \right) - 2M_F^4 \log \left(\frac{M_F^2}{Q^2} \right) - \frac{3}{2} (M_S^4 - 2M_F^4) \right] \quad (\text{A.7})$$

$$\Delta V_{1\text{-loop}} = \frac{\kappa^4 M^4 y^4}{144 \pi^2 a^2 \gamma^2 \gamma_o^{4/3}} \left[F(y) - \left(\frac{1}{54 a^2 \alpha^2 \gamma^2} - \frac{3}{2} \gamma_o^{8/3} \right) \right], \quad (\text{A.8})$$

where

$$F(y) = \frac{1}{81 a^2 \alpha^2 \gamma^2} \log \left(\frac{\kappa^2 M^2 y^2}{27 a^2 \alpha \gamma^2 Q^2 \gamma_o^{2/3}} \right) - \gamma_o^{8/3} \log \left(\frac{\kappa^2 M^2 y^2 \gamma_o^{2/3}}{3 a \gamma Q^2} \right). \quad (\text{A.9})$$

References

- [1] E. Palti, “The Swampland: Introduction and Review,” *Fortsch. Phys.* **67** (2019) no.6, 1900037 [[arXiv:1903.06239](#) [hep-th]].
- [2] N. B. Agmon, A. Bedroya, M. J. Kang and C. Vafa, “Lectures on the string landscape and the Swampland,” [[arXiv:2212.06187](#) [hep-th]].
- [3] T. Van Riet and G. Zoccarato, “Beginners lectures on flux compactifications and related Swampland topics,” [[arXiv:2305.01722](#) [hep-th]].
- [4] V. Balasubramanian, P. Berglund, J. P. Conlon and F. Quevedo, “Systematics of moduli stabilization in Calabi-Yau flux compactifications,” *JHEP* **03** (2005), 007 [[arXiv:hep-th/0502058](#) [hep-th]].
- [5] M. Berg, M. Haack and E. Pajer, “Jumping Through Loops: On Soft Terms from Large Volume Compactifications,” *JHEP* **09** (2007), 031 [[arXiv:0704.0737](#) [hep-th]].
- [6] M. Reece and W. Xue, “SUSY’s Ladder: reframing sequestering at Large Volume,” *JHEP* **04** (2016), 045 [[arXiv:1512.04941](#) [hep-ph]].
- [7] M. Cicoli, I. García-Etxebarria, C. Mayrhofer, F. Quevedo, P. Shukla and R. Valandro, “Global Orientifolded Quivers with Inflation,” *JHEP* **11** (2017), 134 [[arXiv:1706.06128](#) [hep-th]].
- [8] M. Cicoli, I. G. Etxebarria, F. Quevedo, A. Schachner, P. Shukla and R. Valandro, “The Standard Model quiver in de Sitter string compactifications,” *JHEP* **08** (2021), 109 [[arXiv:2106.11964](#) [hep-th]].
- [9] X. Gao, A. Hebecker, S. Schreyer and G. Venken, “The LVS parametric tadpole constraint,” *JHEP* **07** (2022), 056 [[arXiv:2202.04087](#) [hep-th]].
- [10] I. Antoniadis, Y. Chen and G. K. Leontaris, “Perturbative moduli stabilization in type IIB/F-theory framework,” *Eur. Phys. J. C* **78** (2018) no.9, 766 [[arXiv:1803.08941](#) [hep-th]].
- [11] I. Antoniadis, Y. Chen and G. K. Leontaris, “Logarithmic loop corrections, moduli stabilization and de Sitter vacua in string theory,” *JHEP* **01**, 149 (2020) [[arXiv:1909.10525](#) [hep-th]].
- [12] V. Basiouris and G. K. Leontaris, “Note on de Sitter vacua from perturbative and non-perturbative dynamics in type IIB/F-theory compactifications,” *Phys. Lett. B* **810** (2020), 135809 [[arXiv:2007.15423](#) [hep-th]].
- [13] V. Basiouris and G. K. Leontaris, “Remarks on the Effects of Quantum Corrections on Moduli Stabilization and de Sitter Vacua in Type IIB String Theory,” *Fortsch. Phys.* **70** (2022) no.2-3, 2100181 [[arXiv:2109.08421](#) [hep-th]].
- [14] M. Cicoli, J. P. Conlon, A. Maharana, S. Parameswaran, F. Quevedo and I. Zavala, “String cosmology: From the early universe to today,” *Phys. Rept.* **1059** (2024), 1-155 [[arXiv:2303.04819](#) [hep-th]].

- [15] G. K. Leontaris and P. Shukla, “Seeking de Sitter vacua in the string landscape,” PoS **CORFU2022** (2023), 058 [[arXiv:2303.16689](#) [hep-th]].
- [16] C. P. Burgess, R. Kallosh and F. Quevedo, “De Sitter string vacua from supersymmetric D terms,” JHEP **10**, 056 (2003) [[arXiv:hep-th/0309187](#) [hep-th]].
- [17] I. Antoniadis, O. Lacombe and G. K. Leontaris, “Hybrid inflation and waterfall field in string theory from D7-branes,” JHEP **01** (2022), 011 [[arXiv:2109.03243](#) [hep-th]].
- [18] I. Antoniadis, O. Lacombe and G. K. Leontaris, “Type IIB moduli stabilization, inflation, and waterfall fields,” Int. J. Mod. Phys. A **37** (2022) no.34, 2244001
- [19] A. D. Linde, “Hybrid inflation,” Phys. Rev. D **49** (1994), 748-754 [[arXiv:astro-ph/9307002](#)].
G. Lazarides and C. Panagiotakopoulos, “Smooth hybrid inflation,” Phys. Rev. D **52** (1995), R559-R563 [[arXiv:hep-ph/9506325](#) [hep-ph]].
- [20] S. Gukov, C. Vafa and E. Witten, “CFT’s from Calabi-Yau four folds,” Nucl. Phys. B **584** (2000), 69-108 [[arXiv:hep-th/9906070](#) [hep-th]].
- [21] I. Antoniadis, Y. Chen and G. K. Leontaris, “Moduli stabilization and inflation in type IIB/F-theory,” PoS **CORFU2018** (2019), 068 [[arXiv:1901.05075](#) [hep-th]].
- [22] G. K. Leontaris and P. Shukla, “Stabilising all Kähler moduli in perturbative LVS,” JHEP **07** (2022), 047 [[arXiv:2203.03362](#) [hep-th]].
- [23] M. Kreuzer and H. Skarke, “Complete classification of reflexive polyhedra in four-dimensions,” Adv. Theor. Math. Phys. **4** (2000), 1209-1230 [[arXiv:hep-th/0002240](#) [hep-th]].
- [24] R. Altman, J. Gray, Y. H. He, V. Jejjala and B. D. Nelson, “A Calabi-Yau Database: Threefolds Constructed from the Kreuzer-Skarke List,” JHEP **02** (2015), 158 [[arXiv:1411.1418](#) [hep-th]].
- [25] R. Blumenhagen, J. P. Conlon, S. Krippendorff, S. Moster and F. Quevedo, “SUSY Breaking in Local String/F-Theory Models,” JHEP **09** (2009), 007 doi:10.1088/1126-6708/2009/09/007 [[arXiv:0906.3297](#) [hep-th]].
- [26] L. Aparicio, D. G. Cerdeno and L. E. Ibanez, “Modulus-dominated SUSY-breaking soft terms in F-theory and their test at LHC,” JHEP **07** (2008), 099 doi:10.1088/1126-6708/2008/07/099 [[arXiv:0805.2943](#) [hep-ph]].
- [27] D. Lüst, S. Reffert and S. Stieberger, “Flux-induced soft supersymmetry breaking in chiral type IIB orientifolds with D3 / D7-branes,” Nucl. Phys. B **706** (2005), 3-52 [[arXiv:hep-th/0406092](#) [hep-th]].
- [28] K. Becker, M. Becker, M. Haack and J. Louis, “Supersymmetry breaking and alpha-prime corrections to flux induced potentials,” JHEP **06**, 060 (2002) [[arXiv:hep-th/0204254](#) [hep-th]].
- [29] A. Hebecker, P. Mangat, F. Rompineve and L. T. Witkowski, “Dark Radiation predictions from general Large Volume Scenarios,” JHEP **09** (2014), 140 doi:10.1007/JHEP09(2014)140 [[arXiv:1403.6810](#) [hep-ph]].

- [30] J. P. Conlon, D. Cremades and F. Quevedo, “Kahler potentials of chiral matter fields for Calabi-Yau string compactifications,” JHEP **01** (2007), 022 [[arXiv:hep-th/0609180](#) [hep-th]].
- [31] M. Haack, D. Krefl, D. Lüst, A. Van Proeyen and M. Zagermann, “Gaugino Condensates and D-terms from D7-branes,” JHEP **01** (2007), 078 [[arXiv:hep-th/0609211](#) [hep-th]].
- [32] S. Kachru, R. Kallosh, A. Linde and S. P. Trivedi, “de Sitter vacua in string theory,” Phys. Rev. D **68**, 046005 (2003) [[arXiv:hep-th/0301240](#) [hep-th]].
- [33] A. Achucarro, B. de Carlos, J. A. Casas and L. Doplicher, “De Sitter vacua from uplifting D-terms in effective supergravities from realistic strings,” JHEP **06** (2006), 014 [[arXiv:hep-th/0601190](#) [hep-th]].
- [34] I. Antoniadis, O. Lacombe and G. K. Leontaris, “Inflation near a metastable de Sitter vacuum from moduli stabilization,” Eur. Phys. J. C **80**, no.11, 1014 (2020) [[arXiv:2007.10362](#) [hep-th]].
- [35] A. Brignole, L. E. Ibanez and C. Munoz, “Towards a theory of soft terms for the supersymmetric Standard Model,” Nucl. Phys. B **422**, 125-171 (1994) [erratum: Nucl. Phys. B **436**, 747-748 (1995)] [[arXiv:hep-ph/9308271](#) [hep-ph]].
- [36] J. P. Conlon, S. S. Abdussalam, F. Quevedo and K. Suruliz, “Soft SUSY Breaking Terms for Chiral Matter in IIB String Compactifications,” JHEP **01**, 032 (2007) [[arXiv:hep-th/0610129](#) [hep-th]].
- [37] M. Demirtas, M. Kim, L. McAllister and J. Moritz, “Conifold Vacua with Small Flux Superpotential,” Fortsch. Phys. **68** (2020), 2000085 [[arXiv:2009.03312](#) [hep-th]].
- [38] S.R. Coleman and E.J. Weinberg, “Radiative Corrections as the Origin of Spontaneous Symmetry Breaking,” Phys. Rev. D **7**, 1888 (1973) [[arXiv:hep-ph/](#)].
- [39] Y. Akrami *et al.* [Planck], “Planck 2018 results. X. Constraints on inflation,” Astron. Astrophys. **641** (2020), A10 [[arXiv:1807.06211](#) [astro-ph.CO]].
- [40] M. Cicoli, C. P. Burgess and F. Quevedo, “Anisotropic Modulus Stabilisation: Strings at LHC Scales with Micron-sized Extra Dimensions,” JHEP **10**, 119 (2011) [[arXiv:1105.2107](#) [hep-th]].
- [41] G. F. Giudice and A. Masiero, “A Natural Solution to the μ Problem in Supergravity Theories,” Phys. Lett. B **206** (1988), 480-484
- [42] S. Angus, J. P. Conlon, U. Haisch and A. J. Powell, “Loop corrections to ΔN_{eff} in large volume models,” JHEP **12** (2013), 061 [[arXiv:1305.4128](#) [hep-ph]].
- [43] M. Cicoli, K. Sinha and R. Wiley Deal, “The dark universe after reheating in string inflation,” JHEP **12** (2022), 068 [[arXiv:2208.01017](#) [hep-th]].