L_p -SOLVABILITY AND HÖLDER REGULARITY FOR STOCHASTIC TIME FRACTIONAL BURGERS' EQUATIONS DRIVEN BY MULTIPLICATIVE SPACE-TIME WHITE NOISE

BEOM-SEOK HAN

ABSTRACT. We present the L_p -solvability for stochastic time fractional Burgers' equations driven by multiplicative space-time white noise:

$$\partial_t^{\alpha} u = a^{ij} u_{x^i x^j} + b^i u_{x^i} + cu + \bar{b}^i u u_{x^i} + \partial_t^{\beta} \int_0^t \sigma(u) dW_t, \ t > 0; \ u(0, \cdot) = u_0,$$

where $\alpha \in (0,1)$, $\beta < 3\alpha/4 + 1/2$, and $d < 4 - 2(2\beta - 1)_+/\alpha$. The operators ∂_t^{α} and ∂_t^{β} are the Caputo fractional derivatives of order α and β , respectively. The process W_t is an $L_2(\mathbb{R}^d)$ -valued cylindrical Wiener process, and the coefficients a^{ij}, b^i, c and $\sigma(u)$ are random.

In addition to the existence and uniqueness of a solution, we also suggest the Hölder regularity of the solution. For example, for any constant $T < \infty$, small $\varepsilon > 0$, and almost sure $\omega \in \Omega$, we have

$$\sup_{x \in \mathbb{R}^d} |u(\omega, \cdot, x)|_{C^{\left[\frac{\alpha}{2}\left(\left(2-(2\beta-1)_+/\alpha-d/2\right)\wedge 1\right)+\frac{(2\beta-1)_-}{2}\right]\wedge 1-\varepsilon}([0,T])} < \infty$$

and

$$\sup_{t \leq T} |u(\omega, t, \cdot)|_{C^{\left(2-(2\beta-1)_{+}/\alpha - d/2\right) \wedge 1-\varepsilon}(\mathbb{R}^{d})} < \infty.$$

The Hölder regularity of the solution in time changes behavior at $\beta = 1/2$. Furthermore, if $\beta \ge 1/2$, then the Hölder regularity of the solution in time is $\alpha/2$ times the one in space.

1. INTRODUCTION

This article investigates the existence, uniqueness, L_p -regularity, and maximal Hölder regularity of a solution to stochastic time fractional Burgers' equations (STFBEs) driven by space-time white noise. We consider

$$\partial_t^{\alpha} u = Lu + \bar{b}^i u u_{x^i} + \partial_t^{\beta} \int_0^t \sigma(u) dW_t, \quad (\omega, t, x) \in \Omega \times (0, \infty) \times \mathbb{R}^d; \quad u(0, \cdot) = u_0, \quad (1.1)$$

where $\alpha \in (0,1)$, $\beta < \frac{3}{4}\alpha + \frac{1}{2}$, and $d < 4 - \frac{2(2\beta-1)_+}{\alpha}$. The operators ∂_t^{α} and ∂_t^{β} are the Caputo fractional derivatives of order α and β , and the operator L is the second order *random* differential operator defined as follows:

$$(Lu)(\omega, t, x) = a^{ij}(\omega, t, x)u_{x^ix^j} + b^i(\omega, t, x)u_{x^i} + c(\omega, t, x)u.$$

²⁰²⁰ Mathematics Subject Classification. 35R11, 26A33, 60H15, 35R60.

Key words and phrases. Stochastic partial differential equation, Time fractional derivatives, Stochastic Burgers' equation, Time fractional Burgers' equation, Space-time white noise, Hölder regularity.

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. NRF-2021R1C1C2007792) and the BK21 Fostering Outstanding Universities for Research (FOUR) funded by the Ministry of Education (MOE, Korea) and the National Research Foundation of Korea (NRF).

BEOM-SEOK HAN

The random coefficients a^{ij} , b^i , and c are predictable, differentiable (or continuous), and bounded functions. The diffusion coefficient $\sigma(u) = \sigma(\omega, t, x, u)$ is a predictable and measurable function satisfying growth conditions and Lipschitz continuity in u. The detailed conditions on a^{ij} , b^i , c, and σ are described in Assumptions 3.1 and 3.3. The random measure dW_t is induced from an $L_2(\mathbb{R}^d)$ -valued cylindrical Wiener process W_t .

When $\alpha = \beta = 1$ in equation (1.1), the equation is said to be a stochastic Burgers' equation (SBE) of form

$$\partial_t u = Lu + buu_x + \sigma(u)\dot{W}, \quad (\omega, t, x) \in \Omega \times (0, \infty) \times \mathbb{R}; \quad u(0, \cdot) = u_0, \tag{1.2}$$

where \dot{W} is the space-time white noise. Numerous studies have been conducted on the equation (1.2), but we only refer to the reader to [13, 14, 29]. In [13], the author proved the uniqueness, existence, and continuity of a solution to a semilinear equation, including an equation of type (1.2) on the unit interval (0, 1). Additionally, the same properties of a solution on \mathbb{R} were obtained in [14] when the L_2 bounded conditions on $\sigma(u)$ were imposed. In [29], the authors investigated the Hölder regularity and moment estimates of the random field solution to (1.2) with $L = \Delta$ and $\bar{b} = -1$.

In contrast, (deterministic) partial differential equations with Caputo fractional derivatives have been used in many fields, such as electrochemical processes [5, 19], dielectric polarization [33], viscoelastic materials [32], biology [31], and physics [11, 18]. Especially, equation (1.1) with $\alpha \in (0, 1)$ and $\sigma(u) = 0$ is called a time fractional Burgers' equation (TFBE), which describes the propagation of waves through viscous media ([1, 2]). Indeed, various researches have been conducted on numerical analysis for the TFBE (see [3, 9, 10, 20, 30]). From a mathematical standpoint, it is reasonable to wonder whether it is possible to demonstrate the uniqueness and existence of a solution to STFBE (1.1), and also to obtain the Hölder regularity of the solution. To the best of our knowledge, [36] is the only study that answers this question. The authors of [36] demonstrate the existence, uniqueness, and regularity of the *mild* solution to SBEs with fractional derivatives in time and space on a bounded domain $\mathcal{D} \subset \mathbb{R}^d$.

In this paper, we provide the L_p uniqueness, existence, and regularity of a *strong* solution to equation (1.1) with *random* second order differential operator L on the *whole spatial domain* \mathbb{R}^d . Additionally, we achieve the Hölder regularity of the solution in time and space. In detail, if $u(\omega, t, x)$ denotes the solution to equation (1.1), then for any bounded stopping time $\tau \leq T$ and small constant $\varepsilon > 0$, almost surely,

$$\sup_{x \in \mathbb{R}^d} \frac{|u(\omega, \cdot, x)|}{C^{\left[\frac{\alpha}{2}\left((2-(2\beta-1)_+/\alpha-d/2)\wedge 1\right)+\frac{(2\beta-1)_-}{2}\right]\wedge 1-\varepsilon}([0,\tau])}}{\sup_{t \le \tau} |u(\omega, t, \cdot)|_{C^{\left(2-(2\beta-1)_+/\alpha-d/2)\wedge 1-\varepsilon}(\mathbb{R}^d)} < \infty.}$$
(1.3)

where $a_+ = (|a| + a)/2$, $a_- = (|a| - a)/2$, and $C^{\gamma}(\mathcal{D})$ is the Hölder spaces. Observe that the behavior of the Hölder regularity of the solution in time changes from $\beta = 1/2$. For example, if $\beta \geq 1/2$, then the Hölder regularity of the solution in time is $\alpha/2$ times that of the regularity in space. Additionally, we can recover the the Hölder regularity results of SBEs by letting $\alpha, \beta \uparrow 1$. These results are consistent with the well-known results of stochastic heat equations driven by space-time white noise (e.g. [26, Remark 8.7] or [16, Corollary 3.1]). In contrast, if $\beta < 1/2$, the Hölder regularity in time gains additional regularity by as much as $1/2 - \beta$. (Remark 3.11). Several remarks about the proof are made. The proof strategy for the main theorem (Theorem 3.5) is based on [16]. However, some differences exist because since it is not certain that Itô's formula and the maximum principle hold for STFBE (1.1).

Thus, the proof proceeds as follows. As in [16], we focus on proving the uniqueness and existence of the L_p solution in each (large) p > 2, and the main difficulty is to demonstrating the existence of the solutions. Hence, we consider the cut-off form of equation (1.1) to obtain local solutions. Afterward, we construct a global solution candidate u by pasting the local solutions (Lemma 4.3 and Remark 4.4). A uniform L_p bound of u is required to show that our the candidate u is a global solution; thus, we divide the local solution into two parts: the noise-dominating and the nonlinear-dominating part. To estimate the noise-dominating parts, we employ the L_p bound of the diffusion coefficient $\sigma(u)$ (Lemma 4.5). In contrast, to control the nonlinear-dominating part (Lemma 4.8), we employ an inequality similar to the chain rule (Lemma 4.6) and a version of the Grönwall inequality including the Caputo fractional derivatives (Theorem 4.7).

To obtain the maximal Hölder regularity of the solution to equation (1.1), we require two components: the Hölder embedding theorem for the solution space $\mathcal{H}_p^{\gamma}(\tau)$ (Theorem 2.16) and the uniqueness of the solution in p (Theorem 3.9). Indeed, when the L_p existence and uniqueness of a solution are given, we have the Hölder regularity of the solution in each (large) p > 2 by employing the Hölder embedding theorem for the solution space (Theorem 2.16 and Theorem 3.5). The Hölder regularity of the solution becomes larger as a large pis chosen; thus, we have to select p that is as large as possible. Therefore, we require the uniqueness of solutions in p because p varies.

This article is organized as follows. Section 2 introduces the definitions and properties of space-time white noise, fractional calculus, and stochastic Banach spaces. Additionally, we present the Hölder embedding theorem for the solution space $\mathcal{H}_p^{\gamma}(\tau)$. Section 3 states the main results of this article and suggests some remarks. The proof of the main results is presented in Section 4. Next, Section 5 proves the Hölder embedding theorem for the solution space $\mathcal{H}_p^{\gamma}(\tau)$.

We finish this section with an introduction to the notation used in this paper. The sets \mathbb{N} and \mathbb{R} are sets of natural and real numbers, respectively. The set \mathbb{R}^d denotes the *d*-dimensional Euclidean space of points $x = (x^1, \ldots, x^d)$ for $x^i \in \mathbb{R}$. Throughout this paper, we assume Einstein's summation convention on $i, j, k \in \mathbb{N}$. We use := to denote a definition. For a real-valued function f, we set the following:

$$f_+ := \frac{|f| + f}{2}$$
 and $f_- := \frac{|f| - f}{2}$.

For a normed space F, a measure space (X, \mathcal{M}, μ) , and $p \in [1, \infty)$, a space $L_p(X, \mathcal{M}, \mu; F)$ is a set of F-valued \mathcal{M}^{μ} -measurable functions such that

$$||u||_{L_p(X,\mathcal{M},\mu;F)} := \left(\int_X ||u(x)||_F^p \mu(dx)\right)^{1/p} < \infty$$

A set \mathcal{M}^{μ} is the completion of \mathcal{M} with respect to the measure μ . For $\gamma \in (0,1]$ and $k = 0, 1, 2, \ldots$, a set $C^{k+\gamma}(\mathbb{R}^d)$ is the set of \mathbb{R} -valued continuous functions u = u(x) such that

$$|u|_{C^{\gamma+k}(\mathbb{R}^d)} := \sup_{x \in \mathbb{R}^d, |\boldsymbol{\beta}|=k} \left| D^{\boldsymbol{\beta}} u(x) \right| + \sup_{\substack{x, y \in \mathbb{R}^d, x \neq y \\ |\boldsymbol{\beta}|=k}} \frac{\left| D^{\boldsymbol{\beta}} u(x) - D^{\boldsymbol{\beta}} u(y) \right|}{|x - y|^{\gamma}} < \infty,$$

BEOM-SEOK HAN

where β is a multi-index. Similarly, for $\gamma \in (0, 1]$ and $0 < T < \infty$, the set $C^{\gamma}([0, T]; F)$ is the set of *F*-valued continuous functions *u* such that

$$|u|_{C^{\gamma}([0,T];F)} := \sup_{t \in [0,T]} |u(t)|_{F} + \sup_{\substack{t,s \in [0,T], \\ s \neq t}} \frac{|u(t) - u(s)|_{F}}{|t - s|^{\gamma}} < \infty.$$

For $a, b \in \mathbb{R}$, we set $a \wedge b := \min\{a, b\}$ and $a \vee b := \max\{a, b\}$. Let $S = S(\mathbb{R}^d)$ denote the set of Schwartz functions on \mathbb{R}^d . Let $N = N(a_1, a_2, ..., a_k)$ be a generic constant if N depends only on $a_1, a_2, ..., a_k$. The constant N can vary line by line. For functions depending on ω, t , and x, the argument $\omega \in \Omega$ is omitted. Finally, for $x \in \mathbb{R}^d$, $\bar{x}^i := (x^1, \ldots, x^{i-1}, x^{i+1}, \ldots, x^d)$.

2. Preliminaries

In this section, we introduce the definitions and properties of space-time white noise, fractional calculus, and stochastic Banach spaces. Throughout this paper, (Ω, \mathcal{F}, P) is a complete probability space equipped with a filtration $\{\mathcal{F}_t\}_{t\geq 0}$. Let $\{\mathcal{F}_t\}_{t\geq 0}$ denote a filtration satisfying the usual conditions. Let \mathcal{P} be the predictable σ -field related to $\{\mathcal{F}_t\}_{t\geq 0}$.

First, we present the space-time white noise W to understand the stochastic part of (1.1).

Definition 2.1 (Space-time white noise). A generalized random field \dot{W} is said to be the space-time white noise if it is a centered Gaussian random field such that its covariance is given by

$$\mathbb{E}\dot{W}(h)\dot{W}(g) = \int_0^\infty \int_{\mathbb{R}^d} h(t,x)g(t,x)dxdt, \quad \forall h,g \in L_2((0,\infty) \times \mathbb{R}^d).$$

Remark 2.2. We employ a series of Itô's stochastic integral to interpret the stochastic part of equation (1.1). More precisely, let $\{\eta^k : k \in \mathbb{N}\}$ be an orthonormal basis on $L_2(\mathbb{R}^d)$. If we define

$$w_t^k := \int_0^t \int_{\mathbb{R}^d} \eta^k(x) \dot{W}(ds, dx)$$

using the Walsh integral (see [35]), then $\{w_t^k : k \in \mathbb{N}\}$ is a set of one dimensional independent Wiener processes. Then, if we set (see [26, Section 8.3], and [23, Section 7])

$$W_t := \sum_{k=1}^{\infty} \eta^k w_t^k,$$

then W_t is an $L_2(\mathbb{R}^d)$ -valued cylindrical Wiener process and $dW_t = \sum_k \eta^k dw_t^k$. Thus, equation (1.1) can be rewritten as

$$\partial_t^{\alpha} u = Lu + \bar{b}^i u u_{x^i} + \partial_t^{\beta} \int_0^t \sigma(u) \eta^k dw_t^k, \quad (\omega, t, x) \in \Omega \times (0, \infty) \times \mathbb{R}^d; \quad u(0, \cdot) = u_0.$$

Next, we review the facts of fractional calculus. For more information, we refer to the reader to [6, 17, 21, 32].

Definition 2.3. Let $\alpha > 0$, and for $\varphi \in L_1((0,T))$, the Riemann-Liouville fractional integral of the order α is defined as follows:

$$I_t^{\alpha}\varphi(t) := (I_t^{\alpha}\varphi)(t) := \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1}\varphi(s)ds \quad \text{for all} \quad t \in (0,T),$$

where $\Gamma(\alpha) := \int_0^\infty t^{\alpha-1} e^{-t} dt$.

Remark 2.4. For any $q \in [1, \infty]$, by Jensen's inequality

$$\|I^{\alpha}\varphi\|_{L_{q}((0,T))} \leq N(\alpha, p, T)\|\varphi\|_{L_{q}((0,T))}.$$
(2.1)

Therefore, $I_t^{\alpha}\varphi(t)$ is well-defined and finite for almost all $t \leq T$. Additionally, Fubini's theorem implies that, for $\alpha, \beta \geq 0$, we have

$$I^{\alpha+\beta}\varphi(t) = I^{\alpha}I^{\beta}\varphi(t).$$

Definition 2.5. For $\alpha > 0$, let $n \in \mathbb{N}$ be a nonnegative integer such that $n - 1 \leq \alpha < n$. Suppose $\varphi(t)$ is a real-valued function on [0,T] such that φ is (n-1)-times differentiable and $(\frac{d}{dt})^{n-1}\varphi$ is absolutely continuous on [0,T].

(i) The Riemann-Liouville fractional derivative $D_t^{\alpha}\varphi$ is defined as

$$D_t^{\alpha}\varphi(t) := \frac{1}{\Gamma(n-\alpha)} \frac{d^n}{dt^n} \int_0^t (t-s)^{n-\alpha-1} \varphi(s) ds.$$

(ii) The Caputo fractional derivative $\partial_t^{\alpha} \varphi$ is defined as

$$\partial_t^{\alpha} \varphi := \frac{1}{\Gamma(n-\alpha)} \int_0^t (t-s)^{n-\alpha-1} \varphi^{(n)}(s) ds$$
$$:= \frac{1}{\Gamma(n-\alpha)} \frac{d}{dt} \int_0^t (t-s)^{n-\alpha-1} \left[\varphi^{(n-1)}(s) - \varphi^{(n-1)}(0) \right] ds$$

Remark 2.6. (i) For any $\alpha, \beta \geq 0$, $D_t^{\alpha} D_t^{\beta} \varphi = D_t^{\alpha+\beta} \varphi$ and

$$D_t^{\alpha} I_t^{\beta} \varphi = D_t^{\alpha - \beta} \varphi \mathbf{1}_{\alpha > \beta} + I_t^{\beta - \alpha} \varphi \mathbf{1}_{\alpha \le \beta}.$$

Additionally, if $\alpha \in (0,1)$, $I_t^{1-\alpha}\varphi$ is absolutely continuous, and $I_t^{1-\alpha}\varphi(0) = 0$, then the following equality holds:

$$I_t^{\alpha} D_t^{\alpha} \varphi(t) = \varphi(t).$$

(ii) By the definition of fractional derivatives, if $\varphi(0) = \varphi^{(1)}(0) = \cdots = \varphi^{(n-1)}(0) = 0$, then $D_t^{\alpha} \varphi = \partial_t^{\alpha} \varphi$.

Below we recall the definitions and properties of stochastic Banach spaces (for more detail, see [12, 25, 26, 27]). The solution space $\mathcal{H}_p^{\gamma}(T)$ and embedding theorems for $\mathcal{H}_p^{\gamma}(T)$ are suggested.

Definition 2.7. Let p > 1 and $\gamma \in \mathbb{R}$. The space $H_p^{\gamma} = H_p^{\gamma}(\mathbb{R}^d)$ is the set of all tempered distributions u on \mathbb{R} such that

$$\|u\|_{H_p^{\gamma}} := \left\| (1-\Delta)^{\gamma/2} u \right\|_{L_p} = \left\| \mathcal{F}^{-1} \left[(1+|\xi|^2)^{\gamma/2} \mathcal{F}(u)(\xi) \right] \right\|_{L_p} < \infty.$$

Similarly, $H_p^{\gamma}(l_2) = H_p^{\gamma}(\mathbb{R}^d; l_2)$ is a space of l_2 -valued functions $g = (g^1, g^2, \cdots)$ such that

$$\|g\|_{H_p^{\gamma}(l_2)} := \left\| \left| (1-\Delta)^{\gamma/2} g \right|_{l_2} \right\|_{L_p} = \left\| \left| \mathcal{F}^{-1} \left[(1+|\xi|^2)^{\gamma/2} \mathcal{F}(g)(\xi) \right] \right|_{l_2} \right\|_{L_p} < \infty.$$

Remark 2.8. Let $d \in \mathbb{N}$ and $\gamma \in (0, \infty)$. A nonnegative smooth function $R_{\gamma}(x)$ exists on \mathbb{R}^d such that, for $u \in C_c^{\infty}(\mathbb{R}^d)$,

$$\left((1-\Delta)^{-\gamma/2}u\right)(x) = \int_{\mathbb{R}^d} R_{\gamma}(y)u(x-y)dy$$

and

$$|R_{\gamma}(x)| \le NA_{\gamma,d}(x)\mathbf{1}_{|x|\le 2} + Ne^{-|x|/2}\mathbf{1}_{|x|\ge 2},$$

where $N = N(\gamma, d)$ is a positive constant and

$$A_{\gamma,d}(x) = \begin{cases} |x|^{\gamma-d} + 1 + O(|x|^{\gamma-d+2}) & \text{for } 0 < \gamma < d, \\ \log(2/|x|) + 1 + O(|x|^2) & \text{for } \gamma = d, \\ 1 + O(|x|^{\gamma-d}) & \text{for } \gamma > d. \end{cases}$$

For more detail, see [12, Proposition 1.2.5].

We introduce the space of point-wise multipliers in H_p^{γ} .

Definition 2.9. Fix $\gamma \in \mathbb{R}$ and $\alpha \in [0,1)$ such that $\alpha = 0$ if $\gamma \in \mathbb{Z}$ and $\alpha > 0$ if $|\gamma| + \alpha$ is not an integer. Define

$$B^{|\gamma|+\alpha} = \begin{cases} B(\mathbb{R}) & \text{if } \gamma = 0, \\ C^{|\gamma|-1,1}(\mathbb{R}) & \text{if } \gamma \text{ is a nonzero integer}, \\ C^{|\gamma|+\alpha}(\mathbb{R}) & \text{otherwise}, \end{cases}$$
$$B^{|\gamma|+\alpha}(\ell_2) = \begin{cases} B(\mathbb{R}, \ell_2) & \text{if } \gamma = 0, \\ C^{|\gamma|-1,1}(\mathbb{R}, \ell_2) & \text{if } \gamma \text{ is a nonzero integer}, \\ C^{|\gamma|+\alpha}(\mathbb{R}, \ell_2) & \text{otherwise}, \end{cases}$$

where $B(\mathbb{R})$ is the space of bounded Borel functions on \mathbb{R} , $C^{|\gamma|-1,1}(\mathbb{R})$ represents the space of $|\gamma| - 1$ times continuous differentiable functions whose derivatives of the $(|\gamma| - 1)$ th order derivative are Lipschitz continuous, and $C^{|\gamma|+\alpha}$ is the real-valued Hölder spaces. The space $B(\ell_2)$ denotes a function space with ℓ_2 -valued functions instead of real-valued function spaces.

Below we collect the properties of Bessel potential spaces.

Lemma 2.10. Let $\gamma \in \mathbb{R}$ and p > 1.

- (i) The space $C_c^{\infty}(\mathbb{R}^d)$ is dense in H_p^{γ} .
- (ii) Let $\gamma d/p = n + \nu$ for some $n = 0, 1, \cdots$ and $\nu \in (0, 1]$. Then, for any $k \in$ $\{0, 1, \dots, n\}, we have$

$$|D^{k}u|_{C(\mathbb{R}^{d})} + |D^{n}u|_{\mathcal{C}^{\nu}(\mathbb{R}^{d})} \le N ||u||_{H_{p}^{\gamma}},$$
(2.2)

where $\mathcal{C}^{\nu}(\mathbb{R}^d)$ is the Zygmund space. (iii) The operator $D_i: H_p^{\gamma} \to H_p^{\gamma+1}$ is bounded. Moreover, for any $u \in H_p^{\gamma+1}$,

$$\left\|D^{i}u\right\|_{H_{p}^{\gamma}} \leq N\|u\|_{H_{p}^{\gamma+1}}$$

where $N = N(\gamma, p)$.

(iv) For $\gamma_1, \gamma_2 \in \mathbb{R}$, and $u \in H_p^{\gamma_1 + \gamma_2}$, we have

$$\|\Delta^{\gamma_1/2} u\|_{H_p^{\gamma_2}} \le N \|u\|_{H_p^{\gamma_1+\gamma_2}},$$

where $N = N(\gamma_1, \gamma_2)$.

(v) For $\gamma \in (0, 1)$, and $u \in H_p^{\gamma}$, we have

$$\|(1 - \Delta^{\gamma})u\|_{L_p} \le N\left(\|u\|_{L_p} + \|(-\Delta)^{\gamma}u\|_{L_p}\right),$$

where $N = N(\gamma, p)$.

(vi) For any $\mu, \gamma \in \mathbb{R}$, the operator $(1 - \Delta)^{\mu/2} : H_p^{\gamma} \to H_p^{\gamma-\mu}$ is an isometry.

(vii) Let

$$\varepsilon \in [0,1], \quad p_i \in (1,\infty), \quad \gamma_i \in \mathbb{R}, \quad i = 0, 1,$$

$$\gamma = \varepsilon \gamma_1 + (1-\varepsilon)\gamma_0, \quad 1/p = \varepsilon/p_1 + (1-\varepsilon)/p_0.$$

Then, we have

$$||u||_{H_p^{\gamma}} \le ||u||_{H_{p_1}^{\gamma_1}}^{\varepsilon} ||u||_{H_{p_0}^{\gamma_0}}^{1-\varepsilon}.$$

(viii) Let $u \in H_p^{\gamma}$. Then, we have

$$\|au\|_{H_{p}^{\gamma}} \leq N \|a\|_{B^{|\gamma|+\alpha}} \|u\|_{H_{p}^{\gamma}} \quad and \quad \|bu\|_{H_{p}^{\gamma}(\ell_{2})} \leq N \|b\|_{B^{|\gamma|+\alpha}(\ell_{2})} \|u\|_{H_{p}^{\gamma}},$$

where $N = N(\gamma, p)$ and $B^{|\gamma|+\alpha}, B^{|\gamma|+\alpha}(\ell_2)$ are introduced in Definition 2.9.

Proof. The above results are well known. For (i), (iii), (vi), and (vii), see Theorems 13.3.7 (i), 13.8.1, 13.3.7 (ii), and Exercise 13.3.20 of [27], respectively. In the case of (ii) and (iv), see [34]. For (v), see Theorems 1.3.6 and 1.3.8 of [12]. For (viii), we refer the reader to [26, Lemma 5.2].

Definition 2.11 (Stochastic Banach spaces). Let $\tau \leq T$ be a bounded stopping time, $p \geq 2$, and $\gamma \in \mathbb{R}$. Set $(0, \tau] := \{(\omega, t) : 0 < t \leq \tau(\omega)\}$ and define

$$\begin{aligned} \mathbb{H}_{p}^{\gamma}(\tau) &:= L_{p}\left(\left(0, \tau\right]\right), \mathcal{P}, dP \times dt; H_{p}^{\gamma}\right), \\ \mathbb{H}_{p}^{\gamma}(\tau, l_{2}) &:= L_{p}\left(\left(0, \tau\right]\right), \mathcal{P}, dP \times dt; H_{p}^{\gamma}(l_{2})\right), \\ U_{p}^{\gamma} &:= L_{p}\left(\Omega, \mathcal{F}_{0}, H_{p}^{\gamma}\right). \end{aligned}$$

We write $u \in \mathbb{H}_p^{\gamma}$ if $u \in \mathbb{H}_p^{\gamma}(\tau)$ exists for any bounded stopping time τ . Additionally, if $\gamma = 0$, then we use \mathbb{L} instead of \mathbb{H} , $\|f\|_{\mathbb{L}_p(\tau)} := \|f\|_{\mathbb{H}_p^0(\tau)}$. The norm of each space is defined naturally, for example,

$$||f||_{\mathbb{H}_{p}^{\gamma}(\tau)} := \left(\mathbb{E}\int_{0}^{\tau} ||f(t,\cdot)||_{H_{p}^{\gamma}}^{p} dt\right)^{1/p}.$$

Lemma 2.12 exhibits the relation between the stochastic and fractional integrals, which is employed when I_t^{α} or D_t^{α} is applied to the stochastic part of the SPDEs.

Lemma 2.12. Let $T < \infty$ be a constant.

(i) Let $\alpha \geq 0$ and $h \in L_2(\Omega \times [0,T], \mathcal{P}; l_2)$. Then, the equality

$$I^{\alpha}\left(\sum_{k=1}^{\infty}\int_{0}^{\cdot}h^{k}(s)dw_{s}^{k}\right)(t) = \sum_{k=1}^{\infty}\left(I^{\alpha}\int_{0}^{\cdot}h^{k}(s)dw_{s}^{k}\right)(t)$$

holds for all $t \leq T$ almost surely and in $L_2(\Omega \times [0,T])$, where the series on both sides converge in probability.

(ii) If $\alpha \geq 0$ and $h_n \to h$ in $L_2(\Omega \times [0,T], \mathcal{P}; l_2)$ as $n \to \infty$, then

$$\sum_{k=1}^{\infty} \left(I^{\alpha} \int_{0}^{\cdot} h_{n}^{k} dw_{s}^{k} \right)(t) \to \sum_{k=1}^{\infty} \left(I^{\alpha} \int_{0}^{\cdot} h^{k} dw_{s}^{k} \right)(t)$$

in probability uniformly on [0, T].

(iii) If $\alpha > 1/2$ and $h \in L_2(\Omega \times [0,T], \mathcal{P}; l_2)$, then $\left(I^{\alpha} \sum_{k=1}^{\infty} \int_0^{\cdot} h^k(s) dw_s^k\right)(t)$ is differentiable in t and

$$\frac{\partial}{\partial t} \left(I^{\alpha} \sum_{k=1}^{\infty} \int_{0}^{\cdot} h^{k}(s) dw_{s}^{k} \right)(t) = \frac{1}{\Gamma(\alpha)} \sum_{k=1}^{\infty} \int_{0}^{t} (t-s)^{\alpha-1} h^{k}(s) dw_{s}^{k}$$
(a.e.) on $\Omega \times [0,T]$.

Proof. See Lemmas 3.1 and 3.3 of [7].

Fix a small $\kappa_0 > 0$. For $\alpha \in (0, 1)$ and $\beta < \alpha + 1/2$, set

$$c_0 := \frac{(2\beta - 1)_+}{\alpha} + \kappa_0 1_{\beta = 1/2}.$$
(2.3)

Next, we introduce the solution spaces (for more detail, see Definitions 2.9 and 2.12 in [25]).

Definition 2.13 (Solution spaces). Let $\tau \leq T$ be a bounded stopping time, $\alpha \in (0,1)$, $\beta < \alpha + 1/2, \gamma \in \mathbb{R}$, and $p \geq 2$.

(i) For $u \in \mathbb{H}_p^{\gamma}(\tau)$, we write $u \in \mathcal{H}_p^{\gamma}(\tau)$ if $u_0 \in U_p^{\gamma}$, $f \in \mathbb{H}_p^{\gamma-2}(\tau)$, and $g \in \mathbb{H}_p^{\gamma-2+c_0}(\tau, l_2)$ such that

$$\partial_t^{\alpha} u(t,x) = f(t,x) + \partial_t^{\beta} \int_0^t g^k(s,x) dw_s^k, \quad 0 < t \le \tau; \quad u(0,\cdot) = u_0$$

in the sense of distribution. In other words, for any $\phi \in S$, the equality

$$(u(t,\cdot),\phi) = (u_0,\phi) + I_t^{\alpha}(f,\phi) + I_t^{\alpha-\beta} \sum_{k=1}^{\infty} \int_0^t (g^k(s,\cdot),\phi) dw_s^k$$
(2.4)

holds for a.e. $(\omega, t) \in \Omega \times [0, \tau]$. If $\alpha - \beta \in (-1/2, 0)$, we regard $I_t^{\alpha - \beta}$ as $\frac{\partial}{\partial t} I_t^{\alpha - \beta + 1}$. The norm in $\mathcal{H}_p^{\gamma}(\tau)$ is defined as follows:

$$\|u\|_{\mathcal{H}_{p}^{\gamma}(\tau)} := \|u\|_{\mathbb{H}_{p}^{\gamma}(\tau)} + \|u_{0}\|_{U_{p}^{\alpha,\gamma}} + \inf_{f,g} \left\{ \|f\|_{\mathbb{H}_{p}^{\gamma-2}(\tau)} + \|g\|_{\mathbb{H}_{p}^{\gamma-2+c_{0}}(\tau,l_{2})} \right\}.$$
 (2.5)

(ii) We say $u \in \mathcal{H}_{p,loc}^{\gamma}(\tau)$ if there exists a sequence $\tau_n \uparrow \tau$ such that $u \in \mathcal{H}_p^{\gamma}(\tau_n)$ for each n. We write u = v in $\mathcal{H}_{p,loc}^{\gamma}(\tau)$ if a sequence of bounded stopping times $\tau_n \uparrow \tau$ exists such that u = v in $\mathcal{H}_p^{\gamma}(\tau_n)$ for each n. We omit τ if $\tau = \infty$. In other words, $\mathcal{H}_{p,loc}^{\gamma} = \mathcal{H}_{p,loc}^{\gamma}(\infty)$.

Remark 2.14. If $\alpha - \beta \ge 0$, the stochastic part of (2.4) is considered

$$I_t^{\alpha-\beta}\sum_{k=1}^{\infty}\int_0^t (g^k(s,\cdot),\phi)dw_s^k = \sum_{k=1}^{\infty}I_t^{\alpha-\beta}\int_0^t (g^k(s,\cdot),\phi)dw_s^k.$$

Otherwise, if $\alpha - \beta \in (-1/2, 0)$, we regard $I_t^{\alpha - \beta}$ as $\frac{\partial}{\partial t}I_t^{\alpha - \beta + 1}$. Then, by Lemma 2.12 (iii), the stochastic part of (2.4) is

$$\begin{split} I_t^{\alpha-\beta}\left(\sum_{k=1}^{\infty}\int_0^t (g^k(s,\cdot),\phi)dw_s^k\right) &= \frac{\partial}{\partial t}\left(I^{\alpha-\beta+1}\sum_{k=1}^{\infty}\int_0^t (g^k(s,\cdot),\phi)dw_s^k\right) \\ &= \frac{1}{\Gamma(\alpha-\beta+1)}\sum_{k=1}^{\infty}\int_0^t (t-s)^{\alpha-\beta+1}(g^k(s,\cdot),\phi)dw_s^k. \end{split}$$

Below, we provide the properties of the solution space $\mathcal{H}_{p}^{\gamma}(\tau)$.

Theorem 2.15. Let $\tau \leq T$ be a bounded stopping time.

- (i) For $\nu \in \mathbb{R}$, the map $(1 \Delta)^{\nu/2} : \mathcal{H}_p^{\gamma+2}(\tau) \to \mathcal{H}_p^{\gamma-\nu+2}(\tau)$ is an isometry. (ii) If $\gamma \in \mathbb{R}$, $\alpha \in (0,1)$, $\beta < \alpha + 1/2$, and $p \ge 2$, then $\mathcal{H}_p^{\gamma}(\tau)$ is a Banach space with the norm $\|\cdot\|_{\mathcal{H}^{\gamma}_{n}(\tau)}$.

Proof. Despite the fact that the definition of U_p^{γ} is distinct, the proof is a repeat of [25, Theorem 2.14] with τ instead of T.

Next, we suggest the Hölder embedding theorems for $u \in \mathcal{H}_p^{\gamma}(\tau)$. The proof of Theorem 2.16 is contained in Section 5.

Theorem 2.16. Let $\tau \leq T$ be the bounded stopping time, $\gamma \in \mathbb{R}$, $\alpha \in (0,1)$, $\beta < \alpha + 1/2$, and

$$p > 2 \vee \frac{1}{\alpha} \vee \frac{1}{\alpha - \beta + 1/2}.$$
(2.6)

Suppose $u \in \mathcal{H}_p^{\gamma}(\tau)$. Assume α, β, μ , and ν satisfy

$$\frac{1}{\alpha p} < \mu < \frac{(\alpha(\nu + c_0/2) - \beta) \wedge 1/2 + 1/2}{\alpha} \quad and \quad \frac{1}{\alpha p} < \nu < 1 - \frac{c_0}{2}, \tag{2.7}$$

where c_0 is the constant introduced in (2.3). Then, for $u \in C^{\alpha\mu-1/p}([0,\tau]; H_p^{\gamma-2\nu})$ almost surely and

$$\mathbb{E} \|u\|_{C^{\alpha\mu-1/p}\left([0,\tau];H_p^{\gamma-2\nu}\right)}^p \le N \|u\|_{\mathcal{H}_p^{\gamma}(\tau)}^p,\tag{2.8}$$

where $N = N(\alpha, \beta, \gamma, d, p, T)$.

Remark 2.17. Theorem 2.16 is consistent with the previous results ([26, Theorem 7.2]). In other words, if we let $\alpha, \beta \uparrow 1$ in Theorem 2.16, conditions (2.6) and (2.7), and the result (2.8) approach those of the case of $\alpha = \beta = 1$.

By combining Lemma 2.10 (ii) and Theorem 2.16, we have the Hölder embedding results of solution space $\mathcal{H}_p^{(2-c_0-d/2)\wedge 1}(\tau)$ which is a preparation to obtain the maximum Hölder regularity of solutions.

Corollary 2.18. Let $\tau \leq T$ be a bounded stopping time, $\alpha \in (0,1)$, $\beta < \alpha + 1/2$, and $0 < \gamma < (2 - c_0 - d/2) \wedge 1$, where c_0 is introduced in (2.3). Suppose p satisfies (2.6) and $u \in \mathcal{H}_p^{\gamma}(\tau)$. If $\alpha, \beta, \gamma, \nu, d$, and p satisfy

$$\frac{1}{\alpha p} < \mu < \frac{(\alpha(\nu + c_0/2) - \beta) \wedge 1/2 + 1/2}{\alpha} \quad and \quad \frac{1}{\alpha p} < \nu < \left(\frac{\gamma}{2} - \frac{d}{2p}\right) \wedge \left(1 - \frac{c_0}{2}\right), \quad (2.9)$$

then $u \in C^{\alpha\mu-1/p}([0,\tau]; C^{\gamma-2\nu-d/p})$ almost surely and

$$\mathbb{E} \|u\|_{C^{\alpha\mu-1/p}\left([0,\tau];C^{\gamma-2\nu-d/p}(\mathbb{R}^d)\right)}^p \le N \|u\|_{\mathcal{H}_p^{\gamma}(\tau)}^p,$$

where $N = N(\alpha, \beta, \gamma, d, p, T)$.

Proof. For the proof, we employ Lemma 2.10 (ii) and Theorem 2.16. Then, we have

$$\mathbb{E} \|u\|_{C^{\alpha\mu-1/p}([0,\tau];C^{\gamma-2\nu-d/p}(\mathbb{R}^d))}^p \le \mathbb{E} \|u\|_{C^{\alpha\mu-1/p}([0,\tau];H_p^{\gamma-2\nu}(\mathbb{R}^d))}^p \le N \|u\|_{\mathcal{H}_p^{\gamma}(\tau)}^p.$$

Thus, the corollary is proved.

BEOM-SEOK HAN

3. Main Results

This section presents the uniqueness, existence, L_p -regularity, and Hölder regularity of the solution to the following equation:

$$\partial_t^{\alpha} u = Lu + \bar{b}^i u u_{x^i} + \partial_t^{\beta} \sum_k \int_0^t \sigma(u) \eta^k dw_s^k, \quad t > 0; \quad u(0, \cdot) = u_0, \tag{3.1}$$

where $Lu = a^{ij}u_{x^ix^j} + b^iu_{x^i} + cu$. The coefficients a^{ij} , b^i , and c are $\mathcal{P} \times \mathcal{B}(\mathbb{R}^d)$ -measurable, \bar{b}^i is $\mathcal{P} \times \mathcal{B}(\mathbb{R}^{d-1})$ -measurable, and a^{ij} , b^i , c, and \bar{b}^i (and their derivatives) are uniformly bounded (see Assumption 3.1). Additionally, we assume the coefficient \bar{b}^i is independent of x^i . Indeed, because \bar{b}^i is independent of x^i , we can employ the fundamental theorem of calculus to control the nonlinear term $\bar{b}^i u u_{x^i}$ (see Remark 3.2). Moreover, the diffusion coefficient $\sigma(u)$ is dominated by an \mathbb{L}_p function h (see Assumption 3.3) and it is used to obtain a uniform L_p bound of the local solutions (see Remark 3.4).

In Theorem 3.5, we obtain the existence and uniqueness of a solution in \mathcal{H}_p^{γ} , where $\gamma \in (0, 2 - c_0 - d/2) \wedge 1$. The components of equation (3.1) affect the properties of the solution u. For example, if α , β , d, and p are given, the regularity γ is determined. Remarks 3.6, 3.7, and 3.8 provide explanations for these relations.

Additionally, we have the maximal Hölder regularity of the solution in Corollary 3.10. We employ the Hölder embedding theorem for solution spaces (Corollary 2.18). Furthermore, depending on the range of β , the behavior of the Hölder regularity of the solution in time varies. In detail, when $\beta \geq 1/2$, then the Hölder regularity of the solution in space is $\alpha/2$ times of the Hölder regularity of the solution in time. Moreover, if we consider the case $\alpha, \beta \uparrow 1$, then the Hölder regularity in time and space approaches 1/4 and 1/2, which are the results of the SPDEs driven by space-time white noise (e.g. [26, Remark 8.7] or [16, Corollary 3.1]). In the case of $\beta < 1/2, 1/2 - \beta$ of the Hölder regularity in time is obtained due to the regularity of the stochastic integral (Remark 3.11).

The following are assumptions on coefficients.

Assumption 3.1. (i) The coefficients $a^{ij} = a^{ij}(t,x)$, $b^i = b^i(t,x)$, and c = c(t,x) are $\mathcal{P} \times \mathcal{B}(\mathbb{R}^d)$ -measurable.

(ii) The coefficient $\bar{b}^i(t, \bar{x}^i) = \bar{b}^i(t, x^1, \dots, x^{i-1}, x^{i+1}, \dots, x^d)$ is $\mathcal{P} \times \mathcal{B}(\mathbb{R}^{d-1})$ -measurable. (iii) There exists K > 0 such that

$$K^{-1}|\xi|^2 \le a^{ij}(t,x)\xi^i\xi^j \le K|\xi|^2 \quad for \ all \quad (\omega,t,x) \in \Omega \times [0,\infty) \times \mathbb{R}^d, \quad \xi \in \mathbb{R}^d, \tag{3.2}$$

Ì

$$\sum_{i,j} \left| a^{ij}(t,\cdot) \right|_{C^2(\mathbb{R}^d)} + \sum_i \left| b^i(t,\cdot) \right|_{C^2(\mathbb{R}^d)} + |c(t,\cdot)|_{C^2(\mathbb{R}^d)} + \sum_i \left| \bar{b}^i(t,\cdot) \right|_{C^2(\mathbb{R}^{d-1})} \le K \quad (3.3)$$

for all $(\omega,t) \in \Omega \times [0,\infty)$.

Remark 3.2. To prove the existence of a global solution, we need to acquire a uniform L_p bound of the local solutions. Thus, we separate the local solutions into two parts: noise-dominating and nonlinear-dominating parts. In this remark, we consider the nonlinear-dominating parts related to $\bar{b}^i u u_{x^i}$.

If coefficient \bar{b}^i is independent of x^i , coefficient \bar{b}^i can be taken out of the integral for x^i . Then, by the fundamental theorem of calculus to x^i , the nonlinear term $\bar{b}^i u u_{x^i}$ is eliminated in the L_p estimate of the nonlinear-dominating part of the local solutions. Thus, the

nonlinear-dominating parts are controlled by the initial data and diffusion coefficient $\sigma(u)$ (for more information, see Lemma 4.8).

To introduce the assumptions on the diffusion coefficient, we may assume $p \geq 2$.

Assumption 3.3 (p). (i) The coefficient $\sigma(t, x, u)$ is $\mathcal{P} \times \mathcal{B}(\mathbb{R}^d) \times \mathcal{B}(\mathbb{R})$ -measurable. (ii) There exists a constant K such that

$$|\sigma(t, x, u) - \sigma(t, x, v)| \le K|u - v| \quad for \ all \quad (\omega, t, x) \in \Omega \times [0, \infty) \times \mathbb{R}^d, \quad u, v \in \mathbb{R}.$$

(iii) There exists a $\mathcal{P} \times \mathcal{B}(\mathbb{R}^d)$ -measurable function $h \in \mathbb{L}_p$ such that

$$|\sigma(t,x,u)| \le |h(t,x)| \quad for \ all \quad (\omega,t,x) \in \Omega \times [0,\infty) \times \mathbb{R}^d, \quad u \in \mathbb{R}.$$
(3.4)

Remark 3.4. As mentioned in Remark 3.2, we divide the local solutions into two parts, and the nonlinear-dominating parts are controlled by the initial data u_0 and diffusion coefficients $\sigma(u)$. Then, to deal with the noise-dominating term and the terms including $\sigma(u)$, we employ the function h(t, x) introduced in Assumption 3.3 (p) (iii). Indeed, the terms related to the diffusion coefficient $\sigma(u)$ are controlled by the initial data and h so that a uniform L_p bound of u is obtained (see Lemmas 4.5 and 4.8).

Next, we introduce the main results.

Theorem 3.5. Let

$$\alpha \in (0,1), \quad \beta < \frac{3}{4}\alpha + \frac{1}{2}, \quad d < 4 - 2c_0, \quad 0 < \gamma < (2 - c_0 - d/2) \land 1$$
 (3.5)

and

$$p = 2^k \quad for \ some \quad k \in \mathbb{N} \quad and \quad p > 2 \lor \frac{1}{\alpha} \lor \frac{1}{\alpha - \beta + 1/2} \lor \frac{2 + \alpha d}{\alpha \gamma} \lor \frac{d}{1 - \gamma}, \tag{3.6}$$

where c_0 are the constants introduced in (2.3). Suppose Assumptions 3.1 and 3.3 (p) hold. If $u_0 \in U_p^{\gamma}$, then there exists a unique solution $u \in \mathcal{H}_{p,loc}^{\gamma}$ satisfying (3.1). Furthermore, for μ and ν satisfying (2.9), and bounded stopping time $\tau \leq T$, we have

$$u \in C^{\alpha \mu - 1/p}([0, \tau]; C^{\gamma - 2\nu - d/p}) \quad and \quad \|u\|_{C^{\alpha \mu - \frac{1}{p}}([0, \tau]; C^{\gamma - 2\nu - d/p})} < \infty$$
(3.7)

almost surely.

Proof. See **Proof of Theorem 3.5** in Section 4.

Remark 3.6. (i) We assume

 $\alpha \in (0,1)$

because an inequality acting like the chain rule is employed to deal with the nonlineardominating part of the local solution (see Lemma 4.6).

(ii) The conditions

$$\beta < 3\alpha/4 + 1/2$$
 and $d < 4 - 2c_0$

are expected to obtain the uniqueness and existence of solutions to SPDEs with Caputo time fractional derivatives and space-time white noise even for the semilinear case. For example, see [23, Section 7]. Additionally, observe that the choice of α and β allows d = 1, 2, 3, where c_0 is the constant introduced in (2.3).

Remark 3.7. (i) For the existence and uniqueness of local solutions, we impose

$$\gamma \in (0, 2 - c_0 - d/2). \tag{3.8}$$

Heuristically, if u is a measurable, continuous, and bounded solution to equation (3.1), then for given $T < \infty$, we can define a bounded stopping time as follows:

$$\tau_m := \inf\left\{t \ge 0 : \sup_{x \in \mathbb{R}^d} |u(t, x)| \ge m\right\} \wedge T.$$

Then, the solution u satisfies the localized version of equation (3.1) on $(0, \tau_m)$. In other words,

$$\partial_t^{\alpha} u = Lu + \frac{1}{2} \bar{b}^i \left((|u| \wedge m)^2 \right)_{x^i} + \partial_t^{\beta} \sum_k \int_0^t \sigma(u) \eta^k dw_s^k$$
(3.9)

holds on $0 < t < \tau_m$ with $u(0, \cdot) = u_0$. Then, as (3.9) is a semilinear equation, (3.8) has to be satisfied by [23, Theorem 7.1] (for more detail, see [23, Section 7] and [25, Section 5].

(ii) The following condition

$$\gamma \in (0,1) \tag{3.10}$$

is assumed due to the nonlinear term $\overline{b}^i u u_{x^i}$ lowering the regularity of the solution. Even for SBEs ($\alpha = \beta = 1$), the condition in (3.10) is required (for more information, see [13, 14, 15, 16, 29]).

Remark 3.8. (i) To obtain the local solution, we employ the L_p theory for the semilinear equation (see [26, Theorem 5.1]). When we control the nonlinear term $\bar{b}^i u u_{x^i}$ in the L_p estimate, the kernel of $(1 - \Delta)^{-\frac{\gamma-1}{2}}$ has to be controlled. Hence,

$$p > \frac{d}{1 - \gamma}$$

is imposed (see Lemma 4.3).

(ii) We require \mathbb{R} -valued continuous solutions to consider the cut-off version of equation (3.1). Therefore, we assume

$$p > 2 \wedge \frac{1}{\alpha} \wedge \frac{1}{\alpha - \beta + 1/2} \wedge \frac{2 + \alpha d}{\alpha \gamma}$$

which is required to apply the Hölder embedding theorem for \mathcal{H}_p^{γ} (see Theorem 2.16 and Corollary 2.18).

 (iii) As mentioned in Remark 3.6 (i), we employ an inequality similar to the chain rule. To apply (4.13) instead of chain rule for the Caputo fractional derivative, we assume

 $p = 2^k$

for some $k \in \mathbb{N}$.

To achieve the maximal Hölder regularity, we require the uniqueness of the solution in p.

Theorem 3.9. Suppose all the conditions of Theorem 3.5 hold. Let $u \in \mathcal{H}_{p,loc}^{\gamma}$ be the solution of equation (3.1) introduced in Theorem 3.5. If q > p, $u_0 \in U_q^{\alpha,\gamma}$, and Assumption 3.3 (q) hold, then $u \in \mathcal{H}_{q,loc}^{\gamma}$.

Proof. See **Proof of Theorem 3.9** in Section 4.

Finally, we obtain the maximal Hölder regularity of the solution by combining Theorems 3.5 and 3.9. Recall that c_0 is introduced in (2.3).

Corollary 3.10. Suppose α , β , d, and γ satisfy (3.5), and $u_0 \in \bigcap_{p>2} U_p^{(2-c_0-d/2)\wedge 1}$, and $h \in \bigcap_{p>2} \mathbb{L}_p$ satisfies (3.4). Then, for any bounded stopping time $\tau \leq T$, (1.3) holds almost surely.

Proof. When α , β , d, and γ are given in (3.5), we choose p as in (3.6). For each p, there exists a unique solution $u_p \in \mathcal{H}_{p,loc}^{\gamma}$ to equation (3.1). Due to Theorem 3.9, $u_p \in \mathcal{H}_{q,loc}^{\gamma}$ for any $q \geq p$ so that we write u instead of u_p and u is independent of p. Thus, by letting p large in (3.7), we have (1.3). Thus, the corollary is proved.

- **Remark 3.11.** (i) If $1/2 \leq \beta < \alpha + 1/2$, the Hölder regularity in space is $\alpha/2$ times that in time. Furthermore, we can recover the Hölder regularity results of SBEs ($\alpha = \beta = 1$) by considering the case $\alpha, \beta \uparrow 1$. We cite [29, Proposition 5.1] or [16, Corollary 3.1] for reader's convenience.
- (ii) If $\beta < 1/2$, then the Hölder regularity in time obtains additional regularity by as much as $1/2 - \beta$. This phenomenon is caused by the stochastic integral of equation (3.1) adding the Hölder regularity of noise in time almost 1/2, and ∂_t^β reducing the regularity of the noise by β .

4. Proof of Theorems 3.5 and 3.9

We assume that all conditions in Theorem 3.5 hold for the remainder of this section.

To establish the existence of a global solution, we need to obtain the uniqueness and existence of local solutions (Lemma 4.3). With these local solutions, we build a candidate for a global solution. More precisely, we paste the local solutions and demonstrate that the local existence time explodes almost surely (Lemma 4.9). To prove that the local existence time explodes almost surely, we demonstrate that a uniform L_p bound of local solutions exists. In detail, we separate the local solution into noise- and nonlinear-dominating parts. The noise-dominating part is affected by the stochastic part of the equation, and the other part is influenced by the nonlinear term $b^i u u_{x^i}$. When we deal with the noise-dominating part of the solution, the dominating function of the diffusion coefficient provides a uniform L_p bound for the noise-dominating part of the local solutions (see Assumption 3.3 (p) (iii) and Lemma 4.5). The other part is controlled by employing a version of the chain rule and Grönwall inequality (see Lemmas 4.6 and 4.8 and Theorem 4.7).

First, we introduce the uniqueness and existence theorem for semilinear SPDEs.

Assumption 4.1 (τ). (*i*) The functions f(t, x, u) and $g^k(t, x, u)$ are $\mathcal{P} \times \mathcal{B}(\mathbb{R}^d) \times \mathcal{B}(\mathbb{R})$ measurable functions satisfying the following:

$$f(t,x,0) \in \mathbb{H}_p^{\gamma}(\tau)$$
 and $g(t,x,0) = (g^1(t,x,0), g^2(t,x,0), \dots) \in \mathbb{H}_p^{\gamma+1}(\tau, l_2).$

(ii) For any $\varepsilon > 0$, there exists a constant N_{ε} such that for any $u, v \in \mathbb{H}_p^{\gamma}(\tau)$,

$$\|f(u) - f(v)\|_{\mathbb{H}_{p}^{\gamma-2}(\tau)}^{p} + \|g(u) - g(v)\|_{\mathbb{H}_{p}^{\gamma-2+c_{0}}(\tau,l_{2})}^{p} \leq \varepsilon \|u - v\|_{\mathbb{H}_{p}^{\gamma}(\tau)}^{p} + N_{\varepsilon}\|u - v\|_{\mathbb{H}_{p}^{\gamma-2}(\tau)}^{p},$$

where c_0 is the constant introduced in (2.3).

Lemma 4.2. Let $\tau \leq T$ be a bounded stopping time. Suppose Assumption 4.1 (τ) hold. Then, for initial data $u_0 \in U_p^{\gamma}$, the following equation:

$$\partial_t^{\alpha} u = Lu + f(u) + \partial_t^{\beta} \int_0^t g^k(u) dw_t^k, \quad 0 < t \le \tau; \quad u(0, \cdot) = u_0$$
(4.1)

has a unique solution $u \in \mathcal{H}_p^{\gamma}(\tau)$. Moreover,

$$\|u\|_{\mathcal{H}_{p}^{\gamma}(\tau)}^{p} \leq N\left(\|u_{0}\|_{U_{p}^{\gamma}}^{p} + \|f(0)\|_{\mathbb{H}_{p}^{\gamma-2}(\tau)}^{p} + \|g(0)\|_{\mathbb{H}_{p}^{\gamma-2+c_{0}}(\tau,l_{2})}^{p}\right),\tag{4.2}$$

where $N = N(\alpha, \beta, \gamma, d, p, K, T)$ and c_0 is the constant introduced in (2.3).

Proof. Theorem 5.1 of [26] is the motivation of the proof. The case $\tau \equiv T$ is obtained by [25, Theorem 2.18]; thus, we only consider the case $\tau \leq T$.

(Step 1). (Existence) Set

$$f(t,u) := 1_{t \le \tau} f(t,u)$$
 and $\bar{g}(t,u) := 1_{t \le \tau} g(t,u)$

Additionally, $\bar{f}(u)$ and $\bar{g}(u)$ satisfy Assumption 4.1 (*T*). Then, by [25, Theorem 2.18], there exists a unique solution $u \in \mathcal{H}_p^{\gamma}(T)$ such that u satisfies equation (4.1) with \bar{f} and \bar{g} , instead of f and g, respectively. As $\tau \leq T$, we have $u \in \mathcal{H}_p^{\gamma}(\tau)$ and u satisfies equation (4.1) and estimate (4.2) with f and g.

(Step 2). (Uniqueness) Let $u, v \in \mathcal{H}_p^{\gamma}(\tau)$ be two solutions of equation (4.1). Then, [25, Theorem 2.18] yields there exists a unique solution $\bar{v} \in \mathcal{H}_p^{\gamma}(T)$ satisfying

$$\partial_t^{\alpha} \bar{v} = L\bar{v} + \bar{f}(v) + \sum_{k=1}^{\infty} \partial_t^{\beta} \int_0^t \bar{g}^k(v) dw_t^k, \quad 0 < t \le T; \quad \bar{v}(0, \cdot) = u_0.$$
(4.3)

Notice that in (4.3), $\bar{f}(v)$ and $\bar{g}(v)$ are used instead of $\bar{f}(\bar{v})$ and $\bar{g}(\bar{v})$, respectively. Set $\tilde{v} := v - \bar{v}$. Then, for fixed $\omega \in \Omega$, we have

$$\partial_t^{\alpha} \tilde{v} = L \tilde{v}, \quad 0 < t \le \tau; \quad \tilde{v}(0, \cdot) = 0.$$

By the deterministic version of [25, Theorem 2.18], we have $\tilde{v} = 0$ in $L_p((0,\tau] \times \mathbb{R}^d)$ almost surely. Additionally, it implies $v(t, \cdot) = \bar{v}(t, \cdot)$ in $L_p((0,\tau] \times \mathbb{R}^d)$ almost surely. Thus, in equation (4.3), we can replace $\bar{f}(v)$ and $\bar{g}(v)$ with $\bar{f}(\bar{v})$ and $\bar{g}(\bar{v})$. Therefore, $\bar{v} \in \mathcal{H}_p^{\gamma}(T)$ satisfies equation (4.1) on (0,T] with \bar{f}, \bar{g} instead of f, g, respectively. Similarly, by following word for word, there exists $\bar{u} \in \mathcal{H}_p^{\gamma}(T)$ such that \bar{u} satisfies equation (4.1) on (0,T] with \bar{f} and \bar{g} instead of f and g. Thus, by the uniqueness result in $\mathcal{H}_p^{\gamma}(T)$, we have $\bar{u} = \bar{v}$ in $\mathcal{H}_p^{\gamma}(T)$, which implies u = v in $\mathcal{H}_p^{\gamma}(\tau)$. Thus, the lemma is proved. \Box

Next, we provide the uniqueness and existence of a local solution to equation (3.1). As an auxiliary function, we choose $\rho(\cdot) \in C_c^{\infty}(\mathbb{R})$ such that $\rho(z) \ge 0$ on $z \in (-\infty, \infty)$, $\rho(z) = 1$ on $|z| \le 1$, $\rho(z) = 0$ on $|z| \ge 2$, and $\frac{d}{dz}\rho(z) \le 0$ on $z \ge 0$. We define the following:

$$\rho_m(z) := \rho(z/m). \tag{4.4}$$

Lemma 4.3. Let $\tau \leq T$ be a bounded stopping time. For $m \in \mathbb{N}$, there exists $u_m \in \mathcal{H}_p^{\gamma}(\tau)$ such that

$$\partial_t^{\alpha} u_m = L u_m + \bar{b}^i \left(u_m^2 \rho_m(u_m) \right)_{x^i} + \partial_t^{\beta} \int_0^t \sigma(t, x, u_m) \eta^k(x) dw_t^k, \ 0 < t \le \tau; \ u_m(0, \cdot) = u_0,$$

where ρ_m is the function introduced in (4.4). Furthermore, $u_m \in C([0,\tau]; C(\mathbb{R}^d))$ almost surely and

$$\mathbb{E}\sup_{t\leq\tau}\sup_{x\in\mathbb{R}^d}|u_m(t,x)|^p\leq N\|u_m\|_{\mathcal{H}_p^{\gamma}(\tau)}^p<\infty$$
(4.5)

almost surely.

Proof. Due to Lemma 4.2 and Corollary 2.18, it suffices to show that Assumption 4.1 (τ) holds. Because $\sigma(t, x, 0) \leq h(t, x)$ for all ω, t, x and $h \in \mathbb{L}_p$, Assumption 4.1 (i) is satisfied.

In the case of Assumption 4.1 (ii), notice that for $u, v \in \mathbb{R}$, we have

$$\left|u^{2}\rho_{m}(u)-v^{2}\rho_{m}(v)\right|\leq N_{m}|u-v|$$

Then, for $u, v \in \mathbb{H}_p^{\gamma}(\tau)$, by Remark 2.8 and Lemmas 2.10 (viii) and (iii), we have

$$\begin{split} &\|\bar{b}^{i}\left((u(t,\cdot))^{2}\rho_{m}(u(t,\cdot))-(v(t,\cdot))^{2}\rho_{m}(v(t,\cdot))\right)_{x^{i}}\|_{H_{p}^{\gamma-2}}^{p} \\ &\leq N\left\|(u(t,\cdot))^{2}\rho_{m}(u(t,\cdot))-(v(t,\cdot))^{2}\rho_{m}(v(t,\cdot))\right\|_{H_{p}^{\gamma-1}}^{p} \\ &\leq N\int_{\mathbb{R}^{d}}\left(\int_{\mathbb{R}^{d}}|R_{1-\gamma}(x-y)|\left((u(\cdot))^{2}\rho_{m}(\cdot,u(\cdot))-(v(\cdot))^{2}\rho_{m}(\cdot,v(\cdot))\right)(t,y)dy\right)^{p}dx \quad (4.6) \\ &\leq N_{m}\left(\int_{\mathbb{R}^{d}}|R_{1-\gamma}(x)|dx\right)^{p}\int_{\mathbb{R}^{d}}|u(t,x)-v(t,x)|^{p}dx \end{split}$$

and

$$\begin{split} \|\sigma(u)\boldsymbol{\eta} - \sigma(v)\boldsymbol{\eta}\|_{H_{p}^{\gamma-2+c_{0}}(l_{2})}^{p} \\ &\leq \int_{\mathbb{R}^{d}} \left(\sum_{k} \left(\int_{\mathbb{R}^{d}} |R_{-\gamma+2-c_{0}}(x-y)| \left(\sigma(\cdot,u(\cdot)) - \sigma(\cdot,v(\cdot))\right)(t,y)\boldsymbol{\eta}^{k}(y)dy \right)^{2} \right)^{p/2} dx \\ &\leq \int_{\mathbb{R}^{d}} \left(\int_{\mathbb{R}^{d}} |R_{-\gamma+2-c_{0}}(x-y)|^{2} \left(\sigma(t,y,u(t,y)) - \sigma(t,y,v(t,y))\right)^{2} dy \right)^{p/2} dx \\ &\leq K^{p} \int_{\mathbb{R}^{d}} \left(\int_{\mathbb{R}^{d}} |R_{-\gamma+2-c_{0}}(y)|^{2} \left(u(t,x-y) - v(t,x-y)\right)^{2} dy \right)^{p/2} dx \\ &\leq K^{p} \left(\int_{\mathbb{R}^{d}} |R_{-\gamma+2-c_{0}}(y)|^{2} dy \right)^{p/2} \int_{\mathbb{R}^{d}} |u(t,x) - v(t,x)|^{p} dx \end{split}$$
(4.7)

on almost every $(\omega, t) \in (0, \tau]$. Due to Remark 2.8, we have

$$\int_{\mathbb{R}^d} |R_{1-\gamma}(y)| \, dy + \int_{\mathbb{R}^d} |R_{-\gamma+2-c_0}(y)|^2 \, dy < \infty.$$

By integrating with respect to (ω, t) to (4.6) and (4.7), employing Lemma 2.10 (vii), and Young's inequality, we have

$$\begin{aligned} \bar{b}^{i} \left(u^{2} \rho_{m}(u) - v^{2} \rho_{m}(v) \right)_{x^{i}} \big\|_{\mathbb{H}^{\gamma-2}_{p}(\tau)}^{p} + \|\sigma(u)\eta - \sigma(v)\eta\|_{\mathbb{H}^{\gamma-2+c_{0}}_{p}(\tau,l_{2})}^{p} \\ &\leq N_{m} \|u - v\|_{\mathbb{L}^{p}(\tau)}^{p} \\ &\leq \varepsilon \|u - v\|_{\mathbb{H}^{\gamma}_{p}(\tau)}^{p} + N_{m} \|u - v\|_{\mathbb{H}^{\gamma-2}_{p}(\tau)}^{p}. \end{aligned}$$

$$(4.8)$$

To obtain the second assertion, we employ Corollary 2.18. The lemma is proved.

BEOM-SEOK HAN

Remark 4.4. We introduce a candidate for a global solution. Let $T < \infty$. For $m \in \mathbb{N}$, let $u_m \in \mathcal{H}_p^{\gamma}(T)$ be the solution introduced in Lemma 4.3. Then, for $R \in \{1, 2, ..., m\}$, define a stopping time τ_m^R

$$\tau_m^R := \inf\left\{t \ge 0 : \sup_{x \in \mathbb{R}} |u_m(t, x)| \ge R\right\} \wedge T.$$
(4.9)

Observe that

$$\tau_R^R \le \tau_m^m \tag{4.10}$$

Indeed, if R = m, (4.10) is obvious. If R < m, we have $u_m \wedge m = u_m \wedge m \wedge R = u_m \wedge R$ for $t \leq \tau_m^R$. Therefore, u_m and u_R are solutions to equation

$$\partial_t^{\alpha} u = Lu + \overline{b}^i \left(u^2 \rho_R(u) \right)_{x^i} + \sigma(u) \eta^k dw_t^k, \quad 0 < t \le \tau_m^R; \quad u(0, \cdot) = u_0.$$

In contrast, $u_R \wedge R = u_R \wedge R \wedge m = u_R \wedge m$ for $t \leq \tau_R^R$. Thus, u_m and u_R are solutions to equation

$$\partial_t^{\alpha} u = Lu + \bar{b} \left(u^2 \rho_m(u) \right)_{x^i} + \sigma(u) \eta^k dw_t^k, \quad 0 < t \le \tau_R^R; \quad u(0, \cdot) = u_0.$$

Observe that the uniqueness and continuity results in Lemma 4.3 yields that $u_m = u_R$ for all $t \leq (\tau_m^R \vee \tau_R^R)$. Therefore, for $t \leq \tau_m^R$,

$$\sup_{s \le t} \sup_{x \in \mathbb{R}} |u_R(s, x)| = \sup_{s \le t} \sup_{x \in \mathbb{R}} |u_m(s, x)| \le R,$$

and this implies $\tau_m^R \leq \tau_R^R$. Similarly, $\tau_m^R \geq \tau_R^R$; thus,

$$\tau_R^R = \tau_m^R$$

almost surely. Moreover, we have $\tau_m^R \leq \tau_m^m$ because m > R. Therefore, we have (4.10). Further, we define

$$u(t,x) := u_m(t,x) \quad on \quad t \le \tau_m^m$$

and set

$$\tau_{\infty} := \limsup_{m \to \infty} \limsup_{T \to \infty} \tau_m^m. \tag{4.11}$$

It should be remarked that u(t,x) is well-defined on $\Omega \times [0,\infty) \times \mathbb{R}^d$ and the nontrivial domain of u is $\Omega \times [0,\tau_{\infty}) \times \mathbb{R}^d$.

To obtain a uniform L_p bound of the local solution u_m , we separate u_m into noise- and nonlinear-dominating parts. Lemma 4.5 provides the existence, uniqueness, and estimate of noise-dominating parts of u_m .

Lemma 4.5. Let $T < \infty$. Then, there exists $v \in \mathcal{H}_p^{\gamma}(T)$ such that

$$\partial_t^{\alpha} v = Lv + \partial_t^{\beta} \int_0^t \sigma(s, x, u) \eta^k(x) dw_s^k, \quad 0 < t \le T, \quad u(0, \cdot) = u_0$$

Furthermore, $v \in C([0,T]; C(\mathbb{R}^d))$ almost surely, and

$$\mathbb{E}\sup_{t \le T} \sup_{x \in \mathbb{R}^d} |v(t,x)|^p + \mathbb{E}\sup_{t \le T} ||v(t,\cdot)||_{L_p}^p \le N ||v||_{\mathcal{H}_p^{\gamma}(T)}^p \le N ||u_0||_{U_p^{\gamma}}^p + N ||h||_{\mathbb{L}_p(\tau)}^p,$$

where $N = N(\alpha, \beta, \gamma, d, p, K, T)$.

Proof. Similar to the proof of Lemma 4.3, it is enough to show that Assumption 4.1 (τ) holds. Set $\eta = (\eta^1, \eta^2, ...)$. Then, by Remark 2.8, for $t \leq T$

$$\begin{aligned} &\|\sigma(t,\cdot,u(t,\cdot))\boldsymbol{\eta}\|_{H_{p}^{\gamma-2+c_{0}}(l_{2})}^{p} \\ &= \int_{\mathbb{R}^{d}} \left(\sum_{k=1}^{\infty} \left(\int_{\mathbb{R}^{d}} R_{-\gamma+2-c_{0}}(x-y)\sigma(t,y,u(t,y))\eta^{k}(y)dy \right)^{2} \right)^{p/2} dx \\ &= \left(\int_{\mathbb{R}^{d}} |R_{-\gamma+2-c_{0}}(x)|^{2} dx \right)^{p/2} \int_{\mathbb{R}^{d}} |\sigma(t,y,u(t,y))|^{p} dy \\ &\leq \left(\int_{\mathbb{R}^{d}} |R_{-\gamma+2-c_{0}}(x)|^{2} dx \right)^{p/2} \int_{\mathbb{R}^{d}} |h(t,y)|^{p} dy \\ &\leq N \|h(t,\cdot)\|_{L_{p}}^{p}. \end{aligned}$$
(4.12)

Therefore,

$$\|\sigma(u)\boldsymbol{\eta}\|_{\mathbb{H}_{p}^{\gamma-2+c_{0}}(T,l_{2})}^{p} \leq \mathbb{E}\int_{0}^{T}\|\sigma(t,\cdot,u(t,\cdot))\boldsymbol{\eta}\|_{H_{p}^{\gamma-2+c_{0}}(l_{2})}^{p}dt \leq N\|h\|_{\mathbb{L}_{p}}^{p}.$$

Thus, the lemma is proved by Lemma 4.2.

Next, we control the nonlinear-dominating parts of the local solutions. The following two lemmas are crucial in obtaining uniform L_p bounds. Lemma 4.6 functions as a chain rule, and Theorem 4.7 is a version of the Grönwall inequality.

Lemma 4.6. Suppose $\alpha \in (0,1)$ and $k \in \mathbb{N}$. For any $\psi \in C_c^{\infty}((0,\infty) \times \mathbb{R}^d)$, we have

$$\partial_t^{\alpha}(\psi(\cdot,x))^{2^k}(t) \le 2^k \psi(t,x) |\psi(t,x)|^{2^k-2} \partial_t^{\alpha} \psi(t,x), \tag{4.13}$$

for all $(t, x) \in (0, \infty) \times \mathbb{R}^d$.

Proof. We employ the mathematical induction. The results and proof are motivated by (4.2) of [8].

(Step 1). First, we consider the case k = 1. Although the proof is in the proof of [8, Proposition 4.1], we include the proof for the completeness of this paper.

Let $\psi \in C_c^{\infty}((0,\infty) \times \mathbb{R}^d)$ and $t \in (0,\infty)$ and $x \in \mathbb{R}^d$. For $s \in (0,t]$, set

$$F_1(s) := \frac{1}{2} |\psi(s, x)|^2, \quad F_2(s) := \psi(s, x)\psi(t, x),$$

and

$$F(s) := \frac{1}{2} \left(|\psi(s,x)|^2 - |\psi(t,x)|^2 \right) - (\psi(s,x) - \psi(t,x))\psi(t,x) + \frac{1}{2} \left(|\psi(s,x)|^2 - |\psi(t,x)|^2 \right) - \frac{1}{2} \left(|\psi(s,x)|^2 - |\psi(t,x)|^2 \right) + \frac{1}{2} \left(|\psi(t,x)|^2 - |\psi(t,x)|^2 \right) + \frac{1}{2} \left($$

Further,

$$F(s) = \frac{1}{2} |\psi(s, x) - \psi(t, x)|^2 \ge 0$$

on $s \leq t$, and the equality holds for s = t. Notice that the integration by parts implies that

$$\int_0^t (t-s)^{-\alpha} (F_1'(s) - F_2'(s)) ds = \int_0^t (t-s)^{-\alpha} F'(s) ds \le 0.$$

Then, by the definition of ∂_t^{α} (Definition 2.5), we have (4.13) with k = 1.

(Step 2). Let $n \in \mathbb{N}$ and assume that the results hold for $k = 1, 2, \ldots, n-1$. Set $\tilde{\psi}(t,x) := (\psi(t,x))^2$. Since $\tilde{\psi}(t,x) \in C_c^{\infty}((0,\infty) \times \mathbb{R}^d)$, we have

$$\begin{aligned} \partial_t^{\alpha}(\psi(\cdot,x))^{2^n}(t) &= \partial_t^{\alpha}(\tilde{\psi}(\cdot,x))^{2^{n-1}}(t) \\ &\leq 2^{n-1}\tilde{\psi}(t,x) \left| \tilde{\psi}(t,x) \right|^{2^{n-1}-2} \partial_t^{\alpha} \tilde{\psi}(t,x) \\ &= 2^{n-1} \left| \psi(t,x) \right|^{2^n-2} \partial_t^{\alpha}(\psi(t,x))^2 \\ &\leq 2^n \psi(t,x) \left| \psi(t,x) \right|^{2^n-2} \partial_t^{\alpha} \psi(t,x). \end{aligned}$$

The lemma is proved.

Theorem 4.7 (Theorem 8 of [4]). Let $\psi(t)$ be a nonnegative integrable function on [0,T]. For a constant N_1 , if the function ψ satisfies

$$\psi(t) \le \psi_0 + N_1 I_t^{\alpha} \psi$$

on $t \in [0, T]$, then

$$\psi(t) \le \left(1 + \sum_{k=0}^{\infty} \frac{N_1^k}{\Gamma(k\alpha)} \frac{(\Gamma(\alpha)t^{\alpha})^k}{k\alpha}\right) \psi_0$$

on $t \in [0, T]$.

We consider following lemma to control the remainder of the local solution u_m .

Lemma 4.8. Let $u_m \in \mathcal{H}_p^{\gamma}(T)$ and $v \in \mathcal{H}_p^{\gamma}(T)$ be functions introduced in Lemmas 4.3 and 4.5, and τ_m^m be the stopping time introduced in (4.9). Then,

 $\|u_m(t,\cdot)-v(t,\cdot)\|_{L_n}^p$

$$\leq N \sup_{t \leq T} \sup_{x \in \mathbb{R}^d} |v(t,x)|^p \sup_{t \leq T} \|v(s,\cdot)\|_{L_p}^p \left[1 + \sum_{k=0}^{\infty} \frac{\left(1 + \sup_{s \leq t, x \in \mathbb{R}^d} |v(s,x)|^2\right)^k}{\Gamma(k\alpha)} \frac{(\Gamma(\alpha)T^{\alpha})^k}{k\alpha} \right]$$

for all $t \leq \tau_m^m$ almost surely, where N = N(p, K).

Proof. Set

$$w_m := u_m - v$$
 and $f_m := Lw_m + \overline{b}^i((u_m)^2 \rho_m(u_m))_{x^i}$.

Then, we have $f_m \in \mathbb{H}_p^{\gamma-2}(T)$ since $w_m, u_m \in \mathbb{H}_p^{\gamma}(T)$ and estimates similar to (4.8). Additionally, $\partial_t^{\alpha} w_m = f_m$. Let $(\omega, t) \in (0, \tau_m^m]$. Due to [23, Remark 2.9], there exists $w_m^n \in C_c^{\infty}((0, \infty) \times \mathbb{R}^d)$ such that $w_m^n \to w_m$ in $L_p((0, t); H_p^{\gamma})$, and $\partial_t^{\alpha} w_m^n$ is a Cauchy in $L_p((0, t); H_p^{\gamma-2})$. Define

$$f_m^n := \partial_t^\alpha w_m^n.$$

Moreover, f_m is the limit of f_m^n as $n \to \infty$ in $L_p((0,t); H_p^{\gamma-2})$ (see [23, Remark 2.9]). Choose a nonnegative function $\zeta \in C_c^{\infty}(\mathbb{R}^d)$ with a unit integral and set $\zeta_{\varepsilon}(x) := \varepsilon^{-d} \zeta(x/\varepsilon)$ for $\varepsilon > 0$. For $h \in L_{1,loc}(\mathbb{R}^d)$, set $h^{(\varepsilon)}(x) := \int_{\mathbb{R}^d} h(y)\zeta_{\varepsilon}(x-y)dy$. Next, let $\varepsilon > 0$ and $x \in \mathbb{R}^d$. Since $w_m^{n(\varepsilon)} \in C_c^{\infty}((0,\infty) \times \mathbb{R}^d)$ and $p = 2^k$, Lemma 4.6

yields

$$\frac{1}{p}\partial_t^{\alpha} \left(w_m^{n(\varepsilon)}(\cdot, x) \right)^p(t) \le f_m^{n(\varepsilon)}(t, x) w_m^{n(\varepsilon)}(t, x) \left| w_m^{n(\varepsilon)}(t, x) \right|^{p-2}$$
(4.14)

on $t \in (0, \infty)$. Additionally, as $w_m^n(0, x) = 0$, we have

$$w_m^{n(\varepsilon)}(0,x) = 0.$$
 (4.15)

Thus, if we take stochastic integral I_t^{α} on both sides of (4.14), we have

$$\frac{1}{p} \left| w_m^{n(\varepsilon)}(t,x) \right|^p \le I_t^{\alpha} \left[f_m^{n(\varepsilon)}(\cdot,x) w_m^{n(\varepsilon)}(\cdot,x) \left| w_m^{n(\varepsilon)}(\cdot,x) \right|^{p-2} \right]$$
(4.16)

due to $(w_m^{n(\varepsilon)})^p \in C_c^{\infty}((0,\infty) \times \mathbb{R}^d)$, (4.15), and Remark 2.6. Observe that (2.1) with $q = \infty$ and the Hölder inequality imply that

$$\begin{aligned} \left\| I_{\cdot}^{\alpha} \left[f_{m}^{n(\varepsilon)}(\cdot,x) w_{m}^{n(\varepsilon)}(\cdot,x) \left| w_{m}^{n(\varepsilon)}(\cdot,x) \right|^{p-2} - f_{m}^{(\varepsilon)}(\cdot,x) w_{m}^{(\varepsilon)}(\cdot,x) \left| w_{m}^{(\varepsilon)}(\cdot,x) \right|^{p-2} \right] \right\|_{L_{1}((0,t))} \\ &\leq \int_{0}^{t} \left| f_{m}^{n(\varepsilon)}(s,x) w_{m}^{n(\varepsilon)}(s,x) \left| w_{m}^{n(\varepsilon)}(s,x) \right|^{p-2} - f_{m}^{(\varepsilon)}(s,x) w_{m}^{(\varepsilon)}(s,x) \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-2} \right| ds \\ &\leq N \int_{0}^{t} \left| f_{m}^{n(\varepsilon)}(s,x) - f_{m}^{(\varepsilon)}(s,x) \right| \left| w_{m}^{n(\varepsilon)}(s,x) \right|^{p-1} ds \\ &+ N \int_{0}^{t} \left| f_{m}^{(\varepsilon)}(s,x) \left[w_{m}^{n(\varepsilon)}(s,x) \left| w_{m}^{n(\varepsilon)}(s,x) \right|^{p-2} - w_{m}^{(\varepsilon)}(s,x) \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-2} \right] \right| ds \\ &\leq N \left[A_{n} \left\| w_{m}^{n(\varepsilon)}(\cdot,x) \right\|_{L_{p}(0,t)}^{2} + B_{n}C_{n} \left\| f_{m}^{(\varepsilon)}(\cdot,x) \right\|_{L_{p}(0,t)}^{2} \right] \left\| w_{m}^{n(\varepsilon)}(\cdot,x) \right\|_{L_{p}(0,t)}^{p-3}, \end{aligned}$$

$$(4.17)$$

where

$$A_n = \left\| f_m^{n(\varepsilon)}(\cdot, x) - f_m^{(\varepsilon)}(\cdot, x) \right\|_{L_p(0,t)} \quad B_n = \left\| w_m^{n(\varepsilon)}(\cdot, x) - w_m^{(\varepsilon)}(\cdot, x) \right\|_{L_p(0,t)},$$

and

$$C_n = \left\| w_m^{n(\varepsilon)}(\cdot, x) \right\|_{L_p(0,t)} + \left\| w_m^{(\varepsilon)}(\cdot, x) \right\|_{L_p(0,t)}.$$

Moreover,

$$A_n, B_n \to 0 \quad \text{and} \quad C_n \to 2 \left\| w_m^{(\varepsilon)}(\cdot, x) \right\|_{L_p(0,t)} \quad \text{as} \quad n \to \infty$$
 (4.18)

since $w_m^n \to w_m$ and $f_m^n \to f_m$ in $L_p((0,t); H_p^{\gamma})$. Then, by applying (4.18) to (4.17), we have

$$\left\| I^{\alpha}_{\cdot} \left[f^{n(\varepsilon)}_{m}(\cdot, x) w^{n(\varepsilon)}_{m}(\cdot, x) \left| w^{n(\varepsilon)}_{m}(\cdot, x) \right|^{p-2} - f^{(\varepsilon)}_{m}(\cdot, x) w^{(\varepsilon)}_{m}(\cdot, x) \left| w^{(\varepsilon)}_{m}(\cdot, x) \right|^{p-2} \right] \right\|_{L_{1}((0,t))} \to 0$$

as $n \to \infty$. Therefore, there exists a sequence n_l such that $w_m^{n_l(\varepsilon)}(\cdot, x) \to w_m^{(\varepsilon)}(\cdot, x)$ and $I_{\cdot}^{\alpha} \left[f_m^{n_l(\varepsilon)} w_m^{n_l(\varepsilon)} \left| w_m^{n_l(\varepsilon)} \right|^{p-2} \right] \to I_{\cdot}^{\alpha} \left[f_m^{(\varepsilon)} w_m^{(\varepsilon)} \left| w_m^{(\varepsilon)} \right|^{p-2} \right]$ almost everywhere on [0, t]. Furthermore, the convergence holds everywhere on [0, t] due to the continuity in t. Then, by considering sequence n_l instead of n and letting $l \to \infty$ for (4.16), we have

$$\frac{1}{p} \left| w_m^{(\varepsilon)}(t,x) \right|^p \le I_t^{\alpha} \left[f_m^{(\varepsilon)}(\cdot,x) w_m^{(\varepsilon)}(\cdot,x) \left| w_m^{(\varepsilon)}(\cdot,x) \right|^{p-2} \right].$$

BEOM-SEOK HAN

Since $t \leq \tau_m^m$, $\rho_m(u_m) = 1$. By integrating with respect to x, we have

$$\frac{\Gamma(\alpha)}{p} \int_{\mathbb{R}^d} \left| w_m^{(\varepsilon)}(t,x) \right|^p dx \\
\leq \int_0^t (t-s)^{\alpha-1} \int_{\mathbb{R}^d} (Lw_m(s,\cdot))^{(\varepsilon)}(x) w_m^{(\varepsilon)}(s,x) \left| w_m^{(\varepsilon)}(s,x) \right|^{p-2} dx ds \\
+ \int_0^t (t-s)^{\alpha-1} \int_{\mathbb{R}^d} \left[\bar{b}^i(s,\bar{x}^i) \left(|w_m(s,\cdot) + v(s,\cdot)|^2 \right)_{x^i}^{(\varepsilon)}(x) \right] w_m^{(\varepsilon)}(s,x) \left| w_m^{(\varepsilon)}(s,x) \right|^{p-2} dx ds. \tag{4.19}$$

Furthermore, by integration by parts, we obtain

$$\int_{\mathbb{R}^{d}} \left[(Lw_{m}(s,\cdot))^{(\varepsilon)}(x) + \bar{b}^{i} \left(|w_{m}(s,\cdot) + v(s,\cdot)|^{2} \right)_{x^{i}}^{(\varepsilon)}(x) \right] w_{m}^{(\varepsilon)}(s,x) \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-2} dx
\leq -(p-1) \int_{\mathbb{R}^{d}} \left(a^{ij}w_{m} \right)_{x^{j}}^{(\varepsilon)}(s,x) \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-2} w_{mx^{i}}^{(\varepsilon)}(s,x) dx
+ (p-1) \int_{\mathbb{R}^{d}} \left(\left(2a^{ij}_{x^{j}} - b^{i} \right) w_{m} \right)^{(\varepsilon)}(s,x) \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-2} w_{mx^{i}}^{(\varepsilon)}(s,x) dx
+ \int_{\mathbb{R}^{d}} \left(\left(a^{ij}_{x^{i}x^{j}} - b^{i}_{x^{i}} + c \right) w_{m} \right)^{(\varepsilon)}(s,x) w_{m}^{(\varepsilon)}(s,x) \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-2} dx
- (p-1) \int_{\mathbb{R}^{d}} \bar{b}^{i}(s,\bar{x}^{i}) \left((w_{m}(s,\cdot) + v(s,\cdot))^{2} \right)^{(\varepsilon)}(x) \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-2} w_{mx^{i}}^{(\varepsilon)}(s,x) dx.$$
(4.20)

Additionally, observe that

$$\begin{aligned} \left| \left(a^{ij} w_m \right)_{x^j}^{(\varepsilon)}(s,x) - a^{ij}(s,x) w_{mx^j}^{(\varepsilon)}(s,x) \right| \\ &= \varepsilon^{-1} \left| \int_{\mathbb{R}^d} \left(a^{ij}(s,x-\varepsilon y) - a^{ij}(s,x) \right) w_m(s,x-\varepsilon y) \zeta_{y^j}(y) dy \right| \\ &\leq N(K) \int_{\mathbb{R}^d} |w_m(s,x-\varepsilon y)| |y| |\zeta_y(y)| dy, \end{aligned}$$

$$(4.21)$$

and by (3.2),

$$-\int_{\mathbb{R}^d} a^{ij}(s,x) w_{mx^i}^{(\varepsilon)}(s,x) w_{mx^j}^{(\varepsilon)}(s,x) \left| w_m^{(\varepsilon)}(s,x) \right|^{p-2} dx$$

$$\leq -K^{-1} \int_{\mathbb{R}^d} \left| w_m^{(\varepsilon)}(s,x) \right|^{p-2} \left| w_{mx}^{(\varepsilon)}(s,x) \right|^2 dx.$$
(4.22)

Thus, by combining (4.21) and (4.22)

$$-\int_{\mathbb{R}^{d}} \left(a^{ij}w_{m}\right)_{x^{j}}^{(\varepsilon)}(s,x) \left|w_{m}^{(\varepsilon)}(s,x)\right|^{p-2} w_{mx^{i}}^{(\varepsilon)}(s,x)dx$$

$$= -\int_{\mathbb{R}^{d}} \left|w_{m}^{(\varepsilon)}(s,x)\right|^{p-2} w_{mx^{i}}^{(\varepsilon)}(s,x) \left[\left(aw_{m}\right)_{x^{j}}^{(\varepsilon)}(s,x) - a(s,x)w_{mx^{j}}^{(\varepsilon)}(s,x)\right] dx$$

$$-\int_{\mathbb{R}^{d}} a^{ij}(s,x)w_{mx^{i}}^{(\varepsilon)}(s,x)w_{mx^{j}}^{(\varepsilon)}(s,x) \left|w_{m}^{(\varepsilon)}(s,x)\right|^{p-2} dx$$

$$\leq N\int_{\mathbb{R}^{d}} \left|w_{m}^{(\varepsilon)}(s,x)\right|^{p-2} \left|w_{mx^{i}}^{(\varepsilon)}(s,x)\right|^{2} dx$$

$$-K^{-1}\int_{\mathbb{R}^{d}} \left|w_{m}^{(\varepsilon)}(s,x)\right|^{p-2} \left(\int_{\mathbb{R}^{d}} \left|w_{m}(s,x-\varepsilon y)\right| |y| |\zeta_{y}(y)| dy\right)^{2} dx$$

$$\leq N\int_{\mathbb{R}^{d}} \left|w_{m}^{(\varepsilon)}(s,x)\right|^{p-2} \left(\int_{\mathbb{R}^{d}} \left|w_{m}(s,x-\varepsilon y)\right| |y| |\zeta_{y}(y)| dy\right)^{2} dx$$

$$-\frac{1}{2}K^{-1}\int_{\mathbb{R}^{d}} \left|w_{m}^{(\varepsilon)}(s,x)\right|^{p-2} \left|w_{mx}^{(\varepsilon)}(s,x)\right|^{2} dx,$$

where N = N(K). Moreover,

$$\left| \left(\left(2a_{x^{j}}^{ij} - b^{i} \right) w_{m} \right)^{(\varepsilon)}(s, x) \right| = \left| \int_{\mathbb{R}^{d}} \left(2a_{y^{j}}^{ij}(s, y) - b^{i}(s, y) \right) w_{m}(s, y) \zeta_{\varepsilon}(x - y) dy \right|$$

$$\leq K \int_{\mathbb{R}^{d}} |w_{m}(s, y)| \zeta_{\varepsilon}(x - y) dy$$

$$= K(|w_{m}(s, \cdot)|)^{(\varepsilon)}(x)$$

$$(4.24)$$

and

$$\left| \left(\left(a_{x^i x^j}^{ij} - b_{x^i}^i + c \right) w_m \right)^{(\varepsilon)}(s, x) \right| \le K(|w_m(s, \cdot)|)^{(\varepsilon)}(x).$$

$$(4.25)$$

Thus, by applying Hölder's inequality, (4.23), (4.24), and (4.25) to (4.20), we have

$$\int_{\mathbb{R}^{d}} \left[(Lw_{m}(s,\cdot))^{(\varepsilon)}(x) + \bar{b}^{i} \left(|w_{m}(s,\cdot) + v(s,\cdot)|^{2} \right)_{x^{i}}^{(\varepsilon)}(x) \right] w_{m}^{(\varepsilon)}(s,x) \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-2} dx$$

$$\leq N \int_{\mathbb{R}} \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-2} \left(\int_{\mathbb{R}} |w_{m}(s,x - \varepsilon y)| |y| |\zeta_{y}(y)| dy \right)^{2} dx$$

$$- \frac{p-1}{4K} \sum_{i} \int_{\mathbb{R}} \int_{\mathbb{R}} \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-2} \left| w_{mx^{i}}^{(\varepsilon)}(s,x) \right|^{2} dx$$

$$+ N \sum_{i} \int_{\mathbb{R}^{d}} \left((|w_{m}(s,\cdot)|)^{(\varepsilon)}(x) \right)^{2} \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-2} dx$$

$$+ N \int_{\mathbb{R}^{d}} (|w_{m}(s,\cdot)|)^{(\varepsilon)}(x) \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-1} dx.$$

$$- (p-1) \sum_{i} \int_{\mathbb{R}^{d}} \bar{b}^{i}(s,\bar{x}^{i}) \left((w_{m}(s,\cdot) + v(s,\cdot))^{2} \right)^{(\varepsilon)}(x) \left| w_{m}^{(\varepsilon)}(s,x) \right|^{p-2} w_{mx^{i}}^{(\varepsilon)}(s,x) dx,$$

$$(4.26)$$

where N = N(K). Furthermore, note that

$$\int_{\mathbb{R}^d} \left| w_m^{(\varepsilon)}(s,x) \right|^p w_{mx^i}^{(\varepsilon)}(s,x) dx = 0 \quad \text{for} \quad s \le t.$$
(4.27)

Indeed, take a nonnegative smooth function $\phi \in C_c^{\infty}(\mathbb{R}^d)$ such that $\phi(x) = 1$ on |x| < 1, $\phi(x) = 0$ on |x| > 2, and $\sup_{x \in \mathbb{R}^d} |\phi'(x)| \le 2$. Then, integration by parts yields

$$\begin{split} \int_{\mathbb{R}^d} \left| w_m^{(\varepsilon)}(s,x) \right|^p w_{mx^i}^{(\varepsilon)}(s,x) \phi(x/n) dx \\ &= -p \int_{\mathbb{R}^d} \left| w_m^{(\varepsilon)}(s,x) \right|^p w_{mx^i}^{(\varepsilon)}(s,x) \phi(x/n) dx - \frac{1}{n} \int_{\mathbb{R}^d} \left| w_m^{(\varepsilon)}(s,x) \right|^p w_m^{(\varepsilon)}(s,x) \phi'(x/n) dx. \end{split}$$

Thus, we have

$$\limsup_{n \to \infty} \left| \int_{\mathbb{R}^d} \left| w_m^{(\varepsilon)}(s,x) \right|^p w_{mx^i}^{(\varepsilon)}(s,x) \phi(x/n) dx \right| \le \limsup_{n \to \infty} \frac{2}{n(p+1)} \int_{\mathbb{R}^d} \left| w_m^{(\varepsilon)}(s,x) \right|^{p+1} dx = 0$$
(4.28)

and (4.28) yields (4.27). Then, from the last term of (4.26), by applying (4.27) and the Hölder's inequality, we have

$$\begin{split} \left| \sum_{i} \int_{\mathbb{R}^{d-1}} \bar{b}^{i}(s,\bar{x}^{i}) \int_{\mathbb{R}} \left(|w_{m}(t,\cdot) + v(t,\cdot)|^{2} \right)^{(\varepsilon)}(x) \left| w_{m}^{(\varepsilon)}(t,x) \right|^{p-2} w_{mx^{i}}^{(\varepsilon)}(t,x) dx^{i} d\bar{x}^{i} \right| \\ &\leq N \sum_{i} \int_{\mathbb{R}^{d}} \left| \left(|w_{m}(t,\cdot) + v(t,\cdot)|^{2} \right)^{(\varepsilon)}(x) - \left| w_{m}^{(\varepsilon)}(t,x) \right|^{2} \right| \left| w_{m}^{(\varepsilon)}(t,x) \right|^{p-2} \left| w_{mx^{i}}^{(\varepsilon)}(t,x) \right| dx \\ &\leq N \int_{\mathbb{R}^{d}} \left(\left(|w_{m}(t,\cdot) + v(t,\cdot)|^{2} \right)^{(\varepsilon)}(x) - \left| w_{m}^{(\varepsilon)}(t,x) \right|^{2} \right)^{2} \left| w_{m}^{(\varepsilon)}(t,x) \right|^{p-2} dx \\ &+ \frac{1}{8KN} \sum_{i} \int_{\mathbb{R}^{d}} \left| w_{mx^{i}}^{(\varepsilon)}(t,x) \right|^{2} \left| w_{m}^{(\varepsilon)}(t,x) \right|^{p-2} dx, \end{split}$$

$$(4.29)$$

where N = N(K). Then, by applying (4.26) and (4.29) to (4.19), we have

$$\begin{split} &\int_{\mathbb{R}^d} \left| w_m^{(\varepsilon)}(t,x) \right|^p dx \\ &\leq N I_t^{\alpha} \int_{\mathbb{R}} \left| w_m^{(\varepsilon)}(s,x) \right|^{p-2} \left(\int_{\mathbb{R}} |w_m(s,x-\varepsilon y)| |y| |\zeta_y(y)| dy \right)^2 dx \\ &+ N I_t^{\alpha} \int_{\mathbb{R}^d} \left| (|w_m(s,\cdot)|)^{(\varepsilon)}(x) \right|^2 \left| w_m^{(\varepsilon)}(s,x) \right|^{p-2} + (|w_m(s,\cdot)|)^{(\varepsilon)}(x) \left| w_m^{(\varepsilon)}(s,x) \right|^{p-1} dx \\ &+ N I_t^{\alpha} \int_{\mathbb{R}^d} \left(\left(|w_m(t,\cdot)+v(t,\cdot)|^2 \right)^{(\varepsilon)}(x) - \left| w_m^{(\varepsilon)}(t,x) \right|^2 \right)^2 \left| w_m^{(\varepsilon)}(t,x) \right|^{p-2} dx, \end{split}$$

where N = N(p, K). By letting $\varepsilon \downarrow 0$, we have

$$\begin{split} \|w_{m}(t,\cdot)\|_{L_{p}}^{p} &\leq NI_{t}^{\alpha} \int_{\mathbb{R}} |w_{m}(\cdot,x)|^{p} \, dx + NI_{t}^{\alpha} \int_{\mathbb{R}^{d}} \left(|w_{m}(\cdot,x) + v(\cdot,x)|^{2} - |w_{m}(\cdot,x)|^{2} \right)^{2} |w_{m}(\cdot,x)|^{p-2} \, dx \\ &\leq NI_{t}^{\alpha} \int_{\mathbb{R}} |w_{m}(\cdot,x)|^{p} \, dx + NI_{t}^{\alpha} \int_{\mathbb{R}^{d}} |v(\cdot,x)|^{2} \, |w_{m}(\cdot,x)|^{p} + |v(\cdot,x)|^{4} \, |w_{m}(\cdot,x)|^{p-2} \, dx \\ &\leq N \left(1 + \sup_{s \leq t, x \in \mathbb{R}^{d}} |v(s,x)|^{2} \right) I_{t}^{\alpha} \, \|w_{m}(\cdot,\cdot)\|_{L_{p}}^{p} + N \sup_{s \leq t} \|v(s,\cdot)\|_{L_{2p}}^{2p}. \end{split}$$

for all $t \leq \tau_m^m.$ Then, by Theorem 4.7, we obtain

$$\|w_m(t,\cdot)\|_{L_p}^p \le N \sup_{s \le t} \|v(s,\cdot)\|_{L_{2p}}^{2p} \left[1 + \sum_{k=0}^{\infty} \frac{\left(1 + \sup_{s \le t, x \in \mathbb{R}^d} |v(s,x)|^2\right)^k}{\Gamma(k\alpha)} \frac{(\Gamma(\alpha)T^{\alpha})^k}{k\alpha}\right]$$

for all $t \leq \tau_m^m$. The lemma is proved.

Finally, we demonstrate that the global solution candidate does not explode in a finite time.

Lemma 4.9. For any $T < \infty$, we have

$$\lim_{R \to \infty} P\left(\left\{\omega \in \Omega : \sup_{t \le T, x \in \mathbb{R}^d} |u(t, x)| > R\right\}\right) = 0.$$

Proof. Let v be the function introduced in Lemma 4.5. Define

$$\tau^{1}(S) := \inf \left\{ t \ge 0 : \|v(t, \cdot)\|_{L_{p}} \ge S \right\} \wedge T, \quad \tau^{2}(S) := \inf \left\{ t \ge 0 : \sup_{x \in \mathbb{R}^{d}} |v(t, x)| \ge S \right\} \wedge T,$$

and

$$\tau_m^0(S) := \tau_m^m \wedge \tau^1(S) \wedge \tau^2(S),$$

BEOM-SEOK HAN

where τ_m^m is the stopping time introduced in (4.9). Set $r := \frac{p}{p-1}$. Then, by Lemmas 4.2 and 2.10, (viii), Hölder inequality, and Minkowski inequality, we have

$$\begin{split} \|u_{m}\|_{\mathcal{H}_{p}^{\gamma}(\tau_{m}^{0}(S))}^{p} - N\|u_{0}\|_{U_{p}^{\gamma}}^{p} \\ &\leq N \left\|\sum_{i} \bar{b}^{i}(u_{m}^{2}\rho_{m}(u_{m}))_{x^{i}}\right\|_{\mathbb{H}_{p}^{\gamma-2}(\tau_{m}^{0}(S))}^{p} + N\|\sigma(u_{m})\eta\|_{\mathbb{H}_{p}^{\gamma-2+c_{0}}(\tau_{m}^{0}(S),l_{2})}^{p} \\ &\leq N \|u_{m}^{2}\|_{\mathbb{H}_{p}^{\gamma-1}(\tau_{m}^{0}(S))}^{p} + N\|\sigma(u_{m})\eta\|_{\mathbb{H}_{p}^{\gamma-2+c_{0}}(\tau_{m}^{0}(S),l_{2})}^{p} \\ &\leq N \mathbb{E} \int_{0}^{\tau_{m}^{0}(S)} \int_{\mathbb{R}^{d}} \left|\int_{\mathbb{R}^{d}} R_{1-\gamma}(y)|u_{m}(s,x-y)|^{2}dy\right|^{p} dxds \\ &+ N \mathbb{E} \int_{0}^{\tau_{m}^{0}(S)} \int_{\mathbb{R}^{d}} \left|\int_{\mathbb{R}^{d}} |R_{-\gamma+2-c_{0}}(y)|^{2} |u_{m}(s,x-y)|^{2}dy\right|^{p/2} dxds \\ &\leq N \mathbb{E} \int_{0}^{\tau_{m}^{0}(S)} \int_{\mathbb{R}^{d}} \left|\int_{\mathbb{R}^{d}} |R_{1-\gamma}(y)|^{r} |u_{m}(s,x-y)|^{r}dy\right|^{p/r} dx \int_{\mathbb{R}^{d}} |u_{m}(s,x)|^{p} dxds \\ &+ N \mathbb{E} \int_{0}^{\tau_{m}^{0}(S)} \left(\int_{\mathbb{R}^{d}} |R_{-\gamma+2-c_{0}}(x)|^{2} dx\right)^{p/2} \int_{\mathbb{R}^{d}} |u_{m}(s,x)|^{p} dxds \\ &\leq N_{0} \mathbb{E} \int_{0}^{\tau_{m}^{0}(S)} \left[1 + \int_{\mathbb{R}^{d}} |u_{m}(s,x)|^{p} dx\right] \int_{\mathbb{R}^{d}} |u_{m}(s,x)|^{p} dxds, \end{split}$$

where $N_0 = N(\alpha, \beta, \gamma, d, p, K, T) \left[\left(\int_{\mathbb{R}^d} |R_{1-\gamma}(x)|^r dx \right)^{p/r} + \left(\int_{\mathbb{R}^d} |R_{-\gamma+2-c_0}(x)|^2 dx \right)^{p/2} \right]$. Note that $N_0 < \infty$ due to $r < \frac{d}{d+\gamma-1}$ and Remark 2.8. Then, by Lemma 4.8 and the definitions of $\tau_1(S)$ and $\tau_2(S)$,

$$\begin{split} &\int_{\mathbb{R}^d} |u_m(t,x)|^p dx \\ &\leq N \int_{\mathbb{R}^d} |u_m(t,x) - v(t,x)|^p + |v(t,x)|^p dx \\ &\leq N \sup_{s \leq t} \sup_{x \in \mathbb{R}^d} |v(t,x)| \sup_{s \leq t} \|v(s,\cdot)\|_p^p \left[1 + \sum_{k=0}^{\infty} \frac{\left(1 + \sup_{s \leq t, x \in \mathbb{R}^d} |v(t,x)|^2\right)^k}{\Gamma(k\alpha)} \frac{\left(\Gamma(\alpha)T^{\alpha}\right)^k}{k\alpha} \right] \\ &+ \int_{\mathbb{R}^d} |v(t,x)|^p dx \\ &< N(p,S,K). \end{split}$$

$$(4.31)$$

Therefore, by combining (4.30) and (4.31), we have

$$\|u_m\|_{\mathcal{H}_p^{\gamma}(\tau_m^0(S))}^p \le N + N \|u_0\|_{U_p^{\gamma}}^p, \tag{4.32}$$

where $N = N(\alpha, \beta, \gamma, d, p, S, K, T)$. It should be noted that the right-hand side of (4.32) is independent of m. Therefore, by Chebyshev's inequality and Lemma 4.3, we have

$$P\left(\sup_{t \le \tau_m^0(S)} \sup_{x \in \mathbb{R}^d} |u(t,x)| > R\right) \le \frac{1}{R^p} \mathbb{E} \sup_{t \le \tau_m^0, x \in \mathbb{R}^d} |u(t,x)|^p$$
$$\le \frac{1}{R^p} \mathbb{E} \sup_{t \le \tau_m^0, x \in \mathbb{R}^d} |u_m(t,x)|^p$$
$$\le \frac{1}{R^p} ||u_m||_{\mathcal{H}_p^\gamma(\tau_m^0)}^p$$
$$\le \frac{N}{R^p},$$

where $N = N(u_0, \alpha, \beta, \gamma, d, p, S, K, T)$. In contrast, by Lemma 4.5,

$$P\left(\tau^{1}(S) < T\right) + P\left(\tau^{2}(S) < T\right)$$

$$\leq P\left(\sup_{t \leq T} \|v(t, \cdot)\|_{L_{p}} > S\right) + P\left(\sup_{t \leq T} \sup_{x \in \mathbb{R}^{d}} |v(t, x)| > S\right)$$

$$\leq \frac{1}{S^{p}} \mathbb{E}\sup_{t \leq T} \|v(t, \cdot)\|_{L_{p}}^{p} + \frac{1}{S^{p}} \mathbb{E}\sup_{t \leq T, x \in \mathbb{R}^{d}} |v(t, x)|^{p}$$

$$\leq \frac{1}{S^{p}} N(u_{0}, h, \alpha, \beta, \gamma, d, p, K, T).$$

Thus,

$$\begin{split} &P\left(\sup_{t \leq T, x \in \mathbb{R}^d} |u(t, x)| > R\right) \\ &\leq \liminf_{m \to \infty} P\left(\sup_{t \leq \tau_m^0(S), x \in \mathbb{R}^d} |u(t, x)| > R\right) + P\left(\tau^1(S) < T\right) + P\left(\tau^2(S) < T\right) \\ &\leq \frac{N_1}{R^p} + \frac{N_2}{S^p}, \end{split}$$

where $N_1 = N_1(u_0, \alpha, \beta, \gamma, d, p, S, K, T)$ and $N_2 = N_2(u_0, h, \alpha, \beta, \gamma, d, p, K, T)$. The lemma is proved by letting $R \to \infty$ and $S \to \infty$ in order.

Proof of Theorem 3.5. Step 1. (Uniqueness). Suppose $u, \bar{u} \in \mathcal{H}_{p,loc}^{\gamma}$ are nonnegative solutions of equation (3.1). By Definition 2.13, there are bounded stopping times τ_n $(n = 1, 2, \cdots)$ such that

$$\tau_n \uparrow \infty$$
 and $u, \bar{u} \in \mathcal{H}_p^{\gamma}(\tau_n)$.

Fix $n \in \mathbb{N}$. Note that $u, \bar{u} \in C([0, \tau_n]; C(\mathbb{R}^d))$ almost surely and

$$\mathbb{E}\sup_{t\leq\tau_n}\sup_{x\in\mathbb{R}^d}|u(t,x)|^p + \mathbb{E}\sup_{t\leq\tau_n}\sup_{x\in\mathbb{R}^d}|\bar{u}(t,x)|^p < \infty.$$
(4.33)

Then, for $m \in \mathbb{N}$, define

$$\tau_{m,n}^{1} := \inf \left\{ t \ge 0 : \sup_{x \in \mathbb{R}^{d}} |u(t,x)| > m \right\} \wedge \tau_{n},$$

$$\tau_{m,n}^{2} := \inf \left\{ t \ge 0 : \sup_{x \in \mathbb{R}^{d}} |\bar{u}(t,x)| > m \right\} \wedge \tau_{n},$$

and

$$\tau_{m,n} := \tau_{m,n}^1 \wedge \tau_{m,n}^2. \tag{4.34}$$

Due to (4.33), $\tau_{m,n}^1$ and $\tau_{m,n}^2$ are well-defined stopping times; thus, $\tau_{m,n}$ is a stopping time. Observe that $u, \bar{u} \in \mathcal{H}_p^{\gamma}(\tau_{m,n})$ and $\tau_{m,n} \uparrow \tau_n$ as $m \to \infty$ almost surely. Fix $m \in \mathbb{N}$. Notice that $u, \bar{u} \in \mathcal{H}_p^{\gamma}(\tau_{m,n})$ are solutions to equation

$$\partial_t^{\alpha} v = Lv + \bar{b}^i \left(v^2 \rho_m(v) \right)_{x^i} + \partial_t^{\beta} \int_0^t \sigma(v) dW_t, \quad 0 < t \le \tau_{m,n}; \quad v(0, \cdot) = u_0,$$

where $Lv = a^{ij}v_{x^ix^j} + b^iv_{x^i} + cv$. By the uniqueness result in Lemma 4.3, we conclude that $u = \bar{u}$ in $\mathcal{H}_p^{\gamma}(\tau_{m,n})$ for each $m \in \mathbb{N}$. The monotone convergence theorem yields $u = \bar{u}$ in $\mathcal{H}_p^{\gamma}(\tau_n)$, which implies $u = \bar{u}$ in $\mathcal{H}_{p,loc}^{\gamma}$.

Step 2 (Existence.). Let $T < \infty$. For $m \in \mathbb{N}$, define τ_m^m and u as in Remark 4.4. Observe that

$$P\left(\tau_m^m < T\right) \le P\left(\sup_{t \le T, x \in \mathbb{R}^d} |u(t, x)| \ge m\right).$$

Indeed, if $\tau_m^m < T$, then $\sup_{t \le \tau_m^m, x \in \mathbb{R}^d} |u(t, x)| = \sup_{t \le \tau_m^m, x \in \mathbb{R}^d} |u_m(t, x)| = m$ almost surely. Then, by Lemma 4.9, we have

$$\limsup_{m \to \infty} P\left(\tau_m^m < T\right) \le \limsup_{m \to \infty} P\left(\sup_{t \le T, x \in \mathbb{R}^d} |u(t, x)| \ge m\right) = 0$$

Since $T < \infty$ is arbitrary, $\tau_m^m \to \infty$ in probability. In addition, we conclude that $\tau_m^m \uparrow \infty$ almost surely, because τ_m^m is increasing in m.

Last, set $\tau_m := \tau_m^m \wedge m$. Note that (see Remark 4.4)

$$u(t,x) = u_m(t,x) \quad \text{for} \quad t \in [0,\tau_m].$$

Observe that $\sup_{x \in \mathbb{R}^d} |u_m(t,x)| \leq m$ for $t \in [0, \tau_m]$; thus, u_m satisfies (3.1) almost everywhere $t \in [0, \tau_m]$ almost surely. Because $u = u_m$ for $t \in [0, \tau_m]$ and $u_m \in \mathcal{H}_p^{\gamma}(\tau_m)$, it follows that $u \in \mathcal{H}_p^{\gamma}(\tau_m)$ and u satisfies (3.1) for all $t \leq \tau_m$ almost surely. We have $u \in \mathcal{H}_{p,loc}^{\gamma}$ because $\tau_m \uparrow \infty$ as $m \to \infty$ almost surely.

To obtain Hölder regularity, we employ Corollary 2.18. The theorem is proved.

Proof of Theorem 3.9. The proof of Theorem 3.9 is motivated by [26, Corollarly 5.11]. Since q > p, by Theorem 3.5, there exists a unique solution $\bar{u} \in \mathcal{H}_{q,loc}^{\gamma}$ satisfying equation (3.1). By Definition 2.13, there exists τ_n such that $\tau_n \to \infty$ almost surely as $n \to \infty$, $u \in \mathcal{H}_p^{\gamma}(\tau_n)$ and $\bar{u} \in \mathcal{H}_q^{\gamma}(\tau_n)$. Fix $n \in \mathbb{N}$. Because $\frac{2+\alpha d}{\alpha \gamma} , we can define <math>\tau_{m,n}$ $(m \in \mathbb{N})$ as in (4.34). Notice that for any $p_0 > p$, we have

$$u \in \mathbb{L}_{p_0}(\tau_{m,n})$$

since

$$\mathbb{E}\int_{0}^{\tau_{m,n}}\int_{\mathbb{R}}|u(t,x)|^{p_{0}}dxdt \leq m^{p_{0}-p}\mathbb{E}\int_{0}^{\tau_{m,n}}\int_{\mathbb{R}}|u(t,x)|^{p}dxdt < \infty$$

Observe that $\bar{b}^i u u_{x^i} \in \mathbb{H}_q^{\gamma-2}(\tau_{m,n}) \subset \mathbb{H}_q^{-2}(\tau_{m,n})$. Indeed, similar to (4.6),

$$\mathbb{E}\int_0^{\tau_{m,n}} \left\|\frac{1}{2}\bar{b}^i(s,\cdot)\left((u(s,\cdot))^2\right)_{x^i}\right\|_{H^{\gamma-2}_q}^q ds \le N\mathbb{E}\int_0^{\tau_{m,n}}\int_{\mathbb{R}}|u(s,x)|^{2q}dxds < \infty.$$

Additionally, we have

$$au_{xx} \in \mathbb{H}_q^{-2}(\tau_{m,n}), \quad bu_x \in \mathbb{H}_q^{-1}(\tau_{m,n}), \quad \text{and} \quad cu \in \mathbb{L}_q(\tau_{m,n}).$$

Therefore, because $\mathbb{L}_q(\tau_{m,n}) \subset \mathbb{H}_q^{-1}(\tau_{m,n}) \subset \mathbb{H}_q^{-2}(\tau_{m,n})$,

$$a^{ij}u_{x^{i}x^{j}} + b^{i}u_{x^{i}} + cu + \bar{b}^{i}uu_{x^{i}} \in \mathbb{H}_{q}^{-2}(\tau_{m,n}).$$
(4.35)

Similar to (4.12), we have

$$\|\sigma(u)\boldsymbol{\eta}\|_{\mathbb{H}^{\gamma-2+c_0}_q(\tau_{m,n},l_2)}^q \le N \int_0^{\tau_{m,n}} \|h(t,\cdot)\|_{L_q}^q \, dt < \infty.$$
(4.36)

Thus, we have

$$\sigma(u)\boldsymbol{\eta} \in \mathbb{H}_q^{\gamma-2+c_0}(\tau_{m,n}, l_2) \subset \mathbb{H}_q^{-2+c_0}(\tau_{m,n}, l_2).$$

$$(4.37)$$

Due to (4.35), (4.37), and Lemma 4.2, u is in $\mathcal{L}_q(\tau_{m,n})$ and u satisfies (3.1) for almost everywhere $t \leq \tau_{m,n}$ almost surely. On the other hand, since $\bar{b}^i u u_{x^i} \in \mathbb{H}_q^{\gamma-2}(\tau_{m,n})$ and $\sigma(u)\boldsymbol{\eta} \in \mathbb{H}_q^{\gamma-2+c_0}(\tau_{m,n}, l_2)$, Lemma 4.2 implies that there exists $v \in \mathcal{H}_q^{\gamma}(\tau_{m,n})$ satisfying

$$\partial_t^{\alpha} v = Lv + \bar{b}^i u u_{x^i} + \partial_t^{\alpha} \int_0^t \sigma(u) \eta_k dw_t^k, \quad 0 < t \le \tau_{m,n}; \quad v(0, \cdot) = u_0, \tag{4.38}$$

where $Lv = a^{ij}v_{x^ix^j} + b^iv_{x^i} + cv$. In (4.38), note that $\bar{b}^i u u_{x^i}$ and $\sigma^k(u)$ are used instead of $\bar{b}^i v v_{x^i}$ and $\sigma^k(v)$. Moreover, because $u \in \mathcal{L}_q(\tau_{m,n})$ satisfies equation (4.38), $\bar{v} := u - v \in \mathcal{L}_q(\tau_{m,n})$ satisfies

$$\partial_t^{\alpha} \bar{v} = a^{ij} \bar{v}_{x^i x^j} + b^i \bar{v}_{x^i} + c \bar{v}, \quad 0 < t \le \tau_{m,n}; \quad \bar{v}(0, \cdot) = 0$$

By the deterministic version of Lemma 4.2, we have $\bar{v} = 0$ in $\mathcal{L}_q(\tau_{m,n})$; thus, u = v in $\mathcal{L}_q(\tau_{m,n})$. Therefore, u is in $\mathcal{H}_q^{\gamma}(\tau_{m,n})$. As $\bar{u} \in \mathcal{H}_q^{\gamma}(\tau_{m,n})$ and \bar{u} satisfies equation (3.1), by Lemma 4.3, we have $u = \bar{u}$ in $\mathcal{H}_q^{\gamma}(\tau_{m,n})$. The theorem is proved.

5. Proof of Theorem 2.16

This section provides a proof of the embedding theorem for solution spaces $\mathcal{H}_p^{\gamma}(\tau)$. Consider the following fractional diffusion equation

$$\partial_t^{\alpha} u = \Delta u \quad t > 0; \quad u(0, \cdot) = u_0(\cdot), \tag{5.1}$$

where $\alpha \in (0,1)$ and $u_0(\cdot) \in C_c^{\infty}(\mathbb{R}^d)$. It turns out that a fundamental solution p(t,x) exists such that

$$p(t, \cdot) \in L_1(\mathbb{R}^d)$$
 and $\mathcal{F}(p(t, \cdot))(\xi) = E_\alpha(-t^\alpha |\xi|^2)$

(e.g., [24, Theorem 2.1]), and the solution of (5.1) is given by

$$u(t,x) = (u_0 * p(t,\cdot))(x) = \int_{\mathbb{R}^d} u_0(y)p(t,x-y)dy$$

For convenience, define

$$q_{\alpha,\beta}(t,x) := \begin{cases} I^{\alpha-\beta}p(t,x) & \text{if} \quad \alpha \ge \beta \\ D^{\beta-\alpha}p(t,x) & \text{if} \quad \alpha < \beta \end{cases}$$

We gather some facts related to p(t, x) and $q_{\alpha,\beta}(t, x)$ (for more information, see [22, 23, 24]).

Lemma 5.1. Let $d \in \mathbb{N}$, $\alpha \in (0,1)$, $\beta < \alpha + 1/2$, $\gamma \in [0,2)$, and $\sigma \in \mathbb{R}$.

(i) For all $t \neq 0$ and $x \neq 0$,

$$\partial_t^{\alpha} p(t,x) = \Delta p(t,x) \quad and \quad \partial_t p(t,x) = \Delta q_{\alpha,1}(t,x).$$

Additionally, for each $x \neq 0$, $\frac{\partial}{\partial t}p(t,x) \to 0$ as $t \downarrow 0$. Moreover, $\frac{\partial}{\partial t}p(t,x)$ is integrable in \mathbb{R}^d uniformly on $t \in [\delta, T]$ for any $\delta > 0$.

(ii) There exist constants $c = c(\alpha, d)$ and $N = N(\alpha, d)$ such that if $|x|^2 \ge t^{\alpha}$,

$$|p(t,x)| \le N|x|^{-d} \exp\left\{-c|x|^{\frac{2}{2-\alpha}}t^{-\frac{\alpha}{2-\alpha}}\right\}.$$

(iii) Let $n \in \mathbb{N}$. Then, there exists $N = N(\alpha, \gamma, n)$ such that

$$\left| D_t^{\sigma} D_x^n (-\Delta)^{\gamma/2} q_{\alpha,\beta}(1,x) \right| \le N \left(|x|^{-d+2-\gamma-n} \wedge |x|^{-d-\gamma-n} \right).$$

(iv) The scaling properties hold. In other words,

$$q_{\alpha,\beta}(t,x) = t^{-\frac{\alpha a}{2} + \alpha - \beta} q_{\alpha,\beta}(1,xt^{-\frac{\alpha}{2}}),$$
$$(-\Delta)^{\gamma/2} q_{\alpha,\beta}(t,x) = t^{-\frac{\alpha(d+\gamma)}{2} + \alpha - \beta} (-\Delta)^{\gamma/2} q_{\alpha,\beta}(1,xt^{-\frac{\alpha}{2}})$$

Proof. To see (i), (ii), and (iii) follow from Theorems 2.1 and 2.3 of [23]. For (iv), see (5.2) in [23]. \Box

Remark 5.2. To prove Theorem 2.16, we define the operators. Let $\phi \in C_c^{\infty}(\mathbb{R}^d)$ and $f \in C_c^{\infty}((0,\infty) \times \mathbb{R}^d)$. Take a function $g = (g^1, g^2, ...)$ satisfying the form

$$g^{k}(t,x) = \begin{cases} \sum_{i=1}^{n} 1_{(\tau_{i-1},\tau_{i}]}(t)g^{ik}(x) & \text{for } k = 1, 2, \dots, n, \\ 0 & \text{for } k = n+1, \dots \end{cases}$$
(5.2)

for some $n \in \mathbb{N}$, where τ_i is the bounded stopping time and $g^{ik} \in C_c^{\infty}(\mathbb{R}^d)$. Further, we set

$$T_t^1\phi(x) := \int_{\mathbb{R}^d} p(t, x - y)\phi(y)dy, \qquad (5.3)$$

$$T_t^2 f(t,x) := \int_0^t \int_{\mathbb{R}^d} q_{\alpha,1}(t-r,x-y) f(r,y) dy ds,$$
(5.4)

$$T_t^3 g(t,x) := \sum_k \int_0^t \int_{\mathbb{R}^d} q_{\alpha,\beta}(t-r,x-y) g^k(r,y) dy dw_s^k.$$
 (5.5)

It is well-known that $T_t^1\phi$, T_t^2f , and T_t^3g are solutions to

$$\partial_t^{\alpha} u^1 = \Delta u^1; \quad u^1(0, \cdot) = \phi, \tag{5.6}$$

$$\begin{split} \partial_t^{\alpha} u^2 &= \Delta u^2 + f; \quad u^2(0, \cdot) = 0, \\ \partial_t^{\alpha} u^3 &= \Delta u^3 + \partial_t^{\beta} \int_0^t g^k dw_s^k; \quad u^3(0, \cdot) = 0, \end{split}$$

respectively (for more information, see [22, 23, 24]).

First, we provide a smoothing effect of T_t^1 , which implies the continuity of $T_t^1\phi$ in t.

Lemma 5.3. Let $\tau \leq T$ be a bounded stopping time and $\alpha \in (0,1)$. If $p \in (1,\infty)$, $\theta \in [0,\alpha)$, $\phi \in C_c^{\infty}(\mathbb{R}^d)$, and $t \in (0,T)$, we have

$$e^{-t} \left\| T_t^1 \phi \right\|_{H_p^{\frac{2\theta}{\alpha}}} \le N t^{-\theta} \|\phi\|_{L_p}$$

$$(5.7)$$

where $N = N(\alpha, \theta, d, p, T)$.

Proof. In the case of $\theta = 0$, by Minköwski's inequality, we have

$$\|T_t^1\phi\|_{L_p} \le \|p(t,\cdot)*\phi\|_{L_p} \le \|p(t,\cdot)\|_{L_1}\|\phi\|_{L_p} \le \|\phi\|_{L_p}.$$
(5.8)

Thus, we have (5.7). For $\theta \in (0, \alpha)$, observe that

$$\|e^{-t}T_t^1\phi\|_{H_p^{2\theta/\alpha}} = \|(1-\Delta)^{\theta/\alpha}(e^{-t}T_t^1\phi)\|_{L_p} \le \|e^{-t}T_t^1\phi\|_{L_p} + \|(-\Delta)^{\theta/\alpha}(e^{-t}T_t^1\phi)\|_{L_p},$$

where the last inequality follows from Lemma 2.10 (v). As $e^{-t} \leq (Nt^{-\theta} \wedge 1)$, we have

$$\left\|e^{-t}T_t^1\phi\right\|_{H_p^{2\theta/\alpha}} \le N\left(t^{-\theta} \left\|T_t^1\phi\right\|_{L_p} + \left\|(-\Delta)^{\theta/\alpha}T_t^1\phi\right\|_{L_p}\right).$$
(5.9)

By inequality (5.8), we have

$$t^{-\theta} \|T_t^1 \phi\|_{L_p} \le t^{-\theta} \|\phi\|_{L_p}.$$

$$(5.10)$$

On the other hand, Minkowski's inequality yields

$$\|(-\Delta)^{\theta/\alpha} T_t^1 h\|_{L_p} = \|(-\Delta)^{\theta/\alpha} (p(t,\cdot) * \phi)\|_{L_p} \le \|((-\Delta)^{\theta/\alpha} p)(t,\cdot)\|_{L_1} \|\phi\|_{L_p}.$$
 (5.11)
litionally Lemma 5.1 (iv) (ii) and (iii) imply

$$\|((-\Delta)^{\theta/\alpha}p)(t,\cdot)\|_{L_{1}} = \int_{\mathbb{R}^{d}} |((-\Delta)^{\theta/\alpha}p)(t,x)|dx$$

$$\leq t^{-\theta} \int_{\mathbb{R}^{d}} |((-\Delta)^{\theta/\alpha}p)(1,x)|dx$$

$$\leq N(\alpha,\theta,d,p)t^{-\theta}.$$

(5.12)

Then, by applying (5.12) to (5.11), we have

$$\|(-\Delta)^{\theta/\alpha} T_t^1 \phi\|_{L_p} \le N t^{-\theta} \|\phi\|_{L_p}.$$
(5.13)

Thus, by plugging in (5.10) and (5.13) into (5.9), we have (5.7). The lemma is proved. \Box

To proceed further, we introduce one of the embedding theorems for Slobodetskii's spaces.

Lemma 5.4. If $\mu p > 1$ and $p \ge 1$, for any continuous L_p -valued function ϕ and $\gamma \le \rho$, we have the following:

$$\|\phi(\rho) - \phi(\gamma)\|_{L_p}^p \le N(\rho - \gamma)^{\mu p - 1} \int_{\gamma}^{\rho} \int_{\gamma}^{\rho} 1_{t > s} \frac{\|\phi(t) - \phi(s)\|_{L_p}^p}{|t - s|^{1 + \mu p}} ds dt \quad \left(\frac{0}{0} := 0\right), \quad (5.14)$$

where $N = N(\mu, p)$. In particular,

$$\mathbb{E} \sup_{0 \le s < t \le T} \frac{\|\phi(t) - \phi(s)\|_{L_p}^p}{|t - s|^{\mu p - 1}} \le N \int_0^T \int_0^T \mathbf{1}_{t > s} \frac{\mathbb{E} \|\phi(t) - \phi(s)\|_{L_p}^p}{|t - s|^{1 + \mu p}} ds dt.$$
(5.15)

With the help of Lemma 5.4, we obtain the Hölder continuity of $T_t^1 \phi$, $T_t^2 f$, and $T_t^3 g$ on $t \in [0, T]$.

Lemma 5.5. *Let* $\alpha \in (0, 1)$ *.*

(i) If $p \in (1, \infty)$, $\theta \in (0, \alpha)$, and $\mu \in (0, 1)$ satisfy

$$p > \frac{1}{\alpha - \theta}, \quad \mu \in \left(\frac{1}{\alpha p}, \frac{\alpha - \theta}{\alpha}\right)$$
 (5.16)

and $\phi \in C_c^{\infty}(\mathbb{R}^d)$, then for $t, s \in (0, T)$,

$$\left\| T_t^1 f - T_s^1 f \right\|_{H_p^{\frac{2\theta}{\alpha}-2}}^p \le N |t-s|^{\alpha \mu p-1} \left\| \phi \right\|_{L_p}^p,$$
(5.17)

where $N = N(\alpha, \theta, d, p, T)$. Additionally,

$$\sup_{0 \le s < t \le T} \frac{\left\| T_t^1 \phi - T_s^1 \phi \right\|_{H_p^{\frac{2\theta}{\alpha} - 2}}^p}{|t - s|^{\alpha \mu p - 1}} \le N \left\| \phi \right\|_{L_p}^p$$
(5.18)

where $N = N(\alpha, \theta, d, p, T)$.

(ii) If $p \in (1, \infty)$, $\theta \in (0, \alpha)$, and $\mu \in (0, 1)$ satisfy (5.16) and $f \in C_c^{\infty}((0, T) \times \mathbb{R}^d)$, then for $t, s \in (0, T)$,

$$\left\|T_t^2 f - T_s^2 f\right\|_{H_p^{2\theta/\alpha}}^p \le N|t-s|^{\alpha\mu p-1} \int_0^T \|f(r,\cdot)\|_{L_p}^p dr$$
(5.19)

where $N = N(\alpha, \theta, d, p, T)$. Additionally,

$$\sup_{0 \le s < t \le T} \frac{\left\| T_t^2 f - T_s^2 f \right\|_{H_p^{2\theta/\alpha}}^p}{|t - s|^{\alpha \mu p - 1}} \le N \int_0^T \left\| f(r, \cdot) \right\|_{L_p}^p dr,\tag{5.20}$$

where $N = N(\alpha, \theta, d, p, T)$.

(iii) Let $\tau \leq T$ be a bounded stopping time. If $\beta < \alpha + \frac{1}{2}$, $p \in [2, \infty)$, $\theta \in (0, \alpha - \beta + 1/2)$, and $\mu \in (0, 1)$ satisfy

$$p > \frac{1}{(\alpha - \beta - \theta) \wedge 1/2 + 1/2}, \quad \mu \in \left(\frac{1}{\alpha p}, \frac{(\alpha - \beta - \theta) \wedge 1/2 + 1/2}{\alpha}\right),$$

and $g = (g^1, g^2, ...)$ is a function of type (5.2), then, for $t, s \in (0, T)$, we have

$$\mathbb{E} \left\| T_{t\wedge\tau}^3 g - T_{s\wedge\tau}^3 g \right\|_{H_p^{2\theta/\alpha}(l_2)}^p \le N |t-s|^{\alpha\mu p-1} \|g\|_{\mathbb{L}_p(\tau,l_2)}^p, \tag{5.21}$$

where $N = N(\alpha, \beta, \theta, d, p, T)$. Additionally,

$$\mathbb{E} \sup_{0 \le s < t \le \tau} \frac{\left\| T_t^3 g - T_s^3 g \right\|_{H_p^{2\theta/\alpha}(l_2)}^p}{|t - s|^{\alpha \mu p - 1}} \le N \|g\|_{\mathbb{L}_p(\tau, l_2)}^p, \tag{5.22}$$

where $N = N(\alpha, \beta, \theta, d, p, T)$.

Proof of (i). By (5.14), we have

$$\begin{aligned} \left\| T_{\rho}^{1}\phi - T_{\gamma}^{1}\phi \right\|_{H_{p}^{\frac{2\theta}{\alpha}-2}}^{p} &\leq N|\rho - \gamma|^{\alpha\mu p-1} \int_{0}^{T} \int_{0}^{T} \int_{0}^{T} 1_{t>s} \frac{\left\| T_{t}^{1}\phi - T_{s}^{1}\phi \right\|_{H_{p}^{\frac{2\theta}{\alpha}-2}}^{p}}{|t-s|^{1+\alpha\mu p}} ds dt \\ &\leq N|\rho - \gamma|^{\alpha\mu p-1} \int_{0}^{T} \int_{0}^{T-t} t^{-1-\alpha\mu p} \left\| T_{t+s}^{1}\phi - T_{s}^{1}\phi \right\|_{H_{p}^{\frac{2\theta}{\alpha}-2}}^{p} ds dt. \end{aligned}$$
(5.23)

Since $T_t^1 \phi$ is a solution to (5.6), Lemma 5.3 yields

$$\begin{aligned} |T_{t+s}^{1}\phi - T_{s}^{1}\phi||_{H_{p}^{\frac{2\theta}{\alpha}-2}} \\ &\leq \left\| (T_{t+s}^{1}\phi - \phi) - (T_{s}^{1}\phi - \phi) \right\|_{H_{p}^{\frac{2\theta}{\alpha}-2}} \\ &\leq \left\| \int_{0}^{t+s} (t+s-r)^{\alpha-1} \Delta T_{r}^{1}\phi dr - \int_{0}^{s} (s-r)^{\alpha-1} \Delta T_{r}^{1}\phi dr \right\|_{H_{p}^{\frac{2\theta}{\alpha}-2}} \\ &\leq \int_{s}^{t+s} (t+s-r)^{\alpha-1} \left\| T_{r}^{1}\phi \right\|_{H_{p}^{\frac{2\theta}{\alpha}}} dr \\ &+ \int_{0}^{s} \left| (t+s-r)^{\alpha-1} - (s-r)^{\alpha-1} \right| \left\| T_{r}^{1}\phi \right\|_{H_{p}^{\frac{2\theta}{\alpha}}} dr \\ &\leq N(2t^{\alpha}+s^{\alpha}-(t+s)^{\alpha})t^{-\theta} \|\phi\|_{L_{p}}. \end{aligned}$$
(5.24)

Since $0 \le 2t^{\alpha} + s^{\alpha} - (t+s)^{\alpha} \le 2t^{\alpha}$, by applying (5.24) to (5.23),

$$\begin{split} \left\| T_{t}^{1}\phi - T_{s}^{1}\phi \right\|_{H_{p}^{\frac{2\theta}{\alpha}-2}}^{p} &\leq N |\rho - \gamma|^{\alpha\mu p-1} \int_{0}^{T} t^{-1-\alpha\mu p+\alpha p-\theta p} dt \|\phi\|_{L_{p}}^{p} \\ &\leq N |\rho - \gamma|^{\alpha\mu p-1} \|\phi\|_{L_{p}}^{p}. \end{split}$$

Thus, we have (5.17).

To obtain (5.18), use (5.15) instead of (5.14) and repeat the proof.

Proof of (ii). Let $\rho > 0$ and $\gamma > 0$. Notice that (5.14) yields

$$\left\|T_{\rho}^{2}f - T_{\gamma}^{2}f\right\|_{H_{p}^{2\theta/\alpha}}^{p} \leq N|\rho - \gamma|^{\alpha\mu p - 1} \int_{0}^{T} \int_{0}^{T} 1_{t > s} \frac{\left\|T_{t}^{2}f - T_{s}^{2}f\right\|_{H_{p}^{2\theta/\alpha}}^{p}}{|t - s|^{1 + \alpha\mu p}} ds dt.$$
(5.25)

Then, by definition of $T_t^2 f$ (see (5.4)),

$$\begin{aligned} \left\| T_{t}^{2} f - T_{s}^{2} f \right\|_{H_{p}^{2\theta/\alpha}} \\ &\leq \left\| \int_{s}^{t} \int_{\mathbb{R}^{d}} q_{\alpha,1}(t-r,y) f(r,\cdot-y) dy dr \right\|_{H_{p}^{2\theta/\alpha}} \\ &+ \left\| \int_{0}^{s} \int_{\mathbb{R}^{d}} \left(q_{\alpha,1}(t-r,y) - q_{\alpha,1}(s-r,y) \right) f(r,\cdot-y) dy dr \right\|_{H_{p}^{2\theta/\alpha}}. \end{aligned}$$
(5.26)

 Set

$$I_{1} := \int_{0}^{T} \int_{0}^{T} 1_{t>s} \frac{\left\| \int_{s}^{t} \int_{\mathbb{R}^{d}} q_{\alpha,1}(t-r,y) f(r,\cdot-y) dy dr \right\|_{H_{p}^{2\theta/\alpha}}^{p}}{|t-s|^{1+\alpha\mu p}} ds dt,$$

$$I_{2} := \int_{0}^{T} \int_{0}^{T} 1_{t>s} \frac{\left\| \int_{0}^{s} \int_{\mathbb{R}^{d}} \left(q_{\alpha,1}(t-r,y) - q_{\alpha,1}(s-r,y) \right) f(r,\cdot-y) dy dr \right\|_{H_{p}^{2\theta/\alpha}}^{p}}{|t-s|^{1+\alpha\mu p}} ds dt.$$
(5.27)

Then, apply (5.26) and (5.27) to (5.25),

$$\left\|T_t^2 f - T_s^2 f\right\|_{H_p^{2\theta/\alpha}}^p \le |t - s|^{\alpha\mu p - 1} (I_1 + I_2).$$
(5.28)

To deal with I_1 , we employ Minkowski's inequality, the change of variable, and Lemma 2.10 (v). Then,

$$I_{1} \leq \int_{0}^{T} t^{-1-\alpha\mu p} \left(\int_{0}^{t} \int_{\mathbb{R}^{d}} \left| \left((1-\Delta)^{\theta/\alpha} q_{\alpha,1} \right) (r,y) \right| dy dr \right)^{p} dt \int_{0}^{T} \|f(s,\cdot)\|_{L_{p}}^{p} ds$$

$$\leq I_{11} + I_{12},$$
(5.29)

where

$$I_{11} := \int_0^T t^{-1-\alpha\mu p} \left(\int_0^t \int_{\mathbb{R}^d} |q_{\alpha,1}(r,y)| \, dy dr \right)^p dt \int_0^T \|f(s,\cdot)\|_{L_p}^p \, ds,$$

$$I_{12} := \int_0^T t^{-1-\alpha\mu p} \left(\int_0^t \int_{\mathbb{R}^d} \left| \left(\Delta^{\theta/\alpha} q_{\alpha,1} \right) (r,y) \right| \, dy dr \right)^p dt \int_0^T \|f(s,\cdot)\|_{L_p}^p \, ds.$$

Because $\mu < 1$, Lemma 5.1 (iv) and (iii) yield

$$I_{11} = \int_0^T t^{-1-\alpha\mu p + \alpha p} dt \left(\int_{\mathbb{R}^d} |q_{\alpha,1}(1,y)| \, dy \right)^p \int_0^T \|f(s,\cdot)\|_{L_p}^p \, ds$$

$$\leq N \int_0^T \|f(s,\cdot)\|_{L_p}^p \, ds.$$
(5.30)

Similarly, since $\mu < 1 - \theta/\alpha$,

$$I_{12} = \int_0^T t^{-1-\alpha\mu p + \alpha p - \theta p} dt \left(\int_{\mathbb{R}^d} \left| \left(\Delta^{\theta/\alpha} q_{\alpha,1} \right) (1,y) \right| dy \right)^p \int_0^T \|f(s,\cdot)\|_{L_p}^p ds$$

$$\leq N \int_0^T \|f(s,\cdot)\|_{L_p}^p ds.$$
(5.31)

Thus, by applying (5.30) and (5.31) to (5.29), we have

$$I_1 \le N \int_0^T \|f(s, \cdot)\|_{L_p}^p ds.$$
(5.32)

Next, we address I_2 . Similar to the case for I_1 , we have

$$I_{2} \leq \int_{0}^{T} \frac{\int_{0}^{T-t} \left(\int_{0}^{s} \int_{\mathbb{R}^{d}} \left| (1-\Delta)^{\theta/\alpha} \left(q_{\alpha,1}(t+r,y) - q_{\alpha,1}(r,y) \right) \right| dy \| f(s-r,\cdot) \|_{L_{p}} dr \right)^{p} ds}{t^{1+\alpha\mu p}} dt$$

$$\leq I_{21} + I_{22}, \tag{5.33}$$

where

$$I_{21} := \int_0^T \frac{\int_0^{T-t} \left(\int_0^s \int_{\mathbb{R}^d} |q_{\alpha,1}(t+r,y) - q_{\alpha,1}(r,y)| \, dy \|f(s-r,\cdot)\|_{L_p} dr \right)^p \, ds}{t^{1+\alpha\mu p}} dt,$$

$$I_{22} := \int_0^T \frac{\int_0^{T-t} \left(\int_0^s \int_{\mathbb{R}^d} \left| \Delta^{\theta/\alpha} \left(q_{\alpha,1}(t+r,y) - q_{\alpha,1}(r,y) \right) \right| \, dy \|f(s-r,\cdot)\|_{L_p} dr \right)^p \, ds}{t^{1+\alpha\mu p}} dt.$$

Since $\mu < 1$, by Minkowski's inequality and the fundamental theorem of calculus, we have

$$\begin{split} I_{21} &\leq \int_{0}^{T} t^{-1-\alpha\mu p} \left(\int_{0}^{T-t} \int_{\mathbb{R}^{d}} |q_{\alpha,1}(t+r,y) - q_{\alpha,1}(r,y)| \, dy dr \right)^{p} dt \int_{0}^{T} \|f(s,\cdot)\|_{L_{p}(\mathbb{R}^{d})}^{p} ds \\ &\leq \int_{0}^{T} t^{-1-\alpha\mu p} \left(\int_{0}^{T-t} \int_{r}^{t+r} \int_{\mathbb{R}^{d}} |q_{\alpha,2}(s,y)| \, dy ds dr \right)^{p} dt \int_{0}^{T} \|f(s,\cdot)\|_{L_{p}(\mathbb{R}^{d})}^{p} ds \\ &\leq N \int_{0}^{T} t^{-1-\alpha\mu p} \left(\int_{0}^{T-t} r^{\alpha-1} - (t+r)^{\alpha-1} dr \right)^{p} dt \int_{0}^{T} \|f(s,\cdot)\|_{L_{p}(\mathbb{R}^{d})}^{p} ds \\ &\leq N \int_{0}^{T} t^{-1-\alpha\mu p+\alpha p} dt \int_{0}^{T} \|f(s,\cdot)\|_{L_{p}(\mathbb{R}^{d})}^{p} ds \\ &\leq N \int_{0}^{T} \|f(s,\cdot)\|_{L_{p}(\mathbb{R}^{d})}^{p} ds. \end{split}$$

$$(5.34)$$

Additionally, since $\mu < 1 - \theta/\alpha$,

$$I_{22} \leq \int_{0}^{T} t^{-1-\alpha\mu p} \left(\int_{0}^{T-t} \int_{\mathbb{R}^{d}} \int_{r}^{t+r} \left| (\Delta^{\theta/\alpha} q_{\alpha,2}(s,y) \right| ds dy dr \right)^{p} dt \int_{0}^{T} \|f(s,\cdot)\|_{L_{p}}^{p} ds$$

$$\leq N \int_{0}^{T} t^{-1-\alpha\mu p+\alpha p-\theta p} dt \int_{0}^{T} \|f(s,\cdot)\|_{L_{p}}^{p} ds$$

$$\leq N \int_{0}^{T} \|f(s,\cdot)\|_{L_{p}}^{p} ds.$$
(5.35)

Therefore, by employing (5.34) and (5.35) to (5.33), we have

$$I_2 \le N \int_0^T \|f(s, \cdot)\|_{L_p}^p ds,$$
(5.36)

and thus by combining (5.32) and (5.36) to (5.28), we have (5.19).

To obtain (5.20), employ (5.15) instead of (5.14) and repeat the proof word for word. *Proof of* (iii). By (5.14), we have

$$\mathbb{E} \left\| T_{\rho}^{3}g - T_{\gamma}^{3}g \right\|_{H_{p}^{2\theta/\alpha}(l_{2})}^{p} \leq N|\rho - \gamma|^{\alpha\mu p - 1} \int_{0}^{T} \int_{0}^{T} 1_{t > s} \frac{\mathbb{E} \left\| T_{t}^{3}g - T_{s}^{3}g \right\|_{H_{p}^{2\theta/\alpha}(l_{2})}^{p}}{|t - s|^{1 + \alpha\mu p}} ds dt.$$

Notice that the Burkholder-Davis-Gundy and Minkowski's inequalities imply that

$$\begin{split} & \mathbb{E} \left\| T_{t}^{3}g - T_{s}^{3}g \right\|_{H_{p}^{p}(l_{2})}^{p} \\ & \leq N \int_{\mathbb{R}^{d}} \mathbb{E} \left(\sum_{k} \left| \int_{s}^{t} \int_{\mathbb{R}^{d}} ((1-\Delta)^{\theta/\alpha} q_{\alpha,\beta})(t-r,y)g^{k}(r,x-y)dydw_{r}^{k} \right|^{2} \right)^{p/2} \\ & + \left(\sum_{k} \left| \int_{0}^{s} \int_{\mathbb{R}^{d}} \left[(1-\Delta)^{\theta/\alpha} \left(q_{\alpha,\beta}(t-r,y) - q_{\alpha,\beta}(s-r,y) \right) \right] g^{k}(r,x-y)dydw_{r}^{k} \right|^{2} \right)^{p/2} dx \\ & \leq N \int_{\mathbb{R}^{d}} \mathbb{E} \left(\int_{0}^{t-s} \left(\int_{\mathbb{R}^{d}} \left| ((1-\Delta)^{\theta/\alpha} q_{\alpha,\beta})(t-s-r,y) \right| \left| g(s+r,x-y) \right|_{l_{2}} dy \right)^{2} dr \right)^{p/2} \\ & + \left(\int_{0}^{s} \left(\int_{\mathbb{R}^{d}} \left| (1-\Delta)^{\theta/\alpha} \left(q_{\alpha,\beta}(t-r,y) - q_{\alpha,\beta}(s-r,y) \right) \right| \left| g(r,x-y) \right|_{L_{p}(l_{2})} dr \right)^{p/2} dx \\ & \leq N \mathbb{E} \left(\int_{0}^{t-s} \left(\int_{\mathbb{R}^{d}} \left| ((1-\Delta)^{\theta/\alpha} q_{\alpha,\beta})(t-s-r,y) \right| dy \right)^{2} \left\| g(s+r,\cdot) \right\|_{L_{p}(l_{2})}^{2} dr \right)^{p/2} dx \\ & \leq N \mathbb{E} \left(\int_{0}^{s} \left(\int_{\mathbb{R}^{d}} \left| (1-\Delta)^{\theta/\alpha} \left(q_{\alpha,\beta}(t-r,y) - q_{\alpha,\beta}(s-r,y) \right) \right| dy \right)^{2} \left\| g(r,\cdot) \right\|_{L_{p}(l_{2})}^{2} dr \right)^{p/2}. \end{split}$$

Then, set

$$\begin{split} I_3 &:= \int_0^T \int_0^T \mathbf{1}_{t>s} |t-s|^{-1-\alpha\mu p} \mathbb{E}\left(\int_0^{t-s} A(t,s,r) \|g(s+r,\cdot)\|_{L_p(l_2)}^2 dr\right)^{p/2} ds dt, \\ I_4 &:= \int_0^T \int_0^T \mathbf{1}_{t>s} |t-s|^{-1-\alpha\mu p} \mathbb{E}\left(\int_0^s B(t,s,r) \|g(r,\cdot)\|_{L_p(l_2)}^2 dr\right)^{p/2} ds dt, \end{split}$$

where

$$\begin{aligned} A(t,s,r) &= \left(\int_{\mathbb{R}^d} |(1-\Delta)^{\theta/\alpha} q_{\alpha,\beta}(t-s-r,y)| dy \right)^2, \\ B(t,s,r) &= \left(\int_{\mathbb{R}^d} \left| (1-\Delta)^{\theta/\alpha} \left(q_{\alpha,\beta}(t-r,y) - q_{\alpha,\beta}(s-r,y) \right) \right| dy \right)^2. \end{aligned}$$

Note that Minkowski's inequality and Lemma 2.10 (v) imply that

$$I_{3} \leq \int_{0}^{T} t^{-1-\alpha\mu p} \left(\int_{0}^{t} \left(\int_{\mathbb{R}^{d}} \left| ((1-\Delta)^{\theta/\alpha} q_{\alpha,\beta})(t-r,y) \right| dy \right)^{2} dr \right)^{p/2} dt \|g\|_{\mathbb{L}_{p}(T,l_{2})}^{p} \quad (5.37)$$

$$\leq I_{31} + I_{32},$$

where

$$I_{31} := \int_0^T t^{-1-\alpha\mu p} \left(\int_0^t \left(\int_{\mathbb{R}^d} |q_{\alpha,\beta}(r,y)| dy \right)^2 dr \right)^{p/2} dt \|g\|_{\mathbb{L}_p(T,l_2)}^p,$$

$$I_{32} := \int_0^T t^{-1-\alpha\mu p} \left(\int_0^t \left(\int_{\mathbb{R}^d} \left| \left(\Delta^{\theta/\alpha} q_{\alpha,\beta} \right)(r,y) \right| dy \right)^2 dr \right)^{p/2} dt \|g\|_{\mathbb{L}_p(T,l_2)}^p.$$

Since $\frac{1}{\alpha} \left(\alpha - \beta + \frac{1}{2} \right) > \mu$, by Lemma 5.1 (iii), we have

$$I_{31} = \int_{0}^{T} t^{-1-\alpha\mu p} \left(\int_{0}^{t} \left(\int_{\mathbb{R}^{d}} |q_{\alpha,\beta}(r,y)| dy \right)^{2} dr \right)^{p/2} dt ||g||_{\mathbb{L}_{p}(T,l_{2})}^{p} \\ = \int_{0}^{T} t^{-1-\alpha\mu p} \left(\int_{0}^{t} r^{2(\alpha-\beta)} dr \right)^{p/2} dt \left(\int_{\mathbb{R}^{d}} |q_{\alpha,\beta}(1,y)| dy \right)^{p} ||g||_{\mathbb{L}_{p}(T,l_{2})}^{p} \\ = \int_{0}^{T} t^{-1+(\alpha-\beta+1/2-\alpha\mu)p} dt \left(\int_{\mathbb{R}^{d}} |q_{\alpha,\beta}(1,y)| dy \right)^{p} ||g||_{\mathbb{L}_{p}(T,l_{2})}^{p} \\ \leq N ||g||_{\mathbb{L}_{p}(T,l_{2})}^{p}.$$
(5.38)

Similarly, as
$$\frac{1}{\alpha} \left(\alpha - \beta - \theta + \frac{1}{2} \right) > \mu$$
,

$$I_{32} \leq \int_{0}^{T} t^{-1 + (-\alpha\mu + \alpha - \beta - \theta + 1/2)p} dt \left(\int_{\mathbb{R}^{d}} |(\Delta^{\theta/\alpha} q_{\alpha,\beta})(1,y)| dy \right)^{p} ||g||_{\mathbb{L}_{p}(T,l_{2})}^{p}$$

$$\leq N ||g||_{\mathbb{L}_{p}(T,l_{2})}^{p}.$$
(5.39)

Therefore, by employing (5.38) and (5.39) to (5.37), we have

$$I_3 \le N \|g\|_{\mathbb{L}_p(T, l_2)}^p.$$
(5.40)

In the case of I_4 , by Minkowski's inequality and Lemma 2.10 (v), we have

$$I_4 \le I_{41} + I_{42}.$$

Further,

$$\begin{split} I_{41} &:= \int_0^T t^{-1-\alpha\mu p} \mathbb{E} \int_0^{T-t} \left(\int_0^s \left(\int_{\mathbb{R}^d} |C(t,r,y)| dy \right)^2 \|g(s-r,\cdot)\|_{L_p(l_2)}^2 dr \right)^{p/2} ds dt, \\ I_{42} &:= \int_0^T t^{-1-\alpha\mu p} \mathbb{E} \int_0^{T-t} \left(\int_0^s \left(\int_{\mathbb{R}^d} |\Delta^{\theta/\alpha} C(t,r,y)| dy \right)^2 \|g(s-r,\cdot)\|_{L_p(l_2)}^2 dr \right)^{p/2} ds dt, \end{split}$$

where

$$C(t, r, y) = q_{\alpha, \beta}(t + r, y) - q_{\alpha, \beta}(r, y).$$

We address I_{41} . By Minkowski's inequality, the fundamental theorem of calculus, and Lemma 5.1, we have

$$I_{41} \leq N \int_{0}^{T} t^{-1-\alpha\mu p} \left(\int_{0}^{T-t} \left(\int_{\mathbb{R}^{d}} |q_{\alpha,\beta}(t+r,y) - q_{\alpha,\beta}(r,y)| dy \right)^{2} dr \right)^{p/2} dt ||g||_{\mathbb{L}_{p}(T,l_{2})}^{p}$$

$$\leq N \int_{0}^{T} t^{-1-\alpha\mu p} \left(\int_{0}^{T-t} \left(\int_{r}^{t+r} \int_{\mathbb{R}^{d}} |q_{\alpha,\beta+1}(s,y)| dy ds \right)^{2} dr \right)^{p/2} dt ||g||_{\mathbb{L}_{p}(T,l_{2})}^{p}$$

$$\leq N H_{1} ||g||_{\mathbb{L}_{p}(T,l_{2})}^{p}, \qquad (5.41)$$

where

$$H_1 := \int_0^T t^{-1-\alpha\mu p} \left(\int_0^{T-t} \left(\int_r^{t+r} s^{\alpha-\beta-1} ds \right)^2 dr \right)^{p/2} dt$$
(5.42)

and $N = N(\alpha, \beta, d, p, T)$. Next, we claim that

$$H_1 < \infty \tag{5.43}$$

To demonstrate (5.43), set

$$\chi(t) := \int_0^{T-t} \left(\int_r^{t+r} s^{\alpha-\beta-1} ds \right)^2 dr.$$

Furthermore,

$$H_1 = \int_0^T t^{-1 - \alpha \mu p} (\chi(t))^{p/2} dt$$
(5.44)

Depending on the range of $\alpha - \beta$, we consider the following five cases.

(Case 1.) $-1/2 < \alpha - \beta < 0$ For $t \in (0, T)$, we have

$$0 \le \chi(t) \le N \int_0^{T-t} r^{2(\alpha-\beta)} - (t+r)^{2(\alpha-\beta)} dr \le N t^{2(\alpha-\beta)+1},$$
(5.45)

where $N = N(\alpha, \beta)$. Then, since $\frac{1}{\alpha}(\alpha - \beta + 1/2) > \mu$, by combining (5.44) and (5.45),

$$H_1 \le N(\alpha, \beta, p) \int_0^T t^{-1+p(-\alpha\mu+\alpha-\beta+1/2)} dt < \infty.$$

(Case 2.) $\alpha - \beta = 0$

Notice that

$$\chi(t) = \int_0^{T-t} \left(\int_r^{t+r} s^{-1} ds \right)^2 dr = \int_0^{T-t} \left(\log\left(\frac{t+r}{r}\right) \right)^2 dr.$$

Obviously, $\chi(0) = 0$. Note that

$$\chi'(t) \le 2\int_0^\infty \frac{1}{1+r} \log\left(\frac{1+r}{r}\right) dr = 2\int_0^\infty x(e^x - 1)^{-1} dx = \pi^2/3$$

on $t \in (0, T/2)$. Thus,

$$0 \le \chi(t) = \chi(t) - \chi(0) = \int_0^t \chi'(s) ds \le \frac{\pi^2}{3}t$$
(5.46)

on $t \in (0, T/2)$. Additionally, $\chi(t) \leq N$ on $t \in (T/2, T)$. Therefore,

$$H_1 \le N \int_0^{T/2} t^{-1 + (-\alpha \mu + 1/2)p} dt + N < \infty,$$

where $N = N(\alpha, \beta, p)$.

(*Case 3.*) $0 < \alpha - \beta < 1/2$

Observe that χ is twice continuously differentiable, and

$$\chi'(t) = (\alpha - \beta)^{-2} \left[-T^{2(\alpha - \beta)} - (T - t)^{2(\alpha - \beta)} + 2T^{\alpha - \beta}(T - t)^{\alpha - \beta} \right] + 2(\alpha - \beta)^{-1} \int_0^{T - t} (t + r)^{2(\alpha - \beta) - 1} - (t + r)^{\alpha - \beta - 1} r^{\alpha - \beta} dr,$$

and

$$\chi''(t) = 2(\alpha - \beta)^{-1}(T - t)^{2(\alpha - \beta) - 1} - 2(\alpha - \beta)^{-1}T^{\alpha - \beta}(T - t)^{\alpha - \beta - 1} + 2(\alpha - \beta)^{-1}T^{\alpha - \beta - 1}(T - t)^{\alpha - \beta} - 2(\alpha - \beta)^{-1}t^{2(\alpha - \beta) - 1} - 2(\alpha - \beta)^{-1}(\alpha - \beta - 1)\int_{0}^{T - t}(t + r)^{\alpha - \beta - 2}r^{\alpha - \beta}dr.$$
(5.47)

In addition, $\chi(0) = \chi'(0) = 0$. Then, by using the fundamental theorem of calculus and $\alpha - \beta \in (0, 1/2)$, we obtain

$$\begin{split} \chi(t) &= \int_{0}^{t} \int_{0}^{s} \chi''(\rho) d\rho ds \\ &\leq 2(\alpha - \beta)^{-1} \int_{0}^{t} \int_{0}^{s} (T - \rho)^{2(\alpha - \beta) - 1} d\rho ds \\ &- 2(\alpha - \beta)^{-1} (\alpha - \beta - 1) \int_{0}^{t} \int_{0}^{s} \int_{0}^{T - \rho} (\rho + r)^{\alpha - \beta - 2} r^{\alpha - \beta} dr d\rho ds \\ &\leq N \int_{0}^{t} \left(T^{2(\alpha - \beta)} - (T - s)^{2(\alpha - \beta)} \right) ds + N \int_{0}^{t} \int_{0}^{s} \int_{0}^{T - \rho} (\rho + r)^{2(\alpha - \beta) - 2} dr d\rho ds \\ &\leq N \int_{0}^{t} s^{2(\alpha - \beta)} ds + N \int_{0}^{t} \int_{0}^{s} T^{2(\alpha - \beta) - 1} dr d\rho ds \\ &\leq N t^{2(\alpha - \beta) + 1}, \end{split}$$
(5.48)

where $N = N(\alpha, \beta, T)$. Thus,

$$H_1 \le N(\alpha, \beta, p, T) \int_0^T t^{-1+p(-\alpha\mu+\alpha-\beta+1/2)} dt < \infty.$$

(Case 4.)
$$\alpha - \beta = 1/2$$

Because $\chi(0) = \chi'(0) = 0$, by the fundamental theorem of calculus, we have
 $\chi(t) = \int_0^t \int_0^s \chi''(\rho) d\rho ds$
 $\leq N \int_0^t \int_0^s (T - \rho)^{1/2} + N \int_0^{T-\rho} (\rho + r)^{-3/2} r^{1/2} dr d\rho ds$
 $\leq N t^2 (1 + |\log t|),$

where N = N(T). Therefore,

$$H_1 \le N(p,T) \int_0^T t^{-1+p(-\alpha\mu+1)} (1+|\log t|)^{p/2} dt < \infty.$$

(*Case 5.*) $\alpha - \beta > 1/2$

Similar to before, $\chi(0) = \chi'(0) = 0$. Additionally, as in (5.48), we have

$$\begin{split} \chi(t) &= \int_0^t \int_0^s \chi''(r) dr ds \\ &\leq N \int_0^t \left(T^{2(\alpha-\beta)} - (T-s)^{2(\alpha-\beta)} \right) ds + N \int_0^t \int_0^s \int_0^{T-\rho} (\rho+r)^{2(\alpha-\beta)-2} dr d\rho ds \\ &\leq N \int_0^t s ds + N \int_0^t \int_0^s d\rho ds \\ &\leq N t^2, \end{split}$$

where $N = N(\alpha, \beta, T)$. Therefore,

$$H_1 \le N(\alpha, \beta, p, T) \int_0^T t^{-1+p(-\alpha\mu+1)} dt < \infty.$$

Thus, we have (5.43). Then, by combining (5.43) and (5.41), we have $I_{41} \leq N ||g||_{\mathbb{L}_p(T,l_2)}^p$.

Next, we deal with I_{42} . Minkowski's inequality, the fundamental theorem of calculus, and Lemma 5.1 yield

$$\begin{split} I_{42} &\leq N \int_0^T t^{-1-\alpha\mu p} \left(\int_0^{T-t} \left(\int_{\mathbb{R}^d} |\Delta^{\theta/\alpha} q_{\alpha,\beta}(t+r,y) - \Delta^{\theta/\alpha} q_{\alpha,\beta}(r,y)| dy \right)^2 dr \right)^{p/2} dt \|g\|_{\mathbb{L}_p(T,l_2)}^p \\ &\leq N \int_0^T t^{-1-\alpha\mu p} \left(\int_0^{T-t} \left(\int_r^{t+r} \int_{\mathbb{R}^d} |\partial_s \Delta^{\theta/\alpha} q_{\alpha,\beta}(s,y)| dy ds \right)^2 dr \right)^{p/2} dt \|g\|_{\mathbb{L}_p(T,l_2)}^p \\ &\leq N H_2 \|g\|_{\mathbb{L}_p(T,l_2)}^p, \end{split}$$

where $N = N(\alpha, \beta, d, p, T)$ and $H_2 := \int_0^T t^{-1-\alpha\mu p} \left(\int_0^{T-t} \left(\int_r^{t+r} s^{\alpha-\beta-\theta-1} ds \right)^2 dr \right)^{p/2} dt.$ Similar to the case of H_1 (see (5.42) and (5.43)), we demonstrate that

 $H_2 < \infty$

by considering five cases for $\alpha - \beta - \theta$ instead of $\alpha - \beta$. Then, we have $I_{42} \leq N \|g\|_{\mathbb{L}_p(T, l_2)}^p$. The lemma is proved.

Proof of Theorem 2.16. It suffices to show that the assertion holds for $\tau = T$. Indeed, assume the results holds for $\mathcal{H}_p^{\gamma}(T)$, and let $\tau \leq T$ be a bounded stopping time and $u \in \mathcal{H}_p^{\gamma}(\tau)$. Then, by Definition 2.4 for $\varepsilon > 0$, there exists $(f,g) \in \mathbb{H}_p^{\gamma-2}(\tau) \times \mathbb{H}_p^{\gamma-2+c_0}(\tau, l_2)$ such that

$$\partial_t^{\alpha} u = f + \partial_t^{\beta} \int_0^t g^k dw_t^k; \quad u(0, \cdot) = u_0(\cdot)$$

and

$$\|u\|_{\mathbb{H}_{p}^{\gamma}(\tau)} + \|u_{0}\|_{U_{p}^{\gamma}} + \|f\|_{\mathbb{H}_{p}^{\gamma-2}(\tau)} + \|g\|_{\mathbb{H}_{p}^{\gamma-2+c_{0}}(\tau,l_{2})} \leq \|u\|_{\mathcal{H}_{p}^{\gamma}(\tau)} + \varepsilon.$$

Set $\bar{f} := (f - \Delta u) \mathbb{1}_{t \leq \tau}$ and $\bar{g} := g \mathbb{1}_{t \leq \tau}$; thus, u satisfies

$$\partial_t^{\alpha} u = \Delta u + \bar{f} + \partial_t^{\beta} \int_0^t \bar{g}^k dw_t^k, \quad 0 < t \le \tau \,; \quad u(0, \cdot) = u_0(\cdot). \tag{5.49}$$

In contrast, by [25, Theorem 2.18], there exists $v \in \mathcal{H}_p^{\gamma}(T)$ such that v satisfies

$$\partial_t^{\alpha} v = \Delta v + \bar{f} + \partial_t^{\beta} \int_0^t \bar{g}^k dw_t^k, \quad 0 < t \le \tau \,; \quad v(0, \cdot) = u_0(\cdot).$$

Additionally,

$$\begin{aligned} \|v\|_{\mathcal{H}_{p}^{\gamma}(T)} &\leq N\left(\left\|\bar{f}\right\|_{\mathbb{H}_{p}^{\gamma-2}(T)} + \|\bar{g}\|_{\mathbb{H}_{p}^{\gamma-2+c_{0}}(T,l_{2})} + \|u_{0}\|_{U_{p}^{\gamma}}\right) \\ &\leq N\left(\|u\|_{\mathbb{H}_{p}^{\gamma}(\tau)} + \|u_{0}\|_{U_{p}^{\gamma}} + \|f\|_{\mathbb{H}_{p}^{\gamma-2}(\tau)} + \|g\|_{\mathbb{H}_{p}^{\gamma-2+c_{0}}(\tau,l_{2})}\right) \\ &\leq N\|u\|_{\mathcal{H}_{p}^{\gamma}(\tau)} + N\varepsilon, \end{aligned}$$
(5.50)

where N is independent of ε . Because $v \in \mathcal{H}_p^{\gamma}(T), v \in C^{\alpha \mu - 1/p}\left([0,T]; H_p^{\gamma - 2\nu}\right)$ almost surely and

$$\mathbb{E} \|v\|_{C^{\alpha\mu-1/p}([0,T];H_p^{\gamma-2\nu})}^p \le N \|v\|_{\mathcal{H}_p^{\gamma}(T)}^p \tag{5.51}$$

by the hypothesis. Therefore, due to $\tau \leq T$, (5.50) and (5.51) yield

$$\mathbb{E}\|u\|_{C^{\alpha\mu-1/p}([0,\tau];H_p^{\gamma-2\nu})}^p \le \mathbb{E}\|v\|_{C^{\alpha\mu-1/p}([0,T];H_p^{\gamma-2\nu})}^p \le N\|v\|_{\mathcal{H}_p^{\gamma}(T)}^p \le N\|u\|_{\mathcal{H}_p^{\gamma}(\tau)}^p + N\varepsilon,$$
(5.52)

where N is independent of ε . Note that $\overline{u} := u - v$ satisfies

$$\partial_t^\alpha \bar{u} = \Delta \bar{u}, \quad 0 < t \le \tau; \quad \bar{u}(0, \cdot) = 0.$$

Then, by the deterministic version of [25, Theorem 2.18], we have $u(t, \cdot) = v(t, \cdot)$ for almost every $(\omega, t) \in \Omega \times [0, \tau]$. Thus, v is an $H_p^{\gamma-2\nu}$ -valued continuous version of u. Additionally, from (5.52)

$$\mathbb{E} \|u\|_{C^{\alpha\mu-1/p}([0,\tau];H_p^{\gamma-2\nu})}^p \le N \|u\|_{\mathcal{H}_p^{\gamma}(\tau)}^p + N\varepsilon.$$

In addition, as $\varepsilon > 0$ is arbitrary and N is independent of ε , we have (2.8). Additionally, notice that (5.50) allows us to prove the assertions with $\|f\|_{\mathbb{H}_p^{\gamma-2}(T)} + \|g\|_{\mathbb{H}_p^{\gamma-2+c_0}(T,l_2)} + \|u_0\|_{U_p^{\gamma}}$ instead of $\|u\|_{\mathcal{H}_p^{\gamma}(T)}$.

Due to Lemma 2.10 (vi), we only consider the case $\gamma = 2\nu$, where $\nu \in (0, 1)$ satisfies (2.7). Moreover, by using the approximation to the identity, we may assume that u_0 is infinitely differentiable and compactly supported in x. Furthermore, we also assume that f and $g = (g^1, g^2, ...)$ denotes the function of the form satisfying (5.2) (see [26, Theorem 3.10]). Additionally, it should be remarked that u can be written as

$$u(t,x) = T_t^1 u_0 + T_t^2 f + T_t^3 g$$

since u satisfies

$$\partial_t^{\alpha} u = \Delta u + f + \partial_t^{\beta} \int_0^t g^k dw_t^k \, ; \quad u(0, \cdot) = u_0(\cdot)$$

for almost every $(\omega, t) \in (0, \tau]$ (see [22, 23, 24]).

Now, observe that $u \in C^{\alpha\mu-1/p}([0,\tau]; H_p^{\gamma-2\nu})$ almost surely. Indeed, by Lemma 5.5, we have

$$\mathbb{E} \left\| T_t^1 f - T_s^1 f \right\|_{H_p^{\gamma-2\nu}}^{p} \leq N |t-s|^{\alpha\mu p-1} \|\phi\|_{H_p^{\gamma}}^{p}, \\ \mathbb{E} \left\| T_t^2 f - T_s^2 f \right\|_{H_p^{\gamma-2\nu}}^{p} \leq N |t-s|^{\alpha\mu p-1} \|f\|_{\mathbb{H}_p^{\gamma-2}(\tau)}^{p} dr, \\ \mathbb{E} \left\| T_{t\wedge\tau}^3 g - T_{s\wedge\tau}^3 g \right\|_{H_p^{\gamma-2\nu}(l_2)}^{p} \leq N |t-s|^{\alpha\mu p-1} \|g\|_{\mathbb{H}_p^{\gamma-2+c_0}(\tau,l_2)}^{p}.$$

Thus, Kolmogorov continuity theorem (e.g., [28, Theorem 1.4.8]) and the above inequalities again, we have the continuity of u.

Next we prove (2.8). Choose μ and ν satisfy (2.7). Observe that Lemma 2.10 (vi) and (5.17) with $\theta = \alpha - \alpha \nu$ imply that

$$\mathbb{E} \sup_{0 \le s < t \le T} \frac{\|T_t^1 u_0 - T_s^1 u_0\|_{H_p^{\gamma - 2\nu}}^p}{|t - s|^{\alpha \mu p - 1}} \le N \mathbb{E} \|u_0\|_{H_p^{\gamma}}^p$$

As (2.7) is assumed, Lemma 2.10 (vi) and (5.20) with $\theta = \alpha - \alpha \nu$ yield the following:

$$\mathbb{E} \sup_{0 \le s < t \le \tau} \frac{\|T_t^2 f - T_2^2 f\|_{H_p^{\gamma-2\nu}}^p}{|t-s|^{\alpha\mu p-1}} \le N \|f\|_{\mathbb{H}_p^{\gamma-2}(\tau)}^p.$$

Furthermore, by (2.7), Lemma 2.10 (vi), and (5.22) with $\theta = \alpha(1 - c_0/2) - \alpha \nu$, we have

$$\mathbb{E} \sup_{0 \le s < t \le T} \frac{\left\| T_t^3 g - T_s^3 g \right\|_{H_p^{\gamma - 2\nu}(l_2)}^p}{|t - s|^{\alpha \mu p - 1}} \le N \|g\|_{\mathbb{H}_p^{\gamma - 2 + c_0}(\tau, l_2)}^p$$

The theorem is proved.

References

- Garra, Roberto. Fractional-calculus model for temperature and pressure waves in fluid-saturated porous rocks. *Physical Review E*, 84(3):036605, 2011.
- [2] Keller, Jakob J. Propagation of simple non-linear waves in gas filled tubes with friction. Zeitschrift für angewandte Mathematik und Physik ZAMP, 32(2):170–181, 1981.
- [3] Tayyaba Akram, Muhammad Abbas, Muhammad Riaz, Ahmad Ismail, and Norhashidah Ali. An efficient numerical technique for solving time fractional burgers equation. *Alexandria Engineering Journal*, 59(4):2201–2220, 2020.
- [4] Ricardo Almeida. A Gronwall inequality for a general caputo fractional operator. Mathematical Inequalities Applications, 20(4), 2017.
- [5] Ronald Bagley and Peter Torvik. Fractional calculus in the transient analysis of viscoelastically damped structures. AIAA journal, 23(6):918–925, 1985.
- [6] Dumitru Baleanu, Kai Diethelm, Enrico Scalas, and Juan Trujillo. Fractional calculus: models and numerical methods, volume 3. World Scientific, 2012.
- [7] Zhen-Qing Chen, Kyeong-Hun Kim, and Panki Kim. Fractional time stochastic partial differential equations. Stochastic Processes and their Applications, 125(4):1470–1499, 2015.
- [8] Hongjie Dong and Doyoon Kim. L_p-estimates for time fractional parabolic equations with coefficients measurable in time. Advances in Mathematics, 345:289–345, 2019.
- [9] Talaat El-Danaf and Adel Hadhoud. Parametric spline functions for the solution of the one time fractional burgers' equation. Applied Mathematical Modelling, 36(10):4557-4564, 2012.
- [10] Alaattin Esen and Orkun Tasbozan. Numerical solution of time fractional burgers equation. Acta Universitatis Sapientiae, Mathematica, 7(2):167–185, 2015.
- [11] Massimiliano Giona and Hector Eduardo Roman. Fractional diffusion equation for transport phenomena in random media. *Physica A: Statistical Mechanics and its Applications*, 185(1-4):87–97, 1992.
- [12] Loukas Grafakos. Modern fourier analysis, volume 250. Springer, 2009.
- [13] István Gyöngy. Existence and uniqueness results for semilinear stochastic partial differential equations. Stochastic Processes and their Applications, 73(2):271–299, 1998.
- [14] István Gyöngy and David Nualart. On the stochastic Burgers' equation in the real line. The Annals of Probability, 27(2):782–802, 1999.
- [15] Beom-Seok Han. L_p -regularity theory for semilinear stochastic partial differential equations with multiplicative white noise. Journal of Mathematical Analysis and Applications, page 126366, 2022.
- [16] Beom-Seok Han. A regularity theory for stochastic generalized Burgers' equation driven by a multiplicative space-time white noise. Stochastics and Partial Differential Equations: Analysis and Computations, pages 1–41, 2022.
- [17] Richard Herrmann. Fractional calculus: an introduction for physicists. World Scientific, 2011.

- [18] Rudolf Hilfer. Applications of fractional calculus in physics. World scientific, 2000.
- [19] Mitsunojo Ichise, Yutaka Nagayanagi, and Tsugio Kojima. An analog simulation of non-integer order transfer functions for analysis of electrode processes. Journal of Electroanalytical Chemistry and Interfacial Electrochemistry, 33(2):253–265, 1971.
- [20] Mustafa Inc. The approximate and exact solutions of the space-and time-fractional Burgers equations with initial conditions by variational iteration method. *Journal of Mathematical Analysis and Applica*tions, 345(1):476–484, 2008.
- [21] Anatoly Kilbas, Oleg Marichev, and Stefan Samko. Fractional integrals and derivatives (theory and applications), 1993.
- [22] Ildoo Kim, Kyeong-Hun Kim, and Sungbin Lim. An $L_q(L_p)$ -theory for the time fractional evolution equations with variable coefficients. Advances in Mathematics, 306:123–176, 2017.
- [23] Ildoo Kim, Kyeong-Hun Kim, Sungbin Lim. A sobolev space theory for stochastic partial differential equations with time-fractional derivatives. *The Annals of Probability*, 47(4):2087–2139, 2019.
- [24] Kyeong-Hun Kim and Sungbin Lim. Asymptotic behaviors of fundamental solution and its derivatives related to space-time fractional differential equations. *Journal of the Korean Mathematical Society*, 53(4):929–967, 2015.
- [25] Kyeong-Hun Kim and Daehan Park. A sobolev space theory for the time-fractional stochastic partial differential equations driven by levy processes. arXiv preprint arXiv:2006.05050, 2020.
- [26] Nicolai Krylov. An analytic approach to SPDEs. Stochastic partial differential equations: six perspectives, 64:185–242, 1999.
- [27] Nicolai Krylov. Lectures on elliptic and parabolic equations in Sobolev spaces, volume 96. American Mathematical Society, 2008.
- [28] Nicolai Krylov. Introduction to the theory of random processes, volume 43. American Mathematical Society, 2002.
- [29] Peter Lewis and David Nualart. Stochastic Burgers' equation on the real line: regularity and moment estimates. *Stochastics*, 90(7):1053–1086, 2018.
- [30] Dongfang Li, Chengjian Zhang, and Maohua Ran. A linear finite difference scheme for generalized time fractional Burgers' equation. Applied Mathematical Modelling, 40(11-12):6069–6081, 2016.
- [31] Richard Magin. Fractional calculus in bioengineering, part 1. Critical Reviews[™] in Biomedical Engineering, 32(1), 2004.
- [32] Igor Podlubny. Fractional differential equations: an introduction to fractional derivatives, fractional differential equations, to Methods of their solution and some of their applications. ISSN. Elsevier Science, 1998.
- [33] Hun H Sun, Banu Onaral, and Yuan-Ying Tso. Application of the positive reality principle to metal electrode linear polarization phenomena. *IEEE Transactions on Biomedical Engineering*, (10):664–674, 1984.
- [34] Hans Triebel. Theory of Function Spaces. Modern Birkhäuser Classics. Springer Basel, 2010.
- [35] John Walsh. An introduction to stochastic partial differential equations. In École d'Été de Probabilités de Saint Flour XIV-1984, pages 265–439. Springer, 1986.
- [36] Guang-an Zou and Bo Wang. Stochastic Burgers' equation with fractional derivative driven by multiplicative noise. Computers & Mathematics with Applications, 74(12):3195–3208, 2017.

DEPARTMENT OF MATHEMATICS, POHANG UNIVERSITY OF SCIENCE AND TECHNOLOGY, 77, CHEONGAM-RO, NAM-GU, POHANG, GYEONGBUK, 37673, REPUBLIC OF KOREA

Email address: hanbeom@postech.ac.kr