Tribaryons with lattice QCD and one-boson exchange potentials

Tian-Wei Wu, ^{1,2} Si-Qiang Luo, ^{3,4} Ming-Zhu Liu, ^{5,6} Li-Sheng Geng, ^{6,7,8,4,9,*} and Xiang Liu^{3,10,4,11,†}

¹School of Science, Shenzhen Campus of Sun Yat-sen University, Shenzhen 518107, China

²School of Fundamental Physics and Mathematical Sciences,

Hangzhou Institute for Advanced Study, UCAS, Hangzhou 310024, China

³School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China

⁴Lanzhou Center for Theoretical Physics, Lanzhou University, Beijing 102206, China

⁵School of Space and Environment, Beihang University, Beijing 102206, China

⁶School of Physics, Beihang University, Beijing 102206, China

⁷Peng Huanwu Collaborative Center for Research and Education, Beihang University, Beijing 100191, China

⁸Beijing Key Laboratory of Advanced Nuclear Materials and Physics, Beihang University, Beijing 102206, China

⁹Southern Center for Nuclear-Science Theory (SCNT), Institute of Modern Physics,

Chinese Academy of Sciences, Huizhou 516000, China

¹⁰Research Center for Hadron and CSR Physics, Lanzhou University and Institute of Modern Physics of CAS, Lanzhou 730000, China

(¶Dated: November 23, 2023)

Motivated by the existence of two-body hadronic molecules composed of $\Omega\Omega$, $\Omega_{ccc}\Omega_{ccc}$ and $\Omega_{bbb}\Omega_{bbb}$ predicted by lattice QCD simulations, we use the Gaussian expansion method to investigate whether three-body systems composed of $\Omega\Omega\Omega$, $\Omega_{ccc}\Omega_{ccc}\Omega_{ccc}$ and $\Omega_{bbb}\Omega_{bbb}\Omega_{bbb}$ can bind with the two-body 1S_0 interactions provided by lattice QCD. Our results show that none of the three-body systems bind. On the other hand, we find that with the one-boson exchange potentials the $\Omega\Omega\Omega$ system develops a bound state, for which the 5S_2 interaction plays an important role. Our studies support the existence of the $\frac{3}{2}^+$ $\Omega\Omega\Omega$ bound state and the nonexistence of the $\frac{3}{2}^+$ $\Omega_{ccc}\Omega_{ccc}\Omega_{ccc}$ and $\Omega_{bbb}\Omega_{bbb}\Omega_{bbb}$ bound states, due to the suppressed 5S_2 interactions in heavier systems.

Introduction.—The quark model, as a classification scheme for light-flavor hadrons, was proposed by Gell-Mann [1] and Zweig [2] in 1964, which was established when the predicted Ω baryon with the highest strangeness number was observed experimentally [3]. It is often viewed as the first stage in hadron physics. Since 2003, we have witnessed a new stage in hadron physics with the observation of many new hadronic states, such as the charmoniumlike XYZ states and the pentaquark states [4-12], which have stimulated extensive studies, both theoretically and experimentally. Although remarkable progress has been made, a unified understanding of exotic hadronic states is still missing. At present, it is widely acknowledged that one should pay more attention to new configurations, exotic quantum numbers, and special systems in order to better understand the nature of exotic hadronic matter and the nonperturbative strong interaction.

In recent years, fully strange and fully heavy dibaryon systems have attracted considerable attention. With increasing computational power, lattice QCD has become the primary force to derive hadron-hadron interactions in a quantitative way from first principles. In Ref. [13], the authors investigated the $\Omega\Omega$ interaction in the 1S_0 channel, and concluded that there exists a weakly bound state regardless of the Coulomb interaction, which is even shallower than the deuteron. In Ref. [14], the existence of a 1S_0 $\Omega_{ccc}\Omega_{ccc}$ shallow bound state is predicted while it disappears once the Coulomb interaction is taken into account. Very recently, the existence of a deeply bound 1S_0 $\Omega_{bbb}\Omega_{bbb}$ state was also predicted [15]. For the $\Omega\Omega$, $\Omega_{ccc}\Omega_{ccc}$, and $\Omega_{bbb}\Omega_{bbb}$ systems, some of us developed an extended one-boson-exchange (OBE) model to derive their interactions in Ref. [16], and obtained results consistent with

those of lattice QCD [13–15]. In Ref. [17], the authors found the existence of fully heavy dibaryon bound states, $\Omega_{ccc}\Omega_{ccc}$ and $\Omega_{bbb}\Omega_{bbb}$, in the constituent quark model, while the corresponding fully heavy hexaquark states are found to be above the $\Omega_{ccc}\Omega_{ccc}$ and $\Omega_{bbb}\Omega_{bbb}$ mass thresholds in both the constituent quark models [18–20] and the QCD sum rules [21].

On the experimental side, studies of fully heavy multiquarks have made important breakthroughs. In 2020, the LHCb Collaboration reported the observation of the first fully heavy tetraquark state, X(6900) [22]. It was later confirmed by the CMS Collaboration with a statistical significance of 9.4σ , and in addition, two new states X(6600) and X(7200) were observed [23]. The ATLAS Collaboration further confirmed the discovery of the LHCb Collaboration [24]. Clearly, the existence of fully heavy multiquark states can be considered as firmly established.

It is a plausible expectation that tribaryon systems exist, given the predicted existence of dibaryon systems. We note that experimental and theoretical studies of tribaryon systems other than atomic nuclei and hypernuclei have continued for many years without conclusive results [25–46]. For example, in Refs. [27–31], the authors studied the possible existence of nonstrange tribaryons including NNN [28, 31], $NN\Delta$ [28, 31], $N\Delta\Delta$ [29, 31] and $\Delta\Delta\Delta$ [27, 30, 31] systems with the relevant two-body potentials. The strange tribaryons have also been studied, including single strangeness ΛNN and ΣNN [32–34], double strangeness $\Lambda \Lambda N$ and ΣNN [35–39], and multi strangeness $N\Xi\Xi$ [39], ΣNN and ΣNN [35–39], and multi strangeness $\Sigma N\Xi$ [39], ΣNN and ΣNN and ΣNN and ΣNN systems, the ΣNN potential derived from lattice QCD simulations, which can form bound states [47], plays

an important role [40, 41, 48]. On the experimental side, a strange tribaryon $S^0(3115)$ was reported in the ⁴He (stopped K^- , p) reaction, which mainly decays into ΣNN [49]. In Ref. [42], this strange tribaryon is explained as a nonaquark state. Other searches for strange tribaryons have also been performed [43–46].

In this paper, motivated by the remarkable progress achieved on studies of the fully heavy multiquark states from both lattice QCD [13–15] and experiments [22–24], we study the $\Omega\Omega\Omega$ system, as well as the $\Omega_{ccc}\Omega_{ccc}\Omega_{ccc}$ and $\Omega_{bbb}\Omega_{bbb}\Omega_{bbb}$ systems. Notice that this study differ from the previous works. Regarding the works we mentioned above, they have either different species of baryons (ΩNN , $\Omega\Omega N$, ...) or different flavors or charges $(pnn, \Delta\Delta\Delta, \Xi\Xi\Xi, ...)$. To the best of our knowledge, it is the first time that three-body systems that are composed of fully identical flavored baryons have been studied. The systems we study contain only one species of baryons that are composed of only one species of quarks and, therefore, have the highest symmetries. This means that we need only the interactions of a pair of identical baryons and the number of allowed configurations is much reduced as well, thus allowing for more robust predictions.

Most strange tribaryon.—We adopt the Gaussian expansion method (GEM) [50–52] to study the $\Omega\Omega\Omega$ system. To solve the Schrödinger equation with GEM, one needs to derive the two-body interactions and construct the three-body wave functions. We note that three-body interactions may play an important role in many-body systems, such as the nucleus. Unfortunately, no empirical information on the three-body interactions is available for the three identical baryons we have studied. Thus, in this exploratory work, we only consider two-body interactions.

The $\Omega\Omega$ interaction has been derived in lattice QCD [13], where it was shown that the S-wave $\Omega\Omega$ system can bind with a binding energy of $1.6(6)_{-0.6}^{+0.7}$ MeV (without taking into account the Coulomb interaction). In addition to lattice QCD, other methods such as the extended OBE model can also provide the $\Omega\Omega$ interaction [16]. The light meson exchange, including the pseudoscalar(π), scalar(σ) and $vector(\rho, \omega)$ mesons, can well describe many hadron-hadron interactions, which is naively extended to the $\Omega_{ccc}\Omega_{ccc}$ system by invoking the exchange of the charmonium states η_c , χ_{c0} and J/ψ . The couplings between Ω_{ccc} and the charmonium states are assumed to be proportional to the couplings between the nucleon and the light mesons utilizying the quark model. We emphasize that although the OBE model constructed in this way suffers from relatively large uncertainties, these can be minimized by fitting to the lattice QCD binding energies of the $\Omega\Omega$, $\Omega_{ccc}\Omega_{ccc}$, and $\Omega_{bbb}\Omega_{bbb}$ bound states. In this work, we utilize both interactions to study the $\Omega\Omega\Omega$ three-body system, see Fig. 1 for the potentials.

The $\Omega\Omega$ lattice QCD potentials for the 1S_0 channel are expressed with three Gaussian functions $V_L^{^1S_0}(r)=\sum_{i=1}^3 a_i e^{-b_i r^2}$ [13]. Since the Ω baryon is charged, the Coulomb interaction plays an important role in the $\Omega\Omega$ and




FIG. 1. OBE and lattice QCD potentials for the $\Omega\Omega$ system. The blue dashed, orange dashed and green solid lines denote the 1S_0 OBE, the 5S_2 OBE and the 1S_0 lattice QCD potentials, respectively.

 $\Omega\Omega\Omega$ systems. The Coulomb potential between a pair of $\Omega\Omega$ is $V_C(r) = -\alpha/r$, where $\alpha = 1/137$ is the electromagnetic fine structure constant. The lattice QCD simulations provided only the 1S_0 potential between the $\Omega\Omega$ pair. As we see later, the 5S_2 potential plays an important role in the three-body system as well.

In GEM, a three-body system is studied by solving the three-body Schrödinger equation with the three-body wave functions and the Hamiltonian in Jacobi coordinates. For the $\Omega\Omega\Omega$ three-body system, the Schrödinger equation is as follows

$$[T + V_{\Omega\Omega}(r_1) + V_{\Omega\Omega}(r_2) + V_{\Omega\Omega}(r_3) - E]\Psi_J(\vec{r_c}, \vec{R_c}) = 0, (1)$$

where c=1-3 denote the three Jacobi channels, $r_c(R_c)$ are the Jacobi coordinates. T is the kinetic-energy operator and $V_{\Omega\Omega}$ is the two-body $\Omega\Omega$ interaction. For the details on how to construct the Jacobi coordinates and the three-body kinetic-energy operator, please refer to Ref. [51].

The $\Omega\Omega\Omega$ three-body wave function can be written as a sum of three Jacobi channels

$$\Psi_J(\vec{r}_c, \vec{R}_c) = \sum A_\alpha^c \Phi_J^c(\vec{r}_c, \vec{R}_c), \tag{2}$$

where A^c_{α} is the expansion coefficients and α is the set of quantum numbers characterizing the wave function in each Jacobi channel. The wave function of each Jacobi channel reads as

$$\begin{split} &\Phi_J^1(\vec{r}_1,\vec{R}_1) = \left[[[\chi_3\chi_2]_{s_1}\chi_1]_S \otimes [\psi_{l_1}(\vec{r}_1)\phi_{L_1}(\vec{R}_1)]_\Lambda \right]_J, \\ &\Phi_J^2(\vec{r}_2,\vec{R}_2) = \left[[[\chi_1\chi_3]_{s_2}\chi_2]_S \otimes [\psi_{l_2}(\vec{r}_2)\phi_{L_2}(\vec{R}_2)]_\Lambda \right]_J, \\ &\Phi_J^3(\vec{r}_3,\vec{R}_3) = \left[[[\chi_2\chi_1]_{s_3}\chi_3]_S \otimes [\psi_{l_3}(\vec{r}_3)\phi_{L_3}(\vec{R}_3)]_\Lambda \right]_J, \end{split}$$

where χ_i is the spin wave function of the *i*th particle, $H_{s,S}^c = [[\chi_i \chi_j]_s \chi_k]_S$ is the spin wave function of Jacobi channel c,

 $\psi(r_i)\phi(R_i)$ is the spatial wave function, s is the spin of the sub- $\Omega\Omega$ two-body system, S=3/2 is the total spin of $\Omega\Omega\Omega$, l_i (L_i) is the orbit angular momentum corresponding to $r_i(R_i)$, Λ is the total orbit angular momentum built from l and L, and J is the total angular momentum built from Λ and S.

Fermi-Dirac statistics dictates that only the 1S_0 and 5S_2 interactions contribute to the formation of an $\Omega\Omega\Omega$ $\frac{3}{2}^+$ state. The spin coupling coefficients of different spin configurations between Jacobi channels i and j for $i \neq j$ are shown in Table I. Note that for i = j, the matrix is orthogonal.

TABLE I. Coupling coefficients of different spin configurations between Jacobi channels i and j ($i \neq j$). Here, $H_{s,S}^c$ is the spin function, $s = \{0,2\}$ are alternative spin values of $\Omega\Omega$, and S = 3/2 is the total spin of $\Omega\Omega\Omega$.

	$H^{c=i}_{0,\frac{3}{2}}$	$H_{2,\frac{3}{2}}^{c=i}$
$H_{0,\frac{3}{2}}^{c=j}$	$-\frac{1}{4}$	$-\frac{\sqrt{5}}{4}$
$H^{c=j}_{2,\frac{3}{2}}$	$-\frac{\sqrt{5}}{4}$	$\frac{3}{4}$

It is important to point out that for the $\Omega\Omega\Omega$ system, the 5S_2 potential can play a very important role, even more important than the 1S_0 potential. This is because the 5S_2 partial wave is more strongly coupled to the three-body spin-3/2 state than the 1S_0 partial wave. As shown in Table I, the spin coupling coefficient of different Jacobi channels i and j in the 5S_2 partial wave is $\langle H_{2,3/2}^{c=i}|H_{2,3/2}^{c=j}\rangle_{i\neq j}=3/4$ while that in 1S_0 is $\langle H_{0,3/2}^{c=i}|H_{0,3/2}^{c=j}\rangle_{i\neq j}=-1/4$, which means that in the spin space, the coupling between channels i and j in the 5S_2 partial wave is 9 times larger than that in the 1S_0 partial wave.

Once the wave functions are obtained , with either the lattice QCD or OBE $\Omega\Omega$ interactions, one can adopt the GEM [51] to obtain the binding energies and root-mean-square (rms) radius of the $\Omega\Omega\Omega$ system.

The results for the two-body $\Omega\Omega$ system are summarized in Table II, which show that the binding energies and rms radii obtained with the OBE potentials are consistent with those of lattice QCD. With both lattice QCD and OBE potentials, the $\Omega\Omega$ system can bind with a binding energy of $1.4^{+0.9}_{-0.4}$ MeV. The uncertainties are determined by multiplying a scaling factor to the lattice QCD potential so that the binding energy varies from 1.0 to 2.3 MeV, consistent with the lattice QCD result $1.6^{+0.7}_{-0.6}$ MeV [13].

From the analysis given above, we know that both 1S_0 and 5S_2 interactions contribute to the $3/2~\Omega\Omega\Omega$ system. Given that the lattice QCD provided only the 1S_0 interaction, we first consider only the 1S_0 two-body interaction and find that the $\Omega\Omega\Omega$ system does not bind. But this result should not be taken too seriously since the 5S_2 partial wave plays an important role in the spin configuration $(\langle H^{c=i}_{2,3/2}|H^{c=j}_{2,3/2}\rangle_{i\neq j}=\frac{3}{4})$ and has a significant correlation with the 1S_0 partial wave $(\langle H^{c=i}_{0,3/2}|H^{c=j}_{2,3/2}\rangle_{i\neq j}=\frac{\sqrt{5}}{4})$ in the three-body case. Actually, with the OBE 1S_0 and 5S_2 potentials, we find that the three-

body $\Omega\Omega\Omega$ system binds with a binding energy of $5.8^{+2.5}_{-1.2}$ MeV and rms radius $1.9^{+0.1}_{-0.2}$ fm.

Note that the binding energy per baryon of the $\Omega\Omega$ system is larger than that of the $\Omega\Omega$ system, and consequently its rms radius is smaller than that of the $\Omega\Omega$ bound state. This is understandable because for the $\Omega\Omega\Omega$ system the 5S_2 potential plays an important role, while only the 1S_0 potential is relevant for the $\Omega\Omega$ system.

The weights of partial waves and Hamiltonian expectation values of the predicted $\Omega\Omega\Omega$ bound state are given in Table III, which clearly show that the 5S_2 interaction plays a significantly important role in the $\Omega\Omega\Omega$ system. More specifically, the weights of the 1S_0 and 5S_2 partial waves are about 22% and 78%, respectively.

As we mentioned above, since the Ω baryon is charged, the impact of the Coulomb interaction is worth discussing. We find that the Coulomb interaction in this three-body system affects the binding energy by about 2-3 MeV but does not change the conclusion. Considering the Coulomb interaction, the binding energy and rms radius of the $\Omega\Omega\Omega$ bound state predicted by the OBE model are 2.0 MeV and 2.3 fm, respectively.

It is important to discuss where to search for the predicted $\Omega\Omega$ and $\Omega\Omega\Omega$ bound states. In Ref. [48], the production yield of the $\Omega\Omega$ bound state was estimated using a dynamical coalescence mechanism for the relativistic heavy-ion collisions at $\sqrt{s_{NN}}=200$ GeV and 2.76 TeV, which turn out to be of the order of 10^{-6} . In Ref. [41], the production yields of $NN\Omega$ and $N\Omega\Omega$ were estimated to be 10^{-7} and 10^{-9} , respectively. Comparing these results, one can estimate the $\Omega\Omega\Omega$ production rate for the relativistic heavy-ion collisions at $\sqrt{s_{NN}}=200$ GeV and 2.76 TeV, which is of the order of 10^{-11} .

TABLE II. Binding energies (BE) and root-mean-square radii ($\langle r \rangle$) of the $\Omega\Omega$ and $\Omega\Omega\Omega$ bound states obtained with lattice QCD (with only 1S_0) and OBE potentials (with both 1S_0 and 5S_2). BE in MeV and radius $\langle r \rangle$ in fm.

	ΩΩ(ΒΕ)	$\Omega\Omega(\langle r \rangle)$	ΩΩΩ(ΒΕ)	$\Omega\Omega\Omega(\langle r \rangle)$
LQCD	$1.41^{+0.89}_{-0.41}$	$3.45^{+0.52}_{-0.62}$		
OBE	$1.41^{+0.89}_{-0.41}$	$3.33^{+0.51}_{-0.62}$	$5.84^{+2.48}_{-1.22}$	$1.86^{+0.13}_{-0.19}$

TABLE III. Weights of the partial waves and Hamiltonian expectation values (units in MeV) of the $\frac{3}{2}^+$ $\Omega\Omega\Omega$ bound state.

	$\langle \Psi_{3/2}^{^{1}S_{0}} \Psi_{3/2}^{^{1}S_{0}} angle$	$\langle \Psi_{3/2}^{^{5}S_{2}} \Psi_{3/2}^{^{5}S_{2}} angle$	$\langle T \rangle$	$\langle V^{^1S_0} \rangle$	$\langle V^{^5S_2} \rangle$
OBE	22%	78%	52.72	-15.71	-42.84

Most charming and beautiful tribaryons.—It is straightforward to extend the above study to the $\Omega_{ccc}\Omega_{ccc}$ and $\Omega_{bbb}\Omega_{bbb}$ systems, for which the lattice QCD simulations already provided




FIG. 2. OBE and lattice QCD potentials of the $\Omega_{ccc}\Omega_{ccc}$ (top) and $\Omega_{bbb}\Omega_{bbb}$ (bottom) systems. The blue dashed, orange dashed and green solid lines denote the 1S_0 OBE, the 5S_2 OBE and the 1S_0 lattice QCD potentials, respectively.

the 1S_0 potentials [14, 15] and their OBE counterparts also exist [16]. Note that in Ref. [15] no analytic form of the $\Omega_{bbb}\Omega_{bbb}$ potential was provided. We fitted the lattice QCD potential with a sum of three Gaussian functions as done in Ref. [14]. All the lattice QCD potentials and the corresponding OBE potentials are shown in Fig. 2. We note that although the interaction strengths of the lattice QCD potential and those of the OBE potentials are different, the positions where they become the most attractive are almost the same. The same can be said about the $\Omega\Omega$ potentials shown in Fig. 1. Such a coincidence indicates that the OBE model must have captured some essential features of the baryon-baryon potentials.

With the above lattice QCD and the OBE potentials, we can study the two-body and three-body systems composed of Ω_{ccc} and Ω_{bbb} . As shown in Table IV, with only the strong interaction the $\Omega_{ccc}\Omega_{ccc}$ bound system can be formed, but it dissolves once the Coulomb interaction is taken into account. On the other hand, the $\Omega_{bbb}\Omega_{bbb}$ system is always bound regardless of the Coulomb interaction. Furthermore, we note that

the results obtained with the lattice QCD potentials and those with the OBE potentials are similar. Nonetheless, none of the $\Omega_{ccc}\Omega_{ccc}\Omega_{ccc}$ and $\Omega_{bbb}\Omega_{bbb}\Omega_{bbb}$ three-body systems can bind, mainly because of the much weaker 5S_2 interactions, which are nontrivial predictions of the present work.

TABLE IV. Binding energies (BE) and root-mean-square radii ($\langle r \rangle$) of the $\Omega_{ccc}\Omega_{ccc}$ and $\Omega_{bbb}\Omega_{bbb}$ bound states obtained with OBE and LQCD potentials (BE in MeV and radius $\langle r \rangle$ in fm.). NC means that the Coulomb interaction is not taken into account, while C means that the Coulomb interaction is considered.

		$\Omega_{ccc}\Omega_{ccc}$ (NC)	$\Omega_{ccc}\Omega_{ccc}$ (C)	$\Omega_{bbb}\Omega_{bbb}$ (NC)	$\Omega_{bbb}\Omega_{bbb}$ (C)
LQCD	BE	5.54		88.7	79.9
	$\langle r \rangle$	1.14		0.240	0.245
OBE	BE	5.52	• • •	88.6	78.4
	$\langle r \rangle$	1.05		0.198	0.202

Summary.— Motivated by the existence of $\Omega\Omega$, $\Omega_{ccc}\Omega_{ccc}$, and $\Omega_{bbb}\Omega_{bbb}$ bound states predicted by lattice QCD simulations, we studied the $\frac{3}{2}^+$ $\Omega\Omega\Omega$, $\Omega_{ccc}\Omega_{ccc}\Omega_{ccc}$, and $\Omega_{bbb}\Omega_{bbb}\Omega_{bbb}$ three-body systems with the lattice QCD and OBE potentials. We found that the $\Omega\Omega$, $\Omega_{ccc}\Omega_{ccc}$, and $\Omega_{bbb}\Omega_{bbb}$ systems can also bind with the OBE potentials, with binding energies and rms radii consistent with those of lattice QCD simulations. The repulsive Coulomb interactions plays an important role in these systems especially in the $\Omega_{ccc}\Omega_{ccc}$ system, which is strong enough to break the $\Omega_{ccc}\Omega_{ccc}$ pair bound by the strong force.

For the three-body systems, we find that the 5S_2 partial wave plays a very important role in forming the $\frac{3}{2}^+$ three-body state. With only the 1S_0 lattice QCD potentials, the $\Omega\Omega\Omega$, $\Omega_{ccc}\Omega_{ccc}\Omega_{ccc}$, and $\Omega_{bbb}\Omega_{bbb}\Omega_{bbb}$ three-body systems do not bind. With the OBE potentials both in the 1S_0 and 5S_2 partial waves, the $\Omega\Omega\Omega$ system becomes bound, while the $\Omega_{ccc}\Omega_{ccc}\Omega_{ccc}$ and $\Omega_{bbb}\Omega_{bbb}\Omega_{bbb}$ systems remain unbound mainly due to the much suppressed attractive 5S_2 interaction in the two-body $\Omega_{ccc}\Omega_{ccc}$ and $\Omega_{bbb}\Omega_{bbb}$ systems. To verify the existence of the $\Omega\Omega$ bound state, lattice QCD studies of the 5S_2 interactions of the $\Omega\Omega$ system will be the key. We hope that the predicted $\Omega\Omega\Omega$ bound state can be searched for in present and future hadron-hadron colliders.

A particularly interesting discovery of the present work is that even the two-body interactions are attractive and strong enough to form two-body bound states, the three-body systems do not necessarily bind. This is because in three-body systems, spin-spin interactions can play an important role. The three highly symmetric systems studied in the present work provide an ideal platform to understand the relevance of spin-spin interactions in forming few-body bound states. *Acknowledgement.*—This work is partly supported by the National Natural Science Foundation of China under Grants No. 11735003, No. 11975041, No. 11961141004, and the fundamental Research Funds for the Central Universities. X.L.

is supported by the China National Funds for Distinguished Young Scientists under Grant No. 11825503, National Key Research and Development Program of China under Contract No. 2020YFA0406400, the 111 Project under Grant No. B20063, the National Natural Science Foundation of China under Grant No. 12247101, the fundamental Research Funds for the Central Universities, and the project for top-notch innovative talents of Gansu province. M.-Z. L. acknowledges support from the National Natural Science Foundation of China under Grant No. 12105007 and China Postdoctoral Science Foundation under Grants No. 2022M710317, and No. 2022T150036. T.-W. W. acknowledges support from the National Natural Science Foundation of China under Grant No. 12147152 and China Postdoctoral Science Foundation under Grant No. 2022M723119.

- lisheng.geng@buaa.edu.cn xiangliu@lzu.edu.cn
- [1] M. Gell-Mann, Phys. Lett. 8, 214 (1964).
- [2] G. Zweig, "An SU(3) model for strong 2," tion symmetry and its breaking. Version in edited by D. B. Lichtenberg and S. P. Rosen (1964) pp. 22-101.
- [3] V. E. Barnes et al., Phys. Rev. Lett. **12**, 204 (1964).
- [4] N. Brambilla et al., Eur. Phys. J. C 71, 1534 (2011), arXiv:1010.5827 [hep-ph].
- [5] X. Liu, Chin. Sci. Bull. **59**, 3815 (2014), arXiv:1312.7408 [hep-
- [6] H.-X. Chen, W. Chen, X. Liu, and S.-L. Zhu, Phys. Rept. 639, 1 (2016), arXiv:1601.02092 [hep-ph].
- [7] H.-X. Chen, W. Chen, X. Liu, Y.-R. Liu, and S.-L. Zhu, Rept. Prog. Phys. 80, 076201 (2017), arXiv:1609.08928 [hep-ph].
- [8] E. Oset et al., Int. J. Mod. Phys. E 25, 1630001 (2016), arXiv:1601.03972 [hep-ph].
- [9] F.-K. Guo, C. Hanhart, U.-G. Meißner, O. Wang, O. Zhao, and B.-S. Zou, Rev. Mod. Phys. 90, 015004 (2018), [Erratum: Rev.Mod.Phys. 94, 029901 (2022)], arXiv:1705.00141 [hepph].
- [10] Y.-R. Liu, H.-X. Chen, W. Chen, X. Liu, and S.-L. Zhu, Prog. Part. Nucl. Phys. 107, 237 (2019), arXiv:1903.11976 [hep-ph].
- [11] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C.-P. Shen, C. E. Thomas, A. Vairo, and C.-Z. Yuan, Phys. Rept. 873, 1 (2020), arXiv:1907.07583 [hep-ex].
- [12] H.-X. Chen, W. Chen, X. Liu, Y.-R. Liu, and S.-L. Zhu, (2022), arXiv:2204.02649 [hep-ph].
- [13] S. Gongyo et al., Phys. Rev. Lett. 120, 212001 (2018), arXiv:1709.00654 [hep-lat].
- [14] Y. Lyu, H. Tong, T. Sugiura, S. Aoki, T. Doi, T. Hatsuda, J. Meng, and T. Miyamoto, Phys. Rev. Lett. 127, 072003 (2021), arXiv:2102.00181 [hep-lat].
- [15] N. Mathur, M. Padmanath, and D. Chakraborty, (2022), arXiv:2205.02862 [hep-lat].
- [16] M.-Z. Liu and L.-S. Geng, Chin. Phys. Lett. 38, 101201 (2021), arXiv:2107.04957 [hep-ph].
- [17] H. Huang, J. Ping, X. Zhu, and F. Wang, (2020),arXiv:2011.00513 [hep-ph].
- [18] J. M. Alcaraz-Pelegrina and M. C. Gordillo, (2022),arXiv:2205.13886 [hep-ph].

- [19] Q.-F. Lü, D.-Y. Chen, and Y.-B. Dong, (2022),arXiv:2208.03041 [hep-ph].
- [20] X.-Z. Weng and S.-L. Zhu, (2022), arXiv:2207.05505 [hep-ph].
- [21] Z.-G. Wang, (2022), arXiv:2201.02955 [hep-ph].
- [22] R. Aaij et al. (LHCb), Sci. Bull. 65, 1983 (2020), arXiv:2006.16957 [hep-ex].
- [23] Kai Yi on behalf of the CMS Collaboration, Recent CMS results on exotic resonance, Proceedings at ICHEP 2022, https://agenda.infn.it/event/28874/contributions/170300/.
- [24] Evelina Bouhova-Thacker on behalf of LAS Collaboration, **ATLAS** results on exotic hadronic resonances, Proceedings at ICHEP 2022, https://agenda.infn.it/event/28874/contributions/170298/.
- [25] A. Valcarce, H. Garcilazo, F. Fernandez, and P. Gonzalez, Rept. Prog. Phys. 68, 965 (2005), arXiv:hep-ph/0502173.
- [26] A. Park, W. Park, and S. H. Lee, Phys. Rev. D 98, 034001 (2018), arXiv:1801.10350 [hep-ph].
- [27] H. Garcilazo, F. Fernandez, A. Valcarce, and R. D. Mota, Phys. Rev. C 56, 84 (1997).
- [28] H. Garcilazo, A. Valcarce, and F. Fernández, Phys. Rev. C 60, 044002 (1999).
- [29] R. D. Mota, H. Garcilazo, A. Valcarce, and F. Fernandez, Phys. Rev. C 59, 46 (1999).
- [30] A. Valcarce, H. Garcilazo, R. D. Mota, and F. Fernandez, J. Phys. G 27, L1 (2001).
- [31] R. D. Mota, A. Valcarce, F. Fernandez, D. R. Entem, and DEVELOPMENTS IN THE QUARK THEORY OF HADRONS. VOL. 1H196arci 12978, Phys. Rev. C 65, 034006 (2002), arXiv:nuclth/0112059.
 - [32] H. Garcilazo, T. Fernandez-Carames, and A. Valcarce, Phys. Rev. C 75, 034002 (2007), arXiv:hep-ph/0701275.
 - [33] T. Fernandez-Carames, A. Valcarce, H. Garcilazo, and P. Gonzalez, Phys. Rev. C 73, 034004 (2006), arXiv:hep-ph/0601252.
 - [34] H. Garcilazo and A. Valcarce, Symmetry 14, 2381 (2022), arXiv:2211.13970 [nucl-th].
 - [35] H. Garcilazo and A. Valcarce, Phys. Rev. C 92, 014004 (2015), arXiv:1507.03733 [hep-ph].
 - [36] H. Garcilazo, Phys. Rev. C 93, 024001 (2016).
 - [37] H. Garcilazo and A. Valcarce, Phys. Rev. C 93, 034001 (2016), arXiv:1605.04108 [hep-ph].
 - [38] H. Garcilazo and A. Valcarce, Chin. Phys. C 44, 104104 (2020), arXiv:2006.16567 [nucl-th].
 - [39] H. Garcilazo, A. Valcarce, and J. Vijande, Phys. Rev. C 94, 024002 (2016), arXiv:1608.05192 [nucl-th].
 - [40] H. Garcilazo and A. Valcarce, Phys. Rev. C 99, 014001 (2019), arXiv:1901.05678 [hep-ph].
 - [41] L. Zhang, S. Zhang, and Y.-G. Ma, Eur. Phys. J. C 82, 416 (2022), arXiv:2112.02766 [hep-ph].
 - [42] Y. Maezawa, T. Hatsuda, and S. Sasaki, Prog. Theor. Phys. 114, 317 (2005), arXiv:hep-ph/0412025.
 - [43] M. Sato (KEK-PS E471/E549), AIP Conf. Proc. 842, 480 (2006).
 - [44] M. Sato et al., Phys. Lett. B 659, 107 (2008), arXiv:0708.2968 [nucl-ex].
 - [45] M. Sato et al. (KEK-PS E549), Int. J. Mod. Phys. A 24, 442 (2009).
 - [46] H. Yim et al., Phys. Lett. B **688**, 43 (2010).
 - [47] T. Iritani et al. (HAL QCD), Phys. Lett. B 792, 284 (2019), arXiv:1810.03416 [hep-lat].
 - S. Zhang and Y.-G. Ma, Phys. Lett. B 811, 135867 (2020), arXiv:2007.11170 [hep-ph].
 - [49] T. Suzuki et al., Phys. Lett. B 597, 263 (2004).
 - [50] M. Kamimura, Phys. Rev. A 38, 621 (1988).
 - [51] E. Hiyama, Y. Kino, and M. Kamimura, Prog. Part. Nucl. Phys. **51**, 223 (2003).

[52] T.-W. Wu and L.-S. Geng, Few Body Syst. 62, 89 (2021),

arXiv:2105.09703 [hep-ph].