A new insight on positivity and contractivity of the Crank-Nicolson scheme for the heat equation^{*}

I. HIGUERAS¹, <u>T. ROLDÁN¹</u>

¹INSTITUTE FOR ADVANCED MATERIALS AND MATHEMATICS PUBLIC UNIVERSITY OF NAVARRE E-MAILS: higuerasQunavarra.es, teoQunavarra.es.

Abstract

In this paper we study numerical positivity and contractivity in the infinite norm of Crank-Nicolson method when it is applied to the diffusion equation with homogeneous Dirichlet boundary conditions. For this purpose, the amplification matrices are written in terms of three kinds of Chebyshev-like polynomials, and necessary and sufficient bounds to preserve the desired qualitative properties are obtained. For each spatial mesh, we provide the equations that must be solved as well as the intervals that contain these bounds; consequently, they can be easily obtained by a bisection process. Besides, differences between numerical positivity and contractivity are highlighted. This problem has also been addressed by some other authors in the literature and some known results are recovered in our study. Our approach gives a new insight on the problem that completes the panorama and that can be used to study qualitative properties for other problems.

Keywords: Positivity, Maximum norm contractivity, Monotonicity, Crank-Nicolson, Heat equation

1 Introduction

We consider the numerical solution of the one-dimensional heat equation

$$\frac{\partial u(t,x)}{\partial t} = d \,\frac{\partial^2 u(t,x)}{\partial x^2}, \quad x \in [0,1], \, t \ge 0,$$
(1.1)

$$u(0,t) = u(1,t) = 0, \quad t \ge 0,$$
 (1.2)

$$u(x,0) = u_0(x), \quad x \in [0,1].$$
 (1.3)

where u_0 in (1.3) is a given function on [0, 1]. For the sake of simplicity, we consider the diffusion term d = 1 in the rest of the paper. Solutions u(x,t) for the linear parabolic

^{*}Research project PID2019-109045GB-C31 funded by Agencia Estatal de Investigación, Ministerio de Ciencia e Innovación, Spain.

problem (1.1)-(1.3) have several qualitative properties that are relevant in the context of the physical model. In particular, the problem is positivity preserving, that is, for $t \ge 0$,

$$u_0(x) \ge 0 \quad \Rightarrow \quad u(x,t) \ge 0,$$
 (2)

and the solutions are monotonically decreasing in time, i.e., for $t_2 \ge t_1 \ge 0$,

$$\max_{0 \le x \le 1} |u(t_2, x)| \le \max_{0 \le x \le 1} |u(t_1, x)|.$$
(3)

In order to obtain numerical approximations with physical sense, properties (2)-(3) should be preserved in the discretization process. In this paper, we consider the Crank-Nicolson (CN) method, a method of lines approach where second order central finite differences in space are followed by a second order time-stepping method. Spatial discretization of (1.1)-(1.3) with second order central finite differences and mesh width h = 1/(m + 1), gives a semi-discrete linear differential system of the form

$$w'(t) = B_h w(t), \qquad w(0) = w_0, t \ge 0,$$
(4)

where B_h is a matrix of dimension m, that is positivity preserving with monotonically decreasing (in the maximum norm) solutions (see section 2 for details). Next, a time stepping method is used to obtain numerical approximations $w_n \approx w(t_n)$, where $t_n = n\tau$, and τ is the constant time stepsize used. In this paper, we consider approximations of the form

$$w_n = A_m w_{n-1}, n \ge 0,$$
 (5)

where w_0 is a known value, and A_m is a matrix of dimension m that depends on the time stepping method. In particular, for Runge-Kutta methods, $A_m = \phi(\tau B_h)$, where ϕ is the stability function of the scheme.

There is a vast list of references in the analysis of positivity and monotonicity decreasing time stepping schemes (see, e.g., [21, 26, 14, 16, 11, 4, 8, 12, 2, 6, 18]; see too [26, 6, 13] and the references therein). Depending on the context, monotonicity decreasing methods are also known as contractive, SSP (Strong Stability Preserving) or TVD (Total Variation Diminishing) schemes (see, e.g., [23, 21, 14, 8, 4, 6]). In this setting, stepsize restrictions of the form

$$\tau \le C \,\tau_{_{FE}}\,,\tag{6}$$

are obtained, where C denotes the monotonicity threshold factor (also known as SSPcoefficient, radius of absolute monotonicity, contractivity radius, ...) of the time stepping method (see e.g., [15, 27, 5, 23, 4]), and τ_{FE} is the stepsize restriction for the given qualitative property when forward Euler scheme is used to solve the specific ODE problem (see, e.g. [23, p. 379], [8, p. 201], [6, pp. 52-53]). In particular, for problem (4), $\tau_{FE} = h^2/2$ for both positivity and contractivity (see, e.g., [26, p. 22]), and C = 1 for forward Euler method (see section 2 for details). Observe that with this approach, the stepsize restriction (6) is the same for all linear systems with the same τ_{FE} and for some problems this is not a sharp bound.

A well known time stepping method is the θ -method, defined as

$$w_n = (I - \theta \tau B_h)^{-1} (I + (1 - \theta) \tau B_h) w_{n-1}, \qquad \theta \in [0, 1].$$
(7)

Observe that this method can be understood as the composition of a $(1 - \theta)\tau$ -step with forward Euler method and a $\theta\tau$ -step with backward Euler scheme. In particular, for $\theta = 1/2$, CN scheme is obtained. The radius of absolute monotonicity for the θ -method applied to linear problems is $C_{\theta} = 1/(1 - \theta)$ [9]. Consequently, numerical positivity and contractivity can be ensured under the restriction (see [26, p. 135])

$$\frac{\tau}{h^2} \le \frac{1}{2(1-\theta)} \,.$$

However, If we look closer at the iteration matrix A_m in (7), a sharper bound is possible. Different authors have studied numerical preservation of positivity and monotonicity for problem (1.1)-(1.3) with the θ -method (see, e.g., [16, 11, 2]). In [2, p. 72], the authors obtain that the numerical solution is positive if and only if

$$\frac{\tau}{h^2} \le \frac{1 - \sqrt{1 - \theta}}{\theta (1 - \theta)} \,, \tag{8}$$

whereas in [16, Remark 7.1] and [11, p. 456] it is shown that the numerical solution is contractive if and only if

$$\frac{\tau}{h^2} \le \frac{2-\theta}{4(1-\theta)^2} \,. \tag{9}$$

On the following we will denote $s = \tau/h^2$ to the CFL coefficient. The contractivity result in [16] is obtained for the pure initial value problem

$$\frac{\partial u(t,x)}{\partial t} = d \frac{\partial^2 u(t,x)}{\partial x^2}, \quad x \in \mathbb{R}, t \ge 0,$$
$$u(x,0) = u_0(x), \quad x \in \mathbb{R}.$$

In particular, bound (9) for contractivity is valid for all $m \ge 1$ [16, Theorem 4.1] [11]. The bound $\tau/h^2 \le 2/3$ has also been obtained in [3, Theorem 1] in the analysis of the stability of CN method. In [2] and [11] results are based on the shape of the inverse matrix $(I - \theta \tau B_h)^{-1}$ of dimension m, whose entries can be expressed in terms of hyperbolic functions [20]; stepsize restrictions are obtained for each value of m, and bound (9) is valid for all m.

Contributions of the paper

The approach followed in this paper consists on the representation of the iteration matrix A_m in (5) for the CN method in terms of three classes of polynomials, $P_n(x)$, $C_n(x)$ and $U_n(x)$, defined by iterations (26), (28) and (33), respectively; in particular, $U_n(x)$ are the Chebyshev polynomials of the second kind. In the three cases the iteration process is the same, but different initial values are considered. This formalism gives a new insight on the problem that allows us to improve some results in the literature.

The contributions of this paper are the following ones. With regard to positivity, for any number of grid points m, we have obtained that:

1. Crank-Nicolson method is positive if and only if $s = \tau/h^2 \in (0, s_m^{(p)}]$, with $s_m^{(p)} = 1/(\cosh \omega_m^{(p)} - 1)$, where $\omega_m^{(p)}$ is the unique positive root of equation (18). This root

lies in the narrow interval $(\log(2 + \sqrt{2}), \log(2 + \sqrt{3})] \approx (1.22795, 1.31696]$. Thus $\omega_m^{(p)}$ can be easily computed by solving (18) with bisection method. Proposition 2 shows the connection between the bounds $s_m^{(p)}$ and polynomials $P_n(x)$. Some values of $s_m^{(p)}$ are shown in Table 1.

- 2. The sequence of bounds $(s_m^{(p)})$ is strictly monotonically increasing with all the terms in the interval $[1, 2(2 \sqrt{2}))$. As a consequence, the CN method does not preserve positivity when the spatial mesh is refined (keeping s constant).
- 3. In the limit case, when m tends to infinite, we recover the known bound, $s \leq 1.17$ for positivity ([13, p. 126],[2, Table 1]).

With regard to contractivity, we have computed the value $||A_m(s)||_{\infty}$ for any number of grid points m (see Figure 3), and we have obtained that:

- 1. Crank-Nicolson method is contractive if and only if $s = \tau/h^2 \in (0, s_m^{(c)}]$, with $s_m^{(c)} = \infty$ for m = 1, 2, 3. For $m \ge 4$, $s_m^{(c)} = 1/(\cosh \omega_m^{(c)} 1)$, where $\omega_m^{(c)}$ is the unique positive root of equation (21) or (22), depending on the parity of m. This root lies in the interval $[\log ((3+\sqrt{5}+\sqrt{-2}+6\sqrt{5})/4), \log 3) \approx (0.767197, 1.09861]$. Thus $\omega_m^{(c)}$ can be easily computed by solving (21) or (22) with the bisection method. Some values of $s_m^{(p)}$ are shown in Tables 2 and 3 for odd and even values of m, respectively. Some of these values can also be seen in Figure 3.
- 2. $||A_m(s)||_{\infty} < 1$ for $s \in (0, s_m^{(c)})$ and $||A_m(s_m^{(c)})||_{\infty} = 1$ (see Figure 3), that resembles property (14) of the linear system (10).
- 3. For $m \ge 4$, the sequence $(s_m^{(c)})$ is strictly monotonically decreasing with all the terms in the interval $(3/2, 1 + \sqrt{5}]$. As a consequence, CN method preserves contractivity when the spatial mesh is refined (keeping s constant).
- 4. In the limit case, when m tends to infinite, we recover the known bound, $\tau/h^2 \leq 3/2$ for contractivity [16, Th. 4.1(Q3); Section 7.1], [11, Eq. (14)].

The results in this paper complete and improve some results in the literature [2, 11]. From equations (18) and (21)-(22), and the associated intervals, the computation of $s_m^{(p)}$ and $s_m^{(c)}$ for any value of m is straightforward with the bisection method. Besides, we obtain equations to compute bounds $s_m^{(c)}$ both for odd and even values of m, whereas in [11], bounds $s_m^{(c)}$ are only given for even m. From our approach we also get the correct value for $s_3^{(c)}$. Figure 1 illustrates the differences between positivity and contractivity of CN method for the heat problem (1): if the scheme is positive for a given grid mesh m, then it is also contractive for any grid mesh.

Scope of the paper

The rest of the paper is organized as follows. In Section 2, we explain the CN discretization process; notation and definitions are also given in this section. In Section 3 we show the main results of the paper, namely: Theorems 1 and 2; Table 1, containing upper bounds

 $s_m^{(p)}$ for positivity; Tables 2 and 3 containing upper bounds $s_m^{(c)}$ for contractivity (odd and even case); and Figure 1 showing sequences $(s_m^{(p)})$ and $(s_m^{(c)})$. An illustrative example is also given in Section 3. Section 4 contains some conclusions and ideas for future work. The proof of main results are given in Section 6. Previously, some technical material, needed for the proofs in Section 6, is included in Section 5.

2 Crank-Nicolson method for the heat equation

In this paper we consider the Crank-Nicolson method, a method of lines approach where second order central finite differences in space are followed by a second order time-stepping method. Spatial discretization of heat equation (1) with second order central finite differences and mesh width h = 1/(m + 1), gives the semi-discrete linear differential system

$$w'(t) = B_h w(t), \qquad w(0) = w_0, t \ge 0,$$
(10)

where $B_h = (d/h^2)$ tridiag(1, -2, 1) is a matrix of dimension m, $w(t) \approx (u(x_i, t))_{i=1}^m$, $w_0 = (u_0(x_i))_{i=1}^m$, and $x_i = ih$, i = 1, ..., m, are the grid points.

As the diffusion problem (1.1)-(1.3) is positivity preserving (2) and monotonically decreasing (3), in order to obtain numerical approximations with these qualitative properties, problem (10) should also be positivity preserving and contractive in the maximum norm.

An initial value problem

$$w'(t) = f(t, w(t)), \qquad w(t_0) = w_0, t \ge 0,$$
(11)

is called positivity preserving (positive for short) if $w_0 \ge 0$ implies that $w(t) \ge 0$ for $t \ge 0$, where the inequalities should be understood component-wise. Problem (11) is said to be contractive in the maximum norm if its solution w(t) satisfy

$$||w(t_2)||_{\infty} \le ||w(t_1)||_{\infty}$$
 for $t_2 \ge t_1 \ge 0$.

It is well known that a linear problem,

$$w'(t) = Aw(t), \qquad w(t_0) = w_0, t \ge 0,$$
(12)

where $A = (a_{ij})$ is an $m \times m$ matrix, is positive if and only if $a_{ij} \ge 0$ for all $i \ne j$ [13, Theorem 7.2]. Matrix B_h in (10) satisfies this condition and thus problem (10) is positive. Observe that other spatial discretizations do not preserve positivity; indeed, there is an order barrier ($q \le 2$) from the requirement of positivity [13, p. 119].

Contractivity of solutions of the linear problem (12) can be proven by using the concept of logarithmic norm of matrix A. This concept is an extremely useful tool to analyze the growth of solutions to ordinary differential equations because it can take negative values. The solutions of problem (12) are of the form $w(t) = e^{At}w(0)$. If we consider a vector norm and its subordinate matrix norm, both denoted by $\|\cdot\|$, then

$$\|w(t)\| = \|e^{tA}w(0)\| \le \|e^{tA}\| \|w(0)\|,$$
(13)

and contractivity is obtained if and only if $||e^{tA}|| \leq 1$. Given the set

$$\mathcal{M} = \left\{ \delta \in \mathbb{R} \, | \, \| e^{tA} \| \le e^{t\delta} \, , t \ge 0 \right\},\,$$

it can be proven that $\mu_{\|\cdot\|}[A] = \min(\mathcal{M})$, where $\mu_{\|\cdot\|}[A]$ stands for the logarithmic norm of matrix A in the norm $\|\cdot\|$ [22, Proposition 2.1] (see, e.g., [22, 24, 1] and the references therein for the definition and properties of logarithmic norms).

From (13) and the definition of \mathcal{M} , we get the inequalities

$$\|w(t)\| \le e^{t\mu_{\|\cdot\|}[A]} \|w(0)\|, \quad t \ge 0, \qquad \|e^{tA}\| \le e^{t\mu_{\|\cdot\|}[A]} \quad t \ge 0.$$

Thus, if $\mu_{\|\cdot\|}A \leq 0$, the zero solution is stable and $\|e^{tA}\| \leq 1$ for $t \geq 0$; if $\mu_{\|\cdot\|}[A] < 0$, then the zero solution is exponentially stable and $\|e^{tA}\| < 1$ for t > 0 [22, p. 634], [16, p. 2].

For the maximum norm, the logarithmic norm of a matrix $A = (a_{ij})$ is given by

$$\mu_{\infty}[A] = \max_{1 \le i \le n} \left(a_{ii} + \sum_{j=1}^{n} |a_{ij}| \right) \,.$$

In particular, for matrix B_h in (10), as

$$a_{ii} + \sum_{j=1}^{n} |a_{ij}| = \begin{cases} -1, & i = 1, m, \\ 0, & i = 2, \dots, m-1, \end{cases}$$

we get $\mu_{\infty}[B_h] = 0$, and thus

$$||e^{tB_h}||_{\infty} \le 1, \qquad t \ge 0,$$
 (14)

that ensures that problem (10) is contractive in the maximum norm, that is,

$$||w(t)||_{\infty} \le ||w(0)||_{\infty}, \quad t \ge 0.$$

The time stepping process in Crank-Nicolson method with constant time step τ , gives the iteration

$$w_n = \phi(\tau B_h) w_{n-1} \,, \quad n \ge 1 \,,$$

where

$$\phi(z) = \frac{1 + \frac{1}{2}z}{1 - \frac{1}{2}z} \tag{15}$$

is the stability function of the time integrator. On the following, we denote $A_m = \phi(\tau B_h)$ to the Crank-Nicolson iteration matrix of dimension m, that is,

$$A_{m} = \left(I_{m} - \frac{\tau}{2}B_{h}\right)^{-1}\left(I_{m} + \frac{\tau}{2}B_{h}\right) = \begin{pmatrix} 1+s & -\frac{s}{2} & & \\ -\frac{s}{2} & 1+s & \ddots & \\ & \ddots & \ddots & -\frac{s}{2} \\ & & -\frac{s}{2} & 1+s \end{pmatrix}^{-1} \begin{pmatrix} 1-s & \frac{s}{2} & & \\ & \frac{s}{2} & 1-s & \ddots & \\ & & \ddots & \ddots & \frac{s}{2} \\ & & & \frac{s}{2} & 1-s \end{pmatrix} .$$
(16)

Observe that the two matrices in (16), corresponding to half step with forward Euler and half step with backward Euler, commute because of the the linearity of the system (10). Besides, positivity and contractivity can be studied by analyzing these properties for forward and backward Euler separately. Although there are no restrictions for positivity and contractivity with backward Euler applied to system (10), the restriction for positivity and contractivity with forward Euler is $s \leq 1$. This stepsize restriction for positivity is not sharp for problem (10); numerical experiments in [13, p.126] show that numerical positivity can be obtained for $s \leq 1.17$. As it has been pointed out above, a closer look at the iteration matrix A_m in (7) or (16), gives sharper bounds.

3 Main results

In this section we show the main results of the paper concerning stepsize restrictions for positivity and contractivity in the maximum norm for the *m*-dimensional system (10). The proofs require some preliminary material about the structure of matrix A_m and are given in section 6.

On the following theorems, the positivity of the matrix A_m means that all the entries of the matrix are non-negative; similarly, the contractivity in the maximum norm of the matrix A_m means $||A_m||_{\infty} \leq 1$.

Theorem 1. (Positivity of Crank Nicolson method)

1. For $m \in \mathbb{N}$, the matrix $A_m(s)$ in (16) is positive if and only if

$$s \le s_m^{(p)} := \frac{1}{\cosh \omega_m^{(p)} - 1},$$
(17)

where $\omega_m^{(p)} \in (\log(2+\sqrt{2}), \log(2+\sqrt{3})]$ is the unique positive root of equation

$$\coth(m\omega)\sinh\omega = 3\cosh\omega - 4. \tag{18}$$

2. The sequence $(s_m^{(p)})$ is strictly monotonically increasing with all the terms in the narrow interval $[1, 2(2 - \sqrt{2}))$. As a consequence, Crank Nicolson method preserves positivity when the spatial mesh is refined (keeping s constant).

Remark 1.

1. The sequence $(s_m^{(p)})$ increasingly converges to the limit value $s_{\infty}^{(p)} := 2(2 - \sqrt{2})$ (see Table 1 and Figure 1). This value was also obtained in [2] with other techniques. Consequently, if

$$s < s_{\infty}^{(p)} = 2(2 - \sqrt{2}) \approx 1.17$$
, (19)

then there exists a natural number m_0 such that the matrix A_m is positive for any value of $m \ge m_0$.

2. As $\omega_m^{(p)} \in (\log(2+\sqrt{2}), \log(2+\sqrt{3})] \approx (1.22795, 1.31696]$, an approximated value can be easily computed by the bisection method.

In Table 1 below we show the roots $\omega_m^{(p)}$ of equation (18) and the CFL restrictions $s_m^{(p)}$ for positivity in (17) for different values of m.

m	$\omega_m^{(p)}$	$x_m^{(p)} = \cosh \omega_m^{(p)}$	$s_m^{(p)} = 1/(x_m^{(p)} - 1)$
1	$\log(2+\sqrt{3})$	2	1
2	1.23590	$1 + \sqrt{3}/2 \approx 1.86603$	$2/\sqrt{3} \approx 1.15470$
3	1.22864	1.85464	1.17009
4	1.22801	1.85365	1.17144
÷		:	:
7	1.22795	1.85355	1.17157
:	•		
∞	$\log(2+\sqrt{2})$	$(6+\sqrt{2})/4$	$2(2-\sqrt{2}) \approx 1.171572875$

Table 1: Roots $\omega_m^{(p)}$ of (18) and CFL restrictions $s_m^{(p)}$ in (17) for positivity.

Next, we give the results for contractivity in the infinite norm. Observe that the symmetry of matrix A_m makes $||A_m||_{\infty} = ||A_m||_1$, and the result is also valid for the 1-norm.

Theorem 2. (Contractivity of Crank Nicolson method)

- 1. For $m \in \{1, 2, 3\}$ the matrix $A_m(s)$ in (16) is contractive in the maximum norm for any value of s > 0.
- 2. For $m \in \mathbb{N}$, $m \ge 4$, the matrix $A_m(s)$ in (16) is contractive in the maximum norm if and only if

$$s \le s_m^{(c)} := \frac{1}{\cosh \omega_m^{(c)} - 1},$$
(20)

where $\omega_m^{(c)} \in [\log((3 + \sqrt{5} + \sqrt{-2 + 6\sqrt{5}})/4), \log 3) \approx [0.767197, 1.09861)$ is the unique positive root of equation

$$2\sinh\frac{(m-1)\omega}{4}\sinh\frac{(m+1)\omega}{4} = \sinh\frac{\omega}{2}\sinh\frac{(m+1)\omega}{2},\qquad(21)$$

if m is odd, or equation

$$\sinh^{2}\left(\frac{\omega}{2}\right)\sinh\frac{m\omega}{2}\left(\sinh\frac{(m+2)\omega}{2}-\sinh\frac{m\omega}{2}\right)=\sinh\omega\,\sinh\frac{(m+1)\omega}{2}\sinh\frac{m\omega}{4}\sinh\frac{(m-2)\omega}{4}\,,$$
(22)

if m is even.

3. The sequence $(s_m^{(c)})$ is strictly monotonically decreasing with all the terms in the interval $(3/2, 1 + \sqrt{5}]$. As a consequence, Crank Nicolson method does not preserve contractivity when the spatial mesh is refined (keeping s constant).

Remark 2. The sequence $(s_m^{(c)})$ decreasingly converges to the limit value $s_{\infty}^{(c)} := 3/2$ (see Figures 1 and 3). Consequently, the matrix $A_m(s)$ is contractive for all m if and only if $s \in (0, 3/2]$. For infinite matrices the bound $s_{\infty}^{(c)} := 3/2$ has been obtained with different techniques in [3, 16].

In Tables 2 and 3 we give the CFL restrictions $s_m^{(c)}$ in (20) for contractivity in the infinite norm. The values in Table 2 (odd case) and Table 3 (even case) have been obtained from equations (21) and (22), respectively. Observe that both, the roots $\omega_m^{(c)}$ of equation (21) (odd case) and the roots $\omega_m^{(c)}$ of equation (22) (even case), increasingly converge to the limit value log 3. Consequently these roots are in the narrow interval $\left[\log\left((3+\sqrt{5}+\sqrt{-2+6\sqrt{5}})/4\right),\log 3\right) \approx [0.767197, 1.09861)$ and can be easily obtained with bisection method. The numeric values shown in tables 2 and 3 were obtained after 10 iterations with bisection method.



Figure 1: Sequences $s_m^{(p)}$ and $s_m^{(c)}$ with the restrictions over CFL coefficient s for positivity and contractivity.

Corollary 1. Consider the numerical integration of the m-dimensional problem (10) with the Crank-Nicolson method. Then if the method is positive for a given CFL coefficient, then it is also contractive in the infinity and 1 norms.

Proof. It is straightforward from Theorems 1 and 2.

The converse to the previous Corollary is not true as we can see in the following trivial example, where contractivity is preserved while positivity is violated.

Example 1. Consider the diffusion equation in (1) with initial function

$$u(x,0) = \begin{cases} 0 & \text{for } 0 < x < \frac{7}{8} \,, \\ 1 & \text{for } \frac{7}{8} \le x < 1 \,, \end{cases}$$

giving discontinuities at x = 7/8 and x = 1 for t=0. From second order central differences with h = 1/8 we get approximations $\omega(t) = (\omega^1(t), \ldots, \omega^7(t)) \approx (u(x_1, t), \ldots, u(x_7, t))$. Application with $\tau = 0.025$ of one Crank-Nicolson step, $w_1 = A_7 w_0$, gives the vector $w_1 \approx \omega(\tau)$

$$w_1 = (0.0013, 0.0041, 0.0120, 0.0356, 0.1019, 0.2961, -0.1397),$$
(23)

where A_7 is the Crank-Nicolson iteration matrix in (16) for the case m = 7 and w_0 is the initial profile $w_0 = (0, 0, 0, 0, 0, 0, 1)$. Observe that $||w_1||_{\infty} = 0.2961 < ||w_0||_{\infty} = 1$.

In this example the CFL coefficient $s = \tau/h^2 = 1.6$ is greater than the positivity bound $s_7^{(p)} = 1.17157$ (see Table 1), but it is lower than the contrativity one $s_7^{(c)} = 1.61803$ (see Table 2). Consequently, contractivity is preserved while we cannot ensure positivity. Actually, as we can see in vector w_1 in (23), negativity is not preserved.

m	$\omega_m^{(c)}$	$x_m^{(c)} = \cosh \omega_m^{(c)}$	$s_m^{(c)} = 1/(x_m^{(c)} - 1)$
3			∞
5	$2 \operatorname{arccsch} 2 \approx 0.962424$	3/2	2
7	$\log \frac{\left(1 + \sqrt{5} + \sqrt{2(1 + \sqrt{5})}\right)}{2} \approx 1.06131$	$(1 + \sqrt{5})/2$	$(1+\sqrt{5})/2 \approx 1.61803$
9	1.08707	1.65139	1.53518
:		:	
∞	$\log 3 \approx 1.09861$	5/3	3/2

Table 2: Positive root of (21) and bounds for contractivity (odd case).

m	$\omega_m^{(c)}$	$x_m^{(c)} = \cosh \omega_m^{(c)}$	$s_m^{(c)} = 1/(x_m^{(c)} - 1)$
4	$\log(\frac{1}{4}(3+\sqrt{5}+\sqrt{-2+6\sqrt{5}}))$	$\frac{1}{4}(3+\sqrt{5})$	$1+\sqrt{5}$
	pprox 0.767197	≈ 1.30902	≈ 3.23607
:			:
10	1.09110	1.65669	1.52278
:			:
20	1.09855	1.66658	1.5002
:			:
∞	$\log 3 \approx 1.09861$	5/3	3/2

Table 3: Positive root of (22) and bounds for contractivity (even case).

4 Conclusions and future work

In this paper we have studied CFL restrictions when the Crank-Nicolson method is used to solve the heat equation (1) with Dirichlet boundary conditions. We have obtained bounds $s_m^{(p)}$ for positivity and bounds $s_m^{(c)}$ for contractivity for any value of the spatial discretization parameter m. To get these bounds we have represented the Crank-Nicolson iteration matrix A_m in terms of some Chebyshev-like polynomials (26,28,33). We have obtained bounds for the θ -method (7) for the particular case $\theta = 1/2$, but similar bounds can be obtained for other values of the parameter following the same ideas.

We have seen that the positivity of matrix A_m is determined by the largest root of polynomial $P_m(x)$, and we have provided a narrow interval where this root can be found. Similarly, we have considered these polynomials to analyze the contractivity and we have

provided a narrow interval to get the corresponding bounds, both in the odd and even case.

As far as we know, polynomials $P_n(x)$ and $C_n(x)$ have not been used previously in the literature. The strength of this idea can be used to prove qualitative properties for other problems. Furthermore, this approach can also be used for other discretizations of the heat equation (1.1) [10].

5 Preliminary material for the proofs of the main results

In this section we introduce the notation, definitions and some results needed to prove the main results of the paper. In subsection 5.1 we express the Crank-Nicolson iteration matrix (16) in terms of rational functions. These functions can be written easily with the help of some Chebyshev-like polynomials U_m , P_m and C_m . In subsection 5.2 we give the definition of these polynomials and we also add some results that will be used in the proofs of Section 6.

5.1 The Crank-Nicolson matrix A_m in terms of rational functions

A direct computation of the product $(I_m - \frac{\tau}{2}B_h)^{-1}(I_m + \frac{\tau}{2}B_h)$ in (16) gives us the entries of matrix A_m expressed as rational functions, where the polynomials involved can be obtained recursively. These simplified closed expressions will make it easier to get bounds for positivity and contractivity.

Example 2. For m = 3, a direct computation of the symmetric matrix A_3 in (16) gives

$$A_{3}(s) = \begin{pmatrix} \frac{2+2s-2s^{2}-s^{3}}{2+6s+5s^{2}+s^{3}} & \frac{2s}{2+4s+s^{2}} & \frac{s^{2}}{2+6s+5s^{2}+s^{3}} \\ \frac{2s}{2+4s+s^{2}} & \frac{2-s^{2}}{2+4s+s^{2}} & \frac{2s}{2+4s+s^{2}} \\ \frac{s^{2}}{2+6s+5s^{2}+s^{3}} & \frac{2s}{2+4s+s^{2}} & \frac{2+2s-2s^{2}-s^{3}}{2+6s+5s^{2}+s^{3}} \end{pmatrix}.$$

Remember $s = \tau/h^2$ denotes CFL coefficient. This matrix can be written even simpler if we consider the new variable x = 1 + 1/s. Observe that x > 1 when s > 0. With the help of a new kind of polynomials $U_n(x)$, $P_n(x)$ and $C_n(x)$, we can write $A_3(x)$ as

$$A_{3}(x) = \begin{pmatrix} \frac{2x^{3}-4x^{2}+1}{2x^{3}-x} & \frac{2(x-1)}{2x^{2}-1} & \frac{x-1}{2x^{3}-x} \\ \frac{2(x-1)}{2x^{2}-1} & \frac{2x^{2}-4x+1}{2x^{2}-1} & \frac{2(x-1)}{2x^{2}-1} \\ \frac{x-1}{2x^{3}-x} & \frac{2(x-1)}{2x^{2}-1} & \frac{2x^{3}-4x^{2}+1}{2x^{3}-x} \end{pmatrix} = \frac{1}{U_{3}(x)} \begin{pmatrix} P_{3}(x) & C_{2}(x) & C_{1}(x) \\ C_{2}(x) & C_{1}(x) + P_{3}(x) & C_{2}(x) \\ C_{1}(x) & C_{2}(x) & P_{3}(x) \end{pmatrix}.$$

Observe that $A_3(x)$ has been written just in terms of $U_3(x)$, $P_3(x)$, $C_1(x)$ and $C_2(x)$. We give the definition and all the details about these polynomials $U_n(x)$, $P_n(x)$ and $C_n(x)$ in the next subsection. Before, we extend the ideas in this simple example to the more general case of the matrix $A_m(x)$ for any value of m, although we have to distinguish between the odd case and the even case.

Proposition 1. Matrix $A_m(x)$ can be written in terms of polynomials $U_m(x)$, $P_m(x)$ and $C_n(x)$, $n = 1, \ldots, m-1$. If m is an odd number, Crank Nicolson matrix can be reduced to

$$A_{m}(x) = \frac{1}{U_{m}} \begin{pmatrix} P_{m} & C_{m-1} & \dots & C_{\frac{m+1}{2}} & \dots & C_{2} & C_{1} \\ C_{m-1} & P_{m} + C_{m-2} & \dots & C_{\frac{m-1}{2}} + C_{\frac{m+3}{2}} & \dots & C_{1} + C_{3} & C_{2} \\ \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \vdots \\ C_{\frac{m+1}{2}} & C_{\frac{m-1}{2}} + C_{\frac{m+3}{2}} & \dots & P_{m} + \sum_{n=1}^{\frac{m-1}{2}} C_{2n-1} & \dots & C_{\frac{m-1}{2}} + C_{\frac{m+3}{2}} & C_{\frac{m+1}{2}} \\ \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \vdots \\ C_{2} & C_{1} + C_{3} & \dots & C_{\frac{m-1}{2}} + C_{\frac{m+3}{2}} & \dots & P_{m} + C_{m-2} & C_{m-1} \\ C_{1} & C_{2} & \dots & C_{\frac{m+1}{2}} & \dots & C_{m-1} & P_{m} \end{pmatrix}, \quad (24)$$

where all the polynomials are evaluated at x = 1 + 1/s. If m is an even number, we write

$$A_{m}(x) = \frac{1}{U_{m}} \begin{pmatrix} P_{m} & C_{m-1} & \dots & C_{\frac{m+2}{2}} & C_{\frac{m}{2}} & \dots & C_{2} & C_{1} \\ C_{m-1} & P_{m} + C_{m-2} & \dots & C_{\frac{m}{2}} + C_{\frac{m+4}{2}} & C_{\frac{m-2}{2}} + C_{\frac{m+2}{2}} & \dots & C_{1} + C_{3} & C_{2} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ C_{\frac{m+2}{2}} & C_{\frac{m}{2}} + C_{\frac{m+4}{2}} & \dots & P_{m} + \sum_{n=1}^{\frac{m-2}{2}} C_{2n} & \sum_{n=1}^{\frac{m}{2}} C_{2n-1} & \dots & C_{\frac{m-2}{2}} + C_{\frac{m+2}{2}} & C_{\frac{m}{2}} \\ C_{\frac{m}{2}} & C_{\frac{m-2}{2}} + C_{\frac{m+2}{2}} & \dots & \sum_{n=1}^{\frac{m}{2}} C_{2n-1} & P_{m} + \sum_{n=1}^{\frac{m-2}{2}} C_{2n} & \dots & C_{\frac{m}{2}} + C_{\frac{m+4}{2}} & C_{\frac{m+2}{2}} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ C_{2} & C_{1} + C_{3} & \dots & C_{\frac{m-2}{2}} + C_{\frac{m+2}{2}} & C_{\frac{m}{2}} + C_{\frac{m+4}{2}} & \dots & P_{m} + C_{m-2} & C_{m-1} \\ C_{1} & C_{2} & \dots & C_{\frac{m}{2}} & C_{\frac{m+2}{2}} & \dots & C_{m-1} & P_{m} \end{pmatrix}.$$

$$(25)$$

Proof. It is straightforward from the computation of the product $(I_m - \frac{\tau}{2}B_h)^{-1}(I_m + \frac{\tau}{2}B_h)$ in (16) and the use of polynomials $U_n(x)$, $P_n(x)$ and $C_n(x)$, $n = 1, \ldots, m-1$, defined in the next subsection.

Observe that $A_m(x)$ is bisymmetric, that is, it is symmetric on both diagonals. This implies that $A_m(x)$ is also centrosymmetric. Then the entries a_{ij} satisfy $a_{ij} = a_{n-i+1,n-j+1}$, for $1 \leq i, j \leq n$. Consequently, if m is odd, the number of different entries in matrix A_m is $1+3+5+\cdots+m = (m+1)^2/4$, and, if m is even, this number is $2+4+6+\cdots+m = (m/2+1)m/2$. For example, the number of different elements in matrix A_3 in Example 2 is 4, while this number is 6 for matrix A_4 in Example 3 below.

Observe also that the numerator of each entry a_{ij} in matrix $A_m(x)$ is a sum of some polynomials $P_n(x)$, $C_n(x)$, n = 1, ..., m-1, and the number of polynomials in this sum is equal to min $\{i, j, m - i + 1, m - j + 1\}$. Properties of polynomials $U_n(x)$, $P_n(x)$ and $C_n(x)$ will allow us to analyze positivity and contractivity of Crank Nicolson method in a quite simple way. In the next section, we study these properties. **Example 3.** In Example 2 we have considered the odd case m = 3. Here, for completeness, we consider the even case m = 4. A direct computation of the symmetric matrix A_4 in (16) gives

$$A_4(s) = \frac{1}{u_4(s)} \begin{pmatrix} p_4(s) & 4s(4+8s+3s^2) & 8s^2(1+s) & 4s^3\\ 4s(4+8s+3s^2) & 16+32s+4s^2-16s^3-5s^4 & 16s(1+s)^2 & 8s^2(1+s)\\ 8s^2(1+s) & 16s(1+s)^2 & 16+32s+4s^2-16s^3-5s^4 & 4s(4+8s+3s^2)\\ 4s^3 & 8s^2(1+s) & 4s(4+8s+3s^2) & p_4(s) \end{pmatrix},$$

where $p_4(s) = -5s^4 - 24s^3 - 4s^2 + 32s + 16$ and $u_4(s) = 5s^4 + 40s^3 + 84s^2 + 64s + 16$. Now, with the help of variable x = 1 + 1/s, we can write

$$A_4(x) = \frac{1}{U_4(x)} \begin{pmatrix} P_4(x) & C_3(x) & C_2(x) & C_1(x) \\ C_3(x) & P_4(x) + C_2(x) & C_1(x) + C_3(x) & C_2(x) \\ C_2(x) & C_1(x) + C_3(x) & P_4(x) + C_2(x) & C_3(x) \\ C_1(x) & C_2(x) & C_3(x) & P_4(x) \end{pmatrix}$$

Observe that there are two central rows in the even case, but just one in the odd case.

5.2 Polynomials U_n , P_n and C_n

In this section we define the new polynomials $P_n(x)$ and $C_n(x)$. Together with the help of Chebyshev polynomials of the second kind $U_n(x)$ [7, 17, 19], we have got a simple way of writing Crank-Nicolson matrix $A_m(x)$. Besides, here we give some results concerning these polynomials, with particular interest in the distribution of their roots. Chebyshev polynomials of the second kind $U_n(x)$ belong to a general class of orthogonal polynomials and there are many works about the behaviour of their zeros [7, 17]. However, polynomials $P_n(x)$ and $C_n(x)$ do not belong to this class of orthogonal polynomials and, as far as we know, nothing is known about their roots.

Chebyshev polynomials of second kind

Chebyshev polynomial of second kind of degree $n \ge 0$ is defined as

$$U_n(x) = \frac{\sin((n+1)\arccos x)}{\sin(\arccos x)}, \quad x \in [-1,1],$$

or, in the angle variable ω , $U_n(\cos \omega) = \sin((n+1)\omega)/\sin \omega$, $\omega \in [0,\pi]$. These polynomials can also be defined for any value of $x \in \mathbb{R}$ by the recurrence relation

$$U_0(x) = 1,$$

$$U_1(x) = 2x,$$

$$U_n(x) = 2x U_{n-1}(x) - U_{n-2}(x).$$
(26)

It is possible to write the recurrence relation (26) in terms of the determinant of the tridiagonal matrix tridiag(1, 2x, 1) of dimension n.

$$U_n(x) = \begin{vmatrix} 2x & 1 & & \\ 1 & 2x & \ddots & \\ & \ddots & \ddots & 1 \\ & & 1 & 2x \end{vmatrix}$$
(27)

Recall that each polynomial $U_n(x)$ has n roots $x_i^n = \cos(i\pi/(n+1))$, $i = 1, \ldots, n$, in the interval [-1, 1]. These roots are uniformly distributed in the angle variable $\omega = \arccos x$ in the interval $[0, \pi]$. Notice that polynomials $U_n(x)$ defined in (26) are positive for x > 1.

Polynomials P_n

If we change the first two elements in the recursive relation (26), then a new family of polynomials can be defined

$$P_0(x) = -1,$$

$$P_1(x) = 2x - 4,$$

$$P_n(x) = 2x P_{n-1}(x) - P_{n-2}(x),$$
(28)

where $P_n(x)$ denotes the polynomial of degree n. In this case the recurrence relation (28) can also be written in terms of the determinant of a matrix of dimension n.

$$P_n(x) = \begin{vmatrix} 2x - 4 & -1 & & \\ 1 & 2x & 1 & & \\ & \ddots & \ddots & \ddots & \\ & & 1 & 2x & 1 \\ & & & 1 & 2x \end{vmatrix} .$$
(29)

In the next proposition we analyze the roots of each polynomial $P_n(x)$. In the proof, shown in Section 6, it is relevant the fact that each polynomial $P_n(x)$ can be written in terms of Chebyshev polynomials of second kind

$$P_n(x) = 2U_{n-2}(x) - 4U_{n-1}(x) + U_n(x).$$
(30)

This equality is obtained by writing the determinant (29) in terms of the determinant (27), and the use of the recurrence relation (26).

Proposition 2. The polynomial $P_n(x)$ defined in (28) has exactly n-1 roots x_i^n , $i = 1, \ldots, n-1$, in the interval (-1, 1), and an additional isolated root $x_n := x_n^n$ in the interval $(\frac{6+\sqrt{2}}{4}, 2]$. Furthermore, the isolated root is $x_n = \cosh \omega_n$, where ω_n is the unique root of the equation

$$\coth(n\omega) = \frac{3\cosh\omega - 4}{\sinh\omega}, \qquad \omega \in (0,\infty).$$
(31)

Besides, $x_1 = 2$ and the sequence of isolated roots (x_n) decreasingly converges to the limit value $x_{\infty} = (6 + \sqrt{2})/4$.



Figure 2: Polynomials $P_n(x)$, n = 1, ..., 4. Each polynomial has an isolated root in the interval $\left(\frac{6+\sqrt{2}}{4}, 2\right]$.

As it is shown in the proof (see Section 6.3), for any value of n, the unique root ω_n of equation (31) lies in the interval $(\omega_{\infty}, \omega_1] = (\log(2 + \sqrt{2}), \log(2 + \sqrt{3})]$. Consequently, the isolated root $x_n = \cosh \omega_n$ of polynomial $P_n(x)$ lies in the narrow interval $(\frac{6+\sqrt{2}}{4}, 2]$. Having the root well located makes it easy to approach it by any numerical method. In Table 1 we show some of these roots after 10 steps with bisection method.

Polynomials C_n

The polynomial C_n of degree n is defined as

$$C_n(x) = P_n(x) + U_n(x), \qquad n \in \mathbb{N},$$
(32)

where P_n is the polynomial of degree *n* defined above and U_n is the Chebyshev polynomial of second kind of degree *n*. Consequently, all the properties of C_n are consequence of this definition, including its recursive definition

$$C_0(x) = 0,$$

$$C_1(x) = 4(x - 1),$$

$$C_n(x) = 2x C_{n-1}(x) - C_{n-2}(x).$$
(33)

Observe that the recursive formula (33) is the same as (28) for P_n and (26) for U_n , with just different starting values C_0 and C_1 . As in previous cases, it is worth writing $C_n(x)$ in terms of a determinant

$$C_n(x) = \begin{vmatrix} 4(x-1) & 0 & & \\ 1 & 2x & 1 & & \\ & \ddots & \ddots & \ddots & \\ & & 1 & 2x & 1 \\ & & & 1 & 2x \end{vmatrix} .$$
(34)

Proposition 3. Each polynomial $C_n(x)$ has exactly n-1 roots x_i^n , i = 1, ..., n-1, in the interval (-1, 1), and the additional isolated root $x_n = 1$.

Proof. It is straightforward if we use the determinant (34), where we get

$$C_n(x) = 4(x-1)U_{n-1}(x)$$
(35)

Consequently the roots of polynomial C_n are the isolated root $x_n = 1$, and the n-1 roots of the Chebyshev polynomial of second kind of degree n-1.

Corollary 2. If x > 1, then $C_n(x) > 0 \ \forall n \in \mathbb{N}$.

Proof. It is straightforward from the previous proposition.

In the following Lemma we give some technical properties of polynomials $C_n(x)$ that we will need in Section 6.

Lemma 3. For the polynomial C_n of degree n defined in (32) or (33) the following properties hold:

- 1. $C_n(1) = 0, \forall n \in \mathbb{N}$.
- 2. $C_n(x) = 2U_{n-2}(x) 4U_{n-1}(x) + 2U_n(x), \forall n \in \mathbb{N}.$
- 3. If x > 1, then $0 < C_n(x) \le C_{n+1}(x)$, $\forall n \in \mathbb{N}$.
- 4. If x > 1, then $2C_n(x) \le C_{n-1}(x) + C_{n+1}(x)$, $\forall n \in \mathbb{N}$.

Proof. Part 1 is straightforward. Part 2 is also straightforward if we use the relationship (30) and definition (32).

To prove part 3, as x > 1, we set $x = \cosh \omega$ in (35), to obtain

$$C_n(\cosh\omega) = \frac{4(\cosh\omega - 1)\sinh(n\omega)}{\sinh\omega} > 0.$$
(36)

As $\sinh \omega$ is an increasing function, it holds $\sinh(n\omega) < \sinh((n+1)\omega)$, and consequently $C_n(x) \le C_{n+1}(x)$.

Finally, to prove 4, as

$$\sinh((n-1)\omega) + \sinh((n+1)\omega) = 2\cosh\omega\sinh(n\omega) \ge 2\sinh(n\omega),$$

we can use again (36), for x > 1, to obtain that $2C_n(x) \le C_{n-1}(x) + C_{n+1}(x) \quad \forall n \in \mathbb{N}$. \Box

6 Proofs of theorems in Sections 3 and 5

In this section we give the proofs of the main results in the paper.

6.1 Positivity of the Crank-Nicolson method

Recall that the Crank Nicolson method is positive if and only if all the elements in matrix A_m (24)-(25) are positive. The following lemma simplifies the proof of Theorem 1.

Lemma 4. If x > 1, then all the elements in matrix $A_m(x)$ (24)-(25) are non-negative if and only if polynomial $P_m(x)$ is non-negative.

Proof. Recall that polynomials $U_n(x)$ are positive for x > 1. Polynomials $C_n(x)$ are also positive for x > 1 (see Corollary 2). Consequently, all extra-diagonal elements are positive.

For the diagonal elements $A_m^{(i,i)}(x)$, the positivity of polynomials $U_n(x)$ and $C_n(x)$ for all $n \in \mathbb{N}$ implies that

$$\min_{i} A_{m}^{(i,i)}(x) = A_{m}^{(1,1)}(x) = A_{m}^{(m,m)}(x) = \frac{P_{m}(x)}{U_{m}(x)}$$

Then the analysis of the positivity of the elements in $A_m(x)$ is reduced to the positivity of polynomial $P_m(x)$.

Proof of Theorem 1

- 1. From Lemma 4, we just have to study the positivity of polynomials $P_m(x)$ for x > 1. From Proposition 2, $P_m(x)$ has exactly m - 1 roots in the interval (-1, 1) and an additional isolated real root $x_m^{(p)} = \cosh(\omega_m^{(p)})$ in the interval $(\frac{6+\sqrt{2}}{4}, 2]$, where $\omega_m^{(p)}$ is the unique root of equation (31), that is the same as (18). As x = 1 + 1/s, then the CFL coefficient is s = 1/(x-1), and inequality (17) is obtained.
- 2. In the limit case, Proposition 2 gives us the limit value $x_{\infty}^{(p)} = (6 + \sqrt{2})/4$. Then for the CFL coefficient we get the bound $s_{\infty}^{(p)} = 1/(x_{\infty}^{(p)} 1) = 2(2 \sqrt{2})$, and inequality (19) is obtained.

6.2 Contractivity of the Crank-Nicolson method.

Before writing the proof of Theorem 2, we need two technical lemmas. In Lemma 5 below, we compute the maximum norm of matrix $A_m(x)$ in terms of polynomials $U_n(x)$, $P_n(x)$ and $C_n(x)$. Then, in Lemma 6 we get the inequalities needed for the contractivity condition $||A_m||_{\infty} \leq 1$. In contrast to positivity, in the analysis of the contractivity it is necessary to distinguish between the even and odd cases.

Lemma 5.

1. If $m \geq 3$ is a natural odd number, then the maximum norm of Crank-Nicolson matrix is

$$||A_m||_{\infty} = \frac{1}{U_m} \left(|P_m + \sum_{n=1}^{\frac{m-1}{2}} C_{2n-1}| + 2\sum_{i=0}^{\frac{m-3}{2}} \sum_{n=1}^{\frac{m-(2i+1)}{2}} C_{2n+i} \right).$$
(37)

2. If $m \geq 2$ is a natural even number, then the norm of Crank-Nicolson matrix is

$$||A_m||_{\infty} = \frac{1}{U_m} \left(|P_m + \sum_{n=1}^{\frac{m}{2}-1} C_{2n}| + \sum_{i=1}^{\frac{m}{2}-1} \sum_{n=1+i}^{m-i} C_n + \sum_{n=1}^{m/2} C_{2n-1} \right).$$
(38)

Proof. Note that the norm in (37) is obtained from the sum of the elements in the central row of A_m (odd case), while the norm in (38) is obtained from the sum of the elements

in any of the two symmetric central rows (even case). To get this result we will proof the following inequalities

$$\sum_{j=1}^{m} |A_m^{(i,j)}| \le \sum_{j=1}^{m} |A_m^{(i+1,j)}|, \quad i = 1, \dots, \frac{m-1}{2} (\text{odd case}) \, \frac{m-2}{2} (\text{even case}).$$

For the sake of simplicity we will denote A_m^i to the sum $\sum_{j=1}^m |A_m^{(i,j)}| U_m$. Observe that the symmetry of matrices (24-25) makes $A_m^i = A_m^{m+1-i}$, $i = 1, \ldots, (m-1)/2$ (odd case) or $i = 1, \ldots, m/2$ (even case). Then, we will proof that

$$A_m^1 \le A_m^2 \le \dots \le A_m^{\frac{m-1}{2}} \le A_m^{\frac{m+1}{2}} \qquad \text{(odd case)}$$
$$A_m^1 \le A_m^2 \le \dots \le A_m^{\frac{m-2}{2}} \le A_m^{\frac{m}{2}} = A_m^{\frac{m+2}{2}} \qquad \text{(even case)}$$

Consequently, $||A_m||_{\infty}$ is obtained from the sum of the elements in the central row (odd case) or from the sum of the elements in any of the two central rows (even case).

Recall that for a given matrix A with positive extra-diagonal elements the following equality trivially holds

$$||A||_{\infty} = \max_{i} \left(\sum_{j=1}^{m} |a_{ij}| \right) = \max_{i} \left(|a_{ii}| + \sum_{j \neq i} a_{ij} \right).$$
(39)

In our case $C_n(x) \ge 0$ for $x \ge 1$ (see Proposition 3). Consequently all extra-diagonal elements of matrices (24) and (25) are non-negative and we can use (39) to compute $||A_m||_{\infty}$.

For the first and the second row of matrix (24) or (25) we get the sums

$$A_m^1 = |P_m| + \sum_{i=1}^{m-1} C_i, \qquad A_m^2 = |P_m + C_{m-2}| + \sum_{i=1}^{m-1} C_i + \sum_{i=1}^{m-1} C_i - C_{m-2}.$$

Then, after cancelling terms, the difference $A_m^1 - A_m^2$ is

$$A_m^1 - A_m^2 = |P_m| - |P_m + C_{m-2}| - \sum_{2}^{m-1} C_i + C_{m-2}$$

Adding and subtracting C_{m-2} in the term $|P_m|$, we can write

$$A_m^1 - A_m^2 \le |P_m + C_{m-2}| + C_{m-2} - |P_m + C_{m-2}| - \sum_{2}^{m-1} C_i + C_{m-2}$$
$$= 2C_{m-2} - \sum_{2}^{m-1} C_i \le C_{m-3} + C_{m-1} - \sum_{2}^{m-1} C_i = -\sum_{2}^{m-4} C_i - C_{m-2} \le 0,$$

where we have used $2C_{m-2} \leq C_{m-3} + C_{m-1}$ from item (4) in Lemma 3. Observe that property (3) in Lemma 3 allows us to finally write

$$A_m^1 - A_m^2 \le -\sum_{2}^{m-3} C_i \le 0.$$
(40)

If we proceed in the same way for the difference $A_m^2 - A_m^3$, after cancelling terms, and after adding and subtracting C_{m-4} in the term $|P_m + C_{m-2}|$, we get the following inequality

$$A_m^2 - A_m^3 \le 2C_{m-4} - \sum_{3}^{m-2} C_i$$
.

Again, the use of properties (4) and (3), in this order, from Lemma 3 makes it possible to write an inequality analogous to (40)

$$A_m^2 - A_m^3 \le C_{m-5} + C_{m-3} - \sum_{3}^{m-2} C_i \le -\sum_{3}^{m-4} C_i \le 0.$$

The proof follows in the same way for the odd and even case up to the last step when we achieve the central row (odd case) or the two central rows (even case). Then we have to consider two different cases:

1. If m is an odd number, the last step consists in studying the difference $A_m^{\frac{m-1}{2}} - A_n^{\frac{m+1}{2}}$, where $A_n^{\frac{m+1}{2}}$ represents the sum of the elements in the central row. After cancelling terms, we can write

$$A_m^{\frac{m-1}{2}} - A_m^{\frac{m+1}{2}} = |P_m + \sum_{n=2}^{\frac{m-1}{2}} C_{2n-1}| + C_1 - |P_m + \sum_{n=1}^{\frac{m-1}{2}} C_{2n-1}| - C_{\frac{m+1}{2}}$$

Adding and subtracting C_1 in the term $|P_m + \sum_{n=2}^{\frac{m-1}{2}} C_{2n-1}|$, we get

$$A_m^{\frac{m-1}{2}} - A_m^{\frac{m+1}{2}} \le |P_m + \sum_{n=1}^{\frac{m-1}{2}} C_{2n-1}| + 2C_1 - |P_m + \sum_{n=1}^{\frac{m-1}{2}} C_{2n-1}| - C_{\frac{m+1}{2}} = 2C_1 - C_{\frac{m+1}{2}} \le C_0 + C_2 - C_{\frac{m+1}{2}} = C_2 - C_{\frac{m+1}{2}} \le 0,$$

where we have used $2C_1 \leq C_0 + C_2$ from property (4) in Lemma 3.

Consequently, if $m \ge 3$ is an odd number, the maximum value of A_m^i is obtained in the central row $A_m^{\frac{m+1}{2}}$ and we can conclude

$$||A_m||_{\infty} = \frac{A_m^{\frac{m+1}{2}}}{U_m} = \frac{1}{U_m} \left(|P_m + \sum_{n=1}^{\frac{m-1}{2}} C_{2n-1}| + 2\sum_{i=0}^{\frac{m-3}{2}} \sum_{n=1}^{\frac{m-(2i+1)}{2}} C_{2n+i} \right).$$

2. If m is an even number, then there is not a central row but two central symmetric rows $A_m^{\frac{m}{2}}$ and $A_m^{\frac{m+2}{2}}$, and the maximum value is obtained at any of these two files. Now, in the last step of the proof, if $m \ge 4$, we have to write the difference $A_m^{\frac{m-2}{2}} - A_m^{\frac{m}{2}}$. Note that for the simple case m = 2, it holds $A_m^1 = A_m^2$. After cancelling terms, we can write

$$A_m^{\frac{m-2}{2}} - A_m^{\frac{m}{2}} = |P_m + \sum_{n=2}^{\frac{m}{2}-1} C_{2n}| + C_2 - |P_m + \sum_{n=1}^{\frac{m}{2}-1} C_{2n}| - C_{\frac{m}{2}} - C_{\frac{m+2}{2}}$$

Adding and subtracting C_2 in the term $|P_m + \sum_{n=2}^{\frac{m}{2}-1} C_{2n}|$, we get

$$A_m^{\frac{m-2}{2}} - A_m^{\frac{m}{2}} \le |P_m + \sum_{n=1}^{\frac{m}{2}-1} C_{2n}| + 2C_2 - |P_m + \sum_{n=1}^{\frac{m}{2}-1} C_{2n}| - C_{\frac{m}{2}} - C_{\frac{m+2}{2}}$$
$$= 2C_2 - C_{\frac{m}{2}} - C_{\frac{m+2}{2}} \le C_1 + C_3 - C_{\frac{m}{2}} - C_{\frac{m+2}{2}} \le 0,$$

where we have used $2C_2 \leq C_1 + C_3$ from property (4) in Lemma 3.

Consequently, if m is an even number, the maximum value of A_m^i is obtained in the row $A_m^{\frac{m}{2}}$ and we can conclude

$$\|A_m\|_{\infty} = \max_{i} \left(|a_{ii}| + \sum_{j \neq i} a_{ij} \right) = \frac{1}{U_m} \left(|P_m + \sum_{n=1}^{\frac{m}{2}-1} C_{2n}| + \sum_{i=1}^{\frac{m}{2}-1} \sum_{n=1+i}^{m-i} C_n + \sum_{n=1}^{m/2} C_{2n-1} \right) \quad \Box$$

Once we have got the maximum norm of matrix A_m in terms of polynomials U_m , P_m and C_m , we can get bounds $s_m^{(c)}$ for contractivity for any value of m if we are able to solve the corresponding inequality $||A_m||_{\infty} \leq 1$. This is done in the following lemma. In Figure 3 we have plot $||A_m(s)||_{\infty}$ for some values of m, and we have also added some contractivity bounds $s_m^{(c)}$.



Figure 3: $||A_m(s)||$ for different values of $m \in \{1, 2, 3, 4, 5, 7, 9, 21\}$. When $m \in \{1, 2, 3\}$ it holds $||A_m(s)|| < 1$, for all s > 0. For $m \ge 4$, $||A_m(s)||$ cuts the line s = 1 at $s_m^{(c)}$. The sequence $(s_m^{(c)})$ is strictly monotonically decreasing with all the terms in the interval $(3/2, 1 + \sqrt{5}]$.

Lemma 6.

1. (Ood case) If m is a natural odd number, then for the maximum norm of Crank-Nicolson matrix we have

$$||A_m||_{\infty} \le 1 \iff \frac{2\sinh\frac{(m-1)\omega}{4}\sinh\frac{(m+1)\omega}{4}}{\sinh\frac{(m+1)\omega}{2}} \le \sinh\frac{\omega}{2}$$

where $\omega = \operatorname{arccosh}(1+1/s)$. In the limit, when $m \to \infty$, we get contractivity if and only if

$$e^{-\omega/2} \le \sinh \frac{\omega}{2}$$
 (41)

2. (Even case) If m is a natural even number, then we have

$$||A_m||_{\infty} \le 1 \iff \frac{\sinh \frac{m\omega}{2} \left(\sinh \frac{(m+2)\omega}{2} - \sinh \frac{m\omega}{2}\right)}{\sinh \frac{(m+1)\omega}{2} \sinh \frac{m\omega}{4} \sinh \frac{(m-2)\omega}{4}} \ge \frac{\sinh \omega}{\sinh^2 \frac{\omega}{2}}$$

In the limit, when $m \to \infty$, we get contractivity if and only if

$$2\left(-1+e^{\omega}\right) \ge \frac{\sinh\omega}{\sinh^2\frac{\omega}{2}} \tag{42}$$

If $\omega > 0$, inequalities (41) and (42) are equivalent, and they are true iff $\omega \ge \log 3$. In the variable s this is equivalent to the known restriction $s \le 3/2$.

Proof.

1. Remember that, in the odd case, the norm in (37) is obtained from the sum of the elements in the central row of A_m . In that case, the diagonal element in the central row $P_m + \sum_{n=1}^{\frac{m-1}{2}} C_{2n-1}$ can be written in closed form as

$$P_m + \sum_{n=1}^{\frac{m-1}{2}} C_{2n-1} = -U_m + \sum_{n=1}^{\frac{m+1}{2}} C_{2n-1} = -\frac{\sinh((m+1)\omega)}{\sinh\omega} + \frac{4(x-1)\sinh^2\frac{(m+1)\omega}{2}}{\sinh^2\omega},$$

where we have changed $P_m = C_m - U_m$ and we have considered the angle variable $\omega = \operatorname{arccosh} x$. For any value of m, the unique positive root ω_m of this diagonal element lies in the interval $(\log 3, \log(2 + \sqrt{3})]$. This root can easily be obtained from the simplified equation in the variable $s = (\cosh \omega - 1)^{-1}$

$$\sqrt{1+2s} = 2\tanh\frac{(1+m)\operatorname{arccosh}(1+1/s)}{2}$$

In this variable, the unique positive root s_m lies in the interval [1, 3/2). Observe that the sequence of roots (s_m) increasingly converges to the limit value $s_{\infty} = 3/2$. When $s \in (0, s_m)$ the diagonal element $P_m + \sum_{n=1}^{\frac{m-1}{2}} C_{2n-1}$ is positive, and, from (37), we easily obtain $||A_m||_{\infty} < 1$.

On the other hand, when $s \in [s_m, \infty)$, we have $P_m + \sum_{n=1}^{\frac{m-1}{2}} C_{2n-1} \leq 0$, and, from (37), the inequality for contractivity is

$$||A_m||_{\infty} = \frac{1}{U_m} \left(-P_m - \sum_{n=1}^{\frac{m-1}{2}} C_{2n-1} + 2 \sum_{i=0}^{\frac{m-3}{2}} \sum_{n=1}^{\frac{m-(2i+1)}{2}} C_{2n+i} \right) \le 1.$$

This is equivalent to

$$-\sum_{n=1}^{\frac{m-1}{2}} C_{2n-1} + 2\sum_{i=0}^{\frac{m-3}{2}} \sum_{n=1}^{\frac{m-(2i+1)}{2}} C_{2n+i} \le C_m \,,$$

or

$$\sum_{n=1}^{\frac{m+1}{2}} C_{2n-1} - 2 \sum_{i=0}^{\frac{m-3}{2}} \sum_{n=1}^{\frac{m-(2i+1)}{2}} C_{2n+i} \ge 0.$$

As x > 1, in the angle variable $\omega = \operatorname{arccosh} x > 0$, we can write $C_m(\cosh \omega) = 4(x-1)\sinh(m\omega)/\sinh\omega$. Consequently, the previous inequality is reduced to

$$\sum_{n=1}^{\frac{m+1}{2}} \sinh((2n-1)\omega) - 2\sum_{i=0}^{\frac{m-3}{2}} \sum_{n=1}^{\frac{m-(2i+1)}{2}} \sinh((2n+i)\omega) \ge 0.$$

If we use the closed formulas for the expansions $\sum_k \sinh(k\omega)$, $\sum_k \sinh(2k\omega)$, and $\sum_k \sinh((2k-1)\omega)$, then we can write the previous inequality as

$$\frac{4(x-1)\sinh\frac{(m+1)\omega}{2}}{\sinh^2\omega}\left(\sinh\frac{(m+1)\omega}{2} - 2\frac{\sinh\frac{(m-1)\omega}{4}\sinh\frac{(m+1)\omega}{4}}{\sinh\frac{\omega}{2}}\right) \ge 0.$$

And, for x > 1, this is true if and only if

$$\frac{2\sinh\frac{(m-1)\omega}{4}\sinh\frac{(m+1)\omega}{4}}{\sinh\frac{(m+1)\omega}{2}} \le \sinh\frac{\omega}{2}.$$
(43)

In this way, for any value of m, we get contractivity if and only if $\omega \geq \omega_m^{(c)}$, where $\omega_m^{(c)}$ is the unique positive root of the corresponding equality equation. Finally, going back to variable s, we get contractivity if and only if $s \leq s_m^{(c)} := 1/(\cosh \omega_m^{(c)} - 1)$. Computing the limit in (43), when $m \to \infty$, we get contractivity if and only if

$$e^{-\omega/2} \le \sinh \frac{\omega}{2}$$
.

And this is true if and only if $\omega \ge \log 3$. This is $x = \cosh \omega \ge 5/3$ or $s \le 3/2$.

2. In the even case, the norm in (38) is obtained from the sum of the elements in any of the two central symmetric rows. The diagonal element $P_m + \sum_{n=1}^{\frac{m}{2}-1} C_{2n}$ in any of this central rows can be written in closed form as

$$P_m + \sum_{n=1}^{\frac{m}{2}-1} C_{2n} = -U_m + \sum_{n=1}^{\frac{m}{2}} C_{2n} = -\frac{\sinh((m+1)\omega)}{\sinh\omega} + \frac{4(x-1)\sinh\frac{m\omega}{2}\sinh\frac{(m+2)\omega}{2}}{\sinh^2\omega},$$

where $\omega = \operatorname{arccosh} x$ and x = 1 + 1/s. Again, the unique positive root s_m of this central diagonal element lies in the interval [1, 3/2). For any value of m, this root can easily be obtained from the simplified equation

$$\sqrt{1+2s} = 4 \frac{\sinh \frac{m\omega}{2} \sinh \frac{(m+2)\omega}{2}}{\sinh((m+1)\omega)}$$

Observe that this sequence of roots $(s_m)_m$ increasingly converges to the limit value $s_{\infty} = 3/2$. When $s \in (0, s_m)$ the diagonal element $P_m + \sum_{n=1}^{\frac{m}{2}-1} C_{2n}$ is positive, and,

from (38), we easily obtain $||A_m||_{\infty} < 1$. On the other hand, when $s \in [s_m, \infty)$, the diagonal element is negative, and, from (38), the inequality for contractivity is

$$||A_m||_{\infty} = \frac{1}{U_m} \left(-P_m - \sum_{n=1}^{\frac{m}{2}-1} C_{2n} + \sum_{i=1}^{\frac{m}{2}-1} \sum_{n=1+i}^{m-i} C_n + \sum_{n=1}^{m/2} C_{2n-1} \right) \le 1.$$

This is equivalent to

$$-\sum_{n=1}^{\frac{m}{2}-1} C_{2n} + \sum_{i=1}^{\frac{m}{2}-1} \sum_{n=1+i}^{m-i} C_n + \sum_{n=1}^{m/2} C_{2n-1} \le C_m \,,$$

or

$$\sum_{n=1}^{\frac{m}{2}} C_{2n} - \sum_{i=1}^{\frac{m}{2}-1} \sum_{n=1+i}^{m-i} C_n - \sum_{n=1}^{m/2} C_{2n-1} \ge 0$$

As in the odd case, now we can use the angle variable ω to reduce the previous inequality.

$$\sum_{n=1}^{\frac{m}{2}}\sinh(2n\omega) - \sum_{i=1}^{\frac{m}{2}-1}\sum_{n=1+i}^{m-i}\sinh(n\omega) - \sum_{n=1}^{m/2}\sinh((2n-1)\omega) \ge 0.$$

Finally, the closed formulas for the expansions $\sum_k \sinh(k\omega)$, $\sum_k \sinh(2k\omega)$, and $\sum_k \sinh((2k-1)\omega)$, allow us to reduce the inequality to

$$\frac{4(x-1)}{\sinh\omega} \left(\frac{\sinh\frac{m\omega}{2}}{\sinh\omega} \left(\sinh\frac{(m+2)\omega}{2} - \sinh\frac{m\omega}{2} \right) - \frac{\sinh\frac{(m+1)\omega}{2}\sinh\frac{m\omega}{4}\sinh\frac{(m-2)\omega}{4}}{\sinh^2\frac{\omega}{2}} \right) \ge 0.$$

And, for x > 1, this is true if and only if

$$\frac{\sinh\frac{m\omega}{2}\left(\sinh\frac{(m+2)\omega}{2} - \sinh\frac{m\omega}{2}\right)}{\sinh\frac{(m+1)\omega}{2}\sinh\frac{m\omega}{4}\sinh\frac{(m-2)\omega}{4}} \ge \frac{\sinh\omega}{\sinh^2\frac{\omega}{2}}.$$
(44)

In this way, for any value of m, we get contractivity if and only if $\omega \ge \omega_m^{(c)}$, where $\omega_m^{(c)}$ is the unique positive root of the corresponding equality equation in (44). Finally, going back to variable s, we get contractivity if and only if $s \leq 1/(\cosh \omega_m^{(c)} - 1)$. С

Computing the limit in (44), when
$$m \to \infty$$
, we get contractivity if and only if

$$2\left(-1+e^{\omega}\right) \geq \frac{\sinh\omega}{\sinh^2\frac{\omega}{2}}$$

And this is true if and only if $\omega \ge \log 3$. This is $x = \cosh \omega \ge 5/3$ or $s \le 3/2$.

Proof of Theorem 2

1. For $m \in \{1, 2, 3\}$ the proof is straightforward.

2. For $m \in \mathbb{N}$, m > 3, the proof is straightforward from the previous lemma. The inequality for contractivity in the variable $s = 1/(\cosh \omega - 1)$ is

$$s \le s_m^{(c)} := \frac{1}{\cosh \omega_m^{(c)} - 1}$$

where $\omega_m^{(c)}$ is the unique positive root of the equation from (43), if *m* is odd, or from (44) if *m* is even.

3. It is also straightforward from the previous lemma. In the limit, when $m \to \infty$, we get inequality (41) from (43), and inequality (42) from (44). These two inequalities (41) and (42) are equivalent if $\omega > 0$, and they are true if and only if $\omega \ge \log 3$. In the variable s this is equivalent to the known restriction $s \le 3/2$.

6.3 Proof of the remaining results

Proof of Proposition 2

We divide the proof into two parts, the case $x \in (-1, 1)$ and the case x > 1.

1. If $x \in (-1, 1)$, then the angular variable $\omega = \arccos x, \omega \in (0, \pi)$, and equality (30), allow us to write the polynomial P_m in closed form as

$$P_m(\cos\omega) = \frac{2\sin((m-1)\omega) - 4\sin(m\omega) + \sin((m+1)\omega)}{\sin\omega}$$
(45)

If we convert all the angles in the numerator to the angle $m\omega$, we can write

$$=\frac{(3\cos\omega-4)\sin(m\omega)}{\sin\omega}-\cos(m\omega)\,,\tag{46}$$

Then, from (46), we get that the roots of $P_m(\cos \omega)$ in $(0, \pi)$ are the roots of the equation

$$\tan(m\omega) = \frac{\sin\omega}{3\cos\omega - 4}, \quad \omega \in (0,\pi).$$
(47)

The function on the right hand side, $f(\omega) := \sin \omega/(3 \cos \omega - 4)$, is continuous and bounded in the interval $[0, \pi]$. It is decreasing in the interval $(0, 2 \arctan(1/\sqrt{7}))$ and increasing in the interval $(2 \arctan(1/\sqrt{7}), \pi)$. Its maximum value $f(0) = f(\pi) = 0$ is obtained in the boundary, while the minimum value is $f(2 \arctan(1/\sqrt{7})) =$ $-1/\sqrt{7}$ (see Figure 4). Thus, we can affirm that $\tan(m\omega)$ meets m-1 times the function $f(\omega)$, and consequently there are m-1 roots $0 < \omega_{m-1}^m < \omega_{m-2}^m < \cdots < \omega_1^m < \pi$, in the interval $(0, \pi)$. Now, going back to the variable $x = \cos \omega$, we can affirm that the polynomial $P_m(x)$ has m-1 roots $x_i^m = \cos \omega_i^m$ in the interval (-1, 1). As the cosine function is decreasing in the interval $(0, \pi)$, we can write the m-1 roots of $P_m(x)$ as

$$-1 < x_1^m < x_2^m < \dots < x_{m-1}^m < 1$$
.



Figure 4: Roots $0 < \omega_{m-1}^m < \omega_{m-2}^m < \cdots < \omega_1^m < \pi$, of equation (47)

2. If x > 1, we consider the variable $\omega = \operatorname{arccosh} x$, $\omega \in (0, \infty)$. Again, with the help of equality (30), we can write the polynomial P_m in closed form as

$$P_m(\cosh(\omega)) = \frac{2\sinh((m-1)\omega) - 4\sinh(m\omega) + \sinh((m+1)\omega)}{\sinh\omega}$$
(48)

$$=\frac{(3\cosh\omega-4)\sinh(m\omega)}{\sinh\omega}-\cosh(m\omega)\,,\tag{49}$$

where, as in the previous case, we have rewritten the numerator in the terms of the angle $m\omega$. Then, from (49), the roots of $P_m(\cosh(\omega))$ in $(0,\infty)$ are the roots of the equation

$$\coth(m\omega) = \frac{3\cosh\omega - 4}{\sinh\omega}, \qquad \omega \in (0,\infty).$$
(50)

The function on the right, $g(\omega) := (3 \cosh \omega - 4) / \sinh \omega$, is continuous and increasing in the interval $(0, \infty)$ to the limit value of 3. On the left, for any value of m, the function $\coth(m\omega)$ is continuous and decreasing to the limit value of 1 (see Figure 5). Thus, we can affirm that, for any value of m, equation (50) has a unique root ω_m in the interval $(0, \infty)$. Now, going back to the variable $x = \cosh \omega$, we can affirm that the polynomial $P_m(x)$ has a unique root $x_m = \cosh \omega_m$ in the interval $(1, \infty)$.



Figure 5: Left hand side and right hand side of equation (50)

In the limit, when m tends to infinity, for any value of $\omega > 0$, we have $\operatorname{coth}(m\omega) \searrow 1$. Consequently, from (50), the sequence of roots (ω_m) decreasingly converges to the limit value $\omega_{\infty} = \log(2 + \sqrt{2})$, this is the positive solution of the limit equation $3 \cosh \omega - 4 = \sinh \omega$. Now, in the variable $x = \cosh \omega$, we can affirm that each polynomial $P_m(x)$ has a positive root $x_m = \cosh \omega_m$ in the interval $(1, \infty)$. As the cosh function is increasing in the interval $(0, \infty)$, the sequence of roots (x_m) decreasingly converges to the limit value $x_{\infty} = (6 + \sqrt{2})/4$.

$$x_{\infty} = \cosh \omega_{\infty} = \cosh \log(2 + \sqrt{2}) = (6 + \sqrt{2})/4 \approx 1.85355.$$

References

- DESOER, C., AND HANEDA, H. The measure of a matrix as a tool to analyze computer algorithms for circuit analysis. *IEEE Transactions on Circuit Theory 19*, 5 (1972), 480–486.
- [2] FARAGÓ, I., KOROTOV, S., AND SZABÓ, T. Non-negativity preservation of the discrete nonstationary heat equation in 1d and 2d. Aplimat-Journal of applied mathematics 3, 2 (2010), 61.
- [3] FARAGÓ, I., AND PALENCIA, C. Sharpening the estimate of the stability constant in the maximum-norm of the crank-nicolson scheme for the one-dimensional heat equation. Applied numerical mathematics 42, 1-3 (2002), 133-140.
- [4] FERRACINA, L., AND SPIJKER, M. N. Stepsize restrictions for the total-variationdiminishing property in general runge-kutta methods. SIAM Journal on Numerical Analysis 42, 3 (2004), 1073–1093.
- [5] GOTTLIEB, S., AND GOTTLIEB, L.-A. J. Strong stability preserving properties of runge-kutta time discretization methods for linear constant coefficient operators. *Journal of Scientific Computing* 18, 1 (2003), 83–109.
- [6] GOTTLIEB, S., KETCHESON, D. I., AND SHU, C.-W. Strong stability preserving Runge-Kutta and multistep time discretizations. World Scientific, 2011.
- [7] GROSS, J. L., MANSOUR, T., TUCKER, T. W., AND WANG, D. G. Root geometry of polynomial sequences ii: Type (1, 0). Journal of Mathematical Analysis and Applications 441, 2 (2016), 499–528.
- [8] HIGUERAS, I. On strong stability preserving time discretization methods. Journal of Scientific Computing 21, 2 (2004), 193–223.
- [9] HIGUERAS, I., AND ROLDÁN, T. Strong stability preserving properties of composition runge-kutta schemes. *Journal of Scientific Computing 80*, 2 (2019), 784–807.
- [10] HIGUERAS, I., AND ROLDÁN, T. On the positivity and contractivity of crank-nicolson method applied to the periodic diffusion problem. ArXiv (2022).
- [11] HORVÁTH, R. Maximum norm contractivity in the numerical solution of the onedimensional heat equation. Applied numerical mathematics 31, 4 (1999), 451–462.
- [12] HORVÁTH, Z. On the positivity step size threshold of runge-kutta methods. Applied Numerical Mathematics 53, 2-4 (2005), 341–356.

- [13] HUNDSDORFER, W., AND VERWER, J. Numerical Solution of Time-Dependent Advection-Diffusion-Reaction Equations. Springer, 2003.
- [14] KRAAIJEVANGER, J. Contractivity of Runge-Kutta methods. BIT Numerical Mathematics 31, 3 (1991), 482–528.
- [15] KRAAIJEVANGER, J. F. B. M. Contractivity of runge-kutta methods. BIT Numerical Mathematics 31, 3 (1991), 482–528.
- [16] KRAAIJEVANGER, J. F. B. M. Maximum norm contractivity of discretization schemes for the heat equation. Applied numerical mathematics 9, 6 (1992), 475–492.
- [17] LIU, L. L., AND WANG, Y. A unified approach to polynomial sequences with only real zeros. Advances in Applied Mathematics 38, 4 (2007), 542–560.
- [18] NÜSSLEIN, S., RANOCHA, H., AND KETCHESON, D. I. Positivity-preserving adaptive runge-kutta methods. *Communications in Applied Mathematics and Computational Science* 16, 2 (2021), 155–179.
- [19] QI, F., CERNANOVÁ, V., AND SEMENOV, Y. S. Some tridiagonal determinants related to central delannoy numbers, the chebyshev polynomials, and the fibonacci polynomials. *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys* 81, 1 (2019), 123–136.
- [20] Rózsa, P. Linear Algebra and its Applications. Muszaki Konyvkiado, (in Hungarian), 1976.
- [21] SHU, C.-W. Total-variation-diminishing time discretizations. SIAM Journal on Scientific and Statistical Computing 9, 6 (1988), 1073–1084.
- [22] SÖDERLIND, G. The logarithmic norm. history and modern theory. BIT Numerical Mathematics 46, 3 (2006), 631–652.
- [23] SPIJKER, M. N. Stepsize restrictions for stability of one-step methods in the numerical solution of initial value problems. *Mathematics of Computation* 45, 172 (1985), 377–392.
- [24] STRÖM, T. On logarithmic norms. SIAM Journal on Numerical Analysis 12, 5 (1975), 741–753.
- [25] SZABÓ, T. Qualitative properties of some discretized partial differential equations and reliable fuel cell modelling. PhD thesis, Eötvös Loránd University, Budapest, Hungary, 2011.
- [26] THOMÉE, V. Finite difference methods for linear parabolic equations. Handbook of numerical analysis 1 (1990), 5–196. in: P. Ciarlet, J. Lions (Eds.), Handbook of Numerical Analysis, Vol. 1, 1990, North-Holland, Amsterdam.
- [27] VAN DE GRIEND, J., AND KRAAIJEVANGER, J. Absolute monotonicity of rational functions occurring in the numerical solution of initial value problems. *Numerische Mathematik* 49, 4 (1986), 413–424.

[28] WANG, D. G., AND ZHANG, J. Root geometry of polynomial sequences iii: Type (1,1) with positive coefficients. arXiv preprint arXiv:1712.06105 (2017).