A gallery of diagonal stability conditions with structured matrices (and review papers)

Zhiyong Sun

Control Systems Group, Department of EE, TU Eindhoven, the Netherlands

Abstract

This note presents a summary and review of various conditions and characterizations for matrix stability (in particular diagonal matrix stability) and matrix stabilizability.

Keywords: Matrix stability, matrix stabilizability, diagonal stability.

1. Definitions and notations

- A square real matrix is a **Z-matrix** if it has nonpositive off-diagonal elements.
- A Metzler matrix is a real matrix in which all the off-diagonal components are nonnegative (equal to or greater than zero).
- A Z-matrix with positive principal minors is an M-matrix.
 - Note: There are numerous equivalent characterizations for M-matrix (Fiedler and Ptak, 1962; Plemmons, 1977). A more commonly-used condition is the following: A matrix $A \in \mathbb{R}^{n \times n}$ is called an M-matrix, if its non-diagonal entries are non-positive and its eigenvalues have positive real parts.
- A square matrix A is (**positive**) stable if all its eigenvalues have positive parts. Equivalently, a square matrix A is (positive) stable iff there exists a positive definite matrix D such that $AD + DA^T$ is positive definite.
 - Note: in control system theory we often define stable matrix as the set of square matrices whose eigenvalues have negative real parts (a.k.a. Hurwitz matrix). The two definitions of stable matrices will be distinguished in the context.
- A square complex matrix is a **P-matrix** if it has positive principal minors.
- A square complex matrix is a P_0^+ -matrix if it has nonnegative principal minors and at least one principal minor of each order is positive.
- A real square matrix A is **multiplicative D-stable** (in short, **D-stable**) if DA is stable for every positive diagonal matrix D.

- A square matrix A is called **totally stable** if any principal submatrix of A is D-stable.
- A real square matrix A is said to be **additive D-stable** if A + D is stable for every nonnegative diagonal matrix D.
- A real square matrix A is said to be Lyapunov diagonally stable if there exists a positive diagonal matrix D such that $AD + DA^T$ is positive definite.
 - Note: Lyapunov diagonally stable matrices are often referred to as just diagonally stable matrices or as Volterra–Lyapunov stable, or as Volterra dissipative in the literature (see e.g., (Logofet, 2005)).
- A matrix A = {a_{ij}} ∈ ℝ^{n×n} is generalized row-diagonally dominant, if there exists x = (x₁, x₂, ···, x_n) ∈ ℝⁿ with x_i > 0, ∀i, such that

$$|a_{ii}|x_i > \sum_{j=1, j \neq i}^n |a_{ij}|x_j, \forall i = 1, 2, \cdots, n.$$
(1)

A matrix A = {a_{ij}} ∈ ℝ^{n×n} is generalized *column*-diagonally dominant, if there exists x = (x₁, x₂, ··· , x_n) ∈ ℝⁿ with x_i > 0, ∀i, such that

$$|a_{jj}|x_j > \sum_{i=1, i \neq j}^n |a_{ij}|x_i, \forall j = 1, 2, \cdots, n.$$
 (2)

- Note: the set of generalized *column*-diagonally dominant matrices is equivalent to the set of generalized *row*-diagonally dominant matrices (Varga, 1976; Sun et al., 2021). They are also often referred to as **quasi-diagonally dominant** matrices (Kaszkurewicz and Bhaya, 2012).
- For a real matrix A = {a_{ij}} ∈ ℝ^{n×n}, we associate it with a comparison matrix M_A = {m_{ij}} ∈ ℝ^{n×n}, defined by

$$m_{ij} = \begin{cases} |a_{ij}|, & \text{if } j = i; \\ -|a_{ij}|, & \text{if } j \neq i. \end{cases}$$

A given matrix A is called an **H-matrix** if its comparison matrix M_A is an M-matrix.

- The set of H-matrix is equivalent to the set of quasi-diagonally dominant matrices (Kaszkurewicz and Bhaya, 2012; Sun et al., 2021).
- A square matrix A is **diagonally stabilizable** if there exists a diagonal matrix D such that DA is stable.

Note: Many definitions above for real matrices also carry over to complex matrices; the distinction between real and complex matrices will be made clear in the context.

2. Conditions for diagonally stabilizable matrices

A key motivating question: *Given a square matrix A, can we find a diagonal matrix D such that the matrix DA is stable?*

Fisher and Fuller (Fisher and Fuller, 1958) proved the following result:

Theorem 1. (*Fisher and Fuller, 1958*) If P is a real $n \times n$ matrix fulfilling the condition:

• (A): P has at least one sequence of non-zero principal minors M_k of every order $k = 1, 2, \dots, n$, such that M_{k-1} is one of the k first principal minors of M_k ;

then there exists a real diagonal matrix D such that the characteristic equation of DP is stable.

The Fisher-Fuller theorem is also formulated as the following alternative version:

Theorem 2. Let P be an $n \times n$ real matrix all of whose leading principal minors are positive. Then there is an $n \times n$ positive diagonal matrix D such that all the roots of DP are positive and simple.

Fisher later gave a simple proof for a similar yet stronger result (Fisher, 1972).

Theorem 3. (Fisher, 1972) If P is an $n \times n$ real matrix that has at least one nested set of principal minors, M_k , such that $(-1)^k M_k > 0, \forall k = 1 \cdots, n$, then there exists a real diagonal matrix D with positive diagonal elements such that the characteristic roots of DP are all real, negative, and distinct.

Remark 1. Some remarks on the conditions of diagonally stabilizable matrices are in order.

- The above theorems involve determining the sign of (at least) one nested set of principal minors. In (Johnson et al., 1997), sufficient conditions are determined for an n-by-n zero-nonzero pattern to allow a nested sequence of nonzero principal minors. In particular, a method is given to sign such a pattern so that it allows a nested sequence of k-by-k principal minors with $sign(-1)^k$ for $k = 1, \dots, n$.
- The condition in the Fisher-Fuller theorem appears as a sufficient condition for matrix diagonal stabilizability. A necessary condition for matrix diagonal stabilizability is: for each order $k = 1 \cdots$, n, at least one $k \times k$ principal minor of P is non-zero. It is unclear what would be **the** necessary and sufficient condition.

Ballantine (Ballantine, 1970) extended the above Fisher-Fuller theorem to the complex matrix case.

Theorem 4. (*Ballantine*, 1970) Let A be an $n \times n$ complex matrix all of whose leading principal minors are nonzero. Then there is an $n \times n$ complex diagonal matrix D such that all the roots of DA are positive and simple.

Remark 2. It is shown in (Hershkowitz, 1992) the above Ballantine theorem cannot be strengthened by replacing "complex diagonal matrix D" by "positive diagonal matrix D". A counterexample is shown in (Hershkowitz, 1992) involving a 2×2 complex matrix A with positive leading principal minors that there exists no positive diagonal matrix D such that the eigenvalues of DA are positive.

A related problem to characterize diagonal stabilizable matrix is the **Inverse Eigen-value Problem** (IEP), and Friedland (Friedland, 1977) proved the following theorem.

Theorem 5. (*Friedland, 1977*) Let A be a given $n \times n$ complex valued matrix. Assume that all the principal minors of A are different from zero. Then for any specified set $\lambda = \{\lambda, \dots, \lambda_n\} \in \mathbb{C}^n$ there exists a diagonal complex valued matrix D such that the spectrum of AD is the set λ . The number of such D is finite and does not exceed n!. Moreover, for almost all λ the number of the diagonal matrices D is exactly n!.

Remark 3. The Friedland theorem of the IEP problem in the complex matrix case cannot be directly carried over to the real case. Further, it is easy to show with a counterexample of a 2×2 matrix that eigenvalue positionability in the real case cannot always be guaranteed, even with nonzero principal minors.

In (Hershkowitz, 1992) the following two theorems are proved.

Theorem 6. (*Hershkowitz, 1992*) Let A be a **complex** square matrix with positive leading principal minors, and let ϵ be any positive number. Then there exists a positive diagonal matrix D such that the eigenvalues of DA are simple, and the argument of every such eigenvalue is less in absolute value than ϵ .

Theorem 7. (*Hershkowitz, 1992*) Let A be a complex square matrix with real principal minors and positive leading principal minors. Then there exists a positive diagonal matrix D such that DA has simple positive eigenvalues.

Remark 4. The above theorems all present certain sufficient conditions to characterize diagonally stabilizable matrix and the IEP problem, and they are not necessary. A necessary condition for the diagonal matrix D to exist is that for each order i, at least one $i \times i$ principal minor of A is nonzero. However, a full characterization (with necessary and sufficient condition) for diagonally stabilizable matrix still remains an open problem.

A variation of the diagonal matrix stabilization problem is the following:

• Problem (*): Given a real square matrix G, find a real diagonal matrix D such that the product GD is Hurwitz together with all its principal submatrices.

Surprisingly, a necessary and sufficient condition exists for solving the above problem as shown in (Locatelli and Schiavoni, 2012). Let $\mathcal{M} := \{1, 2, \dots, m\}$ and $\mathcal{F} := \{f | f \subset \mathcal{M}\}$. Further, for any $m \times m$ matrix Δ , denote by $\Delta(f)$ the principal submatrix obtained from it after removing the rows and columns with indexes in $f, f \in \mathcal{F}$. The main result of (Locatelli and Schiavoni, 2012) proves the following: **Theorem 8.** (Locatelli and Schiavoni, 2012) Problem (*) admits a solution if and only if

$$det(G(f))det(G_D(f)) > 0, \forall f \in \mathcal{F}$$
(3)

where $G_D = diag\{g_{ii}\}$. Moreover, if the above condition is satisfied, then there exists $\bar{\epsilon} > 0$ such that, for any given $\epsilon \in (0, \bar{\epsilon})$, the matrix

$$D := G_D Z(\epsilon), Z(\epsilon) := -diag\{\epsilon^i\}$$
(4)

solves the stabilization problem (*).

3. Conditions for diagonally stable matrices

We give a short summary of available conditions for diagonally stable matrices (excerpts from (Barker et al., 1978), (Cross, 1978) and (Hershkowitz, 2006)).

- (Barker et al., 1978) Lyapunov diagonally stable matrices are P-matrices.
- (Barker et al., 1978) A matrix A being Lyapunov diagonally stable is equivalent to that there exists a positive diagonal matrix D such that $x^T DAx > 0$ for all nonzero vectors x.
- (Barker et al., 1978) A 2 × 2 real matrix is Lyapunov diagonally stable if and only if it is a P-matrix.
- (Cross, 1978) For a given Lyapunov diagonally stable matrix *P*, all principal submatrices of *P* are Lyapunov diagonally stable.
- (Barker et al., 1978) A real square matrix A is Lyapunov diagonally stable if and only if for every nonzero real symmetric positive semidefinite matrix H the matrix HA has at least one positive diagonal element.
 - Note: this result is termed the BBP theorem, and is proved again in (Shorten et al., 2009) with a simpler proof.
- (Cross, 1978) The set of Lyapunov diagonally stable matrices is a strict subset of multiplicative D-stable matrices.
- (Cross, 1978) The set of Lyapunov diagonally stable matrices is a strict subset of additive D-stable matrices.
 - Note: Multiplicative D-stable and additive D-stable matrices are not necessarily diagonally stable.
- A Z-matrix is Lyapunov diagonally stable if and only if it is a P -matrix (that is, it is an M-matrix).
- A non-singular H-matrix with nonnegative diagonal parts is Lyapunov diagonally stable.
- A quasi-diagonal dominant matrix with nonnegative diagonal parts is Lyapunov diagonally stable. Note the equivalence of Hurwitz H-matrix and quasi-diagonal dominant matrix (Sun et al., 2021).

The following facts are shown in (Cross, 1978) and (Kaszkurewicz and Bhaya, 2012):

- For normal matrices and within the set Z, D-stability, additive D-stability, and diagonal stability are all equivalent to matrix stability.
- If a matrix A is Hurwitz stable, D-stable, or diagonally stable, then the matrices A^T and A^{-1} also have the corresponding properties.

In (Shorten and Narendra, 2009) Shorten and Narendra showed the following necessary and sufficient condition for matrix diagonal stability (an alternative proof of the theorem of Redheffer via the KYP lemma):

Theorem 9. (Shorten and Narendra, 2009) and (Redheffer, 1985) Let $A \in \mathbb{R}^{n \times n}$ be a Hurwitz matrix with negative diagonal entries. Let A_{n-1} denote the $[n - 1 \times n - 1]$ leading sub-matrix of A, and B_{n-1} denote the corresponding block of A^{-1} . Then, the matrix A is diagonally stable, if and only if there is a common diagonal Lyapunov function for the LTI systems $\Sigma_{A_{n-1}}$ and $\Sigma_{B_{n-1}}$.

The above theorem involves finding a common diagonal Lyapunov function for a set of LTI systems, which may be restrictive and computationally demanding in practical applications especially when the dimension of the matrix A is large.

Theorem 1.	
a. In general	
	$\mathcal{L} \Leftarrow \mathcal{A} \mathrel{\Rightarrow} \mathcal{P} \mathrel{\Rightarrow} \mathcal{S},$
b. For Z-matrices, i.e., matrices with nonpositive off-diagonal entries,	
	$\mathcal{A} \Leftrightarrow \mathcal{L} \Leftrightarrow \mathcal{P} \Leftrightarrow \mathcal{S}.$
c. For symmetric matrices	
	$\mathcal{A} \Leftrightarrow \mathcal{L} \Leftrightarrow \mathcal{P} \mathrel{\Rightarrow} \mathcal{G}.$
d. For triangular matrices	
	$\mathcal{A} \Leftrightarrow \mathcal{L} \Leftrightarrow \mathcal{P} \mathrel{\Rightarrow} \mathcal{S}.$
e. for normal matrices	
	$\mathcal{A} \Leftrightarrow \mathcal{L} \mathrel{\Rightarrow} \mathcal{P} \mathrel{\Rightarrow} \mathcal{S}.$
The absence of an implication in the above relations means that a counterexample exists.	

Figure 1: Relations of matrix stability under different matrix types: the main theorem in (Berman and Hershkowitz, 1983)

4. Relations of matrix stability and diagonal stability

The paper (Berman and Hershkowitz, 1983) characterizes the relations of certain special matrices for matrix diagonal stability. They define

- $\mathscr{A} = \{A : \text{there exists a positive definite diagonal matrix } D \text{ such that } AD + DA^T \text{ is positive definite}\};$ i.e., \mathscr{A} denotes the set of diagonally stable matrices;
- $\mathscr{L} = \{A : \text{there exists a positive definite matrix } D \text{ such that } AD + DA^T \text{ is positive definite}\};$ i.e., \mathscr{L} denotes the set of (positive) stable matrices;
- \$\mathcal{P}\$ = {A : the principle minors of A are positive};
 i.e., \$\mathcal{P}\$ denotes the set of P-matrices;
- $\mathscr{S} = \{A : \text{there exists a positive vector } x \text{ such that } Ax \text{ is positive}\};$ i.e., \mathscr{S} denotes the set of semipositive matrices.

The main result of (Berman and Hershkowitz, 1983) is cited and shown in Fig. 1. In general, these different sets of structured matrices are not equivalent, and the set \mathscr{A} is a subset of the other sets. However, for Z-matrices, these sets are equivalent. In particular, for the case of Z-matrices, the characterizations of these sets give equivalent conditions for M-matrices (upon a sign change). Note there are yet many more conditions to characterize M-matrices; see e.g., (Plemmons, 1977).

The review paper (Hershkowitz, 1992) presents the implication relations between matrix stability conditions, and the equivalent relations of matrix stabilities for Z-matrices, as cited in Figs. 2 and 3. Again, as shown in Figs. 3, for Z-matrices, all the stability types are equivalent.



Figure 2: The implication relations between matrix stability conditions, cited from (Hershkowitz, 1992)



Figure 3: For Z-matrices, all the stability types are equivalent. Cited from (Hershkowitz, 1992)



Figure 4: Relations among matrix stabilities. Cited from (Logofet, 2005, Fig.2)

The survey paper (Logofet, 2005) presents some beautiful flower-shaped characterizations of the relations among matrix stabilities, as cited in Figs. 4 and 5.



Figure 5: Petals of sign-stable matrices within the Flower. Cited from (Logofet, 2005, Fig.4)

5. Stability conditions with submatrices and Schur complement

Stability conditions of 'structured' matrices often involve stability properties of submatrices, which employ block submatrices and their Schur complements to determine stability.

In (Narendra and Shorten, 2010), Narendra and Shorten presented necessary and sufficient conditions to characterize if a given Metzler matrix is Hurwitz, based on the fact that a Hurwitz Metzler matrix is diagonally stable. These conditions are generalized in (Souza et al., 2017). We recall some main stability criteria from (Souza et al., 2017).

Lemma 1. Let $A \in \mathbb{R}^{n \times n}$ be a Metzler matrix partitioned in blocks of compatible dimensions as $A = [A_{11}, A_{12}; A_{21}, A_{22}]$ with A_{11} and A_{22} being square matrices. Then the following statements are equivalent.

- A is Hurwitz stable.
- A₁₁ and its Schur complement A/A₁₁ := A₂₂ A₂₁A₁₁⁻¹A₁₂ are Hurwitz stable Metzler matrices.
- A₂₂ and its Schur complement A/A₂₂ := A₁₁ A₁₂A₂₂⁻¹A₂₁ are Hurwitz stable Metzler matrices

Remark 5. Some remarks are in order.

- For a structured matrix, the property that its Schur complements also preserve the same stability and structure properties is termed Schur complement stability property. Other types of structured matrices that have Schur complement stability property include symmetric matrices, triangular matrices, and Schwarz matrices. See (Souza et al., 2017).
- The result on M-matrix in Lemma 1 can be generalized to H-matrix: Let A be a H-matrix partitioned in blocks of compatible dimensions as $A = [A_{11}, A_{12}; A_{21}, A_{22}]$ with A_{11} and A_{22} being square matrices. If A is Hurwitz stable, then A_{11} and its Schur complement $A/A_{11} := A_{22} A_{21}A_{11}^{-1}A_{12}$ are Hurwitz stable H matrices, or A_{22} and its Schur complement $A/A_{22} := A_{11} A_{12}A_{22}^{-1}A_{21}$ are Hurwitz stable H matrices.
- Schur complement and its closure property for several structured matrices (including diagonal matrices, triangular matrices, symmetric matrices, P-matrices, diagonal dominant matrices, M-matrices etc.) are discussed in (Zhang, 2006, Chap. 4).

6. Application examples of matrix diagonal stability conditions

The Fisher-Fuller theorem on diagonal matrix stabilizability (Theorem 1 and its variations) has been rediscovered several times by the control system community, and has been applied in solving distributed stabilization and decentralized control problems in practice. This section reviews two application examples.

6.1. Conditions for decentralized stabilization

- In (Corfmat and Morse, 1973) Corfmat and Morse solved the following problem:
- For given and fixed real matrices A and P, find (if possible) a non-singular diagonal matrix D such that I + ADP is Schur stable (i.e., all eigenvalues of I + ADP are located within the unit circle in the complex plane.

To solve the above problem they proved the following:

Theorem 10. If A is an $n \times n$ strongly non-singular matrix, then there exists a diagonal matrix D such that (I + DA) is Schur stable.

Note: in (Corfmat and Morse, 1973) a matrix is termed **strongly non-singular**, if its all *n* leading principal minors are nonzero.

Theorem 11. If A is a fixed non-singular matrix, then there exists a permutation matrix P such that PA is strongly non-singular.

Solution to decentralized stabilization: the non-singularity of A is a necessary and sufficient condition for the existence of a permutation matrix P and a non-singular diagonal matrix D such that (I + ADP) is Schur stable.

6.2. Distributed stabilization of persistent formations

In (Yu et al., 2009), the problem on persistent formation stabilization involves studying the stabilizability of the following differential equation

$$\dot{z} = \Delta A z$$

where Δ is a diagonal or possibly block diagonal matrix, and A is a rigidity-like matrix on formation shapes. To solve the formation stabilization problem in (Yu et al., 2009) the following result is employed ((Yu et al., 2009, Theorem 3.2)):

Theorem 12. Suppose A is an $m \times m$ non-singular matrix with every leading principal minor nonzero. Then there exists a diagonal D such that the real parts of the eigenvalues of DA are all negative.

We remark that this is a reformulation of the Fisher-Fuller theorem.

7. A selection of key review papers and books on matrix stability and diagonal stability conditions

- The survey paper (Hershkowitz, 1992) that presents a summary of relevant matrix stability results and the developments, up until 1992.
- The paper (Bhaya et al., 2003) that presents comprehensive discussions and characterizations for various classes of matrix stability conditions.
- The paper (Hershkowitz and Keller, 2003) that studies the relations between positivity of principal minors, sign symmetry and stability of matrices.
- The review paper (Hershkowitz, 2006) that presents an concise overview on matrix stability and inertia.
- The book (Kaszkurewicz and Bhaya, 2012) on matrix diagonal stability in systems and computation.
- The summary paper (Logofet, 2005) that presents a review and some beautiful connections/relations on different matrix stabilities.
- The very long survey paper (Kushel, 2019) that provides a unifying viewpoint on matrix stability, and its historical development.
- The recent book (Johnson et al., 2020) on positive matrix, P-matrix and inverse M-matrix.

References

- Ballantine, C., 1970. Stabilization by a diagonal matrix. Proceedings of the American Mathematical Society 25, 728–734.
- Barker, G., Berman, A., Plemmons, R.J., 1978. Positive diagonal solutions to the Lyapunov equations. Linear and Multilinear Algebra 5, 249–256.
- Berman, A., Hershkowitz, D., 1983. Matrix diagonal stability and its implications. SIAM Journal on Algebraic Discrete Methods 4, 377–382.
- Bhaya, A., Kaszkurewicz, E., Santos, R., 2003. Characterizations of classes of stable matrices. Linear algebra and its applications 374, 159–174.
- Corfmat, J., Morse, A., 1973. Stabilization with decentralized feedback control. IEEE Transactions on Automatic Control 18, 679–682.
- Cross, G., 1978. Three types of matrix stability. Linear algebra and its applications 20, 253–263.
- Fiedler, M., Ptak, V., 1962. On matrices with non-positive off-diagonal elements and positive principal minors. Czechoslovak Mathematical Journal 12, 382–400.

- Fisher, F.M., 1972. A simple proof of the Fisher–Fuller theorem, in: Mathematical Proceedings of the Cambridge Philosophical Society, Cambridge University Press. pp. 523–525.
- Fisher, M.E., Fuller, A., 1958. On the stabilization of matrices and the convergence of linear iterative processes, in: Mathematical Proceedings of the Cambridge Philosophical Society, Cambridge University Press. pp. 417–425.
- Friedland, S., 1977. Inverse eigenvalue problems. Linear Algebra and Its Applications 17, 15–51.
- Hershkowitz, D., 1992. Recent directions in matrix stability. Linear Algebra and its Applications 171, 161–186.
- Hershkowitz, D., 2006. Matrix stability and inertia, in: Handbook of Linear Algebra. Chapman and Hall/CRC, pp. 19–1.
- Hershkowitz, D., Keller, N., 2003. Positivity of principal minors, sign symmetry and stability. Linear algebra and its applications 364, 105–124.
- Johnson, C.R., Maybee, J.S., Olesky, D., Van den Driessche, P., 1997. Nested sequences of principal minors and potential stability. Linear Algebra and its Applications 262, 243–257.
- Johnson, C.R., Smith, R.L., Tsatsomeros, M.J., 2020. Matrix positivity. volume 221. Cambridge University Press.
- Kaszkurewicz, E., Bhaya, A., 2012. Matrix diagonal stability in systems and computation. Springer Science & Business Media.
- Kushel, O.Y., 2019. Unifying matrix stability concepts with a view to applications. SIAM Review 61, 643–729.
- Locatelli, A., Schiavoni, N., 2012. A necessary and sufficient condition for the stabilisation of a matrix and its principal submatrices. Linear algebra and its applications 436, 2311–2314.
- Logofet, D.O., 2005. Stronger-than-Lyapunov notions of matrix stability, or how "flowers" help solve problems in mathematical ecology. Linear Algebra and its Applications 398, 75–100.
- Narendra, K.S., Shorten, R., 2010. Hurwitz stability of metzler matrices. IEEE Transactions on Automatic Control 55, 1484–1487.
- Plemmons, R.J., 1977. M-matrix characterizations. I–nonsingular M-matrices. Linear Algebra and its Applications 18, 175–188.
- Redheffer, R., 1985. Volterra multipliers ii. SIAM Journal on Algebraic and Discrete Methods 6, 612–623.

- Shorten, R., Mason, O., King, C., 2009. An alternative proof of the Barker, Berman, Plemmons (BBP) result on diagonal stability and extensions. Linear Algebra and its Applications 430, 34–40.
- Shorten, R., Narendra, K.S., 2009. On a theorem of Redheffer concerning diagonal stability. Linear algebra and its applications 431, 2317–2329.
- Souza, M., Wirth, F.R., Shorten, R.N., 2017. A note on recursive schur complements, block hurwitz stability of metzler matrices, and related results. IEEE Transactions on Automatic Control 62, 4167–4172.
- Sun, Z., Rantzer, A., Li, Z., Robertsson, A., 2021. Distributed adaptive stabilization. Automatica 129, 109616.
- Varga, R.S., 1976. On recurring theorems on diagonal dominance. Linear Algebra and its Applications 13, 1–9.
- Yu, C., Anderson, B.D.O., Dasgupta, S., Fidan, B., 2009. Control of minimally persistent formations in the plane. SIAM Journal on Control and Optimization 48, 206–233.
- Zhang, F., 2006. The Schur complement and its applications. volume 4. Springer Science & Business Media.