Convergence of Adaptive Mixed Interior Penalty Discontinuous Galerkin Methods for H(curl)-Elliptic Problems

K. Liu¹, M. Tang^{2,}, X. Q. Xing² and L. Q. Zhong²

¹ School of Sciece, East China University of Technology, Nanchang, 330013, China

² School of Mathematical Sciences, South China Normal University, Guangzhou, 510631, China

Abstract. In this paper, we study the convergence of adaptive mixed interior penalty discontinuous Galerkin method for H(curl)-elliptic problems. We first get the mixed model of H(curl)-elliptic problem by introducing a new intermediate variable. Then we discuss the continuous variational problem and discrete variational problem, which based on interior penalty discontinuous Galerkin approximation. Next, we construct the corresponding posteriori error indicator, and prove the contraction of the summation of the energy error and the scaled error indicator. At last, we confirm and illustrate the theoretical result through some numerical experiments.

AMS subject classifications: 65M15,65N12,65N30

Key words: Adaptive mixed interior penalty discontinuous Galerkin methods, Convergence, H(curl)-elliptic problems.

1. Introduction

Let $\Omega \subset \mathbb{R}^3$ be Lipschitz bounded polygonal domain with a single connected boundary $\partial \Omega$. We consider the following H(curl)-elliptic problem

$$\nabla \times \mu \nabla \times \boldsymbol{u} + \kappa \boldsymbol{u} = \boldsymbol{f} \qquad \text{in} \qquad \Omega, \tag{1.1}$$

$$\boldsymbol{u} \times \boldsymbol{n} = 0 \quad \text{on} \quad \partial \Omega, \tag{1.2}$$

where *n* is the unit normal vector of the boundary $\partial \Omega$, $f \in L^2(\Omega)$, μ and κ are piecewise constants is consistent with the initial partition \mathscr{T}_0 for Ω and satisfy $\mu_1 < \mu < \mu_2$ and $\kappa_1 < \kappa < \kappa_2$, here, μ_i and $\kappa_i (i = 1, 2)$ are positive constants. By introducing an auxiliary

^{*}Corresponding author. *Email addresses:* liukai@ecut.edu.cn (K. Liu), mingtang@m.scnu.edu.cn (M. Tang),xingxq@scnu.edu.cn(X. Q. Xing), zhong@m.scnu.edu.cn (L. Q. Zhong)

variable $p = \mu \nabla \times u$, then we get the mixed scheme with the boundary value problem (1.1)-(1.2)

$$\boldsymbol{p} = \boldsymbol{\mu} \nabla \times \boldsymbol{u} \qquad \text{in} \qquad \boldsymbol{\Omega}, \tag{1.3}$$

$$\nabla \times \boldsymbol{p} + \kappa \boldsymbol{u} = \boldsymbol{f} \qquad \text{in} \qquad \Omega, \tag{1.4}$$

$$\boldsymbol{u} \times \boldsymbol{n} = 0 \quad \text{on} \quad \partial \Omega. \tag{1.5}$$

The mixed finite element method is very convenient for processing high-order equations and equations containing two or more unknown functions, which has attracted widespread attention. For mixed finite element method, there are only few research results for Maxwell problem [13] and Maxwell's eigenvalue problem [12, 14, 15].

Adaptive finite element method automatically refines and optimizes meshes according to the singularity of solutions. It is a highly reliable and efficient numerical calculation method. At present, the convergence analysis research of the adaptive mixed finite element method for the elliptic equation is relatively complete. Chen, Holst and Xu [7] proved the convergence analysis of the adaptive mixed finite element algorithm for elliptic equations. Du and Xie [10] proved the convergence analysis of the adaptive mixed finite element algorithm for the convection diffusion equation. However, there are only few research results on the posterior error estimator of Maxwell's equations for the adaptive mixed finite element method. For example, Carstensen and Ma [5] establishes the convergence of adaptive mixed finite element methods for second-order linear non-self-adjoint indefinite elliptic problems. Carstensen, Hoppe, Sharma and Warburton [4] designs and analyzes the posterior error estimation of the adaptive hybrid conforming finite element method of *H*(*curl*)-elliptic problem. Recently, Chung, Yuen and Zhong [8] present a-posteriori error analysis for the staggered discontinuous Galerkin method. As far as we know, there are not any published literatures for the convergence analysis of the adaptive mixed finite element method for the boundary value problem(1.3)-(1.5). Our contributions in this paper are to

- construct a new error estimator, which does not include the negative power of the local mesh size in the jump term for the traditional DG method;
- get the convergence of the Adaptive Mixed Interior Penalty Discontinuous Galerkin (AMIPDG) method by using the similar technique used in [2]. However, this technique in [2] can not be used directly for mixed forms.

We present our main result in the following theorem.

Theorem 1.1. Let $\{\mathcal{T}_k, \mathbf{U}_k, \mathbf{Q}_k, \mathbf{u}_k, \mathbf{p}_k, \eta(\mathbf{u}_k, \mathbf{p}_k; \mathcal{T}_k)\}_{k\geq 0}$ be the sequence of meshes, finite element space, mixed discrete solution and posterior error estimate indicator produced by the **AMIPDG** algorithm. Then there exist constants $\rho > 0$ and $\delta \in (0, 1)$, which depend on marking parameter and the shape regularity of the initial mesh \mathcal{T}_0 , such that

$$\||\boldsymbol{u} - \boldsymbol{u}_{k+1}|\|_{k+1}^2 + \rho \eta^2 (\boldsymbol{u}_{k+1}, \boldsymbol{p}_{k+1}; \mathcal{T}_{k+1}) \le \delta \bigg(\||\boldsymbol{u} - \boldsymbol{u}_k|\|_k^2 + \rho \eta^2 (\boldsymbol{u}_k, \boldsymbol{p}_k; \mathcal{T}_k) \bigg).$$

Therefore, for a given precision, the **AMIPDG** method will terminate after a finite number of operations.

For convenience, we let *C* denote a generic positive constant which may be different at different occurrences and adopt the following notation. The subscripted constant C_i represents a particularly important constant. $a \leq b$ means $a \leq Cb$ for some constants *C* which are independent of mesh sizes.

The rest of this paper is organized as follows. In Section 2, we first present the continuous variational problem, the discrete variational problem, and the procedure of AMIPDG. In Section 3, we first show the upper bound estimate of the error, which is key to the convergence analysis, then we prove the indicator reduction and the convergence of AMIPDG algorithm. In Section 4, we provide some numerical experiments to illustrate the effectiveness of the AMIPDG.

2. Adaptive Mixed interior penalty discontinuous Galerkin method

In this section, we introduce the continuous variational problem, the discrete variational problem of mixed internal penalty discontinuous finite element method, and the procedure of AMIPDG.

2.1. Continuous variational problem

For an open and connected bounded domain $D \subset \mathbb{R}^3$, we denote by $L^2(D)$ (resp. $L^2(D) := (L^2(D))^3$) the spaces of square-integrable functions (resp. vector fields) on D with inner product $(\cdot, \cdot)_{0,D}$. We define the spaces

$$H(curl; D) = \{ u \in L^2(D) : \nabla \times u \in L^2(D) \},\$$

$$H(div; D) = \{ u \in L^2(D) : \nabla \cdot u \in L^2(D) \},\$$

with

$$\begin{aligned} (u,v)_{curl,D} &:= (u,v)_{0,D} + (\nabla \times u, \nabla \times v)_{0,D}, \quad \forall u, v \in H(curl; D), \\ (u,v)_{div,D} &:= (u,v)_{0,D} + (\nabla \cdot u, \nabla \cdot v)_{0,D}, \quad \forall u, v \in H(div; D), \end{aligned}$$

and the induced norm as:

$$\|u\|_{curl,D}^{2} := \|u\|_{0,D}^{2} + \|\nabla \times u\|_{0,D}^{2}, \quad \forall u \in H(curl,D),$$

$$\|u\|_{div,D}^{2} := \|u\|_{0,D}^{2} + \|\nabla \cdot u\|_{0,D}^{2}, \quad \forall u \in H(div,D),$$

respectively, where $\|\cdot\|_{L^2(D)} := (\cdot, \cdot)_D^{1/2}$ denotes the norm of the space $L^2(D)$ or $L^2(D)$. We also define $H_0(curl; D) = \{ v \in H(curl; D) : v \times n = 0 \text{ on } \partial D \}$ in the trace sense.

Next, we first define two space $U := H_0(curl; \Omega), Q := L^2(\Omega)$. Then, the mixed variational problem of the mixed boundary value problem (1.3)-(1.5) reads as: find $(u, p) \in U \times Q$ such that:

$$a(\boldsymbol{p},\boldsymbol{q}) - b(\boldsymbol{u},\boldsymbol{q}) = \ell_1(\boldsymbol{q}), \qquad \forall \boldsymbol{q} \in \boldsymbol{Q},$$
(2.1)

$$d(\mathbf{v}, \mathbf{p}) + c(\mathbf{u}, \mathbf{v}) = \ell_2(\mathbf{v}), \qquad \forall \mathbf{v} \in \mathbf{U}.$$
(2.2)

K Liu et al.

The bilinear forms *a*, *b*, *c* and the functionals $\ell_1(\cdot), \ell_2(\cdot)$ are given by

$$a(\boldsymbol{p},\boldsymbol{q}) := (\boldsymbol{p},\boldsymbol{q}), \tag{2.3}$$

$$b(\boldsymbol{u},\boldsymbol{q}) := (\boldsymbol{\mu} \nabla \times \boldsymbol{u},\boldsymbol{q}), \qquad (2.4)$$

$$c(u,v) := (\kappa u, v), \tag{2.5}$$

$$d(\boldsymbol{\nu}, \boldsymbol{p}) := (\nabla \times \boldsymbol{\nu}, \boldsymbol{p}) \tag{2.6}$$

$$\ell_1(q) := 0, \tag{2.7}$$

$$\ell_2(\mathbf{v}) := (f, \mathbf{v}). \tag{2.8}$$

The operator-theoretic framework involves operator $\mathscr{A}: (U \times Q) \to (U \times Q)^*$ defined by

$$(\mathscr{A}(u,p))(v,q) := a(p,q) - b(u,q) + d(v,p) + c(u,v), \forall u,v \in U, p,q \in Q,$$
(2.9)

where $(\mathbf{Q} \times \mathbf{U})^*$ is the dual spaces of $(\mathbf{Q} \times \mathbf{U})$. Then we can rewrite (2.1)-(2.2) as

$$(\mathscr{A}(u,p))(v,q) = \ell(v,q), \qquad (2.10)$$

with $\ell(v, q) = \ell_1(q) + \ell_2(v)$, and ℓ_i are given by (2.7)-(2.8).

Then, we state the well-posedness of the variational problem (2.1)-(2.2) in the following lemma, and it can be found in section 3 of [3].

Lemma 2.1. Under the assumptions on the problem of (1.1)-(1.2), \mathscr{A} is a continuous and bijective linear operator. Hence, for any $\ell = (\ell_1, \ell_2) \in (\mathbf{Q} \times \mathbf{U})^*$, the mixed variational problem (2.1)-(2.2) has a unique solution $(\mathbf{u}, \mathbf{p}) \in (\mathbf{U} \times \mathbf{Q})$, which satisfy the following continuously

$$\|(u,p)\|_{U\times Q} := (\|u\|_{curl,\Omega}^2 + \|p\|_0^2)^{1/2} \lesssim \|\ell_1\|_{Q^*} + \|\ell_2\|_{U^*}.$$
(2.11)

2.2. Discrete variational problem

We suppose that \mathscr{T}_h is a family of shape regularity, quasi-uniform and conform tetrahedral generation on Ω . Let $h_{\tau} = |\tau|^{1/3}$ denote the mesh size with $|\tau|$ being the volume of $\tau \in \mathscr{T}_h$.

Define the discontinuous finite element function space $\mathbb{V}(\mathcal{T}_h)$ as:

$$\mathbb{V}(\mathcal{T}_h) = \{ \mathbf{v} \in L^2(\Omega) : \mathbf{v}_{\tau} = \mathbf{v}|_{\tau} \in (P_l(\tau))^3, \quad \forall \tau \in \mathcal{T}_h \},\$$

where $P_l(\tau)$ is the set of polynomials defined in the volume τ whose degree does not exceed l, where $l \ge 1$ is an integer.

Let $\mathscr{F}_h, \mathscr{F}_h^0$ and \mathscr{F}_h^∂ denote the set of the all faces of its volumes, and the set of internal faces, and the set of boundary faces, respectively. Thus, $\mathscr{F}_h = \mathscr{F}_h^0 \bigcup \mathscr{F}_h^\partial$. Let $H^1(\Omega; \mathscr{T}_h)$ be the space of piecewise Sobolev functions defined by

$$H^{1}(\Omega; \mathscr{T}_{h}) = \left\{ \boldsymbol{\nu} \in L^{2}(\Omega) : \boldsymbol{\nu}_{\tau} = \boldsymbol{\nu}|_{\tau} \in H^{1}(\tau), \quad \forall \ \tau \in \mathscr{T}_{h} \right\}.$$

and $H^1(\Omega; \mathscr{T}_h) = (H^1(\Omega; \mathscr{T}_h))^3$. Let $L^2(\mathscr{T}_h)$ be the set of L^2 functions defined on \mathscr{T}_h . Moreover, we define the following inner products

$$(\mathbf{v}, \mathbf{w})_{\mathscr{T}'_h} = \sum_{\tau \in \mathscr{T}'_h} \int_{\tau} \mathbf{v} \cdot \mathbf{w} \, \mathrm{d}\mathbf{x}, \quad \forall \mathbf{v}, \mathbf{w} \in L^2(\Omega), \; \forall \mathscr{T}'_h \subset \mathscr{T}_h,$$

 $< \mathbf{v}, \mathbf{w} >_{\mathscr{F}'_h} = \sum_{f \in \mathscr{F}'_h} \int_{f} \mathbf{v} \cdot \mathbf{w} \, \mathrm{d}\mathbf{s}, \quad \forall \mathbf{v}, \mathbf{w} \in L^2(\mathscr{F}_h), \; \forall \mathscr{F}'_h \subset \mathscr{F}_h.$

For $f \in \mathscr{F}_h^0$, we have $\tau_i \in \mathscr{T}_h(i = 1, 2)$, such that $f = \partial \tau_1 \cap \partial \tau_2$. Then we denote the jump and average of ν as:

$$\begin{bmatrix} [\boldsymbol{\nu}] \end{bmatrix} = \boldsymbol{\nu}_1 \times \boldsymbol{n}_1 + \boldsymbol{\nu}_2 \times \boldsymbol{n}_2, \quad \forall \boldsymbol{\nu} \in H^1(\Omega; \mathcal{T}_h),$$

$$\{\{\boldsymbol{\nu}\}\} = \frac{\boldsymbol{\nu}_1 + \boldsymbol{\nu}_2}{2}, \quad \forall \boldsymbol{\nu} \in H^1(\Omega; \mathcal{T}_h),$$

where \mathbf{v}_i denote the values of \mathbf{v} on $\mathbf{v}|_{\tau_i}$ (i = 1, 2) and \mathbf{n}_i denote the out unit normal vectors on f exterior $\mathbf{v}|_{\tau_i}$.

For $f \in \mathscr{F}_h^\partial$, we have $\tau \in \mathscr{T}_h$, such that $f = \partial \tau \cap \partial \Omega$. Then we denote the jump and average of \boldsymbol{v} as:

$$[[\boldsymbol{\nu}]] = \boldsymbol{\nu}_{\tau} \times \boldsymbol{n}_{\partial\Omega}, \quad \{\{\boldsymbol{\nu}\}\} = \boldsymbol{\nu}_{\tau}. \tag{2.12}$$

Next, we give the corresponding discrete scheme of (2.1)-(2.2). Firstly, we define the corresponding discrete space as follow

$$U_h := \{ \boldsymbol{\nu}_h \in \mathbb{V}(\mathcal{T}_h) | \quad [[\boldsymbol{\nu}_h]]|_f = 0, \forall f \in \mathcal{F}_h^\partial \}, \\ \boldsymbol{Q}_h := \mathbb{V}(\mathcal{T}_h).$$

Then, the formulation of the discrete Mixed Interior Penalty Discontinuous Galerkin (MIPDG) method reads: find $(u_h, p_h) \in (U_h, Q_h)$ such that

$$a_{h}(\boldsymbol{p}_{h},\boldsymbol{q}_{h}) - b_{h}(\boldsymbol{u}_{h},\boldsymbol{q}_{h}) = \ell_{1,h}(\boldsymbol{q}_{h}) + d_{1,h}(\boldsymbol{u}_{h},\boldsymbol{q}_{h}), \quad \forall \boldsymbol{q}_{h} \in \boldsymbol{Q}_{h}, \quad (2.13)$$

$$d_{h}(\boldsymbol{v}_{h},\boldsymbol{p}_{h}) + c_{h}(\boldsymbol{u}_{h},\boldsymbol{v}_{h}) = \ell_{2,h}(\boldsymbol{v}_{h}) + d_{2,h}(\boldsymbol{u}_{h},\boldsymbol{v}_{h}), \quad \forall \boldsymbol{v}_{h} \in \boldsymbol{U}_{h}, \quad (2.14)$$

where

$$\begin{aligned} a_{h}(\boldsymbol{p}_{h}, \boldsymbol{q}_{h}) &:= (\boldsymbol{p}_{h}, \boldsymbol{q}_{h})_{\mathcal{F}_{h}}, \\ b_{h}(\boldsymbol{u}_{h}, \boldsymbol{q}_{h}) &:= (\mu \nabla \times \boldsymbol{u}_{h}, \boldsymbol{q}_{h})_{\mathcal{F}_{h}}, \\ c_{h}(\boldsymbol{u}_{h}, \boldsymbol{v}_{h}) &:= (\kappa \boldsymbol{u}_{h}, \boldsymbol{v}_{h})_{\mathcal{F}_{h}}, \\ d_{h}(\boldsymbol{v}_{h}, \boldsymbol{p}_{h}) &:= (\nabla \times \boldsymbol{v}_{h}, \boldsymbol{p}_{h})_{\mathcal{F}_{h}}, \\ \ell_{1,h}(\boldsymbol{q}_{h}) &:= 0, \\ \ell_{2,h}(\boldsymbol{v}_{h}) &:= (\boldsymbol{f}, \boldsymbol{v}_{h})_{\mathcal{F}_{h}}, \\ d_{1,h}(\boldsymbol{u}_{h}, \boldsymbol{q}_{h}) &:= - < \{\{\mu \boldsymbol{q}_{h}\}\}, [[\boldsymbol{u}_{h}]] >_{\mathcal{F}_{h}}, \\ d_{2,h}(\boldsymbol{u}_{h}, \boldsymbol{v}_{h}) &:= < (\{\{\mu \nabla \times \boldsymbol{u}_{h}\}\} - \alpha h_{f}^{-1}[[\boldsymbol{u}_{h}]]), [[\boldsymbol{v}_{h}]] >_{\mathcal{F}_{h}}, \end{aligned}$$

here the constant $\alpha > 0$ denote the penalty parameter, h_f denote the diameter of the circumcircle of f. Thus $h_\tau \approx h_f$.

Remark 2.1. The calculation of $\nabla \times u_h$ in the bilinear terms are piecewise derivations.

The standard symmetric Interior Penalty Discontinuous Galerkin (IPDG) method of the boundary value problem (1.1)-(1.2) is to find $u_h \in U_h$, such that

$$a_{IP}(\boldsymbol{u}_{h},\boldsymbol{v}_{h}) = (\kappa \boldsymbol{u}_{h},\boldsymbol{v}_{h})_{\mathcal{T}_{h}} + (\mu \nabla \times \boldsymbol{u}_{h}, \nabla \times \boldsymbol{v}_{h})_{\mathcal{T}_{h}} - \langle \{ \{\mu \nabla \times \boldsymbol{v}_{h} \} \}, [[\boldsymbol{u}_{h}]] \rangle_{\mathcal{T}_{h}} - \langle \{ \{\mu \nabla \times \boldsymbol{u}_{h} \} \}, [[\boldsymbol{v}_{h}]] \rangle_{\mathcal{T}_{h}} + \alpha h_{f}^{-1} \langle [[\boldsymbol{u}_{h}]], [[\boldsymbol{v}_{h}]] \rangle_{\mathcal{T}_{h}}$$

$$= (\boldsymbol{f}, \boldsymbol{v}_{h})_{\mathcal{T}_{h}}.$$

$$(2.15)$$

The following lemma shows that the discrete variational problems (2.13)-(2.14) and (2.15) are equivalent.

Lemma 2.2. [[3], Theorem 4.1] The formulations (2.13)-(2.14) and (2.15) are formally equivalent in the following sense. If $(\mathbf{u}_h, \mathbf{p}_h) \in (\mathbf{U}_h, \mathbf{Q}_h)$ are the solution of discrete variational problem (2.13)-(2.14), then $\mathbf{u}_h \in \mathbf{U}_h$ solves (2.15). Conversely, if $\mathbf{u}_h \in \mathbf{U}_h$ solves (2.15), then there exists some $\mathbf{p}_h \in \mathbf{Q}_h$ such that $(\mathbf{u}_h, \mathbf{p}_h) \in (\mathbf{U}_h, \mathbf{Q}_h)$ are the solution of (2.13)-(2.14).

Ayuso de Dios, Hiptmair and Pagliantini proved the well-posedness of (2.15) in section 2 of [1]. Therefore, by combining Lemma 2.2, we obtain the well-posedness of discrete variational problems (2.13)-(2.14).

2.3. Adaptive Mixed Interior Penalty Discontinuous Galerkin method(AMIPDG)

Our adaptive cycle can be implemented by the following algorithm: Next, we will discuss each step in AEFEM in detail.

2.3.1. Procedure SOLVE

For $f \in L^2(\Omega)$, and a shape regular mesh \mathcal{T}_k , Let $(\boldsymbol{u}_k, \boldsymbol{p}_k)$ be the exact MIPDG solution of (2.13)-(2.14). Here, we assume that the solutions $(\boldsymbol{u}_k, \boldsymbol{p}_k)$ can be solved accurately.

2.3.2. Procedure ESTIMATE

A posteriori error indicator is an essential ingredient of adaptivity. They are computable quantities depending on the computed solution(s) and data that provide information about the quality of approximation and may consequently be used to make judicious mesh modifications. Here, we design a new posteriori error estimation indicator for equations (2.13)-(2.14), which is similar to that in [20]. For $\tau \in \mathcal{T}_h$, $f \in \mathcal{F}_h$ and $(\mathbf{v}_h, \mathbf{q}_h) \in \mathbf{U}_h \times \mathbf{Q}_h$, the residual a posteriori error estimator for the symmetric AMIPDG method is given by

$$\eta^{2}(\boldsymbol{\nu}_{h},\boldsymbol{q}_{h};\tau): = \|R_{1}(\boldsymbol{\nu}_{h},\boldsymbol{q}_{h})\|_{L^{2}(\tau)}^{2} + h_{\tau}^{2} (\|R_{2}(\boldsymbol{\nu}_{h},\boldsymbol{q}_{h})\|_{L^{2}(\tau)}^{2} + \|R_{3}(\boldsymbol{\nu}_{h})\|_{L^{2}(\tau)}^{2}) + \sum_{f \in \partial \tau} h_{f} (\|J_{1}(\boldsymbol{q}_{h})\|_{L^{2}(f)}^{2} + \|J_{2}(\boldsymbol{\nu}_{h})\|_{L^{2}(f)}^{2}).$$

$$(2.16)$$

Algorithm 2.1 Adaptive Mixed Interior Penalty Discontinuous Galerkin Method (AMIPDG) cycle

Input initial triangulation \mathscr{T}_0 ; data f; tolerance tol; marking parameter $\theta \in (0, 1)$. **Output** a triangulation \mathscr{T}_J ; MIPDG solution (u_J, p_J) .

 $\eta = 1; k = 0;$

while $\eta \ge tol$

SOLVE solve discrete varational problem (2.13)-(2.14) on \mathscr{T}_k to get the solution $(\boldsymbol{u}_k, \boldsymbol{p}_k)$; **ESTIMATE** compute the posterior error estimator $\eta = \eta(\boldsymbol{u}_k, \boldsymbol{p}_k, \mathscr{T}_k)$ by using (2.17); **MARK** seek a minimum cardinality $\mathscr{M}_k \subset \mathscr{T}_k$ such that

$$\eta^2(\boldsymbol{u}_k, \boldsymbol{p}_k, \mathscr{M}_k) \geq \theta \eta^2(\boldsymbol{u}_k, \boldsymbol{p}_k, \mathscr{T}_k);$$

REFINE bisect elements in \mathcal{M}_k and the neighboring elements to form a conforming \mathcal{T}_{k+1} ; k = k + 1;

 $\boldsymbol{u}_J = \boldsymbol{u}_k; \ \boldsymbol{p}_J = \boldsymbol{p}_k; \ \mathcal{T}_J = \mathcal{T}_k;$

They consist of the element residuals and face jump residuals as

$$R_{1}(\boldsymbol{v}_{h}, \boldsymbol{q}_{h})|_{\tau} := \boldsymbol{q}_{h}|_{\tau} - \mu \nabla \times \boldsymbol{v}_{h}|_{\tau},$$

$$R_{2}(\boldsymbol{v}_{h}, \boldsymbol{q}_{h})|_{\tau} := \boldsymbol{f}|_{\tau} - (\nabla \times \boldsymbol{q}_{h} + \kappa \boldsymbol{v}_{h})|_{\tau},$$

$$R_{3}(\boldsymbol{v}_{h})|_{\tau} := \nabla \cdot (\boldsymbol{f}|_{\tau} - \kappa \boldsymbol{v}_{h}|_{\tau}),$$

$$J_{1}(\boldsymbol{q}_{h})|_{f} := [[\boldsymbol{q}_{h}]],$$

$$J_{2}(\boldsymbol{v}_{h})|_{f} := [[(\boldsymbol{f} - \kappa \boldsymbol{v}_{h})]].$$

where h_f denote the diameter of the circumcircle of f, and $h_\tau \approx h_f$.

For any set $\mathscr{T}'_h \subseteq \mathscr{T}_h$, the error indicator is defined as

$$\eta^{2}(\boldsymbol{\nu}_{h},\boldsymbol{q}_{h};\mathscr{T}_{h}') = \sum_{\tau\in\mathscr{T}_{h}'} \eta^{2}(\boldsymbol{\nu}_{h},\boldsymbol{q}_{h};\tau).$$
(2.17)

2.3.3. Procedure MARK

We use the Dörfler mark which was proposed by Dörfler [9]. Set marking parameter $\theta \in (0, 1)$, the module MARK outputs a subset of marked elements $\mathcal{M}_k \subset \mathcal{T}_k$ with minimal cardinality, such that

$$\eta^{2}(\boldsymbol{\nu}_{k},\boldsymbol{q}_{k};\mathcal{M}_{k}) \geq \theta \eta^{2}(\boldsymbol{\nu}_{k},\boldsymbol{q}_{k};\mathcal{T}_{k}).$$
(2.18)

2.3.4. Procedure REFINE

Our implementation of **REFINE** uses the longest edge bisection strategy. A detailed introduction about the longest edge bisection strategy was provided in [6]. To avoid confusion, the relationship between the two tetrahedral meshes \mathscr{T}_h and \mathscr{T}_H that are nested into each other is defined as: \mathscr{T}_h is the new mesh division of \mathscr{T}_H after one cycle of the above cycle process, abbreviated as $\mathscr{T}_H \leq \mathscr{T}_h$.

3. Convergence of AMIPDG algorithm

In this section, we establish the upper bound estimate of the error. Subsequently, we demonstrate that the sum of the energy error and the error estimator between two consecutive adaptive loops is a contraction. Finally, we proof that the AMIPDG is convergence.

3.1. The upper bound estimate of the error

In this subsection, before establishing the reliability of a posteriori error estimator, we need to define the corresponding DG norm, for any $(v, q) \in U \times Q$ and $(v_h, q_h) \in U_h \times Q_h$,

$$\|(\boldsymbol{\nu}, \boldsymbol{q}) - (\boldsymbol{\nu}_{h}, \boldsymbol{q}_{h})\|_{DG}^{2} := \|\boldsymbol{q} - \boldsymbol{q}_{h}\|_{L^{2}(\Omega)}^{2} + \|\kappa(\boldsymbol{\nu} - \boldsymbol{\nu}_{h})\|_{L^{2}(\Omega)}^{2} + \sum_{\tau \in \mathscr{T}_{h}} \|\mu \nabla \times (\boldsymbol{\nu} - \boldsymbol{\nu}_{h})\|_{L^{2}(\tau)}^{2} + \sum_{f \in \mathscr{T}_{h}} \alpha h_{f}^{-1} < [[\boldsymbol{\nu}_{h}]], [[\boldsymbol{\nu}_{h}]] >_{f} .$$
(3.1)

Remark 3.1. For any $v \in U$ and $v_h \in U_h$, we have

$$\|[[\mathbf{v}_h]]\|_{L^2(f)}^2 = \|[[(\mathbf{v} - \mathbf{v}_h)]]\|_{L^2(f)}^2, \quad \forall f \in \mathscr{F}_h$$

In fact, $v \in U$ implies that $[[v]]|_f = 0$ (see Chapter 5 of [16]).

We summarize our main result in this subsection as follows.

Theorem 3.1. Let $(u, p) \in U \times Q$ and $(u_h, p_h) \in U_h \times Q_h$ be the solutions of (2.1)-(2.2) and (2.13)-(2.14), respectively. Let $\eta(u_h, p_h; \mathcal{T}_h)$ be the residual error indicator of (2.17). Then we have the following estimate

$$\|(\boldsymbol{u},\boldsymbol{p}) - (\boldsymbol{u}_h,\boldsymbol{p}_h)\|_{DG}^2 \le C_1 \eta^2 (\boldsymbol{u}_h,\boldsymbol{p}_h;\mathcal{T}_h), \qquad (3.2)$$

where the constant C_1 depending on the shape regularity of mesh.

Let $(u_h, p_h) \in U_h \times Q_h$ be the solution of (2.13)-(2.14), similarly to [4], we introduce the nonconformity of the MSIPDG method results in some consistency error:

$$\zeta := \min_{\tilde{\mathbf{v}}_h \in U} \Big(\sum_{\tau \in \mathscr{T}_h} (\|\mathbf{u}_h - \tilde{\mathbf{v}}_h\|_{L^2(\tau)}^2 + \|\nabla \times (\mathbf{u}_h - \tilde{\mathbf{v}}_h)\|_{L^2(\tau)}^2) \Big)^{1/2}.$$
(3.3)

We denote that $\tilde{u}_h \in U$ is the unique minimizer of (3.3), namely

$$\tilde{\zeta} = \Big(\sum_{\tau \in \mathscr{T}_h} (\|\boldsymbol{u}_h - \tilde{\boldsymbol{u}}_h\|_{L^2(\tau)}^2 + \|\nabla \times (\boldsymbol{u}_h - \tilde{\boldsymbol{u}}_h)\|_{L^2(\tau)}^2)\Big)^{1/2}.$$
(3.4)

Lemma 3.1. Let $(u, p) \in U \times Q$ and $(u_h, p_h) \in U_h \times Q_h$ be the solutions of (2.1)-(2.2) and (2.13)-(2.14), respectively, let \tilde{u}_h be the unique minimizer of (3.3), then

$$\|(\boldsymbol{u}-\tilde{\boldsymbol{u}}_{h},\boldsymbol{p}-\boldsymbol{p}_{h})\|_{\boldsymbol{U}\times\boldsymbol{Q}} = (\|\boldsymbol{u}-\tilde{\boldsymbol{u}}_{h}\|_{curl,\Omega}^{2} + \|\boldsymbol{p}-\boldsymbol{p}_{h}\|_{0}^{2})^{1/2} \lesssim \|\tilde{\ell}_{1}\|_{\boldsymbol{Q}^{*}} + \|\tilde{\ell}_{2}\|_{\boldsymbol{U}^{*}},$$

where the residuals $\tilde{\ell}_1 \in \pmb{Q}^*$ and $\tilde{\ell}_2 \in \pmb{U}^*$ defined by

$$\tilde{\ell}_1(\boldsymbol{q}) = \ell_1(\boldsymbol{q}) - a(\boldsymbol{p}_h, \boldsymbol{q}) + b(\tilde{\boldsymbol{u}}_h, \boldsymbol{q}), \quad \forall \boldsymbol{q} \in \boldsymbol{Q},$$

$$\tilde{\tilde{\boldsymbol{u}}} \quad (3.5)$$

$$\ell_2(\mathbf{v}) = \ell_2(\mathbf{v}) - d(\mathbf{v}, \mathbf{p}_h) - c(\tilde{\mathbf{u}}_h, \mathbf{v}), \quad \forall \mathbf{v} \in \mathbf{U}.$$
(3.6)

Proof. For any $q_1, q_2, q \in Q$ and any $v_1, v_2, v \in U$. we have the following property by (2.9)

$$(\mathscr{A}(\mathbf{v}_{1}+\mathbf{v}_{2},\mathbf{q}_{1}+\mathbf{q}_{2}))(\mathbf{v},\mathbf{q})$$

$$= a(\mathbf{q}_{1}+\mathbf{q}_{2},\mathbf{q})-b(\mathbf{v}_{1}+\mathbf{v}_{2},\mathbf{q})+d(\mathbf{v},\mathbf{q}_{1}+\mathbf{q}_{2})+c(\mathbf{v}_{1}+\mathbf{v}_{2},\mathbf{v})$$

$$= a(\mathbf{q}_{1},\mathbf{q})-b(\mathbf{v}_{1},\mathbf{q})+d(\mathbf{v},\mathbf{q}_{1})+c(\mathbf{v}_{1},\mathbf{v})$$

$$+a(\mathbf{q}_{2},\mathbf{q})-b(\mathbf{v}_{2},\mathbf{q})+d(\mathbf{v},\mathbf{q}_{2})+c(\mathbf{v}_{2},\mathbf{v})$$

$$= (\mathscr{A}(\mathbf{v}_{1},\mathbf{q}_{1}))(\mathbf{v},\mathbf{q})+(\mathscr{A}(\mathbf{v}_{2},\mathbf{q}_{2}))(\mathbf{v},\mathbf{q}).$$

Thus,

$$(\mathscr{A}(\boldsymbol{u}-\tilde{\boldsymbol{u}}_h,\boldsymbol{p}-\boldsymbol{p}_h))(\boldsymbol{v},\boldsymbol{q})$$

= $(\mathscr{A}(\boldsymbol{u},\boldsymbol{p}))(\boldsymbol{v},\boldsymbol{q}) - (\mathscr{A}(\tilde{\boldsymbol{u}}_h,\boldsymbol{p}_h))(\boldsymbol{v},\boldsymbol{q})$
= $(\ell_1(\boldsymbol{q}) + \ell_2(\boldsymbol{v})) - (a(\boldsymbol{p}_h,\boldsymbol{q}) - b(\tilde{\boldsymbol{u}}_h,\boldsymbol{q}) + d(\boldsymbol{v},\boldsymbol{p}_h) + c(\tilde{\boldsymbol{u}}_h,\boldsymbol{v}))$
= $\tilde{\ell}_1(\boldsymbol{q}) + \tilde{\ell}_2(\boldsymbol{v}).$

In fact that $(u - \tilde{u}_h, p - p_h) \in U \times Q$ and combining the Lemma 2.1 can concludes the proof.

Next, we will provide upper bounds for $\|\tilde{\ell}_1\|_{Q^*}$ and $\|\tilde{\ell}_2\|_{U^*}$ in Lemmas 3.2 and 3.4, respectively.

Lemma 3.2. Let $(\boldsymbol{u}_h, \boldsymbol{p}_h) \in \boldsymbol{U}_h \times \boldsymbol{Q}_h$ be the solutions of (2.13)-(2.14), and $\tilde{\boldsymbol{u}}_h$ be the unique minimizer of (3.3). Then we get the estimate of the linear functional $\tilde{\ell}_1$ defined in (3.5) as following

$$\|\tilde{\ell}_{1}\|_{\boldsymbol{Q}^{*}} \lesssim \left(\sum_{\tau \in \mathscr{T}_{h}} \|R_{1}(\boldsymbol{u}_{h}, \boldsymbol{p}_{h})\|_{L^{2}(\tau)}^{2}\right)^{1/2} + \left(\sum_{\tau \in \mathscr{T}_{h}} \|\nabla \times (\tilde{\boldsymbol{u}}_{h} - \boldsymbol{u}_{h})\|_{L^{2}(\tau)}^{2}\right)^{1/2}.$$
(3.7)

Proof. For any $q \in Q$, by the definition of $\tilde{\ell}_1$, we have

$$\tilde{\ell}_1(\boldsymbol{q}) = \sum_{\tau \in \mathscr{T}_h} \int_{\tau} \left((\mu \nabla \times \boldsymbol{u}_h - \boldsymbol{p}_h) + \mu \nabla \times (\tilde{\boldsymbol{u}}_h - \boldsymbol{u}_h) \right) \cdot \boldsymbol{q} \, \mathrm{d}\boldsymbol{x}.$$

K Liu et al.

Then applying the Hölder inequality and the Cauchy-Schwarz inequality,

$$\begin{split} |\tilde{\ell}_{1}(\boldsymbol{q})| &\leq \sum_{\tau \in \mathscr{T}_{h}} \|\mu \nabla \times \boldsymbol{u}_{h} - \boldsymbol{p}_{h}\|_{L^{2}(\tau)} \|\boldsymbol{q}\|_{L^{2}(\Omega)} + \sum_{\tau \in \mathscr{T}_{h}} \|\mu \nabla \times (\tilde{\boldsymbol{u}}_{h} - \boldsymbol{u}_{h})\|_{L^{2}(\tau)} \|\boldsymbol{q}\|_{L^{2}(\Omega)} \\ &\lesssim \left(\Big(\sum_{\tau \in \mathscr{T}_{h}} \|R_{1}(\boldsymbol{u}_{h}, \boldsymbol{p}_{h})\|_{L^{2}(\tau)}^{2} \Big)^{1/2} + \Big(\sum_{\tau \in \mathscr{T}_{h}} \|\nabla \times (\tilde{\boldsymbol{u}}_{h} - \boldsymbol{u}_{h})\|_{L^{2}(\tau)}^{2} \Big) \|\boldsymbol{q}\|_{L^{2}(\Omega)}, \end{split}$$

conclude the proof.

Before estimating the term $\|\tilde{\ell}_2\|_{U^*}$, we need to introduce the following interpolation operator with the corresponding approximations.

Lemma 3.3. [[19], Theorem 1] Let $Nd_0^1(\Omega; \mathscr{T}_h)$ be the lowest order edge elements of Nédélec first family. Then there exists an operator $\Pi_h : H_0(curl; \Omega) \to Nd_0^1(\Omega; \mathscr{T}_h)$ with the following properties: For every $\mathbf{v} \in H_0(curl; \Omega)$, there exist $\varphi \in H_0^1(\Omega)$ and $\mathbf{z} \in H_0^1(\Omega)$, such that

$$\boldsymbol{v} - \boldsymbol{\Pi}_h \boldsymbol{v} = \nabla \boldsymbol{\varphi} + \boldsymbol{z}$$

And for any $\tau \in \mathcal{T}_h$ and $f \in \mathcal{F}_h$, we have

$$\begin{aligned} & h_{\tau}^{-1} \|\varphi\|_{L^{2}(\tau)} + \|\nabla\varphi\|_{L^{2}(\tau)} \lesssim h_{\tau} \|\mathbf{v}\|_{L^{2}(\Omega_{\tau})}, \\ & h_{\tau}^{-1} \|\mathbf{z}\|_{L^{2}(\tau)} + \|\nabla\mathbf{z}\|_{L^{2}(\tau)} \lesssim h_{\tau} \|\nabla \times \mathbf{v}\|_{L^{2}(\Omega_{\tau})}, \end{aligned}$$

where $\Omega_{\tau} = \bigcup_{f \in \tau} \Omega_f$, $\Omega_f = \{\tau' \in \mathcal{T}_h, f \in \tau'\}$, and the constants depending on the shape regularity of the mesh.

Lemma 3.4. Let $(\boldsymbol{u}_h, \boldsymbol{p}_h) \in \boldsymbol{U}_h \times \boldsymbol{Q}_h$ be the solution of (2.13)-(2.14), and $\tilde{\boldsymbol{u}}_h$ be the unique solution of (3.3). Then the linear functional $\tilde{\ell}_2$ defined in (3.6) satisfies the following estimate

$$\|\tilde{\ell}_{2}\|_{U^{*}} \lesssim \left(\sum_{\tau \in \mathscr{T}} h_{\tau}^{2} (\|R_{2}(\boldsymbol{u}_{h}, \boldsymbol{p}_{h})\|_{L^{2}(\tau)}^{2} + \|R_{2}(\boldsymbol{u}_{h})\|_{L^{2}(\tau)}^{2}) + \sum_{f \in \mathscr{F}} h_{f} (\|J_{1}(\boldsymbol{p}_{h})\|_{L^{2}(f)}^{2} + \|J_{2}(\boldsymbol{u}_{h})\|_{L^{2}(f)}^{2}) + \sum_{\tau \in \mathscr{T}} \|\boldsymbol{u}_{h} - \tilde{\boldsymbol{u}}_{h}\|_{L^{2}(\tau)}^{2}\right)^{1/2}.$$
 (3.8)

Proof. For any $v \in U$ and Π_h given by Lemma 3.3, we have

$$\boldsymbol{\nu} - \boldsymbol{\Pi}_h \boldsymbol{\nu} = \nabla \boldsymbol{\varphi} + \boldsymbol{z}, \tag{3.9}$$

where $\varphi \in H_0^1(\Omega)$ and $z \in H_0^1(\Omega)$. According to linearity of the operator $\tilde{\ell}_2$ and (3.9), we have

$$\tilde{\ell}_2(\boldsymbol{\nu}) = \tilde{\ell}_2(\Pi_h \boldsymbol{\nu}) + \tilde{\ell}_2(\boldsymbol{\nu} - \Pi_h \boldsymbol{\nu}) = \tilde{\ell}_2(\Pi_h \boldsymbol{\nu}) + \tilde{\ell}_2(\nabla \varphi) + \tilde{\ell}_2(\boldsymbol{z}).$$
(3.10)

We will next estimate the three terms on the right hand side of (3.10).

For the first term $\tilde{\ell}_2(\Pi_h v)$ of (3.10), using the definition of $\tilde{\ell}_2$, we have

$$\tilde{\ell}_2(\Pi_h \boldsymbol{\nu}) = \ell_2(\Pi_h \boldsymbol{\nu}) - d(\Pi_h \boldsymbol{\nu}, \boldsymbol{p}_h) - c(\tilde{\boldsymbol{u}}_h, \Pi_h \boldsymbol{\nu}) = \ell_2(\Pi_h \boldsymbol{\nu}) - d(\Pi_h \boldsymbol{\nu}, \boldsymbol{p}_h) - c(\boldsymbol{u}_h, \Pi_h \boldsymbol{\nu}) + c(\boldsymbol{u}_h - \tilde{\boldsymbol{u}}_h, \Pi_h \boldsymbol{\nu}).$$

Noting that $\Pi_h \mathbf{v} \in Nd_0^1(\Omega; \mathscr{T}_h) \subseteq U_h$ has zero jumps, and combining (2.14), we have

$$\ell_2(\Pi_h \boldsymbol{\nu}) - d(\Pi_h \boldsymbol{\nu}, \boldsymbol{p}_h) - c(\boldsymbol{u}_h, \Pi_h \boldsymbol{\nu}) = \ell_{2,h}(\Pi_h \boldsymbol{\nu}) - d_h(\Pi_h \boldsymbol{\nu}, \boldsymbol{p}_h) - c_h(\boldsymbol{u}_h, \Pi_h \boldsymbol{\nu}) = 0.$$

Thus, we have

$$\begin{split} \tilde{\ell}_2(\Pi_h \boldsymbol{\nu}) &= c(\boldsymbol{\nu}_h - \tilde{\boldsymbol{u}}_h, \Pi_h \boldsymbol{\nu}) \\ &= c(\boldsymbol{\nu}_h - \tilde{\boldsymbol{u}}_h, \boldsymbol{\nu}) + c(\boldsymbol{\nu}_h - \tilde{\boldsymbol{u}}_h, \Pi_h \boldsymbol{\nu} - \boldsymbol{\nu}) \\ &\leq \|\kappa\|_{0,\infty} \|\boldsymbol{\nu}_h - \tilde{\boldsymbol{u}}_h\|_{0,\mathcal{T}_h} (\|\boldsymbol{\nu}\|_{0,\mathcal{T}_h} + \|\Pi_h \boldsymbol{\nu} - \boldsymbol{\nu}\|_{0,\mathcal{T}_h}). \end{split}$$

Then using (3.9), triangle inequality and Lemma 3.3, we get

$$\widetilde{\ell}_{2}(\Pi_{h}\boldsymbol{\nu}) \leq \|\kappa\|_{0,\infty} \|\boldsymbol{\nu}_{h} - \widetilde{\boldsymbol{u}}_{h}\|_{0,\mathscr{T}_{h}} (\|\boldsymbol{\nu}\|_{0,\mathscr{T}_{h}} + \|\nabla\varphi + \boldsymbol{z}\|_{0,\mathscr{T}_{h}}) \\
\leq \|\kappa\|_{0,\infty} \|\boldsymbol{\nu}_{h} - \widetilde{\boldsymbol{u}}_{h}\|_{0,\mathscr{T}_{h}} (\|\boldsymbol{\nu}\|_{0,\mathscr{T}_{h}} + \|\nabla\varphi\|_{0,\mathscr{T}_{h}} + \|\boldsymbol{z}\|_{0,\mathscr{T}_{h}}) \\
\leq \|\kappa\|_{0,\infty} \|\boldsymbol{\nu}_{h} - \widetilde{\boldsymbol{u}}_{h}\|_{0,\mathscr{T}_{h}} \|\boldsymbol{\nu}\|_{curl,\mathscr{T}_{h}}.$$
(3.11)

For the second term $\tilde{\ell}_2(\nabla \varphi)$ of (3.10), using the definition of $\tilde{\ell}_2$, (2.8), (2.4), (2.6) and the fact $\nabla \times \nabla \varphi = 0$, which implies

$$\tilde{\ell}_{2}(\nabla\varphi) = \ell_{2}(\nabla\varphi) - d(\nabla\varphi, \boldsymbol{p}_{h}) - c(\tilde{\boldsymbol{u}}_{h}, \nabla\varphi)
= (f, \nabla\varphi) - (\nabla \times \nabla\varphi, \boldsymbol{p}_{h}) - (\kappa \tilde{\boldsymbol{u}}_{h}, \nabla\varphi)
= (f, \nabla\varphi) - (\kappa \tilde{\boldsymbol{u}}_{h}, \nabla\varphi).$$
(3.12)

By (3.12) and Green's formula, we have

$$\begin{aligned} \tilde{\ell}_2(\nabla\varphi) &= (f, \nabla\varphi) - (\kappa u_h, \nabla\varphi) + (\kappa (u_h - \tilde{u}_h), \nabla\varphi) \\ &\leq \sum_{\tau \in \mathscr{T}_h} (R_3(u_h), \varphi)_{0,\tau} + \sum_{f \in \mathscr{F}_h} < J_2(u_h), \varphi >_{0,f} + (\kappa (u_h - \tilde{u}_h), \nabla\varphi). \end{aligned}$$

Applying the Cauchy-Schwarz inequality, Lemma 3.3 and trace inequality, we have

$$\tilde{\ell}_{2}(\nabla\varphi) \leq \left(\sum_{\tau \in \mathscr{T}_{h}} h_{\tau}^{2} \|R_{3}(\boldsymbol{u}_{h})\|_{0,\tau}^{2} + \sum_{f \in \mathscr{T}_{h}} h_{f} \|J_{2}(\boldsymbol{u}_{h})\|_{0,f}^{2} + \sum_{\tau \in \mathscr{T}_{h}} \|\kappa\|_{0,\infty} \|\boldsymbol{u}_{h} - \tilde{\boldsymbol{u}}_{h}\|_{0,\tau}^{2}\right)^{1/2} \|\boldsymbol{v}\|_{curl,\mathscr{T}_{h}}.$$
(3.13)

Similarly, for the third term $\tilde{\ell}_2(z)$ of (3.10), we have

$$\tilde{\ell}_{2}(\boldsymbol{z}) = (\boldsymbol{f}, \boldsymbol{z}) - (\nabla \times \boldsymbol{z}, \boldsymbol{p}_{h}) - (\kappa \tilde{\boldsymbol{u}}_{h}, \boldsymbol{z})
= (\boldsymbol{f}, \boldsymbol{z}) - (\nabla \times \boldsymbol{z}, \boldsymbol{p}_{h}) - (\kappa \boldsymbol{u}_{h}, \boldsymbol{z}) + (\kappa (\boldsymbol{u}_{h} - \tilde{\boldsymbol{u}}_{h}), \boldsymbol{z})
\leq \left(\sum_{\tau \in \mathscr{T}_{h}} h_{\tau}^{2} \| R_{2}(\boldsymbol{u}_{h}, \boldsymbol{p}_{h}) \|_{0,\tau}^{2} + \sum_{f \in \mathscr{T}_{h}} h_{f} \| J_{1}(\boldsymbol{p}_{h}) \|_{0,f}^{2}
+ \sum_{\tau \in \mathscr{T}_{h}} \| \kappa \|_{0,\infty} \| \boldsymbol{u}_{h} - \tilde{\boldsymbol{u}}_{h} \|_{0,\tau}^{2} \right)^{1/2} \| \boldsymbol{v} \|_{curl,\mathscr{T}_{h}}.$$
(3.14)

Substituting (3.11), (3.13) and (3.14) into (3.10), the proof is completed.

Notice that both (3.7) and (3.8) are related to the terms $\sum_{\tau \in \mathscr{T}_h} \|\nabla \times (\tilde{\boldsymbol{u}}_h - \boldsymbol{u}_h)\|_{L^2(\tau)}^2$ and $\sum_{\tau \in \mathscr{T}} \|\boldsymbol{u}_h - \tilde{\boldsymbol{u}}_h\|_{L^2(\tau)}^2$, which are a part of $\tilde{\zeta}$. Therefore, we prove upper bounds for $\tilde{\zeta}$ in the following Lemma.

Lemma 3.5. Let $(\boldsymbol{u}_h, \boldsymbol{p}_h) \in \boldsymbol{U}_h \times \boldsymbol{Q}_h$ be the solutions of (2.13)-(2.14) and $\tilde{\zeta}$ be consistency error of (3.4), we have

$$\tilde{\zeta}^2 \lesssim \eta^2(\boldsymbol{u}_h, \boldsymbol{p}_h; \mathcal{T}_h). \tag{3.15}$$

Proof. For any $v_h \in U_h$, there exit an interpolation operator $\mathscr{I}_h : H^1(\Omega; \mathscr{T}_h) \to U_h^c$, such that (see Proposition 4.5 of [11])

$$\|\boldsymbol{\nu}_{h} - \mathscr{I}_{h} \boldsymbol{\nu}_{h}\|_{L^{2}(\Omega)}^{2} \lesssim \sum_{f \in \mathscr{F}_{h}} h_{f} \|[[\boldsymbol{\nu}_{h}]]\|_{L^{2}(f)}^{2}, \qquad (3.16)$$

$$\sum_{\tau \in \mathscr{T}_h} \|\nabla \times (\boldsymbol{\nu}_h - \mathscr{I}_h \boldsymbol{\nu}_h)\|_{L^2(\tau)}^2 \lesssim \sum_{f \in \mathscr{F}_h} h_f^{-1} \|[[\boldsymbol{\nu}_h]]\|_{L^2(f)}^2.$$
(3.17)

Then, combining (3.3), (3.4), (3.16), (3.17), and the fact $h_f < 1$, we get

$$\tilde{\zeta}^{2} = \sum_{\tau \in \mathscr{T}_{h}} (\|\boldsymbol{u}_{h} - \tilde{\boldsymbol{u}}_{h}\|_{L^{2}(\tau)}^{2} + \|\nabla \times (\boldsymbol{u}_{h} - \tilde{\boldsymbol{u}}_{h})\|_{L^{2}(\tau)}^{2})
\leq \sum_{\tau \in \mathscr{T}_{h}} (\|\boldsymbol{u}_{h} - \mathscr{I}_{h}\boldsymbol{u}_{h}\|_{L^{2}(\tau)}^{2} + \|\nabla \times (\boldsymbol{u}_{h} - \mathscr{I}_{h}\boldsymbol{u}_{h})\|_{L^{2}(\tau)}^{2})
\lesssim \sum_{f \in \mathscr{T}_{h}} h_{f} \|[[\boldsymbol{u}_{h}]]\|_{L^{2}(f)}^{2} + \sum_{f \in \mathscr{T}_{h}} h_{f}^{-1} \|[[\boldsymbol{u}_{h}]]\|_{L^{2}(f)}^{2}
\lesssim \sum_{f \in \mathscr{T}_{h}} h_{f}^{-1} \|[[\boldsymbol{u}_{h}]]\|_{L^{2}(f)}^{2}.$$
(3.18)

Noting that $(u_h, p_h) \in U_h \times Q_h$ is the solution of discrete variational problem (2.13)-(2.14). Then by using Lemma 2.2, we know that u_h is the solution of discrete variational problem (2.15). Hence, we have (see Lemma 5 of [20])

$$\alpha \|h_f^{-1/2}[[\boldsymbol{u}_h]]\|_{L^2(\mathscr{F}_h)} \lesssim \eta(\boldsymbol{u}_h, \boldsymbol{p}_h; \mathscr{T}_h).$$
(3.19)

12

At last, combining (3.18) and (3.19), we have

$$\tilde{\zeta}^2 \lesssim \eta^2(\boldsymbol{u}_h, \boldsymbol{p}_h; \mathcal{T}).$$

Combining Lemmas 3.1, 3.2, 3.4 and 3.5, we will prove Theorem 3.1.

Proof. [**Proof of** Theorem 3.1:] By using (3.1), the triangle inequality, (3.4), Lemmas 3.1, 3.2, 3.4, 3.5 and (3.19), we get

$$\begin{split} \|(\boldsymbol{u},\boldsymbol{p}) - (\boldsymbol{u}_{h},\boldsymbol{p}_{h})\|_{DG}^{2} \\ \lesssim & \|\boldsymbol{p} - \boldsymbol{p}_{h}\|_{L^{2}(\Omega)}^{2} + \|\kappa(\boldsymbol{u} - \boldsymbol{u}_{h})\|_{L^{2}(\Omega)}^{2} \\ & + \sum_{\tau \in \mathscr{F}_{h}} \|\nabla \times \mu(\boldsymbol{u} - \boldsymbol{u}_{h})\|_{L^{2}(\tau)}^{2} + \sum_{f \in \mathscr{F}_{h}} \alpha h_{f}^{-1} < [[\boldsymbol{u}_{h}]], [[\boldsymbol{u}_{h}]] >_{f} \\ \lesssim & \|\boldsymbol{p} - \boldsymbol{p}_{h}\|_{L^{2}(\Omega)}^{2} + \|\boldsymbol{u} - \tilde{\boldsymbol{u}}_{h}\|_{curl,\Omega}^{2} + \tilde{\zeta}^{2} + \sum_{f \in \mathscr{F}_{h}} \alpha h_{f}^{-1} < [[\boldsymbol{u}_{h}]], [[\boldsymbol{u}_{h}]] >_{f} \\ = & \|(\boldsymbol{u} - \tilde{\boldsymbol{u}}_{h}, \boldsymbol{p} - \boldsymbol{p}_{h})\|_{U \times Q} + \tilde{\zeta}^{2} + \sum_{f \in \mathscr{F}_{h}} \alpha h_{f}^{-1} < [[\boldsymbol{u}_{h}]], [[\boldsymbol{u}_{h}]] >_{f} \\ \lesssim & \|\tilde{\ell}_{1}\|_{Q^{*}}^{2} + \|\tilde{\ell}_{2}\|_{U^{*}}^{2} + \tilde{\zeta}^{2} + \sum_{f \in \mathscr{F}_{h}} \alpha h_{f}^{-1} < [[\boldsymbol{u}_{h}]], [[\boldsymbol{u}_{h}]] >_{f} \\ \leq & C_{1}\eta^{2}(\boldsymbol{u}_{h}, \boldsymbol{p}_{h}; \mathscr{F}_{h}). \end{split}$$

3.2. The error reduces on two successive meshes

For convenience, for any $v \in U$ and $v_h \in U_h$, we denote

$$\||\boldsymbol{\nu} - \boldsymbol{\nu}_{h}\|_{h}^{2} = \|\kappa(\boldsymbol{\nu} - \boldsymbol{\nu}_{h})\|_{L^{2}(\Omega)}^{2} + \sum_{\tau \in \mathscr{T}_{h}} \|\nabla \times \mu(\boldsymbol{\nu} - \boldsymbol{\nu}_{h})\|_{L^{2}(\tau)}^{2} + \sum_{f \in \mathscr{F}_{h}} \alpha h_{f}^{-1} < [[\boldsymbol{\nu}_{h}]], [[\boldsymbol{\nu}_{h}]] >_{f}.$$
(3.20)

Let U_h^c be the H(curl) conforming subspace of U_h given by

$$\boldsymbol{U}_{h}^{c} := \boldsymbol{U}_{h} \cap \mathbf{H}_{0}(curl; \Omega).$$

Then, there is a subspace U_h^{\perp} which can orthogonally decompose U_h under L^2 inner product such that $U_h := U_h^c \oplus U_h^{\perp}$. Especially, if $(u_h, p_h) \in U_h \times Q_h$ is the solution of (2.13)-(2.14), then we have

$$\||\boldsymbol{u}_{h}^{\perp}\|_{h}^{2} \lesssim \alpha \sum_{f \in \partial \tau} \|h_{f}^{-1/2}[[\boldsymbol{u}_{h}]]\|_{L^{2}(f)}^{2}.$$
(3.21)

In fact, from the Lemma 2.2, notice that u_h satisfies the IPDG scheme of (2.15), and according to Lemma 2 in [20], we can obtain (3.21).

In order to easily estimate the jump term of face \mathscr{F}_h , we need to introduce the lifting operators and the corresponding stability estimates, more details are referenced to Proposition 12 in [18].

Let $\mathscr{L}_h : H^1(\Omega; \mathscr{T}_h) \to U_h$ be the lifting operators, which satisfies the following equality

$$\int_{\Omega} \mathcal{L}_h(\boldsymbol{\nu}) \cdot \boldsymbol{w} \, \mathrm{d}\boldsymbol{x} = < [[\boldsymbol{\nu}]], \{\{\boldsymbol{w}\}\} >_{\mathscr{F}_h}, \quad \forall \boldsymbol{w} \in \boldsymbol{U}_h, \tag{3.22}$$

and

$$\|\mathscr{L}_{h}(\mathbf{v})\|_{L^{2}(\Omega)} \leq C_{\mathscr{L}} \|h^{-1/2}[[\mathbf{v}]]\|_{L^{2}(\mathscr{F}_{h})},$$
(3.23)

where the constant $C_{\mathcal{L}}$ depending on the shape regularity of mesh \mathcal{T}_h and the degree of polynomial *l*.

Lemma 3.6. Let $(u, p) \in U \times Q$ and $(u_h, p_h) \in U_h \times Q_h$ be the solutions of (2.1)-(2.2) and (2.13)-(2.14), respectively, we have

$$\|\boldsymbol{p} - \boldsymbol{p}_h\|_{L^2(\Omega)} \lesssim \|\nabla \times (\boldsymbol{u} - \boldsymbol{u}_h)\|_{L^2(\Omega)} + \eta(\boldsymbol{u}_h, \boldsymbol{p}_h; \mathcal{T}_h), \qquad (3.24)$$

$$\|\boldsymbol{p}_{h} - \boldsymbol{p}_{H}\|_{L^{2}(\Omega)} \lesssim \|\nabla \times (\boldsymbol{u}_{h} - \boldsymbol{u}_{H})\|_{L^{2}(\Omega)} + \left(\eta(\boldsymbol{u}_{h}, \boldsymbol{p}_{h}; \mathcal{T}_{h}) + \eta(\boldsymbol{u}_{H}, \boldsymbol{p}_{H}; \mathcal{T}_{H})\right).$$
(3.25)

Proof. Noting that $Q_h \subseteq Q$, and using (2.1), the definition of $R_1(u_h, p_h)$ and (2.16), we have

$$\begin{split} \|\boldsymbol{p} - \boldsymbol{p}_{h}\|_{L^{2}(\mathcal{T}_{h})} &\leq \sup_{\forall \boldsymbol{q} \in \boldsymbol{Q}} \frac{(\boldsymbol{p} - \boldsymbol{p}_{h}, \boldsymbol{q})_{\mathcal{T}_{h}}}{\|\boldsymbol{q}\|_{L^{2}(\mathcal{T}_{h})}} \\ &= \sup_{\forall \boldsymbol{q} \in \boldsymbol{Q}} \frac{(\mu \nabla \times \boldsymbol{u}, \boldsymbol{q})_{\mathcal{T}_{h}} - (R_{1}(\boldsymbol{u}_{h}, \boldsymbol{p}_{h}) + \mu \nabla \times \boldsymbol{u}_{h}, \boldsymbol{q})_{\mathcal{T}_{h}}}{\|\boldsymbol{q}\|_{L^{2}(\mathcal{T}_{h})}} \\ &\leq \sup_{\forall \boldsymbol{q} \in \boldsymbol{Q}} \frac{(\mu \nabla \times (\boldsymbol{u} - \boldsymbol{u}_{h}), \boldsymbol{q})_{\mathcal{T}_{h}} - (R_{1}(\boldsymbol{u}_{h}, \boldsymbol{p}_{h}), \boldsymbol{q})_{\mathcal{T}_{h}}}{\|\boldsymbol{q}\|_{L^{2}(\mathcal{T}_{h})}} \\ &\lesssim \|\nabla \times (\boldsymbol{u} - \boldsymbol{u}_{h})\|_{L^{2}(\mathcal{T}_{h})} + \eta(\boldsymbol{u}_{h}, \boldsymbol{p}_{h}; \mathcal{T}_{h}). \end{split}$$

Similarly, using the definition of $R_1(u_h, p_h)$, (2.13), (3.21)-(3.23), and the fact $[[u_h]] =$

14

$$\begin{split} & [[\boldsymbol{u}_{h}^{c} + \boldsymbol{u}_{h}^{\perp}]] = [[\boldsymbol{u}_{h}^{\perp}]], \text{ we have} \\ & \|\boldsymbol{p}_{h} - \boldsymbol{p}_{H}\|_{L^{2}(\mathcal{T}_{h})} \leq \sup_{\forall \boldsymbol{q}_{h} \in \boldsymbol{Q}_{h}} \frac{(\boldsymbol{p}_{h} - \boldsymbol{p}_{H}, \boldsymbol{q}_{h})_{\mathcal{T}_{h}}}{\|\boldsymbol{q}_{h}\|_{L^{2}(\mathcal{T}_{h})}} \\ & \leq \sup_{\forall \boldsymbol{q}_{h} \in \boldsymbol{Q}_{h}} \frac{(\boldsymbol{p}_{h}, \boldsymbol{q}_{h})_{\mathcal{T}_{h}} - (R_{1}(\boldsymbol{u}_{H}, \boldsymbol{p}_{H}) + \mu \nabla \times \boldsymbol{u}_{H}, \boldsymbol{q}_{h})_{\mathcal{T}_{h}}}{\|\boldsymbol{q}_{h}\|_{L^{2}(\mathcal{T}_{h})}} \\ & \leq \sup_{\forall \boldsymbol{q}_{h} \in \boldsymbol{Q}_{h}} \frac{(\mu \nabla \times \boldsymbol{u}_{h}, \boldsymbol{q}_{h})_{\mathcal{T}_{h}} + <\{\{\boldsymbol{q}_{h}\}\}, [[\mu\boldsymbol{u}_{h}]] >_{\mathcal{T}_{h}} - (R_{1}(\boldsymbol{u}_{H}, \boldsymbol{p}_{H}) + \mu \nabla \times \boldsymbol{u}_{H}, \boldsymbol{q}_{h})_{\mathcal{T}_{h}}}{\|\boldsymbol{q}_{h}\|_{L^{2}(\mathcal{T}_{h})}} \\ & = \sup_{\forall \boldsymbol{q}_{h} \in \boldsymbol{Q}_{h}} \frac{(\mu \nabla \times (\boldsymbol{u}_{h} - \boldsymbol{u}_{H}), \boldsymbol{q}_{h})_{\mathcal{T}_{h}} + <\{\{\boldsymbol{q}_{h}\}\}, [[\mu\boldsymbol{u}_{h}]] >_{\mathcal{T}_{h}} - (R_{1}(\boldsymbol{u}_{H}, \boldsymbol{p}_{H}), \boldsymbol{q}_{h})_{\mathcal{T}_{h}}}{\|\boldsymbol{q}_{h}\|_{L^{2}(\mathcal{T}_{h})}} \\ & \leq \|\nabla \times (\boldsymbol{u}_{h} - \boldsymbol{u}_{H})\|_{L^{2}(\mathcal{T}_{h})} + \|\boldsymbol{h}_{\tau}^{-1/2}[[\boldsymbol{u}_{h}]]\|_{L^{2}(\mathcal{T}_{h})} + \eta(\boldsymbol{u}_{H}, \boldsymbol{p}_{H}; \mathcal{T}_{H})} \\ & \lesssim \|\nabla \times (\boldsymbol{u}_{h} - \boldsymbol{u}_{H})\|_{L^{2}(\mathcal{T}_{h})} + C_{\mathcal{L}}\|\boldsymbol{h}_{\tau}^{-1/2}[[\boldsymbol{u}_{h}^{\perp}]]\|_{L^{2}(\mathcal{T}_{h})} + \eta(\boldsymbol{u}_{H}, \boldsymbol{p}_{H}; \mathcal{T}_{H})} \\ & \lesssim \|\nabla \times (\boldsymbol{u}_{h} - \boldsymbol{u}_{H})\|_{L^{2}(\tau)} + \left(\eta(\boldsymbol{u}_{h}, \boldsymbol{p}_{h}; \mathcal{T}_{h}) + \eta(\boldsymbol{u}_{H}, \boldsymbol{p}_{H}; \mathcal{T}_{H})\right). \end{split}$$

Remark 3.2. Noting that $||(\boldsymbol{u}, \boldsymbol{p}) - (\boldsymbol{u}_h, \boldsymbol{p}_h)||_{DG}^2 + \eta^2(\boldsymbol{u}_h, \boldsymbol{p}_h; \mathcal{T}_h)$ and $|||\boldsymbol{u} - \boldsymbol{u}_h||_h^2 + \eta^2(\boldsymbol{u}_h, \boldsymbol{p}_h; \mathcal{T}_h)$ are equivalent. In fact, by (3.24), we first know that

$$\begin{aligned} \|(\boldsymbol{u},\boldsymbol{p}) - (\boldsymbol{u}_{h},\boldsymbol{p}_{h})\|_{DG}^{2} + \eta^{2}(\boldsymbol{u}_{h},\boldsymbol{p}_{h};\mathcal{T}_{h}) \\ &= \||\boldsymbol{u} - \boldsymbol{u}_{h}\||_{h}^{2} + \|\boldsymbol{p} - \boldsymbol{p}_{h}\|_{L^{2}(\mathcal{T}_{h})}^{2} + \eta^{2}(\boldsymbol{u}_{h},\boldsymbol{p}_{h};\mathcal{T}_{h}) \\ &\lesssim \||\boldsymbol{u} - \boldsymbol{u}_{h}\||_{h}^{2} + \eta^{2}(\boldsymbol{u}_{h},\boldsymbol{p}_{h};\mathcal{T}_{h}). \end{aligned}$$

Secondly, it is shown by the definition of $\|\cdot\|_{DG}$

$$\||\boldsymbol{u}-\boldsymbol{u}_h|\|_h^2+\eta^2(\boldsymbol{u}_h,\boldsymbol{p}_h;\mathcal{T}_h)\leq \|(\boldsymbol{u},\boldsymbol{p})-(\boldsymbol{u}_h,\boldsymbol{p}_h)\|_{DG}^2+\eta^2(\boldsymbol{u}_h,\boldsymbol{p}_h;\mathcal{T}_h).$$

Thus, we next only need to consider the convergence of $|||u - u_h|||_h^2 + \eta^2(u_h, p_h; \mathcal{T}_h)$.

We first show that the error plus some quantity reduces with a fixed factor on two successive meshes.

Lemma 3.7. Given $f \in L^2(\Omega)$ and two tetrahedral mesh \mathscr{T}_h and \mathscr{T}_H , where $\mathscr{T}_H \leq \mathscr{T}_h$. Let $(\boldsymbol{u}, \boldsymbol{p}) \in \boldsymbol{U} \times \boldsymbol{Q}$ be the solution of (2.1)-(2.2), and $(\boldsymbol{u}_h, \boldsymbol{p}_h) \in \boldsymbol{U}_h \times \boldsymbol{Q}_h$, $(\boldsymbol{u}_H, \boldsymbol{p}_H) \in \boldsymbol{U}_H \times \boldsymbol{Q}_H$ be the solutions of (2.13)-(2.14), respectively. Then there exit two constants $\delta_1, \delta_2 \in (0, 1)$, such that

$$\||\boldsymbol{u} - \boldsymbol{u}_{h}|\|_{h}^{2} \leq (1 + \delta_{1}) \||\boldsymbol{u} - \boldsymbol{u}_{H}|\|_{H}^{2} - \frac{1 - \delta_{2}}{2} \||\boldsymbol{u}_{h} - \boldsymbol{u}_{H}|\|_{h}^{2} + \frac{C_{3}}{\delta_{1}\delta_{2}\alpha} \Big(\eta^{2}(\boldsymbol{u}_{h}, \boldsymbol{p}_{h}; \mathcal{T}_{h}) + \eta^{2}(\boldsymbol{u}_{H}, \boldsymbol{p}_{H}; \mathcal{T}_{H}) \Big).$$
(3.26)

where C_3 depending on the $C_{\mathscr{L}}$.

K Liu et al.

Proof. Choosing that $q = \nabla \times v$, and subtracting (2.1) from (2.2), we obtain

$$(\kappa u, v) + (\mu \nabla \times u, \nabla \times v) = (f, v).$$
(3.27)

Subtracting (2.15) from (3.27) with $\mathbf{v} = \mathbf{v}_h = \mathbf{u}_h^c - \mathbf{u}_H^c$, and using $[[\mathbf{u}_h^c - \mathbf{u}_H^c]] = 0$, we have

$$(\kappa(\boldsymbol{u}-\boldsymbol{u}_h),\boldsymbol{u}_h^c-\boldsymbol{u}_H^c)_{0,\mathscr{T}_h}+(\mu\nabla\times(\boldsymbol{u}-\boldsymbol{u}_h),\nabla\times(\boldsymbol{u}_h^c-\boldsymbol{u}_H^c))_{0,\mathscr{T}_h}+<[[\boldsymbol{u}_h]],\{\{\mu\nabla\times(\boldsymbol{u}_h^c-\boldsymbol{u}_H^c)\}\}>_{\mathscr{T}_h}=0,$$

which leads to

$$(\kappa(\boldsymbol{u}-\boldsymbol{u}_{h}),\boldsymbol{u}_{h}^{c}-\boldsymbol{u}_{H}^{c})_{0,\mathscr{T}_{h}}+(\mu\nabla\times(\boldsymbol{u}-\boldsymbol{u}_{h}),\nabla\times(\boldsymbol{u}_{h}^{c}-\boldsymbol{u}_{H}^{c}))_{0,\mathscr{T}_{h}}$$
$$=-<[[\boldsymbol{u}_{h}]],\{\{\mu\boldsymbol{u}_{h}^{c}-\boldsymbol{u}_{H}^{c}\}\}>_{\mathscr{T}_{h}}.$$
(3.28)

Using (3.22) and (3.23), we have

$$<[[\boldsymbol{u}_{h}]], \{\{\nabla \times (\boldsymbol{u}_{h}^{c} - \boldsymbol{u}_{H}^{c})\}\} >_{\mathscr{F}_{h}} = (\mathscr{L}_{h}(\boldsymbol{u}_{h}), \nabla \times (\boldsymbol{u}_{h}^{c} - \boldsymbol{u}_{H}^{c}))_{0,\mathscr{T}_{h}}$$
$$\leq C_{\mathscr{L}} \|h^{-1/2}[[\boldsymbol{u}_{h}]]\|_{0,\mathscr{T}_{h}} \|\nabla \times (\boldsymbol{u}_{h}^{c} - \boldsymbol{u}_{H}^{c})\|_{0,\mathscr{T}_{h}}. \quad (3.29)$$

Let $\boldsymbol{u}_h = \boldsymbol{u}_h^c + \boldsymbol{u}_h^{\perp}$ and $\boldsymbol{u}_H = \boldsymbol{u}_H^c + \boldsymbol{u}_H^{\perp}$, we have

$$u_{h} + u_{H}^{c} - u_{h}^{c} = u_{H} - u_{H}^{\perp} + u_{h}^{\perp}, \qquad (3.30)$$

where $\boldsymbol{u}_{H}^{c} \in \boldsymbol{U}_{H}^{c}, \, \boldsymbol{u}_{h}^{c} \in \boldsymbol{U}_{h}^{c}, \, \boldsymbol{u}_{H}^{\perp} \in \boldsymbol{U}_{H}^{\perp}, \, \boldsymbol{u}_{h}^{\perp} \in \boldsymbol{U}_{h}^{\perp}$. By (3.30), (3.28), (3.29) and Young's inequality, we get

$$\begin{split} \||\boldsymbol{u} - \boldsymbol{u}_{h}|\|_{h}^{2} \\ &= \|\kappa(\boldsymbol{u} - \boldsymbol{u}_{h})\|_{L^{2}(\Omega)}^{2} + \|\nabla \times \mu(\boldsymbol{u} - \boldsymbol{u}_{h})\|_{L^{2}(\Omega)}^{2} \\ &+ \sum_{f \in \mathscr{F}_{h}} \alpha h_{f}^{-1} < [[(\boldsymbol{u} - \boldsymbol{u}_{h})]], [[\boldsymbol{u} - \boldsymbol{u}_{h}]] >_{\mathscr{F}_{h}} \\ &= \||\boldsymbol{u} - \boldsymbol{u}_{h} - \boldsymbol{u}_{H}^{c} + \boldsymbol{u}_{h}^{c}\|\|_{h}^{2} - \||\boldsymbol{u}_{h}^{c} - \boldsymbol{u}_{H}^{c}\|\|_{h}^{2} - 2(\kappa(\boldsymbol{u} - \boldsymbol{u}_{h}), \boldsymbol{u}_{h}^{c} - \boldsymbol{u}_{H}^{c})_{0,\mathscr{F}_{h}} \\ &- 2(\mu \nabla \times (\boldsymbol{u} - \boldsymbol{u}_{h}), \nabla \times (\boldsymbol{u}_{h}^{c} - \boldsymbol{u}_{H}^{c}))_{0,\mathscr{F}_{h}} \\ &- 2\sum_{f \in \mathscr{F}_{h}} \alpha h_{f}^{-1} < [[(\boldsymbol{u} - \boldsymbol{u}_{h})]], [[\boldsymbol{u}_{h}^{c} - \boldsymbol{u}_{H}^{c}]] > \\ \lesssim \||\boldsymbol{u} - \boldsymbol{u}_{H}\|\|_{H}^{2} + 2\||\boldsymbol{u} - \boldsymbol{u}_{H}\|\|_{H}\||\boldsymbol{u}\|_{h}^{1} - \boldsymbol{u}_{H}^{1}\|\|_{h}^{1} + \||\boldsymbol{u}_{h}^{1} - \boldsymbol{u}_{H}^{1}\|\|_{h}^{2} - \||\boldsymbol{u}_{h}^{c} - \boldsymbol{u}_{H}^{c}\|\|_{h}^{2} \\ &+ 2\|h^{-1/2}[[\boldsymbol{u}_{h}]]\|_{0,\mathscr{F}_{h}}\|\nabla \times (\boldsymbol{u}_{h}^{c} - \boldsymbol{u}_{H}^{c}\|)\|_{0,\mathscr{F}_{h}} \\ \leq (1 + \delta_{1})\||\boldsymbol{u} - \boldsymbol{u}_{H}\|\|_{H}^{2} + (1 + \frac{1}{\delta_{1}})\||\boldsymbol{u}_{h}^{1} - \boldsymbol{u}_{H}^{1}\|\|_{h}^{2} - (1 - \hat{\delta}_{2}C_{\mathscr{L}})\||\boldsymbol{u}_{h}^{c} - \boldsymbol{u}_{H}^{c}\|\|_{h}^{2} \\ &+ \frac{C_{\mathscr{L}}}{\hat{\delta}_{2}}\|h^{-1/2}[[\boldsymbol{u}_{h}]]\|_{0,\mathscr{F}_{h}}^{2} \\ = (1 + \delta_{1})\||\boldsymbol{u} - \boldsymbol{u}_{H}\|\|_{H}^{2} + (1 + \frac{1}{\delta_{1}})\||\boldsymbol{u}_{h}^{1} - \boldsymbol{u}_{H}^{1}\|\|_{h}^{2} - (1 - \delta_{2})\||\boldsymbol{u}_{h}^{c} - \boldsymbol{u}_{H}^{c}\|\|_{h}^{2} \\ &+ \frac{C_{\mathscr{L}}}{\delta_{2}}\|h^{-1/2}[[\boldsymbol{u}_{h}]]\|_{0,\mathscr{F}_{h}}^{2}, \end{split}$$

16

where $\delta_2 = \hat{\delta}_2 C_{\mathscr{L}}$. Using $\boldsymbol{u}_H^c = \boldsymbol{u}_H - \boldsymbol{u}_H^{\perp}$, $\boldsymbol{u}_h^c = \boldsymbol{u}_h - \boldsymbol{u}_h^{\perp}$, triangle inequality and average inequality, we have

$$|||\boldsymbol{u}_{h}^{c} - \boldsymbol{u}_{H}^{c}|||_{h}^{2} \ge \frac{1}{2}|||\boldsymbol{u}_{h} - \boldsymbol{u}_{H}|||_{h}^{2} - |||\boldsymbol{u}_{h}^{\perp} - \boldsymbol{u}_{H}^{\perp}|||_{h}^{2}$$

By triangle inequality and (3.21), we obtain

$$\begin{aligned} \||\boldsymbol{u}_{h}^{\perp} - \boldsymbol{u}_{H}^{\perp}||_{h}^{2} &\leq 2(\||\boldsymbol{u}_{h}^{\perp}|\|_{h}^{2} + \||\boldsymbol{u}_{H}^{\perp}|\|_{H}^{2}) \\ &\leq 2\alpha \|h^{-1/2}[[\boldsymbol{u}_{h}^{\perp}]]\|_{0,\mathscr{T}_{h}}^{2} + 2\alpha \|h^{-1/2}[[\boldsymbol{u}_{H}^{\perp}]]\|_{0,\mathscr{T}_{h}}^{2}. \end{aligned}$$

Combining $[[u_H]] = [[u_H^{\perp} + u_H^c]] = [[u_H^{\perp}]]$ and (3.19), we have

$$\begin{aligned} \||\boldsymbol{u} - \boldsymbol{u}_{h}|\|_{h}^{2} &\leq (1 + \delta_{1}) \||\boldsymbol{u} - \boldsymbol{u}_{H}|\|_{H}^{2} - \frac{1 - \delta_{2}}{2} \||\boldsymbol{u}_{h} - \boldsymbol{u}_{H}|\|_{h}^{2} \\ &+ \frac{C_{3}}{\delta_{1}\delta_{2}\alpha} \Big(\eta^{2}(\boldsymbol{u}_{h}, \boldsymbol{p}_{h}; \mathcal{T}_{h}) + \eta^{2}(\boldsymbol{u}_{H}, \boldsymbol{p}_{H}; \mathcal{T}_{H}) \Big). \end{aligned}$$

3.3. Contraction of the error estimator

In this subsection, we prove the reduction of error indicators. Let us first consider the effect of changing the finite element function used in the estimator.

Lemma 3.8. Given $f \in L^2(\Omega)$ and two tetrahedral mesh \mathscr{T}_h , \mathscr{T}_H with $\mathscr{T}_H \leq \mathscr{T}_h$. Let $(\boldsymbol{v}_h, \boldsymbol{q}_h) \in U_h \times \boldsymbol{Q}_h$ and $(\boldsymbol{v}_H, \boldsymbol{q}_H) \in U_H \times \boldsymbol{Q}_H$. For any $\epsilon > 0$, we have

$$\eta^{2}(\boldsymbol{\nu}_{h},\boldsymbol{q}_{h};\mathcal{T}_{h}) \leq (1+\epsilon)\eta^{2}(\boldsymbol{\nu}_{H},\boldsymbol{q}_{H};\mathcal{T}_{h}) + C_{\epsilon} \|(\boldsymbol{\nu}_{h},\boldsymbol{q}_{h}) - (\boldsymbol{\nu}_{H},\boldsymbol{q}_{H})\|_{DG}^{2}, \qquad (3.31)$$

where C_{ϵ} depending on the ϵ , and the mesh size h < 1.

Proof. For any $\tau_* \in \mathscr{T}_h$, we will discuss each of the five components of the mark $\eta^2(\mathbf{v}_h, \mathbf{q}_h; \mathscr{T}_h)$.

Firstly, using the definition of $R_1(v_h, q_h)$ and triangle inequality, we have

$$\begin{aligned} \|R_{1}(\boldsymbol{v}_{h},\boldsymbol{q}_{h})\|_{L^{2}(\tau_{*})} & (3.32) \\ &= \|\boldsymbol{q}_{h} - \mu \nabla \times \boldsymbol{v}_{h}\|_{L^{2}(\tau_{*})} \\ &= \|\boldsymbol{q}_{h} - \boldsymbol{q}_{H} + \mu \nabla \times (\boldsymbol{v}_{H} - \boldsymbol{v}_{h}) + \boldsymbol{q}_{H} - \mu \nabla \times \boldsymbol{v}_{H}\|_{L^{2}(\tau_{*})} \\ &\lesssim \|\boldsymbol{q}_{H} - \nabla \times \boldsymbol{v}_{H}\|_{L^{2}(\tau_{*})} + \|\boldsymbol{q}_{h} - \boldsymbol{q}_{H}\|_{L^{2}(\tau_{*})} + \|\nabla \times (\boldsymbol{v}_{h} - \boldsymbol{v}_{H})\|_{L^{2}(\tau_{*})}. \end{aligned}$$

Secondly, using the definition of $R_2(v_h, q_h)$, triangle inequality and inverse inequality, we get

$$\begin{aligned} h_{\tau_{*}} \|R_{2}(\boldsymbol{v}_{h}, \boldsymbol{q}_{h})\|_{L^{2}(\tau_{*})} & (3.33) \\ &= h_{\tau_{*}}(\|\boldsymbol{f} - \nabla \times \boldsymbol{q}_{h} - \kappa \boldsymbol{v}_{h}\|_{L^{2}(\tau_{*})}) \\ &= h_{\tau_{*}}(\|\boldsymbol{f} - \nabla \times (\boldsymbol{q}_{h} - \boldsymbol{q}_{H}) - \kappa (\boldsymbol{v}_{h} - \boldsymbol{v}_{H}) - \nabla \times \boldsymbol{q}_{H} - \kappa \boldsymbol{v}_{H}\|_{L^{2}(\tau_{*})}) \\ &\leq h_{\tau_{*}}(\|\boldsymbol{f} - \nabla \times \boldsymbol{q}_{H} - \kappa \boldsymbol{v}_{H}\|_{L^{2}(\tau_{*})} + \|\nabla \times (\boldsymbol{q}_{h} - \boldsymbol{q}_{H})\|_{L^{2}(\tau_{*})} + \|\kappa(\boldsymbol{v}_{h} - \boldsymbol{v}_{H})\|_{L^{2}(\tau_{*})}) \\ &\leq h_{\tau_{*}}(\|R_{2}(\boldsymbol{v}_{H}, \boldsymbol{q}_{H})\|_{L^{2}(\tau_{*})} + h_{\tau_{*}}^{-1}\|(\boldsymbol{q}_{h} - \boldsymbol{q}_{H})\|_{L^{2}(\tau_{*})} + \|\kappa(\boldsymbol{v}_{h} - \boldsymbol{v}_{H})\|_{L^{2}(\tau_{*})}) \\ &\lesssim h_{\tau_{*}}\|R_{2}(\boldsymbol{v}_{H}, \boldsymbol{q}_{H})\|_{L^{2}(\tau_{*})} + \|(\boldsymbol{q}_{h} - \boldsymbol{q}_{H})\|_{L^{2}(\tau_{*})} + h_{\tau_{*}}\|\kappa(\boldsymbol{v}_{h} - \boldsymbol{v}_{H})\|_{L^{2}(\tau_{*})}. \end{aligned}$$

Similarly, using the definition of $R_3(v_h)$, triangle inequality and inverse inequality, we get

$$\begin{aligned} h_{\tau_{*}} \|R_{3}(\boldsymbol{v}_{h})\|_{L^{2}(\tau_{*})} & (3.34) \\ &= h_{\tau_{*}} \|\nabla \cdot (\boldsymbol{f} - \kappa \boldsymbol{v}_{h})\|_{L^{2}(\tau_{*})} \\ &= h_{\tau_{*}} \|\nabla \cdot (\boldsymbol{f} - \kappa \boldsymbol{v}_{H} + \kappa \boldsymbol{v}_{H} - \kappa \boldsymbol{v}_{h})\|_{L^{2}(\tau_{*})} \\ &\leq h_{\tau_{*}} (\|\nabla \cdot (\boldsymbol{f} - \kappa \boldsymbol{v}_{H})\|_{L^{2}(\tau_{*})} + \|\nabla \cdot \kappa (\boldsymbol{v}_{H} - \boldsymbol{v}_{h})\|_{L^{2}(\tau_{*})}) \\ &\lesssim h_{\tau_{*}} (\|R_{3}(\boldsymbol{v}_{H})\|_{L^{2}(\tau_{*})} + h_{\tau_{*}}^{-1} \|\kappa (\boldsymbol{v}_{H} - \boldsymbol{v}_{h})\|_{L^{2}(\tau_{*})}) \\ &\lesssim h_{\tau_{*}} \|R_{3}(\boldsymbol{v}_{H})\|_{L^{2}(\tau_{*})} + \|\kappa (\boldsymbol{v}_{H} - \boldsymbol{v}_{h})\|_{L^{2}(\tau_{*})}. \end{aligned}$$

Next, we discuss the jump $J_1(\boldsymbol{q}_h)$ and $J_2(\boldsymbol{v}_h)$. For any $f \in \mathscr{F}(\mathscr{T}_h)$, we let $f = \tau_*^1 \bigcap \tau_*^2$ with $\tau_*^1, \tau_*^2 \in \mathscr{T}_h$. Furthermore, using the definition of $J_1(\boldsymbol{q}_h)$, triangle inequality and trace inequality, we have

$$\begin{aligned} h_{f}^{1/2} \| J_{1}(\boldsymbol{q}_{h}) \|_{L^{2}(f)} & (3.35) \\ &= h_{f}^{1/2} \| [[\boldsymbol{q}_{h}]] \|_{L^{2}(f)} \\ &= h_{f}^{1/2} \| [[\boldsymbol{q}_{H} + \boldsymbol{q}_{h} - \boldsymbol{q}_{H}]] \|_{L^{2}(f)} \\ &\leq h_{f}^{1/2} (\| [[\boldsymbol{q}_{H}]] \|_{L^{2}(f)} + \| [[\boldsymbol{q}_{h} - \boldsymbol{q}_{H}]] \|_{L^{2}(f)}) \\ &\leq h_{f}^{1/2} \| [[\boldsymbol{q}_{H}]] \|_{L^{2}(f)} + h_{f}^{1/2} \| (\boldsymbol{q}_{h} - \boldsymbol{q}_{H}) |_{\tau_{*}^{1}} \|_{L^{2}(f)} + h_{f}^{1/2} \| (\boldsymbol{q}_{h} - \boldsymbol{q}_{H}) |_{\tau_{*}^{2}} \|_{L^{2}(f)} \\ &\lesssim h_{f}^{1/2} \| J_{1}(\boldsymbol{q}_{H}) \|_{L^{2}(f)} + \| (\boldsymbol{q}_{h} - \boldsymbol{q}_{H}) \|_{L^{2}(\tau_{*}^{1} \cup \tau_{*}^{2})}. \end{aligned}$$

Similarly, using the definition of $J_2(v_h)$, triangle inequality and trace inequality, we have

$$\begin{aligned} h_{f}^{1/2} \|J_{2}(\boldsymbol{v}_{h})\|_{L^{2}(f)} & (3.36) \\ &= h_{f}^{1/2} \|[[(f - \kappa \boldsymbol{v}_{h})]]\|_{L^{2}(f)} \\ &= h_{f}^{1/2} \|[[(f - \kappa \boldsymbol{v}_{H} + \kappa \boldsymbol{v}_{H} - \kappa \boldsymbol{v}_{h})]]\|_{L^{2}(f)} \\ &\leq h_{f}^{1/2} (\|[[(f - \kappa \boldsymbol{v}_{H})]]\|_{L^{2}(f)} + \|[[\kappa(\boldsymbol{v}_{H} - \boldsymbol{v}_{h})]]\|_{L^{2}(f)}) \\ &\leq h_{f}^{1/2} \|J_{2}(\boldsymbol{v}_{H})\|_{L^{2}(f)} + h_{f}^{1/2} (\|\kappa(\boldsymbol{v}_{H} - \boldsymbol{v}_{h})|_{\tau_{*}^{1}}\|_{L^{2}(f)} + \|\kappa(\boldsymbol{v}_{H} - \boldsymbol{v}_{h})|_{\tau_{*}^{2}}\|_{L^{2}(f)}) \\ &\lesssim h_{f}^{1/2} \|J_{2}(\boldsymbol{v}_{H})\|_{L^{2}(f)} + \|\kappa \boldsymbol{v}_{H} - \kappa \boldsymbol{v}_{h}\|_{L^{2}(\tau_{*}^{1} \cup \tau_{*}^{2})}. \end{aligned}$$

Finally, the desired result (3.31) is obtained by combining (3.32)-(3.36), Young's inequality and the shape regularity of mesh \mathcal{T}_h .

We then prove the contraction of the error estimator under the assumptions on the problem of (2.13)-(2.14).

Lemma 3.9. Given constant $\theta \in (0, 1)$ and two tetrahedral mesh \mathscr{T}_h , $\mathscr{T}_H(\mathscr{T}_H \leq \mathscr{T}_h)$. Let $(u_H, p_H) \in U_H \times Q_H$ be the solution of (2.13)-(2.14), and $\mathscr{R}_{\mathscr{T}_H \longrightarrow \mathscr{T}_h} = \mathscr{T}_H \setminus (\mathscr{T}_h \cap \mathscr{T}_H)$ be the

18

set of all element refined into \mathcal{T}_h on \mathcal{T}_H . Then, there is a constant $\lambda \in (0,1)$ independent of mesh size, such that

$$\eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathcal{T}_{h}) \leq \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathcal{T}_{H}) - \lambda \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathcal{R}_{\mathcal{T}_{H}} \rightarrow \mathcal{T}_{h}).$$
(3.37)

Proof. Assume that the tetrahedral mesh $\tau \in \mathscr{T}_H$ is divided into two new tetrahedral mesh τ^1_* and τ^2_* with equal volumes, where $\tau^1_*, \tau^2_* \in \mathscr{T}_h$. Thus, $h^3_{\tau^1_*} = |\tau^1_*| = |\tau^2_*| = h^3_{\tau^2_*} = 2^{-1}h^3_{\tau}$ by the shape regularity of mesh, which implies $h_{\tau^1_*} = h_{\tau^2_*} = 2^{-1/3}h_{\tau}$. Then, we have

$$\|R_{1}(\boldsymbol{u}_{H},\boldsymbol{p}_{H})\|_{L^{2}(\tau_{*}^{1})}^{2} + \|R_{1}(\boldsymbol{u}_{H},\boldsymbol{p}_{H})\|_{L^{2}(\tau_{*}^{2})}^{2} \leq \|R_{1}(\boldsymbol{u}_{H},\boldsymbol{p}_{H})\|_{L^{2}(\tau)}^{2},$$
(3.38)

and

$$\begin{aligned} & h_{\tau_{*}^{2}}^{2}(\|R_{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H})\|_{L^{2}(\tau_{*}^{1})}^{2}+\|R_{3}(\boldsymbol{u}_{H})\|_{L^{2}(\tau_{*}^{1})}^{2}) \\ & +h_{\tau_{*}^{2}}^{2}(\|R_{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H})\|_{L^{2}(\tau_{*}^{2})}^{2}+\|R_{3}(\boldsymbol{u}_{H})\|_{L^{2}(\tau_{*}^{2})}^{2}) \\ & \leq 2^{-2/3}h_{\tau}^{2}(\|R_{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H})\|_{L^{2}(\tau)}^{2}+\|R_{3}(\boldsymbol{u}_{H})\|_{L^{2}(\tau)}^{2}). \end{aligned}$$
(3.39)

For any $f \in \partial(\tau_*^1 \cup \tau_*^2)$, which can be divided into three parts;

(1) For the first part, there are two of the faces are constant and belong to au .

(2) For the second part, there are two new faces that overlap and are used to divide the mesh τ . Since $(\boldsymbol{u}_H, \boldsymbol{p}_h) \in \boldsymbol{U}_H \times \boldsymbol{Q}_H$ is a continuous polynomial in the region τ , it follows that the value of $[[\boldsymbol{p}_h]]$ and $[[(\boldsymbol{f} - \kappa \boldsymbol{u}_H)]]$ on this surface is equal to zero.

(3) For the third part, there are four faces that are obtained by dividing the two faces in the τ into two.

Furthermore, we obtain

$$\eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\boldsymbol{\tau}_{*}^{1}) + \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\boldsymbol{\tau}_{*}^{2}) \leq \overline{\gamma}\eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\boldsymbol{\tau}).$$
(3.40)

where constant $\overline{\gamma} \in (0, 1)$ independent of mesh τ .

Next, since $\mathscr{R}_{\mathscr{T}_H \to \mathscr{T}_h}$ represents the part of the set in the tetrahedral set \mathscr{T}_H that will be used to be refined, it follows that $\mathscr{R}_{\mathscr{T}_H \to \mathscr{T}_h} \subset \mathscr{T}_H$. Let $\overline{\mathscr{R}_{\mathscr{T}_H \to \mathscr{T}_h}}$ denote the part of the cell set that has been refined in the tetrahedral set \mathscr{T}_H , we have $\overline{\mathscr{R}_{\mathscr{T}_h \to \mathscr{T}_h}} \in \mathscr{T}_h$. Obviously, $\mathscr{T}_H \setminus \mathscr{R}_{\mathscr{T}_H \to \mathscr{T}_h} = \mathscr{T}_h \setminus \overline{\mathscr{R}_{\mathscr{T}_H \to \mathscr{T}_h}}$. Then combining the (3.40), and the marking strategy (2.18), we have

$$\begin{split} \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathcal{T}_{h}) &= \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathcal{T}_{h}\setminus\overline{\mathscr{R}_{\mathcal{T}_{H}}\rightarrow\mathscr{T}_{h}}) + \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\overline{\mathscr{R}_{\mathcal{T}_{H}}\rightarrow\mathscr{T}_{h}}) \\ &\leq \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathcal{T}_{H}\setminus\mathscr{R}_{\mathcal{T}_{H}}\rightarrow\mathscr{T}_{h}) + \gamma\eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathscr{R}_{\mathcal{T}_{H}}\rightarrow\mathscr{T}_{h}) \\ &\leq \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathcal{T}_{H}) + (\overline{\gamma}-1)\eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathscr{R}_{\mathcal{T}_{H}}\rightarrow\mathscr{T}_{h}) \\ &\leq \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathcal{T}_{H}) - \lambda\eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathscr{R}_{\mathcal{T}_{H}}\rightarrow\mathscr{T}_{h}), \end{split}$$

where $\lambda = 1 - \overline{\gamma} \in (0, 1)$ independent of mesh size.

Now, we combine the Lemmas 3.6, 3.8 and 3.9 to prove the reduction of error indicators.

Lemma 3.10. Given a constant $\theta \in (0, 1)$ and two tetrahedral mesh \mathcal{T}_h , $\mathcal{T}_H(\mathcal{T}_H \leq \mathcal{T}_h)$. Let $(\boldsymbol{u}_h, \boldsymbol{p}_h) \in \boldsymbol{U}_h \times \boldsymbol{Q}_h$ and $(\boldsymbol{u}_H, \boldsymbol{p}_H) \in \boldsymbol{U}_H \times \boldsymbol{Q}_H$ be the solutions of (2.13)-(2.14), respectively. For any $\epsilon > 0$ and $\lambda \in (0, 1)$, we have

$$(1 - \frac{C_{\epsilon}}{\alpha})\eta^{2}(\boldsymbol{u}_{h}, \boldsymbol{p}_{h}; \mathcal{T}_{h}) \leq (1 + \epsilon + \frac{C_{\epsilon}}{\alpha})\eta^{2}(\boldsymbol{u}_{H}, \boldsymbol{p}_{H}; \mathcal{T}_{H}) - (1 + \epsilon)\lambda\eta^{2}(\boldsymbol{u}_{H}, \boldsymbol{p}_{H}; \mathcal{R}_{\mathcal{T}_{H}} \rightarrow \mathcal{T}_{h}) + C_{\epsilon} |||\boldsymbol{u}_{h} - \boldsymbol{u}_{H}|||_{h}^{2}, \quad (3.41)$$

where constant C_{ϵ} depending on the ϵ and mesh size.

Proof. Using the Lemmas 3.6, 3.8 and 3.9, we have

$$\begin{split} \eta^{2}(\boldsymbol{u}_{h},\boldsymbol{p}_{h};\mathscr{T}_{h}) &\leq (1+\epsilon) \bigg(\eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathscr{T}_{H}) - \lambda \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathscr{R}_{\mathscr{T}_{H}} \rightarrow \mathscr{T}_{h}) \bigg) \\ &+ C_{\epsilon} \| (\boldsymbol{u}_{h},\boldsymbol{p}_{h}) - (\boldsymbol{u}_{H},\boldsymbol{p}_{H}) \|_{DG}^{2} \\ &\leq (1+\epsilon) \bigg(\eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathscr{T}_{H}) - \lambda \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathscr{R}_{\mathscr{T}_{H}} \rightarrow \mathscr{T}_{h}) \bigg) \\ &+ C_{\epsilon} \| |\boldsymbol{u}_{h} - \boldsymbol{u}_{H} \| \|_{h}^{2} + \| \boldsymbol{p}_{h} - \boldsymbol{p}_{H} \|_{L^{2}(\Omega)}^{2} \\ &\leq (1+\epsilon) \bigg(\eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathscr{T}_{H}) - \lambda \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathscr{R}_{\mathscr{T}_{H}} \rightarrow \mathscr{T}_{h}) \bigg) \\ &+ C_{\epsilon} \| |\boldsymbol{u}_{h} - \boldsymbol{u}_{H} \| \|_{h}^{2} + \frac{C_{\epsilon}}{\alpha} \bigg(\eta^{2}(\boldsymbol{u}_{h},\boldsymbol{p}_{h};\mathscr{T}_{h}) + \eta^{2}(\boldsymbol{u}_{H},\boldsymbol{p}_{H};\mathscr{T}_{H}) \bigg), \end{split}$$

which completes the proof.

3.4. Convergence result

Now, we proved that the sum of the norm of the error and the scaled error indicator is attenuated.

Theorem 3.2. For a given $\theta \in (0, 1)$, let $\{\mathscr{T}_k, U_k, Q_k, u_k, p_k, \eta(u_k, p_k; \mathscr{T}_k)\}_{k\geq 0}$ be the sequence of meshes, Mixed discrete solution (defined by (2.13)-(2.14)), and the estimate indicator produced by the **AMIPDG** algorithm. Then there exist constants $\rho > 0$, $\delta \in (0, 1)$, which depend on marking parameter θ and the shape regularity of the initial mesh \mathscr{T}_0 , such that

$$\||\boldsymbol{u} - \boldsymbol{u}_{k+1}|\|_{k+1}^2 + \rho \eta^2 (\boldsymbol{u}_{k+1}, \boldsymbol{p}_{k+1}; \mathcal{T}_{k+1}) \le \delta \Big(\||\boldsymbol{u} - \boldsymbol{u}_k\|\|_k^2 + \rho \eta^2 (\boldsymbol{u}_k, \boldsymbol{p}_k; \mathcal{T}_k) \Big)$$

Proof. Setting $\tilde{\rho} = \frac{1-\delta_2}{2C_e}$, then multiply the both sides of the (3.41) inequality by $\tilde{\rho}$, we get

$$\widetilde{\rho}(1 - \frac{C_{\epsilon}}{\alpha})\eta^{2}(\boldsymbol{u}_{k+1}, \boldsymbol{p}_{k+1}; \mathcal{T}_{k+1})$$

$$\leq \widetilde{\rho}(1 + \epsilon + \frac{C_{\epsilon}}{\alpha})\eta^{2}(\boldsymbol{u}_{k}, \boldsymbol{p}_{k}; \mathcal{T}_{k}) - \widetilde{\rho}(1 + \epsilon)\lambda\eta^{2}(\boldsymbol{u}_{k}, \boldsymbol{p}_{k}; \mathcal{R}_{\mathcal{T}_{k} \to \mathcal{T}_{k+1}})$$

$$+ \frac{1 - \delta_{2}}{2} |||\boldsymbol{u}_{k+1} - \boldsymbol{u}_{k}|||_{h}^{2}.$$
(3.42)

Next, by the (3.26) and (3.42), we have

$$\begin{aligned} \||\boldsymbol{u} - \boldsymbol{u}_{k+1}|||_{k+1}^{2} + \widetilde{\rho}(1 - \frac{C_{\epsilon}}{\alpha})\eta^{2}(\boldsymbol{u}_{k+1}, \boldsymbol{p}_{k+1}; \mathscr{T}_{k+1}) \\ &\leq (1 + \delta_{1})\||\boldsymbol{u} - \boldsymbol{u}_{k}|||_{k}^{2} + \frac{C_{3}}{\delta_{1}\delta_{2}\alpha} \Big(\eta^{2}(\boldsymbol{v}_{k+1}, \boldsymbol{q}_{k+1}; \mathscr{T}_{k+1}) + \eta^{2}(\boldsymbol{v}_{k}, \boldsymbol{q}_{k}; \mathscr{T}_{k})\Big) \\ &+ \widetilde{\rho}(1 + \epsilon + \frac{C_{\epsilon}}{\alpha})\eta^{2}(\boldsymbol{u}_{k}, \boldsymbol{p}_{k}; \mathscr{T}_{k}) - \widetilde{\rho}(1 + \epsilon)\lambda\eta^{2}(\boldsymbol{u}_{k}, \boldsymbol{p}_{k}; \mathscr{T}_{k+1}). \end{aligned}$$
(3.43)

First move the term and then according to Dörfler marking strategy (2.18), the Theorem 3.1 and $\||\cdot|\|_h \leq \|\cdot\|_{DG}$, we know $-\eta^2(\boldsymbol{v}_k, \boldsymbol{q}_k; \mathscr{R}_{\mathscr{T}_k \to \mathscr{T}_{k+1}}) \leq -\theta \eta^2(\boldsymbol{v}_k, \boldsymbol{q}_k; \mathscr{T}_k)$, then

$$\begin{split} \||\boldsymbol{u}-\boldsymbol{u}_{k+1}|||_{k+1}^{2} &+ \tilde{\rho}(1-\frac{C_{\epsilon}}{\alpha}-\frac{C_{3}}{\tilde{\rho}\delta_{1}\delta_{2}\alpha})\eta^{2}(\boldsymbol{u}_{k+1},\boldsymbol{p}_{k+1};\mathcal{T}_{k+1}) \\ &\leq (1+\delta_{1})\||\boldsymbol{u}-\boldsymbol{u}_{k}|||_{k}^{2}-\frac{\tilde{\rho}(1+\epsilon)\lambda\theta}{2}\eta^{2}(\boldsymbol{u}_{k},\boldsymbol{p}_{k};\mathcal{T}_{k}) \\ &+ \tilde{\rho}\Big(1+\epsilon+\frac{C_{\epsilon}}{\alpha}+\frac{C_{3}}{\tilde{\rho}\delta_{1}\delta_{2}\alpha}-\frac{(1+\epsilon)\lambda\theta}{2}\Big)\eta^{2}(\boldsymbol{u}_{k},\boldsymbol{p}_{k};\mathcal{T}_{k}) \\ &\leq (1+\delta_{1}-\frac{\tilde{\rho}(1+\epsilon)\lambda\theta C_{1}^{-1}}{2})\||\boldsymbol{u}-\boldsymbol{u}_{k}|||_{k}^{2} \\ &+ \tilde{\rho}\Big(1+\epsilon+\frac{C_{\epsilon}}{\alpha}+\frac{C_{3}}{\tilde{\rho}\delta_{1}\delta_{2}\alpha}-\frac{(1+\epsilon)\lambda\theta}{2}\Big)\eta^{2}(\boldsymbol{u}_{k},\boldsymbol{p}_{k};\mathcal{T}_{k}). \end{split}$$

For convenience, denote

$$\begin{split} \beta_1 &= 1 - \frac{C_{\epsilon}}{\alpha} - \frac{C_3}{\widetilde{\rho} \delta_1 \delta_2 \alpha}, \\ \beta_2 &= 1 + \delta_1 - \frac{\widetilde{\rho} (1 + \epsilon) \lambda \theta C_1^{-1}}{2}, \\ \beta_3 &= (1 + \epsilon) (1 - \frac{\lambda \theta}{2}) + \frac{C_{\epsilon}}{\alpha} + \frac{C_3}{\widetilde{\rho} \delta_1 \delta_2 \alpha}. \end{split}$$

Thus

$$\||\boldsymbol{u} - \boldsymbol{u}_{k+1}|\|_{k+1}^2 + \tilde{\rho}\beta_1\eta^2(\boldsymbol{u}_{k+1}, \boldsymbol{p}_{k+1}; \mathcal{T}_{k+1}) \leq \beta_2\||\boldsymbol{u} - \boldsymbol{u}_k\|\|_k^2 + \tilde{\rho}\beta_3\eta^2(\boldsymbol{u}_k, \boldsymbol{p}_k; \mathcal{T}_k).$$

Next, we firstly choose $\delta_1 = \frac{\tilde{\rho}(1+\epsilon)\lambda\theta C_1^{-1}}{4}$, then select the appropriate δ_2 to make $\tilde{\rho} = \frac{1-\delta_2}{2C_\epsilon}$ smaller to ensure $0 < \delta_1 < 1$, Secondly, we let $\epsilon > 0$ and $(1+\epsilon)(1-\frac{\lambda\theta}{2}) = 1-\frac{\lambda\theta}{4}$ ($\lambda\theta \in (0,1)$), therefore

$$\beta_2 = 1 - \delta_1 \in (0, 1), \ (1 + \epsilon)(1 - \frac{\lambda \theta}{2}) < 1.$$

Furthermore, we choose a sufficiently large penalty parameter α such that

$$\beta_1 > \beta_3.$$

Finally, there is a constant $\delta = \max\{\beta_2, \frac{\beta_1}{\beta_3}\}$. Then, we let $\rho = \tilde{\rho}\beta_1$, and obtain

$$\||\boldsymbol{u} - \boldsymbol{u}_{k+1}|\|_{k+1}^2 + \rho \eta^2 (\boldsymbol{u}_{k+1}, \boldsymbol{p}_{k+1}; \mathscr{T}_{k+1}) \le \delta \Big(\||\boldsymbol{u} - \boldsymbol{u}_k|\|_k^2 + \rho \eta^2 (\boldsymbol{u}_k, \boldsymbol{p}_k; \mathscr{T}_k) \Big).$$

Corollary 3.1. Under the conditions of Theorem 3.2, we have

 $\|(\boldsymbol{u},\boldsymbol{p})-(\boldsymbol{u}_k,\boldsymbol{p}_k)\|_{DG}^2+\rho\eta^2(\boldsymbol{u}_k,\boldsymbol{p}_k;\mathcal{T}_k)\leq\delta^k\widetilde{C}_{\delta}.$

where $\widetilde{C}_{\delta} = C \Big(\|(\boldsymbol{u}, \boldsymbol{p}) - (\boldsymbol{u}_0, \boldsymbol{p}_0)\|_{DG}^2 + \rho \eta^2 (\boldsymbol{u}_0, \boldsymbol{p}_0; \mathcal{T}_0) \Big)$. Therefore, for a given precision, the **AMIPDG** method will terminate after a finite number of operations.

Proof. Using the Remark 3.2 and Theorem 3.2, we have

$$\begin{aligned} \|(\boldsymbol{u},\boldsymbol{p}) - (\boldsymbol{u}_k,\boldsymbol{p}_k)\|_{DG}^2 + \rho \eta^2(\boldsymbol{u}_k,\boldsymbol{p}_k;\mathscr{T}_k) &\leq C \bigg(\||\boldsymbol{u} - \boldsymbol{u}_k\|\|_k^2 + \rho \eta^2(\boldsymbol{u}_k,\boldsymbol{p}_k;\mathscr{T}_k) \bigg) \\ &\leq \delta^k \widetilde{C}_\delta. \end{aligned}$$

4. Numerical experiments

In this section, we test some numerical experiments to show the efficiency and the robustness of AMIPDG. We carry out these numerical experiments by using the MATLAB software package iFEM [6]. In Experiments 4.1 and 4.2, we take $p = \nabla \times u$.

In Example 4.1, we discuss the influence of the penalty parameter α on the error in $\|\cdot\|_{DG}$ norm, and observe the dependency of the condition number of stiffness matrix on α .

Example 4.1. Let $\Omega := [0,1] \times [0,1] \times [0,1]$, we construct the following analytical solution of the model (1.1)-(1.2):

$$u = \begin{pmatrix} x(x-1)y(y-1)z(z-1) \\ \sin(\pi x)\sin(\pi y)\sin(\pi z) \\ (1-e^x)(1-e^{x-1})(1-e^y)(1-e^{y-1})(1-e^z)(1-e^{z-1}) \end{pmatrix}.$$

It is easy to see that the solution **u** satisfies the boundary condition $\mathbf{u} \times \mathbf{n} = 0$ on $\partial \Omega$.

In this example, we get a uniform mesh by partitioning the x-, y- and z-axes into equally distributed $M(M \ge 2)$ subintervals, and then dividing one cube into six tetrahedrons. Let h = 1/M be mesh sizes for different tetrahedrons meshes. We fixed mesh with h = 1/4 and report the error estimates in $\|\cdot\|_{DG}$ norm and condition number of stiffness matrices for different penalty parameters $\alpha = 1, 10, 100, 500$ and 1000 in Table 1. We note that $\|\mathbf{u} - \mathbf{u}_h\|_0$ increases at first and then decreases as the penalty parameter α increases.

α	1	10	100	500	1000
$\frac{\ (\boldsymbol{p}-\boldsymbol{p}_h,\boldsymbol{u}-\boldsymbol{u}_h)\ _{\mathrm{DG}}}{\ \boldsymbol{u}-\boldsymbol{u}_h\ _{\mathrm{DG}}}$	3.949e+00	1.133e-00	8.614e-01	8.649e-01	8.659e-01
Cond	3.235e+04	7.021e+04	5.959e+05	2.995e+06	6.150e+06

Table 1: The error in $\|\cdot\|_{DG}$ norms and condition number of stiffness matrices with h = 1/4.

The condition numbers of stiffness matrices increase with the increase of penalty parameters α .

As a way to balance, in the following numerical tests, we always choose $\alpha = 100$.

Noting that we only consider uniform meshes in Example 4.1. Next we test adaptive meshes.

Example 4.2. Let $\Omega := [0,1] \times [0,1] \times [0,1]$, we construct the following analytical solution of the model (1.1)-(1.2)

$$u = \begin{pmatrix} \frac{x(x-1)y(y-1)z(z-1)}{x^2+y^2+z^2+0.001} \\ \frac{x(x-1)y(y-1)z(z-1)}{x^2+y^2+z^2+0.001} \\ -\frac{x(x-1)y(y-1)z(z-1)}{x^2+y^2+z^2+0.001} \end{pmatrix}.$$

Note that the solution **u** satisfies the condition $\mathbf{u} \times \mathbf{n} = 0$ on $\partial \Omega$.

The right of Figure 1 shows an adaptively refined mesh with marking parameter- θ = 0.7 after *k* = 18. The grid is locally refined near the origin.



Figure 1: Left: the initial mesh with 1152 DoFs. Right: the adaptive mesh($\theta = 0.7$) with 181104 DoFs after 18 refinements.

The Figure 2 shows the curves of $\log N - \log \eta (\boldsymbol{u}_k, \boldsymbol{p}_k; \mathcal{T}_k)$ for parameters $\theta = 0.3, 0.5, 0.7$. The curves indicate the convergence and the quasi-optimality of the adaptive algorithm AMIPDG of $\eta (\boldsymbol{u}_k, \boldsymbol{p}_k; \mathcal{T}_k)$.

Acknowledgment

The first author is supported by the East China University of Technology (DHBK2019209) and Jiangxi Province Education Department (GJJ200755). The second, third and fourth authors are supported by the National Natural Science Foundation of China (Grant No. 12071160). The third author is also supported by the National Natural Science Foundation of China (Grant No. 11901212).

K Liu et al.



Figure 2: Quasi optimality of the AMIPDG of the error $\eta(u_k, p_k; \mathscr{T}_k)$ with different marking parameters θ .

References

- B. AYUSO DE DIOS, R. HIPTMAIR AND C.L. PAGLIANTINI, Auxiliary space preconditioners for SIP-DG discretizations of H(curl)-elliptic problems with discontinuous coefficients. *IMA J. Numer. Anal.* 37(2017), pp, 646-686.
- [2] A. BONITO AND R.H. NOCHETTO, Quasi-optimal convergence rate of an adaptive discontinuous Galerkin method. SIAM J. Numer. Anal. 48(2010), pp. 734–771.
- [3] C. CARSTENSEN AND R.H. HOPPE, Unified framework for an a posteriori error analysis of non-standard finite element approximations of *H*(*curl*)-elliptic problems. *J. Numer. Math.* 17(2009), pp. 27–44.
- [4] C. CARSTENSEN, R.H. HOPPR, N. SHARMA AND T. WARBURTON, Adaptive hybridized interior penalty discontinuous galerkin methods for *H*(*curl*)–elliptic problems. *Numer. Math. Theor. Meth. Appl.* 4(2011), pp. 13–37.
- [5] C. CARSTENSEN AND R. MA, Adaptive mixed finite element methods for non-self-adjoint indefinite second-order elliptic pdes with optimal rates. *SIAM J. Numer. Anal.* 59(2021), pp. 955–982.
- [6] L. CHEN, iFEM: an innovative finite element method package in MATLAB. Technical report, University of California at Irvine (2009).
- [7] L. CHEN, M. HOLST AND J.C. XU, Convergence and optimality of adaptive mixed finite element methods. *Math. Comp.* 78(2009), pp. 35–53.
- [8] E.T. CHUNG, M.C. YUEN AND L.Q. ZHONG, A-posteriori error analysis for a staggered discontinuous Galerkin discretization of the time-harmonic Maxwell's equations. *Appl. Math. Comput.* 237(2014), pp. 613–631.
- [9] L. DÖRFLER, A convergent adaptive algorithm for Poisson's equation. SIAM J. Numer. Anal. 33(1996), pp. 1106–1124.
- [10] S.H. DU AND X.P. XIE, Convergence of an adaptive mixed finite element method for

convection-diffusion-reaction equations. Sci. China Math. 58(2015), pp. 1327-1348.

- [11] P. HOUSTON, I. PERUGIA, A. SCHNEEBELI ADN D. SCHÖTZAU, Interior penalty method for the indefinite time-harmonic Maxwell equations. *Numer. Math.* 100(2005), pp. 485–518.
- [12] W. JIANG, N. LIU, Y. TANG AND Q.H. LIU, Mixed finite element method for 2D vector Maxwell's eigenvalue problem in anisotropic media. *Progress In Electromagnetics Research* 148(2014), pp. 159–170.
- [13] C. JOG AND A. NANDY, Mixed finite elements for electromagnetic. Comput. Math. Appl. 68(2014), pp. 887–902.
- [14] F. KIKUCHI, Mixed and penalty formulations for finite element analysis of an eigenvalue problem in electromagnetism. *Comput. Methods Appl. Mech. Engrg.* **64**(1987), pp. 509–521.
- [15] N. LIU, L. TOBÓN, Y. TANG AND Q.H. LIU, Mixed spectral element method for 2D Maxwell's eigenvalue problem. *Commun. Comput. Phys.* 17(2015), pp. 458–486.
- [16] P. MONK, Finite Element Methods for Maxwell Equations. Numerical Mathematics and Scientific Computation. Oxford University Press, Oxford(2003).
- [17] J.C. NÉDÉLEC, Mixed finite elements in \mathbb{R}^3 . Numer. Math. **35**(1980), pp. 315–341.
- [18] I. PERUGIA, D. SCHÖTZAU AND P. MONK, Stabilized interior penalty methods for the timeharmonic Maxwell equations. *Comput. Methods Appl. Mech. Eng.* 191(2002), pp. 4675–4697.
- [19] J. SCHÖBERL, A posteriori error estimates for Maxwell equations. Math. Comp. 77(2008), pp. 633–649.
- [20] X.Q. XING AND L.Q. ZHONG, A posteriori error estimate of discontinuous Galerkin Method for H(curl)-elliptic problems (in Chinese). *Journal of South China Normal University (Natural Science Edition*). 44(2012), pp. 18–21.