# A Probabilistic Perspective on Feller, Pollard and the Complete Monotonicity of the Mittag-Leffler Function

Nomvelo Karabo Sibisi

sbsnom005@myuct.ac.za

January 19, 2023

#### Abstract

The main contribution of this paper is the use of probability theory to prove that the three-parameter Mittag-Leffler function is the Laplace transform of a distribution and thus completely monotone. Pollard used contour integration to prove the result in the one-parameter case. He also cited personal communication by Feller of a discovery of the result by "methods of probability theory". Feller used the two-dimensional Laplace transform of a bivariate distribution to derive the result. We pursue the theme of probability theory to explore complete monotonicity beyond the contribution due to Feller. Our approach involves an interplay between mixtures and convolutions of stable and gamma densities, together with a limit theorem that leads to a novel integral representation of the three-parameter Mittag-Leffler function (also known as the Prabhakar function).

*Keywords*— Probabilistic reasoning; complete monotonicity; stable & gamma distributions; Mittag-Leffler function; Prabhakar function.

### 1 Introduction

The problem of interest in this paper is the study of the complete monotonicity of the Mittag-Leffler function. Complete monotonicity is an analytic property of functions. Accordingly, Pollard [18] used analytic methods to prove the property in the instance of the Mittag-Leffler function. Pollard also cited personal communication by Feller of a discovery of the result by "methods of probability theory". However, Pollard's comment notwithstanding, the published proof by Feller [7] (XIII.8) also may be regarded as more analytic than probabilistic (we discuss both approaches later in this section). This prompted us to ask the following:

1. What might constitute a "method of probability theory" in proving an analytic property of a function, at least in the context of proving that the Mittag-Leffler function is completely monotone?

2. What additional or complementary insight, if any, might the method of probability theory offer relative to an analytic method?

The strategy of this paper is simple – assign appropriate probability distributions and use the sum and product rules of probability theory to explore analytic attributes of associated functions. Beyond reproducing known analytic results due to Pollard and Feller, we discuss the generalisation that flows from adopting such reasoning. We start with definitions of complete monotonicity and the Mittag-Leffler function.

#### 1.1 Definitions

An infinitely differentiable function  $\varphi(x)$  on x > 0 is completely monotone if its derivatives  $\varphi^{(n)}(x)$  satisfy  $(-1)^n \varphi^{(n)}(x) \ge 0$ ,  $n \ge 0$ . Bernstein's theorem states that  $\varphi(x)$  is completely monotone iff it may be expressed as

$$\varphi(x) = \int_0^\infty e^{-xt} \, dF(t) = \int_0^\infty e^{-xt} f(t) dt \tag{1}$$

for a non-decreasing distribution function F(t) with density f(t), *i.e.*  $F(t) = \int_0^t f(u) du$ . The first integral in (1) is formally called the Laplace-Stieltjes transform of F and the latter the (ordinary) Laplace transform of f. For bounded F(t),  $\varphi(x)$  is defined on  $x \ge 0$ . Integrating (1) by parts in this case gives  $\varphi(x)$  in terms of the ordinary Laplace transform of F:

$$\varphi(x) = x \int_0^\infty e^{-xt} F(t) \, dt = \int_0^\infty e^{-t} F(t/x) \, dt \tag{2}$$

The (one-parameter) Mittag-Leffler function  $E_{\alpha}(x)$  is defined by the infinite series

$$E_{\alpha}(x) = \sum_{k=0}^{\infty} \frac{x^k}{\Gamma(\alpha k + 1)} \quad \alpha \ge 0$$
(3)

For later reference, the Laplace transform of  $E_{\alpha}(-\lambda x^{\alpha})$  ( $\lambda > 0$ ) is

$$\int_0^\infty e^{-sx} E_\alpha(-\lambda x^\alpha) \, dx = \frac{s^{\alpha-1}}{\lambda + s^\alpha} \qquad \operatorname{Re}(s) \ge 0 \tag{4}$$

We shall introduce the two and three-parameter generalisations below. For now, we may turn to the problem of proving the complete monotonicity of  $E_{\alpha}(-x)$ . We discuss the approaches due to Pollard and Feller in turn before turning to our probabilistic perspective.

#### 1.2 Pollard's Method

In a 1948 paper, Pollard [18] led with the opening remark:

"W. Feller communicated to me his discovery – by the methods of probability theory – that if  $0 \le \alpha \le 1$  the function  $E_{\alpha}(-x)$  is completely monotonic for  $x \ge 0$ . This means that it can be written in the form

$$E_{\alpha}(-x) = \int_{0}^{\infty} e^{-xt} dP_{\alpha}(t)$$

where  $P_{\alpha}(t)$  is nondecreasing and bounded. In this note we shall prove this fact directly and determine the function  $P_{\alpha}(t)$  explicitly."

[we use  $P_{\alpha}$  where Pollard used  $F_{\alpha}$ , which we reserve for another purpose]

Having dispensed with  $E_0(-x) = 1/(1+x)$  and  $E_1(-x) = e^{-x}$  since "there is nothing to be proved in these cases", Pollard used a contour integral representation of  $E_{\alpha}(-x)$ :

$$E_{\alpha}(-x) = \frac{1}{2\pi i} \oint_C \frac{s^{\alpha-1}e^s}{x+s^{\alpha}} ds = \frac{1}{2\pi i\alpha} \oint_{C'} \frac{e^{z^{\frac{1}{\alpha}}}}{x+z} dz$$
(5)

to prove that

$$p_{\alpha}(t) \equiv P'_{\alpha}(t) = \frac{1}{\alpha} f_{\alpha}(t^{-1/\alpha}) t^{-1/\alpha - 1} \qquad 0 < \alpha < 1$$
 (6)

where  $f_{\alpha}(t)$  is defined by

$$e^{-s^{\alpha}} = \int_0^{\infty} e^{-st} f_{\alpha}(t) dt \qquad \qquad 0 < \alpha < 1 \tag{7}$$

Pollard [17] had earlier proved that  $f_{\alpha}(t) > 0$ , so that  $p_{\alpha}(t) \ge 0$ , thereby completing his proof that  $E_{\alpha}(-x)$  is completely monotone for  $0 \le \alpha \le 1$ . Pollard stopped at the point of deriving (6), the density  $p_{\alpha}(t) \equiv P'_{\alpha}(t)$ . As per initial task, we proceed to discuss  $P_{\alpha}(t)$  explicitly. We first recognise  $f_{\alpha}(t)$  as the density of the stable distribution  $F_{\alpha}$  on  $[0, \infty)$ 

$$F_{\alpha}(t) = \int_0^t f_{\alpha}(u) \, du \qquad 0 < \alpha < 1 \tag{8}$$

with normalisation  $F_{\alpha}(\infty) = 1$ . In turn, the Pollard distribution  $P_{\alpha}(t)$  is

$$P_{\alpha}(t) = \int_{0}^{t} p_{\alpha}(u) \, du = \frac{1}{\alpha} \int_{0}^{t} f_{\alpha}(u^{-1/\alpha}) \, u^{-1/\alpha - 1} \, du \tag{9}$$

Janson [13] derived  $P_{\alpha}(t)$  as a limiting distribution of a Pólya urn scheme.  $P_{\alpha}(t)$  is known as the Mittag-Leffler distribution in the probabilistic literature (one of two distributions bearing the same name as discussed later).

Setting  $y = u^{-1/\alpha}$  in (9) gives a simple relation between  $P_{\alpha}$  and  $F_{\alpha}$ :

$$P_{\alpha}(t) = \int_{t^{-1/\alpha}}^{\infty} f_{\alpha}(y) \, dy = 1 - \int_{0}^{t^{-1/\alpha}} f_{\alpha}(y) \, dy \equiv 1 - F_{\alpha}(t^{-1/\alpha}) \tag{10}$$

This 'duality' between the Mittag-Leffler and stable distributions is key to the discussion that follows. The Pollard result may accordingly be written in several equivalent forms:

$$E_{\alpha}(-x) = \int_{0}^{\infty} e^{-xt} dP_{\alpha}(t) = \int_{0}^{\infty} e^{-t} P_{\alpha}(t/x) dt$$
  
or 
$$E_{\alpha}(-x^{\alpha}) = \int_{0}^{\infty} e^{-t} P_{\alpha}(x^{-\alpha}t) dt = \int_{0}^{\infty} e^{-t} (1 - F_{\alpha}(xt^{-1/\alpha})) dt$$
(11)

Another representation arising from change of variable in Pollard's original result is

$$\alpha E_{\alpha}(-x^{\alpha}) = \int_{0}^{\infty} e^{-x^{\alpha}u} f_{\alpha}(u^{-1/\alpha}) u^{-1/\alpha - 1} du$$
$$= x \int_{0}^{\infty} e^{-t} f_{\alpha}(xt^{-1/\alpha}) t^{-1/\alpha - 1} dt$$
(12)

Setting aside Pollard's contour integral proof, it is hard to evaluate directly any of the equivalent integral representations above to demonstrate that they do indeed generate  $E_{\alpha}(-x)$ ,  $E_{\alpha}(-x^{\alpha})$ . A method that may be convenient to prove one representation effectively proves all other representations because they are interchangeable ways of stating the Pollard result. In particular, Feller followed an indirect route to prove the representation (11), discussed next.

#### 1.3 Feller's Method

In an illustration of the use of the two-dimensional Laplace transform, Feller [7](p453) considered  $1 - F_{\alpha}(xt^{-1/\alpha})$  as a bivariate distribution over x > 0, t > 0. The Laplace transform over x, followed by that over t gives

$$\int_0^\infty e^{-sx} (1 - F_\alpha(xt^{-1/\alpha})) \, dx = \frac{1}{s} - \frac{e^{-ts^\alpha}}{s} \tag{13}$$

$$\frac{1}{s} \int_0^\infty e^{-\lambda t} \left( 1 - e^{-ts^\alpha} \right) dt = \frac{1}{\lambda} \frac{s^{\alpha - 1}}{\lambda + s^\alpha} \tag{14}$$

By reference to (4), the right hand side of (14) is the Laplace transform of  $E_{\alpha}(-\lambda x^{\alpha})/\lambda$ . Since the two-dimensional Laplace transform equivalently can be evaluated first over t then over x, Feller concluded that

$$E_{\alpha}(-\lambda x^{\alpha}) = \lambda \int_{0}^{\infty} e^{-\lambda t} (1 - F_{\alpha}(xt^{-1/\alpha})) dt$$
(15)

which, for  $\lambda = 1$ , is the Pollard result in the form (11). Feller's proof is based on the interchange of the order of integration (Fubini's theorem) and the uniqueness of Laplace transforms. We represent it by the commutative diagram below, where  $\mathscr{L}_{s|t}$  denotes the one-dimensional Laplace transform of a bivariate source function at fixed t, to give a bivariate function of (s, t) where s is the Laplace variable.

$$1 - F_{\alpha}(xt^{-1/\alpha}) \xrightarrow{\mathcal{L}_{s|t}} \frac{1}{s} - \frac{e^{-ts^{\alpha}}}{s}$$

$$\mathcal{L}_{\lambda|x} \left|_{\text{hard}} \qquad \mathcal{L}_{\lambda|s} \right|_{\text{easy}} \xrightarrow{\mathcal{L}_{\lambda|s}} \frac{1}{s} - \frac{e^{-ts^{\alpha}}}{s}$$

$$\mathcal{L}_{\lambda|x} \left|_{\text{hard}} \qquad \mathcal{L}_{\lambda|s} \right|_{\text{easy}} \xrightarrow{\mathcal{L}_{\lambda|s}} \frac{1}{\lambda} \frac{1}{\lambda} \frac{s^{\alpha-1}}{\lambda + s^{\alpha}}$$

$$(16)$$

The desired proof is the "hard" direct path, which is equivalent to the "easy" indirect path. We will return to commutative diagram representation in a different context later in the paper when we discuss the main theorem.

Feller's concise proof uses "methods of probability theory", as cited by Pollard, only to the extent of choosing the bivariate distribution as input to the two-dimensional Laplace transform. Short of any further insight, the methods by both Pollard and Feller might be described as more analytic than probabilistic.

#### 1.4 Purpose and Scope of Paper

We assign appropriate distribution guided by the task of proving that  $E_{\alpha}(-x)$  is completely monotone. We first cast Feller's argument in such terms before proceeding to a more general probabilistic discussion.

The Mittag-Leffler function is of growing interest in probability theory and physics, with a diversity of applications, notably fractional calculus. A comprehensive study of the properties and applications of the Mittag-Leffler function and its numerous generalisations is beyond the

scope of this paper. We consciously restrict the scope to the theme of complete monotonicity and Mittag-Leffler functions, underpinned by probability theory.

Other studies that explicitly discuss complete monotonicity and Mittag-Leffler functions build upon complex analytic approaches similar to Pollard's rather than the probabilistic underpinning discussed here (de Oliviera et al. [5], Mainardi and Garrappa [14], Górska et al. [10]). These papers comment on the fundamental importance of the complete monotonicity of Mittag-Leffler functions used in the modelling of physical phenomena, such as anomalous dielectric relaxation and viscoelasticity.

Finally, we are keenly aware that there are other views on the interpretation of "methods of probability theory". We comment on this before discussing our probabilistic approach in detail.

#### **1.5** Probabilistic Perspectives

The phrase 'methods of probability theory' used by Pollard may suggest an experiment with random outcomes as a fundamental metaphor. As noted earlier,  $P_{\alpha}$  is derived as a limiting distribution of a Pólya urn scheme in the probabilistic literature.

Diversity of approach is commonplace in probability theory and mathematics more generally. For example, in a context of nonparametric Bayesian analysis, Ferguson [8] constructed the Dirichlet process based on the gamma distribution as the fundamental probabilistic concept, without invoking a random experiment. Blackwell and MacQueen [3] observed that the Ferguson approach "involves a rather deep study of the gamma process" as they proceeded to give an alternate construction based on the metaphor of a generalised Pólya urn scheme. Adopting the one approach is not to deny or diminish the other, but to bring attention to the diversity of thinking in probability theory, even when the end result is the same mathematical object. We look upon this as healthy complementarity rather than undesirable contestation.

For our purpose, we have no need to invoke an underlying random experiment or indeed an explicit random variable, while not denying the latter as an alternative probabilistic approach. Hence, for example, we shall continue to express the Laplace transform of a distribution as an explicit integral rather than as an expectation  $\mathbb{E}\left[e^{-sX}\right]$  for a random variable X.

### 2 A Probabilistic Method

First, we note that the scale change  $s \to t^{1/\alpha} s$  (t > 0) in (7) gives

$$e^{-ts^{\alpha}} = \int_0^{\infty} e^{-sx} f_{\alpha}(x t^{-1/\alpha}) t^{-1/\alpha} dx \equiv \int_0^{\infty} e^{-sx} f_{\alpha}(x|t) dx$$
(17)

where  $f_{\alpha}(x|t) \equiv f_{\alpha}(x t^{-1/\alpha}) t^{-1/\alpha}$  is the stable density conditioned on the scale parameter t, with  $f_{\alpha}(x) \equiv f_{\alpha}(x|1)$ . Correspondingly, the stable distribution conditioned on t is

$$F_{\alpha}(x|t) = \int_{0}^{x} f_{\alpha}(u|t) \, du = \int_{0}^{xt^{-1/\alpha}} f_{\alpha}(u) \, du \equiv F_{\alpha}(xt^{-1/\alpha}) \tag{18}$$

with Laplace transform  $e^{-ts^{\alpha}}/s$ .

We then assign a distribution G(t) to the scale parameter t of  $F_{\alpha}(x|t)$ . Then, by the sum and product rules of probability theory, the unconditional or marginal distribution  $M_{\alpha}(x)$  over x is

$$M_{\alpha}(x) = \int_{0}^{\infty} F_{\alpha}(x|t) dG(t)$$
(19)

with Laplace transform

$$\int_0^\infty e^{-sx} M_\alpha(x) \, dx = \frac{1}{s} \int_0^\infty e^{-ts^\alpha} \, dG(t) \tag{20}$$

 $M_{\alpha}$  is also referred to as a mixture distribution, arising from randomising or mixing the parameter t in  $F_{\alpha}(x|t)$  with G(t). This has the same import as saying that we assign a prior distribution G(t) on t and we shall continue to use the latter language.

G may depend on one or more parameters. A notable example is the gamma distribution  $G(\mu, \lambda)$  with shape and scale parameters  $\mu > 0, \lambda > 0$  respectively:

$$dG(t|\mu,\lambda) = \frac{\lambda^{\mu}}{\Gamma(\mu)} t^{\mu-1} e^{-\lambda t} dt$$
(21)

 $\lambda$  is not fundamental and may be set to  $\lambda = 1$  by change of scale  $t \to \lambda t$ , while  $\mu$  controls the shape of  $G(t|\mu, \lambda)$ . The marginal (19) becomes  $M_{\alpha}(x|\mu, \lambda)$ , with Laplace transform

$$\int_0^\infty e^{-sx} M_\alpha(x|\mu,\lambda) \, dx = \frac{1}{s} \left(\frac{\lambda}{\lambda+s^\alpha}\right)^\mu = \frac{1}{s} \left(1 - \frac{s^\alpha}{\lambda+s^\alpha}\right)^\mu \tag{22}$$

We may now state Feller's approach from a probabilistic perspective.

#### 2.1 A Probabilistic View of Feller's Approach

The case  $\mu = 1$  in (21) gives the exponential distribution  $dG(t|\lambda) = \lambda e^{-\lambda t} dt$ . Then  $M_{\alpha}(x|\lambda) \equiv M_{\alpha}(x|\mu = 1, \lambda)$  is

$$M_{\alpha}(x|\lambda) = \int_{0}^{\infty} F_{\alpha}(x|t) dG(t|\lambda) = \lambda \int_{0}^{\infty} F_{\alpha}(x|t) e^{-\lambda t} dt$$
(23)

The Laplace transform of  $M_{\alpha}(x|\lambda)$ , read from (22) with  $\mu = 1$ , is

$$\int_0^\infty e^{-sx} M_\alpha(x|\lambda) \, dx = \frac{1}{s} - \frac{s^{\alpha-1}}{\lambda + s^\alpha} \tag{24}$$

$$\implies M_{\alpha}(x|\lambda) = 1 - E_{\alpha}(-\lambda x^{\alpha}) \tag{25}$$

$$\implies E_{\alpha}(-\lambda x^{\alpha}) = 1 - M_{\alpha}(x|\lambda) = \lambda \int_{0}^{\infty} (1 - F_{\alpha}(x|t))e^{-\lambda t} dt$$
(26)

This reproduces Feller's result (15) from a probabilistic perspective. The difference is purely a matter of conceptual outlook:

**Feller:** Study the two-dimensional Laplace transform of the bivariate distribution  $1-F_{\alpha}(xt^{-1/\alpha})$ , where  $F_{\alpha}$  is the stable distribution. Deduce that  $E_{\alpha}(-\lambda x^{\alpha})/\lambda$  is the Laplace transform of  $1 - F_{\alpha}(xt^{-1/\alpha})$  over t at fixed x, where  $\lambda$  is the Laplace variable. **Probabilistic:** Assign an exponential prior distribution  $G(t|1,\lambda)$  to the scale factor t of the stable distribution  $F_{\alpha}(x|t) \equiv F_{\alpha}(xt^{-1/\alpha})$ , where  $G(t|\mu,\lambda)$  is the gamma distribution. Marginalise over t to generate the Feller result directly.

Feller himself might also have established the result by the latter reasoning. Under subordination of processes, Feller [7](p451) discussed mixture distributions but he did not specifically discuss the Mittag-Leffler function in this context in his published work. The task fell on Pillai [15] to study  $M_{\alpha}(x|\mu) \equiv M_{\alpha}(x|\mu, \lambda = 1)$ , including its infinite divisibility and the corresponding Mittag-Leffler stochastic process. He also proved that  $M_{\alpha}(x|1) = 1 - E_{\alpha}(-x^{\alpha})$  (as discussed above), which he referred to as the Mittag-Leffler distribution. There are thus two distributions bearing the name "Mittag-Leffler distribution":  $M_{\alpha}(x) = 1 - E_{\alpha}(-x^{\alpha})$  and  $P_{\alpha}(t) = 1 - F_{\alpha}(t^{-1/\alpha})$ .

The natural question arising from the probabilistic approach is whether there might be other choices of  $\mu$  in  $G(\mu, \lambda)$  (or indeed other choices of G altogether) that yield the Pollard result and, if so, what insight they might offer. At face value, there would appear to be nothing further to be said since other choices of  $\mu$  can be expected to lead to different results, beyond the study of the Mittag-Leffler function. The main contribution of this paper is that, in fact, there is a limit relationship that generates the Pollard result for any  $\mu > 0$ , as discussed next.

We first note, given the definition of the conditional stable density

$$f_{\alpha}(x|t) \equiv f_{\alpha}(x t^{-1/\alpha}) t^{-1/\alpha} \implies f_{\alpha}(1|t) \equiv f_{\alpha}(t^{-1/\alpha}) t^{-1/\alpha}$$

that we may write  $P_{\alpha}(t)$  of (9) and the representation (12) of the Pollard result as

$$P_{\alpha}(t) = \int_{0}^{t} p_{\alpha}(u) \, du = \frac{1}{\alpha} \int_{0}^{t} f_{\alpha}(1|u) \, u^{-1} \, du \tag{27}$$

$$\alpha E_{\alpha}(-\lambda x^{\alpha}) = x \int_{0}^{\infty} f_{\alpha}(x|t) t^{-1} e^{-\lambda t} dt \qquad 0 < \alpha < 1$$
(28)

$$u = x^{-\alpha}t: \quad E_{\alpha}(-\lambda x^{\alpha}) = \int_{0}^{\infty} e^{-\lambda x^{\alpha}u} \, dP_{\alpha}(u) \tag{29}$$

The intent is to generate this representation using the general  $G(\mu, \lambda)$  prior distribution, *i.e.* without reference to Pollard's analytic method and without explicit restriction to the  $G(\mu = 1, \lambda)$  case that is equivalent to Feller's approach, as demonstrated above.

### 3 Main Contribution

We first state Theorem 1, which warrants dedicated discussion, even though it is actually a special case of the more general Theorem 3 stated later. We note first that the density of the marginal distribution  $M_{\alpha}(x|\mu,\lambda)$  of Section 2 is

$$m_{\alpha}(x|\mu,\lambda) = \int_{0}^{\infty} f_{\alpha}(x|t) \, dG(t|\mu,\lambda) \qquad \mu > 0, \lambda > 0$$
$$= \frac{\lambda^{\mu}}{\Gamma(\mu)} \int_{0}^{\infty} f_{\alpha}(x|t) \, t^{\mu-1} e^{-\lambda t} \, dt$$
$$= \frac{\mu \lambda^{\mu}}{\Gamma(\mu+1)} \int_{0}^{\infty} f_{\alpha}(x|t) \, t^{\mu-1} e^{-\lambda t} \, dt$$
(30)

where the latter expression follows from the identity  $\mu\Gamma(\mu) = \Gamma(\mu+1)$ .

Theorem 1. The limit

$$\lim_{n \to \infty} \frac{n}{\mu} x \, m_{\alpha}(x|\frac{\mu}{n}, \lambda) = \lim_{n \to \infty} \frac{n}{\mu} x \int_{0}^{\infty} f_{\alpha}(x|t) \, dG(t|\frac{\mu}{n}, \lambda) \tag{31}$$

is finite and independent of  $\mu$  for any  $\mu > 0$ . This limit yields the following integral representation of the Mittag-Leffler function  $E_{\alpha}(-\lambda x^{\alpha})$ 

$$\alpha E_{\alpha}(-\lambda x^{\alpha}) = x \int_{0}^{\infty} f_{\alpha}(x|t) t^{-1} e^{-\lambda t} dt$$
(32)

$$u = x^{-\alpha}t: \quad E_{\alpha}(-\lambda x^{\alpha}) = \int_{0}^{\infty} e^{-\lambda x^{\alpha}u} \, dP_{\alpha}(u) \tag{33}$$

where  $P_{\alpha}(t)$  is the (one-parameter) Pollard distribution

$$P_{\alpha}(t) = \frac{1}{\alpha} \int_{0}^{t} f_{\alpha}(1|u) u^{-1} du$$
$$= \frac{1}{\alpha} \int_{0}^{t} f_{\alpha}(u^{-1/\alpha}) u^{-1/\alpha - 1} du$$

Hence  $E_{\alpha}(-x)$  is completely monotone.

Proof of Theorem 1. The Laplace transform of  $x m_{\alpha}(x|\mu, \lambda)$  is

$$\int_{0}^{\infty} e^{-sx} x \, m_{\alpha}(x|\mu,\lambda) \, dx = \int_{0}^{\infty} e^{-sx} x \int_{0}^{\infty} f_{\alpha}(x|t) \, dG(t|\mu,\lambda) \, dx$$
$$= -\frac{d}{ds} \int_{0}^{\infty} \int_{0}^{\infty} e^{-sx} f_{\alpha}(x|t) \, dx \, dG(t|\mu,\lambda)$$
$$= -\frac{d}{ds} \int_{0}^{\infty} e^{-ts^{\alpha}} \, dG(t|\mu,\lambda)$$
$$= \alpha s^{\alpha-1} \int_{0}^{\infty} t \, e^{-ts^{\alpha}} \, dG(t|\mu,\lambda)$$
$$= \alpha s^{\alpha-1} \frac{\lambda^{\mu}}{\Gamma(\mu)} \int_{0}^{\infty} t^{\mu} e^{-(\lambda+s^{\alpha})t} \, dt$$
$$= \alpha s^{\alpha-1} \frac{\lambda^{\mu}}{\Gamma(\mu)} \frac{\Gamma(\mu+1)}{(\lambda+s^{\alpha})^{\mu+1}}$$
$$= \lambda^{\mu} \mu \alpha \frac{s^{\alpha-1}}{(\lambda+s^{\alpha})^{\mu+1}}$$
$$\implies \lim_{n \to \infty} \frac{n}{\mu} \int_{0}^{\infty} e^{-sx} x \, m_{\alpha}(x|\frac{\mu}{n},\lambda) \, dx = \alpha \frac{s^{\alpha-1}}{\lambda+s^{\alpha}}$$

which is the Laplace transform of  $\alpha E_{\alpha}(-\lambda x^{\alpha})$ . With the aid of (30), it also readily follows that the limit (31) is

$$\lim_{n \to \infty} \frac{n}{\mu} x \, m_{\alpha}(x|\frac{\mu}{n}, \lambda) = x \int_{0}^{\infty} f_{\alpha}(x|t) \, t^{-1} e^{-\lambda t} \, dt$$

The integral representations (32) and (33) of  $E_{\alpha}(-\lambda x^{\alpha})$  follow, hence the conclusion that  $E_{\alpha}(-x)$  is completely monotone.

Pursuing the probabilistic theme, we turn next to Laplace convolution to demonstrate the complete monotonicity of the two and three parameter Mittag-Leffler functions.

# 4 A Convolution Representation

Toward a more general discussion, we first present an alternative representation of  $x f_{\alpha}(x|t)$  using Laplace convolution. The convolution  $\{\rho \star f\}(x)$  of  $\rho(x), f(x)$  is given by

$$\{\rho \star f\}(x) = \int_0^x \rho(x-u)f(u) \, du$$
 (34)

The convolution theorem states that the Laplace transform of  $\{\rho \star f\}$  is a product of the Laplace transforms of  $\rho, f$ .

### 4.1 One Parameter Case

**Proposition 1.** Let  $\rho_{\alpha}(x) = x^{-\alpha}/\Gamma(1-\alpha)$ ,  $0 < \alpha < 1$  with Laplace transform  $s^{\alpha-1}$ . Let  $\{\rho_{\alpha} \star f_{\alpha}(\cdot|t)\}(x)$  be the convolution of  $\rho_{\alpha}(x)$  and  $f_{\alpha}(x|t)$  with Laplace transform  $s^{\alpha-1}e^{-ts^{\alpha}}$ . Then

$$x f_{\alpha}(x|t) = \alpha t \{ \rho_{\alpha} \star f_{\alpha}(\cdot|t) \}(x) = \alpha \{ \rho_{\alpha} \star f_{\alpha} \}(xt^{-1/\alpha})$$
(35)

where  $\{\rho_{\alpha} \star f_{\alpha}\}(x)$  is the convolution of  $\rho_{\alpha}(x)$  and  $f_{\alpha}(x) \equiv f_{\alpha}(x|1)$ . For compatibility with later discussion, we also use the name  $w_{\alpha}(x|t)$  defined by  $\alpha w_{\alpha}(x|t) \equiv x f_{\alpha}(x|t)$ .

Proof of Proposition 1. By the convolution theorem,  $\{\rho_{\alpha} \star f_{\alpha}(\cdot|t)\}(x)$  has Laplace transform

$$s^{\alpha-1}e^{-ts^{\alpha}} = -\frac{1}{\alpha t}\frac{d}{ds}e^{-ts^{\alpha}} = \frac{1}{\alpha t}\int_{0}^{\infty}e^{-sx}xf_{\alpha}(x|t)\,dx$$
$$\implies \alpha t \{\rho_{\alpha} \star f_{\alpha}(\cdot|t)\}(x) = xf_{\alpha}(x|t)$$

The convolution  $\{\rho_{\alpha} \star f_{\alpha}(\cdot|t)\}(x)$  takes the explicit form:

1

$$\{\rho_{\alpha} \star f_{\alpha}(\cdot|t)\}(x) = \int_{0}^{x} \rho_{\alpha}(x-u) f_{\alpha}(u|t) du$$
  

$$= \int_{0}^{x} \rho_{\alpha}(x-u) f_{\alpha}(ut^{-1/\alpha}) t^{-1/\alpha} du$$
  

$$y = ut^{-1/\alpha} := \int_{0}^{xt^{-1/\alpha}} \rho_{\alpha}(x-yt^{1/\alpha}) f_{\alpha}(y) dy$$
  

$$= \int_{0}^{xt^{-1/\alpha}} \rho_{\alpha}(t^{1/\alpha}(xt^{-1/\alpha}-y)) f_{\alpha}(y) dy$$
  

$$= t^{-1} \int_{0}^{xt^{-1/\alpha}} \rho_{\alpha}(xt^{-1/\alpha}-y) f_{\alpha}(y) dy$$
  

$$= t^{-1} \{\rho_{\alpha} \star f_{\alpha}\}(xt^{-1/\alpha})$$

so that  $\alpha w_{\alpha}(x|t) \equiv x f_{\alpha}(x|t) = \alpha t \{ \rho_{\alpha} \star f_{\alpha}(\cdot|t) \}(x) = \alpha \{ \rho_{\alpha} \star f_{\alpha} \}(xt^{-1/\alpha}).$ 

Hence the following are equivalent representations of the Pollard distribution  $P_{\alpha}(t)$ :

$$P_{\alpha}(t) = \int_{0}^{t} w_{\alpha}(1|t) u^{-1} du \equiv \frac{1}{\alpha} \int_{0}^{t} f_{\alpha}(1|u) u^{-1} du$$
  
= 
$$\int_{0}^{t} \{\rho_{\alpha} \star f_{\alpha}(\cdot|u)\}(1) du$$
  
= 
$$\int_{0}^{t} \{\rho_{\alpha} \star f_{\alpha}\}(u^{-1/\alpha}) u^{-1} du$$
 (36)

The motivation for the convolution representation is to facilitate generalisation. Specifically, the Laplace transform  $\alpha t s^{\alpha-1} e^{-ts^{\alpha}}$  of  $x f_{\alpha}(x|t)$  is the derivative of  $-e^{-ts^{\alpha}}$ . However, a more general term like  $ts^{\alpha-\beta}e^{-ts^{\alpha}}$  cannot arise from simple derivatives of  $e^{-ts^{\alpha}}$  for non-integer  $\beta$ . It might be interpreted as a fractional derivative, as can be represented instead by convolutions. Accordingly, we proceed to consider more general convolutions than the convolution form (35) for  $xf_{\alpha}(x|t)$ .

#### 4.2 Two Parameter Case

First, we introduce the two-parameter Mittag-Leffler function

$$E_{\alpha,\beta}(x) = \sum_{k=0}^{\infty} \frac{x^k}{\Gamma(\alpha k + \beta)}$$
(37)

The Laplace transform of  $x^{\beta-1}E_{\alpha,\beta}(-\lambda x^{\alpha})$  is

$$\int_0^\infty e^{-sx} x^{\beta-1} E_{\alpha,\beta}(-\lambda x^\alpha) \, dx = \frac{s^{\alpha-\beta}}{\lambda+s^\alpha} \tag{38}$$

We may now proceed to prove that  $E_{\alpha,\beta}(-x)$  is completely monotone by showing that it is the Laplace transform of a two-parameter variant  $P_{\alpha,\beta}(t)$  of the Pollard distribution. We follow a corresponding two-parameter variant of the convolution argument presented above for the one-parameter case.

**Proposition 2.** Let  $\rho_{\alpha,\beta}(x) = x^{\beta-\alpha-1}/\Gamma(\beta-\alpha)$   $\beta > \alpha$ , with Laplace transform  $s^{\alpha-\beta}$ . Let  $\{\rho_{\alpha,\beta} \star f_{\alpha}(\cdot|t)\}(x)$  be the convolution of  $\rho_{\alpha,\beta}(x)$  and  $f_{\alpha}(x|t)$ . Then

$$w_{\alpha,\beta}(x|t) \equiv t \left\{ \rho_{\alpha,\beta} \star f_{\alpha}(\cdot|t) \right\}(x) = t^{(\beta-1)/\alpha} \left\{ \rho_{\alpha,\beta} \star f_{\alpha} \right\}(xt^{-1/\alpha})$$
(39)

(the name  $w_{\alpha,\beta}(x|t)$  is a shorthand adopted for convenience).

Proof of Proposition 2.

$$\{\rho_{\alpha,\beta} \star f_{\alpha}(\cdot|t)\}(x) = \int_{0}^{x} \rho_{\alpha,\beta}(x-u) f_{\alpha}(u|t) du$$
$$= \int_{0}^{xt^{-1/\alpha}} \rho_{\alpha,\beta}(t^{1/\alpha}(xt^{-1/\alpha}-u)) f_{\alpha}(u) du$$
$$= t^{(\beta-1)/\alpha-1} \int_{0}^{xt^{-1/\alpha}} \rho_{\alpha,\beta}(xt^{-1/\alpha}-u) f_{\alpha}(u) du$$
$$= t^{(\beta-1)/\alpha-1} \{\rho_{\alpha,\beta} \star f_{\alpha}\}(xt^{-1/\alpha})$$

Thus  $w_{\alpha,\beta}(x|t) \equiv t \{ \rho_{\alpha,\beta} \star f_{\alpha}(\cdot|t) \}(x) = t^{(\beta-1)/\alpha} \{ \rho_{\alpha,\beta} \star f_{\alpha} \}(xt^{-1/\alpha}).$ 

**Theorem 2.** The two-parameter Mittag-Leffler function  $E_{\alpha,\beta}(-\lambda x^{\alpha})$  has the integral representation

$$E_{\alpha,\beta}(-\lambda x^{\alpha}) = \int_0^\infty e^{-\lambda x^{\alpha}t} \, dP_{\alpha,\beta}(t) \tag{40}$$

where  $P_{\alpha,\beta}(t)$ , which we refer to as the two-parameter Pollard distribution, is

$$P_{\alpha,\beta}(t) = \int_0^t w_{\alpha,\beta}(1|u) \, u^{-1} \, du$$
  

$$\equiv \int_0^t \{\rho_{\alpha,\beta} \star f_\alpha(\cdot|u)\}(1) \, du$$
  

$$= \int_0^t \{\rho_{\alpha,\beta} \star f_\alpha\}(u^{-1/\alpha}) \, u^{(\beta-1)/\alpha-1} \, du$$
(41)

Hence  $E_{\alpha,\beta}(-x)$  is completely monotone.

*Proof of Theorem 2.* The theorem is a particular case of the more general Theorem 3 below, hence the current proof is deferred to that of the latter theorem.  $\Box$ 

#### 4.3 Three Parameter Case

The three-parameter Mittag-Leffler function, also known as the Prabhakar function, is given by

$$E_{\alpha,\beta}^{\gamma}(x) = \frac{1}{\Gamma(\gamma)} \sum_{k=0}^{\infty} \frac{\Gamma(\gamma+k)}{k! \,\Gamma(\alpha k+\beta)} \, x^k \tag{42}$$

The Laplace transform of  $x^{\beta-1}E^{\gamma}_{\alpha,\beta}(-\lambda x^{\alpha})$  is

$$\int_0^\infty e^{-sx} x^{\beta-1} E_{\alpha,\beta}^\gamma(-\lambda x^\alpha) \, dx = \frac{s^{\alpha\gamma-\beta}}{(\lambda+s^\alpha)^\gamma} \tag{43}$$

We may now proceed to prove that  $E_{\alpha,\beta}^{\gamma}(-x)$  is completely monotone by showing that it is the Laplace transform of a three-parameter variant  $P_{\alpha,\beta}^{\gamma}(t)$  of the Pollard distribution. In principle, we need only have discussed the three-parameter case from the outset because the two and one-parameter instances are the special cases  $\gamma = 1$  and  $\gamma = \beta = 1$  respectively. We chose instead to present in sequential order for clarity of exposition.

We devote a separate section to the three-parameter case, which subsumes all prior discussion, by restating Theorem 1 in the three-parameter context.

### 5 Main Theorem

We start with a proposition required for the general theorem that follows:

**Proposition 3.** Let  $\rho_{\alpha,\beta}^{\gamma}(x) = x^{\beta-\alpha\gamma-1}/\Gamma(\beta-\alpha\gamma)$   $(0 < \alpha < 1, \gamma > 0, \beta > \alpha\gamma)$  and let  $\{\rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}(\cdot|t)\}(x)$  be the convolution of  $\rho_{\alpha,\beta}^{\gamma}(x)$  and the stable density  $f_{\alpha}(x|t)$ . Then

$$w_{\alpha,\beta}^{\gamma}(x|t) \equiv t^{\gamma} \{\rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}(\cdot|t)\}(x) = t^{(\beta-1)/\alpha} \{\rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}\}(xt^{-1/\alpha})$$
(44)

Proof of Proposition 3.

$$\begin{aligned} \{\rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}(\cdot|t)\}(x) &= \int_{0}^{x} \rho_{\alpha,\beta}^{\gamma}(x-u) f_{\alpha}(u|t) \, du \\ &= \int_{0}^{xt^{-1/\alpha}} \rho_{\alpha,\beta}^{\gamma}(t^{1/\alpha}(xt^{-1/\alpha}-u)) f_{\alpha}(u) \, du \\ &= t^{(\beta-1)/\alpha-\gamma} \int_{0}^{xt^{-1/\alpha}} \rho_{\alpha,\beta}^{\gamma}(xt^{-1/\alpha}-u) f_{\alpha}(u) \, du \\ &= t^{(\beta-1)/\alpha-\gamma} \{\rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}\}(xt^{-1/\alpha}) \end{aligned}$$

Thus  $w_{\alpha,\beta}^{\gamma}(x|t) \equiv t^{\gamma} \{ \rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}(\cdot|t) \}(x) = t^{(\beta-1)/\alpha} \{ \rho_{\alpha,\beta}^{\gamma} \star f_{\alpha} \}(xt^{-1/\alpha}).$ 

**Theorem 3.** Let  $\rho_{\alpha,\beta}^{\gamma}(x)$ ,  $w_{\alpha,\beta}^{\gamma}(x|t)$   $(0 < \alpha < 1, \gamma > 0, \beta > \alpha\gamma)$  be as defined in Proposition 3 and let  $G(\mu, \lambda)$  be the gamma distribution with shape and scale parameters  $\mu > 0, \lambda > 0$  respectively. Let the distribution  $M_{\alpha,\beta}^{\gamma}(x|\mu, \lambda)$  have density

$$m_{\alpha,\beta}^{\gamma}(x|\mu,\lambda) = \int_{0}^{\infty} w_{\alpha,\beta}^{\gamma}(x|t) \, dG(t|\mu,\lambda)$$
$$= \frac{\lambda^{\mu}}{\Gamma(\mu)} \int_{0}^{\infty} w_{\alpha,\beta}^{\gamma}(x|t) \, t^{\mu-1} e^{-\lambda t} \, dt$$
(45)

$$\equiv \frac{\lambda^{\mu}}{\Gamma(\mu)} \int_0^\infty \{\rho^{\gamma}_{\alpha,\beta} \star f_{\alpha}(\cdot|t)\}(x) t^{\gamma+\mu-1} e^{-\lambda t} dt$$
(46)

$$= \frac{\lambda^{\mu}}{\Gamma(\mu)} \int_0^\infty \{\rho_{\alpha,\beta}^\gamma \star f_\alpha\}(xt^{-1/\alpha}) t^{(\beta-1)/\alpha+\mu-1} e^{-\lambda t} dt$$
(47)

where the latter two forms follow from Proposition 3. Then the following limit is finite and independent of  $\mu$  for any  $\mu > 0$ 

$$\lim_{n \to \infty} \frac{n}{\mu} m_{\alpha,\beta}^{\gamma}(x|\frac{\mu}{n},\lambda)$$
(48)

This limit yields the following integral representation of the three-parameter Mittag-Leffler or Prabhakar function  $E^{\gamma}_{\alpha,\beta}(-\lambda x^{\alpha})$ 

$$E^{\gamma}_{\alpha,\beta}(-\lambda x^{\alpha}) = \int_0^\infty w^{\gamma}_{\alpha,\beta}(x|t) t^{-1} e^{-\lambda t} dt = \int_0^\infty e^{-\lambda x^{\alpha} t} dP^{\gamma}_{\alpha,\beta}(t)$$
(49)

where  $P^{\gamma}_{\alpha,\beta}(t)$ , which we refer to as the three-parameter Pollard distribution, is

$$P_{\alpha,\beta}^{\gamma}(t) = \int_{0}^{t} w_{\alpha,\beta}^{\gamma}(1|u) u^{-1} du$$
  
$$\equiv \frac{1}{\Gamma(\gamma)} \int_{0}^{t} \{\rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}(\cdot|u)\}(1) u^{\gamma-1} du$$
  
$$= \frac{1}{\Gamma(\gamma)} \int_{0}^{t} \{\rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}\}(u^{-1/\alpha}) u^{(\beta-1)/\alpha-1} du$$
(50)

Hence  $E_{\alpha,\beta}^{\gamma}(-x)$  is completely monotone.

Proof of Theorem 3. The Laplace transform  $\widetilde{m}_{\alpha,\beta}^{\gamma}(s|\mu,\lambda)$  of (45) is

$$\widetilde{m}_{\alpha,\beta}^{\gamma}(s|\mu,\lambda) \equiv \int_{0}^{\infty} e^{-sx} m_{\alpha,\beta}^{\gamma}(x|\mu,\lambda) dx$$
$$= s^{\alpha\gamma-\beta} \frac{\lambda^{\mu}}{\Gamma(\mu)} \int_{0}^{\infty} t^{\gamma+\mu-1} e^{-(\lambda+s^{\alpha})t} dt$$
$$= \lambda^{\mu} \frac{\Gamma(\gamma+\mu)}{\Gamma(\mu)} \frac{s^{\alpha\gamma-\beta}}{(\lambda+s^{\alpha})^{\gamma+\mu}}$$
(51)

$$\implies \lim_{n \to \infty} \frac{n}{\mu} \int_0^\infty e^{-sx} m_{\alpha,\beta}^\gamma(x|\frac{\mu}{n},\lambda) \, dx = \Gamma(\gamma) \frac{s^{\alpha\gamma-\beta}}{(\lambda+s^\alpha)^\gamma} \tag{52}$$

By (43), the right hand side is the Laplace transform of  $\Gamma(\gamma) x^{\beta-1} E^{\gamma}_{\alpha,\beta}(-\lambda x^{\alpha})$ . Given (46) and (47), it also readily follows that the limit (48) is

$$\begin{split} \int_0^\infty t^\gamma \{\rho_{\alpha,\beta}^\gamma \star f_\alpha(\cdot|t)\}(x) t^{-1} e^{-\lambda t} dt &= \int_0^\infty \{\rho_{\alpha,\beta}^\gamma \star f_\alpha\}(xt^{-1/\alpha}) t^{(\beta-1)/\alpha-1} e^{-\lambda t} dt \\ \implies E_{\alpha,\beta}^\gamma(-\lambda x^\alpha) &= \frac{x^{1-\beta}}{\Gamma(\gamma)} \int_0^\infty \{\rho_{\alpha,\beta}^\gamma \star f_\alpha\}(xt^{-1/\alpha}) t^{(\beta-1)/\alpha-1} e^{-\lambda t} dt \\ u &= x^{-\alpha} t := \frac{1}{\Gamma(\gamma)} \int_0^\infty e^{-\lambda x^\alpha u} \{\rho_{\alpha,\beta}^\gamma \star f_\alpha\}(u^{-1/\alpha}) u^{(\beta-1)/\alpha-1} du \\ &= \int_0^\infty e^{-\lambda x^\alpha u} dP_{\alpha,\beta}^\gamma(u) \end{split}$$

Hence  $E_{\alpha,\beta}^{\gamma}(-x)$  is completely monotone.

Theorem 3 may be visually represented by the following commutative diagram, where  $m_{\alpha,\beta}^{\gamma}(x|\mu,\lambda)$ and its Laplace transform  $\tilde{m}_{\alpha,\beta}^{\gamma}(s|\mu,\lambda)$  are given by (45) and (51) respectively. The equivalence of the two routes from the top left node to the bottom left node induces the integral representation of the Mittag-Leffler function.

The representation (49) of  $E^{\gamma}_{\alpha,\beta}(x)$ , with  $P^{\gamma}_{\alpha,\beta}(t)$  given by (50), is equivalent to equation (2.4) in Górska et al. [10]. The difference is one of approach. This paper offers a fundamentally probabilistic argument, while Górska et al. [10] follows a complex analytic route inspired by Pollard [18]. The balance of Górska et al. [10] is devoted to finding an explicit formula for a function  $f^{\gamma}_{\alpha,\beta}(x)$  featuring in the paper in terms of the Meijer *G* function and associated confluent Wright function. In turns out that  $f^{\gamma}_{\alpha,\beta}(x)$  in Górska et al. [10] is identical to  $\{\rho^{\gamma}_{\alpha,\beta} \star f_{\alpha}\}(x)$  in

this paper. We are content to leave it in the conceptually simple convolution form:

$$\{\rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}\}(x) = \int_{0}^{x} \rho_{\alpha,\beta}^{\gamma}(x-u) f_{\alpha}(u) \, du$$
$$= \frac{1}{\Gamma(\beta - \alpha\gamma)} \int_{0}^{x} (x-u)^{\beta - \alpha\gamma - 1} f_{\alpha}(u) \, du$$
(54)

rather than express it in terms of special functions. In our context, we have actually worked with the conditional density

$$w_{\alpha,\beta}^{\gamma}(x|t) \equiv t^{\gamma} \{ \rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}(\cdot|t) \}(x) = t^{(\beta-1)/\alpha} \{ \rho_{\alpha,\beta}^{\gamma} \star f_{\alpha} \}(xt^{-1/\alpha})$$

where we assigned a gamma prior distribution to the scale parameter t. The density  $w_{\alpha,\beta}^{\gamma}(x|t)$  reduces to (54) for the particular choice t = 1.

We have completed the task of proving that the three-parameter Mittag-Leffler function  $E_{\alpha,\beta}^{\gamma}(-x)$  is completely monotone by methods of probability theory, using probabilistic reasoning to derive an explicit form for  $P_{\alpha,\beta}^{\gamma}(t)$ , whose Laplace transform is  $E_{\alpha,\beta}^{\gamma}(-x)$ . Beyond that, we draw conclusions on the complete monotonicity of related functions, notably  $x^{\beta-1}E_{\alpha,\beta}^{\gamma}(-x^{\alpha})$  and  $E_{\alpha,\beta}^{\gamma}(-x^{\alpha})$  in isolation. First, we discuss  $x^{\beta-1}E_{\alpha,\beta}^{\gamma}(-x^{\alpha})$ , the bottom left node of the commutative diagram (53), in the probabilistic context of Theorem 3. The discussion involves an alternative representation of the fundamental probabilistic object – the convolution density  $\{\rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}(\cdot|t)\}(x)$ .

### 6 An Alternative Representation

For  $x^{\beta-1}E^{\gamma}_{\alpha,\beta}(-\lambda x^{\alpha})$  to be completely monotone, there must exist a distribution  $R^{\gamma}_{\alpha,\beta}(u|\lambda)$  defined by the Laplace transform

$$x^{\beta-1}E^{\gamma}_{\alpha,\beta}(-\lambda x^{\alpha}) = \int_0^\infty e^{-xu} \, dR^{\gamma}_{\alpha,\beta}(u|\lambda) \tag{55}$$

In turn, the Laplace transform of (55) is the Stieltjes transform (or iterated Laplace transform) of  $R^{\gamma}_{\alpha,\beta}(u|\lambda)$ :

$$\frac{s^{\alpha\gamma-\beta}}{(\lambda+s^{\alpha})^{\gamma}} = \int_0^\infty \frac{1}{s+u} \, dR^{\gamma}_{\alpha,\beta}(u|\lambda) \tag{56}$$

Then, as de Oliviera et al. [5], Mainardi and Garrappa [14] show, the Stieltjes inversion formula (Titchmarsh [22](11.8, p318), Widder [23](VIII.7, p342)) gives

$$dR^{\gamma}_{\alpha,\beta}(u|\lambda) = \frac{1}{\pi} \operatorname{Im} \left\{ \frac{(e^{-i\pi}u)^{\alpha\gamma-\beta}}{(\lambda+(e^{-i\pi}u)^{\alpha})^{\gamma}} \right\} du$$
(57)

The expression in braces on the RHS of (57) is (56) at  $s = e^{-i\pi}u$ . In particular, for  $\gamma = \beta = 1$ , (57) reduces to

$$dR_{\alpha}(u|\lambda) = \frac{1}{\pi} \frac{\lambda u^{\alpha-1} \sin \pi \alpha}{\lambda^2 + 2\lambda u^{\alpha} \cos \pi \alpha + u^{2\alpha}} du$$
(58)

which has been discussed in various contexts in the fractional calculus and probabilistic literature (e.g. James [12] in the latter context).

We have mentioned (55) for completeness but it was not the core of our probabilistic discussion, whose focus was to determine  $P^{\gamma}_{\alpha,\beta}(t)$ , with Laplace transform  $E^{\gamma}_{\alpha,\beta}(-x)$ . That said, we can offer a 'hybrid' derivation of (55) that combines the core of the probabilistic argument in the form of the convolution density  $\{\rho^{\gamma}_{\alpha,\beta} \star f_{\alpha}(\cdot|t)\}(x)$  with the complex analytic Stieltjes inversion argument presented above.

Assume  $\{\rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}(\cdot|t)\}(x)$  to be the Laplace transform of a distribution  $S_{\alpha,\beta}^{\gamma}(u|t)$ :

$$\{\rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}(\cdot|t)\}(x) = \int_{0}^{\infty} e^{-xu} \, dS_{\alpha,\beta}^{\gamma}(u|t) \tag{59}$$

In turn, the Laplace transform of (59) is the Stieltjes transform of  $S^{\gamma}_{\alpha,\beta}(u|t)$ :

$$s^{\alpha\gamma-\beta}e^{-ts^{\alpha}} = \int_0^\infty \frac{1}{s+u} \, dS^{\gamma}_{\alpha,\beta}(u|t) \tag{60}$$

By the Stieltjes inversion formula:

$$dS^{\gamma}_{\alpha,\beta}(u|t) = \frac{1}{\pi} \operatorname{Im} \left\{ (ue^{-i\pi})^{\alpha\gamma-\beta} e^{-t(ue^{-i\pi})^{\alpha}} \right\} \, du \tag{61}$$

Hence, using the representation (59) in the proof of Theorem 3:

$$\Gamma(\gamma) x^{\beta-1} E_{\alpha,\beta}^{\gamma}(-\lambda x^{\alpha}) = \int_{0}^{\infty} t^{\gamma} \{\rho_{\alpha,\beta}^{\gamma} \star f_{\alpha}(\cdot|t)\}(x) t^{-1} e^{-\lambda t} dt$$

$$= \int_{0}^{\infty} dt t^{\gamma-1} e^{-\lambda t} \int_{0}^{\infty} e^{-xu} dS_{\alpha,\beta}^{\gamma}(u|t)$$

$$= \frac{1}{\pi} \operatorname{Im} \int_{0}^{\infty} du e^{-xu} (u e^{-i\pi})^{\alpha\gamma-\beta} \int_{0}^{\infty} t^{\gamma-1} e^{-(\lambda+(u e^{-i\pi})^{\alpha})t} dt$$

$$= \frac{\Gamma(\gamma)}{\pi} \operatorname{Im} \int_{0}^{\infty} e^{-xu} \frac{(e^{-i\pi}u)^{\alpha\gamma-\beta}}{(\lambda+(e^{-i\pi}u)^{\alpha})^{\gamma}} du$$

$$= \Gamma(\gamma) \int_{0}^{\infty} e^{-xu} dR_{\alpha,\beta}^{\gamma}(u|\lambda)$$
(62)

thereby reproducing (55).

The Stieltjes transform and its complex analytic inverse are not unfamiliar in probability theory. In his study of a family of distributions known as generalised gamma convolutions, Bondesson [4] used the concept under the guise of Pick functions (also known as Nevanlinna functions).

We turn next to the complete monotonicity of  $E_{\alpha,\beta}^{\gamma}(-\lambda x^{\alpha})$ .

## 7 A Further Consequence

There is a well-known property of completely monotone functions (e.g. Schilling et al. [20]) that we state without proof in Proposition 4. We start with a definition:

**Definition 1.** A Bernstein function is a nonnegative function  $\eta(x)$ ,  $x \ge 0$  with a completely monotone derivative, i.e.  $\eta(x) \ge 0$  and  $(-1)^{k-1}\eta^{(k)}(x) \ge 0$ ,  $k \ge 1$ . For example,  $\eta(x|\lambda) = \lambda x^{\alpha}$   $(0 \le \alpha \le 1, \lambda > 0)$  is a Bernstein function.

**Proposition 4.** If  $\varphi(x)$  is completely monotone and  $\eta$  is a Bernstein function,  $\varphi(\eta)$  is completely monotone.

**Theorem 4.** Given a Bernstein function  $\eta$ , the Mittag-Leffler function  $E^{\gamma}_{\alpha,\beta}(-\eta)$  is completely monotone. For example,  $E^{\gamma}_{\alpha,\beta}(-\lambda x^{\alpha})$  is completely monotone.

Proof of Theorem 4. We have already shown that  $E_{\alpha,\beta}^{\gamma}(-x)$  is completely monotone. Hence, by Proposition 4,  $E_{\alpha,\beta}^{\gamma}(-\eta)$  is completely monotone for a Bernstein function  $\eta$ . Specifically,  $\eta(x|\lambda) = \lambda x^{\alpha} \ (0 \le \alpha \le 1, \lambda > 0)$  is a Bernstein function, hence  $E_{\alpha,\beta}^{\gamma}(-\lambda x^{\alpha})$  is completely monotone.

The complete monotonicity of  $E_{\alpha,\beta}^{\gamma}(-\lambda x^{\alpha})$  implies that there exists a distribution  $Q_{\alpha,\beta}^{\gamma}(t|\lambda)$  whose Laplace transform is  $E_{\alpha,\beta}^{\gamma}(-\lambda x^{\alpha})$ :

$$E^{\gamma}_{\alpha,\beta}(-\lambda x^{\alpha}) = \int_0^\infty e^{-xt} \, dQ^{\gamma}_{\alpha,\beta}(t|\lambda) \tag{63}$$

 $Q_{\alpha,\beta}^{\gamma}(t|\lambda)$  is to  $E_{\alpha,\beta}^{\gamma}(-\lambda x^{\alpha})$  what  $P_{\alpha,\beta}^{\gamma}(t)$  is to  $E_{\alpha,\beta}^{\gamma}(-x)$ . However, determining  $Q_{\alpha,\beta}^{\gamma}(t|\lambda)$  appears to be a challenging problem, whether the approach is analytic or probabilistic.

Clearly, (63) and (57) are identical for  $\beta = 1$ , *i.e.*  $Q_{\alpha,1}^{\gamma}(t|\lambda) \equiv R_{\alpha,1}^{\gamma}(t|\lambda)$ . But, to our awareness, determining  $Q_{\alpha,\beta}^{\gamma}(t|\lambda)$  for  $\beta \neq 1$  is an open problem. We shall not pursue it further here. Our primary purpose in this section was to bring attention to Theorem 4 and hence the existence of a distribution  $Q_{\alpha,\beta}^{\gamma}(t|\lambda)$  defined by (63).

### 8 A Different Generalisation

As mentioned in Section 1.5, the Pollard distribution  $P_{\alpha}$  is known as the Mittag-Leffler distribution in probabilistic literature. For completeness, we briefly discuss a different generalisation of  $P_{\alpha}$  that features extensively in such literature. It is known as the generalised Mittag-Leffler distribution  $P_{\alpha,\theta}$  (Pitman [16], p70 (3.27)), also denoted by ML( $\alpha, \theta$ ) (Goldschmidt and Haas [9], Ho et al. [11]).

Despite its name,  $P_{\alpha,\theta}(t)$  is different from the two-parameter Pollard distribution  $P_{\alpha,\beta}(t)$  discussed above, whose Laplace transform is the Mittag-Leffler function  $E_{\alpha,\beta}(-x)$ . Janson [13] showed that  $P_{\alpha,\theta}$  may be constructed as a limiting distribution of a Pólya urn scheme. It is also intimately linked to a concept known as 'polynomial tilting'. For some parameter  $\theta$ ,  $f_{\alpha,\theta}(x) \propto x^{-\theta} f_{\alpha}(x)$  is said to be a polynomially tilted variant of  $f_{\alpha}(x)$  (e.g. Arbel et al. [1], Devroye [6], James [12]). Here, we consider the polynomially tilted density  $f_{\alpha,\theta}(x|t) \propto x^{-\theta} f_{\alpha}(x|t)$  conditioned on a scale factor t > 0. Normalisation gives

$$f_{\alpha,\theta}(x|t) = \frac{\Gamma(\theta+1)}{\Gamma(\theta/\alpha+1)} t^{\theta/\alpha} x^{-\theta} f_{\alpha}(x|t)$$
(64)

so that  $f_{\alpha,\theta}(x|t)$  is defined for  $\theta/\alpha + 1 > 0$ , or  $\theta > -\alpha$ . We then consider a two-parameter

function  $h_{\alpha,\theta}(x|\lambda)$  defined by:

$$\alpha h_{\alpha,\theta}(x|\lambda) = x \int_0^\infty f_{\alpha,\theta}(x|t) t^{-1} e^{-\lambda t} dt$$

$$= \frac{\Gamma(\theta+1)}{\Gamma(\theta/\alpha+1)} x^{1-\theta} \int_0^\infty f_\alpha(x|t) t^{\theta/\alpha-1} e^{-\lambda t} dt$$
(65)

$$u = x^{-\alpha}t: \quad h_{\alpha,\theta}(x|\lambda) = \int_0^\infty e^{-\lambda x^\alpha u} \, dP_{\alpha,\theta}(u) \tag{66}$$

where 
$$P_{\alpha,\theta}(t) = \frac{\Gamma(\theta+1)}{\Gamma(\theta/\alpha+1)} \frac{1}{\alpha} \int_0^t f_\alpha(u^{-1/\alpha}) u^{(\theta-1)/\alpha-1} du$$
 (67)

or 
$$dP_{\alpha,\theta}(t) = \frac{\Gamma(\theta+1)}{\Gamma(\theta/\alpha+1)} t^{\theta/\alpha} dP_{\alpha}(t)$$
 (68)

It is clear from (66) that  $h_{\alpha,\theta}(x|\lambda)$  may be written as  $h_{\alpha,\theta}(\lambda x^{\alpha})$ . It follows that:

- 1.  $h_{\alpha,\theta}(x)$  is completely monotone
- 2.  $\theta = 0$ :  $P_{\alpha,0}(t) = P_{\alpha}(t) \implies h_{\alpha,0}(x) = E_{\alpha}(-x)$ , as directly apparent from comparing (32) and (65).
- 3.  $h_{\alpha,\theta}(\eta)$  is completely monotone where  $\eta$  is a Bernstein function as discussed in Section 7. In particular,  $h_{\alpha,\theta}(\lambda x^{\alpha})$  is completely monotone and thus expressible as the Laplace transform of a corresponding distribution  $Q_{\alpha,\theta}(t|\lambda)$  (distinct from  $Q_{\alpha,\beta}(t|\lambda)$  discussed in Section 7).

We are not aware of a representation of  $h_{\alpha,\theta}$  other than that generated by  $P_{\alpha,\theta}$  in (66). By comparison, the two-parameter Mittag-Leffler function  $E_{\alpha,\beta}$  has a well-established infinite series representation (37), in addition to the representation (40) generated by the two-parameter Pollard distribution  $P_{\alpha,\beta}$ .

### 9 Discussion

The integral representation (49) of  $E^{\gamma}_{\alpha,\beta}(-\lambda x^{\alpha})$  in Theorem 3, arising from the limit (48), contains the Lévy measure  $t^{-1}e^{-\lambda t}dt$  of the infinitely divisible gamma distribution. There is indeed an intimate relationship between completely monotone functions and the theory of infinitely divisible distributions on the nonnegative half-line  $\mathbb{R}_{+} = [0, \infty)$  (Feller [7] (XIII.4, XIII.7), Steutel and van Harn [21] (III)). Sato [19] considers infinitely divisible distributions on  $\mathbb{R}^{d}$ , but the deliberate restriction to  $\mathbb{R}_{+}$  makes for simpler discussion and relates directly to the core concept of complete monotonicity that is of interest here. There is also an intimate link to the generalised gamma convolutions studied by Bondesson [4].

The limit (48) of Theorem 3 is an instance of a limit rule to generate the Lévy measure of an infinitely divisible distribution given in Steutel and van Harn [21] (III(4.7)) and Sato [19] (Corollary 8.9 restricted to  $\mathbb{R}_+$  rather than  $\mathbb{R}^d$ ). Barndorff-Nielsen and Hubalek [2] also cite Sato's Corollary.

Further exploration using the probabilistic machinery of this paper possibly includes the explicit determination of the three-parameter distribution  $Q^{\gamma}_{\alpha,\beta}(t|\lambda)$ , whose Laplace transform is  $E^{\gamma}_{\alpha,\beta}(-\lambda x^{\alpha})$ , as per (63).

# 10 Conclusion

We have presented a probabilistic derivation of the complete monotonicity of the three-parameter Mittag-Leffler function (also known as the Prabhakar function) by expressing it as the Laplace transform of a distribution that we referred to as the three-parameter Pollard distribution. This is a generalisation of a result due to Pollard for the one-parameter case.

### References

- Julyan Arbel, Pierpaolo De Blasi, and Igor Prünster. Stochastic Approximations to the Pitman–Yor Process. *Bayesian Analysis*, 14(4):1201 – 1219, 2019.
- [2] Ole E. Barndorff-Nielsen and Friedrich Hubalek. Probability measures, Lévy measures and analyticity in time. *Bernoulli*, 14(3):764 – 790, 2008.
- [3] David Blackwell and James B. MacQueen. Ferguson distributions via Pólya urn schemes. The Annals of Statistics, 1(2):353–355, 1973.
- [4] Lennart Bondesson. Generalized Gamma Convolutions and Related Classes of Distributions and Densities. Lecture Notes in Statistics, 76. Springer-Verlag, New York, 1992.
- [5] E. Capelas de Oliveira, F. Mainardi, and J. Vaz. Models based on Mittag-Leffler functions for anomalous relaxation in dielectrics. *The European Physical Journal Special Topics*, 193(1):161–171, Mar 2011.
- [6] Luc Devroye. Random variate generation for exponentially and polynomially tilted stable distributions. ACM Trans. Model. Comput. Simul., 19(4), nov 2009.
- [7] William Feller. An Introduction to Probability Theory and its Applications, Vol. II. Wiley, New York, 1971.
- [8] Thomas S. Ferguson. A Bayesian analysis of some nonparametric problems. The Annals of Statistics, 1(2):209–230, 1973.
- [9] Christina Goldschmidt and Bénédicte Haas. A line-breaking construction of the stable trees. *Electronic Journal of Probability*, 20:1 24, 2015.
- [10] K. Górska, Andrzej Horzela, Ambra Lattanzi, and Tibor Pogány. On complete monotonicity of three parameter Mittag-Leffler function. *Applicable Analysis and Discrete Mathematics*, 15:118–128, 04 2021.
- [11] Man-Wai Ho, Lancelot F. James, and John W. Lau. Gibbs partitions, Riemann-Liouville fractional operators, Mittag-Leffler functions, and fragmentations derived from stable subordinators. *Journal of Applied Probability*, 58(2):314–334, 2021.
- [12] Lancelot F. James. Lamperti-type laws. Ann. Appl. Probab., 20(4):1303–1340, 2010.
- [13] Svante Janson. Limit theorems for triangular urn schemes. Probability Theory and Related Fields, 134(3):417–452, Mar 2006.
- [14] Francesco Mainardi and Roberto Garrappa. On complete monotonicity of the Prabhakar function and non-Debye relaxation in dielectrics. *Journal of Computational Physics*, 293:70– 80, 2015. Fractional PDEs.

- [15] R. N. Pillai. On Mittag-Leffler functions and related distributions. Annals of the Institute of Statistical Mathematics, 42(1):157–161, Mar 1990.
- [16] J. Pitman. Combinatorial Stochastic Processes, volume 1875 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2006. Lectures from the 32nd Summer School on Probability Theory held in Saint-Flour, July 7–24, 2002, With a foreword by Jean Picard.
- [17] Harry Pollard. The representation of  $e^{-x^{\lambda}}$  as a Laplace integral. Bulletin of the American Mathematical Society, 52(10):908 910, 1946.
- [18] Harry Pollard. The completely monotonic character of the Mittag-Leffler function  $E_a(-x)$ . Bulletin of the American Mathematical Society, 54(12):1115 – 1116, 1948.
- [19] K. Sato. Lévy Processes and Infinitely Divisible Distributions. Cambridge University Press, Cambridge, 1999.
- [20] René L. Schilling, Renming Song, and Zoran Vondracek. *Bernstein Functions: Theory and Applications*. De Gruyter, Berlin, Boston, 2012.
- [21] F.W. Steutel and K. van Harn. Infinite Divisibility of Probability Distributions on the Real Line. Marcel Dekker, New York, 2003.
- [22] E.C. Titchmarsh. Introduction to the Theory of Fourier Integrals. Clarendon Press, 1948.
- [23] David Vernon Widder. Laplace transform (PMS-6). Princeton university press, 2015.