

Optimal Scaling Results for a Wide Class of Proximal MALA Algorithms

Francesca R. Crucinio^{*1}, Alain Durmus², Pablo Jiménez³, and Gareth O. Roberts⁴

¹CREST, ENSAE Paris

²Centre de Mathématiques Appliquées, Ecole Polytechnique, France, Institut Polytechnique de Paris

³Sorbonne Université and Université Paris Cité, CNRS, Laboratoire de Probabilités, Statistique et Modélisation, F-75005 Paris, France

⁴Department of Statistics, University of Warwick

Abstract

We consider a recently proposed class of MCMC methods which uses proximity maps instead of gradients to build proposal mechanisms which can be employed for both differentiable and non-differentiable targets. These methods have been shown to be stable for a wide class of targets, making them a valuable alternative to Metropolis-adjusted Langevin algorithms (MALA); and have found wide application in imaging contexts. The wider stability properties are obtained by building the Moreau-Yoshida envelope for the target of interest, which depends on a parameter λ . In this work, we investigate the optimal scaling problem for this class of algorithms, which encompasses MALA, and provide practical guidelines for the implementation of these methods.

Contents

1	Introduction	2
2	Proximal MALA Algorithms	5
3	Optimal scaling of Proximal MALA	7
3.1	Differentiable targets	8
3.2	Laplace target	10
4	Practical Implications and Numerical Simulations	13
4.1	Numerical Experiments	14
5	Discussion	16

^{*}Corresponding author: francesca.crucinio@gmail.com

6	Proof of the Result for the Laplace distribution	17
6.1	Proof of Theorem 2	18
6.2	Proof of Proposition 1	19
6.3	Proof of Proposition 2	27
6.4	Proof of Theorem 3	29
A	Proof of Theorem 1	40
A.1	Auxiliary Results for the Proof of Case (a)	41
A.2	Auxiliary Results for the Proof of Case (b)	49
A.3	Auxiliary Results for the Proof of Case (c)	52
A.4	Proof of Theorem 1	55
B	Numerical Experiments	55
B.1	Differentiable Targets	55
B.2	Laplace Target	56
B.3	Mix of a Laplace and differentiable target	56
C	Taylor Expansions for the Results on Differentiable Targets	64
C.1	Coefficients of the Taylor Expansion	64
C.1.1	Case (a)	64
C.1.2	Case (b)	67
C.1.3	Case (c)	68
C.2	Taylor Expansions of the Log-acceptance Ratio	70
C.2.1	R_1	70
C.2.2	R_2	71
C.3	Derivatives of the Proximity Map for Differentiable Targets	75
D	Moments and Integrals for the Laplace Distribution	77
D.1	Moments of Acceptance Ratio for the Laplace Distribution	77
D.2	Bound on Second Moment of Acceptance Ratio for the Laplace Distribution	83
D.3	Additional Integrals for the Laplace Distribution	87
D.4	Integrals for Moment Computations	88
D.4.1	First Moment	88
D.4.2	Second Moment	91
D.4.3	Third Moment	92

1 Introduction

Gradient-based Markov chain Monte Carlo (MCMC) methods have proved to be very successful at sampling from high-dimensional target distributions [9]. The key to their success is that in many cases their mixing time appears to scale better in dimension than competitor algorithms which do not use gradient information (see for example [34]), while their implementation has similar computational cost. Indeed, gradients of target densities can often be computed with computational complexity (in dimension d) which scales no worse than evaluation of the target density itself.

Gradient-based MCMC methods are mainly motivated from stochastic processes constructed to have the target density as limiting distribution [25, 8, 6, 44]. Our analysis will concentrate on the

Metropolis Adjusted Langevin Algorithm (MALA) and its proximal variants which are based on the Langevin diffusion

$$d\mathbf{L}_t = d\mathbf{B}_t + \frac{\nabla \log \pi(\mathbf{L}_t)}{2} dt, \quad (1)$$

where π denotes the target density with respect to the Lebesgue measure and $(\mathbf{B}_t)_{t \geq 0}$ a standard Brownian motion. It is well-known that under appropriate conditions, (1) defines a continuous-time Markov process associated with a Markov semigroup which is reversible with respect to π . From this observation, it has been suggested to use a Euler-Maruyama (EM) approximation of (1). This scheme has been popularized in statistics by [20] and referred to as the *Unadjusted Langevin Algorithm (ULA)* in [36]. Due to time-discretization, ULA typically does not have π as stationary distribution. To address this problem, [39] and independently Besag in his contribution to [20] proposed to add a Metropolis acceptance step at each iteration of the EM scheme, leading to the Metropolis Adjusted Langevin Algorithm (MALA) following [36] who also derive basic stability analysis. The accept/reject step in this algorithm confers two significant advantages: it ensures that the resulting algorithm has exactly the correct invariant distribution, while step sizes can be chosen larger than in the unadjusted case as there is not need to make the step size small to reduce discretization error. On the other hand, MALA algorithms are typically hard to analyze theoretically (see e.g. [7, 13, 16]). However, [34] (see also [5, 32]) have established that MALA has better convergence properties than the Random Walk Metropolis (RWM) algorithm with respect to the dimension d from an optimal scaling perspective (see also [33]).

Whereas gradient-based methods have been successively applied and offer interesting features, they are typically less robust than their vanilla alternatives (for example see [36]); while intuition suggests, and existing underpinning theory requires, that target densities need to be sufficiently smooth for the gradients to be aiding Markov chain convergence. Moreover, while gradient-based MCMC have been successful for smooth densities, there is no reason to believe that they should be effective for densities which are not differentiable at a subset $D \subseteq \mathbb{R}^d$. For non-smooth densities, [30] proposes modified gradient-based algorithms. Their proposed P-MALA algorithm is inspired by the proximal algorithms popular in the optimization literature (e.g. [29]). The main idea is to approximate the (possibly non differentiable but) log-concave target density $\pi \propto \exp(-G)$ by substituting the potential G with its Moreau-Yoshida envelope G^λ (see (3) below for its definition), to obtain a distribution π^λ whose level of smoothness is controlled by the proximal parameter $\lambda > 0$, so that $G^0 = G$. Given this smooth approximation to π one can then build proposals based on time discretizations of the Langevin diffusion targeting π^λ [30, 14]:

$$\xi_{k+1} = \xi_k - \frac{\sigma^2}{2} \nabla G^\lambda(\xi_k) + \sigma Z_{k+1}, \quad (2)$$

where $\sigma^2 > 0$ is a fixed stepsize and $(Z_k)_{k \in \mathbb{N}^*}$ is a sequence of i.i.d. zero-mean Gaussian random variables with identity covariance matrix. Our aims in this paper are broadly to provide theoretical underpinning for a slightly larger family of *proximal MALA* algorithms, analyze how these methods scale with dimension, and to give insights and practical guidance into how they should be implemented supported by the theory we establish.

Proximal optimization and MCMC methods proved to be particularly well-suited for image estimation, where penalties involving the sparsity inducing norms are common [30, 14, 43]. Similar targets are also common in sparse regression contexts [2, 19, 46]. In these situations, the set of non-differentiability points for the target density π is a null set under Lebesgue measure, and, following

[12], we shall focus on this case. However, in contrast to the conclusions of [12] for RWM, we shall demonstrate that optimal scaling of proximal MALA is significantly affected by non-smoothness.

In this work, we first extend the results of [31], considering a wider range of proximal MALA algorithms, as well as a more general class of finite dimensional target distributions. We begin by comparing MALA and its proximal cousin in cases where MALA is well-defined, ie where target densities are sufficiently differentiable. In some cases the proximal operator for a given distribution π is less expensive to compute than $\nabla \log \pi$ [29, 11, 30], so we anticipate that proximal MALA with an appropriately tuned λ might provide a computationally more efficient alternative to MALA, whilst retaining similar scaling properties. In our study, we let both the steps size σ^2 and the regularization parameter λ depend on the dimension d of the target and find that the scaling properties of proximal MALA depend on the relative speed at which λ and σ converge to 0 as $d \rightarrow \infty$. When λ goes to 0 at least as fast as σ^2 , we find that the scaling properties of proximal MALA are equivalent to those of MALA (i.e. σ^2 should decay as $d^{-1/3}$; see Theorem 1–(b), Theorem 1–(c)); when λ converges to 0 more slowly than σ^2 , proximal MALA is less efficient than MALA with σ^2 decaying as $d^{-1/2}$ (Theorem 1–(a)).

We then turn to the optimal scaling of proximal MALA applied to the Laplace distribution $\pi(x) \propto e^{-|x|}$. We focus on this particular non-smooth target since it is the most widely used in applications of proximal MALA, including image deconvolution [30, 14, 43], LASSO, and sparse regression [2, 19, 46]. We establish that non-differentiability of the target even at one point leads to a different optimal scaling than MALA. In particular, the step size has to scale as $d^{-2/3}$ and not as $d^{-1/3}$ (Theorem 2). We thus uncover a new optimal scaling scenario for Metropolis MCMC algorithms which lies in between those of RWM and MALA.

The proof of the result for the differentiable case extends that of [34] for MALA, while the structure of the proof for the Laplace target is similar to that of [12] and constitutes the main element of novelty in this paper. As a special case of the result for the Laplace distribution, we also obtain the optimal scaling for MALA on Laplace targets. We point out that the strategy adopted in the proof of this result is not unique to the Laplace distribution, and could be applied to other distributions provided that the required integrals can be obtained.

To sum up, our main contributions are:

- 1) We extend the result of [31] beyond the Gaussian case, covering all finite dimensional (sufficiently) differentiable targets, and show that, in some cases, proximal MALA affords the same scaling properties of MALA if the proximal parameter λ is chosen appropriately.
- 2) Motivated by applications in imaging and sparse regression applications, we study the scaling of proximal MALA methods for the Laplace target, and show that for values of λ decaying sufficiently fast, the optimal scaling of proximal MALA, i.e. the choice for σ^2 , is different from the one for MALA on differentiable targets and is of order $d^{-2/3}$.
- 3) We use the insights obtained with the aforementioned results to provide practical guidelines for the selection of the proximal parameter λ .

Organization of the paper The paper is structured as follows. In Section 2, we rigorously introduce the class of proximal MALA algorithms that are studied and discuss related works on optimal scaling for MCMC algorithms. In Section 3.1 we state the main result for differentiable targets, showing that the scaling properties of proximal MALA depend on the relative speed at which λ goes to 0 with respect to σ . In Section 3.2 we obtain a scaling limit for proximal MALA when π is a Laplace distribution, as a special case of our result we also obtain the scaling properties of a sub-gradient version of MALA for this target. We collect in Section 4 the main practical

takeaways from these results and discuss possible extensions in Section 5. Finally, in Section 6 we prove the result for the Laplace distribution. The proof of the result for differentiable targets is postponed to Appendix A.

2 Proximal MALA Algorithms

We now introduce the general class of proximal MALA algorithms, first studied in [30]. This class of algorithms aims at sampling from a density with respect to the Lebesgue measure on $\mathbb{R}^d$ of the form $\pi(\mathbf{x}) = \exp(-G(\mathbf{x})) / \int_{\mathbb{R}^d} \exp(-G(\tilde{\mathbf{x}})) d\tilde{\mathbf{x}}$, with G satisfying the following assumption

A0. The function $G : \mathbb{R}^d \rightarrow \mathbb{R}$ is convex, proper and lower semi-continuous.

The main idea behind proximal MALA is to approximate the (possibly non differentiable) target density π by approximating the potential G with its Moreau-Yoshida envelope $G^\lambda : \mathbb{R}^d \rightarrow \mathbb{R}$ defined for $\lambda > 0$ by

$$G^\lambda(\mathbf{x}) = \min_{\mathbf{u} \in \mathbb{R}^d} [G(\mathbf{u}) + \|\mathbf{u} - \mathbf{x}\|^2 / (2\lambda)] . \quad (3)$$

Since G is supposed to be convex, by [38, Theorem 2.26], the Moreau-Yoshida envelope is well-defined, convex and continuously differentiable with

$$\nabla G^\lambda(\mathbf{x}) = \lambda^{-1}(\mathbf{x} - \text{prox}_G^\lambda(\mathbf{x})) , \quad \text{prox}_G^\lambda(\mathbf{x}) = \arg \min_{\mathbf{u} \in \mathbb{R}^d} [G(\mathbf{u}) + \|\mathbf{u} - \mathbf{x}\|^2 / (2\lambda)] . \quad (4)$$

The proximity operator $\mathbf{x} \mapsto \text{prox}_G^\lambda(\mathbf{x})$ behaves similarly to a gradient mapping and moves points in the direction of the minimizers of G . In the limit $\lambda \rightarrow 0$ the quadratic penalty dominates (4) and the proximity operator coincides with the identity operator, i.e. $\text{prox}_G^\lambda(\mathbf{x}) = \mathbf{x}$; in the limit $\lambda \rightarrow \infty$, the quadratic penalty term vanishes and (4) maps all points to the set of minimizers of G .

It was shown in [14, Proposition 1] that, under **A0**, $\int_{\mathbb{R}^d} \exp(-G^\lambda(\mathbf{x})) d\mathbf{x} < \infty$, and therefore the probability density $\pi^\lambda \propto \exp(-G^\lambda)$ is well-defined. In addition, it has been shown that $\|\pi - \pi^\lambda\|_{\text{TV}} \rightarrow 0$ as $\lambda \rightarrow 0$. Based on this observation and since as we have emphasized π^λ is now continuously differentiable, it has been suggested in [30, 14] to use the discretization of the Langevin diffusion associated with π^λ given by (2), which can be rewritten using (4) as

$$\xi_{k+1} = \left(1 - \frac{\sigma^2}{2\lambda}\right) \xi_k + \frac{\sigma^2}{2\lambda} \text{prox}_G^\lambda(\xi_k) + \sigma Z_{k+1} . \quad (5)$$

Similarly to other MCMC methods based on discretizations of the Langevin diffusion (e.g. [36]), one can build unadjusted schemes which target π^λ , expecting draws from these schemes to be close to draws from π for small enough λ , or add a Metropolis-Hastings step to ensure that the resulting algorithm targets π . Unadjusted proximal MCMC methods have been analyzed in [14]; in this paper we focus on Metropolis adjusted proximal MCMC methods and study their scaling properties. More precisely, at each step k and given the current state of the Markov chain X_k , a candidate Y_{k+1} is generated from the transition density associated to (5), $(\mathbf{x}, \mathbf{y}) \mapsto q(\mathbf{x}, \mathbf{y}) = \varphi(\mathbf{y}; [1 - \sigma^2/(2\lambda)]\mathbf{x} + \sigma^2 \text{prox}_G^\lambda(\mathbf{x})/2\lambda, \sigma^2 \mathbf{I}_d)$, where $\varphi(\cdot; \mathbf{u}, \Sigma)$ stands for the d -dimension Gaussian density with mean $\mathbf{u}$ and covariance matrix Σ . Given X_k and Y_{k+1} , Then, the next state is set as:

$$X_{k+1} = Y_{k+1} b_{k+1} + X_k (1 - b_{k+1}) , \quad b_{k+1} = \mathbb{1}_{\mathbb{R}_+} \left(\frac{\pi(Y_{k+1}) q(Y_{k+1}, X_k)}{\pi(X_k) q(X_k, Y_{k+1})} \wedge 1 - U_{k+1} \right) , \quad (6)$$

where $(U_i)_{i \in \mathbb{N}^*}$ is a sequence of i.i.d. uniform random variables on $[0, 1]$.

The value of λ characterizes how close the distribution π^λ is to the original target π and therefore how good the proposal is. Small values of λ provide better approximations to π and therefore better proposals (see [14, Proposition 1]), while larger values of λ provide higher levels of smoothing for non-differentiable distributions (see [30, Figure 1]). In the case $\lambda = \sigma^2/2$ we obtain the special case of proximal MALA referred to as P-MALA in [30].

The main contribution of this paper is to analyze the optimal scaling for proximal MALA defined by (6).

Optimal scaling and related works We briefly summarize here some examples of MCMC algorithms and their optimal scaling results; a full review is out of the scope of this paper and we only mention algorithms to which we will compare proximal MALA in the development of this work.

Popular examples of Metropolis MCMC are RWM and MALA. RWM uses as a proposal the transition density $(\mathbf{x}, \mathbf{y}) \mapsto \varphi(\mathbf{y} ; \mathbf{x}, \sigma^2 \mathbf{I}_d)$, where $\sigma^2 > 0$. The MALA scheme uses as proposal $(\mathbf{x}, \mathbf{y}) \mapsto \varphi(\mathbf{y} ; \mathbf{x} + (\sigma^2/2)\nabla \log \pi(\mathbf{x}), \sigma^2 \mathbf{I}_d)$. As we will show in Section 3.1, proximal MALA can be considered as an extension of MALA.

A natural question to address when implementing Metropolis adjusted algorithms is how to set the parameter σ^2 (variance parameter for RWM, step size parameter for MALA) to maximize the efficiency of the algorithm. Small values of σ^2 result in higher acceptance probability and cause sticky behaviour, while large values of σ^2 result in a high number of rejections with the chain $(X_k)_{k \geq 0}$ moving slowly [35]. Optimal scaling studies aim to address this question by investigating how σ^2 should behave with respect to the dimension d of the support of π in the high dimensional setting $d \rightarrow \infty$, to obtain the best compromise.

The standard optimal scaling set-up considers the case of d -dimensional targets π_d which are product form, i.e.

$$\pi_d(\mathbf{x}^d) = \prod_{i=1}^d \pi(x_i^d), \quad (7)$$

where x_i^d stands for the i -th component of $\mathbf{x}^d$ and π is a one-dimensional probability density with respect to the Lebesgue measure. Under appropriate assumptions on the regularity of π , and assuming that the MCMC algorithm is initialized at stationarity, the optimal value of σ^2 scales as $\ell^2/d^{2\alpha}$ with $\ell > 0$, $2\alpha = 1$ for RWM [33] and $2\alpha = 1/3$ for MALA [34].

By setting α to these values, it is then possible to show that each as $d \rightarrow \infty$ each 1-dimensional component of the Markov chain defined by RWM and MALA, appropriately rescaled in time, converges to the Langevin diffusion

$$dL_t = h(\ell)^{1/2} dB_t - \frac{h(\ell)}{2} [\log \pi]'(x) dt,$$

where $(B_t)_{t \geq 0}$ is a standard Brownian motion and $h(\ell)$, referred to as speed function of the diffusion, is a function of the parameter $\ell > 0$ that we may tune. Indeed, it is well-known that $(L_{h(\ell)t})_{t \geq 0}$ is a solution of the Langevin diffusion (1). As a result, we may identify the values of ℓ maximizing $h(\ell)$ for the algorithms at hand to approximate the fastest version of the Langevin diffusion. The optimal values for ℓ results in an optimal average acceptance probability of 0.234 for RWM and 0.574 for MALA.

The scaling properties allow to get an intuition of the efficiency of the corresponding algorithms: RWM requires $\mathcal{O}(d)$ steps to achieve convergence on a d -dimensional target, i.e. its efficiency is $\mathcal{O}(d^{-1})$, while MALA has efficiency $\mathcal{O}(d^{-1/3})$. While these results are asymptotic in d , the insights obtained by considering the limit case $d \rightarrow \infty$ prove to be useful in practice [35].

In the context of non-smooth and even discontinuous target distributions, studying the simpler RWM algorithm applied to a class of distributions on compact intervals, [27, 28] show that the lack of smoothness affects the optimal scaling of RWM with respect to dimension d . More precisely, they show that for a class of discontinuous densities which includes the uniform distribution on $[0, 1]$, the optimal scaling of RWM is of order $\mathcal{O}(d^{-2})$. On the other hand, in the case where the set of non-differentiability D of π is a null set with respect to the Lebesgue measure, [12] shows that under appropriate conditions, including L^p differentiability, the optimal scaling of RWM is of order $\mathcal{O}(d^{-1})$ still.

The scaling properties of proximal MALA have been partially investigated in [31], which shows that P-MALA, obtained when $\lambda = \sigma^2/2$, has the same scaling properties of MALA for the finite dimensional Gaussian density and for a class of infinite dimensional target measures (Theorem 2.1 and Theorem 5.1 therein, respectively).

3 Optimal scaling of Proximal MALA

We consider the same set up of [34] and briefly recalled above. Given a real-valued function $g : \mathbb{R} \rightarrow \mathbb{R}$ satisfying **A0** we consider the i.i.d. d -dimensional target specified by (7) with

$$\pi(x) \propto \exp(-g(x)) . \quad (8)$$

Since for any $\mathbf{x}^d$, $G(\mathbf{x}^d) = \sum_{i=1}^d g(x_i^d)$, we have by [29, Section 2.1]

$$\text{prox}_G^\lambda(\mathbf{x}^d) = (\text{prox}_g^\lambda(x_1^d), \dots, \text{prox}_g^\lambda(x_d^d))^\top .$$

It follows that the distribution of the proposal with target π_d in (7)-(8) is also product form $q_d(\mathbf{x}^d, \mathbf{y}^d) = \prod_{i=1}^d q(x_i^d, y_i^d)$ with

$$q(x_i^d, y_i^d) = \frac{1}{(2\pi\sigma^2)^{1/2}} \exp \left(-\frac{(y_i^d - (1 - \sigma^2/(2\lambda))x_i^d - \sigma^2 \text{prox}_g^\lambda(x_i^d)/(2\lambda))^2}{2\sigma^2} \right) ,$$

and $\lambda > 0$. For any dimension $d \in \mathbb{N}^*$, we denote by $(X_k^d)_{k \in \mathbb{N}}$ the Markov chain defined by the Metropolis recursion (6) with target distribution π_d and proposal density q_d and associated to the sequence of candidate moves

$$Y_{k+1}^d = \left(1 - \frac{\sigma^2}{2\lambda}\right) X_k^d + \frac{\sigma^2}{2\lambda} \text{prox}_G^\lambda(X_k^d) + \sigma Z_{k+1}^d . \quad (9)$$

As mentioned in the introduction, the focus of this work is on investigating the optimal dependence of the proposal variance σ^2 on the dimension d of the target π . In this section, we make the dependence of the proposal variance on the dimension explicit and let $\sigma_d^2 = \ell^2/d^{2\alpha}$ and $\lambda_d = c^2/2d^{2\beta}$ for some $\alpha, \beta > 0$ and some constants c, ℓ independent on d . Thus, we can write λ_d as a function of σ_d , $\lambda_d = \sigma_d^{2m} r/2$, where we defined $r = c^2/\ell^{2m} > 0$ and $m = \beta/\alpha$. By writing λ_d as a function of σ_d we can decouple the effect of the constants c, ℓ from that of the dependence on d (i.e. α, β). The

value of m controls the relative speed at which σ_d and λ_d converge to 0 as $d \rightarrow \infty$, when $m = 1$, σ_d and λ_d decay to 0 at the same rate, for $m > 1$ the decay of λ_d is faster than that of σ_d and for $m < 1$ the decay of λ_d is slower than that of σ_d . The parameter r allows to refine the comparison between σ_d and λ_d as $\beta = \alpha$. In the case $m = 1, r = 1$ we get the P-MALA algorithm studied in [30, 31], while for all other values of r, m we have a family of proposals whose behaviour depends on r and m .

3.1 Differentiable targets

We start with the case where π is continuously differentiable. Since MALA can be applied to this class of targets, the results obtained in this section allow direct comparison of proximal MALA algorithms with MALA and thus between gradient-based algorithms (MALA) and algorithms that use proximal operator-based approximations of the gradient (proximal MALA). If $G = -\log \pi$ is continuously differentiable, using [3, Corollary 17.6], $\text{prox}_G^\lambda(\mathbf{x}) = -\lambda \nabla G(\text{prox}_G^\lambda(\mathbf{x})) + \mathbf{x}$, and (5) reduces to

$$\xi_{k+1} = \xi_k - \frac{\sigma^2}{2} \nabla G(\text{prox}_G^\lambda(\xi_k)) + \sigma Z_{k+1} . \quad (10)$$

Hence, the value of λ controls how close to ξ_k is the point at which the gradient is evaluated. For $\lambda \rightarrow 0$, the proximal MALA proposal becomes arbitrarily close to that of MALA, while, as λ increases (10) moves away from MALA.

Our main result, Theorem 1 below, shows that the relative speed of decay (i.e. m) influences the optimal scaling of the resulting proximal MALA algorithm, while the constant r influences the speed function of the limiting diffusion.

We make the following assumptions on the regularity of g .

A1. g is a C^8 -function whose derivatives are bounded by some polynomial: there exists $k_0 \in \mathbb{N}$ such that

$$\sup_{x \in \mathbb{R}} \max_{i \in \{0, \dots, 8\}} [g^{(i)}(x)/(1 + |x|^{k_0})] < \infty .$$

Note that under **A0** and **A1**, [14, Lemma A.1] implies that $\int_{\mathbb{R}} x^k \exp(-g(x)) dx < \infty$ for any $k \in \mathbb{N}$. We also assume that the sequence of proximal MALA algorithms is initialized at stationarity.

A2. For any $d \in \mathbb{N}^*$, X_0^d has distribution π_d .

The assumptions above closely resemble those of [34] used to obtain the optimal scaling results for MALA. In particular, **A1** ensures that we can approximate the log-acceptance ratio in (6) with a Taylor expansion, while **A2** avoids technical complications due to the transient phase of the algorithm. We discuss how the latter assumption could be relaxed in Section 5.

For technical reasons, and to allow direct comparisons with the results established in [34] for MALA, we will also consider the following regularity assumption

A3. The function g' is Lipschitz continuous.

We denote by L_t^d the linear interpolation of the first component of the discrete time Markov chain $(X_k^d)_{k \geq 0}$ obtained with the generic proximal MALA algorithm described above

$$L_t^d = (\lceil d^{2\alpha} t \rceil - d^{2\alpha} t) X_{\lfloor d^{2\alpha} t \rfloor, 1}^d + (d^{2\alpha} t - \lfloor d^{2\alpha} t \rfloor) X_{\lceil d^{2\alpha} t \rceil, 1}^d , \quad (11)$$

where $\lfloor \cdot \rfloor$ and $\lceil \cdot \rceil$ denote the lower and upper integer part functions, respectively, and denote by $X_{k,1}^d$ the first component of X_k^d . The following result shows that in the limit $d \rightarrow \infty$ the properties of proximal MALA depend on the relative speed at which $\sigma_d^2 = \ell^2/d^{2\alpha}$ and $\lambda_d = c^2/2d^{2\beta}$ converge to 0. Recall that we set $r = c^2/\ell^{2m} > 0$ and under **A2**, consider for any $d \in \mathbb{N}^*$,

$$a_d(\ell, r) = \mathbb{E} \left[\frac{\pi_d(Y_1^d) q_d(Y_1^d, X_0^d)}{\pi_d(X_0^d) q_d(X_0^d, Y_1^d)} \wedge 1 \right]. \quad (12)$$

Theorem 1. Assume **A0**, **A1** and **A2**. For any $d \in \mathbb{N}^*$, let $\sigma_d^2 = \ell^2/d^{2\alpha}$ and $\lambda_d = c^2/2d^{2\beta}$ with $\alpha, \beta > 0$. Then, the following statements hold.

- (a) If $\alpha = 1/4$, $\beta = 1/8$ and $r > 0$, we have $\lim_{d \rightarrow +\infty} a_d(\ell, r) = 2\Phi(-\ell^2 K_1(r)/2)$, where Φ is the distribution function of a standard normal and

$$K_1^2(r) = \frac{r^2}{4} \mathbb{E} \left[\{g''(X_{0,1}^d) g'(X_{0,1}^d)\}^2 \right].$$

If in addition, **A3** holds.

- (b) If $\alpha = 1/6$, $\beta = 1/6$ and $r > 0$, we have $\lim_{d \rightarrow +\infty} a_d(\ell, r) = 2\Phi(-\ell^3 K_2(r)/2)$, where Φ is the distribution function of a standard normal and

$$\begin{aligned} K_2^2(r) &= \left(\frac{r}{8} + \frac{r^2}{4} \right) \mathbb{E} [\{g''(X_{0,1}^d) g'(X_{0,1}^d)\}^2] + \left(\frac{1}{16} + \frac{r}{8} \right) \mathbb{E} [g''(X_{0,1}^d)^3] \\ &\quad + \frac{5}{48} \mathbb{E} [g'''(X_{0,1}^d)^2]. \end{aligned}$$

- (c) If $\alpha = 1/6$, $\beta > 1/6$ and $r > 0$, we have $\lim_{d \rightarrow +\infty} a_d(\ell, r) = 2\Phi(-\ell^3 K_2(0)/2)$, where Φ is the distribution function of a standard normal.

In addition, in all these cases, as $d \rightarrow \infty$ the process $(L_t^d)_{t \geq 0}$ converges weakly to the Langevin diffusion

$$dL_t = h(\ell, r)^{1/2} dB_t - \frac{h(\ell, r)}{2} g'(x) dt, \quad (13)$$

where $(B_t)_{t \geq 0}$ denotes standard Brownian motion and $h(\ell, r) = \ell^2 a(\ell, r)$ is the speed of the diffusion, setting $a(\ell, r) = \lim_{d \rightarrow \infty} a_d(\ell, r)$. If $\alpha = 1/4$, $\beta = 1/8$, for any $r > 0$, $\ell \mapsto h(\ell, r)$ is maximized at the unique value of ℓ such that $a(\ell, r) = 0.452$; while if $\alpha = 1/6$, $\beta = m/6$ with $m \geq 1$ and $r > 0$, $\ell \mapsto h(\ell, r)$ is maximized at the unique value of ℓ such that $a(\ell, r) = 0.574$.

Proof. The proof follows that of [34, Theorem 1, Theorem 2] and is postponed to Appendix A. $\square$

The theorem above shows that the relative speed at which λ_d converges to 0 influences the scaling of the resulting proximal algorithm. In case (c), $m > 1$ and λ_d decays with d at a faster rate than σ_d^2 . This causes the proximity map (4) to collapse onto the identity and therefore the proposal (10) is arbitrarily close to that of MALA. The resulting scaling limit also coincides with that of MALA established in [34, Theorem 1, Theorem 2].

If λ_d and σ_d^2 decay at the same rate (case (b)), the amount of gradient information provided by the proximity map is controlled by r . Comparing our result for case (b) with [34, Theorem 1] we find that

$$K_2^2(0) = \frac{1}{16} \mathbb{E} [g''(X_{0,1}^d)^3] + \frac{5}{48} \mathbb{E} [g'''(X_{0,1}^d)^2] = K_{\text{MALA}}^2;$$

thus, we have

$$\begin{aligned} K_2^2(r) &= K_2^2(0) + \left(\frac{r}{8} + \frac{r^2}{4}\right) \mathbb{E} [\{g''(X_{0,1}^d)g'(X_{0,1}^d)\}^2] + \frac{r}{8} \mathbb{E} [g''(X_{0,1}^d)^3] \\ &= K_{\text{MALA}}^2 + \left(\frac{r}{8} + \frac{r^2}{4}\right) \mathbb{E} [\{g''(X_{0,1}^d)g'(X_{0,1}^d)\}^2] + \frac{r}{8} \mathbb{E} [g''(X_{0,1}^d)^3] \geq K_{\text{MALA}}^2, \end{aligned}$$

since the convexity of g implies that $g'' \geq 0$. In particular, $K_2^2(r)$ is an increasing function of r achieving its minimum when $r \rightarrow 0$ (i.e. MALA), see Figure 1(a).

In case (a), $m = 1/2$ and λ_d decays more slowly than σ_d^2 . As a consequence, the gradient information provided by the proximity map is smaller than in cases (b)–(c), and the resulting scaling differs from that of MALA. The value of $K_1^2(r)$ is increasing in r and the speed of the corresponding diffusion also depends on r (see Figure 1(a) gray lines and Figure 1(b)).

Example 1 (Gaussian target). Take $g(x) = x^2/2$, $\text{prox}_\lambda^g(x) = x/(1+\lambda)$. In this case, g' is Lipschitz continuous and we have $K_1^2(r) = r^2/4$, $K_2^2(r) = (1 + 4r + 4r^2)/16$ and $K_2^2(0) = K_{\text{MALA}}^2 = 1/16$. The corresponding speeds are given in Figure 1(a). Optimizing for $m = 1, r = 0$ (MALA) and $m = 1, r = 1$ (P-MALA) we obtain

$$h^{\text{MALA}}(\ell, r) = 1.5639, \quad h^{\text{P-MALA}}(\ell, r) = 0.7519,$$

achieved with $\ell^{\text{MALA}} = 1.6503$ and $\ell^{\text{P-MALA}} = 1.1443$, respectively. For Gaussian targets, MALA is geometrically ergodic [13], and therefore the optimal choice in terms of speed of convergence is MALA which is obtained for $r = 0$. The result for $r = 1$ and $m = 1$ are also given in [31, Theorem 2.1].

Example 2 (Target with light tails). Take $g(x) = x^4$, which gives a normalized distribution with normalizing constant $2\Gamma(5/4)$. The proximity map is

$$\text{prox}_g^\lambda(x) = \frac{1}{2} \left[\frac{\sqrt[3]{9\lambda^2x + \sqrt{54\lambda^4x^2 + 3\lambda^3}}}{3^{2/3}\lambda} - \frac{1}{\sqrt[3]{27\lambda^2x + 3\sqrt{54\lambda^4x^2 + 3\lambda^3}}} \right].$$

In this case g' is not Lipschitz continuous and therefore we only consider (a), for which we have $K_1^2(r) = 144r^2\Gamma(11/4)/\Gamma(5/4)$. The corresponding speed is given in Figure 1(b).

3.2 Laplace target

As discussed in the introduction, proximal MALA has been widely used to quantify uncertainty in imaging applications, in which target distributions involving the ℓ^1 norm are particularly common [30, 14, 1, 46].

Here, we consider π_d^{L} to be the product of d i.i.d. Laplace distributions as in (7),

$$\pi_d^{\text{L}}(\mathbf{x}^d) = \prod_{i=1}^d \pi^{\text{L}}(x_i^d), \text{ for } \mathbf{x}^d \in \mathbb{R}^d, \text{ where } \pi^{\text{L}}(x) = 2^{-1} \exp(-|x|) . \quad (14)$$




Figure 1: Value of K for $i = 1, 2$ and speed of the corresponding Langevin diffusion as a function of r for a Gaussian target and a light tail target. We denote by h_1 the speed obtained in case (a), by h_2 that obtained in (b). In case (c) both K_3 and the speed h_3 are constant w.r.t. r and coincide with that of MALA. For the Gaussian target we report the results for case (a)–(c) while for the light tail target we only report (a).

For this particular choice of one-dimensional target distribution, the corresponding potential G is $x \mapsto |x|$ and satisfies **A0**. Then, the proximity map is given by the soft thresholding operator [29, Section 6.1.3]

$$\text{prox}_G^\lambda(x) = (x - \text{sgn}(x)\lambda)\mathbb{1}\{|x| \geq \lambda\}, \quad (15)$$

where $\text{sgn} : \mathbb{R} \rightarrow \{-1, 1\}$ is the sign function, given by $\text{sgn}(x) = -1$ if $x < 0$, $\text{sgn}(0) = 0$, and $\text{sgn}(x) = 1$ otherwise. This operator is a continuous but not continuously differentiable map whose non-differentiability points are the extrema of the interval $[-\lambda, \lambda]$ and are controlled by the value of the proximity parameter λ .

Plugging (15) in (9), the proximal MALA algorithm applied to π_d^L proposes component-wise for $i = 1, \dots, d$

$$Y_{k+1,i}^d = X_{k,i}^d - \frac{\sigma_d^2}{2} \text{sgn}(X_{k,i}^d)\mathbb{1}\{|X_{k,i}^d| \geq \lambda_d\} - \frac{\sigma_d^2}{2\lambda_d} X_{k,i}^d \mathbb{1}\{|X_{k,i}^d| < \lambda_d\} + \sigma_d Z_{k+1,i}^d. \quad (16)$$

For $X_{k,i}^d$ close to 0 (i.e. the point of non-differentiability) the proximal MALA proposal is a biased random walk around $X_{k,i}^d$, while outside the region $[-\lambda_d, \lambda_d]$ the proposal coincides with that of MALA. As $\lambda_d \rightarrow 0$ the region in which the proximal MALA proposal coincide with that of MALA increases and when $\lambda_d \approx 0$ the region $[-\lambda_d, \lambda_d]$ in which the proposal corresponds to a biased random walk is negligible, as confirmed by the asymptotic acceptance rate in Theorem 2.

We also consider the case $\lambda_d = 0$ for any d . Then, the proposal (16) becomes the proposal for the subgradient version of MALA: $Y_{k+1,i}^d = X_{k,i}^d - (\sigma_d^2/2) \text{sgn}(X_{k,i}^d) + \sigma_d Z_{k+1,i}^d$, referred to as sG-MALA.

The proof of the optimal scaling for the Laplace distribution follows the structure of that of [12] for L^p -mean differentiable distributions. We start by characterizing the asymptotic acceptance

ratio of a generic proximal MALA algorithm; contrary to Theorem 1 for differentiable targets, in the limit $d \rightarrow \infty$ the properties of proximal MALA do not depend on the relative speed at which $\sigma_d^2 = \ell^2/d^{2\alpha}$ and $\lambda_d = c^2/2d^{2\beta}$ converge to 0, as long as λ_d decays at least at the same rate as σ_d^2 . In this regime, the region in which the proposal (16) corresponds to a biased random walk proposal is negligible, and therefore we obtain the same scaling obtained with $\lambda_d = 0$ and corresponding to sG-MALA.

Theorem 2. *Assume A2 and consider the sequence of target distributions $\{\pi_d^L\}_{d \in \mathbb{N}^*}$ given in (14). For any $d \in \mathbb{N}^*$, let $\sigma_d^2 = \ell^2/d^{2\alpha}$ and $\lambda_d = c^2/2d^{2\beta}$ with $\alpha = 1/3$ and $\beta = m/3$ for $m \geq 1$. Then, we have $\lim_{d \rightarrow \infty} a_d(\ell, r) = a^L(\ell) = 2\Phi(-\ell^{3/2}/(72\pi)^{1/4})$, where $(a_d(\ell, r))_{d \in \mathbb{N}^*}$ is defined in (12), with $r = c^2/\ell^{2m}$, and Φ is the distribution function of a standard normal.*

Proof. The proof is postponed to Section 6.1. $\square$

Theorem 2 shows that the asymptotic average acceptance rate $a^L(\ell)$ does not depend on r and as a result on c .

Having identified the possible scaling for proximal MALA with Laplace target, we are now ready to show weak convergence to the appropriate Langevin diffusion. To this end, we adapt the proof strategy followed in [22] and [12].

As for the differentiable case, consider the linear interpolation $(L_t^d)_{t \geq 0}$ of the first component of the Markov chain $(X_k^d)_{k \geq 0}$ given in (11). For any $d \in \mathbb{N}^*$, denote by ν_d the law of the process $(L_t^d)_{t \geq 0}$ on the space of continuous functions from $\mathbb{R}_+$ to $\mathbb{R}$, $C(\mathbb{R}^+, \mathbb{R})$, endowed with the topology of uniform convergence over compact sets and its corresponding σ -field. We first show that the sequence $(\nu_d)_{d \in \mathbb{N}^*}$, admits a weak limit point as $d \rightarrow \infty$.

Proposition 1. *Assume A2 and consider the sequence of target distributions $\{\pi_d^L\}_{d \in \mathbb{N}^*}$ given in (14). For any $d \in \mathbb{N}^*$, let $\sigma_d^2 = \ell^2/d^{2\alpha}$ and $\lambda_d = c^2/2d^{2\beta}$ with $\alpha = 1/3$ and $\beta = m/3$. The sequence $(\nu_d)_{d \in \mathbb{N}^*}$ is tight in $M^1(C(\mathbb{R}^+, \mathbb{R}))$, the set of probability measures acting on $C(\mathbb{R}^+, \mathbb{R})$.*

Proof. See Section 6.2. $\square$

By Prokhorov's theorem, the tightness of $(\nu_d)_{d \in \mathbb{N}^*}$ implies existence of a weak limit point ν . In our next result, we give a sufficient condition to show that any limit point of $(\nu_d)_{d \in \mathbb{N}^*}$ coincides with the law of a solution of:

$$dL_t = [h^L(\ell)]^{1/2} dB_t - \frac{h^L(\ell)}{2} \operatorname{sgn}(L_t) dt. \quad (17)$$

To this end, we consider the martingale problem (see [42]) associated with (17), that we now present. Let us denote by $C_c^\infty(\mathbb{R}, \mathbb{R})$ the subset of functions of $C(\mathbb{R}, \mathbb{R})$ which are infinitely many times differentiable and with compact support, and define the generator of (17) for $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$ by

$$LV(x) = \frac{h^L(\ell)}{2} [V''(x) - \operatorname{sgn}(x)V'(x)]. \quad (18)$$

Denote by $(W_t)_{t \geq 0}$ the canonical process on $C(\mathbb{R}_+, \mathbb{R})$, $W_t : \{w_s\}_{s \geq 0} \mapsto w_t$ and the corresponding filtration by $(\mathfrak{F}_t)_{t \geq 0}$. A probability measure ν is said to solve the martingale problem associated

with (17) with initial distribution π^L , if the pushforward of ν by W_0 is π^L and if for all $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$, the process

$$\left(V(W_t) - V(W_0) - \int_0^t LV(W_u) du \right)_{t \geq 0}$$

is a martingale with respect to ν and the filtration $(\mathfrak{F}_t)_{t \geq 0}$. The following proposition gives a sufficient condition to prove that ν is a solution of the martingale problem:

Proposition 2. *Suppose that for any $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$, $m \in \mathbb{N}$, $\rho : \mathbb{R}^m \rightarrow \mathbb{R}$ bounded and continuous, and for any $0 \leq t_1 \leq \dots \leq t_m \leq s \leq t$:*

$$\lim_{d \rightarrow +\infty} \mathbb{E}^{\nu_d} \left[\left(V(W_t) - V(W_s) - \int_s^t LV(W_u) du \right) \rho(W_{t_1}, \dots, W_{t_m}) \right] = 0.$$

Then any limit point of $(\nu_d)_{d \in \mathbb{N}^}$ on $M^1(C(\mathbb{R}^+, \mathbb{R}))$ is a solution to the martingale problem associated with (17).*

Proof. See Section 6.3. □

Finally, we use this sufficient condition to establish that any limit point of $(\nu_d)_{d \in \mathbb{N}^*}$ is a solution of the martingale problem for (17). Uniqueness in law of solutions of (17) allows to conclude that $(L_t^d)_{t \geq 0}$ converges weakly to the Langevin diffusion (17), which establishes our main result.

Theorem 3. *The sequence of processes $\{(L_t^d)_{t \geq 0} : d \in \mathbb{N}^*\}$ converges in distribution towards $(L_t)_{t \geq 0}$, solution of (17) as $d \rightarrow \infty$, with $h^L(\ell) = \ell^2 a^L(\ell)$ and a^L defined in Theorem 2. In addition, h^L is maximized at the unique value of ℓ such that $a^L(\ell) = 0.360$.*

Proof. See Section 6.4. □

4 Practical Implications and Numerical Simulations

The optimal scaling results in Sections 3.1 and 3.2 provide some guidance on the choice of the parameters σ and λ of proximal MALA algorithms, suggesting that smaller values of λ provide better efficiency in terms of number of steps necessary to convergence (Theorem 1).

However, a number of other factors must be taken into account. First, as shown in [26, 37, 36, 21] the convergence properties of Metropolis adjusted algorithms are influenced by the shape of the target distribution and, in particular, by its tail behavior. Secondly, when comparing proximal MALA algorithms with gradient-based methods (e.g. MALA) one must take into account the cost of obtaining the gradients, whether this comes from automatic differentiation algorithms or from evaluating a potentially complicated gradient function. On the other hand, proximity mappings can be quickly found or approximated solving convex optimization problems which have been widely studied in the convex optimization literature (e.g. [29, Chapter 6], [11] and [30, Section 3.2.3]).

In terms of convergence properties, we are usually interested in the family of distributions for which the discrete time Markov chain produced by our algorithm is geometrically ergodic, together with the optimal scaling results briefly recalled in Section 2. Normally, the ergodicity results are given by considering the one-dimensional class of distributions $\mathcal{E}(\beta, \gamma)$ introduced in [36] and defined for $\gamma > 0$ and $0 < \beta < \infty$ by

$$\mathcal{E}(\beta, \gamma) : \{ \pi : \mathbb{R} \rightarrow [0, +\infty) : \pi(x) \propto \exp(-\gamma|x|^\beta), |x| > x_0 \text{ for some } x_0 > 0 \}.$$

As observed by [24], there usually is a trade-off between ergodicity and optimal scaling results, algorithms providing better optimal scaling results tend to be geometrically ergodic for a smaller set of targets (e.g. MALA w.r.t. RWM).

As suggested by Theorem 1, the scaling properties of proximal MALA on differentiable targets are close to those of MALA. This leads to a natural comparison between the two algorithms. First, we observe that A0 rules out targets for which G is not convex and therefore restricts the families $\mathcal{E}(\beta, \gamma)$ to $\beta \geq 1$. To compare MALA with proximal MALA we therefore focus on distributions with $\beta \geq 1$.

It is shown in [36] that MALA is geometrically ergodic for targets in $\mathcal{E}(\beta, \gamma)$ with $1 \leq \beta \leq 2$ (with some caveat for $\beta = 2$). Theorem 1-(b) and (c) show that in this case proximal MALA has the same scaling properties of MALA but in case (b) the asymptotic speed of convergence decays as the constant r increases (Figure 1(a)), with the maximum achieved for $r \rightarrow 0$, for which proximal MALA collapses onto MALA. Since MALA is geometrically ergodic, and achieves better (or equivalent) scaling properties than proximal MALA, it would be natural to prefer MALA to proximal MALA for this set of targets. However, if the gradient is costly to obtain, one might instead consider to use proximal MALA with a small λ , to retain scaling properties as close as possible to that of MALA but to reduce the computational cost of evaluating the gradient.

In the case of differentiable targets with light-tails (i.e. $\beta > 2$), MALA is known not to be geometrically ergodic [36, Section 4.2] while the ergodicity properties of proximal MALA have only been partially studied in [30, Section 3.2.2] for the case $\lambda = \sigma^2/2$ (P-MALA). As shown in [30, Section 2.1], given a distribution $\pi \in \mathcal{E}(\beta, \gamma)$ with $\beta \geq 1$, the distribution π_λ obtained using the potential (3) belongs to $\mathcal{E}(\beta', \gamma')$, where $\beta' = \min(\beta, 2)$ and γ' depending on λ . This suggests that proximal MALA is likely to be geometrically ergodic for appropriate choices of λ ; a first result in this direction is given in [30, Corollary 3.2] for the P-MALA case $\lambda = \sigma^2/2$. Theorem 1-(c) restricts the sets of available λ s showing that for light-tail distributions (for which A3 does not hold) λ should decay at half the speed of σ^2 . Studying the ergodicity properties of proximal MALA in function of the parameter λ is, of course, an interesting problem that we leave for future work.

For the Laplace distribution, Theorem 2 shows that the value of λ does not influence the asymptotic acceptance ratio of proximal MALA, as long as λ decays with d at least as fast as σ^2 . The scaling properties and the asymptotic speed $h(\ell)$ in Theorem 3 do not depend on λ and coincide with that of the sG-MALA (obtained for $\lambda = 0$). Hence, in terms of optimal scaling, there does not seem to be a difference between proximal MALA and sG-MALA for the Laplace distribution.

4.1 Numerical Experiments

To illustrate the results established in Section 3.1 and 3.2 we consider here a small collection of simulation studies. The aim of these studies is to empirically confirm the optimal scalings identified in Theorem 1 and 2, investigate the dimension d at which the asymptotic acceptance ratio $\lim_{d \rightarrow \infty} a_d(\ell, r)$ well approximates the empirical average acceptance ratio and, consequently, for which dimensions d we can expect the optimal asymptotic acceptances in Theorem 1 and 2 to guarantee maximal speed $h(\ell, r)$ (approximated by the expected squared jumping distance, see, e.g. [18]) for the corresponding diffusion. We summarize here our findings, a more detailed discussion can be found in Appendix B.

For the differentiable case, we consider the Gaussian distribution in Example 1 and four algorithmic settings which correspond to the three cases identified in Theorem 1 and MALA. The different values of r and m influence the dimension required to observe convergence to the theoretical limit




Figure 2: Proximal MALA with Laplace target and $m = 1, r = 0$ (sG-MALA). Left: acceptance rate as a function of ℓ for increasing dimension d ; Right: ESJD_d as a function of the acceptance rate $a_d(\ell, r)$.




Figure 3: Proximal MALA with Laplace target and $m = 1, r = 1$ (P-MALA). Left: acceptance rate as a function of ℓ for increasing dimension d ; Right: ESJD_d as a function of the acceptance rate $a_d(\ell, r)$.

in Theorem 1: for $r \rightarrow 0$ and $m = 1$ (MALA) and $m = 1/2, r = 1$ (corresponding to Theorem 1-(a)) the theoretical limit is already achieved for d of order 10^2 , while in the cases $m = 3, r = 2$ and $m = r = 1$ (corresponding to Theorem 1-(c) and (b), respectively) our simulation result match the theoretical limit only for d of order 10^5 or higher.

The results for the Laplace case are similar, with the case $m > 1$ requiring a higher d to observe convergence to the theoretical limit. Figure 2 and Figure 3 provide numerical simulations of the behavior, as d increases, of the mean acceptance ratio $(a_d(\ell, r))_{d \in \mathbb{N}^*}$ as a function of ℓ and $(\text{ESJD}_d)_{d \in \mathbb{N}^*}$ as a function of $(a_d(\ell, r))_{d \in \mathbb{N}^*}$, for sG-MALA ($r = 0$) and P-MALA ($r = 1$) respectively. These confirm our theoretical founding Theorem 2 and Theorem 3.

In general, we find that the optimal average acceptance ratios in Theorem 1 and 3 guarantee maximal speed $h(\ell, r)$ for d sufficiently large (for small d the optimal acceptance ratio often differs from the optimal asymptotic one, see, e.g. [40, Section 2.1]).

To further investigate the scaling of proximal MALA to other non-differentiable densities, we




Figure 4: Proximal MALA for the target (19) and $m = 3, r = 2$, with $\sigma^2 = \ell/d^{2\alpha}$ and $\alpha = 1/3$. Left: acceptance rate as a function of ℓ for increasing dimension d ; Right: ESJD_d as a function of the acceptance rate $a_d(\ell, r)$.

empirically study the case where the sequence of targets is given by: $x^d \in \mathbb{R}^d$,

$$\pi_d^{\text{GL}}(x^d) = \prod_{i=1}^d \exp(-g(x_i^d)), \quad g(x) = |x| + x^2/2, \quad (19)$$

which, like the Laplace distribution in 0, is non-differentiable but convex. The study of such a potential is motivated by Bayesian inverse problems considered in [30, 14], for which the posterior distribution arises from Gaussian observations and sparsity-induced priors like the Laplace distribution. The posterior then has the form (up to a multiplicative constant) $x^d \mapsto \exp(-\|y^d - \mathbf{A}x^d\| - c_r \sum_{i=1}^d |x_i|)$.

For this choice of target (19), the proposal of proximal MALA is given for any $d \in \mathbb{N}^*$, $k \in \mathbb{N}$ and $i \in \{1, \dots, d\}$,

$$Y_{k+1,i}^d = \left(1 - \frac{\sigma_d^2}{2\lambda_d}\right) X_{k,i}^d + \frac{\sigma_d^2}{2\lambda_d} ((X_{k,i}^d - \text{sgn}(X_{k,i}^d)) \mathbb{1}_{\{|X_{k,i}^d| \geq \lambda_d\}} - \lambda_d X_{k,i}^d) + \sigma_d Z_{k+1,i}^d,$$

where $(Z_{k+1,i}^d)_{k \in \mathbb{N}}$ is a sequence of standard normal random variables. We then repeated the same experiments as for the Laplace distribution. The results are shown in Figures 4, 12 and 13. From these figures it is clear that the same scaling holds, i.e. choosing $\sigma_d^2 = \ell/d^{2\alpha}$, $\lambda = \sigma_d^{2m} r/2$ with $\alpha = 1/3$, $r \geq 0$.

5 Discussion

In this work we analyze the scaling properties of a wide class of proximal MALA algorithms introduced in [30, 14] for smooth targets and for the Laplace distribution. We show that the scaling properties of proximal MALA are influenced by the relative speed at which the proximal parameter λ_d and the proposal variance σ_d decay to 0 as $d \rightarrow \infty$ and suggest practical ways to choose λ_d as a function of σ_d to guarantee good results.

In the case of smooth targets, we provide a detailed comparison between proximal MALA and MALA, showing that proximal MALA scales no better than MALA (Theorem 1). In particular,

Theorem 1-(a) shows that if λ_d is too large w.r.t. σ_d then the efficiency of proximal MALA is of order $\mathcal{O}(d^{-1/2})$ and therefore worse than the $\mathcal{O}(d^{-1/3})$ of MALA, suggesting that λ_d should be chosen to decay approximately as σ_d , if possible. If λ_d decays sufficiently fast, then MALA and proximal MALA have similar scaling properties and, in the case in which the proximity map is cheaper to compute than the gradient, one can build proximal MALA algorithms which are as efficient as MALA in terms of scaling but more computationally efficient.

In the case of the Laplace distribution, we show that the scaling of proximal MALA is $\mathcal{O}(d^{-2/3})$ for any λ_d decaying sufficiently fast w.r.t. σ_d and, in the case $\lambda_d = 0$, we obtain a novel optimal scaling result for sG-MALA on Laplace targets.

As discussed in Section 4, our analysis provides some guidance on the choice of the parameters that need to be specified to implement proximal MALA, but this analysis should be complemented by an exploration of the ergodicity properties of proximal MALA to obtain a comprehensive description of the algorithms. We conjecture that for sufficiently large values of λ , proximal MALA applied to light tail distributions will be exponentially ergodic; establishing exactly how large should λ be to guarantee fast convergence is an interesting question that we leave for future work. Obtaining these results would open the doors to adaptive tuning strategies for proximal MALA, which are likely to produce better results than those given by the strategies currently used.

The set up under which we carried out our analysis closely resembles that of [34]; we anticipate that A2 could be relaxed following similar ideas as those in [10, 22] and that our analysis could be extended to d -dimensional targets π_d possessing some dependence structure following the approach of [40, 4, 45]. Finally, the analysis carried out for the Laplace distribution could be extended to other piecewise smooth distributions provided that the moments necessary for the proof in Section 6 can be computed.

6 Proof of the Result for the Laplace distribution

In this section we prove the results in Section 3.2 which give the scaling properties of proximal MALA (and sG-MALA) for the Laplace distribution. We collect technical results (e.g. moment computations, bounds, etc.) in Appendix D.

We recall that $\sigma_d^2 = \ell^2/d^{2\alpha}$ and $\lambda_d = c^2/2d^{2\beta}$ for some $\alpha, \beta > 0$ and some constants c, ℓ independent on d . Thus, we can write λ_d as a function of σ_d , $\lambda_d = \sigma_d^{2m}r/2$, where we define $r = c^2/\ell^{2m} \geq 0$ and $m = \beta/\alpha$.

In order to study the scaling limit of proximal MALA with Laplace target, consider the mapping $b_d : \mathbb{R}^2 \rightarrow \mathbb{R}$ given by

$$b_d : (x, z) \mapsto z - \frac{\sigma_d}{2} \operatorname{sgn}(x) \mathbb{1}\{|x| \geq \sigma_d^{2m}r/2\} - \frac{1}{\sigma_d^{2m-1}r} x \mathbb{1}\{|x| < \sigma_d^{2m}r/2\} , \quad (20)$$

which allows us to write the proposal as $Y_{1,i}^d = X_{0,i}^d + \sigma_d b_d(X_{0,i}^d, Z_{1,i}^d)$, for any $i \in \{1, \dots, d\}$.

We consider also the function $\phi_d : \mathbb{R}^2 \rightarrow \mathbb{R}$, given by

$$\begin{aligned} \phi_d : (x, z) &\mapsto \log \frac{\pi(x + \sigma_d b_d(x, z)) q(x + \sigma_d b_d(x, z), x)}{\pi(x) q(x, x + \sigma_d b_d(x, z))} \\ &= |x| - |x + \sigma_d b_d(x, z)| + \frac{z^2}{2} \\ &\quad - \frac{1}{2\sigma_d^2} \left\{ \frac{\sigma_d^2}{2} \operatorname{sgn}[x + \sigma_d b_d(x, z)] \mathbb{1} \left\{ |x + \sigma_d b_d(x, z)| \geq \frac{\sigma_d^{2m} r}{2} \right\} \right. \\ &\quad \left. - \sigma_d b_d(x, z) \right. \\ &\quad \left. + \frac{1}{\sigma_d^{2(m-1)} r} [x + \sigma_d b_d(x, z)] \mathbb{1} \left\{ |x + \sigma_d b_d(x, z)| < \frac{\sigma_d^{2m} r}{2} \right\} \right\}^2. \end{aligned} \quad (21)$$

6.1 Proof of Theorem 2

The proof of Theorem 2 uses the first three moments of $\phi_{d,1}$, whose computation is postponed to Appendix D.1, and is an application of Lindeberg's central limit theorem. We introduce, for $i \in \{1, \dots, d\}$, $\phi_{d,i} = \phi_d(X_{0,i}^d, Z_{1,i}^d)$ for the sake of conciseness. This allows us to rewrite $a_d(\ell, r)$, defined in (12), in the following way,

$$a_d(\ell, r) = \mathbb{E} \left[\exp \left(\sum_{i=1}^d \phi_{d,i} \right) \wedge 1 \right].$$

Remark 1. Under A2, the families of random variables $(b_d(X_{0,i}^d, Z_{1,i}^d))_{i \in \{1, \dots, d\}}$ and $(\phi_{d,i})_{i \in \{1, \dots, d\}}$ are i.i.d..

To identify the optimal scaling for the Laplace distribution, we look for those values of α such that $\sum_{i=1}^d \mathbb{E}[\phi_{d,i}]$ and $\operatorname{Var}(\sum_{i=1}^d \phi_{d,i})$ converge to a finite value. Using Remark 1, we have that,

$$\sum_{i=1}^d \mathbb{E}[\phi_{d,i}] = d \mathbb{E}[\phi_{d,1}] \quad \text{and} \quad \operatorname{Var} \left(\sum_{i=1}^d \phi_{d,i} \right) = d \operatorname{Var}(\phi_{d,1}). \quad (22)$$

Then, using the integrals in Appendix D.1, we find that the only value of α for which (22) converge to a finite value with the variance strictly positive is $\alpha = 1/3$ as confirmed empirically in Appendix B.2.

Having identified $\alpha = 1/3$, we can then proceed applying Lindeberg's CLT.

Proof of Theorem 2. We start by showing that the acceptance ratio converges to a Gaussian distribution. Define $\mu_d = \mathbb{E}[\phi_{d,1}]$ and $\mathcal{F}_{d,i} = \sigma((X_{0,j}^d, Z_{1,j}^d), 1 \leq j \leq i)$, the natural filtration for $(X_{0,i}^d, Z_{1,i}^d)_{d \in \mathbb{N}, 1 \leq i \leq d}$. The square-integrable martingale sequence

$$\left(\sum_{j=1}^i W_{d,j}, \mathcal{F}_{d,i} \right)_{d \in \mathbb{N}^*, 1 \leq i \leq d}$$

where $W_{d,i} = \phi_{d,i} - \mu_d$, forms a triangular array, to which we can apply the corresponding CLT (e.g. [41, Theorem 4, page 543]). In particular, we have that,

$$\lim_{d \rightarrow \infty} \sum_{i=1}^d \mathbb{E}[W_{d,i}^2 | \mathcal{F}_{d,i-1}] = \lim_{d \rightarrow \infty} d \operatorname{Var}(\phi_{d,1}) = \frac{2\ell^3}{3\sqrt{2\pi}},$$

as shown in Proposition 17 in Appendix D.1. It remains to verify Lindeberg's condition: for $\varepsilon > 0$,

$$\lim_{d \rightarrow \infty} d\mathbb{E} [W_{d,1}^2 \mathbb{1} \{|W_{d,1}| > \varepsilon\}] = 0 .$$

In order to verify Lindeberg's condition we verify the stronger Lyapunov condition: there exists $\epsilon > 0$ such that

$$\lim_{d \rightarrow \infty} d\mathbb{E} [W_{d,1}^{2+\epsilon}] = 0 .$$

Pick $\epsilon = 1$ and expand the cube using $\mu_d = \mathbb{E}[\phi_{d,i}]$,

$$\mathbb{E} [W_{d,1}^3] = \mathbb{E} [\phi_{d,i}^3] - 3\mu_d \mathbb{E} [\phi_{d,i}^2] + 2\mu_d^3 . \quad (23)$$

By Proposition 16 in Appendix D.1, we have $\lim_{d \rightarrow \infty} d\mu_d^3 = 0$, $\lim_{d \rightarrow \infty} \mu_d = 0$, and, by Proposition 17 in Appendix D.1,

$$\lim_{d \rightarrow \infty} d\mathbb{E} [\phi_{d,i}^2] = \frac{2\ell^3}{3\sqrt{2\pi}} .$$

Finally, for the remaining term in (23) we use Proposition 18 in Appendix D.1 to show that $\lim_{d \rightarrow \infty} d\mathbb{E}[\phi_{d,i}^3] = 0$. The above and the fact that, by Proposition 16 in Appendix D.1,

$$\lim_{d \rightarrow \infty} d\mu_d = -\frac{\ell^3}{3\sqrt{2\pi}} ,$$

show, by Lindeberg's CLT, that the acceptance ratio converges in law to a normal random variable $\tilde{Z}$ with mean $-\ell^3/(3\sqrt{2\pi})$ and variance $2\ell^3/(3\sqrt{2\pi})$.

To conclude the proof, we apply the continuous mapping theorem to the bounded and continuous function $x \mapsto e^x \wedge 1$ and obtain

$$\lim_{d \rightarrow \infty} \exp \left(\sum_{i=1}^d \phi_{d,i} \right) \wedge 1 \stackrel{d}{=} e^{\tilde{Z}} \wedge 1 \quad \text{and} \quad \lim_{d \rightarrow \infty} a_d(\ell, r) = \mathbb{E} [e^{\tilde{Z}} \wedge 1] ,$$

where the limit does not depend on r . Defining $a^L(\ell) = \lim_{d \rightarrow \infty} a_d(\ell, r)$ and using [33, Proposition 2.4], we have the result. $\square$

6.2 Proof of Proposition 1

We are interested in the law ν_d of the linear interpolant $(L_t^d)_{t \geq 0}$, defined in (11), of the first component of the chain $(X_k^d)_{k \in \mathbb{N}}$. Let us recall the definition of the chain: assumption **A2** gives the initial distribution π_d , then, for any $k \in \mathbb{N}$, the proposal $Y_{k+1}^d = (Y_{k+1,i}^d)_{1 \leq i \leq d}$ is defined in (16) with $\sigma_d^2 = \ell^2/d^{2\alpha}$, $\lambda_d = \sigma_d^{2m}r/2$ with $\alpha = 1/3$ and $m \geq 1$. The proposal (16) can be written as

$$Y_{k+1,i}^d = X_{k,i}^d + \sigma_d b_d(X_{k,i}^d, Z_{k+1,i}^d) , \quad (24)$$

for any $i \in \{1, \dots, d\}$, where b_d is defined in (20) and $r = c^2/\ell^{2m}$. We further define the acceptance event $A_{k+1}^d = \{b_{k+1}^d = 1\}$ where b_{k+1}^d is as in (6).

We can now expand the expression of the linear interpolant L_t^d using (6), (11) and the definition of A_{k+1}^d ,

$$L_t^d = \begin{cases} X_{\lfloor d^{2\alpha}t \rfloor, 1}^d + (d^{2\alpha}t - \lfloor d^{2\alpha}t \rfloor) \left[\sigma_d Z_{\lceil d^{2\alpha}t \rceil, 1}^d - \frac{\sigma_d^2}{2} \operatorname{sgn}(X_{\lceil d^{2\alpha}t \rceil, 1}^d) \right] \mathbb{1}_{A_{\lceil d^{2\alpha}t \rceil}^d} \\ \quad \text{if } |X_{\lfloor d^{2\alpha}t \rfloor, 1}^d| \geq \frac{\sigma_d^{2m}r}{2} \\ X_{\lfloor d^{2\alpha}t \rfloor, 1}^d + (d^{2\alpha}t - \lfloor d^{2\alpha}t \rfloor) \left[\sigma_d Z_{\lceil d^{2\alpha}t \rceil, 1}^d - \frac{1}{\sigma_d^{2(m-1)}r} X_{\lfloor d^{2\alpha}t \rfloor, 1}^d \right] \mathbb{1}_{A_{\lceil d^{2\alpha}t \rceil}^d} \\ \quad \text{otherwise} \end{cases}, \quad (25)$$

or, equivalently,

$$L_t^d = \begin{cases} X_{\lceil d^{2\alpha}t \rceil, 1}^d - (\lceil d^{2\alpha}t \rceil - d^{2\alpha}t) \left[\sigma_d Z_{\lceil d^{2\alpha}t \rceil, 1}^d - \frac{\sigma_d^2}{2} \operatorname{sgn}(X_{\lceil d^{2\alpha}t \rceil, 1}^d) \right] \mathbb{1}_{A_{\lceil d^{2\alpha}t \rceil}^d} \\ \quad \text{if } |X_{\lceil d^{2\alpha}t \rceil, 1}^d| \geq \frac{\sigma_d^{2m}r}{2} \\ X_{\lceil d^{2\alpha}t \rceil, 1}^d - (\lceil d^{2\alpha}t \rceil - d^{2\alpha}t) \left[\sigma_d Z_{\lceil d^{2\alpha}t \rceil, 1}^d - \frac{1}{\sigma_d^{2(m-1)}r} X_{\lceil d^{2\alpha}t \rceil, 1}^d \right] \mathbb{1}_{A_{\lceil d^{2\alpha}t \rceil}^d} \\ \quad \text{otherwise} \end{cases}.$$

In order to prove Proposition 1, we consider Kolmogorov's criterion for tightness (see [23, Theorem 23.7]): the sequence $(\nu_d)_{d \geq 1}$ is tight if the sequence $(L_0^d)_{d \in \mathbb{N}^*}$ is tight, and

$$\mathbb{E} [(L_t^d - L_s^d)^4] \leq \gamma(t)(t-s)^2,$$

for some non-decreasing positive function γ , all $0 \leq s \leq t$ and all $d \in \mathbb{N}^*$. The condition on $(L_0^d)_{d \in \mathbb{N}^*}$ is straightforward to check, since by A2 the distribution of $L_0^d = X_{0,1}^d$ is π^L for all $d \in \mathbb{N}^*$.

Proof of Proposition 1. Consider $\mathbb{E} [(L_t^d - L_s^d)^4]$, if $\lfloor d^{2\alpha}s \rfloor = \lfloor d^{2\alpha}t \rfloor$, the inequality follows straightforwardly recalling that the moments of normal distributions are bounded: in the case $|X_{\lfloor d^{2\alpha}t \rfloor, 1}^d| = |X_{\lfloor d^{2\alpha}s \rfloor, 1}^d| \geq \sigma_d^{2m}r/2$ it follows directly from the boundedness of the sgn function, while in the case $|X_{\lfloor d^{2\alpha}t \rfloor, 1}^d| = |X_{\lfloor d^{2\alpha}s \rfloor, 1}^d| < \sigma_d^{2m}r/2$ we exploit the boundedness of $X_{\lfloor d^{2\alpha}t \rfloor, 1}^d$ itself.

For all $0 \leq s \leq t$ such that $\lceil d^{2\alpha}s \rceil \leq \lfloor d^{2\alpha}t \rfloor$, we can distinguish three cases.

Case 1 If $|X_{\lfloor d^{2\alpha}t \rfloor, 1}^d| \geq \sigma_d^{2m}r/2$ and $|X_{\lfloor d^{2\alpha}s \rfloor, 1}^d| \geq \sigma_d^{2m}r/2$, then

$$\begin{aligned} L_t^d - L_s^d &= X_{\lfloor d^{2\alpha}t \rfloor, 1}^d - X_{\lfloor d^{2\alpha}s \rfloor, 1}^d \\ &\quad + (d^{2\alpha}t - \lfloor d^{2\alpha}t \rfloor) \left[\sigma_d Z_{\lceil d^{2\alpha}t \rceil, 1}^d - \frac{\sigma_d^2}{2} \operatorname{sgn}(X_{\lceil d^{2\alpha}t \rceil, 1}^d) \right] \mathbb{1}_{A_{\lceil d^{2\alpha}t \rceil}^d} \\ &\quad + (\lceil d^{2\alpha}s \rceil - d^{2\alpha}s) \left[\sigma_d Z_{\lceil d^{2\alpha}s \rceil, 1}^d - \frac{\sigma_d^2}{2} \operatorname{sgn}(X_{\lceil d^{2\alpha}s \rceil, 1}^d) \right] \mathbb{1}_{A_{\lceil d^{2\alpha}s \rceil}^d}. \end{aligned}$$

Using Hölder's inequality and the fact that $0 \leq d^{2\alpha}t - \lfloor d^{2\alpha}t \rfloor \leq 1$ (and similarly for s) we have

$$\begin{aligned} \mathbb{E}[(L_t^d - L_s^d)^4] &\leq C \mathbb{E} \left[\left(X_{\lfloor d^{2\alpha}t \rfloor, 1}^d - X_{\lfloor d^{2\alpha}s \rfloor, 1}^d \right)^4 \right] \\ &\quad + C \frac{(d^{2\alpha}t - \lfloor d^{2\alpha}t \rfloor)^2}{d^{4\alpha}} \mathbb{E} \left[\left(\ell Z_{\lfloor d^{2\alpha}t \rfloor, 1}^d \right)^4 + \frac{\ell^8}{2^4 d^{4\alpha}} \right] \\ &\quad + C \frac{(\lceil d^{2\alpha}s \rceil - d^{2\alpha}s)^2}{d^{4\alpha}} \mathbb{E} \left[\left(\ell Z_{\lceil d^{2\alpha}s \rceil, 1}^d \right)^4 + \frac{\ell^8}{2^4 d^{4\alpha}} \right]. \end{aligned}$$

Recalling that the moments of Z^d are bounded and that $d^{2\alpha}s \leq \lceil d^{2\alpha}s \rceil \leq \lfloor d^{2\alpha}t \rfloor \leq d^{2\alpha}t$, it follows

$$\mathbb{E}[(L_t^d - L_s^d)^4] \leq C \left((t-s)^2 + \mathbb{E} \left[\left(X_{\lfloor d^{2\alpha}t \rfloor, 1}^d - X_{\lfloor d^{2\alpha}s \rfloor, 1}^d \right)^4 \right] \right). \quad (26)$$

Case 2 If $|X_{\lfloor d^{2\alpha}t \rfloor, 1}^d| \geq \sigma_d^{2m}r/2$ and $|X_{\lfloor d^{2\alpha}s \rfloor, 1}^d| < \sigma_d^{2m}r/2$ or $|X_{\lfloor d^{2\alpha}t \rfloor, 1}^d| < \sigma_d^{2m}r/2$ and $|X_{\lfloor d^{2\alpha}s \rfloor, 1}^d| \geq \sigma_d^{2m}r/2$. We only describe the argument for the first case, the second case follows from analogous steps. Take

$$\begin{aligned} L_t^d - L_s^d &= X_{\lfloor d^{2\alpha}t \rfloor, 1}^d + (d^{2\alpha}t - \lfloor d^{2\alpha}t \rfloor) \left[\sigma_d Z_{\lfloor d^{2\alpha}t \rfloor, 1}^d - \frac{\sigma_d^2}{2} \text{sgn}(X_{\lfloor d^{2\alpha}t \rfloor, 1}^d) \right] \mathbb{1}_{\mathcal{A}_{\lfloor d^{2\alpha}t \rfloor}^d} \\ &\quad - X_{\lfloor d^{2\alpha}s \rfloor, 1}^d - (\lceil d^{2\alpha}s \rceil - d^{2\alpha}s) \left(\sigma_d Z_{\lceil d^{2\alpha}s \rceil, 1}^d - \frac{1}{\sigma_d^{2(m-1)}r} X_{\lceil d^{2\alpha}s \rceil, 1}^d \right) \mathbb{1}_{\mathcal{A}_{\lceil d^{2\alpha}s \rceil}^d}. \end{aligned}$$

Proceeding as above, we find that

$$\begin{aligned} \mathbb{E}[(L_t^d - L_s^d)^4] &\leq C \left((t-s)^2 + \mathbb{E} \left[\left(X_{\lfloor d^{2\alpha}t \rfloor, 1}^d - X_{\lfloor d^{2\alpha}s \rfloor, 1}^d \right)^4 \right] \right. \\ &\quad \left. + (\lceil d^{2\alpha}s \rceil - d^{2\alpha}s)^4 \mathbb{E} \left[\left(\frac{1}{\sigma_d^{2(m-1)}r} X_{\lceil d^{2\alpha}s \rceil, 1}^d \right)^4 \right] \right), \end{aligned}$$

and recalling that $|X_{\lfloor d^{2\alpha}s \rfloor, 1}^d| < \sigma_d^{2m}r/2$ we have that $|X_{\lfloor d^{2\alpha}s \rfloor, 1}^d|/(r\sigma_d^{2(m-1)}) < \sigma_d^2/2$. Using this and the same arguments as above, we have

$$\mathbb{E}[(L_t^d - L_s^d)^4] \leq C \left((t-s)^2 + \mathbb{E} \left[\left(X_{\lfloor d^{2\alpha}t \rfloor, 1}^d - X_{\lfloor d^{2\alpha}s \rfloor, 1}^d \right)^4 \right] \right). \quad (27)$$

Case 3 If $|X_{\lfloor d^{2\alpha}t \rfloor, 1}^d| < \sigma_d^{2m}r/2$ and $|X_{\lfloor d^{2\alpha}s \rfloor, 1}^d| < \sigma_d^{2m}r/2$, then

$$\begin{aligned} L_t^d - L_s^d &= X_{\lfloor d^{2\alpha}t \rfloor, 1}^d + (d^{2\alpha}t - \lfloor d^{2\alpha}t \rfloor) \left(\sigma_d Z_{\lfloor d^{2\alpha}t \rfloor, 1}^d - \frac{1}{\sigma_d^{2(m-1)}r} X_{\lfloor d^{2\alpha}t \rfloor, 1}^d \right) \mathbb{1}_{\mathcal{A}_{\lfloor d^{2\alpha}t \rfloor}^d} \\ &\quad - X_{\lfloor d^{2\alpha}s \rfloor, 1}^d + (\lceil d^{2\alpha}s \rceil - d^{2\alpha}s) \left(\sigma_d Z_{\lceil d^{2\alpha}s \rceil, 1}^d - \frac{1}{\sigma_d^{2(m-1)}r} X_{\lceil d^{2\alpha}s \rceil, 1}^d \right) \mathbb{1}_{\mathcal{A}_{\lceil d^{2\alpha}s \rceil}^d}. \end{aligned}$$

Using the boundedness of moments of Gaussian distributions and of $X_{\lfloor d^{2\alpha}t \rfloor, 1}^d, X_{\lfloor d^{2\alpha}s \rfloor, 1}^d$, we have

$$\mathbb{E}[(L_t^d - L_s^d)^4] \leq C \left((t-s)^2 + \mathbb{E} \left[\left(X_{\lfloor d^{2\alpha}t \rfloor, 1}^d - X_{\lfloor d^{2\alpha}s \rfloor, 1}^d \right)^4 \right] \right). \quad (28)$$

Putting (26), (27) and (28) together and using Lemma 1 below we obtain

$$\begin{aligned} \mathbb{E}[(L_t^d - L_s^d)^4] &\leq C \left((t-s)^2 + \sum_{p=2}^4 \frac{(\lfloor d^{2\alpha}t \rfloor - \lfloor d^{2\alpha}s \rfloor)^p}{d^{2\alpha p}} \right) \\ &\leq C(t-s)^2 + C \sum_{p=2}^4 \frac{d^{2\alpha p} (t-s)^p}{d^{2\alpha p}} \leq C(2+t+t^2)(t-s)^2, \end{aligned}$$

which concludes the proof. $\square$

We are now ready to state and prove Lemma 1:

Lemma 1. *There exists $C > 0$ such that for any $k_1, k_2 \in \mathbb{N}$ with $0 \leq k_1 < k_2$,*

$$\mathbb{E}[(X_{k_2, 1}^d - X_{k_1, 1}^d)^4] \leq C \sum_{p=2}^4 \frac{(k_2 - k_1)^p}{d^{2\alpha p}},$$

where $\alpha = 1/3$.

Proof. Recalling the definition of the proposal in (24) and the definition of b_d in (20) we can write

$$\mathbb{E}[(X_{k_2, 1}^d - X_{k_1, 1}^d)^4] = \mathbb{E} \left[\left(\sum_{k=k_1+1}^{k_2} \sigma_d b_d(X_{k-1, 1}^d, Z_{k, 1}^d) \mathbb{1}_{\mathcal{A}_k^d} \right)^4 \right].$$

Then, we expand all acceptance or rejection terms between k_1 and k_2 and use Hölder's inequality to obtain

$$\begin{aligned} \mathbb{E}[(X_{k_2, 1}^d - X_{k_1, 1}^d)^4] &\leq C \left\{ \sigma_d^4 \mathbb{E} \left[\left(\sum_{k=k_1+1}^{k_2} b_d(X_{k-1, 1}^d, Z_{k, 1}^d) \right)^4 \right] \right. \\ &\quad \left. + \sigma_d^4 \mathbb{E} \left[\left(\sum_{k=k_1+1}^{k_2} b_d(X_{k-1, 1}^d, Z_{k, 1}^d) \mathbb{1}_{(\mathcal{A}_k^d)^c} \right)^4 \right] \right\}. \end{aligned}$$

Using again Hölder's inequality, for the first term we have

$$\begin{aligned}
\mathbb{E} \left[\left(\sum_{k=k_1+1}^{k_2} b_d(X_{k-1,1}^d, Z_{k,1}^d) \right)^4 \right] &\leq C \left\{ \mathbb{E} \left[\left(\sum_{k=k_1+1}^{k_2} Z_{k,1}^d \right)^4 \right] \right. \\
&\quad + \frac{\sigma_d^4}{2^4} \mathbb{E} \left[\left(\sum_{k=k_1+1}^{k_2} \operatorname{sgn}(X_{k-1,1}^d) \mathbb{1}_{\{|X_{k-1,1}^d| \geq \sigma_d^{2m} r/2\}} \right)^4 \right] \\
&\quad \left. + \mathbb{E} \left[\left(\sum_{k=k_1+1}^{k_2} \frac{1}{\sigma_d^{2m-1} r} X_{k-1,1}^d \mathbb{1}_{\{|X_{k-1,1}^d| < \sigma_d^{2m} r/2\}} \right)^4 \right] \right\} \\
&\leq C \left[3(k_2 - k_1)^2 + \frac{2\sigma_d^4}{2^4} (k_2 - k_1)^4 \right], \tag{29}
\end{aligned}$$

where the last line follows using the moments of $Z_{k,1}^d$ and the boundedness of $X_{k-1,1}^d$ in the set $\{|X_{k-1,1}^d| < \sigma_d^{2m} r/2\}$.

Using a Binomial expansion of the rejection term, we obtain

$$\mathbb{E} \left[\left(\sum_{k=k_1+1}^{k_2} b_d(X_{k-1,1}^d, Z_{k,1}^d) \mathbb{1}_{(\mathbf{A}_k^d)^c} \right)^4 \right] = \sum \mathbb{E} \left[\prod_{i=1}^4 b_d(X_{m_i-1,1}^d, Z_{m_i,1}^d) \mathbb{1}_{(\mathbf{A}_{m_i}^d)^c} \right], \tag{30}$$

where the sum is over the quadruplets $(m_i)_{1 \leq i \leq 4}$ with $m_i \in \{k_1 + 1, \dots, k_2\}$.

We separate the terms in the sum according to their cardinality, let us denote, for $j \in \{1, \dots, 4\}$,

$$\mathcal{I}_j = \left\{ (m_1, \dots, m_4) \in \{k_1 + 1, \dots, k_2\}^4 : \# \{m_1, \dots, m_4\} = j \right\};$$

and define, for any $(m_1, \dots, m_4) \in \{k_1 + 1, \dots, k_2\}^4$, $\tilde{X}_0^d = X_0^d$ and for any $i \in \{1, \dots, d\}$,

$$\tilde{X}_{k+1,i}^d = \tilde{X}_{k,i}^d + \mathbb{1}_{\{m_1-1, \dots, m_4-1\}^c}(k) \mathbb{1}_{\tilde{\mathbf{A}}_{k+1}^d} \sigma_d b_d(\tilde{X}_{k,i}^d, Z_{k+1,i}^d),$$

where

$$\tilde{\mathbf{A}}_{k+1}^d = \left\{ U_{k+1} \leq \exp \left[\sum_{i=1}^d \phi_d(\tilde{X}_{k,i}^d, Z_{k+1,i}^d) \right] \right\}, \tag{31}$$

and ϕ_d in (21). Denote by $\mathcal{F}$ the σ -algebra generated by the process $(\tilde{X}_k^d)_{k \geq 0}$ and observe that on the event $\bigcap_{j=1}^4 (\mathbf{A}_{m_j}^d)^c$, X_k^d is equal to $\tilde{X}_k^d$.

We consider now the terms in the sum (30).

(i) If $(m_1, \dots, m_4) \in \mathcal{I}_4$, then the m_i s are all distinct and

$$\mathbb{E} \left[\prod_{j=1}^4 b_d(X_{m_j-1,1}^d, Z_{m_j,1}^d) \mathbb{1}_{(\mathbf{A}_{m_j}^d)^c} \middle| \mathcal{F} \right] = \mathbb{E} \left[\prod_{j=1}^4 b_d(\tilde{X}_{m_j-1,1}^d, Z_{m_j,1}^d) \mathbb{1}_{(\tilde{\mathbf{A}}_{m_j}^d)^c} \middle| \mathcal{F} \right].$$

However, $\{b_d(\tilde{X}_{m_j-1,1}^d, Z_{m_j,1}^d) \mathbb{1}_{(\tilde{\mathbf{A}}_{m_j}^d)^c}\}_{j=1,\dots,4}$ are independent conditionally on $\mathcal{F}$. Thus,

$$\begin{aligned} \mathbb{E} \left[\prod_{j=1}^4 b_d \left(\tilde{X}_{m_j-1,1}^d, Z_{m_j,1}^d \right) \mathbb{1}_{(\tilde{\mathbf{A}}_{m_j}^d)^c} \middle| \mathcal{F} \right] &= \prod_{j=1}^4 \mathbb{E} \left[b_d \left(\tilde{X}_{m_j-1,1}^d, Z_{m_j,1}^d \right) \mathbb{1}_{(\tilde{\mathbf{A}}_{m_j}^d)^c} \middle| \mathcal{F} \right] \\ &= \prod_{j=1}^4 \mathbb{E} \left[b_d \left(\tilde{X}_{m_j-1,1}^d, Z_{m_j,1}^d \right) \times \left(1 - \exp \left(\sum_{i=1}^d \phi_d \left(\tilde{X}_{m_j-1,i}^d, Z_{m_j,i}^d \right) \right) \right) \middle| \mathcal{F} \right]_+, \end{aligned}$$

by integrating the uniform variables U_{m_j} in (31).

Recalling the definition of b_d in (20), we can bound the expectation above with

$$\begin{aligned} &\left| \mathbb{E} \left[b_d \left(\tilde{X}_{m_j-1,1}^d, Z_{m_j,1}^d \right) \left\{ 1 - \exp \left(\sum_{i=1}^d \phi_d \left(\tilde{X}_{m_j-1,i}^d, Z_{m_j,i}^d \right) \right) \right\} \middle| \mathcal{F} \right] \right| \quad (32) \\ &\leq \left| \mathbb{E} \left[\left(\frac{\sigma_d}{2} \operatorname{sgn} \left(\tilde{X}_{m_j-1,1}^d \right) \mathbb{1}_{\left\{ |\tilde{X}_{m_j-1,1}^d| \geq \sigma_d^{2m} r/2 \right\}} \right. \right. \right. \\ &\quad \left. \left. - \frac{1}{\sigma_d^{2m-1} r} \tilde{X}_{m_j-1,1}^d \mathbb{1}_{\left\{ |\tilde{X}_{m_j-1,1}^d| < \sigma_d^{2m} r/2 \right\}} \right) \right. \\ &\quad \left. \times \left\{ 1 - \exp \left(\sum_{i=1}^d \phi_d \left(\tilde{X}_{m_j-1,i}^d, Z_{m_j,i}^d \right) \right) \right\} \middle| \mathcal{F} \right] \right| \\ &\quad + \left| \mathbb{E} \left[Z_{m_j,1}^d \left\{ 1 - \exp \left(\sum_{i=1}^d \phi_d \left(\tilde{X}_{m_j-1,i}^d, Z_{m_j,i}^d \right) \right) \right\} \middle| \mathcal{F} \right] \right|. \end{aligned}$$

For the first one, we use the boundedness of the sgn function and of $\tilde{X}_{m_j-1,1}^d$ in the set $\{|\tilde{X}_{m_j-1,1}^d| \leq \sigma_d^{2m} r/2\}$ to obtain

$$\begin{aligned} &\left| \mathbb{E} \left[\left(\frac{\sigma_d}{2} \operatorname{sgn} \left(\tilde{X}_{m_j-1,1}^d \right) \mathbb{1}_{\left\{ |\tilde{X}_{m_j-1,1}^d| \geq \sigma_d^{2m} r/2 \right\}} - \frac{\tilde{X}_{m_j-1,1}^d}{\sigma_d^{2m-1} r} \mathbb{1}_{\left\{ |\tilde{X}_{m_j-1,1}^d| < \sigma_d^{2m} r/2 \right\}} \right) \right. \right. \\ &\quad \left. \times \left\{ 1 - \exp \left(\sum_{i=1}^d \phi_d \left(\tilde{X}_{m_j-1,i}^d, Z_{m_j,i}^d \right) \right) \right\} \middle| \mathcal{F} \right] \right| \\ &\leq \frac{\sigma_d}{2} \mathbb{E} \left[\left| \left\{ 1 - \exp \left(\sum_{i=1}^d \phi_d \left(\tilde{X}_{m_j-1,i}^d, Z_{m_j,i}^d \right) \right) \right\} \right| \middle| \mathcal{F} \right] \leq \frac{\sigma_d}{2}. \quad (33) \end{aligned}$$

We can write the second term as

$$\begin{aligned} &\mathbb{E} \left[Z_{m_j,1}^d \left(1 - \exp \left(\sum_{i=1}^d \phi_d \left(\tilde{X}_{m_j-1,i}^d, Z_{m_j,i}^d \right) \right) \right) \middle| \mathcal{F} \right] \\ &= \mathbb{E} \left[\mathcal{G} \left(\tilde{X}_{m_j-1,1}^d, \sum_{i=2}^d \phi_d \left(\tilde{X}_{m_j-1,i}^d, Z_{m_j,i}^d \right) \right) \middle| \mathcal{F} \right], \end{aligned}$$

where we define $\mathcal{G}(a, b) = \mathbb{E} [Z (1 - \exp(\phi_d(a, Z) + b))_+]]$ with Z a standard Gaussian. Because the function $x \mapsto (1 - \exp(x))_+$ is 1-Lipschitz, we have, using Cauchy-Schwarz and Lemma 3 in Appendix D.2,

$$\begin{aligned} |\mathbb{E} [Z (1 - \exp(\phi_d(a, Z) + b))_+] - \mathbb{E} [Z (1 - \exp(b))_+] | &\leq \mathbb{E} [|Z| |\phi_d(a, Z)|] \\ &\leq \mathbb{E} [Z^2]^{1/2} \mathbb{E} [\phi_d(a, Z)^2]^{1/2} \\ &\leq \mathbb{E} [\phi_d(a, Z)^2]^{1/2} \\ &\leq Cd^{-\alpha}. \end{aligned}$$

However, $\mathbb{E} [Z (1 - \exp(b))_+] = \mathbb{E} [Z] (1 - \exp(b))_+ = 0$, and therefore

$$\left| \mathbb{E} \left[\mathcal{G} \left(\tilde{X}_{m_j-1,1}^d, \sum_{i=2}^d \phi_d \left(\tilde{X}_{m_j-1,i}^d, Z_{m_j,i}^d \right) \right) \middle| \mathcal{F} \right] \right| \leq Cd^{-\alpha}. \quad (34)$$

Combining equations (32), (33) and (34) and recalling that $\sigma_d = \ell d^{-\alpha}$, we have

$$\left| \mathbb{E} \left[b_d \left(\tilde{X}_{m_j-1,1}^d, Z_{m_j,1}^d \right) \left\{ 1 - \exp \left(\sum_{i=1}^d \phi_d \left(\tilde{X}_{m_j-1,i}^d, Z_{m_j,i}^d \right) \right) \right\}_+ \middle| \mathcal{F} \right] \right| \leq Cd^{-\alpha}, \quad (35)$$

from which follows that

$$\begin{aligned} \sum_{(m_1, \dots, m_4) \in \mathcal{I}_4} \left| \mathbb{E} \left[\prod_{i=1}^4 b_d \left(X_{m_i-1,1}^d, Z_{m_i,1}^d \right) \mathbb{1}_{(\Lambda_{m_i}^d)^c} \right] \right| &\leq \sum_{(m_1, \dots, m_4) \in \mathcal{I}_4} \mathbb{E} \left[\prod_{j=1}^4 \frac{C}{d^\alpha} \right] \\ &\leq \binom{k_2 - k_1}{4} \frac{C}{d^{4\alpha}} \leq C \frac{(k_2 - k_1)^4}{d^{4\alpha}}, \quad (36) \end{aligned}$$

using that $|\mathcal{I}_4| = \binom{k_2 - k_1}{4}$.

- (ii) If $(m_1, \dots, m_4) \in \mathcal{I}_3$, only three of the m_i s take distinct values; proceeding as in case (i), we have

$$\begin{aligned} &\left| \mathbb{E} \left[\prod_{j=1}^3 b_d \left(X_{m_j-1,1}^d, Z_{m_j,1}^d \right)^{1+\delta_{1,j}} \mathbb{1}_{(\Lambda_{m_j}^d)^c} \middle| \mathcal{F} \right] \right| \\ &= \prod_{j=1}^3 \left| \mathbb{E} \left[b_d \left(\tilde{X}_{m_j-1,1}^d, Z_{m_j,1}^d \right)^{1+\delta_{1,j}} \left\{ 1 - \exp \left(\sum_{i=1}^d \phi_d \left(\tilde{X}_{m_j-1,i}^d, Z_{m_j,i}^d \right) \right) \right\}_+ \middle| \mathcal{F} \right] \right|, \end{aligned}$$

where $\delta_{1,j}$ denotes a Dirac's delta. For the terms $j \neq 1$, we use (35), while for the term $j = 1$

we bound the indicator function by 1 to obtain

$$\begin{aligned}
& \left| \mathbb{E} \left[\prod_{j=1}^3 b_d \left(X_{m_j-1,1}^d, Z_{m_j,1}^d \right)^{1+\delta_{1,j}} \mathbb{1}_{(\mathbb{A}_{m_j}^d)^c} \middle| \mathcal{F} \right] \right| \\
& \leq \left| \mathbb{E} \left[b_d \left(\tilde{X}_{m_1-1,1}^d, Z_{m_1,1}^d \right)^2 \middle| \mathcal{F} \right] \right| \prod_{j=2}^3 \frac{C}{d^\alpha} \\
& \leq \left(3 + \frac{2\sigma_d^2}{2^2 d^{2\alpha}} \right) \frac{C^2}{d^{2\alpha}} \leq C \frac{1}{d^{2\alpha}},
\end{aligned}$$

where the second-to-last inequality follows using the same approach taken for (29) and recalling that $\sigma_d = \ell d^{-\alpha}$. Hence,

$$\begin{aligned}
& \sum_{(m_1, \dots, m_4) \in \mathcal{I}_3} \left| \mathbb{E} \left[\prod_{i=1}^4 b_d \left(X_{m_i-1,1}^d, Z_{m_i,1}^d \right) \mathbb{1}_{(\mathbb{A}_{m_i}^d)^c} \right] \right| \\
& \leq C \binom{k_2 - k_1}{3} \frac{1}{d^{2\alpha}} \leq C \frac{(k_2 - k_1)^3}{d^{2\alpha}}.
\end{aligned} \tag{37}$$

- (iii) If $(m_1, \dots, m_4) \in \mathcal{I}_2$, we have two different cases: the m_i s take the two values twice or three m_i s have the same value. For the first one, we have, bounding the indicator function with 1,

$$\begin{aligned}
& \mathbb{E} \left[\mathbb{E} \left[\prod_{j=1}^2 b_d \left(X_{m_j-1,1}^d, Z_{m_j,1}^d \right)^2 \mathbb{1}_{(\mathbb{A}_{m_j}^d)^c} \middle| \mathcal{F} \right] \right] \\
& \leq \mathbb{E} \left[\prod_{j=1}^2 \mathbb{E} \left[b_d \left(\tilde{X}_{m_j-1,1}^d, Z_{m_j,1}^d \right)^2 \middle| \mathcal{F} \right] \right].
\end{aligned}$$

Since, conditionally on $\mathcal{F}$, the random variables inside the expectation are normals with bounded mean and variance 1, we have, using the same approach taken for (29),

$$\mathbb{E} \left[\prod_{j=1}^2 \mathbb{E} \left[b_d \left(\tilde{X}_{m_j-1,1}^d, Z_{m_j,1}^d \right)^2 \middle| \mathcal{F} \right] \right] \leq \left(1 + \frac{2\sigma_d^2}{2^2} \right)^2 \leq C.$$

The second case follows similarly

$$\begin{aligned}
& \left| \mathbb{E} \left[\mathbb{E} \left[\prod_{j=1}^2 b_d \left(X_{m_j-1,1}^d, Z_{m_j,1}^d \right)^{1+2\delta_{1,j}} \mathbb{1}_{(\mathbb{A}_{m_j}^d)^c} \middle| \mathcal{F} \right] \right] \right| \\
& \leq \mathbb{E} \left[\mathbb{E} \left[\prod_{j=1}^2 \left| b_d \left(\tilde{X}_{m_j-1,1}^d, Z_{m_j,1}^d \right) \right|^{1+2\delta_{1,j}} \middle| \mathcal{F} \right] \right] \leq C,
\end{aligned}$$

where $\delta_{1,j}$ denotes a Dirac's delta. Therefore,

$$\begin{aligned} & \sum_{(m_1, \dots, m_4) \in \mathcal{I}_2} \left| \mathbb{E} \left[\prod_{i=1}^4 \{b_d(X_{m_i-1,1}^d, Z_{m_i,1}^d) \mathbb{1}_{(\mathbb{A}_{m_i}^d)^c}\} \right] \right| \\ & \leq C \left(\binom{4}{2} + \binom{4}{3} \right) \binom{k_2 - k_1}{2} \leq C(k_2 - k_1)^2. \end{aligned} \quad (38)$$

(iv) If $(m_1, \dots, m_4) \in \mathcal{I}_1$ (i.e. all m_i s take the same value), we bound the indicator function by 1 and, using the same approach taken for (29), we find

$$\mathbb{E} \left[b_d(X_{m_1-1,1}^d, Z_{m_1,1}^d)^4 \mathbb{1}_{(\mathbb{A}_{m_1}^d)^c} \right] \leq C \left(3 + \frac{2\sigma_d^4}{2^4} \right) \leq C,$$

since $\sigma_d = \ell d^{-\alpha}$ and $d \in \mathbb{N}$. Hence,

$$\sum_{(m_1, \dots, m_4) \in \mathcal{I}_1} \left| \mathbb{E} \left[\prod_{i=1}^4 b_d(X_{m_i-1,1}^d, Z_{m_i,1}^d) \mathbb{1}_{(\mathbb{A}_{m_i}^d)^c} \right] \right| \leq C \binom{k_2 - k_1}{1} = C(k_2 - k_1). \quad (39)$$

The result follows combining (36), (37), (38) and (39) in (30). $\square$

6.3 Proof of Proposition 2

We start by proving the following lemma.

Lemma 2. *Let ν be a limit point of the sequence of laws $(\nu_d)_{d \geq 1}$ of $\{(L_t^d)_{t \geq 0} : d \in \mathbb{N}^*\}$. Then for any $t \geq 0$, the pushforward measure of ν by W_t is $\pi^L(dx) = \exp(-|x|)dx/2$.*

Proof. Using (25), we have

$$\begin{aligned} & \mathbb{E} \left[\left| L_t^d - X_{\lfloor d^{2\alpha} t \rfloor, 1}^d \right| \right] \\ & \leq \mathbb{E} \left[\left| (d^{2\alpha} t - \lfloor d^{2\alpha} t \rfloor) \left[\sigma_d Z_{\lfloor d^{2\alpha} t \rfloor, 1}^d - \frac{\sigma_d^2}{2} \operatorname{sgn}(X_{\lfloor d^{2\alpha} t \rfloor, 1}^d) \right] \mathbb{1}_{\{|X_{\lfloor d^{2\alpha} t \rfloor, 1}^d| \geq \sigma_d^{2m} r/2\}} \right| \right] \\ & + \mathbb{E} \left[\left| (d^{2\alpha} t - \lfloor d^{2\alpha} t \rfloor) \left[\sigma_d Z_{\lfloor d^{2\alpha} t \rfloor, 1}^d - \frac{1}{\sigma_d^{2(m-1)} r} X_{\lfloor d^{2\alpha} t \rfloor, 1}^d \right] \mathbb{1}_{\{|X_{\lfloor d^{2\alpha} t \rfloor, 1}^d| < \sigma_d^{2m} r/2\}} \right| \right] \\ & \leq (d^{2\alpha} t - \lfloor d^{2\alpha} t \rfloor) \left(\sigma_d \mathbb{E} \left[|Z_{\lfloor d^{2\alpha} t \rfloor, 1}^d| \right] + \frac{\sigma_d^2}{2} \mathbb{E} \left[|\operatorname{sgn}(X_{\lfloor d^{2\alpha} t \rfloor, 1}^d)| \right] \right. \\ & \quad \left. + \frac{1}{\sigma_d^{2(m-1)} r} \mathbb{E} \left[|X_{\lfloor d^{2\alpha} t \rfloor, 1}^d| \mathbb{1}_{\{|X_{\lfloor d^{2\alpha} t \rfloor, 1}^d| < \sigma_d^{2m} r/2\}} \right] \right) \\ & \leq (d^{2\alpha} t - \lfloor d^{2\alpha} t \rfloor) \left(\frac{\ell}{d^\alpha} \mathbb{E} \left[(Z_{\lfloor d^{2\alpha} t \rfloor, 1}^d)^2 \right]^{1/2} + \frac{\ell^2}{2d^{2\alpha}} + \frac{1}{\sigma_d^{2(m-1)} r} \mathbb{E} \left[\frac{\sigma_d^{2m} r}{2} \right] \right) \\ & \leq (d^{2\alpha} t - \lfloor d^{2\alpha} t \rfloor) \left(\frac{\ell}{d^\alpha} + \frac{\ell^2}{2d^{2\alpha}} + \frac{\ell^2}{2d^{2\alpha}} \right) \leq \frac{C}{d^\alpha}, \end{aligned}$$

where we used Cauchy-Schwarz inequality and the fact that the moments of $Z_{[d^{2\alpha}t],1}^d$ are bounded. The above guarantees that,

$$\lim_{d \rightarrow \infty} \mathbb{E} \left[\left| L_t^d - X_{[d^{2\alpha}t],1}^d \right| \right] = 0 .$$

As $(\nu_d)_{d \geq 1}$ converges weakly towards ν , for any Lipschitz bounded function $\psi : \mathbb{R} \rightarrow \mathbb{R}$,

$$\lim_{d \rightarrow \infty} \mathbb{E} \left[\psi \left(X_{[d^{2\alpha}t],1}^d \right) \right] = \lim_{d \rightarrow \infty} \mathbb{E} \left[\psi \left(L_t^d \right) \right] = \mathbb{E}^\nu \left[\psi(W_t) \right] .$$

The result follows since $X_{[d^{2\alpha}t],1}^d$ is distributed according to $\pi^L(dx) = \exp(-|x|)dx/2$ for any $t \geq 0$ and $d \in \mathbb{N}$. $\square$

We are now ready to prove Proposition 2:

Proof of Proposition 2. Let ν be a limit point of $(\nu_d)_{d \geq 1}$. We start by showing that if for any $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$, $m \in \mathbb{N}$, any bounded and continuous mapping $\rho : \mathbb{R}^m \rightarrow \mathbb{R}$ and any $0 \leq t_1 \leq \dots \leq t_m \leq s \leq t$, ν satisfies

$$\mathbb{E}^\nu \left[\left(V(W_t) - V(W_s) - \int_s^t LV(W_u) du \right) \rho(W_{t_1}, \dots, W_{t_m}) \right] = 0 , \quad (40)$$

then ν is a solution to the martingale problem associated with L .

Let $\mathfrak{F}_s$ denote the σ -algebra generated by

$$\{\rho(W_{t_1}, \dots, W_{t_m}) : m \in \mathbb{N}, \rho : \mathbb{R}^m \rightarrow \mathbb{R} \text{ bounded and continuous, and } 0 \leq t_1 \leq \dots \leq t_m \leq s\} .$$

Then,

$$\mathbb{E}^\nu \left[V(W_t) - V(W_s) - \int_s^t LV(W_u) du \middle| \mathfrak{F}_s \right] = 0 ,$$

showing that the process

$$\left(V(W_t) - V(W_0) - \int_0^t LV(W_u) du \right)_{t \geq 0}$$

is a martingale w.r.t. ν and the filtration $(\mathfrak{F}_t)_{t \geq 0}$.

To prove (40), it is enough to show that for any $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$, $m \in \mathbb{N}$ and any bounded and continuous mapping $\rho : \mathbb{R}^m \rightarrow \mathbb{R}$ and any $0 \leq t_1 \leq \dots \leq t_m \leq s \leq t$, the mapping

$$\Psi_{s,t} : w \mapsto \left(V(w_t) - V(w_s) - \int_s^t LV(w_u) du \right) \rho(w_{t_1}, \dots, w_{t_m}) ,$$

is continuous on a ν -almost sure subset of $C(\mathbb{R}_+, \mathbb{R})$. Let

$$\mathbf{W} = \{w \in C(\mathbb{R}_+, \mathbb{R}) : w_u \neq 0 \text{ for almost any } u \in [s, t]\} .$$

Since $w \in \mathbf{W}^c$ if and only if $\int_s^t \mathbb{1}_{\{0\}}(w_u) du > 0$, using Lemma 2 and the Fubini–Tonelli’s theorem,

$$\mathbb{E}^\nu \left[\int_s^t \mathbb{1}_{\{0\}}(W_u) du \right] = \int_s^t \mathbb{E}^\nu [\mathbb{1}_{\{0\}}(W_u)] du = \int_s^t \pi^L(\{0\}) du = 0 ,$$

and we have that $\nu(\mathbf{W}^c) = 0$.

Since $w \mapsto w_u$ is continuous for any $u \geq 0$, so are $w \mapsto V(w_u)$ and $w \mapsto \rho(w_{t_1}, \dots, w_{t_m})$. Thus, it is enough to prove that the mapping $w \mapsto \int_s^t LV(w_u)du$ is continuous. Let $(w^n)_{n \geq 0}$ be a sequence in $C(\mathbb{R}_+, \mathbb{R})$ that converges to $w \in \mathbf{W}$ in the uniform topology on compact sets. Let u be such that $w_u \neq 0$, therefore, since the sgn function is continuous in a neighbourhood of w_u , $\lim_{n \rightarrow \infty} LV(w_u^n) = LV(w_u)$, thus $\lim_{n \rightarrow \infty} LV(w_u^n) = LV(w_u)$ for almost any $u \in [s, t]$. Finally, using the boundedness of the sequence $(LV(w_u^n))_{n \geq 0}$ and Lebesgue's dominated convergence theorem,

$$\lim_{n \rightarrow \infty} \int_s^t LV(w_u^n)du = \int_s^t LV(w_u)du ,$$

which proves that the mappings $\Psi_{s,t}$ are continuous on $\mathbf{W}$. $\square$

6.4 Proof of Theorem 3

Let us introduce, for any $n \in \mathbb{N}$, $\mathcal{F}_{n,1}^d = \sigma(\{X_{k,1}^d, 0 \leq k \leq n\})$, the σ -algebra generated by the first components of $\{X_k^d \mid 0 \leq k \leq n\}$. We also introduce for any $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$

$$\begin{aligned} M_n^d(V) &= \frac{\ell}{d^\alpha} \sum_{k=0}^{n-1} V'(X_{k,1}^d) \\ &\quad \times \left(b_d(X_{k,1}^d, Z_{k+1,1}^d) \mathbb{1}_{\mathcal{A}_{k+1}^d} - \mathbb{E} \left[b_d(X_{k,1}^d, Z_{k+1,1}^d) \mathbb{1}_{\mathcal{A}_{k+1}^d} \middle| \mathcal{F}_{k,1}^d \right] \right) \\ &\quad + \frac{\ell^2}{2d^{2\alpha}} \sum_{k=0}^{n-1} V''(X_{k,1}^d) \\ &\quad \times \left(b_d(X_{k,1}^d, Z_{k+1,1}^d)^2 \mathbb{1}_{\mathcal{A}_{k+1}^d} - \mathbb{E} \left[b_d(X_{k,1}^d, Z_{k+1,1}^d)^2 \mathbb{1}_{\mathcal{A}_{k+1}^d} \middle| \mathcal{F}_{k,1}^d \right] \right) . \end{aligned} \quad (41)$$

where b_d is defined in (20).

The proof of Theorem 3 follows using the sufficient condition in Proposition 2, the tightness of the sequence $(\nu_d)_{d \geq 1}$ established in Proposition 1 and Proposition 3 below.

Proof. Using Proposition 1, Proposition 2 and Proposition 3 below, it is enough to show that for any $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$, $m \geq 1$, any $0 \leq t_1 \leq \dots \leq t_m \leq s \leq t$ and any bounded and continuous mapping $\rho : \mathbb{R}^m \rightarrow \mathbb{R}$,

$$\lim_{d \rightarrow \infty} \mathbb{E} \left[\left(M_{\lceil d^{2\alpha} t \rceil}^d(V) - M_{\lceil d^{2\alpha} s \rceil}^d(V) \right) \rho(L_{t_1}^d, \dots, L_{t_m}^d) \right] = 0 ,$$

where, for any $n \geq 1$, $M_n^d(V)$ is given by (41). However, this is straightforwardly obtained by taking successively the conditional expectations with respect to $\mathcal{F}_{k,1}^d$ for $k = \lceil d^{2\alpha} t \rceil, \dots, \lceil d^{2\alpha} s \rceil$. $\square$

Proposition 3. *For any $0 \leq s \leq t$, $V \in C_c(\mathbb{R}, \mathbb{R})$ we have*

$$\lim_{d \rightarrow \infty} \mathbb{E} \left[\left| V(L_t^d) - V(L_s^d) - \int_s^t LV(L_u^d) du - \left(M_{\lceil d^{2\alpha} t \rceil}^d(V) - M_{\lceil d^{2\alpha} s \rceil}^d(V) \right) \right| \right] = 0 , \quad (42)$$

where $(L_t^d)_{t \geq 0}$ is defined in (11).

Proof. The process $(L_t^d)_{t \geq 0}$ is piecewise linear, thus it has finite variation. For any $\tau \geq 0$, we define

$$dL_\tau^d = d^{2\alpha} \sigma_d b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right) \mathbb{1}_{A_{\lceil d^{2\alpha} \tau \rceil}^d} d\tau .$$

Since $\sigma_d = \ell d^{-\alpha}$ with $\alpha = 1/3$ and using the fundamental theorem of calculus for piecewise C^1 maps

$$V(L_t^d) - V(L_s^d) = \ell d^\alpha \int_s^t V'(L_\tau^d) b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right) \mathbb{1}_{A_{\lceil d^{2\alpha} \tau \rceil}^d} d\tau , \quad (43)$$

where b_d is defined in (20). A Taylor expansion of V' with Lagrange remainder about $X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d$ gives

$$\begin{aligned} V'(L_\tau^d) &= V' \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) \\ &\quad + \frac{\ell}{d^\alpha} (d^{2\alpha} \tau - \lfloor d^{2\alpha} \tau \rfloor) V'' \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right) \mathbb{1}_{A_{\lceil d^{2\alpha} \tau \rceil}^d} \\ &\quad + \frac{\ell^2}{2d^{2\alpha}} (d^{2\alpha} \tau - \lfloor d^{2\alpha} \tau \rfloor)^2 V^{(3)}(\chi_\tau) b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right) \mathbb{1}_{A_{\lceil d^{2\alpha} \tau \rceil}^d} , \end{aligned}$$

where for any point $\tau \in [s, t]$, there exists $\chi_\tau \in [X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, L_\tau^d]$. Substituting the above into (43) we obtain

$$\begin{aligned} V(L_t^d) - V(L_s^d) &= \ell d^\alpha \int_s^t V' \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right) \mathbb{1}_{A_{\lceil d^{2\alpha} \tau \rceil}^d} d\tau \\ &\quad + \ell^2 \int_s^t (d^{2\alpha} \tau - \lfloor d^{2\alpha} \tau \rfloor) V'' \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right)^2 \mathbb{1}_{A_{\lceil d^{2\alpha} \tau \rceil}^d} d\tau \\ &\quad + \frac{\ell^3}{2d^\alpha} \int_s^t (d^{2\alpha} \tau - \lfloor d^{2\alpha} \tau \rfloor)^2 V^{(3)}(\chi_\tau) b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right)^3 \mathbb{1}_{A_{\lceil d^{2\alpha} \tau \rceil}^d} d\tau . \end{aligned} \quad (44)$$

Since $V^{(3)}$ is bounded, using Fubini-Tonelli's theorem and recalling the definition of b_d in (20), we have that

$$\begin{aligned} &\frac{\ell^3}{2d^\alpha} \mathbb{E} \left[\left| \int_s^t (d^{2\alpha} \tau - \lfloor d^{2\alpha} \tau \rfloor)^2 V^{(3)}(\chi_\tau) b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right)^3 \mathbb{1}_{A_{\lceil d^{2\alpha} \tau \rceil}^d} d\tau \right| \right] \\ &\leq C \frac{\ell^3}{2d^\alpha} \int_s^t \mathbb{E} \left[\left(\left| Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right| + \frac{\ell}{2d^\alpha} \right)^3 \right] d\tau \xrightarrow{d \rightarrow \infty} 0 , \end{aligned}$$

since the moments of $Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d$ are bounded.

For the second term in (44), we observe that most of the integrand is piecewise constant since the process $X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d$ evolves in discrete time. Then, for any integer $d^{2\alpha} s \leq k \leq d^{2\alpha} t - 1$,

$$\begin{aligned} &\int_{k/d^{2\alpha}}^{(k+1)/d^{2\alpha}} (d^{2\alpha} \tau - \lfloor d^{2\alpha} \tau \rfloor) V'' \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right)^2 \mathbb{1}_{A_{\lceil d^{2\alpha} \tau \rceil}^d} d\tau \\ &= \frac{1}{2d^{2\alpha}} V'' \left(X_{k, 1}^d \right) b_d \left(X_{k, 1}^d, Z_{k+1, 1}^d \right)^2 \mathbb{1}_{A_{k+1}^d} \\ &= \frac{1}{2} \int_{k/d^{2\alpha}}^{(k+1)/d^{2\alpha}} V'' \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right)^2 \mathbb{1}_{A_{\lceil d^{2\alpha} \tau \rceil}^d} d\tau . \end{aligned}$$

Thus, we can write

$$\begin{aligned} I &= \int_s^t (d^{2\alpha}\tau - \lfloor d^{2\alpha}\tau \rfloor) V'' \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) b_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d, Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right)^2 \mathbb{1}_{\mathbf{A}_{\lfloor d^{2\alpha}\tau \rfloor}^d} d\tau \\ &= I_1 + I_2, \end{aligned}$$

where we define

$$I_2 := \frac{1}{2} \int_s^t V'' \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) b_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d, Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right)^2 \mathbb{1}_{\mathbf{A}_{\lfloor d^{2\alpha}\tau \rfloor}^d} d\tau,$$

and

$$\begin{aligned} I_1 := \left[\int_s^{\lceil d^{2\alpha}s \rceil / d^{2\alpha}} + \int_{\lfloor d^{2\alpha}t \rfloor / d^{2\alpha}}^t \right] &\left(d^{2\alpha}\tau - \lfloor d^{2\alpha}\tau \rfloor - \frac{1}{2} \right) V'' \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) \\ &\times b_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d, Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right)^2 \mathbb{1}_{\mathbf{A}_{\lfloor d^{2\alpha}\tau \rfloor}^d} d\tau. \end{aligned}$$

In addition, we have

$$\begin{aligned} I_1 &= \frac{1}{2d^{2\alpha}} (d^{2\alpha}s - \lfloor d^{2\alpha}s \rfloor) (\lceil d^{2\alpha}s \rceil - d^{2\alpha}s) V'' \left(X_{\lfloor d^{2\alpha}s \rfloor, 1}^d \right) b_d \left(X_{\lfloor d^{2\alpha}s \rfloor, 1}^d, Z_{\lfloor d^{2\alpha}s \rfloor, 1}^d \right)^2 \mathbb{1}_{\mathbf{A}_{\lfloor d^{2\alpha}s \rfloor}^d} \\ &\quad + \frac{1}{2d^{2\alpha}} (d^{2\alpha}t - \lfloor d^{2\alpha}t \rfloor) (\lceil d^{2\alpha}t \rceil - d^{2\alpha}t) V'' \left(X_{\lfloor d^{2\alpha}t \rfloor, 1}^d \right) b_d \left(X_{\lfloor d^{2\alpha}t \rfloor, 1}^d, Z_{\lfloor d^{2\alpha}t \rfloor, 1}^d \right)^2 \mathbb{1}_{\mathbf{A}_{\lfloor d^{2\alpha}t \rfloor}^d}, \end{aligned}$$

and, since V'' and the moments of $Z_{\lfloor d^{2\alpha}t \rfloor, 1}^d$ are bounded, $\lim_{d \rightarrow \infty} \mathbb{E}[|I_1|] = 0$. Thus,

$$\lim_{d \rightarrow \infty} \mathbb{E}[|V(L_t^d) - V(L_s^d) - I_{s,t}|] = 0,$$

where

$$\begin{aligned} I_{s,t} &= \int_s^t \left\{ \ell d^\alpha V' \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) b_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d, Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) \right. \\ &\quad \left. + \frac{\ell^2}{2} V'' \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) b_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d, Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right)^2 \mathbb{1}_{\mathbf{A}_{\lfloor d^{2\alpha}\tau \rfloor}^d} \right\} d\tau. \end{aligned} \quad (45)$$

Next, we use (18) and write

$$\int_s^t \text{LV}(L_\tau^d) d\tau = \int_s^t \frac{h^L(\ell)}{2} \left[V'' \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) - \text{sgn} \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) V' \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) \right] d\tau - T_3^d, \quad (46)$$

where we define

$$T_3^d = \int_s^t \left(\text{LV} \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) - \text{LV} \left(L_\tau^d \right) \right) d\tau.$$

Finally, we write the difference $M_{\lfloor d^{2\alpha}t \rfloor}^d(V) - M_{\lfloor d^{2\alpha}s \rfloor}^d(V)$ as the integral of a piecewise constant

function

$$\begin{aligned}
M_{\lceil d^{2\alpha} t \rceil}^d(V) - M_{\lceil d^{2\alpha} s \rceil}^d(V) &= I_{s,t} \\
&- \int_s^t \left(\ell d^\alpha V' \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) \mathbb{E} \left[b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right) \mathbb{1}_{\mathbf{A}_{\lceil d^{2\alpha} \tau \rceil}^d} \middle| \mathcal{F}_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right] \right. \\
&\quad \left. + \frac{\ell^2}{2} V'' \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) \mathbb{E} \left[b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right)^2 \mathbb{1}_{\mathbf{A}_{\lceil d^{2\alpha} \tau \rceil}^d} \middle| \mathcal{F}_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right] \right) d\tau \\
&- T_4^d - T_5^d,
\end{aligned} \tag{47}$$

where T_4^d and T_5^d account for the difference between the sum in (41) and the integral, and are defined as

$$\begin{aligned}
T_4^d &= -\frac{\ell}{d^\alpha} (\lceil d^{2\alpha} t \rceil - d^{2\alpha} t) V' \left(X_{\lfloor d^{2\alpha} t \rfloor, 1}^d \right) \left\{ b_d \left(X_{\lfloor d^{2\alpha} t \rfloor, 1}^d, Z_{\lceil d^{2\alpha} t \rceil, 1}^d \right) \mathbb{1}_{\mathbf{A}_{\lceil d^{2\alpha} t \rceil}^d} \right. \\
&\quad \left. - \mathbb{E} \left[b_d \left(X_{\lfloor d^{2\alpha} t \rfloor, 1}^d, Z_{\lceil d^{2\alpha} t \rceil, 1}^d \right) \mathbb{1}_{\mathbf{A}_{\lceil d^{2\alpha} t \rceil}^d} \middle| \mathcal{F}_{\lfloor d^{2\alpha} t \rfloor, 1}^d \right] \right\} \\
&- \frac{\ell^2}{2d^{2\alpha}} (\lceil d^{2\alpha} t \rceil - d^{2\alpha} t) V'' \left(X_{\lfloor d^{2\alpha} t \rfloor, 1}^d \right) \left\{ b_d \left(X_{\lfloor d^{2\alpha} t \rfloor, 1}^d, Z_{\lceil d^{2\alpha} t \rceil, 1}^d \right)^2 \mathbb{1}_{\mathbf{A}_{\lceil d^{2\alpha} t \rceil}^d} \right. \\
&\quad \left. - \mathbb{E} \left[b_d \left(X_{\lfloor d^{2\alpha} t \rfloor, 1}^d, Z_{\lceil d^{2\alpha} t \rceil, 1}^d \right)^2 \mathbb{1}_{\mathbf{A}_{\lceil d^{2\alpha} t \rceil}^d} \middle| \mathcal{F}_{\lfloor d^{2\alpha} t \rfloor, 1}^d \right] \right\},
\end{aligned}$$

and $T_5^d = -T_4^d$ with t substituted by s . Putting (45), (46) and (47) together we obtain

$$I_{s,t} - \int_s^t LV(L_\tau^d) d\tau - \left(M_{\lceil d^{2\alpha} t \rceil}^d(V) - M_{\lceil d^{2\alpha} s \rceil}^d(V) \right) = T_1^d + T_2^d + T_3^d + T_4^d + T_5^d,$$

where T_1^d takes into account all the terms involving $V'(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d)$, and T_2^d the terms involving $V''(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d)$:

$$\begin{aligned}
T_1^d &= \int_s^t V' \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) \\
&\quad \times \left\{ \ell d^\alpha \mathbb{E} \left[b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right) \mathbb{1}_{\mathbf{A}_{\lceil d^{2\alpha} \tau \rceil}^d} \middle| \mathcal{F}_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right] + \frac{h^L(\ell)}{2} \operatorname{sgn} \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) \right\} d\tau, \\
T_2^d &= \int_s^t V'' \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) \\
&\quad \times \left\{ \frac{\ell^2}{2} \mathbb{E} \left[b_d \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d \right)^2 \mathbb{1}_{\mathbf{A}_{\lceil d^{2\alpha} \tau \rceil}^d} \middle| \mathcal{F}_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right] - \frac{h^L(\ell)}{2} \right\} d\tau.
\end{aligned}$$

To obtain (42) it is then sufficient to prove that for any $1 \leq i \leq 5$, $\lim_{d \rightarrow \infty} \mathbb{E} [|T_i^d|] = 0$.

Since V', V'' are bounded and b_d is bounded in expectation because the moments of $Z_{\lceil d^{2\alpha} \tau \rceil, 1}^d$ are bounded, it is easy to show that $\lim_{d \rightarrow \infty} \mathbb{E} [|T_i^d|] = 0$ for $i = 4, 5$. For T_3^d , we write $T_3^d = h^L(\ell)(T_{3,1}^d - T_{3,2}^d)/2$, where

$$\begin{aligned}
T_{3,1}^d &= \int_s^t \left\{ V'' \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) - V''(L_\tau^d) \right\} d\tau, \\
T_{3,2}^d &= \int_s^t \left\{ \operatorname{sgn} \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) V' \left(X_{\lfloor d^{2\alpha} \tau \rfloor, 1}^d \right) - \operatorname{sgn}(L_\tau^d) V'(L_\tau^d) \right\} d\tau.
\end{aligned}$$

Using Fubini-Tonelli's theorem, the convergence of $X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d$ to L_τ^d in Lemma 2 and Lebesgue's dominated convergence theorem we obtain

$$\mathbb{E} [|T_{3,1}^d|] \leq \int_s^t \mathbb{E} \left[\left| V'' \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) - V'' (L_\tau^d) \right| \right] d\tau \xrightarrow{d \rightarrow \infty} 0.$$

We can further decompose $T_{3,2}^d$ as

$$\begin{aligned} T_{3,2}^d &= \int_s^t \left\{ \operatorname{sgn} \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) - \operatorname{sgn} (L_\tau^d) \right\} V' \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) d\tau \\ &\quad + \int_s^t \operatorname{sgn} (L_\tau^d) \left\{ V' \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) - V' (L_\tau^d) \right\} d\tau. \end{aligned}$$

Proceeding as for $T_{3,1}^d$, it is easy to show that the second integral converges to 0 as $d \rightarrow \infty$. We then bound the first integral by

$$\begin{aligned} \mathbb{E} \left[\left| \int_s^t \left\{ \operatorname{sgn} \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) - \operatorname{sgn} (L_\tau^d) \right\} V' \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) d\tau \right| \right] \\ \leq C \int_s^t \mathbb{E} \left[\left| \operatorname{sgn} \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) - \operatorname{sgn} (L_\tau^d) \right| \right] d\tau. \end{aligned}$$

However, since $\{\operatorname{sgn}(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d) \neq \operatorname{sgn}(L_\tau^d)\} \subset \{\operatorname{sgn}(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d) \neq \operatorname{sgn}(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d)\}$, using Lemma 4 in Appendix D.3 we have that

$$\begin{aligned} \mathbb{E} \left[\left| \operatorname{sgn} \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) - \operatorname{sgn} (L_\tau^d) \right| \right] &= 2\mathbb{E} \left[\mathbb{1} \left\{ \operatorname{sgn} \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) \neq \operatorname{sgn} (L_\tau^d) \right\} \right] \\ &= 2\mathbb{E} \left[\mathbb{1} \left\{ \operatorname{sgn} \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) \neq \operatorname{sgn} \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) \right\} \right] \xrightarrow{d \rightarrow \infty} 0. \end{aligned}$$

The above and the dominated converge theorem show that

$$\mathbb{E} \left[\left| \int_s^t \left\{ \operatorname{sgn} \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) - \operatorname{sgn} (L_\tau^d) \right\} V' \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) d\tau \right| \right] \xrightarrow{d \rightarrow \infty} 0.$$

Consider then T_1^d , recalling that the derivatives of V are bounded, we have

$$\begin{aligned} \mathbb{E} [|T_1^d|] &\leq \int_s^t C \mathbb{E} \left[\left| \ell d^\alpha \mathbb{E} \left[b_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d, Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) \mathbb{1}_{\mathcal{A}_{\lfloor d^{2\alpha}\tau \rfloor}^d} \middle| \mathcal{F}_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right] \right. \right. \\ &\quad \left. \left. + \frac{h^L(\ell)}{2} \operatorname{sgn} \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) \right| \right] d\tau \\ &\leq \int_s^t C \left\{ \mathbb{E} \left[\left| D_{1,\tau}^{(1)} \right| \right] + \mathbb{E} \left[\left| D_{2,\tau}^{(1)} \right| \right] \right\} d\tau, \end{aligned}$$

where we define

$$\begin{aligned}
D_{1,\tau}^{(1)} &= \ell d^\alpha \mathbb{E} \left[Z_{\lceil d^{2\alpha}\tau \rceil,1}^d \mathbb{1}_{\mathbf{A}_{\lceil d^{2\alpha}\tau \rceil}^d} \middle| \mathcal{F}_{\lceil d^{2\alpha}\tau \rceil,1}^d \right], \\
D_{2,\tau}^{(1)} &= \frac{h^L(\ell)}{2} \operatorname{sgn} \left(X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d \right) \\
&\quad - \ell d^\alpha \left(\frac{\sigma_d}{2} \operatorname{sgn}(X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d) \mathbb{1}_{|X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d| \geq \sigma_d^{2m} r/2} + \frac{1}{\sigma_d^{2m-1} r} X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d \mathbb{1}_{|X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d| < \sigma_d^{2m} r/2} \right) \\
&\quad \times \mathbb{E} \left[\mathbb{1}_{\mathbf{A}_{\lceil d^{2\alpha}\tau \rceil}^d} \middle| \mathcal{F}_{\lceil d^{2\alpha}\tau \rceil,1}^d \right].
\end{aligned}$$

Let us start with $D_{1,\tau}^{(1)}$:

$$D_{1,\tau}^{(1)} = \ell d^\alpha \mathbb{E} \left[Z_{\lceil d^{2\alpha}\tau \rceil,1}^d \left(1 \wedge \exp \left\{ \sum_{i=1}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor,i}^d, Z_{\lceil d^{2\alpha}\tau \rceil,i}^d \right) \right\} \right) \middle| \mathcal{F}_{\lceil d^{2\alpha}\tau \rceil,1}^d \right],$$

where ϕ_d is given in (21). Then, by independence of the components of $Z_{\lceil d^{2\alpha}\tau \rceil}^d$, we have

$$\begin{aligned}
&\mathbb{E} \left[Z_{\lceil d^{2\alpha}\tau \rceil,1}^d \left(1 \wedge \exp \left\{ \sum_{i=2}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor,i}^d, Z_{\lceil d^{2\alpha}\tau \rceil,i}^d \right) \right\} \right) \middle| \mathcal{F}_{\lceil d^{2\alpha}\tau \rceil,1}^d \right] \\
&= \mathbb{E} \left[Z_{\lceil d^{2\alpha}\tau \rceil,1}^d \right] \mathbb{E} \left[1 \wedge \exp \left\{ \sum_{i=2}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor,i}^d, Z_{\lceil d^{2\alpha}\tau \rceil,i}^d \right) \right\} \middle| \mathcal{F}_{\lceil d^{2\alpha}\tau \rceil,1}^d \right] = 0.
\end{aligned}$$

This allows us to write

$$\begin{aligned}
\mathbb{E} \left[|D_{1,\tau}^{(1)}| \right] &\leq \ell d^\alpha \mathbb{E} \left[|Z_{\lceil d^{2\alpha}\tau \rceil,1}^d| \right. \\
&\quad \left. \left| 1 \wedge \exp \left\{ \sum_{i=1}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor,i}^d, Z_{\lceil d^{2\alpha}\tau \rceil,i}^d \right) \right\} - 1 \wedge \exp \left\{ \sum_{i=2}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor,i}^d, Z_{\lceil d^{2\alpha}\tau \rceil,i}^d \right) \right\} \right| \right].
\end{aligned}$$

However, $x \mapsto 1 \wedge \exp(x)$ is a 1-Lipschitz function, thus

$$\mathbb{E} \left[|D_{1,\tau}^{(1)}| \right] \leq \ell d^\alpha \mathbb{E} \left[|Z_{\lceil d^{2\alpha}\tau \rceil,1}^d| \left| \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d, Z_{\lceil d^{2\alpha}\tau \rceil,1}^d \right) \right| \right],$$

and $D_{1,\tau}^{(1)} \rightarrow 0$ as $d \rightarrow \infty$ by Lemma 5 in Appendix D.3.

For $D_{2,\tau}^{(1)}$, we observe that

$$-\frac{\sigma_d}{2} \mathbb{1}_{|X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d| < \sigma_d^{2m} r/2} \leq \frac{1}{\sigma_d^{2m-1} r} X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d \mathbb{1}_{|X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d| < \sigma_d^{2m} r/2} \leq \frac{\sigma_d}{2} \mathbb{1}_{|X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d| < \sigma_d^{2m} r/2}. \quad (48)$$

Distinguishing between $X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d < 0$ and $X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d \geq 0$, it follows that

$$\begin{aligned}
|D_{2,\tau}^{(1)}| &\leq \left| \operatorname{sgn} \left(X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d \right) \right| \\
&\quad \times \left| \frac{h^L(\ell)}{2} - \ell d^\alpha \left(\frac{\sigma_d}{2} \mathbb{1}_{|X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d| \geq \sigma_d^{2m} r/2} + \frac{\sigma_d}{2} \mathbb{1}_{|X_{\lfloor d^{2\alpha}\tau \rfloor,1}^d| < \sigma_d^{2m} r/2} \right) \mathbb{E} \left[\mathbb{1}_{\mathbf{A}_{\lceil d^{2\alpha}\tau \rceil}^d} \middle| \mathcal{F}_{\lceil d^{2\alpha}\tau \rceil,1}^d \right] \right| \\
&\leq \frac{1}{2} \left| h^L(\ell) - \ell^2 \mathbb{E} \left[\mathbb{1}_{\mathbf{A}_{\lceil d^{2\alpha}\tau \rceil}^d} \middle| \mathcal{F}_{\lceil d^{2\alpha}\tau \rceil,1}^d \right] \right|,
\end{aligned}$$

where we recall that $\sigma_d = \ell d^{-\alpha}$ with $\alpha = 1/3$. Using the triangle inequality we obtain

$$\begin{aligned} 2\mathbb{E} \left[|D_{2,\tau}^{(1)}| \right] &\leq \mathbb{E} \left[\left| h^L(\ell) - \ell^2 \mathbb{E} \left[1 \wedge \exp \left(\sum_{i=1}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) \middle| \mathcal{F}_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right] \right| \right] \\ &\leq \mathbb{E} \left[\left| h^L(\ell) - \ell^2 \mathbb{E} \left[1 \wedge \exp \left(\sum_{i=2}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) \middle| \mathcal{F}_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right] \right| \right] \\ &\quad + \ell^2 \mathbb{E} \left[\left| 1 \wedge \exp \left(\sum_{i=2}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) - 1 \wedge \exp \left(\sum_{i=1}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) \right| \right], \end{aligned}$$

where we used Jensen's inequality to remove the conditional expectation in the last term. Recalling that $x \mapsto 1 \wedge \exp(x)$ is 1-Lipschitz, we can then bound the second term

$$\begin{aligned} &\ell^2 \mathbb{E} \left[\left| 1 \wedge \exp \left(\sum_{i=2}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) - 1 \wedge \exp \left(\sum_{i=1}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) \right| \right] \\ &\leq \ell^2 \mathbb{E} \left[\left| \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha}\tau \rceil, 1}^d \right) \right| \right], \\ &\leq \ell^2 \mathbb{E} \left[\left(\phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha}\tau \rceil, 1}^d \right) \right)^2 \right]^{1/2}, \end{aligned} \tag{49}$$

where the final expectation converges to zero as $d \rightarrow \infty$ by Proposition 17. For the remaining term in $D_{2,\tau}^{(1)}$, since $(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d)_{2 \leq i \leq n}$ is independent of $\mathcal{F}_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d$, we have

$$\begin{aligned} &\ell^2 \mathbb{E} \left[\left| 1 \wedge \exp \left(\sum_{i=2}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) \middle| \mathcal{F}_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right| \right] \\ &= \ell^2 \mathbb{E} \left[\left| 1 \wedge \exp \left(\sum_{i=2}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) \right| \right], \end{aligned}$$

and, using again the fact that $x \mapsto 1 \wedge \exp(x)$ is 1-Lipschitz, we have

$$\begin{aligned} &\left| h^L(\ell) - \ell^2 \mathbb{E} \left[1 \wedge \exp \left(\sum_{i=2}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) \right] \right| \\ &\leq \left| h^L(\ell) - \ell^2 \mathbb{E} \left[1 \wedge \exp \left(\sum_{i=1}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) \right] \right| \\ &\quad + \ell^2 \mathbb{E} \left[\left| \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d, Z_{\lceil d^{2\alpha}\tau \rceil, 1}^d \right) \right| \right]. \end{aligned}$$

The last term goes to 0 as shown in (49), and, as $h^L(\ell) = \ell^2 a^L(\ell)$, with

$$a^L(\ell) = \lim_{d \rightarrow \infty} \mathbb{E} \left[1 \wedge \exp \left(\sum_{i=1}^d \phi_{d,i} \right) \right],$$

by Theorem 2, we obtain

$$\lim_{d \rightarrow \infty} \left| h^L(\ell) - \ell^2 \mathbb{E} \left[1 \wedge \exp \left(\sum_{i=2}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lfloor d^{2\alpha}\tau \rfloor, i}^d \right) \right) \right] \right| = 0 ,$$

showing that $D_{2,\tau}^{(1)} \rightarrow 0$ as $d \rightarrow \infty$. To obtain convergence of T_1^d , we observe that for any $\tau \in [s, t]$, $D_{1,\tau}^{(1)}$ and $D_{2,\tau}^{(1)}$ follow the same distributions as $D_{1,s}^{(1)}$ and $D_{2,s}^{(1)}$, since for any $k \in \mathbb{N}$, X_k^d has distribution π_d^L . Therefore, the convergence towards zero of $\mathbb{E}[|D_{1,\tau}^{(1)}|]$ and $\mathbb{E}[|D_{2,\tau}^{(1)}|]$ is uniform for $\tau \in [s, t]$, which gives us $T_1^d \rightarrow 0$ as $d \rightarrow \infty$.

Finally, consider T_2^d . Using analogous arguments to those used for T_1^d , we obtain

$$\begin{aligned} \mathbb{E}[|T_2^d|] &\leq C \int_s^t \frac{\ell^2}{2} \mathbb{E} \left[\left| \mathbb{E} \left[b_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d, Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right)^2 \mathbb{1}_{\mathcal{A}_{\lfloor d^{2\alpha}\tau \rfloor}^d} \middle| \mathcal{F}_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right] - a^L(\ell, r) \right| \right] d\tau \\ &\leq C \int_s^t \frac{\ell^2}{2} \left\{ \mathbb{E}[|D_{1,\tau}^{(2)}|] + \mathbb{E}[|D_{2,\tau}^{(2)}|] \mathbb{E}[|D_{3,\tau}^{(2)}|] \right\} d\tau , \end{aligned}$$

where we define

$$\begin{aligned} D_{1,\tau}^{(2)} &= \mathbb{E} \left[\left(Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right)^2 \mathbb{1}_{\mathcal{A}_{\lfloor d^{2\alpha}\tau \rfloor}^d} \middle| \mathcal{F}_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right] - a^L(\ell, r) , \\ D_{2,\tau}^{(2)} &= \left(\frac{\sigma_d}{2} \operatorname{sgn}(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d) \mathbb{1}\{|X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d| \geq \sigma_d^{2m} r/2\} \right. \\ &\quad \left. + \frac{1}{\sigma_d^{2m-1} r} X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \mathbb{1}\{|X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d| < \sigma_d^{2m} r/2\} \right)^2 \times \mathbb{E} \left[\mathbb{1}_{\mathcal{A}_{\lfloor d^{2\alpha}\tau \rfloor}^d} \middle| \mathcal{F}_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right] , \\ D_{3,\tau}^{(2)} &= 2 \left(\frac{\sigma_d}{2} \operatorname{sgn}(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d) \mathbb{1}\{|X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d| \geq \sigma_d^{2m} r/2\} \right. \\ &\quad \left. + \frac{1}{\sigma_d^{2m-1} r} X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \mathbb{1}\{|X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d| < \sigma_d^{2m} r/2\} \right) \mathbb{E} \left[Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \mathbb{1}_{\mathcal{A}_{\lfloor d^{2\alpha}\tau \rfloor}^d} \middle| \mathcal{F}_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right] . \end{aligned}$$

Using (48), Cauchy-Schwarz's inequality and the fact that the moments of $Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d$ are bounded we have

$$\mathbb{E}[|D_{2,\tau}^{(2)}|] \leq \frac{\sigma_d^2}{4} \xrightarrow{d \rightarrow \infty} 0 , \quad \mathbb{E}[|D_{3,\tau}^{(2)}|] \leq C \sigma_d \xrightarrow{d \rightarrow \infty} 0 ,$$

since $\sigma_d = \ell d^{-\alpha}$ with $\alpha = 1/3$. The remaining term is bounded similarly to $D_{2,\tau}^{(1)}$, using the fact that $x \mapsto 1 \wedge \exp(x)$ is 1-Lipschitz, we have

$$\begin{aligned} &\mathbb{E}[|D_{3,\tau}^{(2)}|] \\ &\leq \mathbb{E} \left[\left| \mathbb{E} \left[\left(Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right)^2 \left(1 \wedge \exp \left(\sum_{i=2}^d \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, i}^d, Z_{\lfloor d^{2\alpha}\tau \rfloor, i}^d \right) \right) \right) \right] \middle| \mathcal{F}_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right] - a^L(\ell, r) \right| \right] \\ &\quad + \mathbb{E} \left[\left(Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right)^2 \left| \phi_d \left(X_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d, Z_{\lfloor d^{2\alpha}\tau \rfloor, 1}^d \right) \right| \right] . \end{aligned}$$

The second expectation is bounded as (49) using Cauchy-Schwarz's inequality and Proposition 17. For the first expectation, we use the conditional independence of the components of $Z_{\lceil d^{2\alpha}\tau \rceil}^d$ and write

$$\begin{aligned} & \mathbb{E} \left[\left(Z_{\lceil d^{2\alpha}\tau \rceil, 1}^d \right)^2 \left(1 \wedge \exp \left(\sum_{i=2}^d \phi_d \left(X_{\lceil d^{2\alpha}\tau \rceil, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) \right) \middle| \mathcal{F}_{\lceil d^{2\alpha}\tau \rceil, 1}^d \right] \\ &= \mathbb{E} \left[\left(Z_{\lceil d^{2\alpha}\tau \rceil, 1}^d \right)^2 \right] \mathbb{E} \left[\left(1 \wedge \exp \left(\sum_{i=2}^d \phi_d \left(X_{\lceil d^{2\alpha}\tau \rceil, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) \right) \right] \\ &= \mathbb{E} \left[\left(1 \wedge \exp \left(\sum_{i=2}^d \phi_d \left(X_{\lceil d^{2\alpha}\tau \rceil, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) \right) \right] . \end{aligned}$$

It follows that $\mathbb{E}[|D_{3,\tau}^{(2)}|] \rightarrow 0$ as $d \rightarrow \infty$ since, by Theorem 2,

$$\left| \mathbb{E} \left[\left(1 \wedge \exp \left(\sum_{i=2}^d \phi_d \left(X_{\lceil d^{2\alpha}\tau \rceil, i}^d, Z_{\lceil d^{2\alpha}\tau \rceil, i}^d \right) \right) \right) \right] - a^L(\ell, r) \right| \rightarrow 0 .$$

Combining the results for T_i^d , $i = 1, \dots, 5$ we obtain the result. $\square$

Acknowledgments

F.R.C. and G.O.R. acknowledge support from the EPSRC (grant # EP/R034710/1). G.O.R. acknowledges further support from the EPSRC (grant # EP/R018561/1) and the Alan Turing Institute. A.D. acknowledges support from the Lagrange Mathematics and Computing Research Center.

The authors would like to thank Éric Moulines for helpful discussions.

For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

References

- [1] S. Agrawal, D. Vats, K. Łatuszyński, and G. O. Roberts. Optimal scaling of MCMC beyond Metropolis. *arXiv preprint arXiv:2104.02020*, 2021.
- [2] Y. F. Atchadé. A Moreau-Yosida approximation scheme for a class of high-dimensional posterior distributions. *arXiv preprint arXiv:1505.07072*, 2015.
- [3] H. H. Bauschke and P. L. Combettes. *Convex analysis and monotone operator theory in Hilbert spaces*, volume 408. Springer, 2011.
- [4] M. Bédard and J. S. Rosenthal. Optimal scaling of Metropolis algorithms: Heading toward general target distributions. *Canadian Journal of Statistics*, 36(4):483–503, 2008.
- [5] A. Beskos, G. Roberts, and A. Stuart. Optimal scalings for local Metropolis–Hastings chains on nonproduct targets in high dimensions. *Annals of Applied Probability*, 19(3):863 – 898, 2009.

- [6] J. Bierkens, P. Fearnhead, and G. Roberts. The zig-zag process and super-efficient sampling for Bayesian analysis of big data. *Annals of Statistics*, 47(3):1288–1320, 2019.
- [7] N. Bou-Rabee and M. Hairer. Nonasymptotic mixing of the MALA algorithm. *IMA Journal of Numerical Analysis*, 33(1):80–110, 2013.
- [8] A. Bouchard-Côté, S. J. Vollmer, and A. Doucet. The bouncy particle sampler: A nonreversible rejection-free Markov chain Monte Carlo method. *Journal of the American Statistical Association*, 113(522):855–867, 2018.
- [9] S. Brooks, A. Gelman, G. Jones, and X.-L. Meng. *Handbook of Markov chain Monte Carlo*. CRC press, 2011.
- [10] O. F. Christensen, G. O. Roberts, and J. S. Rosenthal. Scaling limits for the transient phase of local Metropolis–Hastings algorithms. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 67(2):253–268, 2005.
- [11] P. L. Combettes and J.-C. Pesquet. Proximal splitting methods in signal processing. In H. H. Bauschke, R. S. Burachik, P. L. Combettes, V. Elser, D. R. Luke, and H. Wolkowicz, editors, *Fixed-Point Algorithms for Inverse Problems in Science and Engineering*, pages 185–212. Springer, 2011.
- [12] A. Durmus, S. Le Corff, E. Moulines, and G. O. Roberts. Optimal scaling of the random walk Metropolis algorithm under L_p mean differentiability. *Journal of Applied Probability*, 54(4):1233–1260, 2017.
- [13] A. Durmus and E. Moulines. On the geometric convergence for MALA under verifiable conditions. *arXiv preprint arXiv:2201.01951*, 2022.
- [14] A. Durmus, E. Moulines, and M. Pereyra. Efficient Bayesian computation by proximal Markov chain Monte Carlo: when Langevin meets Moreau. *SIAM Journal on Imaging Sciences*, 11(1):473–506, 2018.
- [15] R. Durrett. *Probability: Theory and Examples*, volume 49. Cambridge University Press, 2019.
- [16] R. Dwivedi, Y. Chen, M. J. Wainwright, and B. Yu. Log-concave sampling: Metropolis-Hastings algorithms are fast! In S. Bubeck, V. Perchet, and P. Rigollet, editors, *Proceedings of the 31st Conference On Learning Theory*, volume 75 of *Proceedings of Machine Learning Research*, pages 793–797. PMLR, 06–09 Jul 2018.
- [17] S. N. Ethier and T. G. Kurtz. *Markov processes: characterization and convergence*, volume 282. John Wiley & Sons, 2009.
- [18] A. Gelman, G. O. Roberts, and W. Gilks. Efficient Metropolis jumping rules. *Bayesian Statistics*, 1996.
- [19] J. V. Goldman, T. Sell, and S. S. Singh. Gradient-based Markov chain Monte Carlo for Bayesian inference with non-differentiable priors. *Journal of the American Statistical Association*, pages 1–12, 2021.
- [20] U. Grenander and M. I. Miller. Representations of knowledge in complex systems. *Journal of the Royal Statistical Society: Series B (Methodological)*, 56(4):549–581, 1994.

- [21] S. F. Jarner and G. O. Roberts. Convergence of heavy-tailed Monte Carlo Markov chain algorithms. *Scandinavian Journal of Statistics*, 34(4):781–815, 2007.
- [22] B. Jourdain, T. Lelièvre, and B. Miasojedow. Optimal scaling for the transient phase of Metropolis Hastings algorithms: the longtime behavior. *Bernoulli*, 20(4):1930–1978, 2014.
- [23] O. Kallenberg. *Foundations of Modern Probability*. Springer, 2021.
- [24] S. Livingstone and G. Zanella. The Barker proposal: Combining robustness and efficiency in gradient-based MCMC. *Journal of the Royal Statistical Society Series B*, 84(2):496–523, 2022.
- [25] Y.-A. Ma, T. Chen, and E. Fox. A complete recipe for stochastic gradient MCMC. *Advances in Neural Information Processing Systems*, 28, 2015.
- [26] K. L. Mengersen and R. L. Tweedie. Rates of convergence of the Hastings and Metropolis algorithms. *Annals of Statistics*, 24(1):101–121, 1996.
- [27] P. Neal and G. Roberts. Optimal scaling for random walk Metropolis on spherically constrained target densities. *Methodology and Computing in Applied Probability*, 10(2):277–297, 2008.
- [28] P. Neal, G. Roberts, W. K. Yuen, et al. Optimal scaling of random walk metropolis algorithms with discontinuous target densities. *Annals of Applied Probability*, 22(5):1880–1927, 2012.
- [29] N. Parikh and S. Boyd. Proximal algorithms. *Foundations and Trends in Optimization*, 1(3):127–239, 2014.
- [30] M. Pereyra. Proximal Markov chain Monte Carlo algorithms. *Statistics and Computing*, 26(4):745–760, 2016.
- [31] N. S. Pillai. Optimal scaling for the proximal Langevin algorithm in high dimensions. *arXiv preprint arXiv:2204.10793*, 2022.
- [32] N. S. Pillai, A. M. Stuart, and A. H. Thiéry. Optimal scaling and diffusion limits for the Langevin algorithm in high dimensions. *Annals of Applied Probability*, 22(6):2320 – 2356, 2012.
- [33] G. O. Roberts, A. Gelman, and W. R. Gilks. Weak convergence and optimal scaling of random walk Metropolis algorithms. *Annals of Applied Probability*, 7(1):110–120, 1997.
- [34] G. O. Roberts and J. S. Rosenthal. Optimal scaling of discrete approximations to Langevin diffusions. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 60(1):255–268, 1998.
- [35] G. O. Roberts, J. S. Rosenthal, et al. Optimal scaling for various Metropolis-Hastings algorithms. *Statistical Science*, 16(4):351–367, 2001.
- [36] G. O. Roberts and R. L. Tweedie. Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli*, 2(4):341 – 363, 1996.
- [37] G. O. Roberts and R. L. Tweedie. Geometric convergence and central limit theorems for multidimensional Hastings and Metropolis algorithms. *Biometrika*, 83(1):95–110, 1996.

- [38] R. T. Rockafellar and R. J.-B. Wets. *Variational Analysis*, volume 317 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Berlin, Heidelberg, 1998.
- [39] P. J. Rossky, J. D. Doll, and H. L. Friedman. Brownian dynamics as smart Monte Carlo simulation. *The Journal of Chemical Physics*, 69(10):4628–4633, 1978.
- [40] C. Sherlock, G. Roberts, et al. Optimal scaling of the random walk Metropolis on elliptically symmetric unimodal targets. *Bernoulli*, 15(3):774–798, 2009.
- [41] A. N. Shiryaev. *Probability*, volume 25. Springer, 1996.
- [42] D. W. Stroock and S. Varadhan. *Multidimensional Diffusion Processes*. Springer, 1979.
- [43] M. Vono, N. Dobigeon, and P. Chainais. Bayesian image restoration under Poisson noise and log-concave prior. In *ICASSP 2019-2019 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, pages 1712–1716. IEEE, 2019.
- [44] J. Vorstrup Goldman, T. Sell, and S. S. Singh. Gradient-based Markov chain Monte Carlo for Bayesian inference with non-differentiable priors. *Journal of the American Statistical Association*, 0(0):1–12, 2021.
- [45] J. Yang, G. O. Roberts, and J. S. Rosenthal. Optimal scaling of random-walk Metropolis algorithms on general target distributions. *Stochastic Processes and their Applications*, 130(10):6094–6132, 2020.
- [46] X. Zhou, E. C. Chi, and H. Zhou. Proximal MCMC for Bayesian inference of constrained and regularized estimation. *arXiv preprint arXiv:2205.07378*, 2022.

A Proof of Theorem 1

The proof of Theorem 1 follows that of [34, Theorem 1, Theorem 2] and consists of four propositions showing convergence of the log-acceptance probability to a normal random variable and (weak) convergence of the process (11) to a Langevin diffusion.

We start by recalling and defining a number of quantities that we will use in the following proofs. Recall that $\sigma_d = \ell/d^\alpha$, that $\lambda_d = \sigma_d^{2m}r/2$ where $m \geq 1/2$ and $r > 0$ are to be chosen according to the different cases in Theorem 1. Recalling the expression of the proposal given in (9) and using the simplification given in (10), we define the proposal with starting point $\mathbf{x}^d \in \mathbb{R}^d$,

$$\mathbf{y}^d(\mathbf{x}^d, \mathbf{z}^d) = \mathbf{x}^d - \frac{\sigma_d^2}{2} \nabla G \left(\text{prox}_{G^{\sigma_d^{2m}r/2}}(\mathbf{x}^d) \right) + \sigma_d \mathbf{z}^d,$$

where $\mathbf{z}^d \in \mathbb{R}^d$. Since $G(\mathbf{x}^d) = \sum_{i=1}^d g(x_i^d)$, the i -th component of the proposal only depends on the i -th components of $\mathbf{x}^d$ and $\mathbf{z}^d$. Thus, for any $x, z \in \mathbb{R}$ we denote

$$y_d(x, z) = x - \frac{\sigma_d^2}{2} g' \left(\text{prox}_{g^{\sigma_d^{2m}r/2}}(x) \right) + \sigma_d z.$$

The proposal for the chain $(X_k^d)_{k \geq 0}$ is then given by $Y_k^d = \mathbf{y}^d(X_k^d, Z_{k+1}^d) = (y_d(X_{k,i}^d, Z_{k+1,i}^d))_{i \in \{1, \dots, d\}}$. Let us define the generator of the discrete process $(X_k^d)_{k \geq 0}$ for all $V \in C_c^\infty(\mathbb{R}^d, \mathbb{R})$, i.e. infinitely differentiable $\mathbb{R}$ -valued multivariate functions with compact support, and any $\mathbf{x}^d \in \mathbb{R}^d$,

$$\begin{aligned} L_d V(\mathbf{x}^d) &= d^{2\alpha} \mathbb{E} \left[[V(\mathbf{y}^d(\mathbf{x}^d, Z_1^d)) - V(\mathbf{x}^d)] \frac{\pi_d(\mathbf{y}^d(\mathbf{x}^d, Z_1^d)) q_d(\mathbf{y}^d(\mathbf{x}^d, Z_1^d), \mathbf{x}^d)}{\pi_d(\mathbf{x}^d) q_d(\mathbf{x}^d, \mathbf{y}^d(\mathbf{x}^d, Z_1^d))} \wedge 1 \right] \\ &= d^{2\alpha} \mathbb{E} \left[[V(\mathbf{y}^d(\mathbf{x}^d, Z_1^d)) - V(\mathbf{x}^d)] \prod_{i=1}^d \exp(\phi_d(x_i^d, Z_{1,i}^d)) \wedge 1 \right], \end{aligned}$$

where the expectation is w.r.t. $Z_1^d = (Z_{1,i}^d)_{i \in \{1, \dots, d\}}$, a d -dimensional standard normal random variable, and where we defined

$$\begin{aligned} \phi_d(x, z) &= \log \frac{\pi(y_d(x, z)) q(y_d(x, z), x)}{\pi(x) q(x, y_d(x, z))} \\ &= g(x) - g(y_d(x, z)) + \log q(y_d(x, z), x) - \log q(x, y_d(x, z)). \end{aligned} \quad (50)$$

In the remainder we will work with one-dimensional functions $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$ applied to the first component of $\mathbf{x}^d$ so that

$$\begin{aligned} L_d V(\mathbf{x}^d) &= d^{2\alpha} \mathbb{E} \left[[V(y_d(x_1^d, Z_{1,1}^d)) - V(x_1^d)] \frac{\pi_d(\mathbf{y}^d(\mathbf{x}^d, Z_1^d)) q_d(\mathbf{y}^d(\mathbf{x}^d, Z_1^d), \mathbf{x}^d)}{\pi_d(\mathbf{x}^d) q_d(\mathbf{x}^d, \mathbf{y}^d(\mathbf{x}^d, Z_1^d))} \wedge 1 \right] \\ &= d^{2\alpha} \mathbb{E} \left[[V(y_d(x_1^d, Z_{1,1}^d)) - V(x_1^d)] \prod_{i=1}^d \exp(\phi_d(x_i^d, Z_{1,i}^d)) \wedge 1 \right]. \end{aligned} \quad (51)$$

We also define $\tilde{L}_d$ to be a variant of L_d in which the first component of the acceptance ratio is omitted:

$$\tilde{L}_d V(\mathbf{x}^d) = d^{2\alpha} \mathbb{E} \left[[V(y_d(x_1^d, Z_{1,1}^d)) - V(x_1^d)] \prod_{i=2}^d \exp(\phi_d(x_i^d, Z_{1,i}^d)) \wedge 1 \right]. \quad (52)$$

We further define the generator of the Langevin diffusion (13)

$$LV(x) = \frac{h(\ell, r)}{2} [V''(x) - g'(x)V'(x)], \quad (53)$$

where $h(\ell, r) = \ell^2 a(\ell, r)$ is the speed of the diffusion and $a(\ell, r) = \lim_{d \rightarrow \infty} a_d(\ell, r)$ is given in Theorem 1.

We will make use of the derivatives of g in (8) up to order 8, which we denote by g', g'', g''' and $g^{(k)}$ for all $k = 4, \dots, 8$. We recall that $(g^\lambda)'$ is Lipschitz continuous with Lipschitz constant λ^{-1} [38, Proposition 12.19] and that $(g^\lambda)'(x) = \lambda^{-1}(\text{prox}_g^\lambda(x) - x)$, hence prox_g^λ is Lipschitz continuous with Lipschitz constant 1.

A.1 Auxiliary Results for the Proof of Case (a)

First, we characterize the limit behaviour of the acceptance ratio (12).

Proposition 4. Under **A0**, **A1** and **A2**, if $\alpha = 1/4$, $\beta = 1/8$ and $r > 0$, then

(i) the log-acceptance ratio (50), when $d \rightarrow \infty$, satisfies the following Taylor expansion

$$\phi_d(x, z) = d^{-1/2}C_2(x, z) + d^{-3/4}C_3(x, z) + d^{-1}C_4(x, z) + C_5(x, z, \sigma_d) ,$$

where $C_2(x, z)$ is given in (57), C_3 and C_4 are polynomials in z and the derivatives of g , such that $\mathbb{E}[C_3(X_{0,1}^d, Z_{1,1}^d)] = 0$ and $\mathbb{E}[C_2(X_{0,1}^d, Z_{1,1}^d)^2] = -2\mathbb{E}[C_4(X_{0,1}^d, Z_{1,1}^d)]$;

(ii) there exists sets $F_d \subseteq \mathbb{R}^d$ with $d^{2\alpha}\pi_d(F_d^c) \rightarrow 0$ such that

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in F_d} \mathbb{E} \left[\left| \sum_{i=2}^d \phi_d(x_i^d, Z_{1,i}^d) - d^{-1/2} \sum_{i=2}^d C_2(x_i^d, Z_{1,i}^d) + \frac{\ell^4 K_1(r)^2}{2} \right| \right] = 0 , \quad (54)$$

where $K_1(r)$ is given in Theorem 1-(a).

Proof. Take one component of the log-acceptance ratio

$$\phi_d(x, z) = g(x) - g(y_d(x, z)) + \log q(y_d(x, z), x) - \log q(x, y_d(x, z)) ,$$

with $y_d(x, z) = x - \sigma_d^2 g'(\text{prox}_g^{\sigma_d^{2m} r/2}(x))/2 + \sigma_d z$. We have that $\phi_d(x, z) = R_1(x, z, \sigma_d) + R_2(x, z, \sigma_d)$, where

$$\begin{aligned} R_1(x, z, \sigma) &= -g \left[x - \frac{\sigma^2}{2} g' \left(\text{prox}_g^{\sigma^{2m} r/2}(x) \right) + \sigma z \right] + g(x) , \\ R_2(x, z, \sigma) &= \frac{1}{2} z^2 - \frac{1}{2} \left(z - \frac{\sigma}{2} g' \left(\text{prox}_g^{\sigma^{2m} r/2} \left[x + \sigma z - \frac{\sigma^2}{2} g' \left[\text{prox}_g^{\sigma^{2m} r/2}(x) \right] \right] \right) \right. \\ &\quad \left. - \frac{\sigma}{2} g' \left[\text{prox}_g^{\sigma^{2m} r/2}(x) \right] \right)^2 . \end{aligned} \quad (55)$$

Following the approach of [34] we approximate $\phi_d(x, z)$ with a Taylor expansion about $\sigma_d \rightarrow 0$.

(i) Using a Taylor expansion of order 5, we obtain

$$\phi_d(x, z) = d^{-1/2}C_2(x, z) + d^{-3/4}C_3(x, z) + d^{-1}C_4(x, z) + C_5(x, z, \sigma_d) , \quad (56)$$

where

$$C_2(x, z) = \frac{\ell^2}{2} (-r z g''(x) g'(x)) , \quad (57)$$

$C_3(x, z)$ and $C_4(x, z)$ are given in Appendix C.1.1 and we use the integral form for the remainder

$$C_5(x, z, \sigma_d) = \int_0^{\sigma_d} \frac{\partial^5}{\partial \sigma^5} R(x, z, \sigma) \Big|_{\sigma=u} \frac{(\sigma_d - u)^4}{4!} du ,$$

with u between 0 and σ_d and the derivatives of R_1 and R_2 given in Appendix C.2. In addition, integrating by parts and using the moments of $Z_{1,1}^d$ we find that $\mathbb{E}[C_2(X_{0,1}^d, Z_{1,1}^d)] = \mathbb{E}[C_3(X_{0,1}^d, Z_{1,1}^d)] = 0$ and

$$2\mathbb{E}[C_4(X_{0,1}^d, Z_{1,1}^d)] + \mathbb{E}[C_2(X_{0,1}^d, Z_{1,1}^d)^2] = 0 .$$

(ii) To construct the sets F_d , consider, for $j = 3, 4$, $F_{d,j} = F_{d,j}^1 \cap F_{d,j}^2$ where we define

$$F_{d,j}^1 = \left\{ \mathbf{x}^d \in \mathbb{R}^d : \left| \sum_{i=2}^d \mathbb{E} [C_j(x_i^d, Z_{1,i}^d) - C_j(X_{0,i}^d, Z_{1,i}^d)] \right| \leq d^{5/8} \right\},$$

and

$$F_{d,j}^2 = \left\{ \mathbf{x}^d \in \mathbb{R}^d : \left| \sum_{i=2}^d V_j(x_i^d) - \mathbb{E} [V_j(X_{0,i}^d)] \right| \leq d^{6/5} \right\},$$

where $V_j(x) := \text{Var}(C_j(x, Z_{1,1}^d))$. Using Markov's inequality and the fact that C_j, V_j are bounded by polynomials since g and its derivatives are bounded by polynomials, it is easy to show that $d^{1/2}\pi_d((F_{d,j}^1)^c) \rightarrow 0$ and $d^{1/2}\pi_d((F_{d,j}^2)^c) \rightarrow 0$, from which follows $d^{1/2}\pi_d(F_{d,j}^c) \rightarrow 0$ as $d \rightarrow \infty$. To prove L^1 convergence of C_j for $j = 3, 4$, observe that

$$\begin{aligned} & \mathbb{E} \left[\left(\sum_{i=2}^d C_j(x_i^d, Z_{1,i}^d) - \mathbb{E} [C_j(X_{0,1}^d, Z_{1,1}^d)] \right)^2 \right] \\ &= \sum_{i=2}^d V_j(x_i^d) + \left(\sum_{i=2}^d \mathbb{E} [C_j(x_i^d, Z_{1,i}^d) - C_j(X_{0,1}^d, Z_{1,1}^d)] \right)^2, \end{aligned}$$

and that, for $\mathbf{x}^d \in F_{d,j}$, we have

$$\mathbb{E} \left[\left(\sum_{i=2}^d C_j(x_i^d, Z_{1,i}^d) - \mathbb{E} [C_j(X_{0,1}^d, Z_{1,1}^d)] \right)^2 \right] \leq \mathbb{E} [V_j(x_1^d)] (d-1) + d^{6/5} + d^{5/4}.$$

Thus, the third and fourth term in the Taylor expansion (56) converge in L^1 to 0 and $-\ell^4 K_1^2(r)/2$ respectively. Now, consider $C_5(x_i^d, Z_{1,i}^d, \sigma_d)$. We can bound $\frac{\partial^5 R}{\partial \sigma^5}(x, z, \sigma)$ with the derivatives of g evaluated at

$$x + \frac{\sigma^2}{2} \text{prox}_g^{\sigma^{2m} r/2}(x) + \sigma z \quad \text{and} \quad \text{prox}_g^{\sigma^{2m} r/2}(x).$$

Under our assumptions, the derivatives of g are bounded by polynomials M_0 , it follows that there exist polynomials p of the form

$$A \left(1 + \left[\text{prox}_g^{\sigma^{2m} r/2}(x) \right]^N \right) (1 + z^N) (1 + x^N) (1 + \sigma^N),$$

for sufficiently large A and sufficiently large even integer N , such that

$$\begin{aligned} & \left| g^{(k)} \left[x + \frac{\sigma^2}{2} \text{prox}_g^{\sigma^{2m} r/2}(x) + \sigma_d z \right] \right| \vee \left| g^{(k)} \left[\text{prox}_g^{\sigma^{2m} r/2}(x) \right] \right| \\ & \leq p(\text{prox}_g^{\sigma^{2m} r/2}(x), x, z, \sigma_d). \end{aligned}$$

In addition, $|\text{prox}_g^{\sigma^{2m}r/2}(x)| \leq C(1 + |x|)$ for some $C \geq 1$, and we can bound

$$p(\text{prox}_g^{\sigma^{2m}r/2}(x), x, z, \sigma) \leq A(1 + z^N)(1 + x^{2N})(1 + \sigma^N) .$$

Therefore, we have

$$\begin{aligned} \mathbb{E} [|C_5(x_i^d, Z_{1,i}^d, \sigma_d)|] &\leq A \mathbb{E} [1 + (Z_{1,i}^d)^N] (1 + (x_i^d)^{2N}) \int_0^{\sigma_d} (1 + u^N) \frac{(\sigma_d - u)^4}{4!} du \\ &\leq A \mathbb{E} [1 + (Z_{1,i}^d)^N] (1 + (x_i^d)^{2N}) d^{-5/2} \\ &\leq A(1 + (x_i^d)^{2N}) d^{-5/2} , \end{aligned}$$

where the last inequality follows since all the moments of Z_1^d are bounded. Let us denote $p(x) = A(1 + x^{2N})$ and

$$F_{d,5} = \left\{ \mathbf{x}^d \in \mathbb{R}^d : \left| d^{-1} \sum_{i=1}^d p(x_i^d) - \mathbb{E} [p(X_{0,i}^d)] \right| < 1 \right\} .$$

By Chebychev's inequality we have $\pi_d(F_{d,5}^c) \leq \text{Var}(p(X_{0,1}^d))d^{-1}$. Additionally, for all $\mathbf{x}^d \in F_{d,5}$,

$$\begin{aligned} \sum_{i=2}^d \mathbb{E} [|C_5(x_i^d, Z_{1,i}^d, \sigma_d)|] &\leq \sum_{i=2}^d d^{-5/2} (\mathbb{E} [p(X_{0,1}^d)] + d^{-1}) \\ &\leq d^{-3/2} (\mathbb{E} [p(X_{0,1}^d)] + 1) . \end{aligned}$$

Finally, set $F_d = \cap_{j=3}^5 F_{d,j}$. On F_d the last three terms of (56) converge uniformly in L^1 , and (54) follows using the triangle inequality.

□

Next, we compare the generator L_d and $\tilde{L}_d$ in (51) and (52) respectively.

Proposition 5. *Under **A0**, **A1** and **A2**, if $\alpha = 1/4$, $\beta = 1/8$ and $r > 0$, there exists sets $S_d \subseteq \mathbb{R}^d$ with $d^{2\alpha}\pi_d(S_d^c) \rightarrow 0$ such that for any $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$*

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in S_d} |L_d V(\mathbf{x}^d) - \tilde{L}_d V(\mathbf{x}^d)| = 0 ,$$

and

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in S_d} \mathbb{E} \left[\left| \left(\exp \left(\sum_{i=1}^d \phi_d(x_i^d, Z_{1,i}^d) \right) \wedge 1 \right) - \left(\exp \left(\sum_{i=2}^d \phi_d(x_i^d, Z_{1,i}^d) \right) \wedge 1 \right) \right| \right] = 0 .$$

Proof. The function $x \mapsto \exp(x) \wedge 1$ is Lipschitz continuous with Lipschitz constant 1, hence

$$|L_d V(\mathbf{x}^d) - \tilde{L}_d V(\mathbf{x}^d)| \leq d^{2\alpha} \mathbb{E} [|V(y_d(x_1^d, Z_{1,1}^d)) - V(x_1^d)| |R(x_1^d, Z_{1,1}^d, \sigma_d)|] ,$$

where $R(x, z, \sigma) = R_1(x, z, \sigma) + R_2(x, z, \sigma)$ as in (55). Using a Taylor expansion of order 1 about $\sigma = 0$ with integral remainder:

$$R(x, z, \sigma) = R(x, z, 0) + \left. \frac{\partial R}{\partial \sigma}(x, z, \sigma) \right|_{\sigma=0} \sigma + \int_0^\sigma \left. \frac{\partial^2 R}{\partial \sigma^2}(x, z, \sigma) \right|_{\sigma=u} (\sigma - u) du ,$$

we obtain

$$R(x, z, \sigma) = \int_0^\sigma \left. \frac{\partial^2 R}{\partial \sigma^2}(x, z, \sigma) \right|_{\sigma=u} (\sigma - u) du ,$$

where $\frac{\partial^2 R}{\partial \sigma^2}(x, z, \sigma)$ is bounded by the derivatives of g evaluated at

$$x + \frac{\sigma^2}{2} \text{prox}_g^{\sigma^{2m} r/2}(x) + \sigma z \quad \text{and} \quad \text{prox}_g^{\sigma^{2m} r/2}(x) .$$

Under our assumptions, the derivatives of g are bounded by polynomials M_0 , it follows that there exist polynomials p of the form

$$A \left(1 + \left[\text{prox}_g^{\sigma^{2m} r/2}(x) \right]^N \right) (1 + z^N) (1 + x^N) (1 + \sigma^N) ,$$

for sufficiently large A and sufficiently large even integer N , such that

$$\left| g^{(k)} \left[x + \frac{\sigma^2}{2} \text{prox}_g^{\sigma^{2m} r/2}(x) + \sigma z \right] \right| \vee \left| g^{(k)} \left[\text{prox}_g^{\sigma^{2m} r/2}(x) \right] \right| \leq p(\text{prox}_g^{\sigma^{2m} r/2}(x), x, z, \sigma) .$$

Proceeding as in Proposition 4, we can bound

$$p(\text{prox}_g^{\sigma^{2m} r/2}(x), x, z, \sigma) \leq A (1 + z^N) (1 + x^{2N}) (1 + \sigma^N) .$$

Therefore, we have

$$\begin{aligned} |R(x_1^d, Z_{1,1}^d, \sigma_d)| &\leq A (1 + (Z_{1,1}^d)^N) (1 + (x_1^d)^{2N}) \\ &\quad \times \int_0^{\sigma_d} (1 + u^N) (\sigma_d - u) du \leq A (1 + (Z_{1,1}^d)^N) (1 + (x_1^d)^{2N}) \frac{\sigma_d^2}{2} . \end{aligned} \quad (58)$$

Since $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$, there exists a constant C such that

$$\begin{aligned} |V(y_d(x_1^d, Z_{1,1}^d)) - V(x_1^d)| &\leq C |y_d(x_1^d, Z_{1,1}^d) - x_1^d| \\ &\leq C \left(\sigma_d |Z_{1,1}^d| + \frac{\sigma_d^2}{2} \left| g' \left(\text{prox}_g^{\sigma_d^{2m} r/2}(x_1^d) \right) \right| \right) . \end{aligned}$$

Recalling that $g'(\text{prox}_g^\lambda(x)) = (g^\lambda)'(x)$ with $(g^\lambda)'$ Lipschitz continuous with Lipschitz constant λ^{-1} , we have

$$\left| g' \left(\text{prox}_g^{\sigma_d^{2m} r/2}(x_1^d) \right) \right| \leq \frac{2}{\sigma_d^{2m} r} (1 + |x_1^d|) ,$$

and

$$\begin{aligned} |V(y_d(x_1^d, Z_{1,1}^d)) - V(x_1^d)| &\leq C \left(\sigma_d |Z_{1,1}^d| + \frac{\sigma_d^{2-2m}}{r} (1 + |x_1^d|) \right) \\ &\leq C \sigma_d \left(|Z_{1,1}^d| + \frac{1}{r} (1 + |x_1^d|) \right), \end{aligned} \quad (59)$$

since $m = 1/2$. Combining (58) and (59) we obtain

$$\begin{aligned} d^{2\alpha} |V(y_d(x_1^d, Z_{1,1}^d)) - V(x_1^d)| |R(x_1^d, Z_{1,1}^d, \sigma_d)| \\ \leq C \sigma_d (1 + (Z_{1,1}^d)^N) (1 + (x_1^d)^{2N}) \left(|Z_{1,1}^d| + \frac{1}{r} (1 + |x_1^d|) \right), \end{aligned}$$

for some $C > 0$.

Set S_d to be the set in which $1 + (x_1^d)^{2N+1} \leq d^{\alpha/2}$, applying Markov's inequality we obtain

$$d^{2\alpha} \pi_d(S_d^c) = d^{2\alpha} \pi_d \left((1 + (x_1^d)^{2N+1})^5 \geq d^{5\alpha/2} \right) \leq d^{-\alpha/2} \mathbb{E} \left[(1 + (x_1^d)^{2N+1})^5 \right] \xrightarrow{d \rightarrow \infty} 0.$$

Recalling that $|Z_{1,1}^d|$ and $1 + (Z_{1,1}^d)^N$ are bounded, we have that

$$\sup_{\mathbf{x}^d \in S_d} \left| \mathbf{L}_d V(\mathbf{x}^d) - \tilde{\mathbf{L}}_d V(\mathbf{x}^d) \right| \leq C d^{\alpha/2} \frac{\ell}{d^\alpha} \xrightarrow{d \rightarrow \infty} 0.$$

The second results follows from (58) using the same argument. $\square$

The following result considers the convergence to the generator of the Langevin diffusion (53).

Proposition 6. *Under **A0**, **A1** and **A2**, if $\alpha = 1/4$, $\beta = 1/8$ and $r > 0$, there exists sets $T_d \subseteq \mathbb{R}^d$ with $d^{2\alpha} \pi_d(T_d^c) \rightarrow 0$ as $d \rightarrow \infty$, such that for any $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$*

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in T_d} \left| d^{2\alpha} \mathbb{E} [V(y_d(x_1^d, Z_{1,1}^d)) - V(x_1^d)] - \frac{\ell^2}{2} (V''(x_1^d) + g'(x_1^d) V'(x_1^d)) \right| = 0.$$

Proof. Take

$$y_d(x_1^d, Z_{1,1}^d) = x_1^d + \frac{\sigma_d^2}{2} g' \left(\text{prox}_g^{\sigma_d^{2m} r/2}(x_1^d) \right) + \sigma_d Z_{1,1}^d,$$

and use a Taylor expansion of order 2 of

$$W(x, z, \sigma) = V \left[x + \frac{\sigma^2}{2} g' \left(\text{prox}_g^{\sigma^{2m} r/2}(x) \right) + \sigma z \right],$$

about $\sigma = 0$ with integral remainder:

$$\begin{aligned} W(x, z, \sigma) &= W(x, z, 0) + \frac{\partial W}{\partial \sigma}(x, z, \sigma) \Big|_{\sigma=0} \sigma + \frac{1}{2} \frac{\partial^2 W}{\partial \sigma^2}(x, z, \sigma) \Big|_{\sigma=0} \sigma^2 \\ &\quad + \int_0^\sigma \frac{\partial^3 W}{\partial \sigma^3}(x, z, u) \Big|_{\sigma=u} \frac{(\sigma - u)^2}{2} du. \end{aligned}$$

Using the derivatives

$$\begin{aligned} W(x, z, 0) &= V(x) , \quad \frac{\partial W}{\partial \sigma}(x, z, \sigma) \Big|_{\sigma=0} = V'(x)z , \\ \frac{\partial^2 W}{\partial \sigma^2}(x, z, \sigma) &= V''(x)z^2 + V'(x)g'(x) , \end{aligned}$$

and recalling that $\mathbb{E}[Z_{1,1}^d] = 0$, $\mathbb{E}[(Z_{1,1}^d)^2] = 1$, we have

$$\begin{aligned} \mathbb{E}[V(y_d(x_1^d, Z_{1,1}^d)) - V(x_1^d)] &= \frac{\sigma_d^2}{2} [V''(x_1^d) + V'(x_1^d)g'(x_1^d)] \\ &\quad + \mathbb{E} \left[\int_0^{\sigma_d} \frac{\partial^3 W}{\partial \sigma^3}(x_1^d, Z_{1,1}^d, \sigma) \Big|_{\sigma=u} \frac{(\sigma_d - u)^2}{2} du \right] . \end{aligned}$$

Proceeding as in the previous proposition, we can bound

$$\left| \int_0^{\sigma_d} \frac{\partial^3 W}{\partial \sigma^3}(x_1^d, Z_{1,1}^d, \sigma) \Big|_{\sigma=u} \frac{(\sigma_d - u)^2}{2} du \right| \leq A (1 + (Z_{1,1}^d)^N) (1 + (x_1^d)^{2N}) d^{-3\alpha} .$$

Setting T_d to be the set in which $(1 + (x_1^d)^{2N}) \leq d^{\alpha/2}$, the result follows by applying Markov's inequality as in Proposition 5. $\square$

Before proceeding to stating and proving the last auxiliary result, let us denote by $\psi_1 : \mathbb{R} \rightarrow [0, +\infty)$ the characteristic function of the distribution $N(0, \ell^4 K_1^2(r))$, where $K_1^2(r)$ is given in Theorem 1-(a),

$$\psi_1(t) = \exp(-t^2 \ell^4 K_1^2(r)/2) ,$$

and by $\psi_1^d(\mathbf{x}^d; t) = \int \exp(itw) \mathcal{Q}_1^d(\mathbf{x}^d; dw)$ the characteristic functions associated with the law

$$\mathcal{Q}_1^d(\mathbf{x}^d; \cdot) = \mathcal{L} \left\{ d^{-1/2} \sum_{i=2}^d C_2(x_i^d, Z_{1,i}^d) \right\} .$$

Proposition 7. *Under A0, A1 and A2, if $\alpha = 1/4$, $\beta = 1/8$ and $r > 0$, there exists a sequence of sets $H_d \subseteq \mathbb{R}^d$ such that*

- (i) $\lim_{d \rightarrow \infty} d^{2\alpha} \pi_d(H_d^c) = 0$,
- (ii) for all $t \in \mathbb{R}$, $\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in H_d} |\psi_1^d(\mathbf{x}^d; t) - \psi_1(t)| = 0$,
- (iii) for all bounded continuous function $\chi : \mathbb{R} \rightarrow \mathbb{R}$,

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in H_d} \left| \int_{\mathbb{R}} \mathcal{Q}_1^d(\mathbf{x}^d; du) \chi(u) - (2\pi \ell^4 K_1^2(r))^{-1/2} \int_{\mathbb{R}} \chi(u) e^{-u^2/(2\ell^4 K_1^2(r))} du \right| = 0 ,$$

(iv) in particular,

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in H_d} \left| \mathbb{E} \left[1 \wedge \exp \left(d^{-1/2} \sum_{i=2}^d C_2(x_i^d, Z_{1,i}^d) - \frac{\ell^4 K_1^2(r)}{2} \right) \right] - 2\Phi \left(-\frac{\ell^2 K_1(r)}{2} \right) \right| = 0 ,$$

where Φ is the distribution function of the standard normal random variable.

Proof. (i) Define the functions $h_j(x) = [-g''(x)g'(x)]^j$ with $j = 1, \dots, 4$ and let $H_d = H_{d,1} \cap H_{d,2}$ where

$$H_{d,1} = \left\{ \mathbf{x}^d \in \mathbb{R}^d : \left| \frac{1}{d} \sum_{i=2}^d h_j(x_i^d) - \int_{\mathbb{R}} h_j(u) \pi(u) du \right| \leq d^{1/3} \text{ for } j = 1, \dots, 4 \right\},$$

$$H_{d,2} = \left\{ \mathbf{x}^d \in \mathbb{R}^d : |h_j(x_i^d)| \leq d^{2/3} \text{ for } i = 1, \dots, d \text{ and } j = 1, \dots, 4 \right\}.$$

Using Chebychev's inequality, the fact that the derivatives of g are bounded by polynomials and that π has finite moments, we have $d^{1/2} \pi_d((H_{d,1})^c) \rightarrow 0$ as $d \rightarrow \infty$. Similarly, by Markov's inequality we have $d^{1/2} \pi_d((H_{d,2})^c) \rightarrow 0$ as $d \rightarrow \infty$.

(ii) We follow [34, Lemma 3(b)] and decompose

$$\begin{aligned} |\psi_1^d(\mathbf{x}^d; t) - \psi_1(t)| &\leq \left| \psi_1^d(\mathbf{x}^d; t) - \prod_{i=2}^d \left(1 - \frac{t^2}{2d} v(x_i^d) \right) \right| \\ &\quad + \left| \prod_{i=2}^d \left(1 - \frac{t^2}{2d} v(x_i^d) \right) - \prod_{i=2}^d \exp \left(-t^2 \frac{v(x_i^d)}{2d} \right) \right| \\ &\quad + \left| \prod_{i=2}^d \exp \left(-t^2 \frac{v(x_i^d)}{2d} \right) - \exp \left(-t^2 \frac{\ell^4 K_1(r)^2}{2} \right) \right|, \end{aligned} \quad (60)$$

where $v(x_i^d) = \text{Var}(C_2(x_i^d, Z_{1,i}^d)) = \mathbb{E}[C_2(x_i^d, Z_{1,i}^d)^2]$, where the expectation is taken w.r.t. $Z_{1,i}^d$. For the first term, decompose the characteristic function $\psi_1^d(\mathbf{x}^d; t) = \prod_{i=2}^d \theta_i^d(x_i^d; t)$ as the product of the characteristic functions of $d^{-1/2} W_i$ where we define $W_i = C_2(x_i^d, Z_{1,i}^d)$, using [15, equation (3.3.3)] as in the proof of [15, Theorem 3.4.10] we obtain

$$\begin{aligned} \left| \theta_i^d(x_i^d; t) - \left(1 - \frac{t^2}{2d} v(x_i^d) \right) \right| &\leq \mathbb{E} \left[\frac{|t|^3}{d^{3/2}} \frac{|W_i|^3}{3!} \wedge \frac{2|t|^2}{d} \frac{|W_i|^2}{2!} \right] \\ &\leq \mathbb{E} \left[\frac{|t|^3}{d^{3/2}} \frac{|W_i|^3}{3!}; |W_i| \leq d^{1/2} \varepsilon \right] + \frac{t^2}{d} \mathbb{E} \left[|W_i|^2; |W_i| > d^{1/2} \varepsilon \right] \\ &\leq \frac{\varepsilon |t|^3}{6d} \mathbb{E} [|W_i|^2] + \frac{t^2}{\varepsilon^2 d^2} \mathbb{E} [|W_i|^4], \end{aligned}$$

for any $\varepsilon > 0$. For sufficiently large d , we have that $t^2 v(x_i^d)/(2d) \leq 1$ for $\mathbf{x} \in H_{d,2}$, and we can use [15, Lemma 3.4.3]

$$\begin{aligned} \left| \psi_j^d(\mathbf{x}^d; t) - \prod_{i=2}^d \left(1 - \frac{t^2}{2d} v(x_i^d) \right) \right| &\leq \sum_{i=2}^d \left(\frac{\varepsilon |t|^3}{6d} \mathbb{E} [|W_i|^2] + \frac{t^2}{\varepsilon^2 d^2} \mathbb{E} [|W_i|^4] \right) \\ &\leq \frac{\varepsilon \ell^4 |t|^3}{6} (K_1^2(r) + D_1 d^{-1/3}) + \frac{t^2}{\varepsilon^2 d} (\mathbb{E} [|W_i|^4] + D_2 \ell^8 d^{-1/4}), \end{aligned}$$

where the last inequality follows from the fact that $\mathbf{x}^d \in H_{d,2}$ and D_1, D_2 are positive constants. For any $\delta > 0$ we can chose ε small enough so that the first term in the above is less

than $\delta/2$ and we can chose d sufficiently large to make the second term less than $\delta/2$. Thus, for any $\delta > 0$ we can find $\varepsilon > 0$ and $d \in \mathbb{N}$ such that

$$\left| \psi_1^d(\mathbf{x}^d; t) - \prod_{i=2}^d \left(1 - \frac{t^2}{2d} v(x_i^d) \right) \right| < \delta ,$$

the uniform convergence then follows. The second term in (60) converges to 0 uniformly for all $\mathbf{x}^d \in H_{d,1}$; while for the third term in (60) we use again [15, Lemma 3.4.3]

$$\left| \prod_{i=2}^d \exp \left(-t^2 \frac{v(x_i^d)}{2d} \right) - \exp \left(-t^2 \frac{\ell^4 K_1(r)^2}{2} \right) \right| \leq \sum_{i=2}^d \frac{t^4 v(x_i^d)^2}{4d^2} ,$$

which goes to zero when $d \rightarrow \infty$, for all $\mathbf{x}^d \in H_{d,2}$. The result then follows.

- (iii) Let $\chi : \mathbb{R} \rightarrow \mathbb{R}$ be a bounded and continuous function. Define the sequence $\{\mathbf{x}^d : d \in \mathbb{N}^*\}$, where, for any $d \in \mathbb{N}^*$, $\mathbf{x}^d \in H_d$ satisfies,

$$\begin{aligned} & \sup_{y^d \in H_d} \left| \int_{\mathbb{R}} \mathcal{Q}_1^d(y_i^d; du) \chi(u) - (2\pi\ell^4 K_1^2(r))^{-1/2} \int_{\mathbb{R}} \chi(u) e^{-u^2/(2\ell^4 K_1^2(r))} du \right| \\ & \leq \left| \int_{\mathbb{R}} \mathcal{Q}_1^d(x_i^d; du) \chi(u) - (2\pi\ell^4 K_1^2(r))^{-1/2} \int_{\mathbb{R}} \chi(u) e^{-u^2/(2\ell^4 K_1^2(r))} du \right| + d^{-1} . \end{aligned}$$

Then, using (ii) and Lévy's continuity theorem (e.g. [41, Theorem 1, page 322]), we obtain

$$\lim_{d \rightarrow \infty} \left| \int_{\mathbb{R}} \mathcal{Q}_1^d(x_i^d; du) \chi(u) - (2\pi\ell^4 K_1^2(r))^{-1/2} \int_{\mathbb{R}} \chi(u) e^{-u^2/(2\ell^4 K_1^2(r))} du \right| = 0 .$$

Combining this limit with the definition of the sequence $\{\mathbf{x}^d : d \in \mathbb{N}^*\}$, we conclude the proof.

- (iv) This statement follows directly from (iii) and [33, Proposition 2.4]. □

A.2 Auxiliary Results for the Proof of Case (b)

First, we characterize the limit behaviour of the acceptance ratio (12). The following result is an extension of [34, Lemma 1].

Proposition 8. *Under A0, A1, A2 and A3, if $\alpha = 1/6$, $\beta = 1/6$ and $r > 0$, then*

- (i) *the log-acceptance ratio (50), when $d \rightarrow \infty$, satisfies the following Taylor expansion*

$$\begin{aligned} \phi_d(x, z) &= d^{-1/2} C_3(x, z) + d^{-2/3} C_4(x, z) \\ &\quad + d^{-5/6} C_5(x, z) + d^{-1} C_6(x, z) + C_7(x, z, \sigma_d) , \end{aligned}$$

where C_3 is given in (61), C_4, C_5, C_6 are polynomials in z and the derivatives of g , such that $\mathbb{E}[C_j(X_{0,1}^d, Z_{1,1}^d)] = 0$ for $j = 3, 4, 5$ and $\mathbb{E}[C_3(X_{0,1}^d, Z_{1,1}^d)^2] = -2\mathbb{E}[C_6(X_{0,1}^d, Z_{1,1}^d)]$,

(ii) there exists sets $F_d \subseteq \mathbb{R}^d$ with $d^{2\alpha}\pi_d(F_d^c) \rightarrow 0$ such that

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in F_d} \mathbb{E} \left[\left| \sum_{i=2}^d \phi_d(x_i^d, Z_{1,i}^d) - d^{-1/2} \sum_{i=2}^d C_3(x_i^d, Z_{1,i}^d) + \frac{\ell^6 K_2(r)^2}{2} \right| \right] = 0 ,$$

where $K_2(r)$ is given in Theorem 1-(b).

Proof. Take one component of the log-acceptance ratio

$$\phi_d(x, z) = g(x) - g(y_d(x, z)) + \log q(y_d(x, z), x) - \log q(x, y_d(x, z)) ,$$

with $y_d(x, z) = x - \sigma_d^2 g'(\text{prox}_g^{\sigma_d^{2m} r/2}(x)) + \sigma_d z$. Proceeding as in the proof for case (a), we have that $\phi_d(x, z) = R_1(x, z, \sigma_d) + R_2(x, z, \sigma_d)$ where R_1, R_2 are given in (55). Following the approach of [34] we approximate $\phi_d(x, z)$ with a Taylor expansion about $\sigma_d = 0$.

(i) Using a Taylor expansion of order 7, we find that

$$\begin{aligned} \phi_d(x, z) &= d^{-1/2} C_3(x, z) + d^{-2/3} C_4(x, z) + d^{-5/6} C_5(x, z) \\ &\quad + d^{-1} C_6(x, z) + C_7(x, z, \sigma_d) , \end{aligned}$$

where

$$C_3(x, z) = \frac{\ell^3}{6} \left(\frac{1}{2} g'''(x) z^3 - \frac{3}{2} g''(x) g'(x) z (1 + 2r) \right) , \quad (61)$$

$C_4(x, z), C_5(x, z)$ and $C_6(x, z)$ are given in Appendix C.1.2 and integral form of the remainder

$$C_7(x, z, \sigma_d) = \int_0^{\sigma_d} \frac{\partial^7}{\partial \sigma^7} R(x, z, \sigma) \Big|_{\sigma=u} \frac{(\sigma_d - u)^6}{6!} du ,$$

with u between 0 and σ_d and the derivatives of R_1 and R_2 are given in Appendix C.2. In addition, integrating by parts and using the moments of the standard normal $Z_{1,1}^d$, we find that $\mathbb{E}[C_3(X_{0,1}^d, Z_{1,1}^d)] = \mathbb{E}[C_4(X_{0,1}^d, Z_{1,1}^d)] = \mathbb{E}[C_5(X_{0,1}^d, Z_{1,1}^d)] = 0$ and

$$\begin{aligned} \mathbb{E}[C_6(X_{0,1}^d, Z_{1,1}^d)] &= -\frac{\ell^6}{16} (r + 2r^2) \mathbb{E} \left[(g''(X_{0,1}^d) g'(X_{0,1}^d))^2 \right] \\ &\quad - \frac{\ell^6}{16} \left(\frac{1}{2} + r \right) \mathbb{E} [g''(X_{0,1}^d)^3] - \frac{5\ell^6}{96} \mathbb{E} [g'''(X_{0,1}^d)^2] , \end{aligned}$$

which shows that $\mathbb{E}[C_3(X_{0,1}^d, Z_{1,1}^d)^2 + 2C_6(X_{0,1}^d, Z_{1,1}^d)] = 0$.

(ii) The proof of this result follows using the same steps as case (a) and is analogous to that of [34, Lemma 1]. □

Next, we compare the generator L_d and $\tilde{L}_d$ in (51) and (52) respectively, extending [34, Theorem 3].

Proposition 9. Under **A0**, **A1**, **A2** and **A3**, if $\alpha = 1/6$, $\beta = 1/6$ and $r > 0$, there exists sets $S_d \subseteq \mathbb{R}^d$ with $d^{2\alpha}\pi_d(S_d^c) \rightarrow 0$ such that for any $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in S_d} \left| L_d V(\mathbf{x}^d) - \tilde{L}_d V(\mathbf{x}^d) \right| = 0 ,$$

and

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in S_d} \mathbb{E} \left[\left| \left(\exp \left(\sum_{i=1}^d \phi_d(x_i^d, Z_{1,i}^d) \right) \wedge 1 \right) - \left(\exp \left(\sum_{i=2}^d \phi_d(x_i^d, Z_{1,i}^d) \right) \wedge 1 \right) \right| \right] = 0 .$$

Proof. Proceeding as in Proposition 5, for any $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$, there exists a constant C such that

$$\begin{aligned} |V(y_d(x_1^d, Z_{1,1}^d)) - V(x_1^d)| &\leq C |y_d(x_1^d, Z_{1,1}^d) - x_1^d| \\ &\leq C \left(\sigma_d |Z_{1,1}^d| + \frac{\sigma_d^2}{2} \left| g' \left(\text{prox}_g^{\sigma_d^{2m} r/2}(x_1^d) \right) \right| \right) . \end{aligned}$$

Under **A3**, g' is Lipschitz continuous and we have, for some $C \geq 1$,

$$\left| g' \left(\text{prox}_g^{\sigma_d^{2m} r/2}(x_1^d) \right) \right| \leq C \left(1 + \left| \text{prox}_g^{\sigma_d^{2m} r/2}(x_1^d) \right| \right) \leq C(1 + |x_1^d|),$$

where we used the fact that prox_g^λ is 1-Lipschitz continuous for all $\lambda > 0$. The result then follows similarly to Proposition 5 and [34, Theorem 3]. $\square$

The following result considers the convergence to the generator of the Langevin diffusion (53) and is a generalization of [34, Lemma 2].

Proposition 10. Under **A0**, **A1**, **A2** and **A3**, if $\alpha = 1/6$, $\beta = 1/6$ and $r > 0$, there exists sets $T_d \subseteq \mathbb{R}^d$ with $d^{2\alpha}\pi_d(T_d^c) \rightarrow 0$ as $d \rightarrow \infty$ such that for any $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in T_d} \left| d^{2\alpha} \mathbb{E} [V(y_d(x_1^d, Z_{1,1}^d)) - V(x_1^d)] - \frac{\ell^2}{2} (V''(x_1^d) + g'(x_1^d) V'(x_1^d)) \right| = 0 .$$

Proof. The proof is identical to that of Proposition 6. $\square$

Before proceeding to stating and proving the last auxiliary result, let us denote by $\psi_2 : \mathbb{R} \rightarrow [0, +\infty)$ the characteristic function of the distribution $N(0, \ell^6 K_2^2(r))$, where $K_2^2(r)$ is given in Theorem 1-(b),

$$\psi_2(t) = \exp(-t^2 \ell^6 K_2^2(r)/2) ,$$

and by $\psi_2^d(\mathbf{x}^d; t) = \int \exp(itw) \mathcal{Q}_2^d(\mathbf{x}^d; dw)$ the characteristic functions associated with the law

$$\mathcal{Q}_2^d(\mathbf{x}^d; \cdot) = \mathcal{L} \left\{ d^{-1/2} \sum_{i=2}^d C_3(x_i^d, Z_{1,i}^d) \right\} .$$

The following result extends [34, Lemma 3].

Proposition 11. Under **A0**, **A1**, **A2** and **A3**, if $\alpha = 1/6$, $\beta = 1/6$ and $r > 0$, there exists a sequence of sets $H_d \subseteq \mathbb{R}^d$ such that

- (i) $\lim_{d \rightarrow \infty} d^{2\alpha} \pi_d(H_d^c) = 0$,
- (ii) for all $t \in \mathbb{R}$, $\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in H_d} |\psi_2^d(\mathbf{x}^d; t) - \psi_2(t)| = 0$,
- (iii) for all bounded continuous function $\chi : \mathbb{R} \rightarrow \mathbb{R}$,

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in H_d} \left| \int_{\mathbb{R}} \mathcal{Q}_2^d(\mathbf{x}^d; du) \chi(u) - (2\pi \ell^6 K_2^2(r))^{-1/2} \int_{\mathbb{R}} \chi(u) e^{-u^2/(2\ell^6 K_2^2(r))} du \right| = 0,$$

- (iv) in particular,

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in H_d} \left| \mathbb{E} \left[1 \wedge \exp \left(d^{-1/2} \sum_{i=2}^d C_3(x_i^d, Z_{1,i}^d) - \frac{\ell^6 K_2^2(r)}{2} \right) \right] - 2\Phi \left(-\frac{\ell^3 K_2(r)}{2} \right) \right| = 0,$$

where Φ is the distribution function of the standard normal random variable.

Proof. (i) The proof is analogous to that of Proposition 7 and follows the same steps of that of [34, Lemma 3(a)].

- (ii) The proof is analogous to that of Proposition 7 and follows the same steps of that of [34, Lemma 3(b)].

- (iii) Following the steps of (iii) in Proposition 7 and the Lévy's continuity Theorem (e.g. [41, Theorem 1, page 322]) bring the result.

- (iv) This statement follows directly from (iii) and [33, Proposition 2.4].

□

A.3 Auxiliary Results for the Proof of Case (c)

First, we characterize the limit behaviour of the acceptance ratio (12).

Proposition 12. Under **A0**, **A1**, **A2** and **A3** and, if $\alpha = 1/6$, $\beta = m/6$ for $m > 1$ and $r > 0$, then

- (i) the log-acceptance ratio (50), when $d \rightarrow \infty$, satisfies the following Taylor expansion

$$\begin{aligned} \phi_d(x, z) &= d^{-1/2} C_3(x, z) + d^{-2/3} C_4(x, z) \\ &\quad + d^{-5/6} C_5(x, z) + d^{-1} C_6(x, z) + C_7(x, z, \sigma_d), \end{aligned}$$

where C_3 is given in (61), C_4, C_5, C_6 are polynomials in z and the derivatives of g , such that $\mathbb{E}[C_j(X_{0,1}^d, Z_{1,1}^d)] = 0$ for $j = 3, 4, 5$ and $\mathbb{E}[C_3(X_{0,1}^d, Z_{1,1}^d)^2] = -2\mathbb{E}[C_6(X_{0,1}^d, Z_{1,1}^d)]$,

- (ii) there exists sets $F_d \subseteq \mathbb{R}^d$ with $d^{2\alpha} \pi_d(F_d^c) \rightarrow 0$ such that

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in F_d} \mathbb{E} \left[\left| \sum_{i=2}^d \phi_d(x_i^d, Z_{1,i}^d) - d^{-1/2} \sum_{i=2}^d C_3(x_i^d, Z_{1,i}^d) + \frac{\ell^6 K_2^2(0)}{2} \right| \right] = 0,$$

where K_2 is given in Theorem 1-(b).

Proof. Take one component of the log-acceptance ratio

$$\phi_d(x, z) = g(x) - g(y_d(x, z)) + \log q(y_d(x, z), x) - \log q(x, y_d(x, z)) ,$$

with $y_d(x, z) = x - \sigma_d^2 g'(\text{prox}_g^{\sigma_d^{2m} r/2}(x)) + \sigma_d z$. Proceeding as in the proof for case (a), we have that $\phi_d(x, z) = R_1(x, z, \sigma_d) + R_2(x, z, \sigma_d)$ where R_1, R_2 are given in (55). Following the approach of [34] we approximate $\phi_d(x, z)$ with a Taylor expansion about $\sigma_d = 0$.

(i) Using a Taylor expansion of order 7, we find that

$$\begin{aligned} \phi_d(x, z) &= d^{-1/2} C_3(x, z) + d^{-2/3} C_4(x, z) + d^{-5/6} C_5(x, z) \\ &\quad + d^{-1} C_6(x, z) + C_7(x, z, \sigma_d) , \end{aligned}$$

where

$$C_3(x, z) = \frac{\ell^3}{6} \left(\frac{1}{2} g'''(x) z^3 - \frac{3}{2} g''(x) g'(x) z \right) ,$$

$C_4(x, z), C_5(x, z)$ and $C_6(x, z)$ are given in Appendix C.1.3 and integral form of the remainder

$$C_7(x, z, \sigma_d) = \int_0^{\sigma_d} \frac{\partial^7}{\partial \sigma^7} R(x, z, \sigma) \Big|_{\sigma=u} \frac{(\sigma_d - u)^6}{6!} du ,$$

with u between 0 and σ_d and the derivatives of R_1 and R_2 are given in Appendix C.2. In addition, integrating by parts and using the moments of $Z_{1,1}^d$ we find that $\mathbb{E}[C_3(X_{0,1}^d, Z_{1,1}^d)] = \mathbb{E}[C_4(X_{0,1}^d, Z_{1,1}^d)] = \mathbb{E}[C_5(X_{0,1}^d, Z_{1,1}^d)] = 0$ and

$$\mathbb{E}[C_6(X_{0,1}^d, Z_{1,1}^d)] = \ell^6 \left(-\frac{1}{32} \mathbb{E}[g''(X_{0,1}^d)^3] - \frac{5}{96} \mathbb{E}[g'''(X_{0,1}^d)^2] \right) ,$$

which shows that $\mathbb{E}[C_3(X_{0,1}^d, Z_{1,1}^d)^2 + 2C_6(X_{0,1}^d, Z_{1,1}^d)] = 0$.

(ii) The proof of this result follows using the same steps as case (a) and is analogous to that of [34, Lemma 1].

□

Next, we compare the generator L_d and $\tilde{L}_d$ in (51) and (52) respectively.

Proposition 13. *Under A0, A1, A2 and A3, if $\alpha = 1/6$, $\beta = m/6$ for $m > 1$ and $r > 0$, there exists sets $S_d \subseteq \mathbb{R}^d$ with $d^{2\alpha} \pi_d(S_d^c) \rightarrow 0$ as $d \rightarrow \infty$ such that for any $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$*

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in S_d} \left| L_d V(\mathbf{x}^d) - \tilde{L}_d V(\mathbf{x}^d) \right| = 0 ,$$

and

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in S_d} \mathbb{E} \left[\left| \left(\exp \left(\sum_{i=1}^d \phi_d(x_i^d, Z_{1,i}^d) \right) \wedge 1 \right) - \left(\exp \left(\sum_{i=2}^d \phi_d(x_i^d, Z_{1,i}^d) \right) \wedge 1 \right) \right| \right] = 0 .$$

Proof. The proof is identical to that of Proposition 9. $\square$

The following result considers the convergence to the generator of the Langevin diffusion (53).

Proposition 14. *Under A0, A1, A2 and A3, if $\alpha = 1/6$, $\beta = m/6$ for $m > 1$ and $r > 0$, there exists sets $T_d \subseteq \mathbb{R}^d$ with $d^{2\alpha}\pi_d(T_d^c) \rightarrow 0$ as $d \rightarrow \infty$ such that for any $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$*

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in T_d} \left| d^{2\alpha} \mathbb{E} [V(y_d(x_1^d, Z_{1,1}^d)) - V(x_1^d)] - \frac{\ell^2}{2} (V''(x_1^d) + g'(x_1^d)V'(x_1^d)) \right| = 0 .$$

Proof. The proof is identical to that of Proposition 6. $\square$

Before proceeding to stating and proving the last auxiliary result, let us denote by $\psi_3 : \mathbb{R} \rightarrow [0, +\infty)$ the characteristic function of the distribution $N(0, \ell^6 K_2^2(0))$, where K_2^2 is given in Theorem 1-(b),

$$\psi_3(t) = \exp(-t^2 \ell^6 K_2^2(0)/2) ,$$

and by $\psi_3^d(\mathbf{x}^d; t) = \int \exp(itw) \mathcal{Q}_3^d(\mathbf{x}^d; dw)$ the characteristic functions associated with the law

$$\mathcal{Q}_3^d(\mathbf{x}^d; \cdot) = \mathcal{L} \left\{ d^{-1/2} \sum_{i=2}^d C_3(x_i^d, Z_{1,i}^d) \right\} .$$

Proposition 15. *Under A0, A1, A2 and A3, if $\alpha = 1/6$, $\beta = m/6$ for $m > 1$ and $r > 0$, there exists a sequence of sets $H_d \subseteq \mathbb{R}^d$ such that*

- (i) $\lim_{d \rightarrow \infty} d^{2\alpha} \pi_d(H_d^c) = 0$,
- (ii) for all $t \in \mathbb{R}$, $\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in H_d} |\psi_3^d(\mathbf{x}^d; t) - \psi_3(t)| = 0$,
- (iii) for all bounded continuous function $\chi : \mathbb{R} \rightarrow \mathbb{R}$,

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in H_d} \left| \int_{\mathbb{R}} \mathcal{Q}_3^d(\mathbf{x}^d; du) \chi(u) - (2\pi \ell^6 K_2^2(0))^{-1/2} \int_{\mathbb{R}} \chi(u) e^{-u^2/(2\ell^6 K_2^2(0))} du \right| = 0 ,$$

(iv) in particular,

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in H_d} \left| \mathbb{E} \left[1 \wedge \exp \left(d^{-1/2} \sum_{i=2}^d C_3(x_i^d, Z_{1,i}^d) - \frac{\ell^6 K_2^2(0)}{2} \right) \right] - 2\Phi \left(-\frac{\ell^3 K_2(0)}{2} \right) \right| = 0 ,$$

where Φ is the distribution function of the standard normal random variable.

Proof. (i) The proof is analogous to that of Proposition 7 and follows the same steps of that of [34, Lemma 3(a)].

(ii) The proof is analogous to that of Proposition 7 and follows the same steps of that of [34, Lemma 3(b)].

(iii) Following the steps of (iii) in Proposition 7 and the Lévy's continuity Theorem (e.g. [41, Theorem 1, page 322]) bring the result.

(iv) This statement follows directly from (iii) and [33, Proposition 2.4]. $\square$

A.4 Proof of Theorem 1

Proof of Theorem 1. (a) The asymptotic acceptance rate follows by combining Propositions 4–6 with part (iv) of Proposition 7 as in the proof of [34, Theorem 1]. To prove the weak convergence of the process it suffices to show that there exists events $F_d^* \in \mathbb{R}^d$ such that for all $t > 0$

$$\lim_{d \rightarrow \infty} \mathbb{P}(L_t^d \in F_d^* \text{ for all } 0 \leq s \leq t) = 1 ,$$

and

$$\lim_{d \rightarrow \infty} \sup_{\mathbf{x}^d \in F_d^*} |L_d V(\mathbf{x}^d) - LV(\mathbf{x}^d)| .$$

for all $V \in C_c^\infty(\mathbb{R}, \mathbb{R})$ [17, Chapter 4, Corollary 8.7]. We take $F_d^* = F_d \cap S_d \cap T_d \cap H_d$. Then, $d^{2\alpha} \pi_d((F_d^*)^c) \rightarrow 0$ and

$$\lim_{d \rightarrow \infty} \mathbb{P}(L_t^d \in F_d^* \text{ for all } 0 \leq s \leq t) = 1 ,$$

for all fixed t . Combining Propositions 4–7 we obtain convergence of the generators.

To obtain the value of $a(\ell, r)$ maximizing the speed, we observe that $K_1^2(r)$ is a function of the ratio $r = c^2/\ell^{2m} = c^2/\ell$ only, we can take $c \propto \ell^{1/2}$ so that $K_1^2(r)$ is constant for given c . Using the same substitution as in [34, Theorem 2] we find that $h(\ell, r)$ is maximized at the unique value of ℓ such that $a(\ell, r) = 0.452$.

- (b) The proof is analogous to that of case (a) replacing Propositions 4, 5, 6 and 7 with Propositions 8, 9, 10 and 11. To obtain the value of $a(\ell, r)$ maximizing the speed, we observe that $K_2^2(r)$ is a function of the ratio $r = c^2/\ell^{2m} = c^2/\ell^2$ only, we can take $c \propto \ell$ so that $K_2^2(r)$ is constant for given c . Using the same substitution as in [34, Theorem 2] we find that $h(\ell, r)$ is maximized at the unique value of ℓ such that $a(\ell, r) = 0.574$.
- (c) The proof is analogous to that of case (a) replacing Propositions 4, 5, 6 and 7 with Propositions 12, 13, 14 and 15. To obtain the value of $a(\ell, r)$ maximizing the speed, we observe that $K_2^2(0)$ is constant w.r.t. r , we can use the same substitution as in [34, Theorem 2] we find that $h(\ell, r)$ is maximized at the unique value of ℓ such that $a(\ell, r) = 0.574$.

□

B Numerical Experiments

B.1 Differentiable Targets

We collect here a number of numerical experiments confirming the results in Section 3.2. To do so, we consider the Gaussian distribution in Example 1 and four algorithmic settings summarized in Table 1 which correspond to the three cases identified in Theorem 1 and MALA.

The first plot in Figure 5–8 show that for values of α different from those identified in Theorem 1 the acceptance ratio $a_d(\ell, r)$ becomes degenerate as d increases. For the values of α identified in Theorem 1 we analyze the influence of ℓ on the acceptance $a_d(\ell, r)$ (second plot), obtaining for

Case	Figure	Algorithm	α	β	m	r
(a)	5	Proximal MALA	1/4	1/8	1/2	1
(b)	6	P-MALA	1/6	1/6	1	1
(c)	7	Proximal MALA	1/6	1/2	3	2
—	8	MALA	1/6	1/6	1	≈ 0

Table 1: Algorithm setting for the simulation study on the Gaussian target.

$d \rightarrow +\infty$ the expression given by Theorem 1-(a) for Figure 5, the expression in Theorem 1-(b) for Figures 6 and 8 and that in Theorem 1-(c) for Figure 7.

Finally, we consider the relationship between acceptance ratio $a_d(\ell, r)$ and the speed of the diffusion $h(\ell, r)$ approximated by the expected squared jumping distance (see, e.g. [18])

$$\text{ESJD}_d := d^{2\alpha} \mathbb{E} [(X_0^d - X_1^d)^2] .$$

Looking at the last plot in Figure 5–8 we find that, even for relatively small values of d , the shape of the plot of ESJD_d as a function of the acceptance $a_d(\ell, r)$ is similar to that of the theoretical limit. This suggests that tuning the acceptance ratio to be approximately 0.452 when $\alpha = 1/4, \beta = 1/8$ and approximately 0.574 when $\alpha = 1/6, \beta = m/6$ with $m \geq 1$ should generally guarantee high efficiency.

B.2 Laplace Target

We collect here a number of numerical experiments confirming the results for the Laplace distribution in Section 3.2. Similarly to Appendix B.1 we consider three algorithmic settings, summarized in Table 2.

Figure	Algorithm	α	β	m	r
9	sG-MALA	1/3	1/3	1	0
10	P-MALA	1/3	1/3	1	1
11	Proximal MALA	1/3	1	3	2

Table 2: Algorithm setting for the simulation study on the Laplace target.

The first plot in Figures 9–11 shows that for $\alpha \neq 1/3$ the acceptance ratio $a_d(\ell, r)$ becomes degenerate as d increases; while the second plot shows that $(a_d(\ell, r))_{d \in \mathbb{N}^*}$ and $(\text{ESJD}_d)_{d \in \mathbb{N}^*}$ converge to $a^L(\ell)$ and $h^L(\ell)$ given in Theorems 2 and 3, respectively. In the case $m = 3, r = 2$ in Figure 11, we find that the behaviour for low values of d significantly differs from the limiting one. For values of d lower than 130 the ESJD_d achieves its maximum at a value of the acceptance $a_d(\ell, r)$ different from that predicted by Theorem 3. In practice, this might mean that for low dimensional settings the recommended choice of $a(\ell, c) = 0.360$ is far from optimal. Similar behaviours for small d have also been observed in the case of RWM and MALA (e.g., [40, Section 2.1]).

B.3 Mix of a Laplace and differentiable target

We collect here a the rest of the numerical experiments illustrating the scaling the results for the density defined in (19), discussed in Section 4.1. Similarly to Appendix B.2 we consider three




Figure 5: **Case (a)**: Proximal MALA with Gaussian target and $m = 1/2, r = 1$. Average acceptance rate for different choices of α (first); acceptance rate as a function of ℓ for increasing dimension d (second); ESJD_d as a function of the acceptance rate $a_d(\ell, r)$ (third).




Figure 6: **Case (b)**: Proximal MALA with Gaussian target and $m=1, r=1$ (P-MALA). Average acceptance rate for different choices of α (first); acceptance rate as a function of ℓ for increasing dimension d (second); ESJD_d as a function of the acceptance rate $a_d(\ell, r)$ (third).




Figure 7: **Case (c)**: Proximal MALA with Gaussian target and $m = 3, r = 2$. Average acceptance rate for different choices of α (first); acceptance rate as a function of ℓ for increasing dimension d (second); ESJD_d as a function of the acceptance rate $a_d(\ell, r)$ (third).




Figure 8: Proximal MALA with Gaussian target and $m = 1, r \rightarrow 0$ (MALA). Average acceptance rate for different choices of α (first); acceptance rate as a function of ℓ for increasing dimension d (second); ESJD_d as a function of the acceptance rate $a_d(\ell, r)$ (third).




Figure 9: Proximal MALA with Laplace target and $m = 1, r = 0$ (sG-MALA). Average acceptance rate for different choices of α (first); acceptance rate as a function of ℓ for increasing dimension d (second); ESJD_d as a function of the acceptance rate $a_d(\ell, r)$ (third).




Figure 10: Proximal MALA with Laplace target and $m = 1, r = 1$ (P-MALA). Average acceptance rate for different choices of α (first); acceptance rate as a function of ℓ for increasing dimension d (second); ESJD_d as a function of the acceptance rate $a_d(\ell, r)$ (third).




Figure 11: Proximal MALA with Laplace target and $m = 3, r = 2$. Average acceptance rate for different choices of α (first); acceptance rate as a function of ℓ for increasing dimension d (second); ESJD_d as a function of the acceptance rate $a_d(\ell, r)$ (third).

algorithmic settings, summarized in Table 3.

Figure	Algorithm	α	β	m	r
12	sG-MALA	1/3	1/3	1	0
13	P-MALA	1/3	1/3	1	1
4	Proximal MALA	1/3	1	3	2

Table 3: Algorithm setting for the simulation study on the mixed Laplace-normal target.

The first plot in Figures 13–12 shows that for $\alpha \neq 1/3$ the acceptance ratio $a_d(\ell, r)$ becomes degenerate as d increases; while the second plot shows that $(a_d(\ell, r))_{d \in \mathbb{N}^*}$ and $(\text{ESJD}_d)_{d \in \mathbb{N}^*}$ converge as $d \rightarrow \infty$.

It is interesting to note that the convergence of these plots happens faster with respect to the dimension d , compared to the Laplace distribution. It is also surprising that out of the three settings tested for the parameters r and m , the one that seems to converge faster, and that looks most stable, is $m = 3$ and $r = 2$, which was never the case for the other densities. This may suggest that for such distributions, a general Proximal MALA algorithm has better properties of convergence than sG-MALA and P-MALA.

C Taylor Expansions for the Results on Differentiable Targets

C.1 Coefficients of the Taylor Expansion

We collect here the coefficients of the Taylor expansions in Propositions 4, 8 and 12.

C.1.1 Case (a)

If $\alpha = 1/4$, $\beta = 1/8$ and $r > 0$, then the log-acceptance ratio (50) satisfies

$$\phi_d(x, z) = d^{-1/2}C_2(x, z) + d^{-3/4}C_3(x, z) + d^{-1}C_4(x, z) + C_5(x, z, \sigma_d) ,$$

where

$$C_2(x, z) = \frac{\ell^2}{2} (-rzg''(x)g'(x)) ,$$




Figure 12: Proximal MALA with mixed Laplace-normal target and $m = 1, r = 0$ (sG-MALA). Average acceptance rate for different choices of α (first); acceptance rate as a function of ℓ for increasing dimension d (second); ESJD_d as a function of the acceptance rate $a_d(\ell, r)$ (third).




Figure 13: Proximal MALA with mixed Laplace-normal target and $m = 1, r = 1$ (P-MALA). Average acceptance rate for different choices of α (first); acceptance rate as a function of ℓ for increasing dimension d (second); ESJD_d as a function of the acceptance rate $a_d(\ell, r)$ (third).

and

$$\begin{aligned}
C_3(x, z) &= \frac{\ell^3}{6} \left(\frac{z^3}{2} g'''(x) - \frac{3}{2} z^2 r g'(x) g'''(x) - \frac{3}{2} r z^2 [g''(x)]^2 + \frac{3}{4} z r^2 g'''(x) [g'(x)]^2 \right. \\
&\quad \left. - \frac{3}{2} z g'(x) g''(x) + \frac{3}{2} r [g'(x)]^2 g''(x) + 3 r^2 z [g'(x)]^2 g''(x) \right), \\
C_4(x, z) &= \frac{\ell^4}{24} \left\{ g^{(4)}(x) \left(z^4 - \frac{z r^3}{2} [g'(x)]^3 - 3 z^3 r g'(x) + \frac{3}{2} z^2 r^2 [g'(x)]^2 \right) \right. \\
&\quad + g'''(x) \left(-6 z^2 g'(x) - 9 r z^3 g''(x) + 9 z^2 r^2 g'(x) g''(x) + 9 r z [g'(x)]^2 \right. \\
&\quad \left. \left. - \frac{9}{2} z r^3 [g'(x)]^2 g''(x) - \frac{3}{2} r^2 [g'(x)]^3 \right) \right. \\
&\quad + 12 r z g'(x) [g''(x)]^2 + 3 g''(x) [g'(x)]^2 - 3 z^2 [g''(x)]^2 + 3 z^2 r^2 [g''(x)]^3 \\
&\quad \left. \left. - 6 r^2 [g'(x) g''(x)]^2 - 3 z r^3 g'(x) [g''(x)]^3 \right\},
\end{aligned}$$

and we use the integral form for the remainder

$$C_5(x, z, \sigma_d) = \int_0^{\sigma_d} \frac{\partial^5}{\partial \sigma^5} R(x, z, \sigma) \Big|_{\sigma=u} \frac{(\sigma_d - u)^4}{4!} du,$$

with u between 0 and σ_d and the derivatives of R_1 and R_2 given in Appendix C.2.

C.1.2 Case (b)

If $\alpha = 1/6$, $\beta = 1/6$ and $r > 0$, the log-acceptance ratio (50) satisfies

$$\begin{aligned}
\phi_d(x, z) &= d^{-1/2} C_3(x, z) + d^{-2/3} C_4(x, z) \\
&\quad + d^{-5/6} C_5(x, z) + d^{-1} C_6(x, z) + C_7(x, z, \sigma_d),
\end{aligned}$$

where

$$C_3(x, z) = \frac{\ell^3}{6} \left(\frac{1}{2} g'''(x) z^3 - \frac{3}{2} g''(x) g'(x) z (1 + 2r) \right),$$

and

$$\begin{aligned}
C_4(x, z) &= \frac{\ell^4}{24} \left(z^4 g^{(4)}(x) - 6z^2 g'''(x) g'(x) (1+r) \right. \\
&\quad \left. - 3z^2 [g''(x)]^2 (1+2r) + 3g''(x) [g'(x)]^2 (1+2r) \right), \\
C_5(x, z) &= \frac{\ell^5}{120} \left(\frac{3}{2} z^5 g^{(5)}(x) - 15z^3 g^{(4)}(x) g'(x) (1+r) + 15z [g'(x)]^2 g'''(x) \left(\frac{3}{2} + 3r + r^2 \right) \right. \\
&\quad \left. + 15z(1+4r+2r^2) g'(x) [g''(x)]^2 - 15z^3 g''(x) g'''(x) (1+3r) \right), \\
C_6(x, z) &= \frac{\ell^6}{720} \left(2z^6 g^{(6)}(x) - 30(1+r) z^4 g'(x) g^{(5)}(x) + 45(2+4r+r^2) z^2 [g'(x)]^2 g^{(4)}(x) \right. \\
&\quad + 90(r+r^2) z^2 [g''(x)]^3 - 15(2+6r+3r^2) g'''(x) [g'(x)]^3 \\
&\quad - 30(1+4r) z^4 g''(x) g^{(4)}(x) + 45(3+16r+6r^2) z^2 g'(x) g''(x) g'''(x) \\
&\quad \left. - \frac{45}{2} (1+4r) z^4 [g'''(x)]^2 - \frac{45}{2} (1+8r+8r^2) [g'(x) g''(x)]^2 \right),
\end{aligned}$$

and integral form of the remainder

$$C_7(x, z, \sigma_d) = \int_0^{\sigma_d} \frac{\partial^7}{\partial \sigma^7} R(x, z, \sigma) \Big|_{\sigma=u} \frac{(\sigma_d - u)^6}{6!} du,$$

with u between 0 and σ_d and the derivatives of R_1 and R_2 are given in Appendix C.2.

C.1.3 Case (c)

If $\alpha = 1/6$, $\beta = m/6$ for $m > 1$ and $r > 0$, then the log-acceptance ratio (50) satisfies

$$\begin{aligned}
\phi_d(x, z) &= d^{-1/2} C_3(x, z) + d^{-2/3} C_4(x, z) \\
&\quad + d^{-5/6} C_5(x, z) + d^{-1} C_6(x, z) + C_7(x, z, \sigma_d),
\end{aligned}$$

where

$$C_3(x, z) = \frac{\ell^3}{6} \left(\frac{1}{2} g'''(x) z^3 - \frac{3}{2} g''(x) g'(x) z \right)$$

and

$$\begin{aligned}
C_4(x, z) &= \frac{\ell^4}{24} \begin{cases} z^4 g^{(4)}(x) - 6z^2 g'''(x) g'(x) - 3z^2 [g''(x)]^2 + 3g''(x) [g'(x)]^2 \\ \quad - 12zr g'(x) g''(x) & \text{if } m = 3/2 \\ z^4 g^{(4)}(x) - 6z^2 g'''(x) g'(x) - 3z^2 [g''(x)]^2 + 3g''(x) [g'(x)]^2 & \text{otherwise} \end{cases}, \\
C_5(x, z) &= \frac{\ell^5}{120} \begin{cases} \frac{3}{2} z^5 g^{(5)}(x) - 15z^3 g^{(4)}(x) g'(x) + \frac{45}{2} z [g'(x)]^2 g'''(x) + 15z g'(x) [g''(x)]^2 \\ \quad - 15z^3 g''(x) g'''(x) - 30z^2 r [g''(x)]^2 + 30r [g'(x)]^2 g''(x) \\ \quad - 30r z^2 g'(x) g'''(x) & \text{if } m = 3/2 \\ \frac{3}{2} z^5 g^{(5)}(x) - 15z^3 g^{(4)}(x) g'(x) + \frac{45}{2} z [g'(x)]^2 g'''(x) + 15z g'(x) [g''(x)]^2 \\ \quad - 15z^3 g''(x) g'''(x) - 60zr g'(x) g''(x) & \text{if } m = 2 \\ \frac{3}{2} z^5 g^{(5)}(x) - 15z^3 g^{(4)}(x) g'(x) + \frac{45}{2} z [g'(x)]^2 g'''(x) \\ \quad + 15z g'(x) [g''(x)]^2 - 15z^3 g''(x) g'''(x) & \text{otherwise} \end{cases}, \\
C_6(x, z) &= \frac{\ell^6}{720} \begin{cases} 2z^6 g^{(6)}(x) - 30z^4 g'(x) g^{(5)}(x) + 90z^2 [g'(x)]^2 g^{(4)}(x) - 30g'''(x) [g'(x)]^3 \\ \quad - \frac{45}{2} [g'(x) g''(x)]^2 - 30z^4 g''(x) g^{(4)}(x) + 135z^2 g'(x) g''(x) g'''(x) \\ \quad - \frac{45}{2} z^4 [g'''(x)]^2 - 90z^3 r g'(x) g^{(4)}(x) + 540r z g'(x) [g''(x)]^2 \\ \quad + 180r z g'(x) g''(x) + 270r z g'''(x) [g'(x)]^2 - 45zr g''(x) g'''(x) \\ \quad + 90r z^3 g'''(x) & \text{if } m = 3/2 \\ 2z^6 g^{(6)}(x) - 30z^4 g'(x) g^{(5)}(x) + 90z^2 [g'(x)]^2 g^{(4)}(x) - 30g'''(x) [g'(x)]^3 \\ \quad - \frac{45}{2} [g'(x) g''(x)]^2 - 30z^4 g''(x) g^{(4)}(x) + 135z^2 g'(x) g''(x) g'''(x) \\ \quad - \frac{45}{2} z^4 [g'''(x)]^2 - 180z^2 r g'(x) g'''(x) - 180z^2 r [g''(x)]^2 \\ \quad + 180r g''(x) [g'(x)]^2 & \text{if } m = 2 \\ 2z^6 g^{(6)}(x) - 30z^4 g'(x) g^{(5)}(x) + 90z^2 [g'(x)]^2 g^{(4)}(x) - 30g'''(x) [g'(x)]^3 \\ \quad - \frac{45}{2} [g'(x) g''(x)]^2 - 30z^4 g''(x) g^{(4)}(x) + 135z^2 g'(x) g''(x) g'''(x) \\ \quad - \frac{45}{2} z^4 [g'''(x)]^2 - 360zr g'(x) g''(x) & \text{if } m = 5/2 \\ 2z^6 g^{(6)}(x) - 30z^4 g'(x) g^{(5)}(x) + 90z^2 [g'(x)]^2 g^{(4)}(x) - 30g'''(x) [g'(x)]^3 \\ \quad - \frac{45}{2} [g'(x) g''(x)]^2 - 30z^4 g''(x) g^{(4)}(x) + 135z^2 g'(x) g''(x) g'''(x) \\ \quad - \frac{45}{2} z^4 [g'''(x)]^2 & \text{otherwise} \end{cases},
\end{aligned}$$

and integral form of the remainder

$$C_7(x, z, \sigma_d) = \int_0^{\sigma_d} \frac{\partial^7}{\partial \sigma^7} R(x, z, \sigma) \Big|_{\sigma=u} \frac{(\sigma_d - u)^6}{6!} du,$$

with u between 0 and σ_d and the derivatives of R_1 and R_2 are given in Appendix C.2.

C.2 Taylor Expansions of the Log-acceptance Ratio

C.2.1 R_1

Recall that $R_1(x, z, \sigma) = -g \left[x - \frac{\sigma^2}{2} g' \left(\text{prox}_g^{\sigma^{2m} r/2}(x) \right) + \sigma z \right] + g(x)$. We compute the derivatives of R_1 w.r.t. σ evaluated at 0:

$$\begin{aligned}
R_1(x, z, 0) &= 0, \\
\frac{\partial R_1}{\partial \sigma}(x, z, \sigma)|_{\sigma=0} &= -g'(x)z, \\
\frac{\partial^2 R_1}{\partial \sigma^2}(x, z, \sigma)|_{\sigma=0} &= -z^2 g''(x) + [g'(x)]^2, \\
\frac{\partial^3 R_1}{\partial \sigma^3}(x, z, \sigma)|_{\sigma=0} &= -z^3 g'''(x) + 3g'(x)g''(x) \left[z + \frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right], \\
\frac{\partial^4 R_1}{\partial \sigma^4}(x, z, \sigma)|_{\sigma=0} &= -z^4 g^{(4)}(x) + 6z^2 g'''(x)g'(x) - 3g''(x)[g'(x)]^2 \\
&\quad + 12z [g''(x)]^2 \frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \\
&\quad + 6g'(x) \\
&\quad \times \left[g''(x) \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} + \left(\frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right)^2 g'''(x) \right].
\end{aligned}$$

In addition, for $m > 1/2$ we will also use

$$\begin{aligned}
\frac{\partial^5 R_1}{\partial \sigma^5}(x, z, \sigma)|_{\sigma=0} &= -z^5 g^{(5)}(x) + 10z^3 g^{(4)}(x)g'(x) - 15zg'''(x)[g'(x)]^2 \\
&\quad + 30z [g''(x)]^2 \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \\
&\quad + 10g'(x)g''(x) \frac{\partial^3}{\partial \sigma^3} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0}, \\
\frac{\partial^6 R_1}{\partial \sigma^6}(x, z, \sigma)|_{\sigma=0} &= -z^6 g^{(6)}(x) + 15z^4 g^{(5)}(x)g'(x) - 45z^2 g^{(4)}(x)[g'(x)]^2 + 15g'''(x)[g'(x)]^3 \\
&\quad - 90g'(x)[g''(x)]^2 \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \\
&\quad - 60zg''(x) \frac{\partial^3}{\partial \sigma^3} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \\
&\quad + 90g''(x)g'''(x) \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \\
&\quad + 15g'(x) \\
&\quad \times \left(g''(x) \frac{\partial^4}{\partial \sigma^4} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} + 3 \left[\frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right]^2 g'''(x) \right).
\end{aligned}$$

C.2.2 R_2

Recall that

$$R_2(x, z, \sigma) = \frac{1}{2}z^2 - \frac{1}{2} \left(z - \frac{\sigma}{2} g' \left(\text{prox}_g^{\sigma^{2m}r/2} \left[x + \sigma z - \frac{\sigma^2}{2} g' \left[\text{prox}_g^{\sigma^{2m}r/2}(x) \right] \right] \right) - \frac{\sigma}{2} g' \left[\text{prox}_g^{\sigma^{2m}r/2}(x) \right] \right)^2.$$

We compute the derivatives of R_2 w.r.t. σ evaluated at 0:

$$\begin{aligned}
R_2(x, z, 0) &= 0, \\
\frac{\partial R_2}{\partial \sigma}(x, z, \sigma)|_{\sigma=0} &= zg'(x), \\
\frac{\partial^2 R_2}{\partial \sigma^2}(x, z, \sigma)|_{\sigma=0} &= -[g'(x)]^2 + zg''(x) \left[z + 2 \frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right], \\
\frac{\partial^3 R_2}{\partial \sigma^3}(x, z, \sigma)|_{\sigma=0} &= -3g'(x)g''(x) \left[z + 2 \frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] \\
&\quad + \frac{3}{2}z \left(\left[z + \frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right]^2 g'''(x) \right. \\
&\quad + \left[-g'(x) + 2z \frac{\partial^2}{\partial \sigma \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} + \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] g''(x) \\
&\quad \left. + \left[\frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right]^2 g'''(x) + \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} g''(x) \right), \\
\frac{\partial^4 R_2}{\partial \sigma^4}(x, z, \sigma)|_{\sigma=0} &= -3[g''(x)]^2 \left[z + 2 \frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right]^2 \\
&\quad - 6g'(x) \left(\left[z + \frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right]^2 g'''(x) \right. \\
&\quad + \left[-g'(x) + 2z \frac{\partial^2}{\partial \sigma \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} + \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] g''(x) \\
&\quad + \left[\frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right]^2 g'''(x) + \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} g''(x) \Big) \\
&\quad + 2zg^{(4)}(x) \left[z + \frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right]^3 \\
&\quad + 6zg'''(x) \left[z + \frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] \\
&\quad \times \left[-g'(x) + 2z \frac{\partial^2}{\partial \sigma \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} + \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] \\
&\quad + 2zg''(x) \left[-3g''(x) \frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} - 3g'(x) \frac{\partial^2}{\partial \sigma \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right. \\
&\quad + 3z^2 \frac{\partial^3}{\partial \sigma \partial x^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} + 3z \frac{\partial^3}{\partial \sigma^2 \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \\
&\quad \left. + \frac{\partial^3}{\partial \sigma^3} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] \\
&\quad + 2zg^{(4)}(x) \left[\frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right]^3 + 2zg''(x) \frac{\partial^3}{\partial \sigma^3} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \\
&\quad + 6z \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} g'''(x).
\end{aligned}$$

We then proceed to get the derivatives needed for $m > 1/2$:

$$\begin{aligned}
\frac{\partial^5 R_2}{\partial \sigma^5}(x, z, \sigma)|_{\sigma=0} = & -15zg''(x) \left(z^2 g'''(x) + \left[-g'(x) + 2 \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] g''(x) \right) \\
& - 5g'(x) \left(2z^3 g^{(4)}(x) + 6zg'''(x) \left[-g'(x) + \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] \right. \\
& \quad \left. + 2g''(x) \left[3z \frac{\partial^3}{\partial \sigma^2 \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} + \frac{\partial^3}{\partial \sigma^3} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] \right. \\
& \quad \left. + 2g''(x) \frac{\partial^3}{\partial \sigma^3} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right) \\
& + \frac{5}{2} g^{(5)}(x) z^5 + \frac{5}{2} z g''(x) \frac{\partial^4}{\partial \sigma^4} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \\
& + 15g^{(4)}(x) z^3 \left[-g'(x) + \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] \\
& + \frac{15}{2} z g'''(x) \left[-g'(x) + \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right]^2 \\
& + \frac{15}{2} z g'''(x) \left[\frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right]^2 \\
& + 10z^2 g'''(x) \left(3z \frac{\partial^3}{\partial \sigma^2 \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} + \frac{\partial^3}{\partial \sigma^3} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right) \\
& + \frac{5}{2} z g''(x) \left(\frac{\partial^4}{\partial \sigma^4} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} - 6g''(x) \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right. \\
& \quad \left. - 6g'(x) \frac{\partial^3}{\partial \sigma^2 \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} + 6z^2 \frac{\partial^4}{\partial \sigma^2 \partial x^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right. \\
& \quad \left. + 4z \frac{\partial^4}{\partial \sigma^3 \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right)
\end{aligned}$$

and

$$\begin{aligned}
\frac{\partial^6 R_2}{\partial \sigma^6}(x, z, \sigma)|_{\sigma=0} = & -\frac{45}{2} \left(z^2 g'''(x) + \left[-g'(x) + 2 \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] g''(x) \right)^2 \\
& - 15zg''(x) \left(2z^3 g^{(4)}(x) + 6zg'''(x) \left[-g'(x) + \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] \right. \\
& \quad \left. + 2g''(x) \left[3z \frac{\partial^3}{\partial \sigma^2 \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} + \frac{\partial^3}{\partial \sigma^3} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right] \right. \\
& \quad \left. + 2g''(x) \frac{\partial^3}{\partial \sigma^3} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} \right) \\
& + 6g'(x) A^{(5)} - zA^{(6)},
\end{aligned}$$

with

$$\begin{aligned}
A^{(5)} = & -\frac{5}{2}g^{(5)}(x)z^4 - \frac{5}{2}g''(x)\frac{\partial^4}{\partial\sigma^4}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \\
& - 15g^{(4)}(x)z^2 \left[-g'(x) + \frac{\partial^2}{\partial\sigma^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right] \\
& - \frac{15}{2}g'''(x) \left[-g'(x) + \frac{\partial^2}{\partial\sigma^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right]^2 - \frac{15}{2}g'''(x) \left[\frac{\partial^2}{\partial\sigma^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right]^2 \\
& - 10zg'''(x) \left(3z\frac{\partial^3}{\partial\sigma^2\partial x}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} + \frac{\partial^3}{\partial\sigma^3}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right) \\
& - \frac{5}{2}g''(x) \left(\frac{\partial^4}{\partial\sigma^4}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} - 6g''(x)\frac{\partial^2}{\partial\sigma^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right. \\
& \quad - 6g'(x)\frac{\partial^3}{\partial\sigma^2\partial x}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \\
& \quad \left. + 6z^2\frac{\partial^4}{\partial\sigma^2\partial x^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} + 4z\frac{\partial^4}{\partial\sigma^3\partial x}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right), \\
A^{(6)} = & -3 \left(10g'''(x)\frac{\partial^2}{\partial\sigma^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0}\frac{\partial^3}{\partial\sigma^3}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right. \\
& + g''(x)\frac{\partial^5}{\partial\sigma^5}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} + g^{(6)}(x)z^5 \\
& + 10g^{(5)}(x)z^3 \left[-g'(x) + \frac{\partial^2}{\partial\sigma^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right] \\
& + 15zg^{(4)}(x) \left[-g'(x) + \frac{\partial^2}{\partial\sigma^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right]^2 \\
& + 10g^{(4)}(x)z^2 \left[\frac{\partial^3}{\partial\sigma^3}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} + 3z\frac{\partial^3}{\partial\sigma^2\partial x}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right] \\
& + 10g'''(x) \left[-g'(x) + \frac{\partial^2}{\partial\sigma^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right] \\
& \quad \times \left[\frac{\partial^3}{\partial\sigma^3}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} + 3z\frac{\partial^3}{\partial\sigma^2\partial x}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right] \\
& + 5g'''(x)z \left[-6g''(x)\frac{\partial^2}{\partial\sigma^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} - 6g'(x)\frac{\partial^3}{\partial\sigma^2\partial x}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right. \\
& \quad + 6z^2\frac{\partial^4}{\partial\sigma^2\partial x^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} + 3z\frac{\partial^4}{\partial\sigma^3\partial x}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \\
& \quad \left. + \frac{\partial^4}{\partial\sigma^4}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right] \\
& + g''(x) \left[-10g''(x)\frac{\partial^3}{\partial\sigma^3}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} - 10g'(x)\frac{\partial^4}{\partial\sigma^3\partial x}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right. \\
& \quad + 5z\frac{\partial^5}{\partial\sigma^4\partial x}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} + 10z^2\frac{\partial^5}{\partial\sigma^3\partial x^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \\
& \quad + 10z^3\frac{\partial^5}{\partial\sigma^2\partial x^3}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \\
& \quad \left. - 30g'(x)z\frac{\partial^4}{\partial\sigma^2\partial x^2}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} + \frac{\partial^5}{\partial\sigma^5}\text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} \right] \Bigg).
\end{aligned}$$

C.3 Derivatives of the Proximity Map for Differentiable Targets

Recall that, in the differentiable case, $\text{prox}_g^{\sigma^{2m}r/2}(x) = -\frac{\sigma^{2m}r}{2}g'(\text{prox}_g^{\sigma^{2m}r/2}(x)) + x$ then

$$\begin{aligned}\frac{\partial}{\partial \sigma} \text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} &= \begin{cases} -\frac{r}{2}g'(x) & \text{if } m = 1/2 \\ 0 & \text{if } m > 1/2 \\ \infty & \text{otherwise} \end{cases} \\ \frac{\partial^2}{\partial \sigma^2} \text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} &= \begin{cases} \frac{r^2}{2}g'(x)g''(x) & \text{if } m = 1/2 \\ -rg'(x) & \text{if } m = 1 \\ 0 & \text{if } m > 1 \\ \infty & \text{otherwise} \end{cases} \\ \frac{\partial^3}{\partial \sigma^3} \text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} &= \begin{cases} -\frac{3r^3}{8}g'''(x)[g'(x)]^2 - \frac{3r^3}{4}g'(x)[g''(x)]^2 & \text{if } m = 1/2 \\ -3rg'(x) & \text{if } m = 3/2 \\ 0 & \text{if } m = 1, m > 3/2 \\ \infty & \text{otherwise} \end{cases} \\ \frac{\partial^4}{\partial \sigma^4} \text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} &= \begin{cases} < \infty & \text{if } m = 1/2 \\ 6r^2g'(x)g''(x) & \text{if } m = 1 \\ -12rg'(x) & \text{if } m = 2 \\ 0 & \text{if } m = 3/2, m > 2 \\ \infty & \text{otherwise} \end{cases} \\ \frac{\partial^5}{\partial \sigma^5} \text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} &= \begin{cases} < \infty & \text{if } m = 1/2 \\ -60rg'(x) & \text{if } m = 5/2 \\ 0 & \text{if } m = 1, m = 3/2, m = 2, m > 5/2 \\ \infty & \text{otherwise} \end{cases}\end{aligned}$$

and

$$\frac{\partial}{\partial x} \text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} = 1, \quad \frac{\partial^{(k)}}{\partial x^{(k)}} \text{prox}_g^{\sigma^{2m}r/2}(x)|_{\sigma=0} = 0,$$

for all integers $k > 1$. For the mixed derivatives we have

$$\begin{aligned}
\frac{\partial^2}{\partial \sigma \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} &= \begin{cases} -\frac{r}{2} g''(x) & \text{if } m = 1/2 \\ 0 & \text{if } m > 1/2 \\ \infty & \text{otherwise} \end{cases} \\
\frac{\partial^3}{\partial \sigma^2 \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} &= \begin{cases} \frac{r^2}{2} [g''(x)]^2 + \frac{r^2}{2} g'(x) g'''(x) & \text{if } m = 1/2 \\ -r g''(x) & \text{if } m = 1 \\ 0 & \text{if } m > 1/2 \\ \infty & \text{otherwise} \end{cases} \\
\frac{\partial^4}{\partial \sigma^3 \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} &= \begin{cases} < \infty & \text{if } m = 1/2 \\ -3r g''(x) & \text{if } m = 3/2 \\ 0 & \text{if } m = 1, m > 3/2 \\ \infty & \text{otherwise} \end{cases} \\
\frac{\partial^5}{\partial \sigma^4 \partial x} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} &= \begin{cases} < \infty & \text{if } m = 1/2 \\ 6r^2 (g'(x) g'''(x) + [g''(x)]^2) & \text{if } m = 1 \\ -12r g''(x) & \text{if } m = 2 \\ 0 & \text{if } m = 3/2, m > 2 \\ \infty & \text{otherwise} \end{cases}
\end{aligned}$$

and

$$\begin{aligned}
\frac{\partial^3}{\partial \sigma \partial x^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} &= \begin{cases} -\frac{r}{2} g'''(x) & \text{if } m = 1/2 \\ 0 & \text{if } m > 1/2 \\ \infty & \text{otherwise} \end{cases} \\
\frac{\partial^4}{\partial \sigma \partial x^3} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} &= \begin{cases} < \infty & \text{if } m = 1/2 \\ 0 & \text{if } m > 1/2 \\ \infty & \text{otherwise} \end{cases} \\
\frac{\partial^4}{\partial \sigma^2 \partial x^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} &= \begin{cases} < \infty & \text{if } m = 1/2 \\ -r g'''(x) & \text{if } m = 1 \\ 0 & \text{if } m > 1/2 \\ \infty & \text{otherwise} \end{cases} \\
\frac{\partial^5}{\partial \sigma^3 \partial x^2} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} &= \begin{cases} < \infty & \text{if } m = 1/2 \\ -3r g'''(x) & \text{if } m = 3/2 \\ 0 & \text{if } m = 1, m > 3/2 \\ \infty & \text{otherwise} \end{cases} \\
\frac{\partial^5}{\partial \sigma^2 \partial x^3} \text{prox}_g^{\sigma^{2m} r/2}(x)|_{\sigma=0} &= \begin{cases} < \infty & \text{if } m = 1/2 \\ -r g^{(4)}(x) & \text{if } m = 1 \\ 0 & \text{if } m > 1 \\ \infty & \text{otherwise} \end{cases}
\end{aligned}$$

D Moments and Integrals for the Laplace Distribution

D.1 Moments of Acceptance Ratio for the Laplace Distribution

The indicator functions in the definition of ϕ_d identify four different regions:

$$\begin{aligned}
R_1 &:= \left\{ (x, z) : |x| < \sigma^{2m} r/2 \wedge \left| \left(1 - \frac{1}{\sigma^{2(m-1)} r} \right) x + \sigma z \right| < \sigma^{2m} r/2 \right\}, \\
R_2 &:= \left\{ (x, z) : |x| \geq \sigma^{2m} r/2 \wedge \left| x - \frac{\sigma^2}{2} \text{sgn}(x) + \sigma z \right| < \sigma^{2m} r/2 \right\}, \\
R_3 &:= \left\{ (x, z) : |x| < \sigma^{2m} r/2 \wedge \left| \left(1 - \frac{1}{\sigma^{2(m-1)} r} \right) x + \sigma z \right| \geq \sigma^{2m} r/2 \right\}, \\
R_4 &:= \left\{ (x, z) : |x| > \sigma^{2m} r/2 \wedge \left| x - \frac{\sigma^2}{2} \text{sgn}(x) + \sigma z \right| > \sigma^{2m} r/2 \right\},
\end{aligned}$$

with corresponding acceptance ratios

$$\begin{aligned}
\phi_d^1(x, z) &= |x| - \left| \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) x + \sigma z \right| + \frac{z^2}{2} \\
&\quad - \frac{1}{2\sigma^2} \left(\left(\frac{2}{\sigma^{2(m-1)}r} - \frac{1}{\sigma^{4(m-1)}r^2} \right) x - \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) \sigma z \right)^2 \\
\phi_d^2(x, z) &= |x| - \left| x - \frac{\sigma^2}{2} \operatorname{sgn}(x) + \sigma z \right| + \frac{z^2}{2} \\
&\quad - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)}r} x + \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) \left(\frac{\sigma^2}{2} \operatorname{sgn}(x) - \sigma z \right) \right)^2 \\
\phi_d^3(x, z) &= |x| - \left| \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) x + \sigma z \right| + \frac{z^2}{2} \\
&\quad - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)}r} x - \sigma z + \frac{\sigma^2}{2} \operatorname{sgn} \left[\left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) x + \sigma z \right] \right)^2 \\
\phi_d^4(x, z) &= |x| - \left| x - \frac{\sigma^2}{2} \operatorname{sgn}(x) + \sigma z \right| + \frac{z^2}{2} \\
&\quad - \frac{1}{2\sigma^2} \left(\frac{\sigma^2}{2} \operatorname{sgn}(x) - \sigma z + \frac{\sigma^2}{2} \operatorname{sgn} \left[x - \frac{\sigma^2}{2} \operatorname{sgn}(x) + \sigma z \right] \right)^2.
\end{aligned}$$

Let us denote

$$\begin{aligned}
A_1 &:= \left\{ x : 0 \leq x < \frac{\sigma^{2m}r}{2} \right\}, \quad A_2 := \left\{ x : -\frac{\sigma^{2m}r}{2} < x \leq 0 \right\}, \\
A_3 &:= \left\{ x : x \geq \frac{\sigma^{2m}r}{2} \right\}, \quad A_4 := \left\{ x : x \leq -\frac{\sigma^{2m}r}{2} \right\},
\end{aligned}$$

and

$$\begin{aligned}
B_1 &:= \left\{ z : 0 \leq \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) x + \sigma z < \frac{\sigma^{2m}r}{2} \right\}, \\
B_2 &:= \left\{ z : -\frac{\sigma^{2m}r}{2} < \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) x + \sigma z \leq 0 \right\}, \\
B_3 &:= \left\{ z : \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) x + \sigma z \geq \frac{\sigma^{2m}r}{2} \right\}, \\
B_4 &:= \left\{ z : \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) x + \sigma z \leq -\frac{\sigma^{2m}r}{2} \right\},
\end{aligned}$$

and

$$\begin{aligned}
C_1 &:= \left\{ (x, z) : 0 \leq x - \frac{\sigma^2}{2} \operatorname{sgn}(x) + \sigma z < \frac{\sigma^{2m} r}{2} \right\}, \\
C_2 &:= \left\{ (x, z) : -\frac{\sigma^{2m} r}{2} < x - \frac{\sigma^2}{2} \operatorname{sgn}(x) + \sigma z \leq 0 \right\} \\
C_3 &:= \left\{ (x, z) : x - \frac{\sigma^2}{2} \operatorname{sgn}(x) + \sigma z \geq \frac{\sigma^{2m} r}{2} \right\}, \\
C_4 &:= \left\{ (x, z) : x - \frac{\sigma^2}{2} \operatorname{sgn}(x) + \sigma z \leq -\frac{\sigma^{2m} r}{2} \right\},
\end{aligned}$$

so that, up to a set of null measure,

$$\begin{aligned}
R_1 &= (A_1 \cup A_2) \cap (B_1 \cup B_2), & R_2 &= (A_3 \cup A_4) \cap (C_1 \cup C_2), \\
R_3 &= (A_1 \cup A_2) \cap (B_3 \cup B_4), & R_4 &= (A_3 \cup A_4) \cap (C_3 \cup C_4).
\end{aligned}$$

Proposition 16. *Take X a Laplace random variable and Z a standard normal random variable independent of X , then if $\sigma^2 = \ell^2 d^{-2/3}$, we have*

$$\lim_{d \rightarrow +\infty} d\mathbb{E}[\phi_d(X, Z)] = -\frac{\ell^3}{3\sqrt{2\pi}}.$$

Proof. Taking expectations of $\phi_d^i \mathbb{1}_{R_i}$ for $i = 1, \dots, 4$ and exploiting the symmetry of the laws of X

and Z , we can write

$$\begin{aligned}
& \mathbb{E} [\phi_d^1(X, Z) \mathbb{1}_{R_1}(X, Z)] \\
&= 2\mathbb{E} \left[\left(\frac{1}{\sigma^{2(m-1)}r} X - \sigma Z \right) \mathbb{1}_{A_1}(X) \mathbb{1}_{B_1}(X, Z) \right] \\
&+ 2\mathbb{E} \left[\left(2X - \frac{1}{\sigma^{2(m-1)}r} X + \sigma Z \right) \mathbb{1}_{A_1}(X) \mathbb{1}_{B_2}(X, Z) \right] \\
&+ 2\mathbb{E} \left[\left(\frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\left(\frac{2}{\sigma^{2(m-1)}r} - \frac{1}{\sigma^{4(m-1)}r^2} \right) X - \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) \sigma Z \right)^2 \right) \right. \\
&\quad \left. \times \mathbb{1}_{A_1}(X) \mathbb{1}_{B_1 \cup B_2}(X, Z) \right], \\
& \mathbb{E} [\phi_d^2(X, Z) \mathbb{1}_{R_2}(X, Z)] \\
&= 2\mathbb{E} \left[\left(\frac{\sigma^2}{2} - \sigma Z \right) \mathbb{1}_{A_3}(X) \mathbb{1}_{C_1}(X, Z) \right] \\
&+ 2\mathbb{E} \left[\left(2X - \frac{\sigma^2}{2} + \sigma Z \right) \mathbb{1}_{A_3}(X) \mathbb{1}_{C_2}(X, Z) \right] \\
&+ 2\mathbb{E} \left[\left(\frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)}r} X + \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) \left(\frac{\sigma^2}{2} - \sigma Z \right) \right)^2 \right) \right. \\
&\quad \left. \times \mathbb{1}_{A_3}(X) \mathbb{1}_{C_1 \cup C_2}(X, Z) \right], \\
& \mathbb{E} [\phi_d^3(X, Z) \mathbb{1}_{R_3}(X, Z)] \\
&= 2\mathbb{E} \left[\left(\frac{1}{\sigma^{2(m-1)}r} X - \sigma Z \right) \mathbb{1}_{A_1}(X) \mathbb{1}_{B_3}(X, Z) \right] \\
&+ 2\mathbb{E} \left[\left(2X - \frac{1}{\sigma^{2(m-1)}r} X + \sigma Z \right) \mathbb{1}_{A_1}(X) \mathbb{1}_{B_4}(X, Z) \right] \\
&+ 2\mathbb{E} \left[\left(\frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)}r} X - \sigma Z + \frac{\sigma^2}{2} \right)^2 \right) \mathbb{1}_{A_1}(X) \mathbb{1}_{B_3}(X, Z) \right] \\
&+ 2\mathbb{E} \left[\left(\frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)}r} X - \sigma Z - \frac{\sigma^2}{2} \right)^2 \right) \mathbb{1}_{A_1}(X) \mathbb{1}_{B_4}(X, Z) \right], \\
& \mathbb{E} [\phi_d^4(X, Z) \mathbb{1}_{R_4}(X, Z)] = 2\mathbb{E} \left[\left(2X - \frac{\sigma^2}{2} + \sigma Z \right) \mathbb{1}_{A_3}(X) \mathbb{1}_{C_4}(X, Z) \right].
\end{aligned}$$

Using the integrals in Appendix D.4 and Lebesgue's dominated convergence theorem, we find that

for $\alpha = \beta = 1/3$ and $r \geq 0$

$$\begin{aligned}
\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d^1(X, Z)] &= 0 \\
\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d^2(X, Z)] &= -2 \frac{\ell^3 r}{4\sqrt{2\pi}} \int_{-\infty}^0 e^{-z^2/2} z dz = \frac{\ell^3 r}{2\sqrt{2\pi}} \\
\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d^3(X, Z)] &= \frac{3\ell^3 r}{8\sqrt{2\pi}} \int_{-\infty}^0 e^{-z^2/2} z dz - \frac{\ell^3 r}{8\sqrt{2\pi}} \int_0^{+\infty} e^{-z^2/2} z dz = -\frac{\ell^3 r}{2\sqrt{2\pi}} \\
\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d^4(X, Z)] &= \frac{\ell^3}{6\sqrt{2\pi}} \int_{-\infty}^0 e^{-z^2/2} z^3 dz = -\frac{\ell^3}{3\sqrt{2\pi}},
\end{aligned}$$

which gives

$$\begin{aligned}
\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d(X, Z)] &= \lim_{d \rightarrow +\infty} d (\mathbb{E} [\phi_d^1(X, Z)] + \mathbb{E} [\phi_d^2(X, Z)] + \mathbb{E} [\phi_d^3(X, Z)] + \mathbb{E} [\phi_d^4(X, Z)]) \\
&= -\frac{\ell^3}{3\sqrt{2\pi}}.
\end{aligned}$$

For $\alpha = 1/3, \beta = m/3$ for $m > 1$ and $r \geq 0$ we have

$$\begin{aligned}
\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d^1(X, Z)] &= 0 \\
\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d^2(X, Z)] &= 0 \\
\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d^3(X, Z)] &= 0 \\
\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d^4(X, Z)] &= \frac{\ell^3}{6\sqrt{2\pi}} \int_{-\infty}^0 e^{-z^2/2} z^3 dz = -\frac{\ell^3}{3\sqrt{2\pi}},
\end{aligned}$$

which gives

$$\begin{aligned}
\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d(X, Z)] &= \lim_{d \rightarrow +\infty} d (\mathbb{E} [\phi_d^1(X, Z)] + \mathbb{E} [\phi_d^2(X, Z)] + \mathbb{E} [\phi_d^3(X, Z)] + \mathbb{E} [\phi_d^4(X, Z)]) \\
&= -\frac{\ell^3}{3\sqrt{2\pi}}.
\end{aligned}$$

□

Proposition 17. Take X a Laplace random variable and Z a standard normal random variable independent of X , then if $\sigma^2 = \ell^2 d^{-2/3}$

$$\lim_{d \rightarrow +\infty} d \text{Var} (\phi_d(X, Z)) = \frac{2\ell^3}{3\sqrt{2\pi}}.$$

Proof. As a consequence of the previous Proposition we have

$$\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d(X, Z)]^2 = 0.$$

Then, because $R_j \cap R_i = \emptyset$ for all $j \neq i$, we have that

$$\begin{aligned}\mathbb{E} [\phi_d(X, Z)^2] &= \mathbb{E} [\phi_d^1(X, Z)^2 R_1(X, Z)] + \mathbb{E} [\phi_d^2(X, Z)^2 R_2(X, Z)] \\ &\quad + \mathbb{E} [\phi_d^3(X, Z)^2 R_3(X, Z)] + \mathbb{E} [\phi_d^4(X, Z)^2 R_4(X, Z)],\end{aligned}$$

and, exploiting again the symmetry of the laws of X and Z , we have

$$\begin{aligned}\mathbb{E} [\phi_d^1(X, Z)^2 R_1(X, Z)] &= 2\mathbb{E} \left[\left(\frac{1}{\sigma^{2(m-1)r}} X - \sigma Z + \frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\left(\frac{2}{\sigma^{2(m-1)r}} - \frac{1}{\sigma^{4(m-1)r^2}} \right) X - \left(1 - \frac{1}{\sigma^{2(m-1)r}} \right) \sigma Z \right)^2 \right)^2 \right. \\ &\quad \times \mathbb{1}_{A_1}(X) \mathbb{1}_{B_1}(X, Z) \Big], \\ &+ 2\mathbb{E} \left[\left(2X - \frac{1}{\sigma^{2(m-1)r}} X - \sigma Z + \frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\left(\frac{2}{\sigma^{2(m-1)r}} - \frac{1}{\sigma^{4(m-1)r^2}} \right) X - \left(1 - \frac{1}{\sigma^{2(m-1)r}} \right) \sigma Z \right)^2 \right)^2 \right. \\ &\quad \times \mathbb{1}_{A_1}(X) \mathbb{1}_{B_2}(X, Z) \Big], \\ \mathbb{E} [\phi_d^2(X, Z)^2 R_2(X, Z)] &= 2\mathbb{E} \left[\left(\frac{\sigma^2}{2} - \sigma Z + \frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)r}} X + \left(1 - \frac{1}{\sigma^{2(m-1)r}} \right) \left(\frac{\sigma^2}{2} - \sigma Z \right) \right)^2 \right)^2 \mathbb{1}_{A_3}(X) \mathbb{1}_{C_1}(X, Z) \right] \\ &+ 2\mathbb{E} \left[\left(2X - \frac{\sigma^2}{2} + \sigma Z + \frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)r}} X + \left(1 - \frac{1}{\sigma^{2(m-1)r}} \right) \left(\frac{\sigma^2}{2} - \sigma Z \right) \right)^2 \right)^2 \right. \\ &\quad \times \mathbb{1}_{A_1}(X) \mathbb{1}_{C_2}(X, Z) \Big], \\ \mathbb{E} [\phi_d^3(X, Z)^2 R_3(X, Z)] &= 2\mathbb{E} \left[\left(\frac{1}{\sigma^{2(m-1)r}} X - \sigma Z + \frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)r}} X - \sigma Z + \frac{\sigma^2}{2} \right)^2 \right)^2 \mathbb{1}_{A_1}(X) \mathbb{1}_{B_3}(X, Z) \right] \\ &+ 2\mathbb{E} \left[\left(2X - \frac{1}{\sigma^{2(m-1)r}} X + \sigma Z + \frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)r}} X - \sigma Z - \frac{\sigma^2}{2} \right)^2 \right)^2 \mathbb{1}_{A_1}(X) \mathbb{1}_{B_4}(X, Z) \right], \\ \mathbb{E} [\phi_d^4(X, Z)^2 R_4(X, Z)] &= 2\mathbb{E} \left[\left(2X - \frac{\sigma^2}{2} + \sigma Z \right)^2 \mathbb{1}_{A_3}(X) \mathbb{1}_{C_4}(X, Z) \right].\end{aligned}$$

Proceeding as for Proposition 16, using the integrals in Appendix D.4 and Lebesgue's dominated convergence theorem we can then show that for $\alpha = 1/3, \beta = m/3$ for $m \geq 1$ and $r \geq 0$

$$\lim_{d \rightarrow +\infty} d \operatorname{Var} (\phi_d(X, Z)) = \frac{2\ell^3}{3\sqrt{2\pi}}.$$

□

Proposition 18. *Take X a Laplace random variable and Z a standard normal random variable independent of X , then if $\sigma^2 = \ell^2 d^{-2/3}$ we have*

$$\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d(X, Z)^3] = 0.$$

Proof. Following the same structure of the previous propositions we have that

$$\begin{aligned} \mathbb{E} [\phi(X, Z)^3] &= \mathbb{E} [\phi_d^1(X, Z)^3 R_1(X, Z)] + \mathbb{E} [\phi_d^3(X, Z)^2 R_2(X, Z)] \\ &\quad + \mathbb{E} [\phi_d^3(X, Z)^3 R_3(X, Z)] + \mathbb{E} [\phi_d^4(X, Z)^3 R_4(X, Z)], \end{aligned}$$

exploiting again the symmetry of the laws of X and Z , using the integrals in Appendix D.4, the dominated convergence theorem we can then show that

$$\lim_{d \rightarrow +\infty} d\mathbb{E} [\phi_d(X, Z)^3] = 0.$$

□

D.2 Bound on Second Moment of Acceptance Ratio for the Laplace Distribution

Lemma 3. *Let Z be a standard normal random variable and $\sigma = \ell/d^\alpha$ for $\alpha = 1/3$. Then, there exists a constant $C > 0$ such that for all $a \in \mathbb{R}$ and $d \in \mathbb{N}$:*

$$\mathbb{E} [\phi_d(a, Z)^2] \leq \frac{C}{d^{2\alpha}}.$$

Proof. We consider the case $a \geq 0$ and $r \geq \sigma^{2(m-1)}$ only, all the other cases follow from identical arguments. As in the derivation of the moments of ϕ_d in Appendix D.1, we distinguish four regions. We recall that $\sigma = \ell/d^\alpha$ for $\alpha = 1/3$ and thus $\sigma^{p+1} \leq \sigma^p$ for all $p \in \mathbb{N}$. Take $r \geq \sigma^{-2(m-1)}$, for R_1 ,

we have, using Hölder's inequality multiple times,

$$\begin{aligned}
\mathbb{E} [\phi_d^1(a, Z)^2 \mathbb{1}_{R_1}(a, Z)] &= \mathbb{E} [\phi_d^1(a, Z)^2 \mathbb{1}_{B_1 \cup B_2}(a, Z)] \\
&\leq C\sigma^2 \int_{-(1-1/\sigma^{2(m-1)}r)a/\sigma}^{\sigma^{2m-1}r/2-(1-1/\sigma^{2(m-1)}r)a} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left[\left(\frac{1}{\sigma^{2m-1}r} a - z \right)^2 \right. \\
&\quad \left. + \left(\frac{z^2}{2\sigma} - \frac{1}{2\sigma^3} \left(\left(\frac{2}{\sigma^{2(m-1)}r} - \frac{1}{\sigma^{4(m-1)}r^2} \right) a - \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) \sigma z \right)^2 \right)^2 \right] dz \\
&+ C\sigma^2 \int_{-\sigma^{2m-1}r/2-(1-1/\sigma^{2(m-1)}r)a}^{-(1-1/\sigma^{2(m-1)}r)a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left[\left(\frac{2a}{\sigma} - \frac{1}{\sigma^{2m-1}r} a + z \right)^2 \right. \\
&\quad \left. + \left(\frac{z^2}{2\sigma} - \frac{1}{2\sigma^3} \left(\left(\frac{2}{\sigma^{2(m-1)}r} - \frac{1}{\sigma^{4(m-1)}r^2} \right) a - \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) \sigma z \right)^2 \right)^2 \right] dz \\
&\leq C\sigma^2 \int_{-\infty}^{+\infty} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left(4 \left(\frac{a}{\sigma} \right)^2 + 2 \left(\frac{1}{\sigma^{2m-1}r} a \right)^2 + 2z^2 \right) dz \\
&+ C\sigma^2 \int_{-\sigma^{2m-1}r/2-(1-1/\sigma^{2(m-1)}r)a}^{\sigma^{2m-1}r/2-(1-1/\sigma^{2(m-1)}r)a} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \\
&\quad \times \left(\frac{z^2}{2\sigma} - \frac{1}{2\sigma^3} \left(\left(\frac{2}{\sigma^{2(m-1)}r} - \frac{1}{\sigma^{4(m-1)}r^2} \right) a - \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) \sigma z \right)^2 \right)^2 dz \\
&\leq C\sigma^2 + C\sigma^2 \int_{-\sigma^{2m-1}r/2-(1-1/\sigma^{2(m-1)}r)a}^{\sigma^{2m-1}r/2-(1-1/\sigma^{2(m-1)}r)a} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \\
&\quad \times \left(\frac{z^4}{4\sigma^2} + \frac{1}{4\sigma^6} \left(\left(\frac{2}{\sigma^{2(m-1)}r} - \frac{1}{\sigma^{4(m-1)}r^2} \right)^4 a^4 + \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right)^4 \sigma^4 z^4 \right) \right) dz \\
&\leq C\sigma^2,
\end{aligned}$$

where we used the fact that the moments of Z are bounded and $a \leq \sigma^{2m}r/2$ for the first term, and the fact that $z \leq \sigma^{2m-1}r/2$ for the second one. Proceeding as above, for R_3 , $a > 0$ and

$r \geq \sigma^{-2(m-1)}$, we have

$$\begin{aligned}
\mathbb{E} [\phi_d^3(a, Z)^2 \mathbb{1}_{R_3}(a, Z)] &= \mathbb{E} [\phi_d^3(a, Z)^2 \mathbb{1}_{B_3 \cup B_4}(a, Z)] \\
&\leq C\sigma^2 \int_{\sigma^{2m-1}r/2 - (1-1/\sigma^{2(m-1)}r)a/\sigma}^{+\infty} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left[\left(\frac{1}{\sigma^{2m-1}r} a - z \right)^2 \right. \\
&\quad \left. + \left(\frac{z^2}{2\sigma} - \frac{1}{2\sigma^3} \left(\frac{1}{\sigma^{2(m-1)}r} a - \sigma z + \frac{\sigma^2}{2} \right)^2 \right)^2 \right] dz \\
&\quad + C\sigma^2 \int_{-\infty}^{-\sigma^{2m-1}r/2 - (1-1/\sigma^{2(m-1)}r)a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left[\left(\frac{2a}{\sigma} - \frac{1}{\sigma^{2m-1}r} a + z \right)^2 \right. \\
&\quad \left. + \left(\frac{z^2}{2\sigma} - \frac{1}{2\sigma^3} \left(\frac{1}{\sigma^{2(m-1)}r} a - \sigma z + \frac{\sigma^2}{2} \right)^2 \right)^2 \right] dz \\
&\leq C\sigma^2 + C\sigma^2 \int_{-\infty}^{+\infty} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left(\frac{1}{2\sigma^3} \left(\frac{a^2}{\sigma^{4(m-1)}r^2} + \frac{\sigma^4}{4} - \sigma^3 z + \frac{a}{\sigma^{2m-4}r} - \frac{2az}{\sigma^{2m-3}r} \right) \right)^2 dz \\
&\leq C\sigma^2,
\end{aligned}$$

where we used again the boundedness of the moments of Z , the fact that $a \leq \sigma^{2m}r/2$ and that $\sigma^{p+1} \leq \sigma^p$. For R_2 and $a > 0$, we have

$$\begin{aligned}
\mathbb{E} [\phi_d^2(a, Z)^2 \mathbb{1}_{R_2}(a, Z)] &= \mathbb{E} [\phi_d^2(a, Z)^2 \mathbb{1}_{C_1 \cup C_2}(a, Z)] \\
&\leq C\sigma^2 \int_{\sigma/2 - a/\sigma}^{\sigma/2 + \sigma^{2m-1}r/2 - a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left[\left(\frac{\sigma^2}{2} - \sigma z \right)^2 \right. \\
&\quad \left. + \left(\frac{z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)}r} a + \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) \left(\frac{\sigma^2}{2} - \sigma z \right) \right)^2 \right)^2 \right] dz \\
&\quad + C\sigma^2 \int_{\sigma/2 - \sigma^{2m-1}r/2 - a/\sigma}^{\sigma/2 - a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left[\left(2a - \frac{\sigma^2}{2} + \sigma z \right)^2 \right. \\
&\quad \left. + \left(\frac{z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)}r} a + \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) \left(\frac{\sigma^2}{2} - \sigma z \right) \right)^2 \right)^2 \right] dz.
\end{aligned}$$

The first integral is bounded using the moments of Z , while for the third one let us denote

$\chi(a, \sigma, z) := a - \sigma^2/2 + \sigma z$, then

$$\begin{aligned}
& \int_{\sigma/2 - \sigma^{2m-1}r/2 - a/\sigma}^{\sigma/2 + \sigma^{2m-1}r/2 - a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left(\frac{z^2}{2\sigma} - \frac{1}{2\sigma^3} \left(\frac{1}{\sigma^{2(m-1)}r} a + \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) \left(\frac{\sigma^2}{2} - \sigma z \right) \right)^2 \right)^2 dz \\
&= \int_{\sigma/2 - \sigma^{2m-1}r/2 - a/\sigma}^{\sigma/2 + \sigma^{2m-1}r/2 - a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left(\frac{z^2}{2\sigma} - \frac{1}{2\sigma^3} \left(\frac{\chi(a, \sigma, z)}{\sigma^{2(m-1)}r} + \frac{\sigma^2}{2} - \sigma z \right)^2 \right)^2 dz \\
&\leq C \int_{\sigma/2 - \sigma^{2m-1}r/2 - a/\sigma}^{\sigma/2 + \sigma^{2m-1}r/2 - a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left(\frac{z^2}{2\sigma} - \frac{1}{2\sigma^3} \left(\frac{\sigma^2}{2} - \sigma z \right)^2 \right)^2 dz \\
&+ C \int_{\sigma/2 - \sigma^{2m-1}r/2 - a/\sigma}^{\sigma/2 + \sigma^{2m-1}r/2 - a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left(\frac{\chi(a, \sigma, z)^2}{2\sigma^{4m-1}r^2} \right)^2 + \left(\frac{\chi(a, \sigma, z)}{r\sigma^{2m-1}} \left(\frac{\sigma^2}{2} - \sigma z \right) \right)^2 dz;
\end{aligned}$$

recalling that in R_2 we have $|\chi(a, \sigma, z)| \leq \sigma^{2m}r/2$, we obtain that this term is also bounded by $C\sigma^2$. For R_4 and $a > 0$, we have

$$\begin{aligned}
\mathbb{E} [\phi_d^4(a, Z)^2 \mathbb{1}_{R_4}(a, Z)] &= \mathbb{E} [\phi_d^4(a, Z)^2 \mathbb{1}_{C_4}(a, Z)] \\
&= \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r/2 - a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left(2a - \frac{\sigma^2}{2} + \sigma z \right)^2 dz \\
&= \sigma^2 \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r/2 - a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left(\frac{2a}{\sigma} - \frac{\sigma}{2} + z \right)^2 dz
\end{aligned}$$

Collecting all the terms together, we obtain

$$\begin{aligned}
\mathbb{E} [\phi_d(a, Z)^2] &= \sum_{i=1}^4 \mathbb{E} [\phi_d^i(a, Z)^2 \mathbb{1}_{T_i}(a, Z)] \\
&\leq C\sigma^2 + C\sigma^2 \int_{-\infty}^{\sigma/2 - a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left(\frac{2a}{\sigma} - \frac{\sigma}{2} + z \right)^2 dz.
\end{aligned}$$

Recall that $\sigma = \ell d^{-1/3}$. To bound the last integral we use Hölder's inequality

$$\begin{aligned}
\int_{-\infty}^{\sigma/2 - a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left(\frac{2a}{\sigma} - \frac{\sigma}{2} + z \right)^2 dz &\leq C \int_{-\infty}^{\sigma/2 - a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left[\left(\frac{\sigma}{2} + z \right)^2 + 4 \left(\frac{\sigma}{2} - \frac{a}{\sigma} \right)^2 \right] dz \\
&\leq C \int_{-\infty}^{\sigma/2 - a/\sigma} \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left[\left(\frac{\ell^2}{4} + z^2 \right) + 4 \left(\frac{\sigma}{2} - \frac{a}{\sigma} \right)^2 \right] dz.
\end{aligned}$$

The first term is bounded since the moments of Z are bounded. For the second term we use an estimate of the Gaussian cumulative distribution function. Let $\kappa(\ell, d, a) := \ell d^{-1/3}/2 - ad^{1/3}/\ell$. When $z < \kappa(\ell, d, a) < 0$, we have $1 < z/\kappa(\ell, d, a)$ and therefore

$$\begin{aligned}
(2\pi)^{-1/2} \kappa(\ell, d, a)^2 \int_{-\infty}^{\kappa(\ell, d, a)} e^{-z^2/2} dz &\leq (2\pi)^{-1/2} \kappa(\ell, d, a) \int_{-\infty}^{\kappa(\ell, d, a)} z e^{-z^2/2} dz, \\
&= (2\pi)^{-1/2} \kappa(\ell, d, a) \exp(-\kappa(\ell, d, a)^2/2).
\end{aligned}$$

However $y \mapsto ye^{-y^2/2}$ is bounded over $\mathbb{R}$, therefore $(a, d) \mapsto (2\pi)^{-1/2} \kappa(\ell, d, a) \exp(-\kappa(\ell, d, a)^2/2)$ is bounded over $\mathbb{R}_+^* \times \mathbb{N}$. If $\kappa(\ell, d, a) \geq 0$, then we still have $\kappa(\ell, d, a) < \ell$ and thus have the inequality

$$(2\pi)^{-1/2} \kappa(\ell, d, a)^2 \int_{-\infty}^{\kappa(\ell, d, a)} e^{-z^2/2} dz \leq (2\pi)^{-1/2} \ell^2 \int_{-\infty}^{+\infty} e^{-z^2/2} dz = \ell^2.$$

The result then follows since $\sigma = \ell d^{-1/3}$. □

D.3 Additional Integrals for the Laplace Distribution

We collect here two auxiliary Lemmata which are used in the proof of Proposition 3.

Lemma 4. *Take X a Laplace random variable and Z a standard normal random variable independent of X . Let $\tilde{X} := X - \frac{1}{\sigma^{2(m-1)}r} X \mathbb{1}\{|X| < \sigma^{2m}r/2\} - \frac{\sigma^2}{2} \text{sgn}(X) \mathbb{1}\{|X| \geq \sigma^{2m}r/2\} + \sigma Z$, then, for $\sigma = \ell d^{-\alpha}$ with $\alpha = 1/3$,*

$$\mathbb{E} \left[\mathbb{1} \left\{ \text{sgn}(X) \neq \text{sgn}(\tilde{X}) \right\} \right] \rightarrow 0$$

if $d \rightarrow \infty$.

Proof. Using the same strategy of Appendix D.1 and the symmetry of the laws of X, Z , we find that

$$\begin{aligned} \mathbb{E} \left[\mathbb{1} \left\{ \text{sgn}(X) \neq \text{sgn}(\tilde{X}) \right\} \right] &= 2\mathbb{E} [\mathbb{1}_{A_1}(X) \mathbb{1}_{B_2}(X, Z)] + 2\mathbb{E} [\mathbb{1}_{A_3}(X) \mathbb{1}_{C_2}(X, Z)] \\ &\quad + 2\mathbb{E} [\mathbb{1}_{A_1}(X) \mathbb{1}_{B_4}(X, Z)] + 2\mathbb{E} [\mathbb{1}_{A_3}(X) \mathbb{1}_{C_4}(X, Z)]. \end{aligned}$$

Using the same strategy used to obtain the moments of ϕ_d in Appendix D.1, we find that

$$\mathbb{E} [\mathbb{1}_{A_1}(X) \mathbb{1}_{B_2}(X, Z)] = o(1),$$

in addition

$$\begin{aligned} \mathbb{E} [\mathbb{1}_{A_3}(X) \mathbb{1}_{C_2}(X, Z)] &= \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r/2} e^{-z^2/2} \int_{\sigma^2/2 - \sigma z}^{\sigma^2/2 - \sigma z + \sigma^{2m}r/2} e^{-x} dx dz + o(1) \\ &= \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r/2} e^{-z^2/2} \left[\frac{\sigma^2 r}{2} \delta_{m1} + \dots \right] dz + o(1), \end{aligned}$$

where δ_{m1} is a Dirac's delta, and

$$\begin{aligned} &\mathbb{E} [\mathbb{1}_{A_1}(X) \mathbb{1}_{B_4}(X, Z)] + \mathbb{E} [\mathbb{1}_{A_3}(X) \mathbb{1}_{C_4}(X, Z)] \\ &= \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r} e^{-z^2/2} \int_0^{\sigma^2/2 - \sigma z - \sigma^{2m}r/2} e^{-x} dx dz + o(1) \\ &= \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r} e^{-z^2/2} [-\sigma z + \dots] dz + o(1). \end{aligned}$$

Since $\sigma = \ell d^{-1/3}$ and the remainder terms of the Taylor expansions are bounded, Lebesgue's dominated convergence theorem gives

$$\begin{aligned}\mathbb{E}[\mathbb{1}_{A_3}(X)\mathbb{1}_{C_2}(X, Z)] &\rightarrow 0, \\ \mathbb{E}[\mathbb{1}_{A_1}(X)\mathbb{1}_{B_4}(X, Z)] + \mathbb{E}[\mathbb{1}_{A_3}(X)\mathbb{1}_{C_4}(X, Z)] &\rightarrow 0\end{aligned}$$

as $d \rightarrow \infty$. □

Lemma 5. *Take X a Laplace random variable and Z a standard normal random variable independent of X . Then,*

$$d^\alpha \mathbb{E}[|Z| |\phi_d(X, Z)|] \rightarrow 0$$

for $\alpha = 1/3$.

Proof. Using Cauchy-Schwarz's inequality we have that

$$\mathbb{E}[|Z| |\phi_d(X, Z)|] \leq \mathbb{E}[Z^2]^{1/2} \mathbb{E}[\phi_d(X, Z)^2]^{1/2};$$

the first expectation is equal to one, and the second one converges to zero at rate $d^{1/2}$ by Proposition 17. The result follows straightforwardly. □

D.4 Integrals for Moment Computations

We distinguish the case $m = 1/2$ and $m \geq 1$ since the integration bounds significantly differ in these two cases. For values between $1/2$ and 1 the integrals are not finite. The expectations below are obtained by integrating w.r.t. x and using a Taylor expansion about $\sigma = 0$ to obtain the leading order terms. Using the Lagrange form of the remainder for the Taylor expansions, we find that the remainder terms are all of the form $\sigma^{1/\alpha+1} f(\gamma(\sigma, z))/(1/\alpha + 1)!$ where $\gamma(\sigma, z)$ is a point between the limits of integration w.r.t. x and $f : x \mapsto p(x)e^{-x}$, where p is a polynomial. Therefore, using the boundedness of the remainder and Lebesgue's dominated convergence theorem, the integrals w.r.t. z of the remainder terms all converge to 0.

D.4.1 First Moment

For simplicity, we only consider the case for $r \geq \sigma^{-2(m-1)}$, the other case follows analogously.

Region R_1 Let us consider ϕ_d^1 first. We have

$$\begin{aligned}A_1 \cap B_1 &= \begin{cases} 0 \leq x \leq \frac{\sigma^{2m}r}{2} & \text{if } 0 \leq z \leq \frac{\sigma}{2} \\ 0 \leq x \leq \left(\frac{\sigma^{2m}r}{2} - \sigma z\right) \left(1 - \frac{1}{\sigma^{2(m-1)}r}\right)^{-1} & \text{if } \frac{\sigma}{2} \leq z \leq \frac{\sigma^{2m-1}r}{2} \\ -\sigma z \left(1 - \frac{1}{\sigma^{2(m-1)}r}\right)^{-1} \leq x \leq \frac{\sigma^{2m}r}{2} & \text{if } \frac{\sigma}{2} - \frac{\sigma^{2m-1}r}{2} \leq z \leq 0 \end{cases}, \\ A_1 \cap B_2 &= \begin{cases} 0 \leq x \leq \frac{\sigma^{2m}r}{2} & \text{if } -\frac{\sigma^{2m-1}r}{2} \leq z \leq \frac{\sigma}{2} - \frac{\sigma^{2m-1}r}{2} \\ 0 \leq x \leq -\sigma z \left(1 - \frac{1}{\sigma^{2(m-1)}r}\right)^{-1} & \text{if } \frac{\sigma}{2} - \frac{\sigma^{2m-1}r}{2} \leq z \leq 0 \\ -\left(\frac{\sigma^{2m}r}{2} + \sigma z\right) \left(1 - \frac{1}{\sigma^{2(m-1)}r}\right)^{-1} \leq x \leq \frac{\sigma^{2m}r}{2} & \text{if } \frac{\sigma}{2} - \sigma^{2m-1}r \leq z \leq -\frac{\sigma^{2m-1}r}{2} \end{cases},\end{aligned}$$

and

$$A_1 \cap (B_1 \cup B_2) = \begin{cases} 0 \leq x \leq \frac{\sigma^{2m}r}{2} & \text{if } -\frac{\sigma^{2m-1}r}{2} \leq z \leq \frac{\sigma}{2} \\ 0 \leq x \leq \left(\frac{\sigma^{2m}r}{2} - \sigma z\right) \left(1 - \frac{1}{\sigma^{2(m-1)}r}\right)^{-1} & \text{if } \frac{\sigma}{2} \leq z \leq \frac{\sigma^{2m-1}r}{2} \\ -\left(\frac{\sigma^{2m}r}{2} + \sigma z\right) \left(1 - \frac{1}{\sigma^{2(m-1)}r}\right)^{-1} \leq x \leq \frac{\sigma^{2m}r}{2} & \\ \text{if } \frac{\sigma}{2} - \sigma^{2m-1}r \leq z \leq -\frac{\sigma^{2m-1}r}{2} \end{cases}.$$

The corresponding expectations are

$$\begin{aligned} & \mathbb{E} \left[\left(\frac{X}{\sigma^{2(m-1)}r} - \sigma Z \right) \mathbb{1}_{A_1}(X) \mathbb{1}_{B_1}(X, Z) \right] \\ &= \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2}^{\sigma^{2m-1}r/2} e^{-z^2/2} [z^{4-2m} \sigma^{4-2m} \xi(r) + \dots] dz \\ &+ \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r/2}^0 e^{-z^2/2} [z^{4-2m} \sigma^{4-2m} \xi(r) + \dots] dz \\ &+ o(1), \\ & \mathbb{E} \left[\left(2X - \frac{X}{\sigma^{2(m-1)}r} + \sigma Z \right) \mathbb{1}_{A_1}(X) \mathbb{1}_{B_2}(X, Z) \right] \\ &= \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r/2}^0 e^{-z^2/2} [z^{4-2m} \sigma^{4-2m} \xi(r) + \dots] dz \\ &+ \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r}^{-\sigma^{2m-1}r/2} e^{-z^2/2} [z^{4-2m} \sigma^{4-2m} \xi(r) + \dots] dz \\ &+ o(1), \end{aligned}$$

where $\xi : [0, +\infty) \rightarrow \mathbb{R}$ is a function of r only which might change from one line to the other, and

$$\begin{aligned} & \mathbb{E} \left[\left(\frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\left(\frac{2}{\sigma^{2(m-1)}r} - \frac{1}{\sigma^{4(m-1)}r^2} \right) X - \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) \sigma Z \right)^2 \right) \mathbb{1}_{A_1}(X) \mathbb{1}_{B_1 \cup B_2}(X, Z) \right] \\ &= \frac{1}{2\sqrt{2\pi}} \int_{-\sigma^{2m-1}r/2}^{\sigma/2} e^{-z^2/2} [+ \dots] dz \\ &+ \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2}^{\sigma^{2m-1}r/2} e^{-z^2/2} [+ \dots] dz \\ &+ \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r}^{-\sigma^{2m-1}r/2} e^{-z^2/2} [z^2 \sigma^2 \xi(r) + \dots] dz, \end{aligned}$$

where $\xi : [0, +\infty) \rightarrow \mathbb{R}$ is a function of r only which might change from one line to the other.

Region R_2 For ϕ_d^2 , we have

$$A_3 \cap C_1 = \begin{cases} \sigma^2/2 - \sigma z \leq x \leq -\sigma z + \sigma^2/2 + \sigma^{2m}r/2 & \text{if } z < \sigma/2 - \sigma^{2m-1}r/2 \\ \sigma^{2m}r/2 < x \leq -\sigma z + \sigma^2/2 + \sigma^{2m}r/2 & \text{if } \sigma/2 - \sigma^{2m-1}r/2 \leq z \leq \sigma/2 \end{cases},$$

$$A_3 \cap C_2 = \begin{cases} \sigma^2/2 - \sigma z - \sigma^{2m}r/2 \leq x \leq \sigma^2/2 - \sigma z & \text{if } z < \sigma/2 - \sigma^{2m-1}r \\ \sigma^{2m}r/2 < x \leq \sigma^2/2 - \sigma z & \text{if } \sigma/2 - \sigma^{2m-1}r < z < \sigma/2 - \sigma^{2m-1}r/2 \end{cases}$$

and

$$A_3 \cap (C_1 \cup C_2) = \begin{cases} \sigma^{2m}r/2 \leq x \leq \sigma^{2m}r/2 + \sigma^2/2 - \sigma z & \text{if } \sigma/2 - \sigma^{2m-1}r \leq z \leq \sigma/2 \\ -\sigma^{2m}r/2 + \sigma^2/2 - \sigma z < x \leq \sigma^{2m}r/2 + \sigma^2/2 - \sigma z & \\ \text{if } z < \sigma/2 - \sigma^{2m-1}r & \end{cases}.$$

The corresponding expectations are

$$\begin{aligned} \mathbb{E} \left[\left(\frac{\sigma^2}{2} - \sigma Z \right) \mathbb{1}_{A_3}(X) \mathbb{1}_{C_1}(X, Z) \right] &= \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r/2} e^{-z^2/2} \left[-\frac{rz}{2} \sigma^{2m+1} + \dots \right] dz \\ &\quad + \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r/2}^{\sigma/2} e^{-z^2/2} [z^2 \sigma^2 + \dots] dz, \\ \mathbb{E} \left[\left(2X - \frac{\sigma^2}{2} + \sigma Z \right) \mathbb{1}_{A_3}(X) \mathbb{1}_{C_2}(X, Z) \right] &= \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r} e^{-z^2/2} \left[-\frac{rz}{2} \sigma^{2m+1} + \dots \right] dz \\ &\quad + \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r}^{\sigma/2 - \sigma^{2m-1}r/2} e^{-z^2/2} \left[-\frac{rz}{2} \sigma^{2m+1} + \dots \right] dz, \end{aligned}$$

and

$$\begin{aligned} \mathbb{E} \left[\left(\frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)}r} X + \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right) \left(\frac{\sigma^2}{2} - \sigma Z \right) \right)^2 \right) \mathbb{1}_{A_3}(X) \mathbb{1}_{C_1 \cup C_2}(X, Z) \right] \\ = \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r} e^{-z^2/2} \left[\frac{rz}{2} \sigma^{2m+1} + \dots \right] dz \\ + \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r}^{\sigma/2} e^{-z^2/2} \left[-\frac{z^3}{2r} \sigma^{3-2m} + \dots \right] dz. \end{aligned}$$

Region R_3 For ϕ_d^3 , we have, in the case $r \geq \sigma^{-2(m-1)}$,

$$A_1 \cap B_3 = \begin{cases} 0 \leq x \leq \frac{\sigma^{2m}r}{2} & \text{if } z > \frac{\sigma^{2m-1}r}{2} \\ \left(\frac{\sigma^{2m}r}{2} - \sigma z \right) \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right)^{-1} < x \leq \frac{\sigma^{2m}r}{2} & \text{if } \frac{\sigma}{2} \leq z \leq \frac{\sigma^{2m-1}r}{2} \end{cases},$$

$$A_1 \cap B_4 = \begin{cases} 0 \leq x \leq \frac{\sigma^{2m}r}{2} & \text{if } z < \frac{\sigma}{2} - \sigma^{2m-1}r \\ 0 \leq x < \left(\frac{\sigma^{2m}r}{2} - \sigma z \right) \left(1 - \frac{1}{\sigma^{2(m-1)}r} \right)^{-1} & \text{if } \frac{\sigma}{2} - \sigma^{2m-1}r \leq z \leq -\frac{\sigma^{2m-1}r}{2} \end{cases}.$$

The corresponding expectations are

$$\begin{aligned}
& \mathbb{E} \left[\left(\frac{1}{\sigma^{2(m-1)}r} X - \sigma Z + \frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)}r} X - \sigma Z + \frac{\sigma^2}{2} \right)^2 \right) \mathbb{1}_{A_1}(X) \mathbb{1}_{B_3}(X, Z) \right] \\
&= \frac{1}{2\sqrt{2\pi}} \int_{\sigma^{2m-1}r/2}^{+\infty} e^{-z^2/2} \left[-\frac{rz}{8} \sigma^{2m+1} + \dots \right] dz \\
&+ \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2}^{\sigma^{2m-1}r/2} e^{-z^2/2} [z^3 - 2m \sigma^{3-2m} \xi(r) + \dots] dz, \\
& \mathbb{E} \left[\left(2X - \frac{1}{\sigma^{2(m-1)}r} X + \sigma Z + \frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)}r} X - \sigma Z - \frac{\sigma^2}{2} \right)^2 \right) \mathbb{1}_{A_1}(X) \mathbb{1}_{B_4}(X, Z) \right] \\
&= \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r/2} e^{-z^2/2} \left[\frac{3}{8} \sigma^{2m+1} + \dots \right] dz \\
&+ \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r/2}^{-\sigma^{2m-1}r/2} e^{-z^2/2} [z^3 \sigma^{3-2m} \xi(r) + \dots] dz,
\end{aligned}$$

where $\xi : [0, +\infty) \rightarrow \mathbb{R}$ is a function of r only which might change from one line to the other.

Region R_4 Finally, for ϕ_d^4 we have

$$A_3 \cap C_4 = \left\{ z < \frac{\sigma}{2} - \sigma^{2m-1}r, \frac{\sigma^{2m}r}{2} < x \leq \frac{\sigma^2}{2} - \sigma z - \frac{\sigma^{2m}r}{2} \right\},$$

and

$$\mathbb{E} \left[\left(2X - \frac{\sigma^2}{2} + \sigma Z \right) \mathbb{1}_{A_3}(X) \mathbb{1}_{C_4}(X, Z) \right] = \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r} e^{-z^2/2} \left[\frac{z^3}{6} \sigma^3 + \dots \right] dz.$$

D.4.2 Second Moment

For simplicity, we only consider the case for $r \geq \sigma^{-2(m-1)}$, the other case follows analogously.

Region R_1 For ϕ_d^1 we have

$$\begin{aligned}
\mathbb{E} [\phi_d^1(X, Z)^2 \mathbb{1}_{A_1}(X) \mathbb{1}_{B_1}(X, Z)] &= \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2}^{\sigma^{2m-1}r/2} e^{-z^2/2} [z^3 \sigma^3 \xi(r) + \dots] dz \\
&+ \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - r\sigma^{2m-1}/2}^0 e^{-z^2/2} [z^3 \sigma^3 \xi(r) + \dots] dz \\
&+ o(1), \\
\mathbb{E} [\phi_d^1(X, Z)^2 \mathbb{1}_{A_1}(X) \mathbb{1}_{B_2}(X, Z)] &= \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r/2}^0 e^{-z^2/2} [z^3 \sigma^3 \xi(r) + \dots] dz \\
&+ \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r}^{-\sigma^{2m-1}r/2} e^{-z^2/2} [z^3 \sigma^3 \xi(r) + \dots] dz \\
&+ o(1),
\end{aligned}$$

where $\xi : [0, +\infty) \rightarrow \mathbb{R}$ is a function of r only which might change from one line to the other.

Region R_2 For ϕ_d^2 we have

$$\begin{aligned}\mathbb{E} [\phi_d^2(X, Z)^2 \mathbb{1}_{A_3}(X) \mathbb{1}_{C_1}(X, Z)] &= \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r/2} e^{-z^2/2} \left[\frac{rz^2}{2} \sigma^{2m+2} + \dots \right] dz \\ &\quad + \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r/2}^{\sigma/2} e^{-z^2/2} [-z^3 \sigma^3 + \dots] dz, \\ \mathbb{E} [\phi_d^2(X, Z)^2 \mathbb{1}_{A_3}(X) \mathbb{1}_{C_2}(X, Z)] &= \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r} e^{-z^2/2} [z^2 \sigma^{2m+2} + \dots] dz \\ &\quad + \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r}^{\sigma/2 - \sigma^{2m-1}r/2} e^{-z^2/2} \left[-\frac{rz^2}{2} \sigma^{2m+2} + \dots \right] dz.\end{aligned}$$

Region R_3 For ϕ_d^3 we have

$$\begin{aligned}\mathbb{E} \left[\left(\frac{1}{\sigma^{2(m-1)}r} X - \sigma Z + \frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)}r} X - \sigma Z + \frac{\sigma^2}{2} \right)^2 \right)^2 \mathbb{1}_{A_1}(X) \mathbb{1}_{B_3}(X, Z) \right] \\ = \frac{1}{2\sqrt{2\pi}} \int_{\sigma^{2m-1}r/2}^{+\infty} e^{-z^2/2} \left[\frac{rz^2}{24} \sigma^{2m+2} + \dots \right] dz \\ + \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2}^{\sigma^{2m-1}r/2} e^{-z^2/2} \left[\frac{rz^2}{24} \sigma^{2m+2} + \dots \right] dz, \\ \mathbb{E} \left[\left(2X - \frac{1}{\sigma^{2(m-1)}r} X + \sigma Z + \frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)}r} X - \sigma Z - \frac{\sigma^2}{2} \right)^2 \right)^2 \mathbb{1}_{A_1}(X) \mathbb{1}_{B_4}(X, Z) \right] \\ = \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r/2} e^{-z^2/2} \left[\frac{7rz^2}{24} \sigma^{2m+2} + \dots \right] dz \\ + \frac{1}{2\sqrt{2\pi}} \int_{\sigma/2 - \sigma^{2m-1}r/2}^{-\sigma^{2m-1}r/2} e^{-z^2/2} \left[\frac{7rz^2}{24} \sigma^{2m+2} + \dots \right] dz,\end{aligned}$$

Region R_4 For ϕ_d^4 we have

$$\mathbb{E} \left[\left(2X - \frac{\sigma^2}{2} + \sigma Z \right)^2 \mathbb{1}_{A_3}(X) \mathbb{1}_{C_4}(X, Z) \right] = \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r} e^{-z^2/2} \left[-\frac{z^3}{3} \sigma^3 + \dots \right] dz.$$

D.4.3 Third Moment

Having established that the only possible scaling is given by $\alpha = 1/3$, $\beta = m/3$ with $m \geq 1$, we now proceed to bound the third moment of ϕ_d in this case. For simplicity, we only consider the case for $r \geq 1$, the other case follows analogously.

Since $m \geq 1$, we find that $\mathbb{E} [\phi_d^1(X, Z)^3 \mathbb{1}_{R_1}(X, Z)] = o(1)$ as $d \rightarrow \infty$ since the limits of integration all converge to 0. Then, using Hölder's inequality for ϕ_d^2 , we have

$$\begin{aligned}
\mathbb{E} [\phi_d^2(X, Z)^3] &\leq C \mathbb{E} \left[\left(\frac{\sigma^2}{2} - \sigma Z \right)^3 \mathbb{1}_{A_3}(X) \mathbb{1}_{C_1}(X, Z) \right] \\
&\quad + C \mathbb{E} \left[\left(2X - \frac{\sigma^2}{2} + \sigma Z \right)^3 \mathbb{1}_{A_3}(X) \mathbb{1}_{C_2}(X, Z) \right] \\
&\quad + C \mathbb{E} \left[\left(\frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)r}} X + \left(1 - \frac{1}{\sigma^{2(m-1)r}} \right) \left(\frac{\sigma^2}{2} - \sigma Z \right) \right)^2 \right)^3 \right. \\
&\quad \quad \left. \times \mathbb{1}_{A_3}(X) \mathbb{1}_{C_1 \cup C_2}(X, Z) \right] \\
&= \frac{C}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r/2} e^{-z^2/2} \left[-\frac{rz^3}{2} \sigma^5 + \dots \right] dz \\
&\quad + \frac{C}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r} e^{-z^2/2} \left[-\frac{rz^3}{2} \sigma^5 + \dots \right] dz \\
&\quad + \frac{C}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r} e^{-z^2/2} [z^3 \sigma^5 \xi(r) + \dots] dz + o(1),
\end{aligned}$$

where $\xi : [0, +\infty) \rightarrow \mathbb{R}$ is a function of r only which might change from one line to the other. For

ϕ_d^3 , we have, using again Hölder's inequality,

$$\begin{aligned}
& \mathbb{E} \left[\left(\frac{1}{\sigma^{2(m-1)r}} X - \sigma Z + \frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)r}} X - \sigma Z + \frac{\sigma^2}{2} \right)^2 \right)^3 \mathbb{1}_{A_1}(X) \mathbb{1}_{B_3}(X, Z) \right] \\
& \leq C \mathbb{E} \left[\left(\frac{1}{\sigma^{2(m-1)r}} X - \sigma Z \right)^3 \mathbb{1}_{A_1}(X) \mathbb{1}_{B_3}(X, Z) \right] \\
& + C \mathbb{E} \left[\left(\frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)r}} X - \sigma Z + \frac{\sigma^2}{2} \right)^2 \right)^3 \mathbb{1}_{A_1}(X) \mathbb{1}_{B_3}(X, Z) \right] \\
& = \frac{1}{2\sqrt{2\pi}} \int_{\sigma^{2m-1}r/2}^{+\infty} e^{-z^2/2} \left[-\frac{z^3 r}{2} \sigma^5 + \dots \right] dz \\
& + \frac{1}{2\sqrt{2\pi}} \int_{\sigma^{2m-1}r/2}^{+\infty} e^{-z^2/2} [z^3 \sigma^5 \xi(r) + \dots] dz + o(1) \\
& \mathbb{E} \left[\left(2X - \frac{1}{\sigma^{2(m-1)r}} X + \sigma Z + \frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)r}} X - \sigma Z - \frac{\sigma^2}{2} \right)^2 \right)^3 \mathbb{1}_{A_1}(X) \mathbb{1}_{B_4}(X, Z) \right] \\
& \leq C \mathbb{E} \left[\left(2X - \frac{1}{\sigma^{2(m-1)r}} X + \sigma Z \right)^3 \mathbb{1}_{A_1}(X) \mathbb{1}_{B_4}(X, Z) \right] \\
& + C \mathbb{E} \left[\left(\frac{Z^2}{2} - \frac{1}{2\sigma^2} \left(\frac{1}{\sigma^{2(m-1)r}} X - \sigma Z - \frac{\sigma^2}{2} \right)^2 \right)^3 \mathbb{1}_{A_1}(X) \mathbb{1}_{B_4}(X, Z) \right] \\
& = \frac{1}{2\sqrt{2\pi}} \int_{\sigma^{2m-1}r/2}^{+\infty} e^{-z^2/2} \left[\frac{z^3 r}{2} \sigma^5 + \dots \right] dz \\
& + \frac{1}{2\sqrt{2\pi}} \int_{\sigma^{2m-1}r/2}^{+\infty} e^{-z^2/2} [z^3 \sigma^5 \xi(r) + \dots] dz + o(1),
\end{aligned}$$

where $\xi : [0, +\infty) \rightarrow \mathbb{R}$ is a function of r only which might change from one line to the other. Finally, for ϕ_d^4 we have

$$\mathbb{E} \left[\left(2X - \frac{\sigma^2}{2} + \sigma Z \right)^3 \mathbb{1}_{A_3}(X) \mathbb{1}_{C_4}(X, Z) \right] = \frac{1}{2\sqrt{2\pi}} \int_{-\infty}^{\sigma/2 - \sigma^{2m-1}r} e^{-z^2/2} \left[\frac{z^3}{10} \sigma^5 + \dots \right] dz.$$