A RESPONSIBILITY VALUE FOR DIGRAPHS

ROSA VAN DEN ENDE AND DYLAN LAPLACE MERMOUD

ABSTRACT. There is an increasing need to hold players responsible for negative or positive impact that take place elsewhere in a value chain or a network. For example, countries or companies are held more and more responsible for their indirect carbon emissions. We introduce a *responsibility value* that allocates the total impact of the value chain among the players, taking into account their direct impact and their indirect impact through the underlying graph. Moreover, we show that the responsibility value satisfies a set of natural yet important properties.

1. INTRODUCTION

We see that there is an increasing urge to account for environmental or social impact throughout a supply chain or network: through the choices players in such network make, for example by choosing to buy or not buy from a certain supplier, they carry some responsibility for not only their own impacts, but also for the impacts that take place upstream of their own economic activities. Examples of impacts for which indirect responsibility could be assigned to players include but are not limited to the emission of greenhouse gases, the underpayment of employees, the disposal of hazardous waste, but possibly also positive impacts such as the transition to renewable energy. Assigning this responsibility and consequently holding these players liable for such impacts, can incentivise the increase of positive impacts or the decrease of negative impacts.

However, as of now, it is not clear how to assign responsibility for indirect impacts to players in a given network. In the case where the considered impacts are the emissions of greenhouse gases (GHG), there is consensus that players carry some responsibility for their indirect emissions. This could be either the energy required to run their processes, known as scope 2 emissions, or for all the other emissions the sources of which were not controlled by the player: the scope 3 emissions [3]. However, to which extent these players are responsible for their scope 2 and 3 emissions, remains unclear. Moreover, in this context, the same unit of emission could be claimed twice; when the emission falls within one scope for one player and under another scope for another player. The double counting of the same reduction efforts

Date: January 10, 2023.

²⁰²⁰ Mathematics Subject Classification. MSC Primary 91B32; Secondary 05C20.

This work has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 956107, "Economic Policy in Complex Environments (EPOC)".

that might subsequently occur is a major policy concern when trying to achieve the emission reduction goals [1].

We thus aim to clarify and quantify the indirect responsibility for impacts in a general manner. Based on the impact of every player and the underlying graph of the players, we introduce a responsibility value that assigns to every player a responsibility for the social or environmental impact of itself and the nodes it benefits from. We show that this value satisfies certain properties, that resemble the axioms that Shapley used in cooperative game theory to define the Shapley value [?]. In section 3 we offer a method to approximate a solution. Subsequently, we discuss the interpretation of the value of the discount factor that plays an important role on the emphasis that is put on the indirect versus direct responsibility.

2. The responsibility value

Let N be a set of finite cardinality n, called the set of players. Let A be a $(n \times n)$ matrix, with all its entries being real numbers between 0 and 1 and with the sum of entries of each row being equal to 1, i.e., $\sum_{j \in N} A_{i,j} = 1$. We denote by \mathcal{R} the set of such matrices. Let G_A be the weighted, directed graph the node set of which is N and whose adjacency matrix is A. If A is symmetric, then G_A is an undirected graph.

In the following, we interpret each edge (i, j), with *i* the source node and *j* the terminal node, as an interaction between players *i* and *j*, with *j* benefiting from actions of player *i*, and consequentially taking partial responsibility for this player. The reader may think of a link (i, j) as a trade between a seller *i* and a buyer *j*, and $A_{i,j}$ being the proportion of purchases of *j* among the total volume of sales of *i*. Note that there is a flow going through the directed graph: a responsibility flow.

We aim to allocate to every player a share of the total responsibility of the grand coalition N in a way that a few basic properties are satisfied:

- (1) the responsibility of a player for several time periods should be the sum of its responsibilities of each single time period,
- (2) two players that are connected in the same way to identical players should be assigned the same amount of indirect responsibility,
- (3) a player that does not benefit from any other player, that is, a player that has no incoming edges, should not carry any indirect responsibility,
- (4) no responsibility flow between two players in G_A should be ignored, even if this flow consists of a considerable amount of edges. Then, we want to introduce some form of discounting in order to assign more responsibility to a terminal node for source nodes of a short flow than for source nodes of a long flow,

(5) the responsibility for the impact of the entire group should be completely shared among the members of the group. We want to ensure that the responsibility for every unit of impact is assigned to some player while simultaneously preventing double counting.

Then, let $\iota : N \to \mathbb{R}$ be a map that associates to every node of G_A a real value, which can be interpreted as a quantity of impact for which we want to assign responsibility. Let (0, 1) denote the set of real values strictly included between 0 and 1. For each $\gamma \in (0, 1)$, we define a map $\rho^{\gamma} : \mathcal{R} \to \mathbb{R}^n$ associating to each matrix $A \in \mathcal{R}$ an *n*-dimensional vector ρ^{γ} , called the *responsibility value* of (A, ι) . For all $i \in N$, the coordinate $\rho_i^{\gamma}(A, \iota)$ corresponds to the total impact for which player *i* should be held responsible, and is defined by

$$\rho_i^{\gamma}(A,\iota) = \gamma \sum_{k \ge 0} (1-\gamma)^k \left(\sum_{p \in N} A_{p,i}^k \iota(p) \right).$$

As A is an adjacency matrix, $A_{p,i}^k$ (the entry of A^k at (p,i)) counts the number of walks from node p to node i of length k. For a given length k, we take into account all the nodes of the graph that are connected to player i by summing over all the players. Then, we sum over all the possible lengths k, with a discount factor weighting the distance, that decreases the transfer of responsibility. So, for k = 0, one obtains the direct responsibility, while $k \ge 1$ yields the indirect responsibility. We can interpret the factor γ as a discount factor, the value of which we discuss in more detail in section 4.

In the following, we will show that the responsibility value satisfies the desired properties. Denote by **0** the *n*-dimensional row vector $(0, \ldots, 0)$ and denote by **1** the *n*-dimensional row vector $(1, \ldots, 1)$. Also, denote by $r_i(A)$ the *i*-th row of A and by $c_j(A)$ the *j*-th column of A. We write $\iota(N)$ for $\sum_{p \in N} \iota(p)$.

Lemma 1. For all $A \in \mathcal{R}$, for all $k \in \mathbb{N}$ and for all $i, j \in N$, we have

$$A_{i,j}^k \leq 1, \qquad \sum_{j \in N} A_{i,j}^k = 1, \quad and \quad \sum_{p \in N} A_{p,i}^k \iota(p) \leq \iota(N).$$

Proof. We prove the two first statements by induction on k. For k = 0, this is true because A^0 is the identity matrix. Then, assume that there exists $k \in \mathbb{N}$ such that, for all $i, j \in N$, we have $A_{i,j}^k \leq 1$ and $\sum_{j \in N} A_{i,j}^k = 1$. Then,

$$A_{i,j}^{k+1} = \left(A^k \cdot A\right)_{i,j} = \left\langle r_i\left(A^k\right), c_j\left(A\right)\right\rangle \le \left\langle r_i\left(A^k\right), \mathbf{1}^\top\right\rangle = 1.$$

Moreover, we have

$$\sum_{j \in N} A_{i,j}^{k+1} = \sum_{j \in N} \left(\sum_{p \in N} A_{i,p}^k A_{p,j} \right) = \sum_{p \in N} \left(A_{i,p}^k \left(\sum_{j \in N} A_{p,j} \right) \right) = \sum_{p \in N} A_{i,p}^k = 1.$$

The last statement directly follows from the first one.

Proposition 1 (Convergence). Let $\gamma \in (0,1)$, let $A \in \mathcal{R}$ and let $\iota : N \to \mathbb{R}$. For all $i \in N$, we have $\rho_i^{\gamma}(A, \iota) < \infty$.

Proof. Let $\gamma \in (0, 1)$. By Lemma 1, for any $i, p \in N$, we have $\sum_{p \in N} A_{p,i}^k \iota(p) \leq \iota(N)$. It leads, for any $A \in \mathcal{R}$ and any map $\iota : N \to \mathbb{R}$, to

$$\rho_i^{\gamma}(A,\iota) \le \gamma \sum_{k\ge 0} (1-\gamma)^k \iota(N).$$

Because $\iota(N)$ does not depend on k, we can put it out of the sum, and let γ and $\sum_{k>0} (1-\gamma)^k$ cancel each other. Then,

$$\rho_i^{\gamma}(A,\iota) \le \iota(N) < \infty.$$

This result shows that even though we take into account all responsibility flows as desired and therefore obtain a sum over all possible lengths k that is finite, the discounting ensures that ρ^{γ} always has finite coordinates.

Proposition 2 (Additivity). Let $\gamma \in (0,1)$ and let $A \in \mathcal{R}$. For all $i \in N$ and for all pairs of maps ι_1 and ι_2 , we have

$$\rho_i^{\gamma}(A,\iota_1) + \rho_i^{\gamma}(1,\iota_2) = \rho_i^{\gamma}(A,\iota_1+\iota_2).$$

Proof. Let $\gamma \in (0, 1)$. For any player $i \in N$, we have

$$\rho_{i}^{\gamma}(A, \iota_{1} + \iota_{2}) = \gamma \sum_{k \ge 0} (1 - \gamma)^{k} \left(\sum_{p \in N} A_{p,i}^{k} \left(\iota_{1} + \iota_{2} \right) (p) \right)$$

$$= \gamma \sum_{k \ge 0} (1 - \gamma)^{k} \left(\sum_{p \in N} \left[A_{p,i}^{k} \iota_{1}(p) + A_{p,i}^{k} \iota_{2}(p) \right] \right)$$

$$= \gamma \sum_{k \ge 0} (1 - \gamma)^{k} \sum_{p \in N} A_{p,i}^{k} \iota_{1}(p) + \gamma \sum_{k \ge 0} (1 - \gamma)^{k} \sum_{p \in N} A_{p,i}^{k} \iota_{2}(p) .$$

Thus, the responsibility of a player for several time periods is indeed the sum of its responsibilities of each single time period, as desired.

Recall that **0** denotes the *n*-dimensional row vector $(0, \ldots, 0)$.

Definition 1. We say that $i \in N$ is an independent player if we have $c_i(A) = \mathbf{0}^{\top}$. **Lemma 2.** For all independent players $i \in N$, and for all $k \in \mathbb{N} \setminus \{0\}$, we have, $c_i(A^k) = \mathbf{0}^{\top}$. *Proof.* We prove it by induction on k. For k = 1, this is true by definition. Then, assume that there exists $k \in \mathbb{N} \setminus \{0\}$ such that $r_i(A^k) = \mathbf{0}$ and $c_i(A^k) = \mathbf{0}^{\top}$. Then, for any $p \in N$, we have,

$$A_{p,i}^{k+1} = \langle r_p(A), c_i(A^k) \rangle = \langle r_p(A), \mathbf{0}^\top \rangle = 0,$$

+1) = $\mathbf{0}^\top.$

then $c_i(A^{k+1}) = \mathbf{0}^\top$.

Proposition 3 (Independent player property). Let $\gamma \in (0,1)$, let $A \in \mathcal{R}$ and let $\iota : N \to \mathbb{R}$. We have

$$\rho_i^{\gamma}(A,\iota) = \gamma\iota(i).$$

Proof. Let $\gamma \in (0, 1)$ and let $i \in N$ be an independent player. By Lemma 2, for any $p \in N$ and any $k \in \mathbb{N} \setminus \{0\}$, we have $A_{p,i}^k = 0$. Then the sum $\sum_{p \in N} A_{p,i}^k \iota(p)$ equals 0 for $k \in \mathbb{N} \setminus \{0\}$, so

$$\rho_i^{\gamma}(A,\iota) = \gamma \sum_{k=0} (1-\gamma)^k \left(\sum_{p \in N} A_{p,i}^k \iota(p) \right) = \gamma \left(\sum_{p \in N} A_{p,i}^0 \iota(p) \right).$$

 A^0 being the identity matrix, we have $\sum_{p \in N} A^0_{p,i} \iota(p) = A^0_{i,i} \iota(i)$, thus

$$\rho_i^{\gamma}(A,\iota) = \gamma\iota(i).$$

We remark that the responsibility value holds the players responsible for a fraction γ of their direct impact. Then, for player $i \in N$, we call the quantity $\gamma \iota(i)$ the direct responsibility. Subsequently, to determine the total responsibility, we add the responsibility of the indirect impacts. Independent players, i.e. players that do not benefit from any other player, do not carry any indirect responsibility and are thus just assigned a fraction γ of their direct impacts.

Definition 2. We say that two players $i, j \in N$ are symmetric w.r.t. G_A if

 $r_i(A) = r_i(A)$ and $c_i(A) = c_i(A)$.

Lemma 3. For two symmetric players $i, j \in N$, we have, for any $k \in \mathbb{N}$,

$$r_i(A^k) = r_j(A^k)$$
 and $c_i(A^k) = c_j(A^k)$.

Proof. We prove it by induction on k. For k = 0, this is true because A^0 is the identity matrix. Then, assume that there exists $k \in \mathbb{N}$ such that $r_i(A^k) = r_j(A^k)$ and $c_i(A^k) = c_j(A^k)$. Then, for any $p \in N$,

$$A_{i,p}^{k+1} = \langle r_i(A^k), c_p(A) \rangle = \langle r_j(A^k), c_p(A) \rangle = A_{j,p}^{k+1},$$

then $r_i(A^{k+1}) = r_j(A^{k+1})$. With similar calculations on the columns, we have

$$A_{p,i}^{k+1} = \langle r_p(A), c_i(A^k) \rangle = \langle r_p(A), c_j(A^k) \rangle = A_{p,j}^{k+1},$$

) = c_i(A^{k+1})

then $c_i(A^{k+1}) = c_j(A^{k+1}).$

Proposition 4 (Symmetry). Let $\gamma \in (0, 1)$, let $A \in \mathcal{R}$ and let $\iota : N \to \mathbb{R}$. For all pairs of symmetric players *i* and *j*, we have

$$\rho_i^{\gamma}(A,\iota) - \gamma\iota(i) = \rho_j^{\gamma}(A,\iota) - \gamma\iota(j).$$

Proof. Let $\gamma \in (0, 1)$ and let $i, j \in N$ be two symmetric players. Then,

$$\rho_i^{\gamma}(A,\iota) - \gamma\iota(i) = \gamma \sum_{k \ge 1} (1-\gamma)^k \left(\sum_{p \in N} A_{p,i}^k \iota(p) \right).$$

Players i and j are symmetric, by Lemma 3 we have $A_{p,i}^k = A_{p,j}^k$, therefore

$$\rho_i^{\gamma}(A,\iota) - \gamma\iota(i) = \gamma \sum_{k \ge 1} (1-\gamma)^k \left(\sum_{p \in N} A_{p,j}^k \iota(p) \right) = \rho_j^{\gamma}(A,\iota) - \gamma\iota(j).$$

Since for player $i \in N$ the assigned direct responsibility is given by $\gamma \iota(i)$, as we saw in Prop. 3, it follows that $\rho_i^{\gamma}(A, \iota) - \gamma \iota(i)$ can be interpreted as the indirect responsibility. Symmetry then implies the notion of fairness that two players that are connected in the same way to identical players are assigned the same amount of indirect responsibility.

Proposition 5 (Efficiency). Let $\gamma \in (0, 1)$, let $A \in \mathcal{R}$ and let $\iota : N \to \mathbb{R}$. We have

$$\sum_{i\in N}\rho_i^\gamma(A,\iota)=\iota(N)$$

Proof. Let $\gamma \in (0, 1)$, let $A \in \mathcal{R}$ and let $\iota : N \to \mathbb{R}$. For all $i \in N$, we have

$$\sum_{i \in N} \rho_i^{\gamma}(A, \iota) = \sum_{i \in N} \gamma \sum_{k \ge 0} (1 - \gamma)^k \left(\sum_{p \in N} A_{p,i}^k \iota(p) \right)$$
$$= \gamma \sum_{k \ge 0} (1 - \gamma)^k \left(\sum_{p \in N} \left[\sum_{i \in N} A_{p,i}^k \right] \iota(p) \right)$$

By Lemma 1, we have that $\sum_{i \in N} A_{p,i}^k = 1$, and then

$$\sum_{i \in N} \rho_i^{\gamma}(A, \iota) = \gamma \sum_{k \ge 0} (1 - \gamma)^k \iota(N) = \iota(N).$$

Efficiency ensures that all the units of emissions are assigned to some player and prevents double counting.

To summarize, besides always taking on finite coordinates, our responsibility value ρ^{γ} always satisfies additivity, symmetry, the independent player property and efficiency. Note that these properties resemble the axioms that Shapley used to define its Shapley value [?] in cooperative game theory.

A RESPONSIBILITY VALUE FOR DIGRAPHS

3. Approximate solutions and numerical computation

3.1. ε -approximations. With the responsibility value, we aim to provide a practical tool. In the sequel, we show that, despite being an infinite sum, a precise approximation for the responsibility value can be reached with just a few computations. Let $\gamma \in (0, 1)$, let $A \in \mathcal{R}$, let $\iota : N \to \mathbb{R}$ and let $i \in N$. Denote by $\rho^{\gamma}(A, \iota)_{|q}$ the truncated responsibility value, defined, for all $i \in N$, by

$$\rho_i^{\gamma}(A,\iota)_{|q} = \gamma \sum_{k=0}^q (1-\gamma)^k \left(\sum_{p \in N} A_{p,i}^k \iota(p)\right).$$

The value $\rho^{\gamma}(A, \iota)_{|q}$ requires only a limited amount of calculations to be computed, mainly q-1 matrix multiplications. Given $\varepsilon > 0$, we define an ε -approximation as any $r \in \mathbb{R}^n$ such that $\|\rho^{\gamma}(A, \iota) - r\|_{\infty} \leq \varepsilon$, with $\|\cdot\|_{\infty}$ the L^{∞} -norm of \mathbb{R}^n . We denote by q_{ε} the smallest integer such that $\rho^{\gamma}(A, \iota)_{|q_{\varepsilon}}$ is an ε -approximation, i.e.,

$$q_{\varepsilon} \coloneqq \min \left\{ q \in \mathbb{N} \mid \forall i \in N, |\rho_i^{\gamma}(A, \iota) - \rho_i^{\gamma}(A, \iota)|_q | \le \varepsilon \right\}.$$

By Lemma 1, for all $i \in N$ we have $\sum_{p \in N} A_{p,i}^k \iota(p) \leq \iota(N)$, and then

$$\|\rho^{\gamma}(A,\iota) - \rho^{\gamma}(A,\iota)|_{q}\|_{\infty} \leq \gamma \iota(N) \sum_{k \geq q+1} (1-\gamma)^{k}.$$

Using the formulae of geometric series and partial geometric series, we have

$$\sum_{k>q} (1-\gamma)^k = \sum_{k\ge 0} (1-\gamma)^k - \sum_{k=0}^q (1-\gamma)^k = \frac{1}{\gamma} - \frac{1-(1-\gamma)^{q+1}}{\gamma} = \frac{(1-\gamma)^{q+1}}{\gamma},$$

and then $\|\rho^{\gamma}(A,\iota) - \rho^{\gamma}(A,\iota)|_{q}\|_{\infty} \leq \iota(N)(1-\gamma)^{q+1}$. Therefore, q_{ε} is determined by

$$(1-\gamma)^{q_{\varepsilon}} \ge \iota(N)\varepsilon \ge (1-\gamma)^{q_{\varepsilon}+1}.$$

Using a computer, it is fairly easy to find q_{ε} , and then to compute an ε -approximation $r := \rho^{\gamma}(A, \iota)_{|q_{\varepsilon}}$. By minimality of q_{ε} , r is the ε -approximation most efficiently computed, in $q_{\varepsilon} + 1$ steps requiring $q_{\varepsilon} - 1$ matrix multiplications.

3.2. Numerical computation. For convenience, we denote $\vec{\iota}$ the *n*-dimensional vector defined, for all $i \in N$, by $\vec{\iota}_i = \iota(i)$. The formula for the responsibility value becomes $\rho_i^{\gamma}(A, \iota) = \gamma \sum_{k\geq 0} (1-\gamma)^k \langle c_i(A^k), \vec{\iota} \rangle$. Denote by $P_{\gamma}^q(A)$ the matrix derived from A defined by $P_{\gamma}^q(A) \coloneqq \sum_{k=0}^q ((1-\gamma)A)^k$. This new matrix only requires q-1 matrix multiplications to be performed. Rearranging some terms, we find a highly efficient formula to calculate approximations of the responsibility value for all the players *simultaneously*:

$$\rho^{\gamma}(A,\iota)_{|q} = \gamma \,\vec{\iota} \cdot P_{\gamma}^{\,q}(A).$$



FIGURE 1. Adjacency matrix A and its corresponding digraph G_A

3.3. **Example.** Let $\gamma = 0.6$, let $N = \{1, 2, 3, 4\}$, let $\iota(i) = i$, for all $i \in N$, and let A be the adjacency matrix defined in Figure 1.

Notice that, in this case, players 1 and 3 are symmetric, and player 2 is an independent player. By Prop. 3, we have that

$$\rho_2^{0.6}(A,\iota) = 0.6 \cdot \iota(2) = 1.2.$$

It means that player 2, as part of the whole group N, is responsible for an impact of 1.2 among the global impact of $\iota(N) = 10$.

Let $\varepsilon := 10^{-3}$. To obtain an ε -approximation of $\rho^{0.6}(A, \iota)$, we have to find q_{ε} that is the first integer to be smaller than or equal to $\iota(N) \cdot \varepsilon = 0.01$, i.e.,

$$(0.4)^{q_{\varepsilon}} \ge 0.01 \ge (0.4)^{q_{\varepsilon}+1},$$

that is

We find $q_e = 5$. Then, we need to compute the matrix $P_{0.6}^5(A)$, which is approximately

$$P_{0.6}^{5}(A) \simeq \begin{pmatrix} 1.129 & 0 & 0.129 & 0.401 \\ 0.160 & 1 & 0.160 & 0.341 \\ 0.129 & 0 & 1.129 & 0.401 \\ 0.251 & 0 & 0.251 & 1.158 \end{pmatrix}$$

We can now finish the calculations:

$$\rho^{0.6}(A,\iota) \simeq \begin{pmatrix} 0.6 & 1.2 & 1.8 & 2.4 \end{pmatrix} \cdot \begin{pmatrix} 1.129 & 0 & 0.129 & 0.401 \\ 0.160 & 1 & 0.160 & 0.341 \\ 0.129 & 0 & 1.129 & 0.401 \\ 0.251 & 0 & 0.251 & 1.158 \end{pmatrix}$$
$$= \begin{pmatrix} 1.704 & 1.2 & 2.904 & 4.151 \end{pmatrix}.$$

We indeed obtained the right value for player 2. We see that player 2 and player 3 carry less responsibility than their direct impact, while player 1 and player 4 are held responsible for less than their direct impacts.

$$\rho_1^{0.6}(A,\iota) - \gamma\iota(1) \simeq 1.704 - 0.6 = 1.104,$$

$$\rho_3^{0.6}(A,\iota) - \gamma\iota(3) \simeq 2.904 - 1.8 = 1.104,$$

satisfying the symmetry property.

4. The parameter γ

The parameter $\gamma \in (0,1)$ plays an important role in the responsibility measure and determines how much emphasis is put on direct responsibility with respect to indirect responsibility. Recall that for player $i \in N$ with direct impact $\iota(i)$, the direct responsibility a player has is given by $\gamma \iota(i)$, to which we subsequently add the indirect responsibility. To get some intuition for the parameter γ , consider the following example in Figure 2, that consists of $N = \{1, 2\}, A'_{1,2} = A'_{2,2} = 1$ and $A'_{1,1} = A'_{2,1} = 0$. Moreover, the impacts are given by $\iota(1) = 1$ and $\iota(2)$. In words, player 1 sells all of her output to player 2, who in return consumes all her output herself. Since $\iota(2)$, the activities of player 2 do not result in any impact. Then, let us evaluate the assigned responsibilities. For player 1, we obtain

$$\rho_1^{\gamma}(A',\iota) = \gamma\iota(1) = \gamma,$$

while for player 2, we find

$$\rho_2^{\gamma}(A',\iota) = (1-\gamma)\iota(1) + \iota(2) = (1-\gamma)$$



FIGURE 2. Adjacency matrix A' and its corresponding digraph $G_{A'}$

Observe that when $\gamma < \frac{1}{2}$, player 2 carries more responsibility for the impact of player 1 than player 1 itself. In fact, even though its own impact is zero, player 2 has a greater total responsibility than the player 1 who does carry direct responsibility. On the one hand, if one believes that a player should be responsible for its own actions, it could therefore be argued that a value of $\gamma < \frac{1}{2}$ is unreasonable. On the

other hand, one could argue that $\gamma < \frac{1}{2}$ is legitimate since it reflects the power or leverage a player has. In the current example, player 2 is the only player to whom player 1 supplies, making player 2 the reason why player 1 has the impacts. If player 2 were to cease buying from player 1, player 1 would, in the given situation, no longer have any incentive to keep producing the impacts. Thus, the power that player 2 has, is reflected in this high responsibility value. These are moral or political questions which are not easy to find a universal answer to. In our model, one can adapt the value of γ in order to be in line with the point of view on these dilemmas one chooses to take.

5. Conclusions

As a response to the increasing need to assign responsibility to a player for impacts that take place elsewhere in the value chain, whether such impacts are negative impacts like the emission of greenhouse gases, the underpayment of employees or positive impacts such as the transition to renewable energies, we developed a method to assign direct as well as indirect responsibility to players, based on the underlying graph and the direct impacts of every player. Since this responsibility value is shown to satisfy a set of desirable properties, it can be applied in many situations and on many different scales, as players can be interpreted as either countries, firms or individuals. For example, if we consider the impacts to be carbon emissions, the responsible value could play a major role in achieving the emission reduction efforts.

References

- Schneider, L., Kollmuss, A., & Lazarus, M. (2015). Addressing the risk of double counting emission reductions under the UNFCCC. Climatic Change, 131(4), 473-486.
- Shapley, L. S. (2016). 17. A value for n-person games. In Contributions to the Theory of Games (AM-28), Volume II (pp. 307-318). Princeton University Press.
- [3] World Resources Institute and World Business Council on Sustainable Development (2004). The greenhouse gas protocol. World Resources Institute, Washington, D.C., DC, Mar.

Centre d'Économie de la Sorbonne, Université Paris I Panthéon-Sorbonne, 106-112 Boulevard de l'Hôpital, 75013, Paris, France & University of Bielefeld, Universitätsstrasse 25, 33615 Bielefeld, Germany

Email address: rosa.ende@gmail.com

Centre d'Économie de la Sorbonne, Université Paris I Panthéon-Sorbonne, 106-112 Boulevard de l'Hôpital, 75013, Paris, France

Email address: dylan.laplace.mermoud@gmail.com